June 2024 Volume 31

Editors:

Kaye Wilson, Doris Chong, & Ayeshah Khan email: enquiry@pharmac.govt.nz
Telephone +64 4 460 4990

Level 9, 40 Mercer Street PO Box 10 254 Wellington

Freephone Information Line 0800 66 00 50 (9am – 5pm weekdays)

Circulation

You can register to have an electronic version of the Pharmaceutical Schedule (link to PDF copy) emailed to your nominated email address each month by subscribing at pharmac.govt.nz/subscribe.

Production

Typeset automatically from XML and T_EX. XML version of the Schedule available from schedule.pharmac.govt.nz/pub/schedule

Programmers

Anrik Drenth

email: texschedule@pharmac.govt.nz @Pharmaceutical Management Agency



ISSN 1179-3686

This work is licensed under the Creative Commons Attribution 4.0 International licence. In essence, you are free to copy, distribute and adapt it, as long as you attribute the work to Pharmac and abide by the other licence terms. To view a copy of this licence, visit: creativecommons.org/licenses/by/4.0/. Attribution to Pharmac should be in written from and not by reproduction of the Pharmac logo. While care has been taken in compiling this Schedule, Pharmac takes no responsibility for any errors or omissions, and shall not be liable for any consequences arising there from.

		_
Section A	General Rules	5
Section B	Alimentary Tract & Metabolism	6
	Blood & Blood Forming Organs	36
	Cardiovascular System	46
	Dermatologicals	68
	Genito Urinary System	77
	Hormone Preparations – Systemic	84
	Infections – Agents For Systemic Use	95
	Musculoskeletal System	116
	Nervous System	122
	Oncology Agents & Immunosuppressants	152
	Respiratory System & Allergies	248
	Sensory Organs	259
	Various	264
Section C	Extemporaneous Compounds (ECPs)	267
Section D	Special Foods	270
Section I	National Immunisation Schedule	293
	Index	307

Introducing Pharmac

Introducing Pharmac

The Pharmaceutical Management Agency (Pharmac) makes decisions that help control Government spending on pharmaceuticals. This includes community pharmaceuticals, hospital pharmaceuticals, vaccines and increasingly, hospital medical devices. Pharmac negotiates prices, sets subsidy levels and conditions, and makes decisions on changes to the subsidised list.

Pharmac's role:

"to secure for eligible people in need of pharmaceuticals, the best health outcomes that are reasonably achievable from pharmaceutical treatment and from within the amount of funding provided."

Pae Ora (Healthy Futures) Act 2022

To ensure our decisions are as fair and robust as possible we use a decision-making process that incorporates clinical, economic and commercial issues. We also seek the views of users and the wider community through consultation. The processes we generally use are outlined in our Operating Policies and Procedures.

Further information about Pharmac and the way we make funding decisions can be found on the Pharmac website at https://pharmac.govt.nz/about.

Purpose of the Pharmaceutical Schedule

The purpose of the Schedule is to list:

- the Community Pharmaceuticals that are subsidised by the Government and to show the amount of the subsidy paid to contractors, as well as the manufacturer's price and any access conditions that may apply;
- the Hospital Pharmaceuticals that may be used in Health NZ Hospitals, as well as any access conditions that may apply; and
- the Pharmaceuticals, including Medical Devices, used in Health NZ Hospitals for which national prices have been negotiated by Pharmac.

The Schedule does not show the final cost to Government of subsidising each Community Pharmaceutical or to Health NZ Hospitals in purchasing each Pharmaceutical, since that will depend on any rebate and other arrangements Pharmac has with the supplier and, for Pharmaceuticals used in Health NZ Hospitals, on any logistics arrangements put in place.

This book contains sections A to D and Section I of the Pharmaceutical Schedule and lists the Pharmaceuticals funded for use in the community, including vaccines, as well as haemophilia and cancer treatments given in Health NZ Hospitals. Section H lists the Pharmaceuticals that that can be used in Health NZ Hospitals and is a separate publication.


The Pharmaceuticals in this book are listed by therapeutic group, which is based on the WHO Anatomical Therapeutic Chemical (ATC) system. The listings are displayed alphabetically under each heading.

The index lists both chemical entities and product brand names.

Explaining pharmaceutical entries

The Pharmaceutical Schedule lists pharmaceuticals subsidised by the Government, the subsidy, the supplier's price and the access conditions that may apply.

Example



Glossary

Units of Measure

gramg kilogramkg international unitiu	mi mi mi
Abbreviations	
AmpouleAmp	Ge
CapsuleCap	Gr
Cream	Inf
DeviceDev	Ini
DispersibleDisp	Lic
EffervescentEff	Lo
EmulsionEmul	Oi
Enteric Coated EC	Sa

microgrammilligrammillilitre	mg
Gelatinous	
Granules	
Infusion	Inf
Injection	Inj
Liquid	Liq
Long Acting	LA
Ointment	Oint
Sachet	Sach

millimoleunit	
Solution	Supp Tab
Trans Dermal Delivery System	TDDS

Read the General Rules: https://pharmac.govt.nz/section-a.

SECTION B: ALIMENTARY TRACT AND METABOLISM

	0.1.11			
	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
Antacids and Antiflatulents				
Antacids and Reflux Barrier Agents				
ALGINIC ACID Sodium alginate 225 mg and magnesium alginate 87.5 mg po		30	√ (Saviscon Infant
SODIUM ALGINATE * Tab 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg - peppermint flavour	1.80 (14.39)	60	Ó	Gaviscon Extra Strength
* Oral liq 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg per 10 ml		500 m		Acidex
Phosphate Binding Agents				
ALUMINIUM HYDROXIDE * Tab 600 mg CALCIUM CARBONATE	12.56	100	✓ µ	llu-Tab
Oral liq 1,250 mg per 5 ml (500 mg elemental per 5 ml) – Subsidy by endorsement		500 m 173 m		Roxane Calcium carbonate PAI 829
Only when prescribed for patients unable to swallow calc inappropriate and the prescription is endorsed according		s or v	vhere calciu	m carbonate tablets are
Antidiarrhoeals				
Agents Which Reduce Motility				
LOPERAMIDE HYDROCHLORIDE – Up to 30 cap available on a * Tab 2 mg * Cap 2 mg	10.75	400 400	-	lodia Diamide Relief
Rectal and Colonic Anti-inflammatories				
BUDESONIDE Cap modified-release 3 mg - Special Authority see SA1886 below - Retail pharmacy		90 alid fo	_	Budesonide Te Arai

Both:

- 1 Mild to moderate ileal, ileocaecal or proximal Crohn's disease; and
- 2 Any of the following:
 - 2.1 Diabetes; or

Subsidy	Ful	y Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per •	Manufacturer	

continued...

- 2.2 Cushingoid habitus; or
- 2.3 Osteoporosis where there is significant risk of fracture; or
- 2.4 Severe acne following treatment with conventional corticosteroid therapy; or
- 2.5 History of severe psychiatric problems associated with corticosteroid treatment; or
- 2.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
- 2.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated).

Initial application — (collagenous and lymphocytic colitis (microscopic colitis)) from any relevant practitioner. Approvals valid for 6 months where patient has a diagnosis of microscopic colitis (collagenous or lymphocytic colitis) by colonoscopy with biopsies.

Initial application — **(gut Graft versus Host disease)** from any relevant practitioner. Approvals valid for 6 months where patient has a gut Graft versus Host disease following allogenic bone marrow transplantation*.

Note: Indication marked with * is an unapproved indication.

Initial application — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has autoimmune hepatitis*: and
- 2 Patient does not have cirrhosis; and
- 3 Any of the following:
 - 3.1 Diabetes: or
 - 3.2 Cushingoid habitus; or
 - 3.3 Osteoporosis where there is significant risk of fracture: or
 - 3.4 Severe acne following treatment with conventional corticosteroid therapy; or
 - 3.5 History of severe psychiatric problems associated with corticosteroid treatment; or
 - 3.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
 - 3.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated); or3.8 Adolescents with poor linear growth (where conventional corticosteroid use may limit further growth).

Note: Indication marked with * is an unapproved indication.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

HYDROCORTISONE ACETATE

Rectal foam 10%, CFC-Free (14 applications)26.55	15 g OP	Colifoam
HYDROCORTISONE ACETATE WITH PRAMOXINE HYDROCHLORIDE		
Topical aerosol foam, 1% with pramoxine hydrochloride 1%26.55	10 g OP	✓ Proctofoam S29
MESALAZINE		
Tab 400 mg49.50	100	✓ Asacol
Tab long-acting 500 mg56.10	100	✓ Pentasa
Tab 800 mg85.50	90	✓ Asacol
		✓ Asacol S29 S29
Modified release granules, 1 g118.10	100 OP	✓ Pentasa
Enema 1 g per 100 ml41.30	7	✓ Pentasa
Suppos 500 mg22.80	20	✓ Asacol
Suppos 1 g50.96	28	✓ Pentasa

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Generic
OLSALAZINE				
Tab 500 mg	56.02	60	✓	Atnahs
				Olsalazine S29
	93.37	100	1	Dipentum
Cap 250 mg	53.00	100	1	Dipentum
PREDNISOLONE SODIUM				
Rectal foam 20 mg per dose (14 applications)	74.10	1 OP	•	Essential
				Prednisolone S29
SODIUM CROMOGLICATE				
Cap 100 mg	113.35	100	1	Ralicrom
SULFASALAZINE				
* Tab 500 mg	16.52	100	1	Salazopyrin
* Tab EC 500 mg	17.86	100		Salazopyrin EN

Local preparations for Anal and Rectal Disorders

Antihaemorrhoidal Preparations

FLUOCORTOLONE CAPROATE WITH FLUOCORTOLONE PIVALAT	E AND CING	CHOCAINE	
Oint 950 mcg, with fluocortolone pivalate 920 mcg, and			
cinchocaine hydrochloride 5 mg per g	.13.05	30 g OP	Ultraproct
Suppos 630 mcg, with fluocortolone pivalate 610 mcg, and			
cinchocaine hydrochloride 1 mg	8.61	12	Ultraproct
HYDROCORTISONE WITH CINCHOCAINE			
Oint 5 mg with cinchocaine hydrochloride 5 mg per g	.15.00	30 g OP	Proctosedyl
Suppos 5 mg with cinchocaine hydrochloride 5 mg per g		12	✓ Proctosedyl

Management of Anal Fissures

GLYCERYL TRINITRATE − Special Authority see SA1329 below − Retail pharmacy

★ Oint 0.2%......22.00 30 g OP

✓ Rectogesic

⇒SA1329 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has a chronic anal fissure that has persisted for longer than three weeks.

Antispasmodics and Other Agents Altering Gut Motility

GLYCOPYRRONIUM BROMIDE		
Inj 200 mcg per ml, 1 ml ampoule - Up to 10 inj available on a	_	
PSO19.00	5	✓ Robinul
HYOSCINE BUTYLBROMIDE		
* Tab 10 mg6.35	100	✓ Buscopan
* Inj 20 mg, 1 ml - Up to 5 inj available on a PSO1.91	5	✓ Spazmol
MEBEVERINE HYDROCHLORIDE		
* Tab 135 mg	90	✓ Colofac

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic S Per ✓ Manufacturer

Antiulcerants

Antisecretory and Cytoprotective

MISOPROSTOL - Wastage claimable

Helicobacter Pylori Eradication

CLARITHROMYCIN

Tab 500 mg − Subsidy by endorsement.......14.58 14 ✓ Klacid

- a) Maximum of 28 tab per prescription
- Subsidised only if prescribed for helicobacter pylori eradication and prescription is endorsed accordingly.
 Note: the prescription is considered endorsed if clarithromycin is prescribed in conjunction with a proton pump inhibitor and either amoxicillin or metronidazole.

H2 Antagonists

FA	MOTIDINE - Only on a prescription			
*	Tab 20 mg	4.91	100	✓ Famotidine
	-			Hovid S29
*	Tab 40 mg	10.32	100	✓ Famotidine
	-			Hovid S29
*	Inj 10 mg per ml, 4 ml - Subsidy by endorsement	CBS	10	✓ Mylan S29
Subsidy by endorsement – Subsidised for patients receiving treatment as part of palliative care.			e care.	

Proton Pump Inhibitors

LAI	NSOPRAZOLE		
*	Cap 15 mg4.20	100	✓ Lanzol Relief
*	Cap 30 mg	100	✓ Lanzol Relief
ON	EPRAZOLE		
	For omeprazole suspension refer Standard Formulae, page 267		
*	Cap 10 mg	90	✓ Omeprazole actavis 10
*	Cap 20 mg	90	✓ Omeprazole actavis 20
*	Cap 40 mg	90	✓ Omeprazole actavis 40
*	Powder – Only in combination42.50 Only in extemporaneously compounded omeprazole suspension.	5 g	✓ Midwest
*	Inj 40 mg ampoule with diluent37.38	5	✓ <u>Dr Reddy's</u> <u>Omeprazole</u>
			✓ Ocicure S29
(00	cicure S29 Inj 40 mg ampoule with diluent to be delisted 1 August 2024)		
PA	NTOPRAZOLE		
*	Tab EC 20 mg1.99	90	✓ Panzop Relief
	Tab EC 40 mg	90	✓ Panzop Relief

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
Site Protective Agents				
COLLOIDAL BISMUTH SUBCITRATE Tab 120 mg	14.51	50	•	Gastrodenol S29
SUCRALFATE Tab 1 g	35.50 (48.28)	120		Carafate
Bile and Liver Therapy				
RIFAXIMIN - Special Authority see SA1461 below - Re Tab 550 mg		56	1	Xifaxan
■ SA1461 Special Authority for Subsidy Initial application only from a gastroenterologist, hepat hepatologist. Approvals valid for 6 months where the patolerated doses of lactulose. Renewal only from a gastroenterologist, hepatologist or hepatologist. Approvals valid without further renewal ur benefiting from treatment.	atient has hepatic encephalop Practitioner on the recommen	athy d	lespite an	adequate trial of maximum roenterologist or
Diabetes				
Hyperglycaemic Agents				
DIAZOXIDE — Special Authority see SA1320 below – R Cap 25 mg Cap 100 mg Oral liq 50 mg per ml	110.00	100 100 0 ml 0	✓ OP ✓	Proglicem \$29 Proglicem \$29 Proglycem \$29
(Proglycem S29 Oral liq 50 mg per ml to be delisted 1 messand	ovals valid for 12 months whe		d for the tr	
Renewal from any relevant practitioner. Approvals validappropriate and the patient is benefiting from treatment. GLUCAGON HYDROCHLORIDE Inj 1 mg syringe kit – Up to 5 kit available on a PSC		1		Glucagen Hypokit
Insulin - Short-acting Preparations	J			and agon Trypomic
INSULIN NEUTRAL Inj human 100 u per ml	25.26 1	0 ml C		Actrapid Humulin R
▲ Inj human 100 u per ml, 3 ml	42.66	5	1	Actrapid Penfill Humulin R
Insulin - Intermediate-acting Preparation	s			
INSULIN ASPART WITH INSULIN ASPART PROTAMI	NE			

▲ Inj 100 iu per ml, 3 ml prefilled pen......52.15

✓ NovoMix 30 FlexPen

5

	Subsidy		Fully Brand or
	(Manufacturer's Pr \$	Per	idised Generic Manufacturer
INSULIN ISOPHANE	· · · · · · · · · · · · · · · · · · ·		
▲ Inj human 100 u per ml	17.68	10 ml OP	✓ Humulin NPH
			✓ Protaphane
▲ Inj human 100 u per ml, 3 ml	29.86	5	✓ Humulin NPH
			Protaphane Penfill
INSULIN ISOPHANE WITH INSULIN NEUTRAL			•
▲ Inj human with neutral insulin 100 u per ml	25.26	10 ml OP	✓ Humulin 30/70
▲ Inj human with neutral insulin 100 u per ml, 3 ml	12.66	5	✓ Mixtard 30 ✓ Humulin 30/70
Inj numan with neutral insulin 100 triper hill, 5 hill	42.00	J	✓ PenMix 30
			✓ PenMix 50
INSULIN LISPRO WITH INSULIN LISPRO PROTAMINE			
▲ Inj lispro 25% with insulin lispro protamine 75% 100 u per ml,			
3 ml	42.66	5	✓ Humalog Mix 25
lacktriangle Inj lispro 50% with insulin lispro protamine 50% 100 u per ml,			_
3 ml	42.66	5	Humalog Mix 50
Insulin - Long-acting Preparations			
insum - Long-acting Freparations			
INSULIN GLARGINE			_
▲ Inj 100 u per ml, 10 ml		1	✓ Lantus
▲ Inj 100 u per ml, 3 ml		5 5	✓ Lantus✓ Lantus SoloStar
a inj 100 u per mi, 5 mi disposable pen	94.50	J	Lantus SoloStai
Insulin - Rapid Acting Preparations			
INSULIN ASPART			
▲ Inj 100 u per ml, 10 ml		1	✓ NovoRapid
Inj 100 u per ml, 3 ml		5	✓ NovoRapid Penfill
▲ Inj 100 u per ml, 3 ml syringe	31.19	5	✓ NovoRapid FlexPen
INSULIN GLULISINE Inj 100 u per ml, 10 ml	27.02	1	√ Anidro
▲ Inj 100 u per ml, 3 ml		5	✓ Apidra✓ Apidra
▲ Inj 100 u per ml, 3 ml disposable pen		5	✓ Apidra SoloStar
INSULIN LISPRO			•
▲ Inj 100 u per ml, 10 ml	34.92	10 ml OP	✓ Humalog
▲ Inj 100 u per ml, 3 ml	59.52	5	✓ Humalog
Alpha Glucosidase Inhibitors			
ACARBOSE			
* Tab 50 mg	8.95	90	✓ Accarb
* Tab 100 mg		90	✓ <u>Accarb</u>
Oral Hypoglycaemic Agents			
GLIBENCLAMIDE			
* Tab 5 mg	7.50	100	✓ Daonil
GLICLAZIDE			
* Tab 80 mg	20.10	500	✓ Glizide
GLIPIZIDE			
* Tab 5 mg	4.58	100	✓ Minidiab
• • • • • • • • • • • • • • • • • • •			

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
METFORMIN HYDROCHLORIDE * Tab immediate-release 500 mg * Tab immediate-release 850 mg		1,000 500		Metformin Viatris Metformin Viatris
PIOGLITAZONE * Tab 15 mg * Tab 30 mg * Tab 45 mg	7.30	90 90 90	✓	<u>Vexazone</u> <u>Vexazone</u> <u>Vexazone</u>
VILDAGLIPTIN Tab 50 mg		60	1	Galvus
VILDAGLIPTIN WITH METFORMIN HYDROCHLORIDE Tab 50 mg with 1,000 mg metformin hydrochloride Tab 50 mg with 850 mg metformin hydrochloride		60 60	_	Galvumet Galvumet

GLP-1 Agonists

DULAGLUTIDE - Special Authority see SA2338 below - Retail pharmacy

Note: Not to be given in combination with a funded SGLT-2 inhibitor or other GLP-1 agonist.

Inj 1.5mg per 0.5 ml prefilled pen115.23 4 **✓ Trulicity**

⇒SA2338 Special Authority for Subsidy

Note: Subsidy for patients with existing approvals prior to 1 May 2024. Approvals valid without further renewal unless notified. No new patients will be granted from 1 May 2024 until further notice.

LIRAGLUTIDE - Special Authority see SA2339 below - Retail pharmacy

- a) Maximum of 9 inj per prescription
 - a) Not to be given in combination with a funded SGLT-2 inhibitor or other GLP-1 agonist.
 - b) Maximum of 1 pack of 3 (6 mg per ml, 3 ml) prefilled pens will be funded per month.

⇒SA2339 Special Authority for Subsidy

Note: Subsidy for patients with existing approvals prior to 1 May 2024. Approvals valid without further renewal unless notified. No new patients will be granted from 1 May 2024 until further notice.

SGLT2 Inhibitors

⇒SA2068 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Fither:

b)

- 1 Patient has previously received an initial approval for a GLP-1 agonist; or
- 2 All of the following:
 - 2.1 Patient has type 2 diabetes; and
 - 2.2 Any of the following:
 - 2.2.1 Patient is Māori or any Pacific ethnicity*; or
 - 2.2.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)*; or
 - 2.2.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator*; or
 - 2.2.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood

(Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer		Subsidy	Fully	Brand or
\$ Per ✔ Manufacturer	(Manufacturer's Price)		Generic
		\$	Per 🗸	Manufacturer

continued...

or as a young adult*; or

- 2.2.5 Patient has diabetic kidney disease (see note b)*; and
- 2.3 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of at least one blood-glucose lowering agent (e.g. metformin, vildagliptin, or insulin) for at least 3 months.

Notes: * Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

- a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack, ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.
- b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m2 in the presence of diabetes, without alternative cause.

EMPAGLIFLOZIN - Special Authority see SA2068 on the previous page - Retail pharmacy

Note: Not to be given in combination with a funded GLP-1 agonist.

*	Tab 10 mg	58.56	30	Jardiance
*	Tab 25 mg	58.56	30	 Jardiance

EMPAGLIFLOZIN WITH METFORMIN HYDROCHLORIDE - Special Authority see SA2068 on the previous page - Retail pharmacy

Note: Not to be given in combination with a funded GLP-1 agonist.

*	Tab 5 mg with 1,000 mg metformin hydrochloride58.56	60	Jardiamet
*	Tab 5 mg with 500 mg metformin hydrochloride58.56	60	Jardiamet
*	Tab 12.5 mg with 1,000 mg metformin hydrochloride58.56	60	Jardiamet
*	Tab 12.5 mg with 500 mg metformin hydrochloride58.56	60	Jardiamet

Diabetes Management

Ketone Testing

BLOOD KETONE DIAGNOSTIC TEST STRIP - Subsidy by endorsement

- a) Not on a BSO
- b) Maximum of 20 strip per prescription
- c) Up to 10 strip available on a PSO
- d) Patient has any of the following:
 - 1) type 1 diabetes; or
 - 2) permanent neonatal diabetes: or
 - 3) undergone a pancreatectomy; or
 - 4) cystic fibrosis-related diabetes; or
 - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer

Dual Blood Glucose and Blood Ketone Testing

DUAL BLOOD GLUCOSE AND BLOOD KETONE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A dual blood glucose and blood ketone diagnostic test meter is subsidised for a patient who has:
 - 1) type 1 diabetes; or
 - 2) permanent neonatal diabetes: or
 - 3) undergone a pancreatectomy; or
 - 4) cystic fibrosis-related diabetes; or
 - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly. Only 1 meter per patient will be subsidised (no repeat prescriptions). For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Meter with 50 lancets, a lancing device and 10 blood glucose

✓ CareSens Dual 1 OP

Blood Glucose Testing

BLOOD GLUCOSE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A diagnostic blood glucose test meter is subsidised for a patient who:
 - 1) is receiving insulin or sulphonylurea therapy; or
 - 2) is pregnant with diabetes; or
 - 3) is on home TPN at risk of hypoglycaemia or hyperglycaemia; or
 - 4) has a genetic or an acquired disorder of glucose homeostasis, excluding type 1 or type 2 diabetes and metabolic syndrome.

The prescription must be endorsed accordingly. Only one CareSens meter per patient will be subsidised (no repeat prescriptions). Patients already using the CareSens N POP meter and CareSens N meter are not eligible for a new meter, unless they have:

- 1) type 1 diabetes; or
- 2) permanent neonatal diabetes; or
- 3) undergone a pancreatectomy; or
- 4) cystic fibrosis-related diabetes.

For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Meter with 50 lancets, a lancing device and 10 diagnostic test

1 OP ✓ CareSens N ✓ CareSens N POP 20.00 ✓ CareSens N Premier

Note: Only 1 meter available per PSO

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	
\$	Per 🗸	Manufacturer

BLOOD GLUCOSE DIAGNOSTIC TEST STRIP - Up to 50 test available on a PSO

The number of test strips available on a prescription is restricted to 50 unless:

- Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the
 prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed; or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly; or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

Test strips	50 test OP	CareSens N
		✓ CareSens PRO

BLOOD GLUCOSE TEST STRIPS (VISUALLY IMPAIRED)

The number of test strips available on a prescription is restricted to 50 unless:

- Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the
 prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed: or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly; or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

Blood glucose test strips3	33.69	50 test OP	•	SensoCard
----------------------------	-------	------------	---	-----------

Insulin Syringes and Needles

Subsidy is available for disposable insulin syringes, needles, and pen needles if prescribed on the same form as the one used for the supply of insulin or liraglutide or when prescribed for a patient and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of insulin or liraglutide.

INSULIN PEN NEEDLES - Maximum of 200 dev per prescription

*	29 g × 12.7 mm10.95	100	✓ B-D Micro-Fine
	31 g × 5 mm12.26	100	✓ B-D Micro-Fine
	31 g × 6 mm9.50	100	✓ Berpu
	31 g × 8 mm	100	✓ B-D Micro-Fine
	32 g x 4 mm 10.95	100	✓ B-D Micro-Fine

		Subsidy		Fully	Brand or
		(Manufacturer's Price)		Subsidised	
_		\$	Per		Manufacturer
INS	SULIN SYRINGES, DISPOSABLE WITH ATTACHED NEEDLE	- Maximum of 200	dev p	oer prescrip	otion
*	Syringe 0.3 ml with 29 g x 12.7 mm needle	13.56	100	√	B-D Ultra Fine
		1.36	10		
		(1.99)			B-D Ultra Fine
*	Syringe 0.3 ml with 31 g × 8 mm needle	13.56	100	✓	B-D Ultra Fine II
		1.30	10		
		(1.99)			B-D Ultra Fine II
*	Syringe 0.5 ml with 29 g x 12.7 mm needle	13.56	100	✓	B-D Ultra Fine
		1.36	10		
		(1.99)			B-D Ultra Fine
*	Syringe 0.5 ml with 31 g × 8 mm needle	13.56	100	✓	B-D Ultra Fine II
		1.36	10		
		(1.99)			B-D Ultra Fine II
*	Syringe 1 ml with 29 g x 12.7 mm needle	13.56	100	✓	B-D Ultra Fine
		1.36	10		
		(1.99)			B-D Ultra Fine
*	Syringe 1 ml with 31 g × 8 mm needle	13.56	100	✓	B-D Ultra Fine II
		1.36	10		
		(1.99)			B-D Ultra Fine II

Insulin Pumps

INSULIN PUMP - Special Authority see SA1603 below - Retail pharmacy

- a) Maximum of 1 dev per prescription
- b) Only on a prescription
- c) Maximum of 1 insulin pump per patient each four year period.

Min basal rate 0.025 U/h		1	✓ MiniMed 770G
Min basal rate 0.1 U/h	4,500.00	1	✓ Tandem t:slim
	·		X2 with Basal-IQ

⇒SA1603 Special Authority for Subsidy

Initial application — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has permanent neonatal diabetes; and
- 2 A MDI regimen trial is inappropriate; and
- 3 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 4 Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 5 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 6 Either:
 - 6.1 Applicant is a relevant specialist; or
 - 6.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- 2 Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician; and
- 3 It has been at least 4 years since the last insulin pump received by the patient or, in the case of patients qualifying under previous pump therapy for the initial application; the pump is due for replacement; and

Subsidy (Manufacturer's Price)	, ,		Brand or Generic	
	Per	•	Manufacturer	

continued...

- 4 Either:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person); and
- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol: and
- 7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 8 Fither:
 - 8.1 Applicant is a relevant specialist; or
 - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events; and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline; and
- 3 Either:
 - 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 3.2 The pump is due for replacement; and
- 4 Fither:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1: and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol; and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Either:
 - 9.1 Applicant is a relevant specialist; or

Subsidy		Fully	Brand or
(Manufacturer's Price)	9	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol: and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline; and
- 3 Fither:
 - 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 3.2 The pump is due for replacement; and
- 4 Either:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump treatment; and
- 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating pump therapy; and
- 4 The patient is continuing to derive benefit from pump therapy; and
- 5 The patient had achieved and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and
- 6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and
- 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and
- 8 Either:
 - 8.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 8.2 The pump is due for replacement; and
- 9 Either:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and
- 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from the time of commencing pump treatment; and
- 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and
- 4 Either:
 - 4.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 4.2 The pump is due for replacement; and
- 5 Either:
 - 5.1 Applicant is a relevant specialist; or
 - 5.2 Applicant is a nurse practitioner working within their vocational scope.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

Insulin Pump Consumables

⇒SA1985 Special Authority for Subsidy

Initial application — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has permanent neonatal diabetes; and
- 2 A MDI regimen trial is inappropriate; and
- 3 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 4 Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 5 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 6 Fither:
 - 6.1 Applicant is a relevant specialist; or
 - 6.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- 2 Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician; and
- 3 Either:
 - 3.1 Applicant is a relevant specialist; or
 - 3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person); and
- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol; and
- 7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 8 Fither:
 - 8.1 Applicant is a relevant specialist: or
 - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — **(severe unexplained hypoglycaemia)** only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events; and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline; and
- 3 Fither:
 - 3.1 Applicant is a relevant specialist; or

Subsidy		Fully	Brand or
(Manufacturer's Price)	9	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1: and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol; and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Either:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol; and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline; and
- 3 Either:
 - 3.1 Applicant is a relevant specialist; or
 - 3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (**Previous use before 1 September 2012**) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump treatment; and
- 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating pump therapy; and
- 4 The patient is continuing to derive benefit from pump therapy; and
- 5 The patient had achieved and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and
- 6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and
- 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and
- 8 Either:
 - 8.1 Applicant is a relevant specialist; or
 - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less

1 OP

1 OP

✓ TruSteel

✓ TruSteel

ALIMENTARY TRACT AND METABOLISM					
	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer	
continued					
than 80 mmol/mol; and The patient's HbA1c has not deteriorated more than 5 mm The patient has not had an increase in severe unexplained Either: 4.1 Applicant is a relevant specialist; or				ne; and	
4.1 Applicant is a relevant specialist, of 4.2 Applicant is a nurse practitioner working within their	r vocational scope.				
INSULIN PUMP CARTRIDGE – Special Authority see SA1985 of a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 packs of cartridge sets will be funded per Cartridge 300 U, t:lock × 10	year.	narma	·	andem Cartridge	
INSULIN PUMP INFUSION SET (STEEL CANNULA) - Special A		on p		•	
a) Maximum of 3 set per prescription b) Only on a prescription					
c) Maximum of 13 infusion sets will be funded per year.					
10 mm steel needle; 60 cm tubing × 10	130.00	1 OF	· ✓ N	liniMed Sure-T MMT-884A	
10 mm steel needle; 80 cm tubing x 10	130.00	1 OF	, ~ N	liniMed Sure-T MMT-886A	
6 mm steel needle; 60 cm tubing × 10	130.00	1 OF	· • • N	liniMed Sure-T MMT-864A	
6 mm steel needle; 80 cm tubing × 10	130.00	1 OF	· • • N	liniMed Sure-T MMT-866A	
8 mm steel needle; 60 cm tubing × 10	130.00	1 OF	, ~ N	liniMed Sure-T MMT-874A	
8 mm steel needle; 80 cm tubing × 10	130.00	1 OF	, ~ N	liniMed Sure-T MMT-876A	
(MiniMed Sure-T MMT-884A 10 mm steel needle; 60 cm tubing x (MiniMed Sure-T MMT-886A 10 mm steel needle; 80 cm tubing x				min oron	
INSULIN PUMP INFUSION SET (STEEL CANNULA, STRAIGHT			•	SA1985 on page 19 –	
Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year. 6 mm steel cannula; straight insertion; 80 cm line x 10 with					
10 needles	130.00	1 OF	, √ ⊥	ruSteel	
8 mm steel cannula; straight insertion; 80 cm line x 10 with 10 needles	130.00	1 OF	• ✓ T	ruSteel	
6 mm steel cannula; straight insertion; 60 cm line × 10 with	100.00	4.05		0 11	

8 mm steel cannula; straight insertion; 60 cm line × 10 with

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per Manufacturer

INSULIN PUMP INFUSION SET (TEFLON CANNULA) - Special Authority see SA1985 on page 19 - Retail pharmacy

- a) Maximum of 3 set per prescription
- b) Only on a prescription

c) Maximum of 13 infusion sets will be funded per year.			
13 mm teflon needle, 110 cm tubing × 10	130.00	1 OP	MiniMed Silhouette MMT-382A
13 mm teflon needle, 45 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-368A
13 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-381A
13 mm teflon needle, 80 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-383A
17 mm teflon needle, 110 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-377A
17 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-378A
17 mm teflon needle, 80 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-384A
6 mm teflon needle, 110 cm tubing × 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-398A
6 mm teflon needle, 45 cm blue tubing × 10	130.00	1 OP	✓ MiniMed Mio MMT-941A
6 mm teflon needle, 45 cm pink tubing × 10	130.00	1 OP	✓ MiniMed Mio MMT-921A
6 mm teflon needle, 60 cm blue tubing × 10	130.00	1 OP	✓ MiniMed Mio MMT-943A
6 mm teflon needle, 60 cm pink tubing × 10	130.00	1 OP	✓ MiniMed Mio MMT-923A
6 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-399A
6 mm teflon needle, 80 cm blue tubing	130.00	1 OP	✓ MiniMed Mio MMT-945A
6 mm teflon needle, 80 cm clear tubing × 10	130.00	1 OP	✓ MiniMed Mio MMT-965A
6 mm teflon needle, 80 cm pink tubing × 10	130.00	1 OP	✓ MiniMed Mio MMT-925A
6 mm teflon needle, 80 cm tubing × 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-387A
9 mm teflon needle, 110 cm tubing × 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-396A
9 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-397A
9 mm teflon needle, 80 cm clear tubing × 10	130.00	1 OP	✓ MiniMed Mio

MMT-975A

	Subsidy (Manufacturer's Price \$		Fully dised	
9 mm teflon needle, 80 cm tubing × 10	130.00	1 OP	1	MiniMed Quick-Set MMT-386A
(MiniMed Silhouette MMT-382A 13 mm teflon needle, 110 cm tub (MiniMed Silhouette MMT-368A 13 mm teflon needle, 45 cm tubir (MiniMed Silhouette MMT-383A 13 mm teflon needle, 80 cm tubir (MiniMed Silhouette MMT-384A 17 mm teflon needle, 80 cm tubir (MiniMed Quick-Set MMT-387A 6 mm teflon needle, 80 cm tubir (MiniMed Quick-Set MMT-386A 9 mm teflon needle, 80 cm tubing	$\log \times 10$ to be delisted by $\log \times 10$ to be delisted by $\log \times 10$ to be delisted $\log \times 10$ to be delisted by $\log \times 10$ to be delisted	ed 1 July 202 ed 1 July 202 ed 1 July 202 I 1 July 2024	24) 24) 24)	
INSULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE IN SA1985 on page 19 – Retail pharmacy a) Maximum of 3 sets per prescription	ISERTION WITH IN	ISERTION [DEVI	CE) – Special Authority see
b) Only on a prescriptionc) Maximum of 13 infusion sets will be funded per year.				
13 mm teflon cannula; angle insertion; insertion device; 110 c line × 10 with 10 needles	140.00	1 OP	1	AutoSoft 30
line × 10 with 10 needles		1 OP	1	AutoSoft 30
INSULIN PUMP INFUSION SET (TEFLON CANNULA, STRAIGH see SA1985 on page 19 – Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year.	T INSERTION WIT	H INSERTIC)N D	EVICE) - Special Authority
6 mm teflon cannula; straight insertion; insertion device; 110 cm line × 10 with 10 needles		1 OP	•	AutoSoft 90
line × 10 with 10 needles	140.00	1 OP	1	AutoSoft 90
9 mm teflon cannula; straight insertion; insertion device; 110 cm line × 10 with 10 needles	140.00	1 OP	1	AutoSoft 90
9 mm teflon cannula; straight insertion; insertion device; 60 cr line × 10 with 10 needles		1 OP	/	AutoSoft 90
INSULIN PUMP RESERVOIR – Special Authority see SA1985 or				Addoon 50
a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 packs of reservoir sets will be funded per		,		
10 × luer lock conversion cartridges 1.8 ml for Paradigm pum Cartridge for 7 series pump; 3.0 ml × 10	ps50.00	1 OP 1 OP		ADR Cartridge 1.8 MiniMed 3.0 Reservoir MMT-332A
Digestives Including Enzymes				
PANCREATIC ENZYME				
Cap pancreatin 150 mg (amylase 8,000 Ph Eur U, lipase 10,000 Ph Eur U, total protease 600 Ph Eur U)	3/1 03	100	,	Creon 10000
Cap pancreatin 300 mg (amylase 18,000 Ph Eur U, lipase 25,000 Ph Eur U, total protease 1,000 Ph Eur U)		100		Creon 25000
Modified release granules pancreatin 60.12 mg (amylase 3,600 Ph Eur U, lipase 5,000 Ph Eur U, protease 200 Ph Eur U)		20 g OP		Creon Micro
URSODEOXYCHOLIC ACID — Special Authority see SA1739 on Cap 250 mg	the next page - Re	-	у	Ursosan

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subs	idised	Generic	
\$	Per	1	Manufacturer	

⇒SA1739 Special Authority for Subsidy

initial application — (Alagille syndrome or progressive familial intrahepatic cholestasis) from any relevant practitioner.

Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 Patient has been diagnosed with Alagille syndrome; or
- 2 Patient has progressive familial intrahepatic cholestasis.

Initial application — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic severe drug induced cholestatic liver injury; and
- 2 Cholestatic liver injury not due to Total Parenteral Nutrition (TPN) use in adults; and
- 3 Treatment with ursodeoxycholic acid may prevent hospital admission or reduce duration of stay.

Initial application — (Primary biliary cholangitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Primary biliary cholangitis confirmed by antimitochondrial antibody titre (AMA) > 1:80, and raised cholestatic liver enzymes with or without raised serum IgM or, if AMA is negative, by liver biopsy; and
- 2 Patient not requiring a liver transplant (bilirubin > 100 umol/l; decompensated cirrhosis).

Initial application — (**Pregnancy**) from any relevant practitioner. Approvals valid for 6 months where the patient diagnosed with cholestasis of pregnancy.

Initial application — (Haematological Transplant) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 Dotio

- 1 Patient at risk of veno-occlusive disease or has hepatic impairment and is undergoing conditioning treatment prior to allogenic stem cell or bone marrow transplantation; and
- 2 Treatment for up to 13 weeks.

Initial application — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Paediatric patient has developed abnormal liver function as indicated on testing which is likely to be induced by Total Parenteral Nutrition (TPN); and
- 2 Liver function has not improved with modifying the TPN composition.

Renewal — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 6 months where the patient continues to benefit from treatment.

Renewal — (Pregnancy/Primary biliary cholangitis) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months where the paediatric patient continues to require TPN and who is benefiting from treatment, defined as a sustained improvement in bilirubin levels.

Laxatives

Bulk-forming Agents

Faecal Softeners

DO	CUSATE SODIUM - Only on a prescription			
*	Tab 50 mg	3.20	100	Coloxyl
*	Tab 120 mg	4.98	100	✓ Coloxyl

	Subsidy (Manufacturer's P \$	rice) Su Per	Fully bsidised	Brand or Generic Manufacturer
DOCUSATE SODIUM WITH SENNOSIDES Tab 50 mg with sennosides 8 mg	3.50	200	✓ <u>L</u>	axsol
POLOXAMER – Only on a prescription Not funded for use in the ear. * Oral drops 10%	4.17	30 ml OP	√ <u>c</u>	<u>Coloxyl</u>
Opioid Receptor Antagonists - Peripheral				

METHYLNALTREXONE BROMIDE - Special Authority	see SA1691 below - Retail p	harmacy	
Inj 12 mg per 0.6 ml vial	36.00	1	✓ Relistor
	246.00	7	✓ Relistor

⇒SA1691 Special Authority for Subsidy

Initial application — (Opioid induced constipation) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Both:

- 1 The patient is receiving palliative care; and
- 2 Either:
 - 2.1 Oral and rectal treatments for opioid induced constipation are ineffective; or
 - 2.2 Oral and rectal treatments for opioid induced constipation are unable to be tolerated.

Osmotic Laxatives

GLYCEROL * Suppos 2.8/4.0 g - Only on a prescription10.39	20	✓ <u>Lax-suppositories</u> <u>Glycerol</u>
LACTULOSE - Only on a prescription		
* Oral liq 10 g per 15 ml	500 ml	✓ <u>Laevolac</u>
MACROGOL 3350 WITH POTASSIUM CHLORIDE, SODIUM BICARBONATE A Powder for oral soln 13.125 g with potassium chloride 46.6 mg,	AND SODIUM (CHLORIDE
sodium bicarbonate 178.5 mg and sodium chloride 350.7 mg8.50	30	✓ Molaxole
SODIUM ACID PHOSPHATE – Only on a prescription Enema 16% with sodium phosphate 8%2.50	1	✓ Fleet Phosphate Enema
SODIUM CITRATE WITH SODIUM LAURYL SULPHOACETATE — Only on a p.	rescription	
Enema 90 mg with sodium lauryl sulphoacetate 9 mg per ml, 5 ml35.89	50	✓ <u>Micolette</u> ✓ Micolette-S29 S29

(Micolette-S29 S29 Enema 90 mg with sodium lauryl sulphoacetate 9 mg per ml, 5 ml to be delisted 1 August 2024)

Stim				

BISACODYL - Only on a prescription			
* Tab 5 mg	5.80	200	 Bisacodyl Viatris
* Suppos 10 mg		10	✓ Lax-Suppositories
SENNA - Only on a prescription			
* Tab, standardised	2.17	100	
	(8.21)		Senokot
	0.43	20	
	(2.06)		Senokot

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price)		Fully dised	Brand or Generic
	` \$	Per	1	Manufacturer
SODIUM PICOSULFATE - Special Authority see SA2053 below	 Retail pharmacy 			
Oral soln 7.5 mg per ml	7.40 30) ml OP	✓ D	ulcolax SP Drop

⇒SA2053 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The patient is a child with problematic constipation despite an adequate trial of other oral pharmacotherapies including macrogol where practicable; and
- 2 The patient would otherwise require a high-volume bowel cleansing preparation or hospital admission.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Metabolic Disorder Agents

ALGLUCOSIDASE ALFA - Special Authority see SA1986 below - Retail pharmacy ✓ Myozyme

⇒SA1986 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient is aged up to 24 months at the time of initial application and has been diagnosed with infantile Pompe disease; and
- 2 Any of the following:
 - 2.1 Diagnosis confirmed by documented deficiency of acid alpha-glucosidase by prenatal diagnosis using chorionic villus biopsies and/or cultured amniotic cells; or
 - 2.2 Documented deficiency of acid alpha-glucosidase, and urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides; or
 - 2.3 Documented deficiency of acid alpha-glucosidase, and documented molecular genetic testing indicating a disease-causing mutation in the acid alpha-glucosidase gene (GAA gene); or
 - 2.4 Documented urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides, and molecular genetic testing indicating a disease-causing mutation in the GAA gene; and
- 3 Patient has not required long-term invasive ventilation for respiratory failure prior to starting enzyme replacement therapy (ERT): and
- 4 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by ERT or might be reasonably expected to compromise a response to ERT; and
- 5 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks.

Renewal only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks; and
- 3 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 4 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by ERT; and
- 5 Patient has not developed another medical condition that might reasonably be expected to compromise a response to
- 6 There is no evidence of life threatening progression of respiratory disease as evidenced by the needed for > 14 days of invasive ventilation: and
- 7 There is no evidence of new or progressive cardiomyopathy.

ARGININE - Special Authority see SA2042 on the next p	page – Retail pharmacy		
Tab 1,000 mg	CBS	90	Clinicians
Cap 500 mg	CBS	50	✓ Solgar
Powder	CBS	400 g	✓ Biomed

Subsidy (Manufacturer's Price)	Sub	Fully	Brand or Generic	
\$	Per	✓	Manufacturer	

⇒SA2042 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to arginine supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to arginine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

⇒SA1987 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has a confirmed diagnosis of homocystinuria; and
- 2 Any of the following:
 - 2.1 A cystathionine beta-synthase (CBS) deficiency; or
 - 2.2 A 5,10-methylene-tetrahydrofolate reductase (MTHFR) deficiency; or
 - 2.3 A disorder of intracellular cobalamin metabolism; and
- 3 An appropriate homocysteine level has not been achieved despite a sufficient trial of appropriate vitamin supplementation.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

		DENZYME Q10 - Special Authority see SA2039 below - Retail pharmacy	COF
Solgar	30	Cap 120 mgCBS	
✓ Go Healthy	60	Cap 160 mgCBS	

⇒SA2039 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to coenzyme Q10 supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to coenzyme Q10 supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

⇒SA1988 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has been diagnosed with mucopolysaccharidosis VI; and
- 2 Either:
 - 2.1 Diagnosis confirmed by demonstration of N-acetyl-galactosamine-4-sulfatase (arylsulfatase B) deficiency by either enzyme activity assay in leukocytes or skin fibroblasts; or
 - 2.2 Detection of two disease causing mutations and patient has a sibling who is known to have mucopolysaccharidosis VI

Renewal only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication

Subsic	dy Ful	y Brand or
(Manufacture		
<u> </u>	Per •	Manufacturer

continued...

and/or adjustment of infusion rates; and

- 3 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by Enzyme Replacement Therapy (ERT); and
- 4 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT.

IDURSULFASE - Special Authority see SA1623 below - Retail pharmacy

✓ Elaprase

⇒SA1623 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hunter Syndrome (mucopolysaccharidosis II); and
- 2 Fither:
 - 2.1 Diagnosis confirmed by demonstration of iduronate 2-sulfatase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts: or
 - 2.2 Detection of a disease causing mutation in the iduronate 2-sulfatase gene; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with idursulfase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT); and
- 5 Idursulfase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 weeks post-HSCT) at doses no greater than 0.5 mg/kg every week.

LARONIDASE - Special Authority see SA1695 below - Retail pharmacy

⇒SA1695 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hurler Syndrome (mucopolysacchardosis I-H); and
- 2 Fither:
 - 2.1 Diagnosis confirmed by demonstration of alpha-L-iduronidase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts; or
 - 2.2 Detection of two disease causing mutations in the alpha-L-iduronidase gene and patient has a sibling who is known to have Hurler syndrome; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with laronidase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT); and
- 5 Laronidase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 post-HSCT) at doses no greater than 100 units/kg every week.

LEVOCARNITINE -	- Special Authority se	e SA2040 on the next	t page – Retail pharmacy
-----------------	------------------------	----------------------	--------------------------

Tab 500 mg	CBS	30	✓ Solgar
Cap 250 mg	CBS	30	✓ Solgar
Cap 500 mg	CBS	60	✓ Balance
Oral liq 1 g per 10 ml	CBS	118 ml	✓ Carnitor \$29 ✓ Novitium Sugar
Oral liq 500 mg per 10 ml	CBS	300 ml	✓ Balance

✓ Aldurazyme

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
<u></u>	Per	✓	Manufacturer

⇒SA2040 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to carnitine supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to carnitine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

RIBOFLAVIN – Special Authority see SA2041 below – Retail pharmacy Tab 100 mgCBS	100	✓ Country Life ✓ Puritan's Pride Vitamin B-2 100 mg \$29
Cap 100 mgCBS	100	✓ Solgar

⇒SA2041 Special Authority for Subsidy

Initial application only from a metabolic physician or neurologist. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to riboflavin supplementation.

Renewal only from a metabolic physician or neurologist. Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to riboflavin supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

⇒SA1989 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 1 month for applications meeting the following criteria: All of the following:

- 1 Patient has phenylketonuria (PKU) and is pregnant or actively planning to become pregnant; and
- 2 Treatment with sapropterin is required to support management of PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

Renewal only from a metabolic physician or any relevant practitioner on the recommendation of a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Following the initial one-month approval, the patient has demonstrated an adequate response to a 2 to 4 week trial of sapropterin with a clinically appropriate reduction in phenylalanine levels to support management of PKU during pregnancy; or
 - 1.2 On subsequent renewal applications, the patient has previously demonstrated response to treatment with sapropterin and maintained adequate phenylalanine levels to support management of PKU during pregnancy; and
- 2 Any of the following:
 - 2.1 Patient continues to be pregnant and treatment with sapropterin will not continue after delivery; or
 - 2.2 Patient is actively planning a pregnancy and this is the first renewal for treatment with sapropterin; or
 - 2.3 Treatment with sapropterin is required for a second or subsequent pregnancy to support management of their PKU during pregnancy; and

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
 \$	Per	1	

continued...

- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

SODIUM BENZOATE - Special Authority see SA1599 below - Retail pharmacy

⇒SA1599 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

SODIUM PHENYLBUTYRATE - Special Authority see SA1990 below - Retail pharmacy
Grans 483 mg per g.......2,016.00 174 g OP

✓ Pheburane

⇒SA1990 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder involving a deficiency of carbamylphosphate synthetase, ornithine transcarbamylase or argininosuccinate synthetase.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

TAURINE - Special Authority see SA2043 below - Retail pharmacy

Cap 500 mg	CBS	50	✓ Solgar
Cap 1,000 mg		90	✓ Life Extension
Powder	CBS	300 a	✓ Life Extension

⇒SA2043 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected specific mitochondrial disorder that may respond taurine supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has confirmed diagnosis of a specific mitochondrial disorder which responds to taurine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

TRIENTINE – Special Authority see SA2324 below – Retail pharmacy
Cap 250 mg.......2,022.00 100 ✓ Trientine Waymade

⇒SA2324 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Wilson disease; and
- 2 Treatment with D-penicillamine has been trialled and discontinued because the person has experienced intolerable side effects or has not received sufficient benefit; and
- 3 Treatment with zinc has been trailled and discontinued because the person has experienced intolerable side effects or has not received sufficient benefit, or zinc is considered clinically inappropriate as the person has symptomatic liver disease and requires copper chelation.

Gaucher's Disease

⇒SA2137 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has a diagnosis of symptomatic type 1 or type 3* Gaucher disease confirmed by the demonstration of specific deficiency of glucocerebrosidase in leukocytes or cultured skin fibroblasts, and genotypic analysis; and
- 2 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by enzyme replacement therapy (ERT) or the disease might be reasonably expected to compromise a response to ERT; and
- 3 Any of the following:
 - 3.1 Patient has haematological complications of Gaucher disease; or
 - 3.2 Patient has skeletal complications of Gaucher disease; or
 - 3.3 Patient has significant liver dysfunction or hepatomegaly attributable to Gaucher disease; or
 - 3.4 Patient has reduced vital capacity from clinically significant or progressive pulmonary disease due to Gaucher disease: or
 - 3.5 Patient is a child and has experienced growth failure with significant decrease in percentile linear growth over a 6-12 month period; and
- 4 Taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units).

Note: Indication marked with * is an unapproved indication

Renewal only from a metabolic physician or any relevant practitioner on the recommendation of a metabolic physician. Approvals valid for 3 years for applications meeting the following criteria:

All of the following:

- 1 Patient has demonstrated a symptomatic improvement and has maintained improvements in the main symptom or symptoms for which therapy was started; and
- 2 Patient has demonstrated a clinically objective improvement or no deterioration in haemoglobin levels, platelet counts and liver and spleen size; and
- 3 Radiological (MRI) signs of bone activity performed at two years since initiation of treatment, and five yearly thereafter, demonstrate no deterioration shown by the MRI, compared with MRI taken immediately prior to commencement of therapy or adjusted dose; and
- 4 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT; and
- 5 Patient is adherent with regular treatment and taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units).

Mouth and Throat

Agents Used in Mouth Ulceration

BENZYDAMINE HYDROCHLORIDE

Soln 0.15% - Higher subsidy of \$22.60 per 500 ml with

Additional subsidy by endorsement for a patient who has oral mucositis as a result of treatment for cancer, and the prescription is endorsed accordingly.

CARMELLOSE SODIUM WITH GELATIN AND PECTIN

Paste	17.20	56 g OP	✓ Stomahesive
	4.55	15 g OP	
	(7.90)	Ü	Orabase
	1.52	5 g OP	
	(3.60)		Orabase
Powder	8.48 [°]	28 g OP	
	(10.95)	Ü	Stomahesive

Difflam

	Subsidy (Manufacturer's P \$	Price) Subs	Fully Brand or sidised Generic ✓ Manufacturer
TRIAMCINOLONE ACETONIDE Paste 0.1%	5.49	5 g OP	✓ Kenalog in Orabase
Oropharyngeal Anti-infectives			
AMPHOTERICIN B Lozenges 10 mg	5.86	20	✓ Fungilin
Oral gel 20 mg per g		40 g OP	✓ <u>Decozol</u>
Oral liq 100,000 u per ml	2.22	24 ml OP	✓ <u>Nilstat</u>
Vitamins			
Vitamin B			
HYDROXOCOBALAMIN ★ Inj 1 mg per ml, 1 ml ampoule – Up to 6 inj available on	a PSO2.46	3	✓ Cobal-B12 \$29 ✓ <u>Hydroxocobalamin</u> Panpharma ✓ Vita-B12
	4.10	5	✓ Cobalin-H S29 ✓ Neo-Cytamen S29 S29
	8.20	10	✓ Vitarubin Depot Injection \$29
PYRIDOXINE HYDROCHLORIDE a) No more than 100 mg per dose b) Only on a prescription			·
* Tab 50 mg * Tab 50 mg		90 500	✓ <u>Vitamin B6 25</u> ✓ Pyridoxine multichem
THIAMINE HYDROCHLORIDE − Only on a prescription * Tab 50 mg	4.65	100	✓ Thiamine multichem
/ITAMIN B COMPLEX * Tab, strong, BPC		500	✓ Bplex
Vitamin C			
ASCORBIC ACID			
a) No more than 100 mg per dose b) Only on a prescription			
* Tab 100 mg	12.50	500	✓ Cvite

	Subsidy (Manufacturer's Price \$	e) Per	Fully Subsidised	Brand or Generic Manufacturer
Vitamin D				
ALFACALCIDOL				
* Cap 0.25 mcg	26.32	100	1	One-Alpha
			1	One-Alpha S29 S29
* Cap 1 mcg	87.98	100	1	One-Alpha
			•	One-Alpha S29 S29
* Oral drops 2 mcg per ml	60.68	20 ml C)P 🗸	One-Alpha
(One-Alpha S29 S29 Cap 1 mcg to be delisted 1 August 2024)				
CALCITRIOL				
* Cap 0.25 mcg		100	✓	Calcitriol-AFT
* Cap 0.5 mcg	13.68	100	1	<u>Calcitriol-AFT</u>
COLECALCIFEROL				
* Cap 1.25 mg (50,000 iu) - Maximum of 12 cap per prescripti	ion3.65	12	✓	Vit.D3
* Oral liq 188 mcg per ml (7,500 iu per ml)	9.00	5 ml O	Ρ 🗸	Clinicians
Multivitamin Preparations				

MULTIVITAMIN RENAL - Special Authority see SA1546 below -	Retail pharmacy		
* Cap	7.28	30	Clinicians Renal Vit

⇒SA1546 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Fither:

- 1 The patient has chronic kidney disease and is receiving either peritoneal dialysis or haemodialysis; or
- 2 The patient has chronic kidney disease grade 5, defined as patient with an estimated glomerular filtration rate of < 15 ml/min/1.73 m² body surface area (BSA).

MULTIVITAMINS - Special Authority see SA1036 below - Retail pharmacy 200 g OP ✓ Paediatric Seravit

⇒SA1036 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has inborn errors of metabolism.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where patient has had a previous approval for multivitamins.

VITAMINS

*	Tab (BPC cap strength)18.5	0 1,000	✓ Mvite
*	Cap (fat soluble vitamins A, D, E, K) - Special Authority see		
	SA1720 below – Retail pharmacy23.4	0 60	✓ Vitabdeck

⇒SA1720 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Patient has cystic fibrosis with pancreatic insufficiency; or
- 2 Patient is an infant or child with liver disease or short gut syndrome; or
- 3 Patient has severe malabsorption syndrome.

		(Manufacturar's Price)		Subsidised	Generic
		(Manufacturer's Price) \$	Per	Subsidised	Manufacturer
		*			
N	linerals				
С	alcium				
CA	LCIUM CARBONATE				
	Tab 1.25 g (500 mg elemental)	7.28	250	✓	Calci-Tab 500
	Tab eff 1.25 g (500 mg elemental) – Subsidy by endorseme		100	1	Calcium 500 mg
	Subsidy by endorsement – Only when prescribed for pa considered unsuitable.	ediatric patients (< 5	years)	where cal	cium carbonate oral liquid is
CA	LCIUM GLUCONATE				
*	Inj 10%, 10 ml ampoule	32.00	10	✓	Max Health -
					Hameln S29
		64.00	20	1	Max Health \$29
lc	odine				
РО	TASSIUM IODATE				
*	Tab 253 mcg (150 mcg elemental iodine)	5.99	90	•	<u>NeuroTabs</u>
lr	on				
FE	RROUS FUMARATE				
*	Tab 200 mg (65 mg elemental)	3.04	100	1	Ferro-tab
FΕ	RROUS FUMARATE WITH FOLIC ACID				
*	Tab 310 mg (100 mg elemental) with folic acid 350 mcg	5.98	100	1	Ferro-F-Tabs
FE	RROUS SULFATE				
*	Tab long-acting 325 mg (105 mg elemental)	2.55	30	✓	Ferrograd
*	Oral liq 30 mg (6 mg elemental) per 1 ml		250 m		Ferro-Liquid
			500 m		<u>Ferodan</u>
IRO	ON (AS FERRIC CARBOXYMALTOSE) - Special Authority s				
	Inj 50 mg per ml, 10 ml vial	150.00	1	•	Ferinject
	SA1840 Special Authority for Subsidy				
Init	tial application — (serum ferritin less than or equal to 20 r	ncq/L) from any rele	vant r	ractitioner	. Approvals valid for 3

Subsidy

Fully

Brand or

Initial application — (serum ferritin less than or equal to 20 mcg/L) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 Patient has been diagnosed with iron-deficiency anaemia with a serum ferritin level of less than or equal to 20 mcg/L; and
- 2 Any of the following:
 - 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
 - 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
 - 2.3 Rapid correction of anaemia is required.

Renewal — (serum ferritin less than or equal to 20 mcg/L) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient continues to have iron-deficiency anaemia with a serum ferritin level of less than or equal to 20 mcg/L; and
- 2 A re-trial with oral iron is clinically inappropriate.

Initial application — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist,

✓ Ferrosia

Subsidy (Manufacturer's Price)	Subsi	Fully dised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with iron-deficiency anaemia; and
- 2 Any of the following:
 - 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
 - 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
 - 2.3 Patient has symptomatic heart failure, chronic kidney disease stage 3 or more or active inflammatory bowel disease and a trial of oral iron is unlikely to be effective; or

34 50

2.4 Rapid correction of anaemia is required.

Renewal — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient continues to have iron-deficiency anaemia; and
- 2 A re-trial with oral iron is clinically inappropriate.

IKC	ON POLYMALTOS	SE .
*	Ini 50 ma ner ml	2 ml amnoula

Tri do my por mi, z mi ampoulo	J	- Torrosig
Magnesium		
MAGNESIUM HYDROXIDE		
Suspension 8%33.60	355 ml	✓ Phillips Milk of
		Magnesia S29
MAGNESIUM SULPHATE		
* Inj 2 mmol per ml, 5 ml ampoule	10	✓ Martindale
* Inj 2 mmol per ml, 10 ml ampoule75.06	10	✓ Inresa S29
Zinc		
ZINC SULPHATE		
* Cap 137.4 mg (50 mg elemental)	100	✓ Zincaps

BLOOD AND BLOOD FORMING ORGANS

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

Antianaemics

Hypoplastic and Haemolytic

⇒SA2266 Special Authority for Subsidy

Initial application — (chronic renal failure) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient in chronic renal failure: and
- 2 Haemoglobin is less than or equal to 100g/L; and
- 3 Any of the following:
 - 3.1 Both:
 - 3.1.1 Patient does not have diabetes mellitus; and
 - 3.1.2 Glomerular filtration rate is less than or equal to 30ml/min; or
 - 3.2 Both:
 - 3.2.1 Patient has diabetes mellitus; and
 - 3.2.2 Glomerular filtration rate is less than or equal to 45ml/min; or
 - 3.3 Patient is on haemodialysis or peritoneal dialysis.

Initial application — (myelodysplasia) from any specialist. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of myelodysplasia (MDS)*; and
- 2 Has had symptomatic anaemia with haemoglobin < 100g/L and is red cell transfusion-dependent; and
- 3 Patient has very low, low or intermediate risk MDS based on the WHO classification based prognostic scoring system for myelodysplastic syndrome (WPSS); and
- 4 Other causes of anaemia such as B12 and folate deficiency have been excluded; and
- 5 Patient has a serum epoetin level of < 500 IU/L; and
- 6 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Note: Indication marked with * is an unapproved indication

Renewal — (chronic renal failure) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (myelodysplasia) from any specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient's transfusion requirement continues to be reduced with erythropoietin treatment; and
- 2 Transformation to acute myeloid leukaemia has not occurred; and
- 3 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Note: Indication marked with * is an unapproved indication

EPOETIN ALFA - Special Authority see SA2266 above - Retail pharmacy

Wastage claimable			
Inj 1,000 iu in 0.5 ml, syringe	250.00	6	Binocrit
Inj 2,000 iu in 1 ml, syringe	100.00	6	Binocrit
Inj 3,000 iu in 0.3 ml, syringe		6	Binocrit
Inj 4,000 iu in 0.4 ml, syringe		6	✓ Binocrit
Inj 5,000 iu in 0.5 ml, syringe		6	✓ Binocrit
Inj 6,000 iu in 0.6 ml, syringe		6	✓ Binocrit
Inj 8,000 iu in 0.8 ml, syringe		6	✓ Binocrit
Inj 10,000 iu in 1 ml, syringe		6	Binocrit
Inj 40,000 iu in 1 ml, syringe		1	✓ Binocrit

	Subsidy (Manufacturer's Price) \$	Subsi Per	Fully dised ✓	Brand or Generic Manufacturer
Megaloblastic				
FOLIC ACID * Tab 0.8 mg	26.60	1,000		olic Acid multichem
* Tab 5 mg Oral liq 50 mcg per ml		100 5 ml OP	_	olic Acid Viatris iomed

Antifibrinolytics, Haemostatics and Local Sclerosants

EFTRENONACOG ALFA [RECOMBINANT FACTOR IX] - [Xpharm]

For patients with haemophilia B receiving prophylaxis treatment. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management group.

Treaters Group in conjunction with the National Haemor	ohilia Management gro	up.	
Inj 250 iu vial	612.50	1	Alprolix
Inj 500 iu vial	1,225.00	1	✓ Alprolix
Inj 1,000 iu vial	2,450.00	1	✓ Alprolix
Inj 2,000 iu vial	4,900.00	1	✓ Alprolix
Inj 3,000 iu vial	7,350.00	1	✓ Alprolix
Inj 4,000 iu vial	9,800.00	1	Alprolix
ELTROMBOPAG - Special Authority see SA1743 below -	Retail pharmacy		
Wastage claimable			
Tab 25 mg	1,550.00	28	✓ Revolade
Tab 50 mg	3,100.00	28	✓ Revolade

⇒SA1743 Special Authority for Subsidy

Initial application — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has had a splenectomy; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Any of the following:
 - 3.1 Patient has a platelet count of 20,000 to 30,000 platelets per microlitre and has evidence of significant mucocutaneous bleeding; or
 - 3.2 Patient has a platelet count of less than or equal to 20,000 platelets per microlitre and has evidence of active bleeding; or
 - 3.3 Patient has a platelet count of less than or equal to 10,000 platelets per microlitre.

Initial application — (idiopathic thrombocytopenic purpura - preparation for splenectomy) only from a haematologist. Approvals valid for 6 weeks where the patient requires eltrombopag treatment as preparation for splenectomy.

Initial application — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a significant and well-documented contraindication to splenectomy for clinical reasons; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Either:
 - 3.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per microliter: or
 - 3.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre

Subsidy		Fully	Brand or
(Manufacturer's Price)		osidised	Generic
<u> </u>	Per		Manufacturer

continued...

and significant mucocutaneous bleeding.

Initial application — (severe aplastic anaemia) only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Two immunosuppressive therapies have been trialled and failed after therapy of at least 3 months duration; and
- 2 Either:
 - 2.1 Patient has severe aplastic anaemia with a platelet count of less than or equal to 20,000 platelets per microliter; or
 - 2.2 Patient has severe aplastic anaemia with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

Renewal — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 12 months where the patient has obtained a response (see Note) from treatment during the initial approval or subsequent renewal periods and further treatment is required.

Note: Response to treatment is defined as a platelet count of > 30,000 platelets per microlitre.

Renewal — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient's significant contraindication to splenectomy remains; and
- 2 The patient has obtained a response from treatment during the initial approval period; and
- 3 Patient has maintained a platelet count of at least 50,000 platelets per microlitre on treatment; and
- 4 Further treatment with eltrombopag is required to maintain response.

Renewal — (severe aplastic anaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has obtained a response from treatment of at least 20,000 platelets per microlitre above baseline during the initial approval period; and
- 2 Platelet transfusion independence for a minimum of 8 weeks during the initial approval period.

EMICIZUMAB - [Xpharm] - Special Authority see \$A2272 below

✓ Hemlibra	1	3,570.00	Inj 30 mg in 1 ml vial
✓ Hemlibra	1	7,138.00	Inj 60 mg in 0.4 ml vial
✓ Hemlibra	1	•	Inj 105 mg in 0.7 ml vial
✓ Hemlibra	1	17,846.00	Inj 150 mg in 1 ml vial

⇒SA2272 Special Authority for Subsidy

Initial application — (Severe Haemophilia A with or without FVIII inhibitors) only from a haematologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has severe congenital haemophilia A with a severe bleeding phenotype (endogenous factor VIII activity less than or equal to 2%); and
- 2 Emicizumab is to be administered at a dose of no greater than 3 mg/kg weekly for 4 weeks followed by the equivalent of 1.5 mg/kg weekly.

EPTACOG ALFA [RECOMBINANT FACTOR VIIA] - [Xpharm]

For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group. Rare Clinical Circumstances Brand of bypassing agent for > 14 days predicted use. Access to funded treatment for > 14 days predicted use is by named patient application to the Haemophilia Treaters Group, subject to access criteria.

Inj 1 mg syringe	1,178.30	1	✓ NovoSeven RT
Inj 2 mg syringe	2,356.60	1	✓ NovoSeven RT
Inj 5 mg syringe	·	1	✓ NovoSeven RT
Inj 8 mg syringe	·	1	✓ NovoSeven RT

	BLOOD AND	DLUUD FU	JRIMING ORGANS
	Subsidy	Fu	,
	(Manufacturer's Price) \$	Subsidise Per	ed Generic ✓ Manufacturer
FACTOR EIGHT INHIBITOR BYPASSING FRACTION - [Xpha	·		- manadaror
For patients with haemophilia. Preferred Brand of bypassin		oredicted use	. Access to funded treatment
is managed by the Haemophilia Treaters Group in conjuncti	on with the National H	aemophilia M	anagement Group.
Inj 500 Ŭ			/ FEIBA NF
Inj 1,000 U	2,630.00		/ FEIBA NF
Inj 2,500 U	6,575.00	1 •	FEIBA NF
MOROCTOCOG ALFA [RECOMBINANT FACTOR VIII] - [Xph			
For patients with haemophilia. Rare Clinical Circumstances			
treatment is managed by the Haemophilia Treaters Group in	n conjunction with the I	National Haer	nophilia Management Group,
subject to criteria.			4 14
Inj 250 iu prefilled syringe			Xyntha
Inj 500 iu prefilled syringe			Xyntha
Inj 1,000 iu prefilled syringe			/ Xyntha
Inj 2,000 iu prefilled syringeInj 3,000 iu prefilled syringe			✓ Xyntha ✓ Xyntha
		'	Аупша
NONACOG GAMMA, [RECOMBINANT FACTOR IX] – [Xpharn			atom One of the continuation
For patients with haemophilia. Access to funded treatment with the National Haemophilia Management Group.	is managed by the Ha	emopnilia i re	aters Group in conjunction
, ,	40E 00	1 •	/ DIVIDIO
Inj 500 iu vial Inj 1,000 iu vial		-	/ RIXUBIS / RIXUBIS
Inj 2,000 iu vial		-	✓ RIXUBIS
Inj 3,000 iu vial	·	-	✓ RIXUBIS
OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (ADVATE) -	•		THAT
For patients with haemophilia. Preferred Brand of short hal		vr VIII. Accord	to funded treatment is
managed by the Haemophilia Treaters Group in conjunction			
Inj 250 iu vial			Advate
Inj 500 iu vial			✓ Advate
Inj 1,000 iu vial		-	✓ Advate
Inj 1,500 iu vial		-	/ Advate
Inj 2,000 iu vial	,	-	✓ Advate
Inj 3,000 iu vial	,		✓ Advate
OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (KOGENATE	FS) - [Xnharm]		
For patients with haemophilia. Rare Clinical Circumstances		e recombinant	t factor VIII. Access to funded
treatment is managed by the Haemophilia Treaters Group in			
subject to criteria.	,		, , , , , , , , , , , , , , , , , , , ,
Inj 250 iu vial	237.50	1 •	✓ Kogenate FS
Inj 500 iu vial	475.00	1 •	✓ Kogenate FS
Inj 1,000 iu vial	950.00	1 •	✓ Kogenate FS
Inj 2,000 iu vial	,		✓ Kogenate FS
Inj 3,000 iu vial	2,850.00	1 •	Kogenate FS
RURIOCTOCOG ALFA PEGOL [RECOMBINANT FACTOR VIII			
For patients with haemophilia A receiving prophylaxis treatn		d treatment is	managed by the Haemophilia
Treaters Group in conjunction with the National Haemophilia	a Management group.		
Inj 250 iu vial			/ Adynovate
Inj 500 iu vial			Adynovate
Inj 1,000 iu vial			Adynovate
Inj 2,000 iu vial	2,400.00	1 •	Adynovate
SODIUM TETRADECYL SULPHATE			
* Inj 3% 2 ml		5	
	(73.00)		Fibro-vein

Subsidy	, 0	Fully	Brand or
(Manufacturer's Price \$	e) Su Per	bsidised ✓	Generic Manufacturer
RANEXAMIC ACID			
Tab 500 mg10.45	60	✓ N	lercury Pharma
45.68	100	✓ 0	yklokapron
Vitamin K			
HYTOMENADIONE			
Inj 2 mg per 0.2 ml - Up to 5 inj available on a PSO8.00	5		onakion MM
Inj 10 mg per ml, 1 ml – Up to 5 inj available on a PSO9.21	5	✓ K	Conakion MM
Antithrombotic Agents			
Antiplatelet Agents			
SPIRIN			
Tab 100 mg12.65	990	✓ <u>E</u>	thics Aspirin EC
LOPIDOGREL			
Tab 75 mg5.07	84	✓ A	rrow - Clopid

(Brilinta Tab 90 mg to be delisted 1 July 2024)

⇒SA1955 Special Authority for Subsidy

Initial application — (acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

90.00

Both:

DIPYRIDAMOI F

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

TICAGRELOR - Special Authority see SA1955 below - Retail pharmacv

Initial application — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

1 Fither

- 1.1 Patient has had a neurological stenting procedure* in the last 60 days; or
- 1.2 Patient is about to have a neurological stenting procedure performed*; and

2 Either:

- 2.1 Patient has demonstrated clopidogrel resistance using the P2Y12 (VerifyNow) assay or another appropriate platelet function assay and requires antiplatelet treatment with ticagrelor; or
- 2.2 Either:
 - 2.2.1 Clopidogrel resistance has been demonstrated by the occurrence of a new cerebral ischemic event; or
 - 2.2.2 Clopidogrel resistance has been demonstrated by the occurrence of transient ischemic attack symptoms referable to the stent.

Initial application — (Percutaneous coronary intervention with stent deployment) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

1 Patient has undergone percutaneous coronary intervention; and

continued...

60

56

✓ Pytazen SR

✓ Brilinta

✓ Ticagrelor Sandoz

Subsidy)	Fully	Brand or
(Manufacturer's Price		Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic**.

Initial application — (Stent thrombosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has experienced cardiac stent thrombosis whilst on clopidogrel.

Renewal — (subsequent acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Renewal — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient is continuing to benefit from treatment; and
- 2 Treatment continues to be clinically appropriate.

Renewal — (Percutaneous coronary intervention with stent deployment) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic**.

Notes: indications marked with * are unapproved indications.

Note: ** Clopidogrel allergy is defined as a history of anaphylaxis, urticaria, generalised rash or asthma (in non-asthmatic patients) developing soon after clopidogrel is started and is considered unlikely to be caused by any other treatment.

Heparin and Antagonist Preparations

Inj 20 mg in 0.2 ml syrir	nge31.2	3 10	Clexane
Inj 40 mg in 0.4 ml syrir	nge42.49	9 10	✓ Clexane
	nge60.6		✓ Clexane
	nge80.89		✓ Clexane
	nge101.30		✓ Clexane
	ringe125.8		✓ Clexane Forte
	nge143.80		✓ Clexane Forte

⇒SA2152 Special Authority for Subsidy

Initial application — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patients pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Initial application — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

1 For the short-term treatment of venous thromboembolism prior to establishing a therapeutic level of oral anti-coagulant treatment; or

Subsidy	F	ully	Brand or
Manufacturer's Price)	Subsid	sed	Generic
 \$	Per	✓	Manufacturer

continued...

- 2 For the prophylaxis and treatment of venous thromboembolism in high risk surgery; or
- 3 To enable cessation/re-establishment of existing oral anticoagulant treatment pre/post surgery; or
- 4 For the prophylaxis and treatment of venous thromboembolism in Acute Coronary Syndrome surgical intervention; or
- 5 To be used in association with cardioversion of atrial fibrillation.

Initial application — (Short-term use during treatment of COVID-19 with nirmatrelvir with ritonavir) from any relevant practitioner. Approvals valid for 2 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is receiving an anticoagulation treatment that has drug/drug interactions with ritonavir that increases risk of bleeding; and
- 2 Patient meets the Access Criteria for COVID-19 antivirals published on the Pharmac website*; and
- 3 Other antiviral treatments for COVID-19 have been considered and are not clinically suitable options.

Renewal — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

HEPARIN SODIUM

- 1 Low molecular weight heparin treatment is required during a patient's pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Renewal — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month where low molecular weight heparin treatment or prophylaxis is required for a second or subsequent event (surgery, ACS, cardioversion, or prior to oral anti-coagulation).

Inj 1,000 iu per ml, 5 ml ampoule	86.11	50	✓ Pfizer
Inj 5,000 iu per ml, 5 ml vial	83.00	10	✓ <u>Heparin Sodium</u>
			Panpharma
Inj 5,000 iu per ml, 1 ml	32.66	5	✓ DBL Heparin
			Sodium S29
	70.33		✓ Hospira
Inj 25,000 iu per ml, 0.2 ml	22.42	5	✓ Hospira
	42.40		✓ Heparin DBL S29
	482.20	50	✓ Heparin DBL S29
HEPARINISED SALINE			
Inj 10 iu per ml, 5 ml	65.48	50	✓ Pfizer

Oral	Anticoagulants
------	-----------------------

DARIGATRAN

2,12,0,111,0,11			
Cap 75 mg - No more than 2 cap per day	27.99	60	Pradaxa
Pradaxa to be Principal Supply on 1 July 2024			
Cap 110 mg	27.99	60	Pradaxa
Pradaxa to be Principal Supply on 1 July 2024			
Cap 150 mg	27.99	60	Pradaxa
Pradaxa to be Principal Supply on 1 July 2024			
RIVAROXABAN			
Tab 10 mg - No more than 1 tab per day	15.60	30	✓ Xarelto
Tab 15 mg - Up to 14 tab available on a PSO	14.56	28	✓ Xarelto
Tab 20 mg	14.56	28	✓ Xarelto

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	Generic
	\$	Per	1	Manufacturer
WARFARIN SODIUM				
Note: Marevan and Coumadin are not interchangeable.				
* Tab 1 mg	3.46	50	✓	Coumadin
•	7.50	100	✓	Marevan
* Tab 2 mg	4.31	50	✓	Coumadin
* Tab 3 mg	12.00	100	✓	Marevan
* Tab 5 mg	5.93	50	✓	Coumadin
-	13.50	100	✓	Marevan
Blood Colony-stimulating Factors				
FILGRASTIM – Special Authority see SA1259 below – Retail ph	,	40		Nii
Inj 300 mcg per 0.5 ml prefilled syringe		10		Nivestim Nivestim
Inj 480 mcg per 0.5 ml prefilled syringe	148.58	10	•	<u>Nivestim</u>

⇒SA1259 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 20%*); or
- 2 Peripheral blood stem cell mobilisation in patients undergoing haematological transplantation; or
- 3 Peripheral blood stem cell mobilisation or bone marrow donation from healthy donors for transplantation; or
- 4 Treatment of severe chronic neutropenia (ANC < 0.5 ×10⁹/L); or
- 5 Treatment of drug-induced prolonged neutropenia (ANC < 0.5 ×10⁹/L).

Note: *Febrile neutropenia risk greater than or equal to 20% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

PEGFILGRASTIM - Special Authority see SA1912 below - Retail pharmacy

⇒SA1912 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where used for prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 5%*). Note: *Febrile neutropenia risk greater than or equal to 5% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

Fluids and Electrolytes

Intravenous Administration

GLUCOSE [DEXTROSE]		
* Inj 50%, 10 ml ampoule – Up to 5 inj available on a PSO34.75	5	✓ Biomed
* Inj 50%, 90 ml bottle - Up to 5 inj available on a PSO17.50	1	✓ Biomed
POTASSIUM CHLORIDE		
* Inj 75 mg per ml, 10 ml65.00	50	Juno
		✓ Pfizer S29

	Subsidy (Manufacturer's Pri \$	ce) Su Per	Fully bsidised	
SODIUM BICARBONATE				
Inj 8.4%, 50 ml	23.52	1	✓	Biomed
a) Up to 5 inj available on a PSO				
b) Not in combination				
Inj 8.4%, 100 ml	24.10	1	✓	Biomed
a) Up to 5 inj available on a PSOb) Not in combination				
SODIUM CHLORIDE				
Not funded for use as a nasal drop. Not funded for nebulise for nebuliser use.	er use except when	used in co	njunctio	n with an antibiotic intended
Inj 0.9%, bag - Up to 2000 ml available on a PSO	1.33	500 ml	✓	Baxter
,	1.36	1,000 ml	✓	Baxter
Only if prescribed on a prescription for renal dialysis, ma for emergency use. (500 ml and 1,000 ml packs)	aternity or post-nat	al care in th	e home	e of the patient, or on a PSC
Inj 23.4% (4 mmol/ml), 20 ml ampoule		5	✓	Biomed
For Sodium chloride oral liquid formulation refer Standar		267	_	
Inj 0.9%, 5 ml ampoule – Up to 5 inj available on a PSO		20		Fresenius Kabi
Inj 0.9%, 10 ml ampoule – Up to 5 inj available on a PSO		50		Fresenius Kabi
Inj 0.9%, 20 ml ampoule	5.00	20	•	Fresenius Kabi
FOTAL PARENTERAL NUTRITION (TPN) Infusion	CBS	1 OP	/	TPN
WATER				
2) On a bulk supply order; or 3) When used in the extemporaneous compounding of ey 4) When used for the dilution of sodium chloride soln 7% Inj 10 ml ampoule – Up to 5 inj available on a PSO	for cystic fibrosis p	patients only	_	
Inj 20 ml ampoule – Up to 5 inj available on a PSO	5.00			Multichem
		20		Multichem Fresenius Kabi
Oral Administration		20		
		20		
		20 300 g OP	•	
CALCIUM POLYSTYRENE SULPHONATE Powder			•	Fresenius Kabi
CALCIUM POLYSTYRENE SULPHONATE Powder	169.85		,	Fresenius Kabi
CALCIUM POLYSTYRENE SULPHONATE PowderCOMPOUND ELECTROLYTES Powder for oral soln — Up to 5 sach available on a PSO	169.85	300 g OP	,	Fresenius Kabi Calcium Resonium
CALCIUM POLYSTYRENE SULPHONATE Powder COMPOUND ELECTROLYTES Powder for oral soln — Up to 5 sach available on a PSO COMPOUND ELECTROLYTES WITH GLUCOSE [DEXTROSE]	169.85	300 g OP	<i>'</i>	Fresenius Kabi Calcium Resonium Electral
CALCIUM POLYSTYRENE SULPHONATE PowderCOMPOUND ELECTROLYTES Powder for oral soln — Up to 5 sach available on a PSO	169.85	300 g OP	<i>'</i>	Fresenius Kabi Calcium Resonium Electral Hydralyte -
CALCIUM POLYSTYRENE SULPHONATE Powder COMPOUND ELECTROLYTES Powder for oral soln — Up to 5 sach available on a PSO COMPOUND ELECTROLYTES WITH GLUCOSE [DEXTROSE] Soln with electrolytes	169.85	300 g OP	<i>'</i>	Fresenius Kabi Calcium Resonium Electral
CALCIUM POLYSTYRENE SULPHONATE Powder COMPOUND ELECTROLYTES Powder for oral soln — Up to 5 sach available on a PSO COMPOUND ELECTROLYTES WITH GLUCOSE [DEXTROSE] Soln with electrolytes	169.85 9.53	300 g OP 50 1,000 ml Ol	<i>y y y y y y y y y y</i>	Calcium Resonium Electral Hydralyte - Lemonade
CALCIUM POLYSTYRENE SULPHONATE Powder	169.85 9.53	300 g OP	<i>y y y y y y y y y y</i>	Fresenius Kabi Calcium Resonium Electral Hydralyte -
CALCIUM POLYSTYRENE SULPHONATE Powder COMPOUND ELECTROLYTES Powder for oral soln - Up to 5 sach available on a PSO COMPOUND ELECTROLYTES WITH GLUCOSE [DEXTROSE] Soln with electrolytes PHOSPHORUS Tab eff 500 mg (16 mmol)	169.85 9.53 6.53	300 g OP 50 1,000 ml OI	<i>y y y y y y y y y y</i>	Fresenius Kabi Calcium Resonium Electral Hydralyte - Lemonade
CALCIUM POLYSTYRENE SULPHONATE Powder		300 g OP 50 1,000 ml Ol	<i>y y y y y y y y y y</i>	Calcium Resonium Electral Hydralyte - Lemonade Phosphate Phebra
CALCIUM POLYSTYRENE SULPHONATE Powder COMPOUND ELECTROLYTES Powder for oral soln — Up to 5 sach available on a PSO COMPOUND ELECTROLYTES WITH GLUCOSE [DEXTROSE] Soln with electrolytes PHOSPHORUS Tab eff 500 mg (16 mmol) POTASSIUM CHLORIDE * Tab eff 548 mg (14 m eq) with chloride 285 mg (8 m eq)		300 g OP 50 1,000 ml Ol 100 60	\tag{2}	Calcium Resonium Electral Hydralyte - Lemonade Phosphate Phebra Chlorvescent
CALCIUM POLYSTYRENE SULPHONATE Powder		300 g OP 50 1,000 ml OI	\tag{2}	Calcium Resonium Electral Hydralyte - Lemonade Phosphate Phebra
CALCIUM POLYSTYRENE SULPHONATE Powder COMPOUND ELECTROLYTES Powder for oral soln — Up to 5 sach available on a PSO COMPOUND ELECTROLYTES WITH GLUCOSE [DEXTROSE] Soln with electrolytes PHOSPHORUS Tab eff 500 mg (16 mmol) POTASSIUM CHLORIDE * Tab eff 548 mg (14 m eq) with chloride 285 mg (8 m eq)		300 g OP 50 1,000 ml Ol 100 60		Calcium Resonium Electral Hydralyte - Lemonade Phosphate Phebra Chlorvescent

	Subsidy (Manufacturer's Price	ce) Subs	Fully sidised	Brand or Generic	
	\$	Per	✓	Manufacturer	
SODIUM POLYSTYRENE SULPHONATE Powder	84.65	454 g OP	✓ R	Resonium-A	

	(Manufacturer's F	Price) Subs	idised	Generic
	\$	Per	·	Manufacturer
Alpha-Adrenoceptor Blockers				
Alpha Adrenoceptor Blockers				
DOXAZOSIN				
₭ Tab 2 mg		500		xazosin Clinect
★ Tab 4 mg	20.94	500	✓ Do	xazosin Clinect
PHENOXYBENZAMINE HYDROCHLORIDE				
₭ Cap 10 mg	65.00	30	✓ BI	M S29
	216.67	100	✓ Di	benzyline S29
PRAZOSIN				
米 Tab 1 mg	5.53	100	✓ Ar	rotex-Prazosin
			;	S29 S29
	9.98			nipress S29
* Tab 2 mg	7.00	100		rotex-Prazosin
			;	S29 S29
	13.29			nipress S29
* Tab 5 mg	11.70	100		rotex-Prazosin
				S29 S29
	22.00			nipress S29
* Cap 1 mg	15.40	100	✓ Pr	azosin Mylan S29
* Cap 2 mg	15.58	100	✓ Pr	azosin Mylan S29
* Cap 5 mg	23.32	100	✓ Pr	azosin Mylan S29
Agents Affecting the Renin-Angiotensin Syst ACE Inhibitors	(CIII			
CAPTOPRIL				
* Oral liq 5 mg per ml Oral liquid restricted to children under 12 years of ag		100 ml OP	✓ <u>DF</u>	P-Captopril
CILAZAPRIL - Subsidy by endorsement				
Subsidy by endorsement – Subsidised for patients who wendorsed accordingly. Pharmacists may annotate the pridispensing of cilazapril.	• •			
* Tab 0.5 mg	2.69	90	✓ Za	pril
★ Tab 2.5 mg	5.79	90	✓ Za	pril
Tab 5 mg	10.05	90	✓ Za	pril
ENALAPRIL MALEATE				
* Tab 5 mg	1.75	90	✓ Ac	etec
* Tab 10 mg		90	✓ <u>Ac</u>	
* Tab 20 mg	2.35	90	✓ <u>Ac</u>	etec
LISINOPRIL			_	
* Tab 5 mg	11.07	90		hics Lisinopril va Lisinopril
* Tab 10 mg	11.67	90	✓ Et	hics Lisinopril va Lisinopril
* Tab 20 mg	14.69	90	✓ Et	hics Lisinopril
			▼ 10	va Lisinopril

Subsidy

Fully

Brand or

	Subsidy (Manufacturer's Price) \$	Per	Fully Brand Subsidised Gene Manu	
PERINDOPRIL				
* Tab 2 mg	1.58	30	✓ Covers	γl
* Tab 4 mg		30	✓ Covers	
* Tab 8 mg	5.02	30	✓ Covers	
QUINAPRIL				
* Tab 5 mg	5.97	90	✓ Arrow-	Quinapril 5
* Tab 10 mg		90		Quinapril 10
* Tab 20 mg		90		Quinapril 20
RAMIPRIL				
* Cap 1.25 mg	6.90	90	✓ Tryzan	
* Cap 2.5 mg		90	✓ Tryzan	
* Cap 5 mg		90	✓ Tryzan	
* Cap 10 mg		90	✓ Tryzan	
ACE Inhibitors with Diuretics				
Subsidy by endorsement – Subsidised for patients who were 2022 and the prescription is endorsed accordingly. Pharma exists a record of prior dispensing of quinapril with hydrochle Tab 10 mg with hydrochlorothiazide 12.5 mg	cists may annotate the prothiazide. 4.10			ed where there
Angiotensin II Antagonists				
CANDESARTAN CILEXETIL				
* Tab 4 mg	2.00	90	✓ Candes	tar
* Tab 8 mg	2.28	90	✓ Candes	tar
* Tab 16 mg	3.31	90	✓ Candes	tar
* Tab 32 mg	5.26	90	✓ Candes	tar
LOSARTAN POTASSIUM				
* Tab 12.5 mg	2.00	84	✓ Losarta	n Actavis
* Tab 25 mg	2.29	84	✓ Losarta	n Actavis
* Tab 50 mg	2.86	84	✓ Losarta	n Actavis
* Tab 100 mg	4.57	84	✓ Losarta	n Actavis
Angiotensin II Antagonists with Diuretics				
CANDESARTAN CILEXETIL WITH HYDROCHLOROTHIAZIDE				
* Tab 16 mg with hydrochlorothiazide 12.5 mg		30		andesartan
	_		_	16/12.5
* Tab 32 mg with hydrochlorothiazide 12.5 mg	5.25	30		andesartan
			HCTZ	32/12.5
LOSARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE				

✓ Arrow-Losartan &

Hydrochlorothiazide

30

* Tab 50 mg with hydrochlorothiazide 12.5 mg......4.00

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Por 🗸	Manufacturer

Angiotensin II Antagonists with Neprilysin Inhibitors

SACUBITRIL WITH VALSARTAN - Special Authority see S	A2302 below - Retail p	harmacy	
Tab 24.3 mg with valsartan 25.7 mg	190.00	56	✓ Entresto 24/26
Tab 48.6 mg with valsartan 51.4 mg	190.00	56	✓ Entresto 49/51
Tab 97.2 mg with valsartan 102.8 mg	190.00	56	✓ Entresto 97/103

⇒SA2302 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has heart failure; and
- 2 Any of the following:
 - 2.1 Patient is in NYHA/WHO functional class II; or
 - 2.2 Patient is in NYHA/WHO functional class III; or
 - 2.3 Patient is in NYHA/WHO functional class IV; and
- 3 Either:
 - 3.1 Patient has a documented left ventricular ejection fraction (LVEF) of less than or equal to 35%; or
 - 3.2 An ECHO is not reasonably practical, and in the opinion of the treating practitioner the patient would benefit from treatment; and
- 4 Patient is receiving concomitant optimal standard chronic heart failure treatments.

For lignocaine hydrochloride refer to NERVOUS SYSTEM, Anaesthetics, Local, page 122

Antiarrhythmics

To lightcame hydrochloride relet to NETTVOOS STOTEM, Anaesthetics, Local,	page 122	
AMIODARONE HYDROCHLORIDE		
▲ Tab 100 mg	30	✓ Aratac
▲ Tab 200 mg4.49	30	✓ Aratac
Inj 50 mg per ml, 3 ml ampoule - Up to 10 inj available on a PSO 9.12	6	✓ Cordarone-X
15.22	10	✓ Max Health
ATROPINE SULPHATE		
* Inj 600 mcg per ml, 1 ml ampoule - Up to 5 inj available on a		
PSO15.09	10	✓ Martindale
DIGOXIN		
* Tab 62.5 mcg – Up to 30 tab available on a PSO	240	✓ Lanoxin PG
* Tab 250 mcg – Up to 30 tab available on a PSO16.90	240	✓ Lanoxin
* Oral liq 50 mcg per ml	60 ml	✓ Lanoxin
		✓ Lanoxin Paediatric Elixir
		✓ Lanoxin S29 S29
DISOPYRAMIDE PHOSPHATE		
▲ Cap 100 mg20.05	84	✓ Rythmodan - Cheplafarm \$29
23.87	100	✓ Rythmodan

Medsurge

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	Generic
	\$	Per		Manufacturer
FLECAINIDE ACETATE				
▲ Tab 50 mg	19.95	60	✓	Flecainide BNM
			✓	Flecatab S29
▲ Cap long-acting 100 mg	35.78	90	✓	<u>Flecainide</u>
				Controlled
				Release Teva
▲ Cap long-acting 200 mg	54.28	90	✓	Flecainide
				Controlled
				Release Teva
Inj 10 mg per ml, 15 ml ampoule	108.16	5		Tambocor
MEXILETINE HYDROCHLORIDE				
▲ Cap 150 mg	162.00	100	✓	Teva S29
▲ Cap 250 mg	202.00	100	1	Teva S29
PROPAFENONE HYDROCHLORIDE				
▲ Tab 150 mg	40.90	50	1	Rytmonorm
_ : 25 : 55 : 19		•••	'	,
Antihypotensives				
• •				
MIDODRINE - Special Authority see SA1474 below - Retail pha	armacy			
Tab 2.5 mg	38.23	100		MAR-Midodrine S29
			✓	<u>Midodrine</u>
				<u>Medsurge</u>
Tab 5 mg	59.98	100	✓	<u>Midodrine</u>

⇒SA1474 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years where patient has disabling orthostatic hypotension not due to drugs.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Beta-Adrenoceptor Blockers

Beta Adrenoceptor Blockers

ENOLOL			
Tab 50 mg	9.33	500	✓ Viatris
Tab 100 mg	14.20	500	✓ Atenolol Viatris
•			Mylan Atenolol
Oral liq 25 mg per 5 ml	21.25	300 ml OP	✓ Atenolol AFT
, -,			S29 S29
	38.20		✓ Essential
			Generics S29
	49.85		✓ Atenolol AFT
	Tab 50 mg	Tab 50 mg 9.33 Tab 100 mg 14.20 Oral liq 25 mg per 5 ml 21.25 38.20 49.85	Tab 50 mg 9.33 500 Tab 100 mg 14.20 500 Oral liq 25 mg per 5 ml 21.25 300 ml OP 38.20 49.85

Restricted to children under 12 years of age.

(Mylan Atenolol Tab 100 mg to be delisted 1 July 2024)

(Atenolol AFT S29 S29 Oral liq 25 mg per 5 ml to be delisted 1 August 2024)

(Essential Generics S29 Oral lig 25 mg per 5 ml to be delisted 1 August 2024)

	Subsidy (Manufacturer's Price)		Fully Brand or Subsidised Generic
	\$	Per	✓ Manufacturer
BISOPROLOL FUMARATE			
★ Tab 2.5 mg	1.36	90	✓ <u>Ipca-Bisoprolol</u>
米 Tab 5 mg	1.91	90	✓ <u>Ipca-Bisoprolol</u>
米 Tab 10 mg	2.71	90	✓ Ipca-Bisoprolol
CARVEDILOL			
* Tab 6.25 mg	2.24	60	✓ Carvedilol Sandoz
* Tab 12.5 mg	2.30	60	✓ Carvedilol Sandoz
* Tab 25 mg	2.95	60	✓ Carvedilol Sandoz
ABETALOL			
* Tab 100 mg	14.50	100	✓ Trandate
* Tab 200 mg		100	✓ Trandate
* Inj 5 mg per ml, 20 ml ampoule		5	
	(88.60)		Trandate
inj 5 mg per ml, 20 ml vial	42.29	1	
, 01	(48.20)		Alvogen S29
Alvogen 👀 inj 5 mg per ml, 20 ml vial to be de	, ,		ŭ
METOPROLOL SUCCINATE	,		
# Tab long-acting 23.75 mg	4.20	90	✓ Myloc CR
* Tab long-acting 23.75 mg*		90	✓ Myloc CR
* Tab long acting 47.5 mg		90	✓ Myloc CR
* Tab long acting 33 mg*		90	✓ Myloc CR
METOPROLOL TARTRATE		00	inyloo ori
	F 66	100	/ IDCA Meterrelel
★ Tab 50 mg		100	✓ IPCA-Metoprolol
* Tab 50 mg* Tab 100 mg	7.55	60	✓ IPCA-Metoprolol
 ★ Tab 50 mg ★ Tab 100 mg ★ Tab long-acting 200 mg 	7.55 23.40	60 28	✓ IPCA-Metoprolol ✓ Slow-Lopresor
Tab 50 mg Tab 100 mg Tab long-acting 200 mg	7.55 23.40	60	✓ <u>IPCA-Metoprolol</u> ✓ Slow-Lopresor ✓ Metoprolol IV Mylan
Tab 50 mg Tab 100 mg Tab long-acting 200 mg Inj 1 mg per ml, 5 ml vial	7.55 23.40	60 28	✓ <u>IPCA-Metoprolol</u> ✓ Slow-Lopresor ✓ Metoprolol IV Mylan
★ Tab 50 mg ★ Tab 100 mg ★ Tab long-acting 200 mg ★ Inj 1 mg per ml, 5 ml vial		60 28 5	✓ IPCA-Metoprolol ✓ Slow-Lopresor ✓ Metoprolol IV Mylan ✓ Metoprolol IV Viatris
★ Tab 50 mg ★ Tab 100 mg ★ Tab long-acting 200 mg ★ Inj 1 mg per ml, 5 ml vial VADOLOL ★ Tab 40 mg		60 28 5	✓ IPCA-Metoprolol ✓ Slow-Lopresor ✓ Metoprolol IV Mylan ✓ Metoprolol IV Viatris ✓ Nadolol BNM
* Tab 50 mg * Tab 100 mg * Tab long-acting 200 mg * Inj 1 mg per ml, 5 ml vial NADOLOL * Tab 40 mg * Tab 80 mg		60 28 5	✓ IPCA-Metoprolol ✓ Slow-Lopresor ✓ Metoprolol IV Mylan ✓ Metoprolol IV Viatris
★ Tab 50 mg ★ Tab 100 mg ★ Tab long-acting 200 mg ★ Inj 1 mg per ml, 5 ml vial NADOLOL ★ Tab 40 mg		60 28 5	✓ IPCA-Metoprolol ✓ Slow-Lopresor ✓ Metoprolol IV Mylan ✓ Metoprolol IV Viatris ✓ Nadolol BNM
* Tab 50 mg * Tab 100 mg * Tab long-acting 200 mg * Inj 1 mg per ml, 5 ml vial NADOLOL * Tab 40 mg * Tab 80 mg PROPRANOLOL * Tab 10 mg		60 28 5 100 100	✓ IPCA-Metoprolol ✓ Slow-Lopresor ✓ Metoprolol IV Mylan ✓ Metoprolol IV Viatris ✓ Nadolol BNM ✓ Nadolol BNM ✓ Drofate
R Tab 50 mg Tab 100 mg Tab long-acting 200 mg Inj 1 mg per ml, 5 ml vial NADOLOL Tab 40 mg Tab 80 mg PROPRANOLOL Tab 10 mg Tab 40 mg		60 28 5 100 100	✓ IPCA-Metoprolol ✓ Slow-Lopresor ✓ Metoprolol IV Mylan ✓ Metoprolol IV Viatris ✓ Nadolol BNM ✓ Nadolol BNM ✓ Drofate ✓ IPCA-Propranolol
* Tab 50 mg * Tab 100 mg * Tab long-acting 200 mg * Inj 1 mg per ml, 5 ml vial NADOLOL * Tab 40 mg PROPRANOLOL * Tab 10 mg * Tab 40 mg * Cap long-acting 160 mg	7.55 23.40 26.50 26.50 19.19 30.39 7.04 8.75 18.17	60 28 5 100 100	✓ IPCA-Metoprolol ✓ Slow-Lopresor ✓ Metoprolol IV Mylan ✓ Metoprolol IV Viatris ✓ Nadolol BNM ✓ Nadolol BNM ✓ Drofate
R Tab 50 mg R Tab 100 mg R Tab long-acting 200 mg Inj 1 mg per ml, 5 ml vial NADOLOL R Tab 40 mg PROPRANOLOL R Tab 10 mg R Tab 40 mg	7.55 23.40 26.50 26.50 19.19 30.39 7.04 8.75 18.17	60 28 5 100 100	✓ IPCA-Metoprolol ✓ Slow-Lopresor ✓ Metoprolol IV Mylan ✓ Metoprolol IV Viatris ✓ Nadolol BNM ✓ Nadolol BNM ✓ Drofate ✓ IPCA-Propranolol ✓ Cardinol LA
* Tab 50 mg * Tab 100 mg * Tab long-acting 200 mg * Inj 1 mg per ml, 5 ml vial NADOLOL * Tab 40 mg PROPRANOLOL * Tab 10 mg * Tab 40 mg * Cap long-acting 160 mg	7.55 23.40 26.50 19.19 30.39 7.04 8.75 18.17 SA1327 below –	60 28 5 100 100	✓ IPCA-Metoprolol ✓ Slow-Lopresor ✓ Metoprolol IV Mylan ✓ Metoprolol IV Viatris ✓ Nadolol BNM ✓ Nadolol BNM ✓ Drofate ✓ IPCA-Propranolol ✓ Cardinol LA

⇒SA1327 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

		, A I I I	DIOVAGGOLATIOTOTEM
	Subsidy (Manufacturer's Price) \$	Per	Fully Brand or Subsidised Generic Manufacturer
SOTALOL * Tab 80 mg * Tab 160 mg		500 100	✓ <u>Mylan</u> ✓ <u>Mylan</u>
Calcium Channel Blockers			
Dihydropyridine Calcium Channel Blockers			
AMLODIPINE			
* Tab 2.5 mg	1.45	90	✓ <u>Vasorex</u>
* Tab 5 mg	1.21	90	✓ <u>Vasorex</u>
* Tab 10 mg	1.31	90	✓ <u>Vasorex</u>
FELODIPINE			
* Tab long-acting 2.5 mg	1.45	30	✓ Plendil ER
* Tab long-acting 5 mg		90	✓ Felo 5 ER
* Tab long-acting 10 mg	4.32	90	✓ Felo 10 ER
NIFEDIPINE			
* Tab long-acting 10 mg - Subsidy by endorsement	19.42	56	✓ Tensipine MR10 S29
Tab long doing to mg Gabsiay by chaolsement	10.72	50	Tensipine witto
endorsed accordingly. Pharmacists may annotate the predispensing of nifedipine tab long-acting 10 mg. * Tab long-acting 20 mg* Tab long-acting 30 mg	17.72	ed wh 100 14	✓ Nyefax Retard ✓ Mylan Italy (24 hr release) \$29
	10.24	30	✓ Nifedipine Viatris S29
	34.10	100	✓ Mylan (24 hr release) S29
* Tab long-acting 60 mg	52.81	100	✓ Mylan (24 hr release) \$29
Other Calcium Channel Blockers			
DILTIAZEM HYDROCHLORIDE			
* Cap long-acting 120 mg	65.35	500	✓ Diltiazem CD Clinect
* Cap long-acting 180 mg		30	✓ Cardizem CD
* Cap long-acting 240 mg		30	✓ Cardizem CD
PERHEXILINE MALEATE			
* Tab 100 mg	62 90	100	✓ Pexsig
	02.00	100	- I chaig
VERAPAMIL HYDROCHLORIDE	7.01	100	√ loantin
* Tab 90 mg		100 100	✓ Isoptin
* Tab long acting 100 mg			✓ Isoptin
* Tab long-acting 120 mg	30.02	100	✓ Isoptin Retard S29
* Tab long-acting 240 mg	15.10	30	✓ Isoptin SR ✓ Isoptin SR
	12.12	30	✓ Isoptin SR
* Inj 2.5 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO	25.00	5	✓ Isoptin

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy		Fully	Brand or
	(Manufacturer's Price		Subsidised	Generic
	\$	Per		Manufacturer
Centrally-Acting Agents				
CLONIDINE				
* Patch 2.5 mg, 100 mcg per day - Only on a prescription	11.70	4	✓ [<u>Mylan</u>
* Patch 5 mg, 200 mcg per day - Only on a prescription		4		<u>Mylan</u>
* Patch 7.5 mg, 300 mcg per day - Only on a prescription	17.90	4	✓ [<u>Mylan</u>
CLONIDINE HYDROCHLORIDE				
* Tab 25 mcg		112		Clonidine Teva
* Tab 150 mcg		100		<u>Catapres</u>
* Inj 150 mcg per ml, 1 ml ampoule	29.68	10	•	<u>Medsurge</u>
METHYLDOPA				
* Tab 250 mg	15.10	100		Methyldopa Mylan
	52.85	500		Methyldopa Viatris Methyldopa Mylan
	52.05	500	• 1	S29 S29
(Mathyldona Mylan Tah 250 mg to ha delicted 1 Sentember 202	4)			329 329
(Methyldopa Mylan Tab 250 mg to be delisted 1 September 202-	•			
(Methyldopa Mylan S29 S29 Tab 250 mg to be delisted 1 Septe	ember 2024)			
Diuretics				
Loop Diuretics				
BUMETANIDE				
* Tab 1 mg	A Q1	30	1	Burinex S29 S29
	16.36	100		Burinex
* Inj 500 mcg per ml, 4 ml vial		5		Burinex
(Burinex S29 S29 Tab 1 mg to be delisted 1 August 2024)				
FUROSEMIDE [FRUSEMIDE]				
Tab 40 mg - Up to 30 tab available on a PSO	8.00	1.000	✓	IPCA-Frusemide
* Tab 500 mg		50		Urex Forte
Č	89.48		✓	Furosemid-
				Ratiopharm \$29
dt 0 11 40 1	44.00			
* Oral liq 10 mg per ml		0 ml C		Lasix Lasix
 Inj 10 mg per ml, 25 ml ampoule Inj 10 mg per ml, 2 ml ampoule – Up to 5 inj available on a 		6 5		Furosemide-Baxter
(Furosemid-Ratiopharm \$29) Tab 500 mg to be delisted 1 Augu		5	• !	rui oseiiiiue-baxtei
rurosemio-naliophami 🥯 Tab 500 mg to be delisted T Augu	ISI 2024)			
Potassium Sparing Diuretics				
AMILORIDE HYDROCHLORIDE				
Oral liq 1 mg per ml	33.71 2	5 ml C)P 🗸 I	Biomed
EPLERENONE – Special Authority see SA1728 below – Retail	pharmacy			
Tab 25 mg		30	✓ [<u>Inspra</u>
Tab 50 mg	25.00	30	✓]	Inspra
⇒SA1728 Special Authority for Subsidy				
nitial application from any relevant practitioner. Approvals val	id without further ren	ewal ui	nless notifie	ed for applications meetir

continued...

		CARDIO	/ASCU	LAR SYSTEM
	Subsidy (Manufacturer's Pr \$	rice) Subsi Per	dised (Brand or Generic Manufacturer
continued the following criteria: Both: 1 Patient has heart failure with ejection fraction less than 4 2 Fither:	0%; and			
2.1 Patient is intolerant to optimal dosing of spironola2.2 Patient has experienced a clinically significant adv		on optimal dos	ng of spi	ronolactone.
SPIRONOLACTONE * Tab 25 mg Tab 100 mg Oral liq 5 mg per ml	10.65	100 100 25 ml OP	✓ <u>Spi</u> ✓ <u>Spi</u> ✓ Bio	ractin
Potassium Sparing Combination Diuretics				
AMILORIDE HYDROCHLORIDE WITH FUROSEMIDE * Tab 5 mg with furosemide 40 mg AMILORIDE HYDROCHLORIDE WITH HYDROCHLOROTHIAZ * Tab 5 mg with hydrochlorothiazide 50 mg	ZIDE	28 50	✓ Fru	
Thiazide and Related Diuretics				
BENDROFLUMETHIAZIDE [BENDROFLUAZIDE] * Tab 2.5 mg - Up to 150 tab available on a PSO	51.50	500	✓ <u>Arro</u>	<u>ow-</u> endrofluazide
May be supplied on a PSO for reasons other than emer * Tab 5 mg		500	✓ <u>Arro</u> <u>B</u>	<u>ow-</u> endrofluazide
CHLOROTHIAZIDE Oral liq 50 mg per ml CHLORTALIDONE [CHLORTHALIDONE]	29.21	25 ml OP	✓ Bio	med
* Tab 25 mg	6.95	50	✓ Hyc	<u>ıroton</u>

90

1

50

✓ Dapa-Tabs

✓ Metolazone \$29

✓ Zaroxolyn S29

Vasopressin	receptor	antagonists
-------------	----------	-------------

Tab 5 mgCBS

INDAPAMIDE

METOLAZONE

TOLVAPTAN - Special Authority see SA2166 on the next page	- Retail pharmacy	•	
Tab 15 mg	873.50	28 OP	Jinarc
Tab 30 mg	873.50	28 OP	Jinarc
Tab 45 mg + 15 mg	1,747.00	56 OP	Jinarc
Tab 60 mg + 30 mg	1,747.00	56 OP	Jinarc
Tab 90 mg + 30 mg		56 OP	Jinarc

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	
<u> </u>	Per 🗸	Manufacturer

⇒SA2166 Special Authority for Subsidy

Initial application — (autosomal dominant polycystic kidney disease) only from a renal physician or any relevant practitioner on the recommendation of a renal physician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of autosomal dominant polycystic kidney disease; and
- 2 Patient has an estimated glomerular filtration rate (eGFR) of greater than or equal to 25 ml/min/1.73 m² at treatment initiation; and
- 3 Either:
 - 3.1 Patient's disease is rapidly progressing, with a decline in eGFR of greater than or equal to 5 mL/min/1.73 m² within one-year; or
 - 3.2 Patient's disease is rapidly progressing, with an average decline in eGFR of greater than or equal to 2.5 mL/min/1.73 m² per year over a five-year period.

Renewal — (autosomal dominant polycystic kidney disease) only from a renal physician or any relevant practitioner on the recommendation of a renal physician. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has not developed end-stage renal disease, defined as an eGFR of less than 15 mL/min/1.73 m²; and
- 2 Patient has not undergone a kidney transplant.

Lipid-Modifying Agents		
Fibrates		
BEZAFIBRATE * Tab 200 mg	90 30	✓ <u>Bezalip</u>✓ <u>Bezalip</u> Retard
Other Lipid-Modifying Agents		
ACIPIMOX * Cap 250 mg21.56 25.44 (Olbetam S29 S29 Cap 250 mg to be delisted 1 August 2024)	30	✓ Olbetam S29 S29 ✓ Olbetam
Resins		
COLESTIPOL HYDROCHLORIDE Grans for oral liq 5 g32.89 (Colestid Grans for oral liq 5 g to be delisted 1 August 2024) COLESTYRAMINE	30	✓ Colestid
Powder for oral suspension 4 g sachet	50	✓ Colestyramine - Mylan S29 ✓ Quantalan sugar free S29
HMG CoA Reductase Inhibitors (Statins)		
ATORVASTATIN * Tab 10 mg	500 500	✓ Lorstat ✓ Lorstat

Lorstat

Lorstat

500

500

	Subsidy (Manufacturer's Price \$) Per	Fully Subsidised	Generic
PRAVASTATIN				
* Tab 20 mg	7.16	100	✓	Clinect
* Tab 40 mg	12.25	100	1	Clinect
ROSUVASTATIN - Special Authority see SA2093 below - Retail	pharmacy			
* Tab 5 mg	1.29	30	1	Rosuvastatin Viatris
Rosuvastatin Viatris to be Principal Supply on 1 October	2024			
* Tab 10 mg	1.69	30	1	Rosuvastatin Viatris
Rosuvastatin Viatris to be Principal Supply on 1 October	2024			
* Tab 20 mg	2.71	30	1	Rosuvastatin Viatris
* Tab 40 mg	4.55	30	✓	Rosuvastatin Viatris
- CACCOO Createl Authority for Cubaids				

⇒SA2093 Special Authority for Subsidy

Initial application — (cardiovascular disease risk) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient is considered to be at risk of cardiovascular disease; and
 - 1.2 Patient is Māori or any Pacific ethnicity; or
- 2 Both:
 - 2.1 Patient has a calculated risk of cardiovascular disease of at least 15% over 5 years; and
 - 2.2 LDL cholesterol has not reduced to less than 1.8 mmol/litre with treatment with the maximum tolerated dose of atoryastatin and/or simyastatin.

Initial application — (familial hypercholesterolemia) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has familial hypercholesterolemia (defined as a Dutch Lipid Criteria score greater than or equal to 6); and
- 2 LDL cholesterol has not reduced to less than 1.8 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initial application — (established cardiovascular disease) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:
Both:

- 1 Any of the following:
 - 1.1 Patient has proven coronary artery disease (CAD); or
 - 1.2 Patient has proven peripheral artery disease (PAD); or
 - 1.3 Patient has experienced an ischaemic stroke; and
- 2 LDL cholesterol has not reduced to less than 1.4 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initial application — (recurrent major cardiovascular events) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has experienced a recurrent major cardiovascular event (defined as myocardial infarction, ischaemic stroke, coronary revascularisation, hospitalisation for unstable angina) in the last 2 years; and
- 2 LDL cholesterol has not reduced to less than 1.0 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

_	Subsidy	,	Fully	
(I	Manufacturer's Pric \$	e) Per	Subsidised •	
IMVASTATIN				
€ Tab 10 mg	1.68	90		Simvastatin Mylan Simvastatin Viatris
★ Tab 20 mg	2.54	90	/	Simvastatin Viatris
← Tab 40 mg	4.11	90	1	Simvastatin Mylan
			•	Simvastatin Viatris
F Tab 80 mg	8.81	90		Simvastatin Mylan Simvastatin Viatris
Simvastatin Mylan Tab 40 mg to be delisted 1 December 2024) Simvastatin Mylan Tab 80 mg to be delisted 1 September 2024)				
Selective Cholesterol Absorption Inhibitors				
ZETIMIBE				
	1.76	30	1	Ezemibe Viatris
•			1	Ezetimibe Sandoz
ZETIMIBE WITH SIMVASTATIN Tab 10 mg with simvastatin 10 mg	5 15	30	1	Zimybe
Tab 10 mg with simvastatin 10 mg		30		Zimybe
Tab 10 mg with simvastatin 40 mg		30	_	Zimybe
Tab 10 mg with simvastatin 80 mg		30		Zimybe
Nitrates				
GLYCERYL TRINITRATE				
♦ Oral pump spray, 400 mcg per dose – Up to 250 dose				
available on a PSO	7.48 2	50 dose	OP 🗸	Nitrolingual Pump
k Datah 05 ma 5 ma parday	15 70	30	./	Spray Nitroderm TTS
 Patch 25 mg, 5 mg per day Patch 50 mg, 10 mg per day 		30		Nitroderm TTS
	10.02	30	•	Milloueilli 113
SOSORBIDE MONONITRATE	00.40	400	,	I
≰ Tab 20 mg		100		Ismo 20
Tab long-acting 40 mg		30	_	Ismo 40 Retard
Tab long-acting 60 mg	13.50	90	•	<u>Duride</u>
Sympathomimetics				
DRENALINE				
Inj 1 in 1,000, 1 ml ampoule - Up to 5 inj available on a PSO	4.98	5		Aspen Adrenaline
	12.65		1	DBL Adrenaline
Inj 1 in 10,000, 10 ml ampoule - Up to 5 inj available on a PSC	27.00	5	_	Hospira
	49.00	10	1	Aspen Adrenaline
Vasodilators				
YDRALAZINE HYDROCHLORIDE				
Tab 25 mg - Special Authority see SA1321 on the next page -	_			
Retail pharmacy		1	1	Hydralazine
•		56		Onelink \$29
		84		AMDIPHARM \$29
		100		Camber \$29
k Inj 20 mg ampoule	25 90	5	_	Apresoline
- iij 20 iiig aiiipoalo	20.00	J	•	Aprogonine

✓ Minovidil Roma \$29

Subsidy (Manufacturer's Price)	Subs	Fully	Brand or Generic
\$	Per	√	Manufacturer

⇒SA1321 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 For the treatment of refractory hypertension; or
- 2 For the treatment of heart failure in combination with a nitrate, in patients who are intolerant or have not responded to ACE inhibitors and/or angiotensin receptor blockers.

47 N4

60

MINOXIDIL A Tab 10 mg

	78.40	100	✓ Loniten
NICORANDIL - Brand switch fee payable (Pharmacode 267790	03) - see page 264	for details	
▲ Tab 10 mg	21.73	60	✓ Max Health
▲ Tab 20 mg	27.44	60	✓ Max Health
PAPAVERINE HYDROCHLORIDE			
* Inj 12 mg per ml, 10 ml ampoule	257.12	5	✓ Hospira
PENTOXIFYLLINE [OXPENTIFYLLINE]			
Tab 400 mg	42.26	50	✓ Trental 400

Endothelin Receptor Antagonists

AMBRISENTAN - Special Authority see SA2253 below -	Retail pharmacy			
Tab 5 mg	200.00	30	•	Ambrisentan Viatris
Tab 10 mg	200.00	30	•	Ambrisentan Viatris

⇒SA2253 Special Authority for Subsidy

Initial application — (PAH monotherapy) only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II. III or IV: and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁻⁵); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH 2022 (see note below for link to these guidelines) †; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
 - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
 - 5.1 Ambrisentan is to be used as PAH monotherapy; and
 - 5.2 Any of the following:
 - 5.2.1 Patient has experienced intolerable side effects with both sildenafil and bosentan; or
 - 5.2.2 Patient has an absolute contraindication to sildenafil and an absolute or relative contraindication to bosentan (e.g. due to current use of a combined oral contraceptive or liver disease); or
 - 5.2.3 Patient is a child with idiopathic PAH or PAH secondary to congenital heart disease.

Initial application — **(PAH dual therapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II. III or IV: and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁵); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH 2022 (see note below for link to these guidelines) †: or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**: or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
 - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
 - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 All of the following:
 - 5.1 Ambrisentan is to be used as PAH dual therapy; and
 - 5.2 Either:
 - 5.2.1 Patient has tried a PAH monotherapy (sildenafil or bosentan) for at least three months and has not experienced an acceptable response to treatment according to a validated risk stratification tool**; or
 - 5.2.2 Patient has tried PAH dual therapy including bosentan and has experienced intolerable side effects on bosentan; and
 - 5.3 Both:
 - 5.3.1 Patient is presenting in NYHA/WHO functional class III or IV, and in the opinion of the treating clinician would benefit from initial dual therapy; and
 - 5.3.2 Patient has an absolute or relative contraindication to bosentan (e.g. due to current use of a combined oral contraceptive or liver disease).

Initial application — (PAH triple therapy) only from a respiratory specialist, cardiologist, rheumatologist or any relevant

Subsidy (Manufacturer's Price)	Subsi	Fully dised	Brand or Generic
 \$	Per	•	Manufacturer

continued...

practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁻⁵); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH 2022 (see note below for link to these guidelines) †; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
 - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
 - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
 - 5.1 Ambrisentan is to be used as PAH triple therapy; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is on the lung transplant list; or
 - 5.2.2 Both:
 - 5.2.2.1 Patient is presenting in NYHA/WHO functional class IV: and
 - 5.2.2.2 Patient has an absolute or relative contraindication to bosentan (e.g. due to current use of a combined oral contraceptive or liver disease); or
 - 5.2.3 Both:
 - 5.2.3.1 Patient has tried PAH dual therapy for at least three months and remains in an unacceptable risk category according to a validated risk stratification tool**; and
 - 5.2.3.2 Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario.

Renewal only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 2 years where the patient is continuing to derive benefit from ambrisentan treatment according to a validated PAH risk stratification tool**.

Notes: † The European Respiratory Journal Guidelines can be found here: 2022 ECS/ERS Guidelines for the diagnosis and treatment of pulmonary hypertension PAH

** the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

BOSENTAN - Special Authority see SA2254 on the next page	ge - Retail pharmacy		
Tab 62.5 mg	119.85	60	✓ Bosentan Dr
			Reddy's
Tab 125 mg	119.85	60	✓ Bosentan Dr
			Reddy's

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

⇒SA2254 Special Authority for Subsidy

Initial application — **(PAH monotherapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁵); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) †; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**: or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
 - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or
 - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
 - 5.1 Bosentan is to be used as PAH monotherapy; and
 - 5.2 Any of the following:
 - 5.2.1 Patient has experienced intolerable side effects on sildenafil; or
 - 5.2.2 Patient has an absolute contraindication to sildenafil: or
 - 5.2.3 Patient is a child with idiopathic PAH or PAH secondary to congenital heart disease.

Initial application — (PAH dual therapy) only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁻⁵); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or

Subsidy		Fully	Brand or	
(Manufacturer's Pric	ce)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) †; or
- 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
- 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
- 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease: or
- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Bosentan is to be used as part of PAH dual therapy; and
- 6 Fither:
 - 6.1 Patient has tried a PAH monotherapy (sildenafil) for at least three months and has experienced an inadequate therapeutic response to treatment according to a validated risk stratification tool**; or
 - 6.2 Patient is presenting in NYHA/WHO functional class III or IV, and in the opinion of the treating clinician would likely benefit from initial dual therapy.

Initial application — (PAH triple therapy) only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁻⁵); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these quidelines) †: or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
 - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or
 - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
 - 5.1 Bosentan is to be used as part of PAH triple therapy; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is on the lung transplant list; or
 - 5.2.2 Patient is presenting in NYHA/WHO functional class IV; or
 - 5.2.3 Both
 - 5.2.3.1 Patient has tried PAH dual therapy for at least three months and has not experienced an acceptable

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	sidised	Generic	
\$	Per	✓	Manufacturer	

continued...

response to treatment according to a validated risk stratification tool**; and

5.2.3.2 Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario.

Renewal only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 2 years where patient is continuing to derive benefit from bosentan treatment according to a validated PAH risk stratification tool**.

Notes: † The European Respiratory Journal Guidelines can be found here: 2022 ECS/ERS Guidelines for the diagnosis and treatment of pulmonary hypertension PAH

** the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

Phosphodiesterase Type 5 Inhibitors

SILDENAFIL - Special Authority see SA2255 below - Retail pharmacy		
Tab 25 mg	85 4	✓ Vedafil
Tab 50 mg	70 4	✓ Vedafil
Tab 100 mg10.2		✓ Vedafil

⇒SA2255 Special Authority for Subsidy

Initial application — (Raynaud's Phenomenon*) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has Raynaud's Phenomenon*; and
- 2 Patient has severe digital ischaemia (defined as severe pain requiring hospital admission or with a high likelihood of digital ulceration; digital ulcers; or gangrene); and
- 3 Patient is following lifestyle management (avoidance of cold exposure, sufficient protection, smoking cessation support, avoidance of sympathomimetic drugs); and
- 4 Patient is being treated with calcium channel blockers and nitrates (or these are contraindicated/not tolerated).

Initial application — (Pulmonary arterial hypertension*) only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*: and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II. III or IV: and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH is confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) of greater than 20 mmHg; and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) that is less than or equal to 15 mmHg; and
 - 4.1.4 Pulmonary vascular resistance (PVR) of at least 2 Wood Units or greater than 160 International Units (dyn s cm⁵); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH is non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) †; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or

Subsidy (Manufacturer's Price)	Fu Subsidis	,
\$	Per	✓ Manufacturer

continued...

- 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or
- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures.

Initial application — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has a documented history of traumatic or non-traumatic spinal cord injury; and
- 2 Patient has erectile dysfunction secondary to spinal cord injury requiring pharmacological treatment.

Renewal — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Notes: Note: Indications marked with * are Unapproved Indications.

† The European Respiratory Journal Guidelines can be found here: 2022 ECS/ERS Guidelines for the diagnosis and treatment of pulmonary hypertension PAH

** the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

Prostacyclin Analogues

		EPOPROSTENOL – Special Authority see SA2256 below – Retail pharmacy
✓ Veletri	1	Inj 500 mcg vial36.61
✓ Veletri	1	Inj 1.5 mg vial73.21

⇒SA2256 Special Authority for Subsidy

Initial application — (PAH dual therapy) only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class III or IV: and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁻⁵); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these quidelines) †: or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**: or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
 - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 All of the following:
 - 5.1 Epoprostenol is to be used as part of PAH dual therapy with either sildenafil or an endothelin receptor antagonist; and
 - 5.2 Patient is presenting in NYHA/WHO functional class IV; and
 - 5.3 Patient has tried a PAH monotherapy for at least three months and remains in an unacceptable risk category according to a validated risk stratification tool.

Initial application — **(PAH triple therapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁻⁵); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these quidelines) †: or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**: or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
 - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
 - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
 - 5.1 Epoprostenol is to be used as PAH triple therapy; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is on the lung transplant list; or
 - 5.2.2 Patient is presenting in NYHA/WHO functional class IV: or
 - 5.2.3 Both:
 - 5.2.3.1 Patient has tried PAH dual therapy for at least three months and has not experienced an acceptable response to treatment according to a validated risk stratification tool; and
 - 5.2.3.2 Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario.

Renewal only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 2 years where patient is continuing to derive benefit from epoprostenol treatment according to a validated PAH risk stratification tool**.

Notes: † The European Respiratory Journal Guidelines can be found here: 2022 ECS/ERS Guidelines for the diagnosis and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

treatment of pulmonary hypertension PAH

** the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

ILOPROST - Special Authority see SA2257 below - Retail pharmacy

⇒SA2257 Special Authority for Subsidy

Initial application — (PAH monotherapy) only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II. III or IV: and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁻⁵); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) † : or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
 - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
 - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and

5 Both:

- 5.1 Iloprost is to be used as PAH monotherapy; and
- 5.2 Either:
 - 5.2.1 Patient has experienced intolerable side effects on sildenafil and both the funded endothelin receptor antagonists (i.e. both bosentan and ambrisentan); or
 - 5.2.2 Patient has an absolute contraindication to sildenafil and an absolute or relative contraindication to endothelin receptor antagonists.

Initial application — **(PAH dual therapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:

				=
Subsidy	F	ully	Brand or	
(Manufacturer's Price)	Subsid	ised	Generic	
\$	Per	1	Manufacturer	

continued...

- 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁵); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) † : or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**: or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
- 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 All of the following:
 - 5.1 lloprost is to be used as PAH dual therapy with either sildenafil or an endothelin receptor antagonist; and
 - 5.2 Either:
 - 5.2.1 Patient has an absolute contraindication to or has experienced intolerable side effects on sildenafil; or
 - 5.2.2 Patient has an absolute or relative contraindication to or experienced intolerable side effects with a funded endothelin receptor antagonist; and
 - 5.3 Either:
 - 5.3.1 Patient has tried a PAH monotherapy for at least three months and remains in an unacceptable risk category according to a validated risk stratification tool**; or
 - 5.3.2 Patient is presenting in NYHA/WHO functional class III or IV, and in the opinion of the treating clinician would benefit from initial dual therapy.

Initial application — **(PAH triple therapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁵); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these quidelines) †; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a

()	Subsidy	Fully		Brand or
	Manufacturer's Price)	Subsidised		Generic
	\$	Per	✓	Manufacturer

continued...

validated risk stratification tool**; or

- 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
- 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
 - 5.1 Iloprost is to be used as PAH triple therapy; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is on the lung transplant list; or
 - 5.2.2 Patient is presenting in NYHA/WHO functional class IV; or
 - 5.2.3 Both:
 - 5.2.3.1 Patient has tried PAH dual therapy for at least three months and has not experienced an acceptable response to treatment according to a validated risk stratification tool**; and
 - 5.2.3.2 Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario.

Renewal only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 2 years where patient is continuing to derive benefit from iloprost treatment according to a validated PAH risk stratification tool**.

Notes: † The European Respiratory Journal Guidelines can be found here: 2022 ECS/ERS Guidelines for the diagnosis and treatment of pulmonary hypertension PAH

** the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer

Antiacne Preparations

For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 95

ADAPALENE

- a) Maximum of 30 g per prescription
- h) Only on a prescription

b) Only on a presemption			
Gel 0.1%	22.89	30 g OP	Differin
ISOTRETINOIN - Special Authority see SA2023 below - Re	etail pharmacy		
Cap 5 mg	11.26	60	Oratane
Cap 10 mg	18.75	120	✓ Oratane
Cap 20 mg	26.73	120	✓ Oratane

⇒SA2023 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around isotretinoin and is competent to prescribe isotretinoin; and
- 3 Either:
 - 3.1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and that they must not become pregnant during treatment and for a period of one month after the completion of treatment; or
 - 3.2 Patient is not of child bearing potential.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and that they must not become pregnant during treatment and for a period of one month after the completion of treatment: or
- 2 Patient is not of child bearing potential.

TRETINOIN

Crm 0.5 mg per g - Maximum of 50 g per prescription15.57 50 g OP ✓ ReTrieve

Antibacterials Topical

For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 95

HYDROGEN PEROXIDE	0.50	40 00	40
* Crm 1%	8.56	10 g OP	Crystaderm
MUPIROCIN			
Oint 2%	6.60	15 g OP	
	(13.00)	-	Bactroban

- a) Only on a prescription
- b) Not in combination

	Subsidy		Fully	Brand or
	(Manufacturer's Pri		Subsidised	
	\$	Per		Manufacturer
SODIUM FUSIDATE [FUSIDIC ACID]			_	
Crm 2%	1.59	5 g OP	•	<u>Foban</u>
a) Maximum of 5 g per prescription				
b) Only on a prescriptionc) Not in combination				
Oint 2%	1 59	5 g OP	1	Foban
a) Maximum of 5 g per prescription		o g o.		<u> </u>
b) Only on a prescription				
c) Not in combination				
SULFADIAZINE SILVER				
Crm 1%	10.80	50 g OF	•	Flamazine
a) Up to 250 g available on a PSO				
b) Not in combination				
Antifungals Topical				
For systemic antifungals, refer to INFECTIONS, Antifungals, pa	ne 102			
AMOROLFINE	.gc 102			
a) Only on a prescription				
b) Not in combination				
Nail soln 5%	21.87	5 ml OF	•	MycoNail
CLOTRIMAZOLE				
* Crm 1%	1.10	20 g OF	•	Clomazol
a) Only on a prescription		ŭ		
b) Not in combination				
* Soln 1%		20 ml Ol	•	
	(7.55)			Canesten
a) Only on a prescription				
b) Not in combination				
ECONAZOLE NITRATE	1.00	00 = 05	,	
Crm 1%	(8.09)	20 g OF	,	Pevaryl
a) Only on a prescription	(0.09)			i evaryi
b) Not in combination				
Foaming soln 1%, 10 ml sachets	9.89	3		
•	(18.64)			Pevaryl
a) Only on a prescription				
b) Not in combination				
MICONAZOLE NITRATE			_	
* Crm 2%	0.90	15 g OF	•	<u>Multichem</u>
a) Only on a prescription				
b) Not in combination	4.26	30 ml Ol		
* Lotn 2%	(10.03)	30 1111 01		Daktarin
a) Only on a prescription	(10.00)			Danum
b) Not in combination				
* Tinct 2%	4.36	30 ml Ol	>	
	(12.10)			Daktarin
a) Only on a prescription				
b) Not in combination				

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

69

DERMATOLOGICALS				
	Subsidy (Manufacturer's Pr \$	ice) Subs	Fully sidised	Brand or Generic Manufacturer
Antipruritic Preparations				
CALAMINE				
a) Only on a prescription				
b) Not in combination				
Crm, aqueous, BP	3.45	100 g	✓ h	ealthE Calamine Aqueous
CROTAMITON				
a) Only on a prescription				
b) Not in combination				
Crm 10%	3.29	20 g OP	✓ <u>I</u> t	ch-Soothe
MENTHOL - Only in combination				
 Only in combination with a dermatological base or p With or without other dermatological galenicals. 	proprietary Topical Co	orticosteriod –	Plain	
Crystals	6.92	25 g	✓ N	/lidWest
-,	29.60	100 g		lidWest
Corticosteroids Topical For systemic corticosteroids, refer to CORTICOSTEROIDS A	ND RELATED AGEN	ITS, page 85		
Corticosteroids - Plain				
BETAMETHASONE DIPROPIONATE				
Crm 0.05%		15 g OP		iprosone
D: D:	36.00	50 g OP	✓ [Diprosone
Diprosone to be Principal Supply on 1 July 2024	0.00	45 × OD	./ -	N
Oint 0.05%	36.00	15 g OP 50 g OP		Diprosone Diprosone
Diprosone to be Principal Supply on 1 July 2024	30.00	30 g Oi	• •	riprosorie
Oint 0.05% in propylene glycol base	4.33	30 g OP	√ [iprosone OV
BETAMETHASONE VALERATE		J		•
* Crm 0.1%	4.53	50 g OP	√ E	Beta Cream
* Oint 0.1%		50 g OP	_	Beta Ointment
* Lotn 0.1%	25.00	50 ml OP	✓ <u>E</u>	Betnovate
CLOBETASOL PROPIONATE				
* Crm 0.05%	2.40	30 g OP	✓ [<u> Permol</u>
* Oint 0.05%	2.33	30 g OP	✓ [ermol
CLOBETASONE BUTYRATE				
Crm 0.05%	5.38	30 g OP		
	(10.00)		Е	umovate
HYDROCORTISONE				
Y Crm 10/ Only on a proportion	1 70	20 ~ OD	./ 5	thian

HYDROCORTISONE	AND PARAFFIN I	IOLIID AND LANOLIN

Lotn 1% with paraffin liquid 15.9% and lanolin 0.6% - Only on

Powder – Only in combination......49.95

✓ DP Lotn HC 250 ml

galenicals

30 g OP

500 g

25 g

20.40

Up to 5% in a dermatological base (not proprietary Topical Corticosteriod - Plain) with or without other dermatological

✓ Ethics

✓ ABM

✓ Noumed

	Subsidy		Fully	Brand or
	(Manufacturer's		sidised	Generic
	\$	Per	1	Manufacturer
HYDROCORTISONE BUTYRATE				
Lipocream 0.1%	4.85	100 g OP	1	Locoid Lipocream
Oint 0.1%		100 g OP		Locoid .
Milky emul 0.1%	12.33	100 ml OP	1	Locoid Crelo
METHYLPREDNISOLONE ACEPONATE				
Crm 0.1%	4.95	15 g OP	1	<u>Advantan</u>
Oint 0.1%	4.95	15 g OP		Advantan
MOMETASONE FUROATE		- 3 -		
Crm 0.1%	1 05	15 g OP	1	Elocon Alcohol Free
OIII 0.1 /0	3.10	50 g OP		Elocon Alcohol Free
Oint 0.1%	00	15 g OP		Elocon
Onk 0.170	2.90	50 g OP		Elocon
Lotn 0.1%		30 ml OP		Elocon
		30 1111 31	-	
TRIAMCINOLONE ACETONIDE Crm 0.02%	6.40	100 ~ 00	./	Ariotocort
Oint 0.02%		100 g OP		Aristocort
OITIL 0.02%	0.34	100 g OP	•	<u>Aristocort</u>
Corticosteroids - Combination BETAMETHASONE VALERATE WITH SODIUM FUSIDATE	TELISIDIC ACIDI			
Crm 0.1% with sodium fusidate (fusidic acid) 2%		15 g OP		Fucicort
a) Maximum of 15 g per prescriptionb) Only on a prescription				
HYDROCORTISONE WITH MICONAZOLE - Only on a pres	cription			
* Crm 1% with miconazole nitrate 2%		15 g OP	1	Micreme H
HYDROCORTISONE WITH NATAMYCIN AND NEOMYCIN	- Only on a prescri	ption		
Oint 1% with natamycin 1% and neomycin sulphate 0.5%		15 g OP	1	Pimafucort
TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOM		•		
		IIIN		
Crm 1 mg with nystatin 100,000 u, neomycin sulphate 2.9 and gramicidin 250 mcg per g - Only on a prescripti		15 a ∩D		
and gramicium 250 meg per g – Omy on a prescripti	(9.28)	15 g OP		Viaderm KC
	(3.20)			viauciiii NO
Barrier Creams and Emollients				
Dainer Oreans and Emonicities				
Barrier Creams				
DIMETHICONE				
* Crm 5% pump bottle		500 ml OP	•	healthE Dimethicone 5%
* Crm 10% pump bottle	4.52	500 ml OP	•	healthE Dimethicone 10%
ZINC AND CASTOR OIL				
* Oint	4.25	500 g	1	Evara

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's I	Price) Subsi	Fully Brand or dised Generic
	(Manufacturer ST	Per Per	✓ Manufacturer
Emollients			
QUEOUS CREAM			
Crm	1.30	100 g	✓ healthE Aqueous Cream SLS Free
	1.73	500 g	✓ Evara✓ GEM AqueousCream
CETOMACROGOL			
₭ Crm BP	1.99	500 g	 Cetomacrogol-AFT
CETOMACROGOL WITH GLYCEROL		E00 : 05	4-
Crm 90% with glycerol 10%	2.13 3.50	500 ml OP 1,000 ml OP	✓ <u>Evara</u> ✓ <u>Evara</u>
EMULSIFYING OINTMENT			
* Oint BP	3.13	500 g	✓ Emulsifying Ointment ADE
OIL IN WATER EMULSION			
* Crm	2.04	500 g	✓ Fatty Cream AFT
PARAFFIN		05	
Oint liquid paraffin 50% with white soft paraffin 50%	4.94	500 g OP	✓ White Soft Liquid Paraffin AFT
JREA			
* Crm 10% NOOL FAT WITH MINERAL OIL – Only on a prescription	1.37	100 g OP	✓ healthE Urea Cream
* Lotn hydrous 3% with mineral oil	5.60	1,000 ml	
	(14.96)		DP Lotion
	(20.53)	050 1 00	Alpha-Keri Lotion
	1.40 (5.87)	250 ml OP	DP Lotion
	5.60	1,000 ml	DF LOUIDIT
	(23.91)	1,000 1111	BK Lotion
	1.40	250 ml OP	
	(7.73)		BK Lotion
Other Dermatological Bases			
PARAFFIN White poft Colly in combination	4.74	450 a	✓ EVADA White Sett
White soft - Only in combination	4./4	450 g	✓ EVARA White Soft Paraffin
	19.00	2,500 g	✓ EVARA White Soft Paraffin

Only in combination with a dermatological galenical or as a diluent for a proprietary Topical Corticosteroid – Plain.

	Subsidy (Manufacturer's Price \$	e) Sub Per	Fully sidised	Brand or Generic Manufacturer
Minor Skin Infections				
POVIDONE IODINE Oint 10%	7.40	65 g OP	√ E	Betadine
Antiseptic Solution 10%		100 ml 15 ml 500 ml	✓ F	Riodine Riodine Riodine
Skin preparation, povidone iodine 10% with 30% alcohol	(3.48)	100 ml	В	Betadine Skin Prep
Skin preparation, povidone iodine 10% with 70% alcohol	(7.78)	100 ml	P	Pfizer
Parasiticidal Preparations DIMETHICONE				

*	Lotn 4%4.25	200 ml OP	✓ <u>healthE</u>
			Dimethicone 4%
			<u>Lotion</u>

IVERMECTIN – Special Authority see SA2294 below – Retail pharmacy

- PSO for institutional use only. Must be endorsed with the name of the institution for which the PSO is required and a valid Special Authority for patient of that institution.
- 2) Ivermectin available on BSO provided the BSO includes a valid Special Authority for a patient of the institution.
- For the purposes of subsidy of ivermectin, institution means age related residential care facilities, disability care facilities or prisons.

⇒SA2294 Special Authority for Subsidy

Initial application — (Scabies) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Fither:

- 1 The person has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
- 2 Both:
 - 2.1 The person has a confirmed diagnosis of scabies or is a close contact of a scabies case; and
 - 2.2 Either:
 - 2.2.1 The person is unable to complete topical therapy: or
 - 2.2.2 Previous treatment with topical therapy has been tried and not cleared the infestation.

Initial application — (Other parasitic infections) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 filariasis; or
- 2 cutaneous larva migrans (creeping eruption); or
- 3 strongyloidiasis.

Renewal — (Scables) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria: Either:

- 1 The person has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
- 2 Both:

DERMATOLOGICALS

(Ma	Subsidy nufacturer's Price)	Sub	Fully	Brand or Generic
<u> </u>	\$	Per	✓	Manufacturer

continued...

- 2.1 The person has a confirmed diagnosis of scabies or is a close contact of a scabies case; and
- 2.2 Fither:
 - 2.2.1 The person is unable to complete topical therapy: or
 - 2.2.2 Previous treatment with topical therapy has been tried and not cleared the infestation.

Renewal — (Other parasitic infections) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 filariasis; or
- 2 cutaneous larva migrans (creeping eruption); or
- 3 strongyloidiasis.

PERMETHRIN

Psoriasis and Eczema Preparations

ACITRETIN - Special Authority see SA2024 below - Retail pharma	асу		
Cap 10 mg	26.20	60	Novatretin
Novatretin to be Principal Supply on 1 July 2024			
Cap 25 mg	57.37	60	✓ Novatretin
Novatretin to be Principal Supply on 1 July 2024			

⇒SA2024 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around acitretin and is competent to prescribe acitretin; and
- 3 Fither:
 - 3.1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and that they must not become pregnant during treatment and for a period of three years after the completion of treatment; or
 - 3.2 Patient is not of child bearing potential.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and that they must not become pregnant during treatment and for a period of three years after the completion of treatment; or
- 2 Patient is not of child bearing potential.

BETAMETHASONE DIPROPIONATE WITH CALCIPOTRIOL

Gel 500 mcg with calcipotriol 50 mcg per g	60 g OP 60 g OP 30 g OP	✓ Enstilar ✓ <u>Daivobet</u> ✓ <u>Daivobet</u>
CALCIPOTRIOL		
Oint 50 mcg per g 40.00	120 a OP	✓ Daiyonex

			/LITIMATOEOGIOALO
	Subsidy		Fully Brand or
	(Manufacturer's Pric	ce) Subs Per	sidised Generic Manufacturer
COAL TAR		-	
Soln BP – Only in combination	36.25	200 ml	✓ Midwest
1) Up to 10% only in combination with a dermatologic	al base or propriet	ary Topical C	Corticosteriod – Plain
2) With or without other dermatological galenicals.			
COAL TAR WITH ALLANTOIN, MENTHOL, PHENOL AND SULI			
Soln 5% with sulphur 0.5%, menthol 0.75%, phenol 0.5% an allantoin crm 2.5%		75 g OP	
unanton on 2.0/	(8.00)	70 9 01	Egopsoryl TA
	3.43	30 g OP	01 ,
	(4.35)	•	Egopsoryl TA
COAL TAR WITH SALICYLIC ACID AND SULPHUR			
Soln 12% with salicylic acid 2% and sulphur 4% oint	4.97	25 g OP	✓ Coco-Scalp
	7.95	40 g OP	✓ Coco-Scalp
PIMECROLIMUS - Special Authority see SA1970 below - Retail	I pharmacy		
a) Maximum of 15 g per prescription			
b) Note: a maximum of 15 g per prescription and no more t			
Cream 1% >> SA1970 Special Authority for Subsidy	33.00	15 g OP	✓ <u>Elidel</u>
Initial application only from a dermatologist, paediatrician, ophtl of a dermatologist, paediatrician or ophthalmologist. Approvals a meeting the following criteria: Both: 1 Patient has atopic dermatitis on the eyelid; and 2 Patient has at least one of the following contraindications documented epidermal atrophy, documented allergy to to	ralid without further	r renewal unle	ess notified for applications rificial dermatitis, rosacea,
pressure.		,	
PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORE		a prescriptior	ı
* Soln 2.3% with trolamine laurilsulfate and fluorescein sodium	15.41	500 ml	✓ Pinetarsol
SALICYLIC ACID			
Powder – Only in combination		250 g	✓ Midwest
 Only in combination with a dermatological base or With or without other dermatological galenicals. 	proprietary Topical	l Corticostero	id – Plain or collodion flexible
SULPHUR			
Precipitated – Only in combination	6.35	100 g	✓ Midwest
 Only in combination with a dermatological base or With or without other dermatological galenicals. 		l Corticostero	oid – Plain
TAODOLIMIO			
TACROLIMUS Oint 0.19/ Special Authority and SA2074 on the payt page			
Oint 0.1% - Special Authority see SA2074 on the next page Retail pharmacy		30 g OP	✓ Zematop
a) Maximum of 30 g per prescription		55 g 51	- <u>Lomatop</u>
b) Note: a maximum of 30 g per prescription and no maximum	ore than one presc	ription per 12	2 weeks.

DERMATOLOGICALS

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

⇒SA2074 Special Authority for Subsidy

Initial application only from a dermatologist, paediatrician or any relevant practitioner on the recommendation of a dermatologist, paediatrician, . Approvals valid without further renewal unless notified for applications meeting the following criteria: Both:

- 1 Patient has atopic dermatitis on the face; and
- 2 Patient has at least one of the following contraindications to topical corticosteroids: periorificial dermatitis, rosacea, documented epidermal atrophy or documented allergy to topical corticosteroids.

Scalp Preparations

BETAMETHASONE VALERATE * Scalp app 0.1%	9 84	100 ml OP	✓ Beta Scalp
CLOBETASOL PROPIONATE		100 1111 01	<u> Dota Gourp</u>
* Scalp app 0.05%	6.26	30 ml OP	✓ Dermol
HYDROCORTISONE BUTYRATE Scalp lotn 0.1%	6.57	100 ml OP	✓ Locoid
KETOCONAZOLE Shampoo 2%	3.23	100 ml OP	✓ Sebizole
a) Maximum of 100 ml per prescription	4.09		✓ <u>Sebizole</u>

a) Maximum of 100 ml per prescription

b) Only on a prescription

Sunscreens

SUNSCREENS, PROPRIETARY - Subsidy by endorsement

Only if prescribed for a patient with severe photosensitivity secondary to a defined clinical condition and the prescription is endorsed accordingly.

200 g OP ✓ Marine Blue Lotion

SPF 50+

Wart Preparations

For salicylic acid preparations refer to PSORIASIS AND ECZEMA PREPARATIONS, page 74

PODOPHYLLOTOXIN

3.5 ml OP ✓ Condyline

a) Maximum of 3.5 ml per prescription

b) Only on a prescription

Other Skin Preparations

Antineoplastics

FLUOROURACIL SODIUM		
Crm 5%6.95	20 g OP	✓ Efudix
IMIQUIMOD		
Crm 5%, 250 mg sachet 21.72	24	✓ Perrigo

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

		Subsidy (Manufacturer's Price)	Dox	Fully Subsidised	Generic
		\$	Per		Manufacturer
	aceptives - Non-hormonal				
ondo	oms				
NDOM	1S				
49 mi	m - Up to 144 dev available on a PSO	11.42	144	1	Moments
53 m	m	0.95	10		Moments
		11.64	144	•	Moments
	Maximum of 60 dev per prescription				
) Up to 60 dev available on a PSO			_	
53 m	m, 0.05 mm thickness		10		Moments
		11.42	144	•	Moments
	u) Up to 60 dev available on a PSO				
	Maximum of 60 dev per prescription	0.05	40	,	
53 mi	m, chocolate, brown		10		Moments
_) Un to 60 day available are a DCO	11.64	144	•	Moments
	u) Up to 60 dev available on a PSO				
	n) Maximum of 60 dev per prescription m, strawberry, red	0.05	10	./	Moments
33 1111	iii, Sirawberry, red	11.64	144		Moments
	u) Up to 60 dev available on a PSO	11.04	144	•	WOMENIS
	Maximum of 60 dev per prescription				
	m	0.97	10	1	Moments
50 1111		11.64	144		Moments
9	Maximum of 60 dev per prescription	11.04	1-1-1	•	Monitorito
	b) Up to 60 dev available on a PSO				
	m, 0.05 mm thickness	2.00	12	1	Gold Knight
	.,	24.10	144		Gold Knight
а	u) Up to 60 dev available on a PSO				3
	n) Maximum of 60 dev per prescription				
	m, 0.05mm thickness (bulk pack)	20.17	144	1	Gold Knight
) Maximum of 60 dev per prescription				· ·
) Up to 60 dev available on a PSO				
	m, 0.08 mm thickness	0.97	10	1	Moments
		11.64	144	1	Moments
а	u) Up to 60 dev available on a PSO				
) Maximum of 60 dev per prescription				
56 m	m, 0.08 mm thickness, red		10		Moments
		11.64	144	/	Moments
	u) Up to 60 dev available on a PSO				
	Maximum of 60 dev per prescription			_	
56 m	m, chocolate		12		Gold Knight
		21.45	144	/	Gold Knight
	u) Up to 60 dev available on a PSO				
) Maximum of 60 dev per prescription	4 70			O-Id K-1 11
56 MI	m, strawberry		12		Gold Knight
). He to 00 days and lable on a BOO	21.45	144	•	Gold Knight
	u) Up to 60 dev available on a PSO				
) Maximum of 60 dev per prescription	4.00	10	.1	Cold Knight VI
ou mi	m		12		Gold Knight XL Gold Knight XL
		21.89	144	•	adia Kiligili AL

✓ 7 MED NSHA Silver/

TT380 Standard

✓ Choice Load 375

Subsidy (Manufacturer's Pric \$	ce) Per	Fully Subsidised	Brand or Generic Manufacturer	
---------------------------------------	------------	---------------------	-------------------------------------	--

- a) Maximum of 60 dev per prescription
- b) Up to 60 dev available on a PSO

Contraceptive Devices

INTRA-UTERINE DEVICE

- a) Up to 40 dev available on a PSO
- b) Only on a PSO

		•	Copper Short
			✓ Choice 380 7med
			Nsha Silver/
			copper Short
			✓ Choice TT380 Short
* IUD 33.6 mm length x 29.9 mm width	29.80	1	✓ Choice

Contraceptives - Hormonal

Combined Oral Contraceptives

⇒SA0500 Special Authority for Alternate Subsidy

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Either:
 - 1.1 Patient is on a Social Welfare benefit: or
 - 1.2 Patient has an income no greater than the benefit: and

* IUD 29.1 mm length x 23.2 mm width 29.80

2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Fither:

- 1 Patient is on a Social Welfare benefit; or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- · on a Social Welfare benefit; or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

ETHINYLOFSTRADIOL WITH DESOGESTREL

★ Tab 20 mcg with desogestrel 150 mcg and 7 inert tab - Up to 84 tab available on a PSO.......10.00 84 Mercilon 28

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		osidised	Generic
	\$	Per		Manufacturer
THINYLOESTRADIOL WITH LEVONORGESTREL				
★ Tab 20 mcg with levonorgestrel 100 mcg and 7 inert tablets	_			
Up to 84 tab available on a PSO	1.50	84	✓ L	o-Oralcon 20 ED
★ Tab 30 mcg with levonorgestrel 150 mcg		63	_	
0 0	(16.50)		N	licrogynon 30
 a) Higher subsidy of \$15.00 per 63 tab with Special Au b) Up to 63 tab available on a PSO Tab 30 mcg with levonorgestrel 150 mcg and 7 inert tablets Up to 84 tab available on a PSO 	-	the prev		ge Oralcon 30 ED
THINYLOESTRADIOL WITH NORETHISTERONE				
★ Tab 35 mcg with norethisterone 1 mg and 7 inert tab — Up t	0			
84 tab available on a PSO		84	✓ B	revinor 1/28
	16.33	112	✓ B	Brevinor-1 28 Day
				lorimin-1 28 Day
Tab 35 mcg with norethisterone 500 mcg and 7 inert tab - U	Un			•
to 84 tab available on a PSO	21.99	84	✓ N	lorimin
Brevinor-1 28 Day Tab 35 mcg with norethisterone 1 mg and 7 Norimin-1 28 Day Tab 35 mcg with norethisterone 1 mg and 7 i	inert tab to be delisted	l 1 Decei	mber 202	24)

Progestogen-only Contraceptives

⇒SA0500 Special Authority for Alternate Subsidy

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Either:
 - 1.1 Patient is on a Social Welfare benefit; or
 - 1.2 Patient has an income no greater than the benefit; and
- 2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Fither:

- 1 Patient is on a Social Welfare benefit; or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- on a Social Welfare benefit; or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

LEVONORGESTREL

*	Tab 30 mcg - Up to 84 tab available on a PSO		84 112	✓ Microlut ✓ Microlut
	•	22.00	112	• Wilcrolut
*	Subdermal implant $(2 \times 75 \text{ mg rods})$ – Up to 3 pack available			
	on a PSO1	06.92	1	✓ <u>Jadelle</u>
ME	DROXYPROGESTERONE ACETATE			
	Inj 150 mg per ml, 1 ml syringe - Up to 5 inj available on a PSO	9.18	1	✓ Depo-Provera

	GENIT	O-URINARY SYSTE	M
Subsidy (Manufacturer's I \$	Price) Sub	Fully Brand or sidised Generic Manufacturer	
NORETHISTERONE Tab 350 mcg - Up to 84 tab available on a PSO12.25	84	✓ Noriday 28	
Emergency Contraceptives			
LEVONORGESTREL * Tab 1.5 mg1.75	1	✓ <u>Levonorgestrel</u> <u>BNM</u>	
a) Maximum of 2 tab per prescriptionb) Up to 5 tab available on a PSOc) Note: Direct Provision by a pharmacist permitted under the provision	ns in Part I of S	Section A.	
Antiandrogen Oral Contraceptives			
 A maximum \$5.00 prescription charge (patient co-payment) may apply. prescription may be written for up to six months supply. Prescriptions coded in any other way are subject to any non contraceptive prescriptions-contraceptive period of supply. ie. Prescriptions may be written for up to the CYPROTERONE ACETATE WITH ETHINYLOESTRADIOL * Tab 2 mg with ethinyloestradiol 35 mcg and 7 inert tabs - Up to 168 tab available on a PSO			
Gynaecological Anti-infectives			
ACETIC ACID WITH HYDROXYQUINOLINE AND RICINOLEIC ACID Jelly with glacial acetic acid 0.94%, hydroxyquinoline sulphate 0.025%, glycerol 5% and ricinoleic acid 0.75% with applicator8.43 (24.87)	100 g OP	Aci-Jel	
CLOTRIMAZOLE	05 ~ OD	/ Olemenal	
* Vaginal crm 1% with applicators3.50* Vaginal crm 2% with applicators3.85	35 g OP 20 g OP	✓ Clomazol✓ Clomazol	
MICONAZOLE NITRATE * Vaginal crm 2% with applicator6.89	40 g OP	✓ Micreme	
NYSTATIN Vaginal crm 100,000 u per 5 g with applicator(s)5.70	75 g OP	✓ <u>Nilstat</u>	
Myometrial and Vaginal Hormone Preparations			
ERGOMETRINE MALEATE			

	Inj 500 mcg per ml, 1 ml ampoule - Up to 5 inj available on a		
	PSO160.00	5	DBL Ergometrine
OE	STRIOL		
*	Crm 1 mg per g with applicator6.95	15 g OP	✓ Ovestin
*	Pessaries 500 mcg7.55	15	✓ Ovestin

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	
OXYTOCIN – Up to 5 inj available on a PSO				
Inj 5 iu per ml, 1 ml ampoule	4.98	5	✓	Oxytocin BNM
Inj 10 iu per ml, 1 ml ampoule	5.98	5	✓	Oxytocin BNM
			✓	Oxytocin GH S29
	11.96	10	•	Oxytocin Panpharma
(Oxytocin GH S29 Inj 10 iu per ml, 1 ml ampoule to be delisted	1 August 2024)			-
OXYTOCIN WITH ERGOMETRINE MALEATE - Up to 5 inj avai	lable on a PSO			
Inj 5 iu with ergometrine maleate 500 mcg per ml, 1 ml ampo		5	•	<u>Syntometrine</u>

Pregnancy Tests - hCG Urine

PREGNANCY TESTS - HCG URINE

- a) Up to 200 test available on a PSO

b) Only on a PSO Cassette	12.00	40 test OP	✓ Smith BioMed Rapid
	16.00		Pregnancy Test ✓ David One Step
			Cassette
			Pregnancy Test

Urinary Agents

For urinary tract Infections refer to INFECTIONS, Antibacterials, page 113

5-Alpha Reductase Inhibitors

FINASTERIDE - Special Authority see SA0928 below - Retail pharmacy 100 ✓ Ricit * Tab 5 mg4.79

⇒SA0928 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- - 2.1 The patient is intolerant of non-selective alpha blockers or these are contraindicated; or
 - 2.2 Symptoms are not adequately controlled with non-selective alpha blockers.

Alpha-1A Adrenoreceptor Blockers

TAMSULOSIN HYDROCHLORIDE - Special Authority see SA1032 below - Retail pharmacy

✓ Tamsulosin-Rex * Cap 400 mcg22.31

⇒SA1032 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 The patient is intolerant of non-selective alpha blockers or these are contraindicated.

	Subsidy (Manufacturer's I \$	Price) Subs Per	idised Ge	and or neric nufacturer
Other Urinary Agents				
OXYBUTYNIN				
* Tab 5 mg	5.42	100	✓ Alche	emy /butynin
POTASSIUM CITRATE				,
Oral liq 3 mmol per ml — Special Authority see SA1083 Retail pharmacy		200 ml OP	✓ Biom	ed
⇒SA1083 Special Authority for Subsidy				
Initial application from any relevant practitioner. Approval Both:	s valid for 12 months	for applications	meeting the	e following criteria:
1 The patient has recurrent calcium oxalate urolithiasis2 The patient has had more than two renal calculi in the		ne application.		
Renewal from any relevant practitioner. Approvals valid for benefitting from the treatment.			s appropria	e and the patient is
SODIUM CITRO-TARTRATE				
* Grans eff 4 g sachets	3.50	28	✓ <u>Ural</u>	
SOLIFENACIN SUCCINATE	0.05			
Tab 5 mg Tab 10 mg		30 30		enacin Viatris enacin Viatris
Tab 10 mg		30	• <u>301116</u>	macini viatris
Detection of Substances in Urine				
ORTHO-TOLIDINE				
* Compound diagnostic sticks		50 test OP		
	(8.25)		Hema	stix
TETRABROMOPHENOL	10.00	100 ++ 00	. / All	
* Blue diagnostic strips	13.92	100 test OP	✓ Albus	STIX
Obstetric Preparations				
Antiprogesterones				
MIFEPRISTONE				
Tab 200 mg - Up to 15 tab available on a PSO		1	✓ Mifeg	•
	180.00	3	✓ Mifeg	yne

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidis	sed	Generic
\$	Per	✓	Manufacturer

Calcium Homeostasis

CALCITONIN * Inj 100 iu per ml, 1 ml ampoule	121.00	5	✓ Miacalcic
CINACALCET - Special Authority see SA2170 below - Retail	pharmacy		
Tab 30 mg - Wastage claimable	42.06	28	✓ Cinacalet Devatis
Tab 60 mg - Wastage claimable	84.12	28	✓ Cinacalet Devatis

⇒SA2170 Special Authority for Subsidy

Initial application — (parathyroid carcinoma or calciphylaxis) only from a nephrologist or endocrinologist. Approvals valid for 6 months for applications meeting the following criteria:

- Either:
 - 1.1 The patient has been diagnosed with a parathyroid carcinoma (see Note); and
 - 1.2 The patient has persistent hypercalcaemia (serum calcium greater than or equal to 3 mmol/L) despite previous first-line treatments including sodium thiosulfate (where appropriate) and bisphosphonates; and
 - 1.3 The patient is symptomatic; or
 - 2 All of the following:

1 All of the following:

- 2.1 The patient has been diagnosed with calciphylaxis (calcific uraemic arteriolopathy); and
- 2.2 The patient has symptomatic (e.g. painful skin ulcers) hypercalcaemia (serum calcium greater than or equal to 3 mmol/L); and
- 2.3 The patient's condition has not responded to previous first-line treatments including bisphosphonates and sodium thiosulfate

Renewal — (parathyroid carcinoma or calciphylaxis) only from a nephrologist or endocrinologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient's serum calcium level has fallen to < 3mmol/L; and
- 2 The patient has experienced clinically significant symptom improvement.

Note: This does not include parathyroid adenomas unless these have become malignant.

Initial application — (**primary hyperparathyroidism**) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has primary hyperparathyroidism; and
- 2 Either:
 - 2.1 Patient has hypercalcaemia of more than 3 mmol/L with or without symptoms; or
 - 2.2 Patient has hypercalcaemia of more than 2.85 mmol/L with symptoms; and
- 3 Surgery is not feasible or has failed; and
- 4 Patient has other comorbidities, severe bone pain, or calciphylaxis.

Initial application — (secondary or tertiary hyperparathyroidism) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient has tertiary hyperparathyroidism and markedly elevated parathyroid hormone (PTH) with hypercalcaemia; or
 - 1.2 Patient has symptomatic secondary hyperparathyroidism and elevated PTH; and
- 2 Patient is on renal replacement therapy; and
- 3 Any of the following:
 - 3.1 Residual parathyroid tissue has not been localised despite repeat unsuccessful parathyroid explorations; or

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
 \$	Per	✓	

continued...

- 3.2 Parathyroid tissue is surgically inaccessible; or
- 3.3 Parathyroid surgery is not feasible.

Renewal — (secondary or tertiary hyperparathyroidism) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Fither:

- 1 The patient has had a kidney transplant, and following a treatment free interval of at least 12 weeks a clinically acceptable parathyroid hormone (PTH) level to support ongoing cessation of treatment has not been reached; or
- 2 The patient has not received a kidney transplant and trial of withdrawal of cinacalcet is clinically inappropriate.

ZOLEDRONIC ACID

Inj 4 mg per 5 ml, vial	18.00	1	✓ Zoledronic acid
			Viatris

Corticosteroids and Related Agents for Systemic Use

DETAMETHAÇONE CODILIM DHOCDHATE WITH DETAMETHAÇONE ACETATE

BE	TAMETHASONE SODIUM PHOSPHATE WITH BETAMETHASONE ACETAT	E		
*	Inj 3.9 mg with betamethasone acetate 3 mg per ml, 1 ml19.20	5		
	(36.96)			Celestone
	(*****)			Chronodose
D.F.	VANAETHACONE			000000
	XAMETHASONE			
*	Tab 0.5 mg – Up to 60 tab available on a PSO1.50	30		<u>Dexmethsone</u>
*	Tab 4 mg - Up to 30 tab available on a PSO2.65	30		<u>Dexmethsone</u>
	Oral liq 1 mg per ml52.80	25 ml OP	1	Biomed
DE	XAMETHASONE PHOSPHATE			
	Dexamethasone phosphate injection will not be funded for oral use.			
*	Inj 4 mg per ml, 1 ml ampoule – Up to 5 inj available on a PSO7.86	10	1	Hameln
	Inj 4 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO 13.10	10		Hameln
		10	•	<u>namem</u>
	JDROCORTISONE ACETATE		_	
*	Tab 100 mcg11.46	100	/	Florinef
HY	DROCORTISONE			
*	Tab 5 mg8.10	100	1	Douglas
*	Tab 20 mg	100		Douglas
	Inj 100 mg vial	1		Solu-Cortef
	a) Not on a BSO	·	-	<u> </u>
	,			
	b) Up to 5 inj available on a PSO			
ME	THYLPREDNISOLONE			
*	Tab 4 mg112.00	100	1	Medrol
*	Tab 100 mg223.10	20	1	Medrol
NAE	-			
IVIE	THYLPREDNISOLONE (AS SODIUM SUCCINATE)	4	./	Solu-Medrol-Act-
	Inj 40 mg vial22.30	1	•	
				O-Vial
	In: 105 mm vial			Solu-Medrol-Act-
	Inj 125 mg vial34.10	1	•	
				O-Vial
	Ini E00 ma vial	4	.,	Solu-Medrol-Act-
	Inj 500 mg vial26.88	1	•	
				O-Vial
	Ini 4 a vial	4		Calv. Madeal
	Inj 1 g vial32.84	1	•	Solu-Medrol

	Subsidy		Fully	
	(Manufacturer's F	Price) Sub Per	sidised	Generic Manufacturer
	Į.	rei		Manufacturer
METHYLPREDNISOLONE ACETATE		_	_	
Inj 40 mg per ml, 1 ml vial	47.06	5	•	Depo-Medrol
PREDNISOLONE				
Oral liq 5 mg per ml - Up to 30 ml available on a PSO Restricted to children under 12 years of age.	6.00	30 ml OP	1	Redipred
PREDNISONE				
* Tab 1 mg	18.58	500	1	Prednisone Clinect
* Tab 2.5 mg	21.04	500	✓	Prednisone Clinect
* Tab 5 mg - Up to 30 tab available on a PSO	19.30	500	✓	Prednisone Clinect
★ Tab 20 mg – Up to 30 tab available on a PSO	50.51	500	✓	Prednisone Clinect
FETRACOSACTRIN				
★ Inj 250 mcg per ml, 1 ml ampoule	86 25	1	1	Synacthen
iii 200 mog por mi, i mi ampoulo	00.20	'		UK Synacthen
* Inj 1 mg per ml, 1 ml ampoule	690.00	1		Synacthen Depot
,g por mi, . mi ampodio		1		Synacthene
			•	Retard \$29
				netaru 929
FRIAMCINOLONE ACETONIDE			_	
Inj 10 mg per ml, 1 ml ampoule		5		Kenacort-A 10
Inj 40 mg per ml, 1 ml ampoule	52.63	5		Kenacort-A 40
Androgen Agonists and Antagonists				
CYPROTERONE ACETATE				
Tab 50 mg	14.37	50	1	Siterone
Tab 100 mg		50		Siterone
TESTOSTERONE				
Gel (transdermal) 16.2 mg per g	52.00	88 g OP	1	Testogel
Testogel to be Principal Supply on 1 July 2024	52.00	00 y OF	•	restoger
Patch 5 mg per day	225.00	30	1	Androderm
	223.00	30	•	Alluloucilli
TESTOSTERONE CIPIONATE	05.00		,	
Inj 100 mg per ml, 10 ml vial		1		Depo-Testosterone
	393.00		•	Taro-
				Testosterone S29
Tara Tantantarana (MA) Ini 100 man any ani 10 mil vial ta ba da	lists of 4 Assessed OO	10.4)		
Taro-Testosterone 29 Inj 100 mg per ml, 10 ml vial to be de	nisiea i August 20	124)		
TESTOSTERONE ESTERS				_
Inj 250 mg per ml, 1 ml	12.98	1	1	Sustanon Ampoules
FESTOSTERONE UNDECANOATE				
Cap 40 mg - Subsidy by endorsement	36.00	100	1	Steril-Gene S29
Subsidy by endorsement – subsidised for patients who				
1 November 2021 and the prescription is endorsed ac	-			. • .
where there exists a record of prior dispensing of testo				
Ini 250 mg ner ml 4 ml vial				

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer Hormone Replacement Therapy - Systemic Oestrogens **OESTRADIOL** 28 OP (11.10)Estrofem 28 OP Estrofem 8 ✓ Estradiol TDP Mylan ✓ Estraderm MX \$29 ✓ Estradot 14.50 a) No more than 2 patch per week b) Only on a prescription Patch 50 mcg per day......10.75 8 ✓ Estradiol TDP Mylan ✓ Estradiol Viatris ✓ Estraderm MX S29 14.50 ✓ Estradiol Sandoz ✓ Estradot a) No more than 2 patch per week b) Only on a prescription Patch 75 mcg per day......11.88 ✓ Estradiol TDP Mylan 8 ✓ Estradiol Viatris ✓ Estradiol Sandoz 14.50 ✓ Estradot a) No more than 2 patch per week b) Only on a prescription 8 ✓ Estradiol TDP Mylan ✓ Estradiol Viatris 14.50 ✓ Estradiol Sandoz ✓ Estradot ✓ Estraderm MX S29 15.50 a) No more than 2 patch per week b) Only on a prescription **OESTRADIOL VALERATE** 84 ✓ Progynova ✓ Progynova 84 **OESTROGENS** Conjugated, equine tab 300 mcg......3.01 28 Premarin (17.50)Conjugated, equine tab 625 mcg......4.12 28 (17.50)Premarin **Progestogens** MEDBOXYPROGESTERONE ACETATE

30

56

56

100

30

8.75

✓ Provera

✓ Provera

✓ Provera✓ Provera

✓ Provera

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturaria Pri	oo) Cub	Fully Brand or soldised Generic
	(Manufacturer's Prio	Per Sub	osidised Generic ✓ Manufacturer
Progestogen and Oestrogen Combined Prepara	ations		
OESTRADIOL WITH NORETHISTERONE			
* Tab 1 mg with 0.5 mg norethisterone acetate		28 OP	
	(18.10)		Kliovance
* Tab 2 mg with 1 mg norethisterone acetate		28 OP	IZP .
M. T. I.O	(18.10)		Kliogest
* Tab 2 mg with 1 mg norethisterone acetate (10), and 2 mg	E 40	00 OD	
oestradiol tab (12) and 1 mg oestradiol tab (6)		28 OP	Trionguana
	(18.10)		Trisequens
Other Oestrogen Preparations			
OESTRIOL THE PARTY OF THE PARTY	7.70	00	1 Occasilla
* Tab 2 mg	7.70	30	✓ <u>Ovestin</u>
Other Progestogen Preparations			
Other Progestogen Preparations			
LEVONORGESTREL			
* Intra-uterine device 52 mg		1	✓ <u>Mirena</u>
* Intra-uterine device 13.5 mg	215.60	1	✓ <u>Jaydess</u>
MEDROXYPROGESTERONE ACETATE			
Tab 100 mg	116.15	100	Provera HD
NORETHISTERONE			
* Tab 5 mg - Up to 30 tab available on a PSO	5.49	30	✓ Primolut N
PROGESTERONE			
* Cap 100 mg	14.85	30	✓ Utrogestan
Thyroid and Antithyroid Agents			
CARBIMAZOLE			
* Tab 5 mg	7 56	100	✓ Neo-Mercazole
•	7.50	100	11CO-MCTGGZOIC
LEVOTHYROXINE * Tab 25 mcg	5 55	90	✓ Synthroid
* Tab 50 mcg		28	✓ Mercury Pharma
7 Tab 50 Mog	5.79	90	✓ Synthroid
	12.86	200	✓ Eltroxin
	64.28	1,000	✓ Eltroxin
* Tab 100 mcg	1.78	28	✓ Mercury Pharma
-	6.01	90	✓ Synthroid
	13.36	200	✓ Eltroxin
	66.78	1,000	✓ Eltroxin
PROPYLTHIOURACIL – Special Authority see SA1199 below –	Retail pharmacy		
Tab 50 mg	35.00	100	✓ PTU S29

⇒SA1199 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 The patient has hyperthyroidism; and
- 2 The patient is intolerant of carbimazole or carbimazole is contraindicated.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

Subsidy	Fu	ly Brand or	
(Manufacturer's Price)	Subsidise	ed Generic	
\$	Per	 Manufacturer 	

Trophic Hormones

Growth Hormones

SO	MATROPIN (OMNITROPE) - Special Authority see SA2032 below	- Retail pharm	acy	
*	Inj 5 mg cartridge	69.75	i	✓ Omnitrope
				✓ Omnitrope S29 S29
*	Inj 10 mg cartridge	69.75	1	✓ Omnitrope
				✓ Omnitrope S29 S29
*	Inj 15 mg cartridge	139.50	1	✓ Omnitrope
				✓ Omnitrope S29 S29

⇒SA2032 Special Authority for Subsidy

Initial application — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

Fither:

- 1 Growth hormone deficiency causing symptomatic hypoglycaemia, or with other significant growth hormone deficient sequelae (e.g. cardiomyopathy, hepatic dysfunction) and diagnosed with GH < 5 mcg/l on at least two random blood samples in the first 2 weeks of life, or from samples during established hypoglycaemia (whole blood glucose < 2 mmol/l using a laboratory device); or</p>
- 2 All of the following:
 - 2.1 Height velocity < 25th percentile for age adjusted for bone age/pubertal status if appropriate over 6 or 12 months using the standards of Tanner and Davies (1985); and</p>
 - 2.2 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
 - 2.3 Peak growth hormone value of < 5.0 mcg per litre in response to two different growth hormone stimulation tests. In children who are 5 years or older, GH testing with sex steroid priming is required; and
 - 2.4 If the patient has been treated for a malignancy, they should be disease free for at least one year based upon follow-up laboratory and radiological imaging appropriate for the malignancy, unless there are strong medical reasons why this is either not necessary or appropriate; and
 - 2.5 Appropriate imaging of the pituitary gland has been obtained.

Renewal — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 2 Height velocity is greater than or equal to 25th percentile for age (adjusted for bone age/pubertal status if appropriate) while on growth hormone treatment, as calculated over six months using the standards of Tanner and Davis (1985); and
- 3 Height velocity is greater than or equal to 2.0 cm per year, as calculated over 6 months; and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a post-natal genotype confirming Turner Syndrome; and
- 2 Height velocity is < 25th percentile over 6-12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is < 14 years.

Renewal — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

Subsidy (Manufacturaria Price)	Ç.	Fully bsidised	Brand or
 (Manufacturer's Price) \$	Per	⊅Sidised ✓	Generic Manufacturer

continued...

- 1 Height velocity is greater than or equal to 50th percentile for age (while on growth hormone calculated over 6 to 12 months using the Ranke's Turner Syndrome growth velocity charts); and
- 2 Height velocity is greater than or equal to 2 cm per year, calculated over six months; and
- 3 A current bone age is 14 years or under; and
- 4 No serious adverse effect that the specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 3 standard deviations below the mean for age or for bone age if there is marked growth acceleration or delay; and
- 2 Height velocity is < 25th percentile for age (adjusted for bone age/pubertal status if appropriate), as calculated over 6 to 12 months using the standards of Tanner and Davies(1985); and
- 3 A current bone age is < 14 years or under (female patients) or < 16 years (male patients); and
- 4 The patient does not have severe chronic disease (including malignancy or recognized severe skeletal dysplasia) and is not receiving medications known to impair height velocity.

Renewal — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred.

Initial application — (short stature due to chronic renal insufficiency) only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 2 standard deviations below the mean; and
- 2 Height velocity is < 25th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and</p>
- 3 A current bone age is to 14 years or under (female patients) or to 16 years or under (male patients); and
- 4 The patient is metabolically stable, has no evidence of metabolic bone disease and absence of any other severe chronic disease; and
- 5 The patient is under the supervision of a specialist with expertise in renal medicine; and
- 6 Either:
 - 6.1 The patient has a GFR less than or equal to 30 ml/min/1.73m² as measured by the Schwartz method (Height(cm)/plasma creatinine (umol/l)) × 40 = corrected GFR (ml/min/1.73m²) in a child who may or may not be receiving dialysis; or
 - 6.2 The patient has received a renal transplant and has received < 5mg/ m²/day of prednisone or equivalent for at least 6 months...

Renewal — (short stature due to chronic renal insufficiency) only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

Subsidy (Manufacturer's Price)	Subs Per	Fully sidised	Brand or Generic Manufacturer
 Ψ	1 01		I VIGITATION CONTRACTOR

continued...

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not experienced significant biochemical or metabolic deterioration confirmed by diagnostic results; and
- 7 The patient has not received renal transplantation since starting growth hormone treatment; and
- 8 If the patient requires transplantation, growth hormone prescription should cease before transplantation and a new application should be made after transplantation based on the above criteria.

Initial application — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of Prader-Willi syndrome that has been confirmed by genetic testing or clinical scoring criteria; and
- 2 The patient is aged six months or older; and
- 3 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
- 4 Sleep studies or overnight eximetry have been performed and there is no obstructive sleep disorder requiring treatment, or if an obstructive sleep disorder is found, it has been adequately treated under the care of a paediatric respiratory physician and/or ENT surgeon; and
- 5 Either:
 - 5.1 Both:
 - 5.1.1 The patient is aged two years or older; and
 - 5.1.2 There is no evidence of type II diabetes or uncontrolled obesity defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months; or
 - 5.2 The patient is aged between six months and two years and a thorough upper airway assessment is planned to be undertaken prior to treatment commencement and at six to 12 weeks following treatment initiation.

Renewal — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not developed type II diabetes or uncontrolled obesity as defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months.

Initial application — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a medical condition that is known to cause growth hormone deficiency (e.g. surgical removal of the pituitary for treatment of a pituitary tumour); and
- 2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
- 3 The patient has severe growth hormone deficiency (see notes); and

Subsidy		Fully	Brand or	
(Manufacturer's I	,	ubsidised	Generic	
\$	Per		Manufacturer	

continued...

- 4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
- 5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

Renewal — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 All of the following:
 - 1.1 The patient has been treated with somatropin for < 12 months; and
 - 1.2 There has been an improvement in Quality of Life defined as a reduction of at least 8 points on the Quality of Life Assessment of Growth Hormone Deficiency in Adults (QoL-AGHDA®) score from baseline; and
 - 1.3 Serum IGF-I levels have been increased within ±1SD of the mean of the normal range for age and sex; and
- 1.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients, or 1 mg per day for female patients; or 2 All of the following:
 - 2.1 The patient has been treated with somatropin for more than 12 months; and
 - 2.2 The patient has not had a deterioration in Quality of Life defined as a 6 point or greater increase from their lowest QoL-AGHDA® score on treatment (other than due to obvious external factors such as external stressors); and
 - 2.3 Serum IGF-I levels have continued to be maintained within ±1SD of the mean of the normal range for age and sex (other than for obvious external factors); and
 - 2.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients or 1 mg per day for female patients; or
- 3 All of the following:
 - 3.1 The patient has had a Special Authority approval for somatropin for childhood deficiency in children and no longer meets the renewal criteria under this indication; and
 - 3.2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
 - 3.3 The patient has severe growth hormone deficiency (see notes); and
 - 3.4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
 - 3.5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

(Ma	Subsidy anufacturer's Price) \$	Per	Fully Subsidised	Generic
continued Dose of somatropin not to exceed 0.7 mg per day for male patients, of the commencement of treatment for hypopituitarism, patients must doses of corticosteroid and levothyroxine.				
GnRH Analogues				
GOSERELIN Implant 3.6 mg, syringe Implant 10.8 mg, syringe		1		Zoladex Zoladex
Additional subsidy by endorsement where the patient is a child o goserelin and the prescription is endorsed accordingly. Inj 3.75 mg prefilled dual chamber syringe — Higher subsidy of	r adolescent and	is una	able to tole	erate administration of
\$221.60 per 1 inj with Endorsement	66.48 (221.60)	1		Lucrin Depot 1-month
Inj 11.25 mg prefilled dual chamber syringe – Higher subsidy of \$591.68 per 1 inj with Endorsement	177.50 (591.68)	1		Lucrin Depot 3-month
Vasopressin Agonists				
DESMOPRESSIN				
Wafer 120 mcg DESMOPRESSIN ACETATE	47.00	30	•	Minirin Melt
Tab 100 mcg		30		Minirin
Tab 200 mcg		30		Minirin
Nasal spray 10 mcg per dose	34.95	6 ml O	P 🗸	<u>PH&T</u>
Inj 4 mcg per ml, 1 ml	67.18	10	✓	Minirin
Other Endocrine Agents				
CABERGOLINE				
Tab 0.5 mg - Maximum of 2 tab per prescription; can be				
waived by Special Authority see SA2070 below		2		Dostinex
	17.94	8	•	Dostinex

17.94 Dostinex

⇒SA2070 Special Authority for Waiver of Rule

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Hyperprolactinemia; or
- 2 Acromegaly*; or
- 3 Inhibition of lactation.

Renewal — (for patients who have previously been funded under Special Authority form SA1031) from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has previously held a valid Special Authority which has expired and the treatment remains appropriate and the patient is benefiting from treatment.

Note: Indication marked with * is an unapproved indication.

		TRATE

CLOMIFENE CITRATE				
Tab 50 mg	29.84	10	✓ Mylan	
			Clominhen \$29	

	Subsidy (Manufacturer's Price) \$	Su Per	Fully bsidised	Brand or Generic Manufacturer
METYRAPONE Cap 250 mg	558.00	50	✓ M	etopirone

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer **Anthelmintics** ALBENDAZOLE - Special Authority see SA1318 below - Retail pharmacy 60 ✓ Fskazole S29 **⇒SA1318** Special Authority for Subsidy Initial application only from an infectious disease specialist or clinical microbiologist. Approvals valid for 6 months where the patient has hydatids. Renewal only from an infectious disease specialist or clinical microbiologist. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefitting from the treatment. MEBENDAZOLE - Only on a prescription 6 Vermox 15 ml (7.83)Vermox **PRAZIQUANTEL** Biltricide **Antibacterials** a) For topical antibacterials, refer to DERMATOLOGICALS, page 68

Cephalosporins and Cephamycins

b) For anti-infective eye preparations, refer to SENSORY ORGANS, page 259

oopyoo			
CEFACLOR MONOHYDRATE			
Cap 250 mg	25.85	100	 Ranbaxy-Cefaclor
Grans for oral liq 125 mg per 5 ml - Wastage claimable	3.75	100 ml	 Ranbaxy-Cefactor
CEFALEXIN			
Cap 250 mg	3.85	20	Cephalexin ABM
Cap 500 mg		20	✓ Cephalexin ABM
Grans for oral liq 25 mg per ml - Wastage claimable	7.88	100 ml	✓ Flynn
Grans for oral liq 50 mg per ml - Wastage claimable	10.38	100 ml	✓ Flynn
	11.75		Cefalexin Sandoz
CEFAZOLIN - Subsidy by endorsement			
Only if prescribed for dialysis or cellulitis in accordance with	a Health NZ Hos	pital approved	protocol and the prescription is
endorsed accordingly.			p p p
Inj 500 mg vial	3.39	5	✓ Cefazolin-AFT
lnj 1 g vial		5	✓ Cefazolin-AFT
Inj 2 g vial	7.09	5	✓ Cefazolin-AFT
CEFTRIAXONE - Subsidy by endorsement			
a) Up to 10 inj available on a PSO			
b) Subsidised only if prescribed for a dialysis or cystic fibros	is patient, or the	treatment of c	gonorrhoea, or the treatment of
pelvic inflammatory disease, or the treatment of suspected	ed meningococca	l disease, and	I the prescription or PSO is
endorsed accordingly.	•		
Inj 500 mg vial	0.79	1	✓ Ceftriaxone-AFT
Inj 1 g vial	3.59	5	✓ Ceftriaxone-AFT
CEFUROXIME AXETIL - Subsidy by endorsement			
Only if prescribed for prophylaxis of endocarditis and the pre	scription is endo	rsed according	alv.
Tab 250 mg		20	✓ Ascend-
•			Cefuroxime S29

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

Macrolides

AZITHROMYCIN – Maximum of 5 days treatment per prescription; can be waived by Special Authority see SA1683 below A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised on Special Authority.

Tab 250 mg	8.19	30	✓ Apo-Azithromycin
Tab 500 mg - Up to 8 tab available on a PSO		2	✓ Zithromax
Grans for oral liq 200 mg per 5 ml (40 mg per ml) - Wastage			
claimable	16.97	15 ml	✓ Zithromax

⇒SA1683 Special Authority for Waiver of Rule

Initial application — (bronchiolitis obliterans syndrome, cystic fibrosis and atypical Mycobacterium infections) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following:

- 1 Patient has received a lung transplant, stem cell transplant, or bone marrow transplant and requires treatment for bronchiolitis obliterans syndrome*; or
- 2 Patient has received a lung transplant and requires prophylaxis for bronchiolitis obliterans syndrome*; or
- 3 Patient has cystic fibrosis and has chronic infection with Pseudomonas aeruginosa or Pseudomonas-related gram negative organisms*; or
- 4 Patient has an atypical Mycobacterium infection.

Note: Indications marked with * are unapproved indications.

Initial application — (non-cystic fibrosis bronchiectasis*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 For prophylaxis of exacerbations of non-cystic fibrosis bronchiectasis*; and
- 2 Patient is aged 18 and under; and
- 3 Fither:
 - 3.1 Patient has had 3 or more exacerbations of their bronchiectasis, within a 12 month period; or
 - 3.2 Patient has had 3 acute admissions to hospital for treatment of infective respiratory exacerbations within a 12 month period.

Note: Indications marked with * are unapproved indications.

Renewal — (non-cystic fibrosis bronchiectasis*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has completed 12 months of azithromycin treatment for non-cystic fibrosis bronchiectasis; and
- 2 Following initial 12 months of treatment, the patient has not received any further azithromycin treatment for non-cystic fibrosis bronchiectasis for a further 12 months, unless considered clinically inappropriate to stop treatment; and
- 3 The patient will not receive more than a total of 24 months' azithromycin cumulative treatment (see note).

The patient must not have had more than 1 prior approval.

Note: No further renewals will be subsidised. A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised. Indications marked with * are unapproved indications

CLARITHROMYCIN - Maximum of 500 mg per prescription; can be waived by Special Authority see SA1857 below

⇒SA1857 Special Authority for Waiver of Rule

Initial application — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years for applications meeting the following criteria:

Fither:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1 Atypical mycobacterial infection; or
 - 2 Mycobacterium tuberculosis infection where there is drug-resistance or intolerance to standard pharmaceutical agents.

Initial application — (Helicobacter pylori eradication) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 For the eradication of helicobacter pylori in a patient unable to swallow tablets; and
- 2 For use only in combination with omeprazole and amoxicillin as part of a triple therapy regimen.

Initial application — (Prophylaxis of infective endocarditis) from any relevant practitioner. Approvals valid for 3 months where prophylaxis of infective endocarditis associated with surgical or dental procedures if amoxicillin is contra-indicated. Renewal — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

ERYTHROMYCIN (AS LACTOBIONATE) Inj 1 g vial	10.00	1	✓ Erythrocin IV
ERYTHROMYCIN ETHYL SUCCINATE			
Tab 400 mg	16.95	100	✓ E-Mycin
a) Up to 20 tab available on a PSOb) Up to 2 x the maximum PSO quantity for RFPP			
Grans for oral liq 200 mg per 5 ml	5.00	100 ml	✓ E-Mycin
a) Up to 300 ml available on a PSO			
b) Up to 2 x the maximum PSO quantity for RFPPc) Wastage claimable			
Grans for oral liq 400 mg per 5 ml	6.77	100 ml	✓ E-Mycin
a) Up to 200 ml available on a PSO			
b) Wastage claimable			
ROXITHROMYCIN Tob 150 mg	10.10	50	Аннани
Tab 150 mg	13.19	50	Arrow- Roxithromycin
Tab 300 mg	25.00	50	✓ Arrow-
			Roxithromycin

	Subsidy (Manufacturer's Price \$	e) Per	Fully Subsidised	
Penicillins				
AMOXICILLIN				
Cap 250 mg	27.50	500	1	Miro-Amoxicillin
	43.45		•	Alphamox
a) Up to 30 cap available on a PSO				
b) Up to 10 x the maximum PSO quantity for RFPP				
c) Miro-Amoxicillin to be Principal Supply on 1 Septemb				
Cap 500 mg		500		Miro-Amoxicillin
	66.44		•	Alphamox
a) Up to 30 cap available on a PSO				
b) Up to 10 x the maximum PSO quantity for RFPP	004			
c) Miro-Amoxicillin to be Principal Supply on 1 August 2 Grans for oral liq 125 mg per 5 ml	2 22	100 m		Alphamox 125
a) Up to 200 ml available on a PSO		100 111	. •	Alphaniox 123
b) Wastage claimable				
Grans for oral liq 250 mg per 5 ml	2.81	100 m	· •	Alphamox 250
a) Up to 300 ml available on a PSO			•	- III
b) Up to 10 x the maximum PSO quantity for RFPP				
c) Wastage claimable				
Inj 250 mg vial	15.97	10	✓	Ibiamox
Inj 500 mg vial		10		Ibiamox
Inj 1 g vial – Up to 5 inj available on a PSO	21.64	10	1	Ibiamox
(Alphamox Cap 250 mg to be delisted 1 September 2024)				
(Alphamox Cap 500 mg to be delisted 1 August 2024)				
AMOXICILLIN WITH CLAVULANIC ACID				
Tab 500 mg with clavulanic acid 125 mg - Up to 30 tab				
available on a PSO		10	1	Curam Duo 500/125
Grans for oral liq amoxicillin 25 mg with clavulanic acid 6.25 r				
per ml	6.50	100 m	· •	Augmentin
a) Up to 200 ml available on a PSO				
b) Wastage claimable				
Grans for oral liq amoxicillin 50 mg with clavulanic acid 12.5 r	mg			_
per ml - Up to 200 ml available on a PSO	2.20 1	100 ml ()P 🗸	Curam
BENZATHINE BENZYLPENICILLIN				
Inj 900 mg (1.2 million units) in 2.3 ml syringe – Up to 5 inj			_	
available on a PSO	375.97	10	/	Bicillin LA
BENZYLPENICILLIN SODIUM [PENICILLIN G]				
Inj 600 mg (1 million units) vial – Up to 5 inj available on a Ps	SO 16.50	10	1	Sandoz

	Subsidy (Manufacturer's Price		Fully Brand or esidised Generic
	\$	Per	✓ Manufacturer
LUCLOXACILLIN			
Cap 250 mg - Up to 30 cap available on a PSO	15.79	250	✓ Flucloxacillin-AFT
Cap 500 mg - Up to 30 cap available on a PSO	52.99	500	✓ Flucloxacillin-AFT
Grans for oral liq 25 mg per ml	3.29	100 ml	✓ <u>AFT</u>
a) Up to 200 ml available on a PSO			
b) Wastage claimable			
Grans for oral liq 50 mg per ml	3.68	100 ml	✓ <u>AFT</u>
a) Up to 200 ml available on a PSO			
b) Wastage claimable			
Inj 250 mg vial	42.60	10	✓ Flucloxin
Flucloxin to be Principal Supply on 1 July 2024			
Inj 500 mg vial	45.63	10	✓ Flucloxin
Flucloxin to be Principal Supply on 1 July 2024			_
Inj 1 g vial – Up to 5 inj available on a PSO	6.00	5	✓ Flucil
HENOXYMETHYLPENICILLIN (PENICILLIN V)			
Cap 250 mg - Up to 30 cap available on a PSO	3.84	50	✓ Cilicaine VK
Cap 500 mg	6.86	50	✓ Cilicaine VK
a) Up to 20 cap available on a PSO			
b) Up to 2 x the maximum PSO quantity for RFPP			
Grans for oral liq 125 mg per 5 ml	3.40	100 ml	✓ AFT
a) Up to 200 ml available on a PSO			
b) Wastage claimable			
Grans for oral liq 250 mg per 5 ml	4.24	100 ml	✓ <u>AFT</u>
a) Up to 300 ml available on a PSO			
b) Up to 2 x the maximum PSO quantity for RFPP			
c) Wastage claimable			
, , , , , , , , , , , , , , , , , , , ,			

Tetracyclines

DOXYCYCLINE

*	Tab 100 mg - Up to 30 tab available on a PSO	64.43	500	✓ Doxine
MII	NOCYCLINE HYDROCHLORIDE			
*	Tab 50 mg - Additional subsidy by Special Authority see			
	SA1355 below – Retail pharmacy	5.79	60	
		(12.05)		Mino-tabs
*	Cap 100 mg	19.32	100	
		(52.04)		Minomycin

⇒SA1355 Special Authority for Manufacturers Price

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has rosacea.

TETRACYCLINE - Special Authority see SA1332 below - Retail pharmacy
Tab 250 mg58.20 28 ✓ Accord S29

⇒SA1332 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 For the eradication of helicobacter pylori following unsuccessful treatment with appropriate first-line therapy; and
- 2 For use only in combination with bismuth as part of a quadruple therapy regimen.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

Other Antibiotics

For topical antibiotics, refer to DERMATOLOGICALS, page 68

CIPROFLOXACIN

Recommended for patients with any of the following:

- i) microbiologically confirmed and clinically significant pseudomonas infection; or
- ii) prostatitis; or
- iii) pyelonephritis; or
- iv) gonorrhoea.

Tab 250 mg - Up to 5 tab available on a PSO	1.95	28	✓ Ipca-Ciprofloxacin
•	2.42		✓ Cipflox
	3.85	10	✓ Ciprofloxacin -
			Torrent S29
Tab 500 mg - Up to 5 tab available on a PSO	3.10	28	✓ Ipca-Ciprofloxacin
	4.25	10	✓ Ciprofloxacin -
			Torrent S29
Tab 750 mg	5.95	28	✓ Cipflox
CLINDAMYCIN			
Cap hydrochloride 150 mg	5.30	24	✓ Dalacin C
Inj 150 mg per ml, 4 ml ampoule	35.10	10	✓ <u>Hameln</u>
COLISTIN SULPHOMETHATE - Retail pharmacy-Specialist - S	ubsidy by endors	ement	
Only if prescribed for dialysis or cystic fibrosis patient and the			ordingly.
Inj 150 mg		1	✓ Colistin-Link
GENTAMICIN SULPHATE			
Inj 10 mg per ml, 1 ml ampoule - Subsidy by endorsement	95.00	5	✓ DBL Gentamicin
Only if prescribed for a dialysis or cystic fibrosis patient of endorsed accordingly.	or complicated uri	nary tract inf	fection and the prescription is
Inj 10 mg per ml, 2 ml ampoule - Subsidy by endorsement	91.00	5	✓ Wockhardt S29
	182.00	10	✓ Teligent S29
Only if prescribed for a dialysis or cystic fibrosis patient of endorsed accordingly.	or complicated uri	nary tract inf	fection and the prescription is
Inj 40 mg per ml, 2 ml ampoule - Subsidy by endorsement	18.38	10	✓ Pfizer
	87.50	50	✓ Pfizer
Only if prescribed for a dialysis or cystic fibrosis patient of endorsed accordingly.	or complicated uri	nary tract inf	fection and the prescription is
(Teligent \$29 Inj 10 mg per ml, 2 ml ampoule to be delisted 1 At	uaust 2024)		
MOXIFLOXACIN – Special Authority see SA1740 below – Retail	,		
INIONITEONACIN — Special Authority see SA1740 below — netali	priarriacy		

⇒SA1740 Special Authority for Subsidy

Tab 400 mg42.00

No patient co-payment payable

Initial application — (Tuberculosis) only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

1 Both:

1.1 Active tuberculosis*: and

continued...

5

✓ Avelox

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic	
\$	Per	✓	Manufacturer	

continued...

- 1.2 Any of the following:
 - 1.2.1 Documented resistance to one or more first-line medications; or
 - 1.2.2 Suspected resistance to one or more first-line medications (tuberculosis assumed to be contracted in an area with known resistance), as part of regimen containing other second-line agents; or
 - 1.2.3 Impaired visual acuity (considered to preclude ethambutol use); or
 - 1.2.4 Significant pre-existing liver disease or hepatotoxicity from tuberculosis medications; or
 - 1.2.5 Significant documented intolerance and/or side effects following a reasonable trial of first-line medications: ٥r
- 2 Mycobacterium avium-intracellulare complex not responding to other therapy or where such therapy is contraindicated.*; or
- 3 Patient is under five years of age and has had close contact with a confirmed multi-drug resistant tuberculosis case.

Note: Indications marked with * are unapproved indications.

Renewal only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Mycoplasma genitalium) only from a sexual health specialist or Practitioner on the recommendation of a sexual health specialist. Approvals valid for 1 month for applications meeting the following criteria: All of the following:

- 1 Has nucleic acid amplification test (NAAT) confirmed Mycoplasma genitalium* and is symptomatic; and
- - 2.1 Has tried and failed to clear infection using azithromycin: or
 - 2.2 Has laboratory confirmed azithromycin resistance; and
 - 3 Treatment is only for 7 days.

Initial application — (Penetrating eye injury) only from an ophthalmologist. Approvals valid for 1 month where the patient requires prophylaxis following a penetrating eye injury and treatment is for 5 days only.

Note: Indications marked with * are unapproved indications.

PAROMOMYCIN - Special Authority see SA1689 below - Retail pharmacy

16 ✓ Humatin S29

⇒SA1689 Special Authority for Subsidy

Initial application only from an infectious disease specialist, clinical microbiologist or gastroenterologist. Approvals valid for 1 month for applications meeting the following criteria: Either:

- 1 Patient has confirmed cryptosporidium infection; or
- 2 For the eradication of Entamoeba histolyica carriage.

Renewal only from an infectious disease specialist, clinical microbiologist or gastroenterologist. Approvals valid for 1 month for applications meeting the following criteria:

Fither:

- 1 Patient has confirmed cryptosporidium infection; or
- 2 For the eradication of Entamoeba histolyica carriage.

PYRIMETHAMINE - Special Authority see SA1328 below - Retail pharmacy

30 ✓ Daraprim S29

⇒SA1328 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 For the treatment of toxoplasmosis in patients with HIV for a period of 3 months; or
- 2 For pregnant patients for the term of the pregnancy; or
- 3 For infants with congenital toxoplasmosis until 12 months of age.

SODIUM FUSIDATE IFUSIDIC ACIDI

36 ✓ Fucidin

ULFADIAZINE SODIUM – Special Authority see SA1331 below Tab 500 mg SA1331 Special Authority for Subsidy iitial application from any relevant practitioner. Approvals valid	543.20	56		Wockhardt S29
»SA1331 Special Authority for Subsidy itial application from any relevant practitioner. Approvals valid				Wockhardt \$29
itial application from any relevant practitioner. Approvals valid	d without further rei	newal unle		
	d without further rei	newal unle		
			ess notif	fied for applications meetin
ne following criteria:				
ny of the following:				
1 For the treatment of toxoplasmosis in patients with HIV for	r a period of 3 mon	ins; or		
2 For pregnant patients for the term of the pregnancy; or3 For infants with congenital toxoplasmosis until 12 months	of ago			
· ·	or age.			
OBRAMYCIN		_		
Inj 40 mg per ml, 2 ml vial – Subsidy by endorsement		5		Tobramycin (Viatris)
Only if prescribed for dialysis or cystic fibrosis patient and	ia the prescription is	s endorse	a accord	aingiy.
Solution for inhalation 60 mg per ml, 5 ml – Subsidy by	205.00	56 dose	./	Tahramusin DNM
endorsement	395.00	oo dose	V	Tobramycin BNM
a) Wastage claimableb) Only if prescribed for a cystic fibrosis patient and the	proporintian is and	araad aaa	ordinalu	
	prescription is end	Jiseu acc	orumgiy	•
RIMETHOPRIM	40.55	50	,	THE
Fab 300 mg – Up to 30 tab available on a PSO		50	•	<u>TMP</u>
RIMETHOPRIM WITH SULPHAMETHOXAZOLE [CO-TRIMOX)	•			
* Tab trimethoprim 80 mg and sulphamethoxazole 400 mg - L				
to 30 tab available on a PSO		500	•	<u>Trisul</u>
FOral liq 8 mg sulphamethoxazole 40 mg per ml — Up to 200 r		4001	,	D
available on a PSO	2.97	100 ml	•	Deprim
ANCOMYCIN – Subsidy by endorsement			, .	
Only if prescribed for a dialysis or cystic fibrosis patient or for			or for tre	eatment of Clostridium
difficile following metronidazole failure and the prescription is Inj 500 mg vial		igiy. 1	1	Mylan
IIIJ 500 IIIg Viai		'		<u>iviyiaii</u>
Antifungals				
Artificatiguis				
For topical antifungals refer to DERMATOLOGICALS, page 69	9			
For topical antifungals refer to GENITO URINARY, page 81				
LUCONAZOLE				
Cap 50 mg	4.10	28	1	Mylan
Cap 150 mg		1	✓	Mylan
Cap 200 mg		28	•	Mylan
Powder for oral suspension 10 mg per ml - Special Authority				
see SA1359 below – Retail pharmacy	129.02	35 ml	•	Diflucan
Wastage claimable				
»SA1359 Special Authority for Subsidy litial application — (Systemic candidiasis) from any relevant				

Initial application — (Systemic candidiasis) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

Initial application — (Immunocompromised) from any relevant practitioner. Approvals valid for 6 months for applications

Subsidy (Manufacturer's Price)	S	Fully ubsidised	Brand or Generic
\$	Per	✓	Manufacturer

continued...

meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Patient is at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

Renewal — (Systemic candidiasis) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

Renewal — (Immunocompromised) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient remains immunocompromised; and
- 2 Patient remains at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

ITRACONAZOLE

Cap 100 mg6.83	15	Itrazole
Oral liq 10 mg per ml - Special Authority see SA1322 below -		
Retail pharmacy141.80	150 ml OP	Sporanox

⇒SA1322 Special Authority for Subsidy

Initial application only from an infectious disease specialist, clinical microbiologist, clinical immunologist or any relevant practitioner on the recommendation of a infectious disease physician, clinical microbiologist or clinical immunologist. Approvals valid for 6 months where the patient has a congenital immune deficiency.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefitting from the treatment.

KETOCONAZOLE

Tab 200 mg - PCTCBS	30 100	✓ Burel \$29 ✓ Strides Shasun \$29 ✓ Taro \$29 ✓ Teva- Ketoconazole \$29
NYSTATIN		
Tab 500,000 u14.16	50	
(17.09)		Nilstat
Cap 500,000 u12.81	50	
(15.47)		Nilstat
POSACONAZOLE - Special Authority see SA1285 below - Retail pharmacy		
Tab modified-release 100 mg206.00	24	✓ Posaconazole Juno
Oral liq 40 mg per ml342.51	105 ml OP	✓ <u>Devatis</u>
OA 4005 On a stall A sub-sub-sub-state		

⇒SA1285 Special Authority for Subsidy

Initial application only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

Either:

Subsidy		Fully	Brand or
(Manufacturer's Price	e)	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation chemotherapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppressive therapy*.

Renewal only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

Fither:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation therapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppression* and requires on going posaconazole treatment.

Note: * Graft versus host disease (GVHD) on significant immunosuppression is defined as acute GVHD, grade II to IV, or extensive chronic GVHD, or if they were being treated with intensive immunosuppressive therapy consisting of either high-dose corticosteroids (1 mg or greater per kilogram of body weight per day for patients with acute GVHD or 0.8 mg or greater per kilogram every other day for patients with chronic GVHD), antithymocyte globulin, or a combination of two or more immunosuppressive agents or types of treatment.

TERBINAFINE

*	Tab 250 mg4.48		42	✓ Apo-Terbinafine S29
	8.97		84	✓ <u>Deolate</u>
VO	RICONAZOLE - Special Authority see SA1273 below - Retail pharmacy			
	Tab 50 mg91.00		56	✓ Vttack
	Tab 200 mg		56	✓ Vttack
	Powder for oral suspension 40 mg per ml – Wastage			
	claimable1,523.22	7	0 ml	✓ Vfend

⇒SA1273 Special Authority for Subsidy

Initial application — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
 - 3.1 Patient has proven or probable invasive aspergillus infection; or
 - 3.2 Patient has possible invasive aspergillus infection; or
 - 3.3 Patient has fluconazole resistant candidiasis; or
 - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Renewal — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
 - 3.1 Patient continues to require treatment for proven or probable invasive aspergillus infection; or
 - 3.2 Patient continues to require treatment for possible invasive aspergillus infection; or
 - 3.3 Patient has fluconazole resistant candidiasis: or
 - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per ✓	Manufacturer

Antimalarials

PRIMAQUINE - Special Authority see SA1684 below - Retail pharmac	у		
Tab 15 mg4	00.00 1	00 🗸	Sanofi
			Primaguine \$29

⇒SA1684 Special Authority for Subsidy

Initial application only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 The patient has vivax or ovale malaria; and
- 2 Primaquine is to be given for a maximum of 21 days.

Renewal only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 The patient has relapsed vivax or ovale malaria; and
- 2 Primaguine is to be given for a maximum of 21 days.

Antitrichomonal Agents

METRONIDAZOLE			
Tab 200 mg - Up to 30 tab available on a PSO	33.15	250	✓ Metrogyl
Tab 400 mg - Up to 15 tab available on a PSO	5.23	21	✓ Metrogyl
Oral liq benzoate 200 mg per 5 ml	25.00	100 ml	✓ Flagyl-S
Suppos 500 mg		10	✓ Flagyl
ORNIDAZOLE			
Tab 500 mg	36.16	10	✓ Arrow-Ornidazole

Antituberculotics and Antileprotics

Note: There is no co-payment charge for all pharmaceuticals listed in the Antituberculotics and Antileprotics group regardless of immigration status.

BEDAQUILINE - Special Authority see SA2244 below - Retail pharmacy

No patient co-payment payable

⇒SA2244 Special Authority for Subsidy

Initial application — (multi-drug resistant tuberculosis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The person has multi-drug resistant tuberculosis (MDR-TB); and
- 2 Ministry of Health's Tuberculosis Clinical Network has reviewed the individual case and recommends bedaquiline as part of the treatment regimen.

CLOFAZIMINE - Retail pharmacy-Specialist

- a) No patient co-payment payable
- b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or dermatologist.
- **★** Cap 50 mg........442.00 100 **✓ Lamprene** \$29

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	Generic
CYCLOSERINE - Retail pharmacy-Specialist	·			
 a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendat respiratory physician. 			_	_
Cap 250 mg	344.00	60	•	Cyclorin S29
DAPSONE – Retail pharmacy-Specialist				
 a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendat dermatologist 	ion of, an infectious d	iseas	e physicia	n, clinical microbiologist or
Tab 25 mg		100	_	Dapsone
Tab 100 mg	329.50	100	/	Dapsone
ETHAMBUTOL HYDROCHLORIDE - Retail pharmacy-Specialis	st			
a) No patient co-payment payableb) Prescriptions must be written by, or on the recommendat respiratory physician		iseas	e physicia	n, clinical microbiologist or
Tab 100 mg		100		EMB Fatol \$29
Tab 400 mg	49.34	56	/	Myambutol S29
ISONIAZID – Retail pharmacy-Specialist				
 a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendat microbiologist, dermatologist or public health physician 	ion of, an internal me	dicine	physician	ı, paediatrician, clinical
* Tab 100 mg	23.00	100	•	<u>PSM</u>
ISONIAZID WITH RIFAMPICIN - Retail pharmacy-Specialist				
 a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendat microbiologist, dermatologist or public health physician * Tab 100 mg with rifampicin 150 mg * Tab 150 mg with rifampicin 300 mg LINEZOLID – Special Authority see SA2234 below – Retail phar 	89.82 179.13	dicine 100 100	· ·	, paediatrician, clinical Rifinah Rifinah
No patient co-payment payable	macy			
Tab 600 mg	276.89	10	1	Zyvox
Oral liq 20 mg per ml		150 m		Zyvox
■ SA2234 Special Authority for Subsidy Initial application — (multi-drug resistant tuberculosis) from applications meeting the following criteria: Both: 1 The person has multi-drug resistant tuberculosis (MDR-Te		ner.	Approvals	valid for 18 months for
Ministry of Health's Tuberculosis Clinical Network has rev the treatment regimen.		ase a	and recom	mends linezolid as part of
PARA-AMINO SALICYLIC ACID – Retail pharmacy-Specialist				
 a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendat respiratory physician 	ion of, an infectious d	iseas	e specialis	st, clinical microbiologist or
Grans for oral liq 4 g sachet	280.00	30	✓	Paser S29
PROTIONAMIDE - Retail pharmacy-Specialist				
a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendat respiratory physician	ion of, an infectious d	iseas	e specialis	st, clinical microbiologist or
Tab 250 mg	305.00	100	•	Peteha S29
A CH CARLON	000 11			

	ı	NFECTIONS - A	CENTS E	∩	SVSTEMIC LISE
	'	NI ECTIONS - A	GLNISI	on c	3131LIMIC USL
		Subsidy (Manufacturer's Price) \$		Fully lised	Brand or Generic Manufacturer
PY	RAZINAMIDE – Retail pharmacy-Specialist				
	a) No patient co-payment payable				
	 Prescriptions must be written by, or on the recommendat respiratory physician 	ion of, an infectious d	lisease phys	ician,	clinical microbiologist or
*	Tab 500 mg	64.95	100	✓ A	FT-Pyrazinamide
RIF	ABUTIN - Retail pharmacy-Specialist				
	a) No patient co-payment payable				
	b) Prescriptions must be written by, or on the recommendat gastroenterologist	ion of, an infectious d	lisease phys	ician,	respiratory physician or
*	Cap 150 mg	353.71	30	✓ M	lycobutin
RIF	FAMPICIN - Subsidy by endorsement				
	a) No patient co-payment payable				
	b) For confirmed recurrent Staphylococcus aureus infection	in combination with o	other effective	e anti	i-staphylococcal
	antimicrobial based on susceptibilities and the prescription				
	Retail pharmacy - Specialist. Specialist must be an inter	nal medicine physicia	ın, clinical m	icrobio	ologist, dermatologist,
	paediatrician, or public health physician.				
*	Cap 150 mg		100	_	Rifadin
*	Cap 300 mg	122.06	100	_	Rifadin
*	Oral lig 100 mg nor 5 ml	10.60	60 ml		lifadin Sanofi lifadin
*	Oral liq 100 mg per 5 ml	12.00	00 1111	• <u>n</u>	<u>maum</u>
Α	ntivirals				
Ear	ave preparations refer to Eve Preparations, Anti-Infective Pre	parations nage 250			

For eye preparations refer to Eye Preparations, Anti-Infective Preparations, page 259

Hepatitis B	Treatment
-------------	-----------

ENTECAVIR			
* Tab 0.5 mg	12.04	30	✓ Entecavir (Rex)
LAMIVUDINE - Special Authority see SA1685 below	 Retail pharmacy 		
Tab 100 mg	12.06	28	✓ Zetlam
Oral liq 5 mg per ml	270.00	240 ml OP	✓ Zeffix

⇒SA1685 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 1 year where used for the treatment or prevention of hepatitis B.

Renewal from any relevant practitioner. Approvals valid for 2 years where used for the treatment or prevention of hepatitis B. TENOFOVIR DISOPROXIL

Tenofovir disoproxil prescribed under endorsement for the treatment of HIV is included in the count of up to 4 subsidised antiretrovirals for the purposes of Special Authority SA2139., page 110

•••	rab 2 to mg (ood mg as a maisais)		 0.00		Viatris
*	Tab 245 mg (300 mg as a maleate)	,	5.00 .3	30	Tenofovir Disoproxil

Herpesvirus Treatments

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price) \$	Su Per	Fully bsidised	Brand or Generic Manufacturer
VALGANCICLOVIR – Special Authority see SA1993 below – Re Tab 450 mg	, ,	60		alganciclovir Viatris

⇒SA1993 Special Authority for Subsidy

Initial application — (transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 3 months where the patient has undergone a solid organ transplant and requires valganciclovir for CMV prophylaxis.

Renewal — (transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has undergone a solid organ transplant and received anti-thymocyte globulin and requires valganciclovir therapy for CMV prophylaxis; and
 - 1.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following anti-thymocyte globulin; or
- 2 Both:
 - 2.1 Patient has received pulse methylprednisolone for acute rejection and requires further valganciclovir therapy for CMV prophylaxis; and
 - 2.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following pulse methylprednisolone.

Initial application — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has undergone a solid organ transplant and received valganciclovir under Special Authority more than 2 years ago (27 months); and
- 2 Patient has received anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Renewal — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months where the patient has received a further course of anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Initial application — (Lung transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone a lung transplant; and
- 2 Fither:
 - 2.1 The donor was cytomegalovirus positive and the patient is cytomegalovirus negative; or
 - 2.2 The recipient is cytomegalovirus positive; and
- 3 Patient has a high risk of CMV disease.

Initial application — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
 - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
 - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
 - 2.3 Patient has cytomegalovirus retinitis.

Renewal — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
 - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or

continued...

- 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
- 2.3 Patient has cytomegalovirus retinitis.

Note: for the purpose of this Special Authority "immunocompromised" includes transplant recipients, patients with immunosuppressive diseases (e.g. HIV) or those receiving immunosuppressive treatment for other conditions.

Hepatitis C Treatment

GLECAPREVIR WITH PIBRENTASVIR - [Xpharm]

Note the supply of treatment is via Pharmac's approved direct distribution supply. Further details can be found on Pharmac's website https://pharmac.govt.nz/maviret

Tab 100 mg with pibrentasvir 40 mg24,750.00 84 OP ✓ Maviret

LEDIPASVIR WITH SOFOSBUVIR - [Xpharm] - Special Authority see SA1605 below

No patient co-payment payable

⇒SA1605 Special Authority for Subsidy

Special Authority approved by the Hepatitis C Treatment Panel (HepCTP)

Notes: By application to the Hepatitis C Treatment Panel (HepCTP).

Applications will be considered by HepCTP and approved subject to confirmation of eligibility.

Application details may be obtained from Pharmac's website http://www.pharmac.govt.nz/maviret or:

The Coordinator, Hepatitis C Treatment Panel

Pharmac, PO Box 10-254, WELLINGTON Tel: (04) 460 4990,

Email: hepcpanel@pharmac.govt.nz

HIV Prophylaxis and Treatment

EMTRICITABINE WITH TENOFOVIR DISOPROXIL - Subsidy by endorsement; can be waived by Special Authority see SA2138 below

- a) Funding for emtricitabine with tenofovir disoproxil for use as PrEP, should be applied using Special Authority SA2138.
- b) Endorsement for treatment of conditions approved via Special Authority SA2139 (antiretrovirals for confirmed HIV, prevention of maternal transmission, post-exposure prophylaxis following exposure to HIV and percutaneous exposure): Prescription is deemed to be endorsed if emtricitabine with tenofovir disoproxil is co-prescribed with another antiretroviral subsidised under Special Authority SA2139 and the prescription is annotated accordingly by the Pharmacist or endorsed by the prescriber.

Note: Emtricitabine with tenofovir disoproxil prescribed under endorsement, for treatment of conditions approved via Special Authority SA2139 (antiretrovirals for confirmed HIV, prevention of maternal transmission, post-exposure prophylaxis following exposure to HIV and percutaneous exposure), is included in the count of up to 4 subsidised antiretrovirals, and counts as two antiretroviral medications, for the purposes of Special Authority SA2139, page 110 There is an approval process to become a named specialist to prescribe antiretroviral therapy in New Zealand. Further information is available on the Pharmac website.

* Tab 200 mg with tenofovir disoproxil 245 mg (300 mg as a

⇒SA2138 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 Patient has tested HIV negative, does not have signs or symptoms of acute HIV infection and has been assessed for HIV seroconversion; and
- 2 The Practitioner considers the patient is at elevated risk of HIV exposure and use of PrEP is clinically appropriate.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer

continued...

Notes: Refer to local health pathways or the Australasian Society for HIV. Viral Hepatitis and Sexual Health Medicine clinical

https://ashm.org.au/HIV/PrEP/

Renewal from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 Patient has tested HIV negative, does not have signs or symptoms of acute HIV infection and has been assessed for HIV seroconversion: and
- 2 The Practitioner considers the patient is at elevated risk of HIV exposure and use of PrEP is clinically appropriate.

Notes: Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical auidelines:

https://ashm.org.au/HIV/PrEP/

COVID-19 Treatments

MOLNUPIRAVIR - [Xpharm] - Subsidy by endorsement

- a) No patient co-payment payable
- b) Treatment is funded only if patient meets access criteria for oral antiviral COVID-19 treatments (as on Pharmac's website) and has been endorsed accordingly by the prescriber. The supply of treatment is via Pharmac's approved distribution process. Refer to the Pharmac website for more information about this and stock availability.

Lagevrio

NIRMATRELVIR WITH RITONAVIR - [Xpharm] - Subsidy by endorsement

- a) No patient co-payment payable
- b) Treatment is funded only if patient meets access criteria for oral antiviral COVID-19 treatments (as on Pharmac's website) and has been endorsed accordingly by the prescriber. The supply of treatment is via Pharmac's approved distribution process. Refer to the Pharmac website for more information about this and stock availability.

Paxlovid

Antiretrovirals

⇒SA2139 Special Authority for Subsidy

Initial application — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the patient has confirmed HIV infection.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Prevention of maternal transmission) only from a named specialist. Approvals valid for 1 year for applications meeting the following criteria:

Fither:

- 1 Prevention of maternal foetal transmission; or
- 2 Treatment of the newborn for up to eight weeks.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
 \$	Per	1	Manufacturer

continued...

purpose of accessing funding to antiretrovirals.

Some antiretrovirals are unapproved or contraindicated for this indication. Practitioners prescribing these medications should exercise their own skill, judgement, expertise and discretion, and make their own prescribing decisions with respect to the use of a Pharmaceutical for an indication for which it is not approved or contraindicated.

Initial application — (post-exposure prophylaxis following exposure to HIV) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had condomless anal intercourse or receptive vaginal intercourse with a known HIV positive person with an unknown or detectable viral load greater than 200 copies per ml; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required; or
 - 2.4 Patient has had condomless anal intercourse with a person from a high HIV prevalence country or risk group whose HIV status is unknown.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines for PEP (https://www.ashm.org.au/hiv/hiv-management/pep/).

Renewal — (second or subsequent post-exposure prophylaxis) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had condomless anal intercourse or receptive vaginal intercourse with a known HIV positive person with an unknown or detectable viral load greater than 200 copies per ml; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required; or
 - 2.4 Patient has had condomless anal intercourse with a person from a high HIV prevalence country or risk group whose HIV status is unknown.

Initial application — (Percutaneous exposure) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Second or subsequent percutaneous exposure) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Non-nucleosides Reverse Transcriptase Inhibitors

EFAVIRENZ - Special Authority see SA2139 on the	orevious page – Retail pharma	су	
Tab 200 mg	190.15	90	✓ Stocrin
Tab 600 mg	63.38	30	✓ Stocrin
· ·	65.38		✓ Efavirenz
			Milpharm S29

	Subsidy	C !	Fully	Brand or
	(Manufacturer's P	rice) Subs Per	idised •	Generic Manufacturer
TRAVIRINE - Special Authority see SA2139 on page 110 - F Tab 200 mg		60	✓	Intelence
IEVIRAPINE – Special Authority see SA2139 on page 110 – I Tab 200 mg		60		Nevirapine Alphapharm Nevirapine Viatris
Oral suspension 10 mg per ml	203.55	240 ml OP		Viramune Suspension
Nevirapine Alphapharm Tab 200 mg to be delisted 1 July 2024	1)			
Nucleosides Reverse Transcriptase Inhibitors				
ABACAVIR SULPHATE — Special Authority see SA2139 on particle Tab 300 mg		60 240 ml OP page 110 – Re	tail ph	
anti-retroviral Special Authority. Tab 600 mg with lamivudine 300 mg	29.50	30	•	Abacavir/ Lamivudine Viatris
TAVIDENZ WITH EMTDICITADINE AND TEMOCOVID DICC				
FAVIRENZ WITH EMTRICITABINE AND TENOFOVIR DISOI iharmacy Note: Efavirenz with emtricitabine and tenofovir disoproxil anti-retroviral Special Authority	counts as three an	•		
harmacy Note: Efavirenz with emtricitabine and tenofovir disoproxil anti-retroviral Special Authority Tab 600 mg with emtricitabine 200 mg and tenofovir disopr 245 mg (300 mg as a maleate)	counts as three an	iti-retroviral me	dicatio	
Narmacy Note: Efavirenz with emtricitabine and tenofovir disoproxil anti-retroviral Special Authority Tab 600 mg with emtricitabine 200 mg and tenofovir disopr 245 mg (300 mg as a maleate)	counts as three an oxil	iti-retroviral me	dicatio	ons for the purposes of th
Narmacy Note: Efavirenz with emtricitabine and tenofovir disoproxil anti-retroviral Special Authority Tab 600 mg with emtricitabine 200 mg and tenofovir disopr 245 mg (300 mg as a maleate)	counts as three and oxil	ati-retroviral me	dication	ons for the purposes of th
Narmacy Note: Efavirenz with emtricitabine and tenofovir disoproxil anti-retroviral Special Authority Tab 600 mg with emtricitabine 200 mg and tenofovir disopr 245 mg (300 mg as a maleate) EMTRICITABINE – Special Authority see SA2139 on page 110 Cap 200 mg AMIVUDINE – Special Authority see SA2139 on page 110 – I Tab 150 mg	counts as three and coxil	30 30 30 30 30 60 240 ml OP	dication	ons for the purposes of th Viatris Emtriva Lamivudine Viatris
Note: Efavirenz with emtricitabine and tenofovir disoproxil anti-retroviral Special Authority Tab 600 mg with emtricitabine 200 mg and tenofovir disoproxil 245 mg (300 mg as a maleate) EMTRICITABINE — Special Authority see SA2139 on page 110 Cap 200 mg AMIVUDINE — Special Authority see SA2139 on page 110 — Tab 150 mg	counts as three and coxil	30 30 30 30 30 40 60 240 ml OP nacy 100 200 ml OP ne 110 – Retail	dication	viatris Emtriva Lamivudine Viatris BTC Retrovir Retrovir
Narmacy Note: Efavirenz with emtricitabine and tenofovir disoproxil anti-retroviral Special Authority Tab 600 mg with emtricitabine 200 mg and tenofovir disoproxil 245 mg (300 mg as a maleate)	counts as three and coxil	30 30 30 30 30 40 60 240 ml OP nacy 100 200 ml OP ne 110 – Retail	dicatic	viatris Emtriva Lamivudine Viatris BTC Retrovir Retrovir

Protease Inhibitors

ATAZANAVIR SULPHATE - Special Authority see SA2139 of	n page 110 – Retail p	oharmacy	
Cap 150 mg	85.00	60	 Atazanavir Mylan
Cap 200 mg	110.00	60	✓ Atazanavir Mylan
			✓ Atazanavir Viatris

(Atazanavir Mylan Cap 200 mg to be delisted 1 December 2024)

	Subsidy (Manufacturer's Price)	Subs	Fully	Brand or Generic
	\$	Per	1	Manufacturer
DARUNAVIR - Special Authority see SA2139 on page 110 - Re	etail pharmacy			
Tab 400 mg		60	✓ [Darunavir Viatris
Tab 600 mg	225.00	60	✓ [Darunavir Viatris
OPINAVIR WITH RITONAVIR – Special Authority see SA2139	on page 110 - Retail	pharmacy	/	
Tab 100 mg with ritonavir 25 mg	150.00	60	✓ <u>L</u>	opinavir/Ritonavir Mylan
Tab 200 mg with ritonavir 50 mg	295.00	120	✓ <u>L</u>	opinavir/Ritonavir Mylan
RITONAVIR – Special Authority see SA2139 on page 110 – Re	tail pharmacy			
Tab 100 mg	43.31	30	✓ N	lorvir
Strand Transfer Inhibitors				

DOLUTEGRAVIR – Special Authority see SA2139 on page	110 – Retail pharmacy	/	
Tab 50 mg	1,090.00	30	Tivicay
DOLUTEGRAVIR WITH LAMIVUDINE - Special Authority s	see SA2139 on page 1	10 – Retail p	oharmacy
Tab 50 mg with lamivudine 300 mg	1,090.00	30	✓ Dovato
RALTEGRAVIR POTASSIUM - Special Authority see SA21	139 on page 110 – Reta	ail pharmacy	/
Tab 400 mg	1,090.00	60	✓ Isentress
Tab 600 mg	1.090.00	60	✓ Isentress HD

Immune Modulators

PEGYLATED INTERFERON ALFA-2A - Special Authority see SA2034 below - Retail pharmacy

Note: Pharmac will consider funding ribavirin for the small group of patients who have a clinical need for ribavirin and meet Special Authority criteria. Please contact the Hepatitis C Coordinator at Pharmac on 0800-023-588 option 4. Inj 180 mcg prefilled syringe......748.50 ✓ Pegasys

⇒SA2034 Special Authority for Subsidy

Initial application — (chronic hepatitis C - genotype 1, 4, 5 or 6 infection or co-infection with HIV or genotype 2 or 3 post liver transplant) from any specialist. Approvals valid for 18 months for applications meeting the following criteria: Both:

- 1 Any of the following:
 - 1.1 Patient has chronic hepatitis C. genotype 1, 4, 5 or 6 infection; or
 - 1.2 Patient has chronic hepatitis C and is co-infected with HIV; or
 - 1.3 Patient has chronic hepatitis C genotype 2 or 3 and has received a liver transplant; and
- 2 Maximum of 48 weeks therapy.

Renewal — (Chronic hepatitis C - genotype 1 infection) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- - 3.1 Patient has responder relapsed; or
 - 3.2 Patient was a partial responder; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application — (Chronic Hepatitis C - genotype 1 infection treatment more than 4 years prior) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Su	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

following criteria:

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Any of the following:
 - 3.1 Patient has responder relapsed; or
 - 3.2 Patient was a partial responder; or
 - 3.3 Patient received interferon treatment prior to 2004; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application — (chronic hepatitis C - genotype 2 or 3 infection without co-infection with HIV) from any specialist. Approvals valid for 12 months for applications meeting the following criteria:

- 1 Patient has chronic hepatitis C, genotype 2 or 3 infection; and
- 2 Maximum of 6 months therapy.

Initial application — (Hepatitis B) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

Both:

- 1 Patient has confirmed Hepatitis B infection (HBsAg positive for more than 6 months); and
- 2 Patient is Hepatitis B treatment-naive; and
- 3 ALT > 2 times Upper Limit of Normal; and
- 4 HBV DNA < 10 log10 IU/ml; and
- 5 Either:
 - 5.1 HBeAg positive; or
 - 5.2 serum HBV DNA greater than or equal to 2.000 units/ml and significant fibrosis (Metavir Stage F2 or greater or moderate fibrosis); and
- 6 Compensated liver disease; and
- 7 No continuing alcohol abuse or intravenous drug use; and
- 8 Not co-infected with HCV, HIV or HDV; and
- 9 Neither ALT nor AST > 10 times upper limit of normal; and
- 10 No history of hypersensitivity or contraindications to pegylated interferon; and
- 11 Maximum of 48 weeks therapy.

Initial application — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 Patient has a cutaneous T cell lymphoma*; or
- 2 All of the following:
 - 2.1 Patient has a myeloproliferative disorder*; and
 - 2.2 Patient is intolerant of hydroxyurea; and
 - 2.3 Treatment with an agrelide and busulfan is not clinically appropriate; or
- 3 Both:
 - 3.1 Patient has a myeloproliferative disorder; and
 - 3.2 Patient is pregnant, planning pregnancy or lactating.

Renewal — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 No evidence of disease progression; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sul	bsidised	Generic	
\$	Per	•	Manufacturer	

continued...

- 2 The treatment remains appropriate and patient is benefitting from treatment; and
- 3 Either
 - 3.1 Patient has a cutaneous T cell lymphoma*: or
 - 3.2 Both:
 - 3.2.1 Patient has a myeloproliferative disorder*; and
 - 3.2.2 Fither:
 - 3.2.2.1 Remains intolerant of hydroxyurea and treatment with anagrelide and busulfan remains clinically inappropriate; or
 - 3.2.2.2 Patient is pregnant, planning pregnancy or lactating.

Note: Indications marked with * are unapproved indications.

Initial application — **(post-allogenic bone marrow transplant)** from any relevant practitioner. Approvals valid for 3 months where patient has received an allogeneic bone marrow transplant* and has evidence of disease relapse.

Renewal — (post-allogenic bone marrow transplant) from any relevant practitioner. Approvals valid for 3 months where patient is responding and ongoing treatment remains appropriate.

Note: Indications marked with * are unapproved indications.

Urinary Tract Infections

METHENAMINE (HEXAMINE) HIPPURATE		
* Tab 1 g	9.95 100	✓ Hiprex
NITROFURANTOIN		
* Tab 50 mg - Up to 30 tab available on a PSO22	2.20 100	✓ Nifuran
* Tab 100 mg	'.50 100	✓ <u>Nifuran</u>
* Cap modified-release 100 mg - Up to 15 cap available on a		
PSO81	.20 100	✓ Macrobid
NORFLOXACIN		
Tab 400 mg - Subsidy by endorsement245		✓ Arrow-Norfloxacin

Only if prescribed for a patient with an uncomplicated urinary tract infection that is unresponsive to a first line agent or with proven resistance to first line agents and the prescription is endorsed accordingly.

	Subsidy		Fully	
	(Manufacturer's Price) \$	Per	Subsidised	Generic Manufacturer
	Ψ	1 01		Manadactic
Anticholinesterases				
NEOSTIGMINE METILSULFATE				
Inj 2.5 mg per ml, 1 ml ampoule	33.81	10	1	Max Health
PYRIDOSTIGMINE BROMIDE				
▲ Tab 60 mg	50.28	100	✓	Mestinon
Non-Steroidal Anti-Inflammatory Drugs				
DICLOFENAC SODIUM				
# Tab EC 25 mg	1 99	50	1	Diclofenac Sandoz
* Tab 50 mg dispersible		20		Voltaren D
* Tab EC 50 mg		50		Diclofenac Sandoz
* Tab long-acting 75 mg		100		Voltaren SR
* Inj 25 mg per ml, 3 ml ampoule – Up to 5 inj available on a F		5		Voltaren
* Suppos 12.5 mg		10		Voltaren
* Suppos 25 mg		10		Voltaren
* Suppos 50 mg - Up to 10 supp available on a PSO		10		Voltaren
* Suppos 100 mg		10		Voltaren
	7.00	10	•	Voltaien
BUPROFEN	04.40	4 000	,	. "
* Tab 200 mg		1,000		Relieve
* Tab long-acting 800 mg		30		Brufen SR
* Oral liq 20 mg per ml	2.25	200 m	· •	<u>Ethics</u>
KETOPROFEN				
* Cap long-acting 200 mg	12.07	28	✓	Oruvail SR
MEFENAMIC ACID				
* Cap 250 mg	1 25	50		
- σαρ 200 mg	(10.82)	00		Ponstan
	0.50	20		1 Olistali
	(7.50)	20		Ponstan
WARRANGE TO THE STATE OF THE ST	(7.50)			runstan
NAPROXEN				
* Tab 250 mg		500		Noflam 250
* Tab 500 mg		250		Noflam 500
* Tab long-acting 750 mg		28		Naprosyn SR 750
* Tab long-acting 1 g	8.62	28	•	Naprosyn SR 1000
TENOXICAM				
* Tab 20 mg	18.50	100	✓	Tilcotil
* Inj 20 mg vial	9.95	1	✓	AFT
NOAID OIL				
NSAIDs Other				
CELECOXIB				
Cap 100 mg	3.45	60		Celebrex
				Celecoxib Pfizer
Cap 200 mg	3.20	30		Celebrex
			1	Celecoxib Pfizer

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsid	dised	Generic	
\$	Per	✓	Manufacturer	

Topical Products for Joint and Muscular Pain

CAPSAICIN

⇒SA1289 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has osteoarthritis that is not responsive to paracetamol and oral non-steroidal anti-inflammatories are contraindicated.

Antirheumatoid Agents

HYDROXYCHLOROQUINE - Subsidy by endorsement

Subsidised only if prescribed for rheumatoid arthritis, systemic or discoid lupus erythematosus, malaria treatment or suppression, relevant dermatological conditions (cutaneous forms of lupus and lichen planus, cutaneous vasculitides and mucosal ulceration)*, sarcoidosis (pulmonary and non-pulmonary)*, and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of hydroxychloroquine. Note: Indication marked with a * is an unapproved indication.

* Tab 200 mg	8.78	100	✓ Plaquenil
LEFLUNOMIDE			
* Tab 10 mg	6.00	30	✓ Arava
* Tab 20 mg		30	✓ Arava
PENICILLAMINE			
Tab 125 mg	67.23	100	D-Penamine
Tab 250 mg	110.12	100	✓ D-Penamine

Drugs Affecting Bone Metabolism

Alendronate for Osteoporosis

ALENDRONATE SODIUM * Tab 70 mg	0 .	4	✓ Fosamax
Fosamax to be Principal Supply on 1 July 2024			
ALENDRONATE SODIUM WITH COLECALCIFEROL			
* Tab 70 mg with colecalciferol 5,600 iu	9 .	4	✓ Fosamax Plus
Fosamax Plus to be Principal Supply on 1 July 2024			

Other Treatments

DENOSUMAB - Special Authority see SA1777 below - Retail pl	harmacy		
Inj 60 mg prefilled syringe	326.00	1	✓ Prolia

⇒SA1777 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

1 The patient has severe, established osteoporosis; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 2 Either:
 - 2.1 The patient is female and postmenopausal; or
 - 2.2 The patient is male or non-binary; and
 - 3 Any of the following:
 - 3.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note): or
 - 3.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons; or
 - 3.3 History of two significant osteoporotic fractures demonstrated radiologically; or
 - 3.4 Documented T-Score less than or equal to -3.0 (see Note); or
 - 3.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
 - 3.6 Patient has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
 - 4 Zoledronic acid is contraindicated because the patient's creatinine clearance is less than 35 mL/min; and
 - 5 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes); and
 - 6 The patient must not receive concomitant treatment with any other funded antiresorptive agent for this condition or teriparatide.

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
 Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with denosumab
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body
- e) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: risedronate sodium tab 35 mg once weekly; alendronate sodium tab 70 mg or tab 70 mg with cholecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy

PAMIDRONATE DISODIUM

Inj 3 mg per ml, 10 ml vial	32.49	1	Pamisol
Inj 6 mg per ml, 10 ml vial	88.11	1	✓ Pamisol
Inj 9 mg per ml, 10 ml vial		1	Pamisol
RALOXIFENE HYDROCHLORIDE - Special Authority see SA1779	on the next pa	ge – Retail	pharmacy
* Tab 60 mg	53.76	28	Evista

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA1779 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Notes); or
- 2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
- 3 History of two significant osteoporotic fractures demonstrated radiologically; or
- 4 Documented T-Score less than or equal to -3.0 (see Notes); or
- 5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Notes); or
- 6 Patient has had a Special Authority approval for zoledronic acid (Underlying cause Osteoporosis) or has had a Special Authority approval for alendronate (Underlying cause - Osteoporosis) prior to 1 February 2019.

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
 Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for raloxifene funding.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis, and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

RISEDRONATE SODIUM

⇒SA1139 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 18 months for applications meeting the following criteria: All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 The patient has a documented T-score less than or equal to -3.0 (see Notes); and
- 3 The patient has had two or more fractures due to minimal trauma: and
- 4 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes).

Notes:

a) The bone mineral density (BMD) measurement used to derive the T-score must be made using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable

Subsidy		Fully	Brand or	
(Manufacturer's Price	e)	Subsidised	Generic	
\$	Per	•	Manufacturer	

continued...

- b) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: alendronate sodium tab 70 mg or tab 70 mg with colecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily; zoledronic acid 5 mg per year. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy.
- c) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.
- d) A maximum of 18 months of treatment (18 cartridges) will be subsidised.

ZOLEDRONIC ACID

Inj 0.05 mg per ml, 100 ml, bag	22.53 100 ml	I OP ✓ Zoledronic Acid
		Viatris

Hyperuricaemia and Antigout

ALLOPURINOL			
* Tab 100 mg	17.99	1,000	✓ Ipca-Allopurinol
* Tab 300 mg		500	✓ Ipca-Allopurinol
BENZBROMARONE - Special Authority see SA196	3 below – Retail pharmacy		
Tab 50 mg	32.00	100	✓ Narcaricin mite \$29
Tab 100 mg	13.50	30	✓ Desuric S29
			✓ Urinorm S29
	45.00	100	 Benzbromaron AL
			100 S29

(Desuric S29 Tab 100 mg to be delisted 1 August 2024) (Urinorm S29 Tab 100 mg to be delisted 1 August 2024)

(Benzbromaron AL 100 S29 Tab 100 mg to be delisted 1 August 2024)

⇒SA1963 Special Authority for Subsidy

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefitting from the treatment; and
- 2 There is no evidence of liver toxicity and patient is continuing to receive regular (at least every three months) liver function tests.

COLCHICINE

* Tab 500 mcg	.6.00	100	✓ Colgout
FEBUXOSTAT - Special Authority see SA2054 below - Retail pharmac	у		
Tab 80 mg	.4.73	28	✓ Febuxostat (Teva)
Tab 120 mg	11.78	28	✓ Febuxostat (Teva)

⇒SA2054 Special Authority for Subsidy

Initial application — (**Gout**) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with gout; and
- 2 Any of the following:
 - 2.1 The patient has a serum urate level greater than 0.36 mmol/l despite treatment with allopurinol at doses of at least

✓ Medsurge

✓ Dantrium ✓ Dantrium S29 S29 ✓ Dantrium

✓ Norflex

100

100

100

	MU	SCULOSKEL	ETAL SYSTEM
	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
continued			
600 mg/day and addition of probenecid at doses of 2.2 The patient has experienced intolerable side effect and serum urate remains greater than 0.36 mmol/l maximum tolerated dose; or	s from allopurinol sucl	n that treatment of	discontinuation is required
2.3 The patient has renal impairment such that proben remains greater than 0.36 mmol/l despite optimal tr2.4 The patient has previously had an initial Special Au	eatment with allopurir	nol (see Note); or	•
Initial application — (Tumour lysis syndrome) only from a had applications meeting the following criteria: Both:			-
Patient is scheduled to receive cancer therapy carrying an Patient has a documented history of allopurinol intolerance	•	risk of tumour lys	is syndrome; and
Renewal — (Gout) from any relevant practitioner. Approvals va patient is benefitting from treatment.	id for 2 years where t	he treatment rem	nains appropriate and the
Renewal — (Tumour lysis syndrome) only from a haematologi treatment remains appropriate and the patient is benefitting from		rovals valid for 6	weeks where the
PROBENECID * Tab 500 mg	66.95	100 ~ P	robenecid-AFT
Muscle Relaxants			
BACLOFEN			
* Tab 10 mg	11.55	1 / L	acifen ioresal Intrathecal e been ineffective or have

Subsidised only for use in a programmable pump in patients where oral antispastic agents have been ineffective or have

caused intolerable side effects and the prescription is endorsed accordingly. Inj 2 mg per ml, 5 ml ampoule - Subsidy by endorsement...........306.82

caused intolerable side effects and the prescription is endorsed accordingly.

DANTROLENE

ORPHENADRINE CITRATE

Subsidy (Manufacturer's Price)

Fully Subsidised Per

Brand or Generic Manufacturer

Agents for Parkinsonism and Related Disorders

AMANTADINE HYDROCHLORIDE			
▲ Cap 100 mg	38.24	60	✓ Symmetrel
	63.73	100	✓ Symmetrel
APOMORPHINE HYDROCHLORIDE			•
▲ Inj 10 mg per ml, 2 ml ampoule	59.50	5	✓ Movapo
▲ Inj 10 mg per ml, 5 ml ampoule		5	✓ Movapo
ENTACAPONE			·
▲ Tab 200 mg	18 04	100	✓ Comtan
LEVODOPA WITH BENSERAZIDE		100	<u>oomtan</u>
* Tab dispersible 50 mg with benserazide 12.5 mg	12.25	100	✓ Madopar Rapid
* Cap 50 mg with benserazide 12.5 mg		100	✓ Madopar 62.5
* Cap 100 mg with benserazide 25 mg		100	✓ Madopar 125
* Cap long-acting 100 mg with benserazide 25 mg		100	✓ Madopar HBS
* Cap 200 mg with benserazide 50 mg		100	✓ Madopar 250
LEVODOPA WITH CARBIDOPA	20.20	100	- madopai 200
	04.44	100	✓ Sinemet
The same same same same same same same sam		100 100	✓ Sinemet CR
* Tab long-acting 200 mg with carbidopa 50 mg Tab 250 mg with carbidopa 25 mg		100	✓ Sinemet
		100	• Silielliet
PRAMIPEXOLE HYDROCHLORIDE			
▲ Tab 0.25 mg		100	✓ Ramipex
▲ Tab 1 mg	18.66	100	✓ Ramipex
RASAGILINE			
* Tab 1 mg	53.50	30	✓ Azilect S29
ROPINIROLE HYDROCHLORIDE			
▲ Tab 0.25 mg	4.05	84	✓ Ropin
▲ Tab 1 mg	4.95	84	✓ Ropin
▲ Tab 2 mg	6.48	84	✓ Ropin
▲ Tab 5 mg		84	✓ Ropin
TOLCAPONE			
▲ Tab 100 mg	152.38	100	✓ Tasmar
,			

Anticholinergics

BENZATROPINE MESYLATE			
Tab 2 mg	9.59	60	✓ Benztrop
Inj 1 mg per ml, 2 ml		5	✓ Phebra
a) Up to 10 inj available on a PSOb) Only on a PSO			
PROCYCLIDINE HYDROCHLORIDE			
Tah 5 mg	7 40	100	✓ Kemadrin

Agents for Essential Tremor, Chorea and Related Disorders

NERVOUS SYSTEM

Subsidy (Manufacturer's I	Price)	Fully Subsidised	Brand or Generic	
\$	Pei		Manufacturer	

⇒SA1403 Special Authority for Subsidy

Initial application only from a neurologist or respiratory specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has amyotrophic lateral sclerosis with disease duration of 5 years or less; and
- 2 The patient has at least 60 percent of predicted forced vital capacity within 2 months prior to the initial application; and
- 3 The patient has not undergone a tracheostomy; and
- 4 The patient has not experienced respiratory failure; and
- 5 Any of the following:
 - 5.1 The patient is ambulatory; or
 - 5.2 The patient is able to use upper limbs; or
 - 5.3 The patient is able to swallow.

Renewal from any relevant practitioner. Approvals valid for 18 months for applications meeting the following criteria: All of the following:

- 1 The patient has not undergone a tracheostomy; and
- 2 The patient has not experienced respiratory failure; and
- 3 Any of the following:
 - 3.1 The patient is ambulatory; or
 - 3.2 The patient is able to use upper limbs; or
 - 3.3 The patient is able to swallow.

TETRABENAZINE

Tab 25 mg	106.59	112	Motetis

Anaesthetics

LIDOCAINE (LIGNOCAINE)

Local

LIDOCAINE [LIGNOCAINE]			
Gel 2%, tube - Subsidy by endorsement	14.50	30 ml	✓ Xylocaine 2% Jelly
a) Up to 150 ml available on a PSO			
b) Subsidised only if prescribed for urethral or cervical admi	nistration and th	e prescription	n is endorsed accordingly.
Gel 2%, 11 ml urethral syringe - Subsidy by endorsement	59.50	10	✓ Instillagel Lido
a) Up to 5 each available on a PSO			
b) Subsidised only if prescribed for urethral, cervical or recta	al administration	and the pres	scription is endorsed
accordingly.		'	•
LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE			
Oral (gel) soln 2%	44.00	200 ml	✓ Mucosoothe
Inj 1%, 5 ml ampoule – Up to 25 inj available on a PSO		25	✓ Lidocaine-Baxter
,,	17.50	50	
	(35.00)		Xylocaine
Inj 2%, 5 ml ampoule - Up to 5 inj available on a PSO	9.00	25	✓ Lidocaine-Baxter
Inj 1%, 20 ml ampoule - Up to 5 inj available on a PSO	12.00	5	
	(20.00)		Xylocaine
Inj 1%, 20 ml vial - Up to 5 inj available on a PSO	6.85	5	✓ Lidocaine-Baxter
Inj 2%, 20 ml vial - Up to 5 inj available on a PSO	7.15	5	✓ Lidocaine-Baxter

Subsidised only for people receiving palliative care services where other analgesic agents haven't been effective.

10

✓ Xvlocard 500 S29

Inj 10%, 5 ml ampoule - Subsidy by endorsementCBS



Subsidy	Fully	Brand or	_
(Manufacturer's Price)	Subsidised	Generic	
\$	Per ✓	Manufacturer	

Topical Local Anaesthetics

⇒SA0906 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years where the patient is a child with a chronic medical condition requiring frequent injections or venepuncture.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

LIDOCAINE [LIGNOCAINE] - Special Authority see \$A0906 above -	- Retail phan	macy	
Crm 4%	5.40	5 g OP	✓ LMX4
	27.00	30 g OP	✓ LMX4
LIDOCAINE [LIGNOCAINE] WITH PRILOCAINE - Special Authority	see SA0906	above – Reta	il pharmacy
Crm 2.5% with prilocaine 2.5%	45.00	30 g OP	✓ EMLA
Crm 2.5% with prilocaine 2.5% (5 g tubes)	45.00	5	✓ EMLA

Analgesics

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 116

Non-opioid Analgesics			
ASPIRIN			
* Tab dispersible 300 mg - Up to 30 tab available on	a PSO5.65	100	 Ethics Aspirin
CAPSAICIN - Subsidy by endorsement			
Subsidised only if prescribed for post-herpetic neuro accordingly.	algia or diabetic periphera	l neuropathy a	nd the prescription is endorsed
Crm 0.075%	11.95	45 g OP	✓ Zostrix HP
	15.14	57 g OP	Rugby Capsaicin
			Topical
			Cream \$29
NEFOPAM HYDROCHLORIDE			
Tah 30 mg	23 40	90	✓ Acupan

NERVOUS SYSTEM

	Subsidy (Manufacturer's Price)	Sub	Fully sidised	Brand or Generic
	\$	Per	✓	Manufacturer
PARACETAMOL Tab 500 mg - blister pack	19.75	1.000	✓ P	acimol

- a) Maximum of 300 tab per prescription; can be waived by endorsement
- b) Up to 30 tab available on a PSO

c)

- Subsidy by endorsement for higher quantities is available for patients with long term conditions who require
 regular daily dosing for one month or greater, and the prescription is annotated accordingly. Pharmacists may
 annotate the prescription as endorsed where dispensing history supports a long-term condition.
- 2) Maximum of 100 tab per dispensing for non-endorsed patients. If quantities prescribed for more than 100 tabs (for non-endorsed patients), then dispense in repeat dispensings not exceeding 100 tab per dispensing.

- Subsidy by endorsement for higher quantities is available for patients with long term conditions who require regular daily dosing for one month or greater, and the prescription is annotated accordingly. Pharmacists may annotate the prescription as endorsed where dispensing history supports a long-term condition.
- Maximum of 100 tab per dispensing for non-endorsed patients. If quantities prescribed for more than 100 tabs (for non-endorsed patients), then dispense in repeat dispensings not exceeding 100 tab per dispensing.

 Oral liq 120 mg per 5 ml
 3.98
 200 ml
 ✓ Paracetamol (Ethics)

 10.50
 200 ml OP
 ✓ Avallon

- a) Maximum of 600 ml per prescription; can be waived by endorsement
- b) Up to 200 ml available on a PSO
- c) Not in combination

d)

- Maximum of 200 ml per dispensing for non-endorsed patients. If quantities prescribed exceed 200 ml (for non-endorsed patients), then dispense in repeat dispensing not exceeding 200 ml per dispensing.
- 2) Subsidy by endorsement for higher quantities is available for patients with long term conditions who require regular daily dosing for one month or greater and the prescription is endorsed or annotated accordingly. Pharmacists may annotate the prescription as endorsed where dispensing history supports a long-term condition.
- Note: 200 ml presentations of paracetamol oral liquid may be supplied on BSO to a Vaccinator (other than a Pharmacist) under the provisions in Part I of Section A
- 4) Note: Direct Provision by a pharmacist of up to 200 ml permitted under the provisions in Part I of Section A in conjunction with immunisation of a child under 2 years of age with meningococcal B multicomponent vaccine.

- a) Maximum of 600 ml per prescription; can be waived by endorsement
- b) Up to 200 ml available on a PSO
- c) Not in combination

d)

- Maximum of 200 ml per dispensing for non-endorsed patients. If quantities prescribed exceed 200 ml (for non-endorsed patients), then dispense in repeat dispensing not exceeding 200 ml per dispensing.
- 2) Subsidy by endorsement for higher quantities is available for patients with long term conditions who require regular daily dosing for one month or greater and the prescription is endorsed or annotated accordingly. Pharmacists may annotate the prescription as endorsed where dispensing history supports a long-term condition.
- Note: 200 ml presentations of paracetamol oral liquid may be supplied on BSO to a Vaccinator (other than a Pharmacist) under the provisions in Part I of Section A
- 4) Note: Direct Provision by a pharmacist of up to 200 ml permitted under the provisions in Part I of Section A in conjunction with immunisation of a child under 2 years of age with meningococcal B multicomponent vaccine.

		Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
: (Suppos 125 mg	4.29	10	✓	Gacet
. (Suppos 250 mg	5.39	10		Gacet
. (Suppos 500 mg	16.55	50	•	Gacet
)p	ioid Analgesics				
DC	EINE PHOSPHATE - Safety medicine; prescriber may det	ermine dispensing fre	equen	су	
-	Гаb 15 mg	5.92	100	✓	Noumed
-	Гаb 30 mg	6.98	100	✓	Aspen
				✓	Noumed
-	Гаb 60 mg	13.89	100	✓	Noumed
	DROCODEINE TARTRATE				
	Fab long-acting 60 mg	8 60	60	/	DHC Continus
			00	•	DITO CONTINUO
	TANYL				
	a) Only on a controlled drug form				
	No patient co-payment payable				
	c) Safety medicine; prescriber may determine dispensing fr				
	nj 50 mcg per ml, 2 ml ampoule		10		Boucher and Muir
I	nj 50 mcg per ml, 10 ml ampoule	9.41	10		Boucher and Muir
F	Patch 12.5 mcg per hour	6.99	5		Fentanyl Sandoz
F	Patch 25 mcg per hour	7.99	5	✓	Fentanyl Sandoz
F	Patch 50 mcg per hour	9.49	5	✓	Fentanyl Sandoz
			_		Fentanyl Sandoz
F	Patch 75 mcg per hour	17.99	5	•	remanyi Sandoz
	Patch 75 mcg per hour Patch 100 mcg per hour		5 5		Fentanyl Sandoz
F	Patch 100 mcg per hour				
T	Patch 100 mcg per hour HADONE HYDROCHLORIDE				
Ι Τ:	Patch 100 mcg per hour HADONE HYDROCHLORIDE a) Only on a controlled drug form				
T T	Patch 100 mcg per hour	18.59			
T T d d	Patch 100 mcg per hour	18.59 equency	5	•	Fentanyl Sandoz
T T d d	Patch 100 mcg per hour	18.59 equency	5	•	Fentanyl Sandoz
T T d d	Patch 100 mcg per hour	18.59 equency reimbursed at the rat	5	•	Fentanyl Sandoz
T:T	Patch 100 mcg per hour	equency reimbursed at the rat	5 e of th	ne cheape	Fentanyl Sandoz
T a b	Patch 100 mcg per hour	equency reimbursed at the rat	5 e of th	ne cheape	Fentanyl Sandoz st form available Methadone BNM
F	Patch 100 mcg per hour	equency reimbursed at the rat formulae, page 2671.456.40	5 e of th 10 200 m	ne cheape	st form available Methadone BNM Biodone
T : T : C : C : C : C : C : C : C : C :	Patch 100 mcg per hour	equency reimbursed at the rate formulae, page 267	5 e of th 10 200 m 200 m	ne cheape	st form available Methadone BNM Biodone Biodone Forte
T	Patch 100 mcg per hour	equency reimbursed at the rat Formulae, page 2671.456.406.407.50	5 e of th 10 200 m 200 m 200 m	ne cheape	st form available Methadone BNM Biodone Biodone Forte Biodone Extra Fort
T : T : C : C : C : C : C : C : C : C :	Patch 100 mcg per hour	equency reimbursed at the rat Formulae, page 2671.456.406.407.50	5 e of th 10 200 m 200 m	ne cheape	st form available Methadone BNM Biodone Biodone Forte
ET : : : : : : : : : : : : : : : : : : :	Patch 100 mcg per hour	equency reimbursed at the rat Formulae, page 2671.456.406.407.50	5 e of th 10 200 m 200 m 200 m	ne cheape	st form available Methadone BNM Biodone Biodone Forte Biodone Extra Fort
T a a a a a a a a a a a a a a a a a a a	Patch 100 mcg per hour	equency reimbursed at the rat Formulae, page 2671.456.406.407.50	5 e of th 10 200 m 200 m 200 m	ne cheape	st form available Methadone BNM Biodone Biodone Forte Biodone Extra Fort
ET a la l	Patch 100 mcg per hour	equency reimbursed at the rate formulae, page 267	5 e of th 10 200 m 200 m 200 m	ne cheape	st form available Methadone BNM Biodone Biodone Forte Biodone Extra Fort
ET ä	Patch 100 mcg per hour	equency reimbursed at the rate formulae, page 267	5 e of th 10 200 m 200 m 200 m	ne cheape	st form available Methadone BNM Biodone Biodone Forte Biodone Extra Fort
ET in it is	Patch 100 mcg per hour	equency reimbursed at the rate formulae, page 267	5 e of th 10 200 m 200 m 200 m	ne cheape	st form available Methadone BNM Biodone Biodone Forte Biodone Extra Fort
H H H H H H H	Patch 100 mcg per hour	equency reimbursed at the rate formulae, page 267	5 e of th 10 200 m 200 m 200 m 10	ne cheape	st form available Methadone BNM Biodone Biodone Forte Biodone Extra Fort
	Patch 100 mcg per hour	equency reimbursed at the rate formulae, page 267	5 e of th 10 200 m 200 m 10	ne cheape	st form available Methadone BNM Biodone Biodone Forte Biodone Extra Fort AFT RA-Morph
	Patch 100 mcg per hour	equency reimbursed at the rate formulae, page 267	5 e of th 10 200 m 200 m 10 200 m 200 m	ne cheape	st form available Methadone BNM Biodone Biodone Forte Biodone Extra Fort AFT RA-Morph RA-Morph
1	Patch 100 mcg per hour	equency reimbursed at the rate formulae, page 2671.456.407.5068.90	5 e of th 10 200 m 200 m 10 200 m 200 m 200 m 200 m	ne cheape	Fentanyl Sandoz st form available Methadone BNM Biodone Biodone Forte Biodone Extra Fort AFT RA-Morph RA-Morph Ordine \$290 RA-Morph
	Patch 100 mcg per hour	equency reimbursed at the rate formulae, page 2671.456.407.5068.90	5 e of th 10 200 m 200 m 10 200 m 200 m	ne cheape	st form available Methadone BNM Biodone Biodone Forte Biodone Extra Fort AFT RA-Morph RA-Morph Ordine \$229

	Subsidy		Fully	Brand or
	(Manufacturer's Price \$) Per	Subsidised <	Generic Manufacturer
MORPHINE SULPHATE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing fre	equency			
Tab immediate-release 10 mg		10		Sevredol
Tab immediate-release 20 mg		10		Sevredol
Cap long-acting 10 mg		10	_	m-Eslon
Cap long-acting 30 mg		10		m-Eslon
Cap long-acting 60 mg		10		m-Eslon
Cap long-acting 100 mg		10		m-Eslon
Oral liq 2 mg per ml		100 m		Wockhardt S29
	29.80			Oramorph
			•	Oramorph CDC
				S29 S29
Inj 5 mg per ml, 1 ml ampoule - Up to 5 inj available on a PS		5	✓	<u>Medsurge</u>
Inj 10 mg per ml, 1 ml ampoule - Up to 5 inj available on a F		5		<u>Medsurge</u>
Inj 15 mg per ml, 1 ml ampoule - Up to 5 inj available on a F		5		<u>Medsurge</u>
Inj 30 mg per ml, 1 ml ampoule – Up to 5 inj available on a F	PSO6.28	5	/	Medsurge
XYCODONE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing fre	equency			
Tab controlled-release 5 mg	2.69	20	1	Oxycodone Sandoz
•	3.77	28	✓	Oxycodone Sandoz
				S29 S29
	4.04	30	1	OxyContin S29
Tab controlled-release 10 mg		20		Oxycodone Sandoz
Tab controlled follower to mg	3.77	28		Oxycodone Sandoz
	U			S29 S29
Tab controlled-release 20 mg	2.40	20	1	Oxycodone Sandoz
Tab controlled-release 20 mg		20		Oxycodone Sandoz
Tab controlled-release 40 mg		20		Oxycodone Sandoz
Cap immediate-release 5 mg		20		OxyNorm
Cap immediate-release 10 mg		20		OxyNorm
Cap immediate-release 10 mg		20		OxyNorm OxyNorm
Oral liq 5 mg per 5 ml		250 m		OxyNorm OxyNorm
Oral liq 1 mg per ml		250 m		Oxycodone Lucis
		_00 111		S29 S29
Inj 10 mg per ml, 1 ml ampoule	5.82	5	1	Hameln
Inj 10 mg per ml, 2 ml ampoule		5		Hameln
Inj 50 mg per ml, 1 ml ampoule		5		Hameln
PARACETAMOL WITH CODEINE – Safety medicine; prescriber		-		
★ Tab paracetamol 500 mg with codeine phosphate 8 mg		ensing 1,000		/ Paracetamol +
r ab paracetamor 500 mg with codelle phosphate 8 mg	27.30	1,000	•	Codoino (Policyo

Codeine (Relieve)

	Subsidy		Fully Brand or
	(Manufacturer's Price) \$	Per	Subsidised Generic Manufacturer
DETUIDING LIVEROCUL ORIDE	Ψ	1 01	- Manadadio
PETHIDINE HYDROCHLORIDE			
a) Only on a controlled drug form b) No patient co-payment payable			
c) Safety medicine; prescriber may determine dispensing	froquency		
Tab 50 mg		10	✓ Noumed Pethidine
Inj 50 mg per ml, 1 ml ampoule – Up to 5 inj available on		5	✓ DBL Pethidine
ing oo mg por mi, i mi ampoulo — op to o mg available on	a 1 0020.00	Ŭ	Hydrochloride
Inj 50 mg per ml, 2 ml ampoule - Up to 5 inj available on	a PSO30.72	5	✓ DBL Pethidine
.,		-	Hydrochloride
FRAMADOL HYDROCHLORIDE			•
Tab sustained-release 100 mg	1.95	20	✓ Tramal SR 100
Tab sustained-release 150 mg		20	✓ Tramal SR 150
Tab sustained-release 200 mg		20	✓ Tramal SR 200
Cap 50 mg		100	✓ Arrow-Tramadol
, ,			
Antidepressants			
Cyclic and Related Agents			
AMITRIPTYLINE - Safety medicine; prescriber may determine	e dispensing frequency		
Tab 10 mg		100	✓ Arrow-Amitriptyline
Tab 25 mg	1.99	100	✓ Arrow-Amitriptyline
Tab 50 mg	3.14	100	Arrow-Amitriptyline
CLOMIPRAMINE HYDROCHLORIDE - Safety medicine; pres	scriber may determine d	lispen	ising frequency
Tab 10 mg		30	✓ Clomipramine Teva
Tab 25 mg	11.99	30	✓ Clomipramine Teva
	39.97	100	✓ Anafranil S29
Cap 10 mg	9.49	28	Clomipramine Teva
Cap 25 mg	11.19	28	Clomipramine Teva
OSULEPIN [DOTHIEPIN] HYDROCHLORIDE - Subsidy by	endorsement		
 a) Safety medicine; prescriber may determine dispensing b) Subsidy by endorsement – Subsidised for patients who 2019 and the prescription is endorsed accordingly. Ph exists a record of prior dispensing of dosulepin [dothie] 	o were taking dosulepin armacists may annotate pin] hydrochloride.		prescription as endorsed where
Tab 75 mg	3.85	30	Dosulepin Viatris
Cap 25 mg	7.83	50	Dosulepin
			Mylan S29
			Dosulepin
			Viatris S29
Dosulepin Mylan 🖘 Cap 25 mg to be delisted 1 October 20	024)		
	•	ensing	g frequency
	er may determine dispe		
MIPRAMINE HYDROCHLORIDE - Safety medicine; prescrib	er may determine dispe		
MIPRAMINE HYDROCHLORIDE - Safety medicine; prescrib	er may determine dispe 5.48 10.96	50	✓ Tofranil
MIPRAMINE HYDROCHLORIDE – Safety medicine; prescrib Tab 10 mg	er may determine dispe 5.48 10.96	50 100	✓ Tofranil✓ Tofranil
MIPRAMINE HYDROCHLORIDE – Safety medicine; prescrib Tab 10 mg	er may determine dispe 5.48 10.96	50 100	✓ Tofranil ✓ Tofranil ✓ Imipramine
MIPRAMINE HYDROCHLORIDE – Safety medicine; prescrib Tab 10 mg Tab 25 mg	er may determine dispe 5.48 10.96 4.93	50 100 28 50	✓ Tofranil ✓ Tofranil ✓ Imipramine Crescent \$29 ✓ Tofranil
ŭ	er may determine dispe 5.48 10.96 4.93 8.80 scriber may determine of	50 100 28 50	✓ Tofranil ✓ Tofranil ✓ Imipramine Crescent \$29 ✓ Tofranil

			INE	HVUUS SYSTEM
	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
Monoamine-Oxidase Inhibitors (MAOIs) - Non S	Selective			
TRANYLCYPROMINE SULPHATE * Tab 10 mg	22.94	50	1	Parnate
Monoamine-Oxidase Type A Inhibitors				
MOCLOBEMIDE * Tab 150 mg * Tab 300 mg		60 60		Aurorix Aurorix
Selective Serotonin Reuptake Inhibitors				
CITALOPRAM HYDROBROMIDE * Tab 20 mg ESCITALOPRAM		84	✓	<u>Celapram</u>
* Tab 10 mg	0.79 1.07	28		Ipca-Escitalopram Escitalopram (Ethics)
* Tab 20 mg FLUOXETINE HYDROCHLORIDE	1.49	28	✓	Ipca-Escitalopram
* Tab dispersible 20 mg, scored – Subsidy by endorsement Subsidised by endorsement	2.50	28	•	Fluox
 When prescribed for a patient who cannot swallow accordingly; or When prescribed in a daily dose that is not a multi endorsed. Note: Tablets should be combined wit 	ple of 20 mg in which	case t	the prescri	iption is deemed to be
Cap 20 mg		30		Brown & Burk S29
PAROXETINE	3.13	90	•	Arrow-Fluoxetine
* Tab 20 mg	4.11	90	✓	<u>Loxamine</u>
SERTRALINE * Tab 50 mg * Tab 100 mg		30 30		Setrona Setrona
Other Antidepressants				
MIRTAZAPINE	0.00	00		Nad
Tab 30 mg		28 30	1	Noumed Noumed
Tab 45 mg	3.45	28 30		Noumed Noumed
VENLAFAXINE	0.00	0.4		Full-four VD
* Cap 75 mg		84 84		Enlafax XR Enlafax XR
* Cap 150 mg		84		Enlafax XR

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per

✓ Manufacturer

Antiepilepsy Drugs

Agents for Control of Status Epilepticus

DIAZEPAM – Safety medicine; prescriber may determine dispensing frequency Inj 5 mg per ml, 2 ml ampoule – Subsidy by endorsement27.92	5	✓ Hospira
a) Up to 5 inj available on a PSO		
b) Only on a PSO		
 c) PSO must be endorsed "not for anaesthetic procedures". 		
Rectal tubes 5 mg - Up to 5 tube available on a PSO54.58	5	✓ <u>Stesolid</u>
PHENYTOIN SODIUM		
* Inj 50 mg per ml, 2 ml ampoule - Up to 5 inj available on a		
PSO104.58	5	Hospira
* Inj 50 mg per ml, 5 ml ampoule - Up to 5 inj available on a		
PSO154.01	5	Hospira

Control of Epilepsy

CARBAMAZEPINE

* Tab 200 mg	14.53	100	✓ Tegretol
* Tab long-acting 200 mg	16.98	100	✓ Tegretol CR
	33.96	200	✓ Tegretol CR
* Tab 400 mg	34.58	100	✓ Tegretol
* Tab long-acting 400 mg	39.17	100	✓ Tegretol CR
* Oral liq 20 mg per ml	26.37	250 ml	✓ Tegretol
CLOBAZAM - Safety medicine; prescriber may determine dis	spensing frequency		
Tab 10 mg	9.12	50	✓ Frisium
CLONAZEPAM - Safety medicine; prescriber may determine	dispensing frequen	cy	
Oral drops 2.5 mg per ml		10 ml OP	✓ Rivotril
ETHOSUXIMIDE			
Cap 250 mg	78.89	56	✓ Essential
•			Ethosuximide S29
	140.88	100	✓ Zarontin
Oral liq 250 mg per 5 ml	56.35	200 ml	✓ Zarontin
GABAPENTIN			
Note: Not subsidised in combination with subsidised preg	nabalin		
* Cap 100 mg	,	100	✓ Nupentin
* Cap 300 mg		100	✓ Nupentin
* Cap 400 mg	10.26	100	✓ Nupentin
LACOSAMIDE - Special Authority see SA2267 on the next p	age – Retail pharma	acv	
▲ Tab 50 mg	•	14	✓ Vimpat
▲ Tab 100 mg	50.06	14	✓ Vimpat
-	200.24	56	✓ Vimpat
▲ Tab 150 mg	75.10	14	✓ Vimpat
	300.40	56	✓ Vimpat
▲ Tab 200 mg	400.55	56	✓ Vimpat

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sul	osidised	Generic	
\$	Per	1	Manufacturer	

⇒SA2267 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Patient has focal epilepsy; and
- 2 Seizures are not adequately controlled by, or patient has experienced unacceptable side effects from, optimal treatment with all of the following: sodium valproate, topiramate, levetiracetam and any two of carbamazepine, lamotrigine and phenytoin sodium (see Note).

Note: Those of childbearing potential are not required to trial phenytoin sodium, sodium valproate, or topiramate. Those who can father children are not required to trial sodium valproate.

Renewal from any relevant practitioner. Approvals valid for 24 months where the patient has demonstrated a significant and sustained improvement in seizure rate or severity and/or quality of life compared with that prior to starting lacosamide treatment.

LAI	MOTRIGINE			
lack	Tab dispersible 2 mg	55.00	30	✓ Lamictal
lack	Tab dispersible 5 mg	50.00	30	✓ Lamictal
*	Tab dispersible 25 mg	4.20	56	✓ Logem
*	Tab dispersible 50 mg		56	✓ Logem
*	Tab dispersible 100 mg	6.75	56	✓ Logem
LE\	/ETIRACETAM			
	Tab 250 mg	5.84	60	✓ Everet
	Tab 500 mg	10.51	60	✓ Everet
	Tab 750 mg	16.71	60	✓ Everet
	Tab 1,000 mg	21.82	60	✓ Everet
	Oral liq 100 mg per ml	44.78	300 ml OP	✓ Levetiracetam-AFT
РΗ	ENOBARBITONE			
	For phenobarbitone oral liquid refer Standard Formulae, page 267	7		
	Tab 15 mg		500	✓ PSM
	•	248.50		✓ Noumed
				Phenobarbitone
	Tab 30 mg	.398.50	500	✓ Noumed
	•			Phenobarbitone
(PS	SM Tab 15 mg to be delisted 1 August 2024)			
PH	ENYTOIN SODIUM			
	Tab 50 mg	75.00	200	✓ Dilantin Infatab
•	Cap 30 mg		200	✓ Dilantin
	Cap 100 mg		200	✓ Dilantin
*	Oral lig 30 mg per 5 ml		500 ml	✓ Dilantin Paediatric
	EGABALIN			
ГΠ	Note: Not subsidised in combination with subsidised gabapentin			
	Cap 25 mg	2 25	56	✓ Pregabalin Pfizer
		7.80		✓ Milpharm \$29
*	Cap 75 mg		56	✓ Pregabalin Pfizer
7,1	oup 70 mg	8.10	00	✓ Milpharm \$29
	Cap 150 mg		56	✓ Lyrica
	Cap 150 mg	4.01	50	✓ Pregabalin Pfizer
		10.44		✓ Milpharm \$29
	Con 200 mg	12.44	56	•
/8.4:	Cap 300 mg	1.30	00	 Pregabalin Pfizer
	pharm S29 Cap 150 mg to be delisted 1 August 2024)			
	MIDONE			4 - -
*	Tab 250 mg	37.35	100	✓ Primidone Clinect

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

NERVOUS SYSTEM

	Subsidy		Fully	Brand or
	(Manufacturer's Price	e)	Subsidised	I Generic
	\$	Per	•	Manufacturer
SODIUM VALPROATE				
Tab 100 mg	13.65	100	✓	Epilim Crushable
Tab 200 mg EC	27.44	100	✓	Epilim
Tab 500 mg EC	52.24	100	✓	Epilim
* Oral liq 200 mg per 5 ml	20.48	300 n	nl 🗸	Epilim S/F Liquid
			✓	Epilim Syrup
* Inj 100 mg per ml, 4 ml	41.50	1	✓	Epilim IV
STIRIPENTOL - Special Authority see SA2268 below - Retail ph	armacy			
Cap 250 mg	•	60	✓	Diacomit
Powder for oral liq 250 mg sachet	509.29	60	1	Diacomit

⇒SA2268 Special Authority for Subsidy

Initial application only from a paediatric neurologist or Practitioner on the recommendation of a paediatric neurologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient has confirmed diagnosis of Dravet syndrome; and
- 2 Seizures have been inadequately controlled by appropriate courses of sodium valproate, clobazam and at least two of the following: topiramate, levetiracetam, ketogenic diet.

Note: Those of childbearing potential are not required to trial sodium valproate or topiramate. Those who can father children are not required to trial sodium valproate.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where the patient continues to benefit from treatment as measured by reduced seizure frequency from baseline.

. •		
A	Tab 25 mg	

TOPIRAMATE

▲ Tab 25 mg	11.07	60	✓ Arrow-Topiramate
-			✓ Topiramate Actavis
	26.04		✓ Topamax
▲ Tab 50 mg	18.81	60	✓ Arrow-Topiramate
· ·			✓ Topiramate Actavis
	44.26		✓ Topamax
▲ Tab 100 mg	31.99	60	Arrow-Topiramate
· ·			✓ Topiramate Actavis
	75.25		✓ Topamax
▲ Tab 200 mg	55.19	60	Arrow-Topiramate
· ·			✓ Topiramate Actavis
	129.85		✓ Topamax
▲ Sprinkle cap 15 mg	20.84	60	✓ Topamax
▲ Sprinkle cap 25 mg		60	✓ Topamax
VIGABATRIN - Special Authority see SA2088 below - Retain	l pharmacy		
▲ Tab 500 mg		100	✓ Sabril
Powder for oral soln 500 mg per sachet		60	✓ Sabril

⇒SA2088 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Any of the following:
 - 1.1 Patient has infantile spasms; or
 - 1.2 Both:
 - 1.2.1 Patient has epilepsy; and
 - 1.2.2 Either:

		NI	ERVOUS SYSTEM
	Subsidy (Manufacturer's Price)	Ful Subsidise Per •	
continued			
1.2.2.1 Seizures are not adequately controlle 1.2.2.2 Seizures are controlled adequately b optimal treatment with other antiepile 1.3 Patient has tuberous sclerosis complex; and Either:	out the patient has ex		
2.1 Patient is, or will be, receiving regular automated v 6-monthly basis thereafter); or		•	•
2.2 It is impractical or impossible (due to comorbid cor Renewal from any relevant practitioner. Approvals valid without following criteria: Both:		•	
The patient has demonstrated a significant and sustained Either:	improvement in seiz	ure rate or sev	erity and or quality of life; and
2.1 Patient is receiving regular automated visual field to f treatment with vigabatrin; or2.2 It is impractical or impossible (due to comorbid cor		,	
Antimigraine Preparations			
For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 2	age 116		
Acute Migraine Treatment			
RIZATRIPTAN Tab orodispersible 10 mg	4.84	30	✓ <u>Rizamelt</u>
SUMATRIPTAN Tab 50 mg Tab 100 mg	22.68	90 90	✓ <u>Sumagran</u> ✓ <u>Sumagran</u>
Inj 12 mg per ml, 0.5 ml prefilled pen – Maximum of 10 inj po prescription		2 OP	<u>Clustran</u>
Prophylaxis of Migraine			
For Beta Adrenoceptor Blockers refer to CARDIOVASCULAR SY	/STEM, page 49		
PIZOTIFEN * Tab 500 mcg	23.21	100	✓ Sandomigran
Antinausea and Vertigo Agents			
For Antispasmodics refer to ALIMENTARY TRACT, page 8			
APREPITANT – Special Authority see SA0987 below – Retail ph Cap 2 × 80 mg and 1 × 125 mg		3 OP •	Emend Tri-Pack
■SA0987 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid	d for 12 months whe	re the patient is	s undergoing highly

Renewal from any relevant practitioner. Approvals valid for 12 months where the patient is undergoing highly emetogenic

emetogenic chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.

chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.

BETAHISTINE DIHYDROCHLORIDE

	Subsidy (Manufacturer's Price) \$	Per	Fully Brand or Subsidised Generic Manufacturer
CYCLIZINE HYDROCHLORIDE Tab 50 mg	0.49	10	✓ <u>Nausicalm</u>
CYCLIZINE LACTATE			
Inj 50 mg per ml, 1 ml ampoule – Up to 10 inj available on a PSO		10	✓ <u>Hameln</u>
DOMPERIDONE			
* Tab 10 mg	4.00	100	✓ <u>Domperidone</u> <u>Viatris</u>
HYOSCINE HYDROBROMIDE			
Inj 400 mcg per ml, 1 ml ampoule	93.00	10	✓ Martindale S29
below – Retail pharmacy	17.70	2	✓ Scopoderm TTS
,	88.50	10	✓ Scopolamine - Mylan
			✓ Scopolamine - Mylan S29 829

⇒SA1998 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Control of intractable nausea, vomiting, or inability to swallow saliva in the treatment of malignancy or chronic disease where the patient cannot tolerate or does not adequately respond to oral anti-nausea agents; or
- 2 Control of clozapine-induced hypersalivation where trials of at least two other alternative treatments have proven ineffective.

Renewal from any relevant practitioner. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

ME	TOCLOPRAMIDE HYDROCHLORIDE		
*	Tab 10 mg - Up to 30 tab available on a PSO1.57	100	✓ <u>Metoclopramide</u> <u>Actavis 10</u>
*	Inj 5 mg per ml, 2 ml ampoule - Up to 5 inj available on a PSO7.00	10	✓ Baxter
ON	DANSETRON		
*	Tab 4 mg2.27	50	✓ Periset
	Tab disp 4 mg - Up to 10 tab available on a PSO0.56	10	✓ Periset ODT
*	Tab 8 mg4.10	50	✓ Periset
	Tab disp 8 mg - Up to 10 tab available on a PSO0.90	10	✓ Periset ODT
PR	OCHLORPERAZINE		
*	Tab 3 mg buccal5.97	50	
	(30.00)		Buccastem
	(30.00)		Max Health \$29
	(30.00)		Prochlorperazine
			Brown & Burk S29
*	Tab 5 mg - Up to 30 tab available on a PSO25.00	250	✓ Nausafix
			✓ Nausafix - S29 S29
*	Inj 12.5 mg per ml, 1 ml - Up to 5 inj available on a PSO25.81	10	✓ Stemetil

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

Antipsychotics

General

AMISULPRIDE - Safety medicine; prescriber may determine disp		•	(0)
Tab 100 mg		30	✓ Sulprix
Tab 200 mg		60	✓ Sulprix
Tab 400 mg		60	✓ Sulprix
ARIPIPRAZOLE – Safety medicine; prescriber may determine dis		•	
Tab 5 mg	10.50	30	✓ Aripiprazole Sandoz✓ Ascend
			Aripiprazole S29
Tab 10 mg	10.50	30	✓ Aripiprazole Sandoz
Tab 15 mg	10.50	30	✓ Aripiprazole Sandoz
Tab 20 mg	10.50	30	✓ Aripiprazole Sandoz
Tab 30 mg	10.50	30	✓ Aripiprazole Sandoz
CHLORPROMAZINE HYDROCHLORIDE - Safety medicine; pre	scriber may dete	ermine dispen	sing frequency
Tab 25 mg - Up to 30 tab available on a PSO		100	✓ Largactil
Tab 100 mg - Up to 30 tab available on a PSO		100	✓ Largactil
Inj 25 mg per ml, 2 ml - Up to 5 inj available on a PSO		10	✓ Largactil
CLOZAPINE – Hospital pharmacy [HP4]			· ·
Safety medicine; prescriber may determine dispensing freque	encv		
Tab 25 mg	•	50	✓ Clopine
740 25 mg		00	✓ Clozaril
	13.37	100	✓ Clopine
			✓ Clozaril
Tab 50 mg	8.67	50	✓ Clopine
· ·	17.33	100	✓ Clopine
Tab 100 mg	17.33	50	✓ Clopine
•			✓ Clozaril
	34.65	100	✓ Clopine
			✓ Clozaril
Tab 200 mg	34.65	50	✓ Clopine
	69.30	100	✓ Clopine
Suspension 50 mg per ml	67.62	100 ml	✓ Versacloz
HALOPERIDOL - Safety medicine; prescriber may determine dis	pensing frequer	псу	
Tab 500 mcg - Up to 30 tab available on a PSO	6.23	100	✓ Serenace
Tab 1.5 mg - Up to 30 tab available on a PSO	9.43	100	✓ Serenace
Tab 5 mg - Up to 30 tab available on a PSO	14.86	50	✓ Serenace
	29.72	100	✓ Serenace
Oral liq 2 mg per ml - Up to 200 ml available on a PSO		100 ml	✓ Serenace
Inj 5 mg per ml, 1 ml ampoule – Up to 5 inj available on a PS	O21.55	10	✓ Serenace
LEVOMEPROMAZINE - Safety medicine; prescriber may determ	nine dispensing t	frequency	
Tab 25 mg (33.8 mg as a maleate)		100	✓ Nozinan (Swiss)
Tab 25 mg as a maleate	16.10	100	✓ Nozinan `
Tab 100 mg (135 mg as a maleate)	41.75	100	✓ Nozinan (Swiss)
Tab 100 mg as a maleate	41.75	100	✓ Nozinan

	Subsidy		Fully Brand or
	(Manufacturer's Price) \$	Sul Per	bsidised Generic Manufacturer
EVOMEPROMAZINE HYDROCHLORIDE - Safety medicine	e: prescriber may deterr	nine disp	ensina freauency
Inj 25 mg per ml, 1 ml ampoule	•	5	✓ Neuraxpharm \$29
11) 25 11g por 111, 1 111 ampoulo		Ü	✓ Nozinan S29 S29
	24.48	10	✓ Wockhardt
Neuraxpharm 👀 Inj 25 mg per ml, 1 ml ampoule to be delis		10	Wookilaiat
Nozinan S29 S29 Inj 25 mg per ml, 1 ml ampoule to be delis	• ,		
	,		
THIUM CARBONATE – Safety medicine; prescriber may de	, ,		/ Balantal
Tab long-acting 400 mg		100	✓ <u>Priadel</u>
Cap 250 mg		100	Douglas
LANZAPINE – Safety medicine; prescriber may determine d			_
Tab 2.5 mg	1.40	30	Zypine
Zypine to be Principal Supply on 1 August 2024			
Tab 5 mg	1.93	30	Zypine
Zypine to be Principal Supply on 1 August 2024			/-
Tab orodispersible 5 mg		28	Zypine ODT
Tab 10 mg	1.93	30	Zypine
Zypine to be Principal Supply on 1 August 2024			4 - 4 - 4
Tab orodispersible 10 mg	2.89	28	✓ Zypine ODT
ERICYAZINE - Safety medicine; prescriber may determine	dispensing frequency		
Tab 2.5 mg	12.49	100	✓ Neulactil
Tab 10 mg	44.45	100	✓ Neulactil
UETIAPINE - Safety medicine; prescriber may determine di	spensing frequency		
Tab 25 mg	2.36	90	✓ Quetapel
Tab 100 mg		90	✓ Quetapel
Tab 200 mg	10.97	90	✓ Quetapel
Tab 300 mg	15.83	90	✓ Quetapel
ISPERIDONE - Safety medicine; prescriber may determine	dispensing frequency		
Tab 0.5 mg		60	✓ Risperidone (Teva)
Tab 1 mg		60	✓ Risperidone (Teva)
Tab 2 mg		60	✓ Risperidone (Teva)
Tab 3 mg		60	✓ Risperidone (Teva)
Tab 4 mg		60	✓ Risperidone (Teva)
Oral liq 1 mg per ml		30 ml	✓ Risperon
	17.80	100 ml	Risperon
PRASIDONE - Safety medicine; prescriber may determine	dispensing frequency		
Cap 20 mg		60	✓ Zusdone
Cap 40 mg		60	✓ Zusdone
Cap 60 mg		60	✓ Zusdone
Cap 80 mg		60	✓ Zusdone
JCLOPENTHIXOL HYDROCHLORIDE - Safety medicine; p			
Tab 10 mg		100	✓ Clopixol
140 15 mg		100	- оторилог
Depot Injections			
RIPIPRAZOLE - Special Authority see SA2312 on the next	nage – Retail nharmacı	,	
Safety medicine; prescriber may determine dispensing fre		'	
		4	Abilify Maintena 200
Inj 300 mg vial	2/3.50	1	✓ Abilify Maintena S29
Inj 400 mg vial	341.96	1	✓ Abilify Maintena S29
"1] 700 mg viai			- Ability Mailitella 020

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sub	sidised	Generic
\$	Per	✓	Manufacturer

⇒SA2312 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 Both:
 - 1.1 Patient has a current Special Authority approval for olanzapine depot injection, risperidone depot injection or paliperidone depot injection; and
 - 1.2 Patient has tried but has experienced an inadequate response to, or intolerable side effects from, prior therapy with olanzapine depot injection, risperidone depot injection or paliperidone depot injection; or
- 2 Patient has been unable to access olanzapine depot injection due to supply issues with olanzapine depot injection, or otherwise would have been initiated on olanzapine depot injection but has been unable to due to supply issues with olanzapine depot injection (see Note below for the olanzapine Special Authority criteria for new olanzapine depot injection patients prior to 1 April 2024).

Notes: The Olanzapine depot injection Special Authority criteria that apply to criterion 2 in this Aripiprazole Special Authority application are as follows:

- The patient has had an initial Special Authority approval for paliperidone depot injection or risperidone depot injection; or
- All of the following:
 - The patient has schizophrenia: and
 - The patient has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of aripiprazole depot injection has been associated with fewer days of intensive intervention than prior to the initiation of an atypical antipsychotic depot injection. FLUPENTHIXOL DECANOATE – Safety medicine; prescriber may determine dispensing frequency

Inj 20 mg per ml, 1 ml – Up to 5 inj available on a PS	O13.14	5	✓ Fluanxol
Inj 20 mg per ml, 2 ml - Up to 5 inj available on a PS	O20.90	5	✓ Fluanxol
Inj 100 mg per ml, 1 ml - Up to 5 inj available on a P	SO40.87	5	✓ Fluanxol
HALOPERIDOL DECANOATE - Safety medicine; prescr	iber may determine disper	nsing frequ	ency
Inj 50 mg per ml, 1 ml - Up to 5 inj available on a PS	O28.39	5	✓ Haldol
Inj 100 mg per ml, 1 ml - Up to 5 inj available on a Ps	SO55.90	5	✓ Haldol Concentrate
			✓ Haldol
			Decanoas \$29
OLANZAPINE - Special Authority see SA2313 below - F	tetail pharmacy		
a) Safety medicine; prescriber may determine dispen	sing frequency		
b) Note - no new patients to be initiated on olanzapir	ne.		
Inj 210 mg vial	252.00	1	✓ Zyprexa Relprevv
Inj 300 mg vial	414.00	1	✓ Zyprexa Relprevv
Inj 405 mg vial	504.00	1	✓ Zyprexa Relprevv

⇒SA2313 Special Authority for Subsidy

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of olanzapine depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

PALIPERIDONE - Special Authority see SA1429 on the next page - Retail pharmacy

Safety medicine; prescriber may determine dispensing fre	equency		
Inj 25 mg syringe	194.25	1	✓ Invega Sustenna
Inj 50 mg syringe	271.95	1	✓ Invega Sustenna
Inj 75 mg syringe	357.42	1	✓ Invega Sustenna
Inj 100 mg syringe	435.12	1	✓ Invega Sustenna
Inj 150 mg syringe		1	✓ Invega Sustenna



Subsidy (Manufacturer's Price) \$ Per

Fully Subsidised Brand or Generic Manufacturer

⇒SA1429 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Fither:

- 1 The patient has had an initial Special Authority approval for risperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 Has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

PALIPERIDONE PALMITATE - Special Authority see SA2167 below - Retail pharmacy

Inj 175 mg syringe	815.85	1	Invega Trinza
	1,072.26	1	✓ Invega Trinza
Inj 350 mg syringe	1,305.36	1	✓ Invega Trinza
Inj 525 mg syringe	1,305.36	1	✓ Invega Trinza

⇒SA2167 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has schizophrenia; and
- 2 The patient has had an initial Special Authority approval for paliperidone once-monthly depot injection.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

RISPERIDONE - Special Authority see SA1427 below - Retail pharmacy

Safety medicine; prescriber may determine dispensing fre	quency		
Inj 25 mg vial	135.98	1	 Risperdal Consta
Inj 37.5 mg vial	178.71	1	✓ Risperdal Consta
Inj 50 mg vial	217.56	1	✓ Risperdal Consta

⇒SA1427 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 Has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of risperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

ZUCLOPENTHIXOL DECANOATE - Safety medicine; prescriber may determine dispensing frequency

Inj 200 mg per ml, 1 ml − Up to 5 inj available on a PSO19.80 5 **Clopixol**

Brand or

Generic

Ativan

Fully

Subsidised

100

	\$	Per	✓ Manufacturer
Anxiolytics			
BUSPIRONE HYDROCHLORIDE			
* Tab 5 mg	18.50	100	✓ Buspirone Viatris
* Tab 10 mg	12.50	100	✓ Buspirone Viatris
CLONAZEPAM - Safety medicine; prescriber may determine disp	ensing frequenc	ev	
Tab 500 mcg		100	✓ Paxam
Tab 2 mg	10.78	100	✓ Paxam
DIAZEPAM - Safety medicine; prescriber may determine dispensi	ing frequency		
Tab 2 mg	95.00	500	✓ Arrow-Diazepam
Tab 5 mg	115.00	500	✓ Arrow-Diazepam
LORAZEPAM - Safety medicine; prescriber may determine disper			
Tab 1 mg		250	✓ <u>Ativan</u>

Subsidy

(Manufacturer's Price)

Multiple Sclerosis Treatments

⇒SA2274 Special Authority for Subsidy

Initial application — (Multiple Sclerosis - dimethyl fumarate, fingolimod, glatiramer acetate, interferon beta-1-alpha, interferon beta-1-beta, natalizumab and teriflunomide) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Fither:

- 1 All of the following:
 - 1.1 Diagnosis of multiple sclerosis (MS) meets the McDonald 2017 diagnostic criteria for MS and has been confirmed by a neurologist; and
 - 1.2 Patient has an EDSS score between 0 6.0; and

- 1.3 Patient has had at least one significant attack of MS in the previous 12 months or two significant attacks in the past 24 months; and
- 1.4 All of the following:
 - 1.4.1 Each significant attack must be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the attack, but the neurologist/physician must be satisfied that the clinical features were characteristic); and
 - 1.4.2 Each significant attack is associated with characteristic new symptom(s)/sign(s) or substantially worsening of previously experienced symptoms(s)/sign(s); and
 - 1.4.3 Each significant attack has lasted at least one week and has started at least one month after the onset of a previous attack (where relevant); and
 - 1.4.4 Each significant attack can be distinguished from the effects of general fatigue; and is not associated with a fever (T> 37.5°C); and
 - 1.4.5 Either:
 - 1.4.5.1 Each significant attack is severe enough to change either the EDSS or at least one of the Kurtze Functional System scores by at least 1 point; or
 - 1.4.5.2 Each significant attack is a recurrent paroxysmal symptom of multiple sclerosis (tonic seizures/spasms, trigeminal neuralgia, Lhermitte's symptom); and
- 1.5 Evidence of new inflammatory activity on an MRI scan within the past 24 months; and
- 1.6 Any of the following:
 - 1.6.1 A sign of that new inflammatory activity on MRI scanning (in criterion 5 immediately above) is a gadolinium enhancing lesion; or



NERV	/008	SYSIE	IVI										
						(1	Subsic Manufacture \$		S Per	Full; Subsidised	d G	Brand or Generic Manufactur	er
continued	1.6.3	A sign o	f that new	inflamm	atory is a	T2 lesion	esion showi with assoc	ciated loc	cal swe	elling; or		nsible for	the clinical
Note: Treatm	t has ar nent on	A sign on active a two or me	f that new pproval foore funde	inflamm or ocreliz d multiple	atory action attention at the action at the	vity is nev does not treatmer		s compa ary prog neously i	red wit ressive s not p	e MS. ermitted	d.		
Renewal — (I beta-1-beta, r nad an EDSS he patient ha Note: Treatm	natalizu score d s walke	Imab an of 0 to 6.0 d 100 me	d teriflun (inclusiv etres or m	omide) e) with o ore with	from any i r without t or without	relevant p he use of aids in th	ractitioner. unilateral one last six r	Approvor bilater nonths).	als val al aids	id for 12 at any t	2 mont time ir	ths where	e patient has
DIMETHYL FU a) Wasta b) Note:	age clai	mable .		•		·	evious pag			,	ermitt	ted	
Cap 120 i Cap 240 i	mg mg				············		520.00 2,000.00		14 56	7	Tect	fidera fidera	
Cap 0.5 n	age clai Treatn	mable nent on tv	vo or mor	e funded	multiple s	sclerosis t	reatments 2,200.00	simultan	eously 28	1	ermitt		
Inj 40 mg	eatmen prefille	t on two o d syringe	or more fu	inded mu	ıltiple scle	rosis trea	tments sim 1,137.48	ultaneou	ısly is ı 12	not perm	Cop	axone	
Inj 6 millio	eatmen	t on two o	or more fu inge	ınded mu	ıltiple scle	rosis trea	the previo tments sim 1,170.00 1,170.00	ultaneou		not perm	nitted. ′ Avo		
Inj 8 millio	eatmen on iu pe	t on two o	or more fu	ınded mu	ıltiple scle	rosis trea	tments sim 1,322.89	ultaneou	ısly is ı 15	not perm	nitted.	aferon	
Inj 20 mg	eatmen per ml,	t on two o	or more fu al	ınded mu	ıltiple scle	rosis trea	tments sim 1,750.00	ultaneou	ısly is ı 1		nitted. Tys a		
TERIFLUNON a) Wasta b) Note:	age clai	nable	•				s page – Re reatments	·	,	is not p	ermitt	ted.	

Multiple Sclerosis Treatments - Other

OCRELIZUMAB - Special Authority see SA2273 on the next page - Retail pharmacy Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted. 1 ✓ Ocrevus

Aubagio

(ie

Subsidy (Manufacturer's Price) \$

Subsidised Per

Fully

Brand or Generic Manufacturer

⇒SA2273 Special Authority for Subsidy

Initial application — (Multiple Sclerosis - ocrelizumab) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 Diagnosis of multiple sclerosis (MS) meets the McDonald 2017 diagnostic criteria for MS and has been confirmed by a neurologist; and
 - 1.2 Patient has an EDSS score between 0 6.0; and
 - 1.3 Patient has had at least one significant attack of MS in the previous 12 months or two significant attacks in the past 24 months; and
 - 1.4 All of the following:
 - 1.4.1 Each significant attack must be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the attack, but the neurologist/physician must be satisfied that the clinical features were characteristic); and
 - 1.4.2 Each significant attack is associated with characteristic new symptom(s)/sign(s) or substantially worsening of previously experienced symptoms(s)/sign(s); and
 - 1.4.3 Each significant attack has lasted at least one week and has started at least one month after the onset of a previous attack (where relevant); and
 - 1.4.4 Each significant attack can be distinguished from the effects of general fatigue; and is not associated with a fever (T> 37.5°C); and
 - 1.4.5 Fither:
 - 1.4.5.1 Each significant attack is severe enough to change either the EDSS or at least one of the Kurtze Functional System scores by at least 1 point; or
 - 1.4.5.2 Each significant attack is a recurrent paroxysmal symptom of multiple sclerosis (tonic seizures/spasms, trigeminal neuralgia, Lhermitte's symptom); and
 - 1.5 Evidence of new inflammatory activity on an MRI scan within the past 24 months; and
 - 1.6 Any of the following:
 - 1.6.1 A sign of that new inflammatory activity on MRI scanning (in criterion 5 immediately above) is a gadolinium enhancing lesion; or
 - 1.6.2 A sign of that new inflammatory activity is a lesion showing diffusion restriction; or
 - 1.6.3 A sign of that new inflammatory is a T2 lesion with associated local swelling; or
 - 1.6.4 A sign of that new inflammatory activity is a prominent T2 lesion that clearly is responsible for the clinical features of a recent attack that occurred within the last 2 years; or
 - 1.6.5 A sign of that new inflammatory activity is new T2 lesions compared with a previous MRI scan; or
- 2 Patient has an active Special Authority approval for either dimethyl fumarate, fingolimod, glatiramer acetate, interferon beta-1-alpha. interferon beta-1-beta. natalizumab or teriflunomide.

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

Renewal — (Multiple Sclerosis - ocrelizumab) from any relevant practitioner. Approvals valid for 12 months where patient has had an EDSS score of 0 to 6.0 (inclusive) with or without the use of unilateral or bilateral aids at any time in the last six months (ie the patient has walked 100 metres or more with or without aids in the last six months).

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

Initial application — (Primary Progressive Multiple Sclerosis) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Diagnosis of primary progressive multiple sclerosis (PPMS) meets the 2017 McDonald criteria and has been confirmed by a neurologist; and
- 2 Patient has an EDSS 2.0 (score equal to or greater than 2 on pyramidal functions) to EDSS 6.5; and



Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

3 Patient has no history of relapsing remitting multiple sclerosis.

Renewal — (Primary Progressive Multiple Sclerosis) from any relevant practitioner. Approvals valid for 12 months where patient has had an EDSS score of less than or equal to 6.5 at any time in the last six months (ie patient has walked 20 metres with bilateral assistance/aids, without rest in the last six months).

Sedatives and Hypnotics

MELATONIN - Special Authority see SA1666 below - Retail pharmacy

Tab modified-release 2 mg - No more than 5 tab per day11.50 30 ✓ Viaisom

Restricted to patients aged 18 years or under.

⇒SA1666 Special Authority for Subsidy

Initial application only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with persistent and distressing insomnia secondary to a neurodevelopmental disorder (including, but not limited to, autism spectrum disorder or attention deficit hyperactivity disorder)*; and
- 2 Behavioural and environmental approaches have been tried and were unsuccessful, or are inappropriate; and
- 3 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day; and
- 4 Patient is aged 18 years or under*.

Renewal only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is aged 18 years or under*; and
- 2 Patient has demonstrated clinically meaningful benefit from funded modified-release melatonin (clinician determined); and
- 3 Patient has had a trial of funded modified-release melatonin discontinuation within the past 12 months and has had a recurrence of persistent and distressing insomnia; and
- 4 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day.

Note: Indications marked with * are unapproved indications.

MIDAZOLAM – Safety medicine; prescriber may determine dispension in 1 mg per ml, 5 ml ampoule	. ,	10	✓ Midazolam-Baxter
Inj 1 mg per ml, 5 ml plastic ampoule – Up to 10 inj available	0.10	10	• WIIUazUlaIII-Daxlei
on a PSOon a PSO	17.28	10	✓ Pfizer
On a PSO for status epilepticus use only. PSO must be er	ndorsed for statu	s epilepticu	is use only.
Inj 5 mg per ml, 3 ml ampoule	5.00	5	✓ Midazolam-Baxter
Inj 5 mg per ml, 3 ml plastic ampoule - Up to 5 inj available or	1		
a PSO	13.09	5	✓ Pfizer
On a PSO for status epilepticus use only. PSO must be er	ndorsed for statu	s epilepticu	is use only.
PHENOBARBITONE SODIUM - Special Authority see SA1386 be	ow – Retail pha	rmacy	
Inj 200 mg per ml, 1 ml ampoule	113.37	10	✓ Max Health S29
⇒SA1386 Special Authority for Subsidy			

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 For the treatment of terminal agitation that is unresponsive to other agents; and
- 2 The applicant is part of a multidisciplinary team working in palliative care.

TEMAZEPAM - Safety medicine; prescriber may determine dispensing frequency

25 ✓ Normison Tab 10 mg1.40

NERVOUS SYSTEM

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer	
ZOPICLONE – Safety medicine; prescriber may determine dispe	nsing frequency				
Tab 7.5 mg	10.80	500	✓ Z	opicione Actavis	

Spinal Muscular Atrophy

NUSINERSEN - PCT only - Special Authority see SA2174 below

⇒SA2174 Special Authority for Subsidy

Initial application — (spinal muscular atrophy (SMA)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has genetic documentation of homozygous SMN1 gene deletion, homozygous SMN1 point mutation, or compound heterozygous mutation; and
- 2 Patient is 18 years of age or under; and
- 3 Either:
 - 3.1 Patient has experienced the defined signs and symptoms of SMA type I, II or IIIa prior to three years of age; or
 - 3.2 Both:
 - 3.2.1 Patient is pre-symptomatic; and
 - 3.2.2 Patient has three or less copies of SMN2.

Renewal — (spinal muscular atrophy (SMA)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 There has been demonstrated maintenance of motor milestone function since treatment initiation; and
- 2 Patient does not require invasive permanent ventilation (at least 16 hours per day) in the absence of a potentially reversible cause while being treated with nusinersen; and
- 3 Nusinersen not to be administered in combination other SMA disease modifying treatments or gene therapy.

RISDIPLAM - [Xpharm] - Special Authority see SA2203 below

Note: the supply of risdiplam is via Pharmac's approved direct distribution supply. Further details can be found on Pharmac's website https://pharmac.govt.nz/risdiplam

Powder for oral soln 750 mcg per ml. 60 mg per bottle......14,100.00 80 ml OP ✓ Evrysdi

⇒SA2203 Special Authority for Subsidy

Initial application — (spinal muscular atrophy (SMA)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has genetic documentation of homozygous SMN1 gene deletion, homozygous SMN1 point mutation, or compound heterozygous mutation; and
- 2 Patient is 18 years of age or under; and
- 3 Either:
 - 3.1 Patient has experienced the defined signs and symptoms of SMA type I, II or IIIa prior to three years of age; or
 - 3.2 Both:
 - 3.2.1 Patient is pre-symptomatic; and
 - 3.2.2 Patient has three or less copies of SMN2.

Renewal — (spinal muscular atrophy (SMA)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 There has been demonstrated maintenance of motor milestone function since treatment initiation; and
- 2 Patient does not require invasive permanent ventilation (at least 16 hours per day) in the absence of a potentially reversible cause while being treated with risdiplam; and
- 3 Risdiplam not to be administered in combination other SMA disease modifying treatments or gene therapy.

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	Generic
	\$	Per	1	Manufacturer
Chimadanta (ADUD Taraturanta				
Stimulants/ADHD Treatments				
ATOMOXETINE				
Cap 10 mg	18.41	28	1	APO-Atomoxetine
				S29 S29
				Generic Partners
ABO 4: "	43.02		•	APO-Atomoxetine
APO-Atomoxetine to be Principal Supply on 1 August 202		28		Generic Partners
Cap 18 mg	45.57	20		APO-Atomoxetine
APO-Atomoxetine to be Principal Supply on 1 August 202			•	Ar O-Atomoxetime
Cap 25 mg		28	1	Generic Partners
3.77 - 3	44.30		1	APO-Atomoxetine
APO-Atomoxetine to be Principal Supply on 1 August 202	24			
Cap 40 mg	29.22	28		Generic Partners
	46.21		/	APO-Atomoxetine
APO-Atomoxetine to be Principal Supply on 1 August 202				
Cap 60 mg	46.51	28	•	APO-Atomoxetine
			_	S29 S29
	E4 04			Generic Partners APO-Atomoxetine
APO-Atomoxetine to be Principal Supply on 1 August 202	51.31		•	APO-Atomoxetine
Cap 80 mg		28	1	APO-Atomoxetine
σαρ σο πης		20	•	S29 S29
			1	Generic Partners
	65.20			APO-Atomoxetine
APO-Atomoxetine to be Principal Supply on 1 August 202				7.1. • 7.1.•
Cap 100 mg		28	✓	APO-Atomoxetine
				S29 S29
			1	Generic Partners
	65.71		✓	APO-Atomoxetine
APO-Atomoxetine to be Principal Supply on 1 August 202	24			
(APO-Atomoxetine S29 S29 Cap 10 mg to be delisted 1 August 2	2024)			
(Generic Partners Cap 10 mg to be delisted 1 August 2024)				
(Generic Partners Cap 18 mg to be delisted 1 August 2024)				
(Generic Partners Cap 25 mg to be delisted 1 August 2024)				
(Generic Partners Cap 40 mg to be delisted 1 August 2024)	2024)			
(APO-Atomoxetine S29 S29 Cap 60 mg to be delisted 1 August 2 (Generic Partners Cap 60 mg to be delisted 1 August 2024)	2024)			
(APO-Atomoxetine S29 S29 Cap 80 mg to be delisted 1 August 2024)	2024)			
(Generic Partners Cap 80 mg to be delisted 1 August 2024)	1024)			
(APO-Atomoxetine S29 S29 Cap 100 mg to be delisted 1 August	2024)			
(Generic Partners Cap 100 mg to be delisted 1 August 2024)	-0-1/			
DEXAMFETAMINE SULFATE - Special Authority see SA1149 or	the next page – Re	tail n	narmacy	
a) Only on a controlled drug form	nom pago Tio	P		
b) Safety medicine; prescriber may determine dispensing free	quency			
Tab 5 mg	' '	100	✓	Noumed
				Dexamfetamine

NERVOUS SYSTEM

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

⇒SA1149 Special Authority for Subsidy

Initial application — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

Renewal — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Renewal — **(ADHD in patients under 5)** only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.



(Manufacturer's Price) Subsidised Generic \$ Per ✓ Manufacturer	Subsidy		Fully	Brand or
\$ Per ✓ Manufacturer	(Manufacturer's Price)	Subs	sidised	Generic
	 \$	Per	•	Manufacturer

METHYLPHENIDATE HYDROCHLORIDE - Special Authority see SA1964 below - Retail pharmacy

- a) Only on a controlled drug form
- b) Safety medicine; prescriber may determine dispensing frequency
- c) Note: Brand Switch Fee applies only to patients who have transferred from Concerta brand due to an out of stock

c) Note: Brand Switch Fee applies only to patients who have	ransterred tron	n Concerta t	rand due to an out of stock.
Tab immediate-release 5 mg	3.20	30	✓ Rubifen
Tab immediate-release 10 mg	3.00	30	✓ Ritalin
v			✓ Rubifen
Tab extended-release 18 mg - Brand switch fee payable			
(Pharmacode 2677822) - see page 264 for details	7.75	30	Methylphenidate ERTeva
Tab immediate-release 20 mg	7.85	30	✓ Rubifen
Tab sustained-release 20 mg		30	Rubifen SR
Tab extended-release 27 mg - Brand switch fee payable			
(Pharmacode 2677822) - see page 264 for details	11.45	30	Methylphenidate ERTeva
Tab extended-release 36 mg - Brand switch fee payable			
(Pharmacode 2677822) - see page 264 for details	15.50	30	Methylphenidate ERTeva
Tab extended-release 54 mg - Brand switch fee payable			
(Pharmacode 2677822) - see page 264 for details	22.25	30	Methylphenidate ERTeva

⇒SA1964 Special Authority for Subsidy

Initial application — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy*) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

Note: *narcolepsy is not a registered indication for Methylphenidate ER – Teva.

Renewal — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been

NERVOUS SYSTEM

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

consulted within the last 2 years and has recommended treatment for the patient in writing.

Renewal — **(ADHD in patients under 5)** only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Narcolepsy*) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

Note: *narcolepsy is not a registered indication for Methylphenidate ER – Teva.

METHYLPHENIDATE HYDROCHLORIDE EXTENDED-RELEASE - Special Authority see SA2305 below - Retail pharmacy

- a) Only on a controlled drug form
- b) Safety medicine: prescriber may determine dispensing frequency

Tab extended-release 18 mg	58.96	30	Concerta
Tab extended-release 27 mg	65.44	30	Concerta
Tab extended-release 36 mg	71.93	30	Concerta
Tab extended-release 54 mg	86.24	30	Concerta
Cap modified-release 10 mg		30	Ritalin LA
Cap modified-release 20 mg		30	Ritalin LA
Cap modified-release 30 mg		30	Ritalin LA
Cap modified-release 40 mg		30	Ritalin LA
. •			

⇒SA2305 Special Authority for Subsidy

Initial application — (ADHD) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder); and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing; and
- 4 Either:
 - 4.1 Patient is taking a currently subsidised formulation of methylphenidate hydrochloride (immediate-release or sustained-release) which has not been effective due to significant administration and/or difficulties with adherence; or
 - 4.2 There is significant concern regarding the risk of diversion or abuse of immediate-release methylphenidate hydrochloride.

Renewal — (ADHD) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

MODAFINIL – Special Authority see SA1999 on the next page -	 Retail pharmacy 		
Tab 100 mg	29.13	60	✓ Modavigil



	Subsidy	Fully	Brand or
(Manufa	, ,	sidised	Generic
	\$ Per	·	Manutacturer

⇒SA1999 Special Authority for Subsidy

Initial application only from a neurologist or respiratory specialist. Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of narcolepsy and has excessive daytime sleepiness associated with narcolepsy occurring almost daily for three months or more: and
- - 2.1 The patient has a multiple sleep latency test with a mean sleep latency of less than or equal to 10 minutes and 2 or more sleep onset rapid eve movement periods; or
 - 2.2 The patient has at least one of: cataplexy, sleep paralysis or hypnagogic hallucinations; and
- 3 Either:
 - 3.1 An effective dose of a subsidised formulation of methylphenidate or dexamfetamine has been trialled and discontinued because of intolerable side effects; or
 - 3.2 Methylphenidate and dexamfetamine are contraindicated.

Renewal only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

Treatments for Dementia

DONEPEZIL HYDROCHLORIDE - Brand	d switch fee payable (Pharmacode 267972)	8) - see <mark>pa</mark>	age 264 for details
* Tab 5 mg	3.70	84	✓ Ipca-Donepezil
* Tab 10 mg		84	✓ Ipca-Donepezil
RIVASTIGMINE - Special Authority see \$	SA1488 below – Retail pharmacy		
Patch 4.6 mg per 24 hour	38.00	30	✓ <u>Rivastigmine Patch</u> <u>BNM 5</u>
	90.00		Exelon Patch 5
Patch 9.5 mg per 24 hour	38.00	30	✓ Rivastigmine Patch BNM 10
	90.00		✓ Exelon Patch 10

⇒SA1488 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The patient has been diagnosed with dementia; and
- 2 The patient has experienced intolerable nausea and/or vomiting from donepezil tablets.

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate; and
- 2 The patient has demonstrated a significant and sustained benefit from treatment.

Treatments for Substance Dependence

BUPRENORPHINE WITH NALOXONE - Special Authority see SA1203 on the next page - Retail pharmacy

- a) No patient co-payment payable
- b) Safety medicine: prescriber may determine dispensing frequency
- Tab sublingual 8 mg with naloxone 2 mg34.00 28

✓ Buprenorphine Naloxone BNM

✓ Buprenorphine

Subsidy (Manufacturer's Price) S \$ Per

Fully Subsidised

Brand or Generic Manufacturer

⇒SA1203 Special Authority for Subsidy

Initial application — (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health...

Initial application — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient will not be receiving methadone; and
- 3 Patient is currently enrolled in an opioid substitution treatment program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient has previously trialled but failed detoxification with buprenorphine with naloxone with relapse back to opioid use and another attempt is planned; and
- 3 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is or has been receiving maintenance therapy with buprenorphine with naloxone (and is not receiving methadone); and
- 2 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health or is a medical practitioner authorised by the service to manage treatment in this patient.

Renewal — (Maintenance treatment where the patient has previously had an initial application for detoxification) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient received but failed detoxification with buprenorphine with naloxone; and
- 2 Maintenance therapy with buprenorphine with naloxone is planned (and patient will not be receiving methadone); and
- 3 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Tab modified-release 150 mg	15.00	30	✓ <u>Zyban</u>
DISULFIRAM Tab 200 mg	236.40	100	✓ Antabuse
NALTREXONE HYDROCHLORIDE - Special Authority se	ee SA1408 on the next p	age – Retail	pharmacy
Tab 50 mg	77.77	28	✓ Naltrexone AOP \$29
Aldress ACR	83.33	30	✓ Naltraccord
(Naltrexone AOP S29) Tab 50 mg to be delisted 1 August	2024)		



Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

⇒SA1408 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient is currently enrolled in a recognised comprehensive treatment programme for alcohol dependence; and
- 2 Applicant works in or with a community Alcohol and Drug Service contracted to Health NZ or accredited against the New Zealand Alcohol and Other Drug Sector Standard or the National Mental Health Sector Standard.

Renewal from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Compliance with the medication (prescriber determined); and
- 2 Any of the following:
 - 2.1 Patient is still unstable and requires further treatment; or
 - 2.2 Patient achieved significant improvement but requires further treatment; or
 - 2.3 Patient is well controlled but requires maintenance therapy.

NICOTINE

a) Nicotine will not be funded in amounts less than 4 weeks of treatment.

b) Note: Direct Provision by a pharmacist permitted under the provisions in	Part I of Section	on A.
Patch 7 mg - Up to 28 patch available on a PSO19.14	28	Habitrol
Patch 7 mg for direct distribution only - [Xpharm]4.13	7	Habitrol
Patch 14 mg - Up to 28 patch available on a PSO21.05	28	Habitrol
Patch 14 mg for direct distribution only - [Xpharm]12.49	7	Habitrol
Patch 21 mg - Up to 28 patch available on a PSO24.12	28	Habitrol
Patch 21 mg for direct distribution only - [Xpharm]13.19	7	Habitrol
Lozenge 1 mg - Up to 216 loz available on a PSO19.76	216	Habitrol
Lozenge 1 mg for direct distribution only - [Xpharm]12.89	36	Habitrol
Lozenge 2 mg - Up to 216 loz available on a PSO21.65	216	Habitrol
Lozenge 2 mg for direct distribution only - [Xpharm]	36	Habitrol
Gum 2 mg (Fruit) - Up to 204 piece available on a PSO21.42	204	Habitrol
Gum 2 mg (Fruit) for direct distribution only - [Xpharm]17.57	96	Habitrol
Gum 2 mg (Mint) - Up to 204 piece available on a PSO21.42	204	Habitrol
Gum 2 mg (Mint) for direct distribution only - [Xpharm]17.57	96	Habitrol
Gum 4 mg (Fruit) - Up to 204 piece available on a PSO24.17	204	Habitrol
Gum 4 mg (Fruit) for direct distribution only - [Xpharm]23.87	96	Habitrol
Gum 4 mg (Mint) - Up to 204 piece available on a PSO24.17	204	Habitrol
Gum 4 mg (Mint) for direct distribution only - [Xpharm]23.87	96	Habitrol
(Habitrol Patch 7 mg for direct distribution only to be delisted 1 August 2024)		

VARENICLINE TARTRATE - Special Authority see SA1845 below - Retail pharmacy

- a) A maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval, including the starter pack
- b) Varenicline will not be funded in amounts less than 4 weeks of treatment.
- c) The 6-month time period in which a patient can receive a funded 12-week course of varenicline tartrate starts from the date the Special Authority is approved.

Tab 0.5 mg × 11 and 1 mg × 42	16.67	53 OP	✓ Varenicline Pfizer
Tab 1 mg	17.62	56	✓ Varenicline Pfizer

⇒SA1845 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and

NERVOUS SYSTEM

(Manu	Subsidy	Fully	Brand or
	facturer's Price)	Subsidised	Generic
	\$ Pe	er 🗸	Manufacturer

continued...

- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 Either:
 - 3.1 The patient has tried but failed to quit smoking after at least two separate trials of nicotine replacement therapy, at least one of which included the patient receiving comprehensive advice on the optimal use of nicotine replacement therapy; or
 - 3.2 The patient has tried but failed to quit smoking using bupropion or nortriptyline; and
- 4 The patient has not had a Special Authority for varenicline approved in the last 6 months; and
- 5 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 6 The patient is not pregnant; and
- 7 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

Renewal from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 It has been 6 months since the patient's previous Special Authority was approved; and
- 4 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 5 The patient is not pregnant; and
- 6 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

The patient must not have had an approval in the past 6 months.

Notes: a maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval.

This includes the 4-week 'starter' pack.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

Chemotherapeutic Agents

Alkylating Agents

BENDAMUSTINE HYDROCHLORIDE - PCT only - Specialist - Special Authority see SA2153 below

Inj 25 mg vial	 	77.00	1	✓ Ribomustin
Inj 100 mg vial	 	308.00	1	✓ Ribomustin
Inj 1 mg for ECP	 	3.23	1 mg	✓ Baxter

⇒SA2153 Special Authority for Subsidy

Initial application — (treatment naive CLL) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has Binet stage B or C, or progressive stage A chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is chemotherapy treatment naive; and
- 3 The patient is unable to tolerate toxicity of full-dose FCR; and
- 4 Patient has ECOG performance status 0-2; and
- 5 Patient has a Cumulative Illness Rating Scale (CIRS) score of < 6; and
- 6 Bendamustine is to be administered at a maximum dose of 100 mg/m² on days 1 and 2 every 4 weeks for a maximum of 6 cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL). Chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initial application — (Indolent, Low-grade lymphomas) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: All of the following:

- 1 The patient has indolent low grade NHL requiring treatment; and
- 2 Patient has a WHO performance status of 0-2; and
- 3 Any of the following:
 - 3.1 Both:
 - 3.1.1 Patient is treatment naive; and
 - 3.1.2 Bendamustine is to be administered for a maximum of 6 cycles (in combination with rituximab when CD20+); or
 - 3.2 Both:
 - 3.2.1 Patient is refractory to or has relapsed within 12 months of a rituximab containing combined chemo-immunotherapy regimen; and
 - 3.2.2 Bendamustine is to be administered in combination with obinutuzumab for a maximum of 6 cycles; or
 - 3.3 All of the following:
 - 3.3.1 The patient has not received prior bendamustine therapy; and
 - 3.3.2 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+): and
 - 3.3.3 Patient has had a rituximab treatment-free interval of 12 months or more; or
 - 3.4 Bendamustine is to be administered as monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Renewal — (Indolent, Low-grade lymphomas) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient is refractory to or has relapsed within 12 months of rituximab in combination with bendamustine; and
 - 1.2 Bendamustine is to be administered in combination with obinutuzumab for a maximum of 6 cycles; or

Subsidy (Manufacturer's Price)	S	Fully ubsidised	Brand or Generic	
\$	Per	✓	Manufacturer	

continued...

- 2 Both:
 - 2.1 Patients have not received a bendamustine regimen within the last 12 months; and
 - 2.2 Either:
 - 2.2.1 Both:
 - 2.2.1.1 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
 - 2.2.1.2 Patient has had a rituximab treatment-free interval of 12 months or more; or
 - 2.2.2 Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Note: 'indolent, low-grade lymphomas' includes follicular, mantle cell, marginal zone and lymphoplasmacytic/ Waldenstrom's macroglobulinaemia.

Initial application — (Hodgkin's lymphoma*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has Hodgkin's lymphoma requiring treatment; and
- 2 Patient has a ECOG performance status of 0-2: and
- 3 Patient has received one prior line of chemotherapy; and
- 4 Patient's disease relapsed or was refractory following prior chemotherapy; and
- 5 Bendamustine is to be administered in combination with gemcitabine and vinorelbine (BeGeV) at a maximum dose of no greater than 90 mg/m2 twice per cycle, for a maximum of four cycles.

Note: Indications marked with * are unapproved indications.

BUSULFAN - PCT - Retail pharmacy-Specialist

.25 100	✓ Myleran
.59 1	✓ DBL Carboplatin
.20	 Carboplatin Ebewe
.50	✓ Carbaccord
.10 1 mg	✓ Baxter
.00 1	✓ BiCNU
	✓ BiCNU S29 S29
.00 100 mg OP	✓ Baxter
.06 25	✓ Leukeran FC
.00 1	 Cisplatin Ebewe
.00 1	Cisplatin Ebewe
.66	DBL Cisplatin
.31 1 mg	✓ Baxter
.00 50	✓ Cyclonex
.65 1	✓ Endoxan
.80 6	✓ Cytoxan
.25 1	✓ Endoxan
.04 1 mg	✓ Baxter
.00 1	✓ Holoxan
.00 1	✓ Holoxan
.10 1 mg	✓ Baxter
	1.59 1 1

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Generic
OMUSTINE - PCT - Retail pharmacy-Specialist				
Cap 10 mg	132.59	20	1	CeeNU
Cap 40 mg	399.15	20	✓	CeeNU
MELPHALAN				
Tab 2 mg - PCT - Retail pharmacy-Specialist	40.70	25	1	Alkeran
Inj 50 mg - PCT only - Specialist	48.25	1	1	Megval S29
			1	Melpha
	67.80		✓	Alkeran
			1	Alkeran S29 S29
Alkeran S29 S29 Inj 50 mg to be delisted 1 August 2024)				
DXALIPLATIN - PCT only - Specialist				
Inj 100 mg vial	25.01	1	•	Oxaliplatin Actavis 100
	110.00		1	Oxaliplatin Ebewe
Inj 5 mg per ml, 20 ml vial	33.35	1	1	Alchemy Oxaliplatin
	46.32		1	Oxaliplatin Accord
Inj 1 mg for ECP	0.35	1 mg	✓	Baxter
HIOTEPA - PCT only - Specialist				
Inj 15 mg vial	CBS	1	1	Bedford S29
, •			1	Max Health \$29
			1	THIO-TEPA \$29
	398.00			Tepadina
Inj 100 mg vial	CBS	1	1	Max Health \$29
, ,	1,800.00		1	Tepadina

Antimetabolites

		PCT only – Specialist – Special Authority see SA2141 below	AZACITIDINE - PCT only -
✓ Azacitidine Dr	1	al75.06	Inj 100 mg vial
Reddy's ✓ Baxter	1 mg	CP	Inj 1 mg for ECP

⇒SA2141 Special Authority for Subsidy

Initial application only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has International Prognostic Scoring System (IPSS) intermediate-2 or high risk myelodysplastic syndrome: or
 - 1.2 The patient has chronic myelomonocytic leukaemia (10%-29% marrow blasts without myeloproliferative disorder);
 - 1.3 The patient has acute myeloid leukaemia with 20-30% blasts and multi-lineage dysplasia, according to World Health Organisation Classification (WHO); and
- 2 The patient has performance status (WHO/ECOG) grade 0-2; and
- 3 The patient has an estimated life expectancy of at least 3 months.

Renewal only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

	Subsidy		Fully	
	(Manufacturer's Price \$	e) Per	Subsidised	I Generic Manufacturer
CALCIUM FOLINATE	•			
Tab 15 mg - PCT - Retail pharmacy-Specialist	135.33	10	✓	DBL Leucovorin Calcium
Inj 3 mg per ml, 1 ml - PCT - Retail pharmacy-Specialist	17.10	5	1	Hospira
Inj 10 mg per ml, 5 ml vial - PCT - Retail pharmacy-Speciali		1	•	Calcium Folinate Sandoz
			•	Calcium Folinate Sandoz S29 S29
	36.48	5	1	Eurofolic S29
Inj 50 mg - PCT - Retail pharmacy-Specialist		10		Leucovorin
.,				Pharmacia S29
Inj 10 mg per ml, 10 ml vial - PCT only - Specialist	9.49	1	✓	Calcium Folinate Sandoz
	47.45	5	1	Eurofolic S29
Inj 100 mg - PCT only - Specialist	7.33	1	•	Calcium Folinate Ebewe
	94.90	10	•	Leucovorin Pharmacia \$29
Inj 300 mg - PCT only - Specialist	22.51	1	•	Calcium Folinate Ebewe
	25.14		✓	Leucovorin DBL S29
Inj 10 mg per ml, 35 ml vial – PCT only – Specialist	25.14	1	•	Calcium Folinate Sandoz
			•	Calcium Folinate Sandoz S29 S29
Inj 1 g - PCT only - Specialist	67.51	1	•	Calcium Folinate Ebewe
Inj 10 mg per ml, 100 ml vial - PCT only - Specialist	72.00	1	•	Calcium Folinate Sandoz
Inj 1 mg for ECP - PCT only - Specialist	0.06	1 mg	1	Baxter
CAPECITABINE - Retail pharmacy-Specialist				
Tab 150 mg	9.80	60	✓	Capecitabine Viatris
Tab 500 mg	46.50	120	1	Capecitabine Viatris
CLADRIBINE - PCT only - Specialist				
Inj 2 mg per ml, 5 ml	749.96	1	•	Litak S29
Inj 1 mg per ml, 10 ml	749.96	1	1	Leustatin
Inj 10 mg for ECP	749.96 1	10 mg (OP 🗸	Baxter
CYTARABINE				
Inj 20 mg per ml, 5 ml vial – PCT – Retail pharmacy-Speciali Inj 100 mg per ml, 20 ml vial – PCT – Retail	st472.00	5	•	Pfizer
pharmacy-Specialist	48.80	1		Cytarabine DBL Pfizer
				Pfizer S29 S29
Inj 1 mg for ECP - PCT only - Specialist	0.29	10 mc	_	Baxter
Inj 100 mg intrathecal syringe for ECP - PCT only - Speciali		00 mg	,	Baxter

	Subsidy		Fully	Brand or
	(Manufacturer's F		sidised	Generic
	\$	Per		Manufacturer
FLUDARABINE PHOSPHATE				
Tab 10 mg - PCT - Retail pharmacy-Specialist	412.00	20	✓	Fludara Oral
Inj 50 mg vial - PCT only - Specialist	126.80	1	✓	Fludarabine
				Sagent S29
	634.00	5	1	Fludarabine Ebewe
Inj 50 mg for ECP - PCT only - Specialist	126.80	50 mg OP	✓	Baxter
FLUOROURACIL		J		
Inj 50 mg per ml, 20 ml vial – PCT only – Specialist	10.51	1	1	Fluorouracil Accord
Inj 50 mg per ml, 50 ml vial – PCT only – Specialist		i		Fluorouracil Accord
Inj 50 mg per ml, 100 ml vial – PCT only – Specialist		i		Fluorouracil Accord
Inj 1 mg for ECP - PCT only - Specialist		100 mg		Baxter
GEMCITABINE HYDROCHLORIDE - PCT only - Specialist		3		
,				
Inj 43.3 mg per ml (equivalent to 38 mg per ml gemcitabine), 26.3 ml vial		1	1	DBL Gemcitabine
		1		Gemcitabine Ebewe
Inj 1 g Inj 1 mg for ECP		1 mg		Baxter
, ,	0.02	ring	•	Daxiei
RINOTECAN HYDROCHLORIDE – PCT only – Specialist	50.55		,	
Inj 20 mg per ml, 5 ml vial		1		Accord
	71.44		•	Irinotecan Actavis 100
	100.00		1	Irinotecan-Rex
Inj 1 mg for ECP	0.54	1 mg	1	Baxter
/ERCAPTOPURINE		ŭ		
Tab 50 mg - PCT - Retail pharmacy-Specialist	25 90	25	1	Puri-nethol
Oral suspension 20 mg per ml – Retail pharmacy-Specialist		20	• !	un nomor
Special Authority see SA1725 below		100 ml OP	1	Allmercap
——————————————————————————————————————		.00 1111 01	- '	oup

⇒SA1725 Special Authority for Subsidy

Initial application only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where the patient requires a total dose of less than one full 50 mg tablet per day.

Renewal only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where patient still requires a total dose of less than one full 50 mg tablet per day.

		Subsidy (Manufacturer's Price)		Fully Subsidised	Generic
=		\$	Per		Manufacturer
	THOTREXATE	0.00	00		Tuescata
*	Tab 2.5 mg - PCT - Retail pharmacy-Specialist		90 90		Trexate Trexate
*	Inj 2.5 mg per ml, 2 ml – PCT – Retail pharmacy-Specialist		5		Methotrexate DBL
•••	mj 2.0 mg por mi, 2 mi - 1 o 1 - Hotali pharmady opodialist		Ŭ		Methotrexate DBL
					S29 S29
*	Inj 7.5 mg prefilled syringe	14.61	1	1	Methotrexate
	., g F , g				Sandoz
*	Inj 10 mg prefilled syringe	14.66	1	1	Methotrexate
	, 31 , 3				Sandoz
*	Inj 15 mg prefilled syringe	14.77	1	1	Methotrexate
	, , , ,				Sandoz
*	Inj 20 mg prefilled syringe	14.88	1	✓	Methotrexate
					Sandoz
*	Inj 25 mg prefilled syringe	14.99	1	1	Methotrexate
					Sandoz
*	Inj 30 mg prefilled syringe	15.09	1	✓	Methotrexate
					Sandoz
*	Inj 25 mg per ml, 2 ml vial - PCT - Retail pharmacy-Specialis	st30.00	5	1	Methotrexate DBL
				_	Onco-Vial
*	Inj 25 mg per ml, 20 ml vial – PCT – Retail pharmacy-Special	ist45.00	1	/	DBL Methotrexate
				_	Onco-Vial
*	Inj 100 mg per ml, 10 ml – PCT – Retail pharmacy-Specialist	25.00	1	•	Methotrexate Ebewe
*	Inj 100 mg per ml, 50 ml vial – PCT – Retail	07.00		,	Mathatana Ebana
	pharmacy-Specialist		1		Methotrexate Ebewe
*	Inj 1 mg for ECP – PCT only – Specialist		1 mg		Baxter Baxter
*	Inj 5 mg intrathecal syringe for ECP – PCT only – Specialist		mg O	F •	Davici
ΡE	METREXED - PCT only - Specialist - Special Authority see S		4		luna Damatuayad
	Inj 100 mg vial		1		Juno Pemetrexed Juno Pemetrexed
	Inj 500 mg vial		1 mg		Baxter
	Inj 1 mg for ECP	0.00	i ilig	•	Daviel

⇒SA1679 Special Authority for Subsidy

Initial application — (mesothelioma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with mesothelioma; and
- 2 Pemetrexed to be administered at a dose of 500 mg/m² every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cycles.

Renewal — (mesothelioma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment; and
- 3 Pemetrexed to be administered at a dose of 500mg/m² every 21 days for a maximum of 6 cycles.

Initial application — (non-small cell lung carcinoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria: Both:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	(Manufacturer's Price) S		Generic	
\$	Per	✓	Manufacturer	

continued...

- 1 Patient has locally advanced or metastatic non-squamous non-small cell lung carcinoma; and
- 2 Fither:
 - 2.1 Both:
 - 2.1.1 Patient has chemotherapy-naïve disease; and
 - 2.1.2 Permetrexed is to be administered at a dose of 500 mg/m² every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cycles; or
 - 2.2 All of the following:
 - 2.2.1 Patient has had first-line treatment with platinum based chemotherapy; and
 - 2.2.2 Patient has not received prior funded treatment with pemetrexed; and
 - 2.2.3 Pemetrexed is to be administered at a dose of 500 mg/m² every 21 days for a maximum of 6 cycles.

Renewal — (non-small cell lung carcinoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment; and
- 3 Pemetrexed is to be administered at a dose of 500mg/m² every 21 days.

THIOGUANINE	PCT – Retail	pharmacy-Special	ist
T-1-40			

Tab 40 mg126.31 25 ✓ Lanvis

Other Cytotoxic Agents

AMSACRINE - PCT only - Specialist			
Inj 50 mg per ml, 1.5 ml ampoule	1,500.00	6	✓ Amsidine S29
	4,736.00		✓ Amsidine S29
Inj 75 mg	1,250.00	5	✓ AmsaLyo S29
ANAGRELIDE HYDROCHLORIDE - PCT - Retail pharmacy-	Specialist		
Cap 0.5 mg	1,175.87	100	✓ Agrylin
ARSENIC TRIOXIDE - PCT only - Specialist			
Inj 1 mg per ml, 10 ml vial	4,817.00	10	✓ Phenasen
Inj 10 mg for ECP	481.70	10 mg OP	✓ Baxter
BLEOMYCIN SULPHATE - PCT only - Specialist			
Inj 15,000 iu, vial	185.16	1	✓ DBL Bleomycin Sulfate
Inj 1,000 iu for ECP	14.32	1,000 iu	✓ Baxter
BORTEZOMIB - PCT only - Specialist - Special Authority se	e SA1889 below		
Inj 3.5 mg vial	74.93	1	✓ DBL Bortezomib
Inj 1 mg for ECP	22.26	1 mg	✓ Baxter

⇒SA1889 Special Authority for Subsidy

Initial application — (multiple myeloma/amyloidosis) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Fither:

- 1 The patient has symptomatic multiple myeloma; or
- 2 The patient has symptomatic systemic AL amyloidosis *.

Note: Indications marked with * are unapproved indications.

	Subsidy	Duis -\		Fully	
	(Manufacturer's \$	Price)	Per	Subsidised	
ACARBAZINE - PCT only - Specialist	*		-		
Inj 200 mg vial	79 11		1	1	DBL Dacarbazine
iiij 200 iiig vidi	580.60		10	_	Dacarbazine
	300.00		10	•	APP S29
Inj 200 mg for ECP	72 11	200	mg	OP 🗸	Baxter
Dacarbazine APP S29 Inj 200 mg vial to be delisted 1 August		200	ilig	01 •	Daxiei
, ,	2024)				
ACTINOMYCIN [ACTINOMYCIN D] – PCT only – Specialist	055.00		4	./	Coomonon
Inj 0.5 mg vial		0.5	1 mg (Cosmegen Baxter
Inj 0.5 mg for ECP	255.00	0.5	ilig (JP V	Daxlei
AUNORUBICIN – PCT only – Specialist					
Inj 2 mg per ml, 10 ml			1		Pfizer
Inj 20 mg vial	1,495.00		10	•	Daunorubicin
					Zentiva S29
Inj 20 mg for ECP	171.93	20	mg (OP 🗸	Baxter
OCETAXEL - PCT only - Specialist					
Inj 20 mg	48.75		1	✓	Docetaxel Sandoz
Inj 10 mg per ml, 8 ml vial	24.91		1	✓	DBL Docetaxel
Inj 20 mg per ml, 4 ml vial	26.95		1	✓	Docetaxel
					Accord S29
Inj 80 mg	195.00		1	✓	Docetaxel Sandoz
Inj 1 mg for ECP	0.35		1 mg	✓	Baxter
OXORUBICIN HYDROCHLORIDE - PCT only - Specialist			•		
Inj 2 mg per ml, 5 ml vial	10.00		1	/	Doxorubicin Ebewe
Inj 2 mg per ml, 25 ml vial			1	_	Doxorubicin Ebewe
.,,	17.00				Arrow-Doxorubicin
Inj 2 mg per ml, 50 ml vial	23.00		1	1	Doxorubicin Ebewe
Inj 2 mg per ml, 100 ml vial			1	✓	Arrow-Doxorubicin
	69.99			/	Accord \$29
					Doxorubicin Ebewe
Inj 1 mg for ECP	0.35		1 mg	1	Baxter
ccord S29 Inj 2 mg per ml, 100 ml vial to be delisted 1 Augu			J		
PIRUBICIN HYDROCHLORIDE - PCT only - Specialist	,				
Inj 2 mg per ml, 5 ml vial	25.00		1	1	Epirubicin Ebewe
Inj 2 mg per ml, 25 ml vial			i		Epirubicin Ebewe
Inj 2 mg per ml, 100 ml vial			1		Epirubicin Ebewe
Inj 1 mg for ECP			1 mg	_	Baxter
ΓOPOSIDE			9		
Cap 50 mg - PCT - Retail pharmacy-Specialist	340 73		20	1	Vepesid
Cap 100 mg - PCT - Retail pharmacy-Specialist			10		Vepesid
Inj 20 mg per ml, 5 ml vial – PCT – Retail pharmacy-Specia			1		Rex Medical
Inj 1 mg for ECP – PCT only – Specialist			1 mg	_	Baxter
			9		
FOPOSIDE PHOSPHATE - PCT only - Specialist Inj 100 mg (of etoposide base)	40.00		1	./	Etopophos
Inj 1 mg (of etoposide base) for ECP				_	Baxter
			1 mg	•	Dantei
YDROXYUREA [HYDROXYCARBAMIDE] – PCT – Retail pha		I	400		D
Cap 500 mg			100	•	<u>Devatis</u>
RUTINIB - Special Authority see SA2168 on the next page -		/			
			~~		
Tab 140 mg Tab 420 mg			30 30		Imbruvica Imbruvica

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

★Three months or six months, as applicable, dispensed all-at-once

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	/	Manufacturer	

⇒SA2168 Special Authority for Subsidy

Initial application — (chronic lymphocytic leukaemia (CLL)) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic lymphocytic leukaemia (CLL) requiring therapy; and
- 2 Patient has not previously received funded ibrutinib: and
- 3 Ibrutinib is to be used as monotherapy; and
- 4 Any of the following:
 - 4.1 Both:
 - 4.1.1 There is documentation confirming that patient has 17p deletion or TP53 mutation; and
 - 4.1.2 Patient has experienced intolerable side effects with venetoclax monotherapy; or
 - 4.2 All of the following:
 - 4.2.1 Patient has received at least one prior immunochemotherapy for CLL; and
 - 4.2.2 Patient's CLL has relapsed within 36 months of previous treatment; and
 - 4.2.3 Patient has experienced intolerable side effects with venetoclax in combination with rituximab regimen; or
 - 4.3 Patient's CLL is refractory to or has relapsed within 36 months of a venetoclax regimen.

Renewal — (chronic lymphocytic leukaemia (CLL)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 No evidence of clinical disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL) and B-cell prolymphocytic leukaemia (B-PLL)*. Indications marked with * are Unapproved indications.

IDARUBICIN HYDROCHLORIDE

Inj 10 mg vial - PCT only - Specialist	233.64	1	Zavedos
Inj 1 mg for ECP - PCT only - Specialist	25.77	1 mg	✓ Baxter
LENALIDOMIDE – Retail pharmacy-Specialist – Special A Wastage claimable	uthority see SA2047 be	low	
Cap 5 mg	5.122.76	28	✓ Revlimid
Cap 10 mg	·	21	✓ Revlimid
	6,207.00	28	Revlimid
Cap 15 mg	5,429.39	21	Revlimid

⇒SA2047 Special Authority for Subsidy

Initial application — (Relapsed/refractory disease) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has relapsed or refractory multiple myeloma with progressive disease; and
- 2 Patient has not previously been treated with lenalidomide; and
- 3 Fither:
 - 3.1 Lenalidomide to be used as third line* treatment for multiple myeloma; or
 - 3.2 Both:
 - 3.2.1 Lenalidomide to be used as second line treatment for multiple myeloma; and
 - 3.2.2 The patient has experienced severe (grade 3 or higher), dose limiting, peripheral neuropathy with either bortezomib or thalidomide that precludes further treatment with either of these treatments; and

7.239.18

28

21

continued...

✓ Zavedos

✓ Revlimid

✓ Revlimid

Subsidy		Fully	Brand or	
(Manufacturer's Price		Subsidised	Generic	
\$	Per	/	Manufacturer	

continued...

4 Lenalidomide to be administered at a maximum dose of 25 mg/day in combination with dexamethasone.

Initial application — (Maintenance following first-line autologous stem cell transplant (SCT)) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has newly diagnosed symptomatic multiple myeloma and has undergone first-line treatment that included an autologous stem cell transplantation; and
- 2 Patient has at least a stable disease response in the first 100 days after transplantation; and
- 3 Lenalidomide maintenance is to be commenced within 6 months of transplantation; and
- 4 Lenalidomide to be administered at a maximum dose of 15 mg/day.

Renewal — (Relapsed/refractory disease) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

Renewal — (Maintenance following first line autologous SCT) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

Note: Indication marked with * is an unapproved indication. A line of treatment is considered to comprise either: a) a known therapeutic chemotherapy regimen and supportive treatments or b) a transplant induction chemotherapy regimen, stem cell transplantation and supportive treatments. Prescriptions must be written by a registered prescriber in the lenalidomide risk management programme operated by the supplier.

MESNA

Tab 400 mg - PCT - Retail pharmacy-Specialist	314.00	50	Uromitexan
Tab 600 mg - PCT - Retail pharmacy-Specialist	448.50	50	✓ Uromitexan
Inj 100 mg per ml, 4 ml ampoule - PCT only - Specialist	177.45	15	Uromitexan
Inj 100 mg per ml, 10 ml ampoule - PCT only - Specialis	st407.40	15	Uromitexan
Inj 1 mg for ECP - PCT only - Specialist	2.96	100 mg	✓ Baxter
MITOMYCIN C - PCT only - Specialist			
Inj 5 mg vial	641.70	1	✓ Accord S29
Inj 20 mg vial	1,250.00	1	✓ Omegapharm \$29
, ,	,		✓ Teva
Inj 1 mg for ECP	269.85	1 mg	✓ Baxter
MITOZANTRONE - PCT only - Specialist			
Inj 2 mg per ml, 10 ml vial	97.50	1	✓ Mitozantrone Ebewe
Inj 1 mg for ECP	5.51	1 mg	✓ Baxter
NIRAPARIB - Special Authority see SA2325 below - Retail p	oharmacy		
Wastage claimable	•		
Cap 100 mg	8,929.84	56	✓ Zejula
•	13,393.50	84	✓ Zejula

⇒SA2325 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1 Patient has advanced high-grade serous* epithelial ovarian, fallopian tube, or primary peritoneal cancer; and
- 2 Patient has received at least one line** of treatment with platinum-based chemotherapy; and
- 3 Patient has experienced a partial or complete response to the preceding treatment with platinum-based chemotherapy;
- 4 Patient has not previously received funded treatment with a PARP inhibitor; and
- 5 Either:
 - 5.1 Treatment will be commenced within 12 weeks of the patient's last dose of the preceding platinum-based regimen;
 or
 - 5.2 Patient commenced treatment with niraparib prior to 1 May 2024; and
- 6 Treatment to be administered as maintenance treatment; and
- 7 Treatment not to be administered in combination with other chemotherapy.

Renewal from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 No evidence of progressive disease; and
- 2 Treatment to be administered as maintenance treatment; and
- 3 Treatment not to be administered in combination with other chemotherapy; and
- 4 Either:
 - 4.1 Treatment with niraparib to cease after a total duration of 36 months from commencement; or
 - 4.2 Treatment with niraparib is being used in the second-line or later maintenance setting.

Notes: * "high-grade serous" includes tumours with high-grade serous features or a high-grade serous component.

**A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments

OLAPARIB - Retail pharmacy-Specialist - Special Authority see SA2163 below

Tab 100 mg3,7	'01.00 5	o6 •	Lynparza
Tab 150 mg3,7	'01.00 5	56	Lynparza

⇒SA2163 Special Authority for Subsidy

Initial application — (Ovarian cancer) only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a high-grade serous* epithelial ovarian, fallopian tube, or primary peritoneal cancer; and
- 2 There is documentation confirming pathogenic germline BRCA1 or BRCA2 gene mutation; and
- 3 Either:
 - 3.1 All of the following:
 - 3.1.1 Patient has newly diagnosed, advanced disease; and
 - 3.1.2 Patient has received one line** of previous treatment with platinum-based chemotherapy; and
 - 3.1.3 Patient's disease must have experienced a partial or complete response to the first-line platinum-based regimen; or
 - 3.2 All of the following:
 - 3.2.1 Patient has received at least two lines** of previous treatment with platinum-based chemotherapy; and
 - 3.2.2 Patient has platinum sensitive disease defined as disease progression occurring at least 6 months after the last dose of the penultimate line** of platinum-based chemotherapy; and
 - 3.2.3 Patient's disease must have experienced a partial or complete response to treatment with the immediately preceding platinum-based regimen; and
 - 3.2.4 Patient has not previously received funded olaparib treatment; and
- 4 Treatment will be commenced within 12 weeks of the patient's last dose of the immediately preceding platinum-based regimen; and

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

- 5 Treatment to be administered as maintenance treatment; and
- 6 Treatment not to be administered in combination with other chemotherapy.

Renewal — (Ovarian cancer) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from treatment; and
- 2 Either:
 - 2.1 No evidence of progressive disease; or
 - 2.2 Evidence of residual (not progressive) disease and the patient would continue to benefit from treatment in the clinician's opinion; and
- 3 Treatment to be administered as maintenance treatment; and
- 4 Treatment not to be administered in combination with other chemotherapy; and
- 5 Either:
 - 5.1 Both:
 - 5.1.1 Patient has received one line** of previous treatment with platinum-based chemotherapy; and
 - 5.1.2 Documentation confirming that the patient has been informed and acknowledges that the funded treatment period of olaparib will not be continued beyond 2 years if the patient experiences a complete response to treatment and there is no radiological evidence of disease at 2 years; or
 - 5.2 Patient has received at least two lines** of previous treatment with platinum-based chemotherapy.

Notes: *Note "high-grade serous" includes tumours with high-grade serous features or a high-grade serous component.
**A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

PACLITAXEL – PCT only – Specialist			
Inj 30 mg	47.30	5	Paclitaxel Ebewe
Inj 6 mg per ml, 16.7 ml vial		1	✓ Anzatax
	24.00		Paclitaxel Ebewe
	91.67		Paclitaxel Actavis
Inj 150 mg	26.69	1	✓ Paclitaxel Ebewe
, ,	137.50		✓ Anzatax
			✓ Paclitaxel Actavis
Inj 6 mg per ml, 50 ml vial	37.89	1	✓ Anzatax
, ,	44.00		✓ Paclitaxel Ebewe
	275.00		✓ Paclitaxel Actavis
Inj 1 mg for ECP	0.17	1 mg	✓ Baxter
PEGASPARGASE - PCT only - Special Authority see S	A1979 below		
Inj 750 iu per ml, 5 ml vial	3,455.00	1	✓ Oncaspar LYO S29

⇒SA1979 Special Authority for Subsidy

Initial application — (Acute lymphoblastic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has newly diagnosed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

Initial application — (Lymphoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months where the patient has lymphoma requiring L-asparaginase containing protocols (e.g. SMII F)

Renewal — (Acute lymphoblastic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of

Subsidy	Fu	lly Brand or	
(Manufacturer's Price)	Subsidis	ed Generic	
\$	Per	 Manufacturer 	

continued...

a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

- 1 The patient has relapsed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

Inj 10 mg	CBS	1	✓ Nipent S29
PROCARBAZINE HYDROCHLORIDE - PCT - Retail pharmacy	-Specialist		
Cap 50 mg	980.00	50	✓ Natulan S29
TEMOZOLOMIDE - Special Authority see SA2275 below - Reta	il pharmacy		
Cap 5 mg	9.13	5	✓ Temaccord
Cap 20 mg	16.38	5	✓ Temaccord
	18.30		✓ Apo-Temozolomide
Cap 100 mg	35.98	5	✓ Temaccord
	40.20		✓ Apo-Temozolomide
Cap 140 mg	50.12	5	✓ Temaccord
Cap 180 mg	620.00	14	✓ Accord S29
Cap 250 mg	86.34	5	✓ Temaccord

(Accord \$29 Cap 180 mg to be delisted 1 August 2024)

⇒SA2275 Special Authority for Subsidy

Initial application — (gliomas) only from a relevant specialist. Approvals valid for 12 months where the patient has a glioma. Renewal — (gliomas) only from a relevant specialist. Approvals valid for 12 months where treatment remains appropriate and patient is benefitting from treatment.

Initial application — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with metastatic or unresectable well-differentiated neuroendocrine tumour*: and
- 2 Temozolomide is to be given in combination with capecitabine; and
- 3 Temozolomide is to be used in 28 day treatment cycles for a maximum of 5 days treatment per cycle at a maximum dose of 200 mg/m² per day; and
- 4 Temozolomide to be discontinued at disease progression.

Renewal — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Initial application — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 9 months where the patient has relapsed/refractory Ewing's sarcoma.

Renewal — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indication marked with a * is an unapproved indication. Temozolomide is not subsidised for the treatment of relapsed high grade glioma.

THALIDOMIDE	 Retail pharmacy-Specialist – Special Authority see SA1124 	on the next page	
Cap 50 mg		28	Thalomid
Cap 100 mg	g756.00	28	Thalomid

Subsidy		Fully	Brand or	
(Manufacturer's Pri	ce)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

⇒SA1124 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Fither:

- 1 The patient has multiple myeloma; or
- 2 The patient has systemic AL amyloidosis*.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where the patient has obtained a response from treatment during the initial approval period.

Notes: Prescription must be written by a registered prescriber in the thalidomide risk management programme operated by the supplier.

Maximum dose of 400 mg daily as monotherapy or in a combination therapy regimen.

Indication marked with * is an unapproved indication.

TRETINOIN

100 ✓ Vesanoid	479.50	Cap 10 mg - PCT - Retail pharmacy-Specialist
I	uthority see SA1868 below	VENETOCLAX - Retail pharmacy-Specialist - Specialist - Sp
42 OP ✓ Venclexta	1,771.86 42	Tab 14×10 mg, 7×50 mg, 21×100 mg
2 OP ✓ Venclexta		Tab 10 mg
7 OP ✓ Venclexta	239.44 7	Tab 50 mg
120 Venclexta	8.209.41	Tab 100 mg - Wastage claimable

⇒SA1868 Special Authority for Subsidy

Initial application — (relapsed/refractory chronic lymphocytic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic lymphocytic leukaemia requiring treatment; and
- 2 Patient has received at least one prior therapy for chronic lymphocytic leukaemia; and
- 3 Patient has not previously received funded venetoclax; and
- 4 The patient's disease has relapsed within 36 months of previous treatment; and
- 5 Venetoclax to be used in combination with six 28-day cycles of rituximab commencing after the 5-week dose titration schedule with venetoclax; and
- 6 Patient has an ECOG performance status of 0-2.

Renewal — (relapsed/refractory chronic lymphocytic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment; and
- 2 Venetoclax is to be discontinued after a maximum of 24 months of treatment following the titration schedule unless earlier discontinuation is required due to disease progression or unacceptable toxicity.

Initial application — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has previously untreated chronic lymphocytic leukaemia; and
- 2 There is documentation confirming that patient has 17p deletion by FISH testing or TP53 mutation by sequencing; and
- 3 Patient has an ECOG performance status of 0-2.

Renewal — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where the treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL)* and B-cell prolymphocytic leukaemia (B-PLL)*. Indications marked with * are Unapproved indications.

Subsidy		Fully	Brand or
(Manufacturer's Price		Subsidised	
<u> </u>	Per		Manufacturer
VINBLASTINE SULPHATE			
Inj 1 mg per ml, 10 ml vial - PCT - Retail pharmacy-Specialist270.37	5		Hospira
Inj 1 mg for ECP - PCT only - Specialist	1 mg	/	Baxter
VINCRISTINE SULPHATE			
Inj 1 mg per ml, 1 ml vial - PCT - Retail pharmacy-Specialist51.37	5	1	DBL Vincristine
			Sulfate
Inj 1 mg per ml, 2 ml vial - PCT - Retail pharmacy-Specialist 102.73	5	1	DBL Vincristine
			Sulfate
Inj 1 mg for ECP - PCT only - Specialist12.60	1 mg	· •	Baxter
VINORELBINE			
Cap 20 mg30.00	1	1	Vinorelbine Te Arai
Cap 30 mg40.00	1	1	Vinorelbine Te Arai
Cap 80 mg60.00	1	✓	Vinorelbine Te Arai
Inj 10 mg per ml, 1 ml vial - PCT only - Specialist12.00	1	1	Navelbine
42.00			Vinorelbine Ebewe
Inj 10 mg per ml, 5 ml vial – PCT only – Specialist56.00	1	•	Navelbine
168.00			Navelbine S29 S29
210.00		✓	Vinorelbine Ebewe
328.65		✓	Sagent S29
Inj 1 mg for ECP - PCT only - Specialist3.80	1 mg	/	Baxter
(Navelbine Inj 10 mg per ml, 1 ml vial to be delisted 1 October 2024)			
(Navelbine Inj 10 mg per ml, 5 ml vial to be delisted 1 October 2024)			
(Sagent S29 Inj 10 mg per ml, 5 ml vial to be delisted 1 August 2024)			

Protein-tyrosine Kinase Inhibitors

ALECTINIB - Retail pharmacy-Specialist - Special Authority see SA1870 below Wastage claimable 224 ✓ Alecensa

⇒SA1870 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-small cell lung cancer; and
- 2 There is documentation confirming that the patient has an ALK tyrosine kinase gene rearrangement using an appropriate ALK test: and
- 3 Patient has an ECOG performance score of 0-2.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of progressive disease according to RECIST criteria; and
- 2 The patient is benefitting from and tolerating treatment.

DASATINIB - Special Authority see SA1805 on the next page - Retail pharmacy

vvastage ciaimable			
Tab 20 mg	3,774.06	60	✓ Sprycel
Tab 50 mg	6,214.20	60	✓ Sprycel
Tab 70 mg	7,692.58	60	✓ Sprycel

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$
Per ✓ Manufacturer

⇒SA1805 Special Authority for Subsidy

Initial application only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 The patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis or accelerated phase; and
 - 1.2 Maximum dose of 140 mg/day; or
- 2 Both:
 - 2.1 The patient has a diagnosis of Philadelphia chromosome-positive acute lymphoid leukaemia (Ph+ ALL); and
 - 2.2 Maximum dose of 140 mg/day; or
- 3 All of the following:
 - 3.1 The patient has a diagnosis of CML in chronic phase; and
 - 3.2 Maximum dose of 100 mg/day; and
 - 3.3 Any of the following:
 - 3.3.1 Patient has documented treatment failure* with imatinib; or
 - 3.3.2 Patient has experienced treatment-limiting toxicity with imatinib precluding further treatment with imatinib; or
 - 3.3.3 Patient has high-risk chronic-phase CML defined by the Sokal or EURO scoring system; or
 - 3.3.4 Patients is enrolled in the KISS study** and requires dasatinib treatment according to the study protocol.

Renewal only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Lack of treatment failure while on dasatinib*; and
- 2 Dasatinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum dasatinib dose of 140 mg/day for accelerated or blast phase CML and Ph+ ALL, and 100 mg/day for chronic phase CML.

Note: *treatment failure for CML as defined by Leukaemia Net Guidelines. **Kinase-Inhibition Study with Sprycel Start-up https://www.cancertrialsnz.ac.nz/kiss/

ERLOTINIB – Retail pharmacy-Specialist – Special Author	rity see SA2115 below		
Tab 100 mg	280.84	30	✓ Alchemy
Tab 150 mg	484.24	30	✓ Alchemy

⇒SA2115 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 There is documentation confirming that the disease expresses activating mutations of EGFR tyrosine kinase; and
- 3 Either
 - 3.1 Patient is treatment naive; or
 - 3.2 Both:
 - 3.2.1 The patient has discontinued gefitinib due to intolerance; and
 - 3.2.2 The cancer did not progress while on gefitinib; and
- 4 Erlotinib is to be given for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

Renewal — (pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subs	sidised	Generic	
\$	Per	•	Manufacturer	

continued...

All of the following:

- 1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 2 Erlotinib to be discontinued at progression; and
- 3 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

GEFITINIB - Retail pharmacy-Specialist - Special Authority see SA2116 below

Tab 250 mg918.00 30 ✓ Iressa

⇒SA2116 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 Either:
 - 2.1 Patient is treatment naive; or
 - 2.2 Both:
 - 2.2.1 The patient has discontinued erlotinib due to intolerance; and
 - 2.2.2 The cancer did not progress whilst on erlotinib; and
- 3 There is documentation confirming that disease expresses activating mutations of EGFR tyrosine kinase; and
- 4 Gefitinib is to be given for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

Renewal — (pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 2 Gefitinib to be discontinued at progression; and
- 3 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

IMATINIB MESILATE

* Cap 100 mg * Cap 400 mg		60 30	✓ Imatinib-Rex ✓ Imatinib-Rex
NILOTINIB – Special Authority see SA2301 below Wastage claimable	- Retail pharmacy		
Cap 150 mg	4.680.00	120	✓ Tasigna
Cap 200 mg	·	120	✓ Tasigna

⇒SA2301 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis, high risk chronic phase, or in chronic phase; and
- 2 Either:
 - 2.1 Patient has documented CML treatment failure* with a tyrosine kinase inhibitor (TKI); or
 - 2.2 Patient has experienced treatment limiting toxicity with a tyrosine kinase inhibitor (TKI) precluding further treatment; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

Note: *treatment failure as defined by Leukaemia Net Guidelines.

	Subsidy	F	ully	Brand or
(Ma	anufacturer's Price)	Subsid	ised	Generic
	\$	Per	1	Manufacturer

continued...

Renewal only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Lack of treatment failure while on nilotinib as defined by Leukaemia Net Guidelines; and
- 2 Nilotinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

PALBOCICLIB - Retail pharmacy-Specialist - Special Authority see SA1894 below

wastage ciaimable			
Tab 75 mg	4,000.00	21	Ibrance
Tab 100 mg	4,000.00	21	✓ Ibrance
Tab 125 mg	4,000.00	21	✓ Ibrance

⇒SA1894 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a Medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has unresectable locally advanced or metastatic breast cancer; and
- 2 There is documentation confirming disease is hormone-receptor positive and HER2-negative; and
- 3 Patient has an ECOG performance score of 0-2; and
- 4 Either:

second or subsequent line setting

- 4.1 Disease has relapsed or progressed during prior endocrine therapy; or
- 4.2 Both:

first line setting

- 4.2.1 Patient is amenorrhoeic, either naturally or induced, with endocrine levels consistent with a postmenopausal state; and
- 4.2.2 Either:
 - 4.2.2.1 Patient has not received prior systemic treatment for metastatic disease; or
 - 4.2.2.2 All of the following:
 - 4.2.2.2.1 Patient commenced treatment with palbociclib in combination with an endocrine agent prior to 1 April 2020; and
 - 4.2.2.2.2 Patient has not received prior systemic endocrine treatment for metastatic disease; and
 - 4.2.2.2.3 There is no evidence of progressive disease; and
- 5 Treatment must be used in combination with an endocrine partner.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a Medical oncologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment must be used in combination with an endocrine partner; and
- 2 No evidence of progressive disease; and
- 3 The treatment remains appropriate and the patient is benefitting from treatment.

PAZOPANIB - Special Authority see SA1190 below - Retail pharmacy

Tab 200 mg	1,334.70	30	✓ Votrient
Tab 400 mg	2,669.40	30	✓ Votrient

⇒SA1190 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist.

Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

(1	Subsidy Manufacturer's Price)	F Subsid	ully	Brand or Generic
· ·	\$	Per	1	Manufacturer

continued...

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
 - 2.1 The patient is treatment naive; or
 - 2.2 The patient has only received prior cytokine treatment; or
 - 2.3 Both:
 - 2.3.1 The patient has discontinued sunitinib within 3 months of starting treatment due to intolerance; and
 - 2.3.2 The cancer did not progress whilst on sunitinib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and The patient has intermediate or poor prognosis defined as:
- 5 Any of the following:
 - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
 - 5.2 Haemoglobin level < lower limit of normal; or
 - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
 - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
 - 5.5 Karnofsky performance score of less than or equal to 70; or
 - 5.6 2 or more sites of organ metastasis; and
- 6 Pazopanib to be used for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Pazopanib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6.

RUXOLITINIB - Special Authority see SA1890 below - Retail pharmacy

Wastage claimable			
Tab 5 mg	2,500.00	56	Jakavi
Tab 10mg	5,000.00	56	Jakavi
Tab 15 mg	5,000.00	56	Jakavi
Tab 20 mg	5,000.00	56	Jakavi

⇒SA1890 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has primary myelofibrosis or post-polycythemia vera myelofibrosis or post-essential thrombocythemia myelofibrosis; and
- 2 Either:
 - 2.1 A classification of risk of intermediate-2 or high-risk myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS; or
 - 2.2 Both:
 - 2.2.1 A classification of risk of intermediate-1 myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS; and
 - 2.2.2 Patient has severe disease-related symptoms that are resistant, refractory or intolerant to available therapy; and

Subsidy (Manufacturer's Price)	Fully Subsidised	
 \$	Per 🗸	Manufacturer

continued...

3 A maximum dose of 20 mg twice daily is to be given.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 A maximum dose of 20 mg twice daily is to be given.

SUNITINIB - Special Authority see SA2117 below - Retail pharmacy

Cap 12.5 mg	38 28	✓ Sunitinib Pfizer
Cap 25 mg	77 28	✓ Sunitinib Pfizer
Cap 50 mg	62 28	✓ Sunitinib Pfizer

⇒SA2117 Special Authority for Subsidy

Initial application — (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
 - 2.1 The patient is treatment naive; or
 - 2.2 The patient has only received prior cytokine treatment; or
 - 2.3 The patient has only received prior treatment with an investigational agent within the confines of a bona fide clinical trial which has Ethics Committee approval; or
 - 2.4 Both:
 - 2.4.1 The patient has discontinued pazopanib within 3 months of starting treatment due to intolerance; and
 - 2.4.2 The cancer did not progress whilst on pazopanib: and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and The patient has intermediate or poor prognosis defined as:
- 5 Any of the following:
 - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
 - 5.2 Haemoglobin level < lower limit of normal: or
 - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
 - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
 - 5.5 Karnofsky performance score of less than or equal to 70; or
 - 5.6 2 or more sites of organ metastasis; and
- 6 Sunitinib to be used for a maximum of 2 cycles.

Initial application — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal tumour (GIST); and
- 2 Either:
 - 2.1 The patient's disease has progressed following treatment with imatinib; or
 - 2.2 The patient has documented treatment-limiting intolerance, or toxicity to, imatinib.

Renewal — (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer \$

continued...

Notes: Sunitinib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6

Renewal — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: Both:

The patient has responded to treatment or has stable disease as determined by Choi's modified CT response evaluation criteria as follows:

- 1 Any of the following:
 - 1.1 The patient has had a complete response (disappearance of all lesions and no new lesions); or
 - 1.2 The patient has had a partial response (a decrease in size of 10% or more or decrease in tumour density in Hounsfield Units (HU) of 15% or more on CT and no new lesions and no obvious progression of non measurable disease); or
 - 1.3 The patient has stable disease (does not meet criteria the two above) and does not have progressive disease and no symptomatic deterioration attributed to tumour progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: It is recommended that response to treatment be assessed using Choi's modified CT response evaluation criteria (J Clin Oncol, 2007, 25:1753-1759). Progressive disease is defined as either: an increase in tumour size of 10% or more and not meeting criteria of partial response (PR) by tumour density (HU) on CT; or: new lesions, or new intratumoral nodules, or increase in the size of the existing intratumoral nodules.

Renewal — (GIST pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal (GIST); and
- 2 The patient is clinically benifiting from treatment and continued treatment remains appropriate; and
- 3 Sunitinib is to be discontinued at progression; and
- 4 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

Endocrine Therapy

For GnRH ANALOGUES - refer to HORMONE PREPARATIONS, Trophic Hormones, page 89

ABIRATERONE ACETATE - Retail pharmacy-Specialist - Special Authority see \$A2118 below

Wastage claimable

Tab 250 mg4,276.19 120 Zytiga

⇒SA2118 Special Authority for Subsidy

Initial application only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has prostate cancer; and
- 2 Patient has metastases: and
- 3 Patient's disease is castration resistant; and
- 4 Either:
 - 4.1 All of the following:
 - 4.1.1 Patient is symptomatic; and
 - 4.1.2 Patient has disease progression (rising serum PSA) after second line anti-androgen therapy; and
 - 4.1.3 Patient has ECOG performance score of 0-1; and
 - 4.1.4 Patient has not had prior treatment with taxane chemotherapy; or

Subsidy		Fully	Brand or	
(Manufacturer's Price)) S	ubsidised	Generic	
	Per	1	Manufacturer	

continued...

- 4.2 All of the following:
 - 4.2.1 Patient's disease has progressed following prior chemotherapy containing a taxane; and
 - 4.2.2 Patient has ECOG performance score of 0-2; and
 - 4.2.3 Patient has not had prior treatment with abiraterone.

Renewal — (abiraterone acetate) only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Significant decrease in serum PSA from baseline; and
- 2 No evidence of clinical disease progression; and
- 3 No initiation of taxane chemotherapy with abiraterone; and
- 4 The treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 2 Abiraterone acetate to be discontinued at progression; and
- 3 No initiation of taxane chemotherapy with abiraterone; and
- 4 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

BICALUTAMIDE

Tab 50 mg	4.18	28	✓ Apo- Bicalutamide ©29
			✓ Binarex
FLUTAMIDE			
Tab 250 mg	107.55	90	✓ Prostacur S29
·	119.50	100	✓ Flutamin
FULVESTRANT - Retail pharmacy-Specialist - Special Authority s	ee SA1895 be	low	
Inj 50 mg per ml, 5 ml prefilled syringe	1,068.00	2	✓ Faslodex
- CA100E Chapiel Authority for Cubaidy			

SA1895 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has oestrogen-receptor positive locally advanced or metastatic breast cancer; and
- 2 Patient has disease progression following prior treatment with an aromatase inhibitor or tamoxifen for their locally advanced or metastatic disease: and
- 3 Treatment to be given at a dose of 500 mg monthly following loading doses; and
- 4 Treatment to be discontinued at disease progression.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains appropriate and patient is benefitting from treatment; and
- 2 Treatment to be given at a dose of 500 mg monthly; and
- 3 There is no evidence of disease progression.

	Subsidy		Fully	
	(Manufacturer's Price) \$	Per	Subsidised	Generic Manufacturer
OCTREOTIDE				
Inj 50 mcg per ml, 1 ml ampoule	27.58	5	✓	Max Health
			1	Octreotide GH S29
Inj 100 mcg per ml, 1 ml ampoule	32.71	5	✓	Max Health
			✓	Octreotide GH S29
			1	Sun Pharma S29
Inj 500 mcg per ml, 1 ml ampoule	113.10	5	1	Max Health
			1	Octreotide GH \$29
			/	Sun Pharma S29
OCTREOTIDE LONG-ACTING - Special Authority see SA2119	below – Retail pharm	acv		
Inj depot 10 mg prefilled syringe		1	✓	Octreotide Depot
, , , , , ,				Teva
	1,152.00		✓	Sandostatin LAR
Inj depot 20 mg prefilled syringe	647.03	1	1	Octreotide Depot
				Teva
	1,539.00		1	Sandostatin LAR
Inj depot 30 mg prefilled syringe	670.80	1	✓	Sandostatin LAR
· · · · · · · · · · · · · · · · · · ·	718.55		✓	Octreotide Depot
				Teva

⇒SA2119 Special Authority for Subsidy

Initial application — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 The patient has nausea* and vomiting* due to malignant bowel obstruction*; and
- 2 Treatment with antiemetics, rehydration, antimuscarinic agents, corticosteroids and analgesics for at least 48 hours has failed: and
- 3 Octreotide to be given at a maximum dose 1500 mcg daily for up to 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 3 months where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 The patient has acromegaly; and
- 2 Any of the following:
 - 2.1 Treatment with surgery, radiotherapy and a dopamine agonist has failed; or
 - 2.2 Treatment with octreotide is for an interim period while awaiting the effects of radiotherapy and a dopamine agonist has failed: or
 - 2.3 The patient is unwilling, or unable, to undergo surgery and/or radiotherapy.

Renewal — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 IGF1 levels have decreased since starting octreotide; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: In patients with Acromegaly octreotide treatment should be discontinued if IGF1 levels have not decreased after 3 months treatment. In patients treated with radiotherapy octreotide treatment should be withdrawn every 2 years, for 1 month, for assessment of remission. Octreotide treatment should be stopped where there is biochemical evidence of remission (normal IGF1 levels) following octreotide treatment withdrawal for at least 4 weeks

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sul	osidised	Generic	
\$	Per	1	Manufacturer	

continued...

Renewal — (Acromegaly - pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has acromegaly: and
- 2 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 3 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

Initial application — (Other Indications) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 VIPomas and Glucagonomas for patients who are seriously ill in order to improve their clinical state prior to definitive surgery; or
- 2 Both:
 - 2.1 Gastrinoma: and
 - 2.2 Either:
 - 2.2.1 Patient has failed surgery; or
 - 2.2.2 Patient in metastatic disease after H2 antagonists (or proton pump inhibitors) have failed; or
- 3 Both:
 - 3.1 Insulinomas: and
 - 3.2 Surgery is contraindicated or has failed; or
- 4 For pre-operative control of hypoglycaemia and for maintenance therapy; or
- 5 Both:
 - 5.1 Carcinoid syndrome (diagnosed by tissue pathology and/or urinary 5HIAA analysis); and
 - 5.2 Disabling symptoms not controlled by maximal medical therapy.

Note: The use of octreotide in patients with fistulae, oesophageal varices, miscellaneous diarrhoea and hypotension will not be funded as a Special Authority item

Renewal — (Other Indications) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment. Initial application — (pre-operative acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

- All of the following:
 - 1 Patient has acromegaly; and
 - 2 Patient has a large pituitary tumour, greater than 10 mm at its widest; and
 - 3 Patient is scheduled to undergo pituitary surgery in the next six months.

Tah 20 mg 5.32

IAI	MOXIFEN CITRATE	
*	Tab 10 mg	15.00

	••	
Aromatase Inhibitors		
ANASTROZOLE	30	✓ Anatrole
EXEMESTANE	30	✓ Pfizer Exemestane
LETROZOLE * Tab 2.5 mg	30	✓ Latrola

✓ Tamoxifen Sandoz

Tamoxifen Sandoz

60

60

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

Immunosuppressants

Cytotoxic Immunosuppressants

AZATHIOPRINE

不	1ab 25 mg	00	▼ <u>Azamun</u>
*	Tab 50 mg8.10	100	✓ <u>Azamun</u>

MYCOPHENOLATE MOFETIL

TOO! TIENOLATE MOTETIE			
Tab 500 mg	35.90	50	Cellcept
Cap 250 mg	35.90	100	Cellcept
Powder for oral liq 1 g per 5 ml - Subsidy by endorsement18		ml OP	Cellcept

Mycophenolate powder for oral liquid is subsidised only for patients unable to swallow tablets and capsules, and when the prescription is endorsed accordingly.

Fusion Proteins

ETANERCEPT - Special Authority see SA2103 below - Retail pharmacy

Inj 25 mg	690.00	4	Enbrel
Inj 25 mg autoinjector		4	✓ Enbrel
Inj 50 mg autoinjector	1,050.00	4	✓ Enbrel
Ini 50 mg prefilled syringe	1.050.00	4	✓ Enbrel

⇒SA2103 Special Authority for Subsidy

Initial application — (adult-onset Still's disease) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Either:
 - 1.1.1 The patient has had an initial Special Authority approval for adalimumab for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a Health NZ Hospital; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or tocilizumab; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or tocilizumab such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal anti-inflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 The patient has a sustained improvement in inflammatory markers and functional status.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	Subsidised	Generic	
\$	Per	1	Manufacturer	

continued...

Initial application — (ankylosing spondylitis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for ankylosing spondylitis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for ankylosing spondylitis; or
- 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis present for more than six months; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.3 Patient has bilateral sacroiliitis demonstrated by plain radiographs, CT or MRI scan; and
 - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
 - 2.5 Either:
 - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
 - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender (see Notes); and
 - 2.6 A Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of initial application.

Average normal chest expansion corrected for age and gender:

18-24 years - Male: 7.0 cm; Female: 5.5 cm 25-34 years - Male: 7.5 cm; Female: 5.5 cm 35-44 years - Male: 6.5 cm; Female: 4.5 cm 45-54 years - Male: 6.0 cm; Female: 5.0 cm 55-64 years - Male: 5.5 cm; Female: 4.0 cm 65-74 years - Male: 3.0 cm; Female: 2.5 cm

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 3 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 4 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (polyarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist.

Subsidy		Fully	Brand or	
(Manufacturer's Pric	ce)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for polyarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
 - 2.3 Any of the following:
 - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose): or
 - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Renewal — (polyarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for oligoarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for oligoarticular course JIA: or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
 - 2.3 Any of the following:
 - 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsidised		Generic
\$	Per	1	Manufacturer

continued...

- 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose); or
- 2.3.3 High disease activity (cJADAS10 score greater than 4) after a 6-month trial of methotrexate.

Renewal — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Fither:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab or secukinumab for psoriatic arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab or secukinumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab or secukinumab to meet the renewal criteria for adalimumab or secukinumab for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
 - 2.4 Either:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints;
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

of the following

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and

Subsidy (Manufacturer's Price)	Sub	Fully sidised	Brand or Generic
 \$	Per	1	Manufacturer

continued...

- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Indications marked with * are unapproved indications.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

- All of the following:
 - 1 Patient has shown clinical improvement; and
 - 2 Patient continues to require treatment; and
 - 3 A maximum of 8 doses.

Initial application — (Arthritis - rheumatoid) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for rheumatoid arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects; or
 - 1.2.2 The patient has received insufficient benefit to meet the renewal criteria for rheumatoid arthritis; or
- 2 All of the following:
 - 2.1 Patient has had rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
 - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.3 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
 - 2.4 Patient has tried and not responded to at least three months of methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses unless contraindicated); and
 - 2.5 Either:
 - 2.5.1 Patient has tried and not responded to at least three months of methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 2.5.2 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with methotrexate; and
 - 2.6 Fither:
 - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
 - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	9	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Renewal — (Arthritis - rheumatoid) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (severe chronic plaque psoriasis) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe chronic plaque psoriasis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe chronic plaque psoriasis; or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
 - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
 - 2.3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
 - 2.4 The most recent PASI or DLQI assessment is no more than 1 month old at the time of application.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a dermatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a dermatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Fither:

Subsidy	Fully	Brand or	Т
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

- 2.1 Both:
 - 2.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 2.1.2 Either:
 - 2.1.2.1 Following each prior etanercept treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-treatment baseline value; or
 - 2.1.2.2 Following each prior etanercept treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline value; or
- 2.2 Both:
 - 2.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 2.2.2 Either:
 - 2.2.2.1 Following each prior etanercept treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 2.2.2.2 Following each prior etanercept treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Note: A treatment course is defined as a minimum of 12 weeks of etanercept treatment

Initial application — (undifferentiated spondyloarthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undifferentiated peripheral spondyloarthritis* with active peripheral joint arthritis in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day (or maximum tolerated dose); and
- 4 Patient has tried and not responded to at least three months of leflunomide at a dose of up to 20 mg daily (or maximum tolerated dose); and
- 5 Any of the following:
 - 5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application: or
 - 5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour measured no more than one month prior to the date of this application; or
 - 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Note: Indications marked with * are unapproved indications.

Renewal — (undifferentiated spondyloarthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg dose every 7 days.

Immune Modulators

ANTITHYMOCYTE GLOBULIN (EQUINE) - PCT only - Speciali	st		
Inj 50 mg per ml, 5 ml	2,774.48	5	✓ ATGAM
BACILLUS CALMETTE-GUERIN (BCG) VACCINE - PCT only -	Specialist		
Subsidised only for bladder cancer.			
Inj 2-8 × 100 million CFU	149.37	1	✓ OncoTICE
Inj 40 mg per ml, vial	176.90	3	✓ SII-Onco-BCG \$29

Monoclonal Antibodies

ADALIMUMAB (AMGEVITA) - Special Authority see SA2178	below - Retail pharma	acy	
Inj 20 mg per 0.4 ml prefilled syringe	190.00	1	✓ Amgevita
Inj 40 mg per 0.8 ml prefilled pen	375.00	2	✓ Amgevita
Inj 40 mg per 0.8 ml prefilled syringe	375.00	2	✓ Amgevita

⇒SA2178 Special Authority for Subsidy

Initial application — (Behcet's disease - severe) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient has severe Behcet's disease* that is significantly impacting the patient's quality of life; and
- 2 Either:
 - 2.1 The patient has severe ocular, neurological, and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s); or
 - 2.2 The patient has severe gastrointestinal, rheumatological, and/or mucocutaneous symptoms and has not responded adequately to two or more treatments appropriate for the particular symptom(s).

Note: Indications marked with * are unapproved indications.

Initial application — (Hidradenitis suppurativa) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has hidradenitis suppurativa Hurley Stage II or Hurley Stage III lesions in distinct anatomic areas; and
- 2 Patient has tried, but had an inadequate response to at least a 90 day trial of systemic antibiotics or has demonstrated intolerance to or has contraindications for systemic antibiotics; and
- 3 Patient has 3 or more active lesions; and
- 4 The patient has a DLQI of 10 or more and the assessment is no more than 1 month old at time of application.

Renewal — (Hidradenitis suppurativa) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline; and
- 2 The patient has a DLQI improvement of 4 or more from baseline.

Initial application — (Plaque psoriasis - severe chronic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

Subsidy (Manufacturer's Pric	ce)	Fully Subsidised	Brand or Generic	
\$	Per	✓	Manufacturer	

continued...

- 1 Both:
 - 1.1 Patient has had an initial Special Authority approval for etanercept for severe chronic plague psoriasis; and
 - 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects; or
 - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for etanercept for severe chronic plaque psoriasis; or
 - 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a PASI score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plague psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
 - 2.2 Patient has tried, but had an inadequate response to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
 - 2.3 A PASI assessment or DLQI assessment has been completed for at least the most recent prior treatment course but no longer than 1 month following cessation of each prior treatment course and is no more than 1 month old at the time of application.

Renewal — (Plague psoriasis - severe chronic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Both:
 - 1.1 Patient had "whole body" severe chronic plague psoriasis at the start of treatment; and
 - 1.2 Either:
 - 1.2.1 The patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre treatment baseline value; or
 - 1.2.2 The patient has a DLQI improvement of 5 or more, when compared with the pre-treatment baseline value; or
- 2 Both:
 - 2.1 Patient had severe chronic plague psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment: and
 - 2.2 Fither:
 - 2.2.1 The patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 2.2.2 The patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre treatment baseline value.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporin, azathioprine, or methotrexate) and has not received an adequate response.

Note: Indications marked with * are unapproved indications.

Initial application — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active Crohn's disease: and
- 2 Any of the following:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 2.1 Patient has a CDAI score of greater than or equal to 300, or HBI score of greater than or equal to 10; or
- 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
- 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection;
- 2.4 Patient has an ileostomy or colostomy and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Renewal — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 CDAI score has reduced by 100 points from the CDAI score, or HBI score has reduced by 3 points, from when the patient was initiated on adalimumab; or
- 2 CDAI score is 150 or less, or HBI is 4 or less; or
- 3 The patient has demonstrated an adequate response to treatment, but CDAI score and/or HBI score cannot be assessed.

Initial application — (Crohn's disease - children) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has active Crohn's disease; and
- 2 Either:
 - 2.1 Patient has a PCDAI score of greater than or equal to 30; or
 - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Renewal — (Crohn's disease - children) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab; or
- 2 PCDAI score is 15 or less: or
- 3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed.

Initial application — (Crohn's disease - fistulising) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Crohn's disease: and
- 2 Any of the following:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
 - 2.2 Patient has one or more rectovaginal fistula(e); or
 - 2.3 Patient has complex peri-anal fistula; and
- 3 A Baseline Fistula Assessment has been completed and is no more than 1 month old at the time of application.

Renewal — (Crohn's disease - fistulising) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 The number of open draining fistulae have decreased from baseline by at least 50%; or
- 2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain.

Initial application — (Ocular inflammation - chronic) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

Subsidy		ully	Brand or
 (Manufacturer's Price)	Subsid	isea	Generic
\$	Per	✓	Manufacturer

continued...

- 1 The patient has had an initial Special Authority approval for infliximab for chronic ocular inflammation; or
- 2 Both:
 - 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
 - 2.2 Any of the following:
 - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective; or
 - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
 - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Renewal — (Ocular inflammation - chronic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 12 weeks' initial treatment; or
- 2 Following each 2 year treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 2 year treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Initial application — (Ocular inflammation - severe) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Patient has had an initial Special Authority approval for infliximab for severe ocular inflammation; or
- 2 Both:
 - 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
 - 2.2 Any of the following:
 - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
 - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
 - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Renewal — (Ocular inflammation - severe) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 2 year treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 2 year treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Initial application — (ankylosing spondylitis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has had an initial Special Authority approval for etanercept for ankylosing spondylitis; and

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects; or
 - 1.2.2 The patient has received insufficient benefit to meet the renewal criteria for ankylosing spondylitis; or
- 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis for more than six months; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.3 Patient has bilateral sacroiliitis demonstrated by radiology imaging; and
 - 2.4 Patient has not responded adequately to treatment with two or more NSAIDs, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
 - 2.5 Either:
 - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following BASMI measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
 - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender; and
 - 2.6 A BASDAI of at least 6 on a 0-10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment and is no more than 1 month old at the time of application.

Renewal — (ankylosing spondylitis) from any relevant practitioner. Approvals valid for 2 years where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less.

Initial application — (Arthritis - oligoarticular course juvenile idiopathic) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for oligoarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 Fither:
 - 1.2.1 Patient has experienced intolerable side effects; or
 - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for oligoarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance: and
 - 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
 - 2.3 Either:
 - 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose).

Renewal — (Arthritis - oligoarticular course juvenile idiopathic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Fither:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - polyarticular course juvenile idiopathic) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per ✓	Manufacturer

continued...

Fither:

- 1 Both:
 - 1.1 Patient has had an initial Special Authority approval for etanercept for polyarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects; or
 - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for polyarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance: and
 - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
 - 2.3 Any of the following:
 - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Renewal — (Arthritis - polyarticular course juvenile idiopathic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - psoriatic) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 Patient has had an initial Special Authority approval for etanercept or secukinumab for psoriatic arthritis; and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects; or
 - 1.2.2 The patient has received insufficient benefit from to meet the renewal criteria for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine or leflunomide at maximum tolerated doses (unless contraindicated); and
 - 2.4 Either:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.5 Any of the following:
 - 2.5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an ESR greater than 25 mm per hour; or

Subsidy		Fully	Brand or	
(Manufacturer's Price	Sı	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (Arthritis - psoriatic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Fither:

- 1 Following initial treatment, the patient has at least a 50% decrease in swollen joint count from baseline and a clinically significant response in the opinion of the physician; or
- 2 Patient demonstrates at least a continuing 30% improvement in swollen joint count from baseline and a clinically significant response in the opinion of the treating physician.

Initial application — (Arthritis - rheumatoid) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for rheumatoid arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for rheumatoid arthritis; or
- 2 All of the following:
 - 2.1 Patient has had rheumatoid arthritis (either confirmed by radiology imaging, or the patient is CCP antibody positive) for six months duration or longer; and
 - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.3 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
 - 2.4 Patient has tried and not responded to at least three months of methotrexate in combination with sulfasalazine and hydroxychloroguine sulphate at maximum tolerated doses (unless contraindicated); and
 - 2.5 Fither:
 - 2.5.1 Patient has tried and not responded to at least three months of methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 2.5.2 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with methotrexate; and
 - 2.6 Either:
 - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
 - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip.

Renewal — (Arthritis - rheumatoid) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Initial application — (Still's disease - adult-onset (AOSD)) only from a rheumatologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Fither:

1 Both:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1.1 The patient has had an initial Special Authority approval for etanercept and/or tocilizumab for AOSD; and
- 1.2 Fither:
- 1.2.1 Patient has experienced intolerable side effects from etanercept and/or tocilizumab; or
 - 1.2.2 Patient has received insufficient benefit from at least a three-month trial of etanercept and/or tocilizumab; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria; and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, NSAIDs and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Initial application — (ulcerative colitis) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active ulcerative colitis: and
- 2 Either:
 - 2.1 Patient's SCCAI score is greater than or equal to 4; or
 - 2.2 Patient's PUCAI score is greater than or equal to 20; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from prior therapy with immunomodulators and systemic corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 The SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on biologic therapy; or
- 2 The PUCAI score has reduced by 10 points or more from the PUCAI score when the patient was initiation on biologic therapy.

Initial application — (undifferentiated spondyloarthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undifferentiated peripheral spondyloarthritis* with active peripheral joint arthritis in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2 Patient has tried and not responded to at least three months of each of methotrexate, sulfasalazine and leflunomide, at maximum tolerated doses (unless contraindicated); and
- 3 Any of the following:
 - 3.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 3.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application; or
 - 3.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Note: Indications marked with * are unapproved indications

Renewal — (undifferentiated spondyloarthritis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or

Subsidy (Manufacturer's Price)	Sı	Fully ubsidised	Brand or Generic
 \$	Per	1	Manufacturer

continued...

2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response in the opinion of the treating physician.

Initial application — (inflammatory bowel arthritis – axial) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has axial inflammatory pain for six months or more; and
- 3 Patient is unable to take NSAIDs; and
- 4 Patient has unequivocal sacroiliitis demonstrated by radiological imaging or MRI; and
- 5 Patient has not responded adequately to prior treatment consisting of at least 3 months of an exercise regime supervised by a physiotherapist; and
- 6 A BASDAI of at least 6 on a 0-10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment.

Renewal — (inflammatory bowel arthritis – axial) from any relevant practitioner. Approvals valid for 2 years where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less.

Initial application — (inflammatory bowel arthritis – peripheral) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has active arthritis in at least four joints from the following: hip, knee, ankle, subtalar, tarsus, forefoot, wrist, elbow, shoulder, sternoclavicular; and
- 3 Patient has tried and not experienced a response to at least three months of methotrexate, or azathioprine at a maximum tolerated dose (unless contraindicated); and
- 4 Patient has tried and not experienced a response to at least three months of sulfasalazine at a maximum tolerated dose (unless contraindicated); and
- 5 Any of the following:
 - 5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 5.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application; or
 - 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (inflammatory bowel arthritis – peripheral) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 Patient has experienced at least a continuing 30% improvement in active joint count from baseline in the opinion of the treating physician.

ADALIMUMAB (HUMIRA - ALTERNATIVE BRAND) - Special Authority see \$A2157 below - Retail pharmacy

Inj 20 mg per 0.2 ml prefilled syringe1,	599.96	2 🗸	Humira
Inj 40 mg per 0.4 ml prefilled pen1,	599.96	2 🗸	HumiraPen
Inj 40 mg per 0.4 ml prefilled syringe1,	599.96	2 🗸	Humira

⇒SA2157 Special Authority for Subsidy

Initial application — (Behcet's disease – severe) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Subsidy		Fully	Brand or	
(Manufacturer's Price	e)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Behcet's disease – severe) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient has had a good clinical response to treatment with measurably improved quality of life; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Hidradenitis suppurativa) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 7 days. Fortnightly dosing has been considered.

Renewal — (Hidradenitis suppurativa) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

- All of the following:
 - 1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline; and
 - 2 The patient has a Dermatology Quality of Life Index improvement of 4 or more from baseline; and
 - 3 Adalimumab is to be administered at doses no greater than 40mg every 7 days. Fortnightly dosing has been considered.

Initial application — (Psoriasis - severe chronic plaque) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

- All of the following:
 - 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
 - 2 Patient has received a maximum of 6 months treatment with Amgevita; and
 - 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
 - 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Psoriasis - severe chronic plaque) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 Fither:

Subsidy	Fu	lly Brand or	
(Manufacturer's Price)	Subsidis	ed Generic	
\$	Per	 Manufacturer 	

continued...

- 1.1 Both:
 - 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 1.1.2 Either:
 - 1.1.2.1 Following each prior adalimumab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-adalimumab treatment baseline value: or
 - 1.1.2.2 Following each prior adalimumab treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline value; or
- 1.2 Both:
 - 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 1.2.2 Either:
 - 1.2.2.1 Following each prior adalimumab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 1.2.2.2 Following each prior adalimumab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-adalimumab treatment baseline value: and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Pyoderma gangrenosum) only from a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 A maximum of 8 doses.

Renewal — (Pyoderma gangrenosum) only from a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient has demonstrated clinical improvement and continues to require treatment; and
- 2 A maximum of 8 doses.

Initial application — (Crohn's disease - adult) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevitat; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
 - 1.3 Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and

	Subsidy	Fully	Brand or
(Manu	facturer's Price)	Subsidised	Generic
	\$ Pe	er 🗸	Manufacturer

continued...

3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Crohn's disease - adult) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on adalimumab; or
 - 1.2 CDAI score is 150 or less: or
 - 1.3 The patient has demonstrated an adequate response to treatment, but CDAI score cannot be assessed; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Crohn's disease - children) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
 - 1.3 Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Crohn's disease - children) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab; or
 - 1.2 PCDAI score is 15 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment, but PCDAI score cannot be assessed; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Crohn's disease - fistulising) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
 - 1.3 Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Crohn's disease - fistulising) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 Fither:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
- 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Ocular inflammation – chronic) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with Amgevita, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
 - 1.3 Patient has uveitis and is considered to be at risk of vision loss if they were to change treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Ocular inflammation – chronic) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has had a good clinical response following 12 weeks' initial treatment; or
 - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
 - 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and</p>
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Ocular inflammation – severe) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with Amgevita, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
 - 1.3 Patient has uveitis and is considered to be at risk of vision loss if they were to change treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Ocular inflammation – severe) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has had a good clinical response following 3 initial doses; or
 - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and</p>
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita); and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Arthritis – oligoarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

Renewal — (Arthritis – oligoarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months where the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - polyarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

Renewal — (Arthritis - polyarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months where the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - psoriatic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Arthritis - psoriatic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — **(Arthritis – rheumatoid)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Either:
 - 4.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or
 - 4.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

Renewal — (Arthritis – rheumatoid) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 2 Either:
 - 2.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or
 - 2.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

Initial application — (Still's disease – adult-onset (AOSD)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

Subsidy	F	ılly	Brand or	
(Manufacturer's Price)	Subsidis	sed	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

Renewal — (Still's disease – adult-onset (AOSD)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months where the patient has demonstrated a sustained improvement in inflammatory markers and functional status.

AFLIBERCEPT - Special Authority see SA1772 below - Retail pharmacy

⇒SA1772 Special Authority for Subsidy

Initial application — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Wet age-related macular degeneration (wet AMD); or
 - 1.1.2 Polypoidal choroidal vasculopathy; or
 - 1.1.3 Choroidal neovascular membrane from causes other than wet AMD; and
 - 1.2 Either:
 - 1.2.1 The patient has developed severe endophthalmitis or severe posterior uveitis following treatment with bevacizumab; or
 - 1.2.2 There is worsening of vision or failure of retina to dry despite three intraocular injections of bevacizumab four weeks apart; and
 - 1.3 There is no structural damage to the central fovea of the treated eye; and
 - 1.4 Patient has not previously been treated with ranibizumab for longer than 3 months; or
- 2 Either:
 - 2.1 Patient has current approval to use ranibizumab for treatment of wAMD and was found to be intolerant to ranibizumab within 3 months: or
 - 2.2 Patient has previously* (*before June 2018) received treatment with ranibizumab for wAMD and disease was stable while on treatment.

Initial application — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has centre involving diabetic macular oedema (DMO); and
- 2 Patient's disease is non responsive to 4 doses of intravitreal bevacizumab when administered 4-6 weekly; and
- 3 Patient has reduced visual acuity between 6/9 6/36 with functional awareness of reduction in vision; and
- 4 Patient has DMO within central OCT (ocular coherence tomography) subfield > 350 micrometers; and
- 5 There is no centre-involving sub-retinal fibrosis or foveal atrophy.

Renewal — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Documented benefit must be demonstrated to continue; and
- 2 Patient's vision is 6/36 or better on the Snellen visual acuity score; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	1	Manufacturer	

continued...

3 There is no structural damage to the central fovea of the treated eye.

Renewal — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 There is stability or two lines of Snellen visual acuity gain; and
- 2 There is structural improvement on OCT scan (with reduction in intra-retinal cysts, central retinal thickness, and sub-retinal fluid); and
- 3 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 4 There is no centre-involving sub-retinal fibrosis or foveal atrophy; and
- 5 After each consecutive 12 months treatment with (2nd line anti-VEGF agent), patient has retrialled with at least one injection of bevacizumab and had no response.

Subsidy Fully Brand or
(Manufacturer's Price) Subsidised Generic
\$ Per Manufacturer

⇒SA2151 Special Authority for Subsidy

Initial application — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Patient must have a diagnosis of severe eosinophilic asthma documented by a respiratory physician or clinical immunologist; and
- 3 Conditions that mimic asthma eg. vocal cord dysfunction, central airway obstruction, bronchiolitis etc. have been excluded: and
- 4 Patient has a blood eosinophil count of greater than 0.5 x 10 9 cells/L in the last 12 months; and
- 5 Patient must be adherent to optimised asthma therapy including inhaled corticosteroids (equivalent to at least 1000 mcg per day of fluticasone propionate) plus long-acting beta-2 agonist, or budesonide/formoterol as part of the anti-inflammatory reliever therapy plus maintenance regimen, unless contraindicated or not tolerated; and
- 6 Either:
 - 6.1 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral corticosteroids; or
 - 6.2 Patient has received continuous oral corticosteroids of at least the equivalent of 10 mg per day over the previous 3 months; and
- 7 Treatment is not to be used in combination with subsidised mepolizumab; and
- 8 Patient has an Asthma Control Test (ACT) score of 10 or less. Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 52 weeks after the first dose to assess response to treatment; and
- 9 Either:
 - 9.1 Patient has not previously received an anti-IL5 biological therapy for their severe eosinophilic asthma; or
 - 9.2 Both:
 - 9.2.1 Patient was refractory or intolerant to previous anti-IL5 biological therapy; and
 - 9.2.2 Patient was not eligible to continue treatment with previous anti-IL5 biological therapy and discontinued within 12 months of commencing treatment.

Renewal — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 Either:
 - 2.1 Exacerbations have been reduced from baseline by 50% as a result of treatment with benralizumab; or
 - 2.2 Reduction in continuous oral corticosteroid use by 50% or by 10 mg/day while maintaining or improving asthma control.

⇒SA2289 Special Authority for Subsidy

Initial application — (relapsed/refractory Hodgkin lymphoma) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Both:
 - 1.1.1 Patient has relapsed/refractory CD30-positive Hodgkin lymphoma after two or more lines of chemotherapy; and

Subsidy Fully (Manufacturer's Price) Subsidised		Brand or Generic
(Manufacturer's Frice)	Per 🗸	Manufacturer

continued...

- 1.1.2 Patient is ineligible for autologous stem cell transplant; or
- 1.2 Both:
 - 1.2.1 Patient has relapsed/refractory CD30-positive Hodgkin lymphoma; and
 - 1.2.2 Patient has previously undergone autologous stem cell transplant; and
- 2 Patient has not previously received funded brentuximab vedotin; and
- 3 Response to brentuximab vedotin treatment is to be reviewed after a maximum of 6 treatment cycles; and
- 4 Brentuximab vedotin to be administered at doses no greater than 1.8 mg/kg every 3 weeks.

Renewal — (relapsed/refractory Hodgkin lymphoma) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has achieved a partial or complete response to brentuximab vedotin after 6 treatment cycles; and
- 2 Treatment remains clinically appropriate and the patient is benefitting from treatment and treatment is being tolerated; and
- 3 Patient is to receive a maximum of 16 total cycles of brentuximab vedotin treatment.

Initial application — (anaplastic large cell lymphoma) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has relapsed/refractory CD30-positive systemic anaplastic large cell lymphoma; and
- 2 Patient has an ECOG performance status of 0-1; and
- 3 Patient has not previously received brentuximab vedotin; and
- 4 Response to brentuximab vedotin treatment is to be reviewed after a maximum of 6 treatment cycles; and
- 5 Brentuximab vedotin to be administered at doses no greater than 1.8 mg/kg every 3 weeks.

Renewal — (anaplastic large cell lymphoma) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has achieved a partial or complete response to brentuximab vedotin after 6 treatment cycles; and
- 2 Treatment remains clinically appropriate and the patient is benefitting from treatment and treatment is being tolerated; and
- 3 Patient is to receive a maximum of 16 total cycles of brentuximab vedotin treatment.

CASIRIVIMAB AND IMDEVIMAB - [Xpharm] - Special Authority see \$A2096 below

⇒SA2096 Special Authority for Subsidy

Initial application — (Treatment of profoundly immunocompromised patients) from any relevant practitioner. Approvals valid for 2 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed (or probable) COVID-19; and
- 2 The patient is in the community with mild to moderate disease severity*; and
- 3 Patient is profoundly immunocompromised** and is at risk of not having mounted an adequate response to vaccination against COVID-19 or is unvaccinated; and
- 4 Patient's symptoms started within the last 10 days; and
- 5 Patient is not receiving high flow oxygen or assisted/mechanical ventilation; and
- 6 Casirivimab and imdevimab is to be administered at a maximum dose of no greater than 2,400 mg.

Notes: * Mild to moderate disease severity as described on the Ministry of Health Website

** Examples include B-cell depletive illnesses or patients receiving treatment that is B-Cell depleting.

CETUXIMAB - PCT only - Specialist - Special Authority see SA1697 on the next page

Erbitux	1	364.00	 ml vial	Inj 5 mg per ml, 2
✓ Erbitux	1	1,820.00	 0 ml vial	Inj 5 mg per ml, 1
✓ Baxter	1 ma	3.82	 	Ini 1 ma for ECP.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA1697 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, non-metastatic, squamous cell cancer of the head and neck; and
- 2 Patient is contraindicated to, or is intolerant of, cisplatin; and
- 3 Patient has good performance status; and
- 4 To be administered in combination with radiation therapy.

GEMTUZUMAB OZOGAMICIN − PCT only − Specialist − Special Authority see SA2269 below Inj 5 mg vial12,973.00 1 ✓ Mylotarg

⇒SA2269 Special Authority for Subsidy

Initial application only from a haematologist, paediatric haematologist or paediatric oncologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has not received prior chemotherapy for this condition; and
- 2 Patient has de novo CD33-positive acute myeloid leukaemia; and
- 3 Patient does not have acute promyelocytic leukaemia; and
- 4 Gemtuzumab ozogamicin will be used in combination with standard anthracycline and cytarabine (AraC); and
- 5 Patient is being treated with curative intent; and
- 6 Patient's disease risk has been assessed by cytogenetic testing to be good or intermediate; and
- 7 Patient must be considered eligible for standard intensive remission induction chemotherapy with standard anthracycline and cytarabine (AraC); and
- 8 Gemtuzumab ozogamicin to be funded for one course only (one dose at 3 mg per m² body surface area or up to 2 vials of 5 mg as separate doses).

Note: Acute myeloid leukaemia excludes acute promyelocytic leukaemia and acute myeloid leukaemia that is secondary to another haematological disorder (eg myelodysplasia or myeloproliferative disorder).

⇒SA2179 Special Authority for Subsidy

Initial application — (**Crohn's disease (adults))** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active Crohn's disease: and
- 2 Any of the following:
 - 2.1 Patient has a CDAI score of greater than or equal to 300 or HBI score of greater than or equal to 10; or
 - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
 - 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but has experienced an inadequate response to, or intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Renewal — (Crohn's disease (adults)) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

1 Any of the following:

Subsidy	Fu	,	
(Manufacturer's Price)	Subsidis	ed Generic	
\$	Per	 Manufacturer 	

continued...

- 1.1 CDAI score has reduced by 100 points from the CDAI score, or HBI score has reduced by 3 points, from when the patient was initiated on infliximab; or
- 1.2 CDAI score is 150 or less, or HBI is 4 or less; or
- 1.3 The patient has demonstrated an adequate response to treatment but CDAI score and/or HBI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — **(Crohn's disease (children))** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has active Crohn's disease; and
- 2 Either:
 - 2.1 Patient has a PCDAI score of greater than or equal to 30; or
 - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but experienced an inadequate response to, or intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Renewal — (Crohn's disease (children)) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on infliximab; or
 - 1.2 PCDAI score is 15 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — **(Graft vs host disease)** from any relevant practitioner. Approvals valid without further renewal unless notified where patient has steroid-refractory acute graft vs. host disease of the gut.

Initial application — (Pulmonary sarcoidosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has life-threatening pulmonary sarcoidosis diagnosed by a multidisciplinary team that is refractory to other treatments.

Initial application — (acute fulminant ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 weeks for applications meeting the following criteria:
Both:

- 1 Patient has acute, fulminant ulcerative colitis; and
- 2 Treatment with intravenous or high dose oral corticosteroids has not been successful.

Initial application — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Following 12 weeks of infliximab treatment, BASDAI has improved by 4 or more points from pre-infliximab baseline on a 10 point scale, or by 50%, whichever is less: and
- 2 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 3 Infliximab to be administered at doses no greater than 5 mg/kg every 6-8 weeks.

Initial application — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for chronic ocular inflammation; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for chronic ocular inflammation; or
- 2 Both:
 - 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
 - 2.2 Any of the following:
 - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective; or
 - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
 - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Renewal — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.</p>

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (fistulising Crohn's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has confirmed Crohn's disease: and
- 2 Any of the following:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
 - 2.2 Patient has one or more rectovaginal fistula(e); or
 - 2.3 Patent has complex peri-anal fistula.

Renewal — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a

Subsidy (Manufacturer's Price)	Fully Subsidised		
(ivialitulacture) 5 Filoe)	Per 🗸	Manufacturer	
Φ	rei •	Manuacturei	

continued...

gastroenterologist. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Either:
 - 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
 - 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline (in the case of adult patients, as demonstrated by a reduction in the Fistula Assessment score), together with less induration and patient reported pain: and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (neurosarcoidosis) only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with neurosarcoiosis by a multidisciplinary team; and
- 2 Patient has CNS involvement; and
- 3 Patient has steroid-refractory disease; and
- 4 Either:
 - 4.1 IV cyclophosphamide has been tried; or
 - 4.2 Treatment with IV cyclophosphamide is clinically inappropriate.

Renewal — (neurosarcoidosis) only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria:

- Either:
 - 1 A withdrawal period has been tried and the patient has relapsed; or
 - 2 All of the following:
 - 2.1 A withdrawal period has been considered but would not be clinically appropriate; and
 - 2.2 There has been a marked reduction in prednisone dose; and
 - 2.3 Either:
 - 2.3.1 There has been an improvement in MRI appearances; or
 - 2.3.2 Marked improvement in other symptomology.

Initial application — (plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab, etanercept or secukinumab for severe chronic plaque psoriasis; and
 - 1.2 Fither:
 - 1.2.1 Patient has experienced intolerable side effects from adalimumab, etanercept or secukinumab; or
 - 1.2.2 Patient has received insufficient benefit from adalimumab, etanercept or secukinumab to meet the renewal criteria for adalimumab. etanercept or secukinumab for severe chronic plaque psoriasis: or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and

Subsidy (Manufacturer's Price)	Sub	Fully sidised	Brand or Generic
\$	Per	1	Manufacturer

continued...

- 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin. or acitretin: and
- 2.3 A PASI assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course: and
- 2.4 The most recent PASI assessment is no more than 1 month old at the time of initiation.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

- 1 Fither:
 - 1.1 Both:
 - 1.1.1 Patient had "whole body" severe chronic plague psoriasis at the start of treatment; and
 - 1.1.2 Following each prior infliximab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-infliximab treatment baseline value; or
 - 1.2 Both:
 - 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 1.2.2 Either:
 - 1.2.2.1 Following each prior infliximab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 1.2.2.2 Following each prior infliximab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-infliximab treatment baseline value: and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient was being treated with infliximab prior to 1 February 2019; and
- 2 Any of the following:
 - 2.1 Rheumatoid arthritis; or
 - 2.2 Ankylosing spondylitis; or
 - 2.3 Psoriatic arthritis; or
 - 2.4 Severe ocular inflammation: or
 - 2.5 Chronic ocular inflammation; or
 - 2.6 Crohn's disease (adults); or
 - 2.7 Crohn's disease (children); or
 - 2.8 Fistulising Crohn's disease: or
 - 2.9 Severe fulminant ulcerative colitis; or
 - 2.10 Severe ulcerative colitis: or
 - 2.11 Plaque psoriasis: or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per ✓	Manufacturer

continued...

- 2.12 Neurosarcoidosis: or
- 2.13 Severe Behcet's disease.

Initial application — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis; and
- 2 Fither:
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept and/or secukinumab; or
 - 2.2 Following 3-4 months' initial treatment with adalimumab and/or etanercept and/or secukinumab, the patient did not meet the renewal criteria for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Roth:

- 1 Either:
 - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior infliximab treatment in the opinion of the treating physician; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Fither:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept: and
- 3 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance.

Renewal — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Infliximab to be administered at doses no greater than 3 mg/kg every 8 weeks.

Initial application — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe Behcet's disease which is significantly impacting the patient's quality of life (see Notes); and
- 2 Fither:

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
 \$	Per	√	

continued...

- 2.1 The patient has severe ocular, neurological and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s) (see Notes); or
- 2.2 The patient has severe gastrointestinal, rheumatologic and/or mucocutaneous symptoms and has not responded adequately to two or more treatment appropriate for the particular symptom(s) (see Notes); and
- 3 The patient is experiencing significant loss of quality of life.

Notes: Behcet's disease diagnosed according to the International Study Group for Behcet's Disease. Lancet 1990:335(8697):1078-80. Quality of life measured using an appropriate quality of life scale such as that published in Gilworth et al J Rheumatol. 2004:31:931-7.

Treatments appropriate for the particular symptoms are those that are considered standard conventional treatments for these symptoms, for example intravenous/oral steroids and other immunosuppressants for ocular symptoms; azathioprine, steroids, thalidomide, interferon alpha and ciclosporin for mucocutaneous symptoms; and colchicine, steroids and methotrexate for rheumatological symptoms.

Renewal — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has had a good clinical response to initial treatment with measurably improved quality of life; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Renewal — (fulminant ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Where maintenance treatment is considered appropriate, infliximab should be used in combination with immunomodulators and reassessed every 6 months; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe ocular inflammation; and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe ocular inflammation: or
- 2 Both:
 - 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
 - 2.2 Any of the following:
 - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
 - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
 - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Renewal — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

1 The patient has had a good clinical response following 3 initial doses: or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per ✓	Manufacturer

continued...

- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (ulcerative colitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active ulcerative colitis: and
- 2 Either:
 - 2.1 Patients SCCAI is greater than or equal to 4; or
 - 2.2 Patients PUCAI score is greater than or equal to 20; and
- 3 Patient has tried but has experienced an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and systemic corticosteroids.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 The SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on infliximab: or
 - 1.2 The PUCAI score has reduced by 10 points or more from the PUCAI score when the patient was initiated on infliximab; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Note: Indications marked with * are unapproved indications.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

Initial application — (inflammatory bowel arthritis – axial) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has had axial inflammatory pain for six months or more; and

Subsidy		Fully	Brand or	
(Manufacturer's Price	e)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 3 Patient is unable to take NSAIDs; and
- 4 Patient has unequivocal sacroiliitis demonstrated by radiological imaging or MRI; and
- 5 Patient's disease has not responded adequately to prior treatment consisting of at least 3 months of an exercise regime supervised by a physiotherapist; and
- 6 Patient has a BASDAI of at least 6 on a 0 10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment.

Renewal — (inflammatory bowel arthritis – axial) from any relevant practitioner. Approvals valid for 2 years where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10-point scale, or an improvement in BASDAI of 50%, whichever is less.

Initial application — (inflammatory bowel arthritis – peripheral) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has active arthritis in at least four joints from the following: hip, knee, ankle, subtalar, tarsus, forefoot, wrist, elbow, shoulder, sternoclavicular; and
- 3 Patient has tried and not experienced a response to at least three months of methotrexate or azathioprine at a maximum tolerated dose (unless contraindicated); and
- 4 Patient has tried and not experienced a response to at least three months of sulfasalazine at a maximum tolerated dose (unless contraindicated); and
- 5 Any of the following:
 - 5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application: or
 - 5.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application; or
 - 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (inflammatory bowel arthritis – peripheral) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, patient has experienced at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 Patient has experienced at least a continuing 30% improvement in active joint count from baseline in the opinion of the treating physician.

MEPOLIZUMAB - Special Authority see SA2331 below - Retail pharmacy

 Inj 100 mg prefilled pen
 1,638.00
 1
 Nucala

 Inj 100 mg vial
 1,638.00
 1
 Nucala

(Nucala Ini 100 mg vial to be delisted 1 August 2024)

⇒SA2331 Special Authority for Subsidy

Initial application — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Patient must have a diagnosis of severe eosinophilic asthma documented by a respiratory physician or clinical immunologist; and
- 3 Conditions that mimic asthma eg. vocal cord dysfunction, central airway obstruction, bronchiolitis etc. have been excluded; and
- 4 Patient has a blood eosinophil count of greater than 0.5 x 10 9 cells/L in the last 12 months; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsidised		Generic	
\$	Per	1	Manufacturer	

continued...

- 5 Patient must be adherent to optimised asthma therapy including inhaled corticosteroids (equivalent to at least 1000 mcg per day of fluticasone propionate) plus long acting beta-2 agonist, or budesonide/formoterol as part of the single maintenance and reliever therapy regimen, unless contraindicated or not tolerated; and
- 6 Fither:
 - 6.1 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral corticosteroids: or
 - 6.2 Patient has received continuous oral corticosteroids of at least the equivalent of 10 mg per day over the previous 3 months: and
- 7 Treatment is not to be used in combination with subsidised benralizumab; and
- 8 Patient has an Asthma Control Test (ACT) score of 10 or less. Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 52 weeks after the first dose to assess response to treatment; and
- 9 Either:
 - 9.1 Patient has not previously received an anti-IL5 biological therapy for their severe eosinophilic asthma; or
 - 9.2 Both:
 - 9.2.1 Patient was refractory or intolerant to previous anti-IL5 biological therapy; and
 - 9.2.2 Patient was not eligible to continue treatment with previous anti-IL5 biological therapy and discontinued within 12 months of commencing treatment.

Renewal — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 Fither:
 - 2.1 Exacerbations have been reduced from baseline by 50% as a result of treatment with mepolizumab; or
 - 2.2 Reduction in continuous oral corticosteroid use by 50% or by 10 mg/day while maintaining or improving asthma control.

Initial application — (eosinophilic granulomatosis with polyangiitis) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has eosinophilic granulomatosis with polyangiitis; and
- 2 The patient has trialled and not received adequate benefit from at least one of the following for at least three months (unless contraindicated to all): azathioprine, cyclophosphamide, leflunomide, methotrexate, mycophenolate, or rituximab; and
- 3 Either:
 - 3.1 The patient has trialled prednisone for a minimum of three months and is unable to maintain disease control at doses below 7.5 mg per day; or
 - 3.2 Corticosteroids are contraindicated.

Renewal — (eosinophilic granulomatosis with polyangiitis) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months where patient has no evidence of clinical disease progression.

OBINUTUZUMAB - PCT only - Specialist - Special Aut	thority see SA2155 on the I	next page	
Inj 25 mg per ml, 40 ml vial	5,910.00	1	✓ Gazyva
Inj 1 mg for ECP	6.21	1 mg	✓ Baxter

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA2155 Special Authority for Subsidy

Initial application — (chronic lymphocytic leukaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C CD20+ chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is obinutuzumab treatment naive; and
- 3 The patient is not eligible for full dose FCR due to comorbidities with a score > 6 on the Cumulative Illness Rating Scale (CIRS) or reduced renal function (creatinine clearance < 70mL/min); and</p>
- 4 Patient has adequate neutrophil and platelet counts* unless the cytopenias are a consequence of marrow infiltration by CLL; and
- 5 Patient has good performance status; and
- 6 Obinutuzumab to be administered at a maximum cumulative dose of 8,000 mg and in combination with chlorambucil for a maximum of 6 cycles.

Notes: Chronic lymphocytic leukaemia includes small lymphocytic lymphoma. Comorbidity refers only to illness/impairment other than CLL induced illness/impairment in the patient. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with obinutuzumab is expected to improve symptoms and improve ECOG score to < 2.

* Neutrophil greater than or equal to 1.5×10^9 /L and platelets greater than or equal to 75×10^9 /L.

Initial application — (follicular / marginal zone lymphoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 Patient has follicular lymphoma; or
 - 1.2 Patient has marginal zone lymphoma; and
- 2 Patient is refractory to or has relapsed within 12 months of a rituximab containing combined chemo-immunotherapy regimen*; and
- 3 Patient has an ECOG performance status of 0-2; and
- 4 Patient has been previously treated with no more than four chemotherapy regimens; and
- 5 Obinutuzumab to be administered at a maximum dose of 1000 mg for a maximum of 6 cycles in combination with chemotherapy*.

Note: * includes unapproved indications

Renewal — (follicular / marginal zone lymphoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 24 months for applications meeting the following criteria: All of the following:

- 1 Patient has no evidence of disease progression following obinutuzumab induction therapy; and
- 2 Obinutuzumab to be administered at a maximum of 1000 mg every 2 months for a maximum of 2 years; and
- 3 Obinutuzumab to be discontinued at disease progression.

OMALIZUMAB - Special Authority see SA1744 below - Retail pharmacy

⇒SA1744 Special Authority for Subsidy

Initial application — (severe asthma) only from a respiratory specialist or clinical immunologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 6 years or older; and
- 2 Patient has a diagnosis of severe asthma; and
- 3 Past or current evidence of atopy, documented by skin prick testing or RAST; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 4 Total serum human immunoglobulin E (IgE) between 76 IU/mL and 1300 IU/ml at baseline; and
- 5 Proven adherence with optimal inhaled therapy including high dose inhaled corticosteroid (budesonide 1,600 mcg per day or fluticasone propionate 1,000 mcg per day or equivalent), plus long-acting beta-2 agonist therapy (at least salmeterol 50 mcg bd or eformoterol 12 mcg bd) for at least 12 months, unless contraindicated or not tolerated; and
- 6 Either:
 - 6.1 Patient has received courses of systemic corticosteroids equivalent to at least 28 days treatment in the past 12 months. unless contraindicated or not tolerated; or
 - 6.2 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral steroids; and
- 7 Patient has an Asthma Control Test (ACT) score of 10 or less; and
- 8 Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 26 weeks after the first dose to assess response to treatment.

Initial application — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient is symptomatic with Urticaria Activity Score 7 (UAS7) of 20 or above; and
 - 2.1.2 Patient has a Dermatology life quality index (DLQI) of 10 or greater; or
 - 2.2 Patient has a Urticaria Control Test (UCT) of 8 or less; and
- 3 Any of the following:
 - 3.1 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and ciclosporin (> 3 mg/kg day) for at least 6 weeks; or
 - 3.2 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and at least 3 courses of systemic corticosteroids (> 20 mg prednisone per day for at least 5 days) in the previous 6 months; or
 - 3.3 Patient has developed significant adverse effects whilst on corticosteroids or ciclosporin; and
- 4 Either:
 - 4.1 Treatment to be stopped if inadequate response* following 4 doses; or
 - 4.2 Complete response* to 6 doses of omalizumab.

Renewal — (severe asthma) only from a clinical immunologist or respiratory specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 A reduction in the maintenance oral corticosteroid dose or number of exacerbations of at least 50% from baseline.

Renewal — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Patient has previously adequately responded* to 6 doses of omalizumab; or
- 2 Both:
 - 2.1 Patient has previously had a complete response* to 6 doses of omalizumab; and
 - 2.2 Patient has relapsed after cessation of omalizumab therapy.

Note: *Inadequate response defined as less than 50% reduction in baseline UAS7 and DLQI score, or an increase in Urticaria Control Test (UCT) score of less than 4 from baseline. Patient is to be reassessed for response after 4 doses of omalizumab. Complete response is defined as UAS7 less than or equal to 6 and DLQI less than or equal to 5; or UCT of 16. Relapse of chronic urticaria on stopping prednisone/ciclosporin does not justify the funding of omalizumab.

	Subsidy (Manufacturer's Pric	ce) Su Per	Fully ibsidised	Brand or Generic Manufacturer	
PERTUZUMAB - PCT only - Specialist - Special Authority see	SA2276 below				
Inj 30 mg per ml, 14 ml vial	3,927.00	1	✓ P	erjeta	
Inj 420 mg for ECP	3,927.00	420 mg OF	• • B	axter	

⇒SA2276 Special Authority for Subsidy

Initial application — (metastatic breast cancer) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Either:
 - 2.1 Patient is chemotherapy treatment naïve; or
 - 2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
- 3 The patient has good performance status (ECOG grade 0-1); and
- 4 Pertuzumab to be administered in combination with trastuzumab; and
- 5 Pertuzumab maximum first dose of 840 mg, followed by maximum of 420 mg every 3 weeks; and
- 6 Pertuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Fither:

....

- 1 Both:
 - 1.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
 - 1.2 The cancer has not progressed at any time point during the previous 12 months whilst on pertuzumab and trastuzumab; or
- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with pertuzumab and trastuzumab for reasons other than severe toxicity or disease progression; and
 - 2.2 Patient has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with pertuzumab and trastuzumab.

RITUXIMAB (MABTHERA) - PCT only - Specialist - Special	Authority see SA19	76 below	
Inj 100 mg per 10 ml vial	1,075.50	2	✓ Mabthera
Inj 500 mg per 50 ml vial	2,688.30	1	✓ Mabthera
Inj 1 mg for ECP	5.64	1 mg	✓ Baxter (Mabthera)

⇒SA1976 Special Authority for Subsidy

Initial application — (rheumatoid arthritis - TNF inhibitors contraindicated) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Treatment with a Tumour Necrosis Factor alpha inhibitor is contraindicated; and
- 2 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
- 5 Any of the following:
 - 5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Su	ıbsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

the maximum tolerated dose of ciclosporin; or

- 5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
- 5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
- 6 Fither:
 - 6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
 - 6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 7 Either:
 - 7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months; and
- 8 Fither:
 - 8.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 8.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 9 Maximum of two 1.000 mg infusions of rituximab given two weeks apart.

Initial application — (rheumatoid arthritis - prior TNF inhibitor use) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Both:
 - 1.1 The patient has had an initial community Special Authority approval for at least one of etanercept and/or adalimumab for rheumatoid arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 1.2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
 - 2.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 2.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 3 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Renewal — (rheumatoid arthritis - re-treatment in 'partial responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 At 4 months following the initial course of rituximab infusions the patient had between a 30% and 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 At 4 months following the second course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.3 At 4 months following the third and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Fither:

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
- 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Renewal — (rheumatoid arthritis - re-treatment in 'responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 At 4 months following the initial course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 At 4 months following the second and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Either:
 - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

RITUXIMAB (RIXIMYO) - PCT only - Specialist - Special Authority see \$A2233 below

Inj 100 mg per 10 ml vial275.3	3 2	✓ Riximyo
Inj 500 mg per 50 ml vial688.2	0 1	✓ Riximyo
Inj 1 mg for ECP1.3	8 1 m(✓ Baxter (Riximyo)

⇒SA2233 Special Authority for Subsidy

Initial application — (ABO-incompatible organ transplant) from any relevant practitioner. Approvals valid without further renewal unless notified where patient is to undergo an ABO-incompatible solid organ transplant*.

Note: Indications marked with * are unapproved indications.

Initial application — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks; and
- 3 Any of the following:
 - 3.1 Induction therapy with daily oral or pulse intravenous cyclophosphamide has failed to achieve significant improvement of disease after at least 3 months; or
 - 3.2 Patient has previously had a cumulative dose of cyclophosphamide > 15 g or a further repeat 3 month induction course of cyclophosphamide would result in a cumulative dose > 15 g; or
 - 3.3 Cyclophosphamide and methotrexate are contraindicated; or
 - 3.4 Patient is a female of child-bearing potential; or
 - 3.5 Patient has a previous history of haemorrhagic cystitis, urological malignancy or haematological malignancy.

Note: Indications marked with * are unapproved indications.

Renewal — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 Patient has previously responded to treatment with rituximab but is now experiencing an acute flare of vasculitis; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of

Subsidy Fully Brand or
(Manufacturer's Price) Subsidised Generic
\$ Per ✓ Manufacturer

continued...

4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (Antibody-mediated organ transplant rejection) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has been diagnosed with antibody-mediated organ transplant rejection*.

Note: Indications marked with * are unapproved indications.

Initial application — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C chronic lymphocytic leukaemia (CLL) requiring treatment; and
- 2 Any of the following:
 - 2.1 The patient is rituximab treatment naive: or
 - 2.2 Either:
 - 2.2.1 The patient is chemotherapy treatment naive; or
 - 2.2.2 Both:
 - 2.2.2.1 The patient's disease has relapsed following no more than three prior lines of chemotherapy treatment: and
 - 2.2.2.2 The patient has had a treatment-free interval of 12 months or more if previously treated with fludarabine and cyclophosphamide chemotherapy; or
 - 2.3 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax; and
- 3 The patient has good performance status; and
- 4 Either:
 - 4.1 The patient does not have chromosome 17p deletion CLL: or
 - 4.2 Rituximab treatment is to be used in combination with funded venetoclax for relapsed/refractory chronic lymphocytic leukaemia; and
- 5 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles; and
- 6 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration), bendamustine or venetoclax.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with rituximab is expected to improve symptoms and improve ECOG score to < 2.

Renewal — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax; or
 - 1.2 All of the following:
 - 1.2.1 The patient's disease has relapsed following no more than one prior line of treatment with rituximab for CLL; and
 - 1.2.2 The patient has had an interval of 36 months or more since commencement of initial rituximab treatment; and
 - 1.2.3 The patient does not have chromosome 17p deletion CLL; and
 - 1.2.4 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration) or bendamustine; and

Subsidy		Fully	Brand or
(Manufacturer's Price	e)	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

2 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initial application — (Neuromyelitis Optica Spectrum Disorder(NMOSD)) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks; and
- 2 Either:
 - 2.1 The patient has experienced a severe episode or attack of NMOSD (rapidly progressing symptoms and clinical investigations supportive of a severe attack of NMOSD); or
 - 2.2 All of the following:
 - 2.2.1 The patient has experienced a breakthrough attack of NMOSD; and
 - 2.2.2 The patient is receiving treatment with mycophenolate; and
 - 2.2.3 The patients is receiving treatment with corticosteroids.

Renewal — (Neuromyelitis Optica Spectrum Disorder) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks; and
- 2 The patients has responded to the most recent course of rituximab; and
- 3 The patient has not received rituximab in the previous 6 months.

Initial application — (Post-transplant) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 2 To be used for a maximum of 8 treatment cycles.

Note: Indications marked with * are unapproved indications.

Renewal — (Post-transplant) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 3 To be used for no more than 6 treatment cycles.

Note: Indications marked with * are unapproved indications.

Initial application — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
- 2 Either:
 - 2.1 Treatment with corticosteroids and at least one other immunosuppressant for at least a period of 12 months has been ineffective; or
 - 2.2 Both:
 - 2.2.1 Treatment with at least one other immunosuppressant for a period of at least 12 months; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

2.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

Renewal — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Either:
 - 3.1 The patient has relapsed despite treatment with corticosteroids and at least one other immunosuppressant for a period of at least 12 months; or
 - 3.2 Both:
 - 3.2.1 The patient's myasthenia gravis has relapsed despite treatment with at least one immunosuppressant for a period of at least 12 months; and
 - 3.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects

Initial application — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is a child with SDNS* or FRNS*: and
- 2 Treatment with steroids for at least a period of 3 months has been ineffective or associated with evidence of steroid toxicity; and
- 3 Treatment with ciclosporin for at least a period of 3 months has been ineffective and/or discontinued due to unacceptable side effects; and
- 4 Treatment with mycophenolate for at least a period of 3 months with no reduction in disease relapses; and
- 5 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is a child with SRNS* where treatment with steroids and ciclosporin for at least 3 months have been ineffective; and
 - 2 Treatment with tacrolimus for at least 3 months has been ineffective; and
 - 3 Genetic causes of nephrotic syndrome have been excluded; and
 - 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidis	sed	Generic
	Per	1	Manufacturer

continued...

Note: Indications marked with * are unapproved indications.

Renewal — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient has treatment naive aggressive CD20 positive NHL; and
 - 1.2 To be used with a multi-agent chemotherapy regimen given with curative intent; and
 - 1.3 To be used for a maximum of 8 treatment cycles; or
- 2 Both:
 - 2.1 The patient has aggressive CD20 positive NHL with relapsed disease following prior chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Renewal — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has relapsed refractory/aggressive CD20 positive NHL; and
- 3 To be used with a multi-agent chemotherapy regimen given with curative intent; and
- 4 To be used for a maximum of 4 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Initial application — (haemophilia with inhibitors) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria:

Any of the following:

- 1 Patient has mild congenital haemophilia complicated by inhibitors; or
- 2 Patient has severe congenital haemophilia complicated by inhibitors and has failed immune tolerance therapy; or
- 3 Patient has acquired haemophilia.

Renewal — (haemophilia with inhibitors) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for haemophilia with inhibitors; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment.

Initial application — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Fither:
 - 1.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per microlitre: or

Subsidy	Fu	,	
(Manufacturer's Price)	Subsidis	ed Generic	
\$	Per	 Manufacturer 	

continued...

- 1.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding; and
- 2 Any of the following:
 - 2.1 Treatment with steroids and splenectomy have been ineffective; or
 - 2.2 Treatment with steroids has been ineffective and splenectomy is an absolute contraindication; or
 - 2.3 Other treatments including steroids have been ineffective and patient is being prepared for elective surgery (e.g. splenectomy); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks

Note: Indications marked with * are unapproved indications.

Renewal — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Fither:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for immune thrombocytopenic purpura*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (indolent, low-grade lymphomas or hairy cell leukaemia*) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has indolent low grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
 - 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
 - 2.1 The patient has indolent, low grade lymphoma or hairy cell leukaemia* requiring first-line systemic chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Renewal — (indolent, low-grade lymphomas or hairy cell leukaemia*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has indolent, low-grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
- 3 To be used for no more than 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Initial application — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient has autoimmune pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder.

Note: Indications marked with * are unapproved indications.

Renewal — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient was previously treated with rituximab for pure red cell aplasia*

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	/	Manufacturer	

continued

associated with a demonstrable B-cell lymphoproliferative disorder and demonstrated an initial response lasting at least

Note: Indications marked with * are unapproved indications.

Initial application — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Patient has cold haemagglutinin disease*; and
- 2 Patient has severe disease which is characterized by symptomatic anaemia, transfusion dependence or disabling circulatory symptoms; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks

Note: Indications marked with * are unapproved indications.

Renewal — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for severe cold haemagglutinin disease*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Both:

- 1 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks: and
- 2 Either:
 - 2.1 Patient has thrombotic thrombocytopenic purpura* and has experienced progression of clinical symptoms or persistent thrombocytopenia despite plasma exchange; or
 - 2.2 Patient has acute idiopathic thrombotic thrombocytopenic purpura* with neurological or cardiovascular pathology.

Note: Indications marked with * are unapproved indications.

Renewal — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

- All of the following:
 - 1 Patient was previously treated with rituximab for thrombotic thrombocytopenic purpura*: and
 - 2 An initial response lasting at least 12 months was demonstrated; and
 - 3 Patient now requires repeat treatment; and
 - 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe, immediately life- or organ-threatening SLE*; and
- 2 The disease has proved refractory to treatment with steroids at a dose of at least 1 mg/kg; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per ✓	Manufacturer

continued...

- 3 The disease has relapsed following prior treatment for at least 6 months with maximal tolerated doses of azathioprine, mycophenolate mofetil and high dose cyclophosphamide, or cyclophosphamide is contraindicated; and
- 4 Maximum of four 1000 mg infusions of rituximab.

Note: Indications marked with * are unapproved indications.

Renewal — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient's SLE* achieved at least a partial response to the previous round of prior rituximab treatment; and
- 2 The disease has subsequently relapsed; and
- 3 Maximum of two 1000 mg infusions of rituximab.

Note: Indications marked with * are unapproved indications.

Initial application — (warm autoimmune haemolytic anaemia (warm AlHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has warm autoimmune haemolytic anaemia*; and
- 2 One of the following treatments has been ineffective: steroids (including if patient requires ongoing steroids at doses equivalent to > 5 mg prednisone daily), cytotoxic agents (e.g. cyclophosphamide monotherapy or in combination), intravenous immunoglobulin: and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (warm autoimmune haemolytic anaemia (warm AIHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for warm autoimmune haemolytic anaemia*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (severe antisynthetase syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed antisynthetase syndrome; and
- 2 Patient has severe, immediately life or organ threatening disease, including interstitial lung disease; and
- 3 Either:
 - 3.1 Treatment with at least 3 immunosuppressants (oral steroids, cyclophosphamide, methotrexate, mycophenolate, ciclosporin, azathioprine) has not be effective at controlling active disease; or
 - 3.2 Rapid treatment is required due to life threatening complications; and
- 4 Maximum of four 1.000mg infusions of rituximab.

Renewal — (severe antisynthetase syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in inflammatory markers, muscle strength and pulmonary function; and

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidis	sed	Generic
	Per	1	Manufacturer

continued...

- 2 The patient has not received rituximab in the previous 6 months; and
- 3 Maximum of two cycles of 2 × 1,000mg infusions of rituximab given two weeks apart.

Initial application — (graft versus host disease) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has refractory graft versus host disease following transplant; and
- 2 Treatment with at least 3 immunosuppressants (oral steroids, ciclosporin, tacrolimus, mycophenolate, sirolimus) has not be effective at controlling active disease; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Initial application — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has severe chronic inflammatory demyelinating polyneuropathy (CIPD); and
- 2 Fither
 - 2.1 Both:
 - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
 - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
 - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function compared to baseline; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Initial application — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe anti-NMDA receptor autoimmune encephalitis: and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
 - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
 - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subs	sidised	Generic
\$	Per	1	Manufacturer

continued...

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 The patient has experienced a relapse and now requires further treatment; and
- 4 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Initial application — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has CD20+ low grade or follicular B-cell NHL with relapsed disease following prior chemotherapy; and
 - 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
 - 2.1 The patient has CD20+ low grade or follicular B-cell NHL requiring first-line systemic chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Renewal — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 Rituximab is to be used for maintenance in CD20+ low grade or follicular B-cell NHL following induction with first-line systemic chemotherapy; and
- 2 Patient is intended to receive rituximab maintenance therapy for 2 years at a dose of 375 mg/m2 every 8 weeks (maximum of 12 cycles).

Initial application — (Membranous nephropathy) only from a nephrologist or any relevant practitioner on the recommendation of a nephrologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient has biopsy-proven primary/idiopathic membranous nephropathy*; or
 - 1.2 Patient has PLA2 antibodies with no evidence of secondary cause, and an eGFR of > 60ml/min/1.73m2; and
- 2 Patient remains at high risk of progression to end-stage kidney disease despite more than 3 months of treatment with conservative measures (see Note): and
- 3 The total rituximab dose would not exceed the equivalent of 375mg/m2 of body surface area per week for a total of 4 weeks

Renewal — (Membranous nephropathy) only from a nephrologist or any relevant practitioner on the recommendation of a nephrologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for membranous nephropathy*; and
- 2 Either:
 - 2.1 Treatment with rituximab was previously successful, but the condition has relapsed, and the patient now requires repeat treatment; or
 - 2.2 Patient achieved partial response to treatment and requires repeat treatment (see Note); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Notes:

- a) Indications marked with * are unapproved indications.
- b) High risk of progression to end-stage kidney disease defined as > 5g/day proteinuria.
- c) Conservative measures include renin-angiotensin system blockade, blood-pressure management, dietary sodium and protein restriction, treatment of dyslipidaemia, and anticoagulation agents unless contraindicated or the patient has

Subsidy	Oute	Fully	Brand or
 (Manufacturer's Price) \$	Per	idised •	Generic Manufacturer

continued...

experienced intolerable side effects.

d) Partial response defined as a reduction of proteinuria of at least 50% from baseline, and between 0.3 grams and 3.5 grams per 24 hours.

Initial application — (B-cell acute lymphoblastic leukaemia/lymphoma*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient has newly diagnosed B-cell acute lymphoblastic leukaemia/lymphoma*; and
- 2 Treatment must be in combination with an intensive chemotherapy protocol with curative intent; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m² per dose for a maximum of 18 doses.

Note: Indications marked with * are unapproved indications.

Initial application — (desensisation prior to transplant) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires desensitisation prior to mismatched allogenic stem cell transplant*; and
- 2 Patient would receive no more than two doses at 375 mg/m2 of body-surface area.

Note: Indications marked with * are unapproved indications.

Initial application — (pemiphigus*) only from a dermatologist or relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 Patient has severe rapidly progressive pemphigus; and
 - 1.2 Is used in combination with systemic corticosteroids (20 mg/day); and
 - 1.3 Any of the following:
 - 1.3.1 Skin involvement is at least 5% body surface area; or
 - 1.3.2 Significant mucosal involvement (10 or more mucosal erosions) or diffuse gingivitis or confluent large erosions; or
 - 1.3.3 Involvement of two or more mucosal sites; or
- 2 Both:
 - 2.1 Patient has pemphigus; and
 - 2.2 Patient has not experienced adequate clinical benefit from systemic corticosteroids (20 mg/day) in combination with a steroid sparing agent, unless contraindicated.

Note: Indications marked with * are unapproved indications.

Renewal — (pemiphigus*) only from a dermatologist or relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has experienced adequate clinical benefit from rituximab treatment, with improvement in symptoms and healing of skin ulceration and reduction in corticosteroid requirement; and
- 2 Patient has not received rituximab in the previous 6 months.

Note: Indications marked with * are unapproved indications.

Initial application — (immunoglobulin G4-related disease (IgG4-RD*)) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed diagnosis of IgG4-RD*; and
- 2 Either:
 - 2.1 Treatment with corticosteroids and/or disease modifying anti-rheumatic drugs for at least 3 months has been ineffective in lowering corticosteroid dose below 5 mg per day (prednisone equivalent) without relapse; or

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic	
\$	Per	1	Manufacturer	

continued...

- 2.2 Treatment with corticosteroids and/or disease modifying anti-rheumatic drugs is contraindicated or associated with evidence of toxicity or intolerance; and
- 3 Total rituximab dose used should not exceed a maximum of two 1000 mg infusions of rituximab given two weeks apart. Note: Indications marked with * are unapproved indications.

Renewal — (immunoglobulin G4-related disease (IgG4-RD*)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Treatment with rituximab for IgG4-RD* was previously successful and patient's disease has demonstrated sustained response, but the condition has relapsed; or
 - 1.2 Patient is receiving maintenance treatment for IgG4-RD*; and
- 2 Rituximab re-treatment not to be given within 6 months of previous course of treatment; and
- 3 Maximum of two 1000 mg infusions of rituximab given two weeks apart.

Note: Indications marked with * are unapproved indications.

SECUKINUMAB - Special Authority see SA2084 below - Retail pharmacy

Inj 150 mg per ml, 1 ml prefilled syringe.......799.50 1 **✓ Cosentyx** 1,599.00 2 **✓ Cosentyx**

⇒SA2084 Special Authority for Subsidy

Initial application — (severe chronic plaque psoriasis – second-line biologic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab or etanercept, or has trialled infliximab in a Health NZ Hospital, for severe chronic plaque psoriasis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
 - 2.2 The patient has received insufficient benefit from adalimumab, etanercept or infliximab; and
- 3 A Psoriasis Area and Severity Index (PASI) assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Initial application — (severe chronic plaque psoriasis – first-line biologic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
- 2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin: and
- 3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Note: A treatment course is defined as a minimum of 12 weeks of treatment. "Inadequate response" is defined as: for whole

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom sub scores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis – first and second-line biologic) only from a dermatologist or medical practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Patient's PASI score has reduced by 75% or more (PASI 75) as compared to baseline PASI prior to commencing secukinumab: or
 - 1.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing secukinumab; and
- 2 Secukinumab to be administered at a maximum dose of 300 mg monthly.

Initial application — (ankylosing spondylitis – second-line biologic) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and
- 2 Fither
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

Renewal — (ankylosing spondylitis – second-line biologic) only from a rheumatologist or medical practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Following 12 weeks initial treatment of secukinumab treatment, BASDAI has improved by 4 or more points from pre-secukinumab baseline on a 10 point scale, or by 50%, whichever is less: and
- 2 Physician considers that the patient has benefitted from treatment and that continued treatment is appropriate; and
- 3 Secukinumab to be administered at doses no greater than 150 mg monthly.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has had an initial Special Authority approval for adalimumab, etanercept or infliximab for psoriatic arthritis; and
 - 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
 - 1.2.2 Patient has received insufficient benefit from adalimumab, etanercept or infliximab to meet the renewal criteria for adalimumab, etanercept or infliximab for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
 - 2.4 Fither:

(Manufacturer's Price) Subsidised Generic	
\$ Per ✓ Manufacture	r

continued...

- 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
- 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 Either:

- 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior secukinumab treatment in the opinion of the treating physician; and
- 2 Secukinumab to be administered at doses no greater than 300 mg monthly.

SILTUXIMAB - Special Authority see SA1596 below - Retail pharmacy

Note: Siltuximab is to be administered at doses no gre	ater than 11 mg/kg every	3 weeks.	
Inj 100 mg vial	770.57	1	Sylvant
Inj 400 mg vial	3,082.33	1	✓ Sylvant

⇒SA1596 Special Authority for Subsidy

Initial application only from a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's Disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Siltuximab is to be administered at doses no greater than 11 mg/kg every 3 weeks.

Renewal only from a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has sustained improvement in inflammatory markers and functional status.

TIXAGEVIMAB WITH CILGAVIMAB - [Xpharm] - Subsidy by endorsement

- a) No patient co-payment payable
- b) Treatment is funded only if patient meets access criteria for tixagevimab with cilgavimab (as per https://pharmac.govt.nz/Evusheld) and has been endorsed accordingly by the prescriber. The supply of treatment is via Pharmac's approved distribution process. Refer to the Pharmac website for more information about this and stock availability.

Inj 100 mg per ml, 1.5 ml vial with cilgavimab 100 mg per ml, 1.5 ml vial	0.00	1	✓ Evusheld
TOCILIZUMAB - PCT only - Special Authority see SA2332 or	the next page		
Inj 20 mg per ml, 4 ml vial	220.00	1	✓ Actemra
Inj 20 mg per ml, 10 ml vial		1	✓ Actemra
Inj 20 mg per ml, 20 ml vial		1	✓ Actemra
Inj 1 mg for ECP		1 mg	✓ Baxter

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA2332 Special Authority for Subsidy

Initial application — (cytokine release syndrome) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient is enrolled in the Children's Oncology Group AALL1731 trial; and
 - 1.2 The patient has developed grade 3 or 4 cytokine release syndrome associated with the administration of blinatumomab for the treatment of acute lymphoblastic leukaemia; and
 - 1.3 Tocilizumab is to be administered at doses no greater than 8 mg/kg IV for a maximum of 3 doses (if less than 30kg, maximum of 12 mg/kg); or
- 2 All of the following:
 - 2.1 The patient is enrolled in the Malaghan Institute of Medical Research ENABLE trial programme; and
 - 2.2 The patient has developed CRS or Immune Effector Cell-Associated Neurotoxicity Syndrome (ICANS) following CAR T-cell therapy for the treatment of relapsed or refractory B-cell non-Hodgkin lymphoma; and
 - 2.3 Tocilizumab is to be administered according to the consensus guidelines for CRS or ICANS for CAR T-cell therapy at doses no greater than 8 mg/kg IV for a maximum of 3 doses.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient was being treated with tocilizumab prior to 1 February 2019; and
- 2 Any of the following:
 - 2.1 rheumatoid arthritis; or
 - 2.2 systemic juvenile idiopathic arthritis: or
 - 2.3 adult-onset Still's disease; or
 - 2.4 polyarticular juvenile idiopathic arthritis; or
 - 2.5 idiopathic multicentric Castleman's disease.

Initial application — (Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Fither:
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
- 3 Either:
 - 3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or
 - 3.2 Both:
 - 3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a Health NZ Hospital; and
 - 3.2.2 Fither:
 - 3.2.2.1 The patient has experienced intolerable side effects from rituximab; or
 - 3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Initial application — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:
All of the following:

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 2 Tocilizumab is to be used as monotherapy; and
- 3 Fither:
 - 3.1 Treatment with methotrexate is contraindicated; or
 - 3.2 Patient has tried and did not tolerate oral and/or parenteral methotrexate; and

4 Either:

- 4.1 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of ciclosporin alone or in combination with another agent; or
- 4.2 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of leflunomide alone or in combination with another agent; and
- 5 Either:
 - 5.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 active, swollen, tender joints; or
 - 5.2 Patient has persistent symptoms of poorly controlled and active disease in at least four active joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 6 Fither:
 - 6.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 6.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Initial application — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient diagnosed with systemic juvenile idiopathic arthritis; and
- 2 Patient has tried and not responded to a reasonable trial of all of the following, either alone or in combination: oral or parenteral methotrexate; non-steroidal anti-inflammatory drugs (NSAIDs); and systemic corticosteroids.

Initial application — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Either:
 - 1.1.1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a Health NZ Hospital; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal antiinflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Initial application — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: Fither:

continued...

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for both etanercept and adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 The patient has experienced intolerable side effects, or has received insufficient benefit from, both etanercept and adalimumab; or
- 2 All of the following:
 - 2.1 Treatment with a tumour necrosis factor alpha inhibitor is contraindicated; and
 - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
 - 2.3 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.4 Any of the following:
 - 2.4.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.4.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.4.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Initial application — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Tocilizumab to be administered at doses no greater than 8 mg/kg IV every 3-4 weeks.

Initial application — (moderate to severe COVID-19) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed (or probable) COVID-19: and
- 2 Oxygen saturation of < 92% on room air, or requiring supplemental oxygen; and
- 3 Patient is receiving adjunct systemic corticosteroids, or systemic corticosteroids are contraindicated; and
- 4 Tocilizumab is to be administered at doses no greater than 8mg/kg IV for a maximum of one dose; and
- 5 Tocilizumab is not to be administered in combination with barcitinib.

Renewal — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Renewal — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Following up to 6 months' initial treatment, the patient has achieved at least an American College of Rheumatology paediatric 30% improvement criteria (ACR Pedi 30) response from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing ACR Pedi 30 response from baseline.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months where the patient has a sustained improvement in inflammatory markers and functional status.

Renewal — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a

Subsidy	;	Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
	Per	✓	Manufacturer

continued...

rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Renewal — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has a sustained improvement in inflammatory markers and functional status.

TRASTUZUMAB (HERZUMA) - PCT only - Special Authority see SA2293 below

Inj 150 mg vial100.00	1	Herzuma
Inj 440 mg vial	1	✓ Herzuma
Inj 1 mg for ECP	1 mg	✓ Baxter

⇒SA2293 Special Authority for Subsidy

Initial application — (early breast cancer) from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria:

Both:

- 1 The patient has early breast cancer expressing HER-2 IHC 3+ or ISH + (including FISH or other current technology); and
- 2 Maximum cumulative dose of 106 mg/kg (12 months' treatment).

Renewal — (early breast cancer*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
 - 1.2 The patient received prior adjuvant trastuzumab treatment for early breast cancer; and
 - 1.3 Any of the following:
 - 1.3.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
 - 1.3.2 The patient discontinued lapatinib within 3 months due to intolerable side effects and the cancer did not progress whilst on lapatinib; or
 - 1.3.3 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
 - 1.4 Either:
 - 1.4.1 Trastuzumab will not be given in combination with pertuzumab; or
 - 1.4.2 All of the following:
 - 1.4.2.1 Trastuzumab to be administered in combination with pertuzumab; and
 - 1.4.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
 - 1.4.2.3 The patient has good performance status (ECOG grade 0-1); and
 - 1.5 Trastuzumab to be discontinued at disease progression; or
- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with trastuzumab in the metastatic setting for reasons other than severe toxicity or disease progression; and

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 2.2 Patient has signs of disease progression; and
- 2.3 Disease has not progressed during previous treatment with trastuzumab.

Note: * For patients with relapsed HER-2 positive disease who have previously received adjuvant trastuzumab for early breast cancer

Initial application — (metastatic breast cancer) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Fither:
 - 2.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
 - 2.2 The patient discontinued lapatinib within 3 months due to intolerable side effects and the cancer did not progress whilst on lapatinib; and
- 3 Fither:
 - 3.1 Trastuzumab will not be given in combination with pertuzumab; or
 - 3.2 All of the following:
 - 3.2.1 Trastuzumab to be administered in combination with pertuzumab; and
 - 3.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer: and
 - 3.2.3 The patient has good performance status (ECOG grade 0-1); and
- 4 Trastuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
 - 1.2 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
 - 1.3 Trastuzumab to be discontinued at disease progression; or
- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with trastuzumab for reasons other than severe toxicity or disease progression; and
 - 2.2 Patient has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with trastuzumab.

Initial application — (gastric, gastro-oesophageal junction and oesophageal cancer) from any relevant practitioner.

Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has locally advanced or metastatic gastric, gastro-oesophageal junction or oesophageal cancer expressing HER-2 IHC 2+ FISH+ or IHC3+ (or other current technology); and
- 2 Patient has an ECOG score of 0-2.

Renewal — (gastric, gastro-oesophageal junction and oesophageal cancer) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Roth:

- 1 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 2 Trastuzumab to be discontinued at disease progression.

	Subsidy (Manufacturer's Price)	Sub Per	Fully sidised	Brand or Generic Manufacturer	
TRASTUZUMAB EMTANSINE - PCT only - Specialist - Specia	J Authority see SA21/			Manuacturer	
Inj 100 mg vial	•	1	✓ Ka	adcyla	
Inj 160 mg vial	,	1		adcyla	
Inj 1 mg for ECP	24.52	1 mg	✓ Ba	axter	

⇒SA2144 Special Authority for Subsidy

Initial application — (early breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has early breast cancer expressing HER2 IHC3+ or ISH+; and
- 2 Documentation of pathological invasive residual disease in the breast and/or auxiliary lymph nodes following completion of surgery; and
- 3 Patient has completed systemic neoadjuvant therapy with trastuzumab and chemotherapy prior to surgery; and
- 4 Disease has not progressed during neoadjuvant therapy; and
- 5 Patient has left ventricular ejection fraction of 45% or greater; and
- 6 Adjuvant treatment with trastuzumab emtansine to be commenced within 12 weeks of surgery; and
- 7 Trastuzumab emtansine to be discontinued at disease progression; and
- 8 Total adjuvant treatment duration must not exceed 42 weeks (14 cycles).

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Patient has previously received trastuzumab and chemotherapy, separately or in combination; and
- 3 Either:
 - 3.1 The patient has received prior therapy for metastatic disease*; or
 - 3.2 The patient developed disease recurrence during, or within six months of completing adjuvant therapy*; and
- 4 Patient has a good performance status (ECOG 0-1); and
- 5 Either:
 - 5.1 Patient does not have symptomatic brain metastases; or
 - 5.2 Patient has brain metastases and has received prior local CNS therapy; and
- 6 Patient has not received prior funded trastuzumab emtansine treatment; and
- 7 Treatment to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The cancer has not progressed at any time point during the previous approval period whilst on trastuzumab emtansine; and
- 2 Treatment to be discontinued at disease progression.

Note: Prior or adjuvant therapy includes anthracycline, other chemotherapy, biological drugs, or endocrine therapy.

USTEKINUMAB - Special Authority see SA2182 below - Retail pharmacy

Inj 90 mg per ml, 1 ml pre-filled syringe........................4,162.00 1 ✓ Stelara

⇒SA2182 Special Authority for Subsidy

Initial application — (**Crohn's disease - adults**) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Patient is currently on treatment with ustekinumab commenced prior to 1 February 2023 and met all remaining criteria (criterion 2) below at the time of commencing treatment; or
- 2 Both:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 2.1 Patient has active Crohn's disease; and
- 2.2 Fither:
 - 2.2.1 Patient has had an initial approval for prior biologic therapy for Crohn's disease and has experienced intolerable side effects or insufficient benefit to meet renewal criteria; or
 - 2.2.2 Both:
 - 2.2.2.1 Patient meets the initiation criteria for prior biologic therapies for Crohn's disease; and 2.2.2.2 Other biologics for Crohn's disease are contraindicated.

Renewal — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points, or HBI score has reduced by 3 points, from when the patient was initiated on biologic therapy; or
 - 1.2 CDAI score is 150 or less, or HBI is 4 or less; or
 - 1.3 The patient has experienced an adequate response to treatment, but CDAI score and/or HBI score cannot be assessed; and
- 2 Ustekinumab to be administered at a dose no greater than 90 mg every 8 weeks.

Initial application — (Crohn's disease - children*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Patient is currently on treatment with ustekinumab commenced prior to 1 February 2023 and met all remaining criteria (criterion 2) below at the time of commencing treatment; or
- 2 Both:
 - 2.1 Patient has active Crohn's disease; and
 - 2.2 Either:
 - 2.2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria: or
 - 2.2.2 Both:
 - 2.2.2.1 Patient meets the initiation criteria for prior biologic therapies for Crohn's disease; and
 - 2.2.2.2 Other biologics for Crohn's disease are contraindicated.

Note: Indication marked with * is an unapproved indication.

Renewal — (Crohn's disease - children*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from when the patient was initiated on biologic therapy; or
 - 1.2 PCDAI score is 15 or less: or
 - 1.3 The patient has experienced an adequate response to treatment, but CDAI score cannot be assessed; and
- 2 Ustekinumab to administered at a dose no greater than 90 mg every 8 weeks.

Note: Indication marked with * is an unapproved indication.

Initial application — (ulcerative colitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Patient is currently on treatment with ustekinumab commenced prior to 1 February 2023 and met all remaining criteria (criterion 2) below at the time of commencing treatment; or
- 2 Both:
 - 2.1 Patient has active ulcerative colitis; and

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

- 2.2 Either:
 - 2.2.1 Patient has had an initial approval for prior biologic therapy for ulcerative colitis and has experienced intolerable side effects or insufficient benefit to meet renewal criteria; or
 - 2.2.2 Both:
 - 2.2.2.1 Patient meets the initiation criteria for prior biologic therapies for ulcerative colitis; and 2.2.2.2 Other biologics for ulcerative colitis are contraindicated.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 The SCCAI score has reduced by 2 points or more from the SCCAI score since initiation on biologic therapy; or
 - 1.2 PUCAI score has reduced by 10 points or more from the PUCAI score since initiation on biologic therapy*; and
- 2 Ustekinumab will be used at a dose no greater than 90 mg intravenously every 8 weeks.

Note: Criterion marked with * is for an unapproved indication.

.00 1

✓ Entvvio

⇒SA2183 Special Authority for Subsidy

Initial application — (**Crohn's disease - adults**) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active Crohn's disease: and
- 2 Any of the following:
 - 2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria (unless contraindicated); or
 - 2.2 Patient has a CDAI score of greater than or equal to 300, or HBI score of greater than or equal to 10; or
 - 2.3 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.4 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
 - 2.5 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Any of the following:
 - 3.1 Patient has tried but experienced an inadequate response to (including lack of initial response and/or loss of initial response) from prior therapy with immunomodulators and corticosteroids; or
 - 3.2 Patient has experienced intolerable side effects from immunomodulators and corticosteroids; or
 - 3.3 Immunomodulators and corticosteroids are contraindicated.

Renewal — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points, or HBI score has reduced by 3 points, from when the patient was initiated on biologic therapy; or
 - 1.2 CDAI score is 150 or less, or HBI is 4 or less; or
 - 1.3 The patient has experienced an adequate response to treatment, but CDAI score and/or HBI score cannot be assessed; and
- 2 Vedolizumab to administered at a dose no greater than 300 mg every 8 weeks.

Initial application — (Crohn's disease - children*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Subsidy	,	Fully	Brand or	
(Manufacturer's Price	·) :	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

All of the following:

- 1 Paediatric patient has active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria (unless contraindicated); or
 - 2.2 Patient has a Paediatric Crohn's Disease Activity Index (PCDAI) score of greater than or equal to 30; or
 - 2.3 Patient has extensive small intestine disease; and
- 3 Any of the following:
 - 3.1 Patient has tried but experienced an inadequate response to (including lack of initial response and/or loss of initial response) from prior therapy with immunomodulators and corticosteroids; or
 - 3.2 Patient has experienced intolerable side effects from immunomodulators and corticosteroids; or
 - 3.3 Immunomodulators and corticosteroids are contraindicated.

Note: Indication marked with * is an unapproved indication.

Renewal — (Crohn's disease - children*) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from when the patient was initiated on biologic therapy; or
 - 1.2 PCDAI score is 15 or less; or
 - 1.3 The patient has experienced an adequate response to treatment, but CDAI score cannot be assessed; and
- 2 Vedolizumab to administered at a dose no greater than 300mg every 8 weeks.

Note: Indication marked with * is an unapproved indication.

Initial application — (ulcerative colitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active ulcerative colitis; and
- 2 Any of the following:
 - 2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria (unless contraindicated); or
 - 2.2 Patient has a SCCAI score is greater than or equal to 4; or
 - 2.3 Patient's PUCAI score is greater than or equal to 20*; and
- 3 Any of the following:
 - 3.1 Patient has tried but experienced an inadequate response to (including lack of initial response and/or loss of initial response) from prior therapy with immunomodulators and corticosteroids; or
 - 3.2 Patient has experienced intolerable side effects from immunomodulators and corticosteroids; or
 - 3.3 Immunomodulators and corticosteroids are contraindicated.

Note: Indication marked with * is an unapproved indication.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 The SCCAI score has reduced by 2 points or more from the SCCAI score since initiation on biologic therapy; or
 - 1.2 The PUCAI score has reduced by 10 points or more from the PUCAI score since initiation on biologic therapy *; and
- 2 Vedolizumab will be used at a dose no greater than 300 mg intravenously every 8 weeks.

Note: Indication marked with * is an unapproved indication.

Subsidy		Fully	Brand or
(Manufacturer's Price)	Su	ıbsidised	Generic
\$	Per	✓	Manufacturer

Programmed Cell Death-1 (PD-1) Inhibitors

ATEZOLIZUMAB - PCT only - Specialist - Special Autho	rity see SA2264 below		
Inj 60 mg per ml, 20 ml vial	9,503.00	1	Tecentriq
Inj 1 mg for ECP	8.08	1 mg	✓ Baxter

⇒SA2264 Special Authority for Subsidy

Initial application — (non-small cell lung cancer second line monotherapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced or metastatic non-small cell lung cancer; and
- 2 Patient has not received prior funded treatment with an immune checkpoint inhibitor for NSCLC; and
- 3 For patients with non-squamous histology there is documentation confirming that the disease does not express activating mutations of EGFR or ALK tyrosine kinase unless not possible to ascertain; and
- 4 Patient has an ECOG 0-2; and
- 5 Patient has documented disease progression following treatment with at least two cycles of platinum-based chemotherapy; and
- 6 Atezolizumab is to be used as monotherapy at a dose of 1200 mg every three weeks (or equivalent) for a maximum of 16 weeks; and
- 7 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Renewal — (non-small cell lung cancer second line monotherapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Any of the following:
 - 1.1 Patient's disease has had a complete response to treatment; or
 - 1.2 Patient's disease has had a partial response to treatment; or
 - 1.3 Patient has stable disease; and
- 2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period: and
- 3 No evidence of disease progression; and
- 4 The treatment remains clinically appropriate and patient is benefitting from treatment; and
- 5 Atezolizumab to be used at a maximum dose of 1200 mg every three weeks (or equivalent); and
- 6 Treatment with atezolizumab to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

DURVALUMAB - PCT only - Specialist - Special Authority see SA2164 below

		of type of the control of the contro
Imfinzi	1	Inj 50 mg per ml, 10 ml vial4,700.00
✓ Imfinzi	1	Inj 50 mg per ml, 2.4 ml vial
✓ Baxter	1 mg	Inj 1 mg for ECP

⇒SA2164 Special Authority for Subsidy

Initial application — (Non-small cell lung cancer) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 Patient has histologically or cytologically documented stage III, locally advanced, unresectable non-small cell lung cancer (NSCLC): and
- 2 Patient has received two or more cycles of platinum-based chemotherapy concurrently with definitive radiation therapy; and
- 3 Patient has no disease progression following the second or subsequent cycle of platinum-based chemotherapy with

				=
Subsidy		Fully	Brand or	
(Manufacturer's Price)	Su	bsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

definitive radiation therapy treatment; and

- 4 Patient has a ECOG performance status of 0 or 1; and
- 5 Patient has completed last radiation dose within 8 weeks of starting treatment with durvalumab; and
- 6 Patient must not have received prior PD-1 or PD-L1 inhibitor therapy for this condition; and
- 7 Either:
 - 7.1 Durvalumab is to be used at a maximum dose of no greater than 10 mg/kg every 2 weeks; or
 - 7.2 Durvalumab is to be used at a flat dose of 1500 mg every 4 weeks; and
- 8 Treatment with durvalumab to cease upon signs of disease progression.

Renewal — (Non-small cell lung cancer) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The treatment remains clinically appropriate and the patient is benefitting from treatment; and
- 2 Either
 - 2.1 Durvalumab is to be used at a maximum dose of no greater than 10 mg/kg every 2 weeks; or
 - 2.2 Durvalumab is to be used at a flat dose of 1500 mg every 4 weeks; and
- 3 Treatment with durvalumab to cease upon signs of disease progression; and
- 4 Total continuous treatment duration must not exceed 12 months.

NIVOLUMAB - PCT only - Specialist - Special Authority see SA2306 below

Inj 10 mg per ml, 4 ml vial	1,051.98	1	Opdivo
Inj 10 mg per ml, 10 ml vial	2,629.96	1	Opdivo
Inj 1 mg for ECP	27.62	1 mg	✓ Baxter

⇒SA2306 Special Authority for Subsidy

Initial application only from a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Baseline measurement of overall tumour burden is documented clinically and radiologically; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Either:
 - 4.1 Patient has not received funded pembrolizumab; or
 - 4.2 Both:
 - 4.2.1 Patient has received an initial Special Authority approval for pembrolizumab and has discontinued pembrolizumab within 12 weeks of starting treatment due to intolerance; and
 - 4.2.2 The cancer did not progress while the patient was on pembrolizumab; and
- 5 Documentation confirming that the patient has been informed and acknowledges that funded treatment with nivolumab will not be continued if their disease progresses.

Renewal — (less than 24 months on treatment) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Patient's disease has had a complete response to treatment; or
 - 1.1.2 Patient's disease has had a partial response to treatment; or
 - 1.1.3 Patient has stable disease; and
 - 1.2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period: and
 - 1.3 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or

Subsidy		Fully	Brand or
(Manufacturer's Price)		idised	Generic
\$	Per	1	Manufacturer

continued...

- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with nivolumab for reasons other than severe toxicity or disease progression; and
 - 2.2 Patient has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with nivolumab.

Renewal — (more than 24 months on treatment) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 Patient has been on treatment for more than 24 months; and
- 2 Either:
 - 2.1 All of the following:
 - 2.1.1 Any of the following:
 - 2.1.1.1 Patient's disease has had a complete response to treatment; or
 - 2.1.1.2 Patient's disease has had a partial response to treatment; or
 - 2.1.1.3 Patient has stable disease; and
 - 2.1.2 Response to treatment in target lesions has been determined by comparable radiologic or clinical assessment following the most recent treatment period; and
 - 2.1.3 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
 - 2.2 All of the following:
 - 2.2.1 Patient has previously discontinued treatment with nivolumab for reasons other than severe toxicity or disease progression; and
 - 2.2.2 Patient has signs of disease progression; and
 - 2.2.3 Disease has not progressed during previous treatment with nivolumab.

Inj 25 mg per ml, 4 ml vial	4,680.00	1	Keytruda
Inj 1 mg for ECP	47.74	1 mg	✓ Baxter

⇒SA2307 Special Authority for Subsidy

Initial application — (unresectable or metastatic melanoma) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Baseline measurement of overall tumour burden is documented clinically and radiologically; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Fither:
 - 4.1 Patient has not received funded nivolumab; or
 - 4.2 Both:
 - 4.2.1 Patient has received an initial Special Authority approval for nivolumab and has discontinued nivolumab within 12 weeks of starting treatment due to intolerance; and
 - 4.2.2 The cancer did not progress while the patient was on nivolumab; and
- 5 Documentation confirming that the patient has been informed and acknowledges that funded treatment with pembrolizumab will not be continued if their disease progresses.

Renewal — (unresectable or metastatic melanoma, less than 24 months on treatment) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

1 All of the following:

Subsidy (Manufacturer's Price)	s	Fully	Brand or Generic
 \$	Per	1	Manufacturer

continued...

- 1.1 Any of the following:
 - 1.1.1 Patient's disease has had a complete response to treatment; or
 - 1.1.2 Patient's disease has had a partial response to treatment; or
 - 1.1.3 Patient has stable disease; and
- 1.2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period; and
- 1.3 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with pembrolizumab for reasons other than severe toxicity or disease progression: and
 - 2.2 Patient has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with pembrolizumab.

Renewal — (unresectable or metastatic melanoma, more than 24 months on treatment) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 Patient has been on treatment for more than 24 months; and
- - 2.1 All of the following:
 - 2.1.1 Any of the following:
 - 2.1.1.1 Patient's disease has had a complete response to treatment; or
 - 2.1.1.2 Patient's disease has had a partial response to treatment; or
 - 2.1.1.3 Patient has stable disease; and
 - 2.1.2 Response to treatment in target lesions has been determined by comparable radiologic or clinical assessment following the most recent treatment period; and
 - 2.1.3 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
 - 2.2 All of the following:
 - 2.2.1 Patient has previously discontinued treatment with pembrolizumab for reasons other than severe toxicity or disease progression; and
 - 2.2.2 Patient has signs of disease progression; and
 - 2.2.3 Disease has not progressed during previous treatment with pembrolizumab.

Initial application — (non-small cell lung cancer first-line monotherapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-small cell lung cancer; and
- 2 Patient has not had chemotherapy for their disease in the palliative setting; and
- 3 Patient has not received prior funded treatment with an immune checkpoint inhibitor for NSCLC; and
- 4 For patients with non-squamous histology there is documentation confirming that the disease does not express activating mutations of EGFR or ALK tyrosine kinase unless not possible to ascertain; and
- 5 Pembrolizumab to be used as monotherapy; and
- 6 Either:
 - 6.1 There is documentation confirming the disease expresses PD-L1 at a level greater than or equal to 50% as determined by a validated test unless not possible to ascertain; or
 - 6.2 Both:
 - 6.2.1 There is documentation confirming the disease expresses PD-L1 at a level greater than or equal to 1% as determined by a validated test unless not possible to ascertain; and

Subsidy (Manufacturer's		Fully lised	Brand or Generic	
\$	Per	•	Manufacturer	

continued...

- 6.2.2 Chemotherapy is determined to be not in the best interest of the patient based on clinician assessment; and
- 7 Patient has an ECOG 0-2; and
- 8 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks; and
- 9 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Renewal — (non-small cell lung cancer first line monotherapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 Patient's disease has had a complete response to treatment; or
 - 1.2 Patient's disease has had a partial response to treatment; or
 - 1.3 Patient has stable disease; and
- 2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period: and
- 3 No evidence of disease progression; and
- 4 The treatment remains clinically appropriate and patient is benefitting from treatment; and
- 5 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 6 Treatment with pembrolizumab to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initial application — (non-small cell lung cancer first-line combination therapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-small cell lung cancer; and
- 2 The patient has not had chemotherapy for their disease in the palliative setting; and
- 3 Patient has not received prior funded treatment with an immune checkpoint inhibitor for NSCLC; and
- 4 For patients with non-squamous histology there is documentation confirming that the disease does not express activating mutations of EGFR or ALK tyrosine kinase unless not possible to ascertain; and
- 5 Pembrolizumab to be used in combination with platinum-based chemotherapy; and
- 6 Patient has an ECOG 0-2: and
- 7 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks; and
- 8 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Renewal — (non-small cell lung cancer first line combination therapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 Patient's disease has had a complete response to treatment; or
 - 1.2 Patient's disease has had a partial response to treatment: or
 - 1.3 Patient has stable disease; and
- 2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period; and
- 3 No evidence of disease progression; and
- 4 The treatment remains clinically appropriate and patient is benefitting from treatment; and
- 5 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 6 Treatment with pembrolizumab to cease after a total duration of 24 months from commencement (or equivalent of

Subsidy		Fully	Brand or	
(Manufacturer's Price)	9	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

35 cycles dosed every 3 weeks).

Other Immunosuppressants

CICLOSPORIN

Cap 25 mg	44.63	50	Neoral
Cap 50 mg	88.91	50	✓ Neoral
Cap 100 mg	177.81	50	✓ Neoral
Oral liq 100 mg per ml		50 ml OP	✓ Neoral

EVEROLIMUS - Special Authority see \$A2008 below - Retail pharmacy

Wastage claimable 30 ✓ Afinitor Tab 5 mg4,555.76 30 ✓ Afinitor

⇒SA2008 Special Authority for Subsidy

Initial application only from a neurologist or oncologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has tuberous sclerosis: and
- 2 Patient has progressively enlarging sub-ependymal giant cell astrocytomas (SEGAs) that require treatment.

Renewal only from a neurologist or oncologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Documented evidence of SEGA reduction or stabilisation by MRI within the last 3 months; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Everolimus to be discontinued at progression of SEGAs.

SIROLIMUS - Special Authority see SA2270 below - Retail pharmacy

Tab 1 mg	749.99	100	Rapamune
Tab 2 mg	1,499.99	100	 Rapamune
Oral liq 1 mg per ml	449.99	60 ml OP	✓ Rapamune

⇒SA2270 Special Authority for Subsidy

Initial application from any medical practitioner. Approvals valid without further renewal unless notified where the drug is to be used for rescue therapy for an organ transplant recipient.

Notes: Rescue therapy defined as unresponsive to calcineurin inhibitor treatment as defined by refractory rejection; or intolerant to calcineurin inhibitor treatment due to any of the following:

- GFR< 30 ml/min; or
- · Rapidly progressive transplant vasculopathy; or
- Rapidly progressive obstructive bronchiolitis; or
- . HUS or TTP: or
- · Leukoencepthalopathy; or
- · Significant malignant disease

Initial application — (severe non-malignant lymphovascular malformations*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe non-malignant lymphovascular malformation*; and
- 2 Any of the following:
 - 2.1 Malformations are not adequately controlled by sclerotherapy and surgery; or
 - 2.2 Malformations are widespread/extensive and sclerotherapy and surgery are not considered clinically appropriate; or

Subsidy	Fully	Brand or	Т
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

- 2.3 Sirolimus is to be used to reduce malformation prior to consideration of surgery; and
- 3 Patient is being treated by a specialist lymphovascular malformation multi-disciplinary team; and
- 4 Patient has measurable disease as defined by RECIST version 1.1 (see Note).

Renewal — (severe non-malignant lymphovascular malformations*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Patient's disease has had either a complete response or a partial response to treatment, or patient has stable disease according to RECIST version 1.1 (see Note); or
 - 1.2 Patient's disease has stabilised or responded clinically and disease response to treatment has been clearly documents in patient notes; and
- 2 No evidence of progressive disease; and
- 3 The treatment remains clinically appropriate and the patient is benefitting from the treatment.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer et al. Eur J Cancer 2009;45:228-47)

Indications marked with * are unapproved indications

Initial application — (renal angiomyolipoma(s) associated with tuberous sclerosis complex*) only from a nephrologist or urologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has tuberous sclerosis complex*; and
- 2 Evidence of renal angiomyolipoma(s) measuring 3 cm or greater and that have shown interval growth.

Renewal — (renal angiomyolipoma(s) associated with tuberous sclerosis complex*) from any relevant practitioner.

Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Documented evidence of renal angiomyolipoma reduction or stability by magnetic resonance imaging (MRI) or ultrasound;
- 2 Demonstrated stabilisation or improvement in renal function; and
- 3 The patient has not experienced angiomyolipoma haemorrhage or significant adverse effects to sirolimus treatment; and
- 4 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indications marked with * are unapproved indications

Initial application — (refractory seizures associated with tuberous sclerosis complex*) only from a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has epilepsy with a background of documented tuberous sclerosis complex; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Vigabatrin has been trialled and has not adequately controlled seizures; and
 - 2.1.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least two of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); or
 - 2.2 Both:
 - 2.2.1 Vigabatrin is contraindicated; and
 - 2.2.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least three of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); and
- 3 Seizures have a significant impact on quality of life; and

Subsidy	F	ully	Brand or	
(Manufacturer's Price)	Subsidis	sed	Generic	
\$	Per	✓	Manufacturer	

continued...

4 Patient has been assessed and surgery is considered inappropriate for this patient, or the patient has been assessed and would benefit from mTOR inhibitor treatment prior to surgery.

Note: Those of childbearing age potential are not required to trial phenytoin sodium, sodium valproate, or topiramate. Those who can father children are not required to trial sodium valproate.

Renewal — (refractory seizures associated with tuberous sclerosis complex*) only from a neurologist. Approvals valid for 12 months where demonstrated significant and sustained improvement in seizure rate (e.g. 50% reduction in seizure frequency) or severity and/or patient quality of life compared with baseline prior to starting sirolimus treatment.

Note: Indications marked with * are unapproved indications

TACROLIMUS - Special Authority see SA2271 below - Retail pharmacy

Cap 0.5 mg	100	✓ Tacrolimus Sandoz
Cap 0.75 mg	100	✓ Tacrolimus Sandoz
Cap 1 mg	100	✓ Tacrolimus Sandoz
Cap 5 mg248.20	50	✓ Tacrolimus Sandoz

⇒SA2271 Special Authority for Subsidy

Initial application — (organ transplant) only from a relevant specialist. Approvals valid without further renewal unless notified where the patient is an organ transplant recipient.

Note: Subsidy applies for either primary or rescue therapy.

Initial application — (non-transplant indications*) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient requires long-term systemic immunosuppression; and
- 2 Either:
 - 2.1 Ciclosporin has been trialled and discontinued treatment because of unacceptable side effects or inadequate clinical response; or
 - 2.2 Patient is a child with nephrotic syndrome*.

Note: Indications marked with * are unapproved indications

JAK inhibitors

⇒SA2079 Special Authority for Subsidy

Initial application — (Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
- 3 Fither:
 - 3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or
 - 3.2 Both:
 - $3.2.1 \ \ The \ patient \ has \ been \ started \ on \ rituximab \ for \ rheumatoid \ arthritis \ in \ a \ Health \ NZ \ Hospital; \ and$

3.2.2 Either:

continued...

- 3.2.2.1 The patient has experienced intolerable side effects from rituximab; or
- 3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Renewal — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Either:

- 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

	Subsidy		Fully	Brand or
(Manufacturer's Price)	5	Subsidised	Generic
	\$	Per	/	Manufacturer

Antiallergy Preparations

Allergic Emergencies

ADRENALINE - Special Authority see SA2185 below - Retail pharmacy

- a) Maximum of 2 inj per prescription
- Additional prescriptions limited to replacement of up to two devices prior to expiry, or replacement of used device for treatment of anaphylaxis.

Inj 0.15 mg per 0.3 ml auto-injector	90.00	1 OP	 Epipen Jr
Inj 0.3 mg per 0.3 ml auto-injector	90.00	1 OP	Epipen

⇒SA2185 Special Authority for Subsidy

Initial application — (anaphylaxis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Patient has experienced an anaphylactic reaction which has resulted in presentation to a hospital or emergency department; or
 - 1.2 Patient has been assessed to be at significant risk of anaphylaxis by a relevant practitioner; and
- 2 Patient is not to be prescribed more than two devices in initial prescription.

⇒SA1558 Special Authority for Subsidy

Initial application only from a clinical immunologist or relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Supply for anticipated emergency treatment of laryngeal/oro-pharyngeal or severe abdominal attacks of acute hereditary angioedema (HAE) for patients with confirmed diagnosis of C1-esterase inhibitor deficiency; and
- 2 The patient has undergone product training and has agreed upon an action plan for self-administration.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Allergy Desensitisation

⇒SA1367 Special Authority for Subsidy

Initial application only from a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 RAST or skin test positive; and
- 2 Patient has had severe generalised reaction to the sensitising agent.

Renewal only from a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

BEE VENOM ALLERGY TREATMENT - Special Authority see SA1367 above - Retail pharmacy Initiation kit - 5 vials freeze dried venom with diluent 305.00 1 OP ✓ VENOX S29 Maintenance kit - 1 vial freeze dried venom with diluent................305.00 1 OP ✓ VENOX S29 Maintenance kit - 6 vials 120 mcg freeze dried venom, with 1 OP ✓ Venomil S29 Treatment kit - 1 vial 550 mcg freeze dried venom, 1 diluent 9 ml, 3 diluent 1.8 ml305.00 1 OP ✓ Albev Treatment kit - 1 vial 550 mcg freeze dried venom, with diluent 305.00 1 OP ✓ Hymenoptera S29

	Subsidy		Fully Brand or
	(Manufacturer's Price)) Subsid	dised Generic ✓ Manufacturer
WASP VENOM ALLERGY TREATMENT - Special Authority see	e SA1367 on the pre	vious page -	 Retail pharmacy
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze			4
dried polister venom, 1 diluent 9 ml, 1 diluent 1.8 ml	305.00	1 OP	✓ Albey
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze			
dried venom, with diluent	305.00	1 OP	✓ Hymenoptera S29
Treatment kit (Paper wasp venom) - 6 vials 120 mcg freeze	005.00	4.00	(V !!
dried venom, with diluent		1 OP	✓ Venomil S29
Treatment kit (Yellow Jacket venom) - 1 vial 550 mcg freeze		1 OD	/ Uhumamambana 000
dried venom, with diluent	305.00	1 OP	✓ Hymenoptera S29
Treatment kit (Yellow jacket venom) - 1 vial 550 mcg freeze	205.00	1 OD	Albay
dried vespula venom, 1 diluent 9 ml, 1 diluent 1.8 ml		1 OP	✓ Albey
Treatment kit (Yellow jacket venom) - 6 vials 120 mcg freeze dried venom, with diluent		1 OP	✓ Venomil S29
diled verioni, with dildent	303.00	TOP	Venonin 329
Antihistamines			
CETIRIZINE HYDROCHLORIDE			4
* Tab 10 mg		100	✓ <u>Zista</u>
* Oral liq 1 mg per ml	2.84	200 ml	✓ <u>Histaclear</u>
CHLORPHENIRAMINE MALEATE			
* Oral liq 2 mg per 5 ml	9.37	500 ml	✓ Histafen
(Histafen Oral liq 2 mg per 5 ml to be delisted 1 August 2024)			
DEXTROCHLORPHENIRAMINE MALEATE			
* Tab 2 mg		40	B
	(8.40)	00	Polaramine
	1.01	20	Polaramine
* Oral liq 2 mg per 5 ml	(5.99)	100 ml	Polaramine
本 Oral liq 2 mg per 5 mi	(10.29)	100 1111	Polaramine
EEVOEENIA DINIE LIVODOCLII ODIDE	(10.23)		1 Olaramino
FEXOFENADINE HYDROCHLORIDE * Tab 60 mg	4.24	20	
* Tab 60 mg	(8.23)	20	Telfast
* Tab 120 mg		10	Tellast
7 Tub 120 Hg	(8.23)	10	Telfast
	14.22	30	Tondot
	(26.44)		Telfast
LORATADINE	, ,		
* Tab 10 mg	1.78	100	✓ Lorafix
* Oral liq 1 mg per ml		100 ml	✓ Haylor syrup
PROMETHAZINE HYDROCHLORIDE			
* Tab 10 mg	1.39	50	✓ Allersoothe
* Tab 25 mg		50	✓ Allersoothe
* Oral liq 1 mg per 1 ml	3.39	100 ml	✓ Allersoothe
* Inj 25 mg per ml, 2 ml ampoule - Up to 5 inj available on a f	PSO21.09	5	✓ Hospira

Powder for inhalation, 50 mcg per dose, breath activated26.25

	\$	Per	✓ Manufacturer
Inhaled Corticosteroids			
BECLOMETHASONE DIPROPIONATE	44.04	000 de - 00	4.00
Aerosol inhaler, 50 mcg per dose		200 dose OP	✓ Qvar
Aerosol inhaler, 50 mcg per dose CFC-free		200 dose OP	✓ Beclazone 50
Aerosol inhaler, 100 mcg per dose		200 dose OP	✓ Qvar
Aerosol inhaler, 100 mcg per dose CFC-free		200 dose OP	✓ Beclazone 100
Aerosol inhaler, 250 mcg per dose CFC-free	22.67	200 dose OP	✓ Beclazone 250
BUDESONIDE			
Powder for inhalation, 100 mcg per dose	17.00	200 dose OP	✓ Pulmicort
φ _γ			Turbuhaler
Powder for inhalation, 200 mcg per dose	19 00	200 dose OP	✓ Pulmicort
Total for analysis, 200 mag per doco		200 0000 0.	Turbuhaler
Powder for inhalation, 400 mcg per dose	22.00	200 dose OP	✓ Pulmicort
1 Owder for initial ation, 400 they per dose	02.00	200 0036 OI	Turbuhaler
			Turbunaler
FLUTICASONE			_
Aerosol inhaler, 50 mcg per dose		120 dose OP	✓ Flixotide
Powder for inhalation, 50 mcg per dose		60 dose OP	 Flixotide Accuhaler
Powder for inhalation, 100 mcg per dose	7.81	60 dose OP	 Flixotide Accuhaler
Aerosol inhaler, 125 mcg per dose	13.60	120 dose OP	✓ Flixotide
Aerosol inhaler, 250 mcg per dose	24.62	120 dose OP	✓ Flixotide
Powder for inhalation, 250 mcg per dose	11.93	60 dose OP	 Flixotide Accuhaler
Inhaled Long-acting Beta-adrenoceptor Agonist	S		
EFORMOTEROL FUMARATE DIHYDRATE			
Powder for inhalation 4.5 mcg per dose, breath activated			
(equivalent to eformoterol fumarate 6 mcg metered dose	10.32	60 dose OP	
(equivalent to distinction familiate of mag meterod dose	(16.90)	00 0000 01	Oxis Turbuhaler
INDAGATEDOL	(10.00)		Onio i dibulialoi
INDACATEROL 150	04.00	00 1 05	401 D 11
Powder for inhalation 150 mcg		30 dose OP	✓ Onbrez Breezhaler
Powder for inhalation 300 mcg	61.00	30 dose OP	 Onbrez Breezhaler
SALMETEROL			

Subsidy

(Manufacturer's Price)

Fully

Subsidised

Brand or

Generic

120 dose OP

60 dose OP

✓ Serevent✓ Serevent Accuhaler

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

Inhaled Corticosteroids with Long-Acting Beta-Adrenoceptor Agonists

,		
BUDESONIDE WITH EFORMOTEROL		
Powder for inhalation 160 mcg with 4.5 mcg eformoterol		
fumarate per dose (equivalent to 200 mcg budesonide with		45 5 6 1
6 mcg eformoterol fumarate metered dose)41.50	120 dose OP	DuoResp Spiromax
Powder for inhalation 320 mcg with 9 mcg eformoterol fumarate		
per dose (equivalent to 400 mcg budesonide with 12 mcg		
eformoterol fumarate metered dose) – No more than 2	100 doos OD	✓ DueDeen Cuiremey
dose per day	120 dose OP	✓ DuoResp Spiromax ✓ Vannair
Aerosol inhaler 100 mcg with eformoterol fumarate 6 mcg18.23 Powder for inhalation 100 mcg with eformoterol fumarate 6 mcg33.74	120 dose OP	• • • • • • • • • • • • • • • • • • • •
Powder for initial attorn 100 micg with elornioteror furnarate 6 micg55.74	120 dose OP	✓ Symbicort Turbuhaler 100/6
Aerosol inhaler 200 mcg with eformoterol fumarate 6 mcg21.40	120 dose OP	✓ Vannair
Powder for inhalation 200 mcg with eformoterol furnarate 6 mcg33.74	120 dose OP	✓ Symbicort
1 owder for initialation 200 mag with elormoteror furnarate of mag35.74	120 0036 OI	Turbuhaler 200/6
Powder for inhalation 400 mcg with eformoterol fumarate		Turbunuici 200/0
12 mcg - No more than 2 dose per day	60 dose OP	✓ Symbicort
72 mag 140 mare than 2 dood por day	00 0000 01	Turbuhaler 400/12
FI UTICASONE FUROATE WITH VII ANTEROI		
Powder for inhalation 100 mcg with vilanterol 25 mcg44.08	30 dose OP	✓ Breo Ellipta
c c	00 d030 O1	V Dico Lilipia
FLUTICASONE WITH SALMETEROL According to the large with salmeteral 25 mag. 25 70.	120 dose OP	✓ Seretide
Aerosol inhaler 50 mcg with salmeterol 25 mcg	120 dose OP	✓ Seretide
Powder for inhalation 100 mcg with salmeterol 50 mcg – No	120 005e OF	• Seletide
more than 2 dose per day33.74	60 dose OP	✓ Seretide Accuhaler
Powder for inhalation 250 mcg with salmeterol 50 mcg – No	00 003 0 01	• Seletide Acculiatei
ů ů	60 dose OP	✓ Seretide Accuhaler
more than 2 dose per day44.08	00 0036 01	- Oci clide Acculiatei

Beta-Adre	TO COMPANY	A maniata
19121610140142	181016121011014	/^\0 0 0

SALBUTAMOL			
Oral liq 400 mcg per ml	40.00	150 ml	✓ Ventolin
Infusion 1 mg per ml, 5 ml	130.00	10	✓ Ventolin
Inj 500 mcg per ml, 1 ml - Up to 5 inj available on a PSO	130.00	5	✓ Ventolin

	Subsidy		Fully	Brand or
	(Manufacturer's		sidised	Generic
	\$	Per		Manufacturer
Inhaled Beta-Adrenoceptor Agonists				
SALBUTAMOL				
Aerosol inhaler, 100 mcg per dose CFC free – Up to 1000 dose available on a PSO	3.80	200 dose OF		Respigen SalAir
Nebuliser soln, 1 mg per ml, 2.5 ml ampoule – Up to 30 neb	(6.20)			/entolin
available on a PSO	8.96	20	_	Asthalin PMS-
				Salbutamol S29
			√ T	eva-Salbutamol Sterinebs P.F. §29
			✓ \	/entolin Nebules §29
Nebuliser soln, 2 mg per ml, 2.5 ml ampoule – Up to 30 neb	51.11		✓ A	Accord S29
available on a PSO	9.43	20	_	Asthalin PMS-
	14.15	30	√ 9	Salbutamol S29 Salbutamol
				Cipla \$29
(Respigen Aerosol inhaler, 100 mcg per dose CFC free to be deli (Accord S29 Nebuliser soln, 1 mg per ml, 2.5 ml ampoule to be TERBUTALINE SULPHATE				
Powder for inhalation, 200 mcg per dose (equivalent to 250 mcg metered dose), breath activated	22.20	120 dose OF	· ✓ E	Bricanyl Turbuhaler
Anticholinergic Agents				
IPRATROPIUM BROMIDE				
Aerosol inhaler, 20 mcg per dose CFC-free – Up to 400 dos available on a PSO	16.20	200 dose OF	· • •	Atrovent
Nebuliser soln, 250 mcg per ml, 2 ml ampoule – Up to 40 ne available on a PSO		20		Inivent Accord \$29
Inhaled Beta-Adrenoceptor Agonists with Antic	holinergic A	Agents		
SALBUTAMOL WITH IPRATROPIUM BROMIDE				
Aerosol inhaler, 100 mcg with ipratropium bromide, 20 mcg p dose CFC-free		200 dose OF	· • •	Ouolin HFA
Nebuliser soln, 2.5 mg with ipratropium bromide 0.5 mg per vial, 2.5 ml ampoule – Up to 20 neb available on a PSO	14.04	00)alin
	33.12	20 60	_	<u>Duolin</u> Duolin Respules ^{S29}

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic

\$ Per

Manufacturer

Long-Acting Muscarinic Antagonists

GLYCOPYRRONIUM - Subsidy by endorsement

- a) Inhaled glycopyrronium treatment will not be subsidised if patient is also receiving treatment with subsidised tiotropium or umeclidinium
- b) Glycopyrronium powder for inhalation 50 mcg per dose is subsidised only for patients who have been diagnosed as having COPD using spirometry if spirometry is possible, and the prescription is endorsed accordingly.

TIOTROPIUM BROMIDE - Subsidy by endorsement

- a) Tiotropium treatment will not be subsidised if patient is also receiving treatment with subsidised inhaled glycopyrronium or umeclidinium.
- b) Tiotropium bromide is subsidised only for patients who have been diagnosed as having COPD using spirometry if spirometry is possible, and the prescription is endorsed accordingly. Patients who had tiotropium dispensed before 1 October 2018 with a valid Special Authority are deemed endorsed.

UMECLIDINIUM - Subsidy by endorsement

- a) Umeclidinium will not be subsidised if patient is also receiving treatment with subsidised inhaled glycopyrronium or tiotropium bromide.
- b) Umeclidinium powder for inhalation 62.5 mcg per dose is subsidised only for patients who have been diagnosed as having COPD using spirometry if spirometry is possible, and the prescription is endorsed accordingly.

Long-Acting Muscarinic Antagonists with Long-Acting Beta-Adrenoceptor Agonists

Combination long acting muscarinic antagonist and long acting beta-2 agonist will not be subsidised if patient is also receiving treatment with a combination inhaled corticosteroid and long acting beta-2 agonist.

⇒SA1584 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Patient has been stabilised on a long acting muscarinic antagonist; and
- 2 The prescriber considers that the patient would receive additional benefit from switching to a combination product.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Patient is compliant with the medication; and
- 2 Patient has experienced improved COPD symptom control (prescriber determined).

TIOTROPIUM BROMIDE WITH OLODATEROL – Special Authority see SA1584 above – Retail pharmacy Soln for inhalation 2.5 mcg with olodaterol 2.5 mcg81.00 60 dose OP ✓ Spiolto Respimat

UMECLIDINIUM WITH VILANTEROL - Special Authority see SA1584 above - Retail pharmacy

Inhaled Corticosteroid with Long-Acting Muscarinic Antagonist and Beta Agonist

FLUTICASONE FUROATE WITH UMECLIDINIUM AND VILANTEROL – Special Authority see SA2326 on the next page – Retail pharmacy

Powder for inhalation fluticasone furoate 100 mcg with

umeclidinium 62.5 mcg and vilanterol 25 mcg......104.24 30 dose OP ✓ Trelegy Ellipta

Subsidy	F	ully	Brand or	
(Manufacturer's Price)	Subsidis	sed	Generic	
\$	Per	•	Manufacturer	

⇒SA2326 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has a diagnosis of COPD confirmed by spirometry or spirometry has been attempted and technically acceptable results are not possible; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient is currently receiving an inhaled corticosteroid with long acting beta-2 agonist (ICS/LABA) or a long acting muscarinic antagonist with long acting beta-2 agonist (LAMA/LABA); and
 - 2.1.2 Any of the following:

Clinical criteria:

- 2.1.2.1 Patient has a COPD Assessment Test (CAT) score greater than 10; or
- 2.1.2.2 Patient has had 2 or more exacerbations in the previous 12 months; or
- 2.1.2.3 Patient has had one exacerbation requiring hospitalisation in the previous 12 months; or
- 2.1.2.4 Patient has had an eosinophil count greater than or equal to 0.3×10^9 cells/L in the previous 12 months; or
- 2.2 Patient is currently receiving multiple inhaler triple therapy (inhaled corticosteroid with long acting muscarinic antagonist and long acting beta-2 agonist ICS/LAMA/LABA) and met at least one of the clinical criteria above prior to commencing multiple inhaler triple therapy.

Antifibrotics

NINTEDANIB - Special Authority see SA2012 below - Retail pharmacy

Note: Nintedanib not subsidised in combination with subsidised pirfenidone.

Cap 100 mg	2,554.00	60 OP	Ofev
Cap 150 mg	3,870.00	60 OP	Ofev

⇒SA2012 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Nintedanib is to be discontinued at disease progression (See Note); and
- 4 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 5 Any of the following:
 - 5.1 The patient has not previously received treatment with pirfenidone; or
 - 5.2 Patient has previously received pirfenidone, but discontinued pirfenidone within 12 weeks due to intolerance; or
 - 5.3 Patient has previously received pirfenidone, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with pirfenidone).

Renewal — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 3 Nintedanib is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

	Subsidy (Manufacturer's Price)		Fully sidised	Brand or Generic Manufacturer	
PIRFENIDONE – Retail pharmacy-Specialist – Special Authority Note: Pirfenidone is not subsidised in combination with subs					
Tab 801 mg		90 OP 90	_	sbriet sbriet	

⇒SA2013 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note); and
- 4 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 5 Any of the following:
 - 5.1 The patient has not previously received treatment with nintedanib; or
 - 5.2 Patient has previously received nintedanib, but discontinued nintedanib within 12 weeks due to intolerance; or
 - 5.3 Patient has previously received nintedanib, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with nintedanib).

Renewal — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

Leukotriene Receptor Antagonists

MC	NIELUKASI			
*	Tab 4 mg	3.10	28	✓ Montelukast Viatris
	Tab 5 mg		28	✓ Montelukast Viatris
	Tab 10 mg		28	✓ Montelukast Viatris

Methylxanthines

AMINOPHYLLINE

*	Inj 25 mg per ml, 10 ml ampoule - Up to 5 inj available	on a		
	PSO	180.00	5	✓ DBL Aminophylline
TH	EOPHYLLINE			
*	Tab long-acting 250 mg	24.90	100	✓ Nuelin-SR
*	Oral lig 80 mg per 15 ml	17 95	500 ml	✓ Nuelin

Mucolytics

DORNASE ALFA – Special Authority see SA1978 below – Re	tail pharmacy		
Nebuliser soln, 2.5 mg per 2.5 ml ampoule	250.00	6	✓ Pulmozyme

⇒SA1978 Special Authority for Subsidy

Initial application — (cystic fibrosis) only from a respiratory physician or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

	Subsidy	Full	/ Brand or
(Manu	ufacturer's Price)	Subsidise	d Generic
	\$ F	Per 🗸	Manufacturer

continued...

All of the following:

- 1 Patient has a confirmed diagnosis of cystic fibrosis; and
- 2 Patient has previously undergone a trial with, or is currently being treated with, hypertonic saline; and
- 3 Any of the following:
 - 3.1 Patient has required one or more hospital inpatient respiratory admissions in the previous 12 month period; or
 - 3.2 Patient has had 3 exacerbations due to CF, requiring oral or intravenous (IV) antibiotics in the previous 12 month period: or
 - 3.3 Patient has had 1 exacerbation due to CF, requiring oral or IV antibiotics in the previous 12 month period and a Brasfield score of < 22/25; or</p>
 - 3.4 Patient has a diagnosis of allergic bronchopulmonary aspergillosis (ABPA).

Renewal — **(cystic fibrosis)** only from a respiratory physician or paediatrician. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient continues to benefit from treatment.

ELEXACAFTOR WITH TEZACAFTOR, IVACAFTOR AND IVACAFTOR - PCT only - Special Authority see \$A2196 below

⇒SA2196 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with cystic fibrosis; and
- 2 Patient is 6 years of age or older; and
- 3 Either:
 - 3.1 Patient has two cystic fibrosis-causing mutations in the cystic fibrosis transmembrane regulator (CFTR) gene (one from each parental allele); or
 - 3.2 Patient has a sweat chloride value of at least 60 mmol/L by quantitative pilocarpine iontophoresis or by Macroduct sweat collection system; and
- 4 Either:
 - 4.1 Patient has a heterozygous or homozygous F508del mutation; or
 - 4.2 Patient has a G551D mutation or other mutation responsive in vitro to elexacaftor/tezacaftor/ivacaftor (see note a); and
- 5 The treatment must be the sole funded CFTR modulator therapy for this condition; and
- 6 Treatment with elexacaftor/tezacaftor/ivacaftor must be given concomitantly with standard therapy for this condition.

Notes:

 a) Eligible mutations are listed in the Food and Drug Administration (FDA) Trikafta prescribing information https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/212273s004lbl.pdf

IVACAFTOR - PCT only - Specialist - Special Author	rity see SA2017 below		
Tab 150 mg	29,386.00	56	✓ Kalydeco
Oral granules 50 mg, sachet	29,386.00	56	✓ Kalydeco
Oral granules 75 mg, sachet	29,386.00	56	Kalydeco

⇒SA2017 Special Authority for Subsidy

Initial application only from a respiratory specialist or paediatrician. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

	RESPIRATO	RY SYSTE	M AN	ID ALLERGIES
	Subsidy (Manufacturer's Prio \$		Fully dised	Brand or Generic Manufacturer
continued 1 Patient has been diagnosed with cystic fibrosis; and 2 Either: 2.1 Patient must have G551D mutation in the cystic fib least 1 allele; or 2.2 Patient must have other gating (class III) mutation and S549R) in the CFTR gene on at least 1 allele; 3 Patients must have a sweat chloride value of at least 60 m sweat collection system; and 4 Treatment with ivacaftor must be given concomitantly with 5 Patient must not have an acute upper or lower respiratory (including antibiotics) for pulmonary disease in the last 4 w 6 The dose of ivacaftor will not exceed one tablet or one sac 7 Applicant has experience and expertise in the management	(G1244E, G1349E and nmol/L by quantita standard therapy infection, pulmona veeks prior to com chet twice daily; ar	o, G178R, G55 tive pilocarpine for this condit ary exacerbation imencing treated	51S, S1 e iontop ion; and on, or c	251N, S1255P, S549N choresis or by Macroduct d changes in therapy
SODIUM CHLORIDE Not funded for use as a nasal drop. Soln 7%	24.50	90 ml OP	✓ Bi	omed
Nasal Preparations				
Allergy Prophylactics				
BUDESONIDE Metered aqueous nasal spray, 50 mcg per dose Metered aqueous nasal spray, 100 mcg per dose		200 dose OP 200 dose OP		eroClear eroClear
FLUTICASONE PROPIONATE Metered aqueous nasal spray, 50 mcg per dose	1.98 1	120 dose OP	_	ixonase Hayfever & Allergy
IPRATROPIUM BROMIDE Aqueous nasal spray, 0.03%	5.23	15 ml OP	✓ Ur	nivent

Respiratory Devices

MASK FOR SPACER DEVICE

- a) Up to 50 dev available on a PSO
- b) Only on a PSO
- c) Only for children aged six years and under Small......2.70
- PEAK FLOW METER

PEAK FLOW WETEN

- a) Up to 25 dev available on a PSO
- b) Only on a PSO

Low range 9.54

1 ✓ Mini-Wright AFS Low Range

Normal range......9.54

Mini-Wright Standard

✓ e-chamber Mask

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
SPACER DEVICE				
a) Up to 50 dev available on a PSO				
b) Only on a PSO				
220 ml (single patient)	3.65	1	✓	e-chamber Turbo
510 ml (single patient)	5.95	1	•	e-chamber La Grande
800 ml	6.50	1	1	Volumatic

CAFFEINE	CITRATE
----------	---------

Oral liq 20 mg per ml (10 mg base per ml)......16.10 25 ml OP ✓ Biomed

	Subsidy		Fully Brand or
	(Manufacturer's Pri		•
	\$	Per	 Manufacturer
Ear Preparations			
ELLIMETA CONE DIVALATE			
FLUMETASONE PIVALATE	4.40	7.5 OD	()
Ear drops 0.02% with clioquinol 1%	4.46	7.5 ml OP	✓ Locacorten-Viaform
			ED's
			✓ Locorten-Vioform
TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYC	IN AND NYSTATI	N	
Ear drops 1 mg with nystatin 100,000 u, neomycin sulphate			
2.5 mg and gramicidin 250 mcg per g	5.16	7.5 ml OP	✓ Kenacomb
Ear/Eye Preparations			
_ai/_yo : reparations			
DEXAMETHASONE WITH FRAMYCETIN AND GRAMICIDIN			
Ear/Eye drops 500 mcg with framycetin sulphate 5 mg and			
gramicidin 50 mcg per ml	4.50	8 ml OP	
g 2	(9.27)		Otodex S29
	(9.27)		Sofradex
FRAMYCETIN SULPHATE	()		
	4.10	0 ml OD	
Ear/Eye drops 0.5%		8 ml OP	Cofromusin
	(8.65)		Soframycin
Evo Dronorationa			
Eye Preparations			
Eye preparations are only funded for use in the eye, unless explic	citly stated otherw	ico	
Lye preparations are only funded for use in the eye, unless expin	citiy stated officiw	136.	
Anti-Infective Preparations			
·			
ACICLOVIR			
* Eye oint 3%	14.88	4.5 g OP	✓ <u>ViruPOS</u>
CHLORAMPHENICOL			
Eye oint 1%	1.09	5 g OP	✓ Devatis
Eye drops 0.5%	1.45	10 ml OP	✓ Chlorsig
Funded for use in the ear*. Indications marked with * ar		cations.	
CIPROFLOXACIN			
Eye drops 0.3% – Subsidy by endorsement	9.73	5 ml OP	✓ Ciprofloxacin Teva
When prescribed for the treatment of bacterial keratitis of			
for the second line treatment of chronic suppurative otitis			
Note: Indication marked with a * is an unapproved indic		and the presen	iption is endorsed accordingly.
	ation.		
SODIUM FUSIDATE [FUSIDIC ACID]	7.00	- 00	
Eye drops 1%	5.29	5 g OP	✓ Fucithalmic
TOBRAMYCIN			
Eye oint 0.3%	10.45	3.5 g OP	✓ Tobrex
Eye drops 0.3%	11.48	5 ml OP	✓ Tobrex
Corticosteroids and Other Anti-Inflammatory Pr	reparations		
DEVAMETHACONE			
DEXAMETHASONE	Г 00	0.5 ~ 0.0	-/ Mayiday
* Eye oint 0.1%		3.5 g OP	✓ Maxidex
* Eye drops 0.1%		5 ml OP	✓ Maxidex
Ocular implant 700 mcg - Special Authority see SA1680 on			
the next page – Retail pharmacy	1,444.50	1	✓ Ozurdex

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.



Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

⇒SA1680 Special Authority for Subsidy

Initial application — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has diabetic macular oedema with pseudophakic lens; and
- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- 3 Either:
 - 3.1 Patient's disease has progressed despite 3 injections with bevacizumab; or
 - 3.2 Patient is unsuitable or contraindicated to treatment with anti-VEGF agents; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Initial application — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has diabetic macular oedema: and
- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- 3 Patient is of child bearing potential and has not yet completed a family; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Patient is of child bearing potential and has not yet completed a family; and
- 3 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

DEXAMETHASONE WITH NEOMYCIN SULPHATE AND POLYMYXIN B SULPHATE

* Eve pint 0.1% with neomycin sulphate 0.35% and polymyvin h

不	sulphate 6,000 u per g	5 30	3.5 g OP	✓ Maxitrol
			3.5 g OF	♥ WaxitiOi
*	Eye drops 0.1% with neomycin sulphate 0.35% and polymyxin	4.50	5 ml OD	/ Massissal
	b sulphate 6,000 u per ml	4.50	5 ml OP	✓ Maxitrol
DIC	CLOFENAC SODIUM			
	Eye drops 0.1%	8.80	5 ml OP	✓ Voltaren Ophtha
(Vo	oltaren Ophtha Eye drops 0.1% to be delisted 1 December 2024)			
FLI	JOROMETHOLONE			
*	Eye drops 0.1%	3.09	5 ml OP	✓ FML
	,	5.20		✓ Flucon
LE'	VOCABASTINE			
	Eye drops 0.5 mg per ml	8.71	4 ml OP	
	, ,	(10.34)		Livostin
LO	DOXAMIDE			
	Eve drops 0.1%	8.71	10 ml OP	✓ Lomide

	Subsidy		Fully	Brand or
	(Manufacturer's P	rice) Sub	sidised	Generic
	\$	Per	✓	Manufacturer
NEPAFENAC				
Eye drops 0.3%	8.80	3 ml OP	✓	evro
PREDNISOLONE ACETATE				
Eve drops 1%	6.92	10 ml OP	✓ P	rednisolone-AFT
,	7.00	5 ml OP	√ P	red Forte
PREDNISOLONE SODIUM PHOSPHATE - Special Author	ority see SA1715 below	– Retail phai	macy	
Eye drops 0.5%, single dose (preservative free)	41.20	20 dose	✓ N	linims Prednisolone

⇒SA1715 Special Authority for Subsidy

Initial application only from an ophthalmologist or optometrist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has severe inflammation; and
- 2 Patient has a confirmed allergic reaction to preservative in eye drops.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

SODIUM CROMOGLICATE Eye drops 2%2.62	10 ml OP	✓ <u>Allerfix</u>
Glaucoma Preparations - Beta Blockers		
# Eye drops 0.25%	5 ml OP 5 ml OP	✓ Betoptic S✓ Betoptic
* Eye drops 0.25%	5 ml OP 5 ml OP	✓ <u>Arrow-Timolol</u> ✓ <u>Arrow-Timolol</u>
Glaucoma Preparations - Carbonic Anhydrase Inhibitors		
ACETAZOLAMIDE * Tab 250 mg	100	✓ Diamox
# Eye drops 1%	5 ml OP	✓ <u>Azopt</u>
* Eye drops 2% with timolol 0.5%	5 ml OP	✓ <u>Dortimopt</u>
Glaucoma Preparations - Prostaglandin Analogues		
BIMATOPROST * Eye drops 0.03%	3 ml OP	✓ Bimatoprost Multichem
LATANOPROST * Eye drops 0.005%	2.5 ml OP	✓ <u>Teva</u>
* Eye drops 0.004%	2.5 ml OP	✓ <u>Travatan</u>

	Subsidy (Manufacturer's I \$	Price) Subs	Fully idised	Brand or Generic Manufacturer
Glaucoma Preparations - Other				
BRIMONIDINE TARTRATE * Eye drops 0.2% BRIMONIDINE TARTRATE WITH TIMOLOL MALEATE	4.29	5 ml OP	✓ <u>Ar</u>	row-Brimonidine
Eye drops 0.2% with timolol maleate 0.5%	18.50	5 ml OP	✓ Co	ombigan
LATANOPROST WITH TIMOLOL * Eye drops 0.005% with timolol 0.5%	4.95	2.5 ml OP	✓ <u>Ar</u>	row - Lattim
PILOCARPINE HYDROCHLORIDE * Eye drops 1% * Eye drops 2% * Eye drops 4% Subsidised for oral use pursuant to the Standard Formu	5.35 7.99	15 ml OP 15 ml OP 15 ml OP	✓ Iso	opto Carpine opto Carpine opto Carpine
PILOCARPINE NITRATE * Eye drops 2% single dose – Special Authority see SA0895 below – Retail pharmacy	34.19	20 dose	✓ Mi	nims Pilocarpine
■ SA0895 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valiether:	d for 2 years for	applications me	eting the	e following criteria:
 Patient has to use an unpreserved solution due to an alle Patient wears soft contact lenses. 	,			
Note: Minims for a general practice are considered to be "tools or Panaural from any relevant practitioner. Approvals valid for 2 years				,

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is

Mydriatics	and C	yclop	legics
------------	-------	-------	--------

benefiting from treatment.

* Eye drops 1%18.27	15 ml OP	✓ Atropt
CYCLOPENTOLATE HYDROCHLORIDE * Eye drops 1%	15 ml OP	✓ Cyclogyl
* Eye drops 1%, single dose (preservative free) - Only on a prescription84.85	20 dose	✓ Minims Cyclopentolate
TROPICAMIDE * Eye drops 0.5%	15 ml OP 15 ml OP	✓ Mydriacyl ✓ Mydriacyl

Preparations for Tear Deficiency

For acetylcysteine eye drops refer Standard Formulae, page 267

HY	PRO	MEL	LO	SE
*	FVO	dror	ne N	50

HY	PROMELLOSE WITH DEXTRAN			
*	Eve drops 0.3% with dextran 0.1%	2.30	15 ml OP	✓ Poly-Tears

15 ml OP

✓ Methopt

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
` \$	Per	/	Manufacturer

Preservative Free Ocular Lubricants

⇒SA2134 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

- 1 Confirmed diagnosis by slit lamp or Schirmer test of severe secretory dry eye; and
- 2 Either:
 - 2.1 Patient is using eye drops more than four times daily on a regular basis; or
 - 2.2 Patient has had a confirmed allergic reaction to preservative in eye drop.

Renewal from any relevant practitioner. Approvals valid for 24 months where the patient continues to require lubricating eye drops and has benefited from treatment.

CARBOMER - Special Authority see SA2134 above - Retail ph	narmacy		
Ophthalmic gel 0.3%, 0.5 g	8.25	30	✓ Poly-Gel
POLYETHYLENE GLYCOL 400 AND PROPYLENE GLYCOL -	- Special Authority s	ee SA2134 a	above – Retail pharmacy
Eye drops 0.4% and propylene glycol 0.3%, 0.8 ml	10.78	30	✓ Systane Unit Dose
SODIUM HYALURONATE [HYALURONIC ACID] - Special Aut	hority see SA2134 a	above – Reta	ail pharmacy
Eye drops 1 mg per ml	13.85	10 ml OP	✓ Hylo-Fresh
Hylo-Fresh has a 6 month expiry after opening. The Pr	narmacy Procedures	Manual res	triction allowing one bottle per
month is not relevant and therefore only the prescribed	dosage to the neare	est OP may b	oe claimed.

Other Eye Preparations

NAPHAZOLINE HYDROCHLORIDE * Eye drops 0.1%4.15	15 ml OP	✓ Naphcon Forte
OLOPATADINE Eye drops 0.1%2.17	5 ml OP	✓ Olopatadine Teva
PARAFFIN LIQUID WITH WOOL FAT * Eye oint 3% with wool fat 3%	3.5 g OP	✓ Poly-Visc
RETINOL PALMITATE Eve oint 138 mcg per g	5 a OP	✓ VitA-POS

	Subsidy (Manufacturer's Prid	ce) Sub	sidised Generic
	\$	Per	✓ Manufacturer
Various			
PHARMACY SERVICES			
* Brand switch fee	4.50	1 fee	 ✓ BSF Ipca-Donepezil ✓ BSF Max Health ✓ BSF Methylphenidate ER - Teva ✓ BSF Teriparatide - Teva
 a) May only be claimed once per patient. b) The Pharmacode for BSF Methylphenidate ER - Te c) The Pharmacode for BSF Max Health is 2677903 - d) The Pharmacode for BSF Teriparatide - Teva is 26 	see also page 57 79701 - see also pa	ge 119	146
e) The Pharmacode for BSF Ipca-Donepezil is 26797 * COVID-19 Services	1 0	48 1 fee	✓ After Hours Med Mgmt 15 min ✓ After Hours Med Mgmt 30 min ✓ After Hours Med Mgmt 45 min ✓ Antivirals Eligibility Review ✓ Compliance Packaging ✓ Med Mgmt 15 min ✓ Med Mgmt 30 min ✓ Med Mgmt 45 min ✓ Med Mgmt 45 min ✓ Medicine Delivery
* Immunisation administration fee	0.00	1 fee	✓ Immunisation Administration
* Immunisation co-administration fee	0.00	1 fee	✓ Immunisation Co-administration
(BSF Ipca-Donepezil Brand switch fee to be delisted 1 Septem (BSF Max Health Brand switch fee to be delisted 1 August 202 (BSF Methylphenidate ER - Teva Brand switch fee to be delisted (BSF Teriparatide - Teva Brand switch fee to be delisted 1 Sep	4) ed 1 August 2024)		CO-auministration
Agents Used in the Treatment of Poisonings			
Antidotes			
ACETYLCYSTEINE Inj 200 mg per ml, 10 ml ampoule NALOXONE HYDROCHLORIDE a) Up to 10 inj available on a PSO b) Only on a PSO	52.88	10	✓ Martindale Pharma
* Inj 400 mcg per ml, 1 ml ampoule	35.26	10	✓ <u>Hameln</u>

Subsidy

Fully

Brand or

				VARIOUS
	Subsidy (Manufacturer's Pri	ce) (Fully Subsidised	
Removal and Elimination				
CHARCOAL * Oral liq 50 g per 250 ml a) Up to 250 ml available on a PSO b) Only on a PSO	43.50	250 ml C	P 🗸	Carbosorb-X
DEFERASIROX – Special Authority see SA1492 below – Retail Wastage claimable Tab 125 mg dispersible		28		Exjade
Tab 250 mg dispersible	552.00	28 28	✓	Exjade Exjade Exjade
▶SA1492 Special Authority for Subsidy Initial application only from a haematologist. Approvals valid fo All of the following:	, ,,		Ū	ŭ
 The patient has been diagnosed with chronic iron overload 2 Deferasirox is to be given at a daily dose not exceeding 4/3 Any of the following: 3.1 Treatment with maximum tolerated doses of defering combination therapy have proven ineffective as media. 3.2 Treatment with deferiprone has resulted in severe 3.3 Treatment with deferiprone has resulted in arthritis 3.4 Treatment with deferiprone is contraindicated due count (ANC) of < 0.5 cells per μL) or recurrent epis 0.5 - 1.0 cells per μL). Renewal only from a haematologist. Approvals valid for 2 years 	o mg/kg/day; and prone monotherap easured by serum persistent vomiting; or to a history of agrasodes (greater than	oy or defe ferritin lev g or diarrh anulocyto n 2 episod	riprone a vels, liver noea; or sis (defin des) of m	nd desferrioxamine or cardiac MRI T2*; or ed as an absolute neutrophil oderate neutropenia (ANC
Either: 1 For the first renewal following 2 years of therapy, the treat improvement in all three parameters namely serum ferritin 2 For subsequent renewals, the treatment has been tolerate in all three parameters namely serum ferritin, cardiac MRI	i, cardiac MRI T2* ed and has resulte	and liver d in clinic	MRI T2* al stabilit	levels; or
DEFERIPRONE - Special Authority see SA1480 below - Retail Tab 500 mg Oral liq 100 mg per 1 ml SA1480 Special Authority for Subsidy Initial application only from a haematologist. Approvals valid wifollowing criteria:	533.17 266.59	100 250 ml C)P 🗸	Ferriprox Ferriprox for applications meeting the
Either: 1 The patient has been diagnosed with chronic iron overload 2 The patient has been diagnosed with chronic iron overload				a; or
DESFERRIOXAMINE MESILATE * Ini 500 mg vial	151 21	10	s	DBL
Tri j 500 filg viai	101.01	10	·	Desferrioxamine Mesylate for Inj

BP

✓ Deferoxamine Pfizer
S29 S29

VARIOUS

	Subsidy (Manufacturer's Price) \$	Sub Per	Fully sidised	Brand or Generic Manufacturer
SODIUM CALCIUM EDETATE				
* Inj 200 mg per ml, 5 ml	53.31	6		
	(156.71)		С	Calcium Disodium Versenate

Standard Formulae

ACETYLCYSTEINE EYE DROPS Acetylcysteine inj 200 mg per ml, 10 ml Suitable eye drop base	qs qs	PHENOBARBITONE ORAL LIQUID Phenobarbitone Sodium Glycerol BP Water	1 g 70 ml to 100 ml
CODEINE LINCTUS (3 mg per 5 ml) Codeine phosphate	60 mg	PHENOBARBITONE SODIUM PAEDIATRIC ORAL	LIQUID (10
Glycerol	40 ml	mg per ml)	,
Preservative	qs	Phenobarbitone Sodium	400 mg
Water	to 100 ml	Glycerol BP	4 ml
CODEINE LINCTUS (15 mg per 5 ml)		Water	to 40 ml
Codeine phosphate	300 mg	PILOCARPINE ORAL LIQUID	
Glycerol	40 ml	Pilocarpine 4% eye drops	qs
Preservative	qs	Preservative	qs
Water	to 100 ml	Water	to 500 ml
		(Preservative should be used if quantity supplied is	for more
FOLINIC MOUTHWASH	4 4 4 4	than 5 days.)	
Calcium folinate 15 mg tab	1 tab	CALIVA CUDCTITUTE FORMULA	
Preservative Water	qs to 500 ml	SALIVA SUBSTITUTE FORMULA	F ~
(Preservative should be used if quantity supplied is		Methylcellulose Preservative	5 g
than 5 days. Maximum 500 ml per prescription.)	ioi illole	Water	qs to 500 ml
than 5 days. Maximum 300 mi per prescription.		(Preservative should be used if quantity supplied is	
METHADONE MIXTURE		than 5 days. Maximum 500 ml per prescription.)	ioi illoic
Methadone powder	qs	man o dayo. Maximum ooo mi por procenpuon.	
Glycerol	qs	SODIUM CHLORIDE ORAL LIQUID	
Water	to 100 ml	Sodium chloride inj 23.4%, 20 ml	qs
METHYL HYDROXYBENZOATE 10% SOLUTION		Water	qs
Methyl hydroxybenzoate	10 g	(Only funded if prescribed for treatment of hyponatr	aemia)
Propylene glycol	to 100 ml	VANCOMYCIN ORAL SOLUTION (25 mg per ml)	
(Use 1 ml of the 10% solution per 100 ml of oral liqu		Vancomycin 500 mg injection	5 vials
	aid illixtulo)	Glycerin with sucrose suspension	37.5 ml
OMEPRAZOLE SUSPENSION		Water	to 100 ml
Omeprazole capsules or powder	qs	(Only funded if prescribed for treatment of Clostridiu	ım difficile
Sodium bicarbonate powder BP	8.4 g	following metronidazole failure)	
Water	to 100 ml		

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer Extemporaneously Compounded Preparations and Galenicals CODEINE PHOSPHATE - Safety medicine; prescriber may determine dispensing frequency (90.09)Douglas Only in extemporaneously compounded codeine linctus. COLLODION FLEXIBLE Note: This product is no longer being manufactured by the supplier and will be delisted from the Schedule at a date to be determined. 100 ml ✓ PSM COMPOUND HYDROXYBENZOATE - Only in combination Only in extemporaneously compounded oral mixtures. 100 ml ✓ Midwest GLYCERIN WITH SODIUM SACCHARIN - Only in combination Only in combination with Ora-Plus or when used in the vancomycin oral Iguuid Standard Formulae. 473 ml ✓ Ora-Sweet SF GLYCERIN WITH SUCROSE - Only in combination Only in combination with Ora-Plus or when used in the vancomycin oral Iguuid Standard Formulae. Suspension......30.95 473 ml ✓ Ora-Sweet GLYCEROL 500 ml ✓ healthE Glycerol BP Only in extemporaneously compounded oral liquid preparations. METHADONE HYDROCHLORIDE a) Only on a controlled drug form b) No patient co-payment payable c) Safety medicine; prescriber may determine dispensing frequency d) Extemporaneously compounded methadone will only be reimbursed at the rate of the cheapest form available (methadone powder, not methadone tablets).

Powder	7.84	1 g	✓ AFT
(AFT Powder to be delisted 1 July 2024)		•	
METHYL HYDROXYBENZOATE			
Powder	8.98	25 g	✓ Midwest
METHYLCELLULOSE			
Powder	36.95	100 g	✓ MidWest
Suspension – Only in combination	30.95	473 ml	Ora-Plus
METHYLCELLULOSE WITH GLYCERIN AND SODIUM SACCH	ARIN - Only in c	ombination	
Suspension	30.95	473 ml	Ora-Blend SF
METHYLCELLULOSE WITH GLYCERIN AND SUCROSE - Or	ly in combination		
Suspension	30.95	473 ml	Ora-Blend
PHENOBARBITONE SODIUM			
Powder - Only in combination	52.50	10 g	✓ MidWest
	325.00	100 g	✓ MidWest
Only in children up to 12 years			
PROPYLENE GLYCOL			
Only in extemporaneously compounded methyl hydroxyben	zoate 10% solutio	٦.	
Liq	11.25	500 ml	✓ Midwest

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

	Subsidy (Manufacturer's P		Fully sidised	Brand or Generic Manufacturer	
SODIUM BICARBONATE Powder BP - Only in combination Only in extemporaneously compounded omeprazole and		500 g spension.		// // // // // // // // // // // // //	
SYRUP (PHARMACEUTICAL GRADE) – Only in combination Only in extemporaneously compounded oral liquid preparatio Liq		500 ml	✓ N	lidwest	
WATER Tap - Only in combination	0.00	1 ml	✓ T	ap water	

Subsidy (Manufacturer's Price)

Fully Subsidised Per

Brand or Generic Manufacturer

Nutrient Modules

Carbohydrate

⇒SA1930 Special Authority for Subsidy

Initial application — (Cystic fibrosis or kidney disease) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria: Fither:

- 1 cystic fibrosis; or
- 2 chronic kidney disease.

Initial application — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 cancer in children: or
- 2 cancers affecting alimentary tract where there are malabsorption problems in patients over the age of 20 years; or
- 3 faltering growth in an infant/child; or
- 4 bronchopulmonary dysplasia; or
- 5 premature and post premature infant; or
- 6 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified where the patient has inborn errors of metabolism. Renewal — (Cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

CARBOHYDRATE SUPPLEMENT - Special Authority see SA1930 above - Hospital pharmacy [HP3]

400 a OP ✓ Polycal Powder6.72

Carbohydrate And Fat

⇒SA1376 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria: Both:

0.1.1		F. "	D 1	
Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1 Infant or child aged four years or under; and
- 2 cystic fibrosis.

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 infant or child aged four years or under; and
- 2 Any of the following:
 - 2.1 cancer in children; or
 - 2.2 faltering growth; or
 - 2.3 bronchopulmonary dysplasia; or
 - 2.4 premature and post premature infants.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Fat

⇒SA2204 Special Authority for Subsidy

Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified where the patient has an inborn error of metabolism. Initial application — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 faltering growth in an infant/child; or
- 2 bronchopulmonary dysplasia; or
- 3 fat malabsorption; or
- 4 lymphangiectasia; or
- 5 short bowel syndrome: or
- 6 infants with necrotising enterocolitis; or
- 7 biliary atresia: or
- 8 for use in a ketogenic diet; or
- 9 chyle leak; or

continued...

Subsidy		Fully	Brand or
(Manufacturer's Price)	S	ubsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 10 ascites: or
- 11 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. Renewal — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

1 The treatment remains appropriate and the patient is benefiting from treatment; and

EAT CLIDDLEMENT Charity and CA2204 on the provious page. Heapital pharmacy [HD2]

2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT SUPPLEMENT — Special Authority see SA2204 0	n the previous page – nos	pilai pilaimacy	[nroj
Emulsion (neutral)	15.38	200 ml OP	 Calogen
	38.44	500 ml OP	✓ Calogen
Emulsion (strawborn)	15 38	200 ml OP	✓ Calogen

Zindioion (diambony)		200 1111 01	- Guiogon
Oil	37.50	500 ml OP	✓ MCT oil (Nutricia)
MCT Emulsion, 250 ml	143.65	4 OP	✓ Liquigen

Protein

⇒SA1524 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 protein losing enteropathy; or
- 2 high protein needs; or
- 3 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

PROTEIN SUPPLEMENT - Special Authority see SA1524 above	- Hospital pha	rmacy [HP3]	
Powder	8.95	227 g OP	✓

✓ Resource Beneprotein

13.82 225 g OP

✓ Protifar

Subsidy (Manufacturer's Price) Per

Subsidised

Fully

Brand or Generic Manufacturer

Oral and Enteral Feeds

Diabetic Products

⇒SA1095 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is a type I or and II diabetic who is suffering weight loss and malnutrition that requires nutritional support. Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

DIABETIC ENTERAL FEED 1KCAL/ML - Special Authority see SA1095 above - Hospital pharmacy [HP3]

Liquid	4.65	500 ml OP	 Glucerna Select
	7.50	1,000 ml OP	✓ Nutrison Advanced
			Diason

(Nutrison Advanced Diason Liquid to be delisted 1 July 2024)

DIABETIC ORAL FEED 1KCAL/ML - Special Authority see SA1095 above - Hospital pharmacy [HP3]

Liquid (strawberry)1.50	200 ml OP	✓ Diasip
Liquid (vanilla)1.50	200 ml OP	✓ Diasip
2.10		✓ Nutren Diabetes

Fat Modified Products

⇒SA2205 Special Authority for Subsidy

Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified where the patient has an inborn error of metabolism. Initial application — (Indications other than errors of inborn metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Patient has a chyle leak; or
- 2 Modified as a modular feed, made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule, for adults,

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT MODIFIED FEED - Special Authority see SA2205 above - Hospital pharmacy [HP3]

400 g OP Monogen

Subsidy (Manufacturer's Price) Fully Subsidised Per 🗸 Brand or Generic Manufacturer

Paediatric Products For Children Awaiting Liver Transplant

⇒SA1098 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) who requires a liver transplant.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL FEED 1KCAL/ML - Special Authority see SA1098 above - Hospital pharmacy [HP3]

Paediatric Products For Children With Chronic Renal Failure

⇒SA1099 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) with acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Paediatric Products

⇒SA1379 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Child is aged one to ten years; and
- 2 Any of the following:
 - 2.1 the child is being fed via a tube or a tube is to be inserted for the purposes of feeding; or
 - 2.2 any condition causing malabsorption; or
 - 2.3 faltering growth in an infant/child; or
 - 2.4 increased nutritional requirements; or
 - 2.5 the child is being transitioned from TPN or tube feeding to oral feeding.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for

	Subsidy (Manufacturer's Price)	Fully Subsidised	Brand or Generic
	\$	Per 🗸	Manufacturer
continued applications meeting the following criteria:			

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

PAEDIATRIC ENTERAL FEED 1.5KCAL/ML – Special Authority see S Liquid		the previous pag 500 ml OP	ge – Hospital pharmacy [HP3] Frebini Energy Nutrini Energy RTH
PAEDIATRIC ENTERAL FEED 1KCAL/ML – Special Authority see SA Liquid		e previous page 500 ml OP	 Hospital pharmacy [HP3] ✓ Pediasure RTH ✓ Nutrini RTH ✓ Frebini Original
PAEDIATRIC ENTERAL FEED WITH FIBRE 1.5KCAL/ML - Special A pharmacy [HP3]	uthority se	e SA1379 on the	e previous page – Hospital
Liquid	7.00 7.14	500 ml OP	✓ Frebini Energy Fibre✓ Nutrini Energy Multi Fibre
PAEDIATRIC ENTERAL FEED WITH FIBRE 1KCAL/ML - Special Aut pharmacy [HP3]	hority see	SA1379 on the	previous page – Hospital
Liquid	7.00	500 ml OP	✓ Frebini Original Fibre
PAEDIATRIC ORAL FEED 1.5KCAL/ML - Special Authority see SA13	79 on the r	orevious page -	Hospital pharmacy [HP3]
Liquid (strawberry)		200 ml OP	✓ Fortini
Liquid (vanilla)	1.90	200 ml OP	✓ Fortini
	8.67	500 ml OP	✓ Pediasure Plus
PAEDIATRIC ORAL FEED 1KCAL/ML - Special Authority see SA1379	on the pre	evious page – H	ospital pharmacy [HP3]
Liquid (chocolate)	1.33	200 ml OP	✓ Pediasure
Liquid (strawberry)	1.33	200 ml OP	✓ Pediasure
Liquid (vanilla)		200 ml OP	✓ Pediasure
	1.66	250 ml OP	✓ Pediasure
PAEDIATRIC ORAL FEED WITH FIBRE 1.5KCAL/ML - Special Author pharmacy [HP3]	rity see SA	1379 on the pre	evious page – Hospital
Liquid (unflavoured)	1.90	200 ml OP	✓ Fortini Multi Fibre
Liquid (chocolate)	1.90	200 ml OP	✓ Fortini Multi Fibre
Liquid (strawberry)	1.90	200 ml OP	✓ Fortini Multi Fibre
Liquid (vanilla)	1.90	200 ml OP	✓ Fortini Multi Fibre
PEPTIDE-BASED ORAL FEED - Special Authority see SA1379 on the	previous	page – Hospital	pharmacy [HP3]
Powder		400 g OP	✓ Peptamen Junior

Renal Products

⇒SA1101 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the

continued...

Subsidy		ully Brand or	
(Manufacturer's \$	Price) Subsidis Per	sed Generic Manufacturer	

continued...

recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

RENAL ENTERAL FEED 1.8 KCAL/ML – Special Authority s Liquid(Nepro HP RTH Liquid to be delisted 1 August 2024)		revious page – 500 ml OP	Hospital pharmacy [HP3] ✓ Nepro HP RTH
RENAL ORAL FEED 1.8 KCAL/ML - Special Authority see S Liquid		ous page – Hos 220 ml OP	spital pharmacy [HP3] Nepro HP (strawberry) Nepro HP (vanilla)
RENAL ORAL FEED 2 KCAL/ML - Special Authority see SA	1101 on the previou	s page – Hosp	ital pharmacy [HP3]
Liquid, 200 ml bottle	11.52	4 OP	, ,, ,
	(13.24)		NovaSource Renal
Liquid (apricot) 125 ml	13.72	4 OP	✓ Renilon 7.5
Liquid (caramel) 125 ml	13.72	4 OP	✓ Renilon 7.5

Specialised And Elemental Products

⇒SA1377 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 malabsorption; or
- 2 short bowel syndrome; or
- 3 enterocutaneous fistulas; or
- 4 eosinophilic oesophagitis; or
- 5 inflammatory bowel disease; or
- 6 patients with multiple food allergies requiring enteral feeding.

Notes: Each of these products is highly specialised and would be prescribed only by an expert for a specific disorder. The alternative is hospitalisation.

Elemental 028 Extra is more expensive than other products listed in this section and should only be used where the alternatives have been tried first and/or are unsuitable.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL SEMI-ELEMENTAL FEED 1.5KCAL/ML - Sp	pecial Authority see	SA1377 above	- Hospital pharmacy [HP3]
Liquid	22.39	1,000 ml OP	✓ Vital
ORAL ELEMENTAL FEED 0.8KCAL/ML - Special Authority se	ee SA1377 above -	- Hospital pharn	nacy [HP3]
Liquid (grapefruit), 250 ml carton	179.46	18 OP	✓ Elemental 028 Extra
Liquid (pineapple & orange), 250 ml carton	179.46	18 OP	✓ Elemental 028 Extra
Liquid (summer fruits), 250 ml carton	179 46	18 OP	✓ Flemental 028 Extra

	Subsidy (Manufacturer's Pri \$	ce) Subs Per	Fully idised	Brand or Generic Manufacturer
ORAL ELEMENTAL FEED 1KCAL/ML – Special Authority see S Powder (unflavoured)		vious page – I 80 g OP		ıl pharmacy [HP3] 'ivonex TEN
SEMI-ELEMENTAL ENTERAL FEED 1KCAL/ML - Special Autr [HP3]	nority see SA1377	on the previou	us page	e – Hospital pharmacy
Liquid	7.47	500 ml OP	✓ N	lutrison Advanced Peptisorb
	9.60		√ S	survimed OPD

Paediatric Products For Children With Low Energy Requirements

⇒SA1196 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Child aged one to eight years; and
- 2 The child has a low energy requirement but normal protein and micronutrient requirements.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Standard Supplements

⇒SA1859 Special Authority for Subsidy

Initial application — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age; and
- 2 Any of the following:
 - 2.1 The patient has a condition causing malabsorption; or
 - 2.2 The patient has failure to thrive: or
 - 2.3 The patient has increased nutritional requirements; and
- 3 Nutrition goal has been set (eg reach a specific weight or BMI).

Renewal — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 A nutrition goal has been set (eg reach a specific weight or BMI).

Initial application — (Children - exclusive enteral nutrition for Crohn's disease) only from a gastroenterologist or dietitian on

continued...

Subsidy		Fully	Brand or
(Manufacturer's Price	e)	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 Dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Renewal — (Children - exclusive enteral nutrition for Crohn's disease) from any relevant practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 General Practitioners and dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Initial application — (Adults) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

1 Any of the following:

Patient is Malnourished

- 1.1 Patient has a body mass index (BMI) of less than 18.5 kg/m²; or
- 1.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 1.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months; and
- 2 Any of the following:

Patient has not responded to first-line dietary measures over a 4 week period by:

- 2.1 Increasing their food intake frequency (eg snacks between meals); or
- 2.2 Using high-energy foods (e.g. milkshakes, full fat milk, butter, cream, cheese, sugar etc); or
- 2.3 Using over the counter supplements (e.g. Complan); and
- 3 A nutrition goal has been set (e.g. to reach a specific weight or BMI).

Renewal — (Adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 A nutrition goal has been set (eg reach a specific weight or BMI); and
- 2 Any of the following:

Patient is Malnourished

- 2.1 Patient has a body mass index (BMI) of less than 18.5 kg/m²; or
- 2.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 2.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months.

Initial application — (Short-term medical condition) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube or a nasogastric tube is to be inserted for feeding; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Is undergoing a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
 - 5.1 Pregnant: and
 - 5.2 Any of the following:

Subsidy	/ Full	/ Brand or
(Manufacturer'	s Price) Subsidise	d Generic
\$	Per 💌	Manufacturer

continued...

- 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or
- 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
- 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

Renewal — (Short-term medical condition) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Has undergone a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
 - 5.1 Pregnant; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or
 - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
 - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

Initial application — (Long-term medical condition) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis: or
- 3 Liver disease; or
- 4 Chronic Renal failure: or
- 5 Inflammatory bowel disease: or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome: or
- 8 Bowel fistula: or
- 9 Severe chronic neurological conditions; or
- 10 Epidermolysis bullosa; or
- 11 AIDS (CD4 count < 200 cells/mm³); or
- 12 Chronic pancreatitis.

Renewal — (Chronic disease OR tube feeding for patients who have previously been funded under Special Authority forms SA0702 or SA0583) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube - refer to specific medical condition criteria); or

continued...

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

- 2 Cystic Fibrosis; or
- 3 Liver disease; or
- 4 Chronic Renal failure; or
- 5 Inflammatory bowel disease; or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome; or
- 8 Bowel fistula; or
- 9 Severe chronic neurological conditions.

9 Severe chronic neurological conditions.			
ENTERAL FEED 1.5KCAL/ML – Special Authority see SA1859 on p Liquid	•	Hospital pharmac 250 ml OP 1,000 ml OP	y [HP3] ✓ Ensure Plus HN ✓ Ensure Plus HN RTH ✓ Nutrison Energy ✓ Fresubin HP Energy
ENTERAL FEED 1KCAL/ML – Special Authority see SA1859 on pag Liquid	•	spital pharmacy 250 ml OP 1,000 ml OP	[HP3] ✓ Isosource Standard ✓ Fresubin Original ✓ Osmolite RTH ✓ Nutrison RTH
ENTERAL FEED WITH FIBRE 0.83 KCAL/ML — Special Authority se Liquid		on page 277 – Ho 1,000 ml OP	ospital pharmacy [HP3] Nutrison 800 Complete Multi Fibre
ENTERAL FEED WITH FIBRE 1 KCAL/ML – Special Authority see S Liquid	6.56 7.00	page 277 – Hosp 1,000 ml OP	ital pharmacy [HP3] ✓ Jevity RTH ✓ Fresubin Original Fibre
ENTERAL FEED WITH FIBRE 1.2KCAL/ML - Special Authority see		page 277 – Hos 1,000 ml OP	✓ Nutrison Multi Fibre pital pharmacy [HP3] ✓ Jevity Plus RTH
ENTERAL FEED WITH FIBRE 1.5KCAL/ML — Special Authority see Liquid		page 277 – Hos 1,000 ml OP	pital pharmacy [HP3] ✓ Jevity HiCal RTH ✓ Nutrison Energy Multi Fibre
	9.80		Fresubin HP Energy Fibre
ENTERAL FEED WITH PROTEIN 1.2KCAL/ML - Special Authority Liquid	9.60	500 ml OP	✓ Fresubin Intensive
ORAL FEED (POWDER) – Special Authority see SA1859 on page 2 Powder (chocolate)		al pharmacy [HP 840 g OP	3]✓ Sustagen Hospital Formula
Powder (vanilla)	26.00 14.00	850 g OP 840 g OP	✓ Ensure✓ Sustagen HospitalFormula Active
	26.00	850 g OP	✓ Ensure

(Mar	Subsidy nufacturer's Price)	Ful Subsidise	,	Brand or Generic
(Iviai			_	
	\$	Per •	/	Manutacturer

ORAL FEED 1.5KCAL/ML - Special Authority see SA1859 on page 277 - Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, who have severe epidermolysis bullosa, or as exclusive enteral nutrition in children under the age of 18 years for the treatment of Crohn's disease, or for patients with COPD and hypercapnia, defined as CO2 value exceeding 55mmHg. The prescription must be endorsed accordingly.

Liquid (banana) – Higher subsidy of up to \$1.56 per 200 ml with Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip
	(1.56)		Ensure Plus
Liquid (chocolate) - Higher subsidy of up to \$1.56 per 200 ml			
with Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip
	(1.56)		Ensure Plus
Liquid (fruit of the forest) - Higher subsidy of \$1.56 per 200 ml			
with Endorsement	0.72	200 ml OP	
	(1.56)		Ensure Plus
Liquid (strawberry) - Higher subsidy of \$1.26 per 200 ml with	, ,		
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip
Liquid (vanilla) - Higher subsidy of up to \$1.65 per 237 ml with	,		·
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip
	0.85	237 ml OP	
	(1.65)		Ensure Plus
	0.72	200 ml OP	
	(1.56)		Ensure Plus
	()		

ORAL FEED WITH FIBRE 1.5 KCAL/ML – Special Authority see SA1859 on page 277 – Hospital pharmacy [HP3]
Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, or who have severe epidermolysis bullosa. The prescription must be endorsed accordingly.

Liquid (chocolate) – Higher subsidy of \$1.26 per 200 ml with

Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip Multi Fibre
Liquid (strawberry) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip Multi Fibre
Liquid (vanilla) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip Multi Fibre

High Calorie Products

⇒SA1195 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

All of the following:

- 1 Cystic fibrosis; and
- 2 other lower calorie products have been tried; and

continued...

SPECIAL FOODS

Subsid (Manufacturer		ully Brand or sed Generic	
\$	Per	✓ Manufactur	er

continued...

3 patient has substantially increased metabolic requirements.

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 any condition causing malabsorption; or
 - 1.2 faltering growth in an infant/child; or
 - 1.3 increased nutritional requirements; or
 - 1.4 fluid restricted: and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements or is fluid restricted.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL FEED 2 KCAL/ML - Special Aut	thority see SA1195 on the previous pag	ge – Hospital pl	narmacy [HP3]
Liquid	6.50	500 ml OP	✓ Fresubin 2kcal HP
·	6.82		✓ Nutrison
			Concentrated
	13.64	1,000 ml OP	✓ Ensure Two Cal HN RTH

ORAL FEED 2 KCAL/ML - Special Authority see SA1195 on the previous page - Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, or who have severe epidermolysis bullosa. The prescription must be endorsed accordingly.

Liquid (vanilla) - Higher subsidy of \$2.34 per 200 ml with

(2.34) Two Cal HN

Food Thickeners

⇒SA1106 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient has motor neurone disease with swallowing disorder.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for

Subsidy (Manufacturer's Price)	Full Subsidise	
\$	Per •	Manufacturer

continued...

applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Gluten Free Foods

The funding of gluten free foods is no longer being actively managed by Pharmac from 1 April 2011. This means that we are no longer considering the listing of new products, or making subsidy, or other changes to the existing listings. As a result we anticipate that the range of funded items will reduce over time. Management of Coeliac disease with a gluten free diet is necessary for good outcomes. A range of gluten free options are available through retail outlets.

⇒SA1729 Special Authority for Subsidy

Initial application — (all patients) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Either:

- 1 Gluten enteropathy has been diagnosed by biopsy; or
- 2 Patient suffers from dermatitis herpetiformis.

Initial application — (paediatric patients diagnosed by ESPGHAN criteria) only from a paediatric gastroenterologist. Approvals valid without further renewal unless notified where the paediatric patient fulfils ESPGHAN criteria for biopsy free diagnosis of coeliac disease.

GLUTEN ERFE BAKING MIX - Special Authority see SA1729 above - Hospital pharmacy [HP3]

GEOTERT FILE DARRING WITH	openia nationty see of 1720 above Troopital priamacy [111 o]	
Powder	2.81 1,000 g OP	
	(5.15)	Healtheries Simple Baking Mix
GLUTEN FREE BREAD MIX -	- Special Authority see SA1729 above – Hospital pharmacy [HP3]	
Powder	3.93 1,000 g OP	
	(7.32)	NZB Low Gluten Bread Mix
	3.51	
	(10.87)	Horleys Bread Mix
GLUTEN FREE FLOUR - Spe	ecial Authority see SA1729 above – Hospital pharmacy [HP3]	
Powder	5.62 2,000 g OP	
	(18.10)	Horleys Flour

	Subsidy		Fully	Brand or
	(Manufacturer's Prices)	ce) Sub Per	sidised	Generic Manufacturer
	*			
GLUTEN FREE PASTA – Special Authority see SA1729 on the			macy [HF	23]
Buckwheat Spirals		250 g OP	_	
	(3.11)		0	rgran
Corn and Vegetable Shells		250 g OP		
	(2.92)		0	rgran
Corn and Vegetable Spirals	2.00	250 g OP		
	(2.92)		0	rgran
Rice and Corn Lasagne Sheets	1.60	200 g OP		
	(3.82)		0	rgran
Rice and Corn Macaroni	2.00	250 g OP		
	(2.92)	-	0	rgran
Rice and Corn Penne	2.00	250 g OP		•
	(2.92)	•	0	rgran
Rice and Maize Pasta Spirals	2.00 [°]	250 g OP		
·	(2.92)	Ū	0	rgran
Rice and Millet Spirals	2.00 [′]	250 g OP		
•	(3.11)	J	0	rgran
Rice and corn spaghetti noodles	` '	375 g OP		3
1	(2.92)		0	rgran
Vegetable and Rice Spirals	` '	250 g OP	_	3 ··
- g - · · · · · · · · · · · · · · · · · ·	(2.92)		0	rgran
Italian long style spaghetti	` '	220 g OP	·	· J· · · ··
	(3.11)		0	rgran
	(3.11)		·	. 3

Foods And Supplements For Inherited Metabolic Disease

⇒SA2300 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 Dietary management of inherited metabolic disease; or
- 2 For use as a supplement to a Ketogenic diet in patients diagnosed with epilepsy.

Supplements For Homocystinuria

AMINOACID FORMULA WITHOUT METHIONINE	- Special Authority see SA2300	above -	Hospital pharmacy [HP3]
Powder, 12.5 g sachets	349.65	30	✓ HCU Explore 5
Powder, 25 g sachets	1,048.95	30	✓ HCU Express 15
Powder	461.94	500 a OF	✓ XMET Maxamum

Supplements For MSUD and short chain enoyl coA hydratase deficiency

AMINOACID FORMULA WITHOUT VALINE, LEUCINE AND ISOLEUCINE - Special Authority see SA2300 above - Hospital pharmacy [HP3]

Powder, 12.5 g sachets	349.65	30	✓ MSUD Explore 5
Powder, 25 g sachets	1,048.95	30	✓ MSUD Express 15
Powder	437.22	500 g OP	✓ MSUD Maxamum

Subsidy (Manufacturer's Price)

Fully Subsidised Per

Brand or Generic Manufacturer

Supplements For PKU

AMINOACID FORMULA WITHOUT PHENYLALANINE - Special Authority see \$A2300 on the previous page - Hospital pharmacy [HP3]

iailliacy [i ii o]			
Tabs	99.00	75 OP	✓ Phlexy 10
Powder (Lemon), 34 g sachets	883.50	30	✓ PKU Express 20
Powder (Neutral), 12.5 g sachets	220.88	30	✓ PKU Explore 5
Powder (Neutral), 34 g sachets	883.50	30	✓ PKU Express 20
Powder (Orange), 25 g sachets	441.75	30	✓ PKU Explore 10
Powder (Orange), 34 g sachets	883.50	30	✓ PKU Express 20
Powder (Raspberry), 25 g sachets		30	✓ PKU Explore 10
Powder (Tropical), 34 g sachets		30	✓ PKU Express 20
Powder (berry) 28 g sachets		30	✓ PKU Lophlex
, ,, ,			Powder
Powder (chocolate) 36 g sachet	393.00	30	PKU Anamix Junior
			Chocolate
Powder (neutral) 28 g sachets	936.00	30	✓ PKU Lophlex
(, 3			Powder
Powder (neutral) 36 g sachets	393.00	30	✓ PKU Anamix Junior
Powder (orange) 28 g sachets	936.00	30	✓ PKU Lophlex
(0 / 0			Powder
Powder (orange) 36 g sachet	393.00	30	✓ PKU Anamix Junior
(3 / 3			Orange
Powder (vanilla) 36 g sachet	393.00	30	✓ PKU Anamix Junior
, ,			Vanilla
Infant formula	174.72	400 g OP	✓ PKU Anamix Infant
Powder (neutral)	178.79	400 g OP	✓ PKU Start
Powder (orange)		500 g OP	✓ XP Maxamum
Powder (unflavoured)		500 g OP	✓ XP Maxamum
Liquid (berry)		125 ml OP	✓ PKU Anamix Junior
4			LQ
Liquid (orange)	13.10	125 ml OP	✓ PKU Anamix Junior
1 (3 /			LQ
Liquid (unflavoured)	13.10	125 ml OP	✓ PKU Anamix Junior
1 (LQ
Liquid (forest berries), 250 ml carton	540.00	18 OP	 Easiphen Liquid
Liquid (juicy tropical) 125 ml		30 OP	✓ PKU Lophlex LQ 20
Oral semi-solid (berries) 109 g		36 OP	✓ PKU Lophlex
, , , ,			Sensation 20
Liquid (juicy berries) 62.5 ml	939.00	60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy citrus) 62.5 ml		60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy orange) 62.5 ml		60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy berries) 125 ml		30 OP	✓ PKU Lophlex LQ 20
Liquid (juicy orange) 125 ml		30 OP	✓ PKU Lophlex LQ 20
			•

	Subsidy		ully	Brand or
	(Manufacturer's Price		ised	Generic
	<u> </u>	Per		Manufacturer
GLYCOMACROPEPTIDE AND AMINO ACID CONTAINS SOME	E PHENYLALANINE	- Special Au	uthor	ity see SA2300 on
page 284 – Hospital pharmacy [HP3]				
Powder (Banana) 35 g sachets	930.00	30	1	PKU
, ,				sphere20 Banana
Powder (Berry), 20 g sachets	440 28	60	1	PKU Restore
Towaer (Derry), 20 g sacriets	443.20	00		Powder
D 1 (0) 11 \ 00 0 1 1	000 50			
Powder (Chocolate) 32 g Sachets	898.56	30	•	PKU Build
				20 Chocolate
Powder (Chocolate) 35 g sachets	930.00	30		PKU
				sphere20 Chocolate
Powder (Lemon) 35 g sachets	930.00	30	/	PKU
				sphere20 Lemon
Powder (Lemonade) 33.4 g sachets	936.00	30	1	PKU GMPro Ultra
				Lemonade
Powder (Neutral), 16 g sachets	440.00	30	1	PKU Build 10
, ,				PKU Restore
Powder (Orange), 20 g sachets	449.28	60	•	
			_	Powder
Powder (Raspberry Lemonade) 32 g Sachets	898.56	30		PKU Build
				20 Raspberry
				Lemonade
Powder (Smooth) 32 g Sachets	898.56	30	1	PKU Build
· · · · · · · · · · · · · · · ·				20 Smooth
Powder (Vanilla) 32 g Sachets	808 56	30	1	PKU Build 20 Vanilla
Powder (varilla), 40 g sachets		30		Camino Pro
rowder (fledital), 40 g Sacriets	073.92	30		
2 1 (2 12) 22				Bettermilk
Powder (Red Berry) 35 g sachets	930.00	30	•	PKU sphere20 Red
				Berry
Powder (Vanilla) 35 g sachets	930.00	30	/	PKU
				sphere20 Vanilla
Liquid (Coffee Mocha), 250 ml carton	684.45	30 OP	1	PKU Glytactin RTD
4 (15 Lite
Liquid (chocolate), 250 ml carton	684.45	30 OP	1	PKU Glytactin RTD
Liquid (Chocolate), 200 mi carton	004.43	30 01		15
1' '1' ' ' ' ' ' ' ' ' ' ' ' ' ' ' ' '	004.45	00.00		
Liquid (neutral), 250 ml carton	684.45	30 OP	•	PKU Glytactin RTD
				15
Liquid (vanilla), 250 ml carton	684.45	30 OP		PKU Glytactin RTD
				15 Lite
Foods				
LOW PROTEIN BAKING MIX - Special Authority see SA2300 o		al pharmacy	[HP3	3]
Powder	8.55	500 g OP		Loprofin Mix
LOW PROTEIN PASTA - Special Authority see SA2300 on pag	ie 284 – Hospital pha	armacy [HP3]		
Animal shapes		500 g OP		Loprofin
Lasagne		250 g OP		Loprofin
Low protein rice pasta		500 g OP		Loprofin
		•		
Macaroni		250 g OP		Loprofin
Penne		500 g OP		Loprofin
Spaghetti		500 g OP		Loprofin
Spirals	12.39	500 g OP	•	Loprofin

Subsidy

Fully

Brand or

	Subsidy (Manufacturer's Price)	Subsid	-ully ised	Brand or Generic
	\$	Per	1	Manufacturer
Supplements for Tyrosinaemia				
AMINOACID FORMULA WITHOUT PHENYLALANINE AND TYF pharmacy [HP3]	ROSINE - Special Au	thority see	SA230	0 on page 284 – Hospita
Powder (Neutral), 12.5 g sachets		30		R Explore 5
GLYCOMACROPEPTIDE AND AMINO ACID CONTAINS SOME SA2300 on page 284 – Hospital pharmacy [HP3]		ENYLALAN	IINE –	Special Authority see
Powder (Red Berry), 35 g sachets		30 30		R Sphere 20
Powder (Vanilla), 35 g sachets	1,398.60	30	V 11	'R Sphere 20
Supplements for Organic Acidaemias				
AMINOACID FORMULA WITHOUT METHIONINE, THREONINE Hospital pharmacy [HP3]		cial Authori	ty see	SA2300 on page 284 –
Powder, 12.5 g sachets		30		MA/PA Explore 5
Powder, 25 g sachets	1,048.95	30	✓ MI	MA/PA Express 15
Supplements for Glutaric Aciduria type 1				
AMINOACID FORMULA WITHOUT LYSINE - Special Authority Powder, 12.5 g sachets		284 – Hosp 30		armacy [HP3] A Explore 5
Supplements for Glycogen Storage Disease				
HIGH AMYLOPECTIN CORN-STARCH - Special Authority see Powder, 60 g sachets		– Hospital 30		acy [HP3] ycosade
Single dose amino acids				
ARGININE – Special Authority see SA2300 on page 284 – Hosp Powder, 4 g sachets		30	✓ Ar	ginine2000
CITRULLINE - Special Authority see SA2300 on page 284 - Ho Powder, 4 g sachets	211.45	30	✓ Cit	trulline1000
ISOLEUCINE – Special Authority see SA2300 on page 284 – Ho Powder, 4 g sachets		30 30	✓ Iso	pleucine50
LEUCINE - Special Authority see SA2300 on page 284 - Hospit Powder, 4 g sachets		30	✓ Le	ucine100
PHENYLALANINE – Special Authority see SA2300 on page 284 Powder, 4 g sachets		[HP3] 30	✓ Ph	enylalanine50
TYROSINE – Special Authority see SA2300 on page 284 – Hosp Powder, 4 g sachets		30	✓ Ту	rosine1000
VALINE - Special Authority see SA2300 on page 284 - Hospital Powder, 4 g sachets		30	✔ Va	line50

Subsidy (Manufacturer's Price) \$

Fully Subsidised Per

Brand or Generic Manufacturer

Infant Formulae

For Williams Syndrome

⇒SA1110 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is an infant suffering from Williams Syndrome and associated hypercalcaemia.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

1 The treatment remains appropriate and the patient is benefiting from treatment; and

AMINO ACID FORMULA - Special Authority see SA2092 below - Hospital pharmacy [HP3]

2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

LOW CALCIUM INFANT FORMULA - Special Authority see SA1110 above - Hospital pharmacy [HP3]
Powder46.18 400 g OP ✓ Locasol

Gastrointestinal and Other Malabsorptive Problems

VIII VO AOID I OI IIVIOLA	opecial Authority see OAZOOZ Delow	i iospitai piiai	macy [mo]	
Powder		43.60	400 g OP	✓ Alfamino
			_	 Alfamino Junior
Powder (unflavoured).		55.61	400 g OP	✓ Neocate Gold
		-	 Neocate Junior Unflavoured 	
				✓ Neocate SYNEO
		65.72		✓ Elecare
				✓ Elecare LCP
Powder (vanilla)		55.61	400 g OP	✓ Neocate Junior Vanilla
		65.72		✓ Elecare

⇒SA2092 Special Authority for Subsidy

Initial application — (Infants under 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 History of anaphylaxis to cow's milk protein formula or dairy products; or
- 2 Eosinophilic oesophagitis; or
- 3 Ultra-short gut: or
- 4 Severe Immune deficiency; or
- 5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
- 6 Both
 - 6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 6.2 Fither:
 - 6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or 6.2.2 Patient has IgE mediated allergy.

Initial application — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sul	osidised	Generic	
\$	Per	1	Manufacturer	

immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or
 - 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
 - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products; or
 - 2.2 Eosinophilic oesophagitis; or
 - 2.3 Ultra-short gut; or
 - 2.4 Severe Immune deficiency; or
 - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
 - 2.6 Both:
 - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 2.6.2 Either:
 - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number: or
 - 2.6.2.2 Patient has IgE mediated allergy.

Renewal — (Infants up to 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has IgE mediated allergy; and
 - 1.2 All of the following:
 - 1.2.1 Patient remains allergic to cow's milk; and
 - 1.2.2 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy or extensively hydrolysed infant formula has been undertaken; and
 - 1.2.3 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
 - 1.2.4 Amino acid formula is required for a nutritional deficit; and
 - 1.2.5 It has been more than three months from the previous approval; or

2 Both:

- 2.1 Patient has non IgE mediated severe gastrointestinal intolerance (including eosinophilic oesophagitis, ultra-short gut and severe immune deficiency); and
- 2.2 All of the following:
 - 2.2.1 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy, or extensively hydrolysed infant formula has been undertaken; and
 - 2.2.2 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
 - 2.2.3 Amino acid formula is required for a nutritional deficit; and
 - 2.2.4 It has been more than three months from the previous approval.

Renewal — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or

continued...

✓ fully subsidised 289

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	/	Manufacturer

- 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
 - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products: or
 - 2.2 Eosinophilic oesophagitis; or
 - 2.3 Ultra-short gut; or
 - 2.4 Severe Immune deficiency; or
 - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
 - 2.6 Both:
 - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 2.6.2 Either:
 - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or
 - 2.6.2.2 Patient has IgE mediated allergy.

Initial application — (for patients who have a current funding under Special Authority form SA1557) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a valid Special Authority approval for extensively hydrolysed formula (SA1557); and
- 2 Extensively hydrolysed formula (Aptamil Gold+ Pepti Junior, AllerPro SYNEO 1 and 2) is unable to be supplied at this time; and
- 3 The approval only applies to funded dispensings of Neocate Gold and Neocate Syneo.

Note: This criteria is short term funding to cover an out-of-stock situation on some extensively hydrolysed formula powder funded under Special Authority form SA1557. There is no renewal criteria under this restriction.

SA1953 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has impaired gastrointestinal function and either cannot tolerate polymeric feeds, or polymeric feeds are unsuitable; and
- 2 Any of the following:
 - 2.1 Severe malabsorption; or
 - 2.2 Short bowel syndrome; or
 - 2.3 Intractable diarrhoea: or
 - 2.4 Biliary atresia; or
 - 2.5 Cholestatic liver diseases causing malabsorption: or
 - 2.6 Cystic fibrosis; or
 - 2.7 Proven fat malabsorption; or
 - 2.8 Severe intestinal motility disorders causing significant malabsorption; or
 - 2.9 Intestinal failure: or
 - 2.10 Both:
 - 2.10.1 The patient is currently receiving funded amino acid formula; and

continued...

Subsidy (Manufacturer's Pric	ce)	Fully Subsidised	Brand or Generic	
\$	Per	✓	Manufacturer	

- 2.10.2 The patient is to be trialled on, or transitioned to, an enteral liquid peptide formula; and
- 3 Either:
 - 3.1 A semi-elemental or partially hydrolysed powdered feed has been reasonably trialled and considered unsuitable; or
 - 3.2 For step down from intravenous nutrition.

Note: A reasonable trial is defined as a 2-4 week trial.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 An assessment as to whether the patient can be transitioned to a cows milk protein or soy infant formula or extensively hydrolysed formula has been undertaken; and
- 2 The outcome of the assessment is that the patient continues to require an enteral liquid peptide formula; and
- 3 General practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

⇒SA1557 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 Cows milk formula is inappropriate due to severe intolerance or allergy to its protein content; and
 - 1.2 Either:
 - 1.2.1 Soy milk formula has been reasonably trialled without resolution of symptoms; or
 - 1.2.2 Soy milk formula is considered clinically inappropriate or contraindicated; or
- 2 Severe malabsorption: or
- 3 Short bowel syndrome: or
- 4 Intractable diarrhoea: or
- 5 Biliary atresia: or
- 6 Cholestatic liver diseases causing malsorption; or
- 7 Cystic fibrosis: or
- 8 Proven fat malabsorption; or
- 9 Severe intestinal motility disorders causing significant malabsorption; or
- 10 Intestinal failure; or
- 11 All of the following:
 - 11.1 For step down from Amino Acid Formula; and
 - 11.2 The infant is currently receiving funded amino acid formula; and
 - 11.3 The infant is to be trialled on, or transitioned to, an extensively hydrolysed formula; and
 - 11.4 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

Note: A reasonable trial is defined as a 2-4 week trial, or signs of an immediate IgE mediated allergic reaction.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

continued...

✓ fully subsidised 291



	Subsidy		Fully	Brand or
(Ma	anufacturer's Price)	Sı	ubsidised	Generic
	\$	Per	✓	Manufacturer

- 1 An assessment as to whether the infant can be transitioned to a cows milk protein or soy infant formula has been undertaken; and
- 2 The outcome of the assessment is that the infant continues to require an extensively hydrolysed infant formula; and
- 3 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

Fluid Restricted

⇒SA1698 Special Authority for Subsidy

Initial application only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is fluid restricted or volume intolerant and has been diagnosed with faltering growth; and
- 2 Patient is under the care of a paediatrician or dietitian who has recommended treatment with a high energy infant formula; and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: "Volume intolerant" patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

Renewal only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient continues to be fluid restricted or volume intolerant and has faltering growth; and
- 2 Patient is under the care of a hospital paediatrician or dietitian who has recommended treatment with a high energy infant formula; and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: "Volume intolerant" patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

Ketogenic Diet

⇒SA1197 Special Authority for Subsidy

Initial application only from a metabolic physician or paediatric neurologist. Approvals valid for 3 months where the patient has intractable epilepsy, pyruvate dehydrogenase deficiency or glucose transported type-1 deficiency and other conditions requiring a ketogenic diet.

Renewal only from a metabolic physician or paediatric neurologist. Approvals valid for 2 years where the patient is on a ketogenic diet and the patient is benefiting from the diet.

HIGH FAT LOW CARBOHYDRATE FORMULA - Special Authority see SA1197 above - Retail pharmacy

Powder (unflavoured)36.92	300 g OP	✓ KetoCal 4:1
		✓ Ketocal 3:1
Powder (vanilla)36.92	300 g OP	✓ KetoCal 4:1

SECTION I: NATIONAL IMMUNISATION SCHEDULE

Subsidy (Manufacturer's Price) \$

Subsidised Per

Fully

Brand or Generic Manufacturer

Vaccinations

BACILLUS CALMETTE-GUERIN VACCINE - [Xpharm]

For infants at increased risk of tuberculosis. Increased risk is defined as:

- 1) living in a house or family with a person with current or past history of TB; or
- 2) having one or more household members or carers who within the last 5 years lived in a country with a rate of TB > or equal to 40 per 100,000 for 6 months or longer; or
- 3) during their first 5 years will be living 3 months or longer in a country with a rate of TB > or equal to 40 per 100,000 Note a list of countries with high rates of TB are available at www.health.govt.nz/tuberculosis (search for downloads) or www.bcgatlas.org/index.php.
- Inj Mycobacterium bovis BCG (Bacillus Calmette-Guerin),

Danish strain 1331, live attenuated, vial with diluent...............................0.00

10

✓ BCG Vaccine

DIPHTHERIA. TETANUS AND PERTUSSIS VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- c)
- A) Funded for any of the following criteria:
 - 1) A single dose for pregnant women in the second or third trimester of each pregnancy; or
 - 2) A single dose for parents or primary caregivers of infants admitted to a Neonatal Intensive Care Unit or Specialist Care Baby Unit for more than 3 days, who had not been exposed to maternal vaccination at least 14 days prior to birth; or
 - A course of up to four doses is funded for children from age 7 up to the age of 18 years inclusive to complete full primary immunisation; or
 - 4) An additional four doses (as appropriate) are funded for (re-)immunisation for patients post haematopoietic stem cell transplantation or chemotherapy; pre or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or
 - 5) A single dose for vaccination of patients aged from 65 years old; or
 - 6) A single dose for vaccination of patients aged from 45 years old who have not had 4 previous tetanus doses; or
 - 7) For vaccination of previously unimmunised or partially immunised patients; or
 - 8) For revaccination following immunosuppression; or
 - 9) For boosting of patients with tetanus-prone wounds.

Notes: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.

- B) Contractors will be entitled to claim payment from the Funder for the supply of diphtheria, tetanus and pertussis vaccine to patients eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the diphtheria, tetanus and pertussis vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraphs 1 – 9 above.
- Inj 2 IU diphtheria toxoid with 20 IU tetanus toxoid, 8 mcg

pertussis toxoid. 8 mcg pertussis filamentous

10

✓ Boostrix

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

DIPHTHERIA. TETANUS. PERTUSSIS AND POLIO VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- c) A) Funded for any of the following:
 - 1) A single dose for children up to the age of 7 who have completed primary immunisation; or
 - 2) A course of four vaccines is funded for catch up programmes for children (to the age of 10 years) to complete full primary immunisation; or
 - 3) An additional four doses (as appropriate) are funded for (re-)immunisation for people post HSCT, or chemotherapy; pre- or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or
 - 4) Five doses will be funded for children requiring solid organ transplantation.
 - B) Contractors will be entitled to claim payment from the Funder for the supply of Diphtheria, tetanus, pertussis and polio vaccine to people eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Diphtheria, tetanus, pertussis and polio vaccine listed in the Pharmaceutical Schedule.
 - C) Contractors may only claim for populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.

Ini 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg pertussis toxoid, 25 mcg pertussis filamentous haemagglutinin, 8 mcg pertactin and 80 D-antigen units

10 ✓ Infanrix IPV

DIPHTHERIA, TETANUS, PERTUSSIS, POLIO, HEPATITIS B AND HAEMOPHILUS INFLUENZAE TYPE B VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- c) A) Funded for children meeting any of the following criteria
 - 1) Up to four doses for children up to and under the age of 10 for primary immunisation; or
 - 2) An additional four doses (as appropriate) are funded for (re-)immunisation for children up to and under the age of 10 who are patients post haematopoietic stem cell transplantation, or chemotherapy; pre or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens: or
 - 3) Up to five doses for children up to and under the age of 10 receiving solid organ transplantation.
 - B) Contractors will be entitled to claim payment from the Funder for the supply of Diphtheria, tetanus, pertussis, polio, hepatitis B and haemophilus influenzae type b vaccine to people eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Diphtheria, tetanus, pertussis, polio, hepatitis B and haemophilus influenzae type b vaccine listed in the Pharmaceutical Schedule.
 - C) Contractors may only claim for populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Note: A course of up-to four vaccines is funded for catch up programmes for children (up to and under the age of 10 years) to complete full primary immunisation. Please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes.

Inj 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg pertussis toxoid, 25 mcg pertussis filamentous haemagglutinin, 8 mcg pertactin, 80 D-Ag U polio virus,

10 mcg hepatitis B surface antigen in 0.5 ml syringe0.00

10 ✓ Infanrix-hexa

lubsidy	Fully	Brand or
cturer's Price) Su	ibsidised	Generic
 \$ Per	✓	

HAEMOPHILUS INFLUENZAE TYPE B VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- c) A) One dose for people meeting any of the following:
 - 1) For primary vaccination in children: or
 - 2) An additional dose (as appropriate) is funded for (re-)immunisation for people post haematopoietic stem cell transplantation, or chemotherapy; functional asplenic; pre or post splenectomy; pre- or post solid organ transplant, pre or post cochlear implants, renal dialysis and other severely immunosuppressive regimens; or
 - For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine
 physician or paediatrician.
 - B) Contractors will be entitled to claim payment from the Funder for the supply of Haemophilus influenzae type b vaccine to people eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Haemophilus influenzae type b vaccine listed in the Pharmaceutical Schedule.
 - C) Contractors may only claim for populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Haemophilus Influenzae type B polysaccharide 10 mcg conjugated to tetanus toxoid as carrier protein 20-40 mcg; prefilled syringe plus vial 0.5 ml	1 ✓ Hiberix
HEPATITIS A VACCINE - [Xpharm]	
Funded for patients meeting any of the following criteria:	
1) Two vaccinations for use in transplant patients; or	
2) Two vaccinations for use in children with chronic liver disease; or	
One dose of vaccine for close contacts of known hepatitis A cases.	
	_
Inj 1440 ELISA units in 1 ml syringe	1 ✓ <u>Havrix</u>
Inj 720 ELISA units in 0.5 ml syringe0.00 1	1 ✓ <u>Havrix Junior</u>

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer	
HEPATITIS B RECOMBINANT VACCINE – [Xpharm] Inj 10 mcg per 0.5 ml prefilled syringe	0.00	1	√ E	ingerix-B	

- Funded for patients meeting any of the following criteria:
 - 1) for household or sexual contacts of known acute hepatitis B patients or hepatitis B carriers; or
 - 2) for children born to mothers who are hepatitis B surface antigen (HBsAq) positive; or
 - for children up to and under the age of 18 years inclusive who are considered not to have achieved a positive serology and require additional vaccination or require a primary course of vaccination; or
 - 4) for HIV positive patients; or
 - 5) for hepatitis C positive patients; or
 - 6) for patients following non-consensual sexual intercourse: or
 - 7) for patients following immunosuppression; or
 - 8) for solid organ transplant patients; or
- 9) for post-haematopoietic stem cell transplant (HSCT) patients; or
- 10) following needle stick injury.

- 1) for household or sexual contacts of known acute hepatitis B patients or hepatitis B carriers; or
- 2) for children born to mothers who are hepatitis B surface antigen (HBsAg) positive; or
- for children up to and under the age of 18 years inclusive who are considered not to have achieved a positive serology and require additional vaccination or require a primary course of vaccination; or
- 4) for HIV positive patients; or
- 5) for hepatitis C positive patients; or
- 6) for patients following non-consensual sexual intercourse; or
- 7) for patients following immunosuppression; or
- 8) for solid organ transplant patients; or
- 9) for post-haematopoietic stem cell transplant (HSCT) patients; or
- 10) following needle stick injury; or
- 11) for dialysis patients; or
- 12) for liver or kidney transplant patients.

HUMAN PAPILLOMAVIRUS (6, 11, 16, 18, 31, 33, 45, 52 AND 58) VACCINE [HPV]

- a) Maximum of 1 inj per prescription
- b) Only on a prescription
- c) No patient co-payment payable
- d)
- a) A) Any of the following:
 - 1) Maximum of two doses for children aged 14 years and under; or
 - 2) Maximum of three doses for patients meeting any of the following criteria:
 - 1) People aged 15 to 26 years inclusive; or
 - 2) Either:

People aged 9 to 26 years inclusive

- 1) Confirmed HIV infection; or
- 2) Transplant (including stem cell) patients: or
- 3) Maximum of four doses for people aged 9 to 26 years inclusive post chemotherapy
- B) Contractors will be entitled to claim payment from the Funder for the supply of Human papillomavirus vaccine to patients eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Human papillomavirus vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraphs A above.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
INFLUENZA VACCINE Inj 60 mcg in 0.5 ml syringe (quadrivalent vaccine)	120.00	10		fluvac Tetra (2024 formulation)

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

- a) Maximum of 1 inj per prescription
- b) Only on a prescription
- c) No patient co-payment payable
- ď

A) INFLUENZA VACCINE

is available each year for patients who meet the following criteria, as set by Pharmac:

- a) all people 65 years of age and over; or
- b) people under 65 years of age who:
 - i) have any of the following cardiovascular diseases:
 - a) ischaemic heart disease, or
 - b) congestive heart failure, or
 - c) rheumatic heart disease, or
 - d) congenital heart disease, or
 - e) cerebo-vascular disease; or
 - ii) have either of the following chronic respiratory diseases:
 - a) asthma, if on a regular preventative therapy, or
 - b) other chronic respiratory disease with impaired lung function; or
 - iii) have diabetes; or
 - iv) have chronic renal disease; or
 - v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
 - vi) have any of the following other conditions:
 - a) autoimmune disease, or
 - b) immune suppression or immune deficiency, or
 - c) HIV, or
 - d) transplant recipients, or
 - e) neuromuscular and CNS diseases/disorders, or
 - f) haemoglobinopathies, or
 - g) are children on long term aspirin, or
 - h) have a cochlear implant, or
 - i) errors of metabolism at risk of major metabolic decompensation, or
 - i) pre and post splenectomy, or
 - k) Down syndrome, or
 - vii) are pregnant; or
- c) children 4 years of age and under who have been hospitalised for respiratory illness or have a history of significant respiratory illness; or
- d) people under 65 years of age who:
 - i) have any of the following serious mental health conditions:
 - a) schizophrenia, or
 - b) major depressive disorder, or
 - c) bipolar disorder, or
 - d) schizoaffective disorder, or
 - ii) are currently accessing secondary or tertiary mental health and addiction services; or

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy,
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Contractors will be entitled to claim payment for the supply of influenza vaccine to patients eligible under the above criteria pursuant to their contract with Health NZ for subsidised immunisation, and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

		Subsidy		Fully	Brand or
		(Manufacturer's Price)	_	Subsidised	Generic
		\$	Per		Manufacturer
MEASL	ES, MUMPS AND RUBELLA VACCINE				
a)	Only on a prescription				
b)	No patient co-payment payable				
c)					
	Measles, mumps and rubella vaccine				
	A maximum of two doses for any patient meeting	the following criteria:			
	 For primary vaccination in children; or 				
	For revaccination following immunosuppress				
	For any individual susceptible to measles, m				
	4) A maximum of three doses for children who				
	Note: Please refer to the Immunisation Handbook				
	price is listed for the vaccine, doctors can still orde	er measles mumps and	rube	lla vaccine fi	ree of charge, as with
	other Schedule vaccines.				
	B) Contractors will be entitled to claim payment for the				
	eligible under the above criteria pursuant to their o				
	only do so in respect of the measles, mumps and				
	C) Contractors can only claim for patient populations		are co	overed by the	eir contract, which may
lni	a sub-set of the population described in paragraph				
Irij,	measles virus 1,000 CCID50, mumps virus 5,012 CCID Rubella virus 1,000 CCID50; prefilled syringe/ampoule				
	diluent 0.5 ml		10	√ D	riorix
			10	<u> </u>	HOHA
	GOCOCCAL (GROUPS A, C, Y AND W-135) CONJUG	ATE VACCINE			
,	Only on a prescription				
,	No patient co-payment payable				
c)					
	A) Any of the following:				
	Up to three doses and a booster every five y				
	functional or anatomic asplenia, HIV, comple	ment deficiency (acqui	rea o	r inneritea),	or pre or post solid orga
	transplant; or				
	One dose for close contacts of meningococc One dose for person who has proviously been			ony arour: a	\r_
	 One dose for person who has previously had A maximum of two doses for bone marrow tr 		e of	any group; c	Л
	4) A Maximum of two doses for done marrow tr	anspiant batterns, or			

B) Both:1) Person is aged between 13 and 25 years, inclusive; and

5) A maximum of two doses for person pre- and post-immunosuppression*; or

- Either:

 2) Either:
 - One dose for individuals who are entering within the next three months, or in their first year of living in boarding school hostels, tertiary education halls of residence, military barracks, Youth Justice residences, or prisons: or
 - 2) One dose for individuals who turn 13 years of age while living in boarding school hostels.
- C) Contractors will be entitled to claim payment from the Funder for the supply of Meningococcal A, C, Y and W-135 vaccine to patients eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Meningococcal A, C, Y and W-135 vaccine listed in the Pharmaceutical Schedule.
- D) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraphs A-B above.

Note: children under seven years of age require two doses 8 weeks apart, a booster dose three years after the primary series and then five yearly.

*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days. Ini 10 mgg of each meningococcal polysaccharide conjugated

Subsidy (Manufacturer's Pri	ice)	Fully Subsidised	Brand or Generic	
\$	Per	✓	Manufacturer	

MENINGOCOCCAL B MULTICOMPONENT VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- c) Any of the following:
 - A) Three doses for children up to 12 months of age (inclusive) for primary immunisation; or
 - B) Up to three doses (dependent on age at first dose) for a catch-up programme for children from 13 months to 59 months of age (inclusive) for primary immunisation, from 1 March 2023 to 31 August 2025; or
 - C) Both:
 - 1) Person is one year of age or over; and
 - 2) Any of the following:
 - i) up to two doses and a booster every five years for patients pre- and post-splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre- or post-solid organ transplant; or
 - ii) up to two doses for close contacts of meningococcal cases of any group; or
 - iii) up to two doses for person who has previously had meningococcal disease of any group; or
 - iv) up to two doses for bone marrow transplant patients; or
 - v) up to two doses for person pre- and post-immunosuppression*; or
 - D) Both:
 - 1) Person is aged between 13 and 25 years (inclusive); and
 - 2) Either:
 - Two doses for individuals who are entering within the next three months, or in their first year of living in boarding school hostels, tertiary education halls of residence, military barracks, Youth Justice residences or prisons; or
 - ii) Two doses for individuals who turn 13 years of age while living in boarding school hostels.
 - E) Contractors will be entitled to claim payment from the Funder for the supply of Meningococcal B multicomponent vaccine to patients eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Meningococcal B multicomponent vaccine listed in the Pharmaceutical Schedule.
 - F) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraphs A-D above.

*Immunosuppression due to corticosteroid or other immunosuppressive therapy must be for a period of greater than 28 days.

MENINGOCOCCAL C CONJUGATE VACCINE - [Xpharm]

Both:

- 1) The child is under 12 months of age; and
- 2) Any of the following:
 - 1) Up to three doses for patients pre- and post splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre or post solid organ transplant; or
 - 2) Two doses for close contacts of meningococcal cases of any group; or
 - 3) Two doses for child who has previously had meningococcal disease of any group; or
 - 4) A maximum of two doses for bone marrow transplant patients; or
 - 5) A maximum of two doses for child pre- and post-immunosuppression*.

Note: children under 12 months of age require two doses 8 weeks apart. Refer to the Immunisation Handbook for recommended booster schedules with meningococcal ACWY vaccine.

*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days.

✓ Synflorix

10

	Subsidy (Manufacturer's Price) \$	Subs Per	Fully idised	Brand or Generic Manufacturer
PNEUMOCOCCAL (PCV10) CONJUGATE VACCINE - [Xpharm	n]			
A primary course of three doses for previously unvaccing Note: please refer to the Immunisation Handbook for the application.	'	•		
Inj 1 mcg of pneumococcal polysaccharide serotypes 1, 5, 6l 7F, 9V, 14 and 23F; 3 mcg of pneumococcal	' '	outon up	program	

polysaccharide serotypes 4, 18C and 19F in 0.5 ml

-					
	Subsidy		Fully	Brand or	
	(Manufacturer's Price)	Su	bsidised	Generic	
	\$	Per	/	Manufacturer	

PNEUMOCOCCAL (PCV13) CONJUGATE VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- c) A) Any of the following:
 - 1) A course of three doses for previously unvaccinated children up to the age of 59 months inclusive; or
 - 2) Two doses are funded for high risk individuals (over the age of 12 months and under 18 years) who have previously received two doses of the primary course of PCV10; or
 - 3) Up to an additional four doses (as appropriate) are funded for the (re)immunisation of high risk children aged under 5 years with any of the following:
 - a) on immunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient immune response; or
 - b) primary immune deficiencies; or
 - c) HIV infection; or
 - d) renal failure, or nephrotic syndrome; or
 - e) who are immune-suppressed following organ transplantation (including haematopoietic stem cell transplant): or
 - f) cochlear implants or intracranial shunts; or
 - g) cerebrospinal fluid leaks; or
 - receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage
 of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily
 dosage of 20 mg or greater; or
 - i) chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or
 - i) pre term infants, born before 28 weeks gestation; or
 - k) cardiac disease, with cyanosis or failure; or
 - I) diabetes: or
 - m) Down syndrome; or

Ini 30.8 mcg of pneumococcal polysaccharide serotypes 1, 3, 4.

- n) who are pre-or post-splenectomy, or with functional asplenia; or
- 4) Up to an additional four doses (as appropriate) are funded for the (re-)immunisation of individuals 5 years and over with HIV, pre or post haematopoietic stem cell transplantation, or chemotherapy; pre- or post splenectomy; functional asplenia, pre- or post- solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, intracranial shunts, cerebrospinal fluid leaks or primary immunodeficiency; or
- 5) For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.
- B) Contractors will be entitled to claim payment from the Funder for the supply of Pneumococcal (PCV13) conjugate vaccine to people eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Pneumococcal (PCV13) conjugate vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

sub-set of the population described in paragraph A above.

Note: please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes

5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and 23F in 0.5ml		
syringe	10	✓ Prevenar 13
, -	1	✓ Prevenar 1:

	Subsidy	Fully	Brand or
	Manufacturer's Price)	Subsidised	Generic
	\$	Per 🗸	Manufacturer
PNEUMOCOCCAL (PPV23) POLYSACCHARIDE VACCINE - [Xp	harm]		
Either:			

- Up to three doses (as appropriate) for patients with HIV, for patients post haematopoietic stem cell transplant, or chemotherapy; pre- or post-splenectomy or with functional asplenia, pre- or post-solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, or primary immunodeficiency; or
- 2) All of the following:
 - a) Patient is a child under 18 years for (re-)immunisation; and
 - b) Treatment is for a maximum of two doses; and
 - c) Any of the following:
 - i) on immunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient immune response; or
 - ii) with primary immune deficiencies; or
 - iii) with HIV infection; or
 - iv) with renal failure, or nephrotic syndrome; or
 - v) who are immune-suppressed following organ transplantation (including haematopoietic stem cell transplant);
 or
 - vi) with cochlear implants or intracranial shunts; or
 - vii) with cerebrospinal fluid leaks; or
 - viii) receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage of 20 mg or greater; or
 - ix) with chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or
 - x) pre term infants, born before 28 weeks gestation; or
 - xi) with cardiac disease, with cyanosis or failure; or
 - xii) with diabetes: or
 - xiii) with Down syndrome; or

Ini 575 mcg in 0.5 ml prefilled syringe (25 mcg of each

xiv) who are pre-or post-splenectomy, or with functional asplenia.

23 pneumococcal serotype)	0.00	1	✓ Pneumovax 23
POLIOMYELITIS VACCINE - [Xpharm]			
Up to three doses for patients meeting either of the following:			
1) For partially vaccinated or previously unvaccinated individuals	s; or		
For revaccination following immunosuppression.			
Note: Please refer to the Immunisation Handbook for appropriate s	schedule for c	catch-up pr	rogrammes.
Inj 80D antigen units in 0.5 ml syringe	0.00	1	✓ IPOL

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	/	Manufacturer	

BOTAVIBUS OBAL VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- c) A) Maximum of two doses for people meeting the following:
 - 1) first dose to be administered in infants aged under 14 weeks of age; and
 - 2) no vaccination being administered to children aged 24 weeks or over.
 - B) Contractors will be entitled to claim payment from the Funder for the supply of Rotavirus oral vaccine to people eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Rotavirus oral vaccine listed in the Pharmaceutical Schedule.
 - C) Contractors may only claim for populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Oral susp live attenuated human rotavirus 1,000,000 CCID50 per dose, squeezable tube	0.00	10	✓ Rotarix
Oral susp live attenuated human rotavirus 1,000,000 CCID50 per dose, prefilled oral applicator	0.00	10	✓ Rotarix

Subsidy (Manufacturer's Price)	Sı	Fully ubsidised	Brand or Generic
\$	Per	✓	Manufacturer

VARICELLA VACCINE [CHICKENPOX VACCINE]

- a) Only on a prescription
- b) No patient co-payment payable
- c) A) Either:
 - 1) Maximum of one dose for primary vaccination for either:
 - a) Any infant born on or after 1 April 2016; or
 - For previously unvaccinated children turning 11 years old on or after 1 July 2017, who have not
 previously had a varicella infection (chickenpox), or
 - 2) Maximum of two doses for any of the following:
 - a) Any of the following for non-immune individuals:
 - i) with chronic liver disease who may in future be candidates for transplantation; or
 - ii) with deteriorating renal function before transplantation; or
 - iii) prior to solid organ transplant; or
 - iv) prior to any elective immunosuppression*; or
 - v) for post exposure prophylaxis who are immune competent inpatients; or
 - b) For individuals at least 2 years after bone marrow transplantation, on advice of their specialist; or
 - c) For individuals at least 6 months after completion of chemotherapy, on advice of their specialist; or
 - d) For HIV positive non immune to varicella with mild or moderate immunosuppression on advice of HIV specialist; or
 - e) For individuals with inborn errors of metabolism at risk of major metabolic decompensation, with no clinical history of varicella; or
 - f) For household contacts of paediatric patients who are immunocompromised, or undergoing a procedure leading to immune compromise where the household contact has no clinical history of varicella; or
 - g) For household contacts of adult patients who have no clinical history of varicella and who are severely immunocompromised, or undergoing a procedure leading to immune compromise where the household contact has no clinical history of varicella.
 - B) Contractors will be entitled to claim payment from the Funder for the supply of Varicella vaccine [Chickenpox vaccine] vaccine to people eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Varicella vaccine [Chickenpox vaccine] listed in the Pharmaceutical Schedule.
 - C) Contractors may only claim for populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraphs A above.

inimunosuppression due to steroid or other inimunosuppressive there	apy musi be i	oi a ileaille	ent penou or greater i	llai
28 days				
Inj 1350 PFU prefilled syringe0	.00	1 •	✓ Varivax	
, , , ,		10	✓ Varivax	

VARICELLA ZOSTER VACCINE [SHINGLES VACCINE]

- a) Only on a prescription
- b) No patient co-payment payable

c)

- A) Funded for patients meeting the following criteria:
 - 1) Two doses for all people aged 65 years
- B) Contractors will be entitled to claim payment from the Funder for the supply of Varicella zoster vaccine (Shingles vaccine) to patients eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Varicella zoster vaccine [Shingles vaccine] listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Inj 50 mcg per 0.5 ml vial plus vial	0.00	1	✓ Shingrix
		10	✓ Shinariy

Subsidy	Full	/ Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per 🗸	Manufacturer	

Diagnostic Agents

- Symbols -		Disorders	122	Amphotericin B	3
3TC	112	Agents Used in the Treatment of		Amsacrine	15
7 MED NSHA Silver/Copper		Poisonings	264	AmsaLyo	15
Short	79	Agrylin	158	Amsidine	15
- A -		Albendazole		Amzoate	
A-Scabies	74	Albey248	3-249	Anaesthetics	12
Abacavir sulphate	112	Albustix		Anafranil	
Abacavir sulphate with		Alchemy Oxaliplatin	154	Anagrelide hydrochloride	
lamivudine	112	Alchemy Oxybutynin	83	Analgesics	12
Abacavir/Lamivudine Viatris	112	Aldurazyme	28	Anastrozole	
Abilify Maintena	136	Alecensa	166	Anatrole	17
Abiraterone acetate	172	Alectinib	166	Androderm	
Acarbose	11	Alendronate sodium	117	Anoro Ellipta	25
Accarb	11	Alendronate sodium with		Antabuse	
Accuretic 10	47	colecalciferol	117	Antacids and Antiflatulents	
Accuretic 20	47	Alfacalcidol	33	Anthelmintics	9
Acetazolamide	261	Alfamino	288	Antiacne Preparations	
Acetec		Alfamino Junior		Antiallergy Preparations	
Acetic acid with hydroxyquinoline		Alginic acid	6	Antianaemics	
ricinoleic acid		Alglucosidase alfa		Antiandrogen Oral	
Acetylcysteine		Alkeran		Contraceptives	8 [.]
Aci-Jel		Alkeran S29		Antiarrhythmics	
Aciclovir		Allerfix		Antibacterials	
Infection	107	Allerpro Syneo 1		Antibacterials Topical	
Sensory		Allerpro Syneo 2		Anticholinergic Agents	
Acidex		Allersoothe		Anticholinesterases	
Acipimox		Allmercap		Antidepressants	
Acitretin		Allopurinol		Antidiarrhoeals	
Actemra		Alpha-Adrenoceptor Blockers		Antiepilepsy Drugs	
Actinomycin D		Alpha-Keri Lotion		Antifibrinolytics, Haemostatics and	
Actrapid		Alphamox		Local Sclerosants	
Actrapid Penfill		Alphamox 125		Antifibrotics	
Acupan		Alphamox 250		Antifungals	
Adalimumab (Amgevita)		Alprolix		Antifungals Topical	
Adalimumab (Humira - Alternative		Alu-Tab		Antihistamines	
brand)		Aluminium hydroxide		Antihypotensives	
Adapalene		Alvogen		Antimalarials	
Adcetris		Amantadine hydrochloride		Antimigraine Preparations	
ADR Cartridge 1.8		Ambrisentan		Antinausea and Vertigo Agents	
Adrenaline		Ambrisentan Viatris		Antipruritic Preparations	
Cardiovascular	56	Amgevita		Antipsychotics	
Respiratory		Amiloride hydrochloride		Antiretrovirals	
Advantan		Amiloride hydrochloride with		Antirheumatoid Agents	
Advate		furosemide	53	Antispasmodics and Other Agents	
Adynovate		Amiloride hydrochloride with		Altering Gut Motility	
Afinitor		hydrochlorothiazide	53	Antithrombotic Agents	
Aflibercept		Aminophylline		Antithymocyte globulin	
AFT-Pyrazinamide		Amiodarone hydrochloride		(equine)	18
After Hours Med Mgmt 15 min		Amisulpride		Antitrichomonal Agents	10
After Hours Med Mgmt 30 min		Amitriptyline		Antituberculotics and	
After Hours Med Mgmt 45 min		Amlodipine		Antileprotics	10!
Agents Affecting the		Amorolfine		Antiulcerants	
Renin-Angiotensin System	46	Amoxicillin		Antivirals	
Agents for Parkinsonism and Rela		Amoxicillin with clavulanic acid		Antivirals Eligibility Review	26
-				· ,	

Anxiolytics	139	Asthalin	252	Bendrofluazide	5
Anzatax	163	Atazanavir Mylan	112	Bendroflumethiazide	
Apidra		Atazanavir sulphate	112	[Bendrofluazide]	50
Apidra SoloStar	11	Atazanavir Viatris		Benralizumab	
APO-Atomoxetine	144	Atenolol	49	Benzathine benzylpenicillin	
APO-Atomoxetine S29		Atenolol AFT		Benzatropine mesylate	
Apo-Azithromycin		Atenolol AFT S29	49	Benzbromaron AL 100	
Apo-Bicalutamide		Atenolol Viatris		Benzbromarone	
APO-Candesartan HCTZ		Atezolizumab		Benztrop	
16/12.5	47	ATGAM		Benzydamine hydrochloride	
APO-Candesartan HCTZ		Ativan		Benzylpenicillin sodium [Penicillin	
32/12.5	47	Atnahs Olsalazine		G]	
Apo-Temozolomide		Atomoxetine		Beta Cream	
Apo-Terbinafine		Atorvastatin		Beta Ointment	
Apomorphine hydrochloride		Atropine sulphate		Beta Scalp	
Aprepitant		Cardiovascular	48	Beta-Adrenoceptor Agonists	
Apresoline		Sensory		Beta-Adrenoceptor Blockers	
Aptamil Feed Thickener		Atropt		Betadine	
		Atrovent		Betadine Skin Prep	
Aqueous cream				Betaferon	
Aratac Arava		Augmentin			
		Augmentin		Betahistine dihydrochloride	
Arginine		AutoCoff 20		Betaine	
Arginine2000		AutoSoft 30		Betamethasone dipropionate	
Aripiprazole		AutoSoft 90		Betamethasone dipropionate with	
Aripiprazole Sandoz		Avallon		calcipotriol	
Aristocort		Avelox		Betamethasone sodium phosphat	
Arrotex-Prazosin S29		Avonex		with betamethasone acetate	
Arrow - Clopid		Avonex Pen		Betamethasone valerate	
Arrow - Lattim		Azacitidine		Betamethasone valerate with sod	
Arrow-Amitriptyline		Azacitidine Dr Reddy's		fusidate [fusidic acid]	
Arrow-Bendrofluazide		Azamun		Betaxolol	
Arrow-Brimonidine		Azathioprine	176	Betnovate	
Arrow-Diazepam	139	Azilect	122	Betoptic	
Arrow-Doxorubicin	159	Azithromycin	96	Betoptic S	26
Arrow-Fluoxetine	129	Azopt	261	Bexsero	300
Arrow-Losartan &		AZT	112	Bezafibrate	54
Hydrochlorothiazide	47	- B -		Bezalip	
Arrow-Norfloxacin	115	B-D Micro-Fine	15	Bezalip Retard	54
Arrow-Ornidazole	105	B-D Ultra Fine	16	Bicalutamide	173
Arrow-Quinapril 10	47	B-D Ultra Fine II	16	Bicillin LA	98
Arrow-Quinapril 20	47	Bacillus Calmette-Guerin (BCG)		BiCNU	150
Arrow-Quinapril 5		vaccine	183	BiCNU S29	150
Arrow-Roxithromycin	97	Bacillus Calmette-Guerin		Bile and Liver Therapy	10
Arrow-Timolol		vaccine	293	Biltricide	9
Arrow-Topiramate		Baclofen	121	Bimatoprost	
Arrow-Tramadol		Bactroban	68	Bimatoprost Multichem	
Arsenic trioxide		Balance		Binarex	
Asacol		Barrier Creams and Emollients		Binocrit	
Asacol S29		BCG Vaccine		Biodone	
Ascend Aripiprazole		Beclazone 100		Biodone Extra Forte	
Ascend-Cefuroxime		Beclazone 250		Biodone Forte	
Ascorbic acid		Beclazone 50		Bisacodyl	
Aspen Adrenaline		Beclomethasone dipropionate		Bisacodyl Viatris	رح
Aspirin		Bedaquiline		Bisoprolol fumarate	
Blood	40	Bee venom allergy treatment		BK Lotion	ان
Nervous		Bendamustine hydrochloride		Bleomycin sulphate	
DELVOUS	144	Denualiusune nvurucinoliue	102	DICUMPCIA SUIDITALE	100

Blood Colony-stimulating	Calcitriol33	Cefazolin9
Factors43	Calcitriol-AFT33	Cefazolin-AFT99
Blood glucose diagnostic test	Calcium 500 mg Hexal34	Ceftriaxone99
meter 14	Calcium carbonate6, 34	Ceftriaxone-AFT99
Blood glucose diagnostic test	Calcium carbonate PAI6	Cefuroxime axetil99
strip15	Calcium Channel Blockers51	Celapram 129
Blood glucose test strips (visually	Calcium Disodium Versenate266	Celebrex110
impaired)15	Calcium folinate	Celecoxib110
Blood Ketone Diagnostic Test	Calcium Folinate Ebewe	Celecoxib Pfizer11
·	Calcium Folinate Sandoz	Celestone Chronodose
Strip 13 Boostrix 293	Calcium Folinate Sandoz S29 155	Cellcept170
Bortezomib	Calcium gluconate	Centrally-Acting Agents
Bosentan	Calcium Homeostasis84	Cephalexin ABM9
Bosentan Dr Reddy's59	Calcium polystyrene sulphonate44	Cetirizine hydrochloride24
Bplex32	Calcium Resonium44	Cetomacrogol72
Brentuximab Vedotin200	Calogen272	Cetomacrogol with glycerol72
Breo Ellipta251	Camber56	Cetomacrogol-AFT72
Brevinor 1/2880	Camino Pro Bettermilk286	Cetuximab20
Brevinor-1 28 Day80	Candesartan cilexetil47	Charcoal26
Bricanyl Turbuhaler252	Candesartan cilexetil with	Chemotherapeutic Agents152
Brilinta40	hydrochlorothiazide47	Chickenpox vaccine30
Brimonidine tartrate262	Candestar47	Chlorambucil153
Brimonidine tartrate with timolol	Canesten	Chloramphenicol25
maleate262	Capecitabine155	Chlorothiazide5
Brinzolamide261	Capecitabine Viatris155	Chlorpheniramine maleate24
Brown & Burk129	Capsaicin	Chlorpromazine hydrochloride138
Brufen SR116	Musculoskeletal117	Chlorsig25
BSF Ipca-Donepezil264	Nervous124	Chlortalidone [Chlorthalidone]53
BSF Max Health264	Captopril	Chlorthalidone5
BSF Methylphenidate ER -		
_ ,,	Carafate	Choice 280 7mod Nobe Silver/sepper
Teva	Carbaccord	Choice 380 7med Nsha Silver/copper
BSF Teriparatide - Teva264	Carbamazepine	Short
Buccastem134	Carbimazole88	Choice Load 37579
Budesonide	Carbomer	Choice TT380 Short79
Alimentary6	Carboplatin 153	Choice TT380 Standard79
Respiratory250, 257	Carboplatin Ebewe 153	Ciclosporin24
Budesonide Te Arai6	Carbosorb-X265	Cilazapril4
Budesonide with eformoterol251	Cardinol LA50	Cilicaine VK9
Bumetanide52	Cardizem CD51	Cinacalcet84
Buprenorphine Naloxone BNM 148	CareSens Dual14	Cinacalet Devatis84
Buprenorphine with naloxone148	CareSens N14-15	Cipflox100
Bupropion hydrochloride149	CareSens N POP14	Ciprofloxacin
Burel103	CareSens N Premier14	Infection100
Burinex52	CareSens PRO15	Sensory25
Burinex S29	Carmellose sodium with gelatin and	Ciprofloxacin - Torrent10
Buscopan8	pectin31	Ciprofloxacin Teva25
Buspirone hydrochloride139	Carmustine	Cisplatin
Buspirone Viatris139	Carnitor	Cisplatin Ebewe
Busulfan	Carvedilol	Citalopram hydrobromide129
- C -		
-	Carvedilol Sandoz	Citrulline 1000
Cabergoline 93	Casirivimab and imdevimab201	Cladribine
Caffeine citrate258	Catapres	Clarithromycin
Calamine70	CeeNU154	Alimentary
Calci-Tab 50034	Cefaclor monohydrate95	Infection90
Calcipotriol74	Cefalexin95	Clexane4
Calcitonin84	Cefalexin Sandoz95	Clexane Forte4

Clindamycin	100	Compound electrolytes with glue	cose	Daonil	1
Clinicians		[Dextrose]		Dapa-Tabs	5
Clinicians Renal Vit	33	Compound hydroxybenzoate		Dapsone	
Clobazam	130	Comtan	122	Daraprim	
Clobetasol propionate	70, 76	Concerta	147	Darunavir	110
Clobetasone butyrate		Condoms	78	Darunavir Viatris	113
Clofazimine		Condyline	76	Dasatinib	166
Clomazol		Contraceptives - Hormonal	79	Daunorubicin	159
Dermatological	69	Contraceptives - Non-hormonal.	77	Daunorubicin Zentiva	159
Genito-Urinary		Copaxone	140	David One Step Cassette Pregn	ancy
Clomifene citrate	93	Cordarone-X	48	Test	82
Clomipramine hydrochloride	128	Corticosteroids and Related Age		DBL Adrenaline	56
Clomipramine Teva		for Systemic Use	85	DBL Aminophylline	
Clonazepam	130, 139	Corticosteroids Topical		DBL Bleomycin Sulfate	158
Clonidine	52	Cosentyx	227	DBL Bortezomib	
Clonidine hydrochloride	52	Cosmegen	159	DBL Carboplatin	
Clonidine Teva		Coumadin	43	DBL Cisplatin	150
Clopidogrel	40	Country Life	29	DBL Dacarbazine	159
Clopine	135	Coversyl	47	DBL Desferrioxamine Mesylate	for Inj
Clopixol	136, 138	Creon 10000	23	BP	
Clotrimazole		Creon 25000	<mark>23</mark>	DBL Docetaxel	159
Dermatological	69	Creon Micro		DBL Ergometrine	
Genito-Urinary	81	Crotamiton	70	DBL Gemcitabine	156
Clozapine	135	Crystaderm	68	DBL Gentamicin	100
Clozaril	135	Curam	98	DBL Heparin Sodium	
Clustran	133	Curam Duo 500/125	98	DBL Leucovorin Calcium	
Co-trimoxazole	102	Cvite	32	DBL Methotrexate Onco-Vial	15
Coal tar	75	Cyclizine hydrochloride	134	DBL Pethidine Hydrochloride	128
Coal tar with allantoin, mentho	ol,	Cyclizine lactate		DBL Vincristine Sulfate	166
phenol and sulphur	75	Cyclogyl	262	Decozol	32
Coal tar with salicylic acid and	t	Cyclonex	153	Deferasirox	
sulphur	75	Cyclopentolate hydrochloride	262	Deferiprone	26
Cobal-B12		Cyclophosphamide	153	Deferoxamine Pfizer S29	26
Cobalin-H	32	Cyclorin	106	Denosumab	
Coco-Scalp	75	Cycloserine	106	Deolate	104
Codeine phosphate		Cyklokapron	40	Deoxycoformycin	
Extemporaneous	268	Cyproterone acetate	86	Depo-Medrol	8
Nervous	126	Cyproterone acetate with		Depo-Provera	
Coenzyme Q10	<mark>27</mark>	ethinyloestradiol	81	Depo-Testosterone	8
Colchicine	120	Cystadane	<mark>27</mark>	Deprim	102
Colecalciferol	33	Cytarabine		Dermol	,
Colestid	54	Cytotec	9	Desferrioxamine mesilate	
Colestipol hydrochloride	54	Cytoxan	153	Desmopressin	
Colestyramine	54	- D -		Desmopressin acetate	9
Colestyramine - Mylan	54	D-Penamine		Desmopressin-PH&T	9
Colgout	120	Dabigatran	42	Desuric	120
Colifoam	7	Dacarbazine		Detection of Substances in	
Colistin sulphomethate	100	Dacarbazine APP		Urine	8
Colistin-Link	100	Dactinomycin [Actinomycin D]		Dexamethasone	
Collodion flexible		Daivobet	74	Hormone	
Colloidal bismuth subcitrate	10	Daivonex	74	Sensory	
Colofac		Daktarin	69	Dexamethasone phosphate	
Coloxyl	24–25	Dalacin C	100	Dexamethasone with framycetin	
Combigan	262	Dantrium	121	gramicidin	
Compliance Packaging		Dantrium S29	121	Dexamethasone with neomycin	
Compound electrolytes		Dantrolene	121	sulphate and polymyxin B	

sulphate260	Domperidone	134	Elaprase	
Dexamfetamine sulfate144	Domperidone Viatris		Elecare	
Dexmethsone85	Donepezil hydrochloride	148	Elecare LCP	
Dextrochlorpheniramine	Dornase alfa	255	Electral	
maleate249	Dortimopt	261	Elelyso	
Dextrose	Dorzolamide with timolol		Elemental 028 Extra	27
DHC Continus126	Dostinex	93	Elexacaftor with tezacaftor, ivaca	
Diabetes10	Dosulepin [Dothiepin]		and ivacaftor	250
Diabetes Management13	hydrochloride		Elidel	
Diacomit	Dosulepin Mylan		Elocon	
Diagnostic Agents306	Dosulepin Viatris		Elocon Alcohol Free	
Diamide Relief6	Dothiepin		Eltrombopag	
Diamox261	Dovato		Eltroxin	
Diasip273	Doxazosin		EMB Fatol	
Diazepam130, 139	Doxazosin Clinect		Emend Tri-Pack	
Diazoxide10	Doxine		Emicizumab	
Dibenzyline46	Doxorubicin Ebewe		EMLA	
Diclofenac Sandoz116	Doxorubicin hydrochloride	159	Empagliflozin	1
Diclofenac sodium	Doxycycline	99	Empagliflozin with metformin	
Musculoskeletal116	DP Lotion	72	hydrochloride	
Sensory260	DP Lotn HC	70	Emtricitabine	11
Differin68	DP-Captopril		Emtricitabine with tenofovir	
Difflam31	Dr Reddy's Omeprazole		disoproxil	10
Diflucan102	Drofate	50	Emtriva	
Digestives Including Enzymes23	Drugs Affecting Bone		Emulsifying ointment	
Digoxin48	Metabolism	117	Emulsifying Ointment ADE	7
Dihydrocodeine tartrate126	Dual blood glucose and blood ke	tone	Enalapril maleate	
Dilantin	diagnostic test meter	14	Enbrel	
Dilantin Infatab131	Dulaglutide		Endocrine Therapy	
Dilantin Paediatric131	Dulcolax SP Drop		Endoxan	
Diltiazem CD Clinect51	Duocal Super Soluble Powder	271	Engerix-B	
Diltiazem hydrochloride51	Duolin		Enlafax XR	
Dimethicone71, 73	Duolin HFA	252	Enoxaparin sodium	4
Dimethyl fumarate140	Duolin Respules	252	Enstilar	
Dipentum8	DuoResp Spiromax	251	Ensure	
Diphtheria, tetanus and pertussis	Duride	56	Ensure Plus	28
vaccine293	Durvalumab	239	Ensure Plus HN	
Diphtheria, tetanus, pertussis and	- E -		Ensure Plus HN RTH	
polio vaccine294	e-chamber La Grande		Ensure Two Cal HN RTH	
Diphtheria, tetanus, pertussis, polio,	e-chamber Mask		Entacapone	
hepatitis B and haemophilus	e-chamber Turbo		Entecavir	
influenzae type B vaccine 294	E-Mycin	97	Entecavir (Rex)	
Diprosone70	e5 Pharma		Entresto 24/26	
Diprosone OV70	Ear Preparations		Entresto 49/51	
Dipyridamole40	Ear/Eye Preparations		Entresto 97/103	
Disopyramide phosphate48	Easiphen Liquid		Entyvio	
Disulfiram149	Econazole nitrate		Epilim	
Diuretics52	Efavirenz		Epilim Crushable	13
Docetaxel159	Efavirenz Milpharm	111	Epilim IV	13
Docetaxel Accord159	Efavirenz with emtricitabine and		Epilim S/F Liquid	13
Docetaxel Sandoz159	tenofovir disoproxil	112	Epilim Syrup	13
Docusate sodium24	Eformoterol fumarate dihydrate	250	Epipen	24
Docusate sodium with	Eftrenonacog alfa [Recombinant		Epipen Jr	24
sennosides25	factor IX]	37	Epirubicin Ebewe	15
Dolutegravir113	Efudix		Epirubicin hydrochloride	15
Dolutegravir with lamivudine113	Egopsoryl TA	75	Eplerenone	5

Epoetin alfa	36	Eylea	198	Flucon	260
Epoprostenol	63	Ezemibe Viatris	56	Fluconazole	102
Eptacog alfa [Recombinant fac	tor	Ezetimibe	56	Fludara Oral	156
VIIa]		Ezetimibe Sandoz	56	Fludarabine Ebewe	156
Erbitux	201	Ezetimibe with simvastatin	56	Fludarabine phosphate	
Ergometrine maleate	81	-F-		Fludarabine Sagent	156
Erlotinib	167	Factor eight inhibitor bypassing		Fludrocortisone acetate	85
Erythrocin IV		fraction	39	Fluids and Electrolytes	
Erythromycin (as lactobionate)	97	Famotidine	9	Flumetasone pivalate	
Erythromycin ethyl succinate		Famotidine Hovid	9	Fluocortolone caproate with	
Esbriet		Fasenra	199	fluocortolone pivalate and	
Escitalopram	129	Faslodex	173	cinchocaine	8
Escitalopram (Ethics)	129	Fatty Cream AFT	72	Fluorometholone	
Eskazole		Febuxostat	120	Fluorouracil	156
Essential Ethosuximide	130	Febuxostat (Teva)	120	Fluorouracil Accord	156
Essential Generics	49	FEIBA NF		Fluorouracil sodium	76
Essential Prednisolone		Felo 10 ER	51	Fluox	
Estraderm MX	87	Felo 5 ER	51	Fluoxetine hydrochloride	129
Estradiol Sandoz		Felodipine		Flupenthixol decanoate	
Estradiol TDP Mylan	87	Fentanyl		Flutamide	
Estradiol Viatris		Fentanyl Sandoz		Flutamin	
Estradot		Ferinject		Fluticasone	250
Estrofem	87	Ferodan		Fluticasone furoate with	
Etanercept	176	Ferriprox		umeclidinium and vilanterol	253
Ethambutol hydrochloride		Ferro-F-Tabs		Fluticasone furoate with	
Ethics Aspirin		Ferro-Liquid		vilanterol	251
Ethics Aspirin EC		Ferro-tab		Fluticasone propionate	
Ethics Lisinopril		Ferrograd		Fluticasone with salmeterol	
Ethinyloestradiol with		Ferrosig		Flynn	95
desogestrel	79	Ferrous fumarate	34	FML	
Ethinyloestradiol with		Ferrous fumarate with folic acid.	34	Foban	69
levonorgestrel	80	Ferrous sulfate	34	Folic acid	37
Ethinyloestradiol with		Fexofenadine hydrochloride	249	Folic Acid multichem	37
norethisterone	80	Fibro-vein		Folic Acid Viatris	
Ethosuximide	130	Filgrastim		Food Thickeners	
Etopophos		Finasteride		Foods And Supplements For	
Etoposide	159	Fingolimod		Inherited Metabolic Disease	284
Etoposide phosphate		Firazyr		Fortini	
Etravirine		Flagyl		Fortini Multi Fibre	275
Eumovate	70	Flagyl-S		Fortisip	281
Eurofolic	155	Flamazine		Fortisip Multi Fibre	
Evara		Flecainide acetate	49	Fosamax	
EVARA White Soft Paraffin	72	Flecainide BNM	49	Fosamax Plus	117
Everet	131	Flecainide Controlled Release		Framycetin sulphate	259
Everolimus	244	Teva	49	Frebini Energy	
Evista	118	Flecatab	49	Frebini Energy Fibre	
Evrysdi	143	Fleet Phosphate Enema	25	Frebini Original	
Evusheld	229	Flixonase Hayfever & Allergy		Frebini Original Fibre	
Exelon Patch 10	148	Flixotide		Fresubin 2kcal HP	282
Exelon Patch 5		Flixotide Accuhaler		Fresubin HP Energy	
Exemestane		Florinef		Fresubin HP Energy Fibre	
Exjade		Fluanxol	137	Fresubin Intensive	
Extemporaneously Compounde		Flucil		Fresubin Original	
Preparations and		Flucloxacillin		Fresubin Original Fibre	
Galenicals	268	Flucloxacillin-AFT		Frisium	
Eye Preparations		Flucloxin		Frumil	

Frusemide	52	Glycosade	287	Humira	191
Fucicort	71	Gold Knight	78	HumiraPen	191
Fucidin	101	Gold Knight XL	78	Humulin 30/70	11
Fucithalmic	259	Goserelin	93	Humulin NPH	1 1
Fulvestrant	173	Gynaecological Anti-infectives	81	Humulin R	10
Fungilin	32	-H-		Hyaluronic acid	263
Furosemid-Ratiopharm	52	Habitrol	150	Hydralazine	56
Furosemide [Frusemide]		Haemophilus influenzae type B		Hydralazine hydrochloride	
Furosemide-Baxter	52	vaccine	. 295	Hydralyte - Lemonade	
fusidic acid		Haldol	137	Hydrocortisone	
Dermatological	69, 71	Haldol Concentrate	137	Dermatological	70
Infection	101	Haldol Decanoas	137	Hormone	
Sensory	259	Haloperidol	. 135	Hydrocortisone acetate	
. G -		Haloperidol decanoate		Hydrocortisone acetate with	
GA Explore 5	287	Harvoni		pramoxine hydrochloride	7
Gabapentin		Havrix	.295	Hydrocortisone and paraffin liquid	
Gacet	126	Havrix Junior	.295	and lanolin	
Galsulfase	27	Haylor syrup	.249	Hydrocortisone butyrate	
Galvumet		HCU Explore 5		Hydrocortisone with cinchocaine	
Galvus		HCU Express 15		Hydrocortisone with miconazole	
Gardasil 9		healthE Aqueous Cream SLS		Hydrocortisone with natamycin an	
Gastrodenol		Free	72	neomycin	
Gaviscon Extra Strength		healthE Calamine Aqueous		Hydrogen peroxide	
Gaviscon Infant		healthE Dimethicone 10%		Hydroxocobalamin	
Gazyva		healthE Dimethicone 4% Lotion	73	Hydroxocobalamin Panpharma	
Gefitinib		healthE Dimethicone 5%		hydroxycarbamide	
GEM Aqueous Cream		healthE Glycerol BP		Hydroxychloroquine	
Gemcitabine Ebewe		healthE Urea Cream		Hydroxyurea	
Gemcitabine hydrochloride		Healtheries Simple Baking Mix	.283	[hydroxycarbamide]	159
Gemtuzumab ozogamicin		Hemastix		Hygroton	
Gentamicin sulphate	100	Hemlibra	38	Hylo-Fresh	
Gilenya		Heparin sodium		Hymenoptera24	18-249
Ginet		Heparin Sodium Panpharma		Hyoscine butylbromide	
Glatiramer acetate		Heparinised saline		Hyoscine hydrobromide	
Glecaprevir with pibrentasvir		Heparon Junior		Hyperuricaemia and Antigout	
Glibenclamide		Hepatitis A vaccine		Hypromellose	
Gliclazide		Hepatitis B recombinant		Hypromellose with dextran	
Glipizide		vaccine	. 296	-1-	
Glizide		Herzuma		Ibiamox	98
Glucagen Hypokit		Hiberix		Ibrance	
Glucagon hydrochloride		Hiprex		Ibrutinib	
Glucerna Select		Histaclear		Ibuprofen	
Glucose [Dextrose]		Histafen		Icatibant	
Gluten Free Foods		Holoxan	. 153	Idarubicin hydrochloride	160
Glycerin with sodium saccharing		Horleys Bread Mix		Idursulfase	
Glycerin with sucrose		Horleys Flour		Ifosfamide	
Glycerol		Hormone Replacement Therapy -		llevro	261
Alimentary	25	Systemic	87	Iloprost	
Extemporaneous		HPV		Imatinib mesilate	
Glyceryl trinitrate		Humalog		Imatinib-Rex	
Alimentary	8	Humalog Mix 25	11	Imbruvica	
Cardiovascular		Humalog Mix 50		Imfinzi	
Glycopyrronium		Human papillomavirus (6, 11, 16, 1		Imipramine Crescent	
Glycopyrronium bromide		31, 33, 45, 52 and 58) vaccine	•	Imipramine hydrochloride	
Glycopyrronium with		[HPV]	. 296	Imiquimod	
indacaterol	253	Humatin		Immune Modulators	

Immunisation Administration	264	Invega Trinza	138	Ketocal 3:1	292
Immunisation		Ipca-Allopurinol	120	KetoCal 4:1	292
Co-administration	264	Ipca-Bisoprolol	50	Ketoconazole	
Immunosuppressants	176	Ipca-Ciprofloxacin		Dermatological	76
Incruse Ellipta		Ipca-Donepezil	148	Infection	
Indacaterol	250	Ipca-Escitalopram	129	Ketogenic Diet	292
Indapamide	.53	IPCA-Frusemide	52	Ketoprofen	116
Infanrix IPV		IPCA-Metoprolol		KetoSens	
Infanrix-hexa	294	IPCA-Propranolol	50	Keytruda	24
Infant Formulae	288	IPOL		Kindergen	
Infatrini	292	Ipratropium bromide	252, 257	Klacid	
Infliximab		Iressa	168	Alimentary	
Influenza vaccine	297	Irinotecan Actavis 100	156	Infection	96
Influvac Tetra		Irinotecan hydrochloride	156	Kliogest	8
(2024 formulation)	297	Irinotecan-Rex		Kliovance	8
Inhaled Corticosteroids		Iron (as ferric carboxymaltos	e)34	Kogenate FS	39
Inhaled Long-acting		Iron polymaltose		Konakion MM	
Beta-adrenoceptor Agonists	250	Isentress		Konsyl-D	
Inresa		Isentress HD		Kuvan	
Inspra		Ismo 20		-L-	
Instillagel Lido		Ismo 40 Retard		Labetalol	50
Insulin aspart		Isoleucine50	287	Lacosamide	
Insulin aspart with insulin aspart		Isoniazid	106	Lactulose	2
protamine	.10	Isoniazid with rifampicin		Laevolac	
Insulin glargine		Isoptin		Lagevrio	110
Insulin glulisine		Isoptin Retard		Lamictal	
Insulin isophane		Isoptin SR		Lamivudine	
Insulin isophane with insulin		Isopto Carpine		Lamivudine Viatris	
neutral	.11	Isosorbide mononitrate		Lamivudine/Zidovudine Viatris	112
Insulin lispro		Isosource Standard	280	Lamotrigine	13 ⁻
Insulin lispro with insulin lispro		Isotretinoin	68	Lamprene	
protamine	.11	Ispaghula (psyllium) husk		Lanoxin	
Insulin neutral		Itch-Soothe		Lanoxin Paediatric Elixir	
Insulin pen needles		Itraconazole		Lanoxin PG	
Insulin pump		Itrazole		Lanoxin S29	
Insulin pump cartridge		Ivacaftor		Lansoprazole	
Insulin pump infusion set (steel		Ivermectin		Lantus	
cannula)	21	- J -		Lantus SoloStar	
Insulin pump infusion set (steel		Jadelle	80	Lanvis	
cannula, straight insertion)	21	Jakavi		Lanzol Relief	
Insulin pump infusion set (teflon		Jardiamet		Largactil	
cannula)	22	Jardiance		Laronidase	
Insulin pump infusion set (teflon		Jaydess		Lasix	
cannula, angle insertion with		Jevity HiCal RTH		Latanoprost	
insertion device)	23	Jevity Plus RTH		Latanoprost with timolol	
Insulin pump infusion set (teflon		Jevity RTH	280	Lax-Suppositories	
cannula, straight insertion with		Jinarc		Lax-suppositories Glycerol	
insertion device)	23	Juno Pemetrexed		Laxatives	
Insulin pump reservoir		- K -		Laxsol	
Insulin syringes, disposable with		Kadcyla	235	Ledipasvir with sofosbuvir	
attached needle	16	Kalydeco		Leflunomide	
Intelence		Kemadrin		Lenalidomide	
Interferon beta-1-alpha		Kenacomb		Letrole	
Interferon beta-1-beta		Kenacort-A 10		Letrozole	
Intra-uterine device		Kenacort-A 40		Leucine100	
Invega Sustenna		Kenalog in Orabase		Leukeran FC	15
	. • .				

Leukotriene Receptor		Loratadine	249	Meningococcal B multicomponent	t
Antagonists	255	Lorazepam	139	vaccine	300
Leuprorelin	93	Lorstat	54	Meningococcal C conjugate	
Leustatin	155	Losartan Actavis	47	vaccine	300
Levetiracetam	131	Losartan potassium	47	MenQuadfi	299
Levetiracetam-AFT	131	Losartan potassium with		Menthol	70
Levocabastine	260	hydrochlorothiazide	47	Mepolizumab	210
Levocarnitine	28	Lovir	107	Mercaptopurine	
Levodopa with benserazide.	122	Loxamine	129	Mercilon 28	
Levodopa with carbidopa	122	Lucrin Depot 1-month	93	Mesalazine	
Levomepromazine		Lucrin Depot 3-month	93	Mesna	16 ⁻
Levomepromazine		Lynparza	162	Mestinon	116
hydrochloride	136	Lyrica	131	Metabolic Disorder Agents	26
Levonorgestrel		- M -		Metformin hydrochloride	12
Genito-Urinary	80–81	m-Eslon	127	Metformin Viatris	
Hormone	88	Mabthera	214	Methadone BNM	126
Levonorgestrel BNM	81	Macrobid	115	Methadone hydrochloride	
Levothyroxine		Macrogol 3350 with potassium		Extemporaneous	268
Lidocaine [Lignocaine]		chloride, sodium bicarbonat	e and	Nervous	
Lidocaine [Lignocaine]		sodium chloride	25	Methenamine (hexamine)	
hydrochloride	123	Madopar 125		hippurate	118
Lidocaine [Lignocaine] with		Madopar 250		Methopt	
prilocaine	124	Madopar 62.5		Methotrexate	
Lidocaine-Baxter		Madopar HBS	122	Methotrexate DBL Onco-Vial	157
Life Extension	30	Madopar Rapid		Methotrexate DBL S29	157
Lignocaine		Magnesium hydroxide		Methotrexate Ebewe	157
Linezolid		Magnesium sulphate		Methotrexate Sandoz	157
Lioresal Intrathecal		Mantoux		Methyl hydroxybenzoate	
Lipid-Modifying Agents		MAR-Midodrine		Methylcellulose	
Liquigen		Marevan		Methylcellulose with glycerin and	
Liraglutide		Marine Blue Lotion SPF 50+		sodium saccharin	268
Lisinopril		Martindale Pharma	264	Methylcellulose with glycerin and	
Litak		Mask for spacer device		sucrose	268
Lithium carbonate		Maviret		Methyldopa	
Livostin		Maxidex		Methyldopa Mylan	
LMX4		Maxitrol		Methyldopa Mylan S29	
Lo-Oralcon 20 ED		MCT oil (Nutricia)		Methyldopa Viatris	52
Locacorten-Viaform ED's		Measles, mumps and rubella		Methylnaltrexone bromide	
Local preparations for Anal a		vaccine	299	Methylphenidate ER - Teva	
Rectal Disorders		Mebendazole		Methylphenidate hydrochloride	
Locasol		Mebeverine hydrochloride		Methylphenidate hydrochloride	
Locoid		Med Mgmt 15 min		extended-release	147
Locoid Crelo		Med Mgmt 30 min		Methylprednisolone	
Locoid Lipocream		Med Mgmt 45 min		Methylprednisolone (as sodium	
Locorten-Vioform		Medicine Delivery		succinate)	85
Lodoxamide		Medrol		Methylprednisolone aceponate	
Logem		Medroxyprogesterone acetate		Methylprednisolone acetate	
Lomide		Genito-Urinary	80	Methylxanthines	
Lomustine		Hormone		Metoclopramide Actavis 10	
Loniten		Mefenamic acid		Metoclopramide hydrochloride	
Loperamide hydrochloride		Megval		Metolazone	
Lopinavir with ritonavir		Melatonin		Metopirone	
Lopinavir/Ritonavir Mylan		Melpha		Metoprolol IV Mylan	
Loprofin		Melphalan		Metoprolol IV Viatris	
Loprofin Mix		Meningococcal (groups A, C, Y		Metoprolol succinate	
Lorafix		W-135) conjugate vaccine		Metoprolol tartrate	

Metrogyl	105	Minims Pilocarpine	262	Mylan (24 hr release)	<u>5</u>
Metronidazole		Minims Prednisolone		Mylan Atenolol	
Metyrapone		Minipress	46	Mylan Clomiphen	
Mexiletine hydrochloride	49	Minirin	93	Mylan Italy (24 hr release)	
Miacalcic		Minirin Melt		Myleran	15
Micolette		Mino-tabs		Myloc CR	
Micolette-S29	25	Minocycline hydrochloride	99	Mylotarg	
Miconazole		Minomycin		Myometrial and Vaginal Hormon	
Miconazole nitrate		Minor Skin Infections		Preparations	
Dermatological	69	Minoxidil		Myozyme	
Genito-Urinary		Minoxidil Roma		- N -	
Micreme		Mirena		Nadolol	5
Micreme H		Miro-Amoxicillin		Nadolol BNM	
Microgynon 30		Mirtazapine		Naglazyme	
Microlut		Misoprostol		Naloxone hydrochloride	
Midazolam		Mitomycin C		Naltraccord	
Midazolam-Baxter		Mitozantrone		Naltrexone AOP	
Midodrine		Mitozantrone Ebewe		Naltrexone hydrochloride	
Mifegyne		Mixtard 30		Naphazoline hydrochloride	
Mifonriotono	00				
Mile barre		MMA/PA Explore 5		Naphcon Forte	
Milpharm		MMA/PA Express 15		Naprosyn SR 1000	
Minerals		Moclobemide		Naprosyn SR 750	11
Mini-Wright AFS Low Range		Modafinil		Naproxen	11
Mini-Wright Standard		Modavigil		Narcaricin mite	
Minidiab	11	Moduretic		Nasal Preparations	
MiniMed 3.0 Reservoir		Molaxole		Natalizumab	
MMT-332A		Molnupiravir		Natulan	
MiniMed 770G		Moments		Nausafix	
MiniMed Mio MMT-921A		Mometasone furoate		Nausafix - S29	
MiniMed Mio MMT-923A	22	Monogen	273	Nausicalm	13
MiniMed Mio MMT-925A	22	Montelukast	255	Navelbine	
MiniMed Mio MMT-941A	22	Montelukast Viatris	255	Navelbine S29	16
MiniMed Mio MMT-943A	22	Moroctocog alfa [Recombinant	t factor	Nefopam hydrochloride	12
MiniMed Mio MMT-945A	22	VIII]	39	Neisvac-C	30
MiniMed Mio MMT-965A	22	Morphine hydrochloride		Neo-Cytamen S29	3
MiniMed Mio MMT-975A	22	Morphine sulphate		Neo-Mercazole	8
MiniMed Quick-Set MMT-386A	23	Motetis		Neocate Gold	
MiniMed Quick-Set MMT-387A		Mouth and Throat		Neocate Junior Unflavoured	
MiniMed Quick-Set MMT-396A		Movapo		Neocate Junior Vanilla	
MiniMed Quick-Set MMT-397A		Moxifloxacin		Neocate SYNEO	
MiniMed Quick-Set MMT-398A		MSUD Explore 5		Neoral	
MiniMed Quick-Set MMT-399A		MSUD Express 15		Neostigmine metilsulfate	
MiniMed Silhouette MMT-368A		MSUD Maxamum		Nepafenac	
MiniMed Silhouette MMT-377A		Mucolytics		Nepro HP (strawberry)	
MiniMed Silhouette MMT-378A		Mucosoothe		Nepro HP (vanilla)	
MiniMed Silhouette MMT-381A		Multiple Sclerosis Treatments		Nepro HP RTH	
MiniMed Silhouette MMT-382A		Multivitamin renal		Neulactil	
		Multivitamins			
MiniMed Silhouette MMT-383A MiniMed Silhouette MMT-384A				Neuraxpharm NeuroTabs	
		Mupirocin			
MiniMed Sure-T MMT-864A		Muscle Relaxants		Nevirapine	
MiniMed Sure-T MMT-866A		Mvite		Nevirapine Alphapharm	
MiniMed Sure-T MMT-874A		Myambutol		Nevirapine Viatris	
MiniMed Sure-T MMT-876A		Mycobutin		Nicorandil	
MiniMed Sure-T MMT-884A		MycoNail	69	Nicotine	
MiniMed Sure-T MMT-886A		Mycophenolate mofetil		Nifedipine	5
Minims Cyclopentolate	262	Mydriacyl	262	Nifedipine Viatris	5

Nifuran	115	Nutren Diabetes	273	Omeprazole actavis 20	9
Nilotinib	168	Nutrient Modules		Omeprazole actavis 40	
Nilstat		Nutrini Energy Multi Fibre	275	Omnitrope	
Alimentary	32	Nutrini Energy RTH	275	Omnitrope S29	89
Genito-Urinary		Nutrini Low Energy Multi Fibre	277	Onbrez Breezhaler	
Infection	103	Nutrini Peptisorb	290	Oncaspar LYO	163
Nintedanib	254	Nutrini Peptisorb Energy	290	OncoTICE	183
Nipent	164	Nutrini RTH	275	Ondansetron	134
Niraparib	161	Nutrison 800 Complete Multi		One-Alpha	33
Nirmatrelvir with ritonavir		Fibre	280	One-Alpha S29	33
Nitrates	<mark>56</mark>	Nutrison Advanced Diason	273	Opdivo	240
Nitroderm TTS	<mark>56</mark>	Nutrison Advanced Peptisorb	277	Ora-Blend	268
Nitrofurantoin	115	Nutrison Concentrated	282	Ora-Blend SF	268
Nitrolingual Pump Spray	56	Nutrison Energy	280	Ora-Plus	268
Nivestim	43	Nutrison Energy Multi Fibre	280	Ora-Sweet	268
Nivolumab	240	Nutrison Multi Fibre	280	Ora-Sweet SF	268
Nodia	6	Nutrison RTH		Orabase	31
Noflam 250		Nyefax Retard	51	Oral and Enteral Feeds	273
Noflam 500		Nystatin		Oralcon 30 ED	80
Non-Steroidal Anti-Inflammate		Alimentary	32	Oramorph	
Drugs		Genito-Urinary		Oramorph CDC S29	
Nonacog gamma, [Recombin		Infection		Oratane	68
Factor IX]		NZB Low Gluten Bread Mix		Ordine	
Norethisterone		- 0 -		Orgran	
Genito-Urinary	81	Obinutuzumab	211	Ornidazole	
Hormone		Obstetric Preparations		Orphenadrine citrate	
Norflex		Ocicure		Ortho-tolidine	
Norfloxacin		Ocrelizumab		Oruvail SR	
Noriday 28		Ocrevus		Osmolite RTH	
Norimin		Octocog alfa [Recombinant factor		Other Endocrine Agents	
Norimin-1 28 Day		VIII] (Advate)		Other Oestrogen Preparations .	
Normison		Octocog alfa [Recombinant factor		Other Progestogen	
Norpress		VIII] (Kogenate FS)		Preparations	88
Nortriptyline hydrochloride		Octreotide		Other Skin Preparations	
Norvir		Octreotide Depot Teva		Otodex	
Noumed Dexamfetamine		Octreotide GH		Ovestin	
Noumed Paracetamol		Octreotide long-acting		Genito-Urinary	81
Noumed Pethidine		Oestradiol		Hormone	88
Noumed Phenobarbitone		Oestradiol valerate		Oxaliplatin	
NovaSource Renal		Oestradiol with norethisterone		Oxaliplatin Accord	154
Novatretin		Oestriol		Oxaliplatin Actavis 100	
Novitium Sugar Free		Genito-Urinary	81	Oxaliplatin Ebewe	
NovoMix 30 FlexPen		Hormone		Oxis Turbuhaler	
NovoRapid		Oestrogens		Oxpentifylline	57
NovoRapid FlexPen		Ofev		Oxybutynin	83
NovoRapid Penfill		Oil in water emulsion		Oxycodone hydrochloride	
NovoSeven RT	38	Olanzapine		Oxycodone Lucis S29	
Nozinan		Olaparib		Oxycodone Sandoz	
Nozinan (Swiss)		Olbetam		Oxycodone Sandoz S29	127
Nozinan S29		Olbetam S29		OxyContin	197
Nucala		Olopatadine		OxyNorm	127
Nuelin		Olopatadine Teva		Oxytocin	
Nuelin-SR		Olsalazine		Oxytocin BNM	02
Nupentin		Omalizumab		Oxytocin GH	
Nusinersen		Omeprazole		Oxytocin Gri	
Nutilis		Omeprazole actavis 10		Oxytocin with ergometrine	02
1 1 G U I I G U I G U I I G U I I G U I G U I I G U I I G U		omopiazoio aolavio 10	• • • • • • • • • • • • • • • • • • • •	Saytooni waa orgonioanio	

maleate	82	Perindopril	47	PKU Glytactin RTD 15	28
Ozurdex	259	Periset		PKU Glytactin RTD 15 Lite	28
- P -		Periset ODT	134	PKU GMPro Ultra Lemonade	
Pacifen	121	Perjeta	214	PKU Lophlex LQ 10	
Pacimol		Permethrin		PKU Lophlex LQ 20	28
Paclitaxel		Perrigo		PKU Lophlex Powder	
Paclitaxel Actavis		Pertuzumab	214	PKU Lophlex Sensation 20	
Paclitaxel Ebewe		Peteha		PKU Restore Powder	
Paediatric Seravit		Pethidine hydrochloride		PKU sphere20 Banana	
Palbociclib		Pevaryl		PKU sphere20 Chocolate	
Paliperidone		Pexsig		PKU sphere20 Lemon	
Paliperidone palmitate		Pfizer Exemestane		PKU sphere20 Red Berry	
Pamidronate disodium		Pfizer S29		PKU sphere20 Vanilla	
Pamisol		Pharmacy Services		PKU Start	
Pamol		Pheburane		Plaquenil	
Pancreatic enzyme		Phenasen		Plendil ER	
Pantoprazole		Phenobarbitone		PMS-Salbutamol	
Panzop Relief		Phenobarbitone sodium		Pneumococcal (PCV10) conjuga	
Papaverine hydrochloride		Extemporaneous	268	vaccine	
Para-amino salicylic acid		Nervous		Pneumococcal (PCV13) conjuga	
Paracetamol		Phenoxybenzamine		vaccine	
Paracetamol (Ethics)		hydrochloride	46	Pneumococcal (PPV23)	
Paracetamol + Codeine	120	Phenoxymethylpenicillin (Per		polysaccharide vaccine	30
(Relieve)	127	V)		Pneumovax 23	
Paracetamol with codeine		Phenylalanine50		Podophyllotoxin	
Paraffin		Phenytoin sodium		Polaramine	
Paraffin liquid with wool fat		Phillips Milk of Magnesia		Poliomyelitis vaccine	30
Parasiticidal Preparations		Phlexy 10		Poloxamer	
Parnate		Phosphate Phebra		Poly-Gel	
Paromomycin		Phosphorus			
Paroxetine				Poly-Tears Poly-Visc	20
Paser		Phytomenadione		Polycal	
Paxam		Pilocarpine hydrochloride			21
Paxlovid		Pilocarpine nitrate		Polyethylene glycol 400 and	06
		Pimecrolimus		propylene glycol Ponstan	
Pazopanib				Posaconazole	
Peak flow meter		Pine tar with trolamine laurils			
Pediasure		and fluorescein		Posaconazole Juno	
Pediasure Plus		Pinetarsol		Potassium chloride	
Pediasure RTH		Pioglitazone		Potassium citrate	
Pegaspargase		Pirfenidone		Potassium iodate	
Pegasys	113	Pizotifen		Povidone iodine	
Pegfilgrastim		PKU Anamix Infant		Pradaxa	
Pegylated interferon alfa-2a		PKU Anamix Junior		Pramipexole hydrochloride	
Pembrolizumab		PKU Anamix Junior Chocolat		Pravastatin	
Pemetrexed		PKU Anamix Junior LQ		Praziquantel	9
Penicillamine		PKU Anamix Junior Orange .		Prazosin	
Penicillin G		PKU Anamix Junior Vanilla		Prazosin Mylan	
PenMix 30		PKU Build 10		Pred Forte	
PenMix 50		PKU Build 20 Chocolate	286	Prednisolone	
Pentasa		PKU Build 20 Raspberry	000	Prednisolone acetate	
Pentostatin [Deoxycoformycin]		Lemonade		Prednisolone sodium	
Pentoxifylline [Oxpentifylline]	57	PKU Build 20 Smooth		Prednisolone sodium	
Peptamen Junior		PKU Build 20 Vanilla		phosphate	26
Pepti-Junior		PKU Explore 10		Prednisolone-AFT	
Perhexiline maleate		PKU Explore 5		Prednisone	
Pericyazine	136	PKU Express 20	285	Prednisone Clinect	8

Pregabalin		Quinapril with		Rituximab (Mabthera)	
Pregabalin Pfizer		hydrochlorothiazide		Rituximab (Riximyo)	216
Pregnancy Tests - hCG Urine		Qvar	250	Rivaroxaban	
Premarin		- R -		Rivastigmine	
Prevenar 13		RA-Morph	126	Rivastigmine Patch BNM 10	148
Priadel		Ralicrom	8	Rivastigmine Patch BNM 5	
Primaquine	105	Raloxifene hydrochloride	118	Rivotril	130
Primidone	131	Raltegravir potassium		Riximyo	216
Primidone Clinect		Ramipex	122	RIXUBIS	39
Primolut N	88	Ramipril		Rizamelt	
Priorix	299	Ranbaxy-Cefaclor	95	Rizatriptan	133
Probenecid	121	Rapamune	244	Robinul	
Probenecid-AFT		Rasagiline	122	Ronapreve	201
Procarbazine hydrochloride	164	Reandron 1000		Ropin	122
Prochlorperazine	134	Recombinant factor IX	37, 39	Ropinirole hydrochloride	122
Prochlorperazine Brown &		Recombinant factor VIIa	38	Rosuvastatin	
Burk	134	Recombinant factor VIII	39	Rosuvastatin Viatris	55
Proctofoam	<mark>7</mark>	Rectogesic	8	Rotarix	304
Proctosedyl	8	Redipred	86	Rotavirus oral vaccine	304
Procyclidine hydrochloride	122	Relieve	116	Roxane-Propranolol	
Progesterone		Relistor		Roxithromycin	
Proglicem		Remicade	202	Rubifen	146
Proglycem		Renilon 7.5	<mark>276</mark>	Rubifen SR	
Progynova		Resonium-A	45	Rugby Capsaicin Topical Cream	1
Prolia	117	Resource Beneprotein	<mark>272</mark>	Musculoskeletal	
Promethazine hydrochloride	249	Respigen		Nervous	
Propafenone hydrochloride		Respiratory Devices		Rurioctocog alfa pegol [Recomb	inant
Propranolol		Respiratory Stimulants		factor VIII]	
Propylene glycol		Retinol palmitate		Ruxolitinib	
Propylthiouracil		ReTrieve		Rythmodan	
Prostacur		Retrovir	112	Rythmodan - Cheplafarm	
Protaphane	11	Revlimid	160	Rytmonorm	
Protaphane Penfill	11	Revolade	37	· - S -	
Protifar		Riboflavin		Sabril	132
Protionamide	106	Ribomustin		Sacubitril with valsartan	
Provera	87	Ricit		Sagent	
Provera HD		Rifabutin		SalAir	
Psoriasis and Eczema		Rifadin		Salazopyrin	
Preparations	74	Rifadin Sanofi		Salazopyrin EN	
PTU		Rifampicin		Salbutamol	
Pulmicort Turbuhaler		Rifaximin		Salbutamol Cipla	
Pulmozyme		Rifinah		Salbutamol with ipratropium	
Puri-nethol		Rilutek	122	bromide	252
Puritan's Pride Vitamin		Riluzole		Salicylic acid	
B-2 100 mg	29	RINVOQ		Salmeterol	
Pyrazinamide		Riodine		Sandomigran	
Pyridostigmine bromide		Risdiplam		Sandostatin LAR	
Pyridoxine hydrochloride		Risedronate Sandoz		Sanofi Primaquine	
Pyridoxine multichem		Risedronate sodium		Sapropterin dihydrochloride	20
Pyrimethamine		Risperdal Consta		Scalp Preparations	
Pytazen SR		Risperidone		Scopoderm TTS	
- Q -		Risperidone (Teva)	136	Scopolamine - Mylan	
Quantalan sugar free	54	Risperon		Scopolamine - Mylan S29	
Quetapel		Ritalin		Sebizole	
Quetiapine		Ritalin LA		Secukinumab	
Quinapril		Ritonavir		Sedatives and Hypnotics	
~~				Dodantoo ana myphonoo	1-72

Seebri Breezhaler	253	Sodium polystyrene sulphonate	45	Symbicort Turbuhaler 400/12	25
Senna	25	Sodium tetradecyl sulphate	39	Symmetrel	
Senokot		Sodium valproate	132	Sympathomimetics	
SensoCard	15	Sofradex	259	Synacthen	8
Serc	133	Soframycin	259	Synacthen Depot	
Serenace	135	Solgar		Synacthene Retard	
Seretide	251	Solifenacin succinate		Synflorix	
Seretide Accuhaler		Solifenacin Viatris		Synthroid	
Serevent	250	Solu-Cortef	85	Syntometrine	
Serevent Accuhaler		Solu-Medrol		Syrup (pharmaceutical grade)	
Sertraline	129	Solu-Medrol-Act-O-Vial	85	Systane Unit Dose	
Setrona	129	Somatropin (Omnitrope)	89	· -Т-	
Sevredol	127	Sotalol	51	Tacrolimus	
Sex Hormones Non		Spacer device	258	Dermatological	7
Contraceptive	86	Span-K	44	Oncology	
Shingles vaccine		Spazmol		Tacrolimus Sandoz	246
Shingrix		Spinal Muscular Atrophy		Taliglucerase alfa	30
SII-Onco-BCG		Spinraza		Tambocor	49
Sildenafil		Spiolto Respimat		Tamoxifen citrate	
Siltuximab		Spiractin		Tamoxifen Sandoz	17
Simvastatin		Spiriva		Tamsulosin hydrochloride	
Simvastatin Mylan		Spiriva Respimat		Tamsulosin-Rex	8
Simvastatin Viatris		Spironolactone		Tandem Cartridge	2
Sinemet		Sporanox		Tandem t:slim X2 with Basal-IQ	
Sinemet CR		Sprycel		Tap water	
Sirolimus		Stelara		Taro	
Sirturo		Stemetil		Taro-Testosterone	
Siterone		Steril-Gene		Tasigna	168
Slow-Lopresor		SteroClear		Tasmar	
Smith BioMed Rapid Pregnancy		Stesolid		Taurine	
Test	82	Stimulants/ADHD Treatments	144	Tecentriq	
Sodibic		Stiripentol		Tecfidera	
Sodium acid phosphate	25	Stocrin		Tegretol	
Sodium alginate		Stomahesive		Tegretol CR	
Sodium benzoate		Strides Shasun		Telfast	249
Sodium bicarbonate		Stromectol		Teligent	
Blood	44	Sucralfate		Temaccord	
Extemporaneous		Sulfadiazine Silver		Temazepam	
Sodium calcium edetate		Sulfadiazine sodium		Temozolomide	164
Sodium chloride		Sulfasalazine		Tenofovir disoproxil	
Blood	44	Sulphur		Tenofovir Disoproxil Emtricitabine	
Respiratory		Sulprix		Viatr	
Sodium citrate with sodium lauryl		Sumagran	133	Tenofovir Disoproxil Viatris	
sulphoacetate	25	Sumatriptan	133	Tenoxicam	
Sodium citro-tartrate		Sunitinib		Tensipine MR10	
Sodium cromoglicate		Sunitinib Pfizer		Tepadina	
Alimentary	8	Sunscreens		Terbinafine	
Sensory		Sunscreens, proprietary		Terbutaline sulphate	
Sodium Fusidate [fusidic acid]		Survimed OPD		Teriflunomide	
Dermatological	69	Sustagen Hospital Formula		Teriparatide	
Infection		Sustagen Hospital Formula		Teriparatide - Teva	
Sensory		Active	280	Testogel	
Sodium hyaluronate [Hyaluronic		Sustanon Ampoules		Testosterone	
acid]	263	Sylvant		Testosterone cipionate	
Sodium phenylbutyrate		Symbicort Turbuhaler 100/6		Testosterone esters	
Sodium picosulfate		Symbicort Turbuhaler 200/6		Testosterone undecanoate	
	-	,	_		_

Tetrabenazine	123	Treatments for Substance		- V -	
Tetrabromophenol	83	Dependence	148	Vaccinations	293
Tetracosactrin	86	Trelegy Ellipta	253	Vaclovir	107
Tetracycline	99	Trental 400		Valaciclovir	107
Teva Lisinopril		Tretinoin		Valganciclovir	108
Teva-Ketoconazole		Dermatological	68	Valganciclovir Viatris	
Teva-Salbutamol Sterinebs		Oncology		Valine50	
P.F	252	Trexate	157	Vancomycin	
Thalidomide	164	Triamcinolone acetonide		Vannair	
Thalomid	164	Alimentary	32	Varenicline Pfizer	
Theophylline		Dermatological		Varenicline tartrate	150
Thiamine hydrochloride	32	Hormone		Varicella vaccine [Chickenpox	
Thiamine multichem	32	Triamcinolone acetonide with		vaccine]	305
THIO-TEPA		gramicidin, neomycin and ny	statin	Varicella zoster vaccine [Shing	
Thioguanine		Dermatological		vaccine]	
Thiotepa		Sensory		Various	
Thyroid and Antithyroid Agents		Trientine		Varivax	
Ticagrelor		Trientine Waymade		Vasodilators	56
Ticagrelor Sandoz		Trikafta		Vasopressin Agonists	
Tilcotil		Trimethoprim		Vasorex	
Timolol		Trimethoprim with		Vebulis	
Tiotropium bromide		sulphamethoxazole		Vedafil	
Tiotropium bromide with		[Co-trimoxazole]	102	Vedolizumab	
olodaterol	253	Trisequens		Veletri	
Tivicay		Trisul		Venclexta	
Tixagevimab with cilgavimab		Trophic Hormones		Venetoclax	
TMP		Tropicamide		Venlafaxine	
Tobramycin		Trulicity		Venomil	
Infection	102	TruSteel		VENOX	
Sensory		Tryzan		Ventolin	.251-252
Tobramycin (Viatris)		Tuberculin PPD [Mantoux] test.		Ventolin Nebules	
Tobramycin BNM		Tubersol		Vepesid	
Tobrex		Two Cal HN		Verapamil hydrochloride	
Tocilizumab		TYR Explore 5		Vermox	
Tofranil		TYR Sphere 20		Versacloz	
Tolcapone		Tyrosine1000		Vesanoid	
Tolvaptan		Tysabri		Vexazone	
Topamax		- U -		Vfend	
Topical Products for Joint and		UK Synacthen	86	Viaderm KC	
Muscular Pain	117	Ultibro Breezhaler		Victoza	
Topiramate		Ultraproct		Vigabatrin	
Topiramate Actavis		Umeclidinium		Vigisom	142
Total parenteral nutrition (TPN)		Umeclidinium with vilanterol		Vildagliptin	
TPN		Univent		Vildagliptin with metformin	
Tramadol hydrochloride		Upadacitinib		hydrochloride	12
Tramal SR 100		Ural		Vimpat	
Tramal SR 150		Urea		Vinblastine sulphate	
Tramal SR 200		Urex Forte		Vincristine sulphate	
Trandate		Urinary Agents		Vinorelbine	
Tranexamic acid		Urinary Tract Infections		Vinorelbine Ebewe	
Tranylcypromine sulphate		Urinorm		Vinorelbine Te Arai	
Trastuzumab (Herzuma)		Uromitexan		Viramune Suspension	
Trastuzumab emtansine		Ursodeoxycholic acid		ViruPOS	250
Travatan		Ursosan		Vit.D3	
Travoprost		Ustekinumab		Vita-B12	
Treatments for Dementia		Utrogestan		Vita-POS	
HOGHINGING FOR DEHICHING	140	oogosiai1		*1011 OU	200

Vitabdeck33
Vital276
Vitamin B complex32
Vitamin B6 2532
Vitamins32–33
Vitarubin Depot Injection32
Vivonex TEN277
Voltaren
Voltaren D
Voltaren Ophtha260
Voltaren SR
Volumatic
Voriconazole
Votrient 169
Vttack
- W -
Warfarin sodium43
Wart Preparations
Wasp venom allergy treatment249
Water
Blood44
Extemporaneous 269
White Soft Liquid Paraffin AFT72
Wool fat with mineral oil
- X -
Xarelto42
Xifaxan
XMET Maxamum284
Xolair
XP Maxamum
Xylocaine 123
Xylocaine 2% Jelly
Xylocard 500
Xyntha39
- Z -
Zapril
Zarontin
Zaroxolyn53
Zavedos
Zeffix
Zejula
Zematop
Zetlam
Ziagen
Zidovudine [AZT]112
Zidovudine [AZT] with
lamivudine
Ziextenzo
Zimybe
Zinc and castor oil71
Zinc sulphate35
Zincaps
•
Zista
Zithromax96

Zoledronic acid	
Hormone	85
Musculoskeletal	120
Zoledronic acid Viatris	
Hormone	85
Musculoskeletal	120
Zopiclone	143
Zopiclone Actavis	
Zostrix	117
Zostrix HP	124
Zuclopenthixol decanoate	
Zuclopenthixol hydrochloride	136
Zusdone	136
Zyban	149
Zypine	
Zypine ODT	
Zyprexa Relprevy	137
Zytiga	172
710/02	106