SA2103 - Etanercept

Arthritis - rheumatoid - Renewal Arthritis - rheumatoid - Initial application	9
Adult-onset Still's disease - Initial application	2
Adult-onset Still's disease - Renewal	
Ankylosing spondylitis - Initial application	3
Ankylosing spondylitis - Renewal	4
Oligoarticular course juvenile idiopathic arthritis - Initial application	5
Oligoarticular course juvenile idiopathic arthritis - Renewal	6
Polyarticular course juvenile idiopathic arthritis - Initial application	4
Polyarticular course juvenile idiopathic arthritis - Renewal	5
Psoriatic arthritis - Initial application	7
Psoriatic arthritis - Renewal	8
Pyoderma gangrenosum - Initial application	8
Pyoderma gangrenosum - Renewal	8
Severe chronic plaque psoriasis - Initial application	11
Severe chronic plaque psoriasis - Renewal	
Undifferentiated spondyloarthritis - Initial application	13
Undifferentiated spondyloarthritis - Renewal	13

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
	Address:	
Fax Number:		Fax Number:

Etanercept

		The patient has had an initial Special Authority approval for adalimumab for adult-onset Still's disease (AOSD)
		or The patient has been started on tocilizumab for AOSD in a Health NZ Hospital
	and	
		The patient has experienced intolerable side effects from adalimumab and/or tocilizumab
		The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or tocilizumab such that they do not meet the renewal criteria for AOSD
or		
	and	Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430)
	[Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal anti-inflammatory drugs (NSAIDs) and methotrexate
	and	Patient has persistent symptoms of disabling poorly controlled and active disease

Renewal — adult-onset Still's disease

Current approval Number (if known):.....

Applications only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months. Prerequisites(tick boxes where appropriate)

	or		Applicant is a rheumatologist
			Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment
and		The p	patient has a sustained improvement in inflammatory markers and functional status

I confirm the above details are correct and that in signing this form I understand I may be audited.

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
	Address:	
Fax Number:		Fax Number:

Etanercept - continued

	and		The patient has had an initial Special Authority approval for adalimumab for ankylosing spondylitis
		or	The patient has experienced intolerable side effects from adalimumab
		-	The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for ankylosing spondylitis
or			
	and		Patient has a confirmed diagnosis of ankylosing spondylitis present for more than six months
	and		Patient has low back pain and stiffness that is relieved by exercise but not by rest
	[Patient has bilateral sacroiliitis demonstrated by plain radiographs, CT or MRI scan
	and		Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis
	and	or	Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right)
		01	Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender (see Notes)
	and		A Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale
sure n	nust be	no i	ust have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI more than 1 month old at the time of initial application.
24 year	rs - Mal	e: 7	.0 cm; Female: 5.5 cm
			.5 cm; Female: 5.5 cm .5 cm; Female: 4.5 cm

Enquiries	to Ministry	of Health
0800 855	066	

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
	Address:	
Fax Number:		Fax Number:
Etanercept - continued		

ent app	proval	Number (if known):
	-	r from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months. Sk boxes where appropriate)
and [and [and	۹ ۹ [] ۹	Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less Physician considers that the patient has benefited from treatment and that continued treatment is appropriate Etanercept to be administered at doses no greater than 50 mg every 7 days
	E	Etanercept to be administered at doses no greater than 50 mg every 7 days
		n — polyarticular course juvenile idiopathic arthritis
	ites(tio	y from a named specialist or rheumatologist. Approvals valid for 6 months. ck boxes where appropriate) The patient has had an initial Special Authority approval for adalimumab for polyarticular course juvenile idiopathic arthritis (JIA)
		ck boxes where appropriate)
	ites(tio	Ck boxes where appropriate) The patient has had an initial Special Authority approval for adalimumab for polyarticular course juvenile idiopathic arthritis (JIA) or The patient has experienced intolerable side effects from adalimumab
	ites(tio	Ck boxes where appropriate) The patient has had an initial Special Authority approval for adalimumab for polyarticular course juvenile idiopathic arthritis (JIA) The patient has experienced intolerable side effects from adalimumab The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for polyarticular
equisi	and	The patient has had an initial Special Authority approval for adalimumab for polyarticular course juvenile idiopathic arthritis (JIA) Image: Contract of the patient has experienced intolerable side effects from adalimumab Image: Contract of the patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for polyarticular course JIA
equisi	ites(tio	The patient has had an initial Special Authority approval for adalimumab for polyarticular course juvenile idiopathic arthritis (JIA) Image: Contract of the patient has experienced intolerable side effects from adalimumab Image: Contract of the patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for polyarticular
equisi	and	 The patient has had an initial Special Authority approval for adalimumab for polyarticular course juvenile idiopathic arthritis (JIA) The patient has experienced intolerable side effects from adalimumab The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for polyarticular course JIA To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance Patient has had polyarticular course JIA for 6 months duration or longer At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose)
equisi	and	The patient has had an initial Special Authority approval for adalimumab for polyarticular course juvenile idiopathic arthritis (JIA) Image: Image
equisi	and	ck boxes where appropriate) The patient has had an initial Special Authority approval for adalimumab for polyarticular course juvenile idiopathic arthritis (JIA or The patient has experienced intolerable side effects from adalimumab or The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for polyarticular course JIA To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance Patient has had polyarticular course JIA for 6 months duration or longer At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum toleracted dose) or Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum toleracted dose)

Enquiries	to Ministry	of Health
0800 855	066	

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
	Address:	
Fax Number:		Fax Number:

Etanercept - continued

Current approval Number (If known):	Renewal	l — poly	yarticular course juvenile idiopathic arthritis
valid for 6 months. Prerequisites(tick boxes where appropriate) and	Current a	approval	Number (if known):
Prerequisites(tick boxes where appropriate) Prerequisites(tick boxes where appropriate) and Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline Or On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline Initial application — oligoarticular course juvenile idiopathic arthritis Applications only from a named specialist or rheumatologist. Approvals valid for 6 months. Prerequisites(tick boxes where appropriate) Initial application — oligoarticular course juvenile idiopathic arthritis Applications only from a named specialist or rheumatologist. Approvals valid for 6 months. Prerequisites(tick boxes where appropriate) Initial application — oligoarticular course juvenile idiopathic arthritis Applications only from a named specialist or rheumatologist. Approvals valid for 6 months. Prerequisites(tick boxes where appropriate) Initial application — oligoarticular course juvenile idiopathic arthritis Application — oligoarticular course juvenile idiopathic arthritis Or Or Or Description The patient has had an initial Special Authority approval for adalimumab for oligoarticular course juvenile idiopathic arthritis (JIA) Or Description The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for oligoarticular course JIA Or Or Or Description And Description To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance Or Description And Description			
and			
or physician's global assessment from baseline or On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline Initial application - oligoarticular course juvenile idiopathic arthritis Applications only from a named specialist or rheumatologist. Approvals valid for 6 months. Prerequisites(tick boxes where appropriate) Image: the patient has had an initial Special Authority approval for adalimumab for oligoarticular course juvenile idiopathic arthritis (JIA) or The patient has experienced intolerable side effects from adalimumab or The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for oligoarticular course JIA or To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance Patient has had oligoarticular course JIA for 6 months duration or longer	an		Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance
Initial application — oligoarticular course juvenile idiopathic arthritis Applications only from a named specialist or rheumatologist. Approvals valid for 6 months. Prerequisites(tick boxes where appropriate) Initial application — oligoarticular course juvenile idiopathic arthritis Applications only from a named specialist or rheumatologist. Approvals valid for 6 months. Prerequisites(tick boxes where appropriate) Image: The patient has had an initial Special Authority approval for adalimumab for oligoarticular course juvenile idiopathic arthritis (JIA) or The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for oligoarticular course JIA or Image: The patient has had on unitial course therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance			
Applications only from a named specialist or rheumatologist. Approvals valid for 6 months. Prerequisites(tick boxes where appropriate)			
Applications only from a named specialist or rheumatologist. Approvals valid for 6 months. Prerequisites(tick boxes where appropriate)			
and or The patient has experienced intolerable side effects from adalimumab or The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for oligoarticular course JIA or and To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance Patient has had oligoarticular course JIA for 6 months duration or longer	Applicati	tions only	y from a named specialist or rheumatologist. Approvals valid for 6 months.
or The patient has experienced intolerable side effects from adalimumab or The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for oligoarticular course JIA or and To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance Patient has had oligoarticular course JIA for 6 months duration or longer			
or The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for oligoarticular course JIA or and Patient has had oligoarticular course JIA for 6 months duration or longer		and	The patient has had an initial Special Authority approval for adalimumab for oligoarticular course juvenile idiopathic arthritis (JIA)
or The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for oligoarticular course JIA or Patient has an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance Patient has had oligoarticular course JIA for 6 months duration or longer			
To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance and Patient has had oligoarticular course JIA for 6 months duration or longer			The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for oligoarticular
and Patient has had oligoarticular course JIA for 6 months duration or longer	or	·	
Patient has had oligoarticular course JIA for 6 months duration or longer		and	To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance
		and	Patient has had oligoarticular course JIA for 6 months duration or longer
At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose)			maximum tolerated dose)
Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose)			Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose)
High disease activity (cJADAS10 score greater than 4) after a 6-month trial of methotrexate			

Enquiries	to Ministry	of Health
0800 855	066	

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
	Address:	
Fax Number:		Fax Number:
Etanercept - continued		

Renewal — oligoarticular course juvenile idiopathic arthritis	
Current approval Number (if known): Applications only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months. Prerequisites (tick boxes where appropriate)	
Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance and	
 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline 	

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
	Address:	
Fax Number:		Fax Number:

Etanercept - continued

Initial application — psoriatic arthritis Applications only from a rheumatologist. Approvals valid for 6 months. Prerequisites(tick boxes where appropriate)				
	an	The patient has had an initial Special Authority approval for adalimumab or secukinumab for psoriatic arthritis		
		The patient has experienced intolerable side effects from adalimumab or secukinumab		
		The patient has received insufficient benefit from adalimumab or secukinumab to meet the renewal criteria for adalimuma or secukinumab for psoriatic arthritis	b	
o	r			
	an	Patient has had severe active psoriatic arthritis for six months duration or longer		
	an	Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose		
	an	Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses)		
		Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints		
		or Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip		
	an		- -	
		Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application or		
		Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or		
		ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months		

Enquiries	to Ministry	of Health
0800 855	066	

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
	Address:	
Fax Number:		Fax Number:

Etanercept - continued

enewal –	– psoriatic arthritis
urrent ap	proval Number (if known):
	s only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months. Ites(tick boxes where appropriate)
	or Applicant is a rheumatologist
	Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment
and	Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician
	The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician
and [Etanercept to be administered at doses no greater than 50 mg every 7 days
rerequisi and and	 ns only from a dermatologist. Approvals valid for 4 months. ites(tick boxes where appropriate) Patient has pyoderma gangrenosum* Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response A maximum of 8 doses cations marked with * are unapproved indications.
enewal –	– pyoderma gangrenosum
urrent ap	proval Number (if known):
-	s only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months. Ites(tick boxes where appropriate)
and	Patient has shown clinical improvement
and	Patient continues to require treatment
	A maximum of 8 doses

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
	Address:	
Fax Number:		Fax Number:

Etanercept - continued

Appl	icatio	ns only	n — Arthritis - rheumatoid from a rheumatologist. Approvals valid for 6 months. k boxes where appropriate)	
		and	The patient has had an initial Special Authority approval for adalimumab for rheumatoid arthritis	
			The patient has experienced intolerable side effects	
			The patient has received insufficient benefit to meet the renewal criteria for rheumatoid arthritis	
	or			
		and	Patient has had rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer	
		and	Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance	
		Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated)		
		 Patient has tried and not responded to at least three months of methotrexate in combination with sulfasalazine and hydroxychloroq sulphate (at maximum tolerated doses unless contraindicated) and 		
			Patient has tried and not responded to at least three months of methotrexate in combination with the maximum tolerated dose of ciclosporin	
			Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with methotrexate	
		and		
			Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints	
			Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip	

Enquiries to Ministry of Health	
0800 855 066	

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
	Address:	
Fax Number:		Fax Number:
Etanercept - continued		
Renewal — Arthritis - rheumatoid Current approval Number (if known): Applications from any relevant practitioner. Approv Prerequisites(tick boxes where appropriate) Treatment is to be used as an adju- intolerance		se of methotrexate is limited by toxicity or
or On subsequent reapplication	ne patient has at least a 50% decrease in active joint opinion of the physician ns, the patient demonstrates at least a continuing 30% ificant response to treatment in the opinion of the phy	% improvement in active joint count from
and Etanercept to be administered at c	loses no greater than 50 mg every 7 days	

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
	Address:	
Fax Number:		Fax Number:

Etanercept - continued

		from a dermatologist. Approvals valid for 4 months. boxes where appropriate)
	and	The patient has had an initial Special Authority approval for adalimumab for severe chronic plaque psoriasis
		The patient has experienced intolerable side effects from adalimumab
		The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe chronic plaque psoriasis
or		
		 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis
	and	Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin
	and	A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course
		The most recent PASI or DLQI assessment is no more than 1 month old at the time of application
while still o nand or fo	on treatr ot, at lea	e response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably nent but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, ast 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following

cessation of the most recent prior treatment.

Enquiries	to Ministry	of Health
0800 855	066	

APPLICANT (stamp or sticker acceptable)		PATIENT NHI:	REFERRER Reg No:
Reg No:		First Names:	First Names:
Name:		Surname:	Surname:
Address:		DOB:	Address:
		Address:	
Fax Number: .			Fax Number:
Etanercept	- continued		
Renewal — s	evere chronic plaque psoriasis		
Applications o	val Number (if known): nly from a dermatologist or Practition (tick boxes where appropriate)	ner on the recommendation of a dermatologist. Appro	ovals valid for 6 months.
and	Applicant is a Practitioner ar continues with etanercept tre Patient had "whole bo and Following each or Following each Following each	d confirms that a dermatologist has provided a letter, eatment dy" severe chronic plaque psoriasis at the start of tree prior etanercept treatment course the patient has a P at this level, when compared with the pre-treatment ba prior etanercept treatment course the patient has a D 5 or more, when compared with the pre-treatment ba	atment ASI score which is reduced by 75% or more, aseline value ermatology Quality of Life Index (DLQI)
	And Patient had severe ch and Following each all 3 of erythems course baseline Following each Following each	ronic plaque psoriasis of the face, or palm of a hand of prior etanercept treatment course the patient has a re a, thickness and scaling, to slight or better, or sustain values prior etanercept treatment course the patient has a re tained at this level, as compared to the pre-treatment	eduction in the PASI symptom subscores for ed at this level, as compared to the treatment eduction of 75% or more in the skin area
and	·	oses no greater than 50 mg every 7 days n of 12 weeks of etanercept treatment	

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
	Address:	
Fax Number:		Fax Number:

Etanercept - continued

			tion — undifferentiated spondyloarthritis nly from a rheumatologist. Approvals valid for 6 months.
			(tick boxes where appropriate)
	and		Patient has undifferentiated peripheral spondyloarthritis* with active peripheral joint arthritis in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip
	and		Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose
	and		Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day (or maximum tolerated dose)
	and [Patient has tried and not responded to at least three months of leflunomide at a dose of up to 20 mg daily (or maximum tolerated dose)
		or	Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application
		or	Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour measured no more than one month prior to the date of this application
		-	ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months
Note	: Indic	atio	ns marked with * are unapproved indications.
Ren	ewal –	– un	ndifferentiated spondyloarthritis
Curr	ent ap	prov	al Number (if known):
Applications only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months. Prerequisites (tick boxes where appropriate)			
		or	Applicant is a rheumatologist
			Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment
	and		
		or	Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician
			The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician
I	and [Etanercept to be administered at doses no greater than 50 mg dose every 7 days