APPLICATION FOR SUBSIDY BY SPECIAL AUTHORITY

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
	Address:	
Fax Number:		Fax Number:

Bendamustine hydrochloride

	Initial application — treatment naive CLL Applications only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months.			
Prerequisit	es(tick boxes where appropriate)			
and	The patient has Binet stage B or C, or progressive stage A chronic lymphocytic leukaemia requiring treatment			
and	The patient is chemotherapy treatment naive			
and _	The patient is unable to tolerate toxicity of full-dose FCR			
and	Patient has ECOG performance status 0-2			
and_	Patient has a Cumulative Illness Rating Scale (CIRS) score of < 6			
	Bendamustine is to be administered at a maximum dose of 100 mg/m ² on days 1 and 2 every 4 weeks for a maximum of 6 cycles			
	nic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL). Chemotherapy treatment is considered to comprise a known prapeutic chemotherapy regimen and supportive treatments.			
Applications	cation — Indolent, Low-grade lymphomas only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months. es(tick boxes where appropriate)			
	The patient has indolent low grade NHL requiring treatment			
and and	Patient has a WHO performance status of 0-2			
	Patient is treatment naive			
	Bendamustine is to be administered for a maximum of 6 cycles (in combination with rituximab when CD20+)			
	or			
	Patient is refractory to or has relapsed within 12 months of a rituximab containing combined chemo-immunotherapy regimen and			
	Bendamustine is to be administered in combination with obinutuzumab for a maximum of 6 cycles			
	or			
	and and and a solution of the			
	Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+)			
	And Patient has had a rituximab treatment-free interval of 12 months or more			
	or Bendamustine is to be administered as monotherapy for a maximum of 6 cycles in rituximab refractory patients			

I confirm the above details are correct and that in signing this form I understand I may be audited.

APPLICATION FOR SUBSIDY **BY SPECIAL AUTHORITY**

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
	Address:	
Fax Number:		Fax Number:

Bendamustine hydrochloride - continued

Applications only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months. Prerequisites(tick boxes where appropriate)		Renewal — Indolent, Low-grade lymphomas			
Prerequisites(tick boxes where appropriate) Image: Patient is refractory to or has relapsed within 12 months of rituximab in combination with bendamustine and Bendamustine is to be administered in combination with obinutuzumab for a maximum of 6 cycles or Patients have not received a bendamustine regimen within the last 12 months and Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+) and Patient has had a rituximab treatment-free interval of 12 months or more or Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients Note: Indolent, low-grade lymphomas' includes follicular, mantle cell, marginal zone and lymphoplasmacytic/ Waldenstrom's macroglobulinaemia. Initial application — Hodgkin's lymphoma* Application of a relevant specialist. Approvals valid for 6 months. Prerequisites(tick boxes where appropriate) Patient has Hodgkin's lymphoma requiring treatment and Patient has Hodgkin's lymphoma requiring treatment and Patient has ECOG performance status of 0-2 and Patient has received one prior line of chemotherapy and Patient's disease relapsed or was refractory following prior chemotherapy	Current approval Number (if known):				
and Bendamustine is to be administered in combination with obinutuzumab for a maximum of 6 cycles or Patients have not received a bendamustine regimen within the last 12 months and Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+) and Patient has had a rituximab treatment-free interval of 12 months or more or Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients Note: 'indolent, low-grade lymphomas' includes follicular, mantle cell, marginal zone and lymphoplasmacytic/ Waldenstrom's macroglobulinaemia. Initial application - Hodgkin's lymphoma* Application or a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months. Prerequisites(tick boxes where appropriate) Patient has Hodgkin's lymphoma requiring treatment and Patient has a ECOG performance status of 0-2 and Patient has received one prior line of chemotherapy and Patient's disease relapsed or was refractory following prior chemotherapy					
or Patient has Hodgkin's lymphoma requiring treatment Patient has Rodgkin's lymphoma requiring treatment Patient has a ECOG performance status of 0-2 and Patient is disease relapsed or was refractory following prior chemotherapy					
Image: Patients have not received a bendamustine regimen within the last 12 months and Patients have not received a bendamustine regimen within the last 12 months and Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+) and Patient has had a rituximab treatment-free interval of 12 months or more or Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients Note: 'indolent, low-grade lymphomas' includes follicular, mantle cell, marginal zone and lymphoplasmacytic/ Waldenstrom's macroglobulinaemia. Initial application — Hodgkin's lymphoma* Applications only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months. Prerequisites(tick boxes where appropriate) Patient has a ECOG performance status of 0-2 and Patient has received one prior line of chemotherapy and Patient has received one prior line of chemotherapy and Patient's disease relapsed or was refractory following prior chemotherapy					
Initial application — Hodgkin's lymphoma* Applications only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months. Prerequisites(tick boxes where appropriate) Patient has Hodgkin's lymphoma requiring treatment and Patient has received one prior line of chemotherapy and Patient has received one prior line of chemotherapy and Patient's disease relapsed or was refractory following prior chemotherapy		Patients have not received a bendamustine regimen within the last 12 months			
Image: Patient has had a rituximab treatment-free interval of 12 months or more or Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients Note: 'indolent, low-grade lymphomas' includes follicular, mantle cell, marginal zone and lymphoplasmacytic/ Waldenstrom's macroglobulinaemia. Initial application - Hodgkin's lymphoma* Applications only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months. Prerequisites(tick boxes where appropriate) Image: Patient has Hodgkin's lymphoma requiring treatment and Patient has a ECOG performance status of 0-2 and Patient has received one prior line of chemotherapy and Patient's disease relapsed or was refractory following prior chemotherapy			when CD20+)		
Image: Sendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients Note: 'indolent, low-grade lymphomas' includes follicular, mantle cell, marginal zone and lymphoplasmacytic/ Waldenstrom's macroglobulinaemia. Initial application — Hodgkin's lymphoma* Applications only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months. Prerequisites(tick boxes where appropriate) Image: Patient has Hodgkin's lymphoma requiring treatment and Patient has a ECOG performance status of 0-2 and Patient has received one prior line of chemotherapy and Patient has received one prior line of chemotherapy and Patient's disease relapsed or was refractory following prior chemotherapy					
Initial application — Hodgkin's lymphoma* Applications only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months. Prerequisites(tick boxes where appropriate) Patient has Hodgkin's lymphoma requiring treatment and Patient has a ECOG performance status of 0-2 and Patient has received one prior line of chemotherapy and Patient's disease relapsed or was refractory following prior chemotherapy and					
Applications only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months. Prerequisites(tick boxes where appropriate) Patient has Hodgkin's lymphoma requiring treatment and Patient has a ECOG performance status of 0-2 and Patient has received one prior line of chemotherapy and Patient's disease relapsed or was refractory following prior chemotherapy	Note:	'indo	olent, low-grade lymphomas' includes follicular, mantle cell, marginal zone and lymphoplasmacytic/ Waldenstrom's macroglobulinaemia.		
and Patient has a ECOG performance status of 0-2 and Patient has received one prior line of chemotherapy and Patient's disease relapsed or was refractory following prior chemotherapy and	Appli	cation	ns only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months.		
Patient has a ECOG performance status of 0-2 and Patient has received one prior line of chemotherapy and Patient's disease relapsed or was refractory following prior chemotherapy and		and			
and Patient's disease relapsed or was refractory following prior chemotherapy and		Patient has a ECOG performance status of 0-2			
and		Patient has received one prior line of chemotherapy			
	Patient's disease relapsed or was refractory following prior chemotherapy				
Bendamustine is to be administered in combination with gemcitabine and vinorelbine (BeGeV) at a maximum dose of no greater than 90 mg/m2 twice per cycle, for a maximum of four cycles		Į	Bendamustine is to be administered in combination with gemcitabine and vinorelbine (BeGeV) at a maximum dose of no greater than 90 mg/m2 twice per cycle, for a maximum of four cycles		

Note: Indications marked with * are unapproved indications.

I confirm the above details are correct and that in signing this form I understand I may be audited.