# December 2024 Volume 31

#### **Editors:**

Kaye Wilson, & Doris Chong email: enquiry@pharmac.govt.nz Telephone +64 4 460 4990 Level 9, 40 Mercer Street PO Box 10 254 Wellington

Freephone Information Line 0800 66 00 50 (9am – 5pm weekdays)

#### Circulation

You can register to have an electronic version of the Pharmaceutical Schedule (link to PDF copy) emailed to your nominated email address each month by subscribing at <a href="mailto:pharmac.govt.nz/subscribe">pharmac.govt.nz/subscribe</a>.

### Production

Typeset automatically from XML and T<sub>E</sub>X. XML version of the Schedule available from schedule.pharmac.govt.nz/pub/schedule

### **Programmers**

Anrik Drenth

email: texschedule@pharmac.govt.nz @Pharmaceutical Management Agency



ISSN 1179-3686

This work is licensed under the Creative Commons Attribution 4.0 International licence. In essence, you are free to copy, distribute and adapt it, as long as you attribute the work to Pharmac and abide by the other licence terms. To view a copy of this licence, visit: creativecommons.org/licenses/by/4.0/. Attribution to Pharmac should be in written from and not by reproduction of the Pharmac logo. While care has been taken in compiling this Schedule, Pharmac takes no responsibility for any errors or omissions, and shall not be liable for any consequences arising there from.

	ind oddonig i narmae i	_
Section A	General Rules	5
Section B	Alimentary Tract & Metabolism	6
	Blood & Blood Forming Organs	38
	Cardiovascular System	48
	Dermatologicals	70
	Genito Urinary System	80
	Hormone Preparations – Systemic	87
	Infections – Agents For Systemic Use	98
	Musculoskeletal System	120
	Nervous System	126
	Oncology Agents & Immunosuppressants	156
	Respiratory System & Allergies	258
	Sensory Organs	268
	Various	273
Section C	Extemporaneous Compounds (ECPs)	276
Section D	Special Foods	278
Section I	National Immunisation Schedule	303
	Index	319

Introducing Pharmac

# **Introducing Pharmac**

The Pharmaceutical Management Agency (Pharmac) makes decisions that help control Government spending on pharmaceuticals. This includes community pharmaceuticals, hospital pharmaceuticals, vaccines and increasingly, hospital medical devices. Pharmac negotiates prices, sets subsidy levels and conditions, and makes decisions on changes to the subsidised list.

#### Pharmac's role:

"to secure for eligible people in need of pharmaceuticals, the best health outcomes that are reasonably achievable from pharmaceutical treatment and from within the amount of funding provided."

Pae Ora (Healthy Futures) Act 2022

To ensure our decisions are as fair and robust as possible we use a decision-making process that incorporates clinical, economic and commercial issues. We also seek the views of users and the wider community through consultation. The processes we generally use are outlined in our Operating Policies and Procedures.

Further information about Pharmac and the way we make funding decisions can be found on the Pharmac website at <a href="https://pharmac.govt.nz/about">https://pharmac.govt.nz/about</a>.

# **Purpose of the Pharmaceutical Schedule**

The purpose of the Schedule is to list:

- the Community Pharmaceuticals that are subsidised by the Government and to show the amount of the subsidy paid to contractors, as well as the manufacturer's price and any access conditions that may apply;
- the Hospital Pharmaceuticals that may be used in Health NZ Hospitals, as well as any access conditions that may apply; and
- the Pharmaceuticals, including Medical Devices, used in Health NZ Hospitals for which national prices have been negotiated by Pharmac.

The Schedule does not show the final cost to Government of subsidising each Community Pharmaceutical or to Health NZ Hospitals in purchasing each Pharmaceutical, since that will depend on any rebate and other arrangements Pharmac has with the supplier and, for Pharmaceuticals used in Health NZ Hospitals, on any logistics arrangements put in place.

This book contains sections A to D and Section I of the Pharmaceutical Schedule and lists the Pharmaceuticals funded for use in the community, including vaccines, as well as haemophilia and cancer treatments given in Health NZ Hospitals. Section H lists the Pharmaceuticals that that can be used in Health NZ Hospitals and is a separate publication.


The Pharmaceuticals in this book are listed by therapeutic group, which is based on the WHO Anatomical Therapeutic Chemical (ATC) system. The listings are displayed alphabetically under each heading.

The index lists both chemical entities and product brand names.

# **Explaining pharmaceutical entries**

The Pharmaceutical Schedule lists pharmaceuticals subsidised by the Government, the subsidy, the supplier's price and the access conditions that may apply.

### Example



# Glossary

### **Units of Measure**

gramg kilogramkg international unitiu	mi mi mi
Abbreviations	
AmpouleAmp	Ge
CapsuleCap	Gr
Cream	Inf
DeviceDev	Ini
DispersibleDisp	Lic
EffervescentEff	Lo
EmulsionEmul	Oi
Enteric Coated EC	Sa

microgrammilligrammillilitre	mg
Gelatinous	
Granules	
Infusion	Inf
Injection	Inj
Liquid	Liq
Long Acting	LA
Ointment	Oint
Sachet	Sach

millimoleunit	
Solution	Supp Tab
Trans Dermal Delivery System	TDDS

Read the General Rules: https://pharmac.govt.nz/section-a.

# SECTION B: ALIMENTARY TRACT AND METABOLISM

SECTION B: ALIMENTARY TRACT AND MET	TABOLISM			
	Subsidy (Manufacturer's Price) \$	Su Per	Fully bsidised	Brand or Generic Manufacturer
Antacids and Antiflatulents				
Antacids and Reflux Barrier Agents				
ALGINIC ACID  Sodium alginate 225 mg and magnesium alginate 87.5 mg pe sachet		30	<b>√</b> G	aviscon Infant
* Tab 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg - peppermint flavour	1.80 (14.39)	60	G	aviscon Extra Strength
Oral liq 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg per 10 ml		500 ml	A	cidex
Phosphate Binding Agents				
ALUMINIUM HYDROXIDE  * Tab 600 mg  CALCIUM CARBONATE  Oral liq 1,250 mg per 5 ml (500 mg elemental per 5 ml) –		100		lu-Tab
Subsidy by endorsement	47.30	500 ml 473 ml	<b>√</b> C	oxane alcium carbonate PAI <sup>829</sup>
Only when prescribed for patients unable to swallow calc inappropriate and the prescription is endorsed accordingl		ts or whe	ere calciur	n carbonate tablets are
Antidiarrhoeals				
Agents Which Reduce Motility				
LOPERAMIDE HYDROCHLORIDE – Up to 30 cap available on a  * Tab 2 mg  * Cap 2 mg	10.75	400 400	✓ N ✓ <u>D</u>	odia iamide Relief
Rectal and Colonic Anti-inflammatories				
BUDESONIDE  Cap modified-release 3 mg - Special Authority see SA1886  below - Retail pharmacy	87.60	90	<b>✓</b> <u>B</u>	udesonide Te Arai

# ⇒SA1886 Special Authority for Subsidy

Initial application — (Crohn's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Mild to moderate ileal, ileocaecal or proximal Crohn's disease; and
- 2 Any of the following:
  - 2.1 Diabetes; or

Subsidy	Ful	y Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per •	Manufacturer	

continued...

- 2.2 Cushingoid habitus; or
- 2.3 Osteoporosis where there is significant risk of fracture; or
- 2.4 Severe acne following treatment with conventional corticosteroid therapy; or
- 2.5 History of severe psychiatric problems associated with corticosteroid treatment; or
- 2.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
- 2.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated).

Initial application — (collagenous and lymphocytic colitis (microscopic colitis)) from any relevant practitioner. Approvals valid for 6 months where patient has a diagnosis of microscopic colitis (collagenous or lymphocytic colitis) by colonoscopy with biopsies.

**Initial application** — **(gut Graft versus Host disease)** from any relevant practitioner. Approvals valid for 6 months where patient has a gut Graft versus Host disease following allogenic bone marrow transplantation\*.

Note: Indication marked with \* is an unapproved indication.

Initial application — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has autoimmune hepatitis\*: and
- 2 Patient does not have cirrhosis; and
- 3 Any of the following:
  - 3.1 Diabetes: or
  - 3.2 Cushingoid habitus; or
  - 3.3 Osteoporosis where there is significant risk of fracture: or
  - 3.4 Severe acne following treatment with conventional corticosteroid therapy; or
  - 3.5 History of severe psychiatric problems associated with corticosteroid treatment; or
  - 3.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
  - 3.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated); or3.8 Adolescents with poor linear growth (where conventional corticosteroid use may limit further growth).

Note: Indication marked with \* is an unapproved indication.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

#### HYDROCORTISONE ACETATE

Rectal foam 10%, CFC-Free (14 applications)26.55	15 g OP	<ul><li>Colifoam</li></ul>
HYDROCORTISONE ACETATE WITH PRAMOXINE HYDROCHLORIDE		
Topical aerosol foam, 1% with pramoxine hydrochloride 1%26.55	10 g OP	✓ Proctofoam S29
MESALAZINE		
Tab 400 mg49.50	100	✓ Asacol
Tab long-acting 500 mg56.10	100	✓ Pentasa
Tab 800 mg85.50	90	✓ Asacol
		✓ Asacol S29 S29
Modified release granules, 1 g118.10	100 OP	✓ Pentasa
Enema 1 g per 100 ml41.30	7	✓ Pentasa
Suppos 500 mg22.80	20	✓ Asacol
Suppos 1 g50.96	28	✓ Pentasa

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
OLSALAZINE				
Tab 500 mg	56.02	60	<b>✓</b>	Atnahs
				Olsalazine S29
	93.37	100	✓ [	Dipentum
Cap 250 mg	53.00	100	✓ [	Dipentum
SODIUM CROMOGLICATE Cap 100 mg	113.35	100	<b>√</b> F	Ralicrom
SULFASALAZINE  * Tab 500 mg	16.52	100	<b>√</b> 9	Salazopyrin
* Tab EC 500 mg		100		Salazopyrin EN

# **Local preparations for Anal and Rectal Disorders**

# **Antihaemorrhoidal Preparations**

FLUOCORTOLONE CAPROATE WITH FLUOCORTOLONE PIVALATE AND CINCHOCAINE				
Oint 950 mcg, with fluocortolone pivalate 920 mcg, and				
cinchocaine hydrochloride 5 mg per g13.05	30 g OP	<ul><li>Ultraproct</li></ul>		
Suppos 630 mcg, with fluocortolone pivalate 610 mcg, and				
cinchocaine hydrochloride 1 mg8.61	12	<ul><li>Ultraproct</li></ul>		
HYDROCORTISONE WITH CINCHOCAINE				
Oint 5 mg with cinchocaine hydrochloride 5 mg per g15.00	30 g OP	✓ Proctosedyl		
Suppos 5 mg with cinchocaine hydrochloride 5 mg per g9.90	12	✓ Proctosedyl		

# **Management of Anal Fissures**

GLYCERYL TRINITRATE − Special Authority see SA1329 below − Retail pharmacy

★ Oint 0.2%......22.00 30 g OP

✓ Rectogesic

# **⇒SA1329** Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has a chronic anal fissure that has persisted for longer than three weeks.

# **Antispasmodics and Other Agents Altering Gut Motility**

Inj 200 mcg per ml, 1 ml ampoule – Up to 10 inj available on a			
PSO	19.00	5	✓ Robinul
HYOSCINE BUTYLBROMIDE			
* Tab 10 mg	2.25	20	<ul><li>✓ Hyoscine Butylbromide (Adiramedica)</li></ul>
	6.35	100	✓ Buscopan
* Inj 20 mg, 1 ml – Up to 5 inj available on a PSO(Buscopan Tab 10 mg to be delisted 1 April 2025)	1.91	5	✓ <u>Spazmol</u>
MEBEVERINE HYDROCHLORIDE			
* Tab 135 mg	8.50	90	✓ Colofac

ALIMENTARY TRACT AND METABOLISM Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer **Antiulcerants** Antisecretory and Cytoprotective MISOPROSTOL - Wastage claimable \* Tab 200 mcg - Up to 120 tab available on a PSO ......47.73 ✓ Cytotec 120 **Helicobacter Pylori Eradication CLARITHROMYCIN** 14 Klacid a) Maximum of 28 tab per prescription b) Subsidised only if prescribed for helicobacter pylori eradication and prescription is endorsed accordingly. Note: the prescription is considered endorsed if clarithromycin is prescribed in conjunction with a proton pump inhibitor and either amoxicillin or metronidazole. **H2 Antagonists** FAMOTIDINE - Only on a prescription \* Tab 20 mg ......4.91 ✓ Famotidine 100 Hovid S29 100 ✓ Famotidine Hovid S29 Inj 10 mg per ml, 4 ml - Subsidy by endorsement ......CBS 10 ✓ Mylan S29 Subsidy by endorsement – Subsidised for patients receiving treatment as part of palliative care. **Proton Pump Inhibitors** LANSOPRAZOLE ✓ Lanzol Relief 100 Lanzol Relief to be Principal Supply on 1 February 2025 100 ✓ Lanzol Relief Lanzol Relief to be Principal Supply on 1 February 2025 **OMEPRAZOLE** For omeprazole suspension refer Standard Formulae, page 276 ✓ Omeprazole Teva 90 Omeprazole actavis 10 ✓ Omeprazole Teva 90 ✓ Omeprazole actavis 20 ✓ Omeprazole Teva 90 ✓ Omeprazole actavis

PANTOPRAZOLE	
¥ Toh EC 00 mg	4

*	Tab EC 20 mg1.9	9 9	0
*	Tab EC 40 mg2.7	4 9	0

<sup>✓</sup> Panzop Relief
✓ Panzop Relief

40

✓ Midwest

✓ <u>Dr Reddy's</u>

<u>Omeprazole</u>
✓ Ocicure \$29

5 q

5

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price \$	e) Per	Fully Subsidised	Brand or Generic Manufacturer
Site Protective Agents				
COLLOIDAL BISMUTH SUBCITRATE  Tab 120 mgSUCRALFATE	14.51	50	<b>✓</b> G	astrodenol \$29
Tab 1 g	35.50 (48.28)	120	С	arafate
Bile and Liver Therapy				
RIFAXIMIN – Special Authority see SA1461 below – Retail pl Tab 550 mg  SA1461 Special Authority for Subsidy	,	56	<b>√</b> <u>X</u>	<u>ifaxan</u>
Initial application only from a gastroenterologist, hepatologist nepatologist. Approvals valid for 6 months where the patient tolerated doses of lactulose.  Renewal only from a gastroenterologist, hepatologist or Pract nepatologist. Approvals valid without further renewal unless repending from treatment.	has hepatic encephalo	oathy d	lespite an ad n of a gastro	lequate trial of maximum enterologist or
Diabetes				
Hyperglycaemic Agents	la a mara a su			
DIAZOXIDE - Special Authority see SA1320 below - Retail p Cap 25 mg Cap 100 mg Oral liq 50 mg per ml  → SA1320 Special Authority for Subsidy	110.00	100 100 30 ml C	<b>✓</b> P	roglicem \$29 roglicem \$29 5 Pharma \$29
nitial application from any relevant practitioner. Approvals hypoglycaemia caused by hyperinsulinism.	valid for 12 months whe	ere use	d for the trea	atment of confirmed
Renewal from any relevant practitioner. Approvals valid with appropriate and the patient is benefiting from treatment.  GLUCAGON HYDROCHLORIDE	out further renewal unle	ess noti	ified where th	he treatment remains
Inj 1 mg syringe kit – Up to 5 kit available on a PSO	32.00	1	<b>✓</b> G	lucagen Hypokit
Insulin - Short-acting Preparations				
NSULIN NEUTRAL ▲ Inj human 100 u per ml	25.26	10 ml C		ctrapid lumulin R
▲ Inj human 100 u per ml, 3 ml	42.66	5	✓ A	ctrapid Penfill umulin R
Insulin - Intermediate-acting Preparations				
NSULIN ASPART WITH INSULIN ASPART PROTAMINE  Inj 100 iu per ml, 3 ml prefilled pen	52.15	5	✓ N	ovoMix 30 FlexPen

	Subsidy		Fully Brand or
	(Manufacturer's F		idised Generic
	\$	Per	✓ Manufacturer
NSULIN ISOPHANE			
Inj human 100 u per ml		10 ml OP	<ul><li>✓ Humulin NPH</li><li>✓ Protaphane</li></ul>
Inj human 100 u per ml, 3 ml	29.86	5	<ul><li>✓ Humulin NPH</li><li>✓ Protaphane Penfill</li></ul>
NSULIN ISOPHANE WITH INSULIN NEUTRAL			
Inj human with neutral insulin 100 u per ml	25.26	10 ml OP	<ul><li>✓ Humulin 30/70</li><li>✓ Mixtard 30</li></ul>
Inj human with neutral insulin 100 u per ml, 3 ml	42.66	5	✓ Humulin 30/70 ✓ PenMix 30 ✓ PenMix 50
Mixtard 30 Inj human with neutral insulin 100 u per ml to be delis PenMix 50 Inj human with neutral insulin 100 u per ml, 3 ml to be			
NSULIN LISPRO WITH INSULIN LISPRO PROTAMINE			
▲ Inj lispro 25% with insulin lispro protamine 75% 100 u per ml, 3 ml		5	✓ Humalog Mix 25
▲ Inj lispro 50% with insulin lispro protamine 50% 100 u per ml, 3 ml		5	✓ Humalog Mix 50
Insulin - Long-acting Preparations			
NSULIN GLARGINE			
Inj 100 u per ml, 10 ml	63.00	1	✓ Lantus
Inj 100 u per ml, 3 ml		5	✓ Lantus
Inj 100 u per ml, 3 ml disposable pen		5	✓ Lantus SoloStar
Insulin - Rapid Acting Preparations			
NSULIN ASPART			
Inj 100 u per ml, 10 ml	30.03	1	✓ NovoRapid
Inj 100 u per ml, 3 ml	51.19	5	✓ NovoRapid Penfill
Inj 100 u per ml, 3 ml syringe	51.19	5	NovoRapid FlexPen
NSULIN GLULISINE			
Inj 100 u per ml, 10 ml	27.03	1	✓ Apidra
Inj 100 u per ml, 3 ml	46.07	5	✓ Apidra
Inj 100 u per ml, 3 ml disposable pen	46.07	5	✓ Apidra SoloStar
NSULIN LISPRO			
Inj 100 u per ml, 10 ml	34.92	10 ml OP	✓ Humalog
Inj 100 u per ml, 3 ml		5	✓ Humalog
Alpha Glucosidase Inhibitors			
ACARBOSE			
★ Tab 50 mgAccarb to be Principal Supply on 1 February 2025	11.20	90	✓ Accarb
* Tab 100 mgAccarb to be Principal Supply on 1 February 2025	17.38	90	✓ Accarb
Oral Hypoglycaemic Agents			
GLIBENCLAMIDE			
<b>≮</b> Tab 5 mg	7.50	100	✓ Daonil

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	
GLICLAZIDE			_	
* Tab 80 mg	20.10	500	•	Glizide
GLIPIZIDE				
* Tab 5 mg	6.86	100	•	Minidiab
Minidiab to be Principal Supply on 1 March 2025				
METFORMIN HYDROCHLORIDE			_	
* Tab immediate-release 500 mg		1,000	_	Metformin Viatris
* Tab immediate-release 850 mg	11.28	500	•	Metformin Viatris
PIOGLITAZONE				
* Tab 15 mg		90	_	<u>Vexazone</u>
* Tab 30 mg		90	/	Vexazone
* Tab 45 mg	12.00	90	•	<u>Vexazone</u>
VILDAGLIPTIN				
Tab 50 mg	35.00	60	•	Galvus
VILDAGLIPTIN WITH METFORMIN HYDROCHLORIDE				
Tab 50 mg with 1,000 mg metformin hydrochloride	35.00	60	•	Galvumet
Tab 50 mg with 850 mg metformin hydrochloride	35.00	60	/	Galvumet

# **GLP-1 Agonists**

DULAGLUTIDE - Special Authority see SA2338 below - Retail pharmacy

Note: Not to be given in combination with a funded SGLT-2 inhibitor or other GLP-1 agonist.

## ⇒SA2338 Special Authority for Subsidy

Note: Subsidy for patients with existing approvals prior to 1 May 2024. Approvals valid without further renewal unless notified. No new patients will be granted from 1 May 2024 until further notice.

LIRAGLUTIDE - Special Authority see SA2339 below - Retail pharmacy

- a) Maximum of 9 inj per prescription
- b)
- a) Not to be given in combination with a funded SGLT-2 inhibitor or other GLP-1 agonist.
- b) Maximum of 1 pack of 3 (6 mg per ml, 3 ml) prefilled pens will be funded per month.

### ⇒SA2339 Special Authority for Subsidy

Note: Subsidy for patients with existing approvals prior to 1 May 2024. Approvals valid without further renewal unless notified. No new patients will be granted from 1 May 2024 until further notice.

### **SGLT2 Inhibitors**

### ⇒SA2408 Special Authority for Subsidy

**Initial application** — (heart failure reduced ejection fraction) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### All of the following:

- 1 Patient has heart failure: and
- 2 Patient is in NYHA functional class II or III or IV; and
- 3 Either:
  - 3.1 Patient has a documented left ventricular ejection fraction (LVEF) of less than or equal to 40%; or

Subsidy (Manufacturer's Price)		Fully	Brand or Generic
 \$	Per	1	Manufacturer

continued...

- 3.2 An ECHO is not reasonably practicable, and in the opinion of the treating practitioner the patient would benefit from treatment; and
- 4 Patient is receiving concomitant optimal standard funded chronic heart failure treatment.

Initial application — (Type 2 Diabetes) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

- Either:
  - 1 Patient has previously received an initial approval for a GLP-1 agonist; or
  - 2 All of the following:
    - 2.1 Patient has type 2 diabetes; and
    - 2.2 Any of the following:
      - 2.2.1 Patient is Maori or any Pacific ethnicity\*; or
      - 2.2.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)\*; or
      - 2.2.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator\*; or
      - 2.2.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult\*; or
      - 2.2.5 Patient has diabetic kidney disease (see note b)\*; and

EMPAGLIFLOZIN - Special Authority see SA2408 on the previous page - Retail pharmacy

2.3 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of at least one blood-glucose lowering agent (e.g. metformin, vildagliptin, or insulin) for at least 3 months.

Notes: \* Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

- a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack, ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.
- b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m2 in the presence of diabetes, without alternative cause.
- c) Funded [empagliflozin / empagliflozin with metformin hydrochloride] treatment is not to be given in combination with a funded GLP-1 unless receiving (empagliflozin / empagliflozin with metformin hydrochloride] for the treatment of heart failure.

	Tab 10 mg		30 <sup>°</sup> 30	<ul><li>✓ Jardiance</li><li>✓ Jardiance</li></ul>
	PAGLIFLOZIN WITH METFORMIN HYDROCHLORIDE rmacy	- Special Authority see	SA2408	on the previous page - Retail
	Tab 5 mg with 1,000 mg metformin hydrochloride	58.56	60	✓ Jardiamet
	Tab 5 mg with 500 mg metformin hydrochloride		60	✓ Jardiamet
	Tab 12.5 mg with 1,000 mg metformin hydrochloride		60	✓ Jardiamet
*	Tab 12.5 mg with 500 mg metformin hydrochloride	58.56	60	<ul><li>Jardiamet</li></ul>

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer

# **Diabetes Management**

# **Ketone Testing**

BLOOD KETONE DIAGNOSTIC TEST STRIP - Subsidy by endorsement

- a) Not on a BSO
- b) Maximum of 20 strip per prescription
- c) Up to 10 strip available on a PSO
- d) Patient has any of the following:
  - 1) type 1 diabetes; or
  - 2) permanent neonatal diabetes: or
  - 3) undergone a pancreatectomy; or
  - 4) cystic fibrosis-related diabetes; or
  - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly.

# **Dual Blood Glucose and Blood Ketone Testing**

DUAL BLOOD GLUCOSE AND BLOOD KETONE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A dual blood glucose and blood ketone diagnostic test meter is subsidised for a patient who has:
  - 1) type 1 diabetes; or
  - 2) permanent neonatal diabetes; or
  - 3) undergone a pancreatectomy; or
  - 4) cystic fibrosis-related diabetes; or
  - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly. Only 1 meter per patient will be subsidised (no repeat prescriptions). For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Meter with 50 lancets, a lancing device and 10 blood glucose

				_
Subsidy	F	ully	Brand or	
(Manufacturer's Price)	Subsid	sed	Generic	
\$	Per	•	Manufacturer	

# **Blood Glucose Testing**

BLOOD GLUCOSE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A diagnostic blood glucose test meter is subsidised for a patient who:
  - 1) is receiving insulin or sulphonylurea therapy; or
  - 2) is pregnant with diabetes: or
  - 3) is on home TPN at risk of hypoglycaemia or hyperglycaemia; or
  - 4) has a genetic or an acquired disorder of glucose homeostasis, excluding type 1 or type 2 diabetes and metabolic syndrome.

The prescription must be endorsed accordingly. Only one CareSens meter per patient will be subsidised (no repeat prescriptions). Patients already using the CareSens N POP meter and CareSens N meter are not eligible for a new meter, unless they have:

- 1) type 1 diabetes; or
- 2) permanent neonatal diabetes: or
- 3) undergone a pancreatectomy; or
- 4) cystic fibrosis-related diabetes.

For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Note: Only 1 meter available per PSO

BLOOD GLUCOSE DIAGNOSTIC TEST STRIP - Up to 50 test available on a PSO

The number of test strips available on a prescription is restricted to 50 unless:

- Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the
  prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed; or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly; or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

Test strips	50 test OP	✓ CareSens N
		✓ CareSens PRO

# BLOOD GLUCOSE TEST STRIPS (VISUALLY IMPAIRED)

The number of test strips available on a prescription is restricted to 50 unless:

- Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the
  prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed; or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly: or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

Blood glucose test strips	33.69	50 test OP	✓ SensoCard
---------------------------	-------	------------	-------------

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

# **Insulin Syringes and Needles**

Subsidy is available for disposable insulin syringes, needles, and pen needles if prescribed on the same form as the one used for the supply of insulin or liraglutide or when prescribed for a patient and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of insulin or liraglutide.

INSULIN PEN NEEDLES - Maximum of 200 dev per prescription

IIV	octivi Livine Edeco — Maximum of 200 dev per prescription	1		
*	29 g × 12.7 mm	10.95	100	✓ B-D Micro-Fine
*	31 g × 5 mm	12.26	100	✓ B-D Micro-Fine
*	31 g × 6 mm	9.50	100	✓ Berpu
*	31 g × 8 mm		100	✓ B-D Micro-Fine
*	32 g × 4 mm		100	✓ B-D Micro-Fine
INS	SULIN SYRINGES, DISPOSABLE WITH ATTACHED NEEDLE	E – Maximum of 2	00 dev per p	orescription
*	Syringe 0.3 ml with 29 g x 12.7 mm needle	13.56	100	✓ B-D Ultra Fine
		1.36	10	
		(1.99)		B-D Ultra Fine
*	Syringe 0.3 ml with 31 g x 8 mm needle	13.56	100	B-D Ultra Fine II
	, ,	1.30	10	
		(1.99)		B-D Ultra Fine II
*	Syringe 0.5 ml with 29 g x 12.7 mm needle	13.56	100	B-D Ultra Fine
	, ,	1.36	10	
		(1.99)		B-D Ultra Fine
*	Syringe 0.5 ml with 31 g x 8 mm needle	13.56	100	B-D Ultra Fine II
	, ,	1.36	10	
		(1.99)		B-D Ultra Fine II
*	Syringe 1 ml with 29 g x 12.7 mm needle	13.56	100	B-D Ultra Fine
	, -	1.36	10	
		(1.99)		B-D Ultra Fine
*	Syringe 1 ml with 31 g × 8 mm needle	13.56	100	B-D Ultra Fine II
	•	1.36	10	
		(1.99)		B-D Ultra Fine II

# **Insulin Pumps**

INSULIN PUMP - Special Authority see SA1603 below - Retail pharmacy

- a) Maximum of 1 dev per prescription
- b) Only on a prescription
- c) Maximum of 1 insulin pump per patient each four year period.

✓ MiniMed 770G Min basal rate 0.025 U/h ......8,800.00

(MiniMed 770G Min basal rate 0.025 U/h to be delisted 1 January 2025)

### ⇒SA1603 Special Authority for Subsidy

Initial application — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has permanent neonatal diabetes; and
- 2 A MDI regimen trial is inappropriate; and
- 3 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 4 Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and

Subsidy	0.1	Fully	Brand or	
(Manufacturer's Price) \$	Per Sul	bsidised •	Generic Manufacturer	

continued...

- 5 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 6 Fither:
  - 6.1 Applicant is a relevant specialist; or
    - 6.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- 2 Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician; and
- 3 It has been at least 4 years since the last insulin pump received by the patient or, in the case of patients qualifying under previous pump therapy for the initial application; the pump is due for replacement; and
- 4 Either:
  - 4.1 Applicant is a relevant specialist; or
  - 4.2 Applicant is a nurse practitioner working within their vocational scope.

**Initial application** — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person); and
- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol: and
- 7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 8 Fither:
  - 8.1 Applicant is a relevant specialist: or
  - 8.2 Applicant is a nurse practitioner working within their vocational scope.

**Renewal** — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events; and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline; and
- 3 Either:
  - 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
  - 3.2 The pump is due for replacement; and
- 4 Fither:
  - 4.1 Applicant is a relevant specialist; or
  - 4.2 Applicant is a nurse practitioner working within their vocational scope.

**Initial application** — **(HbA1c)** only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and

<del></del>			
	Subsidy	Fully	Brand or
	(Manufacturer's Price)	Subsidised	Generic
	\$	Per 🗸	Manufacturer

continued...

- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1: and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol: and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Fither:
  - 9.1 Applicant is a relevant specialist; or
  - 9.2 Applicant is a nurse practitioner working within their vocational scope.

**Renewal — (HbA1c)** only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol: and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline; and
- 3 Fither
  - 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
  - 3.2 The pump is due for replacement; and
- 4 Either:
  - 4.1 Applicant is a relevant specialist; or
  - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump treatment; and
- 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating pump therapy; and
- 4 The patient is continuing to derive benefit from pump therapy; and
- 5 The patient had achieved and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and
- 6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and
- 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and
- 8 Either:
  - 8.1 It has been at least 4 years since the last insulin pump was received by the patient; or
  - 8.2 The pump is due for replacement; and
- 9 Either:
  - 9.1 Applicant is a relevant specialist; or
  - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

Subsidy		ully	Brand or
(Manufacturer's Price)	Subsid	ised	Generic
\$	Per	✓	Manufacturer

#### continued...

- 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and
- 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from the time of commencing pump treatment; and
- 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and
- 4 Either:
  - 4.1 It has been at least 4 years since the last insulin pump was received by the patient; or
  - 4.2 The pump is due for replacement; and
- 5 Either:
  - 5.1 Applicant is a relevant specialist; or
  - 5.2 Applicant is a nurse practitioner working within their vocational scope.

INSULIN PUMP WITH ALGORITHM - Special Authority see SA2367 below - Retail pharmacy

- a) Maximum of 1 dev per prescription
- b) Only on a prescription
- c) Maximum of 1 insulin pump per patient each four year period.

  Min basal rate 0.02 U/h ......8,970.00

- ✓ mylife YpsoPump with CamAPS FX
- ✓ Tandem t:slim
  X2 with Basal-IQ
- ✓ Tandem t:slim

  X2 with Control-IQ

### ⇒SA2367 Special Authority for Subsidy

Initial application — (type 1 diabetes) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 The patient has type 1 diabetes; or
  - 1.2 The patient has permanent neonatal diabetes or specific monogenic diabetes subtypes with insulin deficiency, considered by the treating endocrinologist as likely to benefit; or
  - 1.3 The patient has Type 3c diabetes considered by the treating endocrinologist as likely to benefit (Type 3c diabetes includes insulin deficiency due to pancreatectomy, insulin deficiency secondary to cystic fibrosis or pancreatitis); or
  - 1.4 The patient has atypical inherited forms of diabetes; and
- 2 Patient has been evaluated by a diabetes multidisciplinary team for their suitability for insulin pump therapy; and
- 3 In the opinion of the treating relevant practitioner the patient would benefit from an Automated Insulin Delivery (AID) system.

Renewal — (type 1 diabetes) from any relevant practitioner. Approvals valid for 6 months where the patient is continuing to derive benefit according to the treatment plan agreed at induction.

# **Insulin Pump Consumables**

### ⇒SA2380 Special Authority for Subsidy

**Initial application — (type 1 diabetes)** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 The patient has type 1 diabetes; or

	Subsidy (Manufacturer's Price) \$	Sub:	Fully sidised	Brand or Generic Manufacturer
continued				
1.2 The patient has permanent neonatal diabetes of		betes sub	types w	vith insulin deficiency,
considered by the treating endocrinologist as lil			ما مدنام،	anafit (Ta Oa diabata
1.3 The patient has Type 3c diabetes considered b includes insulin deficiency due to pancreatecto				
1.4 The patient has atypical inherited forms of diab		condary ic	Cystic	iibiosis oi paricieatitis),
Patient has been evaluated by a diabetes multidiscipling	*	ility for ins	ulin nu	mn therany: and
3 In the opinion of the treating relevant practitioner the p	,	,		1 1 7 /
system.				
Renewal — (type 1 diabetes) from any relevant practitioner.	Approvals valid for 2 ye	ears where	e the pa	atient is continuing to
derive benefit according to the treatment plan agreed at induc	tion.		·	· ·
INSULIN PUMP CARTRIDGE - Special Authority see SA238	0 on the previous page -	- Retail pl	narmacy	У
a) Maximum of 5 sets per prescription				
b) Only on a prescription				
c) Maximum of 19 packs of cartridge sets will be funded	per year.			
* Cartridge 300 U, t:lock × 10		1 OP		andem Cartridge
INSULIN PUMP INFUSION SET (STEEL CANNULA) - Spec	ial Authority see SA2380	on the pi	evious	page - Retail pharmacy
a) Maximum of 5 set per prescription				
b) Only on a prescription				
c) Maximum of 19 infusion sets will be funded per year.	100.00	1 OD		Similard Come T
* 6 mm steel needle; 60 cm tubing × 10	130.00	1 OP	• 1	/liniMed Sure-T MMT-864A
* 6 mm steel needle; 80 cm tubing × 10	130.00	1 OP	✓ N	/liniMed Sure-T MMT-866A
* 8 mm steel needle; 60 cm tubing × 10	130.00	1 OP	✓ N	/liniMed Sure-T MMT-874A
* 8 mm steel needle; 80 cm tubing × 10	130.00	1 OP	<b>✓</b> N	MiniMed Sure-T

(MiniMed Sure-T MMT-864A 6 mm steel needle; 60 cm tubing  $\times$  10 to be delisted 1 October 2026) (MiniMed Sure-T MMT-866A 6 mm steel needle; 80 cm tubing  $\times$  10 to be delisted 1 October 2026) (MiniMed Sure-T MMT-874A 8 mm steel needle; 60 cm tubing  $\times$  10 to be delisted 1 October 2026) (MiniMed Sure-T MMT-876A 8 mm steel needle; 80 cm tubing  $\times$  10 to be delisted 1 October 2026)

MMT-876A

Subsidy		Fully	Brand or	
(Manufacturer's Price) \$	Per	Subsidised <	Generic Manufacturer	
	_			=

INSULIN PUMP INFUSION SET (STEEL CANNULA, STRAIGHT INSERTION) - Special Authority see SA2380 on page 19 - Retail pharmacy

a) Maximum of 5 sets per prescription

	b) Only on a prescription			
	c) Maximum of 19 infusion sets will be funded per year.			
	5.5 mm steel cannula; straight insertion; 45 cm line $\times$ 10 with			
	10 needles	136.00	1 OP	mylife Orbit micro
	5.5 mm steel needle; straight insertion; 60 cm line $\times$ 10 with			
	10 needles	136.00	1 OP	mylife Orbit micro
*	5.5 mm steel needle; straight insertion; 80 cm line $\times$ 10 with			
	10 needles	136.00	1 OP	mylife Orbit micro
*	8.5 mm steel needle; straight insertion; 60 cm line × 10 with			
	10 needles	136.00	1 OP	mylife Orbit micro
*	8.5 mm steel needle; straight insertion; 80 cm line × 10 with			
	10 needles	136.00	1 OP	mylife Orbit micro
*	6 mm steel cannula; straight insertion; 80 cm line × 10 with			
	10 needles	182.00	1 OP	✓ TruSteel
*	8 mm steel cannula; straight insertion; 80 cm line $\times$ 10 with			
	10 needles	182.00	1 OP	✓ TruSteel
*	6 mm steel cannula; straight insertion; 60 cm line × 10 with			
	10 needles	182.00	1 OP	✓ TruSteel
*	8 mm steel cannula; straight insertion; 60 cm line $\times$ 10 with			
	10 needles	182.00	1 OP	✓ TruSteel

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

INSULIN PUMP INFUSION SET (TEFLON CANNULA) - Special Authority see SA2380 on page 19 - Retail pharmacy

- a) Maximum of 5 set per prescription
- b) Only on a prescription

	c) Maximum of 19 infusion sets will be funded per year.		
*	13 mm teflon needle, 60 cm tubing × 10130.00	1 OP	<ul><li>MiniMed Silhouette MMT-381A</li></ul>
*	17 mm teflon needle, 110 cm tubing × 10130.00	1 OP	<ul><li>MiniMed Silhouette MMT-377A</li></ul>
*	17 mm teflon needle, 60 cm tubing × 10130.00	1 OP	<ul><li>MiniMed Silhouette MMT-378A</li></ul>
*	6 mm teflon needle, 110 cm tubing x 10130.00	1 OP	MiniMed Quick-Set MMT-398A
*	6 mm teflon needle, 45 cm blue tubing × 10130.00	1 OP	✓ MiniMed Mio MMT-941A
*	6 mm teflon needle, 45 cm pink tubing x 10130.00	1 OP	✓ MiniMed Mio MMT-921A
*	6 mm teflon needle, 60 cm blue tubing × 10130.00	1 OP	✓ MiniMed Mio MMT-943A
*	6 mm teflon needle, 60 cm pink tubing x 10130.00	1 OP	✓ MiniMed Mio MMT-923A
*	6 mm teflon needle, 60 cm tubing × 10	1 OP	<ul><li>MiniMed Quick-Set MMT-399A</li></ul>
*	6 mm teflon needle, 80 cm blue tubing	1 OP	✓ MiniMed Mio MMT-945A
*	6 mm teflon needle, 80 cm clear tubing × 10130.00	1 OP	✓ MiniMed Mio MMT-965A
*	6 mm teflon needle, 80 cm pink tubing × 10130.00	1 OP	✓ MiniMed Mio MMT-925A
*	9 mm teflon needle, 110 cm tubing × 10	1 OP	<ul><li>MiniMed Quick-Set MMT-396A</li></ul>
*	9 mm teflon needle, 60 cm tubing × 10	1 OP	✓ MiniMed Quick-Set MMT-397A
*	9 mm teflon needle, 80 cm clear tubing × 10130.00	1 OP	✓ MiniMed Mio MMT-975A

(MiniMed Silhouette MMT-381A 13 mm teflon needle, 60 cm tubing × 10 to be delisted 1 October 2026) (MiniMed Silhouette MMT-377A 17 mm teflon needle, 110 cm tubing × 10 to be delisted 1 October 2026) (MiniMed Silhouette MMT-378A 17 mm teflon needle, 60 cm tubing × 10 to be delisted 1 October 2026) (MiniMed Quick-Set MMT-398A 6 mm teflon needle, 110 cm tubing × 10 to be delisted 1 October 2026) (MiniMed Mio MMT-941A 6 mm teflon needle, 45 cm blue tubing × 10 to be delisted 1 October 2026) (MiniMed Mio MMT-921A 6 mm teflon needle, 45 cm pink tubing × 10 to be delisted 1 October 2026) (MiniMed Mio MMT-923A 6 mm teflon needle, 60 cm blue tubing × 10 to be delisted 1 October 2026) (MiniMed Mio MMT-943A 6 mm teflon needle, 60 cm bink tubing × 10 to be delisted 1 October 2026) (MiniMed Quick-Set MMT-399A 6 mm teflon needle, 60 cm bink tubing × 10 to be delisted 1 October 2026) (MiniMed Mio MMT-945A 6 mm teflon needle, 80 cm blue tubing to be delisted 1 October 2026) (MiniMed Mio MMT-965A 6 mm teflon needle, 80 cm clear tubing × 10 to be delisted 1 October 2026) (MiniMed Mio MMT-95A 6 mm teflon needle, 80 cm pink tubing × 10 to be delisted 1 October 2026) (MiniMed Quick-Set MMT-396A 9 mm teflon needle, 110 cm tubing × 10 to be delisted 1 October 2026) (MiniMed Quick-Set MMT-397A 9 mm teflon needle, 60 cm tubing × 10 to be delisted 1 October 2026) (MiniMed Quick-Set MMT-397A 9 mm teflon needle, 60 cm tubing × 10 to be delisted 1 October 2026) (MiniMed Mio MMT-975A 9 mm teflon needle, 60 cm tubing × 10 to be delisted 1 October 2026)

		Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
SA	SULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE IN 2380 on page 19 – Retail pharmacy  a) Maximum of 5 sets per prescription  b) Only on a prescription  c) Maximum of 19 infusion sets will be funded per year.		SERTI	ON DEVIC	E) – Special Authority see
	13 mm teflon cannula; angle insertion; insertion device; 110 c	182.00	1 OP	✓ A	utoSoft 30
	13 mm teflon cannula; angle insertion; insertion device; 60 cn line × 10 with 10 needles	182.00	1 OP		utoSoft 30
see	SULIN PUMP INFUSION SET (TEFLON CANNULA, FLEXIBLE 2 SA2380 on page 19 – Retail pharmacy  a) Maximum of 5 set per prescription b) Only on a prescription c) Maximum of 19 infusion sets will be funded per year.		INSEF	RTION DEV	(ICE) – Special Authority
	6 mm teflon cannula; flexible insertion; insertion device; 46 cr line × 10 with 10 needles	157.00	1 OP	<b>✓</b> m	nylife Inset soft
	6 mm teflon cannula; flexible insertion; insertion device; 60 cr line with integrated inserter x 10 with 10 needles	157.00	1 OP	<b>✓</b> m	nylife Inset soft
*	line × 10 with 10 needles	157.00	1 OP	<b>✓</b> m	nylife Inset soft
*	line × 10 with 10 needles	157.00	1 OP	<b>✓</b> m	nylife Inset soft
*	9 mm teflon cannula; flexible insertion; insertion device; 80 cr line $\times$ 10 with 10 needles		1 OP	✓ n	nylife Inset soft
see	SULIN PUMP INFUSION SET (TEFLON CANNULA, STRAIGH SA2380 on page 19 – Retail pharmacy  a) Maximum of 5 sets per prescription b) Only on a prescription c) Maximum of 19 infusion sets will be funded per year. 6 mm teflon cannula; straight insertion; insertion device;	T INSERTION WITH	inse	RTION DE	VICE) – Special Authority
	110 cm line × 10 with 10 needles		1 OP	✓ A	utoSoft 90
	line x 10 with 10 needles		1 OP	✓ A	utoSoft 90
	110 cm line × 10 with 10 needles		1 OP	✓ A	utoSoft 90
*	9 mm teflon cannula; straight insertion; insertion device; 60 cr line × 10 with 10 needles		1 OP	✓ A	utoSoft 90
Re	SULIN PUMP INFUSION SET (TEFLON CANNULA, VARIABLI tail pharmacy  a) Maximum of 5 set per prescription b) Only on a prescription c) Maximum of 19 infusion sets will be funded per year.  13 mm teflon cannula; variable insertion; 60 cm line × 10 with 10 needles	, ,	pecial A	ŕ	e SA2380 on page 19 –

	Subsidy (Manufacturer's Price)		Fully sidised	Brand or Generic	
	\$	Per		Manufacturer	
INSULIN PUMP RESERVOIR - Special Authority see SA2380 on page 19 - Retail pharmacy					

- a) Maximum of 9 sets per prescription
- b) Only on a prescription
- c) Maximum of 36 packs of resevoir sets will be funded per year

*	10 x 1.6 ml glass reservoir for YpsoPump	50.00	1 OP	✓ mylife YpsoPump  Reservoir
*	10 × luer lock conversion cartridges 1.8 ml for Paradigm pumps	50.00	1 OP	✓ ADR Cartridge 1.8

1 OP

✓ MiniMed 3.0 Reservoir MMT-332A

(ADR Cartridge 1.8 10 × luer lock conversion cartridges 1.8 ml for Paradigm pumps to be delisted 1 October 2026) (MiniMed 3.0 Reservoir MMT-332A Cartridge for 7 series pump: 3.0 ml x 10 to be delisted 1 October 2026)

### Continuous Glucose Monitor

CONTINUOUS GLUCOSE MONITOR (INTEROPERABLE) - Special Authority see \$A2371 below - Retail pharmacy

- a) Brand switch fee payable (Pharmacode 2692147) see page 273 for details
- b) Only on a prescription

\* Sensor (9) and transmitter (Dexcom G6) - Maximum of 1 dev 1 OP ✓ Dexcom G6 per prescription......990.00 Maximum of 5 dev will be funded per year. Sensor (Dexcom G7) – Maximum of 9 dev per prescription...........110.00 ✓ Dexcom G7 Maximum of 40 dev will be funded per year. Sensor (Freestyle Libre 3 Plus) – Maximum of 6 dev per 1 ✓ Freestyle Libre 3 Plus

Maximum of 28 dev will be funded per year.

#### ⇒SA2371 Special Authority for Subsidy

Initial application — (type 1 diabetes) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 The patient has type 1 diabetes; or
  - 1.2 The patient has permanent neonatal diabetes or specific monogenic diabetes subtypes with insulin deficiency, considered by the treating endocrinologist or relevant secondary health care professional as practicable, as likely to
  - 1.3 The patient has Type 3c diabetes considered by the treating endocrinologist or relevant secondary health care professional as practicable, as likely to benefit (Type 3c diabetes includes insulin deficiency due to pancreatectomy, insulin deficiency secondary to cystic fibrosis or pancreatitis); or
  - 1.4 The patient has atypical inherited forms of diabetes; and
- 2 In the opinion of the treating relevant practitioner the patient would benefit from an Automated Insulin Delivery (AID) system.

Renewal — (type 1 diabetes) from any relevant practitioner. Approvals valid for 2 years where the patient is continuing to derive benefit according to the treatment plan agreed at induction.

CONTINUOUS GLUCOSE MONITOR (STANDALONE) - Special Authority see SA2370 on the next page - Retail pharmacy

- a) Brand switch fee payable (Pharmacode 2692139) see page 273 for details
- b) Only on a prescription
- \* Sensor (Dexcom ONE+) Maximum of 9 dev per prescription ...... 81.00 ✓ Dexcom ONE+ Maximum of 40 dev will be funded per year.
- \* Sensor (Freestyle Libre 2) Maximum of 7 dev per prescription.....92.83 1 ✓ Freestyle Libre 2 Maximum of 29 dev will be funded per year.

Subsidy (Manufacturer's Price)	Sı	Fully ubsidised	Brand or Generic
 \$	Per	✓	Manufacturer

### ⇒SA2370 Special Authority for Subsidy

**Initial application** — (type 1 diabetes) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 The patient has type 1 diabetes; or
- 2 The patient has permanent neonatal diabetes or specific monogenic diabetes subtypes with insulin deficiency, considered by the treating endocrinologist or relevant secondary health care professional as practicable, as likely to benefit; or
- 3 The patient has Type 3c diabetes considered by the treating endocrinologist or relevant secondary health care professional as practicable, as likely to benefit (Type 3c diabetes includes insulin deficiency due to pancreatectomy, insulin deficiency secondary to cystic fibrosis or pancreatitis); or
- 4 The patient has atypical inherited forms of diabetes.

Renewal — (type 1 diabetes) from any relevant practitioner. Approvals valid for 2 years where the patient is continuing to derive benefit according to the treatment plan agreed at induction.

# **Digestives Including Enzymes**

#### PANCREATIC ENZYME

Cap pancreatin 150 mg (amylase 8,000 Ph Eur U, lipase			
10,000 Ph Eur U, total protease 600 Ph Eur U)	34.93	100	✓ Creon 10000
Cap pancreatin 300 mg (amylase 18,000 Ph Eur U, lipase			
25,000 Ph Eur U, total protease 1,000 Ph Eur U)	94.38	100	✓ Creon 25000
Modified release granules pancreatin 60.12 mg (amylase			
3,600 Ph Eur U, lipase 5,000 Ph Eur U, protease 200 Ph			
Eur U)	34.93	20 g OP	<ul><li>Creon Micro</li></ul>
URSODEOXYCHOLIC ACID - Special Authority see SA1739 below	/ – Retail pha	ırmacy	
Cap 250 mg	33.95	100	✓ <u>Ursosan</u>

### ⇒SA1739 Special Authority for Subsidy

Initial application — (Alagille syndrome or progressive familial intrahepatic cholestasis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 Patient has been diagnosed with Alagille syndrome; or
- 2 Patient has progressive familial intrahepatic cholestasis.

Initial application — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic severe drug induced cholestatic liver injury; and
- 2 Cholestatic liver injury not due to Total Parenteral Nutrition (TPN) use in adults; and
- 3 Treatment with ursodeoxycholic acid may prevent hospital admission or reduce duration of stay.

Initial application — (Primary biliary cholangitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Primary biliary cholangitis confirmed by antimitochondrial antibody titre (AMA) > 1:80, and raised cholestatic liver enzymes with or without raised serum IgM or, if AMA is negative, by liver biopsy; and
- 2 Patient not requiring a liver transplant (bilirubin > 100 umol/l; decompensated cirrhosis).

**Initial application** — (**Pregnancy**) from any relevant practitioner. Approvals valid for 6 months where the patient diagnosed with cholestasis of pregnancy.

**Initial application** — (Haematological Transplant) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

continued...

Both:

- 1 Patient at risk of veno-occlusive disease or has hepatic impairment and is undergoing conditioning treatment prior to allogenic stem cell or bone marrow transplantation; and
- 2 Treatment for up to 13 weeks.

Initial application — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Paediatric patient has developed abnormal liver function as indicated on testing which is likely to be induced by Total Parenteral Nutrition (TPN); and
- 2 Liver function has not improved with modifying the TPN composition.

Renewal — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 6 months where the patient continues to benefit from treatment.

Renewal — (Pregnancy/Primary biliary cholangitis) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months where the paediatric patient continues to require TPN and who is benefiting from treatment, defined as a sustained improvement in bilirubin levels

### Laxatives

# **Bulk-forming Agents**

* Powder for oral soin20.00	500 g OP	▼ Konsyl-D	
Faecal Softeners			
DOCUSATE SODIUM - Only on a prescription         * Tab 50 mg       3.20         * Tab 120 mg       4.98	100 100	✓ <u>Coloxyl</u> ✓ <u>Coloxyl</u>	
DOCUSATE SODIUM WITH SENNOSIDES Tab 50 mg with sennosides 8 mg3.50	200	✓ <u>Laxsol</u>	
POLOXAMER – Only on a prescription  Not funded for use in the ear.  * Oral drops 10%4.17	30 ml OP	✓ Coloxyl	

20 00

E00 ~ OD

# **Opioid Receptor Antagonists - Peripheral**

ISPAGHULA (PSYLLIUM) HUSK - Only on a prescription

METHYLNALTREXONE BROMIDE - Special Authority	see SA1691 below - Retail	pharmacy	
Inj 12 mg per 0.6 ml vial	36.00	1	✓ Relistor
, -,	246.00	7	✓ Relistor

### ⇒SA1691 Special Authority for Subsidy

Initial application — (Opioid induced constipation) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient is receiving palliative care; and
- 2 Either:
  - 2.1 Oral and rectal treatments for opioid induced constipation are ineffective; or
  - 2.2 Oral and rectal treatments for opioid induced constipation are unable to be tolerated.

	Subsidy (Manufacturer's Price \$	) Sub Per	Fully sidised	Brand or Generic Manufacturer
Osmotic Laxatives				
GLYCEROL  * Suppos 2.8/4.0 g - Only on a prescription	10.39	20	<b>√</b> <u>L</u>	.ax-suppositories Glycerol
LACTULOSE – Only on a prescription  * Oral liq 10 g per 15 ml	3.61	500 ml	<b>✓</b> <u>L</u>	.aevolac
MACROGOL 3350 WITH POTASSIUM CHLORIDE, SODIUM BIG Powder for oral soln 13.125 g with potassium chloride 46.6 m		SODIUM	CHLORI	DE
sodium bicarbonate 178.5 mg and sodium chloride 350.7		30	<b>✓</b> <u>N</u>	<u>Nolaxole</u>
SODIUM ACID PHOSPHATE – Only on a prescription Enema 16% with sodium phosphate 8%	2.50	1	<b>√</b> F	Fleet Phosphate Enema
SODIUM CITRATE WITH SODIUM LAURYL SULPHOACETATE Enema 90 mg with sodium lauryl sulphoacetate 9 mg per ml,		ription		
5 ml		50	✓ <u>N</u>	<u>/licolette</u>
Stimulant Laxatives				
BISACODYL - Only on a prescription  * Tab 5 mg  * Suppos 10 mg  Lax-Suppositories to be Principal Supply on 1 February 2	4.14	200 10	_	Bisacodyl Viatris ax-Suppositories
SENNA – Only on a prescription  * Tab, standardised	(8.21)	100	S	Senokot
	0.43 (2.06)	20	S	Senokot
SODIUM PICOSULFATE - Special Authority see SA2053 below Oral soln 7.5 mg per ml		30 ml OP	✓ [	Oulcolax SP Drop
⇒SA2053 Special Authority for Subsidy				•

### SA2053 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The patient is a child with problematic constipation despite an adequate trial of other oral pharmacotherapies including macrogol where practicable; and
- 2 The patient would otherwise require a high-volume bowel cleansing preparation or hospital admission.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

# Metabolic Disorder Agents

ALGLUCOSIDASE ALFA – Special Authority see SA1986 below – Retail pharmacy ✓ Myozyme

# ⇒SA1986 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

1 The patient is aged up to 24 months at the time of initial application and has been diagnosed with infantile Pompe disease;

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsid	dised	Generic	
\$	Per	✓	Manufacturer	

continued...

and

- 2 Any of the following:
  - 2.1 Diagnosis confirmed by documented deficiency of acid alpha-glucosidase by prenatal diagnosis using chorionic villus biopsies and/or cultured amniotic cells; or
  - 2.2 Documented deficiency of acid alpha-glucosidase, and urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides; or
  - 2.3 Documented deficiency of acid alpha-glucosidase, and documented molecular genetic testing indicating a disease-causing mutation in the acid alpha-glucosidase gene (GAA gene); or
  - 2.4 Documented urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides, and molecular genetic testing indicating a disease-causing mutation in the GAA gene; and
- 3 Patient has not required long-term invasive ventilation for respiratory failure prior to starting enzyme replacement therapy (ERT); and
- 4 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by ERT or might be reasonably expected to compromise a response to ERT; and
- 5 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks.

**Renewal** only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks; and
- 3 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 4 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by ERT; and
- 5 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT: and
- 6 There is no evidence of life threatening progression of respiratory disease as evidenced by the needed for > 14 days of invasive ventilation; and
- 7 There is no evidence of new or progressive cardiomyopathy.

ARGININE - Special Authority see SA2042 below - Retail pharm	acy		
Tab 1,000 mg	CBS	90	<ul><li>Clinicians</li></ul>
Cap 500 mg	CBS	50	✓ Solgar
Powder	CBS	400 g	✓ Biomed

### ⇒SA2042 Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to arginine supplementation.

**Renewal** only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to arginine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

# ⇒SA1987 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has a confirmed diagnosis of homocystinuria; and
- 2 Any of the following:

Subsidy	Fully	Brand or
acturer's Price)	Subsidised	Generic
 \$ Pe	er 🗸	

continued...

- 2.1 A cystathionine beta-synthase (CBS) deficiency; or
- 2.2 A 5,10-methylene-tetrahydrofolate reductase (MTHFR) deficiency; or
- 2.3 A disorder of intracellular cobalamin metabolism: and
- 3 An appropriate homocysteine level has not been achieved despite a sufficient trial of appropriate vitamin supplementation.

**Renewal** only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

COENZYME Q10 − Special Authority see SA2039 below − Retail pharmacy
Cap 120 mg......CBS 30
Cap 160 mg.....CBS 60

✓ Solgar
✓ Go Healthy

# **⇒SA2039** Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to coenzyme Q10 supplementation.

**Renewal** only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to coenzyme Q10 supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

# SA1988 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has been diagnosed with mucopolysaccharidosis VI; and
- 2 Either:
  - 2.1 Diagnosis confirmed by demonstration of N-acetyl-galactosamine-4-sulfatase (arylsulfatase B) deficiency by either enzyme activity assay in leukocytes or skin fibroblasts; or
  - 2.2 Detection of two disease causing mutations and patient has a sibling who is known to have mucopolysaccharidosis

Renewal only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 3 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by Enzyme Replacement Therapy (ERT); and
- 4 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT.

### ⇒SA1623 Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hunter Syndrome (mucopolysaccharidosis II); and
- 2 Fither:
  - 2.1 Diagnosis confirmed by demonstration of iduronate 2-sulfatase deficiency in white blood cells by either enzyme

Subsidy	Fı	ully Brand or	
(Manufacturer's F	Price) Subsidis	ed Generic	
\$	Per	✓ Manufacturer	

continued...

assav in cultured skin fibroblasts: or

- 2.2 Detection of a disease causing mutation in the iduronate 2-sulfatase gene; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with idursulfase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT); and
- 5 Idursulfase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 weeks post-HSCT) at doses no greater than 0.5 mg/kg every week.

### ⇒SA1695 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hurler Syndrome (mucopolysacchardosis I-H); and
- 2 Either:
  - 2.1 Diagnosis confirmed by demonstration of alpha-L-iduronidase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts; or
  - 2.2 Detection of two disease causing mutations in the alpha-L-iduronidase gene and patient has a sibling who is known to have Hurler syndrome; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with laronidase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT): and
- 5 Laronidase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 post-HSCT) at doses no greater than 100 units/kg every week.

LEVOCARNITINE - Special Authority see SA2040 below - Retail pharmacy Tab 500 mg ......CBS 30 ✓ Solgar Cap 250 mg.......CBS 30 ✓ Solgar ✓ Balance 60 ✓ Metabolics 300 Oral lig 1 g per 10 ml ......CBS ✓ Carnitor S29 118 ml ✓ Novitium Sugar Free S29 Oral lig 500 mg per 10 ml ......CBS ✓ Balance 300 ml

### ⇒SA2040 Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to carnitine supplementation.

**Renewal** only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to carnitine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per Manufacturer

# ⇒SA2041 Special Authority for Subsidy

**Initial application** only from a metabolic physician or neurologist. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to riboflavin supplementation.

Renewal only from a metabolic physician or neurologist. Approvals valid for 24 months for applications meeting the following criteria:

#### Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to riboflavin supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

### **⇒SA1989** Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 1 month for applications meeting the following criteria: All of the following:

- 1 Patient has phenylketonuria (PKU) and is pregnant or actively planning to become pregnant; and
- 2 Treatment with sapropterin is required to support management of PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

**Renewal** only from a metabolic physician or any relevant practitioner on the recommendation of a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria:

### All of the following:

- 1 Either:
  - 1.1 Following the initial one-month approval, the patient has demonstrated an adequate response to a 2 to 4 week trial of sapropterin with a clinically appropriate reduction in phenylalanine levels to support management of PKU during pregnancy; or
  - 1.2 On subsequent renewal applications, the patient has previously demonstrated response to treatment with sapropterin and maintained adequate phenylalanine levels to support management of PKU during pregnancy; and
- 2 Any of the following:
  - 2.1 Patient continues to be pregnant and treatment with sapropterin will not continue after delivery; or
  - 2.2 Patient is actively planning a pregnancy and this is the first renewal for treatment with sapropterin; or
  - 2.3 Treatment with sapropterin is required for a second or subsequent pregnancy to support management of their PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

#### ⇒SA1599 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder.

**Renewal** only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

SODIUM PHENYLBUTYRATE – Special Authority see SA1990 on the next page – Retail pharmacy Grans 483 mg per g......2,016.00 174 g OP ✓ Pheburane

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer

# ⇒SA1990 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder involving a deficiency of carbamylphosphate synthetase, ornithine transcarbamylase or argininosuccinate synthetase.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

TAURINE - Specia	I Authority see	SA2043 belo	w – Retail pharmacy

Cap 500 mg	CBS	50	✓ Solgar
Cap 1,000 mg		90	✓ Life Extension
Powder		300 g	✓ Life Extension

### ⇒SA2043 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected specific mitochondrial disorder that may respond taurine supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has confirmed diagnosis of a specific mitochondrial disorder which responds to taurine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

TRIENTINE - Special Authority see SA2324 below - Retail pharmacy Cap 250 mg......2,022.00 100 ✓ Trientine Waymade

# ⇒SA2324 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Wilson disease; and
- 2 Treatment with D-penicillamine has been trialled and discontinued because the person has experienced intolerable side effects or has not received sufficient benefit; and
- 3 Treatment with zinc has been trailled and discontinued because the person has experienced intolerable side effects or has not received sufficient benefit, or zinc is considered clinically inappropriate as the person has symptomatic liver disease and requires copper chelation.

### Gaucher's Disease

TALIGLUCERASE ALFA - Special Authority see SA2137 below - Retail pharmacy ✓ Elelvso 1

# ⇒SA2137 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has a diagnosis of symptomatic type 1 or type 3\* Gaucher disease confirmed by the demonstration of specific deficiency of glucocerebrosidase in leukocytes or cultured skin fibroblasts, and genotypic analysis; and
- 2 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by enzyme replacement therapy (ERT) or the disease might be reasonably expected to compromise a response to ERT; and
- 3 Any of the following:
  - 3.1 Patient has haematological complications of Gaucher disease; or
  - 3.2 Patient has skeletal complications of Gaucher disease; or
  - 3.3 Patient has significant liver dysfunction or hepatomegaly attributable to Gaucher disease; or
  - 3.4 Patient has reduced vital capacity from clinically significant or progressive pulmonary disease due to Gaucher

	ALIMENTAF	RY TRACT	T AND	METABOLISM
	Subsidy (Manufacturer's Pric \$	e) Subs Per	Fully idised	Brand or Generic Manufacturer
continued				
disease; or  3.5 Patient is a child and has experienced growth fai  6-12 month period; and	lure with significant o	decrease in p	ercentil	e linear growth over a
4 Taliglucerase alfa is to be administered at a dose no gre whole vial (200 units).	ater than 30 unit/kg	every other v	veek ro	unded to the nearest
Note: Indication marked with * is an unapproved indication <b>Renewal</b> only from a metabolic physician or any relevant practi Approvals valid for 3 years for applications meeting the followin All of the following:	tioner on the recomn g criteria:	nendation of	a metal	bolic physician.
Patient has demonstrated a symptomatic improvement a symptoms for which therapy was started; and     Patient has demonstrated a clinically objective improver				
liver and spleen size; and  Radiological (MRI) signs of bone activity performed at tw demonstrate no deterioration shown by the MRI, compa or adjusted dose; and	vo years since initiati	on of treatme	ent, and	I five yearly thereafter,
Patient has not developed another medical condition that ERT; and     Patient is adherent with regular treatment and taliglucers every other week rounded to the nearest whole vial (200	ase alfa is to be adm	·	·	·
Mouth and Throat				
Agents Used in Mouth Ulceration				
BENZYDAMINE HYDROCHLORIDE Soln 0.15% - Higher subsidy of \$22.60 per 500 ml with Endorsement	9.00	500 ml		
	(22.60)			ifflam
Additional subsidy by endorsement for a patient who h prescription is endorsed accordingly.	as oral mucositis as	a result of tre	eatment	for cancer, and the
CARMELLOSE SODIUM WITH GELATIN AND PECTIN	17.00	50 = OD		******
Paste	4.55 (7.90)	56 g OP 15 g OP		tomahesive Irabase
	1.52	5 g OP	_	
Powder	(3.60) 8.48 (10.95)	28 g OP		rabase tomahesive
TRIAMCINOLONE ACETONIDE Paste 0.1%	, ,	5 g OP	✓ <u>K</u>	enalog in Orabase

<sup>✓</sup> Fungilin 20 MICONAZOLE 40 g OP ✓ Decozol

Decozol to be Principal Supply on 1 February 2025

**Oropharyngeal Anti-infectives** 

AMPHOTERICIN B

Subsidy		Fully Brand or
(Manufacturer's I	Price) Subs	idised Generic  Manufacturer
YSTATIN		
Oral liq 100,000 u per ml2.22	24 ml OP	✓ Nilstat
Vitamins		
Vitamin B		
YDROXOCOBALAMIN		
f Inj 1 mg per ml, 1 ml ampoule - Up to 6 inj available on a PSO2.46	3	✓ Cobal-B12 529 ✓ Hydroxocobalamin Panpharma ✓ Vita-B12
4.10	5	✓ Cobalin-H \$29 ✓ Neo-Cytamen \$29 \$29
8.20	10	✓ Vitarubin Depot Injection S29
/ita-B12 Inj 1 mg per ml, 1 ml ampoule to be delisted 1 July 2025) YRIDOXINE HYDROCHLORIDE  a) No more than 100 mg per dose b) Only on a prescription		·
Tab 25 mg — No patient co-payment payable	90 500	✓ <u>Vitamin B6 25</u> ✓ Pyridoxine
F Tab 50 mg23.45	500	multichem
HIAMINE HYDROCHLORIDE - Only on a prescription		
F Tab 50 mg4.65	100	✓ <u>Thiamine multichem</u>
ITAMIN B COMPLEX	F00	√ Bulay
* Tab, strong, BPC11.25	500	✓ Bplex
Vitamin C		
SCORBIC ACID  a) No more than 100 mg per dose b) Only on a prescription  Tab 100 mg12.50	500	✓ Cvite
·	500	<u> Cvite</u>
Vitamin D		
LFACALCIDOL  € Cap 0.25 mcg26.32	100	✓ One-Alpha
€ Cap 1 mcg87.98	100	<ul><li>✓ One-Alpha S29 S29</li><li>✓ One-Alpha</li></ul>
€ Oral drops 2 mcg per ml	20 ml OP	✓ One-Alpha
ALCITRIOL		
Cap 0.25 mcg	100	✓ Calcitriol-AFT
: Cap 0.5 mcg	100	✓ Calcitriol-AFT ✓ Calcitriol-AFT S29 529
OLECALCIFEROL		•
Cap 1.25 mg (50,000 iu) — Maximum of 12 cap per prescription3.65	12 5 ml OP	✓ <u>Vit.D3</u> ✓ Clinicians
• Oral liq 188 mcg per ml (7,500 iu per ml)9.00	3 IIII UP	- Cillicialis
✓ fully subsidised ©29 Unap	proved medicine s	supplied under Section 29

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer **Multivitamin Preparations** MULTIVITAMIN RENAL - Special Authority see SA1546 below - Retail pharmacy 30 ✓ Clinicians Renal Vit. ⇒SA1546 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Fither: 1 The patient has chronic kidney disease and is receiving either peritoneal dialysis or haemodialysis; or 2 The patient has chronic kidney disease grade 5, defined as patient with an estimated glomerular filtration rate of < 15 ml/min/1.73 m<sup>2</sup> body surface area (BSA). MULTIVITAMINS - Special Authority see SA1036 below - Retail pharmacy 200 g OP ✓ Paediatric Seravit ⇒SA1036 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has inborn errors of metabolism. Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where patient has had a previous approval for multivitamins. **VITAMINS** 1.000 Mvite \* Cap (fat soluble vitamins A, D, E, K) - Special Authority see Vitabdeck ⇒SA1720 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following: 1 Patient has cystic fibrosis with pancreatic insufficiency; or 2 Patient is an infant or child with liver disease or short gut syndrome; or 3 Patient has severe malabsorption syndrome. **Minerals** Calcium **CALCIUM CARBONATE** 250 ✓ Calci-Tab 500 \* Tab eff 1.25 g (500 mg elemental) - Subsidy by endorsement......260.00 100 ✓ Calcium 500 mg Hexal S29 Subsidy by endorsement - Only when prescribed for paediatric patients (< 5 years) where calcium carbonate oral liquid is considered unsuitable. **CALCIUM GLUCONATE** 10 ✓ Max Health -Hameln S29 lodine

90

✓ NeuroTabs

POTASSIUM IODATE

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Generic
Iron				
FERROUS FUMARATE  * Tab 200 mg (65 mg elemental)  Ferro-tab to be Principal Supply on 1 February 2025	3.49	100	•	Ferro-tab
FERROUS FUMARATE WITH FOLIC ACID  * Tab 310 mg (100 mg elemental) with folic acid 350 mcg	5.98	100	•	Ferro-F-Tabs
* Tab long-acting 325 mg (105 mg elemental)  * Oral liq 30 mg (6 mg elemental) per 1 ml	9.25	30 250 m 500 m	nl 🗸	<u>Ferrograd</u> Ferro-Liquid <u>Ferodan</u>
IRON (AS FERRIC CARBOXYMALTOSE) – Special Authority so Inj 50 mg per ml, 10 ml vial		Retail <sub>I</sub> 1		Ferinject

Initial application — (Anaemia) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with anaemia; and
- 2 Any of the following:
  - 2.1 Serum ferritin level is 20 mcg/L or less; or
  - 2.2 Both:
    - 2.2.1 Serum ferritin is between 20 and 50 mcg/L; and
    - 2.2.2 C-Reactive Protein (CRP) is at least 5 mg/L; or
  - 2.3 Patient has chronic inflammatory disease with symptoms of anaemia despite normal iron levels; and
- 3 Any of the following:
  - 3.1 Oral iron treatment has proven ineffective; or
  - 3.2 Oral iron treatment has resulted in dose-limiting intolerance; or
  - 3.3 Rapid correction of anaemia is required.

Renewal — (Anaemia) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

### Both:

- 1 Patient continues to have anaemia with a serum ferritin level of 20 mcg/L, or less or between 20 and 50 mcg/L with CRP of at least 5 mg/L, or has chronic inflammatory disease with symptoms of anaemia despite normal iron levels; and
- 2 A trial (or re-trial) with oral iron is clinically inappropriate.

Initial application — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 Patient has been diagnosed with iron-deficiency anaemia; and
- 2 Any of the following:
  - 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
  - 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
  - 2.3 Patient has symptomatic heart failure, chronic kidney disease stage 3 or more or active inflammatory bowel disease and a trial of oral iron is unlikely to be effective; or
  - 2.4 Rapid correction of anaemia is required.

Renewal — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist.

# ALIMENTARY TRACT AND METABOLISM

100

✓ Zincaps

	Subsidy (Manufacturer's Price \$	) Sub Per	Fully sidised	Brand or Generic Manufacturer
continued Approvals valid for 3 months for applications meeting the followi Both:  1 Patient continues to have iron-deficiency anaemia; and 2 A re-trial with oral iron is clinically inappropriate.	ng criteria:			
IRON POLYMALTOSE  * Inj 50 mg per ml, 2 ml ampoule	34.50	5	<b>√</b> F	errosig
Magnesium				
MAGNESIUM HYDROXIDE Suspension 8%	33.60	355 ml	<b>√</b> P	Phillips Milk of Magnesia 629
MAGNESIUM SULPHATE  * Inj 2 mmol per ml, 5 ml ampoule  * Inj 2 mmol per ml, 10 ml ampoule		10 10		lartindale nresa 829
Zinc				
ZINC SULPHATE				

\* Cap 137.4 mg (50 mg elemental)......11.00

Subsidy	Fu	lly	Brand or	
(Manufacturer's Price)	Subsidis	ed	Generic	
\$	Per	/	Manufacturer	

### **Antianaemics**

### Hypoplastic and Haemolytic

### ⇒SA2266 Special Authority for Subsidy

Initial application — (chronic renal failure) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient in chronic renal failure: and
- 2 Haemoglobin is less than or equal to 100g/L; and
- 3 Any of the following:
  - 3.1 Both:
    - 3.1.1 Patient does not have diabetes mellitus; and
    - 3.1.2 Glomerular filtration rate is less than or equal to 30ml/min; or
  - 3.2 Both:
    - 3.2.1 Patient has diabetes mellitus; and
    - 3.2.2 Glomerular filtration rate is less than or equal to 45ml/min; or
  - 3.3 Patient is on haemodialysis or peritoneal dialysis.

Initial application — (myelodysplasia) from any specialist. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of myelodysplasia (MDS)\*; and
- 2 Has had symptomatic anaemia with haemoglobin < 100g/L and is red cell transfusion-dependent; and
- 3 Patient has very low, low or intermediate risk MDS based on the WHO classification based prognostic scoring system for myelodysplastic syndrome (WPSS); and
- 4 Other causes of anaemia such as B12 and folate deficiency have been excluded; and
- 5 Patient has a serum epoetin level of < 500 IU/L; and
- 6 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Note: Indication marked with \* is an unapproved indication

**Renewal** — **(chronic renal failure)** from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (myelodysplasia) from any specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient's transfusion requirement continues to be reduced with erythropoietin treatment; and
- 2 Transformation to acute myeloid leukaemia has not occurred; and
- 3 The minimum necessary dose of epoetin would be used and will not exceed 80,000 ju per week.

Note: Indication marked with \* is an unapproved indication

#### EPOETIN ALFA - Special Authority see SA2266 above - Retail pharmacy

Wastage claimable			
Inj 1,000 iu in 0.5 ml, syringe	250.00	6	Binocrit
Inj 2,000 iu in 1 ml, syringe		6	Binocrit
Inj 3,000 iu in 0.3 ml, syringe		6	Binocrit
Inj 4,000 iu in 0.4 ml, syringe		6	Binocrit
Inj 5,000 iu in 0.5 ml, syringe		6	<ul><li>Binocrit</li></ul>
Inj 6,000 iu in 0.6 ml, syringe		6	Binocrit
Inj 8,000 iu in 0.8 ml, syringe		6	Binocrit
Inj 10,000 iu in 1 ml, syringe		6	Binocrit
Inj 40,000 iu in 1 ml, syringe		1	Binocrit

	Subsidy (Manufacturer's Price) \$	Su Per	Fully bsidised	Brand or Generic Manufacturer
Megaloblastic				
FOLIC ACID  * Tab 0.8 mg	26.60	1,000		olic Acid multichem
* Tab 5 mg Oral liq 50 mcg per ml		100 5 ml OP	_	olic Acid Viatris iomed

## **Antifibrinolytics, Haemostatics and Local Sclerosants**

### EFTRENONACOG ALFA [RECOMBINANT FACTOR IX] - [Xpharm]

For patients with haemophilia B receiving prophylaxis treatment. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management group.

Treaters Group in conjunction with the National Haemor	ohilia Management gro	up.	
Inj 250 iu vial	612.50	1	Alprolix
Inj 500 iu vial	1,225.00	1	✓ Alprolix
Inj 1,000 iu vial	2,450.00	1	✓ Alprolix
Inj 2,000 iu vial	4,900.00	1	✓ Alprolix
Inj 3,000 iu vial	7,350.00	1	✓ Alprolix
Inj 4,000 iu vial	9,800.00	1	Alprolix
ELTROMBOPAG - Special Authority see SA1743 below -	Retail pharmacy		
Wastage claimable			
Tab 25 mg	1,550.00	28	✓ Revolade
Tab 50 mg	3,100.00	28	✓ Revolade

### ⇒SA1743 Special Authority for Subsidy

Initial application — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 6 weeks for applications meeting the following criteria:

#### All of the following:

- 1 Patient has had a splenectomy; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Any of the following:
  - 3.1 Patient has a platelet count of 20,000 to 30,000 platelets per microlitre and has evidence of significant mucocutaneous bleeding; or
  - 3.2 Patient has a platelet count of less than or equal to 20,000 platelets per microlitre and has evidence of active bleeding; or
  - 3.3 Patient has a platelet count of less than or equal to 10,000 platelets per microlitre.

Initial application — (idiopathic thrombocytopenic purpura - preparation for splenectomy) only from a haematologist. Approvals valid for 6 weeks where the patient requires eltrombopag treatment as preparation for splenectomy. Initial application — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist.

Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a significant and well-documented contraindication to splenectomy for clinical reasons; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Either:
  - 3.1 Patient has immune thrombocytopenic purpura\* with a platelet count of less than or equal to 20,000 platelets per microliter: or
  - 3.2 Patient has immune thrombocytopenic purpura\* with a platelet count of 20,000 to 30,000 platelets per microlitre

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidi	sed	Generic
\$	Per	1	Manufacturer

continued...

and significant mucocutaneous bleeding.

**Initial application — (severe aplastic anaemia)** only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Two immunosuppressive therapies have been trialled and failed after therapy of at least 3 months duration; and
- 2 Either:
  - 2.1 Patient has severe aplastic anaemia with a platelet count of less than or equal to 20,000 platelets per microliter; or
  - 2.2 Patient has severe aplastic anaemia with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

Renewal — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 12 months where the patient has obtained a response (see Note) from treatment during the initial approval or subsequent renewal periods and further treatment is required.

Note: Response to treatment is defined as a platelet count of > 30,000 platelets per microlitre.

Renewal — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

#### All of the following:

- 1 The patient's significant contraindication to splenectomy remains; and
- 2 The patient has obtained a response from treatment during the initial approval period; and
- 3 Patient has maintained a platelet count of at least 50,000 platelets per microlitre on treatment; and
- 4 Further treatment with eltrombopag is required to maintain response.

Renewal — (severe aplastic anaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

### Both:

- 1 The patient has obtained a response from treatment of at least 20,000 platelets per microlitre above baseline during the initial approval period; and
- 2 Platelet transfusion independence for a minimum of 8 weeks during the initial approval period.

#### EMICIZUMAB - [Xpharm] - Special Authority see \$A2272 below

Inj 30 mg in 1 ml vial	3,570.00	1	✓ Hemlibra
Inj 60 mg in 0.4 ml vial	7,138.00	1	✓ Hemlibra
Inj 105 mg in 0.7 ml vial	·	1	✓ Hemlibra
Inj 150 mg in 1 ml vial		1	✓ Hemlibra

### ⇒SA2272 Special Authority for Subsidy

Initial application — (Severe Haemophilia A with or without FVIII inhibitors) only from a haematologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has severe congenital haemophilia A with a severe bleeding phenotype (endogenous factor VIII activity less than or equal to 2%); and
- 2 Emicizumab is to be administered at a dose of no greater than 3 mg/kg weekly for 4 weeks followed by the equivalent of 1.5 mg/kg weekly.

### EPTACOG ALFA [RECOMBINANT FACTOR VIIA] - [Xpharm]

For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group. Rare Clinical Circumstances Brand of bypassing agent for > 14 days predicted use. Access to funded treatment for > 14 days predicted use is by named patient application to the Haemophilia Treaters Group, subject to access criteria.

Inj 1 mg syringe	1	✓ NovoSeven RT
Inj 2 mg syringe	1	✓ NovoSeven RT
Inj 5 mg syringe	1	✓ NovoSeven RT
Inj 8 mg syringe	1	✓ NovoSeven RT

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
 \$	Per 🗸	

#### FACTOR EIGHT INHIBITOR BYPASSING FRACTION - [Xpharm]

For patients with haemophilia. Preferred Brand of bypassing agent for > 14 days predicted use. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

- 1	a managed by the macinopinic	i ricatora di cup ili conjunctioni with the riation	ai i iacinopii	illa iviariagerrierit art
-	nj 500 U	1,315.00	1	✓ FEIBA NF
-	nį 1,000 U	2,630.00	1	✓ FEIBA NF
	• •	6.575.00	1	✓ FEIBA NF

### MOROCTOCOG ALFA [RECOMBINANT FACTOR VIII] - [Xpharm]

For patients with haemophilia. Rare Clinical Circumstances Brand of short half-life recombinant factor VIII. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group, subject to criteria.

subject to criteria.			
Inj 250 iu prefilled syringe	287.50	1	Xyntha
Inj 500 iu prefilled syringe		1	Xyntha
Inj 1,000 iu prefilled syringe		1	Xyntha
Inj 2,000 iu prefilled syringe		1	Xyntha
Inj 3,000 iu prefilled syringe		1	Xyntha

#### NONACOG GAMMA, [RECOMBINANT FACTOR IX] - [Xpharm]

For patients with naemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

Inj 500 iu vial	435.00	1	✓ RIXUBIS
Inj 1,000 iu vial		1	✓ RIXUBIS
Inj 2,000 iu vial	1,740.00	1	✓ RIXUBIS
Inj 3,000 iu vial	2,610.00	1	✓ RIXUBIS

### (RIXUBIS Inj 500 iu vial to be delisted 1 February 2025)

### OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (ADVATE) - [Xpharm]

For patients with haemophilia. Preferred Brand of short half-life recombinant factor VIII. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

Inj 250 iu vial	210.00	1	✓ Advate
lnį 500 iu vial		1	✓ Advate
Inj 1,000 iu vial		1	✓ Advate
Inj 1,500 iu vial		1	✓ Advate
Inj 2,000 iu vial	,	1	✓ Advate
Ini 3.000 iu vial		1	✓ Advate

### (Advate Inj 250 iu vial to be delisted 1 February 2025)

(Advate Inj 1,500 iu vial to be delisted 1 February 2025)

### OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (KOGENATE FS) - [Xpharm]

For patients with haemophilia. Rare Clinical Circumstances Brand of short half-life recombinant factor VIII. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group, subject to criteria

Inj 250 iu vial	237.50	1	✓ Kogenate FS
lnį 500 iu vial	475.00	1	✓ Kogenate FS
lnj 1,000 iu vial		1	✓ Kogenate FS
Inj 2,000 iu vial		1	✓ Kogenate FS
Inj 3,000 iu vial	2,850.00	1	✓ Kogenate FS

illy Brand or	Fully	/	Subsidy
	Subsidised		(Manufacturer's
✓ Manufacturer	er 🗸	Pe	\$
			CTOCOG ALFA PEGOL [RECOMBINANT FACTOR VIII] - [Xpharm]
managed by the Haemophilia	eatment is m	o funded tr	r patients with haemophilia A receiving prophylaxis treatment. Access to
• .		group.	eaters Group in conjunction with the National Haemophilia Management
✓ Adynovate	✓ ,	1	250 iu vial300.00
✓ Adynovate	✓,	1	500 iu vial600.00
✓ Adynovate	✓ ,	1	1,000 iu vial
✓ Adynovate	✓ ,	1	2,000 iu vial2,400.00
			vate Inj 250 iu vial to be delisted 1 February 2025)
			vate Inj 500 iu vial to be delisted 1 February 2025)
			M TETRADECYL SULPHATE
		5	3% 2 ml28.50
Fibro-vein		•	(73.00)
	•		EXAMIC ACID
✓ Mercury Pharma	<b>1</b>	60	
✓ Cyklokapron	-	10	b 500 mg
Cyklokapion	0 •	10	70.00
			nin K
			IIIII IX
			DMENADIONE
✓ Konakion MM	<b>✓</b>	5	2 mg per 0.2 ml - Up to 5 inj available on a PSO8.00
✓ Konakion MM	✓	5	10 mg per ml, 1 ml - Up to 5 inj available on a PSO9.21
			2 mg per 0.2 ml – Up to 5 inj available on a PSO8.00

### **Antiplatelet Agents**

ASPIRIN	990	✓ Ethics Aspirin EC
CLOPIDOGREL  * Tab 75 mg5.07	84	✓ Arrow - Clopid
DIPYRIDAMOLE  * Tab long-acting 150 mg13.93	60	✓ Pytazen SR
TICAGRELOR – Special Authority see SA1955 below – Retail pharmacy  * Tab 90 mg20.35	56	✓ Ticagrelor Sandoz

### **⇒SA1955** Special Authority for Subsidy

**Initial application** — (acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

#### Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Initial application — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

### 1 Either:

- 1.1 Patient has had a neurological stenting procedure\* in the last 60 days; or
- 1.2 Patient is about to have a neurological stenting procedure performed\*; and

(Mar	Subsidy nufacturer's Price)	Subsidis Per	ully sed	Brand or Generic Manufacturer
	Ψ	rei		Manuacturei

continued...

- 2 Either:
  - 2.1 Patient has demonstrated clopidogrel resistance using the P2Y12 (VerifyNow) assay or another appropriate platelet function assay and requires antiplatelet treatment with ticagrelor; or
  - 2.2 Either:
    - 2.2.1 Clopidogrel resistance has been demonstrated by the occurrence of a new cerebral ischemic event; or
    - 2.2.2 Clopidogrel resistance has been demonstrated by the occurrence of transient ischemic attack symptoms referable to the stent.

**Initial application** — (Percutaneous coronary intervention with stent deployment) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic\*\*.

Initial application — (Stent thrombosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has experienced cardiac stent thrombosis whilst on clopidogrel.

**Renewal — (subsequent acute coronary syndrome)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Renewal — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient is continuing to benefit from treatment; and
- 2 Treatment continues to be clinically appropriate.

**Renewal** — (**Percutaneous coronary intervention with stent deployment**) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic\*\*.

Notes: indications marked with \* are unapproved indications.

Note: \*\* Clopidogrel allergy is defined as a history of anaphylaxis, urticaria, generalised rash or asthma (in non-asthmatic patients) developing soon after clopidogrel is started and is considered unlikely to be caused by any other treatment.

Heparin and Antagonist Preparations			
ENOXAPARIN SODIUM - Special Authority see SA2152 below -	Retail pharmac	y	
Inj 20 mg in 0.2 ml syringe	21.90	10	✓ Clexane
Clexane to be Principal Supply on 1 February 2025			
Inj 40 mg in 0.4 ml syringe	29.74	10	✓ Clexane
Clexane to be Principal Supply on 1 February 2025			
Inj 60 mg in 0.6 ml syringe	42.47	10	✓ Clexane
Clexane to be Principal Supply on 1 February 2025			
Inj 80 mg in 0.8 ml syringe	56.62	10	✓ Clexane
Clexane to be Principal Supply on 1 February 2025			
Inj 100 mg in 1 ml syringe	70.91	10	✓ Clexane
Clexane to be Principal Supply on 1 February 2025			
Inj 120 mg in 0.8 ml syringe	88.11	10	Clexane Forte
Clexane Forte to be Principal Supply on 1 February 2025			
Inj 150 mg in 1 ml syringe	100.70	10	Clexane Forte
Clexane Forte to be Principal Supply on 1 February 2025			

Subsidy

(Manufacturer's Price)

Fully

Subsidised

Brand or

Generic Manufacturer

### **⇒SA2152** Special Authority for Subsidy

**Initial application** — (**Pregnancy, Malignancy or Haemodialysis**) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patients pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Initial application — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 For the short-term treatment of venous thromboembolism prior to establishing a therapeutic level of oral anti-coagulant treatment; or
- 2 For the prophylaxis and treatment of venous thromboembolism in high risk surgery; or
- 3 To enable cessation/re-establishment of existing oral anticoagulant treatment pre/post surgery; or
- 4 For the prophylaxis and treatment of venous thromboembolism in Acute Coronary Syndrome surgical intervention; or
- 5 To be used in association with cardioversion of atrial fibrillation.

Initial application — (Short-term use during treatment of COVID-19 with nirmatrelvir with ritonavir) from any relevant practitioner. Approvals valid for 2 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is receiving an anticoagulation treatment that has drug/drug interactions with ritonavir that increases risk of bleeding; and
- 2 Patient meets the Access Criteria for COVID-19 antivirals published on the Pharmac website\*; and
- 3 Other antiviral treatments for COVID-19 have been considered and are not clinically suitable options.

Renewal — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patient's pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Renewal — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month where low molecular weight heparin treatment or prophylaxis is required for a second or subsequent event (surgery, ACS, cardioversion, or prior to oral anti-coagulation).

(I	Subsidy Manufacturer's Price) \$	Per	Fully Subsidised	
HEPARIN SODIUM				
Inj 1,000 iu per ml, 10 ml vial	127.44	25	1	Pfizer S29
Inj 1,000 iu per ml, 5 ml ampoule	25.49	10	✓	Wockhardt S29
	103.70		✓	Wockhardt PSF S29
	127.44	50	✓	Pfizer
Inj 5,000 iu per ml, 5 ml vial	83.00	10	•	Heparin Sodium Panpharma
Inj 5,000 iu per ml, 1 ml	70.33	5	✓	Hospira
Inj 25,000 iu per ml, 0.2 ml	22.42	5	✓	Hospira
	42.40		✓	Heparin DBL S29
	482.20	50	1	Heparin DBL S29
HEPARINISED SALINE Inj 10 iu per ml, 5 ml	96.91	50	•	Pfizer
Oral Anticoagulants				
DABIGATRAN				
Cap 75 mg - No more than 2 cap per day	27.99	60	✓	<u>Pradaxa</u>
Cap 110 mg	27.99	60		<u>Pradaxa</u>
Cap 150 mg	27.99	60	•	<u>Pradaxa</u>
RIVAROXABAN				
Tab 10 mg - No more than 1 tab per day		30		<u>Xarelto</u>
Tab 15 mg - Up to 14 tab available on a PSO	14.56	28		Xarelto
Tab 20 mg	14.56	28	•	Xarelto
WARFARIN SODIUM				
Note: Marevan and Coumadin are not interchangeable.				
* Tab 1 mg	3.46	50	✓	Coumadin
	7.50	100		Marevan
<b>★</b> Tab 2 mg		50		Coumadin
* Tab 3 mg		100		Marevan
* Tab 5 mg		50		Coumadin
	13.50	100	/	Marevan

FILGRASTIM - Special Authority see SA1259 below - Retail pl	harmacy		
Inj 300 mcg per 0.5 ml prefilled syringe	86.60	10	✓ Nivestim
Inj 480 mcg per 0.5 ml prefilled syringe	133.72	10	✓ <u>Nivestim</u>

### ⇒SA1259 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 20%\*); or
- 2 Peripheral blood stem cell mobilisation in patients undergoing haematological transplantation; or
- 3 Peripheral blood stem cell mobilisation or bone marrow donation from healthy donors for transplantation; or
- 4 Treatment of severe chronic neutropenia (ANC < 0.5 ×10<sup>9</sup>/L); or
- 5 Treatment of drug-induced prolonged neutropenia (ANC < 0.5 ×10<sup>9</sup>/L).

Note: \*Febrile neutropenia risk greater than or equal to 20% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

	Subsidy (Manufacturer's Price) \$	Sul Per	Fully bsidised	Brand or Generic Manufacturer	
PEGFILGRASTIM - Special Authority see SA1912 below - Reta	ail pharmacy				
Inj 6 mg per 0.6 ml syringe	65.00	1	✓ Zie	<u>extenzo</u>	
			✓ Zie	extenzo AU S29	

### ⇒SA1912 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where used for prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 5%\*). Note: \*Febrile neutropenia risk greater than or equal to 5% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

# Fluids and Electrolytes

### **Intravenous Administration**

GLUCOSE [DEXTROSE]		
* Inj 50%, 10 ml ampoule – Up to 5 inj available on a PSO34.75	5	✓ Biomed
* Inj 50%, 90 ml bottle – Up to 5 inj available on a PSO17.50	1	✓ Biomed
POTASSIUM CHLORIDE		
* Inj 75 mg per ml, 10 ml65.00	50	✓ Juno
•		✓ LumaCina
		✓ Pfizer S29
SODIUM BICARBONATE		
Inj 8.4%, 50 ml23.52	1	✓ Biomed
a) Up to 5 inj available on a PSO		
b) Not in combination		
lnj 8.4%, 100 ml24.10	1	✓ Biomed
a) Up to 5 inj available on a PSO		
b) Not in combination		
SODIUM CHLORIDE		
Not final address on a second dress. Not final address as built as were assessed and		

Not funded for use as a nasal drop. Not funded for nebuliser use except when used in conjunction with an antibiotic intended for nebuliser use.

	1.58	1,000 ml	✓ Baxter
Only if prescribed on a prescription for renal dialysis, mat	ernity or post-nat	al care in the	e home of the patient, or on a PSO
for emergency use. (500 ml and 1,000 ml packs)			
Inj 23.4% (4 mmol/ml), 20 ml ampoule	38.25	5	✓ Biomed
For Sodium chloride oral liquid formulation refer Standard	d Formulae, <mark>page</mark>	276	

Inj 23.4% (4 mmol/ml), 20 ml ampoule	38.25	5	Biomed
For Sodium chloride oral liquid formulation refer Standa	ard Formulae, page	276	
Inj 0.9%, 5 ml ampoule - Up to 5 inj available on a PSO	4.00	20	✓ Fresenius Kabi
Inj 0.9%, 10 ml ampoule - Up to 5 inj available on a PSO	5.25	50	✓ Fresenius Kabi
Inj 0.9%, 20 ml ampoule	5.00	20	✓ Fresenius Kabi
TOTAL PARENTERAL NUTRITION (TPN)			
Infusion	CBS	1 OP	✓ TPN

500 ml

✓ Baxter

Subsidy (Manufacturer's Price)	Subs	Fully idised	Brand or Generic	
\$	Per	/	Manufacturer	

### WATER

- On a prescription or Practitioner's Supply Order only when on the same form as an injection listed in the Pharmaceutical Schedule requiring a solvent or diluent; or
- 2) On a bulk supply order; or
- 3) When used in the extemporaneous compounding of eye drops; or
- 4) When used for the dilution of sodium chloride soln 7% for cystic fibrosis patients only.

Inj 10 ml ampoule – Up to 5 inj available on a PSO7.60 Inj 20 ml ampoule – Up to 5 inj available on a PSO5.00	50 20	<ul><li>✓ Multichem</li><li>✓ Fresenius Kabi</li></ul>
Oral Administration		
CALCIUM POLYSTYRENE SULPHONATE  Powder	300 g OP	✓ Calcium Resonium
Powder for oral soln — Up to 5 sach available on a PSO9.53	50	✓ <u>Electral</u>
COMPOUND ELECTROLYTES WITH GLUCOSE [DEXTROSE] Soln with electrolytes	1,000 ml OP	✓ <u>Hydralyte -</u> Lemonade
PHOSPHORUS		
Tab eff 500 mg (16 mmol)82.50	100	Phosphate Phebra
POTASSIUM CHLORIDE		
* Tab eff 548 mg (14 m eq) with chloride 285 mg (8 m eq)	60	Chlorvescent
* Tab long-acting 600 mg (8 mmol)15.35	200	✓ Span-K
SODIUM BICARBONATE		
Cap 840 mg8.52	100	<ul><li>✓ Sodibic</li><li>✓ Sodibic</li></ul>
SODIUM POLYSTYRENE SULPHONATE		
Powder84.65	454 g OP	✓ Resonium-A

	Subsidy		Fully	
	(Manufacturer's Price \$	e) Per	Subsidised	
	Ψ	rei		Manuacturer
Alpha-Adrenoceptor Blockers				
Alpha Adrenoceptor Blockers				
DOXAZOSIN	17.05	E00	./	Doxazosin Clinect
* Tab 2 mg * Tab 4 mg		500 500		Doxazosin Clinect
· ·	20.34	500	•	DOXAZOSIII CIIIIECI
PHENOXYBENZAMINE HYDROCHLORIDE			_	
* Cap 10 mg		30	_	BNM \$29
	216.67	100	•	Dibenzyline S29
PRAZOSIN				
* Tab 1 mg	5.53	100	•	Arrotex-Prazosin
				S29 S29
	9.98		1	Minipress S29
* Tab 2 mg	7.00	100	1	Arrotex-Prazosin
				S29 S29
	13.29		1	Minipress S29
* Tab 5 mg	11.70	100		Arrotex-Prazosin
Ç				S29 S29
	22.00		1	Minipress S29
* Cap 1 mg		100		Prazosin Mylan S29
* Cap 2 mg		100		Prazosin Mylan S29
* Cap 5 mg		100		Prazosin Mylan S29
ж оар э під	23.32	100	•	Frazosiii wylaii 323
Agents Affecting the Renin-Angiotensin Syste	m			
- Agome - mooning mo Homin - migrotonism o you	••			
ACE Inhibitors				
CAPTOPRIL				
* Oral liq 5 mg per ml	86.00 1	00 ml (	OP 🗸	DP-Captopril
Oral liquid restricted to children under 12 years of age.				
CILAZAPRIL - Subsidy by endorsement				
Subsidy by endorsement – Subsidised for patients who were				
endorsed accordingly. Pharmacists may annotate the pres	cription as endorsed	where i	there exist	ts a record of prior
dispensing of cilazapril.	0.00		,	
* Tab 0.5 mg		90		Zapril
* Tab 2.5 mg		90		Zapril
Tab 5 mg	10.05	90	•	Zapril
ENALAPRIL MALEATE			_	
* Tab 5 mg		90		Acetec
* Tab 10 mg		90	/	Acetec
* Tab 20 mg	2.35	90	•	Acetec
LISINOPRIL				
* Tab 5 mg	11.07	90		Ethics Lisinopril
				Teva Lisinopril
* Tab 10 mg	11.67	90		Ethics Lisinopril
				Teva Lisinopril
* Tab 20 mg	14.69	90		Ethics Lisinopril
			•	Teva Lisinopril

	Subsidy		Fully	
	(Manufacturer's Price)	Per	Subsidised	I Generic Manufacturer
	\$	Per		Manutacturer
ERINDOPRIL			_	
F Tab 2 mg		30		Coversyl
Fab 4 mg		30		Coversyl
Fab 8 mg	3.94	30	•	Coversyl
UINAPRIL				
F Tab 5 mg	10.24	90	1	Arrow-Quinapril 5
Arrow-Quinapril 5 to be Principal Supply on 1 March 2	025			•
F Tab 10 mg	12.51	90	1	Arrow-Quinapril 10
Arrow-Quinapril 10 to be Principal Supply on 1 March	2025			•
F Tab 20 mg	14.83	90	1	Arrow-Quinapril 20
Arrow-Quinapril 20 to be Principal Supply on 1 March	2025			•
AMIPRIL				
Cap 1.25 mg	17.25	90	1	Tryzan
Tryzan to be Principal Supply on 1 February 2025		00	•	11 y 2 u 11
Cap 2.5 mg	16 50	90	1	Tryzan
Tryzan to be Principal Supply on 1 February 2025	10.00	50	•	11 y 2 a 11
Cap 5 mg	16.88	90	1	Tryzan
Tryzan to be Principal Supply on 1 February 2025		50	•	11 y 2 a 11
Cap 10 mg	17.63	90	1	Tryzan
Tryzan to be Principal Supply on 1 February 2025		50	•	11 y 2 a 11
Angiotensin II Antagonists  ANDESARTAN CILEXETIL				
F Tab 4 mg	2.68	90	✓	Candestar
Candestar to be Principal Supply on 1 February 2025				
F Tab 8 mg	2.67	90	1	Candestar
Candestar to be Principal Supply on 1 February 2025				
Fab 16 mg	4.22	90	✓	Candestar
Candestar to be Principal Supply on 1 February 2025				
F Tab 32 mg	5.24	90	1	Candestar
Candestar to be Principal Supply on 1 February 2025				
OSARTAN POTASSIUM				
Tab 12.5 mg	2 00	84	1	Losartan Actavis
: Tab 25 mg		84		Losartan Actavis
Tab 50 mg		84		Losartan Actavis
Tab 100 mg		84		Losartan Actavis
<u> </u>		-		
Angiotensin II Antagonists with Diuretics	_			
ANDESARTAN CILEXETIL WITH HYDROCHLOROTHIAZID			_	
Tab 16 mg with hydrochlorothiazide 12.5 mg	4.10	30	•	APO-Candesartan HCTZ 16/12.5
Tab 32 mg with hydrochlorothiazide 12.5 mg	5 25	30	1	APO-Candesartan
Tab oz my marnyarodniorodniazido 12.0 my		00	•	HCTZ 32/12.5
OOADTAN DOTAGOUNAMENTU INCORONI COOTI II COOTI				11012 02/12.0
OSARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE	4.00	00		A
Tab 50 mg with hydrochlorothiazide 12.5 mg	4.00	30	•	Arrow-Losartan &
				<u>Hydrochlorothiazide</u>

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subs	idised	Generic	
\$	Per	✓	Manufacturer	

# **Angiotensin II Antagonists with Neprilysin Inhibitors**

	pharmacy	authority see SA2302 below - Retail	SACUBITRIL WITH VALSARTAN - Special Aut
✓ Entresto 24/26	56	190.00	Tab 24.3 mg with valsartan 25.7 mg
✓ Entresto 49/51	56	190.00	Tab 48.6 mg with valsartan 51.4 mg
✓ Entresto 97/103	56	190.00	Tab 97.2 mg with valsartan 102.8 mg

### ⇒SA2302 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has heart failure; and
- 2 Any of the following:
  - 2.1 Patient is in NYHA/WHO functional class II; or
  - 2.2 Patient is in NYHA/WHO functional class III; or
  - 2.3 Patient is in NYHA/WHO functional class IV; and
- 3 Either:
  - 3.1 Patient has a documented left ventricular ejection fraction (LVEF) of less than or equal to 35%; or
  - 3.2 An ECHO is not reasonably practical, and in the opinion of the treating practitioner the patient would benefit from treatment; and
- 4 Patient is receiving concomitant optimal standard chronic heart failure treatments.

a hudraphlarida rafar ta NEDVOLIC SYSTEM. Apparathatian Land. page 100

## **Antiarrhythmics**

For lignocaine hydrochloride refer to NERVOUS SYSTEM, Anaesthetics, Local, page	ge 126	
AMIODARONE HYDROCHLORIDE		
▲ Tab 100 mg3.49	30	✓ Aratac
▲ Tab 200 mg4.49	30	✓ Aratac
Inj 50 mg per ml, 3 ml ampoule - Up to 10 inj available on a PSO 9.12	6	✓ Cordarone-X
15.22	10	✓ Max Health
ATROPINE SULPHATE		
* Inj 600 mcg per ml, 1 ml ampoule - Up to 5 inj available on a		
PSO16.10	10	✓ Juno S29
		✓ Martindale
Martindale to be Principal Supply on 1 February 2025		
DIGOXIN		
* Tab 62.5 mcg - Up to 30 tab available on a PSO	240	✓ Lanoxin PG
* Tab 250 mcg - Up to 30 tab available on a PSO16.90	240	✓ Lanoxin
* Oral liq 50 mcg per ml	60 ml	✓ Lanoxin
		<ul> <li>Lanoxin Paediatric</li> </ul>
		Elixir
		✓ Lanoxin S29 S29
DISOPYRAMIDE PHOSPHATE		
▲ Cap 100 mg23.87	100	✓ Rythmodan
55.90	84	✓ Rythmodan -
00.00	• •	Cheplafarm S29

✓ Midodrine Medsurge

	_			
(1	Subsidy Manufacturer's Price) \$	Per	Fully Subsidised	
FLECAINIDE ACETATE				
▲ Tab 50 mg	19.95	60	✓	Flecainide BNM
▲ Cap long-acting 100 mg	35.78	90	•	Flecainide Controlled Release Teva
▲ Cap long-acting 200 mg	54.28	90	•	Flecainide Controlled Release Teva
Inj 10 mg per ml, 15 ml ampoule	102.79	5	✓	Almarytm \$29
	108.16		✓	Tambocor
MEXILETINE HYDROCHLORIDE				
▲ Cap 150 mg	162.00	100	✓	Teva S29
▲ Cap 250 mg	202.00	100	•	Teva S29
▲ Tab 150 mg	40.90	50	✓	Rytmonorm
Antihypotensives				
MIDODRINE - Special Authority see SA1474 below - Retail pharm	nacy			
Tab 2.5 mg		100		MAR-Midodrine S29 Midodrine Medsurge
Midodrine Medsurge to be Principal Supply on 1 February	2025			
Tab 5 mg	58.88	100	<b>✓</b>	MAR-Midodrine S29

Midodrine Medsurge to be Principal Supply on 1 February 2025

### ⇒SA1474 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 2 years where patient has disabling orthostatic hypotension not due to drugs.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

# **Beta-Adrenoceptor Blockers**

# **Beta Adrenoceptor Blockers**

ATENOLOL		
* Tab 50 mg11.00	500	✓ Viatris
Viatris to be Principal Supply on 1 February 2025		
* Tab 100 mg18.50	500	Atenolol Viatris
Atenolol Viatris to be Principal Supply on 1 February 2025		
* Oral liq 25 mg per 5 ml	300 ml OP	✓ Atenolol AFT
Restricted to children under 12 years of age.		
BISOPROLOL FUMARATE		
* Tab 2.5 mg1.36	90	✓ Ipca-Bisoprolol
* Tab 5 mg1.91	90	✓ Ipca-Bisoprolol
* Tab 10 mg2.71	90	✓ Ipca-Bisoprolol

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy		Fully Brand or
	(Manufacturer's Price) \$	Per	Subsidised Generic  Manufacturer
CARVEDILOL	*		
* Tab 6.25 mg	2.24	60	✓ Carvedilol Sandoz
* Tab 12.5 mg		60	✓ Carvedilol Sandoz
* Tab 25 mg		60	✓ Carvedilol Sandoz
LABETALOL			
* Tab 100 mg	14.50	100	✓ Trandate
* Tab 200 mg	27.00	100	✓ Trandate
* Inj 5 mg per ml, 20 ml ampoule		5	
, ,	(88.60)		Trandate
METOPROLOL SUCCINATE			
* Tab long-acting 23.75 mg	4.20	90	✓ Myloc CR
* Tab long-acting 47.5 mg		90	✓ Myloc CR
* Tab long-acting 95 mg	5.24	90	✓ Myloc CR
* Tab long-acting 190 mg	9.76	90	✓ Myloc CR
METOPROLOL TARTRATE			
* Tab 50 mg	5.66	100	✓ IPCA-Metoprolol
* Tab 100 mg		60	✓ IPCA-Metoprolol
* Tab long-acting 200 mg		28	✓ Slow-Lopresor
* Inj 1 mg per ml, 5 ml vial	26.50	5	✓ Metoprolol IV Mylan
			Metoprolol IV Viatris
NADOLOL			
* Tab 40 mg	19.19	100	✓ Nadolol BNM
* Tab 80 mg		100	✓ Nadolol BNM
PROPRANOLOL			
* Tab 10 mg	7.04	100	✓ Drofate
* Tab 40 mg		100	✓ IPCA-Propranolol
* Cap long-acting 160 mg		100	✓ Cardinol LA
* Oral lig 4 mg per ml - Special Authority see SA1327 belo			
Retail pharmacy		500 m	nl <b>✓ Roxane</b> -
· · · · · · · · · · · · · · · · · · ·			Propranolol S29
			•

### ⇒SA1327 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

### SOTALOL

*	Tab 80 mg	37.50	500	✓ Mylan
*	Tab 160 mg	14.00	100	Mylan

		Subsidy	Fully Bran		Brand or
		(Manufacturer's Price)	Sub	sidised	Generic
_		\$	Per		Manufacturer
	alabara Obarra di Birahara				
C	alcium Channel Blockers				
_	ibudranuridina Calaium Channal Blackara				
D	ihydropyridine Calcium Channel Blockers				
ΑN	ILODIPINE				
*	Tab 2.5 mg	1.45	90	1	Vasorex
	Tab 5 mg		90	-	Vasorex
	Tab 10 mg		90	-	Vasorex
	LODIPINE				
	Tab long-acting 2.5 mg	2 18	30	1	Plendil ER
~	Plendil ER to be Principal Supply on 1 February 2025	2.10	00	•	i iciidii Lii
*	Tab long-acting 5 mg	6 57	90	1	Felo 5 ER
~	Felo 5 ER to be Principal Supply on 1 February 2025	0.57	30	•	I CIO J LII
*	Tab long-acting 10 mg	6 95	90	<b>/</b>	Felo 10 ER
~	Felo 10 ER to be Principal Supply on 1 February 2025	0.33	30	•	I CIO IO LII
	FEDIPINE				
*	Tab long-acting 10 mg - Subsidy by endorsement	19.42	56	•	Tensipine MR10 S29
	Subsidised for patients who were taking nifedipine tab lo	na-acting 10 mg prior	to 1 July	2023 a	and the prescription is
	endorsed accordingly. Pharmacists may annotate the p				
	dispensing of nifedipine tab long-acting 10 mg.	recomplion as endorse	ou whole	uioio o	Aloto a record or prior
*	Tab long-acting 20 mg	17 72	100	<b>✓</b> I	Nyefax Retard
*	Tab long-acting 30 mg		14		Mylan Italy (24 hr
•••	745 1511g 454111g 55 111g		• •		release) \$29
		34.10	100	./	,
		34.10	100	•	Mylan (24 hr
					release) S29
*	Tab long-acting 60 mg	52.81	100	<b>✓</b>	Mylan (24 hr
					release) S29
_					
C	ther Calcium Channel Blockers				
ווח	TIAZEM HYDROCHLORIDE				
וטו	Cap long-acting 120 mg	65.35	500	1	Diltiazem CD Clinect
*	Cap long-acting 120 mg		30		Cardizem CD
	Cap long-acting 700 mg		30	-	Cardizem CD
		9.50	50		Cardizeiii CD
	RHEXILINE MALEATE	00.00	400		D
	Tab 100 mg	62.90	100	•	Pexsig
۷E	RAPAMIL HYDROCHLORIDE				
*	Tab 40 mg	7.01	100	✓	Isoptin
*	Tab 80 mg	11.74	100	✓	Isoptin
*	Tab long-acting 120 mg	36.02	100	✓	Isoptin Retard \$29
				✓	Isoptin SR

Isoptin SR

✓ Isoptin

30

5

Inj 2.5 mg per ml, 2 ml ampoule - Up to 5 inj available on a

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

(N	Subsidy fanufacturer's Price) \$	Per	Fully Subsidised	
Centrally-Acting Agents				
LONIDINE				
Patch 2.5 mg, 100 mcg per day - Only on a prescription	11.70	4	1	Mylan
Patch 5 mg, 200 mcg per day - Only on a prescription		4	✓	Mylan
Patch 7.5 mg, 300 mcg per day - Only on a prescription		4	1	Mylan
LONIDINE HYDROCHLORIDE				
€ Tab 25 mcg	29.32	112	1	Clonidine Teva
F Tab 150 mcg		100		Catapres
Catapres to be Principal Supply on 1 February 2025				•
Inj 150 mcg per ml, 1 ml ampoule	14.10	5	1	Catapres
, , , , ,	29.68	10		Medsurge
Catapres to be Principal Supply on 1 January 2025				-
Medsurge Inj 150 mcg per ml, 1 ml ampoule to be delisted 1 Janua	ry 2025)			
IETHYLDOPA				
	15.10	100	/	Methyldopa Viatris
9				, , , , , , , , , , , , , , , , , , , ,
Diuretics				
Loop Diuretics				
=p =				
UMETANIDE				
€ Tab 1 mg	16.36	100	✓	Burinex
Inj 500 mcg per ml, 4 ml vial	7.95	5	✓	Burinex
UROSEMIDE [FRUSEMIDE]				
Tab 40 mg - Up to 30 tab available on a PSO	12.80	1,000	/	IPCA-Frusemide
IPCA-Frusemide to be Principal Supply on 1 February 2025		.,		
F Tab 500 mg		50	/	Urex Forte
Oral liq 10 mg per ml		0 ml C		Lasix
Inj 10 mg per ml, 25 ml ampoule		6	1	Lasix
Inj 10 mg per ml, 2 ml ampoule - Up to 5 inj available on a PSC		5	1	Furosemide-Baxter
Potassium Sparing Diuretics				
MILORIDE HYDROCHLORIDE				
Tab 5 mg	81.07	100	1	Padagis S29
	171.41	28	1	Wockhardt S29
Oral liq 1 mg per ml	33.71 2	5 ml C	)P 🗸	Biomed
PLERENONE - Special Authority see SA1728 below - Retail pha				
Tab 25 mg		30	1	Inspra
Tab 50 mg		30		Inspra
. as as an inf	20.00	50	•	<b>.</b>

**Initial application** from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

### Both:

- 1 Patient has heart failure with ejection fraction less than 40%; and
- 2 Fither
  - 2.1 Patient is intolerant to optimal dosing of spironolactone; or
  - 2.2 Patient has experienced a clinically significant adverse effect while on optimal dosing of spironolactone.

				_
	Subsidy (Manufacturer's Price) \$		Fully dised	Brand or Generic Manufacturer
SPIRONOLACTONE  * Tab 25 mg  * Tab 100 mg  Oral liq 5 mg per ml	10.65	100 100 5 ml OP	<b>√</b> 9	Spiractin Spiractin Biomed
Potassium Sparing Combination Diuretics				
AMILORIDE HYDROCHLORIDE WITH FUROSEMIDE  * Tab 5 mg with furosemide 40 mg		28	<b>✓</b> I	Frumil
* Tab 5 mg with hydrochlorothiazide 50 mg		50	<b>√</b>	Moduretic
Thiazide and Related Diuretics				
BENDROFLUMETHIAZIDE [BENDROFLUAZIDE]  * Tab 2.5 mg – Up to 150 tab available on a PSO	51.50	500	1	Arrow- Bendrofluazide
May be supplied on a PSO for reasons other than emerg  * Tab 5 mg		500	<b>√</b> <u>i</u>	Arrow- Bendrofluazide
CHLOROTHIAZIDE Oral liq 50 mg per ml	29.21 2	5 ml OP	<b>✓</b> I	Biomed
CHLORTALIDONE [CHLORTHALIDONE]  * Tab 25 mg	6.95	50	<b>✓</b> <u>I</u>	<u>Hygroton</u>
INDAPAMIDE  ★ Tab 2.5 mg  METOLAZONE	16.00	90	<b>√</b> <u>I</u>	Dapa-Tabs
Tab 5 mg	CBS	1 50	-	Metolazone ©29 Zaroxolyn ©29
Vasopressin receptor antagonists				
TOLVAPTAN — Special Authority see SA2166 below — Retail pha Tab 15 mg Tab 30 mg Tab 45 mg + 15 mg Tab 60 mg + 30 mg Tab 90 mg + 30 mg	873.50 873.50 1,747.00 1,747.00	28 OP 28 OP 56 OP 56 OP 56 OP	1	Jinarc Jinarc Jinarc Jinarc Jinarc

**⇒SA2166** Special Authority for Subsidy

Initial application — (autosomal dominant polycystic kidney disease) only from a renal physician or any relevant practitioner on the recommendation of a renal physician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of autosomal dominant polycystic kidney disease; and
- 2 Patient has an estimated glomerular filtration rate (eGFR) of greater than or equal to 25 ml/min/1.73 m² at treatment initiation; and
- 3 Either:

		Subsidy (Manufacturer's Price) \$	Fully Subsidised Per 🗸	Brand or Generic Manufacturer
	Patient's disease is rapidly progressing, with a one-year; or	· ·	•	
3.2	Patient's disease is rapidly progressing, with an		R of greater than	or equal to
Donowol (	2.5 mL/min/1.73 m <sup>2</sup> per year over a five-year p autosomal dominant polycystic kidney disea		voicion or ony rol	avent proctitioner on the
	ion of a renal physician. Approvals valid for 12			
	t has not developed end-stage renal disease, do t has not undergone a kidney transplant.	efined as an eGFR of less	s than 15 mL/min	1/1.73 m <sup>2</sup> ; and
Lipid-Mod	difying Agents			
Fibrates				
BEZAFIBRAT	E			
	mg	22.65	90 🗸 E	Bezalip
	lip to be Principal Supply on 1 March 2025 acting 400 mg	21 54	30 <b>✓ E</b>	Bezalip Retard
-	lip Retard to be Principal Supply on 1 March 20			Sezanp rictard
Other Lip	id-Modifying Agents			
ACIPIMOX			_	
<b>₭</b> Cap 250	mg	38.19	30	Olbetam
Resins				
COLESTYRA			_	
Powder fo	or oral suspension 4 g sachet	61.50	50	Colestyramine -
			./ (	Mylan S29
			• (	Quantalan sugar free S29
HMG CoA	A Reductase Inhibitors (Statins)			
ATORVASTA	TIN			
-	g	5.16	500 <b>✓</b> <u>L</u>	_orstat
	ğ		-	orstat
	g		-	<u>-orstat</u>
	g	∠ე.ᲐᲧ	500 <b>✓</b> <u>L</u>	<u>-orstat</u>
PRAVASTAT ₭ Tab 20 m		7 16	100 🗸 (	Clinect
	g		_	<u>Silnect</u> Clinect
	TIN - Special Authority see SA2093 on the ne		_	, <del>.</del>
	Triv - Special Authority see SA2090 on the fie.		,	Rosuvastatin Viatris
	g			Rosuvastatin Viatris
		0.74	00	No

\* Tab 20 mg .......2.71

30

30

✓ Rosuvastatin Viatris

✓ Rosuvastatin Viatris

Subsidy (Manufacturer's Price)	Su	Fully bsidised	Brand or Generic
 \$	Per	1	Manufacturer

### ⇒SA2093 Special Authority for Subsidy

Initial application — (cardiovascular disease risk) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Either:

- 1 Both:
  - 1.1 Patient is considered to be at risk of cardiovascular disease; and
  - 1.2 Patient is Māori or any Pacific ethnicity; or
- 2 Both:
  - 2.1 Patient has a calculated risk of cardiovascular disease of at least 15% over 5 years; and
  - 2.2 LDL cholesterol has not reduced to less than 1.8 mmol/litre with treatment with the maximum tolerated dose of atoryastatin and/or simyastatin.

Initial application — (familial hypercholesterolemia) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Both:

- 1 Patient has familial hypercholesterolemia (defined as a Dutch Lipid Criteria score greater than or equal to 6); and
- 2 LDL cholesterol has not reduced to less than 1.8 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initial application — (established cardiovascular disease) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 Patient has proven coronary artery disease (CAD); or
  - 1.2 Patient has proven peripheral artery disease (PAD); or
  - 1.3 Patient has experienced an ischaemic stroke; and
- 2 LDL cholesterol has not reduced to less than 1.4 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initial application — (recurrent major cardiovascular events) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has experienced a recurrent major cardiovascular event (defined as myocardial infarction, ischaemic stroke, coronary revascularisation, hospitalisation for unstable angina) in the last 2 years; and
- 2 LDL cholesterol has not reduced to less than 1.0 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

SIN	NVASTATIN			
*	Tab 10 mg	1.68	90	<ul> <li>Simvastatin Mylan</li> </ul>
	ŭ			✓ Simvastatin Viatris
*	Tab 20 mg	2.54	90	✓ Simvastatin Viatris
		4.11	90	✓ Simvastatin Viatris
		8.81	90	✓ Simvastatin Viatris

#### 

(Ezemibe Viatris Tab 10 mg to be delisted 1 July 2025)

✓ Ezetimibe Sandoz

### CARDIOVASCIII AR SYSTEM

CANDIOVASCULAN STSTEW				
	Subsidy (Manufacturer's Pri	ce) Subs	Fully sidised	Brand or Generic Manufacturer
EZETIMIBE WITH SIMVASTATIN				
Tab 10 mg with simvastatin 10 mg	5.15	30	<b>√</b> Z	Zimybe
Tab 10 mg with simvastatin 20 mg	6.15	30	✓ Z	Zimybe
Tab 10 mg with simvastatin 40 mg		30	✓ Z	Zimybe
Tab 10 mg with simvastatin 80 mg	8.15	30	<b>√</b> Z	Zimybe
Nitrates				
GLYCERYL TRINITRATE				•
* Oral pump spray, 400 mcg per dose – Up to 250 dose	7 48 2	250 dose OP	<b>✓</b> N	Nitrolingual Pump

*	Oral pump spray, 400 mcg per dose – Up to 250 dose available on a PSO	7.48	250 dose OP	✓ Nitrolingual Pump Spray
*	Patch 25 mg, 5 mg per day	15.73	30	✓ Nitroderm TTS
	Patch 50 mg, 10 mg per day		30	✓ Nitroderm TTS
ISC	SORBIDE MONONITRATE			
*	Tab 20 mg	22.49	100	✓ Ismo 20
*	Tab long-acting 40 mg	9.80	30	✓ Ismo 40 Retard

Tab long-acting 60 mg......13.50

# **Sympathomimetics**

ADRENALINE		
Inj 1 in 1,000, 1 ml ampoule - Up to 5 inj available on a PSO4.98	5	✓ Aspen Adrenaline
13.27		✓ DBL Adrenaline
25.30	10	✓ Hameln S29
Inj 1 in 10,000, 10 ml ampoule - Up to 5 inj available on a PSO27.00	5	✓ Hospira
49.00	10	Aspen Adrenaline

## **Vasodilators**

	DITALAZINE ITI DITOOTEOTIDE		
*	Tab 25 mg - Special Authority see SA1321 below - Retail		
	pharmacyCBS	1	<ul><li>Hydralazine</li></ul>
		56	✓ Onelink S29
		84	✓ AMDIPHARM \$29
		100	✓ Camber S29

### ⇒SA1321 Special Authority for Subsidy

HYDRAL AZINE HYDROCHLORIDE

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

### Either:

1 For the treatment of refractory hypertension; or

\* Inj 20 mg ampoule......25.90

2 For the treatment of heart failure in combination with a nitrate, in patients who are intolerant or have not responded to ACE inhibitors and/or angiotensin receptor blockers.

#### MINOXIDIL

▲ Tab 10 mg	47.04 78.40	60 100	✓ Minoxidil Roma S29 ✓ Loniten
NICORANDIL			
▲ Tab 10 mg	21.73	60	✓ Max Health
▲ Tab 20 mg		60	✓ Max Health

✓ Duride

✓ Apresoline

90

5

✓ Ambrisentan Viatris

✓ Ambrisentan Viatris

30

30

(	Subsidy (Manufacturer's Price) \$	Su Per	Fully bsidised	Brand or Generic Manufacturer
PAPAVERINE HYDROCHLORIDE				
* Inj 12 mg per ml, 10 ml ampoule	257.12	5	<b>✓</b> H	lospira
PENTOXIFYLLINE [OXPENTIFYLLINE]				
Tab 400 mg	44.37	50	<b>✓</b> T	rental 400
Endothelin Receptor Antagonists				
AMBRISENTAN - Special Authority see SA2253 below - Retail pl	harmacy			

### 

Initial application — (PAH monotherapy) only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

#### All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and

- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH has been confirmed by right heart catheterisation; and
    - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
    - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
    - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm<sup>-5</sup>); and
    - 4.1.5 Any of the following:
      - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH 2022 (see note below for link to these guidelines) †; or
      - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool\*\*; or
      - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
  - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
  - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
  - 5.1 Ambrisentan is to be used as PAH monotherapy; and
  - 5.2 Any of the following:
    - 5.2.1 Patient has experienced intolerable side effects with both sildenafil and bosentan; or
    - 5.2.2 Patient has an absolute contraindication to sildenafil and an absolute or relative contraindication to bosentan (e.g. due to current use of a combined oral contraceptive or liver disease); or
    - 5.2.3 Patient is a child with idiopathic PAH or PAH secondary to congenital heart disease.

**Initial application** — **(PAH dual therapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

### All of the following:

1 Patient has pulmonary arterial hypertension (PAH); and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per ✓	Manufacturer

continued...

- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH has been confirmed by right heart catheterisation; and
    - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
    - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
    - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm<sup>5</sup>); and
    - 4.1.5 Any of the following:
      - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH 2022 (see note below for link to these guidelines) †; or
      - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool\*\*; or
      - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
  - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
  - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 All of the following:
  - 5.1 Ambrisentan is to be used as PAH dual therapy; and
  - 5.2 Either
    - 5.2.1 Patient has tried a PAH monotherapy (sildenafil or bosentan) for at least three months and has not experienced an acceptable response to treatment according to a validated risk stratification tool\*\*; or
    - 5.2.2 Patient has tried PAH dual therapy including bosentan and has experienced intolerable side effects on bosentan; and
  - 5.3 Both:
    - 5.3.1 Patient is presenting in NYHA/WHO functional class III or IV, and in the opinion of the treating clinician would benefit from initial dual therapy; and
    - 5.3.2 Patient has an absolute or relative contraindication to bosentan (e.g. due to current use of a combined oral contraceptive or liver disease).

**Initial application** — **(PAH triple therapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH has been confirmed by right heart catheterisation; and
    - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
    - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
    - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm<sup>5</sup>); and
    - 4.1.5 Any of the following:

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
 \$	Per	1	

continued...

- 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH 2022 (see note below for link to these guidelines) †; or
- 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool\*\*; or
- 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
- 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
  - 5.1 Ambrisentan is to be used as PAH triple therapy; and
  - 5.2 Any of the following:
    - 5.2.1 Patient is on the lung transplant list; or
    - 522 Both
      - 5.2.2.1 Patient is presenting in NYHA/WHO functional class IV; and
      - 5.2.2.2 Patient has an absolute or relative contraindication to bosentan (e.g. due to current use of a combined oral contraceptive or liver disease); or
    - 5.2.3 Both:
      - 5.2.3.1 Patient has tried PAH dual therapy for at least three months and remains in an unacceptable risk category according to a validated risk stratification tool\*\*; and
      - 5.2.3.2 Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario.

**Renewal** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 2 years where the patient is continuing to derive benefit from ambrisentan treatment according to a validated PAH risk stratification tool\*\*.

Notes: † The European Respiratory Journal Guidelines can be found here: 2022 ECS/ERS Guidelines for the diagnosis and treatment of pulmonary hypertension PAH

\*\* the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

BOSENTAN - Special Authority see SA2254 below - Retail pharmacy		
Tab 62.5 mg100.00	60	✓ Bosentan Dr Reddy's
Bosentan Dr Reddy's to be Principal Supply on 1 January 2025		
Tab 125 mg100.00	60	✓ Bosentan Dr Reddy's

Bosentan Dr Reddy's to be Principal Supply on 1 January 2025

### ⇒SA2254 Special Authority for Subsidy

**Initial application** — **(PAH monotherapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)\*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and

Subsidy		Fully	Brand or	Т
(Manufacturer's Price)	5	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH has been confirmed by right heart catheterisation; and
    - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
    - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
    - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm<sup>5</sup>); and
    - 4.1.5 Any of the following:
      - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these quidelines) †: or
      - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool\*\*; or
      - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
  - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or
  - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
  - 5 Both:
    - 5.1 Bosentan is to be used as PAH monotherapy; and
    - 5.2 Any of the following:
      - 5.2.1 Patient has experienced intolerable side effects on sildenafil; or
      - 5.2.2 Patient has an absolute contraindication to sildenafil; or
      - 5.2.3 Patient is a child with idiopathic PAH or PAH secondary to congenital heart disease.

**Initial application** — **(PAH dual therapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)\*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH has been confirmed by right heart catheterisation; and
    - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
    - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
    - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm<sup>-5</sup>); and
    - 4.1.5 Any of the following:
      - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these quidelines) †: or
      - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool\*\*: or
      - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
  - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or

Su	ubsidy F	ully	Brand or
(Manufact	turer's Price) Subsidi	sed	Generic
·	\$ Per	✓	Manufacturer

continued...

- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Bosentan is to be used as part of PAH dual therapy; and
- 6 Either:
  - 6.1 Patient has tried a PAH monotherapy (sildenafil) for at least three months and has experienced an inadequate therapeutic response to treatment according to a validated risk stratification tool\*\*; or
  - 6.2 Patient is presenting in NYHA/WHO functional class III or IV, and in the opinion of the treating clinician would likely benefit from initial dual therapy.

**Initial application** — **(PAH triple therapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)\*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH has been confirmed by right heart catheterisation; and
    - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
    - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
    - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm<sup>-5</sup>); and
    - 4.1.5 Any of the following:
      - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these quidelines) † : or
      - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool\*\*: or
      - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
  - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or
  - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
  - 5.1 Bosentan is to be used as part of PAH triple therapy; and
  - 5.2 Any of the following:
    - 5.2.1 Patient is on the lung transplant list: or
    - 5.2.2 Patient is presenting in NYHA/WHO functional class IV: or
    - 5.2.3 Both:
      - 5.2.3.1 Patient has tried PAH dual therapy for at least three months and has not experienced an acceptable response to treatment according to a validated risk stratification tool\*\*: and
      - 5.2.3.2 Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario.

**Renewal** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 2 years where patient is continuing to derive benefit from bosentan treatment according to a validated PAH risk stratification tool\*\*.

Notes: † The European Respiratory Journal Guidelines can be found here: 2022 ECS/ERS Guidelines for the diagnosis and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsidised		Generic	
\$	Per	✓	Manufacturer	

continued...

### treatment of pulmonary hypertension PAH

\*\* the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

### Phosphodiesterase Type 5 Inhibitors

SILDENAFIL – Special Authority see SA2255 below – Retail phar	macy		
Tab 25 mg	0.72	4	✓ Vedafil
Tab 50 mg	1.45	4	✓ Vedafil
Tab 100 mg		12	✓ Vedafil

### ⇒SA2255 Special Authority for Subsidy

Initial application — (Raynaud's Phenomenon\*) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has Ravnaud's Phenomenon\*: and
- 2 Patient has severe digital ischaemia (defined as severe pain requiring hospital admission or with a high likelihood of digital ulceration; digital ulcers; or gangrene); and
- 3 Patient is following lifestyle management (avoidance of cold exposure, sufficient protection, smoking cessation support, avoidance of sympathomimetic drugs); and
- 4 Patient is being treated with calcium channel blockers and nitrates (or these are contraindicated/not tolerated).

Initial application — (Pulmonary arterial hypertension\*) only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)\*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II. III or IV: and
- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH is confirmed by right heart catheterisation; and
    - 4.1.2 A mean pulmonary artery pressure (PAPm) of greater than 20 mmHg; and
    - 4.1.3 A pulmonary capillary wedge pressure (PCWP) that is less than or equal to 15 mmHg; and
    - 4.1.4 Pulmonary vascular resistance (PVR) of at least 2 Wood Units or greater than 160 International Units (dyn s cm<sup>-5</sup>); and
    - 4.1.5 Any of the following:
      - 4.1.5.1 PAH is non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) †; or
      - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool\*\*: or
      - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
  - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or
  - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures.

Initial application — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:
Both:

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic	
\$	Per	•	Manufacturer	

continued...

- 1 Patient has a documented history of traumatic or non-traumatic spinal cord injury; and
- 2 Patient has erectile dysfunction secondary to spinal cord injury requiring pharmacological treatment.

Renewal — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Notes: Note: Indications marked with \* are Unapproved Indications.

† The European Respiratory Journal Guidelines can be found here: 2022 ECS/ERS Guidelines for the diagnosis and treatment of pulmonary hypertension PAH

\*\* the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

### **Prostacyclin Analogues**

		EPOPROSTENOL – Special Authority see SA2256 below – Retail pharmacy
✓ Veletri	1	Inj 500 mcg vial36.61
✓ Veletri	1	Inj 1.5 mg vial73.21

### ⇒SA2256 Special Authority for Subsidy

**Initial application** — **(PAH dual therapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class III or IV; and
- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH has been confirmed by right heart catheterisation; and
    - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
    - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
    - 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm<sup>-5</sup>); and
    - 4.1.5 Any of the following:
      - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these quidelines) †: or
      - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool\*\*; or
      - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
  - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
  - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 All of the following:
  - 5.1 Epoprostenol is to be used as part of PAH dual therapy with either sildenafil or an endothelin receptor antagonist; and
  - 5.2 Patient is presenting in NYHA/WHO functional class IV; and
  - 5.3 Patient has tried a PAH monotherapy for at least three months and remains in an unacceptable risk category according to a validated risk stratification tool.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

**Initial application** — **(PAH triple therapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class III or IV; and
- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH has been confirmed by right heart catheterisation; and
    - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
    - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
    - 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm<sup>5</sup>); and
    - 4.1.5 Any of the following:
      - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) †; or
      - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool\*\*; or
      - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
  - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
  - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and

#### 5 Both:

- 5.1 Epoprostenol is to be used as PAH triple therapy; and
- 5.2 Any of the following:
  - 5.2.1 Patient is on the lung transplant list; or
  - 5.2.2 Patient is presenting in NYHA/WHO functional class IV: or
  - 5.2.3 Both:
    - 5.2.3.1 Patient has tried PAH dual therapy for at least three months and has not experienced an acceptable response to treatment according to a validated risk stratification tool; and
    - 5.2.3.2 Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario.

**Renewal** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 2 years where patient is continuing to derive benefit from epoprostenol treatment according to a validated PAH risk stratification tool\*\*.

Notes: † The European Respiratory Journal Guidelines can be found here: 2022 ECS/ERS Guidelines for the diagnosis and treatment of pulmonary hypertension PAH

\*\*\* the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

30

✓ Vebulis

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic 
\$ Per ✔ Manufacturer

⇒SA2257 Special Authority for Subsidy

**Initial application** — **(PAH monotherapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH has been confirmed by right heart catheterisation; and
    - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
    - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
    - 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm<sup>-5</sup>); and
    - 4.1.5 Any of the following:
      - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) †; or
      - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool\*\*: or
      - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
  - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
  - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
  - 5.1 Iloprost is to be used as PAH monotherapy; and
  - 5.2 Either:
    - 5.2.1 Patient has experienced intolerable side effects on sildenafil and both the funded endothelin receptor antagonists (i.e. both bosentan and ambrisentan); or
    - 5.2.2 Patient has an absolute contraindication to sildenafil and an absolute or relative contraindication to endothelin receptor antagonists.

**Initial application** — **(PAH dual therapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH has been confirmed by right heart catheterisation; and
    - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
    - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
    - 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm<sup>-5</sup>); and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$ P	Per	Manufacturer

continued...

- 4.1.5 Any of the following:
  - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) †; or
  - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool\*\*: or
  - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
- 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 All of the following:
  - 5.1 lloprost is to be used as PAH dual therapy with either sildenafil or an endothelin receptor antagonist; and
  - 5.2 Either:
    - 5.2.1 Patient has an absolute contraindication to or has experienced intolerable side effects on sildenafil; or
    - 5.2.2 Patient has an absolute or relative contraindication to or experienced intolerable side effects with a funded endothelin receptor antagonist; and
  - 5.3 Either:
    - 5.3.1 Patient has tried a PAH monotherapy for at least three months and remains in an unacceptable risk category according to a validated risk stratification tool\*\*; or
    - 5.3.2 Patient is presenting in NYHA/WHO functional class III or IV, and in the opinion of the treating clinician would benefit from initial dual therapy.

Initial application — (PAH triple therapy) only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH has been confirmed by right heart catheterisation; and
    - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
    - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
    - 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm<sup>-5</sup>); and
    - 4.1.5 Any of the following:
      - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these quidelines) †: or
      - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool\*\*; or
      - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
  - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
  - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and

Subsidy (Manufacturer's Pric	ce)	Fully Subsidised	Brand or Generic	
<u> </u>	Per	✓	Manufacturer	

#### continued...

- 5 Both:
  - 5.1 Iloprost is to be used as PAH triple therapy; and
  - 5.2 Any of the following:
    - 5.2.1 Patient is on the lung transplant list; or
    - 5.2.2 Patient is presenting in NYHA/WHO functional class IV; or
    - 5.2.3 Both:
      - 5.2.3.1 Patient has tried PAH dual therapy for at least three months and has not experienced an acceptable response to treatment according to a validated risk stratification tool\*\*; and
      - 5.2.3.2 Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario.

**Renewal** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 2 years where patient is continuing to derive benefit from iloprost treatment according to a validated PAH risk stratification tool\*\*.

Notes: † The European Respiratory Journal Guidelines can be found here: 2022 ECS/ERS Guidelines for the diagnosis and treatment of pulmonary hypertension PAH

\*\* the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.



Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

# **Antiacne Preparations**

For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 98

#### **ADAPALENE**

- a) Maximum of 30 g per prescription
- h) Only on a prescription

b) Only on a presemption			
Gel 0.1%	22.89	30 g OP	<ul><li>Differin</li></ul>
ISOTRETINOIN - Special Authority see SA2023 below - Rei	tail pharmacy		
Cap 5 mg	11.26	60	<ul><li>Oratane</li></ul>
Cap 10 mg	18.75	120	✓ Oratane
Cap 20 mg	26.73	120	✓ Oratane

#### ⇒SA2023 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around isotretinoin and is competent to prescribe isotretinoin; and
- 3 Either:
  - 3.1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and that they must not become pregnant during treatment and for a period of one month after the completion of treatment; or
  - 3.2 Patient is not of child bearing potential.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and that they must not become pregnant during treatment and for a period of one month after the completion of treatment: or
- 2 Patient is not of child bearing potential.

### **TRETINOIN**

Crm 0.5 mg per g - Maximum of 50 g per prescription	16.82	50 g OP	✓ ReTrieve
ReTrieve to be Principal Supply on 1 February 2025			

# **Antibacterials Topical**

For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 98

HYDROGEN PEROXIDE			
* Crm 1%	8.56	10 g OP	<ul><li>Crystaderm</li></ul>
MUPIROCIN			
Oint 2%	6.60	15 g OP	
	(13.00)	· ·	Bactroban

- a) Only on a prescription
- b) Not in combination

			ERM	ATOLOGICALS
	Subsidy (Manufacturer's F \$	Price) Subs	Fully sidised	Brand or Generic Manufacturer
SODIUM FUSIDATE [FUSIDIC ACID]				
Crm 2%	1.69	5 g OP	<b>✓</b> F	oban
a) Maximum of 5 g per prescription b) Only on a prescription c) Not in combination d) Foban to be Principal Supply on 1 February 2025 Oint 2% a) Maximum of 5 g per prescription b) Only on a prescription c) Not in combination	1.69	5 g OP	<b>√</b> F	oban
d) Foban to be Principal Supply on 1 February 2025				
SULFADIAZINE SILVER				
Crm 1%		50 g OP	_	lamazine
	15.44		<b>✓</b>	Ascend \$29
<ul><li>a) Up to 250 g available on a PSO</li><li>b) Not in combination</li></ul>				
Antifungals Topical				
For systemic antifungals, refer to INFECTIONS, Antifungals, pa	ge 105			
AMOROLFINE				
a) Only on a prescription				
b) Not in combination	04.07	Г I ОП		ANI.
Nail soln 5%	21.07	5 ml OP	<u> </u>	<u>lycoNail</u>
CLOTRIMAZOLE  * Crm 1%	1 10	20 g OP	./ (	Clomazol
a) Only on a prescription	1.10	20 g OF	• •	JOIIIazoi
b) Not in combination				
* Soln 1%	4.36	20 ml OP		
	(7.55)		(	Canesten
<ul><li>a) Only on a prescription</li><li>b) Not in combination</li></ul>				
ECONAZOLE NITRATE				
Crm 1%	1.00 (8.09)	20 g OP	F	<sup>o</sup> evaryl
a) Only on a prescription				
b) Not in combination				
	0.00	^		

3

Pevaryl

(18.64)

a) Only on a prescriptionb) Not in combination

Foaming soln 1%, 10 ml sachets......9.89

# **DERMATOLOGICALS**

	Subsidy (Manufacturer's \$	Price) Per	Fully Subsidised	
MICONAZOLE NITRATE  * Crm 2%  a) Only on a prescription	0.90	15 g (	OP 🗸	<u>Multichem</u>
b) Not in combination  * Lotn 2%	4.36 (10.03)	30 ml (	OP	Daktarin
a) Only on a prescription b) Not in combination  * Tinct 2%	4.36 (12.10)	30 ml (	OP	Daktarin
a) Only on a prescription     b) Not in combination				
Antipruritic Preparations				
CALAMINE  a) Only on a prescription b) Not in combination Crm, aqueous, BP	3.45	100	g 🗸	healthE Calamine
CROTAMITON				Aqueous
a) Only on a prescription b) Not in combination  Crm 10%  Itch-Soothe to be Principal Supply on 1 February 2025	3.49	20 g (	OP ✓	Itch-Soothe
MENTHOL – Only in combination				
<ol> <li>Only in combination with a dermatological base or propri</li> <li>With or without other dermatological galenicals.</li> </ol>	ietary Topical (	Corticoster	riod – Plain	
Crystals	6.92 29.60	25 g 100 g	,	MidWest MidWest
Corticosteroids Topical				
For systemic corticosteroids, refer to CORTICOSTEROIDS AND F	RELATED AGE	ENTS, pag	e 88	
Corticosteroids - Plain				
BETAMETHASONE DIPROPIONATE				

BETAMETHASONE DIPROPIONATE		
Crm 0.05%2.96	15 g OP	✓ Diprosone
36.00	50 g OP	✓ Diprosone
Oint 0.05%2.96	15 g OP	✓ Diprosone
36.00	50 g OP	✓ Diprosone
Oint 0.05% in propylene glycol base4.33	30 g OP	Diprosone OV
BETAMETHASONE VALERATE		
* Crm 0.1%	50 g OP	✓ Beta Cream
Beta Cream to be Principal Supply on 1 February 2025		
* Oint 0.1%7.90	50 g OP	✓ Beta Ointment
Beta Ointment to be Principal Supply on 1 February 2025		
<b>*</b> Lotn 0.1%	50 ml OP	✓ Betnovate

	Subsidy		Fully	
	(Manufacturer's Price \$	e) Subsid Per	dised •	Generic Manufacturer
OLOPETAGOL PROPIONATE	Ψ	rei	<u> </u>	Manuacturer
CLOBETASOL PROPIONATE  * Crm 0.05%	0.40	30 g OP	./	Dermol
* Oint 0.05%		30 g OP		Dermol
	2.00	30 y Oi	٠	Defiliol
CLOBETASONE BUTYRATE  Crm 0.05%	E 20	20 a OB		
GIII 0.05%	(10.00)	30 g OP		Eumovate
HYDROCORTISONE	(10.00)			Lumovato
* Crm 1% - Only on a prescription	1 70	30 g OP	_	Ethics
* Citil 1 % - Only of a prescription	20.40	500 g		Noumed
* Powder – Only in combination		25 g		ABM
Up to 5% in a dermatological base (not proprietary Topic		•		
galenicals			••••	iout outor dominatorogical
HYDROCORTISONE AND PARAFFIN LIQUID AND LANOLIN				
Lotn 1% with paraffin liquid 15.9% and lanolin 0.6% – Only o	n			
a prescription		250 ml	1	DP Lotn HC
HYDROCORTISONE BUTYRATE		200		<u></u>
Lipocream 0.1%	4 85	100 g OP	1	Locoid Lipocream
Oint 0.1%		100 g OP		Locoid
Milky emul 0.1%		00 ml OP	1	Locoid Crelo
METHYLPREDNISOLONE ACEPONATE				
Crm 0.1%	4.95	15 g OP	1	Advantan
Oint 0.1%		15 g OP	_	Advantan
MOMETASONE FUROATE		- 3 -		
Crm 0.1%	2.25	15 g OP	1	Elocon Alcohol Free
<b> </b>	3.50	50 g OP		Elocon Alcohol Free
Elocon Alcohol Free to be Principal Supply on 1 February	/ 2025	J		
Oint 0.1%	2.25	15 g OP	1	Elocon
	3.50	50 g OP	1	Elocon
Elocon to be Principal Supply on 1 February 2025			_	
Lotn 0.1%	4.99	30 ml OP	/	Elocon
Elocon to be Principal Supply on 1 February 2025				
TRIAMCINOLONE ACETONIDE			_	
Crm 0.02%		100 g OP		Aristocort
Oint 0.02%	6.54	100 g OP	•	<u>Aristocort</u>
Corticosteroids - Combination				
BETAMETHASONE VALERATE WITH SODIUM FUSIDATE [FUS	-	45 00		
Crm 0.1% with sodium fusidate (fusidic acid) 2%		15 g OP		F
a) Maximum of 15 a new proportion	(10.45)			Fucicort
a) Maximum of 15 g per prescription     b) Only on a prescription				
, , , , ,				
HYDROCORTISONE WITH MICONAZOLE — Only on a prescript		15 a OD	.,	Miorama H
Crm 1% with miconazole nitrate 2%	2.85	15 g OP	•	Micreme H
HYDROCORTISONE WITH NATAMYCIN AND NEOMYCIN — Or	, , ,		.,	Pimafucort
Oint 1% with natamycin 1% and neomycin sulphate 0.5%	3.35	15 g OP	•	rinialucort

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's \$		Fully Brand or dised Generic  Manufacturer
TRIAMCINOLONE ACETONIDE WITH GRAMICIDE Crm 1 mg with nystatin 100,000 u, neomycin su		TIN	
and gramicidin 250 mcg per g - Only on a	prescription3.49 (9.28)	15 g OP	Viaderm KC
Barrier Creams and Emollients			
Barrier Creams			
DIMETHICONE  * Crm 5% pump bottle	4.30	500 ml OP	✓ <u>healthE</u> Dimethicone 5%
* Crm 10% pump bottle	4.52	500 ml OP	✓ healthE  Dimethicone 10%
ZINC AND CASTOR OIL  * Oint	4.25	500 g	✓ Evara
Emollients			
AQUEOUS CREAM Crm	1.30	100 g	✓ healthE Aqueous Cream SLS Free
	1.65 1.73	500 g	✓ Evara ✓ GEM Aqueous Cream
(healthE Aqueous Cream SLS Free Crm to be delis (GEM Aqueous Cream Crm to be delisted 1 March			
CETOMACROGOL  * Crm BP  Cetomacrogol-AFT to be Principal Supply of		500 g	✓ Cetomacrogol-AFT
CETOMACROGOL WITH GLYCEROL Crm 90% with glycerol 10%	2.13 3.50	500 ml OP 1,000 ml OP	✓ <u>Evara</u> ✓ <u>Evara</u>
EMULSIFYING OINTMENT  * Oint BP	3.13	500 g	✓ Emulsifying Ointment ADE
OIL IN WATER EMULSION  * Crm		500 g	✓ Fatty Cream AFT
(Fatty Cream AFT Crm to be delisted 1 April 2025)	2.10		✓ Fatty Emulsion Cream (Evara)
PARAFFIN			
Oint liquid paraffin 50% with white soft paraffin	50%4.94	500 g OP	✓ White Soft Liquid  Paraffin AFT
WREA	1.37	100 g OP	✓ healthE Urea Cream

✓ EVARA White Soft Paraffin

	Subsidy		Fully Brand or
	(Manufacturer's I \$	Price) Subsic Per	dised Generic  Manufacturer
WOOL FAT WITH MINERAL OIL - Only on a prescription			
* Lotn hydrous 3% with mineral oil	5.60	1,000 ml	
	(14.96)		DP Lotion
	(20.53)		Alpha-Keri Lotion
	1.40	250 ml OP	
	(5.87)		DP Lotion
	5.60	1,000 ml	
	(23.91)		BK Lotion
	1.40	250 ml OP	
	(7.73)		BK Lotion
Other Dermatological Bases			
PARAFFIN			
White soft - Only in combination	4.74	450 g	✓ EVARA White Soft  Paraffin

Only in combination with a dermatological galenical or as a diluent for a proprietary Topical Corticosteroid – Plain.

19.00

2.500 a

Minor Skin Infections			
POVIDONE IODINE			
Oint 10%	7.40	65 g OP	<ul><li>Betadine</li></ul>
a) Maximum of 130 g per prescription			
b) Only on a prescription			
Antiseptic Solution 10%	4.99	100 ml	✓ Riodine
Antiseptic soln 10%	3.83	15 ml	✓ Riodine
	6.99	500 ml	✓ Riodine
Skin preparation, povidone iodine 10% with 30% alcohol	1.63	100 ml	
	(3.48)		Betadine Skin Prep

# **Parasiticidal Preparations**

DIMETHICONE

* Lotn 4%	.25 200 ml OP	✓ <u>healthE</u> <u>Dimethicone 4%</u> <u>Lotion</u>
IVERMECTIN - Special Authority see SA2294 below - Retail pharmacy		
Tab 3 mg - Up to 100 tab available on a PSO17.	.20 4	✓ Stromectol

- 1) PSO for institutional use only. Must be endorsed with the name of the institution for which the PSO is required and a valid Special Authority for patient of that institution.
- 2) Ivermectin available on BSO provided the BSO includes a valid Special Authority for a patient of the institution.
- For the purposes of subsidy of ivermectin, institution means age related residential care facilities, disability care facilities or prisons.

### ⇒SA2294 Special Authority for Subsidy

Initial application — (Scables) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

Either:

- 1 The person has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
- 2 Both:
  - 2.1 The person has a confirmed diagnosis of scabies or is a close contact of a scabies case; and
  - 22 Fither
    - 2.2.1 The person is unable to complete topical therapy; or
    - 2.2.2 Previous treatment with topical therapy has been tried and not cleared the infestation.

**Initial application — (Other parasitic infections)** from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 filariasis: or
- 2 cutaneous larva migrans (creeping eruption); or
- 3 strongyloidiasis.

**Renewal — (Scables)** from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria: Either:

- 1 The person has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
- 2 Both:
  - 2.1 The person has a confirmed diagnosis of scabies or is a close contact of a scabies case; and
  - 2.2 Fither:
    - 2.2.1 The person is unable to complete topical therapy; or
    - 2.2.2 Previous treatment with topical therapy has been tried and not cleared the infestation.

Renewal — (Other parasitic infections) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 filariasis: or
- 2 cutaneous larva migrans (creeping eruption); or
- 3 strongyloidiasis.

#### **PERMETHRIN**

# **Psoriasis and Eczema Preparations**

ACITRETIN - Special Authority see SA2024 below - Retail p	narmacy		
Cap 10 mg	26.20	60	Novatretin
Cap 25 mg		60	✓ Novatretin

### ⇒SA2024 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around acitretin and is competent to prescribe acitretin; and
- 3 Either:
  - 3.1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and that they must not become pregnant during treatment and for a period of three years after the completion of treatment; or

Subsidy		Fully	Brand or
(Manufacturer's Price) Subs		lised	Generic
 \$	Per	✓	

continued...

3.2 Patient is not of child bearing potential.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Fither:

- 1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and that they must not become pregnant during treatment and for a period of three years after the completion of treatment; or
- 2 Patient is not of child bearing potential.

### BETAMETHASONE DIPROPIONATE WITH CALCIPOTRIOL

Foam spray 500 mcg with calcipotriol 50 mcg per g	40.92	60 g OP 60 g OP 30 g OP	✓ Enstilar ✓ <u>Daivobet</u> ✓ <u>Daivobet</u>
CALCIPOTRIOL Oint 50 mcg per g	40.00	120 g OP	✓ Daivonex
COAL TAR Soln BP - Only in combination	36.25	200 ml	✓ Midwest
1) I In to 100/ only in combination with a dermatalogical I	haaa ar nranr	intani Taninal C	'articoptoriod D

- 1) Up to 10% only in combination with a dermatological base or proprietary Topical Corticosteriod Plain
- 2) With or without other dermatological galenicals.

#### COAL TAR WITH ALLANTOIN, MENTHOL, PHENOL AND SULPHUR

Soln 5% with sulphur 0.5%, menthol 0.75%, phenol 0.5% and			
allantoin crm 2.5%	6.59	75 g OP	
	(8.00)	•	Egopsoryl TA
	3.43	30 g OP	
	(4.35)		Egopsoryl TA
COAL TAR WITH SALICYLIC ACID AND SULPHUR			
Soln 12% with salicylic acid 2% and sulphur 4% oint	4.97	25 g OP	✓ Coco-Scalp
• ,	7.95	40 g OP	✓ Coco-Scalp

PIMECROLIMUS - Special Authority see SA1970 below - Retail pharmacy

- a) Maximum of 15 g per prescription
- b) Note: a maximum of 15 g per prescription and no more than one prescription per 12 weeks.

### ⇒SA1970 Special Authority for Subsidy

**Initial application** only from a dermatologist, paediatrician, ophthalmologist or any relevant practitioner on the recommendation of a dermatologist, paediatrician or ophthalmologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Both:

- 1 Patient has atopic dermatitis on the eyelid; and
- 2 Patient has at least one of the following contraindications to topical corticosteroids: periorificial dermatitis, rosacea, documented epidermal atrophy, documented allergy to topical corticosteroids, cataracts, glaucoma, or raised intraocular pressure.

#### PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORESCEIN - Only on a prescription

\* Soln 2.3% with trolamine laurilsulfate and fluorescein sodium..........5.41 500 ml / Pinetarsol

	Subsidy (Manufacturer's Pric	ce) Sub Per	Fully sidised	Brand or Generic Manufacturer
SALICYLIC ACID Powder – Only in combination	18.88	250 g	✓ N	lidwest
<ol> <li>Only in combination with a dermatological base or</li> <li>With or without other dermatological galenicals.</li> </ol>	proprietary Topical	Corticoster	oid – Pla	ain or collodion flexible
SULPHUR				
Precipitated - Only in combination		100 g		lidwest
<ol> <li>Only in combination with a dermatological base or</li> <li>With or without other dermatological galenicals.</li> </ol>	proprietary Topical	l Corticoster	oid – Pla	ain
TACROLIMUS				
Oint 0.1% - Special Authority see SA2074 below - Retail pharmacy	33.00	30 g OP	<b>√</b> <u>Z</u>	<u>'ematop</u>
<ul><li>a) Maximum of 30 g per prescription</li><li>b) Note: a maximum of 30 g per prescription and no m</li></ul>	ore than one presc	ription per 1	2 weeks	S.
⇒SA2074 Special Authority for Subsidy	p. 000			•
<b>Initial application</b> only from a dermatologist, paediatrician or an paediatrician, . Approvals valid without further renewal unless n				
Both:				
<ul><li>1 Patient has atopic dermatitis on the face; and</li><li>2 Patient has at least one of the following contraindications</li></ul>	to tonical corticost	eroids: neri	orificial d	dermatitis rosacea
documented epidermal atrophy or documented allergy to			ormoiar (	Joinnattio, roodood,
Scalp Preparations				
BETAMETHASONE VALERATE				
* Scalp app 0.1%	12.95	100 ml OP	<b>✓</b> E	Beta Scalp
Beta Scalp to be Principal Supply on 1 February 2025				
CLOBETASOL PROPIONATE			_	
* Scalp app 0.05%	6.26	30 ml OP	✓ [	<u> Permol</u>
HYDROCORTISONE BUTYRATE				
Scalp lotn 0.1%	6.57	100 ml OP	✓ L	ocoid
KETOCONAZOLE				
Shampoo 2%		100 ml OP	_	<u>Sebizole</u>
a) Maximum of 100 mlinti	4.09		<b>✓</b> S	<u>Sebizole</u>
<ul><li>a) Maximum of 100 ml per prescription</li><li>b) Only on a prescription</li></ul>				
Sunscreens				
SUNSCREENS, PROPRIETARY – Subsidy by endorsement				
Only if prescribed for a patient with severe photosensitivity sendorsed accordingly.	econdary to a defir	ned clinical c	ondition	and the prescription is

✓ Marine Blue Lotion SPF 50+

200 g OP

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic S Per ✓ Manufacturer

# **Wart Preparations**

For salicylic acid preparations refer to PSORIASIS AND ECZEMA PREPARATIONS, page 76

#### **PODOPHYLLOTOXIN**

- a) Maximum of 3.5 ml per prescription
- b) Only on a prescription

# **Other Skin Preparations**

# **Antineoplastics**

# **IMIQUIMOD**

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	•	Manufacturer

Subsidy Fully Brand or
(Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

# **Contraceptives - Non-hormonal**

# **Condoms**

	NDOMS			_
	49 mm - Up to 144 dev available on a PSO		144	✓ Moments
•	53 mm		10	✓ Moments
		14.25	144	✓ Moments
	a) Maximum of 60 dev per prescription			
	b) Up to 60 dev available on a PSO			
÷	53 mm, 0.05 mm thickness		10	✓ Moments
		14.25	144	✓ Moments
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription		40	
•	53 mm, chocolate, brown		10	✓ Moments
	\	14.25	144	✓ Moments
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription	4.45	10	/ Namenta
+	53 mm, strawberry, red		10	✓ Moments
	a) The to CO day available are a DOO	14.25	144	✓ Moments
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription	1.15	10	✓ Moments
÷	56 mm	1.15 14.50	10 144	✓ Moments ✓ Moments
	a) Manianum of CO days are a significan	14.50	144	▼ Infoments
	a) Maximum of 60 dev per prescription			
	b) Up to 60 dev available on a PSO	0.00	10	✓ Cold Maint
+	56 mm, 0.05 mm thickness	2.00 24.10	12 144	✓ Gold Knight ✓ Gold Knight
	a) Unite 60 day available are a DOO	24.10	144	• Gold Killgill
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription	20.17	144	Cold Knight
•	56 mm, 0.05mm thickness (bulk pack)	20.17	144	✓ Gold Knight
	a) Maximum of 60 dev per prescription			
	b) Up to 60 dev available on a PSO	1.15	10	✓ Moments
+	56 mm, 0.08 mm thickness		10 144	✓ Moments ✓ Moments
	a) The ta CO day available as a DCO	14.25	144	▼ Woments
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription	1.15	10	./ Mamanta
+	56 mm, 0.08 mm thickness, red		10 144	✓ Moments ✓ Moments
	a) The to CO day available are a DOO	14.25	144	▼ Infoments
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription	1.70	10	✓ Cold Maint
÷	56 mm, chocolate	1./9 21.45	12 144	✓ Gold Knight ✓ Gold Knight
	a) The ta CO day available as a DCO	∠1.45	144	• Gold Knight
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription	1.70	12	✓ Cold Knight
+	56 mm, strawberry	1.79 21.45	12 144	✓ Gold Knight
	a) The ta CO day available as a DCO	∠1.45	144	✓ Gold Knight
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription	1.00	10	✓ Cold Knight VI
÷	60 mm	1.82	12 144	<ul><li>✓ Gold Knight XL</li><li>✓ Gold Knight XL</li></ul>
	a) Manianum of CO days are a significan	∠1.09	144	▼ Gold Kilight XL
	a) Maximum of 60 dev per prescription			

Subsidy (Manufacturer's Price)	S	Fully subsidised	Brand or Generic	
\$	Per	✓	Manufacturer	

- a) Maximum of 60 dev per prescription
- b) Up to 60 dev available on a PSO

## Contraceptive Devices

#### INTRA-UTERINE DEVICE

- a) Up to 40 dev available on a PSO
- b) Only on a PSO

*	IUD 29.1 mm length × 23.2 mm width	1	✓ Choice 380 7med
			Nsha Silver/
			copper Short
*	IUD 33.6 mm length × 29.9 mm width26.80	1	✓ TCu 380 Plus
			<u>Normal</u>
*	IUD 35.5 mm length × 19.6 mm width	1	<ul> <li>Cu 375 Standard</li> </ul>

# **Contraceptives - Hormonal**

# **Combined Oral Contraceptives**

# **⇒SA0500** Special Authority for Alternate Subsidy

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Either:
  - 1.1 Patient is on a Social Welfare benefit; or
  - 1.2 Patient has an income no greater than the benefit; and
- 2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient is on a Social Welfare benefit; or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- · on a Social Welfare benefit; or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

#### ETHINYLOESTRADIOL WITH DESOGESTREL

\* Tab 20 mcg with desogestrel 150 mcg and 7 inert tab - Up to ✓ Mercilon 28 84 tab available on a PSO......10.00 84

			Fulls Described
	Subsidy		Fully Brand or
	(Manufacturer's Price)	Subsi	
	\$	Per	✓ Manufacturer
THINYLOESTRADIOL WITH LEVONORGESTREL			
Tab 20 mcg with levonorgestrel 100 mcg and 7 inert tablets -			
Up to 84 tab available on a PSO	1.50	84	✓ Lo-Oralcon 20 ED
Fab 30 mcg with levonorgestrel 150 mcg	6.62	63	
	(16.50)		Microgynon 30
<ul> <li>a) Higher subsidy of \$15.00 per 63 tab with Special Author</li> <li>b) Up to 63 tab available on a PSO</li> <li>Tab 30 mcg with levonorgestrel 150 mcg and 7 inert tablets – Up to 84 tab available on a PSO</li> </ul>		the previo	ous page  ✓ Oralcon 30 ED
THINYLOESTRADIOL WITH NORETHISTERONE			
Tab 35 mcg with norethisterone 1 mg and 7 inert tab	12.25	84	✓ Alyacen S29
•			✓ Brevinor 1/28
<ul> <li>a) Brand switch fee payable (Pharmacode 2692112) - se</li> <li>b) Up to 84 tab available on a PSO</li> <li>Tab 35 mcg with norethisterone 500 mcg and 7 inert tab - Up</li> </ul>		ls	
to 84 tab available on a PSO		84	✓ Norimin

# **Progestogen-only Contraceptives**

#### ⇒SA0500 Special Authority for Alternate Subsidy

**Initial application** from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Fither:
  - 1.1 Patient is on a Social Welfare benefit; or
  - 1.2 Patient has an income no greater than the benefit; and
- 2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Fither:

- 1 Patient is on a Social Welfare benefit: or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- on a Social Welfare benefit: or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

### LEVONORGESTREL

*	Tab 30 mcg - Up to 84 tab available on a PSO16.	50	84	✓ Microlut
	22.	00 1	112	✓ Microlut
*	Subdermal implant (2 $\times$ 75 mg rods) – Up to 3 pack available			
	on a PSO106.	92	1	✓ <u>Jadelle</u>
ME	DROXYPROGESTERONE ACETATE			
	Inj 150 mg per ml, 1 ml syringe - Up to 5 inj available on a PSO9.	18	1	✓ Depo-Provera

	Subsidy (Manufacturer's Price) \$	Sub Per	Fully sidised	
NORETHISTERONE – Brand switch fee payable (Pharmacode 2 Tab 350 mcg – Up to 84 tab available on a PSO	,	273 for de 84	1	Norethinderone - CDC Noriday Noriday 28

# **Emergency Contraceptives**

LEVONORGESTREL

- - a) Maximum of 2 tab per prescription
  - b) Up to 5 tab available on a PSO
  - c) Note: Direct Provision by a pharmacist permitted under the provisions in Part I of Section A.

# **Antiandrogen Oral Contraceptives**

Prescribers may code prescriptions "contraceptive" (code "O") when used as indicated for contraception. The period of supply and prescription charge will be as per other contraceptives, as follows:

- A maximum \$5.00 prescription charge (patient co-payment) may apply.
- prescription may be written for up to six months supply.

Prescriptions coded in any other way are subject to any non contraceptive prescription charges that apply, and the non-contraceptive period of supply. ie. Prescriptions may be written for up to three months supply.

#### CYPROTERONE ACETATE WITH ETHINYLOESTRADIOL

# **Gynaecological Anti-infectives**

ACETIC ACID WITH HYDROXYQUINOLINE AND RICINOLEIC ACID		
Jelly with glacial acetic acid 0.94%, hydroxyquinoline sulphate		
0.025%, glycerol 5% and ricinoleic acid 0.75% with applicator8.43	100 g OP	
(24.87)		Aci-Jel
CLOTRIMAZOLE		
* Vaginal crm 1% with applicators	35 g OP	✓ Clomazol
* Vaginal crm 2% with applicators	20 g OP	✓ Clomazol
MICONAZOLE NITRATE		
* Vaginal crm 2% with applicator	40 g OP	✓ Micreme
NYSTATIN		
Vaginal crm 100,000 u per 5 g with applicator(s)5.70	75 g OP	✓ Nilstat

# Myometrial and Vaginal Hormone Preparations

EK	JOMETRINE MALEATE			
	Inj 500 mcg per ml, 1 ml ampoule - Up to 5 inj available on a			
	PSO	160.00	5	✓ DBL Ergometrine
OE:	STRIOL			-
*	Crm 1 mg per g with applicator	6.95	15 g OP	✓ Ovestin
*	Pessaries 500 mcg	7.55	15	✓ Ovestin

Cassette Pregnancy Test

	Subsidy (Manufacturer's Pr \$	ice) Subs	Fully sidised	Brand or Generic Manufacturer
OXYTOCIN – Up to 5 inj available on a PSO				
Inj 5 iu per ml, 1 ml ampoule	4.98	5	<b>√</b> (	Oxytocin BNM
Inj 10 iu per ml, 1 ml ampoule		5		Dxytocin BNM
	11.96	10	<b>√</b> 0	Oxytocin Panpharma
OXYTOCIN WITH ERGOMETRINE MALEATE – Up to 5 inj ava Inj 5 iu with ergometrine maleate 500 mcg per ml, 1 ml ampo		5	<b>√</b> <u>S</u>	Syntometrine
Pregnancy Tests - hCG Urine				
BETA-HCG LOW SENSITIVITY URINE TEST KIT – Up to 15 te Note: For use in abortion services only.	est available on a F	PSO PSO		
Midstream	16.28	1 test OP	✓ 0	CheckTop
PREGNANCY TESTS - HCG URINE				
a) Up to 200 test available on a PSO				
b) Only on a PSO				
Cassette	12.00	40 test OP	<b>√</b> S	Smith BioMed Rapid Pregnancy Test
	16.00		<b>✓</b> D	avid One Step

(Smith BioMed Rapid Pregnancy Test Cassette to be delisted 1 March 2025)

# **Urinary Agents**

For urinary tract Infections refer to INFECTIONS, Antibacterials, page 117

## 5-Alpha Reductase Inhibitors

FINASTERIDE – Special Authority see SA0928 below – Retail pharmacy

★ Tab 5 mg .......4.79 100 ✓ Ricit

# ⇒SA0928 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 Fither:
  - 2.1 The patient is intolerant of non-selective alpha blockers or these are contraindicated; or
  - 2.2 Symptoms are not adequately controlled with non-selective alpha blockers.

# Alpha-1A Adrenoreceptor Blockers

TAMSULOSIN HYDROCHLORIDE - Special Authority see SA1032 below - Retail pharmacy

# **⇒SA1032** Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 The patient is intolerant of non-selective alpha blockers or these are contraindicated.

	Subsidy (Manufacturer's I \$	Price) Subs Per	Fully Brand or idised Generic ✓ Manufacturer
Other Urinary Agents			
OXYBUTYNIN * Tab 5 mg	5.42	100	✓ Alchemy Oxybutynin
POTASSIUM CITRATE			
Oral liq 3 mmol per ml – Special Authority see SA1083 be Retail pharmacy		200 ml OP	✓ Biomed
SA1083 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals v Both:	alid for 12 months	for applications	meeting the following criteria:
1 The patient has recurrent calcium oxalate urolithiasis; a		a application	
2 The patient has had more than two renal calculi in the t Renewal from any relevant practitioner. Approvals valid for 2 benefitting from the treatment.		• • •	s appropriate and the patient is
SODIUM CITRO-TARTRATE  * Grans eff 4 g sachets	3.50	28	✓ <u>Ural</u>
SOLIFENACIN SUCCINATE			
Tab 5 mg Tab 10 mg		30 30	<ul><li>✓ Solifenacin Viatris</li><li>✓ Solifenacin Viatris</li></ul>
Detection of Substances in Urine			
ORTHO-TOLIDINE			
* Compound diagnostic sticks	7.50 (8.25)	50 test OP	Hemastix
TETRABROMOPHENOL			
* Blue diagnostic strips	13.92	100 test OP	✓ Albustix
Obstetric Preparations			
Antiprogesterones			

Tab 200 mg - Up to 15 tab available on a PSO......79.90

✓ Mifegyne✓ Mifegyne

3

180.00

MIFEPRISTONE

Subsidy

Fully

Brand or

	(Manufacturer's Price)	Subsidised		Generic	
	\$	Per		Manufacturer	
Calcium Homeostasis					
CALCITONIN					

CALCITONIN			
* Inj 100 iu per ml, 1 ml ampoule	121.00	5	✓ Miacalcic
CINACALCET - Special Authority see SA2170 below - Retail	pharmacy		
Tab 30 mg - Wastage claimable	25.24	28	✓ Cinacalet Devatis
Tab 60 mg - Wastage claimable	50.47	28	✓ Cinacalet Devatis

⇒SA2170 Special Authority for Subsidy

Initial application — (parathyroid carcinoma or calciphylaxis) only from a nephrologist or endocrinologist. Approvals valid for 6 months for applications meeting the following criteria: Either:

- 1 All of the following:
  - 1.1 The patient has been diagnosed with a parathyroid carcinoma (see Note); and
  - 1.2 The patient has persistent hypercalcaemia (serum calcium greater than or equal to 3 mmol/L) despite previous first-line treatments including sodium thiosulfate (where appropriate) and bisphosphonates; and
  - 1.3 The patient is symptomatic; or
- 2 All of the following:
  - 2.1 The patient has been diagnosed with calciphylaxis (calcific uraemic arteriolopathy); and
  - 2.2 The patient has symptomatic (e.g. painful skin ulcers) hypercalcaemia (serum calcium greater than or equal to 3 mmol/L): and
  - 2.3 The patient's condition has not responded to previous first-line treatments including bisphosphonates and sodium

Renewal — (parathyroid carcinoma or calciphylaxis) only from a nephrologist or endocrinologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient's serum calcium level has fallen to < 3mmol/L; and
- 2 The patient has experienced clinically significant symptom improvement.

Note: This does not include parathyroid adenomas unless these have become malignant.

Initial application — (primary hyperparathyroidism) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### All of the following:

- 1 Patient has primary hyperparathyroidism; and
- 2 Either:
  - 2.1 Patient has hypercalcaemia of more than 3 mmol/L with or without symptoms; or
  - 2.2 Patient has hypercalcaemia of more than 2.85 mmol/L with symptoms; and
- 3 Surgery is not feasible or has failed; and
- 4 Patient has other comorbidities, severe bone pain, or calciphylaxis.

Initial application — (secondary or tertiary hyperparathyroidism) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 Patient has tertiary hyperparathyroidism and markedly elevated parathyroid hormone (PTH) with hypercalcaemia:
  - 1.2 Patient has symptomatic secondary hyperparathyroidism and elevated PTH; and
- 2 Patient is on renal replacement therapy; and
- 3 Any of the following:
  - 3.1 Residual parathyroid tissue has not been localised despite repeat unsuccessful parathyroid explorations; or

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per		Manufacturer

continued...

- 3.2 Parathyroid tissue is surgically inaccessible; or
- 3.3 Parathyroid surgery is not feasible.

**Renewal — (secondary or tertiary hyperparathyroidism)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

#### Fither:

- 1 The patient has had a kidney transplant, and following a treatment free interval of at least 12 weeks a clinically acceptable parathyroid hormone (PTH) level to support ongoing cessation of treatment has not been reached; or
- 2 The patient has not received a kidney transplant and trial of withdrawal of cinacalcet is clinically inappropriate.

# ZOLEDRONIC ACID

# Corticosteroids and Related Agents for Systemic Use

BETAMETHASONE SODIUM PHOSPHATE WITH BETAMETHASONE ACETAT	ΓΕ	
* Inj 3.9 mg with betamethasone acetate 3 mg per ml, 1 ml	5	
(36.96)		Celestone
()		Chronodose
DEXAMETHASONE		
	00	✓ Dexmethsone
* Tab 0.5 mg – Up to 60 tab available on a PSO1.80	30	Dexmetnsone
Dexmethsone to be Principal Supply on 1 February 2025		4.5
* Tab 4 mg - Up to 30 tab available on a PSO	30	<ul><li>Dexmethsone</li></ul>
Dexmethsone to be Principal Supply on 1 February 2025		
Oral liq 1 mg per ml52.80	25 ml OP	✓ Biomed
DEXAMETHASONE PHOSPHATE		
Dexamethasone phosphate injection will not be funded for oral use.		
* Inj 4 mg per ml, 1 ml ampoule – Up to 5 inj available on a PSO7.86	10	✓ Hameln
* Inj 4 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO 13.10	10	✓ Hameln
FLUDROCORTISONE ACETATE		
	100	./ Flavinof
* Tab 100 mcg11.46	100	✓ <u>Florinef</u>
HYDROCORTISONE		
* Tab 5 mg8.10	100	Douglas
* Tab 20 mg	100	<ul><li>Douglas</li></ul>
* Inj 100 mg vial	1	✓ Solu-Cortef
a) Not on a BSO		
b) Up to 5 inj available on a PSO		
METHYLPREDNISOLONE		
	100	✓ Medrol
· · · · · · · · · · · · · · · · · · ·	20	✓ Medrol
* Tab 100 mg223.10	20	• iviedroi
METHYLPREDNISOLONE (AS SODIUM SUCCINATE)		
Inj 40 mg vial22.30	1	✓ Solu-Medrol-Act-
		O-Vial
Inj 125 mg vial34.10	1	✓ Solu-Medrol-Act-
		O-Vial
lat 500 avantal		Cooks Marked Ask
Inj 500 mg vial26.88	1	✓ Solu-Medrol-Act-
		O-Vial
lai 1 a viol	4	✓ Solu-Medrol
Inj 1 g vial32.84	1	♥ Solu-Medroi

	Subsidy		Fully Brand or
	(Manufacturer's Price \$	e) S Per	Subsidised Generic  Manufacturer
	Ψ	rei	Wallulaciulei
IETHYLPREDNISOLONE ACETATE	47.00	-	( Danie Maduel
Inj 40 mg per ml, 1 ml vial	47.06	5	✓ Depo-Medrol
REDNISOLONE			
<ul> <li>Oral liq 5 mg per ml – Up to 30 ml available on a PSO Restricted to children under 12 years of age.</li> </ul>	6.00	30 ml OF	
REDNISONE			
F Tab 1 mg	18.58	500	✓ Prednisone Clinect
Tab 2.5 mg	21.04	500	✓ Prednisone Clinect
Tab 5 mg - Up to 30 tab available on a PSO	19.30	500	✓ Prednisone Clinect
Tab 20 mg - Up to 30 tab available on a PSO		500	✓ Prednisone Clinect
ETRACOSACTRIN			
F Inj 250 mcg per ml, 1 ml ampoule	86 25	1	✓ Synacthen
ing 200 may per mi, i mi ampoule	00.20	1	✓ UK Synacthen
Inj 1 mg per ml, 1 ml ampoule	600.00	1	✓ Synacthen Depot
ing ing permi, rim ampoule	090.00	1	✓ Synacthene
			Retard S29
RIAMCINOLONE ACETONIDE			
Inj 10 mg per ml, 1 ml ampoule		5	✓ Kenacort-A 10
Inj 40 mg per ml, 1 ml ampoule	52.63	5	✓ Kenacort-A 40
Sex Hormones Non Contraceptive			
Sex Hormones Non Contraceptive			
Androgen Agonists and Antagonists			
YPROTERONE ACETATE			
Tab 50 mg	14.37	50	✓ Siterone
Tab 100 mg	28.03	50	✓ Siterone
ESTOSTERONE			
Gel (transdermal) 16.2 mg per g	52.00	88 g OP	✓ Testogel
		55 g 51	<u> </u>
ESTOSTERONE CIPIONATE	05.00	4	/ Dama Tastast
Inj 100 mg per ml, 10 ml vial	85.00	1	✓ Depo-Testosterone
ESTOSTERONE ESTERS			
Inj 250 mg per ml, 1 ml	12.98	1	<ul> <li>Sustanon Ampoules</li> </ul>
ESTOSTERONE UNDECANOATE			
Cap 40 mg - Subsidy by endorsement	36.00	100	✓ Steril-Gene S29
Subsidy by endorsement – subsidised for patients who		rone und	decanoate cap 40mg prior to
1 November 2021 and the prescription is endorsed ac			
where there exists a record of prior dispensing of testo			
	00.00		

Inj 250 mg per ml, 4 ml vial......86.00

✓ Reandron 1000

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic 

\$ Per ✔ Manufacturer

# Hormone Replacement Therapy - Systemic

# **Oestrogens**

OE	STRADIOL			
*	Tab 1 mg	4.12	28 OP	
		(11.10)		Estrofem
*	Tab 2 mg	4.12	28 OP	
		(11.10)		Estrofem
	Gel (transdermal) 0.06% (750 mcg/actuation)	14.25	80 g OP	✓ Estrogel
	Patch 25 mcg per day	9.85	8	<ul><li>Estradiol TDP Mylan</li></ul>
		13.50		✓ Estraderm MX S29
		14.50		✓ Estradot
		21.35		✓ Lyllana
	a) No more than 2 patch per week			•
	b) Only on a prescription			
	Patch 50 mcg per day	10.75	8	✓ Estradiol TDP Mylan
			-	✓ Estradiol Viatris
		14.50		✓ Estraderm MX \$29
		14.50		✓ Estradiol Sandoz
				✓ Estradot
		21.55		✓ Lyllana
	a) No mare than 2 noted nor wools	21.00		Lylialia
	a) No more than 2 patch per week			
	b) Only on a prescription	11.00	0	✓ Estrodial TDD Mides
	Patch 75 mcg per day	11.00	8	<ul><li>✓ Estradiol TDP Mylan</li><li>✓ Estradiol Viatris</li></ul>
		14.50		✓ Estradiol Viatris ✓ Estradiol Sandoz
		14.50		✓ Estradioi Sandoz ✓ Estradot
		00.07		
	\	22.37		✓ Lyllana
	a) No more than 2 patch per week			
	b) Only on a prescription	40.05	•	4 5 · "   TDD
	Patch 100 mcg per day	12.95	8	✓ Estradiol TDP Mylan
		44.50		✓ Estradiol Viatris
		14.50		✓ Estradiol Sandoz
				✓ Estradot
		15.50		✓ Estraderm MX S29
		22.77		<ul><li>Lyllana</li></ul>
	<ul> <li>a) No more than 2 patch per week</li> </ul>			
	b) Only on a prescription			
OE	STRADIOL VALERATE			
*	Tab 1 mg	12.36	84	✓ Progynova
	Tab 2 mg		84	✓ Progynova
	STROGENS			<b>~.</b>
*		2 01	28	
~	Ourjugated, equilie tab 500 fileg	(19.25)	20	Premarin
*	Conjugated, equine tab 625 mcg		28	i iemami
~	Ourjugated, equilie tab 020 mcg	(19.25)	20	Premarin
		(19.20)		FICHIANN

	Outstale.		Fulls Decades
	Subsidy (Manufacturer's Price)	Subs	Fully Brand or sidised Generic
	\$	Per	✓ Manufacturer
Progestogens			
EDROXYPROGESTERONE ACETATE			
F Tab 2.5 mg	6.56	30	✓ Provera
	8.75	56	✓ Provera
Tab 5 mg		56	✓ Provera
Tab 10 mg	20.13	100	✓ Provera
Tab 10 mg	10.28	30	✓ Provera
Progestogen and Oestrogen Combined Prepara	ntions		
ESTRADIOL WITH NORETHISTERONE			
Tab 1 mg with 0.5 mg norethisterone acetate		28 OP	
Tab 2 mg with 1 mg norethisterone acetate	(18.10)		Kliovance
Tab 2 mg with 1 mg norethisterone acetate		28 OP	IZP I
T10 314 315 55 55	(18.10)		Kliogest
Tab 2 mg with 1 mg norethisterone acetate (10), and 2 mg	F 40	00.05	
oestradiol tab (12) and 1 mg oestradiol tab (6)		28 OP	Tricoguese
	(18.10)		Trisequens
Other Oestrogen Preparations			
ESTRIOL			
Tab 2 mg	7.70	30	✓ Ovestin
Other Progestogen Preparations			
EVONORGESTREL			
Intra-uterine device 52 mg	269 50	1	✓ Mirena
Intra-uterine device 13.5 mg		1	✓ Jaydess
EDROXYPROGESTERONE ACETATE			,
Tab 100 mg	133.57	100	✓ Provera HD
DRETHISTERONE - Brand switch fee payable (Pharmacode 2			
Tab 5 mg - Up to 30 tab available on a PSO		30	✓ Primolut N
ROGESTERONE	- · · •		
Cap 100 mg	14.85	30	✓ Utrogestan
			<u> </u>
hyroid and Antithyroid Agents			
ARBIMAZOLE			
Tab 5 mg	7.56	100	✓ Neo-Mercazole
-		.00	1100 morouzoro
VOTHYROXINE Tab 25 mcg	5 55	90	✓ Synthroid
Tab 50 mcg		28	✓ Mercury Pharma
Tab 50 mcg	5.79	90	✓ Synthroid
	64.28	1,000	✓ Eltroxin
Tablet 50 mcg		200	✓ Eltroxin
Tab 100 mcg		28	✓ Mercury Pharma
-	6.01	90	✓ Synthroid
	66.78	1,000	✓ Eltroxin
Tablet 100 mcg	13.36	200	✓ Eltroxin

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
PROPYLTHIOURACIL - Special Authority see SA1199 below -	Retail pharmacy			
Tab 50 mg	35.00	100	<b>√</b> P	TU \$29
TO CA1100 Chaniel Authority for Cubaidy				

**⇒SA1199** Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 The patient has hyperthyroidism; and
- 2 The patient is intolerant of carbimazole or carbimazole is contraindicated.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

# **Trophic Hormones**

#### **Growth Hormones**

SOMATROPIN (OMNITROPE) – Special Authority see SA2032 be	•	armacy	/ Omnitusus
* Inj 5 mg cartridge	80.21	ı	Omnitrope
			✓ Omnitrope S29 S29
Omnitrope to be Principal Supply on 1 February 2025			
* Inj 10 mg cartridge	80.21	1	✓ Omnitrope
			✓ Omnitrope S29 S29
Omnitrope to be Principal Supply on 1 February 2025			•
* Inj 15 mg cartridge	139.50	1	✓ Omnitrope
, ,			✓ Omnitrope S29 S29
Omnitrope to be Principal Supply on 1 February 2025			•
(Omnitrope S29 S29 Inj 5 mg cartridge to be delisted 1 February 2	2025)		
(Omnitrope S29 S29 Inj 10 mg cartridge to be delisted 1 February	(2025)		

#### ⇒SA2032 Special Authority for Subsidy

(Omnitrope S29 S29 Inj 15 mg cartridge to be delisted 1 February 2025)

Initial application — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

Fither:

- 1 Growth hormone deficiency causing symptomatic hypoglycaemia, or with other significant growth hormone deficient sequelae (e.g. cardiomyopathy, hepatic dysfunction) and diagnosed with GH < 5 mcg/l on at least two random blood samples in the first 2 weeks of life, or from samples during established hypoglycaemia (whole blood glucose < 2 mmol/l using a laboratory device); or</p>
- 2 All of the following:
  - 2.1 Height velocity < 25th percentile for age adjusted for bone age/pubertal status if appropriate over 6 or 12 months using the standards of Tanner and Davies (1985); and
  - 2.2 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
  - 2.3 Peak growth hormone value of < 5.0 mcg per litre in response to two different growth hormone stimulation tests. In children who are 5 years or older, GH testing with sex steroid priming is required; and</p>
  - 2.4 If the patient has been treated for a malignancy, they should be disease free for at least one year based upon follow-up laboratory and radiological imaging appropriate for the malignancy, unless there are strong medical reasons why this is either not necessary or appropriate; and
  - 2.5 Appropriate imaging of the pituitary gland has been obtained.

Renewal — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

(Manufacturer <sup>i</sup> s Price) Subsidised Generic \$ Per ✔ Manufacturer
---

continued...

- 1 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 2 Height velocity is greater than or equal to 25th percentile for age (adjusted for bone age/pubertal status if appropriate) while on growth hormone treatment, as calculated over six months using the standards of Tanner and Davis (1985); and
- 3 Height velocity is greater than or equal to 2.0 cm per year, as calculated over 6 months; and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a post-natal genotype confirming Turner Syndrome; and
- 2 Height velocity is < 25th percentile over 6-12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is < 14 years.

**Renewal — (Turner syndrome)** only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile for age (while on growth hormone calculated over 6 to 12 months using the Ranke's Turner Syndrome growth velocity charts); and
- 2 Height velocity is greater than or equal to 2 cm per year, calculated over six months; and
- 3 A current bone age is 14 years or under; and
- 4 No serious adverse effect that the specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 3 standard deviations below the mean for age or for bone age if there is marked growth acceleration or delay; and
- 2 Height velocity is < 25th percentile for age (adjusted for bone age/pubertal status if appropriate), as calculated over 6 to 12 months using the standards of Tanner and Davies(1985); and
- 3 A current bone age is < 14 years or under (female patients) or < 16 years (male patients); and
- 4 The patient does not have severe chronic disease (including malignancy or recognized severe skeletal dysplasia) and is not receiving medications known to impair height velocity.

**Renewal** — **(short stature without growth hormone deficiency)** only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred.

Initial application — (short stature due to chronic renal insufficiency) only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

1 The patient's height is more than 2 standard deviations below the mean; and

Subsidy	,	Fully	Brand or	
(Manufacturer's Price	·) :	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 2 Height velocity is < 25th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and</p>
- 3 A current bone age is to 14 years or under (female patients) or to 16 years or under (male patients); and
- 4 The patient is metabolically stable, has no evidence of metabolic bone disease and absence of any other severe chronic disease; and
- 5 The patient is under the supervision of a specialist with expertise in renal medicine; and
- 6 Either
  - 6.1 The patient has a GFR less than or equal to 30 ml/min/1.73m² as measured by the Schwartz method (Height(cm)/plasma creatinine (umol/l)) × 40 = corrected GFR (ml/min/1.73m²) in a child who may or may not be receiving dialysis; or
  - 6.2 The patient has received a renal transplant and has received < 5mg/ m²/day of prednisone or equivalent for at least 6 months...

Renewal — (short stature due to chronic renal insufficiency) only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not experienced significant biochemical or metabolic deterioration confirmed by diagnostic results; and
- 7 The patient has not received renal transplantation since starting growth hormone treatment; and
- 8 If the patient requires transplantation, growth hormone prescription should cease before transplantation and a new application should be made after transplantation based on the above criteria.

**Initial application — (Prader-Willi syndrome)** only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of Prader-Willi syndrome that has been confirmed by genetic testing or clinical scoring criteria;
- 2 The patient is aged six months or older; and
- 3 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
- 4 Sleep studies or overnight oximetry have been performed and there is no obstructive sleep disorder requiring treatment, or if an obstructive sleep disorder is found, it has been adequately treated under the care of a paediatric respiratory physician and/or ENT surgeon; and
- 5 Either:
  - 5.1 Both:
    - 5.1.1 The patient is aged two years or older; and
    - 5.1.2 There is no evidence of type II diabetes or uncontrolled obesity defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months; or
  - 5.2 The patient is aged between six months and two years and a thorough upper airway assessment is planned to be undertaken prior to treatment commencement and at six to 12 weeks following treatment initiation.

Renewal — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

Subsidy (Manufacturer's Price)	Sub	Fully	Brand or Generic	
 \$	Per	1	Manufacturer	

continued...

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not developed type II diabetes or uncontrolled obesity as defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months.

Initial application — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a medical condition that is known to cause growth hormone deficiency (e.g. surgical removal of the pituitary for treatment of a pituitary tumour); and
- 2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
- 3 The patient has severe growth hormone deficiency (see notes); and
- 4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
- 5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

Renewal — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 All of the following:
  - 1.1 The patient has been treated with somatropin for < 12 months; and
  - 1.2 There has been an improvement in Quality of Life defined as a reduction of at least 8 points on the Quality of Life Assessment of Growth Hormone Deficiency in Adults (QoL-AGHDA®) score from baseline; and
  - 1.3 Serum IGF-I levels have been increased within ±1SD of the mean of the normal range for age and sex; and
  - 1.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients, or 1 mg per day for female patients; or
- 2 All of the following:
  - 2.1 The patient has been treated with somatropin for more than 12 months; and
  - 2.2 The patient has not had a deterioration in Quality of Life defined as a 6 point or greater increase from their lowest QoL-AGHDA® score on treatment (other than due to obvious external factors such as external stressors); and
  - 2.3 Serum IGF-I levels have continued to be maintained within ±1SD of the mean of the normal range for age and sex (other than for obvious external factors); and
  - 2.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients or 1 mg per day for female patients; or

Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer	

continued...

- 3 All of the following:
  - 3.1 The patient has had a Special Authority approval for somatropin for childhood deficiency in children and no longer meets the renewal criteria under this indication; and
  - 3.2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
  - 3.3 The patient has severe growth hormone deficiency (see notes); and
  - 3.4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
  - 3.5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

of \$591.68 per 1 inj with Endorsement.......177.50

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

## **GnRH Analogues**

GOSEREI IN

Implant 3.6 mg, syringe	66.48	1	✓ Zoladex
Implant 10.8 mg, syringe	138.23	1	✓ Zoladex
LEUPRORELIN			
Additional subsidy by endorsement where the patient is a child or goserelin and the prescription is endorsed accordingly.	r adolescent a	and is unable	to tolerate administration of
Inj 3.75 mg prefilled dual chamber syringe - Higher subsidy of			
\$221.60 per 1 inj with Endorsement	66.48	1	
	(221.60)		Lucrin Depot 1-month
Inj 11.25 mg prefilled dual chamber syringe - Higher subsidy			

(591.68)

Vasopressin	Adonists
v asopi cssiii	Agomoto

DESMOPRESSIN			
Wafer 120 mcg	47.00	30	<ul><li>Minirin Melt</li></ul>
DESMOPRESSIN ACETATE			
Tab 100 mcg	25.00	30	✓ Minirin
Tab 200 mcg	54.45	30	✓ Minirin
▲ Nasal spray 10 mcg per dose	34.95	6 ml OP	✓ Desmopressin-
			PH&T
Ini 4 mcg per ml. 1 ml	67 18	10	✓ Minirin

Lucrin Depot 3-month

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	/	Manufacturer

# **Other Endocrine Agents**

#### **CABERGOLINE**

### ⇒SA2070 Special Authority for Waiver of Rule

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Hyperprolactinemia; or
- 2 Acromegaly\*; or
- 3 Inhibition of lactation.

Renewal — (for patients who have previously been funded under Special Authority form SA1031) from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has previously held a valid Special Authority which has expired and the treatment remains appropriate and the patient is benefiting from treatment.

Note: Indication marked with \* is an unapproved indication.

# CLOMIFENE CITRATE

Tab 50 mg	29.84	10	✓ Mylan Clomiphen S29
METYRAPONE Cap 250 mg	558.00	50	✓ Metopirone

	Subsidy		Fully Brand or	
	(Manufacturer's Price)		sidised Generic	
	\$	Per	✓ Manufacturer	
Anthelmintics				
ALBENDAZOLE - Special Authority see SA1318 below - Retail p	oharmacy			
Tab 400 mg		60	✓ Eskazole S29	
	403.20	00	LSRAZOIC 023	
■ SA1318 Special Authority for Subsidy Initial application only from an infectious disease specialist or cli patient has hydatids.	nical microbiologist.	Approval	s valid for 6 months where the	
Renewal only from an infectious disease specialist or clinical micr	rohiologist Annroys	ale valid fo	6 months where the treatment	
remains appropriate and the patient is benefitting from the treatme		alo valla lo	o montho where the treatment	
MEBENDAZOLE – Only on a prescription	J110.			
Tab 100 mg	5 10	6	√ Vormov	
Oral liq 100 mg per 5 ml		15 ml	✓ <u>Vermox</u>	
Oraciliq 100 mg per 5 mi	(7.83)	13 1111	Vermox	
PR 1 7 10 11 11 17 17 17 17 17 17 17 17 17 17 17	(7.00)		Vermox	
PRAZIQUANTEL	00.00	0	/ Dilledatela	
Tab 600 mg	68.00	8	✓ Biltricide	
Antibactoriala				
Antibacterials				
a) For topical antibacterials, refer to DERMATOLOGICALS, page	270			
b) For anti-infective eye preparations, refer to SENSORY ORGAI				
	10, page 200			
Cephalosporins and Cephamycins				
CEFACLOR MONOHYDRATE				
Cap 250 mg	25.85	100	✓ Ranbaxy-Cefaclor	
Grans for oral liq 125 mg per 5 ml – Wastage claimable		100 ml	✓ Ranbaxy-Cefactor	
CEFALEXIN			<u></u>	
Cap 250 mg	2.05	20	Conhalovin APM	
Cap 500 mg		20	<ul> <li>✓ Cephalexin ABM</li> <li>✓ Cephalexin ABM</li> </ul>	
Grans for oral lig 25 mg per ml – Wastage claimable		100 ml	✓ Flynn	
Grans for oral liq 50 mg per ml — Wastage claimable		100 ml	✓ Flynn	
Grans for oral liq 50 mg per mi – wastage claimable	11.75	100 1111	✓ Cefalexin Sandoz	
OFFATOLINI Och cid decent decent and	11.75		ocialexiii oailuoz	
CEFAZOLIN – Subsidy by endorsement				
Only if prescribed for dialysis or cellulitis in accordance with a	i Health INZ Hospital	approved	protocol and the prescription is	
endorsed accordingly. Inj 500 mg vial	2 20	5	✓ Cefazolin-AFT	
Inj 1 g vial		5	✓ Cefazolin-AFT	
Inj 2 g vial		5	✓ Cefazolin-AFT	
, 3	7.00	3	OCIAZOIII-AI I	
CEFTRIAXONE – Subsidy by endorsement				
a) Up to 10 inj available on a PSO				
b) Subsidised only if prescribed for a dialysis or cystic fibrosis patient, or the treatment of gonorrhoea, or the treatment of				
pelvic inflammatory disease, or the treatment of suspected	a meningococcai dis	sease, and	the prescription of PSO is	
endorsed accordingly. Inj 500 mg vial	0.70	1	✓ Ceftriaxone-AFT	
Inj 500 mg viai		1 5	✓ Ceftriaxone-AFT	
. •		J	• GEILLIANOLIE-AF I	
CEFUROXIME AXETIL – Subsidy by endorsement				
Only if prescribed for prophylaxis of endocarditis and the pres	scription is endorsed	according	lly.	

Tab 250 mg .......CBS

✓ Ascend-

Cefuroxime \$29

20

Subsidy	Fu	lly Brand or	
(Manufacturer's Price)	Subsidise	ed Generic	
\$	Per	<ul> <li>Manufacturer</li> </ul>	

#### **Macrolides**

AZITHROMYCIN – Maximum of 5 days treatment per prescription; can be waived by Special Authority see SA1683 below A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised on Special Authority.

Tab 250 mg8.19	30	✓ Apo-Azithromycin
Tab 500 mg - Up to 8 tab available on a PSO2.57		✓ Zithromax
Grans for oral liq 200 mg per 5 ml (40 mg per ml) - Wastage		
claimable	15 ml	✓ Zithromax

### ⇒SA1683 Special Authority for Waiver of Rule

Initial application — (bronchiolitis obliterans syndrome, cystic fibrosis and atypical Mycobacterium infections) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following:

- 1 Patient has received a lung transplant, stem cell transplant, or bone marrow transplant and requires treatment for bronchiolitis obliterans syndrome\*; or
- 2 Patient has received a lung transplant and requires prophylaxis for bronchiolitis obliterans syndrome\*; or
- 3 Patient has cystic fibrosis and has chronic infection with Pseudomonas aeruginosa or Pseudomonas-related gram negative organisms\*; or
- 4 Patient has an atypical Mycobacterium infection.

Note: Indications marked with \* are unapproved indications.

Initial application — (non-cystic fibrosis bronchiectasis\*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 For prophylaxis of exacerbations of non-cystic fibrosis bronchiectasis\*; and
- 2 Patient is aged 18 and under; and
- 3 Either:
  - 3.1 Patient has had 3 or more exacerbations of their bronchiectasis, within a 12 month period; or
  - 3.2 Patient has had 3 acute admissions to hospital for treatment of infective respiratory exacerbations within a 12 month period.

Note: Indications marked with \* are unapproved indications.

Renewal — (non-cystic fibrosis bronchiectasis\*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has completed 12 months of azithromycin treatment for non-cystic fibrosis bronchiectasis; and
- 2 Following initial 12 months of treatment, the patient has not received any further azithromycin treatment for non-cystic fibrosis bronchiectasis for a further 12 months, unless considered clinically inappropriate to stop treatment; and
- 3 The patient will not receive more than a total of 24 months' azithromycin cumulative treatment (see note).

The patient must not have had more than 1 prior approval.

Note: No further renewals will be subsidised. A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised. Indications marked with \* are unapproved indications

CLARITHROMYCIN – Maximum of 500 mg per prescription; can be waived by Special Authority see SA1857 below

**⇒SA1857** Special Authority for Waiver of Rule

Initial application — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years for applications meeting the following criteria:

Fither:

Subsidy	Fi	ully	Brand or
(Manufacturer's Price)	Subsidis	sed	Generic
\$	Per	✓	Manufacturer

continued...

- 1 Atypical mycobacterial infection; or
- 2 Mycobacterium tuberculosis infection where there is drug-resistance or intolerance to standard pharmaceutical agents.

**Initial application — (Helicobacter pylori eradication)** from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 For the eradication of helicobacter pylori in a patient unable to swallow tablets; and
- 2 For use only in combination with omeprazole and amoxicillin as part of a triple therapy regimen.

Initial application — (Prophylaxis of infective endocarditis) from any relevant practitioner. Approvals valid for 3 months where prophylaxis of infective endocarditis associated with surgical or dental procedures if amoxicillin is contra-indicated. Renewal — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

ERYTHROMYCIN (AS LACTOBIONATE) Inj 1 g vial	10.00	1	✓ Erythrocin IV
ERYTHROMYCIN ETHYL SUCCINATE			
Tab 400 mg	16.95	100	✓ E-Mycin
a) Up to 20 tab available on a PSO     b) Up to 2 x the maximum PSO quantity for RFPP Grans for oral liq 200 mg per 5 ml	5.00	100 ml	✓ E-Mycin
<ul> <li>a) Up to 300 ml available on a PSO</li> <li>b) Up to 2 x the maximum PSO quantity for RFPP</li> <li>c) Wastage claimable</li> </ul>			
Grans for oral liq 400 mg per 5 mla) Up to 200 ml available on a PSO b) Wastage claimable	6.77	100 ml	✓ E-Mycin
ROXITHROMYCIN			
Tab 150 mg	13.19	50	✓ <u>Arrow-</u> <u>Roxithromycin</u>
Tab 300 mg	25.00	50	✓ <u>Arrow-</u> <u>Roxithromycin</u>

	Subsidy		Fully	Brand or
	(Manufacturer's		idised	Generic
	\$	Per		Manufacturer
Penicillins				
AMOVICILLIN				
AMOXICILLIN	07.50	500		Mina Amazziaillia
Cap 250 mg	27.50	500	•	Miro-Amoxicillin
a) Up to 30 cap available on a PSO				
b) Up to 10 x the maximum PSO quantity for RFPP			_	
Cap 500 mg	41.00	500	•	<u> Miro-Amoxicillin</u>
<ul> <li>a) Up to 30 cap available on a PSO</li> </ul>				
<li>b) Up to 10 x the maximum PSO quantity for RFPP</li>				
Grans for oral liq 125 mg per 5 ml	2.22	100 ml	1	Alphamox 125
a) Up to 200 ml available on a PSO				
b) Wastage claimable				
Grans for oral liq 250 mg per 5 ml	2.81	100 ml	1	Alphamox 250
a) Up to 300 ml available on a PSO				
b) Up to 10 x the maximum PSO quantity for RFPP				
c) Wastage claimable				
Inj 250 mg vial	15 07	10	1	lbiamox
Inj 500 mg vial		10		Ibiamox
Inj 1 g vial – Up to 5 inj available on a PSO		10		Ibiamox
	21.04	10	•	IDIAIIIOX
AMOXICILLIN WITH CLAVULANIC ACID				
Tab 500 mg with clavulanic acid 125 mg - Up to 30 tab				
available on a PSO	1.59	10	1	Curam Duo 500/125
Grans for oral liq amoxicillin 25 mg with clavulanic acid 6.25	mg			
per ml		100 ml	1	Augmentin
a) Up to 200 ml available on a PSO				· ·
b) Wastage claimable				
Grans for oral liq amoxicillin 50 mg with clavulanic acid 12.5	ma			
per ml – Up to 200 ml available on a PSO		100 ml OP	1	Curam
por nii op to 200 nii available on a 1 00	5.61	100 1111 01		Amoxiclav Devatis
	5.01		•	Forte
(Curam Grana for aral lia amaviaillin 50 ma with alavulania anid 1	10 E ma nor ml t	a ha daliatad 1	luna a	
(Curam Grans for oral liq amoxicillin 50 mg with clavulanic acid 1	2.5 mg per mi u	o de delisted i c	iui i <del>e</del> 2	023)
BENZATHINE BENZYLPENICILLIN				
Inj 900 mg (1.2 million units) in 2.3 ml syringe - Up to 5 inj				
available on a PSO	375.97	10	✓	Bicillin LA
BENZYLPENICILLIN SODIUM [PENICILLIN G]				
Inj 600 mg (1 million units) vial – Up to 5 inj available on a P	SO 16.50	10	1	Sandoz
, , , ,	10.00	10		<u> </u>
FLUCLOXACILLIN	45.70	050	,	Floritorio (III) AFT
Cap 250 mg – Up to 30 cap available on a PSO		250		Flucloxacillin-AFT
Cap 500 mg – Up to 30 cap available on a PSO		500		Flucloxacillin-AFT
Grans for oral liq 25 mg per ml	4.89	100 ml	•	AFT
a) Up to 200 ml available on a PSO				
b) Wastage claimable				
<ul> <li>c) AFT to be Principal Supply on 1 February 2025</li> </ul>				
Grans for oral liq 50 mg per ml	5.89	100 ml	✓.	AFT
<ul> <li>a) Up to 200 ml available on a PSO</li> </ul>				
b) Wastage claimable				
c) AFT to be Principal Supply on 1 February 2025				
Inj 250 mg vial	42.60	10	1	Flucloxin
Inj 500 mg vial	45.63	10		Flucloxin
Inj 1 g vial – Up to 5 inj available on a PSO		5		Flucil
				_

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price) \$	) Si Per	Fully ubsidised	Brand or Generic Manufacturer
PHENOXYMETHYLPENICILLIN (PENICILLIN V)  Cap 250 mg – Up to 30 cap available on a PSO  Cilicaine VK to be Principal Supply on 1 February 2025	7.68	50	<b>√</b> (	Cilicaine VK
Cap 500 mg		50	✔ (	Cilicaine VK
Grans for oral liq 125 mg per 5 mla) Up to 200 ml available on a PSO b) Wastage claimable	3.40	100 ml	✓ <u>I</u>	<u>AFT</u>
Grans for oral liq 250 mg per 5 ml	4.24	100 ml	✓ <u>I</u>	<u>AFT</u>

# **Tetracyclines**

DO	XYCYCLINE			
*	Tab 100 mg - Up to 30 tab available on a PSO	64.43	500	Doxine
MΙΝ	NOCYCLINE HYDROCHLORIDE			
*	Tab 50 mg - Additional subsidy by Special Authority see			
	SA1355 below – Retail pharmacy	5.79	60	
		(12.05)		Mino-tabs
*	Cap 100 mg	19.32	100	
		(52.04)		Minomycin

# **⇒SA1355** Special Authority for Manufacturers Price

**Initial application** from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has rosacea.

TETRACYCLINE – Special Authority see SA1332 below – Retail pharmacy
Tab 250 mg ......58.20

### ⇒SA1332 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 For the eradication of helicobacter pylori following unsuccessful treatment with appropriate first-line therapy; and
- 2 For use only in combination with bismuth as part of a quadruple therapy regimen.

# Other Antibiotics

For topical antibiotics, refer to DERMATOLOGICALS, page 70

#### **CIPROFLOXACIN**

Recommended for patients with any of the following:

- i) microbiologically confirmed and clinically significant pseudomonas infection; or
- ii) prostatitis; or
- iii) pyelonephritis; or
- iv) gonorrhoea.

Tab 250 mg - Up to 5 tab available on a PSO	1.95	28	✓ Ipca-Ciprofloxacin
Tab 500 mg - Up to 5 tab available on a PSO	3.10	28	✓ Ipca-Ciprofloxacin
Tab 750 mg	4.80	28	✓ Ipca-Ciprofloxacin

	Subsidy		Fully Brand or	
	(Manufacturer's Price)		Subsidised Generic	
	\$	Per	✓ Manufacture	er
CLINDAMYCIN				
Cap hydrochloride 150 mg	4.94	24	✓ Dalacin C	
Inj 150 mg per ml, 4 ml ampoule	35.10	10	✓ <u>Hameln</u>	
COLISTIN SULPHOMETHATE - Retail pharmacy-Specialist - S	Subsidy by endorseme	nt		
Only if prescribed for dialysis or cystic fibrosis patient and the			accordingly.	
Inj 150 mg		1	Colistin-Link	
GENTAMICIN SULPHATE				
Inj 40 mg per ml, 2 ml vial - Subsidy by endorsement	36.70	5	✓ Cidomycin	
, , ,			P/Free S29	
Only if prescribed for a dialysis or cystic fibrosis patient of endorsed accordingly.	or complicated urinary	trac	t infection and the prescr	iption is
Inj 10 mg per ml, 1 ml ampoule - Subsidy by endorsement	95.00	5	✓ DBL Gentam	icin
Only if prescribed for a dialysis or cystic fibrosis patient of endorsed accordingly.			t infection and the prescr	iption is
Inj 10 mg per ml, 2 ml ampoule - Subsidy by endorsement	91.00	5	✓ Wockhardt	29
Only if prescribed for a dialysis or cystic fibrosis patient of endorsed accordingly.		trac	t infection and the prescr	iption is
Inj 40 mg per ml, 2 ml ampoule - Subsidy by endorsement	18.38	10	✓ Pfizer	
	91.90	50	<ul><li>Gentamicin</li></ul>	
			Noridem S2	9
Only if prescribed for a dialysis or cystic fibrosis patient of endorsed accordingly.	or complicated urinary	trac	t infection and the prescr	iption is
MOXIFLOXACIN - Special Authority see SA1740 below - Retail No patient co-payment payable	I pharmacy			
Tab 400 mg	42.00	5	✓ Avelox	

# ⇒SA1740 Special Authority for Subsidy

Initial application — (Tuberculosis) only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

#### 1 Both:

- 1.1 Active tuberculosis\*: and
- 1.2 Any of the following:
  - 1.2.1 Documented resistance to one or more first-line medications; or
  - 1.2.2 Suspected resistance to one or more first-line medications (tuberculosis assumed to be contracted in an area with known resistance), as part of regimen containing other second-line agents; or
  - 1.2.3 Impaired visual acuity (considered to preclude ethambutol use); or
  - 1.2.4 Significant pre-existing liver disease or hepatotoxicity from tuberculosis medications; or
  - 1.2.5 Significant documented intolerance and/or side effects following a reasonable trial of first-line medications; or
- 2 Mycobacterium avium-intracellulare complex not responding to other therapy or where such therapy is contraindicated.\*; or
- 3 Patient is under five years of age and has had close contact with a confirmed multi-drug resistant tuberculosis case.

Note: Indications marked with \* are unapproved indications.

**Renewal** only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Mycoplasma genitalium) only from a sexual health specialist or Practitioner on the recommendation of a sexual health specialist. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

Subsidy		Fully	Brand or	_
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1 Has nucleic acid amplification test (NAAT) confirmed Mycoplasma genitalium\* and is symptomatic; and
- 2 Either
  - 2.1 Has tried and failed to clear infection using azithromycin: or
  - 2.2 Has laboratory confirmed azithromycin resistance; and
- 3 Treatment is only for 7 days.

Initial application — (Penetrating eye injury) only from an ophthalmologist. Approvals valid for 1 month where the patient requires prophylaxis following a penetrating eye injury and treatment is for 5 days only.

Note: Indications marked with \* are unapproved indications.

PAROMOMYCIN - Special Authority see SA1689 below - Retail pharmacy

Cap 250 mg......126.00 16 **✔ Humatin** \$29

# ⇒SA1689 Special Authority for Subsidy

**Initial application** only from an infectious disease specialist, clinical microbiologist or gastroenterologist. Approvals valid for 1 month for applications meeting the following criteria:

#### Either:

- 1 Patient has confirmed cryptosporidium infection; or
- 2 For the eradication of Entamoeba histolyica carriage.

Renewal only from an infectious disease specialist, clinical microbiologist or gastroenterologist. Approvals valid for 1 month for applications meeting the following criteria:

#### Either:

- 1 Patient has confirmed cryptosporidium infection; or
- 2 For the eradication of Entamoeba histolyica carriage.

PYRIMETHAMINE - Special Authority see SA1328 below - Retail pharmacy

### ⇒SA1328 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 For the treatment of toxoplasmosis in patients with HIV for a period of 3 months; or
- 2 For pregnant patients for the term of the pregnancy; or
- 3 For infants with congenital toxoplasmosis until 12 months of age.

#### SODIUM FUSIDATE [FUSIDIC ACID]

#### SA1331 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 For the treatment of toxoplasmosis in patients with HIV for a period of 3 months; or
- 2 For pregnant patients for the term of the pregnancy; or
- 3 For infants with congenital toxoplasmosis until 12 months of age.

Subsidy		Fully	Brand or
•	-,		Generic Manufacturer
Ψ	1 01		Wandacturer
15.50	5	1	Tobramycin (Viatris)
a and prodompalom a	5 01140100	u uooo.u	9.7.
395.00	56 dose	1	Tobramycin BNM
orescription is end	orsed acc	ordingly.	
27.83	50	/	TMP
AZOLE]			
р			
115.74	500	1	Trisul
	100		Dameim
5.00	100 MI	•	Deprim
		(	atus and of Ola atuialissus
		or for the	aument of Clostridium
3.38	1919.	✓	<u>Mylan</u>
	(Manufacturer's Pric	(Manufacturer's Price)         S Per           \$         \$ Per	(Manufacturer's Price) Subsidised Per Subsidiary Su

# **Antifungals**

- a) For topical antifungals refer to DERMATOLOGICALS, page 71
- b) For topical antifungals refer to GENITO URINARY, page 84

#### **FLUCONAZOLE**

Cap 50 mg4.10	28	✓ Mylan
Cap 150 mg0.45	1	✓ Mylan
Cap 200 mg8.90	28	✓ Mylan
Powder for oral suspension 10 mg per ml - Special Authority		
see SA1359 below – Retail pharmacy129.02	35 ml	<ul><li>Diflucan</li></ul>
Wastage claimable		

#### ⇒SA1359 Special Authority for Subsidy

**Initial application** — **(Systemic candidiasis)** from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

#### Both:

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

**Initial application — (Immunocompromised)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

### All of the following:

- 1 Patient is immunocompromised; and
- 2 Patient is at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

**Renewal — (Systemic candidiasis)** from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

	Subsidy		Fully	Brand or
(Ma	anufacturer's Price)	Subsid	dised	Generic
	\$	Per	✓	Manufacturer

#### continued...

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

**Renewal — (Immunocompromised)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

#### All of the following:

- 1 Patient remains immunocompromised: and
- 2 Patient remains at moderate to high risk of invasive fungal infection; and

Tab 200 mg - PCT ......CBS

3 Patient is unable to swallow capsules.

#### **ITRACONAZOLE**

Cap 100 mg6.83	15	Itrazole
Oral lig 10 mg per ml - Special Authority see SA1322 below -		
Retail pharmacy141.80	150 ml OP	✓ Kent S29
		Sporanox

# ⇒SA1322 Special Authority for Subsidy

**Initial application** only from an infectious disease specialist, clinical microbiologist, clinical immunologist or any relevant practitioner on the recommendation of a infectious disease physician, clinical microbiologist or clinical immunologist. Approvals valid for 6 months where the patient has a congenital immune deficiency.

**Renewal** from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefitting from the treatment.

30

✓ Burel S29

#### **KETOCONAZOLE**

	100	✓ Strides Shasun \$29 ✓ Taro \$29 ✓ Teva- Ketoconazole \$29
NYSTATIN		
Tab 500,000 u14.16	50	
(17.09)		Nilstat
Cap 500,000 u12.81	50	
(15.47)		Nilstat
POSACONAZOLE - Special Authority see SA2383 below - Retail pharmacy		
Tab modified-release 100 mg206.00	24	✓ Posaconazole Juno
Oral liq 40 mg per ml342.51	105 ml OP	✓ <u>Devatis</u>

# ⇒SA2383 Special Authority for Subsidy

**Initial application** only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

#### Either:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation chemotherapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppressive therapy\*.

**Renewal** only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

Either:

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation therapy: or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppression\* and requires on going posaconazole treatment.

Note: \* Graft versus host disease (GVHD) on significant immunosuppression is defined as acute GVHD, grade II to IV, or extensive chronic GVHD, or if they were being treated with intensive immunosuppressive therapy consisting of either high-dose corticosteroids (1 mg or greater per kilogram of body weight per day for patients with acute GVHD or 0.8 mg or greater per kilogram every other day for patients with chronic GVHD), antithymocyte globulin, or a combination of two or more immunosuppressive agents or types of treatment.

Initial application — (Invasive fungal infection prophylaxis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient is at risk of invasive fungal infection; and
- 2 Either:
  - 2.1 Posaconazole is prescribed by, or recommended by a haematologist, transplant physician, infectious disease specialist, paediatric haematologist or paediatric oncologist; or
  - 2.2 Prescribing posaconazole is in accordance with a protocol or guideline that has been endorsed by the Health New Zealand - Te Whatu Ora Hospital in the specific settings where there is a greater than 10% risk of invasive fungal infection (IFI).

**Renewal — (Invasive fungal infection prophylaxis)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient is at risk of invasive fungal infection; and
- 2 Either:
  - 2.1 Posaconazole is prescribed by, or recommended by a haematologist, transplant physician, infectious disease specialist, paediatric haematologist or paediatric oncologist; or
  - 2.2 Prescribing posaconazole is in accordance with a protocol or guideline that has been endorsed by the Health New Zealand - Te Whatu Ora Hospital in the specific settings where there is a greater than 10% risk of invasive fungal infection (IFI).

#### **TERBINAFINE**

* Tab 250 mg8.9	97 84	✓ Deolate
VORICONAZOLE - Special Authority see SA2384 below - Retail pharmacy	/	
Tab 50 mg91.0	00 56	✓ Vttack
Tab 200 mg350.0	00 56	✓ Vttack
Powder for oral suspension 40 mg per ml - Wastage		
claimable1,523.2	22 70 ml	✓ Vfend

#### ⇒SA2384 Special Authority for Subsidy

Initial application — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
  - 3.1 Patient has proven or probable invasive aspergillus infection; or
  - 3.2 Patient has possible invasive aspergillus infection; or
  - 3.3 Patient has fluconazole resistant candidiasis; or

rsidy F Irer's Price) Subsidi	. ,	rand or eneric
 \$ Per	✓ Ma	anufacturer

continued...

3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

**Renewal — (invasive fungal infection)** only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
  - 3.1 Patient continues to require treatment for proven or probable invasive aspergillus infection; or
  - 3.2 Patient continues to require treatment for possible invasive aspergillus infection; or
  - 3.3 Patient has fluconazole resistant candidiasis; or
  - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Initial application — (Invasive fungal infection prophylaxis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient is at risk of invasive fungal infection; and
- 2 Either:
  - 2.1 Voriconazole is prescribed by, or recommended by a haematologist, transplant physician, infectious disease specialist, paediatric haematologist or paediatric oncologist; or
  - 2.2 Prescribing voriconazole is in accordance with a protocol or guideline that has been endorsed by the Health New Zealand - Te Whatu Ora Hospital in the specific settings where there is a greater than 10% risk of invasive fungal infection (IFI).

Renewal — (Invasive fungal infection prophylaxis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient is at risk of invasive fungal infection; and
- 2 Fither:
  - 2.1 Voriconazole is prescribed by, or recommended by a haematologist, transplant physician, infectious disease specialist, paediatric haematologist or paediatric oncologist; or
  - 2.2 Prescribing voriconazole is in accordance with a protocol or guideline that has been endorsed by the Health New Zealand - Te Whatu Ora Hospital in the specific settings where there is a greater than 10% risk of invasive fungal infection (IFI).

#### **Antimalarials**

### ⇒SA1684 Special Authority for Subsidy

**Initial application** only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 The patient has vivax or ovale malaria; and
- 2 Primaguine is to be given for a maximum of 21 days.

**Renewal** only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 The patient has relapsed vivax or ovale malaria; and
- 2 Primaquine is to be given for a maximum of 21 days.

	Subsidy		Fully	Brand or
	(Manufacturer's F	Price)	Subsidised	Generic
	\$	Per		Manufacturer
Antitrichomonal Agents				
METRONIDAZOLE				
Tab 200 mg - Up to 30 tab available on a PSO	25.86	250	1	Metronidamed
1 ab 200 mg	33.15			Metrogyl
Metronidamed to be Principal Supply on 1 March 2025				
Tab 400 mg - Up to 15 tab available on a PSO	4.29	21	1	Metronidamed
	5.23		1	Metrogyl
Metronidamed to be Principal Supply on 1 March 2025				<b>.</b> ,
Oral lig benzoate 200 mg per 5 ml	25.00	100 ml	1	Flagyl-S
Suppos 500 mg	24.48	10		Flagyl
(Metrogyl Tab 200 mg to be delisted 1 March 2025)				•
(Metrogyl Tab 400 mg to be delisted 1 March 2025)				
ORNIDAZOLE				
Tab 500 mg	36 52	10	J	Arrow-Ornidazole
Arrow-Ornidazole to be Principal Supply on 1 March 202		10	•	Allow-Ollidazoie
Arrow-Ornidazole to be i filicipal Supply of i i March 202	3			
<b>Antituberculotics and Antileprotics</b>				
Antituber culotics and Antileprotics				
Note: There is no co-payment charge for all pharmaceuticals liste	ed in the Antitub	erculotics a	nd Antile	protics group regardless of
immigration status.				, 9 J 9
BEDAQUILINE - Special Authority see SA2244 below - Retail pl	harmacy			
No patient co-payment payable	паппасу			
Tab 100mg	3 084 51	24 OP	1	Sirturo
	0,004.01	2401	•	Ontaro
⇒SA2244 Special Authority for Subsidy			·	
Initial application — (multi-drug resistant tuberculosis) from	any relevant pra	actitioner. /	Approvais	valid for 6 months for
applications meeting the following criteria:				
Both:				
1 The person has multi-drug resistant tuberculosis (MDR-TE	, :			
2 Ministry of Health's Tuberculosis Clinical Network has revi	ewed the individ	iuai case ai	nd recomi	mends bedaquiline as part
of the treatment regimen.				
CLOFAZIMINE – Retail pharmacy-Specialist				
a) No patient co-payment payable				
b) Prescriptions must be written by, or on the recommendation	on of, an infection	ous disease	physicia	n, clinical microbiologist or
dermatologist.			. ,	
* Cap 50 mg	442.00	100	1	Lamprene S29
CYCLOSERINE - Retail pharmacy-Specialist				•
a) No patient co-payment payable				
b) Prescriptions must be written by, or on the recommendation	on of an infectio	nue diepaee	nhveicia	n clinical microhiologist or
respiratory physician.	on or, an imedic	Jus uiscasc	priysiciai	ii, ciiilicai microbiologist oi
Cap 250 mg	344.00	60	1	Cyclorin \$29
		00	•	Oycioiiii 🚥
DAPSONE - Retail pharmacy-Specialist				
a) No patient co-payment payable				
b) Prescriptions must be written by, or on the recommendation	on of, an infection	ous disease	physiciai	n, clinical microbiologist or
dermatologist			_	_
Tab 25 mg		100		Dapsone
Tab 100 mg	329.50	100	/	Dapsone

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	
ETHAMBUTOL HYDROCHLORIDE - Retail pharmacy-Specialis	t			
a) No patient co-payment payable     b) Prescriptions must be written by, or on the recommendati respiratory physician		disease	e physiciar	n, clinical microbiologist or
Tab 100 mg		100	✓	EMB Fatol \$29
Tab 400 mg	49.34	56	/	Myambutol \$29
ISONIAZID - Retail pharmacy-Specialist				
<ul> <li>a) No patient co-payment payable</li> <li>b) Prescriptions must be written by, or on the recommendati microbiologist, dermatologist or public health physician</li> </ul>	on of, an internal me	dicine	physician	, paediatrician, clinical
* Tab 100 mg	23.00	100	1	PSM
	94.50			Isoniazid Teva S29
(POM Tab 400 area to be delicated 4 Mars 2005)	327.41		•	Noumed Isoniazid
(PSM Tab 100 mg to be delisted 1 May 2025)				
ISONIAZID WITH RIFAMPICIN - Retail pharmacy-Specialist				
<ul> <li>a) No patient co-payment payable</li> <li>b) Prescriptions must be written by, or on the recommendati microbiologist, dermatologist or public health physician</li> </ul>	on of, an internal me	dicine	physician	, paediatrician, clinical
* Tab 100 mg with rifampicin 150 mg	89.82	100	✓	Rifinah
Rifinah to be Principal Supply on 1 February 2025				
* Tab 150 mg with rifampicin 300 mg Rifinah to be Principal Supply on 1 February 2025	179.13	100	•	Rifinah
LINEZOLID – Special Authority see SA2234 below – Retail phar No patient co-payment payable	macy			
Tab 600 mg	194 60	10	/	Zyvox
Oral lig 20 mg per ml		150 m		Zyvox
■ SA2234 Special Authority for Subsidy Initial application — (multi-drug resistant tuberculosis) from applications meeting the following criteria: Both:  1 The person has multi-drug resistant tuberculosis (MDR-TE		oner. A	Approvals	valid for 18 months for
2 Ministry of Health's Tuberculosis Clinical Network has revi the treatment regimen.		case a	nd recomr	mends linezolid as part of
PARA-AMINO SALICYLIC ACID - Retail pharmacy-Specialist				
<ul> <li>a) No patient co-payment payable</li> <li>b) Prescriptions must be written by, or on the recommendation respiratory physician</li> </ul>				_
Grans for oral liq 4 g sachet	280.00	30	•	Paser S29
PROTIONAMIDE - Retail pharmacy-Specialist				
<ul> <li>a) No patient co-payment payable</li> <li>b) Prescriptions must be written by, or on the recommendati respiratory physician</li> </ul>	on of, an infectious o	disease	e specialis	t, clinical microbiologist or
Tab 250 mg	305.00	100	✓	Peteha S29
PYRAZINAMIDE - Retail pharmacy-Specialist				
<ul> <li>a) No patient co-payment payable</li> <li>b) Prescriptions must be written by, or on the recommendati respiratory physician</li> </ul>	on of, an infectious o	disease	e physiciar	n, clinical microbiologist or
* Tab 500 mg	64.95	100	✓	AFT-Pyrazinamide
* 180 500 mg	64.95	100	•	AF I-Pyrazinamide

		INFECTIONS -	AGENTS	FOR S	SYSTEMIC USE
_		Subsidy (Manufacturer's Pri	ce) Subs	Fully idised	Brand or Generic Manufacturer
RII	-ABUTIN - Retail pharmacy-Specialist				
	a) No patient co-payment payable     b) Prescriptions must be written by, or on the recommend gastroenterologist	ation of, an infectiou	ıs disease phy	/sician,	respiratory physician or
*	Cap 150 mg	353.71	30	✓ N	lycobutin
RII	FAMPICIN - Subsidy by endorsement				
*	<ul> <li>a) No patient co-payment payable</li> <li>b) For confirmed recurrent Staphylococcus aureus infectio antimicrobial based on susceptibilities and the prescrip Retail pharmacy - Specialist. Specialist must be an integrate page of payable payable.</li> <li>Cap 150 mg</li></ul>	tion is endorsed accernal medicine phys	ordingly; can ician, clinical i 100	be waiv microbio	ved by endorsement - ologist, dermatologist,
*	Cap 300 mg	122.06	100	_	lifadin
*	Oral liq 100 mg per 5 ml	12.60	60 ml		lifadin Sanofi l <u>ifadin</u>
A	ntivirals				
Fo	r eye preparations refer to Eye Preparations, Anti-Infective F	Preparations, page 2	68		
H	epatitis B Treatment				
	TECAVIR				
	Tab 0.5 mg		30	<b>✓</b> <u>E</u>	ntecavir (Rex)
LA	MIVUDINE – Special Authority see SA1685 below – Retail   Tab 100 mg Oral liq 5 mg per ml	12.06	28 240 ml OP		<u>etlam</u> effix
	SA1685 Special Authority for Subsidy				
	tial application only from a relevant specialist or medical pr provals valid for 1 year where used for the treatment or prev			of a re	levant specialist.
	newal from any relevant practitioner. Approvals valid for 2 y	•		nt or pre	evention of hepatitis B.
TE	NOFOVIR DISOPROXIL  Tenofovir disoproxil prescribed under endorsement for the antiretrovirals for the purposes of Special Authority SA213:		included in the	e count	of up to 4 subsidised
*	Tab 245 mg (300 mg as a maleate)		30	<b>✓</b> <u>T</u>	enofovir Disoproxil Viatris
Н	erpesvirus Treatments				
AC	ICLOVIR				
*	Tab dispersible 200 mg	1.78	25	<b>✓</b> <u>L</u>	<u>ovir</u>
*	Tab dispersible 400 mg		56	✓ <u>L</u>	
	Tab dispersible 800 mg	6.46	35	✓ <u>L</u>	<u>ovir</u>
VA	LACICLOVIR Tab 500 mg Vaclovir to be Principal Supply on 1 February 2025	9.64	30	<b>√</b> ∨	aclovir
	Tab 1 000 mg	17 70	20	√ V	oolovir

Tab 450 mg ......140.89

Valganciclovir Viatris to be Principal Supply on 1 February 2025

Tab 1,000 mg .......17.78

VALGANCICLOVIR - Special Authority see SA1993 on the next page - Retail pharmacy

Vaclovir to be Principal Supply on 1 February 2025

30

60

✓ Vaclovir

✓ Valganciclovir

Viatris

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

			_
Subsidy	Fully	Brand or	_
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

### ⇒SA1993 Special Authority for Subsidy

**Initial application** — **(transplant cytomegalovirus prophylaxis)** only from a relevant specialist. Approvals valid for 3 months where the patient has undergone a solid organ transplant and requires valganciclovir for CMV prophylaxis.

Renewal — (transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

#### Either:

- 1 Both:
  - 1.1 Patient has undergone a solid organ transplant and received anti-thymocyte globulin and requires valganciclovir therapy for CMV prophylaxis; and
  - 1.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following anti-thymocyte globulin; or
- 2 Both:
  - 2.1 Patient has received pulse methylprednisolone for acute rejection and requires further valganciclovir therapy for CMV prophylaxis; and
  - 2.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following pulse methylprednisolone.

Initial application — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has undergone a solid organ transplant and received valganciclovir under Special Authority more than 2 years ago (27 months); and
- 2 Patient has received anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Renewal — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months where the patient has received a further course of anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Initial application — (Lung transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone a lung transplant; and
- 2 Either:
  - 2.1 The donor was cytomegalovirus positive and the patient is cytomegalovirus negative; or
  - 2.2 The recipient is cytomegalovirus positive; and
- 3 Patient has a high risk of CMV disease.

Initial application — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

#### Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
  - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
  - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
  - 2.3 Patient has cytomegalovirus retinitis.

Renewal — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

#### Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
  - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
  - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
  - 2.3 Patient has cytomegalovirus retinitis.

Note: for the purpose of this Special Authority "immunocompromised" includes transplant recipients, patients with immunosuppressive diseases (e.g. HIV) or those receiving immunosuppressive treatment for other conditions.

Subsidy (Manufacturer's Price)

Subsidised Per 🗸

Fully

Brand or Generic Manufacturer

## **Hepatitis C Treatment**

GLECAPREVIR WITH PIBRENTASVIR - [Xpharm]

Note the supply of treatment is via Pharmac's approved direct distribution supply. Further details can be found on Pharmac's website https://pharmac.govt.nz/maviret

Tab 100 mg with pibrentasvir 40 mg ......24,750.00 84 OP ✓ Maviret

LEDIPASVIR WITH SOFOSBUVIR - [Xpharm] - Special Authority see SA1605 below

No patient co-payment payable

Tab 90 mg with sofosbuvir 400 mg.......24,363.46 28 **✓ Harvoni** 

### ⇒SA1605 Special Authority for Subsidy

Special Authority approved by the Hepatitis C Treatment Panel (HepCTP)

Notes: By application to the Hepatitis C Treatment Panel (HepCTP).

Applications will be considered by HepCTP and approved subject to confirmation of eligibility.

Application details may be obtained from Pharmac's website http://www.pharmac.govt.nz/mayiret or:

The Coordinator, Hepatitis C Treatment Panel

Pharmac, PO Box 10-254, WELLINGTON Tel: (04) 460 4990.

Email: hepcpanel@pharmac.govt.nz

## **HIV Prophylaxis and Treatment**

EMTRICITABINE WITH TENOFOVIR DISOPROXIL — Subsidy by endorsement; can be waived by Special Authority see SA2138 below

- a) Funding for emtricitabine with tenofovir disoproxil for use as PrEP, should be applied using Special Authority SA2138.
- b) Endorsement for treatment of conditions approved via Special Authority SA2139 (antiretrovirals for confirmed HIV, prevention of maternal transmission, post-exposure prophylaxis following exposure to HIV and percutaneous exposure): Prescription is deemed to be endorsed if emtricitabine with tenofovir disoproxil is co-prescribed with another antiretroviral subsidised under Special Authority SA2139 and the prescription is annotated accordingly by the Pharmacist or endorsed by the prescriber.

Note: Emtricitabine with tenofovir disoproxil prescribed under endorsement, for treatment of conditions approved via Special Authority SA2139 (antiretrovirals for confirmed HIV, prevention of maternal transmission, post-exposure prophylaxis following exposure to HIV and percutaneous exposure), is included in the count of up to 4 subsidised antiretrovirals, and counts as two antiretroviral medications, for the purposes of Special Authority SA2139, page 114 There is an approval process to become a named specialist to prescribe antiretroviral therapy in New Zealand. Further information is available on the Pharmac website.

\* Tab 200 mg with tenofovir disoproxil 245 mg (300 mg as a

### ⇒SA2138 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria:

Both:

1 Patient has tested HIV negative, does not have signs or symptoms of acute HIV infection and has been assessed for HIV seroconversion; and

30

✓ Teva

2 The Practitioner considers the patient is at elevated risk of HIV exposure and use of PrEP is clinically appropriate.

Notes: Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines:

https://ashm.org.au/HIV/PrEP/

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	1	Manufacturer

continued...

Renewal from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria:

- 1 Patient has tested HIV negative, does not have signs or symptoms of acute HIV infection and has been assessed for HIV seroconversion; and
- 2 The Practitioner considers the patient is at elevated risk of HIV exposure and use of PrEP is clinically appropriate.

Notes: Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical quidelines:

https://ashm.org.au/HIV/PrEP/

### **COVID-19 Treatments**

MOLNUPIRAVIR - [Xpharm] - Subsidy by endorsement

- a) No patient co-payment payable
- b) Treatment is funded only if patient meets access criteria for oral antiviral COVID-19 treatments (as on <u>Pharmac's website</u>) and has been endorsed accordingly by the prescriber. The supply of treatment is via Pharmac's approved distribution process. Refer to the Pharmac website for more information about this and stock availability.

NIRMATRELVIR WITH RITONAVIR - [Xpharm] - Subsidy by endorsement

- a) No patient co-payment payable
- b) Treatment is funded only if patient meets access criteria for oral antiviral COVID-19 treatments (as on <u>Pharmac's website</u>) and has been endorsed accordingly by the prescriber. The supply of treatment is via Pharmac's approved distribution process. Refer to the Pharmac website for more information about this and stock availability.

### Antiretrovirals

### ⇒SA2139 Special Authority for Subsidy

Initial application — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the patient has confirmed HIV infection.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Prevention of maternal transmission) only from a named specialist. Approvals valid for 1 year for applications meeting the following criteria:

Fither:

- 1 Prevention of maternal foetal transmission: or
- 2 Treatment of the newborn for up to eight weeks.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Some antiretrovirals are unapproved or contraindicated for this indication. Practitioners prescribing these medications should exercise their own skill, judgement, expertise and discretion, and make their own prescribing decisions with respect to the use of a Pharmaceutical for an indication for which it is not approved or contraindicated.

Subsidy	Fu	ılly	Brand or
(Manufacturer's Price)	Subsidis	ed	Generic
 \$	Per	•	

continued...

Initial application — (post-exposure prophylaxis following exposure to HIV) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

- Both:
  - 1 Treatment course to be initiated within 72 hours post exposure; and
  - 2 Any of the following:
    - 2.1 Patient has had condomless anal intercourse or receptive vaginal intercourse with a known HIV positive person with an unknown or detectable viral load greater than 200 copies per ml; or
    - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
    - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required; or
    - 2.4 Patient has had condomless anal intercourse with a person from a high HIV prevalence country or risk group whose HIV status is unknown.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines for PEP (https://www.ashm.org.au/hiv/hiv-management/pep/).

Renewal — (second or subsequent post-exposure prophylaxis) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
  - 2.1 Patient has had condomless anal intercourse or receptive vaginal intercourse with a known HIV positive person with an unknown or detectable viral load greater than 200 copies per ml; or
  - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
  - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required; or
  - 2.4 Patient has had condomless anal intercourse with a person from a high HIV prevalence country or risk group whose HIV status is unknown.

Initial application — (Percutaneous exposure) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Second or subsequent percutaneous exposure) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

## Non-nucleosides Reverse Transcriptase Inhibitors

EFAVIRENZ - Special Authority see SA2139 on the previous page -	Retail pha	rmacy	
Tab 600 mg	65.38	30	✓ Efavirenz
			Milpharm S29
ETRAVIRINE - Special Authority see SA2139 on the previous page	- Retail pha	armacy	
Tab 200 mg	770.00	60	✓ Intelence
NEVIRAPINE - Special Authority see SA2139 on the previous page	- Retail pha	armacy	
Tab 200 mg	198.25	60	✓ Nevirapine Viatris
Nevirapine Viatris to be Principal Supply on 1 February 2025			
Oral suspension 10 mg per ml	203.55	240 ml OP	✓ Viramune Suspension

	Subsidy (Manufacturer's Price \$	e) Subs	Fully sidised	Brand or Generic Manufacturer
Nucleosides Reverse Transcriptase Inhibitors				
ABACAVIR SULPHATE - Special Authority see SA2139 on pag Tab 300 mg		macy 60	<b>√</b> Z	/iagen
ABACAVIR SULPHATE WITH LAMIVUDINE – Special Authority Note: abacavir with lamivudine (combination tablets) counts anti-retroviral Special Authority.				
Tab 600 mg with lamivudine 300 mg	29.50	30	✓ ½	<u>Abacavir/</u> <u>Lamivudine</u> Viatris
EFAVIRENZ WITH EMTRICITABINE AND TENOFOVIR DISOPI pharmacy Note: Efavirenz with emtricitabine and tenofovir disoproxil of anti-retroviral Special Authority Tab 600 mg with emtricitabine 200 mg and tenofovir disoprox	ounts as three anti-	•		
245 mg (300 mg as a maleate)  EMTRICITABINE – Special Authority see SA2139 on page 114		30	<b>✓</b> \	/iatris
Cap 200 mg	307.20	30	<b>✓</b> E	Emtriva
LAMIVUDINE – Special Authority see SA2139 on page 114 – Re Tab 150 mg Oral liq 10 mg per ml	98.00	60 240 ml OP	✓ <u>L</u>	amivudine Viatris
ZIDOVUDINE [AZT] – Special Authority see SA2139 on page 11 Cap 100 mgOral liq 10 mg per ml	152.25	y 100 200 ml OP		Retrovir Retrovir
ZIDOVUDINE [AZT] WITH LAMIVUDINE – Special Authority see Note: zidovudine [AZT] with lamivudine (combination tablets the anti-retroviral Special Authority.	, ,			•
Tab 300 mg with lamivudine 150 mg	92.40	60	<b>√</b> L	amivudine/ Zidovudine Viatris
Protease Inhibitors				
ATAZANAVIR SULPHATE – Special Authority see SA2139 on p Cap 150 mg Cap 200 mg	85.00	60 60	_	Atazanavir Mylan Atazanavir Viatris
DARUNAVIR – Special Authority see SA2139 on page 114 – Re Tab 400 mg Tab 600 mg	150.00	60 60	_	Darunavir Viatris Darunavir Viatris
LOPINAVIR WITH RITONAVIR — Special Authority see SA2139 Tab 100 mg with ritonavir 25 mg		ail pharmac 60	•	.opinavir/Ritonavir Mylan
Tab 200 mg with ritonavir 50 mg	875.00	120	<b>✓</b> L	.opinavir/Ritonavir Mylan
Lopinavir/Ritonavir Mylan to be Principal Supply on 1 Fe	•			•
RITONAVIR – Special Authority see SA2139 on page 114 – Ret Tab 100 mg		30	<b>✓</b> N	lorvir

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	osidised	Generic	
\$	Per	✓	Manufacturer	

### Strand Transfer Inhibitors

DOLUTEGRAVIR – Special Authority see SA2139 on page 1	14 – Retail pharmacy		
Tab 50 mg	1,090.00	30	✓ Tivicay
DOLUTEGRAVIR WITH LAMIVUDINE - Special Authority se	ee SA2139 on page 114	– Retail p	harmacy
Tab 50 mg with lamivudine 300 mg	1,090.00	30	✓ Dovato
RALTEGRAVIR POTASSIUM - Special Authority see SA213	89 on page 114 – Retail	pharmacy	
Tab 400 mg	1,090.00	60	✓ Isentress
Tab 600 mg	1 090 00	60	✓ Isentress HD

### **Immune Modulators**

PEGYLATED INTERFERON ALFA-2A - Special Authority see SA2034 below - Retail pharmacy

Note: Pharmac will consider funding ribavirin for the small group of patients who have a clinical need for ribavirin and meet Special Authority criteria. Please contact the Hepatitis C Coordinator at Pharmac on 0800-023-588 option 4. Inj 180 mcg prefilled syringe......748.50

### ⇒SA2034 Special Authority for Subsidy

Initial application — (chronic hepatitis C - genotype 1, 4, 5 or 6 infection or co-infection with HIV or genotype 2 or 3 post liver transplant) from any specialist. Approvals valid for 18 months for applications meeting the following criteria: Both:

- 1 Any of the following:
  - 1.1 Patient has chronic hepatitis C, genotype 1, 4, 5 or 6 infection; or
  - 1.2 Patient has chronic hepatitis C and is co-infected with HIV; or
  - 1.3 Patient has chronic hepatitis C genotype 2 or 3 and has received a liver transplant; and
- 2 Maximum of 48 weeks therapy.

Renewal — (Chronic hepatitis C - genotype 1 infection) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria: All of the following:

- - 1 Patient has chronic hepatitis C, genotype 1; and
  - 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
  - 3 Fither:
    - 3.1 Patient has responder relapsed; or
    - 3.2 Patient was a partial responder; and
  - 4 Patient is to be treated in combination with boceprevir; and
  - 5 Maximum of 48 weeks therapy.

Initial application — (Chronic Hepatitis C - genotype 1 infection treatment more than 4 years prior) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Any of the following:
  - 3.1 Patient has responder relapsed; or
  - 3.2 Patient was a partial responder; or
  - 3.3 Patient received interferon treatment prior to 2004; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Subsidy		Fully	Brand or
(Manufacturer's Price	e)	Subsidised	Generic
\$	Per	•	Manufacturer

continued...

Initial application — (chronic hepatitis C - genotype 2 or 3 infection without co-infection with HIV) from any specialist.

Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has chronic hepatitis C, genotype 2 or 3 infection; and
- 2 Maximum of 6 months therapy.

Initial application — (Hepatitis B) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Hepatitis B infection (HBsAg positive for more than 6 months); and
- 2 Patient is Hepatitis B treatment-naive; and
- 3 ALT > 2 times Upper Limit of Normal; and
- 4 HBV DNA < 10 log10 IU/ml; and
- 5 Either:
  - 5.1 HBeAg positive; or
  - 5.2 serum HBV DNA greater than or equal to 2,000 units/ml and significant fibrosis (Metavir Stage F2 or greater or moderate fibrosis); and
- 6 Compensated liver disease: and
- 7 No continuing alcohol abuse or intravenous drug use; and
- 8 Not co-infected with HCV. HIV or HDV: and
- 9 Neither ALT nor AST > 10 times upper limit of normal; and
- 10 No history of hypersensitivity or contraindications to pegylated interferon; and
- 11 Maximum of 48 weeks therapy.

Initial application — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 Patient has a cutaneous T cell lymphoma\*; or
- 2 All of the following:
  - 2.1 Patient has a myeloproliferative disorder\*; and
  - 2.2 Patient is intolerant of hydroxyurea; and
  - 2.3 Treatment with an agrelide and busulfan is not clinically appropriate; or
- 3 Both:
  - 3.1 Patient has a myeloproliferative disorder; and
  - 3.2 Patient is pregnant, planning pregnancy or lactating.

Renewal — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment; and
- 3 Either:
  - 3.1 Patient has a cutaneous T cell lymphoma\*; or
  - 3.2 Both:
    - 3.2.1 Patient has a myeloproliferative disorder\*; and
    - 3.2.2 Either:
      - 3.2.2.1 Remains intolerant of hydroxyurea and treatment with anagrelide and busulfan remains clinically inappropriate; or
      - 3.2.2.2 Patient is pregnant, planning pregnancy or lactating.

Note: Indications marked with \* are unapproved indications.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	9	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued

Initial application — (post-allogenic bone marrow transplant) from any relevant practitioner. Approvals valid for 3 months where patient has received an allogeneic bone marrow transplant\* and has evidence of disease relapse.

Renewal — (post-allogenic bone marrow transplant) from any relevant practitioner. Approvals valid for 3 months where patient is responding and ongoing treatment remains appropriate.

Note: Indications marked with \* are unapproved indications.

## **Urinary Tract Infections**

FOSFOMYCIN - Special Authority see SA2406 below - Retail pharmacy ✓ UroFos 

### ⇒SA2406 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 months for applications meeting the following criteria: Both:

- 1 Patient has an acute, symptomatic, bacteriologically-proven uncomplicated urinary tract infection (UTI)/cystitis with Escherichia Coli; and
- 2 Fither:
  - 2.1 Microbiological testing confirms the pathogen is resistant to all of: trimethoprim, nitrofurantoin, amoxicillin, cefaclor. cefalexin, amoxicillin with clavulanic acid, and norfloxacin; or
  - 2.2 The patient has a contraindication or intolerance to all of: trimethoprim, nitrofurantoin, amoxicillin, cefaclor, cefalexin, amoxicillin with clavulanic acid, and norfloxacin that the pathogen is susceptible to.

Renewal from any relevant practitioner. Approvals valid for 2 months for applications meeting the following criteria: Both:

- 1 Patient has an acute, symptomatic, bacteriologically-proven uncomplicated urinary tract infection (UTI)/cystitis with Escherichia Coli: and
- 2 Fither:
  - 2.1 Microbiological testing confirms the pathogen is resistant to all of: trimethoprim, nitrofurantoin, amoxicillin, cefaclor, cefalexin, amoxicillin with clavulanic acid, and norfloxacin; or
  - 2.2 The patient has a contraindication or intolerance to all of: trimethoprim, nitrofurantoin, amoxicillin, cefaclor, cefalexin, amoxicillin with clavulanic acid, and norfloxacin that the pathogen is susceptible to.

METHENAMINE (HEXAMINE) HIPPURATE			
* Tab 1 g	19.95	100	✓ Hiprex
NITROFURANTOIN			
* Tab 50 mg - Up to 30 tab available on a PSO	22.20	100	✓ Nifuran
* Tab 100 mg	37.50	100	✓ Nifuran
* Cap modified-release 100 mg - Up to 15 cap available on a			
PSO	81.20	100	✓ Macrobid
NORFLOXACIN			
Tab 400 mg - Subsidy by endorsement	245.00	100	✓ Arrow-Norfloxacin
Only if prescribed for a patient with an uncomplicated urin			

٥r with proven resistance to first line agents and the prescription is endorsed accordingly.

	Subsidy		Fully Brand or	
	(Manufacturer's Price	e) Sub	sidised Generic	
	\$	Per	<ul> <li>Manufacturer</li> </ul>	
Anticholinesterases				
NEOSTIGMINE METILSULFATE				
Inj 2.5 mg per ml, 1 ml ampoule	48.25	10	✓ Max Health	
Max Health to be Principal Supply on 1 February 2025		. •		
,				
PYRIDOSTIGMINE BROMIDE			_	
▲ Tab 60 mg	50.28	100	Mestinon	
Non-Steroidal Anti-Inflammatory Drugs				
,				
DICLOFENAC SODIUM				
* Tab EC 25 mg	2.19	50	✓ Diclofenac Sandoz	
Diclofenac Sandoz to be Principal Supply on 1 February				
		20	✓ Voltaren D	
the second are personal and are the second are the				
* Tab EC 50 mg		50	<ul> <li>Diclofenac Sandoz</li> </ul>	
Diclofenac Sandoz to be Principal Supply on 1 February				
* Tab long-acting 75 mg	19.60	100	✓ Voltaren SR	
* Inj 25 mg per ml, 3 ml ampoule – Up to 5 inj available on a P	SO 13.20	5	✓ Voltaren	
* Suppos 12.5 mg	2.04	10	✓ Voltaren	
* Suppos 25 mg	2.44	10	✓ Voltaren	
* Suppos 50 mg - Up to 10 supp available on a PSO		10	✓ Voltaren	
* Suppos 100 mg		10	✓ Voltaren	
0	7.00	10	Voltaren	
IBUPROFEN				
* Tab 200 mg	21.40	1,000	✓ Relieve	
* Tab long-acting 800 mg	3.05	30	✓ Brufen SR	
• •	3.65		✓ Ibuprofen SR BNM	
* Oral liq 20 mg per ml	2.85	200 ml	✓ Ethics	
(Brufen SR Tab long-acting 800 mg to be delisted 1 April 2025)			00	
, , ,				
KETOPROFEN			_	
* Cap long-acting 200 mg	12.07	28	Oruvail SR	
MEFENAMIC ACID				
* Cap 250 mg	1.25	50		
* Cap 250 mg		50	Danatan	
	(10.82)	00	Ponstan	
	0.50	20		
	(7.50)		Ponstan	
NAPROXEN				
* Tab 250 mg	39 23	500	✓ Noflam 250	
Noflam 250 to be Principal Supply on 1 February 2025	00.20	000	· Honain 200	
,	04.45	050	✓ Noflam 500	
* Tab 500 mg	34.45	250	• Nonam 500	
Noflam 500 to be Principal Supply on 1 February 2025				
* Tab long-acting 750 mg		28	Naprosyn SR 750	
Naprosyn SR 750 to be Principal Supply on 1 February 2				
* Tab long-acting 1 g	11.50	28	Naprosyn SR 1000	
Naprosyn SR 1000 to be Principal Supply on 1 February			• •	
TENOXICAM	10.50	100	/ Tileatil	
* Tab 20 mg		100	✓ <u>Tilcotil</u>	
* Inj 20 mg vial	9.95	1	✓ AFT	

	MI	JSCUL	.USKEL	LETAL SYSTEM
A)	Subsidy Manufacturer's Price	) Si Per	Fully ubsidised	Brand or Generic Manufacturer
NSAIDs Other				
CELECOXIB				
Cap 100 mg	3.45	60	-	Celebrex Celecoxib Pfizer
Cap 200 mg	3.20	30	_	Celebrex
			<b>√</b> <u>C</u>	Celecoxib Pfizer
Topical Products for Joint and Muscular Pain				
CAPSAICIN				
Crm 0.025% - Special Authority see SA1289 below - Retail				
pharmacy	9.75	45 g OP		Co-Rub Osteo S29
	13.00	60 g OP		ostrix Rugby Capsaicin
		3		Topical
SA1289 Special Authority for Subsidy				Cream S29
steoarthritis that is not responsive to paracetamol and oral non-ste  Antirheumatoid Agents  IYDROXYCHLOROQUINE SULPHATE  Tab 200 mg		100		oca-
				Hydroxychloroquine
Plaquenil Tab 200 mg to be delisted 1 May 2025)	8.78		<b>✓</b> F	Plaquenil
EFLUNOMIDE				
₭ Tab 10 mg	6.00	30	<b>✓</b> <u>P</u>	<u>Irava</u>
<b>k</b> Tab 20 mg	6.00	30	<b>✓</b> <u>F</u>	<u>Irava</u>
PENICILLAMINE Tab 125 mg	67.23	100	<b>√</b> Γ	)-Penamine
Tab 125 mg		100		)-Penamine
Drugs Affecting Bone Metabolism				
-				
Alendronate for Osteoporosis				
ALENDRONATE SODIUM	0.40		, -	
* Tab 70 mg	3.10	4	<b>✓</b> <u>F</u>	osamax
LENDRONATE SODIUM WITH COLECALCIFEROL  ★ Tab 70 mg with colecalciferol 5,600 iu	1.99	4	<b>√</b> F	osamax Plus
Other Treatments			_	
DENOSUMAB - Special Authority see SA1777 on the next page -	Hetail pharmacy			

✓ Prolia

Inj 60 mg prefilled syringe......326.00

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

## **⇒SA1777** Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 Fither:
  - 2.1 The patient is female and postmenopausal; or
  - 2.2 The patient is male or non-binary; and
- 3 Any of the following:
  - 3.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
  - 3.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons; or
  - 3.3 History of two significant osteoporotic fractures demonstrated radiologically; or
  - 3.4 Documented T-Score less than or equal to -3.0 (see Note); or
  - 3.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
  - 3.6 Patient has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 4 Zoledronic acid is contraindicated because the patient's creatinine clearance is less than 35 mL/min; and
- 5 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes); and
- 6 The patient must not receive concomitant treatment with any other funded antiresorptive agent for this condition or teriparatide.

#### Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
   Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with denosumab
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body
- e) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: risedronate sodium tab 35 mg once weekly; alendronate sodium tab 70 mg or tab 70 mg with cholecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy

	DISODIUM

Inj 3 mg per ml, 10 ml vial	32.49	1	Pamisol
Inj 6 mg per ml, 10 ml vial	88.11	1	✓ Pamisol
Inj 9 mg per ml, 10 ml vial	94.34	1	✓ Pamisol
DALOVIEENE UVEDOOUU ODIDE O LIA II II OAAT	<b>70</b>		

Si	Subsidy	Fully	Brand or
(Manufac	cturer's Price) Subsi	dised	Generic
	\$ Per	1	Manufacturer

### ⇒SA1779 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Notes); or
- 2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
- 3 History of two significant osteoporotic fractures demonstrated radiologically: or
- 4 Documented T-Score less than or equal to -3.0 (see Notes); or
- 5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Notes); or
- 6 Patient has had a Special Authority approval for zoledronic acid (Underlying cause Osteoporosis) or has had a Special Authority approval for alendronate (Underlying cause - Osteoporosis) prior to 1 February 2019.

#### Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
   Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for raloxifene funding.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis, and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

#### RISEDRONATE SODIUM

Tab 35 mg2.50	4	✓ Risedronate Sandoz
TERIPARATIDE – Special Authority see SA1139 below – Retail pharmacy		
Inj 250 mcg per ml, 2.4 ml	1	✓ Teriparatide - Teva

### ⇒SA1139 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 18 months for applications meeting the following criteria: All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 The patient has a documented T-score less than or equal to -3.0 (see Notes); and
- 3 The patient has had two or more fractures due to minimal trauma; and
- 4 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes).

#### Notes:

- a) The bone mineral density (BMD) measurement used to derive the T-score must be made using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: alendronate sodium tab 70 mg or tab 70 mg with colecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily; zoledronic acid 5 mg per year. If an intolerance of a severity necessitating permanent treatment withdrawal develops

Subsidy		Fully	Brand or	
(Manufacturer's Price	e)	Subsidised	Generic	
\$	Per	•	Manufacturer	

#### continued...

during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy.

- c) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.
- d) A maximum of 18 months of treatment (18 cartridges) will be subsidised.

#### **ZOLEDRONIC ACID**

## Hyperuricaemia and Antigout

ALLOPURINOL			
* Tab 100 mg	17.99	1,000	✓ Ipca-Allopurinol
* Tab 300 mg	22.50	500	✓ Ipca-Allopurinol
BENZBROMARONE - Special Authority see SA1963 below -	Retail pharmacy		
Tab 50 mg	32.00	100	✓ Narcaricin mite S29

### ⇒SA1963 Special Authority for Subsidy

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

- 1 The treatment remains appropriate and the patient is benefitting from the treatment; and
- 2 There is no evidence of liver toxicity and patient is continuing to receive regular (at least every three months) liver function tests

#### COLCHICINE

* Tab 500 mcg6.00	100	✓ Colgout
FEBUXOSTAT - Special Authority see SA2054 below - Retail pharmacy		
Tab 80 mg4.73	28	✓ Febuxostat (Teva)
Tab 120 mg11.78	28	✓ Febuxostat (Teva)

#### ⇒SA2054 Special Authority for Subsidy

**Initial application** — (Gout) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

#### Both:

- 1 Patient has been diagnosed with gout: and
- 2 Any of the following:
  - 2.1 The patient has a serum urate level greater than 0.36 mmol/l despite treatment with allopurinol at doses of at least 600 mg/day and addition of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
  - 2.2 The patient has experienced intolerable side effects from allopurinol such that treatment discontinuation is required and serum urate remains greater than 0.36 mmol/l despite use of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
  - 2.3 The patient has renal impairment such that probenecid is contraindicated or likely to be ineffective and serum urate remains greater than 0.36 mmol/l despite optimal treatment with allopurinol (see Note); or
  - 2.4 The patient has previously had an initial Special Authority approval for benzbromarone for treatment of gout...

Initial application — (Tumour lysis syndrome) only from a haematologist or oncologist. Approvals valid for 6 weeks for applications meeting the following criteria:

#### Both:

1 Patient is scheduled to receive cancer therapy carrying an intermediate or high risk of tumour lysis syndrome; and

Intrathecal

	Subsidy (Manufacturer's Price)	Fully Subsidised		Brand or Generic	
	\$	Per	✓	Manufacturer	
continued					

2 Patient has a documented history of allopurinol intolerance.

Renewal — (Gout) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from treatment.

Renewal — (Tumour lysis syndrome) only from a haematologist or oncologist. Approvals valid for 6 weeks where the treatment remains appropriate and the patient is benefitting from treatment.

**PROBENECID** 

100 ✓ Probenecid-AFT

## **Muscle Relaxants**

BA	CLOFEN			
*	Tab 10 mg	3.70	100	✓ Pacifen
	Inj 0.05 mg per ml, 1 ml ampoule - Subsidy by endorsement1	1.55	1	✓ Lioresal Intrathecal
	Subsidised only for use in a programmable pump in patients whe	re oral antisp	astic agent	s have been ineffective or have
	caused intolerable side effects and the prescription is endorsed a	ccordingly.	•	
	Inj 2 mg per ml, 5 ml ampoule - Subsidy by endorsement306		5	✓ Medsurge
	490	0.91	10	✓ Sintetica Baclofen

- a) Subsidised only for use in a programmable pump in patients where oral antispastic agents have been ineffective or have caused intolerable side effects and the prescription is endorsed accordingly.
- b) Sintetica Baclofen Intrathecal to be Principal Supply on 1 March 2025

(Medsurge Inj 2 mg per ml, 5 ml ampoule to be delisted 1 March 2025)

### DANTROI ENF

Cap 25 mg	112.13	100	✓ Dantrium ✓ Dantrium S29 \$29
Cap 50 mg	77.00	100	✓ Dantrium
ORPHENADRINE CITRATE			
Tab 100 mg	23.25	100	✓ Norflex
Norflex to be Principal Supply on 1 February 2025			

Subsidy (Manufacturer's Price) Fully Subsidised Per

Brand or Generic Manufacturer

# **Agents for Parkinsonism and Related Disorders**

Dopamine Agonists and	Related Agents
-----------------------	----------------

AMANTADINE HYDROCHLORIDE		
▲ Cap 100 mg38.24	60	✓ Symmetrel
63.73	100	✓ Symmetrel
APOMORPHINE HYDROCHLORIDE		·
▲ Inj 10 mg per ml, 2 ml ampoule	5	✓ Movapo
▲ Inj 10 mg per ml, 5 ml ampoule	5	✓ Movapo
ENTACAPONE	·	
▲ Tab 200 mg	100	✓ Comtan
Ÿ	100	Volitali
LEVODOPA WITH BENSERAZIDE	400	4.1. 5.11
* Tab dispersible 50 mg with benserazide 12.5 mg	100	✓ Madopar Rapid
* Cap 50 mg with benserazide 12.5 mg	100	✓ Madopar 62.5
* Cap 100 mg with benserazide 25 mg	100	✓ Madopar 125
* Cap long-acting 100 mg with benserazide 25 mg22.85	100	✓ Madopar HBS
* Cap 200 mg with benserazide 50 mg26.25	100	Madopar 250
LEVODOPA WITH CARBIDOPA		
* Tab 100 mg with carbidopa 25 mg	100	Sinemet
Sinemet to be Principal Supply on 1 February 2025		
* Tab long-acting 200 mg with carbidopa 50 mg44.99	100	Sinemet CR
Sinemet CR to be Principal Supply on 1 February 2025		
* Tab 250 mg with carbidopa 25 mg	100	Sinemet
Sinemet to be Principal Supply on 1 February 2025		
PRAMIPEXOLE HYDROCHLORIDE		
▲ Tab 0.25 mg	100	✓ Ramipex
▲ Tab 1 mg	100	✓ Ramipex
RASAGILINE		<del></del>
* Tab 1 mg53.50	30	✓ Azilect S29
<b>Ç</b>	50	AZIIGUL
ROPINIROLE HYDROCHLORIDE	0.4	45 .
▲ Tab 0.25 mg	84	Ropin
▲ Tab 1 mg	84	✓ Ropin
▲ Tab 2 mg	84	Ropin
▲ Tab 5 mg14.50	84	✓ <u>Ropin</u>
TOLCAPONE		
▲ Tab 100 mg152.38	100	✓ Tasmar
Anticholineraics		

## **Anticholinergics**

BENZATROPINE MESYLATE			
Tab 2 mg	9.59	60	✓ Benztrop
Inj 1 mg per ml, 2 ml	95.00	5	✓ Phebra
a) Up to 10 inj available on a PSO			
b) Only on a PSO			
PROCYCLIDINE HYDROCHLORIDE			

✓ Kemadrin

100

### **NERVOUS SYSTEM**

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

## Agents for Essential Tremor, Chorea and Related Disorders

RILUZOLE - Special Authority see SA1403 below - Retail pharmacy

Wastage claimable

Tab 50 mg .......117.00 56 **✓ Rilutek** 

Rilutek to be Principal Supply on 1 February 2025

### ⇒SA1403 Special Authority for Subsidy

**Initial application** only from a neurologist or respiratory specialist. Approvals valid for 6 months for applications meeting the following criteria:

### All of the following:

- 1 The patient has amyotrophic lateral sclerosis with disease duration of 5 years or less; and
- 2 The patient has at least 60 percent of predicted forced vital capacity within 2 months prior to the initial application; and
- 3 The patient has not undergone a tracheostomy; and
- 4 The patient has not experienced respiratory failure; and
- 5 Any of the following:
  - 5.1 The patient is ambulatory; or
  - 5.2 The patient is able to use upper limbs; or
  - 5.3 The patient is able to swallow.

**Renewal** from any relevant practitioner. Approvals valid for 18 months for applications meeting the following criteria: All of the following:

- 1 The patient has not undergone a tracheostomy; and
- 2 The patient has not experienced respiratory failure; and
- 3 Any of the following:
  - 3.1 The patient is ambulatory; or
  - 3.2 The patient is able to use upper limbs; or
  - 3.3 The patient is able to swallow.

#### **TETRABENAZINE**

### **Anaesthetics**

### Local

#### LIDOCAINE [LIGNOCAINE]

a) Up to 150 ml available on a PSO

b) Subsidised only if prescribed for urethral or cervical administration and the prescription is endorsed accordingly.

Gel 2%, 11 ml urethral syringe − Subsidy by endorsement...............59.50 10 ✓ Instillagel Lido

- a) Up to 5 each available on a PSO
- Subsidised only if prescribed for urethral, cervical or rectal administration and the prescription is endorsed accordingly.

	Subsidy		Fully	Brand or
	(Manufacturer's Price	e)	Subsidised	Generic
	\$	Per	•	Manufacturer
IDOCAINE [LIGNOCAINE] HYDROCHLORIDE				
Oral (gel) soln 2%	44.00	200 m	<b>✓</b>	Mucosoothe
Inj 1%, 5 ml ampoule - Up to 25 inj available on a PSO	9.50	25	✓	Lidocaine-Baxter
	17.50	50		
	(35.00)			Xylocaine
Inj 2%, 5 ml ampoule - Up to 5 inj available on a PSO	9.00 <sup>°</sup>	25	✓	Lidocaine-Baxter
Inj 1%, 20 ml ampoule - Up to 5 inj available on a PSO	12.00	5		
	(20.00)			Xylocaine
Inj 1%, 20 ml vial - Up to 5 inj available on a PSO	6.85	5	✓	Lidocaine-Baxter
Inj 2%, 20 ml vial - Up to 5 inj available on a PSO	7.15	5	✓	Lidocaine-Baxter
Inj 10%, 5 ml ampoule - Subsidy by endorsement	CBS	10	1	Xylocard 500 S29
Subsidised only for people receiving palliative care services	vices where other and	algesic	agents ha	ven't been effective.

## **Topical Local Anaesthetics**

## **⇒SA0906** Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 2 years where the patient is a child with a chronic medical condition requiring frequent injections or venepuncture.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

LIDOCAINE [LIGNOCAINE] - Special Authority:	see <mark>SA0906 above –</mark> Retail ph	narmacy	
Crm 4%	5.40	5 g OP	✓ LMX4
	27.00	30 g OP	✓ LMX4
LIDOCAINE [LIGNOCAINE] WITH PRILOCAINE	- Special Authority see SA09	906 above – Reta	il pharmacy
Crm 2.5% with prilocaine 2.5%	45.00	30 g OP	✓ EMLA
Crm 2.5% with prilocaine 2.5% (5 g tubes)	45.00	5	EMLA

## **Analgesics**

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 120

## **Non-opioid Analgesics**

ASPIRIN  * Tab dispersible 300 mg - Up to 30 tab available on a PSO	5.65	100	✓ Ethics Aspirin
CAPSAICIN – Subsidy by endorsement Subsidised only if prescribed for post-herpetic neuralgia or diabetic paccordingly.	peripheral ne	europathy and	the prescription is endorsed
Crm 0.075%	11.95	45 g OP	✓ Zo-Rub HP S29 ✓ Zostrix HP
	15.14	57 g OP	✓ Rugby Capsaicin Topical Cream S29
NEFOPAM HYDROCHLORIDE Tab 30 mg	23.40	90	✓ Acupan

## NERVOUS SYSTEM

				VOCOCIOILM
	Subsidy (Manufacturer's Price) \$	) Sub	Fully sidised	Brand or Generic Manufacturer
PARACETAMOL				
Tab 500 mg - blister pack	19.75	1,000	<b>✓</b> P	acimol
<ul> <li>a) Maximum of 300 tab per prescription; can be waiv</li> <li>b) Up to 30 tab available on a PSO</li> <li>c)</li> </ul>	ed by endorsement			
Subsidy by endorsement for higher quantitie     regular daily dosing for one month or greater     annotate the prescription as endorsed where     Maximum of 100 tab per dispensing for non-     (for non-endorsed patients), then dispense in  Tab 500 mg - bottle pack — Maximum of 300 tab per	r, and the prescription is e dispensing history sup endorsed patients. If q	s annotate ports a lo uantities p	d according-term or orescribe	lingly. Pharmacists may condition. d for more than 100 tabs
prescription; can be waived by endorsement	17.92	1,000	✓ N	oumed Paracetamol
<ol> <li>Subsidy by endorsement for higher quantities is daily dosing for one month or greater, and the p prescription as endorsed where dispensing hist</li> <li>Maximum of 100 tab per dispensing for non-end non-endorsed patients), then dispense in repeat</li> </ol>	rescription is annotated bry supports a long-tern lorsed patients. If quan	l according n condition ntities pres	gly. Pha n. cribed fo	rmacists may annotate the
Oral liq 120 mg per 5 ml	3.98	200 ml	<b>✓</b> <u>P</u>	aracetamol (Ethics)
<ul> <li>a) Maximum of 600 ml per prescription; can be waive</li> <li>b) Up to 200 ml available on a PSO</li> <li>c) Not in combination</li> <li>d)</li> </ul>	ed by endorsement			(=====)
<ol> <li>Maximum of 200 ml per dispensing for non-enon-endorsed patients), then dispense in reg</li> <li>Subsidy by endorsement for higher quantitie regular daily dosing for one month or greater Pharmacists may annotate the prescription a condition.</li> <li>Note: 200 ml presentations of paracetamol Pharmacist) under the provisions in Part I of</li> <li>Note: Direct Provision by a pharmacist of up</li> </ol>	peat dispensing not exc s is available for patien r and the prescription is as endorsed where disp oral liquid may be supp Section A o to 200 ml permitted ur	eeding 20 ts with lon endorsed ensing his lied on BS	0 ml per g term co l or anno story sup 60 to a V	dispensing. onditions who require tated accordingly. ports a long-term accinator (other than a
conjunction with immunisation of a child und Oral liq 250 mg per 5 ml	3.35	neningoco 200 ml	_	amol
<ul> <li>a) Maximum of 600 ml per prescription; can be waive</li> <li>b) Up to 200 ml available on a PSO</li> <li>c) Not in combination</li> <li>d)</li> </ul>	ed by endorsement			
Maximum of 200 ml per dispensing for non-endorsed patients), then dispense in reg     Subsidy by endorsement for higher quantitie regular daily dosing for one month or greater	peat dispensing not exc s is available for patien	eeding 20 ts with lon	0 ml per g term c	dispensing. onditions who require

- regular daily dosing for one month or greater and the prescription is endorsed or annotated accordingly. Pharmacists may annotate the prescription as endorsed where dispensing history supports a long-term
- 3) Note: 200 ml presentations of paracetamol oral liquid may be supplied on BSO to a Vaccinator (other than a Pharmacist) under the provisions in Part I of Section A
- 4) Note: Direct Provision by a pharmacist of up to 200 ml permitted under the provisions in Part I of Section A in conjunction with immunisation of a child under 2 years of age with meningococcal B multicomponent vaccine.
- 10 ✓ Gacet

		Subsidy		Fully	
		(Manufacturer's Price \$	e) Per	Subsidised	Generic Manufacturer
÷	Suppos 250 mg	5.39	10	1	Gacet
÷	Suppos 500 mg		50	✓	Gacet
0	pioid Analgesics				
Ю	DEINE PHOSPHATE - Safety medicine; prescriber may dete	ermine dispensing fr	equen	су	
	Tab 15 mg	5.92	100	′ 🗸	Noumed
	Tab 30 mg	6.98	100	✓	Aspen
	-			✓	Noumed
	Tab 60 mg	13.89	100	1	Noumed
λIF	YDROCODEINE TARTRATE				
- 11	Tab long-acting 60 mg	8 60	60	1	DHC Continus
,			00	•	
Εĺ	NTANYL				
	a) Only on a controlled drug form				
	b) No patient co-payment payable				
	c) Safety medicine; prescriber may determine dispensing fre		4.0	,	
	Inj 50 mcg per ml, 2 ml ampoule		10		Boucher and Muir
	Inj 50 mcg per ml, 10 ml ampoule		10	_	Boucher and Muir
	Patch 12.5 mcg per hour		5		Fentanyl Sandoz
	Patch 25 mcg per hour		5		Fentanyl Sandoz
	Patch 50 mcg per hour		5		Fentanyl Sandoz
	Patch 75 mcg per hour		5		Fentanyl Sandoz
	Patch 100 mcg per hour	16.37	5	•	Fentanyl Sandoz
ΛE	THADONE HYDROCHLORIDE				
	a) Only on a controlled drug form				
	b) No patient co-payment payable				
	b) 140 patient de payment payable				
	c) Safety medicine; prescriber may determine dispensing fre	equency			
	<ul> <li>c) Safety medicine; prescriber may determine dispensing free</li> <li>d) Extemporaneously compounded methadone will only be remained to the compounded methadone.</li> </ul>		ite of th	ne cheape	st form available
	<ul> <li>c) Safety medicine, prescriber may determine dispensing fred)</li> <li>d) Extemporaneously compounded methadone will only be remethadone powder, not methadone tablets).</li> </ul>	reimbursed at the ra			
	c) Safety medicine, prescriber may determine dispensing fred) Extemporaneously compounded methadone will only be r (methadone powder, not methadone tablets).  Tab 5 mg	reimbursed at the ra	10	·	Methadone BNM
	c) Safety medicine; prescriber may determine dispensing fred) Extemporaneously compounded methadone will only be r (methadone powder, not methadone tablets).  Tab 5 mg	reimbursed at the ra		·	
	c) Safety medicine; prescriber may determine dispensing fred) Extemporaneously compounded methadone will only be r (methadone powder, not methadone tablets).  Tab 5 mg	eimbursed at the ra 1.45 7.80	10 200 m	nl /	Methadone BNM Biodone
	c) Safety medicine; prescriber may determine dispensing fred) Extemporaneously compounded methadone will only be r (methadone powder, not methadone tablets).  Tab 5 mg  Oral liq 2 mg per ml  Biodone to be Principal Supply on 1 February 2025  Oral liq 5 mg per ml	eimbursed at the ra1.45 7.80	10	nl /	Methadone BNM
	c) Safety medicine; prescriber may determine dispensing fred) Extemporaneously compounded methadone will only be refuncted in the control of	eimbursed at the ra1.45 7.80 7.80	10 200 m 200 m		Methadone BNM Biodone Biodone Forte
	c) Safety medicine; prescriber may determine dispensing fred) Extemporaneously compounded methadone will only be r (methadone powder, not methadone tablets).  Tab 5 mg  Oral liq 2 mg per ml  Biodone to be Principal Supply on 1 February 2025  Oral liq 5 mg per ml  Biodone Forte to be Principal Supply on 1 February 2025  Oral liq 10 mg per ml	eimbursed at the ra	10 200 m		Methadone BNM Biodone Biodone Forte
	c) Safety medicine; prescriber may determine dispensing fred) Extemporaneously compounded methadone will only be refuncted in the control of		10 200 m 200 m 200 m		Methadone BNM Biodone Biodone Forte Biodone Extra Forte
	c) Safety medicine; prescriber may determine dispensing fred) Extemporaneously compounded methadone will only be r (methadone powder, not methadone tablets).  Tab 5 mg  Oral liq 2 mg per ml  Biodone to be Principal Supply on 1 February 2025  Oral liq 5 mg per ml  Biodone Forte to be Principal Supply on 1 February 2025  Oral liq 10 mg per ml		10 200 m 200 m		Methadone BNM Biodone Biodone Forte
10	c) Safety medicine; prescriber may determine dispensing fred) Extemporaneously compounded methadone will only be refuncted in the control of		10 200 m 200 m 200 m		Methadone BNM Biodone Biodone Forte Biodone Extra Forte
ИΟ	c) Safety medicine, prescriber may determine dispensing fre d) Extemporaneously compounded methadone will only be r (methadone powder, not methadone tablets).  Tab 5 mg  Oral liq 2 mg per ml  Biodone to be Principal Supply on 1 February 2025  Oral liq 5 mg per ml  Biodone Forte to be Principal Supply on 1 February 2025  Oral liq 10 mg per ml  Biodone Extra Forte to be Principal Supply on 1 February 10 mg per ml  Biodone Extra Forte to be Principal Supply on 1 February 10 mg per ml , 1 ml  RPHINE HYDROCHLORIDE		10 200 m 200 m 200 m		Methadone BNM Biodone Biodone Forte Biodone Extra Forte
ИΟ	c) Safety medicine, prescriber may determine dispensing fre d) Extemporaneously compounded methadone will only be r (methadone powder, not methadone tablets).  Tab 5 mg  Oral liq 2 mg per ml  Biodone to be Principal Supply on 1 February 2025  Oral liq 5 mg per ml  Biodone Forte to be Principal Supply on 1 February 2025  Oral liq 10 mg per ml  Biodone Extra Forte to be Principal Supply on 1 February 10 mg per ml  Biodone Extra Forte to be Principal Supply on 1 February 10 mg per ml , 1 ml  RPHINE HYDROCHLORIDE a) Only on a controlled drug form		10 200 m 200 m 200 m		Methadone BNM Biodone Biodone Forte Biodone Extra Forte
ИΟ	c) Safety medicine, prescriber may determine dispensing fre d) Extemporaneously compounded methadone will only be r (methadone powder, not methadone tablets).  Tab 5 mg  Oral liq 2 mg per ml  Biodone to be Principal Supply on 1 February 2025  Oral liq 5 mg per ml  Biodone Forte to be Principal Supply on 1 February 2025  Oral liq 10 mg per ml  Biodone Extra Forte to be Principal Supply on 1 February 10 mg per ml  Biodone Extra Forte to be Principal Supply on 1 February 10 mg per ml  RPHINE HYDROCHLORIDE  a) Only on a controlled drug form b) No patient co-payment payable	-eimbursed at the ra 	10 200 m 200 m 200 m		Methadone BNM Biodone Biodone Forte Biodone Extra Forte
ΛΟ	c) Safety medicine, prescriber may determine dispensing fred) Extemporaneously compounded methadone will only be referred (methadone powder, not methadone tablets).  Tab 5 mg	-eimbursed at the ra 	10 200 m 200 m 200 m		Methadone BNM Biodone Biodone Forte Biodone Extra Forte AFT
МΟ	c) Safety medicine, prescriber may determine dispensing fred) Extemporaneously compounded methadone will only be referred (methadone powder, not methadone tablets).  Tab 5 mg	eimbursed at the ra	10 200 m 200 m 200 m 10		Methadone BNM Biodone Biodone Forte Biodone Extra Forte AFT
МО	c) Safety medicine, prescriber may determine dispensing fred) Extemporaneously compounded methadone will only be referred (methadone powder, not methadone tablets).  Tab 5 mg	eimbursed at the ra	10 200 m 200 m 200 m 10		Methadone BNM Biodone Biodone Forte Biodone Extra Forte AFT

	Subsidy (Manufacturer's Price	) S	Fully ubsidised	Brand or Generic
	\$	Per	✓	Manufacturer
ORPHINE SULPHATE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing	frequency			
Tab immediate-release 10 mg		10	<b>√</b> 9	Sevredol
Tab immediate-release 20 mg	5.52	10	✓ 9	Sevredol
Cap long-acting 10 mg	3.00	10	<b>✓</b> n	n-Eslon
Cap long-acting 30 mg	4.30	10	<b>√</b> r	n-Eslon
Cap long-acting 60 mg	9.00	10	✓ r	n-Eslon
Cap long-acting 100 mg	10.50	10	✓ n	n-Eslon
Oral lig 2 mg per ml		100 ml	✓ V	Vockhardt \$29
0.8 = g p 0	29.80			Dramorph
	20.00			Dramorph CDC
			•	S29 S29
Ini E ma nor ml. 1 ml amnaula Un to E ini available on a	DCO 5 20	5	./ 1	
Inj 5 mg per ml, 1 ml ampoule – Up to 5 inj available on a		5 5		<u>Nedsurge</u> Nedsurge
Inj 10 mg per ml, 1 ml ampoule – Up to 5 inj available on a		5 5	_	
Inj 15 mg per ml, 1 ml ampoule – Up to 5 inj available on a		5 5		<u>Nedsurge</u>
Inj 30 mg per ml, 1 ml ampoule - Up to 5 inj available on a	a PSU0.28	Э	<u> </u>	<u>Nedsurge</u>
(YCODONE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing	frequency			
Tab controlled-release 5 mg	2.49	20		Oxycodone Sando
	3.77	28	✓ (	Oxycodone Sando
				S29 S29
	4.04	30	10	OxyContin S29
Tab immediate-release 5 mg		100		Oxycodone Amne
Tab controlled-release 10 mg		20		Oxycodone Sand
- 42 00 m o no 4 n o 10 m o n o 10 m	3.77	28		Oxycodone Sand
	•	_0		S29 S29
Tab immediate release 10 mg	10 77	100		
Tab immediate-release 10 mg		20		Oxycodone Amne
Tab controlled-release 20 mg				Oxycodone Sando
Tab immediate-release 20 mg		100		Oxycodone Amne
Tab controlled-release 40 mg		20	_	Oxycodone Sando
Tab controlled-release 80 mg		20	_	Oxycodone Sando
Cap immediate-release 20 mg		20		OxyNorm
Oral liq 1 mg per ml	37.08	250 ml	✔ (	Oxycodone Lucis
				S29 S29
Inj 10 mg per ml, 1 ml ampoule		5	_	<u>łameln</u>
Inj 10 mg per ml, 2 ml ampoule	8.62	5	_	<u>lameln</u>
Inj 50 mg per ml, 1 ml ampoule	14.90	5	<b>✓</b> <u>F</u>	<u>lameln</u>
xyNorm Cap immediate-release 20 mg to be delisted 1 Marc	ch 2025)			
RACETAMOL WITH CODEINE - Safety medicine; prescrib	ner may determine disr	ensina f	requency	
Tab paracetamol 500 mg with codeine phosphate 8 mg		1,000		Paracetamol +

Paracetamol + Codeine (Relieve)

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	
PETHIDINE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
<ul> <li>Safety medicine; prescriber may determine dispensing free</li> </ul>			_	
Tab 50 mg		10		Noumed Pethidine
Inj 50 mg per ml, 1 ml ampoule - Up to 5 inj available on a P	SO29.88	5	•	DBL Pethidine
lei 50 manual Ombananda - Hata 5 ini analahla ana B	000 00 70	_	,	Hydrochloride
Inj 50 mg per ml, 2 ml ampoule – Up to 5 inj available on a P	SO30.72	5	•	DBL Pethidine
FRAMAROL LIVEROCUL ORIDE				Hydrochloride
FRAMADOL HYDROCHLORIDE	4.05	00		Trom al CD 400
Tab sustained-release 100 mg		20		Tramal SR 100
Tab sustained release 150 mg		20 20		Tramal SR 150 Tramal SR 200
Tab sustained-release 200 mg Cap 50 mg		100		Arrow-Tramadol
Oap 30 mg		100		Allow-Halliadol
Antidepressants				
Cyclic and Related Agents				
AMITRIPTYLINE - Safety medicine; prescriber may determine d	ispensing frequency			
Tab 10 mg		100	/	Arrow-Amitriptyline
Tab 25 mg		100		Arrow-Amitriptyline
Tab 50 mg		100	1	Arrow-Amitriptyline
CLOMIPRAMINE HYDROCHLORIDE - Safety medicine; prescri		isper	sina freau	encv
Tab 10 mg		30		Clomipramine Teva
Tab 25 mg		30		Clomipramine Teva
	39.97	100		Anafranil S29
Cap 10 mg		28		Clomipramine Teva
Cap 25 mg		28		Clomipramine Teva
DOSULEPIN [DOTHIEPIN] HYDROCHLORIDE - Subsidy by en				•
a) Safety medicine; prescriber may determine dispensing fre				
Subsidy by endorsement – Subsidised for patients who we 2019 and the prescription is endorsed accordingly. Pharr exists a record of prior dispensing of dosulepin [dothiepin]	ere taking dosulepin macists may annotate			
Tab 75 mg		30	1	Dosulepin Viatris
Cap 25 mg		50		Dosulepin
•				Viatris S29
MIPRAMINE HYDROCHLORIDE - Safety medicine; prescriber	may determine dispe	nsino	frequency	1
Tab 10 mg		50		Tofranil
•	10.96	100	1	Tofranil
Tab 25 mg	4.93	28	1	Imipramine
•				Crescent S29
	8.80	50	✓	Tofranil
NORTRIPTYLINE HYDROCHLORIDE - Safety medicine; prescr	riber may determine o	lispe	nsing freat	iency
Tab 10 mg	•	100		Norpress
Tab 25 mg		180	_	Norpress
Monoamine-Oxidase Inhibitors (MAOIs) - Non S	elective			
FRANYLCYPROMINE SULPHATE				
CHANGE OF HOWING OUR HATE				
* Tab 10 mg	22.94	50	/	Parnate

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
Monoamine-Oxidase Type A Inhibitors				
MOCLOBEMIDE				
* Tab 150 mg Aurorix to be Principal Supply on 1 February 2025	23.60	60	•	Aurorix
* Tab 300 mg	38.50	60	•	Aurorix
Selective Serotonin Reuptake Inhibitors				
CITALOPRAM HYDROBROMIDE				
* Tab 20 mg	2.86	84	•	Celapram
ESCITALOPRAM			_	
* Tab 10 mg	0.79 1.07	28		Ipca-Escitalopram Escitalopram
	1.07		•	(Ethics)
★ Tab 20 mg	1.49	28	1	Ipca-Escitalopram
LUOXETINE HYDROCHLORIDE				
Tab dispersible 20 mg, scored – Subsidy by endorsement Subsidised by endorsement	2.50	28	•	Fluox
<ol> <li>When prescribed for a patient who cannot swallow accordingly; or</li> <li>When prescribed in a daily dose that is not a multipendorsed. Note: Tablets should be combined with</li> </ol>	ble of 20 mg in which	case	the prescr	iption is deemed to be
₭ Cap 20 mg	3.13	90	1	Arrow-Fluoxetine
PAROXETINE				
₭ Tab 20 mg SERTRALINE	4.11	90	•	<u>Loxamine</u>
₭ Tab 50 mg	0.99	30	✓	Setrona
Fab 100 mg	1.74	30	/	<u>Setrona</u>
Other Antidepressants				
MIRTAZAPINE				
Tab 30 mg	2.60	28		Noumed
Tab 45 mm	0.45	30		Noumed
Tab 45 mg	3.45	28 30		Noumed Noumed
/ENLAFAXINE		00	•	
k Cap 37.5 mg	8.29	84	/	Enlafax XR
* Cap 75 mg	10.32	84		Enlafax XR
* Cap 150 mg	13.95	84	/	Enlafax XR

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

# **Antiepilepsy Drugs**

## **Agents for Control of Status Epilepticus**

DIAZEPAM – Safety medicine; prescriber may determine dispensing frequency Inj 5 mg per ml, 2 ml ampoule – Subsidy by endorsement27.92	5	✓ Hospira
a) Up to 5 inj available on a PSO		
b) Only on a PSO		
<ul> <li>c) PSO must be endorsed "not for anaesthetic procedures".</li> </ul>		
Rectal tubes 5 mg - Up to 5 tube available on a PSO54.58	5	✓ <u>Stesolid</u>
PHENYTOIN SODIUM		
* Inj 50 mg per ml, 2 ml ampoule - Up to 5 inj available on a		
PSO104.58	5	<ul><li>Hospira</li></ul>
* Inj 50 mg per ml, 5 ml ampoule - Up to 5 inj available on a		
PSO154.01	5	<ul><li>Hospira</li></ul>

## **Control of Epilepsy**

CARBAMAZEPINE			
* Tab 200 mg	14.53	100	✓ Tegretol
			✓ Tegretol AU
* Tab long-acting 200 mg	16.98	100	Tegretol CR
	33.96	200	✓ Tegretol CR
* Tab 400 mg	34.58	100	✓ Tegretol
* Tab long-acting 400 mg	39.17	100	Tegretol CR
* Oral liq 20 mg per ml	26.37	250 ml	✓ Tegretol
CLOBAZAM - Safety medicine; prescriber may determine of	ispensing frequency		
Tab 10 mg	9.12	50	✓ Frisium
CLONAZEPAM - Safety medicine; prescriber may determin	e dispensing frequen	cy	
Oral drops 2.5 mg per ml	7.38	10 ml OP	✓ Rivotril
ETHOSUXIMIDE			
Cap 250 mg	78.89	56	✓ Essential
			Ethosuximide S29
	140.88	100	✓ Zarontin
Oral liq 250 mg per 5 ml	56.35	200 ml	✓ Zarontin
GABAPENTIN			
Note: Not subsidised in combination with subsidised pro	egabalin		
* Cap 100 mg	6.45	100	✓ Nupentin
* Cap 300 mg	8.45	100	✓ Nupentin
* Cap 400 mg	10.26	100	✓ Nupentin
LACOSAMIDE - Special Authority see SA2267 on the next	page – Retail pharma	acy	
▲ Tab 50 mg	25.04	14	✓ Vimpat
▲ Tab 100 mg	50.06	14	✓ Vimpat
-	200.24	56	✓ Vimpat
▲ Tab 150 mg	75.10	14	✓ Vimpat
	300.40	56	✓ Vimpat
▲ Tab 200 mg	400.55	56	✓ Vimpat

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	/	Manufacturer

### ⇒SA2267 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Patient has focal epilepsy; and
- 2 Seizures are not adequately controlled by, or patient has experienced unacceptable side effects from, optimal treatment with all of the following: sodium valproate, topiramate, levetiracetam and any two of carbamazepine, lamotrigine and phenytoin sodium (see Note).

Note: Those of childbearing potential are not required to trial phenytoin sodium, sodium valproate, or topiramate. Those who can father children are not required to trial sodium valproate.

**Renewal** from any relevant practitioner. Approvals valid for 24 months where the patient has demonstrated a significant and sustained improvement in seizure rate or severity and/or quality of life compared with that prior to starting lacosamide treatment.

			•		ŭ
	MOTRIGINE			_	
•	Tab dispersible 2 mg		30		Lamictal
•	Tab dispersible 5 mg		30		Lamictal
*	Tab dispersible 25 mg		56		Logem
*	Tab dispersible 50 mg		56		Logem
*	Tab dispersible 100 mg	6.75	56	•	Logem
LE	VETIRACETAM				
	Tab 250 mg	5.84	60	1	Everet
	Tab 500 mg	10.51	60	1	Everet
	Tab 750 mg	16.71	60	1	Everet
	Tab 1,000 mg	21.82	60	1	Everet
	Oral liq 100 mg per ml	44.78	300 ml OP	1	Levetiracetam-AFT
	Inj 100 mg per ml, 5 ml vial	38.95	10	1	Levetiracetam-AFT
PΗ	IENOBARBITONE				
	For phenobarbitone oral liquid refer Standard Formulae, page 2	276			
	Tab 15 mg		500	1	Noumed
	10 10 mg	240.00	300	٠	Phenobarbitone
	Tab 20 mg	200 50	500	./	Noumed
	Tab 30 mg	396.30	500	•	Phenobarbitone
					PHEHODAIDILOHE
	IENYTOIN SODIUM			_	
*	Tab 50 mg		200		Dilantin Infatab
	Cap 30 mg		200		Dilantin
	Cap 100 mg		200		Dilantin
*	Oral liq 30 mg per 5 ml	22.03	500 ml		Dilantin Paediatric
PF	REGABALIN				
	Note: Not subsidised in combination with subsidised gabapent	tin			
*	Cap 25 mg	2.25	56	1	Pregabalin Pfizer
	•	7.80		1	Milpharm S29
*	Cap 75 mg	2.65	56		Pregabalin Pfizer
		8.10			Milpharm S29
*	Cap 150 mg		56		Lyrica
7,1	σαρ 100 mg		00		Pregabalin Pfizer
*	Cap 300 mg	7.38	56		Pregabalin Pfizer
			00	-	ogubumi i medi
	RIMIDONE	07.05	100		Deinsidene Olineet
*	Tab 250 mg	37.35	100	•	Primidone Clinect

## **NERVOUS SYSTEM**

Price)	Subsidised	
Pei		
	•	Manufacturer
100	<b>/</b>	Epilim Crushable
100	<b>/</b>	Épilim
100	<b>/</b>	Épilim
300 r	nl 🗸	Epilim S/F Liquid
	✓	Epilim Syrup
1	✓	Epilim IV
60	1	Diacomit
60	•	Diacomit
	100 100 300 r 1	100 100 300 ml

### ⇒SA2268 Special Authority for Subsidy

**TOPIRAMATE** 

Initial application only from a paediatric neurologist or Practitioner on the recommendation of a paediatric neurologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 Patient has confirmed diagnosis of Dravet syndrome; and

▲ Tab 25 mg .......11.07

2 Seizures have been inadequately controlled by appropriate courses of sodium valproate, clobazam and at least two of the following: topiramate, levetiracetam, ketogenic diet.

Note: Those of childbearing potential are not required to trial sodium valproate or topiramate. Those who can father children are not required to trial sodium valproate.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where the patient continues to benefit from treatment as measured by reduced seizure frequency from baseline.

· ·			✓ Topiramate Actavis
	26.04		✓ Topamax
▲ Tab 50 mg	18.81	60	Arrow-Topiramate
· ·			✓ Topiramate Actavis
	44.26		✓ Topamax
▲ Tab 100 mg	31.99	60	✓ Arrow-Topiramate
•			✓ Topiramate Actavis
	75.25		✓ Topamax
▲ Tab 200 mg	55.19	60	✓ Arrow-Topiramate
•			✓ Topiramate Actavis
	129.85		✓ Topamax
▲ Sprinkle cap 15 mg	20.84	60	✓ Topamax
▲ Sprinkle cap 25 mg	26.04	60	✓ Topamax

### **⇒SA2088** Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Any of the following:
  - 1.1 Patient has infantile spasms: or
  - 1.2 Both:
    - 1.2.1 Patient has epilepsy: and

VIGABATRIN − Special Authority see SA2088 below − Retail pharmacy

Tab 500 mg ......119.30

▲ Powder for oral soln 500 mg per sachet.......71.58

1.2.2 Either:

continued...

60

100

60

Sabril

✓ Sabril

✓ Arrow-Topiramate

Subs	,	ully	Brand or
(Manufactur		sed	Generic
\$	S Per	✓	Manufacturer

continued...

- 1.2.2.1 Seizures are not adequately controlled with optimal treatment with other antiepilepsy agents; or
- 1.2.2.2 Seizures are controlled adequately but the patient has experienced unacceptable side effects from optimal treatment with other antiepilepsy agents; or
- 1.3 Patient has tuberous sclerosis complex; and
- 2 Either:
  - 2.1 Patient is, or will be, receiving regular automated visual field testing (ideally before starting therapy and on a 6-monthly basis thereafter): or
  - 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields...

**Renewal** from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient has demonstrated a significant and sustained improvement in seizure rate or severity and or quality of life; and
- 2 Either
  - 2.1 Patient is receiving regular automated visual field testing (ideally every 6 months) on an ongoing basis for duration of treatment with vigabatrin; or
  - 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields...

## **Antimigraine Preparations**

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 120

## **Acute Migraine Treatment**

RIZATRIPTAN			
Tab orodispersible 10 mg	4.84	30	✓ Rizamelt
SUMATRIPTAN			
Tab 50 mg	14.41	90	<ul><li>Sumagran</li></ul>
Tab 100 mg	22.68	90	✓ Sumagran
Inj 12 mg per ml, 0.5 ml prefilled pen - Maximum of 10 inj per			
prescription	29.80	2 OP	✓ Clustran

### **Prophylaxis of Migraine**

For Beta Adrenoceptor Blockers refer to CARDIOVASCULAR SYSTEM, page 51

**PIZOTIFEN** 

**★** Tab 500 mcg......23.21 100 **✓ Sandomigran** 

## **Antinausea and Vertigo Agents**

For Antispasmodics refer to ALIMENTARY TRACT, page 8

APREPITANT - Special Authority see SA0987 below - Retail pharmacy

Cap 2 × 80 mg and 1 × 125 mg......21.90 3 OP **✓ Emend Tri-Pack** 

Emend Tri-Pack to be Principal Supply on 1 January 2025

### ⇒SA0987 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 12 months where the patient is undergoing highly emetogenic chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.

**Renewal** from any relevant practitioner. Approvals valid for 12 months where the patient is undergoing highly emetogenic chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.

BETAHISTINE DIHYDROCHLORIDE

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
CYCLIZINE HYDROCHLORIDE  Tab 50 mg  Nausicalm to be Principal Supply on 1 February 2025	0.66	10	<b>✓</b> I	Nausicalm
CYCLIZINE LACTATE				
Inj 50 mg per ml, 1 ml ampoule – Up to 10 inj available on a PSO  DOMPERIDONE	16.36	10	_	<u>Hameln</u>
* Tab 10 mg	4.00	100	<b>✓</b> <u>I</u>	<u>Domperidone</u>
HYOSCINE HYDROBROMIDE				<u>Viatris</u>
* Inj 400 mcg per ml, 1 ml ampoule	93.00	10	<b>✓</b>	Martindale S29
Patch 1 mg per 72 hours - Special Authority see SA1998				
below - Retail pharmacy	17.70	2	<b>√</b> 9	Scopoderm TTS
	88.50	10	✓ 9	Scopolamine - Mylan
			✓ 9	Scopolamine -
				Mylan S29 S29

(Scopoderm TTS Patch 1 mg per 72 hours to be delisted 1 January 2025)

(Scopolamine - Mylan S29 S29 Patch 1 mg per 72 hours to be delisted 1 February 2025)

## **⇒SA1998** Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Control of intractable nausea, vomiting, or inability to swallow saliva in the treatment of malignancy or chronic disease where the patient cannot tolerate or does not adequately respond to oral anti-nausea agents; or
- 2 Control of clozapine-induced hypersalivation where trials of at least two other alternative treatments have proven ineffective.

**Renewal** from any relevant practitioner. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

METOCLOPRAMIDE HYDROCHLORIDE		
* Tab 10 mg - Up to 30 tab available on a PSO1.57	100	✓ <u>Metoclopramide</u> <u>Actavis 10</u>
* Inj 5 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO7.00	10	✓ Baxter
ONDANSETRON		
* Tab 4 mg2.27	50	✓ Periset
Tab disp 4 mg - Up to 10 tab available on a PSO0.56	10	✓ Periset ODT
* Tab 8 mg4.10	50	✓ Periset
Tab disp 8 mg - Up to 10 tab available on a PSO0.90	10	✓ Periset ODT
PROCHLORPERAZINE		
* Tab 3 mg buccal5.97	50	
(30.00)		Buccastem
(30.00)		Max Health S29
(30.00)		Prochlorperazine
		Brown & Burk S29
* Tab 5 mg - Up to 30 tab available on a PSO25.00	250	✓ Nausafix
* Inj 12.5 mg per ml, 1 ml - Up to 5 inj available on a PSO25.81	10	✓ Stemetil

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sub	sidised	Generic
\$	Per	1	Manufacturer

## **Antipsychotics**

## General

AMISULPRIDE - Safety medicine; prescriber may determine dispe	ensing frequen	су	
Tab 100 mg	5.84	30	✓ Sulprix
Tab 200 mg	14.47	60	✓ Sulprix
Tab 400 mg	35.06	60	✓ Sulprix
ARIPIPRAZOLE – Safety medicine; prescriber may determine disp	nensina freaue	ncv	
Tab 5 mg		30	✓ Aripiprazole Sandoz
1 ab 3 mg	10.50	00	✓ Ampriprozoic Gandoz  ✓ Ascend
T 1 40	40.50		Aripiprazole S29
Tab 10 mg		30	✓ Aripiprazole Sandoz
Tab 15 mg		30	✓ Aripiprazole Sandoz
Tab 20 mg		30	✓ Aripiprazole Sandoz
Tab 30 mg		30	Aripiprazole Sandoz
CHLORPROMAZINE HYDROCHLORIDE - Safety medicine; pres		ermine dispen	
Tab 25 mg - Up to 30 tab available on a PSO	15.62	100	<ul><li>Largactil</li></ul>
Tab 100 mg - Up to 30 tab available on a PSO	36.73	100	<ul><li>Largactil</li></ul>
Inj 25 mg per ml, 2 ml - Up to 5 inj available on a PSO	30.79	10	<ul><li>Largactil</li></ul>
CLOZAPINE - Hospital pharmacy [HP4]			
Safety medicine; prescriber may determine dispensing frequer	ncv		
Tab 25 mg		50	✓ Clopine
740 25 mg		00	✓ Clozaril
	13.37	100	✓ Clopine
	10.07	100	✓ Clozaril
Tab 50 mg	8.67	50	✓ Clopine
7 45 55 11g	17.33	100	✓ Clopine
Tab 100 mg		50	✓ Clopine
1 3 3 1 3 1 1 3 1 1 3 1 1 1 1 1 1 1 1 1			✓ Clozaril
	34.65	100	✓ Clopine
			✓ Clozaril
Tab 200 mg	34.65	50	✓ Clopine
	69.30	100	✓ Clopine
Suspension 50 mg per ml		100 ml	✓ Versacloz
HALOPERIDOL – Safety medicine; prescriber may determine disp		201	
Tab 500 mcg – Up to 30 tab available on a PSO		100	✓ Serenace
Tab 1.5 mg - Up to 30 tab available on a PSO		100	✓ Serenace
		50	✓ Serenace
Tab 5 mg - Up to 30 tab available on a PSO	29.72	100	✓ Serenace
Oval lia 0 ma nov ml Un to 200 ml available on a DCO		100 100 ml	✓ Serenace
Oral liq 2 mg per ml — Up to 200 ml available on a PSO		100 1111	✓ Serenace
Inj 5 mg per ml, 1 ml ampoule - Up to 5 inj available on a PSC		• •	Serenace
LEVOMEPROMAZINE - Safety medicine; prescriber may determi			
Tab 25 mg (33.8 mg as a maleate)		100	✓ Nozinan (Swiss)
Tab 25 mg as a maleate		100	✓ Nozinan
Tab 100 mg (135 mg as a maleate)		100	✓ Nozinan (Swiss)
Tab 100 mg as a maleate	41.75	100	✓ Nozinan
LEVOMEPROMAZINE HYDROCHLORIDE - Safety medicine; pre	escriber may de	etermine dispe	nsing frequency
Inj 25 mg per ml, 1 ml ampoule		10	✓ Nozinan S29 S29
, 01: ,	-	-	✓ Wockhardt

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	Generic
	\$	Per	1	Manufacturer
LITHIUM CARBONATE - Safety medicine; prescriber may det	ormina dienoneina froa	uono	,	
				Duindal
Tab long-acting 400 mg	82.80	100	•	Priadel
Priadel to be Principal Supply on 1 February 2025			_	
Cap 250 mg	22.36	100	•	Douglas
OLANZAPINE - Safety medicine; prescriber may determine di	spensing frequency			
Tab 2.5 mg		30	1	Zypine
Tab 5 mg		30		Zypine
Tab orodispersible 5 mg		28		Zypine ODT
		30		Zypine
Tab 10 mg				
Tab orodispersible 10 mg	2.89	28	•	Zypine ODT
PERICYAZINE - Safety medicine; prescriber may determine d	ispensing frequency			
Tab 2.5 mg	13.61	100	✓	Neulactil
Tab 10 mg	48.45	100	1	Neulactil
· ·				
QUETIAPINE – Safety medicine; prescriber may determine dis			,	
Tab 25 mg		90		Quetapel
Tab 100 mg		90		Quetapel
Tab 200 mg	10.97	90	/	Quetapel
Tab 300 mg	15.83	90	✓	Quetapel
RISPERIDONE – Safety medicine; prescriber may determine of	dienancina fraguancy			
		20	./	Risperdal
Tab 0.5 mg				•
	2.17	60	_	Risperidone (Teva)
	4.01		•	Risperidone
				Sandoz S29
Tab 1 mg	2.44	60	✓	Risperdal
· ·			1	Risperidone (Teva)
	3.68			Risperidone
	0.00			Sandoz S29
Tale O as a	0.70	00	,	
Tab 2 mg	2.72	60		Risperdal
				Risperidone (Teva)
	5.38		•	Risperidone
				Sandoz S29
Tab 3 mg	4.50	60	1	Risperdal
				Risperidone (Teva)
	8.57			Risperidone
	0.07		-	•
			_	Sandoz S29
Tab 4 mg		60		Risperidone (Teva)
Oral liq 1 mg per ml	10.29	30 m	•	Risperon
ZIPRASIDONE - Safety medicine; prescriber may determine d	lispensina frequency			
Cap 20 mg		60	1	Zusdone
Cap 40 mg		60		Zusdone
_ '		60		Zusdone
Cap 80 mg		60		Zusdone
Cap 80 mg				
ZUCLOPENTHIXOL HYDROCHLORIDE - Safety medicine; po	rescriber may determin	e disp	ensing fre	equency
Tab 10 mg	31.45	100	✓	Clopixol
•				•

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsid	dised	Generic
<b>\$</b>	Per	•	Manufacturer

## **Depot Injections**

ARIPIPRAZOLE – Special Authority see SA2395 below – Retail pharmacy Safety medicine; prescriber may determine dispensing frequency		
Inj 300 mg vial	1	<ul> <li>✓ Abilify Maintena</li> <li>✓ Abilify Maintena</li> <li>S29 S29</li> </ul>
Inj 400 mg vial341.96	1	✓ Abilify Maintena ✓ Abilify Maintena S29 S29

### ⇒SA2395 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Either:

- 1 Either:
  - 1.1 The patient has had an initial Special Authority approval for risperidone depot injection, paliperidone depot injection or olanzapine depot injection; or
  - 1.2 All of the following:
    - 1.2.1 The patient has schizophrenia or other psychotic disorder; and

ELLIDENTHIVOL DECANOATE Cofety medicine: properitor may determine dispension fractions

- 1.2.2 The patient has received treatment with oral atypical antipsychotic agents but has been unable to adhere; and
- 1.2.3 The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months; or
- 2 Patient has been unable to access olanzapine depot injection due to supply issues with olanzapine depot injection, or otherwise would have been started on olanzapine depot injection but has been unable to due to supply issues with olanzapine depot injection.

Notes: The Olanzapine depot injection Special Authority criteria that apply to criterion 2 in this Aripiprazole Special Authority application are as follows:

- The patient has had an initial Special Authority approval for paliperidone depot injection or risperidone depot injection; or
- All of the following:
  - The patient has schizophrenia; and
  - The patient has not been able to adhere with treatment using oral atypical antipsychotic agents; and
  - The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months.

FLUPENTHIXOL DECANOATE — Safety medicine; prescriber m	ay determine disp	ensing treq	uency
Inj 20 mg per ml, 1 ml - Up to 5 inj available on a PSO	13.14	5	✓ Fluanxol
Inj 20 mg per ml, 2 ml - Up to 5 inj available on a PSO	20.90	5	✓ Fluanxol
Inj 100 mg per ml, 1 ml - Up to 5 inj available on a PSO	40.87	5	✓ Fluanxol
HALOPERIDOL DECANOATE - Safety medicine; prescriber ma	y determine dispe	nsing frequ	ency
Inj 50 mg per ml, 1 ml - Up to 5 inj available on a PSO	28.39	5	✓ Haldol
Inj 100 mg per ml, 1 ml - Up to 5 inj available on a PSO	55.90	5	<ul> <li>Haldol Concentrate</li> </ul>
			✓ Haldol
			Decanoas \$29
OLANZAPINE - Special Authority see SA2313 on the next page	- Retail pharmac	y	
a) Safety medicine; prescriber may determine dispensing from	equency		
b) Note - no new patients to be initiated on olanzapine.			
Inj 210 mg vial	252.00	1	✓ Zyprexa Relprevv
Inj 300 mg vial	414.00	1	✓ Zyprexa Relprevv
Inj 405 mg vial	504.00	1	✓ Zyprexa Relprevv



Sub	bsidy	Fully	Brand or
(Manufactu	turer's Price) Subs	sidised	Generic
•	\$ Per	✓	Manufacturer

### ⇒SA2313 Special Authority for Subsidy

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of olanzapine depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

PALIPERIDONE - Special Authority see SA2396 below - Retail pharmacy

Safety medicine; prescriber may determine dispensing frequency

Inj 25 mg syringe	194.25	1	✓ Invega Sustenna
Inj 50 mg syringe	271.95	1	✓ Invega Sustenna
Inj 75 mg syringe	357.42	1	✓ Invega Sustenna
Inj 100 mg syringe	435.12	1	✓ Invega Sustenna
Inj 150 mg syringe		1	✓ Invega Sustenna

### ⇒SA2396 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for risperidone depot injection or olanzapine depot injection or aripiprazole depot injection; or
- 2 All of the following:
  - 2.1 The patient has schizophrenia or other psychotic disorder; and
  - 2.2 Has been unable to adhere to treatment using oral atypical antipsychotic agents; and
  - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

**Renewal** from any relevant practitioner. Approvals valid for 12 months where the initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

## PALIPERIDONE PALMITATE - Special Authority see SA2167 below - Retail pharmacy

Inj 175 mg syringe	815.85	1	Invega Trinza
	1,072.26	1	✓ Invega Trinza
Inj 350 mg syringe	1,305.36	1	✓ Invega Trinza
Inj 525 mg syringe	1,305.36	1	✓ Invega Trinza

### ⇒SA2167 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has schizophrenia: and
- 2 The patient has had an initial Special Authority approval for paliperidone once-monthly depot injection.

**Renewal** from any relevant practitioner. Approvals valid for 12 months where the initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

RISPERIDONE - Special Authority see SA2397 below - Retail pharmacy

### ⇒SA2397 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Fither:

1 The patient has had an initial Special Authority approval for paliperidone depot injection or olanzapine depot injection or aripiprazole depot injection; or

Subs	sidy Full	/ Brand or
(Manufactur	rer's Price) Subsidise	d Generic
\$	Per 🗸	Manufacturer

continued...

- 2 All of the following:
  - 2.1 The patient has schizophrenia or other psychotic disorder; and
  - 2.2 Has not been able to adhere with treatment using oral atypical antipsychotic agents; and
  - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of risperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

ZUCLOPENTHIXOL DECANOATE - Safety medicine; prescriber may determine dispensing frequency ✓ Clopixol

## **Anxiolytics**

BUSPIRONE HYDROCHLORIDE			
* Tab 5 mg	13.95	100	✓ Buspirone Viatris
* Tab 10 mg	12.50	100	✓ Buspirone Viatris
CLONAZEPAM - Safety medicine; prescriber may determine disp	pensing frequency		
Tab 500 mcg	5.64	100	✓ Paxam
Tab 2 mg	10.78	100	✓ Paxam
DIAZEPAM - Safety medicine; prescriber may determine dispens	sing frequency		
Tab 2 mg	95.00	500	✓ Arrow-Diazepam
Tab 5 mg	115.00	500	✓ Arrow-Diazepam
LORAZEPAM - Safety medicine; prescriber may determine dispe	ensing frequency		
Tab 1 mg	10.20	250	✓ Ativan
Ativan to be Principal Supply on 1 February 2025			
Tab 2.5 mg	13.13	100	✓ Ativan
Ativan to be Principal Supply on 1 February 2025			

## **Multiple Sclerosis Treatments**

### ⇒SA2274 Special Authority for Subsidy

Initial application — (Multiple Sclerosis - dimethyl fumarate, fingolimod, glatiramer acetate, interferon beta-1-alpha, interferon beta-1-beta, natalizumab and teriflunomide) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 All of the following:
  - 1.1 Diagnosis of multiple sclerosis (MS) meets the McDonald 2017 diagnostic criteria for MS and has been confirmed by a neurologist; and
  - 1.2 Patient has an EDSS score between 0 6.0; and
  - 1.3 Patient has had at least one significant attack of MS in the previous 12 months or two significant attacks in the past 24 months: and
  - 1.4 All of the following:
    - 1.4.1 Each significant attack must be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the attack, but the neurologist/physician must be satisfied that the clinical features were characteristic); and
    - 1.4.2 Each significant attack is associated with characteristic new symptom(s)/sign(s) or substantially worsening

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	•	Manufacturer

continued...

of previously experienced symptoms(s)/sign(s); and

- 1.4.3 Each significant attack has lasted at least one week and has started at least one month after the onset of a previous attack (where relevant); and
- 1.4.4 Each significant attack can be distinguished from the effects of general fatigue; and is not associated with a fever (T> 37.5°C); and
- 1.4.5 Either:
  - 1.4.5.1 Each significant attack is severe enough to change either the EDSS or at least one of the Kurtze Functional System scores by at least 1 point; or
  - 1.4.5.2 Each significant attack is a recurrent paroxysmal symptom of multiple sclerosis (tonic seizures/spasms, trigeminal neuralgia, Lhermitte's symptom); and
- 1.5 Evidence of new inflammatory activity on an MRI scan within the past 24 months; and
- 1.6 Any of the following:
  - 1.6.1 A sign of that new inflammatory activity on MRI scanning (in criterion 5 immediately above) is a gadolinium enhancing lesion; or
  - 1.6.2 A sign of that new inflammatory activity is a lesion showing diffusion restriction; or
  - 1.6.3 A sign of that new inflammatory is a T2 lesion with associated local swelling; or
  - 1.6.4 A sign of that new inflammatory activity is a prominent T2 lesion that clearly is responsible for the clinical features of a recent attack that occurred within the last 2 years; or
  - 1.6.5 A sign of that new inflammatory activity is new T2 lesions compared with a previous MRI scan; or
- 2 Patient has an active approval for ocrelizumab and does not have primary progressive MS.

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

Renewal — (Multiple Sclerosis - dimethyl fumarate, fingolimod, glatiramer acetate, interferon beta-1-alpha, interferon beta-1-beta, natalizumab and teriflunomide) from any relevant practitioner. Approvals valid for 12 months where patient has had an EDSS score of 0 to 6.0 (inclusive) with or without the use of unilateral or bilateral aids at any time in the last six months (ie the patient has walked 100 metres or more with or without aids in the last six months).

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

DIMETHYL FUMARATE - Special Authority see SA2274 on the previous page - Retail pharmacy

- a) Wastage claimable
- b) Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

Cap 120 mg	520.00	14	Tecfidera
Cap 240 mg	2,000.00	56	✓ Tecfidera

FINGOLIMOD - Special Authority see SA2274 on the previous page - Retail pharmacy

- a) Wastage claimable
- b) Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

Cap 0.5 mg	2.200.00	28	Gilenva

GLATIRAMER ACETATE - Special Authority see SA2274 on the previous page - Retail pharmacy

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

INTERFERON BETA-1-ALPHA - Special Authority see SA2274 on the previous page - Retail pharmacy

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

INTERFERON BETA-1-BETA - Special Authority see SA2274 on the previous page - Retail pharmacy

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

NATALIZUMAB - Special Authority see SA2274 on the previous page - Retail pharmacy

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

# **NERVOUS SYSTEM**

Subsidy	Fully		Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	/	Manufacturer	

TERIFLUNOMIDE - Special Authority see SA2274 on page 143 - Retail pharmacy

a) Wastage claimable

b) Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

659.90 **✓ Aubagio** 

(Aubagio Tab 14 mg to be delisted 1 April 2025)

# **Multiple Sclerosis Treatments - Other**

OCRELIZUMAB - Special Authority see SA2273 below - Retail pharmacy

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

#### ⇒SA2273 Special Authority for Subsidy

Initial application — (Multiple Sclerosis - ocrelizumab) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Fither:

- 1 All of the following:
  - 1.1 Diagnosis of multiple sclerosis (MS) meets the McDonald 2017 diagnostic criteria for MS and has been confirmed by a neurologist; and
  - 1.2 Patient has an EDSS score between 0 6.0; and
  - 1.3 Patient has had at least one significant attack of MS in the previous 12 months or two significant attacks in the past 24 months; and
  - 1.4 All of the following:
    - 1.4.1 Each significant attack must be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the attack, but the neurologist/physician must be satisfied that the clinical features were characteristic); and
    - 1.4.2 Each significant attack is associated with characteristic new symptom(s)/sign(s) or substantially worsening of previously experienced symptoms(s)/sign(s); and
    - 1.4.3 Each significant attack has lasted at least one week and has started at least one month after the onset of a previous attack (where relevant); and
    - 1.4.4 Each significant attack can be distinguished from the effects of general fatigue; and is not associated with a fever (T> 37.5°C); and
    - 1.4.5 Fither:
      - 1.4.5.1 Each significant attack is severe enough to change either the EDSS or at least one of the Kurtze Functional System scores by at least 1 point; or
      - 1.4.5.2 Each significant attack is a recurrent paroxysmal symptom of multiple sclerosis (tonic seizures/spasms, trigeminal neuralgia, Lhermitte's symptom); and
  - 1.5 Evidence of new inflammatory activity on an MRI scan within the past 24 months; and
  - 1.6 Any of the following:
    - 1.6.1 A sign of that new inflammatory activity on MRI scanning (in criterion 5 immediately above) is a gadolinium enhancing lesion; or
    - 1.6.2 A sign of that new inflammatory activity is a lesion showing diffusion restriction; or
    - 1.6.3 A sign of that new inflammatory is a T2 lesion with associated local swelling; or
    - 1.6.4 A sign of that new inflammatory activity is a prominent T2 lesion that clearly is responsible for the clinical features of a recent attack that occurred within the last 2 years; or
    - 1.6.5 A sign of that new inflammatory activity is new T2 lesions compared with a previous MRI scan; or
- 2 Patient has an active Special Authority approval for either dimethyl fumarate, fingolimod, glatiramer acetate, interferon beta-1-alpha, interferon beta-1-beta, natalizumab or teriflunomide.



Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

Renewal — (Multiple Sclerosis - ocrelizumab) from any relevant practitioner. Approvals valid for 12 months where patient has had an EDSS score of 0 to 6.0 (inclusive) with or without the use of unilateral or bilateral aids at any time in the last six months (ie the patient has walked 100 metres or more with or without aids in the last six months).

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

Initial application — (Primary Progressive Multiple Sclerosis) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Diagnosis of primary progressive multiple sclerosis (PPMS) meets the 2017 McDonald criteria and has been confirmed by a neurologist; and
- 2 Patient has an EDSS 2.0 (score equal to or greater than 2 on pyramidal functions) to EDSS 6.5; and
- 3 Patient has no history of relapsing remitting multiple sclerosis.

Renewal — (Primary Progressive Multiple Sclerosis) from any relevant practitioner. Approvals valid for 12 months where patient has had an EDSS score of less than or equal to 6.5 at any time in the last six months (ie patient has walked 20 metres with bilateral assistance/aids, without rest in the last six months).

# **Sedatives and Hypnotics**

# ⇒SA1666 Special Authority for Subsidy

**Initial application** only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with persistent and distressing insomnia secondary to a neurodevelopmental disorder (including, but not limited to, autism spectrum disorder or attention deficit hyperactivity disorder)\*; and
- 2 Behavioural and environmental approaches have been tried and were unsuccessful, or are inappropriate; and
- 3 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day; and
- 4 Patient is aged 18 years or under\*.

**Renewal** only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is aged 18 years or under\*; and
- 2 Patient has demonstrated clinically meaningful benefit from funded modified-release melatonin (clinician determined); and
- 3 Patient has had a trial of funded modified-release melatonin discontinuation within the past 12 months and has had a recurrence of persistent and distressing insomnia; and
- 4 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day.

Note: Indications marked with \* are unapproved indications.

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	
	\$	Per		Manufacturer
MIDAZOLAM – Safety medicine; prescriber may determine dispe	ensing frequency			
Inj 1 mg per ml, 5 ml ampoule	. ,	10	1	Midazolam-Baxter
, 31 , 1	16.75		1	Midazolam Viatris
Inj 1 mg per ml, 5 ml plastic ampoule - Up to 10 inj available	i			
on a PSO		10	1	Pfizer
On a PSO for status epilepticus use only. PSO must be	endorsed for status e	pilept	ticus use	only.
Inj 5 mg per ml, 1 ml plastic ampoule - Up to 10 inj available	i			•
on a PSO		10	1	Midazolam-Pfizer
On a PSO for status epilepticus use only. PSO must be	endorsed for status e	pilept	ticus use	only.
Inj 5 mg per ml, 3 ml ampoule - Brand switch fee payable				•
(Pharmacode 2695863) - see page 273 for details	4.75	5	1	Midazolam-Baxter
, , ,	5.50		1	Midazolam Viatris
Inj 5 mg per ml, 3 ml plastic ampoule - Up to 5 inj available	on			
a PSO		5	1	Pfizer
On a PSO for status epilepticus use only. PSO must be	endorsed for status e	pilept	ticus use	only.
PHENOBARBITONE SODIUM - Special Authority see SA1386 b				•
Inj 200 mg per ml, 1 ml ampoule		10	1	Max Health \$29
SA1386 Special Authority for Subsidy	110.07	10	•	IVIAA I ICAILII 029
NSA1386  Special Authority for Subsidy				

# **⇒SA1386** Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Both:

- 1 For the treatment of terminal agitation that is unresponsive to other agents; and
- 2 The applicant is part of a multidisciplinary team working in palliative care.

TEMAZEPAM - Safety medicine; prescriber may determine di	spensing frequency		
Tab 10 mg	1.40	25	✓ Normison
ZOPICLONE - Safety medicine; prescriber may determine dis	pensing frequency		
Tab 7.5 mg	21.85	500	Zopiclone Actavis
Zopiclone Actavis to be Principal Supply on 1 Februar	y 2025		-

# **Spinal Muscular Atrophy**

NUSINERSEN − PCT only − Special Authority see SA2174 below
Inj 12 mg per 5 ml vial .......120,000.00 1 ✓ Spinraza

#### ⇒SA2174 Special Authority for Subsidy

Initial application — (spinal muscular atrophy (SMA)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has genetic documentation of homozygous SMN1 gene deletion, homozygous SMN1 point mutation, or compound heterozygous mutation; and
- 2 Patient is 18 years of age or under; and
- 3 Either:
  - 3.1 Patient has experienced the defined signs and symptoms of SMA type I, II or IIIa prior to three years of age; or
  - 3.2 Both:
    - 3.2.1 Patient is pre-symptomatic; and
    - 3.2.2 Patient has three or less copies of SMN2.

Renewal — (spinal muscular atrophy (SMA)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:



Subsidy	Subsidy Fully		Brand or
(Manufacturer's Price)		Subsidised	Generic
 \$	Per	1	Manufacturer

continued...

All of the following:

- 1 There has been demonstrated maintenance of motor milestone function since treatment initiation; and
- 2 Patient does not require invasive permanent ventilation (at least 16 hours per day) in the absence of a potentially reversible cause while being treated with nusinersen; and
- 3 Nusinersen not to be administered in combination other SMA disease modifying treatments or gene therapy.

#### RISDIPLAM - [Xpharm] - Special Authority see SA2203 below

Note: the supply of risdiplam is via Pharmac's approved direct distribution supply. Further details can be found on Pharmac's website https://pharmac.govt.nz/risdiplam

Powder for oral soln 750 mcg per ml, 60 mg per bottle......14,100.00 80 ml OP ✓ Evrysdi

## **⇒SA2203** Special Authority for Subsidy

Initial application — (spinal muscular atrophy (SMA)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has genetic documentation of homozygous SMN1 gene deletion, homozygous SMN1 point mutation, or compound heterozygous mutation; and
- 2 Patient is 18 years of age or under; and
- 3 Either:
  - 3.1 Patient has experienced the defined signs and symptoms of SMA type I, II or IIIa prior to three years of age; or
  - 3.2 Both:
    - 3.2.1 Patient is pre-symptomatic; and
    - 3.2.2 Patient has three or less copies of SMN2.

Renewal — (spinal muscular atrophy (SMA)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 There has been demonstrated maintenance of motor milestone function since treatment initiation; and
- 2 Patient does not require invasive permanent ventilation (at least 16 hours per day) in the absence of a potentially reversible cause while being treated with risdiplam; and
- 3 Risdiplam not to be administered in combination other SMA disease modifying treatments or gene therapy.

# **Stimulants/ADHD Treatments**

ATOMOXETINE			
Cap 10 mg	43.02	28	✓ APO-Atomoxetine
Cap 18 mg	45.57	28	✓ APO-Atomoxetine
Cap 25 mg	44.30	28	✓ APO-Atomoxetine
Cap 40 mg	46.21	28	✓ APO-Atomoxetine
Cap 60 mg	51.31	28	✓ APO-Atomoxetine
Cap 80 mg	65.20	28	✓ APO-Atomoxetine
Cap 100 mg	65.71	28	✓ APO-Atomoxetine
DEXAMFETAMINE SULFATE – Special Authority see SA24  a) Only on a controlled drug form b) Safety medicine; prescriber may determine dispensin		Retail phar	macy
Tab 5 mg	29.80	100	✓ <u>Noumed</u> Dexamfetamine

Subsidy		Fully
(Manufacturer's Price)		Subsidised
\$	Per	✓

Brand or Generic Manufacturer

## ⇒SA2410 Special Authority for Subsidy

Initial application — (ADHD in patients aged 5 years or over) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid without further renewal unless notified for applications meeting the following criteria: All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) in patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
  - 3.1 Applicant is a paediatrician or psychiatrist; or
  - 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients aged under 5 years) only from a paediatrician or psychiatrist. Approvals valid without further renewal unless notified for applications meeting the following criteria: Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) in patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid without further renewal unless notified where the patient suffers from narcolepsy.

LISDEXAMFETAMINE DIMESILATE - Special Authority see SA2415 below - Retail pharmacy

- a) Only on a controlled drug form
- b) Safety medicine; prescriber may determine dispensing frequency

Cap 30 mg - No more than 1 cap per day	60.00	30	Vyvanse
Cap 50 mg	60.00	30	✓ Vyvanse
Cap 70 mg	60.00	30	✓ Vyvanse

#### ⇒SA2415 Special Authority for Subsidy

**Initial application** only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid without further renewal unless notified for applications meeting the following criteria: Either:

- 1 Patient is currently on treatment with lisdexamfetamine dimesilate and met all remaining criteria prior to commencing treatment: or
- 2 All of the following:
  - 2.1 ADHD (Attention Deficit and Hyperactivity Disorder); and
  - 2.2 Diagnosed according to DSM-V or ICD 11 criteria; and
  - 2.3 Either:
    - 2.3.1 Applicant is a paediatrician or psychiatrist; or
    - 2.3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing; and
  - 2.4 Any of the following:
    - 2.4.1 Patient is taking a currently subsidised formulation of atomoxetine or methylphenidate hydrochloride (extended-release) and has not received sufficient benefit or has experienced intolerable side effects; or
    - 2.4.2 Patient is taking a currently subsidised formulation of dexamfetamine sulfate (immediate-release) which has not been effective due to significant administration and/or treatment adherence difficulties; or
    - 2.4.3 There is significant concern regarding the risk of diversion or abuse of immediate release dexamfetamine sulfate: or
    - 2.4.4 Patient is taking a currently subsidised formulation of methylphenidate hydrochloride (immediate-release or sustained release) which has not been effective due to significant administration and/or treatment



Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Por 🗸	Manufacturer

continued...

adherence difficulties: or

- 2.4.5 There is significant concern regarding the risk of diversion or abuse of immediate release methylphenidate hydrochloride; or
- 2.4.6 Both:
  - 2.4.6.1 Patient would have been prescribed a subsidised formulation of methylphenidate hydrochloride (extended-release) but has been unable to access due to supply issues with methylphenidate hydrochloride (extended-release); and
  - 2.4.6.2 Other alternative stimulant presentations (methylphenidate or dexamfetamine) are not appropriate;
- 2.5 Lisdexamfetamine dimesilate is not to be used in combination with another funded methylphenidate presentation.

# METHYLPHENIDATE HYDROCHLORIDE - Special Authority see SA2411 below - Retail pharmacy

- a) Only on a controlled drug form
- b) Safety medicine; prescriber may determine dispensing frequency 30 ✓ Rubifen ✓ Ritalin Tab immediate-release 10 mg......3.00 30 ✓ Rubifen ✓ Methylphenidate ER 30 - Teva ✓ Rubifen 30 30 ✓ Rubifen SR ✓ Methylphenidate ER 30 - Teva ✓ Methylphenidate ER 30 - Teva Tab extended-release 54 mg......22.25 30 Methylphenidate ER - Teva

# ⇒SA2411 Special Authority for Subsidy

Initial application — (ADHD in patients aged 5 years or over) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid without further renewal unless notified for applications meeting the following criteria: All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) in patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Fither:
  - 3.1 Applicant is a paediatrician or psychiatrist; or
  - 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients aged under 5 years) only from a paediatrician or psychiatrist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

- Both:
  - 1 ADHD (Attention Deficit and Hyperactivity Disorder) in patients under 5 years of age; and
  - 2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy\*) only from a neurologist or respiratory specialist. Approvals valid without further renewal unless notified where the patient suffers from narcolepsy.

Note: \*narcolepsy is not a registered indication for Methylphenidate ER – Teva.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	/	Manufacturer	

METHYLPHENIDATE HYDROCHLORIDE EXTENDED-RELEASE - Special Authority see SA2412 below - Retail pharmacy

- a) Only on a controlled drug form
- b) Safety medicine; prescriber may determine dispensing frequency

b) Salety medicine, prescriber may determine dispensing in	equency		
Tab extended-release 18 mg	58.96	30	Concerta
Tab extended-release 27 mg	65.44	30	<ul><li>Concerta</li></ul>
Tab extended-release 36 mg	71.93	30	<ul><li>Concerta</li></ul>
Tab extended-release 54 mg	86.24	30	<ul><li>Concerta</li></ul>
Cap modified-release 10 mg		30	Ritalin LA
Cap modified-release 20 mg		30	Ritalin LA
Cap modified-release 30 mg		30	Ritalin LA
Cap modified-release 40 mg		30	✓ Ritalin LA

# ⇒SA2412 Special Authority for Subsidy

Initial application — (ADHD) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder); and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Fither:
  - 3.1 Applicant is a paediatrician or psychiatrist; or
  - 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing; and
- 4 Either:
  - 4.1 Patient is taking a currently subsidised formulation of methylphenidate hydrochloride (immediate-release or sustained-release) which has not been effective due to significant administration and/or difficulties with adherence; or
  - 4.2 There is significant concern regarding the risk of diversion or abuse of immediate-release methylphenidate hydrochloride.

MODAFINIL – Special Authority see SA2413 below – Retail pharma	acy		
Tab 100 mg	14.27	30	Modafinil Max
•			Health
	29.13	60	✓ Modaviqil

(Modavigil Tab 100 mg to be delisted 1 May 2025)

# ⇒SA2413 Special Authority for Subsidy

**Initial application** only from a neurologist or respiratory specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

# All of the following:

- 1 The patient has a diagnosis of narcolepsy and has excessive daytime sleepiness associated with narcolepsy occurring almost daily for three months or more; and
- 2 Either:
  - 2.1 The patient has a multiple sleep latency test with a mean sleep latency of less than or equal to 10 minutes and 2 or more sleep onset rapid eye movement periods; or
  - 2.2 The patient has at least one of: cataplexy, sleep paralysis or hypnagogic hallucinations; and
- 3 Either:
  - 3.1 An effective dose of a subsidised formulation of methylphenidate or dexamfetamine has been trialled and discontinued because of intolerable side effects; or
  - 3.2 Methylphenidate and dexamfetamine are contraindicated.



	(Manufacturer's Price) \$	Per	Subsidised 🗸	Generic Manufacturer
Treatments for Dementia				
DONEPEZIL HYDROCHLORIDE				
* Tab 5 mg	3.70	84	✓	Ipca-Donepezil
* Tab 10 mg	5.50	84	✓	Ipca-Donepezil
RIVASTIGMINE - Special Authority see SA1488 below - Retail	pharmacv			
Patch 4.6 mg per 24 hour		30	•	Rivastigmine Patch BNM 5
	90.00		✓	Exelon Patch 5
Rivastigmine Patch BNM 5 to be Principal Supply on 1 N	larch 2025			
Patch 9.5 mg per 24 hour	49.40	30	✓	Rivastigmine Patch BNM 10
	90.00		✓	Exelon Patch 10
Rivastigmine Patch BNM 10 to be Principal Supply on 1	March 2025			
(Exelon Patch 5 Patch 4.6 mg per 24 hour to be delisted 1 March	2025)			
(Exelon Patch 10 Patch 9.5 mg per 24 hour to be delisted 1 Marc	,			
OA4400 On a dal Authority for Outsalds				

Subsidy

Fully

Brand or

## ⇒SA1488 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The patient has been diagnosed with dementia; and
- 2 The patient has experienced intolerable nausea and/or vomiting from donepezil tablets.

**Renewal** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate; and
- 2 The patient has demonstrated a significant and sustained benefit from treatment.

# Treatments for Substance Dependence

BUPRENORPHINE WITH NALOXONE - Special Authority see SA1203 below - Retail pharmacy

- a) No patient co-payment payable
- b) Safety medicine; prescriber may determine dispensing frequency

Naloxone BNM

# ⇒SA1203 Special Authority for Subsidy

**Initial application — (Detoxification)** from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health..

**Initial application — (Maintenance treatment)** from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient will not be receiving methadone; and

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidi	sed	Generic
\$	Per	•	Manufacturer

continued...

- 3 Patient is currently enrolled in an opioid substitution treatment program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient has previously trialled but failed detoxification with buprenorphine with naloxone with relapse back to opioid use and another attempt is planned; and
- 3 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is or has been receiving maintenance therapy with buprenorphine with naloxone (and is not receiving methadone); and
- 2 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health or is a medical practitioner authorised by the service to manage treatment in this patient.

Renewal — (Maintenance treatment where the patient has previously had an initial application for detoxification) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient received but failed detoxification with buprenorphine with naloxone; and
- 2 Maintenance therapy with buprenorphine with naloxone is planned (and patient will not be receiving methadone); and
- 3 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Tab modified-release 150 mg	15.00	30	✓ <u>Zyban</u>
DISULFIRAM			
Tab 200 mg	236.40	100	✓ Antabuse
NALTREXONE HYDROCHLORIDE - Special Authority see \$	SA1408 below – Reta	il pharmacy	
Tab 50 mg	77.77	28	✓ Naltrexone AOP S29
	83.33	30	✓ Naltraccord
	102.60		✓ Naltrexone Max
			Health S29
	138.88	50	✓ Revia S29

#### ⇒SA1408 Special Authority for Subsidy

BLIBBODION HADDOCHI ODIDE

**Initial application** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient is currently enrolled in a recognised comprehensive treatment programme for alcohol dependence; and
- 2 Applicant works in or with a community Alcohol and Drug Service contracted to Health NZ or accredited against the New Zealand Alcohol and Other Drug Sector Standard or the National Mental Health Sector Standard.

Renewal from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

1 Compliance with the medication (prescriber determined); and



Subsidy		Fully	Brand or
(Manufacturer's Price)	Sı	ubsidised	Generic
\$	Per	/	Manufacturer

continued...

- 2 Any of the following:
  - 2.1 Patient is still unstable and requires further treatment; or
  - 2.2 Patient achieved significant improvement but requires further treatment; or
  - 2.3 Patient is well controlled but requires maintenance therapy.

#### NICOTINE

- a) Nicotine will not be funded in amounts less than 4 weeks of treatment.
- b) Note: Direct Provision by a pharmacist permitted under the provisions in Part I of Section A. Patch 7 mg - Up to 28 patch available on a PSO .......19.62 ✓ Habitrol Patch 14 mg - Up to 28 patch available on a PSO ......21.57 28 ✓ Habitrol Patch 14 mg for direct distribution only - [Xpharm]......12.49 7 ✓ Habitrol Patch 21 mg - Up to 28 patch available on a PSO ......24.72 28 ✓ Habitrol Patch 21 mg for direct distribution only - [Xpharm]......13.19 7 ✓ Habitrol Lozenge 1 mg - Up to 216 loz available on a PSO......22.53 216 ✓ Habitrol 36 ✓ Habitrol ✓ Habitrol Lozenge 2 mg - Up to 216 loz available on a PSO......24.68 216 36 ✓ Habitrol Gum 2 mg (Fruit) - Up to 204 piece available on a PSO ......23.02 204 ✓ Habitrol Gum 2 mg (Fruit) for direct distribution only - [Xpharm].......17.57 96 ✓ Habitrol Gum 2 mg (Mint) - Up to 204 piece available on a PSO......23.02 204 ✓ Habitrol Gum 2 mg (Mint) for direct distribution only - [Xpharm]......17.57 96 ✓ Habitrol Gum 4 mg (Fruit) - Up to 204 piece available on a PSO ......25.98 204 ✓ Habitrol Gum 4 mg (Fruit) for direct distribution only - [Xpharm]......23.87 ✓ Habitrol 96 Gum 4 mg (Mint) – Up to 204 piece available on a PSO......25.98 204 ✓ Habitrol Gum 4 mg (Mint) for direct distribution only - [Xpharm]......23.87 96 ✓ Habitrol

VARENICLINE TARTRATE - Special Authority see SA1845 below - Retail pharmacy

- a) A maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval, including the starter pack
- b) Varenicline will not be funded in amounts less than 4 weeks of treatment.
- c) The 6-month time period in which a patient can receive a funded 12-week course of varenicline tartrate starts from the date the Special Authority is approved.

Tab 0.5 mg × 11 and 1 mg × 42	16.67	53 OP	✓ Varenicline Pfizer
Tab 1 mg	17.62	56	✓ Varenicline Pfizer

#### **⇒SA1845** Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 Either:
  - 3.1 The patient has tried but failed to quit smoking after at least two separate trials of nicotine replacement therapy, at least one of which included the patient receiving comprehensive advice on the optimal use of nicotine replacement therapy; or
  - 3.2 The patient has tried but failed to guit smoking using bupropion or nortriptyline; and
- 4 The patient has not had a Special Authority for varenicline approved in the last 6 months; and
- 5 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and

# NERVOUS SYSTEM

\$ Per ✔ Manufacturer		Subsidy (Manufacturer's Price)	Subsi Per	Fully dised	Brand or Generic Manufacturer
-----------------------	--	-----------------------------------	--------------	----------------	-------------------------------------

continued...

- 6 The patient is not pregnant; and
- 7 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

**Renewal** from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 It has been 6 months since the patient's previous Special Authority was approved; and
- 4 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 5 The patient is not pregnant; and
- 6 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

The patient must not have had an approval in the past 6 months.

Notes: a maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval.

This includes the 4-week 'starter' pack.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

# **Chemotherapeutic Agents**

# Alkylating Agents

BENDAMUSTINE HYDROCHLORIDE – PCT only – Specialist – Special A		2398 below  Bendamustine
Inj 25 mg vial50.	U5 I	Sandoz
77.	00	✓ Ribomustin
Inj 100 mg vial200.	20 1	<ul><li>Bendamustine Sandoz</li></ul>
308.	00	✓ Ribomustin
Inj 1 mg for ECP3.	23 1 mg	g ✓ Baxter

# ⇒SA2398 Special Authority for Subsidy

Initial application — (CLL\*) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has chronic lymphocytic leukaemia requiring treatment; and
- 2 Patient has ECOG performance status of 0-2; and
- 3 Bendamustine is to be administered at a maximum dose of 100 mg/m² on days 1 and 2 every 4 weeks for a maximum of 6 cycles.

Note: Indication marked with a \* includes indications that are unapproved. 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL).

Initial application — (Indolent, Low-grade lymphomas) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has indolent low grade NHL requiring treatment; and
- 2 The patient has ECOG performance status of 0-2; and
- 3 Any of the following:
  - 3.1 Both:
    - 3.1.1 Patient is treatment naive; and
    - 3.1.2 Bendamustine is to be administered for a maximum of 6 cycles (in combination with rituximab when CD20+); or
  - 3.2 Both:
    - 3.2.1 Patient is refractory to or has relapsed within 12 months of a rituximab containing combined chemo-immunotherapy regimen; and
    - 3.2.2 Bendamustine is to be administered in combination with obinutuzumab for a maximum of 6 cycles; or
  - 3.3 All of the following:
    - 3.3.1 The patient has not received prior bendamustine therapy; and
    - 3.3.2 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
    - 3.3.3 Patient has had a rituximab treatment-free interval of 12 months or more; or
  - 3.4 Bendamustine is to be administered as monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Renewal — (Indolent, Low-grade lymphomas) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: Either:

- 1 Both:
  - 1.1 Patient is refractory to or has relapsed within 12 months of rituximab in combination with bendamustine; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sul	bsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1.2 Bendamustine is to be administered in combination with obinutuzumab for a maximum of 6 cycles; or
- 2 Both:
  - 2.1 Patients have not received a bendamustine regimen within the last 12 months; and
  - 2.2 Fither:
    - 2.2.1 Both:
      - 2.2.1.1 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
      - 2.2.1.2 Patient has had a rituximab treatment-free interval of 12 months or more; or
    - 2.2.2 Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Note: 'indolent, low-grade lymphomas' includes follicular, mantle cell, marginal zone and lymphoplasmacytic/ Waldenstrom's macroglobulinaemia.

Initial application — (Hodgkin's lymphoma\*) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has Hodgkin's lymphoma requiring treatment; and
- 2 Patient has a ECOG performance status of 0-2; and
- 3 Patient has received one prior line of chemotherapy; and
- 4 Patient's disease relapsed or was refractory following prior chemotherapy; and
- 5 Bendamustine is to be administered in combination with gemcitabine and vinorelbine (BeGeV) at a maximum dose of no greater than 90 mg/m2 twice per cycle, for a maximum of four cycles.

Note: Indications marked with \* are unapproved indications.

Tab 2 mg	89.25	100	✓ Myleran
CARBOPLATIN - PCT only - Specialist			-
Inj 10 mg per ml, 45 ml vial	25.73	1	✓ Carboplatin Accord
, ,	32.59		✓ DBL Carboplatin
	48.50		✓ Carbaccord
Inj 1 mg for ECP	0.06	1 mg	✓ Baxter
CARMUSTINE - PCT only - Specialist			
Inj 100 mg vial	710.00	1	✓ BiCNU
, •			✓ BiCNU S29 S29
			✓ Novadoz S29
Inj 100 mg for ECP	710.00	100 mg OP	✓ Baxter
CHLORAMBUCIL - PCT - Retail pharmacy-Specialist		-	
Tab 2 mg	29.06	25	✓ Leukeran FC
CISPLATIN – PCT only – Specialist			
Inj 1 mg per ml, 50 ml vial	9.45	1	✓ Cisplatin Accord
ing i mg por mi, oo mi vidi	15.00	•	✓ Cisplatin Ebewe
Inj 1 mg per ml, 100 ml vial		1	✓ Cisplatin Accord
.,	21.00	•	✓ Cisplatin Ebewe
	29.66		✓ DBL Cisplatin
Inj 1 mg for ECP	0.31	1 mg	✓ Baxter

	Subsidy		Fully	
(	(Manufacturer's Price)	Per	Subsidised	
DVOLORUGORUANIDE	Ψ	1 61		Manuacturei
CYCLOPHOSPHAMIDE	145.00			Ovelenev
Tab 50 mg - PCT - Retail pharmacy-Specialist		50		Cyclonex
Inj 1 g vial - PCT - Retail pharmacy-Specialist		1		Endoxan
Fadeven to be Driveinal County on 1 February 2005	127.80	6	•	Cytoxan
Endoxan to be Principal Supply on 1 February 2025	05.00			Fadavan
Inj 2 g vial – PCT only – Specialist	95.06	1	•	Endoxan
Endoxan to be Principal Supply on 1 February 2025	0.05	4		Davidan
Inj 1 mg for ECP - PCT only - Specialist	0.05	1 mg	•	Baxter
FOSFAMIDE - PCT only - Specialist			_	
lnj 1 g		1		Holoxan
Inj 2 g		1		Holoxan
Inj 1 mg for ECP	0.10	1 mg	•	Baxter
OMUSTINE - PCT - Retail pharmacy-Specialist				
Cap 10 mg	132.59	20	1	CeeNU
Cap 40 mg		20		CeeNU
1 0	880.00		/	Medac S29
CeeNU Cap 10 mg to be delisted 1 January 2025)				
CeeNU Cap 40 mg to be delisted 1 January 2025)				
, ,				
MELPHALAN	40.70	0.5	,	A II
Tab 2 mg - PCT - Retail pharmacy-Specialist		25		Alkeran
Inj 50 mg - PCT only - Specialist	48.25	1		Megval S29
				Melpha
	67.80		•	Alkeran
DXALIPLATIN - PCT only - Specialist				
Inj 100 mg vial	25.01	1	✓	Oxaliplatin Actavis
				100
	110.00		✓	Oxaliplatin Ebewe
Inj 5 mg per ml, 20 ml vial	33.35	1	✓	<b>Alchemy Oxaliplatin</b>
	46.32		✓	Oxaliplatin Accord
Inj 1 mg for ECP	0.35	1 mg	✓	Baxter
THIOTEPA - PCT only - Specialist				
Inj 15 mg vial	CBS	1	1	Bedford S29
ing to mg viai				Max Health S29
	200.00			THIO-TEPA S29
het 400 mm stell	398.00			Tepadina
Inj 100 mg vial		1		Max Health \$29
	1,800.00		•	Tepadina
Antimetabolites				
AZACITIDINE - PCT only - Specialist - Special Authority see SA	2141 below			
Inj 100 mg vial		1	1	Azacitidine Dr
ing 100 mg viai			,	Reddy's
Inj 1 mg for ECP	0.54	1 mg	J	Baxter
SA21/11 Special Authority for Subsidy		i iliy	•	DUALCI

**⇒SA2141** Special Authority for Subsidy

Initial application only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsi	dised	Generic	
\$	Per	✓	Manufacturer	

continued...

# All of the following:

- 1 Any of the following:
  - 1.1 The patient has International Prognostic Scoring System (IPSS) intermediate-2 or high risk myelodysplastic syndrome; or
  - 1.2 The patient has chronic myelomonocytic leukaemia (10%-29% marrow blasts without myeloproliferative disorder); or
  - 1.3 The patient has acute myeloid leukaemia with 20-30% blasts and multi-lineage dysplasia, according to World Health Organisation Classification (WHO); and
- 2 The patient has performance status (WHO/ECOG) grade 0-2; and
- 3 The patient has an estimated life expectancy of at least 3 months.

**Renewal** only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

OALOUIN FOLINATE		
CALCIUM FOLINATE Tab 15 mg - PCT - Retail pharmacy-Specialist135.33	10	✓ DBL Leucovorin Calcium
Inj 3 mg per ml, 1 ml - PCT - Retail pharmacy-Specialist	5	✓ Hospira
Inj 10 mg per ml, 5 ml vial - PCT - Retail pharmacy-Specialist7.28	1	<ul><li>Calcium Folinate Sandoz</li></ul>
		✓ Calcium Folinate Sandoz S29 \$29
36.48	5	✓ Eurofolic S29
Inj 50 mg - PCT - Retail pharmacy-Specialist72.80	10	✓ Leucovorin
,		Pharmacia \$29
Inj 10 mg per ml, 10 ml vial - PCT only - Specialist9.49	1	✓ Calcium Folinate Sandoz
47.45	5	✓ Eurofolic S29
Inj 100 mg - PCT only - Specialist7.33	1	<ul><li>Calcium Folinate Ebewe</li></ul>
94.90	10	✓ Leucovorin  Pharmacia S29
Inj 300 mg - PCT only - Specialist21.55	1	✓ Leucovorin DBL S29
22.51		✓ Calcium Folinate Ebewe
Inj 10 mg per ml, 35 ml vial - PCT only - Specialist25.14	1	<ul><li>Calcium Folinate Sandoz</li></ul>
		✓ Calcium Folinate Sandoz S29 S29
Inj 1 g - PCT only - Specialist67.51	1	✓ Calcium Folinate Ebewe
Inj 10 mg per ml, 100 ml vial - PCT only - Specialist72.00	1	✓ Calcium Folinate Sandoz
		✓ Eurofolic S29
Inj 1 mg for ECP - PCT only - Specialist0.06	1 mg	✓ Baxter

	Subsidy		Fully Brand or
	(Manufacturer's		idised Generic
	\$	Per	✓ Manufacturer
CAPECITABINE – Retail pharmacy-Specialist			
Tab 150 mg		60	✓ Capecitabine Viatris
Tab 500 mg	46.50	120	<ul> <li>Capecitabine Viatris</li> </ul>
CLADRIBINE - PCT only - Specialist			
Inj 1 mg per ml, 10 ml		1	✓ Leustatin
Inj 10 mg for ECP	749.96	10 mg OP	✓ Baxter
YTARABINE			
Inj 20 mg per ml, 5 ml vial - PCT - Retail pharmacy-Speci	alist472.00	5	✓ Pfizer
Inj 100 mg per ml, 20 ml vial - PCT - Retail			
pharmacy-Specialist	48.80	1	<ul> <li>Cytarabine DBL</li> </ul>
			✓ Pfizer
			✓ Pfizer S29 S29
Inj 1 mg for ECP - PCT only - Specialist	0.29	10 mg	✓ Baxter
Inj 100 mg intrathecal syringe for ECP - PCT only - Speci		100 mg OP	✓ Baxter
LUDARABINE PHOSPHATE		· ·	
Tab 10 mg - PCT - Retail pharmacy-Specialist	412.00	20	✓ Fludara Oral
Inj 50 mg vial – PCT only – Specialist		1	✓ Fludarabine
.,		•	Sagent S29
	634.00	5	✓ Fludarabine Ebewe
Inj 50 mg for ECP - PCT only - Specialist		50 mg OP	✓ Baxter
, ,	120.00	oo mg or	Duxio
LUOROURACIL	10.51		/ Fluorenmanii Annord
Inj 50 mg per ml, 20 ml vial – PCT only – Specialist Inj 50 mg per ml, 50 ml vial – PCT only – Specialist		1 1	<ul> <li>✓ Fluorouracil Accord</li> <li>✓ Fluorouracil Accord</li> </ul>
Inj 50 mg per ml, 100 ml vial – PCT only – Specialist		1	✓ Fluorouracil Accord
Inj 1 mg for ECP – PCT only – Specialist	19.30 0.41	100 mg	✓ Baxter
	0.41	100 mg	Daxiei
EMCITABINE HYDROCHLORIDE – PCT only – Specialist			
Inj 43.3 mg per ml (equivalent to 38 mg per ml gemcitabine	e),		
26.3 ml vial		1	✓ DBL Gemcitabine
Inj 1 g		1	✓ Gemcitabine Ebewe
Inj 1 mg for ECP	0.02	1 mg	✓ Baxter
INOTECAN HYDROCHLORIDE - PCT only - Specialist			
Inj 20 mg per ml, 5 ml vial		1	✓ Accord
	71.44		✓ Irinotecan Actavis
			100
	100.00		✓ Irinotecan-Rex
Inj 1 mg for ECP	0.54	1 mg	✓ Baxter
ERCAPTOPURINE			
Tab 50 mg - PCT - Retail pharmacy-Specialist	25.90	25	✓ Puri-nethol
Oral suspension 20 mg per ml - Retail pharmacy-Speciali	st –		
Special Authority see SA1725 below		100 ml OP	✓ Allmercap
•			✓ Xaluprine S29

# **⇒SA1725** Special Authority for Subsidy

Initial application only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where the patient requires a total dose of less than one full 50 mg tablet per day.

**Renewal** only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where patient still requires a total dose of less than one full 50 mg tablet per day.

Subsidy (Magnifectured a	Orion) Colle	Fully Brand or
(Manufacturer's F \$	Per Subs	sidised Generic  ✓ Manufacturer
ETHOTREXATE		
Tab 2.5 mg - PCT - Retail pharmacy-Specialist7.80	90	✓ Trexate
Fab 10 mg - PCT - Retail pharmacy-Specialist	90	✓ Trexate
Inj 2.5 mg per ml, 2 ml – PCT – Retail pharmacy-Specialist56.05	5	✓ Methotrexate DBL
.,g p,	-	✓ Methotrexate DBL
		<b>S29</b> S29
k Inj 7.5 mg prefilled syringe29.17	1	✓ Methotrexate
Tily 7.0 mg promod dynngd	•	Sandoz
Methotrexate Sandoz to be Principal Supply on 1 February 2025		
F Inj 10 mg prefilled syringe	1	✓ Methotrexate
The first of the promote dynage	•	Sandoz
Methotrexate Sandoz to be Principal Supply on 1 February 2025		
Inj 15 mg prefilled syringe24.53	1	✓ Methotrexate
,, p	•	Sandoz
Methotrexate Sandoz to be Principal Supply on 1 February 2025		
Inj 20 mg prefilled syringe16.64	1	✓ Methotrexate
,g F 9)ge.	•	Sandoz
Methotrexate Sandoz to be Principal Supply on 1 February 2025		- · · · <del></del>
Inj 25 mg prefilled syringe	1	✓ Methotrexate
, - 3, , 3-		Sandoz
Methotrexate Sandoz to be Principal Supply on 1 February 2025		
k Inj 30 mg prefilled syringe	1	✓ Methotrexate
, 31 , 3		Sandoz
Methotrexate Sandoz to be Principal Supply on 1 February 2025		
Inj 25 mg per ml, 2 ml vial - PCT - Retail pharmacy-Specialist30.00	5	✓ Methotrexate DBL
		Onco-Vial
Inj 25 mg per ml, 20 ml vial - PCT - Retail pharmacy-Specialist45.00	1	✓ DBL Methotrexate
		Onco-Vial
Inj 100 mg per ml, 10 ml - PCT - Retail pharmacy-Specialist25.00	1	✓ Methotrexate Ebewe
Inj 100 mg per ml, 50 ml vial - PCT - Retail		
pharmacy-Specialist	1	✓ Methotrexate Ebewe
Finj 1 mg for ECP - PCT only - Specialist	1 mg	✓ Baxter
Inj 5 mg intrathecal syringe for ECP – PCT only – Specialist4.73	5 mg OP	✓ Baxter
EMETREXED - PCT only - Specialist		
Inj 100 mg vial	1	✓ Pemetrexed-AFT
60.89		✓ Juno Pemetrexed
Inj 500 mg vial29.99	1	✓ Pemetrexed-AFT
217.77		Juno Pemetrexed
Inj 1 mg for ECP	1 mg	✓ Baxter
HIOGUANINE - PCT - Retail pharmacy-Specialist	-	
Tab 40 mg	25	✓ Lanvis
I J		
Other Cytotoxic Agents		
MSACRINE - PCT only - Specialist		
Inj 50 mg per ml, 1.5 ml ampoule	6	✓ Amsidine \$29
4.736.00	-	✓ Amsidine S29
Inj 75 mg	5	✓ AmsaLyo S29
, ,	J	•
6,218.00		✓ AmsaLyo S29

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

	Subsidy	D: \	Fully Brand or
	(Manufacturer's I \$	Price) Subs Per	idised Generic  ✓ Manufacturer
NAGRELIDE HYDROCHLORIDE - PCT - Retail pharmacy	, Specialist	1 01	- Managara
Cap 0.5 mg		100	✓ Agrylin
· · · · · · · · · · · · · · · · · · ·		100	• Agryllii
RSENIC TRIOXIDE - PCT only - Specialist	4 917 00	10	✓ Phenasen
Inj 1 mg per ml, 10 ml vial Inj 10 mg for ECP		10 10 mg OP	✓ Baxter
	401.70	10 mg Oi	Daxiei
LEOMYCIN SULPHATE – PCT only – Specialist	105 16	1	✓ DBL Bleomycin
Inj 15,000 iu, vial	103.10	1	Sulfate
Inj 1,000 iu for ECP	1/1 32	1,000 iu	✓ Baxter
• •		1,000 10	Daxiei
ORTEZOMIB - PCT only - Specialist - Special Authority so			✓ DBL Bortezomib
Inj 3.5 mg vial Inj 1 mg for ECP		1 1 ma	✓ Baxter
	22.20	1 mg	Daxiei
SA2355 Special Authority for Subsidy		and the second of the second o	Street Coate and a second coate
tial application — (plasma cell dyscrasia) from any rele			
tified where the patient has plasma cell dyscrasia, not inclu	ung waidenstrom	macrogiobulinae	ernia, requiring treatment
ACARBAZINE – PCT only – Specialist Inj 200 mg vial	70 11	4	/ DDI Dasaubarina
Inj 200 mg for ECP		1 200 mg OP	✓ DBL Dacarbazine ✓ Baxter
, -		200 Hig OF	<b>▼</b> Daxlei
ACTINOMYCIN [ACTINOMYCIN D] - PCT only - Specialis			
Inj 0.5 mg vial		1	✓ Cosmegen
Inj 0.5 mg for ECP	255.00	0.5 mg OP	✓ Baxter
AUNORUBICIN – PCT only – Specialist			4
Inj 2 mg per ml, 10 ml		1	✓ Pfizer
Inj 20 mg for ECP	171.93	20 mg OP	✓ Baxter
OCETAXEL - PCT only - Specialist			_
Inj 20 mg		1	✓ Docetaxel Sando
Inj 10 mg per ml, 8 ml vial		1	✓ DBL Docetaxel
Inj 20 mg per ml, 4 ml vial	26.95	1	✓ Docetaxel
			Accord S29
Inj 80 mg		_ 1	✓ Docetaxel Sando
Inj 1 mg for ECP		1 mg	✓ Baxter
OXORUBICIN HYDROCHLORIDE - PCT only - Specialist			
Inj 2 mg per ml, 5 ml vial		1	Doxorubicin Ebev
Inj 2 mg per ml, 25 ml vial		1	✓ Doxorubicin Ebev
lai O ara a caract. FO artistict	17.00	_	✓ Arrow-Doxorubic
Inj 2 mg per ml, 50 ml vial		1 1	✓ Doxorubicin Ebev
Inj 2 mg per ml, 100 ml vial	69.99	ı	<ul> <li>✓ Arrow-Doxorubic</li> <li>✓ Doxorubicin Ebev</li> </ul>
Inj 1 mg for ECP		1 mg	✓ Baxter
		i ilig	Duxici
PIRUBICIN HYDROCHLORIDE – PCT only – Specialist	05.00		/ Fuluubiain Fhans
Inj 2 mg per ml, 5 ml vial	25.00	1 1	<ul><li>✓ Epirubicin Ebewe</li><li>✓ Epirubicin Ebewe</li></ul>
Inj 2 mg per ml, 25 ml vial Inj 2 mg per ml, 100 ml vial		1	✓ Epirubicin Ebewe
Inj 1 mg for ECP		1 mg	✓ Baxter
, -		ı mg	- Duxiei
TOPOSIDE  Can 50 mg PCT Potail pharmacy Specialist	240.72	20	✓ Vepesid
Cap 50 mg - PCT - Retail pharmacy-Specialist Cap 100 mg - PCT - Retail pharmacy-Specialist		20 10	✓ Vepesid ✓
		10	✓ Vepesid ✓ Rex Medical
Ini 20 ma nor mi 5 mi vial = DCT = Rotail nharmany Sha			
Inj 20 mg per ml, 5 ml vial – PCT – Retail pharmacy-Spe Inj 1 mg for ECP – PCT only – Specialist		1 mg	✓ Baxter

	Subsidy (Manufacturer's Price)	) Per	Fully Subsidised	
ETOPOSIDE PHOSPHATE – PCT only – Specialist Inj 100 mg (of etoposide base) Inj 1 mg (of etoposide base) for ECP		1 1 mg		Etopophos Baxter
HYDROXYUREA [HYDROXYCARBAMIDE] – PCT – Retail phart	, ,	100	/	<u>Devatis</u>
IBRUTINIB – Special Authority see SA2168 below – Retail pharm Tab 140 mg Tab 420 mg	3,217.00	30 30		Imbruvica Imbruvica

### ⇒SA2168 Special Authority for Subsidy

Initial application — (chronic lymphocytic leukaemia (CLL)) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

#### All of the following:

- 1 Patient has chronic lymphocytic leukaemia (CLL) requiring therapy; and
- 2 Patient has not previously received funded ibrutinib; and
- 3 Ibrutinib is to be used as monotherapy; and
- 4 Any of the following:
  - 4.1 Both:
    - 4.1.1 There is documentation confirming that patient has 17p deletion or TP53 mutation; and
    - 4.1.2 Patient has experienced intolerable side effects with venetoclax monotherapy; or
  - 4.2 All of the following:
    - 4.2.1 Patient has received at least one prior immunochemotherapy for CLL; and
    - 4.2.2 Patient's CLL has relapsed within 36 months of previous treatment; and
    - 4.2.3 Patient has experienced intolerable side effects with venetoclax in combination with rituximab regimen; or
  - 4.3 Patient's CLL is refractory to or has relapsed within 36 months of a venetoclax regimen.

**Renewal — (chronic lymphocytic leukaemia (CLL))** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

## Both:

- 1 No evidence of clinical disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL) and B-cell prolymphocytic leukaemia (B-PLL)\*. Indications marked with \* are Unapproved indications.

#### IDARUBICIN HYDROCHLORIDE

Inj 5 mg vial - PCT only - Specialist	109.74	1	Zavedos
Inj 10 mg vial - PCT only - Specialist	233.64	1	✓ Zavedos
Inj 1 mg for ECP - PCT only - Specialist	25.77	1 mg	✓ Baxter
ENALIDOMIDE (DEVILIMID) - Datail pharmagy Chanielist	Charial Authority and	CA0047 on	the next need

# LENALIDOMIDE (REVLIMID) - Retail pharmacy-Specialist - Special Authority see SA2047 on the next page Wastage claimable

Cap 5 mg	5,122.76	28	✓ Revlimid
Cap 10 mg		28	✓ Revlimid
Cap 15 mg	7,239.18	28	✓ Revlimid
Can 25 mg	7 627 00	21	✓ Revlimid

(Revlimid Cap 5 mg to be delisted 1 February 2025)

(Revlimid Cap 10 mg to be delisted 1 February 2025)

(Revlimid Cap 15 mg to be delisted 1 February 2025)

(Revlimid Cap 25 mg to be delisted 1 February 2025)

		Fully	Brand or
(Manufacturer's Price) \$	Subside Per	aisea 🗸	Generic Manufacturer

# ⇒SA2047 Special Authority for Subsidy

Initial application — (Relapsed/refractory disease) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has relapsed or refractory multiple myeloma with progressive disease; and
- 2 Patient has not previously been treated with lenalidomide; and
- 3 Either
  - 3.1 Lenalidomide to be used as third line\* treatment for multiple myeloma; or
  - 3.2 Both:
    - 3.2.1 Lenalidomide to be used as second line treatment for multiple myeloma; and
    - 3.2.2 The patient has experienced severe (grade 3 or higher), dose limiting, peripheral neuropathy with either bortezomib or thalidomide that precludes further treatment with either of these treatments; and
- 4 Lenalidomide to be administered at a maximum dose of 25 mg/day in combination with dexamethasone.

**Initial application** — (Maintenance following first-line autologous stem cell transplant (SCT)) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has newly diagnosed symptomatic multiple myeloma and has undergone first-line treatment that included an autologous stem cell transplantation; and
- 2 Patient has at least a stable disease response in the first 100 days after transplantation; and
- 3 Lenalidomide maintenance is to be commenced within 6 months of transplantation; and
- 4 Lenalidomide to be administered at a maximum dose of 15 mg/day.

Renewal — (Relapsed/refractory disease) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

Renewal — (Maintenance following first line autologous SCT) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

Note: Indication marked with \* is an unapproved indication. A line of treatment is considered to comprise either: a) a known therapeutic chemotherapy regimen and supportive treatments or b) a transplant induction chemotherapy regimen, stem cell transplantation and supportive treatments. Prescriptions must be written by a registered prescriber in the lenalidomide risk management programme operated by the supplier.

LENALIDOMIDE (VIATRIS) – Special Authority see SA2353 on the next page – Re Cap 5 mg76.92	etail pharma 21	cy ✓ Lenalidomide
		Viatris
Lenalidomide Viatris to be Principal Supply on 1 February 2025		
Cap 10 mg50.30	21	✓ Lenalidomide Viatris
Lenalidomide Viatris to be Principal Supply on 1 February 2025		
Cap 15 mg	21	✓ Lenalidomide Viatris
Lenalidomide Viatris to be Principal Supply on 1 February 2025		
Cap 25 mg65.09	21	<ul><li>Lenalidomide Viatris</li></ul>
Lenalidomide Viatris to be Principal Supply on 1 February 2025		

Subsidy		ully	Brand or
(Manufacturer's Price)	Subsid	ised	Generic
\$	Per	✓	Manufacturer

# ⇒SA2353 Special Authority for Subsidy

Initial application — (Plasma cell dyscrasia) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Both:

- 1 Patient has plasma cell dyscrasia, not including Waldenström macroglobulinaemia, requiring treatment; and
- 2 Patient is not refractory to prior lenalidomide use.

Initial application — (Myelodysplastic syndrome) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

#### Both:

- 1 Patient has low or intermediate-1 risk myelodysplastic syndrome (based on IPSS or an IPSS-R score of less than 3.5) associated with a deletion 5g cytogenetic abnormality; and
- 2 Patient has transfusion-dependent anaemia.

Renewal — (Myelodysplastic syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

### Both:

- 1 Patient has not needed a transfusion in the last 4 months; and
- 2 No evidence of disease progression.

#### **MESNA**

MESINA			
Tab 400 mg - PCT - Retail pharmacy-Specialist	314.00	50	✓ Uromitexan
Tab 600 mg - PCT - Retail pharmacy-Specialist	448.50	50	✓ Uromitexan
Inj 100 mg per ml, 4 ml ampoule - PCT only - Specialis		15	✓ Uromitexan
Inj 100 mg per ml, 10 ml ampoule – PCT only – Speciali		15	✓ Uromitexan
Inj 1 mg for ECP – PCT only – Specialist		100 mg	✓ Baxter
MITOMYCIN C - PCT only - Specialist			
Inj 5 mg vial	526.00	1	✓ Mitomycin
			(Sagent) S29
	577.50		✓ Mitomycin
			(Fresenius
			Kabi) S29
	641.70		✓ Accord S29
Inj 20 mg vial	1.250.00	1	✓ Omegapharm S29
,	,		✓ Teva
Inj 1 mg for ECP	269.85	1 mg	✓ Baxter
MITOZANTRONE - PCT only - Specialist			
Inj 2 mg per ml, 10 ml vial	97.50	1	✓ Mitozantrone Ebewe
Inj 1 mg for ECP		1 mg	✓ Baxter
NIRAPARIB - Special Authority see SA2325 below - Retail	oharmacv		
Wastage claimable	,		
Tab 100 mg	13.393.50	84	✓ Zejula
Cap 100 mg		56	✓ Zejula
	13,393.50	84	✓ Zejula
	-,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,		

# ⇒SA2325 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has advanced high-grade serous\* epithelial ovarian, fallopian tube, or primary peritoneal cancer; and
- 2 Patient has received at least one line\*\* of treatment with platinum-based chemotherapy; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsi	dised	Generic	
\$	Per	1	Manufacturer	

continued...

- 3 Patient has experienced a partial or complete response to the preceding treatment with platinum-based chemotherapy; and
- 4 Patient has not previously received funded treatment with a PARP inhibitor; and
- 5 Either
  - 5.1 Treatment will be commenced within 12 weeks of the patient's last dose of the preceding platinum-based regimen;
  - 5.2 Patient commenced treatment with niraparib prior to 1 May 2024; and
- 6 Treatment to be administered as maintenance treatment; and
- 7 Treatment not to be administered in combination with other chemotherapy.

**Renewal** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 No evidence of progressive disease: and
- 2 Treatment to be administered as maintenance treatment; and
- 3 Treatment not to be administered in combination with other chemotherapy; and
- 4 Either
  - 4.1 Treatment with niraparib to cease after a total duration of 36 months from commencement; or
  - 4.2 Treatment with niraparib is being used in the second-line or later maintenance setting.

Notes: \* "high-grade serous" includes tumours with high-grade serous features or a high-grade serous component.

\*\*A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments

OLAPARIB – Retail pharmacy-Specialist – Special Authority	see SA2163 below		
Tab 100 mg	3,701.00	56	<ul><li>Lynparza</li></ul>
Tab 150 mg	3,701.00	56	<ul><li>Lynparza</li></ul>

#### ⇒SA2163 Special Authority for Subsidy

Initial application — (Ovarian cancer) only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a high-grade serous\* epithelial ovarian, fallopian tube, or primary peritoneal cancer; and
- 2 There is documentation confirming pathogenic germline BRCA1 or BRCA2 gene mutation; and
- 3 Fither:
  - 3.1 All of the following:
    - 3.1.1 Patient has newly diagnosed, advanced disease; and
    - 3.1.2 Patient has received one line\*\* of previous treatment with platinum-based chemotherapy; and
    - 3.1.3 Patient's disease must have experienced a partial or complete response to the first-line platinum-based regimen; or
  - 3.2 All of the following:
    - 3.2.1 Patient has received at least two lines\*\* of previous treatment with platinum-based chemotherapy; and
    - 3.2.2 Patient has platinum sensitive disease defined as disease progression occurring at least 6 months after the last dose of the penultimate line\*\* of platinum-based chemotherapy; and
    - 3.2.3 Patient's disease must have experienced a partial or complete response to treatment with the immediately preceding platinum-based regimen; and
    - 3.2.4 Patient has not previously received funded olaparib treatment; and
- 4 Treatment will be commenced within 12 weeks of the patient's last dose of the immediately preceding platinum-based regimen; and
- 5 Treatment to be administered as maintenance treatment; and
- 6 Treatment not to be administered in combination with other chemotherapy.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Su	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Renewal — (Ovarian cancer) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from treatment; and
- 2 Either:
  - 2.1 No evidence of progressive disease; or
  - 2.2 Evidence of residual (not progressive) disease and the patient would continue to benefit from treatment in the clinician's opinion; and
- 3 Treatment to be administered as maintenance treatment; and
- 4 Treatment not to be administered in combination with other chemotherapy; and
- 5 Either:
  - 5.1 Both:
    - 5.1.1 Patient has received one line\*\* of previous treatment with platinum-based chemotherapy; and
    - 5.1.2 Documentation confirming that the patient has been informed and acknowledges that the funded treatment period of olaparib will not be continued beyond 2 years if the patient experiences a complete response to treatment and there is no radiological evidence of disease at 2 years; or
  - 5.2 Patient has received at least two lines\*\* of previous treatment with platinum-based chemotherapy.

Notes: \*Note "high-grade serous" includes tumours with high-grade serous features or a high-grade serous component.
\*\*A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

PACLITAXEL – PCT only – Specialist			
Inj 30 mg	47.30	5	✓ Paclitaxel Ebewe
Inj 6 mg per ml, 16.7 ml vial	19.59	1	✓ Anzatax
	24.00		✓ Paclitaxel Ebewe
	91.67		✓ Paclitaxel Actavis
Inj 150 mg	26.69	1	✓ Paclitaxel Ebewe
	137.50		✓ Anzatax
			✓ Paclitaxel Actavis
Inj 6 mg per ml, 50 ml vial	37.89	1	✓ Anzatax
	44.00		✓ Paclitaxel Ebewe
	275.00		✓ Paclitaxel Actavis
Inj 1 mg for ECP	0.17	1 mg	✓ Baxter
PEGASPARGASE - PCT only - Special Authority se	e SA1979 below		
Inj 750 iu per ml, 5 ml vial	3,973.25	1	<ul><li>Oncaspar LYO</li></ul>

#### ⇒SA1979 Special Authority for Subsidy

Initial application — (Acute lymphoblastic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has newly diagnosed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

Initial application — (Lymphoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months where the patient has lymphoma requiring L-asparaginase containing protocols (e.g. SMILE).

Renewal — (Acute lymphoblastic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has relapsed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

	Subsidy (Manufacturer's Price)		Fully Subsidised	
	(Manufacturer's Price)	Per		
PENTOSTATIN [DEOXYCOFORMYCIN] - PCT only - Special	list			
Inj 10 mg	CBS	1	1	Nipent \$29
POMALIDOMIDE - Special Authority see SA2354 below - Ret	ail pharmacy			
Cap 1 mg	47.45	14	✓	Pomolide
	71.18	21	✓	<u>Pomolide</u>
Cap 2 mg	94.90	14	1	<u>Pomolide</u>
	142.35	21	✓	<u>Pomolide</u>
Cap 3 mg	142.35	14	✓	<u>Pomolide</u>
	213.53	21	✓	Pomolide
Cap 4 mg	189.81	14	✓	Pomolide
	284.71	21	✓	Pomolide

# ⇒SA2354 Special Authority for Subsidy

Initial application — (Relapsed/refractory plasma cell dyscrasia) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has relapsed or refractory plasma cell dyscrasia, not including Waldenström macroglobulinaemia, requiring treatment; and
- 2 Patient has not received prior funded pomalidomide.

Renewal — (Relapsed/refractory plasma cell dyscrasia) from any relevant practitioner. Approvals valid for 12 months where there is no evidence of disease progression.

PROCARBAZINE HYDROCHLORIDE – PCT – Retail pharmar Cap 50 mg		50	✓ Natulan S29
TEMOZOLOMIDE - Special Authority see SA2275 below - Re	tail pharmacy		
Cap 5 mg		5	✓ Temaccord
•			✓ Temozolomide-
			Taro \$29
Cap 20 mg	16.38	5	✓ Temaccord
, ,	18.30		✓ Apo-Temozolomide
Cap 100 mg	35.98	5	✓ Temaccord
,	40.20		✓ Apo-Temozolomide
Cap 140 mg	50.12	5	✓ Temaccord
Cap 250 mg	86.34	5	✓ Temaccord

# ⇒SA2275 Special Authority for Subsidy

Initial application — (gliomas) only from a relevant specialist. Approvals valid for 12 months where the patient has a glioma. Renewal — (gliomas) only from a relevant specialist. Approvals valid for 12 months where treatment remains appropriate and patient is benefitting from treatment.

**Initial application**—(neuroendocrine tumours) only from a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with metastatic or unresectable well-differentiated neuroendocrine tumour\*; and
- 2 Temozolomide is to be given in combination with capecitabine; and
- 3 Temozolomide is to be used in 28 day treatment cycles for a maximum of 5 days treatment per cycle at a maximum dose of 200 mg/m² per day; and
- 4 Temozolomide to be discontinued at disease progression.

Renewal — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

(Manu	Subsidy	Fully	Brand or
	facturer's Price)	Subsidised	Generic
<u> </u>	\$ Pe	er 🗸	Manufacturer

continued...

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

**Initial application** — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 9 months where the patient has relapsed/refractory Ewing's sarcoma.

Renewal — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indication marked with a \* is an unapproved indication. Temozolomide is not subsidised for the treatment of relapsed high grade glioma.

THALIDOMIDE - Retail pharmacy-Specialist - Special Au	uthority see SA2356 below		
Cap 50 mg	378.00	28	Thalomid
Cap 100 mg	756.00	28	✓ Thalomid

## ⇒SA2356 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 12 months where the patient has plasma cell dyscrasia, not including Waldenström macroglobulinaemia, requiring treatment.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has obtained a response from treatment during the initial approval period.

Notes: Prescription must be written by a registered prescriber in the thalidomide risk management programme operated by the supplier.

Maximum dose of 400 mg daily as monotherapy or in a combination therapy regimen.

#### TRETINOIN

Cap 10 mg - PCT - Retail pharmacy-Specialist479.50	100	<ul><li>Vesanoid</li></ul>
VENETOCLAX - Retail pharmacy-Specialist - Special Authority see SA1868	B below	
Tab 14 × 10 mg, 7 × 50 mg, 21 × 100 mg	42 OP	✓ Venclexta
Tab 10 mg13.68		✓ Venclexta
Tab 50 mg239.44	7 OP	✓ Venclexta
Tab 100 mg - Wastage claimable8,209.41	120	✓ Venclexta

# ⇒SA1868 Special Authority for Subsidy

Initial application — (relapsed/refractory chronic lymphocytic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic lymphocytic leukaemia requiring treatment; and
- 2 Patient has received at least one prior therapy for chronic lymphocytic leukaemia; and
- 3 Patient has not previously received funded venetoclax; and
- 4 The patient's disease has relapsed within 36 months of previous treatment; and
- 5 Venetoclax to be used in combination with six 28-day cycles of rituximab commencing after the 5-week dose titration schedule with venetoclax; and
- 6 Patient has an ECOG performance status of 0-2.

Renewal — (relapsed/refractory chronic lymphocytic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: Both:

1 Treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment; and

Subsidy		Fully	Brand or
(Manufacturer's Price)	Su	bsidised	Generic
\$	Per	•	Manufacturer

continued...

2 Venetoclax is to be discontinued after a maximum of 24 months of treatment following the titration schedule unless earlier discontinuation is required due to disease progression or unacceptable toxicity.

Initial application — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation\*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has previously untreated chronic lymphocytic leukaemia; and
- 2 There is documentation confirming that patient has 17p deletion by FISH testing or TP53 mutation by sequencing; and
- 3 Patient has an ECOG performance status of 0-2.

Renewal — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation\*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where the treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL)\* and B-cell prolymphocytic leukaemia (B-PLL)\*. Indications marked with \* are Unapproved indications.

#### VINBLASTINE SULPHATE

Inj 1 mg per ml, 10 ml vial – PCT – Retail pharmacy-Specialist270.37	5	✓ Hospira
Inj 1 mg for ECP - PCT only - Specialist	1 mg	✓ Baxter
VINCRISTINE SULPHATE		
Inj 1 mg per ml, 1 ml vial - PCT - Retail pharmacy-Specialist74.52	5	✓ DBL Vincristine Sulfate
Inj 1 mg per ml, 2 ml vial - PCT - Retail pharmacy-Specialist102.73	5	<ul><li>DBL Vincristine Sulfate</li></ul>
Inj 1 mg for ECP - PCT only - Specialist12.60	1 mg	✓ Baxter
VINORELBINE		
Cap 20 mg30.00	1	✓ Vinorelbine Te Arai
Cap 30 mg40.00	1	✓ Vinorelbine Te Arai
Cap 80 mg60.00	1	✓ Vinorelbine Te Arai
Inj 10 mg per ml, 1 ml vial - PCT only - Specialist42.00	1	✓ Vinorelbine Ebewe
Inj 10 mg per ml, 5 ml vial - PCT only - Specialist168.00	1	✓ Navelbine S29 S29
210.00		✓ Vinorelbine Ebewe
Inj 1 mg for ECP - PCT only - Specialist	1 mg	✓ Baxter

# **Protein-tyrosine Kinase Inhibitors**

/ LEEO 1 11 11 10	riotan priarriacy	Opoolanot	opoolai riailionity ooo or	i i o i o boioii		
Wastage	claimable					
Cap 150	mg			7,935.00	224	Alecensa

#### ⇒SA1870 Special Authority for Subsidy

**Initial application** only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

1 Patient has locally advanced, or metastatic, unresectable, non-small cell lung cancer; and

ALECTINIB - Retail pharmacy-Specialist - Special Authority see SA1870 below

- 2 There is documentation confirming that the patient has an ALK tyrosine kinase gene rearrangement using an appropriate ALK test; and
- 3 Patient has an ECOG performance score of 0-2.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid

(Manufacturer's Price) Subsidised Generic \$ Per ✓ Manufacturer	Sul	ıbsidy Fu	lly Brand or
\$ Per ✓ Manufacturer	(Manufact		
		\$ Per	Manufacturer

continued...

for 6 months for applications meeting the following criteria:

#### Roth:

- 1 No evidence of progressive disease according to RECIST criteria; and
- 2 The patient is benefitting from and tolerating treatment.

# DASATINIB - Special Authority see SA2385 below - Retail pharmacy

Wastage claimable

Tab 20 mg	132.88	60	✓ Dasatinib-Teva
	3,774.06		✓ Sprycel
Dasatinib-Teva to be Principal Supply on 1 March 202	25		
Tab 50 mg	304.13	60	Dasatinib-Teva
·	6,214.20		✓ Sprycel
Dasatinib-Teva to be Principal Supply on 1 March 202	25		
Tab 70 mg	415.75	60	Dasatinib-Teva
·	7,692.58		✓ Sprycel

Dasatinib-Teva to be Principal Supply on 1 March 2025

(Sprycel Tab 20 mg to be delisted 1 March 2025) (Sprycel Tab 50 mg to be delisted 1 March 2025)

(Sprycel Tab 70 mg to be delisted 1 March 2025)

# ⇒SA2385 Special Authority for Subsidy

Initial application only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis or accelerated phase; or
- 2 The patient has a diagnosis of Philadelphia chromosome-positive acute lymphoid leukaemia (Ph+ ALL); or
- 3 Both:
  - 3.1 The patient has a diagnosis of CML in chronic phase; and
  - 3.2 Any of the following:
    - 3.2.1 Patient has documented treatment failure\* with imatinib; or
    - 3.2.2 Patient has experienced treatment-limiting toxicity with imatinib precluding further treatment with imatinib; or
    - 3.2.3 Patient has high-risk chronic-phase CML defined by the Sokal or EURO scoring system.

**Renewal** only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

#### Both:

- 1 Lack of treatment failure while on dasatinib\*; and
- 2 Dasatinib treatment remains appropriate and the patient is benefiting from treatment.

Note: \*treatment failure for CML as defined by Leukaemia Net Guidelines.

ERLOTINIB - Retail pharmacy-Specialist - Special Authority see SA2115 below

Tab 100 mg	280.84	30	Alchemy
Tab 150 mg	181 21	30	✓ Alchemy

#### ⇒SA2115 Special Authority for Subsidy

**Initial application** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 There is documentation confirming that the disease expresses activating mutations of EGFR tyrosine kinase; and
- 3 Fither:

Subsidy		Fully	Brand or	
(Manufacturer's Pric	ce)	Subsidised	Generic	
\$	Per	1	Manufacturer	

continued...

- 3.1 Patient is treatment naive; or
- 3.2 Both:
  - 3.2.1 The patient has discontinued defitinib due to intolerance; and
  - 3.2.2 The cancer did not progress while on gefitinib; and
- 4 Erlotinib is to be given for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

Renewal — (pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 2 Erlotinib to be discontinued at progression; and
- 3 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

GEFITINIB – Retail pharmacy-Specialist – Special Authority see SA2116 below Tab 250 mg .......918.00

30 ✓ Iressa

# ⇒SA2116 Special Authority for Subsidy

**Initial application** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 Fither:
  - 2.1 Patient is treatment naive; or
  - 2.2 Both:
    - 2.2.1 The patient has discontinued erlotinib due to intolerance; and
    - 2.2.2 The cancer did not progress whilst on erlotinib; and
- 3 There is documentation confirming that disease expresses activating mutations of EGFR tyrosine kinase; and
- 4 Gefitinib is to be given for a maximum of 3 months.

**Renewal** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

Renewal — (pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 2 Gefitinib to be discontinued at progression; and
- 3 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

#### **IMATINIB MESILATE**

* Cap 400 mg		60 30	✓ <u>Imatinib-Rex</u> ✓ <u>Imatinib-Rex</u>
LENVATINIB – Special Authority see SA2407 below Wastage claimable	ow - Retail pharmacy		
Cap 4 mg	3,407.40	30	✓ Lenvima
Cap 10 mg	3,407.40	30	✓ Lenvima

### ⇒SA2407 Special Authority for Subsidy

**Initial application — (thyroid cancer)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

Fither:

- 1 Patient is currently on treatment with lenvatinib and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
  - 2.1 The patient has locally advanced or metastatic differentiated thyroid cancer; and
  - 2.2 Fither
    - 2.2.1 Patient must have symptomatic progressive disease prior to treatment; or
    - 2.2.2 Patient must progressive disease at critical anatomical sites with a high risk of morbidity or mortality where local control cannot be achieved by other measures; and
  - 2.3 Any of the following:
    - 2.3.1 A lesion without iodine uptake in a RAI scan: or
    - 2.3.2 Receiving cumulative RAI greater than or equal to 600 mCi; or
    - 2.3.3 Experiencing disease progression after a RAI treatment within 12 months; or
    - 2.3.4 Experiencing disease progression after two RAI treatments administered within 12 months of each other; and
  - 2.4 Patient has thyroid stimulating hormone (TSH) adequately supressed; and
  - 2.5 Patient is not a candidate for radiotherapy with curative intent; and
  - 2.6 Surgery is clinically inappropriate; and
  - 2.7 Patient has an ECOG performance status of 0-2.

Renewal — (thyroid cancer) from any relevant practitioner. Approvals valid for 6 months where there is no evidence of disease progression.

Initial application — (unresectable hepatocellular carcinoma) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has unresectable hepatocellular carcinoma; and
- 2 Patient has preserved liver function (Childs-Pugh A); and
- 3 Transarterial chemoembolisation (TACE) is unsuitable; and
- 4 Patient has an ECOG performance status of 0-2; and
- 5 Patient has not received prior systemic therapy for their disease in the palliative setting.

Renewal — (unresectable hepatocellular carcinoma) from any relevant practitioner. Approvals valid for 6 months where there is no evidence of disease progression.

**Initial application** — **(renal cell carcinoma)** from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 All of the following:
  - 1.1 The patient has metastatic renal cell carcinoma; and
  - 1.2 The disease is of predominant clear-cell histology; and
  - 1.3 The patient has documented disease progression following one previous line of treatment; and
  - 1.4 The patient has an ECOG performance status of 0-2; and
  - 1.5 Lenvatinib is to be used in combination with everolimus; or
- 2 All of the following:
  - 2.1 Patient has received funded treatment with nivolumab for the second line treatment of metastatic renal cell carcinoma; and
  - 2.2 Patient has experienced treatment limiting toxicity from treatment with nivolumab; and
  - 2.3 Lenvatinib is to be used in combination with everolimus; and
  - 2.4 There is no evidence of disease progression.

Renewal — (renal cell carcinoma) from any relevant practitioner. Approvals valid for 4 months where there is no evidence of disease progression.

MIDOSTAURIN - PCT only - Special Authority see SA2342 on the next page

Cap 25 mg......10,981.00

Rydapt

56

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Por 🗸	Manufacturer

# ⇒SA2342 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria: All of the following:

- 1 Patient has a diagnosis of acute myeloid leukaemia; and
- 2 Condition must be FMS tyrosine kinase 3 (FLT3) mutation positive; and
- 3 Patient must not have received a prior line of intensive chemotherapy for acute myeloid leukaemia; and
- 4 Patient is to receive standard intensive chemotherapy in combination with midostaurin only; and
- 5 Midostaurin to be funded for a maximum of 4 cycles.

## NILOTINIB - Special Authority see SA2301 below - Retail pharmacy

Wastage claimable

Cap 150 mg	4,680.00	120	Tasigna
Cap 200 mg	6,532.00	120	Tasigna

#### ⇒SA2301 Special Authority for Subsidy

**Initial application** only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis, high risk chronic phase, or in chronic phase; and
- 2 Fither:
  - 2.1 Patient has documented CML treatment failure\* with a tyrosine kinase inhibitor (TKI); or
  - 2.2 Patient has experienced treatment limiting toxicity with a tyrosine kinase inhibitor (TKI) precluding further treatment; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

Note: \*treatment failure as defined by Leukaemia Net Guidelines.

Renewal only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

#### All of the following:

- 1 Lack of treatment failure while on nilotinib as defined by Leukaemia Net Guidelines; and
- 2 Nilotinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

#### PALBOCICLIB - Special Authority see SA2345 below - Retail pharmacy

## Wastage claimable

***************************************			
Tab 75 mg	4,000.00	21	Ibrance
Tab 100 mg	4,000.00	21	✓ Ibrance
Tah 125 mg	4 000 00	21	✓ Ibrance

#### ⇒SA2345 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Either:

- 1 All of the following:
  - 1.1 Patient has unresectable locally advanced or metastatic breast cancer; and
  - 1.2 There is documentation confirming disease is hormone-receptor positive and HER2-negative; and
  - 1.3 Patient has an ECOG performance score of 0-2; and
  - 1.4 Fither:
    - 1.4.1 Disease has relapsed or progressed during prior endocrine therapy; or
    - 1.4.2 Both:
      - 1.4.2.1 Patient is amenorrhoeic, either naturally or induced, with endocrine levels consistent with a postmenopausal or without menstrual-potential state; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	bsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1.4.2.2 Patient has not received prior systemic treatment for metastatic disease; and
- 1.5 Treatment must be used in combination with an endocrine partner; and
- 1.6 Patient has not received prior funded treatment with a CDK4/6 inhibitor; or
- 2 All of the following:
  - 2.1 Patient has an active Special Authority approval for ribociclib; and
  - 2.2 Patient has experienced a grade 3 or 4 adverse reaction to ribociclib that cannot be managed by dose reductions and requires treatment discontinuation; and
  - 2.3 Treatment must be used in combination with an endocrine partner; and
  - 2.4 There is no evidence of progressive disease since initiation of ribociclib.

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 Treatment must be used in combination with an endocrine partner; and
- 2 There is no evidence of progressive disease since initiation of palbociclib.

PAZOPANIB - Special Authority see SA1190 below - Retail pharmacy

Tab 200 mg	172.88	30	Pazopanib Teva
<b>3</b>	1,334.70		✓ Votrient
Tab 400 mg	464.00	30	Pazopanib Teva
· ·	2,669.40		✓ Votrient

(Votrient Tab 200 mg to be delisted 1 May 2025)

(Votrient Tab 400 mg to be delisted 1 May 2025)

# ⇒SA1190 Special Authority for Subsidy

**Initial application** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
  - 2.1 The patient is treatment naive; or
  - 2.2 The patient has only received prior cytokine treatment; or
  - 2.3 Both:
    - 2.3.1 The patient has discontinued sunitinib within 3 months of starting treatment due to intolerance; and
    - 2.3.2 The cancer did not progress whilst on sunitinib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and

The patient has intermediate or poor prognosis defined as:

- 5 Any of the following:
  - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
  - 5.2 Haemoglobin level < lower limit of normal; or
  - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
  - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
  - 5.5 Karnofsky performance score of less than or equal to 70; or
  - 5.6 2 or more sites of organ metastasis; and
- 6 Pazopanib to be used for a maximum of 3 months.

**Renewal** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

1 No evidence of disease progression; and

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Pazopanib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6.

RIBOCICLIB - Special Authority see SA2343 below - Retail pharmacy

Wastage claimable

✓ Kisqali	21	Tab 200 mg1,883.00
✓ Kisqali	42	3,767.00
✓ Kisqali	63	5.650.00

# ⇒SA2343 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Fither:

- 1 All of the following:
  - 1.1 Patient has unresectable locally advanced or metastatic breast cancer; and
  - 1.2 There is documentation confirming disease is hormone-receptor positive and HER2-negative; and
  - 1.3 Patient has an ECOG performance score of 0-2; and
  - 1.4 Any of the following:
    - 1.4.1 Disease has relapsed or progressed during prior endocrine therapy; or
    - 1.4.2 Both:
      - 1.4.2.1 Patient is amenorrhoeic, either naturally or induced, with endocrine levels consistent with a postmenopausal or without menstrual-potential state; and
      - 1.4.2.2 Patient has not received prior systemic endocrine treatment for metastatic disease; or
    - 1.4.3 Both:
      - 1.4.3.1 Patient commenced treatment with ribociclib in combination with an endocrine partner prior to 1 July 2024; and
      - 1.4.3.2 There is no evidence of progressive disease; and
  - 1.5 Treatment to be used in combination with an endocrine partner; and
  - 1.6 Patient has not received prior funded treatment with a CDK4/6 inhibitor; or
- 2 All of the following:
  - 2.1 Patient has an active Special Authority approval for palbociclib; and
  - 2.2 Patient has experienced a grade 3 or 4 adverse reaction to palbociclib that cannot be managed by dose reductions and requires treatment discontinuation; and
  - 2.3 Treatment must be used in combination with an endocrine partner; and
  - 2.4 There is no evidence of progressive disease since initiation of palbociclib.

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 Treatment must be used in combination with an endocrine partner; and
- 2 There is no evidence of progressive disease since initiation of ribociclib.

RUXOLITINIB – Special Authority see SA1890 on the next page – Retail pharmacy Wastage claimable

Tab 5 mg Tab 10mg		56 56	✓ Jakavi ✓ Jakavi
Tab 15 mg	5,000.00	56	✓ Jakavi
Tab 20 mg	5,000.00	56	Jakavi

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subs	idised	Generic
\$	Per	1	Manufacturer

# **⇒SA1890** Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has primary myelofibrosis or post-polycythemia vera myelofibrosis or post-essential thrombocythemia myelofibrosis; and
- 2 Fither:
  - 2.1 A classification of risk of intermediate-2 or high-risk myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS; or
  - 2.2 Both
    - 2.2.1 A classification of risk of intermediate-1 myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS; and
    - 2.2.2 Patient has severe disease-related symptoms that are resistant, refractory or intolerant to available therapy; and
- 3 A maximum dose of 20 mg twice daily is to be given.

**Renewal** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 A maximum dose of 20 mg twice daily is to be given.

SUNITINIB - Special Authority see SA2117 below - Reta	il pharmacy		
Cap 12.5 mg	208.38	28	<ul> <li>Sunitinib Pfizer</li> </ul>
Cap 25 mg	416.77	28	<ul> <li>Sunitinib Pfizer</li> </ul>
Cap 50 mg	694.62	28	<ul> <li>Sunitinib Pfizer</li> </ul>

# ⇒SA2117 Special Authority for Subsidy

Initial application — (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
  - 2.1 The patient is treatment naive; or
  - 2.2 The patient has only received prior cytokine treatment; or
  - 2.3 The patient has only received prior treatment with an investigational agent within the confines of a bona fide clinical trial which has Ethics Committee approval; or
  - 2.4 Both:
    - 2.4.1 The patient has discontinued pazopanib within 3 months of starting treatment due to intolerance; and 2.4.2 The cancer did not progress whilst on pazopanib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and

The patient has intermediate or poor prognosis defined as:

- 5 Any of the following:
  - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
  - 5.2 Haemoglobin level < lower limit of normal; or
  - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
  - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
  - 5.5 Karnofsky performance score of less than or equal to 70; or

(Manufacturer <sup>'</sup> s Price) Subsidised Generic \$ Per ✔ Manufacturer
---

continued...

- 5.6 2 or more sites of organ metastasis; and
- 6 Sunitinib to be used for a maximum of 2 cycles.

Initial application — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal tumour (GIST); and
- 2 Fither:
  - 2.1 The patient's disease has progressed following treatment with imatinib: or
  - 2.2 The patient has documented treatment-limiting intolerance, or toxicity to, imatinib.

**Renewal — (RCC)** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Sunitinib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6

Renewal — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

The patient has responded to treatment or has stable disease as determined by Choi's modified CT response evaluation criteria as follows:

- 1 Any of the following:
  - 1.1 The patient has had a complete response (disappearance of all lesions and no new lesions); or
  - 1.2 The patient has had a partial response (a decrease in size of 10% or more or decrease in tumour density in Hounsfield Units (HU) of 15% or more on CT and no new lesions and no obvious progression of non measurable disease); or
  - 1.3 The patient has stable disease (does not meet criteria the two above) and does not have progressive disease and no symptomatic deterioration attributed to tumour progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: It is recommended that response to treatment be assessed using Choi's modified CT response evaluation criteria (J Clin Oncol, 2007, 25:1753-1759). Progressive disease is defined as either: an increase in tumour size of 10% or more and not meeting criteria of partial response (PR) by tumour density (HU) on CT; or: new lesions, or new intratumoral nodules, or increase in the size of the existing intratumoral nodules.

Renewal — (GIST pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal (GIST); and
- 2 The patient is clinically benifiting from treatment and continued treatment remains appropriate; and
- 3 Sunitinib is to be discontinued at progression; and
- 4 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

# **Endocrine Therapy**

For GnRH ANALOGUES - refer to HORMONE PREPARATIONS, Trophic Hormones, page 92

ABIRATERONE ACETATE – Retail pharmacy-Specialist – Special Authority see SA2118 on the next page Wastage claimable

Tab 250 mg .......4,276.19 120 **✓ Zytiga** 

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

# ⇒SA2118 Special Authority for Subsidy

Initial application only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has prostate cancer; and
- 2 Patient has metastases: and
- 3 Patient's disease is castration resistant; and
- 4 Fither:
  - 4.1 All of the following:
    - 4.1.1 Patient is symptomatic; and
    - 4.1.2 Patient has disease progression (rising serum PSA) after second line anti-androgen therapy; and
    - 4.1.3 Patient has ECOG performance score of 0-1; and
    - 4.1.4 Patient has not had prior treatment with taxane chemotherapy; or
  - 4.2 All of the following:
    - 4.2.1 Patient's disease has progressed following prior chemotherapy containing a taxane; and
    - 4.2.2 Patient has ECOG performance score of 0-2; and
    - 4.2.3 Patient has not had prior treatment with abiraterone.

Renewal — (abiraterone acetate) only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Significant decrease in serum PSA from baseline; and
- 2 No evidence of clinical disease progression; and
- 3 No initiation of taxane chemotherapy with abiraterone; and
- 4 The treatment remains appropriate and the patient is benefiting from treatment.

**Renewal — (pandemic circumstances)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 2 Abiraterone acetate to be discontinued at progression; and
- 3 No initiation of taxane chemotherapy with abiraterone; and
- 4 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

BIC	ΑL	U	TAM	IDE
	_			

4.18	28	✓ Binarex
107.55	90	✓ Prostacur S29
119.50	100	✓ Flutamin
thority see SA1895 bel	ow	
1,068.00	2	✓ Faslodex
	107.55 119.50 hority see SA1895 bel	

# ⇒SA1895 Special Authority for Subsidy

**Initial application** only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has oestrogen-receptor positive locally advanced or metastatic breast cancer; and
- 2 Patient has disease progression following prior treatment with an aromatase inhibitor or tamoxifen for their locally advanced or metastatic disease: and

Subsidy (Manufacturer's Price)	Fı Subsidis	ully	Brand or Generic
\$	Per	1	Manufacturer

continued...

- 3 Treatment to be given at a dose of 500 mg monthly following loading doses; and
- 4 Treatment to be discontinued at disease progression.

**Renewal** only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

#### All of the following:

- 1 Treatment remains appropriate and patient is benefitting from treatment; and
- 2 Treatment to be given at a dose of 500 mg monthly; and
- 3 There is no evidence of disease progression.

#### OCTREOTIDE

00111201122			
Inj 50 mcg per ml, 1 ml vial	27.58	5	✓ Omega S29
Inj 100 mcg per ml, 1 ml vial		5	✓ Omega S29
Inj 500 mcg per ml, 1 ml vial	113.10	5	✓ Omega S29
Inj 50 mcg per ml, 1 ml ampoule		5	✓ Max Health
,			✓ Octreotide GH S29
Inj 100 mcg per ml, 1 ml ampoule	32.71	5	✓ Max Health
, , ,			✓ Octreotide GH S29
			✓ Sun Pharma \$29
Inj 500 mcg per ml, 1 ml ampoule	113.10	5	✓ Max Health
, , ,			✓ Octreotide GH S29
			✓ Sun Pharma S29
OCTREOTIDE LONG-ACTING - Special Authority see SA2119 be	olow – Retail pha	rmacy	
Inj depot 10 mg prefilled syringe		í	<ul> <li>Sandostatin LAR</li> </ul>
Inj depot 20 mg prefilled syringe	583.70	1	✓ Sandostatin LAR
Inj depot 30 mg prefilled syringe		1	✓ Sandostatin LAR

#### ⇒SA2119 Special Authority for Subsidy

Initial application — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 The patient has nausea\* and vomiting\* due to malignant bowel obstruction\*; and
- 2 Treatment with antiemetics, rehydration, antimuscarinic agents, corticosteroids and analgesics for at least 48 hours has failed; and
- 3 Octreotide to be given at a maximum dose 1500 mcg daily for up to 4 weeks.

Note: Indications marked with \* are unapproved indications.

**Renewal — (Malignant Bowel Obstruction)** from any relevant practitioner. Approvals valid for 3 months where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has acromegaly; and
- 2 Any of the following:
  - 2.1 Treatment with surgery, radiotherapy and a dopamine agonist has failed; or
  - 2.2 Treatment with octreotide is for an interim period while awaiting the effects of radiotherapy and a dopamine agonist has failed; or
  - 2.3 The patient is unwilling, or unable, to undergo surgery and/or radiotherapy.

Renewal — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer
--

continued...

Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 IGF1 levels have decreased since starting octreotide; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: In patients with Acromegaly octreotide treatment should be discontinued if IGF1 levels have not decreased after 3 months treatment. In patients treated with radiotherapy octreotide treatment should be withdrawn every 2 years, for 1 month, for assessment of remission. Octreotide treatment should be stopped where there is biochemical evidence of remission (normal IGF1 levels) following octreotide treatment withdrawal for at least 4 weeks

Renewal — (Acromegaly - pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has acromegaly; and
- 2 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 3 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

Initial application — (Other Indications) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

- Any of the following:
  - 1 VIPomas and Glucagonomas for patients who are seriously ill in order to improve their clinical state prior to definitive surgery; or
  - 2 Both:
    - 2.1 Gastrinoma: and
    - 2.2 Either:
      - 2.2.1 Patient has failed surgery; or
      - 2.2.2 Patient in metastatic disease after H2 antagonists (or proton pump inhibitors) have failed; or
  - 3 Both:
    - 3.1 Insulinomas: and
    - 3.2 Surgery is contraindicated or has failed; or
  - 4 For pre-operative control of hypoglycaemia and for maintenance therapy; or
  - 5 Both:
    - 5.1 Carcinoid syndrome (diagnosed by tissue pathology and/or urinary 5HIAA analysis); and
    - 5.2 Disabling symptoms not controlled by maximal medical therapy.

Note: The use of octreotide in patients with fistulae, oesophageal varices, miscellaneous diarrhoea and hypotension will not be funded as a Special Authority item

Renewal — (Other Indications) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment. Initial application — (pre-operative acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

- All of the following:
  - 1 Patient has acromegaly; and
  - 2 Patient has a large pituitary tumour, greater than 10 mm at its widest; and
  - 3 Patient is scheduled to undergo pituitary surgery in the next six months.

#### TAMOXIFEN CITRATE

*	Tab 10 mg15.00	60	✓ <u>Tamoxifen Sandoz</u>
*	Tab 20 mg5.32	60	✓ Tamoxifen Sandoz

#### **Aromatase Inhibitors**

AN.	ASTROZOLE			
*	Tab 1 mg	4.39	30	✓ Anatrole

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
EXEMESTANE  * Tab 25 mg	9.86	30	<b>√</b> <u>P</u>	fizer Exemestane
LETROZOLE  * Tab 2.5 mg	4.67	30	<b>√</b> <u>L</u>	etrole

# **Immunosuppressants**

# Cytotoxic Immunosuppressants

AZATHIOPRINE		
* Tab 25 mg7.36	60	<ul><li>Azamun</li></ul>
* Tab 50 mg8.10	100	✓ Azamun
MYCOPHENOLATE MOFETIL		
Tab 500 mg35.90	50	<ul><li>Cellcept</li></ul>
Cap 250 mg35.90	100	✓ Cellcept
Powder for oral liq 1 g per 5 ml - Subsidy by endorsement187.25	165 ml OP	✓ Cellcept

Mycophenolate powder for oral liquid is subsidised only for patients unable to swallow tablets and capsules, and when the prescription is endorsed accordingly.

#### **Fusion Proteins**

ETANERCEPT – Special Authority see SA2399 below – Retail pharmacy					
Inj 25 mg	690.00	4	<ul><li>Enbrel</li></ul>		
Inj 25 mg autoinjector	690.00	4	<ul><li>Enbrel</li></ul>		
Inj 50 mg autoinjector	1,050.00	4	<ul><li>Enbrel</li></ul>		
Inj 50 mg prefilled syringe	1,050.00	4	<ul><li>Enbrel</li></ul>		

#### ⇒SA2399 Special Authority for Subsidy

Initial application — (adult-onset Still's disease) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

#### Either:

- 1 Both:
  - 1.1 Fither:
    - 1.1.1 The patient has had an initial Special Authority approval for adalimumab for adult-onset Still's disease (AOSD); or
    - 1.1.2 The patient has been started on tocilizumab for AOSD in a Health NZ Hospital; and
  - 1.2 Fither:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or tocilizumab; or
    - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or tocilizumab such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
  - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
  - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal anti-inflammatory drugs (NSAIDs) and methotrexate; and
  - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 Fither:

Subsidy		Fully	Brand or
(Manufacturer's Price)		idised	Generic
<b>\$</b>	Per	1	Manufacturer

continued...

- 1.1 Applicant is a rheumatologist; or
- 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 The patient has a sustained improvement in inflammatory markers and functional status.

**Initial application** — **(ankylosing spondylitis)** only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for ankylosing spondylitis; and
  - 1.2 Fither:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for ankylosing spondylitis; or
- 2 All of the following:
  - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis present for more than six months; and
  - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
  - 2.3 Patient has bilateral sacroiliitis demonstrated by plain radiographs, CT or MRI scan; and
  - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
  - 2.5 Either:
    - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
    - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender (see Notes); and
  - 2.6 A Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of initial application.

Average normal chest expansion corrected for age and gender:

18-24 years - Male: 7.0 cm; Female: 5.5 cm 25-34 years - Male: 7.5 cm; Female: 5.5 cm 35-44 years - Male: 6.5 cm; Female: 4.5 cm 45-54 years - Male: 6.0 cm; Female: 5.0 cm 55-64 years - Male: 5.5 cm; Female: 4.0 cm 65-74 years - Male: 4.0 cm; Female: 4.0 cm

75+ years - Male: 3.0 cm; Female: 2.5 cm

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
  - 1.1 Applicant is a rheumatologist; or
  - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less;

Su	ubsidy Fu	lly Brand or
(Manufact	cturer's Price) Subsidise	ed Generic
	\$ Per	Manufacturer

continued...

and

- 3 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 4 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (polyarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

# 1 Both:

- 1.1 The patient has had an initial Special Authority approval for adalimumab for polyarticular course juvenile idiopathic arthritis (JIA): and
- 1.2 Either:
  - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
  - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for polyarticular course JIA; or

#### 2 All of the following:

- 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
- 2.3 Any of the following:
  - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
  - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose): or
  - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Renewal — (polyarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

# Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Fither:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
  - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

#### Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for oligoarticular course juvenile idiopathic arthritis (JIA); and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for oligoarticular course JIA; or
- 2 All of the following:
  - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and

Subsidy		Fully	Brand or
(Manufacturer's Price)	Su	bsidised	Generic
\$	Per	1	Manufacturer

continued...

- 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
- 2.3 Any of the following:
  - 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
  - 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose); or
  - 2.3.3 High disease activity (cJADAS10 score greater than 4) after a 6-month trial of methotrexate.

Renewal — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Fither:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
  - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab or secukinumab for psoriatic arthritis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab or secukinumab; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab or secukinumab to meet the renewal criteria for adalimumab or secukinumab for psoriatic arthritis; or
- 2 All of the following:
  - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
  - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
  - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
  - 2.4 Either:
    - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints;
    - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
  - 2.5 Any of the following:
    - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
    - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
    - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	sidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1 Either:
  - 1.1 Applicant is a rheumatologist; or
  - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

**Initial application — (pyoderma gangrenosum)** only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum\*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Indications marked with \* are unapproved indications.

**Renewal — (pyoderma gangrenosum)** only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement: and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

Initial application — (Arthritis - rheumatoid) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for rheumatoid arthritis; and
  - 1.2 Fither:
    - 1.2.1 The patient has experienced intolerable side effects; or
    - 1.2.2 The patient has received insufficient benefit to meet the renewal criteria for rheumatoid arthritis; or
- 2 All of the following:
  - 2.1 Patient has had rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
  - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
  - 2.3 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
  - 2.4 Patient has tried and not responded to at least three months of methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses unless contraindicated); and
  - 2.5 Either:
    - 2.5.1 Patient has tried and not responded to at least three months of methotrexate in combination with the maximum tolerated dose of ciclosporin; or
    - 2.5.2 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with methotrexate; and
  - 2.6 Fither:

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	•	Manufacturer	

continued...

- 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
- 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip.

**Renewal — (Arthritis - rheumatoid)** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
  - 2.1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

**Initial application — (severe chronic plaque psoriasis)** only from a dermatologist or any relevant practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria: Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe chronic plaque psoriasis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe chronic plaque psoriasis; or
- 2 All of the following:
  - 2.1 Any of the following:
    - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
    - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; or
    - 2.1.3 Patient has severe chronic localised genital or flexural plaque psoriasis where the plaques or lesions have been present for at least 6 months from the time of initial diagnosis, and with a Dermatology Life Quality Index (DLQI) score greater than 10; and
  - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
  - 2.3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
  - 2.4 The most recent PASI or DLQI assessment is no more than 1 month old at the time of application.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand, foot, genital or flexural areas at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and for the face, palm of a hand or sole of a foot the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis) from any relevant practitioner. Approvals valid for 6 months for applications

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

continued...

meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 Both:
    - 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
    - 1.1.2 Either:
      - 1.1.2.1 Following each prior etanercept treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-treatment baseline value; or
      - 1.1.2.2 Following each prior etanercept treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline value; or
  - 1.2 Both:
    - 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
    - 1.2.2 Either:
      - 1.2.2.1 Following each prior etanercept treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
      - 1.2.2.2 Following each prior etanercept treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; or
  - 1.3 Both:
    - 1.3.1 Patient had severe chronic localised genital or flexural plague psoriasis at the start of treatment; and
    - 1.3.2 Either:
      - 1.3.2.1 The patient has experienced a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; or
      - 1.3.2.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing etanercept; and
- 2 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Note: A treatment course is defined as a minimum of 12 weeks of etanercept treatment

Initial application — (undifferentiated spondyloarthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undifferentiated peripheral spondyloarthritis\* with active peripheral joint arthritis in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day (or maximum tolerated dose); and
- 4 Patient has tried and not responded to at least three months of leflunomide at a dose of up to 20 mg daily (or maximum tolerated dose); and
- 5 Any of the following:
  - 5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
  - 5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour measured no more than one month prior to the date of this application; or
  - 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Note: Indications marked with \* are unapproved indications.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	9	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Renewal — (undifferentiated spondyloarthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Fither:
  - 1.1 Applicant is a rheumatologist; or
  - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Fither:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg dose every 7 days.

#### Immune Modulators

ANTITHYMOCYTE GLOBULIN (EQUINE) – PCT only –	- Specialist		
Inj 50 mg per ml, 5 ml	2,774.48	5	✓ ATGAM
BACILLUS CALMETTE-GUERIN (BCG) VACCINE - PC	CT only – Specialist		
Subsidised only for bladder cancer.			
Inj 2-8 × 100 million CFU	149.37	1	✓ OncoTICE
Inj 40 mg per ml, vial	176.90	3	✓ SII-Onco-BCG S29

# Monoclonal Antibodies

	су	SA2400 below – Retali pharm	ADALIMUMAB (AMGEVITA) - Special Authority see 5
✓ Amgevita	1	190.00	Inj 20 mg per 0.4 ml prefilled syringe
✓ Amgevita	2	375.00	Inj 40 mg per 0.8 ml prefilled pen
✓ Amgevita	2	375.00	Inj 40 mg per 0.8 ml prefilled syringe

#### ⇒SA2400 Special Authority for Subsidy

Initial application — (Behcet's disease - severe) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Both:

- 1 The patient has severe Behcet's disease\* that is significantly impacting the patient's quality of life; and
- 2 Fither:
  - 2.1 The patient has severe ocular, neurological, and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s); or
  - 2.2 The patient has severe gastrointestinal, rheumatological, and/or mucocutaneous symptoms and has not responded adequately to two or more treatments appropriate for the particular symptom(s).

Note: Indications marked with \* are unapproved indications.

Initial application — (Hidradenitis suppurativa) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has hidradenitis suppurativa Hurley Stage II or Hurley Stage III lesions in distinct anatomic areas; and
- 2 Patient has tried, but had an inadequate response to at least a 90 day trial of systemic antibiotics or has demonstrated intolerance to or has contraindications for systemic antibiotics; and
- 3 Patient has 3 or more active lesions; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

4 The patient has a DLQI of 10 or more and the assessment is no more than 1 month old at time of application.

Renewal — (Hidradenitis suppurativa) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline; and
- 2 The patient has a DLQI improvement of 4 or more from baseline.

Initial application — (Plaque psoriasis - severe chronic) only from a dermatologist or any relevant practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

- 1 Both:
  - 1.1 Patient has had an initial Special Authority approval for etanercept for severe chronic plaque psoriasis; and
  - 12 Fither
    - 1.2.1 Patient has experienced intolerable side effects: or
    - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for etanercept for severe chronic plaque psoriasis; or
- 2 All of the following:
  - 2.1 Any of the following:
    - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a PASI score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
    - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; or
    - 2.1.3 Patient has severe chronic localised genital or flexural plaque psoriasis where the plaques or lesions have been present for at least 6 months from the time of initial diagnosis, and with a Dermatology Life Quality Index (DLQI) score greater than 10; and
  - 2.2 Patient has tried, but had an inadequate response to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin. or acitretin: and
  - 2.3 A PASI assessment or DLQI assessment has been completed for at least the most recent prior treatment course but no longer than 1 month following cessation of each prior treatment course and is no more than 1 month old at the time of application.

Renewal — (Plaque psoriasis - severe chronic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 Both:
  - 1.1 Patient had "whole body" severe chronic plague psoriasis at the start of treatment; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced a 75% or more reduction in PASI score, or is sustained at this level, when compared with the pre-treatment baseline value; or
    - 1.2.2 The patient has a DLQI improvement of 5 or more, when compared with the pre-treatment baseline value; or
- 2 Both:
  - 2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
  - 2.2 Either:
    - 2.2.1 The patient has experienced reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

2.2.2 The patient has experienced reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre treatment baseline value; or

- 3 Both:
  - 3.1 Patient had severe chronic localised genital or flexural plaque psoriasis at the start of treatment; and
  - 3.2 Fither
    - 3.2.1 The patient has experienced a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; or
    - 3.2.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing adalimumab.

**Initial application — (pyoderma gangrenosum)** only from a dermatologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has pyoderma gangrenosum\*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporin, azathioprine, or methotrexate) and has not received an adequate response.

Note: Indications marked with \* are unapproved indications.

Initial application — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active Crohn's disease: and
- 2 Any of the following:
  - 2.1 Patient has a CDAI score of greater than or equal to 300, or HBI score of greater than or equal to 10; or
  - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
  - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
  - 2.4 Patient has an ileostomy or colostomy and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Renewal — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 CDAI score has reduced by 100 points from the CDAI score, or HBI score has reduced by 3 points, from when the patient was initiated on adalimumab; or
- 2 CDAI score is 150 or less, or HBI is 4 or less; or
- 3 The patient has demonstrated an adequate response to treatment, but CDAI score and/or HBI score cannot be assessed.

Initial application — (Crohn's disease - children) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has active Crohn's disease; and
- 2 Either:
  - 2.1 Patient has a PCDAI score of greater than or equal to 30; or
  - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Renewal — (Crohn's disease - children) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

Subsidy		Fully	Brand or	
(Manufacturer's	Price) Su	ıbsidised	Generic	
\$	Per	/	Manufacturer	

continued...

- 1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab; or
- 2 PCDAI score is 15 or less; or
- 3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed.

**Initial application** — **(Crohn's disease - fistulising)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Crohn's disease; and
- 2 Any of the following:
  - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
  - 2.2 Patient has one or more rectovaginal fistula(e); or
  - 2.3 Patient has complex peri-anal fistula; and
- 3 A Baseline Fistula Assessment has been completed and is no more than 1 month old at the time of application.

Renewal — (Crohn's disease - fistulising) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 The number of open draining fistulae have decreased from baseline by at least 50%; or
- 2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain.

**Initial application — (Ocular inflammation - chronic)** from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 The patient has had an initial Special Authority approval for infliximab for chronic ocular inflammation; or
- 2 Both:
  - 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
  - 2.2 Any of the following:
    - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective: or
    - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
    - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Renewal — (Ocular inflammation - chronic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 12 weeks' initial treatment; or
- 2 Following each 2 year treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
- 3 Following each 2 year treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

**Initial application — (Ocular inflammation - severe)** from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 Patient has had an initial Special Authority approval for infliximab for severe ocular inflammation; or
- 2 Both:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
- 2.2 Any of the following:
  - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
  - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
  - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Renewal — (Ocular inflammation - severe) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 2 year treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 2 year treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Initial application — (ankylosing spondylitis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 Patient has had an initial Special Authority approval for etanercept for ankylosing spondylitis; and
  - 1.2 Fither:
    - 1.2.1 The patient has experienced intolerable side effects; or
    - 1.2.2 The patient has received insufficient benefit to meet the renewal criteria for ankylosing spondylitis; or
- 2 All of the following:
  - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis for more than six months; and
  - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
  - 2.3 Patient has bilateral sacroiliitis demonstrated by radiology imaging; and
  - 2.4 Patient has not responded adequately to treatment with two or more NSAIDs, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
  - 2.5 Either:
    - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following BASMI measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
    - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender; and
  - 2.6 A BASDAI of at least 6 on a 0-10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment and is no more than 1 month old at the time of application.

Renewal — (ankylosing spondylitis) from any relevant practitioner. Approvals valid for 2 years where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less.

Initial application — (Arthritis - oligoarticular course juvenile idiopathic) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for etanercept for oligoarticular course juvenile idiopathic arthritis (JIA); and

Subsidy	Fı	ılly	Brand or
(Manufacturer's Price)	Subsidis	ed	Generic
 \$	Per	✓	

continued...

- 1.2 Either:
  - 1.2.1 Patient has experienced intolerable side effects; or
  - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for oligoarticular course JIA; or
- 2 All of the following:
  - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
  - 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
  - 2.3 Either:
    - 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
    - 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose).

Renewal — (Arthritis - oligoarticular course juvenile idiopathic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

#### Fither:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - polyarticular course juvenile idiopathic) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 Patient has had an initial Special Authority approval for etanercept for polyarticular course juvenile idiopathic arthritis (JIA): and
  - 1.2 Fither:
    - 1.2.1 Patient has experienced intolerable side effects; or
    - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for polyarticular course JIA; or
- 2 All of the following:
  - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
  - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
  - 2.3 Any of the following:
    - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
    - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
    - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Renewal — (Arthritis - polyarticular course juvenile idiopathic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

#### Fither:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - psoriatic) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

Either:

#### 1 Both:

- 1.1 Patient has had an initial Special Authority approval for etanercept or secukinumab for psoriatic arthritis; and
- 1.2 Either:
  - 1.2.1 The patient has experienced intolerable side effects; or
  - 1.2.2 The patient has received insufficient benefit from to meet the renewal criteria for psoriatic arthritis; or

#### 2 All of the following:

- 2.1 Patient has had active psoriatic arthritis for six months duration or longer; and
- 2.2 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
- 2.3 Patient has tried and not responded to at least three months of sulfasalazine or leflunomide at maximum tolerated doses (unless contraindicated); and
- 2.4 Either:
  - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
  - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2.5 Any of the following:
  - 2.5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
  - 2.5.2 Patient has an ESR greater than 25 mm per hour; or
  - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

**Renewal — (Arthritis - psoriatic)** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Fither:

- 1 Following initial treatment, the patient has at least a 50% decrease in swollen joint count from baseline and a clinically significant response in the opinion of the physician; or
- 2 Patient demonstrates at least a continuing 30% improvement in swollen joint count from baseline and a clinically significant response in the opinion of the treating physician.

Initial application — (Arthritis - rheumatoid) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for etanercept for rheumatoid arthritis; and
  - 1.2 Fither:
    - 1.2.1 The patient has experienced intolerable side effects; or
    - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for rheumatoid arthritis: or
- 2 All of the following:
  - 2.1 Patient has had rheumatoid arthritis (either confirmed by radiology imaging, or the patient is CCP antibody positive) for six months duration or longer; and
  - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
  - 2.3 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
  - 2.4 Patient has tried and not responded to at least three months of methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate at maximum tolerated doses (unless contraindicated); and

Subsidy (Manufacturer's Price	)	Fully Subsidised	Brand or Generic	
\$	Per	✓	Manufacturer	

continued...

- 2.5 Either:
  - 2.5.1 Patient has tried and not responded to at least three months of methotrexate in combination with the maximum tolerated dose of ciclosporin; or
  - 2.5.2 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with methotrexate; and
- 2.6 Fither:
  - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
  - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip.

Renewal — (Arthritis - rheumatoid) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Initial application — (Still's disease - adult-onset (AOSD)) only from a rheumatologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for etanercept and/or tocilizumab for AOSD; and
  - 1.2 Fither:
    - 1.2.1 Patient has experienced intolerable side effects from etanercept and/or tocilizumab; or
    - 1.2.2 Patient has received insufficient benefit from at least a three-month trial of etanercept and/or tocilizumab; or
- 2 All of the following:
  - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria; and
  - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, NSAIDs and methotrexate; and
  - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

**Initial application — (ulcerative colitis)** from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active ulcerative colitis; and
- 2 Either:
  - 2.1 Patient's SCCAI score is greater than or equal to 4; or
  - 2.2 Patient's PUCAI score is greater than or equal to 20; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from prior therapy with immunomodulators and systemic corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Fither:

- 1 The SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on biologic therapy; or
- 2 The PUCAI score has reduced by 10 points or more from the PUCAI score when the patient was initiation on biologic therapy.

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sı	ubsidised	Generic
\$	Per	✓	Manufacturer

continued...

Initial application — (undifferentiated spondyloarthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undifferentiated peripheral spondyloarthritis\* with active peripheral joint arthritis in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2 Patient has tried and not responded to at least three months of each of methotrexate, sulfasalazine and leflunomide, at maximum tolerated doses (unless contraindicated); and
- 3 Any of the following:
  - 3.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application: or
  - 3.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application: or
  - 3.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Note: Indications marked with \* are unapproved indications

**Renewal — (undifferentiated spondyloarthritis)** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response in the opinion of the treating physician.

Initial application — (inflammatory bowel arthritis – axial) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has axial inflammatory pain for six months or more; and
- 3 Patient is unable to take NSAIDs: and
- 4 Patient has unequivocal sacroillitis demonstrated by radiological imaging or MRI; and
- 5 Patient has not responded adequately to prior treatment consisting of at least 3 months of an exercise regime supervised by a physiotherapist; and
- 6 A BASDAI of at least 6 on a 0-10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment.

**Renewal — (inflammatory bowel arthritis – axial)** from any relevant practitioner. Approvals valid for 2 years where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less.

**Initial application — (inflammatory bowel arthritis – peripheral)** only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has active arthritis in at least four joints from the following: hip, knee, ankle, subtalar, tarsus, forefoot, wrist, elbow, shoulder, sternoclavicular; and
- 3 Patient has tried and not experienced a response to at least three months of methotrexate, or azathioprine at a maximum tolerated dose (unless contraindicated); and
- 4 Patient has tried and not experienced a response to at least three months of sulfasalazine at a maximum tolerated dose (unless contraindicated); and
- 5 Any of the following:
  - 5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this

Subsidy		Fully	Brand or	
(Manufacturer's Price	,	Subsidised	Generic	
\$	Per		Manutacturer	

continued...

application; or

- 5.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application; or
- 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

**Renewal — (inflammatory bowel arthritis – peripheral)** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 Patient has experienced at least a continuing 30% improvement in active joint count from baseline in the opinion of the treating physician.

ADALIMUMAB (HUMIRA - ALTERNATIVE BRAND) - Special Authority see SA2157 below - Retail pharmacy

Inj 20 mg per 0.2 ml prefilled syringe1,	599.96	2	Humira
Inj 40 mg per 0.4 ml prefilled pen1,	599.96	2	HumiraPen
Inj 40 mg per 0.4 ml prefilled syringe1,	599.96	2	' Humira

## ⇒SA2157 Special Authority for Subsidy

Initial application — (Behcet's disease – severe) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Behcet's disease – severe) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient has had a good clinical response to treatment with measurably improved quality of life; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Hidradenitis suppurativa) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 7 days. Fortnightly dosing has been considered.

**Renewal — (Hidradenitis suppurativa)** only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline: and
- 2 The patient has a Dermatology Quality of Life Index improvement of 4 or more from baseline; and
- 3 Adalimumab is to be administered at doses no greater than 40mg every 7 days. Fortnightly dosing has been considered.

Initial application — (Psoriasis - severe chronic plaque) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Either:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Psoriasis - severe chronic plaque) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Either:
  - 1.1 Both:
    - 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
    - 1.1.2 Either:
      - 1.1.2.1 Following each prior adalimumab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-adalimumab treatment baseline value; or
      - 1.1.2.2 Following each prior adalimumab treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline value; or
  - 1.2 Both:
    - 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment: and
    - 1.2.2 Fither:
      - 1.2.2.1 Following each prior adalimumab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
      - 1.2.2.2 Following each prior adalimumab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-adalimumab treatment baseline value: and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Pyoderma gangrenosum) only from a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and

	Subsidy		Fully	Brand or
(1	Manufacturer's Price)	Subs	sidised	Generic
	\$	Per	1	Manufacturer

continued...

- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 A maximum of 8 doses.

**Renewal — (Pyoderma gangrenosum)** only from a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient has demonstrated clinical improvement and continues to require treatment; and
- 2 A maximum of 8 doses.

Initial application — (Crohn's disease - adult) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:
All of the following:

- 1 Any of the following:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevitat; or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
  - 1.3 Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Crohn's disease - adult) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on adalimumab; or
  - 1.2 CDAI score is 150 or less: or
  - 1.3 The patient has demonstrated an adequate response to treatment, but CDAI score cannot be assessed; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Crohn's disease - children) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
  - 1.3 Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Crohn's disease - children) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab; or
  - 1.2 PCDAI score is 15 or less: or

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsid	dised	Generic	
\$	Per	1	Manufacturer	

continued...

- 1.3 The patient has demonstrated an adequate response to treatment, but PCDAI score cannot be assessed; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Crohn's disease - fistulising) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
  - 1.3 Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Crohn's disease - fistulising) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
  - 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Ocular inflammation – chronic) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with Amgevita, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
  - 1.3 Patient has uveitis and is considered to be at risk of vision loss if they were to change treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Ocular inflammation – chronic) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 The patient has had a good clinical response following 12 weeks' initial treatment; or
  - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
  - 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and</p>
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Subsidy (Manufacturer's Price)	Fully Subsidised	
\$	Per 🗸	Manufacturer

continued...

Initial application — (Ocular inflammation – severe) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with Amgevita, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
  - 1.3 Patient has uveitis and is considered to be at risk of vision loss if they were to change treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Ocular inflammation – severe) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 The patient has had a good clinical response following 3 initial doses; or
  - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
  - 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and</p>
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita): and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

**Renewal — (ankylosing spondylitis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Arthritis – oligoarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or

Subsidy	F	ully	Brand or	_
(Manufacturer's Price)	Subsidi	sed	Generic	
\$	Per	1	Manufacturer	

continued...

- 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

Renewal — (Arthritis – oligoarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months where the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - polyarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

Renewal — (Arthritis - polyarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months where the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - psoriatic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Arthritis - psoriatic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Arthritis – rheumatoid) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	sidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Fither:
  - 4.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or
  - 4.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

Renewal — (Arthritis – rheumatoid) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 2 Fither:
  - 2.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or
  - 2.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

Initial application — (Still's disease – adult-onset (AOSD)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

Renewal — (Still's disease – adult-onset (AOSD)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months where the patient has demonstrated a sustained improvement in inflammatory markers and functional status.

AFLIBERCEPT - Special Authority see SA1772 below - Retail pharmacy

#### ⇒SA1772 Special Authority for Subsidy

Initial application — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 3 months for applications meeting the following criteria:

Fither:

- 1 All of the following:
  - 1.1 Any of the following:
    - 1.1.1 Wet age-related macular degeneration (wet AMD); or
    - 1.1.2 Polypoidal choroidal vasculopathy; or
    - 1.1.3 Choroidal neovascular membrane from causes other than wet AMD; and
  - 1.2 Either:
    - 1.2.1 The patient has developed severe endophthalmitis or severe posterior uveitis following treatment with bevacizumab; or
    - 1.2.2 There is worsening of vision or failure of retina to dry despite three intraocular injections of bevacizumab four weeks apart; and

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

- 1.3 There is no structural damage to the central fovea of the treated eye; and
- 1.4 Patient has not previously been treated with ranibizumab for longer than 3 months; or
- 2 Fither:
  - 2.1 Patient has current approval to use ranibizumab for treatment of wAMD and was found to be intolerant to ranibizumab within 3 months; or
  - 2.2 Patient has previously\* (\*before June 2018) received treatment with ranibizumab for wAMD and disease was stable while on treatment.

**Initial application — (diabetic macular oedema)** only from an ophthalmologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has centre involving diabetic macular oedema (DMO); and
- 2 Patient's disease is non responsive to 4 doses of intravitreal bevacizumab when administered 4-6 weekly; and
- 3 Patient has reduced visual acuity between 6/9 6/36 with functional awareness of reduction in vision; and
- 4 Patient has DMO within central OCT (ocular coherence tomography) subfield > 350 micrometers; and
- 5 There is no centre-involving sub-retinal fibrosis or foveal atrophy.

Renewal — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Documented benefit must be demonstrated to continue; and
- 2 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 3 There is no structural damage to the central fovea of the treated eye.

Renewal — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 There is stability or two lines of Snellen visual acuity gain; and
- 2 There is structural improvement on OCT scan (with reduction in intra-retinal cysts, central retinal thickness, and sub-retinal fluid); and
- 3 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 4 There is no centre-involving sub-retinal fibrosis or foveal atrophy; and
- 5 After each consecutive 12 months treatment with (2nd line anti-VEGF agent), patient has retrialled with at least one injection of bevacizumab and had no response.

BENRALIZUMAB - Special Authority see SA2151 below - Retail pharmacy

⇒SA2151 Special Authority for Subsidy

Initial application — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Patient must have a diagnosis of severe eosinophilic asthma documented by a respiratory physician or clinical immunologist; and
- 3 Conditions that mimic asthma eg. vocal cord dysfunction, central airway obstruction, bronchiolitis etc. have been excluded; and
- 4 Patient has a blood eosinophil count of greater than 0.5 × 10^9 cells/L in the last 12 months; and
- 5 Patient must be adherent to optimised asthma therapy including inhaled corticosteroids (equivalent to at least 1000 mcg per day of fluticasone propionate) plus long-acting beta-2 agonist, or budesonide/formoterol as part of the anti-inflammatory reliever therapy plus maintenance regimen, unless contraindicated or not tolerated; and

	Subsidy	F	ully	Brand or
(M	Ianufacturer's Price)	Subsidised		Generic
	\$	Per	•	Manufacturer

continued...

- 6 Either:
  - 6.1 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral corticosteroids: or
  - 6.2 Patient has received continuous oral corticosteroids of at least the equivalent of 10 mg per day over the previous 3 months; and
- 7 Treatment is not to be used in combination with subsidised mepolizumab; and
- 8 Patient has an Asthma Control Test (ACT) score of 10 or less. Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 52 weeks after the first dose to assess response to treatment; and
- 9 Either
  - 9.1 Patient has not previously received an anti-IL5 biological therapy for their severe eosinophilic asthma; or
  - 9.2 Both:
    - 9.2.1 Patient was refractory or intolerant to previous anti-IL5 biological therapy; and
    - 9.2.2 Patient was not eligible to continue treatment with previous anti-IL5 biological therapy and discontinued within 12 months of commencing treatment.

Renewal — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 Either:
  - 2.1 Exacerbations have been reduced from baseline by 50% as a result of treatment with benralizumab; or
  - 2.2 Reduction in continuous oral corticosteroid use by 50% or by 10 mg/day while maintaining or improving asthma control

⇒SA2289 Special Authority for Subsidy

**Initial application — (relapsed/refractory Hodgkin lymphoma)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 Both:
    - 1.1.1 Patient has relapsed/refractory CD30-positive Hodgkin lymphoma after two or more lines of chemotherapy;
    - 1.1.2 Patient is ineligible for autologous stem cell transplant; or
  - 1.2 Both:
    - 1.2.1 Patient has relapsed/refractory CD30-positive Hodgkin lymphoma; and
    - 1.2.2 Patient has previously undergone autologous stem cell transplant; and
- 2 Patient has not previously received funded brentuximab vedotin; and
- 3 Response to brentuximab vedotin treatment is to be reviewed after a maximum of 6 treatment cycles; and
- 4 Brentuximab vedotin to be administered at doses no greater than 1.8 mg/kg every 3 weeks.

**Renewal — (relapsed/refractory Hodgkin lymphoma)** from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has achieved a partial or complete response to brentuximab vedotin after 6 treatment cycles; and
- 2 Treatment remains clinically appropriate and the patient is benefitting from treatment and treatment is being tolerated; and

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer	sidised Generic
--	-----------------

continued...

3 Patient is to receive a maximum of 16 total cycles of brentuximab vedotin treatment.

**Initial application — (anaplastic large cell lymphoma)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has relapsed/refractory CD30-positive systemic anaplastic large cell lymphoma; and
- 2 Patient has an ECOG performance status of 0-1; and
- 3 Patient has not previously received brentuximab vedotin; and
- 4 Response to brentuximab vedotin treatment is to be reviewed after a maximum of 6 treatment cycles; and
- 5 Brentuximab vedotin to be administered at doses no greater than 1.8 mg/kg every 3 weeks.

Renewal — (anaplastic large cell lymphoma) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has achieved a partial or complete response to brentuximab vedotin after 6 treatment cycles; and
- 2 Treatment remains clinically appropriate and the patient is benefitting from treatment and treatment is being tolerated; and
- 3 Patient is to receive a maximum of 16 total cycles of brentuximab vedotin treatment.

#### CASIRIVIMAB AND IMDEVIMAB - [Xpharm] - Special Authority see SA2096 below

#### ⇒SA2096 Special Authority for Subsidy

Initial application — (Treatment of profoundly immunocompromised patients) from any relevant practitioner. Approvals valid for 2 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed (or probable) COVID-19; and
- 2 The patient is in the community with mild to moderate disease severity\*; and
- 3 Patient is profoundly immunocompromised\*\* and is at risk of not having mounted an adequate response to vaccination against COVID-19 or is unvaccinated; and
- 4 Patient's symptoms started within the last 10 days; and
- 5 Patient is not receiving high flow oxygen or assisted/mechanical ventilation; and
- 6 Casirivimab and imdevimab is to be administered at a maximum dose of no greater than 2,400 mg.

Notes: \* Mild to moderate disease severity as described on the Ministry of Health Website

\*\* Examples include B-cell depletive illnesses or patients receiving treatment that is B-Cell depleting.

#### CETUXIMAB - PCT only - Specialist - Special Authority see SA2401 below

Inj 5 mg per ml, 20 ml vial	364.00	1	Erbitux
Inj 5 mg per ml, 100 ml vial	1,820.00	1	Erbitux
Inj 1 mg for ECP	3.82	1 mg	✓ Baxter

#### ⇒SA2401 Special Authority for Subsidy

Initial application — (head and neck cancer, locally advanced) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, non-metastatic, squamous cell cancer of the head and neck; and
- 2 Cisplatin is contraindicated or has resulted in intolerable side effects; and
- 3 Patient has an ECOG performance score of 0-2; and
- 4 To be administered in combination with radiation therapy.

Initial application — (colorectal cancer, metastatic) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Por 🗸	Manufacturer

continued...

- 1 Patient has metastatic colorectal cancer located on the left side of the colon (see Note); and
- 2 There is documentation confirming disease is RAS and BRAF wild-type; and
- 3 Patient has an ECOG performance score of 0-2; and
- 4 Patient has not received prior funded treatment with cetuximab; and
- 5 Either:
  - 5.1 Cetuximab is to be used in combination with chemotherapy; or
  - 5.2 Chemotherapy is determined to not be in the best interest of the patient based on clinician assessment.

Renewal — (colorectal cancer, metastatic) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where there is no evidence of disease progression.

Note: Left-sided colorectal cancer comprises of the distal one-third of the transverse colon, the splenic flexure, the descending colon, the sigmoid colon, or the rectum.

## ⇒SA2269 Special Authority for Subsidy

**Initial application** only from a haematologist, paediatric haematologist or paediatric oncologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has not received prior chemotherapy for this condition; and
- 2 Patient has de novo CD33-positive acute myeloid leukaemia; and
- 3 Patient does not have acute promyelocytic leukaemia; and
- 4 Gemtuzumab ozogamicin will be used in combination with standard anthracycline and cytarabine (AraC); and
- 5 Patient is being treated with curative intent; and
- 6 Patient's disease risk has been assessed by cytogenetic testing to be good or intermediate; and
- 7 Patient must be considered eligible for standard intensive remission induction chemotherapy with standard anthracycline and cytarabine (AraC); and
- 8 Gemtuzumab ozogamicin to be funded for one course only (one dose at 3 mg per m² body surface area or up to 2 vials of 5 mg as separate doses).

Note: Acute myeloid leukaemia excludes acute promyelocytic leukaemia and acute myeloid leukaemia that is secondary to another haematological disorder (eg myelodysplasia or myeloproliferative disorder).

# **⇒SA2402** Special Authority for Subsidy

**Initial application — (Crohn's disease (adults))** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active Crohn's disease; and
- 2 Any of the following:
  - 2.1 Patient has a CDAI score of greater than or equal to 300 or HBI score of greater than or equal to 10; or
  - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
  - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
  - 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but has experienced an inadequate response to, or intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Renewal — (Crohn's disease (adults)) from any relevant practitioner. Approvals valid for 2 years for applications meeting the

Subsidy	Full	/ Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per 🗸	Manufacturer Manufacturer	

continued...

following criteria:

Both:

- 1 Any of the following:
  - 1.1 CDAI score has reduced by 100 points from the CDAI score, or HBI score has reduced by 3 points, from when the patient was initiated on infliximab; or
  - 1.2 CDAI score is 150 or less, or HBI is 4 or less; or
  - 1.3 The patient has demonstrated an adequate response to treatment but CDAI score and/or HBI score cannot be assessed: and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (Crohn's disease (children)) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has active Crohn's disease; and
- 2 Either:
  - 2.1 Patient has a PCDAI score of greater than or equal to 30; or
  - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but experienced an inadequate response to, or intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Renewal — (Crohn's disease (children)) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on infliximab; or
  - 1.2 PCDAI score is 15 or less; or
  - 1.3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

**Initial application** — **(Graft vs host disease)** from any relevant practitioner. Approvals valid without further renewal unless notified where patient has steroid-refractory acute graft vs. host disease of the qut.

Initial application — (Pulmonary sarcoidosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has life-threatening pulmonary sarcoidosis diagnosed by a multidisciplinary team that is refractory to other treatments.

Initial application — (acute fulminant ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 weeks for applications meeting the following criteria:
Both:

- 1 Patient has acute, fulminant ulcerative colitis; and
- 2 Treatment with intravenous or high dose oral corticosteroids has not been successful.

Initial application — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and
- 2 Fither:

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
\$	Per	✓	

continued...

- 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
- 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

**Renewal — (ankylosing spondylitis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Following 12 weeks of infliximab treatment, BASDAI has improved by 4 or more points from pre-infliximab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 3 Infliximab to be administered at doses no greater than 5 mg/kg every 6-8 weeks.

**Initial application — (chronic ocular inflammation)** from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for chronic ocular inflammation; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for chronic ocular inflammation; or
- 2 Both:
  - 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
  - 2.2 Any of the following:
    - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective: or
    - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
    - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

**Renewal — (chronic ocular inflammation)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.</p>

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (fistulising Crohn's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has confirmed Crohn's disease: and
- 2 Any of the following:
  - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

- 2.2 Patient has one or more rectovaginal fistula(e); or
- 2.3 Patent has complex peri-anal fistula.

Renewal — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
  - 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline (in the case of adult patients, as demonstrated by a reduction in the Fistula Assessment score), together with less induration and patient reported pain; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (neurosarcoidosis) only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with neurosarcoiosis by a multidisciplinary team; and
- 2 Patient has CNS involvement; and
- 3 Patient has steroid-refractory disease; and
- 4 Either:
  - 4.1 IV cyclophosphamide has been tried; or
  - 4.2 Treatment with IV cyclophosphamide is clinically inappropriate.

**Renewal — (neurosarcoidosis)** only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria:

Either:

- 1 A withdrawal period has been tried and the patient has relapsed; or
- 2 All of the following:
  - 2.1 A withdrawal period has been considered but would not be clinically appropriate; and
  - 2.2 There has been a marked reduction in prednisone dose; and
  - 2.3 Either:
    - 2.3.1 There has been an improvement in MRI appearances; or
    - 2.3.2 Marked improvement in other symptomology.

Initial application — (plaque psoriasis) only from a dermatologist or any relevant practitioner on the recommendation of a dermatologist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab, etanercept or secukinumab for severe chronic plaque psoriasis; and
  - 1.2 Either:
    - 1.2.1 Patient has experienced intolerable side effects from adalimumab, etanercept or secukinumab; or
    - 1.2.2 Patient has received insufficient benefit from adalimumab, etanercept or secukinumab to meet the renewal criteria for adalimumab, etanercept or secukinumab for severe chronic plaque psoriasis; or
- 2 All of the following:
  - 2.1 Any of the following:
    - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial

Subsidy (Manufacturer's Price)	Subs	Fully	Brand or Generic	
(Manufacturer's Frice)	Per	siuiseu •	Manufacturer	

continued...

- diagnosis; or
- 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; or
- 2.1.3 Patient has severe chronic localised genital or flexural plaque psoriasis where the plaques or lesions have been present for at least 6 months from the time of initial diagnosis, and with a Dermatology Life Quality Index (DLQI) score greater than 10; and
- 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
- 2.3 A PASI assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course: and
- 2.4 The most recent PASI assessment is no more than 1 month old at the time of initiation.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand, foot, genital or flexural areas at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and for the face, palm of a hand or sole of a foot the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

**Renewal — (plaque psoriasis)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 Any of the following:

#### 1.1 Both:

- 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
- 1.1.2 Following each prior infliximab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-infliximab treatment baseline value; or

## 1.2 Both:

- 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
- 1.2.2 Either:
  - 1.2.2.1 Following each prior infliximab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
  - 1.2.2.2 Following each prior infliximab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-infliximab treatment baseline value: or
- 1.3 Both:
  - 1.3.1 Patient had severe chronic localised genital or flexural plaque psoriasis at the start of treatment; and
  - 1.3.2 Either:
    - 1.3.2.1 The patient has experienced a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; or
    - 1.3.2.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing infliximab; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Su	bsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Both:

- 1 Patient was being treated with infliximab prior to 1 February 2019; and
- 2 Any of the following:
  - 2.1 Rheumatoid arthritis: or
  - 2.2 Ankylosing spondylitis; or
  - 2.3 Psoriatic arthritis; or
  - 2.4 Severe ocular inflammation: or
  - 2.5 Chronic ocular inflammation: or
  - 2.6 Crohn's disease (adults); or
  - 2.7 Crohn's disease (children): or
  - 2.8 Fistulising Crohn's disease; or
  - 2.9 Severe fulminant ulcerative colitis; or
  - 2.10 Severe ulcerative colitis: or
  - 2.11 Plaque psoriasis; or
  - 2.12 Neurosarcoidosis; or
  - 2.13 Severe Behcet's disease.

Initial application — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis; and
- 2 Fither:
  - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept and/or secukinumab; or
  - 2.2 Following 3-4 months' initial treatment with adalimumab and/or etanercept and/or secukinumab, the patient did not meet the renewal criteria for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior infliximab treatment in the opinion of the treating physician; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 1 The patient 2 Either:
  - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
  - 2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept; and
- 3 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance.

Renewal — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	/	Manufacturer

continued...

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Infliximab to be administered at doses no greater than 3 mg/kg every 8 weeks.

**Initial application — (severe Behcet's disease)** from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe Behcet's disease which is significantly impacting the patient's quality of life (see Notes); and
- 2 Fither
  - 2.1 The patient has severe ocular, neurological and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s) (see Notes); or
  - 2.2 The patient has severe gastrointestinal, rheumatologic and/or mucocutaneous symptoms and has not responded adequately to two or more treatment appropriate for the particular symptom(s) (see Notes); and
- 3 The patient is experiencing significant loss of quality of life.

Notes: Behcet's disease diagnosed according to the International Study Group for Behcet's Disease. Lancet 1990;335(8697):1078-80. Quality of life measured using an appropriate quality of life scale such as that published in Gilworth et al J Rheumatol. 2004;31:931-7.

Treatments appropriate for the particular symptoms are those that are considered standard conventional treatments for these symptoms, for example intravenous/oral steroids and other immunosuppressants for ocular symptoms; azathioprine, steroids, thalidomide, interferon alpha and ciclosporin for mucocutaneous symptoms; and colchicine, steroids and methotrexate for rheumatological symptoms.

Renewal — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has had a good clinical response to initial treatment with measurably improved quality of life; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Renewal — (fulminant ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Where maintenance treatment is considered appropriate, infliximab should be used in combination with immunomodulators and reassessed every 6 months; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

**Initial application — (severe ocular inflammation)** from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe ocular inflammation; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe ocular inflammation: or

- 2 Both:
  - 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
  - 2.2 Any of the following:
    - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
    - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
    - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

**Renewal — (severe ocular inflammation)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (ulcerative colitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active ulcerative colitis; and
- 2 Either:
  - 2.1 Patients SCCAI is greater than or equal to 4: or
  - 2.2 Patients PUCAI score is greater than or equal to 20; and
- 3 Patient has tried but has experienced an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and systemic corticosteroids.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 The SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on infliximals or
  - 1.2 The PUCAI score has reduced by 10 points or more from the PUCAI score when the patient was initiated on infliximab; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

**Initial application — (pyoderma gangrenosum)** only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

1 Patient has pyoderma gangrenosum\*; and

Subsidy (Manufacturer's I	Price) Sub	Fully	Brand or Generic	
\$	Per	•	Manufacturer	

continued...

- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Note: Indications marked with \* are unapproved indications.

**Renewal — (pyoderma gangrenosum)** only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

Initial application — (inflammatory bowel arthritis – axial) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has had axial inflammatory pain for six months or more; and
- 3 Patient is unable to take NSAIDs: and
- 4 Patient has unequivocal sacroiliitis demonstrated by radiological imaging or MRI; and
- 5 Patient's disease has not responded adequately to prior treatment consisting of at least 3 months of an exercise regime supervised by a physiotherapist; and
- 6 Patient has a BASDAI of at least 6 on a 0 10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment.

Renewal — (inflammatory bowel arthritis – axial) from any relevant practitioner. Approvals valid for 2 years where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10-point scale, or an improvement in BASDAI of 50%, whichever is less.

Initial application — (inflammatory bowel arthritis – peripheral) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has active arthritis in at least four joints from the following: hip, knee, ankle, subtalar, tarsus, forefoot, wrist, elbow, shoulder, sternoclavicular; and
- 3 Patient has tried and not experienced a response to at least three months of methotrexate or azathioprine at a maximum tolerated dose (unless contraindicated); and
- 4 Patient has tried and not experienced a response to at least three months of sulfasalazine at a maximum tolerated dose (unless contraindicated); and
- 5 Any of the following:
  - 5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
  - 5.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application; or
  - 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

**Renewal — (inflammatory bowel arthritis – peripheral)** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Fither:

- 1 Following initial treatment, patient has experienced at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 Patient has experienced at least a continuing 30% improvement in active joint count from baseline in the opinion of the

Nucala

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

treating physician.

MEPOLIZUMAB - Special Authority see SA2331 below - Retail pharmacy

# ⇒SA2331 Special Authority for Subsidy

Initial application — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Patient must have a diagnosis of severe eosinophilic asthma documented by a respiratory physician or clinical immunologist; and
- 3 Conditions that mimic asthma eg. vocal cord dysfunction, central airway obstruction, bronchiolitis etc. have been excluded: and
- 4 Patient has a blood eosinophil count of greater than 0.5 × 10<sup>9</sup> cells/L in the last 12 months; and
- 5 Patient must be adherent to optimised asthma therapy including inhaled corticosteroids (equivalent to at least 1000 mcg per day of fluticasone propionate) plus long acting beta-2 agonist, or budesonide/formoterol as part of the single maintenance and reliever therapy regimen, unless contraindicated or not tolerated; and
- 6 Either:
  - 6.1 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral corticosteroids: or
  - 6.2 Patient has received continuous oral corticosteroids of at least the equivalent of 10 mg per day over the previous 3 months: and
- 7 Treatment is not to be used in combination with subsidised benralizumab; and
- 8 Patient has an Asthma Control Test (ACT) score of 10 or less. Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 52 weeks after the first dose to assess response to treatment; and
- 9 Either:
  - 9.1 Patient has not previously received an anti-IL5 biological therapy for their severe eosinophilic asthma; or
  - 9.2 Both:
    - 9.2.1 Patient was refractory or intolerant to previous anti-IL5 biological therapy; and
    - 9.2.2 Patient was not eligible to continue treatment with previous anti-IL5 biological therapy and discontinued within 12 months of commencing treatment.

**Renewal — (Severe eosinophilic asthma)** only from a respiratory physician or clinical immunologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 Either:
  - 2.1 Exacerbations have been reduced from baseline by 50% as a result of treatment with mepolizumab; or
  - 2.2 Reduction in continuous oral corticosteroid use by 50% or by 10 mg/day while maintaining or improving asthma control.

Initial application — (eosinophilic granulomatosis with polyangiitis) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has eosinophilic granulomatosis with polyangiitis; and
- 2 The patient has trialled and not received adequate benefit from at least one of the following for at least three months (unless contraindicated to all): azathioprine, cyclophosphamide, leflunomide, methotrexate, mycophenolate, or rituximab;

 Subsidy		Fully	Brand or
(Manufacturer's Price)	_	Subsidised	Generic
\$	Per		Manufacturer

continued...

and 3 Either:

- 3.1 The patient has trialled prednisone for a minimum of three months and is unable to maintain disease control at doses below 7.5 mg per day; or
- 3.2 Corticosteroids are contraindicated.

Renewal — (eosinophilic granulomatosis with polyangiitis) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months where patient has no evidence of clinical disease progression.

# ⇒SA2155 Special Authority for Subsidy

**Initial application — (chronic lymphocytic leukaemia)** only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C CD20+ chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is obinutuzumab treatment naive: and
- 3 The patient is not eligible for full dose FCR due to comorbidities with a score > 6 on the Cumulative Illness Rating Scale (CIRS) or reduced renal function (creatinine clearance < 70mL/min); and
- 4 Patient has adequate neutrophil and platelet counts\* unless the cytopenias are a consequence of marrow infiltration by CLL; and
- 5 Patient has good performance status; and
- 6 Obinutuzumab to be administered at a maximum cumulative dose of 8,000 mg and in combination with chlorambucil for a maximum of 6 cycles.

Notes: Chronic lymphocytic leukaemia includes small lymphocytic lymphoma. Comorbidity refers only to illness/impairment other than CLL induced illness/impairment in the patient. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with obinutuzumab is expected to improve symptoms and improve ECOG score to < 2.

\* Neutrophil greater than or equal to  $1.5 \times 10^9$ /L and platelets greater than or equal to  $75 \times 10^9$ /L.

Initial application — (follicular / marginal zone lymphoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: All of the following:

- 1 Either:
  - 1.1 Patient has follicular lymphoma; or
  - 1.2 Patient has marginal zone lymphoma; and
- 2 Patient is refractory to or has relapsed within 12 months of a rituximab containing combined chemo-immunotherapy regimen\*; and
- 3 Patient has an ECOG performance status of 0-2; and
- 4 Patient has been previously treated with no more than four chemotherapy regimens; and
- 5 Obinutuzumab to be administered at a maximum dose of 1000 mg for a maximum of 6 cycles in combination with chemotherapy\*.

Note: \* includes unapproved indications

**Renewal — (follicular / marginal zone lymphoma)** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 24 months for applications meeting the following criteria: All of the following:

- 1 Patient has no evidence of disease progression following obinutuzumab induction therapy; and
- 2 Obinutuzumab to be administered at a maximum of 1000 mg every 2 months for a maximum of 2 years; and
- 3 Obinutuzumab to be discontinued at disease progression.

	Subsidy		Fully	Brand or	
	(Manufacturer's Price)		Subsidised	Generic	
	\$	Per	1	Manufacturer	
OMALIZUMAB - Special Authority see SA1744 below - Retail ph	armacy				
Inj 150 mg prefilled syringe	450.00	1	<b>√</b> )	(olair	
, 01 , 0			<b>✓</b> )	(olair AU	
Inj 150 mg vial	450.00	1	_	Colair	

# ⇒SA1744 Special Authority for Subsidy

Initial application — (severe asthma) only from a respiratory specialist or clinical immunologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 6 years or older; and
- 2 Patient has a diagnosis of severe asthma; and
- 3 Past or current evidence of atopy, documented by skin prick testing or RAST; and
- 4 Total serum human immunoglobulin E (IgE) between 76 IU/mL and 1300 IU/ml at baseline; and
- 5 Proven adherence with optimal inhaled therapy including high dose inhaled corticosteroid (budesonide 1,600 mcg per day or fluticasone propionate 1,000 mcg per day or equivalent), plus long-acting beta-2 agonist therapy (at least salmeterol 50 mcg bd or eformoterol 12 mcg bd) for at least 12 months, unless contraindicated or not tolerated; and
- 6 Fither:
  - 6.1 Patient has received courses of systemic corticosteroids equivalent to at least 28 days treatment in the past 12 months. unless contraindicated or not tolerated; or
  - 6.2 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral steroids; and
- 7 Patient has an Asthma Control Test (ACT) score of 10 or less; and
- 8 Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 26 weeks after the first dose to assess response to treatment.

Initial application — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

- All of the following:
  - 1 Patient must be aged 12 years or older; and
  - 2 Fither:
    - 2.1 Both:
      - 2.1.1 Patient is symptomatic with Urticaria Activity Score 7 (UAS7) of 20 or above; and
      - 2.1.2 Patient has a Dermatology life quality index (DLQI) of 10 or greater; or
    - 2.2 Patient has a Urticaria Control Test (UCT) of 8 or less; and
  - 3 Any of the following:
    - 3.1 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and ciclosporin (> 3 mg/kg day) for at least 6 weeks; or
    - 3.2 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and at least 3 courses of systemic corticosteroids (> 20 mg prednisone per day for at least 5 days) in the previous 6 months; or
    - 3.3 Patient has developed significant adverse effects whilst on corticosteroids or ciclosporin; and
  - 4 Either:
    - 4.1 Treatment to be stopped if inadequate response\* following 4 doses; or
    - 4.2 Complete response\* to 6 doses of omalizumab.

**Renewal — (severe asthma)** only from a clinical immunologist or respiratory specialist. Approvals valid for 2 years for applications meeting the following criteria:

- Both:
  - 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
  - 2 A reduction in the maintenance oral corticosteroid dose or number of exacerbations of at least 50% from baseline.

Subsidy	y Full	/ Brand or
(Manufacturer's	s Price) Subsidise	d Generic
\$	Per 🗸	Manufacturer

continued...

Renewal — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

# Either:

- 1 Patient has previously adequately responded\* to 6 doses of omalizumab; or
- 2 Both:
  - 2.1 Patient has previously had a complete response\* to 6 doses of omalizumab; and
  - 2.2 Patient has relapsed after cessation of omalizumab therapy.

Note: \*Inadequate response defined as less than 50% reduction in baseline UAS7 and DLQI score, or an increase in Urticaria Control Test (UCT) score of less than 4 from baseline. Patient is to be reassessed for response after 4 doses of omalizumab. Complete response is defined as UAS7 less than or equal to 6 and DLQI less than or equal to 5; or UCT of 16. Relapse of chronic urticaria on stopping prednisone/ciclosporin does not justify the funding of omalizumab.

PERTUZUMAB - PCT only - Specialist - Special Authority see SA2276 below

Inj 30 mg per ml, 14 ml vial	3,927.00	1	✓ Perjeta
Inj 420 mg for ECP	3,927.00	420 mg OP	✓ Baxter

# ⇒SA2276 Special Authority for Subsidy

**Initial application — (metastatic breast cancer)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Fither:
  - 2.1 Patient is chemotherapy treatment naïve; or
  - 2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
- 3 The patient has good performance status (ECOG grade 0-1); and
- 4 Pertuzumab to be administered in combination with trastuzumab; and
- 5 Pertuzumab maximum first dose of 840 mg, followed by maximum of 420 mg every 3 weeks; and
- 6 Pertuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

#### Either:

- 1 Both:
  - 1.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
  - 1.2 The cancer has not progressed at any time point during the previous 12 months whilst on pertuzumab and trastuzumab; or
- 2 All of the following:
  - 2.1 Patient has previously discontinued treatment with pertuzumab and trastuzumab for reasons other than severe toxicity or disease progression; and
  - 2.2 Patient has signs of disease progression; and
  - 2.3 Disease has not progressed during previous treatment with pertuzumab and trastuzumab.

RITUXIMAB (MABTHERA) - PCT only -	ecialist – Special Authority see SA1976 on the next page
-----------------------------------	--

Inj 100 mg per 10 ml vial	1,075.50	2	<ul><li>Mabthera</li></ul>
Inj 500 mg per 50 ml vial	2,688.30	1	✓ Mabthera
Inj 1 mg for ECP	5.64	1 ma	✓ Baxter (Mabthera)

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

# ⇒SA1976 Special Authority for Subsidy

Initial application — (rheumatoid arthritis - TNF inhibitors contraindicated) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Treatment with a Tumour Necrosis Factor alpha inhibitor is contraindicated: and
- 2 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
- 5 Any of the following:
  - 5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or
  - 5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
  - 5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
- 6 Either:
  - 6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
  - 6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 7 Either:
  - 7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
  - 7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months; and
- 8 Fither:
  - 8.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
  - 8.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 9 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Initial application — (rheumatoid arthritis - prior TNF inhibitor use) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Both:
  - 1.1 The patient has had an initial community Special Authority approval for at least one of etanercept and/or adalimumab for rheumatoid arthritis: and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
    - 1.2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept for rheumatoid arthritis: and
- 2 Fither:
  - 2.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
  - 2.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 3 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

5	Subsidy	Fully	Brand or
(Manufa	acturer's Price)	Subsidised	Generic
	\$ Per	•	Manufacturer

continued...

Renewal — (rheumatoid arthritis - re-treatment in 'partial responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 At 4 months following the initial course of rituximab infusions the patient had between a 30% and 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 1.2 At 4 months following the second course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 1.3 At 4 months following the third and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Either:
  - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
- 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Renewal — (rheumatoid arthritis - re-treatment in 'responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Either:
  - 1.1 At 4 months following the initial course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 1.2 At 4 months following the second and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Either:
  - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
  - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

RITUXIMAB (RIXIMYO) - PCT only - Specialist - Special Au	uthority see SA2233 b	pelow	
Inj 100 mg per 10 ml vial	275.33	2	✓ Riximyo
Inj 500 mg per 50 ml vial	688.20	1	✓ Riximyo
Inj 1 mg for ECP	1.38	1 mg	✓ Baxter (Riximyo)

#### ⇒SA2233 Special Authority for Subsidy

Initial application — (ABO-incompatible organ transplant) from any relevant practitioner. Approvals valid without further renewal unless notified where patient is to undergo an ABO-incompatible solid organ transplant\*.

Note: Indications marked with \* are unapproved indications.

**Initial application** — **(ANCA associated vasculitis)** from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis\*; and
- 2 The total rituximab dose would not exceed the equivalent of 375 mg/m<sup>2</sup> of body-surface area per week for a total of 4 weeks; and
- 3 Any of the following:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 3.1 Induction therapy with daily oral or pulse intravenous cyclophosphamide has failed to achieve significant improvement of disease after at least 3 months; or
- 3.2 Patient has previously had a cumulative dose of cyclophosphamide > 15 g or a further repeat 3 month induction course of cyclophosphamide would result in a cumulative dose > 15 g; or
- 3.3 Cyclophosphamide and methotrexate are contraindicated; or
- 3.4 Patient is a female of child-bearing potential; or
- 3.5 Patient has a previous history of haemorrhagic cystitis, urological malignancy or haematological malignancy.

Note: Indications marked with \* are unapproved indications.

Renewal — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis\*; and
- 2 Patient has previously responded to treatment with rituximab but is now experiencing an acute flare of vasculitis; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m<sup>2</sup> of body-surface area per week for a total of 4 weeks.

Note: Indications marked with \* are unapproved indications.

**Initial application — (Antibody-mediated organ transplant rejection)** from any relevant practitioner. Approvals valid without further renewal unless notified where patient has been diagnosed with antibody-mediated organ transplant rejection\*.

Note: Indications marked with  $^{\star}$  are unapproved indications.

Initial application — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C chronic lymphocytic leukaemia (CLL) requiring treatment; and
- 2 Any of the following:
  - 2.1 The patient is rituximab treatment naive; or
  - 2.2 Either:
    - 2.2.1 The patient is chemotherapy treatment naive; or
    - 2.2.2 Both:
      - 2.2.2.1 The patient's disease has relapsed following no more than three prior lines of chemotherapy treatment: and
      - 2.2.2.2 The patient has had a treatment-free interval of 12 months or more if previously treated with fludarabine and cyclophosphamide chemotherapy; or
  - 2.3 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax; and
- 3 The patient has good performance status; and
- 4 Either:
  - 4.1 The patient does not have chromosome 17p deletion CLL; or
  - 4.2 Rituximab treatment is to be used in combination with funded venetoclax for relapsed/refractory chronic lymphocytic leukaemia; and
- 5 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles; and
- 6 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration), bendamustine or venetoclax.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with rituximab is expected to improve symptoms and improve ECOG score to < 2.

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

**Renewal — (Chronic lymphocytic leukaemia)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax; or
  - 1.2 All of the following:
    - 1.2.1 The patient's disease has relapsed following no more than one prior line of treatment with rituximab for CLL;
    - 1.2.2 The patient has had an interval of 36 months or more since commencement of initial rituximab treatment; and
    - 1.2.3 The patient does not have chromosome 17p deletion CLL; and
    - 1.2.4 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration) or bendamustine; and
- 2 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initial application — (Neuromyelitis Optica Spectrum Disorder(NMOSD)) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks; and
- 2 Either:
  - 2.1 The patient has experienced a severe episode or attack of NMOSD (rapidly progressing symptoms and clinical investigations supportive of a severe attack of NMOSD); or
  - 2.2 All of the following:
    - 2.2.1 The patient has experienced a breakthrough attack of NMOSD; and
    - 2.2.2 The patient is receiving treatment with mycophenolate; and
    - 2.2.3 The patients is receiving treatment with corticosteroids.

Renewal — (Neuromyelitis Optica Spectrum Disorder) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks; and
- 2 The patients has responded to the most recent course of rituximab; and
- 3 The patient has not received rituximab in the previous 6 months.

Initial application — (Post-transplant) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has B-cell post-transplant lymphoproliferative disorder\*; and
- 2 To be used for a maximum of 8 treatment cycles.

Note: Indications marked with \* are unapproved indications.

Renewal — (Post-transplant) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

Subsidy	Fu	,	
(Manufacturer's Price)	Subsidis	ed Generic	
\$	Per	<ul> <li>Manufacturer</li> </ul>	

continued...

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has B-cell post-transplant lymphoproliferative disorder\*; and
- 3 To be used for no more than 6 treatment cycles.

Note: Indications marked with \* are unapproved indications.

Initial application — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
- 2 Fither:
  - 2.1 Treatment with corticosteroids and at least one other immunosuppressant for at least a period of 12 months has been ineffective; or
  - 2.2 Both:
    - 2.2.1 Treatment with at least one other immunosuppressant for a period of at least 12 months; and
    - 2.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

Renewal — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Either:
  - 3.1 The patient has relapsed despite treatment with corticosteroids and at least one other immunosuppressant for a period of at least 12 months; or
  - 3.2 Both:
    - 3.2.1 The patient's myasthenia gravis has relapsed despite treatment with at least one immunosuppressant for a period of at least 12 months; and
    - 3.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

Initial application — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is a child with SDNS\* or FRNS\*; and
- 2 Treatment with steroids for at least a period of 3 months has been ineffective or associated with evidence of steroid toxicity; and
- 3 Treatment with ciclosporin for at least a period of 3 months has been ineffective and/or discontinued due to unacceptable side effects; and
- 4 Treatment with mycophenolate for at least a period of 3 months with no reduction in disease relapses; and
- 5 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with \* are unapproved indications.

Renewal — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1 Patient who was previously treated with rituximab for nephrotic syndrome\*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m<sup>2</sup> of body surface area per week for a total of 4 weeks.

Note: Indications marked with \* are unapproved indications.

Initial application — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Patient is a child with SRNS\* where treatment with steroids and ciclosporin for at least 3 months have been ineffective: and
- 2 Treatment with tacrolimus for at least 3 months has been ineffective; and
- 3 Genetic causes of nephrotic syndrome have been excluded; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m<sup>2</sup> of body surface area per week for a total of 4 weeks.

Note: Indications marked with \* are unapproved indications.

Renewal — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome\*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m<sup>2</sup> of body surface area per week for a total of 4 weeks.

Note: Indications marked with \* are unapproved indications.

Initial application — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Fither:

- 1 All of the following:
  - 1.1 The patient has treatment naive aggressive CD20 positive NHL; and
  - 1.2 To be used with a multi-agent chemotherapy regimen given with curative intent; and
  - 1.3 To be used for a maximum of 8 treatment cycles; or
- 2 Both:
  - 2.1 The patient has aggressive CD20 positive NHL with relapsed disease following prior chemotherapy; and
  - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Renewal — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has relapsed refractory/aggressive CD20 positive NHL; and
- 3 To be used with a multi-agent chemotherapy regimen given with curative intent; and
- 4 To be used for a maximum of 4 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Initial application — (haemophilia with inhibitors) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria: Any of the following:

Subsidy	Fu	,	
(Manufacturer's Price)	Subsidis	ed Generic	
\$	Per	<ul> <li>Manufacturer</li> </ul>	

continued...

- 1 Patient has mild congenital haemophilia complicated by inhibitors; or
- 2 Patient has severe congenital haemophilia complicated by inhibitors and has failed immune tolerance therapy; or
- 3 Patient has acquired haemophilia.

Renewal — (haemophilia with inhibitors) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for haemophilia with inhibitors; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment.

Initial application — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Fither:
  - 1.1 Patient has immune thrombocytopenic purpura\* with a platelet count of less than or equal to 20,000 platelets per
  - 1.2 Patient has immune thrombocytopenic purpura\* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding; and
- 2 Any of the following:
  - 2.1 Treatment with steroids and splenectomy have been ineffective; or
  - 2.2 Treatment with steroids has been ineffective and splenectomy is an absolute contraindication; or
  - 2.3 Other treatments including steroids have been ineffective and patient is being prepared for elective surgery (e.g. splenectomy); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with \* are unapproved indications.

Renewal — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m<sup>2</sup> weekly for 4 weeks) is now planned; or
- 2 All of the following:
  - 2.1 Patient was previously treated with rituximab for immune thrombocytopenic purpura\*; and
  - 2.2 An initial response lasting at least 12 months was demonstrated; and
  - 2.3 Patient now requires repeat treatment.

Note: Indications marked with \* are unapproved indications.

Initial application — (indolent, low-grade lymphomas or hairy cell leukaemia\*) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

Fither:

- 1 Both:
  - 1.1 The patient has indolent low grade NHL or hairy cell leukaemia\* with relapsed disease following prior chemotherapy; and
  - 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
  - 2.1 The patient has indolent, low grade lymphoma or hairy cell leukaemia\* requiring first-line systemic chemotherapy;
  - 2.2 To be used for a maximum of 6 treatment cycles.

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. \*Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Renewal — (indolent, low-grade lymphomas or hairy cell leukaemia\*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has indolent, low-grade NHL or hairy cell leukaemia\* with relapsed disease following prior chemotherapy; and
- 3 To be used for no more than 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. \*Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Initial application — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient has autoimmune pure red cell aplasia\* associated with a demonstrable B-cell lymphoproliferative disorder.

Note: Indications marked with \* are unapproved indications.

Renewal — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient was previously treated with rituximab for pure red cell aplasia\* associated with a demonstrable B-cell lymphoproliferative disorder and demonstrated an initial response lasting at least 12 months.

Note: Indications marked with \* are unapproved indications.

Initial application — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Patient has cold haemagglutinin disease\*; and
- 2 Patient has severe disease which is characterized by symptomatic anaemia, transfusion dependence or disabling circulatory symptoms; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of

Note: Indications marked with \* are unapproved indications.

Renewal — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m<sup>2</sup> weekly for 4 weeks) is now planned; or
- 2 All of the following:
  - 2.1 Patient was previously treated with rituximab for severe cold haemagglutinin disease\*; and
  - 2.2 An initial response lasting at least 12 months was demonstrated; and
  - 2.3 Patient now requires repeat treatment.

Note: Indications marked with \* are unapproved indications.

Initial application — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Both:

- 1 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks: and
- 2 Fither:
  - 2.1 Patient has thrombotic thrombocytopenic purpura\* and has experienced progression of clinical symptoms or persistent thrombocytopenia despite plasma exchange; or
  - 2.2 Patient has acute idiopathic thrombotic thrombocytopenic purpura\* with neurological or cardiovascular pathology.

Note: Indications marked with \* are unapproved indications.

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

**Renewal — (thrombotic thrombocytopenic purpura (TTP))** only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for thrombotic thrombocytopenic purpura\*; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with \* are unapproved indications.

Initial application — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe, immediately life- or organ-threatening SLE\*; and
- 2 The disease has proved refractory to treatment with steroids at a dose of at least 1 mg/kg; and
- 3 The disease has relapsed following prior treatment for at least 6 months with maximal tolerated doses of azathioprine, mycophenolate mofetil and high dose cyclophosphamide, or cyclophosphamide is contraindicated; and
- 4 Maximum of four 1000 mg infusions of rituximab.

Note: Indications marked with \* are unapproved indications.

Renewal — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient's SLE\* achieved at least a partial response to the previous round of prior rituximab treatment; and
- 2 The disease has subsequently relapsed; and
- 3 Maximum of two 1000 mg infusions of rituximab.

Note: Indications marked with \* are unapproved indications.

Initial application — (warm autoimmune haemolytic anaemia (warm AlHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has warm autoimmune haemolytic anaemia\*: and
- 2 One of the following treatments has been ineffective: steroids (including if patient requires ongoing steroids at doses equivalent to > 5 mg prednisone daily), cytotoxic agents (e.g. cyclophosphamide monotherapy or in combination), intravenous immunoglobulin; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with \* are unapproved indications.

Renewal — (warm autoimmune haemolytic anaemia (warm AIHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Fither:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
  - 2.1 Patient was previously treated with rituximab for warm autoimmune haemolytic anaemia\*; and
  - 2.2 An initial response lasting at least 12 months was demonstrated; and
  - 2.3 Patient now requires repeat treatment.

Note: Indications marked with \* are unapproved indications.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

**Initial application — (severe antisynthetase syndrome)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed antisynthetase syndrome; and
- 2 Patient has severe, immediately life or organ threatening disease, including interstitial lung disease; and
- 3 Either
  - 3.1 Treatment with at least 3 immunosuppressants (oral steroids, cyclophosphamide, methotrexate, mycophenolate, ciclosporin, azathioprine) has not be effective at controlling active disease; or
  - 3.2 Rapid treatment is required due to life threatening complications; and
- 4 Maximum of four 1,000mg infusions of rituximab.

**Renewal — (severe antisynthetase syndrome)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in inflammatory markers, muscle strength and pulmonary function; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 Maximum of two cycles of 2 x 1,000mg infusions of rituximab given two weeks apart.

Initial application — (graft versus host disease) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has refractory graft versus host disease following transplant; and
- 2 Treatment with at least 3 immunosuppressants (oral steroids, ciclosporin, tacrolimus, mycophenolate, sirolimus) has not be effective at controlling active disease; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks

Initial application — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has severe chronic inflammatory demyelinating polyneuropathy (CIPD); and
- 2 Either:
  - 2.1 Both:
    - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
    - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
  - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function compared to baseline; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

Initial application — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe anti-NMDA receptor autoimmune encephalitis: and
- 2 Fither
  - 2.1 Both:
    - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
    - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
  - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 The patient has experienced a relapse and now requires further treatment; and
- 4 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Initial application — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has CD20+ low grade or follicular B-cell NHL with relapsed disease following prior chemotherapy; and
  - 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
  - 2.1 The patient has CD20+ low grade or follicular B-cell NHL requiring first-line systemic chemotherapy; and
  - 2.2 To be used for a maximum of 6 treatment cycles.

Renewal — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 Rituximab is to be used for maintenance in CD20+ low grade or follicular B-cell NHL following induction with first-line systemic chemotherapy; and
- 2 Patient is intended to receive rituximab maintenance therapy for 2 years at a dose of 375 mg/m2 every 8 weeks (maximum of 12 cycles).

Initial application — (Membranous nephropathy) only from a nephrologist or any relevant practitioner on the recommendation of a nephrologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Fither:
  - 1.1 Patient has biopsy-proven primary/idiopathic membranous nephropathy\*; or
  - 1.2 Patient has PLA2 antibodies with no evidence of secondary cause, and an eGFR of > 60ml/min/1.73m2; and
- 2 Patient remains at high risk of progression to end-stage kidney disease despite more than 3 months of treatment with conservative measures (see Note): and

Subsidy (Manufacturer's Pr	ico)	Fully Subsidised	Brand or Generic	
(Mahuactuler S FI			Generic	
\$	Pe	r 🗸	Manufacturer	

continued...

3 The total rituximab dose would not exceed the equivalent of 375mg/m2 of body surface area per week for a total of 4 weeks

**Renewal — (Membranous nephropathy)** only from a nephrologist or any relevant practitioner on the recommendation of a nephrologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for membranous nephropathy\*; and
- 2 Either:
  - 2.1 Treatment with rituximab was previously successful, but the condition has relapsed, and the patient now requires repeat treatment; or
  - 2.2 Patient achieved partial response to treatment and requires repeat treatment (see Note); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

#### Notes:

- a) Indications marked with \* are unapproved indications.
- b) High risk of progression to end-stage kidney disease defined as > 5g/day proteinuria.
- c) Conservative measures include renin-angiotensin system blockade, blood-pressure management, dietary sodium and protein restriction, treatment of dyslipidaemia, and anticoagulation agents unless contraindicated or the patient has experienced intolerable side effects.
- d) Partial response defined as a reduction of proteinuria of at least 50% from baseline, and between 0.3 grams and 3.5 grams per 24 hours.

Initial application — (B-cell acute lymphoblastic leukaemia/lymphoma\*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient has newly diagnosed B-cell acute lymphoblastic leukaemia/lymphoma\*; and
- 2 Treatment must be in combination with an intensive chemotherapy protocol with curative intent; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m² per dose for a maximum of 18 doses.

Note: Indications marked with \* are unapproved indications.

**Initial application** — (desensisation prior to transplant) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires desensitisation prior to mismatched allogenic stem cell transplant\*; and
- 2 Patient would receive no more than two doses at 375 mg/m2 of body-surface area.

Note: Indications marked with \* are unapproved indications.

Initial application — (pemiphigus\*) only from a dermatologist or relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

#### Either:

- 1 All of the following:
  - 1.1 Patient has severe rapidly progressive pemphigus; and
  - 1.2 Is used in combination with systemic corticosteroids (20 mg/day); and
  - 1.3 Any of the following:
    - 1.3.1 Skin involvement is at least 5% body surface area; or
    - 1.3.2 Significant mucosal involvement (10 or more mucosal erosions) or diffuse gingivitis or confluent large erosions; or
    - 1.3.3 Involvement of two or more mucosal sites; or
- 2 Both:
  - 2.1 Patient has pemphigus; and

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

2.2 Patient has not experienced adequate clinical benefit from systemic corticosteroids (20 mg/day) in combination with a steroid sparing agent, unless contraindicated.

Note: Indications marked with \* are unapproved indications.

Renewal — (pemiphigus\*) only from a dermatologist or relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has experienced adequate clinical benefit from rituximab treatment, with improvement in symptoms and healing of skin ulceration and reduction in corticosteroid requirement; and
- 2 Patient has not received rituximab in the previous 6 months.

Note: Indications marked with \* are unapproved indications.

Initial application — (immunoglobulin G4-related disease (IgG4-RD\*)) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed diagnosis of IgG4-RD\*; and
- 2 Either
  - 2.1 Treatment with corticosteroids and/or disease modifying anti-rheumatic drugs for at least 3 months has been ineffective in lowering corticosteroid dose below 5 mg per day (prednisone equivalent) without relapse; or
  - 2.2 Treatment with corticosteroids and/or disease modifying anti-rheumatic drugs is contraindicated or associated with evidence of toxicity or intolerance; and
- 3 Total rituximab dose used should not exceed a maximum of two 1000 mg infusions of rituximab given two weeks apart. Note: Indications marked with \* are unapproved indications.

Renewal — (immunoglobulin G4-related disease (IgG4-RD\*)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 Treatment with rituximab for IgG4-RD\* was previously successful and patient's disease has demonstrated sustained response, but the condition has relapsed; or
  - 1.2 Patient is receiving maintenance treatment for IgG4-RD\*; and
- 2 Rituximab re-treatment not to be given within 6 months of previous course of treatment; and
- 3 Maximum of two 1000 mg infusions of rituximab given two weeks apart.

Note: Indications marked with \* are unapproved indications.

SECUKINUMAB – Special Authority see SA2403 below – Retail pharmacy

# ⇒SA2403 Special Authority for Subsidy

Initial application — (severe chronic plaque psoriasis – second-line biologic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab or etanercept, or has trialled infliximab in a Health NZ Hospital, for severe chronic plaque psoriasis; and
- 2 Fither:
  - 2.1 The patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
  - 2.2 The patient has received insufficient benefit from adalimumab, etanercept or infliximab; and
- 3 A Psoriasis Area and Severity Index (PASI) assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and

Subs (Manufactur		
\$	Per	Manufacturer

continued...

4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Initial application — (severe chronic plaque psoriasis – first-line biologic) only from a dermatologist or any relevant practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
  - 1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; or
  - 1.3 Patient has severe chronic localised genital or flexural plaque psoriasis where the plaques or lesions have been present for at least 6 months from the time of initial diagnosis, and with a Dermatology Life Quality Index (DLQI) score greater than 10; and
- 2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
- 3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Note: A treatment course is defined as a minimum of 12 weeks of treatment. "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand. foot, genital or flexural areas, at least 2 of the 3 PASI symptom sub scores for erythema, thickness and scaling are rated as severe or very severe, and for the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis – first and second-line biologic) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 Either:
    - 1.1.1 Patient's PASI score has reduced by 75% or more (PASI 75) as compared to baseline PASI prior to commencing secukinumab: or
    - 1.1.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing secukinumab; or
  - 1.2 Both:
    - 1.2.1 Patient had severe chronic localised genital or flexural plague psoriasis at the start of treatment; and
    - 1.2.2 Either:
      - 1.2.2.1 The patient has experienced a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; or
      - 1.2.2.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing secukinumab; and
- 2 Secukinumab to be administered at a maximum dose of 300 mg monthly.

Initial application — (ankylosing spondylitis – second-line biologic) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	1	Manufacturer	

continued...

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and
- 2 Either:
  - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
  - 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

Renewal — (ankylosing spondylitis – second-line biologic) only from a rheumatologist or medical practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Following 12 weeks initial treatment of secukinumab treatment, BASDAI has improved by 4 or more points from pre-secukinumab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefitted from treatment and that continued treatment is appropriate; and
- 3 Secukinumab to be administered at doses no greater than 150 mg monthly.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
  - 1.1 Patient has had an initial Special Authority approval for adalimumab, etanercept or infliximab for psoriatic arthritis; and
  - 1.2 Fither:
    - 1.2.1 Patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
    - 1.2.2 Patient has received insufficient benefit from adalimumab, etanercept or infliximab to meet the renewal criteria for adalimumab, etanercept or infliximab for psoriatic arthritis; or
- 2 All of the following:
  - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
  - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
  - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
  - 2.4 Either:
    - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints;
    - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
  - 2.5 Any of the following:
    - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
    - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
    - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior secukinumab treatment in the opinion of the treating physician; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

2 Secukinumab to be administered at doses no greater than 300 mg monthly.

#### SILTUXIMAB - Special Authority see SA1596 below - Retail pharmacy

Note: Siltuximab is to be administered at doses no greater than 11 mg/kg every 3 weeks.

Inj 100 mg vial	770.57	1	Sylvant
Inj 400 mg vial	3,082.33	1	Sylvant

# ⇒SA1596 Special Authority for Subsidy

Initial application only from a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's Disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Siltuximab is to be administered at doses no greater than 11 mg/kg every 3 weeks.

**Renewal** only from a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has sustained improvement in inflammatory markers and functional status.

TIXAGEVIMAB WITH CILGAVIMAB - [Xpharm] - Subsidy by endorsement

- a) No patient co-payment payable
- b) Treatment is funded only if patient meets access criteria for tixagevimab with cilgavimab (as per https://pharmac.govt.nz/Evusheld) and has been endorsed accordingly by the prescriber. The supply of treatment is via Pharmac's approved distribution process. Refer to the Pharmac website for more information about this and stock availability.
- Inj 100 mg per ml, 1.5 ml vial with cilgavimab 100 mg per ml.1.5 ml vial.....

ml,1.5 ml vial	0.00	1	<ul><li>Evusheld</li></ul>
TOCILIZUMAB - PCT only - Special Authority see SA24	104 below		
Inj 20 mg per ml, 4 ml vial	220.00	1	✓ Actemra
Inj 20 mg per ml, 10 ml vial	550.00	1	✓ Actemra
Inj 20 mg per ml, 20 ml vial	1,100.00	1	✓ Actemra
Inj 1 mg for ECP		1 mg	<ul><li>Baxter</li></ul>

#### ⇒SA2404 Special Authority for Subsidy

**Initial application — (cytokine release syndrome)** from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Either:

- 1 Both:
  - 1.1 The patient has developed grade 3 or 4 cytokine release syndrome associated with the administration of blinatumomab for the treatment of acute lymphoblastic leukaemia; and
  - 1.2 Tocilizumab is to be administered at doses no greater than 8 mg/kg IV for a maximum of 3 doses (if less than 30kg, maximum of 12 mg/kg); or
- 2 All of the following:
  - 2.1 The patient is enrolled in the Malaghan Institute of Medical Research ENABLE trial programme; and
  - 2.2 The patient has developed CRS or Immune Effector Cell-Associated Neurotoxicity Syndrome (ICANS) following CAR T-cell therapy for the treatment of relapsed or refractory B-cell non-Hodgkin lymphoma; and
  - 2.3 Tocilizumab is to be administered according to the consensus guidelines for CRS or ICANS for CAR T-cell therapy at doses no greater than 8 mg/kg IV for a maximum of 3 doses.

**Initial application — (previous use)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

Subsidy		Fully	Brand or
(Manufacturer's Price)	9	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 1 Patient was being treated with tocilizumab prior to 1 February 2019; and
- 2 Any of the following:
  - 2.1 rheumatoid arthritis: or
  - 2.2 systemic juvenile idiopathic arthritis; or
  - 2.3 adult-onset Still's disease; or
  - 2.4 polyarticular juvenile idiopathic arthritis; or
  - 2.5 idiopathic multicentric Castleman's disease.

Initial application — (Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Fither
  - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
  - 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
- 3 Fither:
  - 3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or
  - 3.2 Both:
    - 3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a Health NZ Hospital; and
    - 3.2.2 Either:
      - 3.2.2.1 The patient has experienced intolerable side effects from rituximab; or
      - 3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Initial application — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 2 Tocilizumab is to be used as monotherapy; and
- 3 Fither:
  - 3.1 Treatment with methotrexate is contraindicated; or
  - 3.2 Patient has tried and did not tolerate oral and/or parenteral methotrexate; and
- 4 Either:
  - 4.1 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of ciclosporin alone or in combination with another agent; or
  - 4.2 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of leflunomide alone or in combination with another agent; and
- 5 Fither:
  - 5.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 active, swollen, tender joints; or
  - 5.2 Patient has persistent symptoms of poorly controlled and active disease in at least four active joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 6 Either:
  - 6.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	/	Manufacturer

continued...

6.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Initial application — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient diagnosed with systemic juvenile idiopathic arthritis; and
- 2 Patient has tried and not responded to a reasonable trial of all of the following, either alone or in combination: oral or parenteral methotrexate; non-steroidal anti-inflammatory drugs (NSAIDs); and systemic corticosteroids.

Initial application — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
  - 1.1 Fither:
    - 1.1.1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for adult-onset Still's disease (AOSD); or
    - 1.1.2 The patient has been started on tocilizumab for AOSD in a Health NZ Hospital; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
    - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for AOSD: or
- 2 All of the following:
  - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
  - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal antiinflammatory drugs (NSAIDs) and methotrexate; and
  - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Initial application — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for both etanercept and adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
  - 1.2 The patient has experienced intolerable side effects, or has received insufficient benefit from, both etanercept and adalimumab; or
- 2 All of the following:
  - 2.1 Treatment with a tumour necrosis factor alpha inhibitor is contraindicated; and
  - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
  - 2.3 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
  - 2.4 Any of the following:
    - 2.4.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
    - 2.4.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
    - 2.4.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Initial application — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

continued...

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Tocilizumab to be administered at doses no greater than 8 mg/kg IV every 3-4 weeks.

Initial application — (moderate to severe COVID-19) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed (or probable) COVID-19; and
- 2 Oxygen saturation of < 92% on room air, or requiring supplemental oxygen; and
- 3 Patient is receiving adjunct systemic corticosteroids, or systemic corticosteroids are contraindicated; and
- 4 Tocilizumab is to be administered at doses no greater than 8mg/kg IV for a maximum of one dose; and
- 5 Tocilizumab is not to be administered in combination with barcitinib.

Renewal — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Renewal — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Following up to 6 months' initial treatment, the patient has achieved at least an American College of Rheumatology paediatric 30% improvement criteria (ACR Pedi 30) response from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing ACR Pedi 30 response from baseline.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months where the patient has a sustained improvement in inflammatory markers and functional status. Renewal — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
  - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Renewal — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has a sustained improvement in inflammatory markers and functional status.

 TRASTUZUMAB (HERZUMA) – PCT only – Special Authority see SA2293 below

 Inj 150 mg vial
 100.00
 1
 ✓ Herzuma

 Inj 440 mg vial
 293.35
 1
 ✓ Herzuma

 Inj 1 mg for ECP
 0.70
 1 mg
 ✓ Baxter

⇒SA2293 Special Authority for Subsidy

Initial application — (early breast cancer) from any relevant practitioner. Approvals valid for 15 months for applications

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

meeting the following criteria:

Both:

- 1 The patient has early breast cancer expressing HER-2 IHC 3+ or ISH + (including FISH or other current technology); and
- 2 Maximum cumulative dose of 106 mg/kg (12 months' treatment).

Renewal — (early breast cancer\*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 All of the following:
  - 1.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
  - 1.2 The patient received prior adjuvant trastuzumab treatment for early breast cancer; and
  - 1.3 Any of the following:
    - 1.3.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
    - 1.3.2 The patient discontinued lapatinib within 3 months due to intolerable side effects and the cancer did not progress whilst on lapatinib; or
    - 1.3.3 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
  - 1.4 Fither:
    - 1.4.1 Trastuzumab will not be given in combination with pertuzumab; or
    - 1.4.2 All of the following:
      - 1.4.2.1 Trastuzumab to be administered in combination with pertuzumab; and
      - 1.4.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
      - 1.4.2.3 The patient has good performance status (ECOG grade 0-1); and
  - 1.5 Trastuzumab to be discontinued at disease progression; or
- 2 All of the following:
  - 2.1 Patient has previously discontinued treatment with trastuzumab in the metastatic setting for reasons other than severe toxicity or disease progression; and
  - 2.2 Patient has signs of disease progression; and
  - 2.3 Disease has not progressed during previous treatment with trastuzumab.

Note: \* For patients with relapsed HER-2 positive disease who have previously received adjuvant trastuzumab for early breast cancer

**Initial application — (metastatic breast cancer)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Either:
  - 2.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
  - 2.2 The patient discontinued lapatinib within 3 months due to intolerable side effects and the cancer did not progress whilst on lapatinib; and
- 3 Either:
  - 3.1 Trastuzumab will not be given in combination with pertuzumab; or
  - 3.2 All of the following:
    - 3.2.1 Trastuzumab to be administered in combination with pertuzumab; and
    - 3.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast

Subsidy	S	Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

cancer: and

- 3.2.3 The patient has good performance status (ECOG grade 0-1); and
- 4 Trastuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

#### Either:

- 1 All of the following:
  - 1.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
  - 1.2 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
  - 1.3 Trastuzumab to be discontinued at disease progression; or
- 2 All of the following:
  - 2.1 Patient has previously discontinued treatment with trastuzumab for reasons other than severe toxicity or disease progression; and
  - 2.2 Patient has signs of disease progression; and
  - 2.3 Disease has not progressed during previous treatment with trastuzumab.

Initial application — (gastric, gastro-oesophageal junction and oesophageal cancer) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has locally advanced or metastatic gastric, gastro-oesophageal junction or oesophageal cancer expressing HER-2 IHC 2+ FISH+ or IHC3+ (or other current technology); and
- 2 Patient has an ECOG score of 0-2.

Renewal — (gastric, gastro-oesophageal junction and oesophageal cancer) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 2 Trastuzumab to be discontinued at disease progression.

# TRASTUZUMAB EMTANSINE - PCT only - Specialist - Special Authority see SA2144 below

Inj 100 mg vial	2,320.00	1	Kadcyla
Inj 160 mg vial	3,712.00	1	✓ Kadcyla
Inj 1 mg for ECP	24.52	1 mg	✓ Baxter

#### ⇒SA2144 Special Authority for Subsidy

Initial application — (early breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has early breast cancer expressing HER2 IHC3+ or ISH+; and
- 2 Documentation of pathological invasive residual disease in the breast and/or auxiliary lymph nodes following completion of surgery; and
- 3 Patient has completed systemic neoadjuvant therapy with trastuzumab and chemotherapy prior to surgery; and
- 4 Disease has not progressed during neoadjuvant therapy; and
- 5 Patient has left ventricular ejection fraction of 45% or greater; and
- 6 Adjuvant treatment with trastuzumab emtansine to be commenced within 12 weeks of surgery; and
- 7 Trastuzumab emtansine to be discontinued at disease progression; and
- 8 Total adjuvant treatment duration must not exceed 42 weeks (14 cycles).

**Initial application** — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer \$ Per

continued...

All of the following:

- 1 Patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Patient has previously received trastuzumab and chemotherapy, separately or in combination; and
- 3 Fither:
  - 3.1 The patient has received prior therapy for metastatic disease\*; or
  - 3.2 The patient developed disease recurrence during, or within six months of completing adjuvant therapy\*; and
- 4 Patient has a good performance status (ECOG 0-1); and
- 5 Fither:
  - 5.1 Patient does not have symptomatic brain metastases; or
  - 5.2 Patient has brain metastases and has received prior local CNS therapy; and
- 6 Patient has not received prior funded trastuzumab emtansine treatment; and
- 7 Treatment to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The cancer has not progressed at any time point during the previous approval period whilst on trastuzumab emtansine:
- 2 Treatment to be discontinued at disease progression.

Note: Prior or adjuvant therapy includes anthracycline, other chemotherapy, biological drugs, or endocrine therapy.

USTEKINUMAB - Special Authority see SA2182 below - Retail pharmacy

Inj 90 mg per ml, 1 ml pre-filled syringe......4,162.00 ✓ Stelara

#### ⇒SA2182 Special Authority for Subsidy

Initial application — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

#### Either:

- 1 Patient is currently on treatment with ustekinumab commenced prior to 1 February 2023 and met all remaining criteria (criterion 2) below at the time of commencing treatment; or
- 2 Both:
  - 2.1 Patient has active Crohn's disease; and
  - 2.2 Either:
    - 2.2.1 Patient has had an initial approval for prior biologic therapy for Crohn's disease and has experienced intolerable side effects or insufficient benefit to meet renewal criteria; or
    - 2.2.2 Both:
      - 2.2.2.1 Patient meets the initiation criteria for prior biologic therapies for Crohn's disease; and
      - 2.2.2.2 Other biologics for Crohn's disease are contraindicated.

Renewal — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 CDAI score has reduced by 100 points, or HBI score has reduced by 3 points, from when the patient was initiated on biologic therapy; or
  - 1.2 CDAI score is 150 or less, or HBI is 4 or less; or
  - 1.3 The patient has experienced an adequate response to treatment, but CDAI score and/or HBI score cannot be assessed: and
  - 2 Ustekinumab to be administered at a dose no greater than 90 mg every 8 weeks.

Initial application — (Crohn's disease - children\*) from any relevant practitioner. Approvals valid for 6 months for applications

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

meeting the following criteria:

#### Either:

- 1 Patient is currently on treatment with ustekinumab commenced prior to 1 February 2023 and met all remaining criteria (criterion 2) below at the time of commencing treatment; or
- 2 Both:
  - 2.1 Patient has active Crohn's disease; and
  - 2.2 Fither:
    - 2.2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria; or
    - 2.2.2 Both:
      - 2.2.2.1 Patient meets the initiation criteria for prior biologic therapies for Crohn's disease; and
      - 2.2.2.2 Other biologics for Crohn's disease are contraindicated.

Note: Indication marked with \* is an unapproved indication.

Renewal — (Crohn's disease - children\*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 PCDAI score has reduced by 10 points from when the patient was initiated on biologic therapy; or
  - 1.2 PCDAI score is 15 or less; or
  - 1.3 The patient has experienced an adequate response to treatment, but CDAI score cannot be assessed; and
  - 2 Ustekinumab to administered at a dose no greater than 90 mg every 8 weeks.

Note: Indication marked with \* is an unapproved indication.

Initial application — (ulcerative colitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

# Either:

- 1 Patient is currently on treatment with ustekinumab commenced prior to 1 February 2023 and met all remaining criteria (criterion 2) below at the time of commencing treatment; or
- 2 Both:
  - 2.1 Patient has active ulcerative colitis; and
  - 2.2 Either:
    - 2.2.1 Patient has had an initial approval for prior biologic therapy for ulcerative colitis and has experienced intolerable side effects or insufficient benefit to meet renewal criteria; or
    - 2.2.2 Both:
      - 2.2.2.1 Patient meets the initiation criteria for prior biologic therapies for ulcerative colitis; and
      - 2.2.2.2 Other biologics for ulcerative colitis are contraindicated.

**Renewal — (ulcerative colitis)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 The SCCAI score has reduced by 2 points or more from the SCCAI score since initiation on biologic therapy; or
  - 1.2 PUCAI score has reduced by 10 points or more from the PUCAI score since initiation on biologic therapy\*; and
- 2 Ustekinumab will be used at a dose no greater than 90 mg intravenously every 8 weeks.

Note: Criterion marked with \* is for an unapproved indication.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

# ⇒SA2183 Special Authority for Subsidy

Initial application — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active Crohn's disease; and
- 2 Any of the following:
  - 2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria (unless contraindicated); or
  - 2.2 Patient has a CDAI score of greater than or equal to 300, or HBI score of greater than or equal to 10; or
  - 2.3 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
  - 2.4 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection:
  - 2.5 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Any of the following:
  - 3.1 Patient has tried but experienced an inadequate response to (including lack of initial response and/or loss of initial response) from prior therapy with immunomodulators and corticosteroids; or
  - 3.2 Patient has experienced intolerable side effects from immunomodulators and corticosteroids; or
  - 3.3 Immunomodulators and corticosteroids are contraindicated.

Renewal — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 CDAI score has reduced by 100 points, or HBI score has reduced by 3 points, from when the patient was initiated on biologic therapy; or
  - 1.2 CDAI score is 150 or less, or HBI is 4 or less; or
  - 1.3 The patient has experienced an adequate response to treatment, but CDAI score and/or HBI score cannot be assessed; and
- 2 Vedolizumab to administered at a dose no greater than 300 mg every 8 weeks.

Initial application — (Crohn's disease - children\*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has active Crohn's disease; and
- 2 Any of the following:
  - 2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria (unless contraindicated); or
  - 2.2 Patient has a Paediatric Crohn's Disease Activity Index (PCDAI) score of greater than or equal to 30; or
  - 2.3 Patient has extensive small intestine disease; and
- 3 Any of the following:
  - 3.1 Patient has tried but experienced an inadequate response to (including lack of initial response and/or loss of initial response) from prior therapy with immunomodulators and corticosteroids; or
  - 3.2 Patient has experienced intolerable side effects from immunomodulators and corticosteroids; or
  - 3.3 Immunomodulators and corticosteroids are contraindicated.

Note: Indication marked with \* is an unapproved indication.

Renewal — (Crohn's disease - children\*) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

1 Any of the following:

Subsidy (Manufacturer's Price)	Sul	Fully	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

- 1.1 PCDAI score has reduced by 10 points from when the patient was initiated on biologic therapy; or
- 1.2 PCDAI score is 15 or less; or
- 1.3 The patient has experienced an adequate response to treatment, but CDAI score cannot be assessed; and
- 2 Vedolizumab to administered at a dose no greater than 300mg every 8 weeks.

Note: Indication marked with \* is an unapproved indication.

**Initial application — (ulcerative colitis)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active ulcerative colitis: and
- 2 Any of the following:
  - 2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria (unless contraindicated); or
  - 2.2 Patient has a SCCAI score is greater than or equal to 4; or
  - 2.3 Patient's PUCAI score is greater than or equal to 20\*; and
- 3 Any of the following:
  - 3.1 Patient has tried but experienced an inadequate response to (including lack of initial response and/or loss of initial response) from prior therapy with immunomodulators and corticosteroids; or
  - 3.2 Patient has experienced intolerable side effects from immunomodulators and corticosteroids; or
  - 3.3 Immunomodulators and corticosteroids are contraindicated.

Note: Indication marked with \* is an unapproved indication.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 The SCCAI score has reduced by 2 points or more from the SCCAI score since initiation on biologic therapy; or
  - 1.2 The PUCAI score has reduced by 10 points or more from the PUCAI score since initiation on biologic therapy \*; and
- 2 Vedolizumab will be used at a dose no greater than 300 mg intravenously every 8 weeks.

Note: Indication marked with \* is an unapproved indication.

# Programmed Cell Death-1 (PD-1) Inhibitors

		Specialist – Special Authority see SA2264 below	ATEZOLIZUMAB – PCT only – Specialist -
✓ Tecentriq	1	9,503.00	Inj 60 mg per ml, 20 ml vial
✓ Baxter	1 mg	8.08	Inj 1 mg for ECP

#### ⇒SA2264 Special Authority for Subsidy

Initial application — (non-small cell lung cancer second line monotherapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced or metastatic non-small cell lung cancer; and
- 2 Patient has not received prior funded treatment with an immune checkpoint inhibitor for NSCLC; and
- 3 For patients with non-squamous histology there is documentation confirming that the disease does not express activating mutations of EGFR or ALK tyrosine kinase unless not possible to ascertain; and
- 4 Patient has an ECOG 0-2; and
- 5 Patient has documented disease progression following treatment with at least two cycles of platinum-based chemotherapy; and
- 6 Atezolizumab is to be used as monotherapy at a dose of 1200 mg every three weeks (or equivalent) for a maximum of

Subsidy (Manufacturer's Price)	F Subsidi	ully	Brand or Generic
 \$	Per	<b>√</b>	Manufacturer

continued...

16 weeks: and

7 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Renewal — (non-small cell lung cancer second line monotherapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Any of the following:
  - 1.1 Patient's disease has had a complete response to treatment; or
  - 1.2 Patient's disease has had a partial response to treatment; or
  - 1.3 Patient has stable disease; and
- 2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period; and
- 3 No evidence of disease progression; and
- 4 The treatment remains clinically appropriate and patient is benefitting from treatment; and
- 5 Atezolizumab to be used at a maximum dose of 1200 mg every three weeks (or equivalent); and
- 6 Treatment with atezolizumab to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

URVALUMAB – PCT only – Specialist – Special Auth	nority see SA2164 below		
Inj 50 mg per ml, 10 ml vial	4,700.00	1	Imfinzi
Inj 50 mg per ml, 2.4 ml vial	1,128.00	1	🗸 lmfinzi
Inj 1 mg for ECP	9.59	1 mg	✓ Baxter

# ⇒SA2164 Special Authority for Subsidy

Initial application — (Non-small cell lung cancer) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 Patient has histologically or cytologically documented stage III, locally advanced, unresectable non-small cell lung cancer (NSCLC); and
- 2 Patient has received two or more cycles of platinum-based chemotherapy concurrently with definitive radiation therapy; and
- 3 Patient has no disease progression following the second or subsequent cycle of platinum-based chemotherapy with definitive radiation therapy treatment; and
- 4 Patient has a ECOG performance status of 0 or 1; and
- 5 Patient has completed last radiation dose within 8 weeks of starting treatment with durvalumab; and
- 6 Patient must not have received prior PD-1 or PD-L1 inhibitor therapy for this condition; and
- 7 Either:
  - 7.1 Durvalumab is to be used at a maximum dose of no greater than 10 mg/kg every 2 weeks; or
  - 7.2 Durvalumab is to be used at a flat dose of 1500 mg every 4 weeks; and
- 8 Treatment with durvalumab to cease upon signs of disease progression.

Renewal — (Non-small cell lung cancer) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The treatment remains clinically appropriate and the patient is benefitting from treatment; and
- 2 Fither:
  - 2.1 Durvalumab is to be used at a maximum dose of no greater than 10 mg/kg every 2 weeks; or
  - 2.2 Durvalumab is to be used at a flat dose of 1500 mg every 4 weeks; and
- 3 Treatment with durvalumab to cease upon signs of disease progression; and
- 4 Total continuous treatment duration must not exceed 12 months.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
NIVOLUMAB - PCT only - Specialist - Special Authority see SA	2405 below			
Inj 10 mg per ml, 4 ml vial	1,051.98	1	✓	Opdivo
Inj 10 mg per ml, 10 ml vial	2,629.96	1	✓	Opdivo
Inj 1 mg for ECP		1 mg	<b>/</b>	Baxter

# **⇒SA2405** Special Authority for Subsidy

Initial application only from a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Baseline measurement of overall tumour burden is documented clinically and radiologically; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Either:
  - 4.1 Patient has not received funded pembrolizumab; or
  - 4.2 Both:
    - 4.2.1 Patient has received an initial Special Authority approval for pembrolizumab and has discontinued pembrolizumab within 12 weeks of starting treatment due to intolerance; and
    - 4.2.2 The cancer did not progress while the patient was on pembrolizumab; and
- 5 Documentation confirming that the patient has been informed and acknowledges that funded treatment with nivolumab will not be continued if their disease progresses.

Renewal — (less than 24 months on treatment) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 All of the following:
  - 1.1 Any of the following:
    - 1.1.1 Patient's disease has had a complete response to treatment; or
    - 1.1.2 Patient's disease has had a partial response to treatment; or
    - 1.1.3 Patient has stable disease; and
  - 1.2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period: and
  - 1.3 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
- 2 All of the following:
  - 2.1 Patient has previously discontinued treatment with nivolumab for reasons other than severe toxicity or disease progression; and
  - 2.2 Patient has signs of disease progression; and
  - 2.3 Disease has not progressed during previous treatment with nivolumab.

Renewal — (more than 24 months on treatment) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: Both:

- 1 Patient has been on treatment for more than 24 months; and
- 2 Either:
  - 2.1 All of the following:
    - 2.1.1 Any of the following:
      - 2.1.1.1 Patient's disease has had a complete response to treatment; or
      - 2.1.1.2 Patient's disease has had a partial response to treatment; or
      - 2.1.1.3 Patient has stable disease; and
    - 2.1.2 Response to treatment in target lesions has been determined by comparable radiologic or clinical assessment following the most recent treatment period; and
    - 2.1.3 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
  - 2.2 All of the following:

Subsidy	Fu	lly Brand or	
(Manufacturer's Price)	Subsidis	ed Generic	
\$	Per	<ul> <li>Manufacturer</li> </ul>	

continued...

- 2.2.1 Patient has previously discontinued treatment with nivolumab for reasons other than severe toxicity or disease progression; and
- 2.2.2 Patient has signs of disease progression; and
- 2.2.3 Disease has not progressed during previous treatment with nivolumab.

Initial application — (Renal cell carcinoma) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 Patient is currently on treatment with nivolumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
  - 2.1 Patient has metastatic renal-cell carcinoma; and
  - 2.2 The disease is of predominant clear-cell histology; and
  - 2.3 Patient has an ECOG performance score of 0-2; and
  - 2.4 Patient has documented disease progression following one or two previous regimens of antiangiogenic therapy; and
  - 2.5 Nivolumab is to be used as monotherapy at a maximum dose of 240 mg every 2 weeks (or equivalent) and discontinued at disease progression.

Renewal — (Renal cell carcinoma) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 Patient's disease has had a complete response to treatment; or
  - 1.2 Patient's disease has had a partial response to treatment; or
  - 1.3 Patient has stable disease; and
- 2 No evidence of disease progression; and
- 3 Nivolumab is to be used as monotherapy at a maximum dose of 240 mg every 2 weeks (or equivalent) and discontinued at disease progression.

#### 

#### ⇒SA2386 Special Authority for Subsidy

Initial application — (unresectable or metastatic melanoma) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Baseline measurement of overall tumour burden is documented clinically and radiologically; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Fither:
  - 4.1 Patient has not received funded nivolumab; or
  - 4.2 Both:
    - 4.2.1 Patient has received an initial Special Authority approval for nivolumab and has discontinued nivolumab within 12 weeks of starting treatment due to intolerance; and
    - 4.2.2 The cancer did not progress while the patient was on nivolumab; and
- 5 Documentation confirming that the patient has been informed and acknowledges that funded treatment with pembrolizumab will not be continued if their disease progresses.

Renewal — (unresectable or metastatic melanoma, less than 24 months on treatment) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the

Subsidy		Fully	Brand or
(Manufacturer's Price)	9	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

following criteria:

Either:

- 1 All of the following:
  - 1.1 Any of the following:
    - 1.1.1 Patient's disease has had a complete response to treatment; or
    - 1.1.2 Patient's disease has had a partial response to treatment; or
    - 1.1.3 Patient has stable disease; and
  - 1.2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period; and
  - 1.3 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
- 2 All of the following:
  - 2.1 Patient has previously discontinued treatment with pembrolizumab for reasons other than severe toxicity or disease progression; and
  - 2.2 Patient has signs of disease progression; and
  - 2.3 Disease has not progressed during previous treatment with pembrolizumab.

Renewal — (unresectable or metastatic melanoma, more than 24 months on treatment) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

#### Both:

- 1 Patient has been on treatment for more than 24 months; and
- 2 Fither:
  - 2.1 All of the following:
    - 2.1.1 Any of the following:
      - 2.1.1.1 Patient's disease has had a complete response to treatment; or
      - 2.1.1.2 Patient's disease has had a partial response to treatment; or
      - 2.1.1.3 Patient has stable disease; and
    - 2.1.2 Response to treatment in target lesions has been determined by comparable radiologic or clinical assessment following the most recent treatment period; and
    - 2.1.3 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
  - 2.2 All of the following:
    - 2.2.1 Patient has previously discontinued treatment with pembrolizumab for reasons other than severe toxicity or disease progression; and
    - 2.2.2 Patient has signs of disease progression; and
    - 2.2.3 Disease has not progressed during previous treatment with pembrolizumab.

Initial application — (non-small cell lung cancer first-line monotherapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

# All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-small cell lung cancer; and
- 2 Patient has not had chemotherapy for their disease in the palliative setting; and
- 3 Patient has not received prior funded treatment with an immune checkpoint inhibitor for NSCLC; and
- 4 For patients with non-squamous histology there is documentation confirming that the disease does not express activating mutations of EGFR or ALK tyrosine kinase unless not possible to ascertain; and
- 5 Pembrolizumab to be used as monotherapy: and
- 6 Either:
  - 6.1 There is documentation confirming the disease expresses PD-L1 at a level greater than or equal to 50% as determined by a validated test unless not possible to ascertain; or

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer
--

continued...

- 6.2 Both:
  - 6.2.1 There is documentation confirming the disease expresses PD-L1 at a level greater than or equal to 1% as determined by a validated test unless not possible to ascertain; and
  - 6.2.2 Chemotherapy is determined to be not in the best interest of the patient based on clinician assessment; and
- 7 Patient has an ECOG 0-2: and
- 8 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks; and
- 9 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Renewal — (non-small cell lung cancer first line monotherapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Any of the following:
  - 1.1 Patient's disease has had a complete response to treatment; or
  - 1.2 Patient's disease has had a partial response to treatment; or
  - 1.3 Patient has stable disease; and
- 2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period; and
- 3 No evidence of disease progression; and
- 4 The treatment remains clinically appropriate and patient is benefitting from treatment; and
- 5 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 6 Treatment with pembrolizumab to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

**Initial application — (non-small cell lung cancer first-line combination therapy)** only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-small cell lung cancer; and
- 2 The patient has not had chemotherapy for their disease in the palliative setting; and
- 3 Patient has not received prior funded treatment with an immune checkpoint inhibitor for NSCLC; and
- 4 For patients with non-squamous histology there is documentation confirming that the disease does not express activating mutations of EGFR or ALK tyrosine kinase unless not possible to ascertain; and
- 5 Pembrolizumab to be used in combination with platinum-based chemotherapy; and
- 6 Patient has an ECOG 0-2; and
- 7 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks; and
- 8 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Renewal — (non-small cell lung cancer first line combination therapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 Patient's disease has had a complete response to treatment; or
  - 1.2 Patient's disease has had a partial response to treatment: or
  - 1.3 Patient has stable disease; and
- 2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period: and
- 3 No evidence of disease progression; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 4 The treatment remains clinically appropriate and patient is benefitting from treatment; and
- 5 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 6 Treatment with pembrolizumab to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

**Initial application** — (breast cancer, advanced) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: Either:

- 1 Patient is currently on treatment with pembrolizumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
  - 2.1 Fither:
    - 2.1.1 Patient has recurrent or de novo unresectable, inoperable locally advanced triple-negative breast cancer (that does not express ER, PR or HER2 IHC3+ or ISH+ [including FISH or other technology]); or
    - 2.1.2 Patient has recurrent or de novo metastatic triple-negative breast cancer (that does not express ER, PR or HER2 IHC3+ or ISH+ (including FISH or other technology)); and
  - 2.2 Patient is treated with palliative intent: and
  - 2.3 Patient's cancer has confirmed PD-L1 Combined Positive Score (CPS) is greater than or equal to 10; and
  - 2.4 Patient has received no prior systemic therapy in the palliative setting; and
  - 2.5 Patient has an ECOG score of 0-2; and
  - 2.6 Pembrolizumab is to be used in combination with chemotherapy; and
  - 2.7 Baseline measurement of overall tumour burden is documented clinically and radiologically; and
  - 2.8 Pembrolizumab is to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks.

Renewal — (breast cancer, advanced) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 Patient's disease has had a complete response to treatment; or
  - 1.2 Patient's disease has had a partial response to treatment; or
  - 1.3 Patient has stable disease; and
- 2 No evidence of disease progression; and
- 3 Response to treatment in target lesions has been determined by a comparable radiologic assessment following the most recent treatment period; and
- 4 Pembrolizumab is to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 5 Treatment with pembrolizumab is to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initial application — (head and neck squamous cell carcinoma) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria: Fither:

- 1 Patient is currently on treatment with pembrolizumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
  - 2.1 Patient has recurrent or metastatic head and neck squamous cell carcinoma of mucosal origin (excluding nasopharyngeal carcinoma) that is incurable by local therapies; and
  - 2.2 Patient has not received prior systemic therapy in the recurrent or metastatic setting; and
  - 2.3 Patient has a positive PD-L1 combined positive score (CPS) of greater than or equal to 1; and
  - 2.4 Patient has an ECOG performance score of 0-2; and
  - 2.5 Either:
    - 2.5.1 Pembrolizumab to be used in combination with platinum-based chemotherapy; or

Subsidy	Sı	Fully	Brand or
(Manufacturer's Price)		ubsidised	Generic
\$	Per	1	Manufacturer

continued...

- 2.5.2 Pembrolizumab to be used as monotherapy; and
- 2.6 Pembrolizumab is to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks.

Renewal — (head and neck squamous cell carcinoma) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 Patient's disease has had a complete response to treatment; or
  - 1.2 Patient's disease has had a partial response to treatment; or
  - 1.3 Patient has stable disease; and
- 2 No evidence of disease progression; and
- 3 Pembrolizumab is to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 4 Treatment with pembrolizumab is to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initial application — (MSI-H/dMMR advanced colorectal cancer) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria: Either:

- 1 Patient is currently on treatment with pembrolizumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
  - 2.1 Either:
    - 2.1.1 Patient has deficient mismatch repair (dMMR) or microsatellite instability-high (MSI-H) metastatic colorectal cancer: or
    - 2.1.2 Patient has deficient mismatch repair (dMMR) or microsatellite instability-high (MSI-H) unresectable colorectal cancer: and
  - 2.2 Patient is treated with palliative intent; and
  - 2.3 Patient has not previously received funded treatment with pembrolizumab; and
  - 2.4 Patient has an ECOG performance score of 0-2; and
  - 2.5 Baseline measurement of overall tumour burden is documented clinically and radiologically; and
  - 2.6 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks.

Renewal — (MSI-H/dMMR advanced colorectal cancer) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 3 Treatment with pembrolizumab is to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initial application — (Urothelial carcinoma) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria: Fither:

- 1 Patient is currently on treatment with pembrolizumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
  - 2.1 Patient has inoperable locally advanced (T4) or metastatic urothelial carcinoma; and
  - 2.2 Patient has an ECOG performance score of 0-2; and
  - 2.3 Patient has documented disease progression following treatment with chemotherapy; and
  - 2.4 Pembrolizumab to be used as monotherapy at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Renewal — (Urothelial carcinoma) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 Patient's disease has had a complete response to treatment; or
  - 1.2 Patient's disease has had a partial response to treatment; or
  - 1.3 Patient has stable disease; and
- 2 No evidence of disease progression; and
- 3 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 4 Treatment with pembrolizumab is to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initial application — (relapsed/refractory Hodgkin lymphoma) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria: Either:

- 1 Patient is currently on treatment with pembrolizumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
  - 2.1 Either:
    - 2.1.1 Both:
      - 2.1.1.1 Patient has relapsed/refractory Hodgkin lymphoma after two or more lines of chemotherapy; and
      - 2.1.1.2 Patient is ineligible for autologous stem cell transplant; or
    - 2.1.2 Patient has relapsed/refractory Hodgkin lymphoma and has previously undergone an autologous stem cell transplant; and
  - 2.2 Patient has not previously received funded pembrolizumab; and
  - 2.3 Pembrolizumab to be administered at doses no greater than 200 mg once every 3 weeks.

Renewal — (relapsed/refractory Hodgkin lymphoma) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient has received a partial or complete response to pembrolizumab; and
- 2 Treatment with pembrolizumab is to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

# Other Immunosuppressants

CICLOSPORIN			
Cap 25 mg	44.63	50	✓ Neoral
Cap 50 mg		50	✓ Neoral
Cap 100 mg		50	✓ Neoral
Oral liq 100 mg per ml		50 ml OP	✓ Neoral
EVEROLIMUS - Special Authority see SA2414 below - Retail pha	rmacy		
Wastage claimable			
Tab 10 mg	6,512.29	30	✓ Afinitor
Tab 5 mg	4,555.76	30	✓ Afinitor

⇒SA2414 Special Authority for Subsidy

**Initial application** only from a neurologist or oncologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

Subsidy		Fully	Brand or	_
(Manufacturer's Price)	)	Subsidised	Generic	
\$	Per	1	Manufacturer	

continued...

- 1 Patient has tuberous sclerosis; and
- 2 Patient has progressively enlarging sub-ependymal giant cell astrocytomas (SEGAs) that require treatment.

**Renewal** only from a neurologist or oncologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Documented evidence of SEGA reduction or stabilisation by MRI within the last 3 months; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Everolimus to be discontinued at progression of SEGAs.

Initial application — (renal cell carcinoma) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 All of the following:
  - 1.1 The patient has metastatic renal cell carcinoma; and
  - 1.2 The disease is of predominant clear-cell histology; and
  - 1.3 The patient has documented disease progression following one previous line of treatment; and
  - 1.4 The patient has an ECOG performance status of 0-2; and
  - 1.5 Everolimus is to be used in combination with lenvatinib; or
- 2 All of the following:
  - 2.1 Patient has received funded treatment with nivolumab for the second line treatment of metastatic renal cell carcinoma; and
  - 2.2 Patient has experienced treatment limiting toxicity from treatment with nivolumab; and
  - 2.3 Everolimus is to be used in combination with lenvatinib; and
  - 2.4 There is no evidence of disease progression.

Renewal — (renal cell carcinoma) from any relevant practitioner. Approvals valid for 4 months where there is no evidence of disease progression.

SIROLIMUS - Special Authority see SA2270 below - Retail pharmacy

Tab 1 mg	749.99	100	Rapamune
Tab 2 mg	1,499.99	100	✓ Rapamune
Oral lig 1 mg per ml	449.99	60 ml OP	✓ Rapamune

#### ⇒SA2270 Special Authority for Subsidy

**Initial application** from any medical practitioner. Approvals valid without further renewal unless notified where the drug is to be used for rescue therapy for an organ transplant recipient.

Notes: Rescue therapy defined as unresponsive to calcineurin inhibitor treatment as defined by refractory rejection; or intolerant to calcineurin inhibitor treatment due to any of the following:

- GFR< 30 ml/min: or
- Rapidly progressive transplant vasculopathy; or
- · Rapidly progressive obstructive bronchiolitis; or
- . HUS or TTP; or
- Leukoencepthalopathy; or
- · Significant malignant disease

Initial application — (severe non-malignant lymphovascular malformations\*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe non-malignant lymphovascular malformation\*; and
- 2 Any of the following:
  - 2.1 Malformations are not adequately controlled by sclerotherapy and surgery; or
  - 2.2 Malformations are widespread/extensive and sclerotherapy and surgery are not considered clinically appropriate; or

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	1	Manufacturer	

continued...

- 2.3 Sirolimus is to be used to reduce malformation prior to consideration of surgery; and
- 3 Patient is being treated by a specialist lymphovascular malformation multi-disciplinary team; and
- 4 Patient has measurable disease as defined by RECIST version 1.1 (see Note).

Renewal — (severe non-malignant lymphovascular malformations\*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
  - 1.1 Patient's disease has had either a complete response or a partial response to treatment, or patient has stable disease according to RECIST version 1.1 (see Note); or
  - 1.2 Patient's disease has stabilised or responded clinically and disease response to treatment has been clearly documents in patient notes; and
- 2 No evidence of progressive disease; and
- 3 The treatment remains clinically appropriate and the patient is benefitting from the treatment.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer et al. Eur J Cancer 2009;45:228-47)

Indications marked with \* are unapproved indications

Initial application — (renal angiomyolipoma(s) associated with tuberous sclerosis complex\*) only from a nephrologist or urologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has tuberous sclerosis complex\*; and
- 2 Evidence of renal angiomyolipoma(s) measuring 3 cm or greater and that have shown interval growth.

Renewal — (renal angiomyolipoma(s) associated with tuberous sclerosis complex\*) from any relevant practitioner.

Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Documented evidence of renal angiomyolipoma reduction or stability by magnetic resonance imaging (MRI) or ultrasound;
- 2 Demonstrated stabilisation or improvement in renal function; and
- 3 The patient has not experienced angiomyolipoma haemorrhage or significant adverse effects to sirolimus treatment; and
- 4 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indications marked with \* are unapproved indications

Initial application — (refractory seizures associated with tuberous sclerosis complex\*) only from a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has epilepsy with a background of documented tuberous sclerosis complex; and
- 2 Either:
  - 2.1 Both:
    - 2.1.1 Vigabatrin has been trialled and has not adequately controlled seizures; and
    - 2.1.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least two of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); or
  - 2.2 Both:
    - 2.2.1 Vigabatrin is contraindicated; and
    - 2.2.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least three of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note): and
- 3 Seizures have a significant impact on quality of life; and

Subsidy		Fully	Brand or	
(Manufacturer's Pr	rice)	Subsidised	Generic	
\$	Per	•	Manufacturer	

continued...

4 Patient has been assessed and surgery is considered inappropriate for this patient, or the patient has been assessed and would benefit from mTOR inhibitor treatment prior to surgery.

Note: Those of childbearing age potential are not required to trial phenytoin sodium, sodium valproate, or topiramate. Those who can father children are not required to trial sodium valproate.

Renewal — (refractory seizures associated with tuberous sclerosis complex\*) only from a neurologist. Approvals valid for 12 months where demonstrated significant and sustained improvement in seizure rate (e.g. 50% reduction in seizure frequency) or severity and/or patient quality of life compared with baseline prior to starting sirolimus treatment.

Note: Indications marked with \* are unapproved indications

TACROLIMUS - Special Authority see SA2271 below - Retail pharmacy

Cap 0.5 mg	100	✓ Tacrolimus Sandoz
Cap 0.75 mg	100	✓ Tacrolimus Sandoz
Cap 1 mg84.30	100	✓ Tacrolimus Sandoz
Cap 5 mg248.20	50	✓ Tacrolimus Sandoz

#### ⇒SA2271 Special Authority for Subsidy

**Initial application — (organ transplant)** only from a relevant specialist. Approvals valid without further renewal unless notified where the patient is an organ transplant recipient.

Note: Subsidy applies for either primary or rescue therapy.

Initial application — (non-transplant indications\*) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient requires long-term systemic immunosuppression; and
- 2 Either:
  - 2.1 Ciclosporin has been trialled and discontinued treatment because of unacceptable side effects or inadequate clinical response; or
  - 2.2 Patient is a child with nephrotic syndrome\*.

Note: Indications marked with \* are unapproved indications

#### JAK inhibitors

### ⇒SA2079 Special Authority for Subsidy

Initial application — (Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
  - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
  - 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
- 3 Fither:
  - 3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or
  - 3.2 Both:
    - 3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a Health NZ Hospital; and 3.2.2 Fither:

Subsidy	5	Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	1	Manufacturer

continued...

- 3.2.2.1 The patient has experienced intolerable side effects from rituximab; or
- 3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Renewal — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Either:

- 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Subsi	idy Fully Brand or
(Manufacture	er's Price) Subsidised Generic
\$	Per   Manufacturer

# **Antiallergy Preparations**

### Allergic Emergencies

ADRENALINE - Special Authority see SA2185 below - Retail pharmacy

- a) Maximum of 2 ini per prescription
- Additional prescriptions limited to replacement of up to two devices prior to expiry, or replacement of used device for treatment of anaphylaxis.

Inj 0.15 mg per 0.3 ml auto-injector	90.00	1 OP	<ul> <li>Epipen Jr</li> </ul>
Inj 0.3 mg per 0.3 ml auto-injector	90.00	1 OP	<ul><li>Epipen</li></ul>

### ⇒SA2185 Special Authority for Subsidy

Initial application — (anaphylaxis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 Patient has experienced an anaphylactic reaction which has resulted in presentation to a hospital or emergency department; or
  - 1.2 Patient has been assessed to be at significant risk of anaphylaxis by a relevant practitioner; and
- 2 Patient is not to be prescribed more than two devices in initial prescription.

#### ⇒SA1558 Special Authority for Subsidy

**Initial application** only from a clinical immunologist or relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Supply for anticipated emergency treatment of laryngeal/oro-pharyngeal or severe abdominal attacks of acute hereditary angioedema (HAE) for patients with confirmed diagnosis of C1-esterase inhibitor deficiency; and
- 2 The patient has undergone product training and has agreed upon an action plan for self-administration.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

# Allergy Desensitisation

#### ⇒SA1367 Special Authority for Subsidy

Initial application only from a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 RAST or skin test positive; and
- 2 Patient has had severe generalised reaction to the sensitising agent.

**Renewal** only from a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

BEE VENOM ALLERGY TREATMENT - Special Authority see SA1367 above - Retail pharmacy Initiation kit - 5 vials freeze dried venom with diluent 305.00 1 OP ✓ VENOX S29 Maintenance kit - 1 vial freeze dried venom with diluent...............305.00 1 OP ✓ VENOX S29 Maintenance kit - 6 vials 120 mcg freeze dried venom, with 1 OP ✓ Venomil S29 Treatment kit - 1 vial 550 mcg freeze dried venom, 1 diluent 9 ml, 3 diluent 1.8 ml .......334.39 1 OP ✓ Albev Treatment kit - 1 vial 550 mcg freeze dried venom, with diluent ..... 305.00 1 OP ✓ Hymenoptera S29

	Subsidy (Manufacturer's Pr	ice) Subsi	. ,	rand or eneric
	\$	Per		lanufacturer
WASP VENOM ALLERGY TREATMENT - Special Authority se	ee SA1367 on the r	revious page	- Retail pl	narmacy
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze				,
dried polistes venom, 1 diluent 9 ml, 3 diluent 1.8 ml	382.23	1 OP	✓ Albe	y
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze				
dried venom, with diluent	305.00	1 OP	Hym	enoptera S29
Treatment kit (Paper wasp venom) - 6 vials 120 mcg freeze			_	
dried venom, with diluent		1 OP	✓ Vend	omil S29
Treatment kit (Yellow Jacket venom) - 1 vial 550 mcg freez				
dried venom, with diluent		1 OP	✓ Hym	enoptera S29
Treatment kit (Yellow jacket venom) - 1 vial 550 mcg freeze		4.00	/ A11-	
dried vespula venom, 1 diluent 9 ml, 3 diluent 1.8 ml		1 OP	✓ Albe	y
Treatment kit (Yellow jacket venom) - 6 vials 120 mcg freez dried venom, with diluent		1 OP	√ Von	omil S29
unea venom, with anaem		I UF	- ven	JIIII 929
Antihistamines				
CETIRIZINE HYDROCHLORIDE	4 74	100	./ 7:-1	_
* Tab 10 mg		100 200 ml	✓ Zista ✓ Hista	-
* Oral liq 1 mg per ml	2.04	200 1111	▼ nist	aciear
DEXTROCHLORPHENIRAMINE MALEATE	0.00	40		
* Tab 2 mg		40	Polo	ramine
	(8.40) 1.01	20	Fula	rannine
	(5.99)	20	Pola	ramine
* Oral liq 2 mg per 5 ml		100 ml	1 014	
37.	(10.29)		Pola	ramine
FEXOFENADINE HYDROCHLORIDE				
* Tab 60 mg	4.34	20		
	(8.23)		Telfa	ıst
* Tab 120 mg	4.74	10		
	(8.23)		Telfa	ıst
	14.22	30		
	(26.44)		Telfa	ıst
LORATADINE			4.	
* Tab 10 mg		100	Lora	
* Oral liq 1 mg per ml	1.43	100 ml	• нау	or syrup
PROMETHAZINE HYDROCHLORIDE	4.00			
* Tab 10 mg		50	✓ Allei	
* Tab 25 mg  * Oral lig 1 mg per 1 ml		50 100 ml	✓ <u>Aller</u> ✓ Aller	
★ Oralliq Fing per Hill ★ Inj 25 mg per ml, 2 ml ampoule – Up to 5 inj available on a		5	✓ Hos	
III 25 III poi IIII, 2 III ampoulo op to 5 III available off a		•	- 1103	y 11 4
Inhaled Corticosteroids				
BECLOMETHASONE DIPROPIONATE				
Aerosol inhaler, 50 mcg per dose	14.01	200 dose OP	✓ Qva	r
Aerosol inhaler, 50 mcg per dose CFC-free		200 dose OP		azone 50
Aerosol inhaler, 100 mcg per dose		200 dose OP	✓ Qva	
Aerosol inhaler, 100 mcg per dose CFC-free	12.50	200 dose OP	✓ Becl	azone 100

Aerosol inhaler, 250 mcg per dose CFC-free ......22.67

200 dose OP ✓ Beclazone 250

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's		Fully Brand or sidised Generic
	\$	Per	✓ Manufacturer
UDESONIDE			
Powder for inhalation, 100 mcg per dose	17.00	200 dose OP	✓ Pulmicort Turbuhaler
Powder for inhalation, 200 mcg per dose	19.00	200 dose OP	✓ Pulmicort Turbuhaler
Powder for inhalation, 400 mcg per dose	32.00	200 dose OP	
LUTICASONE			Turburiaici
Aerosol inhaler, 50 mcg per dose	7 19	120 dose OP	✓ Flixotide
Powder for inhalation, 50 mcg per dose		60 dose OP	✓ Flixotide Accuhaler
		60 dose OP	✓ Flixotide Accuhaler
Powder for inhalation, 100 mcg per dose			
Aerosol inhaler, 125 mcg per dose		120 dose OP	
Aerosol inhaler, 250 mcg per dose		120 dose OP	
Powder for inhalation, 250 mcg per dose	11.93	60 dose OP	✓ Flixotide Accuhaler
nhaled Long-acting Beta-adrenoceptor Agonis	sts		
FORMOTEROL FUMARATE DIHYDRATE			
Powder for inhalation 4.5 mcg per dose, breath activated			
(equivalent to eformoterol fumarate 6 mcg metered dos	e) 10.32	60 dose OP	
(equivalent to elormoterol lumarate o meg metered dos	(16.90)	OU GUSE OF	Oxis Turbuhaler
	(10.90)		Oxis Turburialer
IDACATEROL			
Powder for inhalation 150 mcg	61.00	30 dose OP	<ul> <li>Onbrez Breezhaler</li> </ul>
Powder for inhalation 300 mcg	61.00	30 dose OP	Onbrez Breezhaler
ALMETEROL			
	00.05	100 dees OD	. Communit
Aerosol inhaler CFC-free, 25 mcg per dose		120 dose OP	• • • • • • • • • • • • • • • • • • • •
Powder for inhalation 50 med per doce breath activated	26.25	60 dose OP	Serevent Accuhaler
Powder for inhalation, 50 mcg per dose, breath activated		00 dose or	
nhaled Corticosteroids with Long-Acting Beta			3
			3
nhaled Corticosteroids with Long-Acting Beta-			3
nhaled Corticosteroids with Long-Acting Beta- UDESONIDE WITH EFORMOTEROL Powder for inhalation 160 mcg with 4.5 mcg eformoterol	-Adrenocept		5
nhaled Corticosteroids with Long-Acting Beta- UDESONIDE WITH EFORMOTEROL Powder for inhalation 160 mcg with 4.5 mcg eformoterol fumarate per dose (equivalent to 200 mcg budesonide v	-Adrenocept	tor Agonists	
nhaled Corticosteroids with Long-Acting Beta- UDESONIDE WITH EFORMOTEROL  Powder for inhalation 160 mcg with 4.5 mcg eformoterol fumarate per dose (equivalent to 200 mcg budesonide of mcg eformoterol fumarate metered dose)	-Adrenocept		
nhaled Corticosteroids with Long-Acting Beta- UDESONIDE WITH EFORMOTEROL  Powder for inhalation 160 mcg with 4.5 mcg eformoterol fumarate per dose (equivalent to 200 mcg budesonide 6 mcg eformoterol fumarate metered dose)	-Adrenocept with41.50 rate	tor Agonists	
nhaled Corticosteroids with Long-Acting Beta- UDESONIDE WITH EFORMOTEROL  Powder for inhalation 160 mcg with 4.5 mcg eformoterol fumarate per dose (equivalent to 200 mcg budesonide 6 mcg eformoterol fumarate metered dose)	-Adrenocept with41.50 rate	tor Agonists	
nhaled Corticosteroids with Long-Acting Beta- UDESONIDE WITH EFORMOTEROL  Powder for inhalation 160 mcg with 4.5 mcg eformoterol fumarate per dose (equivalent to 200 mcg budesonide 6 mcg eformoterol fumarate metered dose)  Powder for inhalation 320 mcg with 9 mcg eformoterol fuma per dose (equivalent to 400 mcg budesonide with 12 mc eformoterol fumarate metered dose) – No more than 2	with	tor Agonists	✓ DuoResp Spiromax
nhaled Corticosteroids with Long-Acting Beta- UDESONIDE WITH EFORMOTEROL  Powder for inhalation 160 mcg with 4.5 mcg eformoterol fumarate per dose (equivalent to 200 mcg budesonide of mcg eformoterol fumarate metered dose)  Powder for inhalation 320 mcg with 9 mcg eformoterol fumare per dose (equivalent to 400 mcg budesonide with 12 mcg eformoterol fumarate metered dose) – No more than 2 dose per day	with	tor Agonists	<ul><li>✓ DuoResp Spiromax</li><li>✓ DuoResp Spiromax</li></ul>
nhaled Corticosteroids with Long-Acting Beta- UDESONIDE WITH EFORMOTEROL  Powder for inhalation 160 mcg with 4.5 mcg eformoterol fumarate per dose (equivalent to 200 mcg budesonide 6 mcg eformoterol fumarate metered dose)  Powder for inhalation 320 mcg with 9 mcg eformoterol fuma per dose (equivalent to 400 mcg budesonide with 12 mc eformoterol fumarate metered dose) – No more than 2	with	tor Agonists	✓ DuoResp Spiromax ✓ DuoResp Spiromax
nhaled Corticosteroids with Long-Acting Beta- UDESONIDE WITH EFORMOTEROL  Powder for inhalation 160 mcg with 4.5 mcg eformoterol fumarate per dose (equivalent to 200 mcg budesonide of mcg eformoterol fumarate metered dose)  Powder for inhalation 320 mcg with 9 mcg eformoterol fuma per dose (equivalent to 400 mcg budesonide with 12 mc eformoterol fumarate metered dose) – No more than 2 dose per day	with	tor Agonists  120 dose OP	✓ DuoResp Spiromax ✓ DuoResp Spiromax ✓ Vannair
nhaled Corticosteroids with Long-Acting Beta- UDESONIDE WITH EFORMOTEROL  Powder for inhalation 160 mcg with 4.5 mcg eformoterol fumarate per dose (equivalent to 200 mcg budesonide of mcg eformoterol fumarate metered dose)  Powder for inhalation 320 mcg with 9 mcg eformoterol fumare per dose (equivalent to 400 mcg budesonide with 12 mcg eformoterol fumarate metered dose) – No more than 2 dose per day	with	120 dose OP 120 dose OP 120 dose OP 120 dose OP	<ul> <li>✓ DuoResp Spiromax</li> <li>✓ DuoResp Spiromax</li> <li>✓ Vannair</li> <li>✓ Symbicort</li> </ul>
nhaled Corticosteroids with Long-Acting Beta- UDESONIDE WITH EFORMOTEROL  Powder for inhalation 160 mcg with 4.5 mcg eformoterol fumarate per dose (equivalent to 200 mcg budesonide of mcg eformoterol fumarate metered dose)  Powder for inhalation 320 mcg with 9 mcg eformoterol fumarate per dose (equivalent to 400 mcg budesonide with 12 mc eformoterol fumarate metered dose) – No more than 2 dose per day	with	120 dose OP	<ul> <li>✓ DuoResp Spiromax</li> <li>✓ DuoResp Spiromax</li> <li>✓ Vannair</li> <li>✓ Symbicort         <ul> <li>Turbuhaler 100/6</li> </ul> </li> </ul>
nhaled Corticosteroids with Long-Acting Beta- UDESONIDE WITH EFORMOTEROL  Powder for inhalation 160 mcg with 4.5 mcg eformoterol fumarate per dose (equivalent to 200 mcg budesonide of mcg eformoterol fumarate metered dose)  Powder for inhalation 320 mcg with 9 mcg eformoterol fumarate per dose (equivalent to 400 mcg budesonide with 12 mcg. eformoterol fumarate metered dose) — No more than 2 dose per day	with	120 dose OP 120 dose OP 120 dose OP 120 dose OP 120 dose OP	✓ DuoResp Spiromax ✓ DuoResp Spiromax ✓ Vannair ✓ Symbicort Turbuhaler 100/6 ✓ Vannair
nhaled Corticosteroids with Long-Acting Beta- UDESONIDE WITH EFORMOTEROL  Powder for inhalation 160 mcg with 4.5 mcg eformoterol fumarate per dose (equivalent to 200 mcg budesonide of mcg eformoterol fumarate metered dose)  Powder for inhalation 320 mcg with 9 mcg eformoterol fumarate per dose (equivalent to 400 mcg budesonide with 12 mc eformoterol fumarate metered dose) – No more than 2 dose per day	with	120 dose OP	✓ DuoResp Spiromax ✓ DuoResp Spiromax ✓ Vannair ✓ Symbicort Turbuhaler 100/6 ✓ Vannair ✓ Symbicort
nhaled Corticosteroids with Long-Acting Beta- UDESONIDE WITH EFORMOTEROL  Powder for inhalation 160 mcg with 4.5 mcg eformoterol fumarate per dose (equivalent to 200 mcg budesonide of mcg eformoterol fumarate metered dose)  Powder for inhalation 320 mcg with 9 mcg eformoterol fumarate per dose (equivalent to 400 mcg budesonide with 12 mcg. eformoterol fumarate metered dose) — No more than 2 dose per day	with	120 dose OP 120 dose OP 120 dose OP 120 dose OP 120 dose OP	✓ DuoResp Spiromax ✓ DuoResp Spiromax ✓ Vannair ✓ Symbicort Turbuhaler 100/6 ✓ Vannair
nhaled Corticosteroids with Long-Acting Beta- UDESONIDE WITH EFORMOTEROL  Powder for inhalation 160 mcg with 4.5 mcg eformoterol fumarate per dose (equivalent to 200 mcg budesonide 6 mcg eformoterol fumarate metered dose)	with	120 dose OP 120 dose OP 120 dose OP 120 dose OP 120 dose OP	✓ DuoResp Spiromax ✓ DuoResp Spiromax ✓ Vannair ✓ Symbicort Turbuhaler 100/6 ✓ Vannair ✓ Symbicort Turbuhaler 200/6
nhaled Corticosteroids with Long-Acting Beta- UDESONIDE WITH EFORMOTEROL  Powder for inhalation 160 mcg with 4.5 mcg eformoterol fumarate per dose (equivalent to 200 mcg budesonide of mcg eformoterol fumarate metered dose)	with	120 dose OP 120 dose OP 120 dose OP 120 dose OP 120 dose OP	✓ DuoResp Spiromax ✓ DuoResp Spiromax ✓ Vannair ✓ Symbicort Turbuhaler 100/6 ✓ Vannair ✓ Symbicort
nhaled Corticosteroids with Long-Acting Beta- UDESONIDE WITH EFORMOTEROL  Powder for inhalation 160 mcg with 4.5 mcg eformoterol fumarate per dose (equivalent to 200 mcg budesonide 6 mcg eformoterol fumarate metered dose)	with	120 dose OP 120 dose OP 120 dose OP 120 dose OP 120 dose OP 120 dose OP	✓ DuoResp Spiromax ✓ DuoResp Spiromax ✓ Vannair ✓ Symbicort Turbuhaler 100/6 ✓ Vannair ✓ Symbicort Turbuhaler 200/6 ✓ Symbicort
nhaled Corticosteroids with Long-Acting Beta- UDESONIDE WITH EFORMOTEROL  Powder for inhalation 160 mcg with 4.5 mcg eformoterol fumarate per dose (equivalent to 200 mcg budesonide 6 mcg eformoterol fumarate metered dose)	with	120 dose OP 120 dose OP 120 dose OP 120 dose OP 120 dose OP 120 dose OP	✓ DuoResp Spiromax ✓ DuoResp Spiromax ✓ Vannair ✓ Symbicort Turbuhaler 100/6 ✓ Vannair ✓ Symbicort Turbuhaler 200/6 ✓ Symbicort
nhaled Corticosteroids with Long-Acting Beta- UDESONIDE WITH EFORMOTEROL  Powder for inhalation 160 mcg with 4.5 mcg eformoterol fumarate per dose (equivalent to 200 mcg budesonide 6 mcg eformoterol fumarate metered dose)	with	120 dose OP 120 dose OP 120 dose OP 120 dose OP 120 dose OP 120 dose OP	✓ DuoResp Spiromax ✓ DuoResp Spiromax ✓ Vannair ✓ Symbicort Turbuhaler 100/6 ✓ Vannair ✓ Symbicort Turbuhaler 200/6

	IILOI IIIA	10111 01012	IN AND ALLENGIES
	Subsidy (Manufacturer's	Price) Subsi Per	Fully Brand or idised Generic Manufacturer
FLUTICASONE WITH SALMETEROL			
Aerosol inhaler 50 mcg with salmeterol 25 mcg		120 dose OP	✓ Seretide
Aerosol inhaler 125 mcg with salmeterol 25 mcg	32.60	120 dose OP	✓ Seretide
Powder for inhalation 100 mcg with salmeterol 50 mcg - No			
more than 2 dose per day	33.74	60 dose OP	Seretide Accuhaler
Powder for inhalation 250 mcg with salmeterol 50 mcg – No	44.00	00 de e OD	Constitute Assessments
more than 2 dose per day	44.08	60 dose OP	Seretide Accuhaler
Beta-Adrenoceptor Agonists			
SALBUTAMOL			
Oral liq 400 mcg per ml	50.00	150 ml	✓ Ventolin
Infusion 1 mg per ml, 5 ml		10	✓ Ventolin
Inj 500 mcg per ml, 1 ml - Up to 5 inj available on a PSO	130.00	5	✓ Ventolin
Inhaled Beta-Adrenoceptor Agonists			
SALBUTAMOL			
Aerosol inhaler, 100 mcg per dose CFC free – Up to 1000			
dose available on a PSO	3.80	200 dose OP	✓ SalAir
	(6.80)	200 0000 0.	Ventolin
Nebuliser soln, 1 mg per ml, 2.5 ml ampoule - Up to 30 neb	, ,		
available on a PSO	8.96	20	✓ Asthalin
Nebuliser soln, 2 mg per ml, 2.5 ml ampoule - Up to 30 neb			
available on a PSO	9.43	20	✓ Asthalin
TERBUTALINE SULPHATE			
Powder for inhalation, 200 mcg per dose (equivalent to			
250 mcg metered dose), breath activated	22.20	120 dose OP	Bricanyl Turbuhaler
Anticholinergic Agents			
IPRATROPIUM BROMIDE			
Aerosol inhaler, 20 mcg per dose CFC-free - Up to 400 dose	е		
available on a PSO		200 dose OP	✓ Atrovent
Nebuliser soln, 250 mcg per ml, 2 ml ampoule - Up to 40 ne			
available on a PSO	5.86	10	✓ Pharmascience S29
	11.73	20	✓ Ipratropium
			IVAX S29
			✓ Univent
	28.20		✓ Accord S29

(Pharmascience S29 Nebuliser soln, 250 mcg per ml, 2 ml ampoule to be delisted 1 May 2025)
(Ipratropium IVAX S29 Nebuliser soln, 250 mcg per ml, 2 ml ampoule to be delisted 1 February 2025)
(Accord S29 Nebuliser soln, 250 mcg per ml, 2 ml ampoule to be delisted 1 January 2025)

Subsi	idy Ful	y Brand or
(Manufacture	er's Price) Subsidise	d Generic
\$	Per •	Manufacturer

### Inhaled Beta-Adrenoceptor Agonists with Anticholinergic Agents

#### SALBUTAMOL WITH IPRATROPIUM BROMIDE

Aerosol inhaler, 100 mcg with ipratropium bromide, 20 mcg per		
dose CFC-free12.19	200 dose OP	✓ Duolin HFA
Nebuliser soln, 2.5 mg with ipratropium bromide 0.5 mg per		
vial, 2.5 ml ampoule - Up to 20 neb available on a PSO 11.04	20	✓ Duolin
		✓ Duolin Cipla S29
33.12	60	✓ Duolin
		Respules \$29

(Duolin Cipla S29 Nebuliser soln, 2.5 mg with ipratropium bromide 0.5 mg per vial, 2.5 ml ampoule to be delisted 1 April 2025) (Duolin Respules S29 Nebuliser soln, 2.5 mg with ipratropium bromide 0.5 mg per vial, 2.5 ml ampoule to be delisted 1 January 2025)

### **Long-Acting Muscarinic Antagonists**

#### GLYCOPYRRONIUM - Subsidy by endorsement

- a) Inhaled glycopyrronium treatment will not be subsidised if patient is also receiving treatment with subsidised tiotropium or umeclidinium.
- b) Glycopyrronium powder for inhalation 50 mcg per dose is subsidised only for patients who have been diagnosed as having COPD using spirometry if spirometry is possible, and the prescription is endorsed accordingly.

### TIOTROPIUM BROMIDE - Subsidy by endorsement

- a) Tiotropium treatment will not be subsidised if patient is also receiving treatment with subsidised inhaled glycopyrronium or umeclidinium.
- b) Tiotropium bromide is subsidised only for patients who have been diagnosed as having COPD using spirometry if spirometry is possible, and the prescription is endorsed accordingly. Patients who had tiotropium dispensed before 1 October 2018 with a valid Special Authority are deemed endorsed.

#### UMECLIDINIUM - Subsidy by endorsement

- a) Umeclidinium will not be subsidised if patient is also receiving treatment with subsidised inhaled glycopyrronium or tiotropium bromide.
- b) Umeclidinium powder for inhalation 62.5 mcg per dose is subsidised only for patients who have been diagnosed as having COPD using spirometry if spirometry is possible, and the prescription is endorsed accordingly.

# Long-Acting Muscarinic Antagonists with Long-Acting Beta-Adrenoceptor Agonists

Combination long acting muscarinic antagonist and long acting beta-2 agonist will not be subsidised if patient is also receiving treatment with a combination inhaled corticosteroid and long acting beta-2 agonist.

#### ⇒SA1584 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Patient has been stabilised on a long acting muscarinic antagonist; and
- 2 The prescriber considers that the patient would receive additional benefit from switching to a combination product.

Subsidy	e)	Fully	Brand or
(Manufacturer's Price		Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

- 1 Patient is compliant with the medication; and
- 2 Patient has experienced improved COPD symptom control (prescriber determined).

GLYCOPYRRONIUM WITH INDACATEROL - Special Authority see SA1584 on the previous page - Retail pharmacy ✓ Ultibro Breezhaler Powder for Inhalation 50 mcg with indacaterol 110 mcg......81.00 30 dose OP TIOTROPIUM BROMIDE WITH OLODATEROL - Special Authority see SA1584 on the previous page - Retail pharmacy

✓ Spiolto Respimat Soln for inhalation 2.5 mcg with olodaterol 2.5 mcg.....81.00 60 dose OP

UMECLIDINIUM WITH VILANTEROL - Special Authority see SA1584 on the previous page - Retail pharmacy Powder for inhalation 62.5 mcg with vilanterol 25 mcg ......77.00 30 dose OP ✓ Anoro Ellipta

### Inhaled Corticosteroid with Long-Acting Muscarinic Antagonist and Beta Agonist

FLUTICASONE FUROATE WITH UMECLIDINIUM AND VILANTEROL - Special Authority see \$A2326 below - Retail pharmacy Powder for inhalation fluticasone furoate 100 mcg with

umeclidinium 62.5 mcg and vilanterol 25 mcg......104.24 ✓ Trelegy Ellipta 30 dose OP

#### ⇒SA2326 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Both:

- 1 Patient has a diagnosis of COPD confirmed by spirometry or spirometry has been attempted and technically acceptable results are not possible; and
  - 2 Either:
    - 2.1 Both:
      - 2.1.1 Patient is currently receiving an inhaled corticosteroid with long acting beta-2 agonist (ICS/LABA) or a long acting muscarinic antagonist with long acting beta-2 agonist (LAMA/LABA); and
      - 2.1.2 Any of the following:

Clinical criteria:

- 2.1.2.1 Patient has a COPD Assessment Test (CAT) score greater than 10; or
- 2.1.2.2 Patient has had 2 or more exacerbations in the previous 12 months; or
- 2.1.2.3 Patient has had one exacerbation requiring hospitalisation in the previous 12 months; or
- 2.1.2.4 Patient has had an eosinophil count greater than or equal to 0.3 x 10°9 cells/L in the previous 12 months: or
- 2.2 Patient is currently receiving multiple inhaler triple therapy (inhaled corticosteroid with long acting muscarinic antagonist and long acting beta-2 agonist - ICS/LAMA/LABA) and met at least one of the clinical criteria above prior to commencing multiple inhaler triple therapy.

# **Antifibrotics**

NINTEDANIB - Special Authority see SA2012 below - Retail pharmacy

Note: Nintedanib not subsidised in combination with subsidised pirfenidone.

60 OP ✓ Ofev 60 OP ✓ Ofev

#### ⇒SA2012 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

Subsidy		Fully	Brand or	
(Manufacturer's Price	,	Subsidised	Generic	
\$	Per		Manufacturer	

continued...

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Nintedanib is to be discontinued at disease progression (See Note); and
- 4 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 5 Any of the following:
  - 5.1 The patient has not previously received treatment with pirfenidone; or
  - 5.2 Patient has previously received pirfenidone, but discontinued pirfenidone within 12 weeks due to intolerance; or
  - 5.3 Patient has previously received pirfenidone, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with pirfenidone).

Renewal — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 3 Nintedanib is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

PIRFENIDONE - Retail pharmacy-Specialist - Special Authority see SA2013 below

Note: Pirfenidone is not subsidised in combination with subsidised nintedanib.

Tab 801 mg	3,645.00	90 OP	<ul><li>Esbriet</li></ul>
Tab 267 mg	1,215.00	90	<ul><li>Esbriet</li></ul>

#### ⇒SA2013 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note); and
- 4 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 5 Any of the following:
  - 5.1 The patient has not previously received treatment with nintedanib; or
  - 5.2 Patient has previously received nintedanib, but discontinued nintedanib within 12 weeks due to intolerance; or
  - 5.3 Patient has previously received nintedanib, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with nintedanib).

Renewal — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

MONTELLIZACT

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

# Leukotriene Receptor Antagonists

✓ fully subsidised

**Principal Supply** 

IVIC	INTELUKAST		
*	Tab 4 mg3.10	28	✓ Montelukast Viatris
	Tab 5 mg	28	✓ Montelukast Viatris
	Tab 10 mg	28	✓ Montelukast Viatris

vlline

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	sidised	Generic	
\$	Per	✓	Manufacturer	

# Methylxanthines

#### **AMINOPHYLLINE**

PSO	180.00	5	✓ DBL Aminophy
THEOPHYLLINE			
* Tab long-acting 250 mg	24.90	100	✓ Nuelin-SR
* Oral lig 80 mg per 15 ml	17 95	500 ml	✓ Nuelin

### Mucolytics

DORNASE ALFA - Special Authority see SA1978 below - Reta	ail pharmacy		
Nebuliser soln, 2.5 mg per 2.5 ml ampoule	250.00	6	✓ Pulmozyme

#### ⇒SA1978 Special Authority for Subsidy

Initial application — (cystic fibrosis) only from a respiratory physician or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 Patient has a confirmed diagnosis of cystic fibrosis; and

\* Ini 25 mg ner ml 10 ml amnoule - I in to 5 ini available on a

- 2 Patient has previously undergone a trial with, or is currently being treated with, hypertonic saline; and
- 3 Any of the following:
  - 3.1 Patient has required one or more hospital inpatient respiratory admissions in the previous 12 month period; or
  - 3.2 Patient has had 3 exacerbations due to CF, requiring oral or intravenous (IV) antibiotics in the previous 12 month period: or
  - 3.3 Patient has had 1 exacerbation due to CF, requiring oral or IV antibiotics in the previous 12 month period and a Brasfield score of < 22/25; or
  - 3.4 Patient has a diagnosis of allergic bronchopulmonary aspergillosis (ABPA).

**Renewal** — **(cystic fibrosis)** only from a respiratory physician or paediatrician. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient continues to benefit from treatment.

ELEXACAFTOR WITH TEZACAFTOR, IVACAFTOR AND IVACAFTOR - PCT only - Special Authority see SA2196 below

Tab elexacaftor 50 mg with tezacaftor 25 mg, ivacaftor 37.5 mg

Tab elexacaftor 100 mg with tezacaftor 50 mg, ivacaftor 75 mg

(56) and ivacaftor 150 mg (28) .......27,647.39 84 OP ✓ **Trikafta** 

#### ⇒SA2196 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with cystic fibrosis; and
- 2 Patient is 6 years of age or older; and
- 3 Fither:
  - 3.1 Patient has two cystic fibrosis-causing mutations in the cystic fibrosis transmembrane regulator (CFTR) gene (one from each parental allele): or
  - 3.2 Patient has a sweat chloride value of at least 60 mmol/L by quantitative pilocarpine iontophoresis or by Macroduct sweat collection system; and
- 4 Either:
  - 4.1 Patient has a heterozygous or homozygous F508del mutation; or
  - 4.2 Patient has a G551D mutation or other mutation responsive in vitro to elexacaftor/tezacaftor/ivacaftor (see note a);

	ubsidy cturer's Price) Subs	Fully	Brand or Generic
·	\$ Per	•	Manufacturer

continued...

and

- 5 The treatment must be the sole funded CFTR modulator therapy for this condition; and
- 6 Treatment with elexacaftor/tezacaftor/ivacaftor must be given concomitantly with standard therapy for this condition.

#### Notes:

 a) Eligible mutations are listed in the Food and Drug Administration (FDA) Trikafta prescribing information https://www.accessdata.fda.gov/drugsatfda\_docs/label/2021/212273s004lbl.pdf

IVACAFTOR - PCT only - Specialist - Special Authorit	ty see SA2017 below		
Tab 150 mg	29,386.00	56	✓ Kalydeco
Oral granules 50 mg, sachet	29,386.00	56	✓ Kalydeco
Oral granules 75 mg, sachet	29,386.00	56	✓ Kalydeco

#### ⇒SA2017 Special Authority for Subsidy

Initial application only from a respiratory specialist or paediatrician. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### All of the following:

- 1 Patient has been diagnosed with cystic fibrosis; and
- 2 Either
  - 2.1 Patient must have G551D mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene on at least 1 allele: or
  - 2.2 Patient must have other gating (class III) mutation (G1244E, G1349D, G178R, G551S, S1251N, S1255P, S549N and S549R) in the CFTR gene on at least 1 allele: and
- 3 Patients must have a sweat chloride value of at least 60 mmol/L by quantitative pilocarpine iontophoresis or by Macroduct sweat collection system; and
- 4 Treatment with ivacaftor must be given concomitantly with standard therapy for this condition; and
- 5 Patient must not have an acute upper or lower respiratory infection, pulmonary exacerbation, or changes in therapy (including antibiotics) for pulmonary disease in the last 4 weeks prior to commencing treatment with ivacaftor; and
- 6 The dose of ivacaftor will not exceed one tablet or one sachet twice daily; and
- 7 Applicant has experience and expertise in the management of cystic fibrosis.

#### SODIUM CHI ORIDE

Not funded for use as a nasal drop.

# **Nasal Preparations**

# **Allergy Prophylactics**

BUDESONIDE		
Metered aqueous nasal spray, 50 mcg per dose2.59	200 dose OP	✓ SteroClear
SteroClear to be Principal Supply on 1 February 2025		
Metered agueous nasal spray, 100 mcg per dose2.89	200 dose OP	✓ SteroClear
SteroClear to be Principal Supply on 1 February 2025	200 0000 01	3.0.00.00.
, ,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,		
FLUTICASONE PROPIONATE		
Metered aqueous nasal spray, 50 mcg per dose	120 dose OP	<ul> <li>Flixonase Hayfever</li> </ul>
		& Allergy
IDDATDODIUM DDOMIDE		3,
IPRATROPIUM BROMIDE		_
Aqueous nasal spray, 0.03%5.23	15 ml OP	Univent

25 ml OP

✓ Biomed

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
Respiratory Devices				
MASK FOR SPACER DEVICE  a) Up to 50 dev available on a PSO b) Only on a PSO c) Only for children aged six years and under				
Small  PEAK FLOW METER  a) Up to 25 dev available on a PSO b) Only on a PSO	2.70	1	<b>√</b> e	-chamber Mask
Low range	9.54	1	✓ N	lini-Wright AFS Low Range
Normal range	9.54	1	✓ N	lini-Wright Standard
SPACER DEVICE				
a) Up to 50 dev available on a PSO     b) Only on a PSO				
220 ml (single patient)	3.65	1	<b>√</b> e	-chamber Turbo
510 ml (single patient)		1	<b>√</b> e	-chamber La Grande
800 ml	6.50	1	<b>√</b> ∨	olumatic
Respiratory Stimulants				
CAFFEINE CITRATE				

Oral liq 20 mg per ml (10 mg base per ml)......16.10



	Subsidy		Fully	Brand or
	(Manufacturer's Pr		idised	Generic
	\$	Per		Manufacturer
Ear Preparations				
FLUMETASONE PIVALATE				
Ear drops 0.02% with clioquinol 1%	4.46	7.5 ml OP	<b>√</b> L	ocacorten-Viaform ED's
			✓ L	.ocorten-Vioform
TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYC	IN AND NYSTATI	N		
Ear drops 1 mg with nystatin 100,000 u, neomycin sulphate				
2.5 mg and gramicidin 250 mcg per g	5.16	7.5 ml OP	✓ K	(enacomb
Ear/Eye Preparations				
DEXAMETHASONE WITH FRAMYCETIN AND GRAMICIDIN				
Ear/Eye drops 500 mcg with framycetin sulphate 5 mg and				
gramicidin 50 mcg per ml		8 ml OP		
	(9.27)			Otodex \$29
	(9.27)		5	Sofradex
FRAMYCETIN SULPHATE	4.40	0 100		
Ear/Eye drops 0.5%		8 ml OP	c	ofrom voin
	(8.65)		3	Soframycin
Eye Preparations				
Eye preparations are only funded for use in the eye, unless expli	icitly stated otherw	vise.		
Anti-Infective Preparations				
ACICLOVIR				
* Eye oint 3%	15.89	4.5 g OP	✓ \	/iruPOS
ViruPOS to be Principal Supply on 1 February 2025				
CHLORAMPHENICOL				
Eye oint 1%		5 g OP	_	<u>Devatis</u>
Eye drops 0.5%Funded for use in the ear*. Indications marked with * ar		10 ml OP	• [	Chlorsig
CIPROFLOXACIN	re unapproved mu	icalions.		
Eye drops 0.3% – Subsidy by endorsement	10.85	5 ml OP	10	Ciprofloxacin Teva
a) When prescribed for the treatment of bacterial kerati				•
or for the second line treatment of chronic suppurativ				
accordingly. Note: Indication marked with a * is an			•	•
b) Ciprofloxacin Teva to be Principal Supply on 1 March	h 2025			
SODIUM FUSIDATE [FUSIDIC ACID]				
Eye drops 1%	5.29	5 g OP	<b>✓</b> F	ucithalmic
TOBRAMYCIN				
Eye oint 0.3%		3.5 g OP		obrex
Eye drops 0.3%	11.48	5 ml OP	<b>✓</b> T	obrex

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

# **Corticosteroids and Other Anti-Inflammatory Preparations**

DE	XAMETHASONE			
*	Eye oint 0.1%	5.86	3.5 g OP	✓ Maxidex
*	Eye drops 0.1%	4.50	5 ml OP	<ul><li>Maxidex</li></ul>
	Ocular implant 700 mcg - Special Authority see SA1680 below			
	- Retail pharmacy1,4	44.50	1	<ul><li>Ozurdex</li></ul>

#### ⇒SA1680 Special Authority for Subsidy

Initial application — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

#### All of the following:

- 1 Patient has diabetic macular oedema with pseudophakic lens: and
- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- 3 Fither
  - 3.1 Patient's disease has progressed despite 3 injections with bevacizumab; or
  - 3.2 Patient is unsuitable or contraindicated to treatment with anti-VEGF agents; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

### Both:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Initial application — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

#### All of the following:

- 1 Patient has diabetic macular oedema; and
- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- 3 Patient is of child bearing potential and has not yet completed a family; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

#### All of the following:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Patient is of child bearing potential and has not yet completed a family; and
- 3 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

#### DEXAMETHASONE WITH NEOMYCIN SULPHATE AND POLYMYXIN B SULPHATE

* Eye oint 0.1% with neomycin sulphate 0.35% and polymyxin b			
sulphate 6,000 u per g	5.39	3.5 g OP	<ul><li>Maxitrol</li></ul>
* Eye drops 0.1% with neomycin sulphate 0.35% and polymyxin			
b sulphate 6,000 u per ml	4.50	5 ml OP	<ul><li>Maxitrol</li></ul>
FLUOROMETHOLONE			
* Eye drops 0.1%	3.09	5 ml OP	✓ FML
	5.20		✓ Flucon
LEVOCABASTINE			
Eye drops 0.5 mg per ml	8.71	4 ml OP	
	(10.34)		Livostin

	Subsidy (Manufacturer's Pr	ice) Subs	Fully sidised	Brand or Generic Manufacturer
LODOXAMIDE Eye drops 0.1%	<del></del>	10 ml OP	✓ L	.omide
NEPAFENAC Eye drops 0.3%	8.80	3 ml OP	<b>✓</b>	levro
PREDNISOLONE ACETATE Eye drops 1%	6.92 7.00	10 ml OP 5 ml OP		Prednisolone-AFT Pred Forte
PREDNISOLONE SODIUM PHOSPHATE – Special Authority se Eye drops 0.5%, single dose (preservative free)		– Retail pharr 20 dose	•	<i>l</i> linims Prednisolone

### **⇒SA1715** Special Authority for Subsidy

**Initial application** only from an ophthalmologist or optometrist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has severe inflammation; and
- 2 Patient has a confirmed allergic reaction to preservative in eye drops.

**Renewal** from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

SODIUM CROMOGLICATE			
Eye drops 2%	2.62	10 ml OP	✓ <u>Allerfix</u>

Glaucoma	Preparations - Beta Blockers	
----------	------------------------------	--

# Eye drops 0.25%	5 ml OP 5 ml OP	✓ Betoptic S ✓ Betoptic
TIMOLOL	5 ml OP	✓ Arrow-Timolol

* Tab 250 mg	100	✓ Diamox
BRINZOLAMIDE  * Eye drops 1%5.11	5 ml OP	✓ Azopt
DORZOLAMIDE WITH TIMOLOL  * Eye drops 2% with timolol 0.5%	5 ml OP	✓ Dortimopt

# Glaucoma Preparations - Prostaglandin Analogues

וווט	MATOFROST			
*	Eye drops 0.03%	5.15	3 ml OP	<ul><li>Lumigan</li></ul>
	,	5.95		✓ Bimatoprost
				Multichem

Lumigan to be Principal Supply on 1 January 2025 (Bimatoprost Multichem Eye drops 0.03% to be delisted 1 January 2025)

5 ml OP

✓ Arrow-Timolol

ACETAZOL AMIDE

DIMATORROST

				_
	Subsidy (Manufacturer's Pri	ce) Subs	Fully sidised	Brand or Generic Manufacturer
LATANOPROST  * Eye drops 0.005%  Teva to be Principal Supply on 1 March 2025	2.08	2.5 ml OP	<b>✓</b> Te	eva
TRAVOPROST   * Eye drops 0.004%	6.80	2.5 ml OP	<b>✓</b> <u>Tr</u>	ravatan_
Glaucoma Preparations - Other				
# Eye drops 0.2%		5 ml OP	<b>✓</b> Aı	rrow-Brimonidine
BRIMONIDINE TARTRATE WITH TIMOLOL MALEATE  * Eye drops 0.2% with timolol maleate 0.5%	7.13	5 ml OP	✓ <u>C</u>	ombigan_
LATANOPROST WITH TIMOLOL  * Eye drops 0.005% with timolol 0.5%	4.95	2.5 ml OP	✓ <u>Aı</u>	rrow - Lattim
PILOCARPINE HYDROCHLORIDE  * Eye drops 1%  * Eye drops 2%  * Eye drops 4%  Subsidised for oral use pursuant to the Standard Formul	5.35 7.99	15 ml OP 15 ml OP 15 ml OP	✓ Is	opto Carpine opto Carpine opto Carpine
PILOCARPINE NITRATE  * Eye drops 2% single dose – Special Authority see SA0895 below – Retail pharmacy	35.90	20 dose	✓ M	inims Pilocarpine

**⇒SA0895** Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient has to use an unpreserved solution due to an allergy to the preservative; or
- 2 Patient wears soft contact lenses.

Note: Minims for a general practice are considered to be "tools of trade" and are not approved as special authority items. **Renewal** from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Mydriatics and Cycloplegics		
ATROPINE SULPHATE		
* Eye drops 1%	15 ml OP	✓ <u>Atropt</u>
CYCLOPENTOLATE HYDROCHLORIDE		
* Eye drops 1%	15 ml OP	✓ Cyclogyl
* Eye drops 1%, single dose (preservative free) – Only on a	00 doos	✓ Minims
prescription84.85	20 dose	Cyclopentolate
TROPICAMIDE		
* Eye drops 0.5%	15 ml OP	✓ Mydriacyl
* Eye drops 1%8.66	15 ml OP	✓ Mydriacyl
Preparations for Tear Deficiency		
For acetylcysteine eye drops refer Standard Formulae, page 276 HYPROMELLOSE		

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

\*Three months or six months, as applicable, dispensed all-at-once

✓ Methopt

15 ml OP



	Subsidy (Manufacturer's Price)		Fully idised	Brand or Generic
	\$	Per	1	Manufacturer
HYPROMELLOSE WITH DEXTRAN  * Eye drops 0.3% with dextran 0.1%	2.30	15 ml OP	<b>✓</b> P	oly-Tears

#### **Preservative Free Ocular Lubricants**

#### ⇒SA2134 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 Confirmed diagnosis by slit lamp or Schirmer test of severe secretory dry eye; and
- 2 Either:
  - 2.1 Patient is using eye drops more than four times daily on a regular basis; or
  - 2.2 Patient has had a confirmed allergic reaction to preservative in eye drop.

**Renewal** from any relevant practitioner. Approvals valid for 24 months where the patient continues to require lubricating eye drops and has benefited from treatment.

# Other Eve Preparations

NAPHAZOLINE HYDROCHLORIDE  * Eye drops 0.1%4.	15 15 ml OP	✓ Naphcon Forte
, ·	65	✓ Albalon
Albalon to be Principal Supply on 1 January 2025		
(Naphcon Forte Eye drops 0.1% to be delisted 1 January 2025)		
OLOPATADINE		
Eye drops 0.1%2.	17 5 ml OP	✓ Olopatadine Teva
PARAFFIN LIQUID WITH WOOL FAT		
* Eye oint 3% with wool fat 3%	63 3.5 g OP	✓ Poly-Visc
RETINOL PALMITATE	•	-
Eye oint 138 mcg per g	80 5 g OP	✓ VitA-POS

					VARIOUS
	(Manu	Subsidy sfacturer's Price)	Sı Per	Fully ubsidised	Brand or Generic Manufacturer
٧	arious				
PH *	ARMACY SERVICES Brand switch fee	4.50	1 fee	✓ E	Midazolam-Baxter BSF Norethinderone
* * *	a) May only be claimed once per patient. b) The Pharmacode for BSF Alyacen is 2692112 - see also per c) The Pharmacode for BSF Norethinderone - CDC is 269212 d) The Pharmacode for BSF Continuous glucose monitor (state) The Pharmacode for BSF Continuous glucose monitor (interpretable) of the Pharmacode for BSF Midazolam-Baxter is 2695863 - sometime in the Pharmacode for BSF Midazolam-Baxter is 2695863 - sometime in the Pharmacode for BSF Midazolam-Baxter is 2695863 - sometime in the Pharmacode for BSF Midazolam-Baxter is 2695863 - sometime in the Pharmacode for BSF Midazolam-Baxter is 2695863 - sometime in the Pharmacode for BSF Midazolam-Baxter is 2695863 - sometime in the Pharmacode for BSF Midazolam-Baxter is 2695863 - sometime in the Pharmacode for BSF Norethinderone - CDC is 269212 d).	0 - see also pandalo is 269213 erope is 269214 ee also page 1 0.00 0.00	39 - see 17 - see	also page	

(BSF Continuous glucose monitor (interope Brand switch fee to be delisted 1 January 2025)

(BSF Continuous glucose monitor (standalo Brand switch fee to be delisted 1 January 2025)

(BSF Midazolam-Baxter Brand switch fee to be delisted 1 February 2025)

(BSF Norethinderone - CDC Brand switch fee to be delisted 1 January 2025)

# **Agents Used in the Treatment of Poisonings**

### **Antidotes**

ACETYL CYSTEINE

Inj 200 mg per ml, 10 ml ampoule	42.99	10	✓ DBL Acetylcysteine
	52.88		Martindale Pharma
(Martindale Pharma Inj 200 mg per ml, 10 ml ampoule to be delis	ted 1 April 2025)		
NALOXONE HYDROCHLORIDE			
a) Up to 10 inj available on a PSO			
b) Only on a PSO			
* Inj 400 mcg per ml, 1 ml ampoule	13.29	5	DBL Naloxone
,			Hydrochloride
	35.26	10	✓ Hameln

(Hameln Inj 400 mcg per ml, 1 ml ampoule to be delisted 1 April 2025)



VARIOUS				
	Subsidy (Manufacturer's F \$	Price) Subs	Fully sidised	Brand or Generic Manufacturer
Removal and Elimination				
CHARCOAL  * Oral liq 50 g per 250 ml  a) Up to 250 ml available on a PSO  b) Only on a PSO	43.50	250 ml OP	✓ C	arbosorb-X
DEFERASIROX – Special Authority see SA1492 below – Retail   Wastage claimable Tab 125 mg dispersible Tab 250 mg dispersible	276.00 552.00	28 28	✓ Ex	xjade xjade
Tab 500 mg dispersible	·	28 dications meeting		xjade  ollowing criteria:
<ul> <li>All of the following: <ol> <li>The patient has been diagnosed with chronic iron overload</li> <li>Deferasirox is to be given at a daily dose not exceeding 40</li> <li>Any of the following: <ol> <li>Treatment with maximum tolerated doses of defering combination therapy have proven ineffective as me</li> <li>Treatment with deferiprone has resulted in severely</li> <li>Treatment with deferiprone is contraindicated due to count (ANC) of &lt; 0.5 cells per μL) or recurrent epis 0.5 - 1.0 cells per μL).</li> </ol> </li> <li>Renewal only from a haematologist. Approvals valid for 2 years Either: <ol> <li>For the first renewal following 2 years of therapy, the treat improvement in all three parameters namely serum ferriting.</li> <li>For subsequent renewals, the treatment has been tolerated in all three parameters namely serum ferritin, cardiac MRI</li> </ol> </li> </ol></li></ul>	o mg/kg/day; and one monothers easured by serur persistent vomitics or to a history of accodes (greater the for applications ment has been to a cardiac MRI T2 dand has result T2* and liver MI	apy or deferipron ferritin levels, ing or diarrhoea granulocytosis (an 2 episodes) meeting the follolerated and have and liver MR ted in clinical st	one and liver or a; or defined of mode lowing c as result I T2* lev	desferrioxamine cardiac MRI T2*; or as an absolute neutrophil erate neutropenia (ANC riteria: ted in clinical vels; or
DEFERIPRONE - Special Authority see SA1480 below - Retail Tab 500 mg Oral liq 100 mg per 1 ml  SA1480 Special Authority for Subsidy	533.17	100 250 ml OP		erriprox erriprox
Initial application only from a haematologist. Approvals valid wi following criteria:  Either:  1 The patient has been diagnosed with chronic iron overload 2 The patient has been diagnosed with chronic iron overload.	d due to congeni	ital inherited an	aemia;	
DESFERRIOXAMINE MESILATE  * Inj 500 mg vial	•	10	<b>✓</b> D	BL Desferrioxamine Mesylate for Inj BP

✓ Deferoxamine Pfizer S29 \$29



	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
SODIUM CALCIUM EDETATE  * Inj 200 mg per ml, 5 ml	53.31 (156.71)	6	С	Calcium Disodium Versenate

# **Standard Formulae**

Otaliaala i Olillalac			
ACETYLCYSTEINE EYE DROPS Acetylcysteine inj 200 mg per ml, 10 ml	qs	PHENOBARBITONE SODIUM PAEDIATRIC ORAL mg per ml)	LIQUID (10
Suitable eye drop base	qs .	Phenobarbitone Sodium	400 mg
, ,	·	Glycerol BP	4 ml
CODEINE LINCTUS (3 mg per 5 ml)		Water	to 40 ml
Codeine phosphate	60 mg		
Glycerol	40 ml	PILOCARPINE ORAL LIQUID	
Preservative	qs	Pilocarpine 4% eye drops	qs
Water	to 100 ml	Preservative	qs
CODEINE LINOTHO (45 5)		Water	to 500 ml
CODEINE LINCTUS (15 mg per 5 ml)	000	(Preservative should be used if quantity supplied is	for more
Codeine phosphate	300 mg	than 5 days.)	
Glycerol	40 ml	SALIVA SUBSTITUTE FORMULA	
Preservative	qs		F ~
Water	to 100 ml	Methylcellulose Preservative	5 g
FOLINIC MOUTHWASH		Water	qs to 500 ml
Calcium folinate 15 mg tab	1 tab		
Preservative	qs	(Preservative should be used if quantity supplied is than 5 days. Maximum 500 ml per prescription.)	ior more
Water	to 500 ml	man 5 days. Maximum 500 mi per prescription.)	
(Preservative should be used if quantity supplied is		SODIUM CHLORIDE ORAL LIQUID	
than 5 days. Maximum 500 ml per prescription.)	ioi illoic	Sodium chloride inj 23.4%, 20 ml	qs
than 5 days. Maximum 500 mi per prescription.		Water	qs
METHYL HYDROXYBENZOATE 10% SOLUTION		(Only funded if prescribed for treatment of hyponatr	
Methyl hydroxybenzoate	10 g		,
Propylene glycol	to 100 ml	VANCOMYCIN ORAL SOLUTION (25 mg per ml)	
(Use 1 ml of the 10% solution per 100 ml of oral liqu	id mixture)	Vancomycin 500 mg injection	5 vials
OMEDDAZOLE OLIODENIOLONI		Glycerin with sucrose suspension	37.5 ml
OMEPRAZOLE SUSPENSION		Water	to 100 ml
Omeprazole capsules or powder	qs	(Only funded if prescribed for treatment of Clostridiu	ım difficile
Sodium bicarbonate powder BP	8.4 g	following metronidazole failure)	
Water	to 100 ml		
PHENOBARBITONE ORAL LIQUID			
Phenobarbitone Sodium	1 g		
Glycerol BP	70 ml		

to 100 ml

Water

# EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic

Per Manufacturer

# **Extemporaneously Compounded Preparations and Galenicals**

CODEINE PHOSPHATE – Safety medicine; prescriber may determin	e dispensina f	requency	
Powder – Only in combination		25 g	
	(90.09)		Douglas
Only in extemporaneously compounded codeine linctus.			
COLLODION FLEXIBLE			
Note: This product is no longer being manufactured by the suppli determined.	er and will be	delisted from 1	ine Schedule at a date to be
Collodion flexible	19.30	100 ml	✓ PSM
COMPOUND HYDROXYBENZOATE – Only in combination	10.00	100 1111	
Only in extemporaneously compounded oral mixtures.			
Soln	30.00	100 ml	✓ Midwest
GLYCERIN WITH SODIUM SACCHARIN - Only in combination			
Only in combination with Ora-Plus or when used in the vancomyc	in oral Iquuid S	Standard Form	nulae.
Suspension	30.95	473 ml	✓ Ora-Sweet SF
GLYCERIN WITH SUCROSE - Only in combination			
Only in combination with Ora-Plus or when used in the vancomyc	in oral Iquuid S		
Suspension	30.95	473 ml	✓ Ora-Sweet
GLYCEROL			•
* Liquid – Only in combination		500 ml	✓ healthE Glycerol BP
Only in extemporaneously compounded oral liquid preparatio	ns.		
METHYL HYDROXYBENZOATE	0.00	05	/ Mishanat
Powder	8.98	25 g	✓ Midwest
METHYLCELLULOSE Powder	26.05	100 ~	✓ MidWoot
Suspension – Only in combination		100 g 473 ml	<ul><li>✓ MidWest</li><li>✓ Ora-Plus</li></ul>
METHYLCELLULOSE WITH GLYCERIN AND SODIUM SACCHARIN			· Old-I lub
Suspension		473 ml	✓ Ora-Blend SF
METHYLCELLULOSE WITH GLYCERIN AND SUCROSE - Only in a		4701111	ora bicina or
Suspension		473 ml	✓ Ora-Blend
PHENOBARBITONE SODIUM			0.0 2.0
Powder – Only in combination	52.50	10 g	✓ MidWest
	325.00	100 g	✓ MidWest
Only in children up to 12 years		-	
PROPYLENE GLYCOL			
Only in extemporaneously compounded methyl hydroxybenzoate			_
Liq	11.25	500 ml	✓ Midwest
SODIUM BICARBONATE			•
Powder BP – Only in combination		500 g	✓ Midwest
Only in extemporaneously compounded omeprazole and lans	soprazoie susp	ension.	
SYRUP (PHARMACEUTICAL GRADE) – Only in combination			
Only in extemporaneously compounded oral liquid preparations.  Liq	14 95	500 ml	✓ Midwest
•	I <del>T</del> J	JUU IIII	- WIIWWEST
WATER Tap – Only in combination	0.00	1 ml	✓ Tap water
rap Only in combination	0.00	1 1111	- Iup water

Subsidy (Manufacturer's Price)

Fully Subsidised Per

Brand or Generic Manufacturer

# **Nutrient Modules**

### Carbohydrate

#### ⇒SA1930 Special Authority for Subsidy

Initial application — (Cystic fibrosis or kidney disease) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria: Fither:

- 1 cystic fibrosis; or
- 2 chronic kidney disease.

Initial application — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 cancer in children: or
- 2 cancers affecting alimentary tract where there are malabsorption problems in patients over the age of 20 years; or
- 3 faltering growth in an infant/child; or
- 4 bronchopulmonary dysplasia; or
- 5 premature and post premature infant; or
- 6 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified where the patient has inborn errors of metabolism. Renewal — (Cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

CARBOHYDRATE SUPPLEMENT - Special Authority see SA1930 above - Hospital pharmacy [HP3]

400 a OP ✓ Polycal Powder .......6.72

# Carbohydrate And Fat

# ⇒SA1376 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria: Both:

Subsidy		Fully	Brand or	
(Manufacturer's Pri	ce)	Subsidised	Generic	
\$	Per	•	Manufacturer	

- 1 Infant or child aged four years or under; and
- 2 cystic fibrosis.

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 infant or child aged four years or under; and
- 2 Any of the following:
  - 2.1 cancer in children; or
  - 2.2 faltering growth; or
  - 2.3 bronchopulmonary dysplasia; or
  - 2.4 premature and post premature infants.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

( )		3 -	Soluble Powder	
Powder (neutral)	71.77	400 a OP •	<ul> <li>Duocal Super</li> </ul>	
CARBOHYDRATE AND FAT SUPPLEMENT - Spec	cial Authority see SA1376 o	on the previous page	– Hospital pharmacy [F	1P3]

#### Fat

# **⇒SA2204** Special Authority for Subsidy

Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified where the patient has an inborn error of metabolism.

Initial application — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 faltering growth in an infant/child; or
- 2 bronchopulmonary dysplasia; or
- 3 fat malabsorption; or
- 4 lymphangiectasia; or
- 5 short bowel syndrome: or
- 6 infants with necrotising enterocolitis; or
- 7 biliary atresia; or
- 8 for use in a ketogenic diet; or
- 9 chyle leak; or

continued...

✓ fully subsidised 279



Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	•	Manufacturer

- 10 ascites: or
- 11 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. Renewal — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT SUPPLEMENT	<ul> <li>Special Authority see SA2204 on the previous</li> </ul>	page – H	ospital pharmacy [	[HP3]
	1)	4 = 00		

Emulsion (neutral)		
38.44	500 ml OP	✓ Calogen
Emulsion (strawberry)15.38	200 ml OP	<ul><li>Calogen</li></ul>
Oil	500 ml OP	<ul><li>MCT oil (Nutricia)</li></ul>
MCT Emulsion, 250 ml143.65	4 OP	<ul><li>Liquigen</li></ul>

#### **Protein**

#### ⇒SA1524 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 protein losing enteropathy; or
- 2 high protein needs; or
- 3 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

	armacy [HP3]	MENT - Special Authority see SA1524 above - Hospital pha	PROTEIN SUPPLEMENT
✓ Resource	227 g OP	8.95	Powder
Beneprotein			
✓ Protifar	225 a OP	13.82	

Subsidy (Manufacturer's Price) \$

Subsidised Per

Fully

Brand or Generic Manufacturer

### **Oral and Enteral Feeds**

#### **Diabetic Products**

#### ⇒SA1095 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is a type I or and II diabetic who is suffering weight loss and malnutrition that requires nutritional support. Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

DIABETIC ENTERAL FEED 1KCAL/ML - Special Authority see	SA1095 above -	Hospital pharm	acy [HP3]
Liquid	4.65	500 ml OP	Glucerna Select
DIABETIC ORAL FEED 1KCAL/ML - Special Authority see SA	1095 above – Hos	pital pharmacy	[HP3]
Liquid (strawberry)	2.25	200 ml OP	✓ Diasip
Liquid (vanilla)	2.10	200 ml OP	✓ Nutren Diabetes
, , ,	2.25		Diasip

### **Fat Modified Products**

### ⇒SA2205 Special Authority for Subsidy

Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified where the patient has an inborn error of metabolism. Initial application — (Indications other than errors of inborn metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Patient has a chyle leak; or
- 2 Modified as a modular feed, made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule, for adults.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

#### Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

✓ fully subsidised 281

Subsidy (Manufacturer's Price)

Fully Subsidised Brand or Generic Manufacturer

### Paediatric Products For Children Awaiting Liver Transplant

#### ⇒SA1098 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) who requires a liver transplant.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

#### Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL FEED 1KCAL/ML - Special Authority see SA1098 above - Hospital pharmacy (HP3)

400 g OP ✓ Heparon Junior 

### Paediatric Products For Children With Chronic Renal Failure

# **⇒SA1099** Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 vears where the patient is a child (up to 18 years) with acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

#### Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL FEED 1KCAL/ML - Special Authority see SA1099 above - Hospital pharmacy [HP3] Powder .......64.26 400 a OP ✓ Kindergen

#### Paediatric Products

### ⇒SA1379 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

#### Both:

- 1 Child is aged one to ten years; and
- 2 Any of the following:
  - 2.1 the child is being fed via a tube or a tube is to be inserted for the purposes of feeding; or
  - 2.2 any condition causing malabsorption; or
  - 2.3 faltering growth in an infant/child; or
  - 2.4 increased nutritional requirements; or
  - 2.5 the child is being transitioned from TPN or tube feeding to oral feeding.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for

			SI	PECIAL FOODS
	Subsidy (Manufacturer's Price \$	e) Subs Per	Fully idised	Brand or Generic Manufacturer
continued applications meeting the following criteria: Both:  1 The treatment remains appropriate and the patient is ber 2 General Practitioners must include the name of the dietiti practitioner and date contacted.			nally re	egistered general
PAEDIATRIC ENTERAL FEED 1.5KCAL/ML - Special Authorit Liquid		e previous pa 500 ml OP	<b>✓</b> F	Hospital pharmacy [HP3] Frebini Energy Jutrini Energy RTH
PAEDIATRIC ENTERAL FEED 1KCAL/ML - Special Authority : Liquid		previous pag 500 ml OP	✓ F	spital pharmacy [HP3] Pediasure RTH Jutrini RTH Frebini Original
PAEDIATRIC ENTERAL FEED WITH FIBRE 1.5KCAL/ML - Sp. pharmacy [HP3]	pecial Authority see \$	SA1379 on t	he prev	vious page – Hospital
Liquíd	7.00 5 7.14	500 ml OP		rebini Energy Fibre Nutrini Energy Multi Fibre
PAEDIATRIC ENTERAL FEED WITH FIBRE 1KCAL/ML - Sperpharmacy [HP3]	cial Authority see SA	1379 on the	previo	ous page – Hospital
Liquid	7.00 5	600 ml OP	<b>√</b> F	rebini Original

PAEDIATRIC ORAL FEED 1.5KCAL/ML - Special Authority	y see SA1379 on the	previous page -	- Hospital pharmacy [HP3]
Liquid (strawberry)	1.90	200 ml OP	✓ Fortini
Liquid (vanilla)	1.90	200 ml OP	✓ Fortini
	8.67	500 ml OP	✓ Pediasure Plus
PAEDIATRIC ORAL FEED 1KCAL/ML - Special Authority s	see SA1379 on the pr	evious page – H	Hospital pharmacy [HP3]
Liquid (chocolate)	1.33	200 ml OP	✓ Pediasure

PAEDIATRIC ORAL FEED WITH FIBRE 1.5KCAL/ML	- Special Authority see SA1379 on the previous page - Hospital
pharmacy [HP3]	

Liquid (strawberry)......1.33

Liquid (unflavoured)	1.90	200 ml OP	✓ Fortini Multi Fibre
Liquid (chocolate)	1.90	200 ml OP	✓ Fortini Multi Fibre
Liquid (strawberry)	1.90	200 ml OP	✓ Fortini Multi Fibre
Liquid (vanilla)	1.90	200 ml OP	✓ Fortini Multi Fibre

PEPTIDE-BASED ORAL FEED - Special Authority see SA1379 on the prev	vious page -	Hospital p	harmacy [HP3]
Powder	60 400	a OP	<ul> <li>Peptamen Junior</li> </ul>

### **Renal Products**

### ⇒SA1101 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the

continued...

Fibre

✓ Pediasure

✓ Pediasure

✓ Pediasure

200 ml OP

200 ml OP

250 ml OP

✓ fully subsidised 283

Subsidy	)	Fully	Brand or
(Manufacturer's Price		Subsidised	Generic
	Per	✓	Manufacturer

recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

#### Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

RENAL ORAL FEED 1.8 KCAL/ML - Special Authority see SA1101 on the previous page - Hospital pharmacy [HP3] ✓ Nepro HP 220 ml OP (strawberry) ✓ Nepro HP (vanilla) RENAL ORAL FEED 2 KCAL/ML - Special Authority see SA1101 on the previous page - Hospital pharmacy [HP3] 4 OP ✓ NovaSource Renal 4 OP ✓ Renilon 7.5 4 OP ✓ Renilon 7.5 

### **Specialised And Elemental Products**

### ⇒SA1377 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 malabsorption: or
- 2 short bowel syndrome; or
- 3 enterocutaneous fistulas: or
- 4 eosinophilic oesophagitis; or
- 5 inflammatory bowel disease: or
- 6 patients with multiple food allergies requiring enteral feeding.

Notes: Each of these products is highly specialised and would be prescribed only by an expert for a specific disorder. The alternative is hospitalisation.

Elemental 028 Extra is more expensive than other products listed in this section and should only be used where the alternatives have been tried first and/or are unsuitable.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

#### Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL SEMI-ELEMENTAL FEED 1.5KCAL/ML - Sp Liquid				
ORAL ELEMENTAL FEED 0.8KCAL/ML - Special Authority se	e SA1377 above	– Hospital phar	macy [HP3]	
Liquid (grapefruit), 250 ml carton	179.46	18 OP	✓ Elemental 028 Extra	
Liquid (pineapple & orange), 250 ml carton	179.46	18 OP	✓ Elemental 028 Extra	
Liquid (summer fruits), 250 ml carton		18 OP	✓ Elemental 028 Extra	
ORAL ELEMENTAL FEED 1KCAL/ML - Special Authority see SA1377 above - Hospital pharmacy [HP3]				
Powder (unflavoured)	4.50	80 a OP	✓ Vivonex TEN	

	Subsidy (Manufacturer's Price \$	e) Subs Per	Fully idised	Brand or Generic Manufacturer
SEMI-ELEMENTAL ENTERAL FEED 1KCAL/ML – Special Autl [HP3] Liquid	·	n the previo	✓ N	lutrison Advanced
	9.60			Peptisorb Survimed OPD

### Paediatric Products For Children With Low Energy Requirements

#### ⇒SA1196 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Child aged one to eight years; and
- 2 The child has a low energy requirement but normal protein and micronutrient requirements.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

# Standard Supplements

#### ⇒SA1859 Special Authority for Subsidy

Initial application — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age; and
- 2 Any of the following:
  - 2.1 The patient has a condition causing malabsorption; or
  - 2.2 The patient has failure to thrive; or
  - 2.3 The patient has increased nutritional requirements; and
- 3 Nutrition goal has been set (eg reach a specific weight or BMI).

Renewal — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 A nutrition goal has been set (eg reach a specific weight or BMI).

Initial application — (Children - exclusive enteral nutrition for Crohn's disease) only from a gastroenterologist or dietitian on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

continued...

✓ fully subsidised 285

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
 <b>`</b> \$	Per	✓	

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 Dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Renewal — (Children - exclusive enteral nutrition for Crohn's disease) from any relevant practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 General Practitioners and dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Initial application — (Adults) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

1 Any of the following:

Patient is Malnourished

- 1.1 Patient has a body mass index (BMI) of less than 18.5 kg/m<sup>2</sup>; or
- 1.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 1.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months; and
- 2 Any of the following:

Patient has not responded to first-line dietary measures over a 4 week period by:

- 2.1 Increasing their food intake frequency (eg snacks between meals); or
- 2.2 Using high-energy foods (e.g. milkshakes, full fat milk, butter, cream, cheese, sugar etc); or
- 2.3 Using over the counter supplements (e.g. Complan); and
- 3 A nutrition goal has been set (e.g. to reach a specific weight or BMI).

Renewal — (Adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 A nutrition goal has been set (eg reach a specific weight or BMI); and
- 2 Any of the following:

Patient is Malnourished

- 2.1 Patient has a body mass index (BMI) of less than 18.5 kg/m<sup>2</sup>; or
- 2.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 2.3 Patient has a BMI of less than 20 kg/m<sup>2</sup> and unintentional weight loss greater than 5% within the last 3-6 months.

**Initial application — (Short-term medical condition)** from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube or a nasogastric tube is to be inserted for feeding; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Is undergoing a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
  - 5.1 Pregnant; and
  - 5.2 Any of the following:
    - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or

Subsidy	Full	y Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per 🗸	Manufacturer	

- 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
- 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

Renewal — (Short-term medical condition) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Has undergone a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
  - 5.1 Pregnant: and
  - 5.2 Any of the following:
    - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis
    - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
    - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

Initial application — (Long-term medical condition) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis: or
- 3 Liver disease: or
- 4 Chronic Renal failure: or
- 5 Inflammatory bowel disease; or
- 6 Chronic obstructive pulmonary disease with hypercapnia: or
- 7 Short bowel syndrome; or
- 8 Bowel fistula: or
- 9 Severe chronic neurological conditions; or
- 10 Epidermolysis bullosa: or
- 11 AIDS (CD4 count < 200 cells/mm3): or
- 12 Chronic pancreatitis.

Renewal — (Chronic disease OR tube feeding for patients who have previously been funded under Special Authority forms SA0702 or SA0583) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis: or
- 3 Liver disease: or

continued...

287 ✓ fully subsidised

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sı	ubsidised	Generic
\$	Per	✓	Manufacturer

- 4 Chronic Renal failure; or
- 5 Inflammatory bowel disease; or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome; or
- 8 Bowel fistula; or

9 Severe chronic neurological conditions.		
ENTERAL FEED 1.5KCAL/ML - Special Authority see SA1859 on page 285 Liquid2.1' 8.6i	7 250 ml OP	[HP3]  ✓ Ensure Plus HN  ✓ Ensure Plus HN  RTH
9.0 9.6		<ul><li>✓ Nutrison Energy</li><li>✓ Fresubin HP Energy</li></ul>
ENTERAL FEED 1KCAL/ML - Special Authority see SA1859 on page 285 - Liquid	4 250 ml OP 0 1,000 ml OP 6	HP3]  ✓ Isosource Standard  ✓ Fresubin Original  ✓ Osmolite RTH  ✓ Nutrison RTH
ENTERAL FEED WITH FIBRE 0.83 KCAL/ML – Special Authority see SA18 Liquid		spital pharmacy [HP3] Nutrison 800 Complete Multi Fibre
ENTERAL FEED WITH FIBRE 1 KCAL/ML - Special Authority see SA1859 Liquid	6 1,000 ml OP	al pharmacy [HP3]  ✓ Jevity RTH  ✓ Fresubin Original  Fibre
7.2  ENTERAL FEED WITH FIBRE 1.2KCAL/ML - Special Authority see \$A1850 Liquid	9 on page 285 – Hosp	✓ Nutrison Multi Fibre ital pharmacy [HP3] ✓ Jevity Plus RTH
ENTERAL FEED WITH FIBRE 1.5KCAL/ML — Special Authority see SA1850 Liquid	9 on page 285 – Hosp	•
9.8	0	✓ Fresubin HP Energy Fibre
ENTERAL FEED WITH PROTEIN 1.2KCAL/ML - Special Authority see SA1 Liquid9.6		ospital pharmacy [HP3]  Fresubin Intensive
ORAL FEED (POWDER) – Special Authority see SA1859 on page 285 – Ho Powder (chocolate)14.00		] ✓ Sustagen Hospital Formula
Powder (vanilla)14.0		<ul><li>✓ Ensure</li><li>✓ Sustagen Hospital Formula Active</li></ul>
26.0	0 850 g OP	✓ Ensure

	Subsidy	Fully	Brand or
(1)	Manufacturer's Price)	Subsidised	Generic
	\$	Per 🗸	Manufacturer

### ORAL FEED 1.5KCAL/ML - Special Authority see SA1859 on page 285 - Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, who have severe epidermolysis bullosa, or as exclusive enteral nutrition in children under the age of 18 years for the treatment of Crohn's disease, or for patients with COPD and hypercapnia, defined as CO2 value exceeding 55mmHg. The prescription must be endorsed accordingly.

Liquid (banana) - Higher subsidy of up to \$1.76 per 200 ml with Endorsement	0.72	200 ml OP	
	(1.56) (1.76)		Ensure Plus Fortisip
Liquid (chocolate) - Higher subsidy of up to \$1.76 per 200 ml	0.70	000 1 OD	
with Endorsement	0.72 (1.56)	200 ml OP	Ensure Plus
	(1.76)		Fortisip
Liquid (fruit of the forest) – Higher subsidy of \$1.56 per 200 ml with Endorsement	0.72	200 ml OP	
Will Endotsonion	(1.56)	200 1111 01	Ensure Plus
Liquid (strawberry) – Higher subsidy of \$1.76 per 200 ml with	0.70	000   OD	
Endorsement	0.72 (1.76)	200 ml OP	Fortisip
Liquid (vanilla) - Higher subsidy of up to \$1.76 per 200 ml with	,		
Endorsement	0.85 (1.65)	237 ml OP	Ensure Plus
	0.72	200 ml OP	Liisule Flus
	(1.56)		Ensure Plus
	(1.76)		Fortisip

ORAL FEED WITH FIBRE 1.5 KCAL/ML — Special Authority see SA1859 on page 285 — Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, or who have severe epidermolysis bullosa. The prescription must be endorsed accordingly.

Endorsement	0.72	200 ml OP	
	(1.76)		Fortisip Multi Fibre
Liquid (strawberry) - Higher subsidy of \$1.76 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.76)		Fortisip Multi Fibre
Liquid (vanilla) - Higher subsidy of \$1.76 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.76)		Fortisip Multi Fibre

# **High Calorie Products**

### ⇒SA1195 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

All of the following:

- 1 Cystic fibrosis; and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements.

Liquid (chocolate) - Higher subsidy of \$1.76 per 200 ml with

continued...

# **SPECIAL FOODS**

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	1	Manufacturer

continued...

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 any condition causing malabsorption; or
  - 1.2 faltering growth in an infant/child; or
  - 1.3 increased nutritional requirements; or
  - 1.4 fluid restricted: and

practitioner and date contacted.

EDAL EEED OLGOAL AAL

- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements or is fluid restricted.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL FEED 2 KCAL/ML - Special Authority see SA1195 on the previous	page – Hospital p	harmacy [HP3]
Liquid6.50	500 ml OP	✓ Fresubin 2kcal HP
6.82		✓ Nutrison
		Concentrated
13.64	1,000 ml OP	✓ Ensure Two Cal HN
		RTH

ORAL FEED 2 KCAL/ML - Special Authority see SA1195 on the previous page - Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, or who have severe epidermolysis bullosa. The prescription must be endorsed accordingly.

Liquid (vanilla) - Higher subsidy of \$2.34 per 200 ml with

(2.34) Two Cal HN

# **Food Thickeners**

# ⇒SA1106 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient has motor neurone disease with swallowing disorder.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

continued...

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sı	ubsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

### **Gluten Free Foods**

The funding of gluten free foods is no longer being actively managed by Pharmac from 1 April 2011. This means that we are no longer considering the listing of new products, or making subsidy, or other changes to the existing listings. As a result we anticipate that the range of funded items will reduce over time. Management of Coeliac disease with a gluten free diet is necessary for good outcomes. A range of gluten free options are available through retail outlets.

### ⇒SA1729 Special Authority for Subsidy

Initial application — (all patients) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Either:

- 1 Gluten enteropathy has been diagnosed by biopsy; or
- 2 Patient suffers from dermatitis herpetiformis.

Initial application — (paediatric patients diagnosed by ESPGHAN criteria) only from a paediatric gastroenterologist. Approvals valid without further renewal unless notified where the paediatric patient fulfils ESPGHAN criteria for biopsy free diagnosis of coeliac disease.

GLUTEN FREE BAKING MIX – Special Authority see SA1729 Powder		pharmacy [HP3] 1,000 g OP	
	(5.15)		Healtheries Simple Baking Mix
GLUTEN FREE BREAD MIX - Special Authority see SA1729	above – Hospital r	oharmacy [HP3]	
Powder		1,000 g OP	
	(7.32)		NZB Low Gluten Bread Mix
	3.51		
	(10.87)		Horleys Bread Mix
GLUTEN FREE FLOUR - Special Authority see SA1729 abov	e – Hospital pharr	nacy [HP3]	
Powder		2,000 g OP	
	(18.10)	. 3	Horleys Flour

	Subsidy (Manufacturer's Pric	ce) Su Per	Fully bsidised	Brand or Generic Manufacturer
GLUTEN FREE PASTA - Special Authority see SA1729 on the	previous page – Ho	ospital pha	rmacv [H	P31
Buckwheat Spirals		250 g OP	, [	-1
•	(3.11)	ŭ		Orgran
Corn and Vegetable Shells	2.00	250 g OP		
	(2.92)			Orgran
Corn and Vegetable Spirals	2.00	250 g OP		
	(2.92)			Orgran
Rice and Corn Lasagne Sheets	1.60	200 g OP		
	(3.82)		C	Orgran
Rice and Corn Macaroni	2.00	250 g OP		
	(2.92)		C	Orgran
Rice and Corn Penne	2.00	250 g OP		
	(2.92)		C	Orgran
Rice and Maize Pasta Spirals	2.00	250 g OP		
	(2.92)		C	Orgran
Rice and Millet Spirals		250 g OP		
	(3.11)			Orgran
Rice and corn spaghetti noodles		375 g OP		_
	(2.92)		C	Orgran
Vegetable and Rice Spirals		250 g OP	_	
	(2.92)		C	Orgran
Italian long style spaghetti		220 g OP	_	
	(3.11)		(	Orgran

# **Foods And Supplements For Inherited Metabolic Disease**

# ⇒SA2357 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid without further renewal unless notified where patient requires dietary management of inherited metabolic disorders.

# **Supplements For Homocystinuria**

AMINOACID FORMULA WITHOUT METHIONINE	<ul> <li>Special Authority see SA2357</li> </ul>	<sup>7</sup> above – H	lospital pharmacy [HP3]
Powder (neutral), 36 g sachets	750.30	30	✓ HCU Anamix Junior
Powder, 12.5 g sachets	349.65	30	✓ HCU Explore 5
Powder, 25 g sachets	1,048.95	30	✓ HCU Express 15
Powder (neutral), can		500 g OP	✓ XMET Maxamum
Powder (unflavoured), can	260.00	400 g OP	<ul> <li>HCU Anamix Infant</li> </ul>
Liquid (juicy berries), 125 ml bottle	1,684.80	30	✓ HCU Lophlex LQ
Liquid (orange), 125 ml bottle		36	<ul> <li>HCU Anamix Junior</li> </ul>
			LQ

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	/	Manufacturer

# Supplements For MSUD and short chain enoyl coA hydratase deficiency

AMINOACID FORMULA WITHOUT VALINE, LEUCINE AND ISOLEUCINE - Special Authority see \$A2357 on the previous page - Hospital pharmacy [HP3]

Powder (neutral) 36 g sachets	750.00	30	<ul><li>MSUD Anamix Junior</li></ul>
Powder, 12.5 g sachets	349.65	30	✓ MSUD Explore 5
Powder, 25 g sachets		30	✓ MSUD Express 15
Powder (neutral), can		500 g OP	✓ MSUD Maxamum
Powder (orange), can		500 g OP	✓ MSUD Maxamum
Powder (unflavoured), can	260.00	400 g OP	<ul><li>MSUD Anamix Infant</li></ul>
Liquid (orange) 125 ml bottles	941.40	36	<ul><li>MSUD Anamix Junior LQ</li></ul>
Liquid (juicy berries) 125 ml pouches	1,684.80	30	✓ MSUD Lophlex LQ 20

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Generic Manufacturer

# **Supplements For PKU**

AMINOACID FORMULA WITHOUT PHENYLALANINE -			
Tabs		75 OP	✓ Phlexy 10
Powder (Lemon), 34 g sachets		30	✓ PKU Express 20
Powder (Neutral), 12.5 g sachets		30	✓ PKU Explore 5
Powder (Neutral), 34 g sachets		30	✓ PKU Express 20
Powder (Orange), 25 g sachets		30	✓ PKU Explore 10
Powder (Orange), 34 g sachets	883.50	30	✓ PKU Express 20
Powder (Raspberry), 25 g sachets	441.75	30	✓ PKU Explore 10
Powder (Tropical), 34 g sachets	883.50	30	✓ PKU Express 20
Powder (berry) 28 g sachets	936.00	30	✓ PKU Lophlex
			Powder
Powder (chocolate) 36 g sachet	393.00	30	✓ PKU Anamix Junior
(- · · · · · · · · · · · · · ·			Chocolate
Powder (neutral) 28 g sachets	936 00	30	✓ PKU Lophlex
1 owder (nedital) 20 g sacriets	950.00	30	Powder
Daviday (navityal) OC y analasta	000.00	00	
Powder (neutral) 36 g sachets		30	✓ PKU Anamix Junior
Powder (orange) 28 g sachets	936.00	30	✓ PKU Lophlex
			Powder
Powder (orange) 36 g sachet	393.00	30	PKU Anamix Junior
			Orange
Powder (unflavoured) 12.5 g sachets	234.00	30	✓ PKU First Spoon
Powder (vanilla) 36 g sachet	393.00	30	PKU Anamix Junior
			Vanilla
Infant formula	174.72	400 g OP	✓ PKU Anamix Infant
Powder (orange)	320.00	500 g OP	✓ XP Maxamum
Powder (unflavoured)		500 g OP	✓ XP Maxamum
Liquid (berry)		125 ml OP	✓ PKU Anamix Junior
_ 1 ( )/			LQ
Liquid (orange)	13 10	125 ml OP	✓ PKU Anamix Junior
Liquid (orange)		123 1111 01	LQ
Lieurial (coeffee consel)	10.10	105 ml OD	
Liquid (unflavoured)	13.10	125 ml OP	✓ PKU Anamix Junior
			LQ.
Liquid (forest berries), 250 ml carton		18 OP	<ul><li>Easiphen Liquid</li></ul>
Liquid (juicy tropical) 125 ml		30 OP	✓ PKU Lophlex LQ 20
Oral semi-solid (berries) 109 g	1,123.20	36 OP	✓ PKU Lophlex
			Sensation 20
Powder (neutral), 400 g can	715.16	4 OP	✓ PKU Start
Liquid (juicy berries) 62.5 ml		60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy citrus) 62.5 ml	939.00	60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy orange) 62.5 ml	939.00	60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy berries) 125 ml		30 OP	✓ PKU Lophlex LQ 20
Liquid (juicy orange) 125 ml		30 OP	✓ PKU Lophlex LQ 20

(PKU Anamix Junior LQ Liquid (unflavoured) to be delisted 1 January 2025) (PKU Lophlex LQ 10 Liquid (juicy citrus) 62.5 ml to be delisted 1 January 2025) (PKU Lophlex LQ 10 Liquid (juicy orange) 62.5 ml to be delisted 1 January 2025)

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	d Generic
GLYCOMACROPEPTIDE AND AMINO ACID CONTAINS SOME	PHENYLALANINE	- Special Author	prity see SA2357 on
page 292 – Hospital pharmacy [HP3]			
Powder (Banana) 35 g sachets	930.00	30	PKU
			sphere20 Banana
Powder (Berry), 20 g sachets	449.28	60	PKU Restore
			Powder
Powder (Chocolate) 32 g sachets	898.56	30	PKU Build
			20 Chocolate
Powder (Chocolate) 35 g sachets	930.00	30	PKU
			sphere20 Chocolate
Develop (Leman) OF a cochete	000.00	00	DIZU
Powder (Lemon) 35 g sachets	930.00	30	PKU
2			sphere20 Lemon
Powder (Lemonade) 33.4 g sachets	936.00	30	PKU GMPro Ultra
		_	Lemonade
Powder (Neutral), 15 g sachets			PKU Build 10
Powder (Orange), 20 g sachets	449.28	60	PKU Restore
			Powder
Powder (Raspberry Lemonade) 31 g sachets	898.56	30	PKU Build
			20 Raspberry
			Lemonade
Powder (Smooth) 31 g sachets	898.56	30	PKU Build
			20 Smooth
Powder (Vanilla) 33 g sachets	898.56	30	PKU Build 20 Vanilla
Powder (neutral), 40 g sachets	673.92	30	Glytactin Bettermilk
Powder (unflavoured) 12.5 g sachets	468.00	30	PKU GMPro Mix-In
Powder (vanilla) 33.4 g sachets	936.00	30	PKU GMPro Ultra
			Vanilla
Powder (Red Berry) 35 g sachets	930.00	30	PKU sphere20 Red
			Berry
Powder (Vanilla) 35 g sachets	930.00	30	PKU
, , ,			sphere20 Vanilla
Liquid (neutral), 250 ml carton	280.80	18	PKU GMPro LQ
Liquid (original), 250 ml carton		30 OP ✓	PKU Glytactin RTD
4 (			15
Liquid (Coffee Mocha), 250 ml carton	684.45	30 OP ✓	PKU Glytactin RTD
q (00.100 .1100.110/),			15 Lite
Liquid (chocolate), 250 ml carton	684 45	30 OP ✓	PKU Glytactin RTD
Eigaia (onocolato), 200 mi carton		00 Oi	15
Liquid (vanilla), 250 ml carton	684.45	30 OP ✓	PKU Glytactin RTD
Liquid (variilla), 200 IIII Gartoii		00 01	15 Lite
			IJ LIIG

# Foods

LOW PROTEIN BAKING MIX - Special Authority see SA2357 on pag	e 292 –	Hospital pharmacy	[HP3]
Powder	8.55	500 g OP	✓ Loprofin Mix

OW PROTEIN PASTA - Special Authority see SA2357 on pag Animal shapesLasagne		Per	idised Generic  Manufacturer
Animal shapesLasagne			
Animal shapesLasagne		pharmacy (HP3	וס
Lasagne	12 20	paao, [	
· ·	1∠.35	500 g OP	✓ Loprofin
	6.19	250 g OP	✓ Loprofin
Low protein rice pasta	12.39	500 g OP	✓ Loprofin
Macaroni	6.19	250 g OP	✓ Loprofin
Penne	12.39	500 g OP	✓ Loprofin
Spaghetti	12.39	500 g OP	✓ Loprofin
Spirals		500 g OP	✓ Loprofin
Supplements for Tyrosinaemia			
AMINOACID FORMULA WITHOUT PHENYLALANINE AND TY	ROSINE - Speci	al Authority see	e SA2357 on page 292 – Hosp
harmacy [HP3]			_
Powder (Neutral), 12.5 g sachets	349.65	30	✓ TYR Explore 5
Powder (neutral) 36 g sachets	471.00	30	TYR Anamix Junior
Powder, can	260.00	400 g OP	TYR Anamix Infant
Liquid (juicy berries) 125 ml pouches	1,684.80	30	✓ TYR Lophlex LQ 20
Liquid (orange) 125 ml bottle	941.40	36	✓ TYR Anamix Junior
			LQ
GLYCOMACROPEPTIDE AND AMINO ACID CONTAINS SOME	E TYROSINE ANI	D PHENYLALA	NINE – Special Authority see
SA2357 on page 292 – Hospital pharmacy [HP3]			
Powder (Red Berry), 35 g sachets		30	✓ TYR Sphere 20
Powder (Vanilla), 35 g sachets	1,398.60	30	✓ TYR Sphere 20
Supplements for Organic Acidaemias	E TUDEONINE		0 114 11 11 01005
AMINOACID FORMULA WITHOUT ISOLEUCINE, METHIONIN on page 292 – Hospital pharmacy [HP3]	E, THREONINE A	AND VALINE -	- Special Authority see SA2357
Powder, can	260.00	400 g OP	<ul><li>MMA/PA Anamix Infant</li></ul>
AMINOACID FORMULA WITHOUT METHIONINE, THREONINE Hospital pharmacy [HP3]	E AND VALINE -	Special Autho	rity see SA2357 on page 292 -
Powder (neutral), 18 g sachets	750.30	30	<ul><li>MMA/PA Anamix Junior</li></ul>
Powder, 12.5 g sachets	349.65	30	✓ MMA/PA Explore 5
Powder, 25 g sachets		30	✓ MMA/PA Express 15
Supplements for Glutaric Aciduria type 1			
AMINOACID FORMULA WITHOUT LYSINE - Special Authority	see SA2357 on i	nage 292 – Hos	spital pharmacy [HP3]
Powder (neutral), 18 g sachets		30	✓ GA1 Anamix Junior
Powder, 12.5 g sachets		30	✓ GA Explore 5
Powder, r2.5 g sacriets		400 g OP	✓ GA1 Anamix Infant
rowder, cari	200.00	400 g OF	V GAT Allallitz illialit
Supplements for Glycogen Storage Disease			
HIGH AMYLOPECTIN CORN-STARCH - Special Authority see Powder, 60 g sachets		292 – Hospita 30	l pharmacy [HP3]  ✓ Glycosade
Single dose amino acids			
ARGININE - Special Authority see SA2357 on page 292 - Hosp	pital pharmacy [H	P3]	

Subsidy (Manufacturer's Price) \$	S Per	Fully Subsidised	Brand or Generic Manufacturer
CITRULLINE – Special Authority see SA2357 on page 292 – Hospital pharmacy [HP3] Powder, 4 g sachets211.45	30	1	Citrulline1000
SOLEUCINE – Special Authority see SA2357 on page 292 – Hospital pharmacy [HP3] Powder, 4 g sachets141.05	] 30	✓	Isoleucine50
LEUCINE – Special Authority see SA2357 on page 292 – Hospital pharmacy [HP3] Powder, 4 g sachets141.05	30	✓	Leucine100
PHENYLALANINE - Special Authority see SA2357 on page 292 - Hospital pharmacy [ Powder, 4 g sachets141.05	[HP3] 30		Phenylalanine50
TYROSINE - Special Authority see SA2357 on page 292 - Hospital pharmacy [HP3] Powder, 4 g sachets211.45	30	✓	Tyrosine1000
VALINE - Special Authority see SA2357 on page 292 - Hospital pharmacy [HP3] Powder, 4 g sachets141.05	30	1	Valine50
Other Fat Modified Products			
ELEMENTAL FEED WITH HIGH MEDIUM CHAIN TRIGLYCERIDES - Special Authori pharmacy [HP3]	ity se	e SA2357	on page 292 – Hospital
Powder (neutral), 100 g sachets47.01	10	1	Emsogen
Carbohydrate and Fat with added vitamins and minerals			
PROTEIN FREE SUPPLEMENT CONTAINING CARBOHYDRATE, FAT WITH ADDED Authority see SA2357 on page 292 – Hospital pharmacy [HP3]	VITA	AMINS AN	ND MINERALS - Special
	g OF	· •	Energivit
Essential Amino Acids			
ESSENTIAL AMINOACID FORMULA - Special Authority see SA2357 on page 292 - F Powder (neutral), can313.73 200	Hospit Og OF		acy [HP3] Essential Amino Acid Mix
Infant Formulae			

### Infant Formulae

# For Williams Syndrome

# ⇒SA1110 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is an infant suffering from Williams Syndrome and associated hypercalcaemia.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

LOW CALCIUM INFANT FORMULA - Special Authority see SA1110 above - Hospital pharmacy [HP3]
Powder .......46.18 400 g OP ✓ Locasol

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

# **Gastrointestinal and Other Malabsorptive Problems**

AMINO ACID FORMULA - Special Authority see SA2092 Powder		nacy [HP3] 400 g OP	✓ Alfamino
Powder (unflavoured)	55.61	400 g OP	<ul> <li>✓ Alfamino Junior</li> <li>✓ Neocate Gold</li> <li>✓ Neocate Junior</li> <li>Unflavoured</li> </ul>
	65.72		<ul><li>✓ Neocate SYNEO</li><li>✓ Elecare</li><li>✓ Elecare LCP</li></ul>
Powder (vanilla)	55.61	400 g OP	✓ Neocate Junior Vanilla
	65.72		✓ Elecare

⇒SA2092 Special Authority for Subsidy

**Initial application** — (Infants under 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

Α

- 1 History of anaphylaxis to cow's milk protein formula or dairy products; or
- 2 Eosinophilic oesophagitis; or
- 3 Ultra-short gut; or
- 4 Severe Immune deficiency; or
- 5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
- 6 Both
  - 6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
  - 6.2 Either:
    - 6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or 6.2.2 Patient has IgE mediated allergy.

Initial application — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or
  - 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
  - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products; or
  - 2.2 Eosinophilic oesophagitis; or
  - 2.3 Ultra-short gut; or
  - 2.4 Severe Immune deficiency: or
  - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
  - 2.6 Both:
    - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
    - 2.6.2 Either:
      - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval

continued...

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

number: or

2.6.2.2 Patient has IgE mediated allergy.

Renewal — (Infants up to 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 Patient has IgE mediated allergy; and
  - 1.2 All of the following:
    - 1.2.1 Patient remains allergic to cow's milk; and
    - 1.2.2 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy or extensively hydrolysed infant formula has been undertaken; and
    - 1.2.3 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
    - 1.2.4 Amino acid formula is required for a nutritional deficit; and
    - 1.2.5 It has been more than three months from the previous approval; or

#### 2 Both:

- 2.1 Patient has non IgE mediated severe gastrointestinal intolerance (including eosinophilic oesophagitis, ultra-short gut and severe immune deficiency); and
- 2.2 All of the following:
  - 2.2.1 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy, or extensively hydrolysed infant formula has been undertaken; and
    - 2.2.2 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
    - 2.2.3 Amino acid formula is required for a nutritional deficit; and
    - 2.2.4 It has been more than three months from the previous approval.

Renewal — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither
  - 1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or
  - 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
  - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products; or
  - 2.2 Eosinophilic oesophagitis; or
  - 2.3 Ultra-short gut; or
  - 2.4 Severe Immune deficiency; or
  - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
  - 2.6 Both:
    - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or alleroy or malabsorotion; and
    - 2.6.2 Either:
      - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or
      - 2.6.2.2 Patient has IgE mediated allergy.

Initial application — (for patients who have a current funding under Special Authority form SA1557) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 months for applications meeting the following criteria:

continued...

ubsidy cturer's Price) Subs	Fully	Brand or Generic
 \$ Per	•	Manufacturer

continued...

All of the following:

- 1 Patient has a valid Special Authority approval for extensively hydrolysed formula (SA1557); and
- 2 Extensively hydrolysed formula (Aptamil Gold+ Pepti Junior, AllerPro SYNEO 1 and 2) is unable to be supplied at this time; and
- 3 The approval only applies to funded dispensings of Neocate Gold and Neocate Syneo.

Note: This criteria is short term funding to cover an out-of-stock situation on some extensively hydrolysed formula powder funded under Special Authority form SA1557. There is no renewal criteria under this restriction.

ENTERAL LIQUID PEPTIDE FORMULA - Special Authority see SA1953 below - Hospital pharmacy [HP3]

Liquid 1 kcal/ml12.44	500 ml OP	✓ Nutrini Peptisorb
Liquid 1.5 kcal/ml	500 ml OP	✓ Nutrini Peptisorb
		Energy

## ⇒SA1953 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has impaired gastrointestinal function and either cannot tolerate polymeric feeds, or polymeric feeds are unsuitable; and
- 2 Any of the following:
  - 2.1 Severe malabsorption; or
  - 2.2 Short bowel syndrome; or
  - 2.3 Intractable diarrhoea; or
  - 2.4 Biliary atresia; or
  - 2.5 Cholestatic liver diseases causing malabsorption; or
  - 2.6 Cystic fibrosis; or
  - 2.7 Proven fat malabsorption; or
  - 2.8 Severe intestinal motility disorders causing significant malabsorption; or
  - 2.9 Intestinal failure: or
  - 2.10 Both:
    - 2.10.1 The patient is currently receiving funded amino acid formula; and
    - 2.10.2 The patient is to be trialled on, or transitioned to, an enteral liquid peptide formula; and
- 3 Either:
  - 3.1 A semi-elemental or partially hydrolysed powdered feed has been reasonably trialled and considered unsuitable; or
  - 3.2 For step down from intravenous nutrition.

Note: A reasonable trial is defined as a 2-4 week trial.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 An assessment as to whether the patient can be transitioned to a cows milk protein or soy infant formula or extensively hydrolysed formula has been undertaken; and
- 2 The outcome of the assessment is that the patient continues to require an enteral liquid peptide formula; and
- 3 General practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

EXTENSIVELY HYDROLYSED FORMULA – Special A	uthority see SA1557 on the	e next page	<ul><li>Hospital pharmacy [HP3]</li></ul>
Powder	18.10	450 g OP	✓ Pepti-Junior
	36.20	900 g OP	✓ Allerpro Syneo 1
			✓ Allerpro Syneo 2

Subsidy (Manufacturer's Price) \$

Fully Subsidised Per

Brand or Generic Manufacturer

### ⇒SA1557 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 Both:
  - 1.1 Cows milk formula is inappropriate due to severe intolerance or allergy to its protein content; and
  - 12 Fither
    - 1.2.1 Soy milk formula has been reasonably trialled without resolution of symptoms; or
    - 1.2.2 Soy milk formula is considered clinically inappropriate or contraindicated; or
- 2 Severe malabsorption; or
- 3 Short bowel syndrome; or
- 4 Intractable diarrhoea; or
- 5 Biliary atresia; or
- 6 Cholestatic liver diseases causing malsorption; or
- 7 Cystic fibrosis: or
- 8 Proven fat malabsorption; or
- 9 Severe intestinal motility disorders causing significant malabsorption; or
- 10 Intestinal failure; or
- 11 All of the following:
  - 11.1 For step down from Amino Acid Formula: and
  - 11.2 The infant is currently receiving funded amino acid formula; and
  - 11.3 The infant is to be trialled on, or transitioned to, an extensively hydrolysed formula; and
  - 11.4 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

Note: A reasonable trial is defined as a 2-4 week trial, or signs of an immediate IgE mediated allergic reaction.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 An assessment as to whether the infant can be transitioned to a cows milk protein or soy infant formula has been undertaken; and
- 2 The outcome of the assessment is that the infant continues to require an extensively hydrolysed infant formula; and
- 3 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

#### Fluid Restricted

PAEDIATRIC ORAL/ENTERAL FEED 1 KCAL/ML - Special Authority see SA1698 below - Hospital pharmacy [HP3] Liquid.......2.80 125 ml OP ✓ Infatrini

#### ⇒SA1698 Special Authority for Subsidy

**Initial application** only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is fluid restricted or volume intolerant and has been diagnosed with faltering growth; and
- 2 Patient is under the care of a paediatrician or dietitian who has recommended treatment with a high energy infant formula; and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

continued...

# SPECIAL FOODS

Subsidy	:)	Fully	Brand or
(Manufacturer's Price		Subsidised	Generic
<u> </u>	Per	✓	Manufacturer

continued...

Note: "Volume intolerant" patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

Renewal only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient continues to be fluid restricted or volume intolerant and has faltering growth: and
- 2 Patient is under the care of a hospital paediatrician or dietitian who has recommended treatment with a high energy infant formula; and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: 'Volume intolerant' patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

# **Ketogenic Diet**

#### ⇒SA1197 Special Authority for Subsidy

**Initial application** only from a metabolic physician or paediatric neurologist. Approvals valid for 3 months where the patient has intractable epilepsy, pyruvate dehydrogenase deficiency or glucose transported type-1 deficiency and other conditions requiring a ketogenic diet.

**Renewal** only from a metabolic physician or paediatric neurologist. Approvals valid for 2 years where the patient is on a ketogenic diet and the patient is benefiting from the diet.

HIGH FAT LOW CARBOHYDRATE FORMULA - Special Authority see SA1197 above - Retail pharmacy

Powder (unflavoured)36.92	300 g OP	✓ KetoCal 4:1
	_	✓ Ketocal 3:1
Powder (vanilla)36.92	300 g OP	✓ KetoCal 4:1

### SECTION I: NATIONAL IMMUNISATION SCHEDULE

Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

Subsidy

# **Vaccinations**

BACILLUS CALMETTE-GUERIN VACCINE - [Xpharm]

For infants at increased risk of tuberculosis. Increased risk is defined as:

- 1) living in a house or family with a person with current or past history of TB; or
- 2) having one or more household members or carers who within the last 5 years lived in a country with a rate of TB > or egual to 40 per 100,000 for 6 months or longer; or
- 3) during their first 5 years will be living 3 months or longer in a country with a rate of TB > or equal to 40 per 100,000 Note a list of countries with high rates of TB are available at www.health.govt.nz/tuberculosis (search for downloads) or www.bcgatlas.org/index.php.

Inj Mycobacterium bovis BCG (Bacillus Calmette-Guerin),

Danish strain 1331, live attenuated, vial with diluent...............................0.00 10 ✓ BCG Vaccine AJV

COVID-19 VACCINE - [Xpharm]

Inj 10 mcg raxtozinameran per 0.3 ml, 0.48 ml vial; paediatric ✓ Comirnaty Omicron 10 (XBB.1.5)

Either:

- 1) One dose for previously unvaccinated children aged 5-11 years old; or
- 2) Up to three doses for immunocompromised children aged 5-11 years old.
- Inj 3 mcg raxtozinameran per 0.2 ml, 0.4 ml vial; infant vaccine, 10 ✓ Comirnaty Omicron (XBB.1.5)

Up to three doses for previously unvaccinated children aged 6 months - 4 years at high risk of severe illness.

Inj 30 mcg raxtozinameran per 0.3 ml, 0.48 ml vial; adult vaccine, light grey cap......0.00

✓ Comirnaty Omicron

(XBB.1.5)

Any of the following:

- 1) One dose for previously unvaccinated people aged 12-15 years old; or
- 2) Up to three doses for immunocompromised people aged 12-15 years old; or
- 3) Up to two doses for previously unvaccinated people 16-29 years old; or
- 4) Up to four doses for people aged 16-29 at high risk of severe illness; or
- 5) One dose for previously unvaccinated people aged 30 and older; or
- 6) One additional dose every 6 months for previously vaccinated people aged 30 years and over additional dose is given at least 6 months after last dose.

Ini 30 mcg raxtozinameran per 0.3 ml. 2.25 ml vial: adult

vaccine, dark grey cap......0.00 10 **Comirnaty Omicron** (XBB.1.5)

Any of the following:

- 1) One dose for previously unvaccinated people aged 12-15 years old; or
- 2) Up to three doses for immunocompromised people aged 12-15 years old; or
- 3) Up to two doses for previously unvaccinated people 16-29 years old; or
- 4) Up to four doses for people aged 16-29 at high risk of severe illness; or
- 5) One dose for previously unvaccinated people aged 30 and older; or
- 6) One additional dose every 6 months for previously vaccinated people aged 30 years and over additional dose is given at least 6 months after last dose.

Subsid	dy Full	/ Brand or
(Manufacture	r's Price) Subsidise	d Generic
\$	Per 💌	Manufacturer

#### DIPHTHERIA. TETANUS AND PERTUSSIS VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- c)
- A) Funded for any of the following criteria:
  - 1) A single dose for pregnant women in the second or third trimester of each pregnancy: or
  - 2) A single dose for parents or primary caregivers of infants admitted to a Neonatal Intensive Care Unit or Specialist Care Baby Unit for more than 3 days, who had not been exposed to maternal vaccination at least 14 days prior to birth; or
  - A course of up to four doses is funded for children from age 7 up to the age of 18 years inclusive to complete full primary immunisation; or
  - 4) An additional four doses (as appropriate) are funded for (re-)immunisation for patients post haematopoietic stem cell transplantation or chemotherapy; pre or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or
  - 5) A single dose for vaccination of patients aged from 65 years old: or
  - 6) A single dose for vaccination of patients aged from 45 years old who have not had 4 previous tetanus doses; or
  - 7) For vaccination of previously unimmunised or partially immunised patients; or
  - 8) For revaccination following immunosuppression; or
  - 9) For boosting of patients with tetanus-prone wounds.

Notes: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.

- B) Contractors will be entitled to claim payment from the Funder for the supply of diphtheria, tetanus and pertussis vaccine to patients eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the diphtheria, tetanus and pertussis vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraphs 1 – 9 above.

Inj 2 IU diphtheria toxoid with 20 IU tetanus toxoid, 8 mcg			
pertussis toxoid, 8 mcg pertussis filamentous			
haemagglutinin and 2.5 mcg pertactin in 0.5 ml prefilled			
syringe	0.00	10	✓ Boostrix

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	1	Manufacturer	

#### DIPHTHERIA, TETANUS, PERTUSSIS AND POLIO VACCINE

- a) Only on a prescription
- b) No patient co-payment payable

c)

- A) Funded for any of the following:
  - 1) A single dose for children up to the age of 7 who have completed primary immunisation; or
  - A course of four vaccines is funded for catch up programmes for children (to the age of 10 years) to complete full primary immunisation; or
  - 3) An additional four doses (as appropriate) are funded for (re-)immunisation for people post HSCT, or chemotherapy; pre- or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or
  - 4) Five doses will be funded for children requiring solid organ transplantation.
- B) Contractors will be entitled to claim payment from the Funder for the supply of Diphtheria, tetanus, pertussis and polio vaccine to people eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Diphtheria, tetanus, pertussis and polio vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.

Inj 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg pertussis toxoid, 25 mcg pertussis filamentous

haemagglutinin, 8 mcg pertactin and 80 D-antigen units

10

✓ Infanrix IPV

#### DIPHTHERIA, TETANUS, PERTUSSIS, POLIO, HEPATITIS B AND HAEMOPHILUS INFLUENZAE TYPE B VACCINE

- a) Only on a prescription
- b) No patient co-payment payable

c)

- A) Funded for children meeting any of the following criteria
  - 1) Up to four doses for children under the age of 10 years for primary immunisation; or
  - An additional four doses (as appropriate) for (re-)immunisation of children under the age of 18 years post haematopoietic stem cell transplantation; or
  - 3) An additional four doses (as appropriate) for (re-)immunisation of children under the age of 10 years who are post chemotherapy; pre or post splenectomy; undergoing renal dialysis and other severely immunosuppressive regimens; or
  - 4) Up to five doses for children under the age of 10 years receiving solid organ transplantation.
- B) Contractors will be entitled to claim payment from the Funder for the supply of Diphtheria, tetanus, pertussis, polio, hepatitis B and haemophilus influenzae type b vaccine to people eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Diphtheria, tetanus, pertussis, polio, hepatitis B and haemophilus influenzae type b vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Note: A course of up-to four vaccines is funded for catch up programmes for children (up to and under the age of 10 years) to complete full primary immunisation. Please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes.

Inj 30IU diphtheria with 40IU tetanus and 25mcg pertussis toxoids, 25mcg pertussis filamentous haemagglutinin, 8mcg pertactin, 80D-AgU polio virus, 10mcg hepatitis B antigen, 10mcg H. influenzae type b with tetanus toxoid

10

✓ Infanrix-hexa

I IN	ATIONAL IMMONISATION SCHEDULE					
		Subsidy		Fully	Brand or	
		(Manufacturer's Price) \$	Su Per	bsidised	Generic Manufacturer	
HAFMO	PHILUS INFLUENZAE TYPE B VACCINE	<u> </u>				_
	Only on a prescription					
	No patient co-payment payable					
-,	A) One dose for people meeting any of the following:					
	<ol> <li>For primary vaccination in children; or</li> </ol>		_			
	2) An additional dose (as appropriate) is funded f	` '				#II
	transplantation, or chemotherapy; functional attransplant, pre or post cochlear implants, rena				,	٦r
	For use in testing for primary immunodeficience					,,
	physician or paediatrician.	,				
	B) Contractors will be entitled to claim payment from the					
	vaccine to people eligible under the above criteria p					
	for subsidised immunisation, and they may only do sin the Pharmaceutical Schedule.	so in respect of the H	aemoph	ilus influe	nzae type b vaccine li	st
	C) Contractors may only claim for populations within th	e criteria that are cov	ered by	their cont	ract, which may be a	
	sub-set of the population described in paragraph A					
lnj 1	0 mcg vial with diluent syringe	0.00	1	✓ <u>A</u>	ct-HIB	
	TIS A VACCINE - [Xpharm]					
Fun	ded for patients meeting any of the following criteria:					
,	Two vaccinations for use in transplant patients; or					
,	Two vaccinations for use in children with chronic liver d	*				
3)	One dose of vaccine for close contacts of known hepati	tis A cases.				

1

✓ <u>Havrix 1440</u> ✓ Havrix Junior

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer	
HEDATITIS B DECOMBINIANT VACCINE [Vaborm]				

#### HEPATITIS B RECOMBINANT VACCINE - [Xpharm]

✓ Engerix-B 

Funded for patients meeting any of the following criteria:

- 1) for household or sexual contacts of known acute hepatitis B patients or hepatitis B carriers; or
- 2) for children born to mothers who are hepatitis B surface antigen (HBsAq) positive; or
- 3) for children up to and under the age of 18 years inclusive who are considered not to have achieved a positive serology and require additional vaccination or require a primary course of vaccination; or
- 4) for HIV positive patients; or
- 5) for hepatitis C positive patients; or
- 6) for patients following non-consensual sexual intercourse; or
- 7) for patients prior to planned immunosuppression for greater than 28 days; or
- 8) for patients following immunosuppression; or
- 9) for solid organ transplant patients; or
- 10) for post-haematopoietic stem cell transplant (HSCT) patients; or
- 11) following needle stick injury.

Inj 20 mcg per 1 ml prefilled syringe........................0.00 ✓ Engerix-B

Funded for patients meeting any of the following criteria:

- 1) for household or sexual contacts of known acute hepatitis B patients or hepatitis B carriers; or
- 2) for children born to mothers who are hepatitis B surface antigen (HBsAg) positive; or
- 3) for children up to and under the age of 18 years inclusive who are considered not to have achieved a positive serology and require additional vaccination or require a primary course of vaccination; or
- 4) for HIV positive patients; or
- 5) for hepatitis C positive patients; or
- 6) for patients following non-consensual sexual intercourse; or
- 7) for patients prior to planned immunosuppression for greater than 28 days; or
- 8) for patients following immunosuppression; or
- 9) for solid organ transplant patients; or
- 10) for post-haematopoietic stem cell transplant (HSCT) patients; or
- 11) following needle stick injury; or
- 12) for dialysis patients; or
- 13) for liver or kidney transplant patients.

	Subsidy		Fully	Brand or	
(Manufacturer's Price) Subsidised Generic	(Manufacturer's Price)	Sub	sidised	Generic	
\$ Per ✔ Manufacturer	\$	Per	1	Manufacturer	

HUMAN PAPILLOMAVIRUS (6, 11, 16, 18, 31, 33, 45, 52 AND 58) VACCINE [HPV]

- a) Maximum of 1 inj per prescription
- b) Only on a prescription
- c) No patient co-payment payable
- d
- a) A) Any of the following:
  - 1) Maximum of two doses for children aged 14 years and under; or
  - 2) Maximum of three doses for people meeting any of the following criteria:
    - 1) People aged 15 to 26 years inclusive; or
    - Either

People aged 9 to 26 years inclusive who have

- 1) Confirmed HIV infection; or
- 2) Received a transplant (including stem cell): or
- 3) Maximum of four doses for people aged 9 to 26 years inclusive post chemotherapy
- B) Contractors will be entitled to claim payment from the Funder for the supply of Human papillomavirus vaccine to people eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Human papillomavirus vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraphs A above.

	Subsidy (Manufacturer's Price) \$	Sı Per	Fully ubsidised	Brand or Generic Manufacturer
INFLUENZA VACCINE Inj 60 mcg in 0.5 ml syringe (quadrivalent vaccine)	120.00	10		fluvac Tetra (2024 formulation)

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

- a) Maximum of 1 inj per prescription
- b) Only on a prescription
- c) No patient co-payment payable
- ď

#### A) INFLUENZA VACCINE

is available each year for patients who meet the following criteria, as set by Pharmac:

- a) all people 65 years of age and over; or
- b) people under 65 years of age who:
  - i) have any of the following cardiovascular diseases:
    - a) ischaemic heart disease, or
    - b) congestive heart failure, or
    - c) rheumatic heart disease, or
    - d) congenital heart disease, or
    - e) cerebo-vascular disease; or
  - ii) have either of the following chronic respiratory diseases:
    - a) asthma, if on a regular preventative therapy, or
    - b) other chronic respiratory disease with impaired lung function; or
  - iii) have diabetes: or
  - iv) have chronic renal disease; or
  - v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
  - vi) have any of the following other conditions:
    - a) autoimmune disease, or
    - b) immune suppression or immune deficiency, or
    - c) HIV, or
    - d) transplant recipients, or
    - e) neuromuscular and CNS diseases/disorders, or
    - f) haemoglobinopathies, or
    - g) are children on long term aspirin, or
    - h) have a cochlear implant, or
    - i) errors of metabolism at risk of major metabolic decompensation, or
    - i) pre and post splenectomy, or
    - k) Down syndrome, or
  - vii) are pregnant; or
- c) children 4 years of age and under who have been hospitalised for respiratory illness or have a history of significant respiratory illness; or
- d) people under 65 years of age who:
  - i) have any of the following serious mental health conditions:
    - a) schizophrenia, or
    - b) major depressive disorder, or
    - c) bipolar disorder, or
    - d) schizoaffective disorder, or
  - ii) are currently accessing secondary or tertiary mental health and addiction services; or

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy,
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Contractors will be entitled to claim payment for the supply of influenza vaccine to patients eligible under the above criteria pursuant to their contract with Health NZ for subsidised immunisation, and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Subsidy (Manufacturer's Price)	,		Brand or Generic
 \$	Per	•	Manufacturer

#### MEASLES, MUMPS AND RUBELLA VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- c)

### A) Measles, mumps and rubella vaccine

A maximum of two doses for any patient meeting the following criteria:

- 1) For primary vaccination in children; or
- 2) For revaccination following immunosuppression; or
- 3) For any individual susceptible to measles, mumps or rubella; or
- 4) A maximum of three doses for children who have had their first dose prior to 12 months.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes. Although a price is listed for the vaccine, doctors can still order measles mumps and rubella vaccine free of charge, as with other Schedule vaccines.

- B) Contractors will be entitled to claim payment for the supply of measles, mumps and rubella vaccine to patients eligible under the above criteria pursuant to their contract with Health NZ for subsidised immunisation, and they may only do so in respect of the measles, mumps and rubella vaccine listed in the Pharmaceutical Schedule.
- C) Contractors can only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Inj, measles virus 1,000 CCID50, mumps virus 5,012 CCID50,		
Rubella virus 1,000 CCID50; prefilled syringe/ampoule of		
diluent 0.5 ml	10	Priorix

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

#### MENINGOCOCCAL (GROUPS A, C, Y AND W-135) CONJUGATE VACCINE

Inj 10 mcg of each meningococcal polysaccharide conjugated

a) Only on a prescription

b) No patient co-payment payable

C,

- A) Any of the following:
  - Up to three doses and a booster every five years for patients pre- and post splenectomy and for patients
    with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre or post
    solid organ transplant; or

MenQuadfi

- 2) One dose for close contacts of meningococcal cases of any group; or
- 3) One dose for person who has previously had meningococcal disease of any group; or
- 4) A maximum of two doses for bone marrow transplant patients; or
- 5) A maximum of two doses for person pre- and post-immunosuppression\*; or
- B) Both:
  - 1) Person is aged between 13 and 25 years, inclusive; and
  - 2) Either:
    - One dose for individuals who are entering within the next three months, or in their first year of living in boarding school hostels, tertiary education halls of residence, military barracks, Youth Justice residences, or prisons; or
    - 2) One dose for individuals who turn 13 years of age while living in boarding school hostels.
- C) Contractors will be entitled to claim payment from the Funder for the supply of Meningococcal A, C, Y and W-135 vaccine to patients eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Meningococcal A, C, Y and W-135 vaccine listed in the Pharmaceutical Schedule.
- D) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraphs A-B above.

Note: children under seven years of age require two doses 8 weeks apart, a booster dose three years after the primary series and then five yearly.

\*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days.

Inj 5 mcg of each meningococcal polysaccharide conjugated to a total of approximately 44 mcg of tetanus toxoid carrier

- A) Both:
  - 1) The child is under 12 months of age; and
  - 2) Any of the following:
    - A maximum of three doses (dependant on age at first dose) for patients pre- and post- splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre- or post- solid organ transplant; or
    - A maximum of three doses (dependant on age at first dose) for close contacts of meningococcal cases of any group; or
    - A maximum of three doses (dependant on age at first dose) for child who has previously had meningococcal disease of any group; or
    - 4) A maximum of three doses (dependant on age at first dose) for bone marrow transplant patients; or
    - A maximum of three doses (dependant on age at first dose) for child pre- and post-immunosuppression\*.

Note: infants from 6 weeks to less than 6 months of age require a 2+1 schedule, infants from 6 months to less than 12 months of age require a 1+1 schedule. Refer to the Immunisation Handbook for recommended booster schedules with meningococcal ACWY vaccine.

\*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	5	Subsidised	Generic	
\$	Per	✓	Manufacturer	

#### MENINGOCOCCAL B MULTICOMPONENT VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- c) Any of the following:
  - A) Three doses for children up to 12 months of age (inclusive) for primary immunisation; or
  - B) Up to three doses (dependent on age at first dose) for a catch-up programme for children from 13 months to 59 months of age (inclusive) for primary immunisation, from 1 March 2023 to 31 August 2025; or
  - C) Both:
    - 1) Person is one year of age or over; and
    - 2) Any of the following:
      - i) up to two doses and a booster every five years for patients pre- and post-splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre- or post-solid organ transplant; or
      - ii) up to two doses for close contacts of meningococcal cases of any group; or
      - iii) up to two doses for person who has previously had meningococcal disease of any group; or
      - iv) up to two doses for bone marrow transplant patients; or
      - v) up to two doses for person pre- and post-immunosuppression\*; or
  - D) Both:
    - 1) Person is aged between 13 and 25 years (inclusive); and
    - 2) Either:
      - Two doses for individuals who are entering within the next three months, or in their first year of living in boarding school hostels, tertiary education halls of residence, military barracks, Youth Justice residences or prisons; or
      - ii) Two doses for individuals who turn 13 years of age while living in boarding school hostels.
  - E) Contractors will be entitled to claim payment from the Funder for the supply of Meningococcal B multicomponent vaccine to patients eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Meningococcal B multicomponent vaccine listed in the Pharmaceutical Schedule.
  - F) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraphs A-D above.

\*Immunosuppression due to corticosteroid or other immunosuppressive therapy must be for a period of greater than 28 days.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Por 🗸	Manufacturer

#### PNEUMOCOCCAL (PCV13) CONJUGATE VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- c)
- A) Any of the following:
  - 1) A course of three doses for previously unvaccinated children up to the age of 59 months inclusive; or
  - Two doses are funded for high risk individuals (over the age of 12 months and under 18 years) who have previously received two doses of the primary course of PCV10: or
  - 3) Up to an additional four doses (as appropriate) are funded for the (re)immunisation of high risk children aged under 5 years with any of the following:
    - a) on immunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient immune response; or
    - b) primary immune deficiencies; or
    - c) HIV infection: or
    - d) renal failure, or nephrotic syndrome; or
    - e) who are immune-suppressed following organ transplantation (including haematopoietic stem cell transplant); or
    - f) cochlear implants or intracranial shunts; or
    - g) cerebrospinal fluid leaks; or
    - n) receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage of 20 mg or greater; or
    - i) chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or
    - j) pre term infants, born before 28 weeks gestation; or
    - k) cardiac disease, with cyanosis or failure; or
    - diabetes; or
    - m) Down syndrome; or
    - n) who are pre-or post-splenectomy, or with functional asplenia; or
  - 4) Up to an additional four doses (as appropriate) are funded for the (re-)immunisation of individuals 5 years and over with HIV, pre or post haematopoietic stem cell transplantation, or chemotherapy; pre- or post splenectomy; functional asplenia, pre- or post- solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, intracranial shunts, cerebrospinal fluid leaks or primary immunodeficiency; or
  - 5) For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.
- B) Contractors will be entitled to claim payment from the Funder for the supply of Pneumococcal (PCV13) conjugate vaccine to people eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Pneumococcal (PCV13) conjugate vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Note: please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes

Inj 30.8 mcg of pneumococcal polysaccharide serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and 23F in 0.5ml

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
PNEUMOCOCCAL (PPV23) POLYSACCHARIDE VACCINE - [ Either:	[Xpharm]		

- 1) Up to three doses (as appropriate) for patients with HIV, for patients post haematopoietic stem cell transplant, or chemotherapy; pre- or post-splenectomy or with functional asplenia, pre- or post-solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, or primary immunodeficiency; or
- 2) All of the following:
  - a) Patient is a child under 18 years for (re-)immunisation; and
  - b) Treatment is for a maximum of two doses; and
  - c) Any of the following:
    - i) on immunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient immune response; or
    - ii) with primary immune deficiencies: or
    - iii) with HIV infection; or
    - iv) with renal failure, or nephrotic syndrome; or
    - v) who are immune-suppressed following organ transplantation (including haematopoietic stem cell transplant);
    - vi) with cochlear implants or intracranial shunts; or
    - vii) with cerebrospinal fluid leaks; or
    - viii) receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage of 20 mg or greater; or
    - ix) with chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or
    - x) pre term infants, born before 28 weeks gestation; or

xi) with diabetes; or xii) with Down syndrome; or xiv) who are pre-or post-splenectomy, or with functional asple	enia.	
Inj 575 mcg in 0.5 ml prefilled syringe (25 mcg of each 23 pneumococcal serotype)	1	✓ Pneumovax 23
POLIOMYELITIS VACCINE - [Xpharm]		
Up to three doses for patients meeting either of the following:		
<ol> <li>For partially vaccinated or previously unvaccinated individuals; or</li> </ol>		
<ol><li>For revaccination following immunosuppression.</li></ol>		
Note: Please refer to the Immunisation Handbook for appropriate schedul	le for catch-up	p programmes.
Inj 80D antigen units in 0.5 ml syringe0.00	1	✓ <u>IPOL</u>

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

#### ROTAVIRUS ORAL VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- c)
- A) Maximum of two doses for people meeting the following:
  - 1) first dose to be administered in infants aged under 14 weeks of age; and
  - 2) no vaccination being administered to children aged 24 weeks or over.
- B) Contractors will be entitled to claim payment from the Funder for the supply of Rotavirus oral vaccine to people eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Rotavirus oral vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Oral susp live attenuated human rotavirus		
1,000,000 CCID50 per dose, squeezable tube0.00	) 10	Rotarix
Oral susp live attenuated human rotavirus		
1,000,000 CCID50 per dose, prefilled oral applicator0.00	) 10	✓ Rotarix

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer	
--	--

### VARICELLA VACCINE [CHICKENPOX VACCINE]

- a) Only on a prescription
- b) No patient co-payment payable
- c)
- A) Either:
  - 1) Maximum of one dose for primary vaccination for either:
    - a) Any infant born on or after 1 April 2016; or
    - For previously unvaccinated children turning 11 years old on or after 1 July 2017, who have not previously had a varicella infection (chickenpox), or
  - 2) Maximum of two doses for any of the following:
    - a) Any of the following for non-immune individuals:
      - i) with chronic liver disease who may in future be candidates for transplantation; or
      - ii) with deteriorating renal function before transplantation; or
      - iii) prior to solid organ transplant; or
      - iv) prior to any elective immunosuppression\*; or
      - v) for post exposure prophylaxis who are immune competent inpatients; or
    - b) For individuals at least 2 years after bone marrow transplantation, on advice of their specialist; or
    - c) For individuals at least 6 months after completion of chemotherapy, on advice of their specialist; or
    - d) For HIV positive non immune to varicella with mild or moderate immunosuppression on advice of HIV specialist; or
    - e) For individuals with inborn errors of metabolism at risk of major metabolic decompensation, with no clinical history of varicella; or
    - f) For household contacts of paediatric patients who are immunocompromised, or undergoing a procedure leading to immune compromise where the household contact has no clinical history of varicella; or
    - g) For household contacts of adult patients who have no clinical history of varicella and who are severely immunocompromised, or undergoing a procedure leading to immune compromise where the household contact has no clinical history of varicella.
- B) Contractors will be entitled to claim payment from the Funder for the supply of Varicella vaccine [Chickenpox vaccine] vaccine to people eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Varicella vaccine [Chickenpox vaccine] listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraphs A above.

\* immunosuppression due to steroid or other immunosuppressive therapy must be for a treatment period of greater than 28 days

	Subsidy (Manufacturer's Price) \$	Sub Per	Fully osidised	Brand or Generic Manufacturer
ARICELLA ZOSTER VACCINE [SHINGLES VACCINE]				
a) Only on a prescription				
b) No patient co-payment payable     c)				
A) Funded for patients meeting the following criteria:				
1) Either:				
1) Two doses for all people aged 65 years	s, or			
<ol><li>Two doses for people 18 years of age of</li></ol>	or older with any of the	following	j:	
<ul> <li>a) pre- and post-haematopoietic ster</li> </ul>	m cell transplant or cell	ular ther	apy; or	
b) pre- or post-solid organ transplan	t; or			
c) haematological malignancies; or				
<ul> <li>d) people living with poorly controlled</li> </ul>	d HIV infection; or			
<ul> <li>e) planned or receiving disease mod</li> </ul>	lifying anti-rheumatic d	rugs (DN	1ARDs –	targeted synthetic,
biologic, or conventional synthetic	c) for polymyalgia rheur	natica, s	ystemic I	upus erythematosus or
rheumatoid arthritis; or	=\			
f) end stage kidney disease (CKD 4	or 5); or			
g) primary immunodeficiency				. (01: 1
B) Contractors will be entitled to claim payment from				, ,
vaccine) to patients eligible under the above criteri	a pursuant to their con	tract with	ı Health I	New ∠ealand (Health N∠)

vaccine) to patients eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Varicella zoster vaccine [Shingles vaccine] listed in the Pharmaceutical Schedule.

C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

# **Diagnostic Agents**

- Symbols -	Albendazole	98	Amzoate	31
3TC116	Albey	258–259	Anaesthetics	127
- A -	Albustix	86	Anafranil	
A-Scabies76	Alchemy Oxaliplatin	158	Anagrelide hydrochloride	
Abacavir sulphate116	Alchemy Oxybutynin		Analgesics	128
Abacavir sulphate with	Aldurazyme	30	Anastrozole	181
lamivudine116	Alecensa	170	Anatrole	181
Abacavir/Lamivudine Viatris116	Alectinib	170	Anoro Ellipta	263
Abilify Maintena141	Alendronate sodium	121	Antabuse	153
Abilify Maintena S29141	Alendronate sodium with		Antacids and Antiflatulents	6
Abiraterone acetate178	colecalciferol	121	Anthelmintics	98
Acarbose11	Alfacalcidol	34	Antiacne Preparations	
Accarb11	Alfamino	298	Antiallergy Preparations	258
Acetazolamide270	Alfamino Junior	298	Antianaemics	
Acetec48	Alginic acid	6	Antiandrogen Oral	
Acetic acid with hydroxyquinoline and	Alglucosidase alfa	<mark>27</mark>	Contraceptives	84
ricinoleic acid84	Alkeran	158	Antiarrhythmics	
Acetylcysteine273	Allerfix	270	Antibacterials	98
Aci-Jel84	Allerpro Syneo 1	300	Antibacterials Topical	
Aciclovir	Allerpro Syneo 2	300	Anticholinergic Agents	
Infection111	Allersoothe	259	Anticholinesterases	
Sensory268	Allmercap	160	Antidepressants	
Acidex6	Allopurinol		Antidiarrhoeals	
Acipimox56	Almarytm	51	Antiepilepsy Drugs	134
Acitretin76	Alpha-Adrenoceptor Blockers		Antifibrinolytics, Haemostatics and	
Act-HIB306	Alpha-Keri Lotion		Local Sclerosants	
Actemra236	Alphamox 125		Antifibrotics	263
Actinomycin D162	Alphamox 250	101	Antifungals	105
Actrapid10	Alprolix		Antifungals Topical	
Actrapid Penfill10	Alu-Tab		Antihistamines	
Acupan128	Aluminium hydroxide	6	Antihypotensives	
Adalimumab (Amgevita)189	Alyacen	83	Antimalarials	
Adalimumab (Humira - Alternative	Amantadine hydrochloride		Antimigraine Preparations	137
brand) 198	Ambrisentan	59	Antinausea and Vertigo Agents	137
Adapalene70	Ambrisentan Viatris	59	Antipruritic Preparations	
Adcetris206	Amgevita	189	Antipsychotics	139
ADR Cartridge 1.824	Amiloride hydrochloride	54	Antiretrovirals	114
Adrenaline	Amiloride hydrochloride with		Antirheumatoid Agents	121
Cardiovascular58	furosemide	<u>55</u>	Antispasmodics and Other Agents	
Respiratory258	Amiloride hydrochloride with		Altering Gut Motility	8
Advantan73	hydrochlorothiazide	<u>55</u>	Antithrombotic Agents	42
Advate41	Aminophylline	265	Antithymocyte globulin	
Adynovate42	Amiodarone hydrochloride	50	(equine)	189
Afinitor253	Amisulpride	139	Antitrichomonal Agents	109
Aflibercept204	Amitriptyline	132	Antituberculotics and	
AFT-Pyrazinamide110	Amlodipine	53	Antileprotics	109
Agents Affecting the	Amorolfine	71	Antiulcerants	g
Renin-Angiotensin System 48	Amoxicillin		Antivirals	
Agents for Parkinsonism and Related	Amoxicillin with clavulanic acid		Anxiolytics	143
Disorders 126	Amoxiclav Devatis Forte	101	Anzatax	
Agents Used in the Treatment of	Amphotericin B		Apidra	
Poisonings273	Amsacrine		Apidra SoloStar	
Agrylin162	AmsaLyo		APO-Atomoxetine	148
Albalon272	Amsidine	161	Apo-Azithromycin	99

APO-Candesartan HCTZ		ATGAM	189	Benzylpenicillin sodium [Penicillin	
16/12.5	49	Ativan	143	G]	101
APO-Candesartan HCTZ		Atnahs Olsalazine	8	Beta Cream	72
32/12.5	49	Atomoxetine	148	Beta Ointment	72
Apo-Temozolomide	168	Atorvastatin	56	Beta Scalp	78
Apomorphine hydrochloride	126	Atropine sulphate		Beta-Adrenoceptor Agonists	261
Aprepitant	137	Cardiovascular	50	Beta-Adrenoceptor Blockers	51
Apresoline	58	Sensory		Beta-hCG low sensitivity urine test	
Aptamil Feed Thickener	291	Atropt	271	kit	85
Aqueous cream		Atrovent		Betadine	75
Aratac		Aubagio	145	Betadine Skin Prep	75
Arava	121	Augmentin	101	Betaferon	144
Arginine	<mark>28</mark>	Aurorix		Betahistine dihydrochloride	137
Arginine2000	296	AutoSoft 30	23	Betaine	28
Aripiprazole	139, 141	AutoSoft 90	23	Betamethasone dipropionate	72
Aripiprazole Sandoz	139	Avelox	103	Betamethasone dipropionate with	
Aristocort		Avonex	144	calcipotriol	77
Arrotex-Prazosin S29	48	Avonex Pen	144	Betamethasone sodium phosphate	
Arrow - Clopid	42	Azacitidine	158	with betamethasone acetate	88
Arrow - Lattim	271	Azacitidine Dr Reddy's	158	Betamethasone valerate7	2, 78
Arrow-Amitriptyline	132	Azamun		Betamethasone valerate with sodiu	m
Arrow-Bendrofluazide	55	Azathioprine	182	fusidate [fusidic acid]	73
Arrow-Brimonidine	271	Azilect	126	Betaxolol	
Arrow-Diazepam	143	Azithromycin	99	Betnovate	72
Arrow-Doxorubicin		Azopt	270	Betoptic	270
Arrow-Fluoxetine	133	AZT	116	Betoptic S	
Arrow-Losartan &		- B -		Bexsero	313
Hydrochlorothiazide	49	B-D Micro-Fine	16	Bezafibrate	56
Arrow-Norfloxacin	119	B-D Ultra Fine	16	Bezalip	56
Arrow-Ornidazole	109	B-D Ultra Fine II	16	Bezalip Retard	56
Arrow-Quinapril 10	49	Bacillus Calmette-Guerin (BCG)		Bicalutamide	
Arrow-Quinapril 20		vaccine	189	Bicillin LA	101
Arrow-Quinapril 5		Bacillus Calmette-Guerin		BiCNU	157
Arrow-Roxithromycin		vaccine	303	BiCNU S29	157
Arrow-Timolol		Baclofen	125	Bile and Liver Therapy	10
Arrow-Topiramate		Bactroban	70	Biltricide	
Arrow-Tramadol		Balance	30	Bimatoprost	270
Arsenic trioxide		Barrier Creams and Emollients	74	Bimatoprost Multichem	
Asacol	7	BCG Vaccine AJV	303	Binarex	
Asacol S29		Beclazone 100	259	Binocrit	38
Ascend	71	Beclazone 250	259	Biodone	130
Ascend Aripiprazole		Beclazone 50	259	Biodone Extra Forte	130
Ascend-Cefuroxime		Beclomethasone dipropionate	259	Biodone Forte	130
Ascorbic acid	34	Bedaquiline		Bisacodyl	
Aspen Adrenaline		Bee venom allergy treatment		Bisacodyl Viatris	
Aspirin		Bendamustine hydrochloride		Bisoprolol fumarate	
Blood	42	Bendamustine Sandoz		BK Lotion	
Nervous	128	Bendrofluazide	55	Bleomycin sulphate	162
Asthalin	261	Bendroflumethiazide		Blood Colony-stimulating	
Atazanavir Mylan	116	[Bendrofluazide]	55	Factors	45
Atazanavir sulphate		Benralizumab		Blood glucose diagnostic test	
Atazanavir Viatris		Benzathine benzylpenicillin		meter	15
Atenolol		Benzatropine mesylate		Blood glucose diagnostic test	
Atenolol AFT		Benzbromarone		strip	15
Atenolol Viatris		Benztrop		Blood glucose test strips (visually	
Atezolizumab		Benzydamine hydrochloride		impaired)	15

Blood Ketone Diagnostic Test	Calcium Folinate Sandoz	159	Cellcept	18
Strip 14	Calcium Folinate Sandoz S29	159	Centrally-Acting Agents	5
Boostrix304	Calcium gluconate	35	Cephalexin ABM	9
Bortezomib	Calcium Homeostasis		Cetirizine hydrochloride	
Bosentan61	Calcium polystyrene sulphona	te47	Cetomacrogol	
Bosentan Dr Reddy's61	Calcium Resonium		Cetomacrogol with glycerol	74
Bplex34	Calogen		Cetomacrogol-AFT	
Brentuximab Vedotin206	Camber		Cetuximab	
Breo Ellipta260	Candesartan cilexetil		Charcoal	
Brevinor 1/2883	Candesartan cilexetil with		CheckTop	
Bricanyl Turbuhaler261	hydrochlorothiazide	49	Chemotherapeutic Agents	
Brimonidine tartrate271	Candestar		Chickenpox vaccine	
Brimonidine tartrate with timolol	Canesten		Chlorambucil	
maleate271	Capecitabine		Chloramphenicol	
Brinzolamide270	Capecitabine Viatris		Chlorothiazide	
Brufen SR	Capsaicin		Chlorpromazine hydrochloride	
BSF Alyacen273	Musculoskeletal	121	Chlorsig	26
BSF Continuous glucose monitor	Nervous		Chlortalidone [Chlorthalidone]	
(interope	Captopril		Chlorthalidone	
BSF Continuous glucose monitor	Carafate		Chlorvescent	
(standalo273	Carbaccord		Choice 380 7med Nsha Silver/cop	
BSF Midazolam-Baxter273	Carbamazepine		Short	
BSF Norethinderone - CDC273	Carbimazole		Ciclosporin	
Buccastem138	Carbomer		Cidomycin P/Free	
Budesonide	Carboplatin		Cilazapril	
Alimentary6	Carboplatin Accord		Cilicaine VK	
Respiratory260, 266	Carbosorb-X		Cinacalcet	
Budesonide Te Arai6	Cardinol LA		Cinacalet Devatis	
Budesonide vith eformoterol260	Cardizem CD		Ciprofloxacin	
Bumetanide54	CareSens Dual		Infection	101
Buprenorphine Naloxone BNM152	CareSens N		Sensory	
Buprenorphine with naloxone	CareSens N POP			
	CareSens N Premier		Ciprofloxacin Teva Cisplatin	
Bupropion hydrochloride				
Burel	CareSens PRO		Cisplatin Accord	
	Carmellose sodium with gelati	n anu	Cisplatin Ebewe	
Buscopan 8	pectin		Citalopram hydrobromide	
Buspirone hydrochloride	Carmustine		Citrulline1000	
Buspirone Viatris	Carnitor		Cladribine	100
Busulfan157	Carvedilol		Clarithromycin	
- C -	Carvedilol Sandoz		Alimentary	
Cabergoline	Casirivimab and imdevimab		Infection	
Caffeine citrate	Catapres		Clexane	
Calamine	CeeNU		Clexane Forte	
Calci-Tab 50035	Cefaclor monohydrate		Clindamycin	
Calcipotriol	Cefalexin		Clinicians	
Calcitonin87	Cefalexin Sandoz		Clinicians Renal Vit	
Calcitriol34	Cefazolin		Clobazam	
Calcitriol-AFT34	Cefazolin-AFT		Clobetasol propionate	73, 7
Calcitriol-AFT S2934	Ceftriaxone		Clobetasone butyrate	7
Calcium 500 mg Hexal35	Ceftriaxone-AFT		Clofazimine	10
Calcium carbonate	Cefuroxime axetil		Clomazol	
Calcium carbonate PAI6	Celapram		Dermatological	7
Calcium Channel Blockers53	Celebrex		Genito-Urinary	84
Calcium Disodium Versenate275	Celecoxib		Clomifene citrate	
Calcium folinate	Celecoxib Pfizer		Clomipramine hydrochloride	
Calcium Folinate Ebewe159	Celestone Chronodose	88	Clomipramine Teva	13

# **INDEX: Generic Chemicals and Brands**

Clonazepam1	34, 143	Corticosteroids and Related Ager	nts	Test	8
Clonidine	54	for Systemic Use		DBL Acetylcysteine	
Clonidine hydrochloride	54	Corticosteroids Topical	72	DBL Adrenaline	58
Clonidine Teva	54	Cosentyx	233	DBL Aminophylline	26
Clopidogrel	42	Cosmegen	162	DBL Bleomycin Sulfate	
Clopine		Coumadin		DBL Bortezomib	
Clopixol1		Country Life	30	DBL Carboplatin	157
Clotrimazole		Coversyl	49	DBL Cisplatin	157
Dermatological	71	COVID-19 vaccine	303	DBL Dacarbazine	162
Genito-Urinary	84	Creon 10000	25	DBL Desferrioxamine Mesylate fo	r Inj
Clozapine		Creon 25000	25	BP	274
Clozaril	139	Creon Micro	25	DBL Docetaxel	
Clustran	137	Crotamiton	72	DBL Ergometrine	84
Co-trimoxazole	105	Crystaderm	70	DBL Gemcitabine	160
Coal tar	77	Cu 375 Standard	82	DBL Gentamicin	103
Coal tar with allantoin, menthol,		Curam	101	DBL Leucovorin Calcium	159
phenol and sulphur	77	Curam Duo 500/125	101	DBL Methotrexate Onco-Vial	
Coal tar with salicylic acid and		Cvite	34	DBL Naloxone Hydrochloride	
sulphur	77	Cyclizine hydrochloride	138	DBL Pethidine Hydrochloride	132
Cobal-B12	34	Cyclizine lactate		DBL Vincristine Sulfate	170
Cobalin-H	34	Cyclogyl	271	Decozol	3
Coco-Scalp	77	Cyclonex	158	Deferasirox	274
Codeine phosphate		Cyclopentolate hydrochloride	271	Deferiprone	274
Extemporaneous	277	Cyclophosphamide	158	Deferoxamine Pfizer S29	274
Nervous	130	Cyclorin		Denosumab	12
Coenzyme Q10	29	Cycloserine	109	Deolate	107
Colchicine	124	Cyklokapron	42	Deoxycoformycin	168
Colecalciferol	34	Cyproterone acetate	89	Depo-Medrol	89
Colestyramine	56	Cyproterone acetate with		Depo-Provera	
Colestyramine - Mylan		ethinyloestradiol	84	Depo-Testosterone	89
Colgout		Cystadane		Deprim	10
Colifoam	7	Cytarabine	160	Dermol	
Colistin sulphomethate	103	Cytotec	9	Desferrioxamine mesilate	274
Colistin-Link	103	Cytoxan	158	Desmopressin	96
Collodion flexible	277	- D -		Desmopressin acetate	
Colloidal bismuth subcitrate	10	D-Penamine	121	Desmopressin-PH&T	90
Colofac	8	Dabigatran	45	Detection of Substances in	
Coloxyl	26	Dacarbazine	162	Urine	86
Combigan	271	Dactinomycin [Actinomycin D]	162	Dexamethasone	
Comirnaty Omicron (XBB.1.5)	303	Daivobet	<mark>77</mark>	Hormone	8
Compound electrolytes	47	Daivonex	<mark>77</mark>	Sensory	269
Compound electrolytes with gluc-	ose	Daktarin	72	Dexamethasone phosphate	
[Dextrose]	47	Dalacin C	103	Dexamethasone with framycetin a	and
Compound hydroxybenzoate	277	Dantrium	125	gramicidin	268
Comtan	126	Dantrium S29		Dexamethasone with neomycin	
Concerta	151	Dantrolene	125	sulphate and polymyxin B	
Condoms	81	Daonil	11	sulphate	269
Condyline	79	Dapa-Tabs	<mark>55</mark>	Dexamfetamine sulfate	148
Continuous glucose monitor		Dapsone	109	Dexcom G6	
(interoperable)	24	Daraprim	104	Dexcom G7	
Continuous glucose monitor		Darunavir		Dexcom ONE+	2
(standalone)	24	Darunavir Viatris	116	Dexmethsone	8
Contraceptives - Hormonal		Dasatinib	171	Dextrochlorpheniramine	
Contraceptives - Non-hormonal		Dasatinib-Teva	171	maleate	259
Copaxone	144	Daunorubicin		Dextrose	
Cordarone-X	50	David One Step Cassette Pregna	ancy	DHC Continus	130

Diabetes10	Dosulepin Viatris132	Elocon	7
Diabetes Management14	Dothiepin132	Elocon Alcohol Free	7
Diacomit136	Dovato117	Eltrombopag	3
Diagnostic Agents318	Doxazosin48	Eltroxin	
Diamide Relief6	Doxazosin Clinect48	EMB Fatol	110
Diamox270	Doxine102	Emend Tri-Pack	13
Diasip281	Doxorubicin Ebewe162	Emicizumab	40
Diazepam134, 143	Doxorubicin hydrochloride162	EMLA	12
Diazoxide10	Doxycycline102	Empagliflozin	13
Dibenzyline48	DP Lotion75	Empagliflozin with metformin	
Diclofenac Sandoz120	DP Lotn HC73	hydrochloride	13
Diclofenac sodium 120	DP-Captopril48	Emsogen	
Differin70	Dr Reddy's Omeprazole9	Emtricitabine	
Difflam33	Drofate52	Emtricitabine with tenofovir	
Diflucan105	Drugs Affecting Bone	disoproxil	11:
Digestives Including Enzymes25	Metabolism 121	Emtriva	110
Digoxin50	Dual blood glucose and blood ketone	Emulsifying ointment	74
Dihydrocodeine tartrate130	diagnostic test meter 14	Emulsifying Ointment ADE	74
Dilantin	Dulaglutide12	Enalapril maleate	4
Dilantin Infatab135	Dulcolax SP Drop27	Enbrel	
Dilantin Paediatric135	Duocal Super Soluble Powder279	Endocrine Therapy	17
Diltiazem CD Clinect53	Duolin262	Endoxan	15
Diltiazem hydrochloride53	Duolin Cipla262	Energivit	29
Dimethicone74–75	Duolin HFA262	Engerix-B	
Dimethyl fumarate144	Duolin Respules262	Enlafax XR	13
Dipentum8	DuoResp Spiromax260	Enoxaparin sodium	
Diphtheria, tetanus and pertussis	Duride58	Enstilar	
vaccine304	Durvalumab246	Ensure	28
V400110			
Diphtheria, tetanus, pertussis and	-E-	Ensure Plus	
			28
Diphtheria, tetanus, pertussis and	-E-	Ensure Plus Ensure Plus HN Ensure Plus HN RTH	289 288
Diphtheria, tetanus, pertussis and polio vaccine305	- E - e-chamber La Grande	Ensure Plus Ensure Plus HN	289 288
Diphtheria, tetanus, pertussis and polio vaccine	- E - e-chamber La Grande	Ensure Plus Ensure Plus HN Ensure Plus HN RTH Ensure Two Cal HN RTH	289 286 290
Diphtheria, tetanus, pertussis and polio vaccine	- E - e-chamber La Grande267 e-chamber Mask267	Ensure Plus Ensure Plus HN Ensure Plus HN RTH	289 286 286 290
Diphtheria, tetanus, pertussis and polio vaccine	- E - e-chamber La Grande	Ensure Plus Ensure Plus HN Ensure Plus HN RTH Ensure Two Cal HN RTH Entacapone	289 286 290 120
Diphtheria, tetanus, pertussis and polio vaccine	- E - e-chamber La Grande	Ensure Plus Ensure Plus HN Ensure Plus HN RTH Ensure Two Cal HN RTH Entacapone Entecavir Entecavir (Rex)	289 288 290 120 11
Diphtheria, tetanus, pertussis and polio vaccine	e-chamber La Grande	Ensure Plus Ensure Plus HN Ensure Plus HN RTH Ensure Two Cal HN RTH Entacapone Entecavir	289 288 290 120 11
Diphtheria, tetanus, pertussis and polio vaccine	- E - e-chamber La Grande	Ensure Plus Ensure Plus HN Ensure Plus HN RTH Ensure Two Cal HN RTH Entacapone Entecavir Entecavir (Rex)	289 288 290 120 11 11
Diphtheria, tetanus, pertussis and polio vaccine	e-chamber La Grande	Ensure Plus Ensure Plus HN Ensure Plus HN RTH Ensure Two Cal HN RTH Entacapone Entecavir Entecavir (Rex) Entresto 24/26 Entresto 49/51	289 286 290 11 11 50
Diphtheria, tetanus, pertussis and polio vaccine	- E - e-chamber La Grande	Ensure Plus	289 286 290 11 11 50 50
Diphtheria, tetanus, pertussis and polio vaccine	e-chamber La Grande	Ensure Plus	289 286 290 11 11 50 50 243 130
Diphtheria, tetanus, pertussis and polio vaccine	e-chamber La Grande	Ensure Plus	289 286 290 11 11 15 50 243 130
Diphtheria, tetanus, pertussis and polio vaccine	- E - e-chamber La Grande	Ensure Plus	288 288 290 120 111 111 50 50 244 130 130
Diphtheria, tetanus, pertussis and polio vaccine	- E - e-chamber La Grande	Ensure Plus	288 288 290 120 111 111 150 50 50 243 130 130 130
Diphtheria, tetanus, pertussis and polio vaccine	e-chamber La Grande	Ensure Plus	288 288 299 110 111 111 50 55 244 130 130 131 131
Diphtheria, tetanus, pertussis and polio vaccine	e-chamber La Grande	Ensure Plus	288 288 288 290 120 111 111 50 50 130
Diphtheria, tetanus, pertussis and polio vaccine	e-chamber La Grande	Ensure Plus	288 288 299 120 111 111 50 50 130
Diphtheria, tetanus, pertussis and polio vaccine	e-chamber La Grande	Ensure Plus	288 288 289 299 120 111 111 55 50 130 130 130 130 130 131
Diphtheria, tetanus, pertussis and polio vaccine	e-chamber La Grande	Ensure Plus	288 288 299 120 111 50 130 130 130 125 255 162 162 155
Diphtheria, tetanus, pertussis and polio vaccine	e-chamber La Grande	Ensure Plus	288 288 289 290 120 111 55 50 130 130 130 130 130 136 156 156 156 157 158
Diphtheria, tetanus, pertussis and polio vaccine	e-chamber La Grande	Ensure Plus	288 288 289 290 120 111 55 50 130 130 130 130 130 136 156 156 156 157 158
Diphtheria, tetanus, pertussis and polio vaccine	e-chamber La Grande	Ensure Plus HN Ensure Plus HN Ensure Plus HN RTH Ensure Two Cal HN RTH Ensure Two Cal HN RTH Entacapone Entecavir (Rex) Entresto 24/26 Entresto 49/51 Entresto 97/103 Entyvio Epilim Crushable Epilim IV Epilim S/F Liquid Epilim Syrup Epipen Epipen Jr Epirubicin Ebewe Epirubicin hydrochloride Eplerenone Epoetin alfa Epoprostenol	288 288 289 290 120 111 55 50 130 130 130 130 130 136 156 156 156 157 158
Diphtheria, tetanus, pertussis and polio vaccine	e-chamber La Grande	Ensure Plus	288 288 289 299 120 111 50 130 130 130 255 166 56 66
Diphtheria, tetanus, pertussis and polio vaccine	e-chamber La Grande	Ensure Plus	288 288 289 299 120 111 50 50 130 130 255 166 56 56 66 66 44
Diphtheria, tetanus, pertussis and polio vaccine	e-chamber La Grande	Ensure Plus HN	
Diphtheria, tetanus, pertussis and polio vaccine	e-chamber La Grande	Ensure Plus HN	

Erythrocin IV	100	fraction	41	Fluids and Electrolytes	46
Erythromycin (as lactobionate)	100	Famotidine	9	Flumetasone pivalate	268
Erythromycin ethyl succinate	100	Famotidine Hovid	9	Fluocortolone caproate with	
Esbriet	264	Fasenra	205	fluocortolone pivalate and	
Escitalopram	133	Faslodex		cinchocaine	8
Escitalopram (Ethics)	133	Fatty Cream AFT	74	Fluorometholone	269
Eskazole	98	Fatty Emulsion Cream (Evara)	74	Fluorouracil	160
Essential Amino Acid Mix	297	Febuxostat	124	Fluorouracil Accord	160
Essential Ethosuximide	134	Febuxostat (Teva)	124	Fluorouracil sodium	79
Estraderm MX	90	FEIBA NF	41	Fluox	133
Estradiol Sandoz	90	Felo 10 ER	53	Fluoxetine hydrochloride	133
Estradiol TDP Mylan		Felo 5 ER	<mark>53</mark>	Flupenthixol decanoate	141
Estradiol Viatris	90	Felodipine	53	Flutamide	179
Estradot	90	Fentanyl	130	Flutamin	179
Estrofem	90	Fentanyl Sandoz		Fluticasone	260
Estrogel	90	Ferinject	36	Fluticasone furoate with	
Etanercept	182	Ferodan	36	umeclidinium and vilanterol	263
Ethambutol hydrochloride	110	Ferriprox		Fluticasone furoate with	
Ethics Aspirin	128	Ferro-F-Tabs	36	vilanterol	260
Ethics Aspirin EC		Ferro-Liquid	36	Fluticasone propionate	266
Ethics Lisinopril		Ferro-tab		Fluticasone with salmeterol	
Ethinyloestradiol with		Ferrograd	36	Flynn	98
desogestrel	82	Ferrosig	37	FML	269
Ethinyloestradiol with		Ferrous fumarate		Foban	
levonorgestrel	83	Ferrous fumarate with folic acid		Folic acid	
Ethinyloestradiol with		Ferrous sulfate		Folic Acid multichem	
norethisterone	83	Fexofenadine hydrochloride		Folic Acid Viatris	39
Ethosuximide		Fibro-vein		Food Thickeners	
Etopophos		Filgrastim		Foods And Supplements For	
Etoposide		Finasteride		Inherited Metabolic Disease	292
Etoposide phosphate		Fingolimod		Fortini	
Etravirine		Firazyr		Fortini Multi Fibre	
Eumovate		Flagyl		Fortisip	
Eurofolic		Flagyl-S		Fortisip Multi Fibre	
Evara		Flamazine		Fosamax	
EVARA White Soft Paraffin		Flecainide acetate		Fosamax Plus	
Everet		Flecainide BNM		Fosfomycin	
Everolimus		Flecainide Controlled Release		Framycetin sulphate	
Evista		Teva	51	Frebini Energy	
Evrysdi		Fleet Phosphate Enema		Frebini Energy Fibre	
Evusheld		Flixonase Hayfever & Allergy		Frebini Original	
Exelon Patch 10		Flixotide		Frebini Original Fibre	
Exelon Patch 5		Flixotide Accuhaler		Freestyle Libre 2	
Exemestane		Florinef		Freestyle Libre 3 Plus	
Exjade		Fluanxol		Fresubin 2kcal HP	
Extemporaneously Compounded		Flucil		Fresubin HP Energy	
Preparations and	•	Flucloxacillin		Fresubin HP Energy Fibre	
Galenicals	277	Flucloxacillin-AFT		Fresubin Intensive	288
Eye Preparations		Flucloxin		Fresubin Original	
Eylea		Flucon		Fresubin Original Fibre	
Ezemibe Viatris		Fluconazole		Frisium	
Ezetimibe		Fludara Oral		Frumil	
Ezetimibe Sandoz		Fludarabine Ebewe		Frusemide	
Ezetimibe sandoz		Fludarabine phosphate		Fucicort	
- F -	00	Fludarabine Sagent		Fucidin	
Factor eight inhibitor bypassing		Fludrocortisone acetate		Fucithalmic	
i actor eight inhibitor bypassing		i iuuiocoilisoile acelale	00	1 uoiti all III0	200

Fulvestrant	179	Gold Knight		Humatin	104
Fungilin	33	Gold Knight XL	81	Humira	19
Furosemide [Frusemide]	54	Goserelin		HumiraPen	19
Furosemide-Baxter	54	Gynaecological Anti-infectives	84	Humulin 30/70	1
fusidic acid		- H -		Humulin NPH	
Dermatological	71, 73	Habitrol	154	Humulin R	10
Infection	104	Haemophilus influenzae type B		Hyaluronic acid	
Sensory		vaccine	306	Hydralazine	5
- G -		Haldol	141	Hydralazine hydrochloride	5
GA Explore 5	296	Haldol Concentrate	141	Hydralyte - Lemonade	4
GA1 Anamix Infant		Haldol Decanoas	141	Hydrocortisone	
GA1 Anamix Junior	296	Haloperidol	139	Dermatological	7
Gabapentin	134	Haloperidol decanoate	141	Hormone	8
Gacet		Harvoni		Hydrocortisone acetate	
Galsulfase	29	Havrix 1440	306	Hydrocortisone acetate with	
Galvumet	12	Havrix Junior	306	pramoxine hydrochloride	
Galvus	12	Haylor syrup	259	Hydrocortisone and paraffin liquid	
Gardasil 9	308	HCU Anamix Infant		and lanolin	7
Gastrodenol	10	HCU Anamix Junior	292	Hydrocortisone butyrate	
Gaviscon Extra Strength	6	HCU Anamix Junior LQ	292	Hydrocortisone with cinchocaine	
Gaviscon Infant		HCU Explore 5		Hydrocortisone with miconazole	
Gazyva		HCU Express 15		Hydrocortisone with natamycin an	
Gefitinib		HCU Lophlex LQ		neomycin	
GEM Aqueous Cream		healthE Aqueous Cream SLS		Hydrogen peroxide	
Gemcitabine Ebewe		Free	74	Hydroxocobalamin	
Gemcitabine hydrochloride		healthE Calamine Aqueous		Hydroxocobalamin Panpharma	
Gemtuzumab ozogamicin		healthE Dimethicone 10%		hydroxycarbamide	
Gentamicin Noridem		healthE Dimethicone 4% Lotion		Hydroxychloroquine sulphate	
Gentamicin sulphate		healthE Dimethicone 5%		Hydroxyurea	
Gilenya		healthE Glycerol BP		[hydroxycarbamide]	16
Ginet		healthE Urea Cream		Hygroton	
Glatiramer acetate		Healtheries Simple Baking Mix		Hylo-Fresh	
Glecaprevir with pibrentasvir.		Hemastix		Hymenoptera25	
Glibenclamide		Hemlibra		Hyoscine butylbromide	
Gliclazide		Heparin sodium		Hyoscine Butylbromide	
Glipizide		Heparin Sodium Panpharma		(Adiramedica)	
Glizide		Heparinised saline		Hyoscine hydrobromide	13
Glucagen Hypokit		Heparon Junior		Hyperuricaemia and Antigout	
Glucagon hydrochloride		Hepatitis A vaccine		Hypromellose	
Glucerna Select		Hepatitis B recombinant		Hypromellose with dextran	27
Glucose [Dextrose]		vaccine	307	- I -	
Gluten Free Foods		Herzuma		lbiamox	10
Glycerin with sodium sacchari		Hiprex		Ibrance	
Glycerin with sucrose		Histaclear		Ibrutinib	
Glycerol		Holoxan		Ibuprofen	
Alimentary	27	Horleys Bread Mix		Ibuprofen SR BNM	
Extemporaneous		Horleys Flour		Icatibant	
Glyceryl trinitrate		Hormone Replacement Therapy		Idarubicin hydrochloride	
Alimentary		Systemic		Idursulfase	
Cardiovascular		HPV		Ifosfamide	
Glycopyrronium		Humalog		llevro	
Glycopyrronium bromide		Humalog Mix 25		lloprost	
Glycopyrronium with	0	Humalog Mix 50		Imatinib mesilate	
indacaterol	262	Human papillomavirus (6, 11, 16		Imatinib-Rex	
Glycosade		31, 33, 45, 52 and 58) vaccine		Imbruvica	
Glytactin Bettermilk				Imfinzi	
Giylaciiii Delleiiiiik	∠∀ე	[HPV]	500		240

Imipramine Crescent	132	cannula, variable insertion)	23	Jevity HiCal RTH	28
Imipramine hydrochloride	132	Insulin pump reservoir	24	Jevity Plus RTH	
Imiquimod	79	Insulin pump with algorithm		Jevity RTH	28
Immune Modulators	117	Insulin syringes, disposable with		Jinarc	
Immunisation - Flu		attached needle	16	Juno Pemetrexed	
Immunisation Flu and Shingles .		Intelence		- K -	
Immunisation Other		Interferon beta-1-alpha		Kadcyla	24
Immunosuppressants		Interferon beta-1-beta		Kalydeco	
Incruse Ellipta		Intra-uterine device		Kemadrin	
Indacaterol		Invega Sustenna		Kenacomb	
Indapamide		Invega Trinza		Kenacort-A 10	
Infanrix IPV		Ipca-Allopurinol		Kenacort-A 40	
Infanrix-hexa		Ipca-Bisoprolol		Kenalog in Orabase	
Infant Formulae		Ipca-Ciprofloxacin		Kent	
Infatrini		Ipca-Donepezil		Ketocal 3:1	
Infliximab		Ipca-Escitalopram		KetoCal 4:1	
Influenza vaccine		IPCA-Frusemide		Ketoconazole	30
	309				7
Influvac Tetra	000	Ipca-Hydroxychloroquine		Dermatological	
(2024 formulation)		IPCA-Metoprolol		Infection	
Inhaled Corticosteroids	259	IPCA-Propranolol		Ketogenic Diet	
Inhaled Long-acting		IPOL		Ketoprofen	
Beta-adrenoceptor Agonists.		Ipratropium bromide2		KetoSens	
Inresa		Ipratropium IVAX		Keytruda	
Inspra		Iressa		Kindergen	
Instillagel Lido		Irinotecan Actavis 100		Kisqali	17
Insulin aspart		Irinotecan hydrochloride		Klacid	
Insulin aspart with insulin aspart	t	Irinotecan-Rex	160	Alimentary	
protamine		Iron (as ferric carboxymaltose)	36	Infection	
Insulin glargine	11	Iron polymaltose	37	Kliogest	9
Insulin glulisine		Isentress	117	Kliovance	
Insulin isophane	11	Isentress HD	117	Kogenate FS	4
Insulin isophane with insulin		Ismo 20	58	Konakion MM	4
neutral	11	Ismo 40 Retard	58	Konsyl-D	
Insulin lispro	11	Isoleucine50	297	Kuvan	3
Insulin lispro with insulin lispro		Isoniazid	110	-L-	
protamine	11	Isoniazid Teva		Labetalol	5
Insulin neutral		Isoniazid with rifampicin		Lacosamide	
Insulin pen needles		Isoptin		Lactulose	
Insulin pump		Isoptin Retard		Laevolac	
Insulin pump cartridge		Isoptin SR		Lagevrio	
Insulin pump infusion set (steel		Isopto Carpine		Lamictal	
cannula)	20	Isosorbide mononitrate		Lamivudine	
Insulin pump infusion set (steel	20	Isosource Standard		Lamivudine Viatris	
cannula, straight insertion)	21	Isotretinoin		Lamivudine/Zidovudine Viatris	
Insulin pump infusion set (teflon		Ispaghula (psyllium) husk		Lamotrigine	
cannula)		Itch-Soothe		Lamprene	
Insulin pump infusion set (teflon		Itraconazole		Lanoxin	
cannula, angle insertion with	00	Itrazole		Lanoxin Paediatric Elixir	
insertion device)		Ivacaftor		Lanoxin PG	
Insulin pump infusion set (teflon		Ivermectin	/5	Lanoxin S29	
cannula, flexible insertion wit		- <b>J -</b>	00	Lansoprazole	
insertion device)		Jadelle		Lantus	
Insulin pump infusion set (teflon		Jakavi		Lantus SoloStar	
cannula, straight insertion wit		Jardiamet		Lanvis	
insertion device)		Jardiance		Lanzol Relief	
Insulin pump infusion set (teflon		Jaydess	91	Largactil	13

Laronidase	30	Locacorten-Viaform ED's	268	Maxidex	26
Lasix		Local preparations for Anal and		Maxitrol	26
Latanoprost	271	Rectal Disorders	8	MCT oil (Nutricia)	280
Latanoprost with timolol	271	Locasol	297	Measles, mumps and rubella	
Lax-Suppositories	<mark>27</mark>	Locoid	.73, 78	vaccine	
Lax-suppositories Glycerol	<mark>27</mark>	Locoid Crelo	73	Mebendazole	
Laxatives	26	Locoid Lipocream	73	Mebeverine hydrochloride	
Laxsol	26	Locorten-Vioform	268	Medac	15
Ledipasvir with sofosbuvir	113	Lodoxamide		Medrol	8
Leflunomide	121	Logem	135	Medroxyprogesterone acetate	
Lenalidomide (Revlimid)	163	Lomide		Genito-Urinary	8
Lenalidomide (Viatris)	164	Lomustine	158	Hormone	9
Lenalidomide Viatris	164	Loniten		Mefenamic acid	120
Lenvatinib	172	Loperamide hydrochloride	6	Megval	15
Lenvima	172	Lopinavir with ritonavir	116	Melatonin	14
Letrole	182	Lopinavir/Ritonavir Mylan	116	Melpha	15
Letrozole	182	Loprofin		Melphalan	15
Leucine100	297	Loprofin Mix		Meningococcal (groups A, C, Y and	d
Leukeran FC	157	Lorafix		W-135) conjugate vaccine	
Leukotriene Receptor		Loratadine	259	Meningococcal B multicomponent	
Antagonists	264	Lorazepam	143	vaccine	313
Leuprorelin		Lorstat		MenQuadfi	
Leustatin		Losartan Actavis	49	Menthol	7
Levetiracetam	135	Losartan potassium	49	Mepolizumab	21
Levetiracetam-AFT	135	Losartan potassium with		Mercaptopurine	
Levocabastine	269	hydrochlorothiazide	49	Mercilon 28	
Levocarnitine	30	Lovir		Mesalazine	
Levodopa with benserazide	126	Loxamine		Mesna	16
Levodopa with carbidopa		Lucrin Depot 1-month	96	Mestinon	
Levomepromazine		Lucrin Depot 3-month		Metabolic Disorder Agents	
Levomepromazine		LumaCina		Metabolics	
hydrochloride	139	Lumigan		Metformin hydrochloride	
Levonorgestrel		Lyllana		Metformin Viatris	
Genito-Urinary	83-84	Lynparza		Methadone BNM	130
Hormone		Lyrica		Methadone hydrochloride	
Levonorgestrel BNM		- M -		Methenamine (hexamine)	
Levothyroxine		m-Eslon	131	hippurate	119
Lidocaine [Lignocaine]		Mabthera		Methopt	
Lidocaine [Lignocaine]		Macrobid		Methotrexate	
hydrochloride	128	Macrogol 3350 with potassium		Methotrexate DBL Onco-Vial	
Lidocaine [Lignocaine] with		chloride, sodium bicarbonate a	and	Methotrexate DBL S29	
prilocaine	128	sodium chloride		Methotrexate Ebewe	
Lidocaine-Baxter		Madopar 125		Methotrexate Sandoz	
Life Extension		Madopar 250		Methyl hydroxybenzoate	
Lignocaine		Madopar 62.5		Methylcellulose	
Linezolid		Madopar HBS		Methylcellulose with glycerin and	
Lioresal Intrathecal		Madopar Rapid		sodium saccharin	27
Lipid-Modifying Agents		Magnesium hydroxide		Methylcellulose with glycerin and	
Liquigen		Magnesium sulphate		sucrose	
Liraglutide		Mantoux		Methyldopa	
Lisdexamfetamine dimesilate .		MAR-Midodrine		Methyldopa Viatris	5
Lisinopril		Marevan		Methylnaltrexone bromide	2
Lithium carbonate	140	Marine Blue Lotion SPF 50+		Methylphenidate ER - Teva	
Livostin		Martindale Pharma		Methylphenidate hydrochloride	150
LMX4		Mask for spacer device		Methylphenidate hydrochloride	10
Lo-Oralcon 20 ED		Maviret		extended-release	15
				JAIOHAGA 1010400	10

Methylprednisolone	88	MiniMed Quick-Set MMT-399A	22	MSUD Anamix Junior LQ	293
Methylprednisolone (as sodium		MiniMed Silhouette MMT-377A	22	MSUD Explore 5	
succinate)	88	MiniMed Silhouette MMT-378A		MSUD Express 15	
Methylprednisolone aceponate		MiniMed Silhouette MMT-381A	22	MSUD Lophlex LQ 20	
Methylprednisolone acetate		MiniMed Sure-T MMT-864A		MSUD Maxamum	
Methylxanthines		MiniMed Sure-T MMT-866A		Mucolytics	
Metoclopramide Actavis 10		MiniMed Sure-T MMT-874A		Mucosoothe	
Metoclopramide hydrochloride		MiniMed Sure-T MMT-876A		Multiple Sclerosis Treatments	
Metolazone		Minims Cyclopentolate	271	Multivitamin renal	
Metopirone		Minims Pilocarpine		Multivitamins	3
Metoprolol IV Mylan		Minims Prednisolone		Mupirocin	
Metoprolol IV Viatris		Minipress		Muscle Relaxants	
Metoprolol succinate		Minirin		Mvite	
Metoprolol tartrate		Minirin Melt		Myambutol	
Metrogyl		Mino-tabs		Mycobutin	
Metronidamed		Minocycline hydrochloride		MycoNail	
Metronidazole		Minomycin		Mycophenolate mofetil	
Metyrapone		Minor Skin Infections		Mydriacyl	
Mexiletine hydrochloride		Minoxidil		Mylan (24 hr release)	
Miacalcic		Minoxidil Roma		Mylan Clomiphen	
Micolette		Mirena		Mylan Italy (24 hr release)	
Miconazole		Miro-Amoxicillin		Myleran	
Miconazole nitrate		Mirtazapine		mylife Inset soft	
Dermatological	72	Misoprostol		mylife Orbit micro	2
Genito-Urinary		Mitomycin (Fresenius Kabi)		mylife YpsoPump Reservoir	
Micreme		Mitomycin (Sagent)		mylife YpsoPump with CamAPS	
Micreme H		Mitomycin C		FX	10
Microgynon 30		Mitozantrone		Myloc CR	
Microlut		Mitozantrone Ebewe		Mylotarg	
Midazolam		Mixtard 30		Myometrial and Vaginal Hormone	
Midazolam Viatris		MMA/PA Anamix Infant		Preparations	
Midazolam-Baxter		MMA/PA Anamix Junior		Myozyme	
Midazolam-Pfizer		MMA/PA Explore 5		- N -	
Midodrine		MMA/PA Express 15		Nadolol	5
Midostaurin		Moclobemide		Nadolol BNM	
Mifegyne		Modafinil		Naglazyme	
Mifepristone		Modafinil Max Health		Naloxone hydrochloride	
Milpharm		Modavigil		Naltraccord	
Minerals		Moduretic		Naltrexone AOP	
Mini-Wright AFS Low Range		Molaxole		Naltrexone hydrochloride	
Mini-Wright Standard		Molnupiravir		Naltrexone Max Health	
Minidiab		Moments		Naphazoline hydrochloride	
MiniMed 3.0 Reservoir		Mometasone furoate		Naphcon Forte	
MMT-332A	24	Monogen		Naprosyn SR 1000	
MiniMed 770G		Montelukast		Naprosyn SR 750	
MiniMed Mio MMT-921A		Montelukast Viatris		Naproxen	120
MiniMed Mio MMT-923A		Moroctocog alfa [Recombinant		Narcaricin mite	12
MiniMed Mio MMT-925A		VIII]		Nasal Preparations	266
MiniMed Mio MMT-923A		Morphine hydrochloride		Natalizumab	14/
MiniMed Mio MMT-943A		Morphine sulphate		Natulan	
MiniMed Mio MMT-945A		Motetis		Nausafix	
MiniMed Mio MMT-965A		Mouth and Throat		Nausicalm	
MiniMed Mio MMT-975A		Movapo		Navelbine S29	
MiniMed Quick-Set MMT-396A.		Moxifloxacin		Nefopam hydrochloride	
MiniMed Quick-Set MMT-397A.		MSUD Anamix Infant		Neo-Cytamen S29	
MiniMed Quick-Set MMT-398A.		MSUD Anamix Junior		Neo-Mercazole	
IVIII IIIVIEU QUICK-JEL IVIIVI I -JYOA .		INIOUD AHAHHIX JUHIUI	∠೨೦	INCU-IVICICAZUIC	🔊

Neocate Gold	298	Noumed Pethidine	132	Oestradiol valerate	90
Neocate Junior Unflavoured	298	Noumed Phenobarbitone	135	Oestradiol with norethisterone	91
Neocate Junior Vanilla	298	Novadoz	157	Oestriol	
Neocate SYNEO	298	NovaSource Renal	284	Genito-Urinary	84
Neoral	253	Novatretin	76	Hormone	91
Neostigmine metilsulfate	120	Novitium Sugar Free	30	Oestrogens	90
Nepafenac	270	NovoMix 30 FlexPen	10	Ofev	263
Nepro HP (strawberry)	284	NovoRapid	11	Oil in water emulsion	
Nepro HP (vanilla)	284	NovoRapid FlexPen		Olanzapine14	40–141
Neulactil	140	NovoRapid Penfill	11	Olaparib	166
NeuroTabs	35	NovoSeven RT		Olbetam	56
Nevirapine	115	Nozinan	139	Olopatadine	
Nevirapine Viatris	115	Nozinan (Swiss)	139	Olopatadine Teva	272
Nicorandil		Nozinan S29	139	Olsalazine	
Nicotine		Nucala	217	Omalizumab	219
Nifedipine		Nuelin	265	Omeprazole	
Nifuran	119	Nuelin-SR	265	Omeprazole actavis 10	9
Nilotinib	174	Nupentin	134	Omeprazole actavis 20	9
Nilstat		Nusinersen	147	Omeprazole actavis 40	9
Alimentary	34	Nutilis	291	Omeprazole Teva	9
Genito-Urinary		Nutren Diabetes	281	Omnitrope	92
Infection	106	Nutrient Modules	278	Omnitrope S29	92
Nimenrix	312	Nutrini Energy Multi Fibre	283	Onbrez Breezhaler	260
Nintedanib	263	Nutrini Energy RTH	283	Oncaspar LYO	167
Nipent	168	Nutrini Low Energy Multi Fibre	285	OncoTICE	189
Niraparib	165	Nutrini Peptisorb	300	Ondansetron	138
Nirmatrelvir with ritonavir	114	Nutrini Peptisorb Energy	300	One-Alpha	34
Nitrates	58	Nutrini RTH		One-Alpha S29	
Nitroderm TTS	58	Nutrison 800 Complete Multi		Opdivo	247
Nitrofurantoin	119	Fibre	288	Ora-Blend	277
Nitrolingual Pump Spray	58	Nutrison Advanced Peptisorb	285	Ora-Blend SF	
Nivestim	45	Nutrison Concentrated	290	Ora-Plus	277
Nivolumab	247	Nutrison Energy	288	Ora-Sweet	277
Nodia	6	Nutrison Energy Multi Fibre	288	Ora-Sweet SF	277
Noflam 250	120	Nutrison Multi Fibre	288	Orabase	33
Noflam 500	120	Nutrison RTH	288	Oral and Enteral Feeds	281
Non-Steroidal Anti-Inflammato	ry	Nyefax Retard	53	Oralcon 30 ED	83
Drugs		Nystatin		Oramorph	131
Nonacog gamma, [Recombina	ant	Alimentary	34	Oramorph CDC S29	131
Factor IX]	41	Genito-Urinary	84	Oratane	
Norethinderone - CDC	84	Infection	106	Orgran	292
Norethisterone		NZB Low Gluten Bread Mix	291	Ornidazole	109
Genito-Urinary	84	- 0 -		Orphenadrine citrate	125
Hormone	91	Obinutuzumab	218	Ortho-tolidine	
Norflex	125	Obstetric Preparations	86	Oruvail SR	
Norfloxacin	119	Ocicure	9	Osmolite RTH	288
Noriday	84	Ocrelizumab	145	Other Endocrine Agents	97
Noriday 28	84	Ocrevus	145	Other Oestrogen Preparations	91
Norimin		Octocog alfa [Recombinant facto	or	Other Progestogen	
Normison		VIII] (Advate)		Preparations	91
Norpress		Octocog alfa [Recombinant factor		Other Skin Preparations	
Nortriptyline hydrochloride		VIII] (Kogenate FS)		Otodex	
Norvir		Octreotide		Ovestin	
Noumed Dexamfetamine		Octreotide GH		Genito-Urinary	84
Noumed Isoniazid		Octreotide long-acting		Hormone	
Noumed Paracetamol		Oestradiol		Oxaliplatin	
				•	

Oxaliplatin Accord	158	Pediasure Plus	283	Pine tar with trolamine laurilsulfate	)
Oxaliplatin Actavis 100	158	Pediasure RTH		and fluorescein	7
Oxaliplatin Ebewe	158	Pegaspargase	167	Pinetarsol	7
Oxis Turbuhaler		Pegasys	117	Pioglitazone	
Oxpentifylline	59	Pegfilgrastim	46	Pirfenidone	264
Oxybutynin		Pegylated interferon alfa-2a	117	Pizotifen	13
Oxycodone Amneal		Pembrolizumab		PKU Anamix Infant	294
Oxycodone hydrochloride	131	Pemetrexed	161	PKU Anamix Junior	294
Oxycodone Lucis S29	131	Pemetrexed-AFT	161	PKU Anamix Junior Chocolate	294
Oxycodone Sandoz	131	Penicillamine	121	PKU Anamix Junior LQ	294
Oxycodone Sandoz S29		Penicillin G	101	PKU Anamix Junior Orange	294
OxyContin	131	PenMix 30	11	PKU Anamix Junior Vanilla	
OxyNorm		PenMix 50	11	PKU Build 10	29
Oxytocin		Pentasa	7	PKU Build 20 Chocolate	29
Oxytocin BNM		Pentostatin [Deoxycoformycin]	168	PKU Build 20 Raspberry	
Oxytocin Panpharma		Pentoxifylline [Oxpentifylline]	59	Lemonade	29
Oxytocin with ergometrine		Peptamen Junior		PKU Build 20 Smooth	
maleate	85	Pepti-Junior		PKU Build 20 Vanilla	
Ozurdex		Perhexiline maleate	53	PKU Explore 10	
- P -		Pericyazine		PKU Explore 5	
Pacifen	125	Perindopril		PKU Express 20	
Pacimol		Periset		PKU First Spoon	
Paclitaxel		Periset ODT		PKU Glytactin RTD 15	
Paclitaxel Actavis		Perjeta		PKU Glytactin RTD 15 Lite	
Paclitaxel Ebewe		Permethrin		PKU GMPro LQ	
Padagis		Perrigo		PKU GMPro Mix-In	
Paediatric Seravit		Pertuzumab		PKU GMPro Ultra Lemonade	
Palbociclib		Peteha		PKU GMPro Ultra Vanilla	
Paliperidone		Pethidine hydrochloride		PKU Lophlex LQ 10	
Paliperidone palmitate		Pevaryl		PKU Lophlex LQ 20	20
Pamidronate disodium		Pexsig		PKU Lophlex Powder	
Pamisol		Pfizer Exemestane		PKU Lophlex Sensation 20	
Pamol		Pfizer S29		PKU Restore Powder	
				PKU sphere20 Banana	
Pancreatic enzyme		Pharmacy Services Pharmascience			
Pantoprazole				PKU sphere20 Chocolate	
Panzop Relief		Phenocen		PKU sphere20 Lemon	
Papaverine hydrochloride		Phenasen		PKU sphere20 Red Berry	
Para-amino salicylic acid		Phenobarbitone	135	PKU sphere20 Vanilla	
Paracetamol		Phenobarbitone sodium	077	PKU Start	
Paracetamol (Ethics)	129	Extemporaneous		Plaquenil	
Paracetamol + Codeine	404	Nervous	147	Plendil ER	
(Relieve)		Phenoxybenzamine	40	Pneumococcal (PCV13) conjugate	
Paracetamol with codeine		hydrochloride		vaccine	314
Paraffin		Phenoxymethylpenicillin (Penici		Pneumococcal (PPV23)	041
Paraffin liquid with wool fat		V)		polysaccharide vaccine	
Parasiticidal Preparations		Phenylalanine50		Pneumovax 23	
Parnate		Phenytoin sodium		Podophyllotoxin	
Paromomycin	104	Phillips Milk of Magnesia		Polaramine	
Paroxetine		Phlexy 10		Poliomyelitis vaccine	
Paser		Phosphate Phebra		Poloxamer	
Paxam		Phosphorus		Poly-Gel	
Paxlovid		Phytomenadione		Poly-Tears	272
Pazopanib	175	Pilocarpine hydrochloride		Poly-Visc	
Pazopanib Teva		Pilocarpine nitrate		Polycal	278
Peak flow meter		Pimafucort		Polyethylene glycol 400 and	
Pediasure	283	Pimecrolimus	<mark>77</mark>	propylene glycol	272

Pomalidomide	168	Protionamide	110	Ricit	8
Pomolide	168	Provera	91	Rifabutin	11
Ponstan	120	Provera HD	91	Rifadin	11
Posaconazole	106	Psoriasis and Eczema		Rifadin Sanofi	
Posaconazole Juno		Preparations	76	Rifampicin	11
Potassium chloride	46–47	PTU	92	Rifaximin	1
Potassium citrate	86	Pulmicort Turbuhaler	260	Rifinah	110
Potassium iodate	35	Pulmozyme	265	Rilutek	12
Povidone iodine	75	Puri-nethol		Riluzole	12
Pradaxa	45	Puritan's Pride Vitamin		RINVOQ	25
Pramipexole hydrochloride	126	B-2 100 mg	30	Riodine	7
Pravastatin	56	Pyrazinamide	110	Risdiplam	148
Praziquantel		Pyridostigmine bromide	120	Risedronate Sandoz	
Prazosin	48	Pyridoxine hydrochloride		Risedronate sodium	12
Prazosin Mylan	48	Pyridoxine multichem	34	Risperdal	
Pred Forte		Pyrimethamine		Risperdal Consta	
Prednisolone	89	Pytazen SR		Risperidone14	
Prednisolone acetate	270	- Q -		Risperidone (Teva)	
Prednisolone sodium		Quantalan sugar free	56	Risperidone Sandoz	
phosphate	270	Quetapel		Risperon	140
Prednisolone-AFT		Quetiapine		Ritalin	
Prednisone		Quinapril		Ritalin LA	
Prednisone Clinect		Qvar		Ritonavir	
Pregabalin	135	- R -		Rituximab (Mabthera)	
Pregabalin Pfizer		RA-Morph	130	Rituximab (Riximyo)	
Pregnancy Tests - hCG Urine		Ralicrom		Rivaroxaban	
Premarin		Raloxifene hydrochloride		Rivastigmine	
Prevenar 13		Raltegravir potassium		Rivastigmine Patch BNM 10	
Priadel	140	Ramipex		Rivastigmine Patch BNM 5	
Primaguine		Ramipril		Rivotril	
Primidone		Ranbaxy-Cefaclor		Riximyo	
Primidone Clinect		Rapamune		RIXUBIS	
Primolut N		Rasagiline	126	Rizamelt	
Priorix		Reandron 1000		Rizatriptan	
Probenecid		Recombinant factor IX		Robinul	
Probenecid-AFT		Recombinant factor VIIa	,	Ronapreve	
Procarbazine hydrochloride		Recombinant factor VIII		Ropin	
Prochlorperazine		Rectogesic		Ropinirole hydrochloride	
Prochlorperazine Brown &		Redipred		Rosuvastatin	
Burk	138	Relieve		Rosuvastatin Viatris	
Proctofoam		Relistor		Rotarix	
Proctosedyl		Remicade		Rotavirus oral vaccine	
Procyclidine hydrochloride		Renilon 7.5		Roxane-Propranolol	5
Progesterone		Resonium-A		Roxithromycin	10
Proglicem		Resource Beneprotein		Rubifen	
Progynova		Respiratory Devices		Rubifen SR	
Prolia		Respiratory Stimulants		Rugby Capsaicin Topical Cream	
Promethazine hydrochloride		Retinol palmitate		Musculoskeletal	
Propafenone hydrochloride		ReTrieve		Nervous	
Propranolol		Retrovir		Rurioctocog alfa pegol [Recombir	
Propylene glycol		Revia		factor VIII]	
Propylthiouracil	۱۱ کـ	Revlimid		Ruxolitinib	
Prostacur		Revolade		Rydapt	
Protaphane		Ribociclib		Rythmodan	
Protaphane Penfill		Riboflavin		Rythmodan - Cheplafarm	
Protifar		Ribomustin		Rytmonorm	5i
ı ıvulal	200	i iiboiiiuaiiii	100	ı ıyunonom	J

-\$-		Sodium alginate		Strides Shasun	10
Sabril	136	Sodium benzoate	31	Stromectol	
Sacubitril with valsartan	50	Sodium bicarbonate		Sucralfate	1
SalAir	261	Blood	46-47	Sulfadiazin-Heyl	10
Salazopyrin	8	Extemporaneous	277	Sulfadiazine Silver	
Salazopyrin EN		Sodium calcium edetate	275	Sulfadiazine sodium	
Salbutamol	261	Sodium chloride		Sulfasalazine	
Salbutamol with ipratropium		Blood	46	Sulphur	7
bromide	262	Respiratory	266	Sulprix	13
Salicylic acid		Sodium citrate with sodium laury	/l	Sumagran	13
Salmeterol	260	sulphoacetate	<mark>27</mark>	Sumatriptan	
Sandomigran	137	Sodium citro-tartrate	86	Sunitinib	
Sandostatin LAR	180	Sodium cromoglicate		Sunitinib Pfizer	17
Sanofi Primaquine	108	Alimentary	8	Sunscreens	7
Sapropterin dihydrochloride		Sensory	270	Sunscreens, proprietary	
Scalp Preparations	78	Sodium Fusidate [fusidic acid]		Survimed OPD	28
Scopoderm TTS		Dermatological	71	Sustagen Hospital Formula	28
Scopolamine - Mylan	138	Infection	104	Sustagen Hospital Formula	
Scopolamine - Mylan S29	138	Sensory	268	Active	28
Sebizole	78	Sodium hyaluronate [Hyaluronic		Sustanon Ampoules	8
Secukinumab	233	acid]	272	Sylvant	23
Sedatives and Hypnotics	146	Sodium phenylbutyrate	31	Symbicort Turbuhaler 100/6	26
Seebri Breezhaler	262	Sodium picosulfate	27	Symbicort Turbuhaler 200/6	26
Senna	27	Sodium polystyrene sulphonate	47	Symbicort Turbuhaler 400/12	26
Senokot	27	Sodium tetradecyl sulphate	42	Symmetrel	12
SensoCard	15	Sodium valproate	136	Sympathomimetics	
Serc	137	Sofradex	268	Synacthen	
Serenace		Soframycin	268	Synacthen Depot	8
Seretide	261	Solgar2	8–30, 32	Synacthene Retard	8
Seretide Accuhaler	261	Solifenacin succinate	86	Synthroid	9
Serevent	260	Solifenacin Viatris	86	Syntometrine	8
Serevent Accuhaler		Solu-Cortef	88	Syrup (pharmaceutical grade)	27
Sertraline	133	Solu-Medrol	88	Systane Unit Dose	27
Setrona	133	Solu-Medrol-Act-O-Vial	88	- T -	
Sevredol	131	Somatropin (Omnitrope)	92	Tacrolimus	
Sex Hormones Non		Sotalol	52	Dermatological	7
Contraceptive	89	Spacer device	267	Oncology	
Shingles vaccine	318	Span-K	47	Tacrolimus Sandoz	
Shingrix		Spazmol		Taliglucerase alfa	
SII-Onco-BCG		Spinal Muscular Atrophy		Tambocor	
Sildenafil		Spinraza		Tamoxifen citrate	
Siltuximab		Spiolto Respimat		Tamoxifen Sandoz	
Simvastatin		Spiractin		Tamsulosin hydrochloride	
Simvastatin Mylan	57	Spiriva		Tamsulosin-Rex	
Simvastatin Viatris	57	Spiriva Respimat		Tandem Cartridge	
Sinemet	126	Spironolactone	55	Tandem t:slim X2 with Basal-IQ	1
Sinemet CR		Sporanox		Tandem t:slim X2 with	
Sintetica Baclofen Intrathecal	125	Sprycel	171	Control-IQ	1
Sirolimus	254	Stelara	242	Tap water	27
Sirturo		Stemetil		Taro	10
Siterone	89	Steril-Gene		Tasigna	17
Slow-Lopresor	52	SteroClear	266	Tasmar	12
Smith BioMed Rapid Pregnancy		Stesolid		Taurine	
Test	85	Stimulants/ADHD Treatments		TCu 380 Plus Normal	
Sodibic	47	Stiripentol	136	Tecentriq	
Sodium acid phosphate	27	Stomahesive		Tecfidera	14

Tegretol	134	Tobrex	268	Two Cal HN	290
Tegretol AU	134	Tocilizumab	236	TYR Anamix Infant	29
Tegretol CR		Tofranil	132	TYR Anamix Junior	29
Telfast		Tolcapone	126	TYR Anamix Junior LQ	29
Temaccord		Tolvaptan		TYR Explore 5	
Temazepam	147	Topamax		TYR Lophlex LQ 20	29
Temozolomide		Topical Products for Joint and		TYR Sphere 20	
Temozolomide-Taro	168	Muscular Pain	121	Tyrosine1000	29
Tenofovir disoproxil	111	Topiramate		Tysabri	14
Tenofovir Disoproxil Emtricitabi		Topiramate Actavis		. U -	
Viatr		Total parenteral nutrition (TPN)		UK Synacthen	8
Tenofovir Disoproxil Viatris	111	TPN	46	Ultibro Breezhaler	
Tenoxicam	120	Tramadol hydrochloride		Ultraproct	
Tensipine MR10		Tramal SR 100		Umeclidinium	
Tepadina		Tramal SR 150	132	Umeclidinium with vilanterol	26
Terbinafine	107	Tramal SR 200		Univent26	1, 26
Terbutaline sulphate		Trandate		Upadacitinib	
Teriflunomide		Tranexamic acid	42	Ural	
Teriflunomide Sandoz		Tranylcypromine sulphate		Urea	
Teriparatide		Trastuzumab (Herzuma)		Urex Forte	
Teriparatide - Teva		Trastuzumab emtansine		Urinary Agents	
Testogel		Travatan		Urinary Tract Infections	
Testosterone		Travoprost		UroFos	
Testosterone cipionate		Treatments for Dementia		Uromitexan	
Testosterone esters		Treatments for Substance		Ursodeoxycholic acid	
Testosterone undecanoate		Dependence	152	Ursosan	
Tetrabenazine		Trelegy Ellipta		Ustekinumab	
Tetrabromophenol		Trental 400		Utrogestan	
Tetracosactrin		Tretinoin		- V -	
Tetracycline		Dermatological	70	Vaccinations	30
Teva Lisinopril		Oncology		Vaclovir	
Teva-Ketoconazole		Trexate		Valaciclovir	
Thalidomide		Triamcinolone acetonide		Valganciclovir	
Thalomid		Alimentary	33	Valganciclovir Viatris	
Theophylline		Dermatological		Valine50	
Thiamine hydrochloride		Hormone		Vancomycin	
Thiamine multichem		Triamcinolone acetonide with		Vannair	
THIO-TEPA		gramicidin, neomycin and nys	tatin	Varenicline Pfizer	
Thioguanine		Dermatological		Varenicline tartrate	
Thiotepa		Sensory		Varicella vaccine [Chickenpox	
Thyroid and Antithyroid Agents		Trientine		vaccine]	31
Ticagrelor		Trientine Waymade		Varicella zoster vaccine [Shingles	
Ticagrelor Sandoz		Trikafta		vaccine]	318
Tilcotil		Trimethoprim		Varilrix	
Timolol		Trimethoprim with		Various	
Tiotropium bromide		sulphamethoxazole		VariSoft	
Tiotropium bromide with		[Co-trimoxazole]	105	Vasodilators	
olodaterol	263	Trisequens		Vasopressin Agonists	
Tivicay		Trisul		Vasorex	5
Tixagevimab with cilgavimab		Trophic Hormones		Vebulis	6
TMP		Tropicamide		Vedafil	
Tobramycin		Trulicity		Vedolizumab	
Infection	105	TruSteel		Veletri	
Sensory		Tryzan		Venclexta	
Tobramycin (Viatris)		Tuberculin PPD [Mantoux] test	318	Venetoclax	
		Tubersol	010	Venlafaxine	
Tobramycin BNM	105	Lubersol	310	venialaxine	1.5

Venomil	258-259	XMET Maxamum	292
VENOX		Xolair	
Ventolin		Xolair AU	
Vepesid		XP Maxamum	
Verapamil hydrochloride		Xylocaine	
		•	
Vermox		Xylocaine 2% Jelly	
Versacloz		Xylocard 500	
Vesanoid		Xyntha	41
Vexazone		-Z-	40
Vfend		Zapril	
Viaderm KC		Zarontin	
Victoza		Zaroxolyn	
Vigabatrin		Zavedos	
Vigisom	146	Zeffix	111
Vildagliptin	12	Zejula	165
Vildagliptin with metformin		Zematop	78
hydrochloride	12	Zetlam	
Vimpat		Ziagen	
Vinblastine sulphate		Zidovudine [AZT]	
Vincristine sulphate		Zidovudine [AZT] with	
Vinorelbine		lamivudine	116
		Ziextenzo	
Vinorelbine Ebewe			
Vinorelbine Te Arai		Ziextenzo AU	
Viramune Suspension		Zimybe	
ViruPOS		Zinc and castor oil	
Vit.D3		Zinc sulphate	
Vita-B12		Zincaps	
VitA-POS		Ziprasidone	
Vitabdeck	35	Zista	259
Vital	284	Zithromax	99
Vitamin B complex	34	Zo-Rub HP	128
Vitamin B6 25		Zo-Rub Osteo	121
Vitamins	34–35	Zoladex	
Vitarubin Depot Injection		Zoledronic acid	
Vivonex TEN		Hormone	88
Voltaren		Musculoskeletal	
Voltaren D		Zoledronic acid Viatris	124
			90
Voltaren SR		Hormone	
Volumatic		Musculoskeletal	
Voriconazole		Zopiclone	
Votrient		Zopiclone Actavis	
Vttack		Zostrix	
Vyvanse	149	Zostrix HP	
- W -		Zuclopenthixol decanoate	143
Warfarin sodium	45	Zuclopenthixol hydrochloride	140
Wart Preparations		Zusdone	
Wasp venom allergy treatmer		Zyban	
Water		Zypine	
Blood	<i>1</i> 7	Zypine ODT	
Extemporaneous		Zyprexa Relprevv	
White Soft Liquid Paraffin AF		Zytiga	
Wool fat with mineral oil	/5	Zyvox	110
- X -			
Xaluprine			
Xarelto	45		
Xifaxan	10		