# January 2025

#### **Editors:**

Kave Wilson & Doris Chong email: enquiry@pharmac.govt.nz Telephone +64 4 460 4990 Level 9, 40 Mercer Street PO Box 10 254 Wellington

**Freephone Information Line** 0800 66 00 50 (9am - 5pm weekdays)

#### Circulation

You can register to have an electronic version of the Pharmaceutical Schedule (link to PDF copy) emailed to your nominated email address each month by subscribing at pharmac.govt.nz/subscribe.

#### Production

Typeset automatically from XML and T<sub>F</sub>X. XML version of the Schedule available from schedule.pharmac.govt.nz/pub/schedule

### **Programmers**

Anrik Drenth

email: texschedule@pharmac.govt.nz ©Pharmaceutical Management Agency



ISSN 1179-3686

This work is licensed under the Creative Commons Attribution 4.0 International licence. In essence, you are free to copy, distribute and adapt it, as long as you attribute the work to Pharmac and abide by the other licence terms. To view a copy of this licence, visit: creativecommons.org/licenses/by/4.0/. Attribution to Pharmac should be in written form and not by reproduction of the Pharmac logo. While care has been taken in compiling this Schedule, Pharmac takes no responsibility for any errors or omissions, and shall not be liable for any consequences arising there from.

	oudonig i namuo	
Section A	General Rules	5
Section B	Alimentary Tract & Metabolism	6
	Blood & Blood Forming Organs	35
	Cardiovascular System	45
	Dermatologicals	67
	Genito Urinary System	77
	Hormone Preparations – Systemic	84
	Infections – Agents For Systemic Use	95
	Musculoskeletal System	117
	Nervous System	123
	Oncology Agents & Immunosuppressants	153
	Respiratory System & Allergies	257
	Sensory Organs	267
	Various	272
Section C	Extemporaneous Compounds (ECPs)	274
Section D	Special Foods	276
Section I	National Immunisation Schedule	301
	Index	318

Introducing Pharmac

## **Introducing Pharmac**

The Pharmaceutical Management Agency (Pharmac) makes decisions that help control Government spending on pharmaceuticals. This includes community pharmaceuticals, hospital pharmaceuticals, vaccines and increasingly, hospital medical devices. Pharmac negotiates prices, sets subsidy levels and conditions, and makes decisions on changes to the subsidised list.

#### Pharmac's role:

"to secure for eligible people in need of pharmaceuticals, the best health outcomes that are reasonably achievable from pharmaceutical treatment and from within the amount of funding provided."

Pae Ora (Healthy Futures) Act 2022

To ensure our decisions are as fair and robust as possible we use a decision-making process that incorporates clinical, economic and commercial issues. We also seek the views of users and the wider community through consultation. The processes we generally use are outlined in our Operating Policies and Procedures.

Further information about Pharmac and the way we make funding decisions can be found on the Pharmac website at <a href="https://pharmac.govt.nz/about">https://pharmac.govt.nz/about</a>.

## **Purpose of the Pharmaceutical Schedule**

The purpose of the Schedule is to list:

- the Community Pharmaceuticals that are subsidised by the Government and to show the amount of the subsidy paid to contractors, as well as the manufacturer's price and any access conditions that may apply;
- the Hospital Pharmaceuticals that may be used in Health NZ Hospitals, as well as any access conditions that may apply; and
- the Pharmaceuticals, including Medical Devices, used in Health NZ Hospitals for which national prices have been negotiated by Pharmac.

The Schedule does not show the final cost to Government of subsidising each Community Pharmaceutical or to Health NZ Hospitals in purchasing each Pharmaceutical, since that will depend on any rebate and other arrangements Pharmac has with the supplier and, for Pharmaceuticals used in Health NZ Hospitals, on any logistics arrangements put in place.

This book contains sections A to D and Section I of the Pharmaceutical Schedule and lists the Pharmaceuticals funded for use in the community, including vaccines, as well as haemophilia and cancer treatments given in Health NZ Hospitals. Section H lists the Pharmaceuticals that that can be used in Health NZ Hospitals and is a separate publication.


The Pharmaceuticals in this book are listed by therapeutic group, which is based on the WHO Anatomical Therapeutic Chemical (ATC) system. The listings are displayed alphabetically under each heading.

The index lists both chemical entities and product brand names.

## **Explaining pharmaceutical entries**

The Pharmaceutical Schedule lists pharmaceuticals subsidised by the Government, the subsidy, the supplier's price and the access conditions that may apply.

### Example



## Glossary

## **Units of Measure**

gramg kilogramkg international unitiu	mi mi mi
Abbreviations	
AmpouleAmp	Ge
CapsuleCap	Gr
Cream	Inf
DeviceDev	Ini
DispersibleDisp	Lic
EffervescentEff	Lo
EmulsionEmul	Oi
Enteric Coated EC	Sa

microgrammilligrammillilitre	mg
Gelatinous	
Granules	
Infusion	Inf
Injection	Inj
Liquid	Liq
Long Acting	LA
Ointment	Oint
Sachet	Sach

millimoleunit	
Solution	Supp Tab
Trans Dermal Delivery System	TDDS

Read the General Rules: https://pharmac.govt.nz/section-a.

## SECTION B: ALIMENTARY TRACT AND METABOLISM

SECTION B: ALIMENTARY TRACT AND MET	TABOLISM			
	Subsidy (Manufacturer's Price) \$	Su Per	Fully bsidised	Brand or Generic Manufacturer
Antacids and Antiflatulents				
Antacids and Reflux Barrier Agents				
ALGINIC ACID  Sodium alginate 225 mg and magnesium alginate 87.5 mg pe sachet		30	<b>√</b> G	aviscon Infant
* Tab 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg - peppermint flavour	1.80 (14.39)	60	G	aviscon Extra Strength
Oral liq 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg per 10 ml		500 ml	A	cidex
Phosphate Binding Agents				
ALUMINIUM HYDROXIDE  * Tab 600 mg  CALCIUM CARBONATE  Oral liq 1,250 mg per 5 ml (500 mg elemental per 5 ml) –		100		lu-Tab
Subsidy by endorsement	47.30	500 ml 473 ml	<b>√</b> C	oxane alcium carbonate PAI <sup>829</sup>
Only when prescribed for patients unable to swallow calc inappropriate and the prescription is endorsed accordingl		ts or whe	ere calciur	n carbonate tablets are
Antidiarrhoeals				
Agents Which Reduce Motility				
LOPERAMIDE HYDROCHLORIDE – Up to 30 cap available on a  * Tab 2 mg  * Cap 2 mg	10.75	400 400	✓ N ✓ <u>D</u>	odia iamide Relief
Rectal and Colonic Anti-inflammatories				
BUDESONIDE  Cap modified-release 3 mg - Special Authority see SA1886  below - Retail pharmacy	87.60	90	<b>✓</b> <u>B</u>	udesonide Te Arai

## ⇒SA1886 Special Authority for Subsidy

Initial application — (Crohn's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Mild to moderate ileal, ileocaecal or proximal Crohn's disease; and
- 2 Any of the following:
  - 2.1 Diabetes; or

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsid	ised	Generic
\$	Per	1	Manufacturer

continued...

- 2.2 Cushingoid habitus; or
- 2.3 Osteoporosis where there is significant risk of fracture; or
- 2.4 Severe acne following treatment with conventional corticosteroid therapy; or
- 2.5 History of severe psychiatric problems associated with corticosteroid treatment; or
- 2.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
- 2.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated).

Initial application — (collagenous and lymphocytic colitis (microscopic colitis)) from any relevant practitioner. Approvals valid for 6 months where patient has a diagnosis of microscopic colitis (collagenous or lymphocytic colitis) by colonoscopy with biopsies.

Initial application — (gut Graft versus Host disease) from any relevant practitioner. Approvals valid for 6 months where patient has a gut Graft versus Host disease following allogenic bone marrow transplantation\*.

Note: Indication marked with \* is an unapproved indication.

Initial application — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has autoimmune hepatitis\*: and
- 2 Patient does not have cirrhosis; and
- 3 Any of the following:
  - 3.1 Diabetes: or
  - 3.2 Cushingoid habitus; or
  - 3.3 Osteoporosis where there is significant risk of fracture; or
  - 3.4 Severe acne following treatment with conventional corticosteroid therapy; or
  - 3.5 History of severe psychiatric problems associated with corticosteroid treatment; or
  - 3.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
  - 3.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated); or3.8 Adolescents with poor linear growth (where conventional corticosteroid use may limit further growth).

Note: Indication marked with \* is an unapproved indication.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

#### HYDROCORTISONE ACETATE Rectal foam 10%, CFC-Free (14 applications)......57.09 15 g OP ✓ Colifoam HYDROCORTISONE ACETATE WITH PRAMOXINE HYDROCHLORIDE Topical aerosol foam, 1% with pramoxine hydrochloride 1%.......26.55 10 g OP ✓ Proctofoam S29 **MESALAZINE** 100 ✓ Asacol 100 ✓ Pentasa Tab 800 mg ......85.50 90 ✓ Asacol ✓ Asacol S29 S29 100 OP ✓ Pentasa ✓ Pentasa Enema 1 g per 100 ml .......41.30 7 20 ✓ Asacol

✓ Pentasa

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
OLSALAZINE				
Tab 500 mg	56.02	60	<b>✓</b>	Atnahs
				Olsalazine S29
	93.37	100	✓ [	Dipentum
Cap 250 mg	53.00	100	✓ [	Dipentum
SODIUM CROMOGLICATE Cap 100 mg	113.35	100	<b>√</b> F	Ralicrom
SULFASALAZINE  * Tab 500 mg	16.52	100	<b>√</b> 9	Salazopyrin
* Tab EC 500 mg		100		Salazopyrin EN

## **Local preparations for Anal and Rectal Disorders**

## **Antihaemorrhoidal Preparations**

FLUOCORTOLONE CAPROATE WITH FLUOCORTOLONE PIVALATE AND CII	NCHOCAINE	
Oint 950 mcg, with fluocortolone pivalate 920 mcg, and		
cinchocaine hydrochloride 5 mg per g13.05	30 g OP	<ul><li>Ultraproct</li></ul>
Suppos 630 mcg, with fluocortolone pivalate 610 mcg, and		
cinchocaine hydrochloride 1 mg8.61	12	<ul><li>Ultraproct</li></ul>
HYDROCORTISONE WITH CINCHOCAINE		
Oint 5 mg with cinchocaine hydrochloride 5 mg per g15.00	30 g OP	✓ Proctosedyl
Suppos 5 mg with cinchocaine hydrochloride 5 mg per g9.90	12	✓ Proctosedyl

## Management of Anal Fissures

GLYCERYL TRINITRATE − Special Authority see SA1329 below − Retail pharmacy

★ Oint 0.2%......22.00 30 g OP

✓ Rectogesic

## ⇒SA1329 Special Authority for Subsidy

CL VCODVDDONII IM BDOMIDE

**Initial application** from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has a chronic anal fissure that has persisted for longer than three weeks.

## **Antispasmodics and Other Agents Altering Gut Motility**

GLYCOPYRHONIUM BROMIDE		
Inj 200 mcg per ml, 1 ml ampoule  – Up to 10 inj available on a PSO19.00	5	✓ Robinul
HYOSCINE BUTYLBROMIDE		
* Tab 10 mg2.25	20	✓ Hyoscine Butylbromide (Adiramedica)
6.35	100	✓ Buscopan
Hyoscine Butylbromide (Adiramedica) to be Principal Supply on 1 April 2025	5	
* Inj 20 mg, 1 ml – Up to 5 inj available on a PSO1.91 (Buscopan Tab 10 mg to be delisted 1 April 2025)	5	✓ <u>Spazmol</u>
MEBEVERINE HYDROCHLORIDE		
* Tab 135 mg8.50	90	✓ Colofac

ALIMENTARY TRACT AND METABOLISM Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer **Antiulcerants** Antisecretory and Cytoprotective MISOPROSTOL - Wastage claimable \* Tab 200 mcg - Up to 120 tab available on a PSO .......47.73 ✓ Cytotec 120 **Helicobacter Pylori Eradication CLARITHROMYCIN** 14 ✓ Klacid a) Maximum of 28 tab per prescription b) Subsidised only if prescribed for helicobacter pylori eradication and prescription is endorsed accordingly. Note: the prescription is considered endorsed if clarithromycin is prescribed in conjunction with a proton pump inhibitor and either amoxicillin or metronidazole. **H2 Antagonists** FAMOTIDINE - Only on a prescription ✓ Famotidine 100 Hovid S29 100 ✓ Famotidine Hovid S29 Inj 10 mg per ml, 4 ml - Subsidy by endorsement ......CBS 10 ✓ Mylan S29 Subsidy by endorsement – Subsidised for patients receiving treatment as part of palliative care. **Proton Pump Inhibitors** 

400	<b>4</b> 1 15 11 4
100	✓ Lanzol Relief
100	✓ Lanzol Relief
90	Omeprazole Teva
	✓ Omeprazole actavis 10
90	✓ Omeprazole Teva
	✓ Omeprazole actavis 20
90	✓ Omeprazole Teva
	✓ Omeprazole actavis 40
5 g	✓ Midwest
· ·	
5	✓ <u>Dr Reddy's</u> <u>Omeprazole</u>
	✓ Ocicure S29
90	✓ Panzop Relief
90	✓ Panzop Relief
	90 90 90 5 g 5

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price \$	e) Per	Fully Subsidised	Brand or Generic Manufacturer
Site Protective Agents				
COLLOIDAL BISMUTH SUBCITRATE  Tab 120 mgSUCRALFATE	14.51	50	<b>✓</b> G	astrodenol \$29
Tab 1 g	35.50 (48.28)	120	С	arafate
Bile and Liver Therapy				
RIFAXIMIN – Special Authority see SA1461 below – Retail pl Tab 550 mg  SA1461 Special Authority for Subsidy	,	56	<b>√</b> <u>X</u>	<u>ifaxan</u>
Initial application only from a gastroenterologist, hepatologist nepatologist. Approvals valid for 6 months where the patient tolerated doses of lactulose.  Renewal only from a gastroenterologist, hepatologist or Pract nepatologist. Approvals valid without further renewal unless repending from treatment.	has hepatic encephalo	oathy d	lespite an ad n of a gastro	lequate trial of maximum enterologist or
Diabetes				
Hyperglycaemic Agents	la a mara a su			
DIAZOXIDE - Special Authority see SA1320 below - Retail p Cap 25 mg Cap 100 mg Oral liq 50 mg per ml  → SA1320 Special Authority for Subsidy	110.00	100 100 30 ml C	<b>✓</b> P	roglicem \$29 roglicem \$29 5 Pharma \$29
nitial application from any relevant practitioner. Approvals hypoglycaemia caused by hyperinsulinism.	valid for 12 months whe	ere use	d for the trea	atment of confirmed
Renewal from any relevant practitioner. Approvals valid with appropriate and the patient is benefiting from treatment.  GLUCAGON HYDROCHLORIDE	out further renewal unle	ess noti	ified where th	he treatment remains
Inj 1 mg syringe kit – Up to 5 kit available on a PSO	32.00	1	<b>✓</b> G	lucagen Hypokit
Insulin - Short-acting Preparations				
NSULIN NEUTRAL ▲ Inj human 100 u per ml	25.26	10 ml C		ctrapid lumulin R
▲ Inj human 100 u per ml, 3 ml	42.66	5	✓ A	ctrapid Penfill umulin R
Insulin - Intermediate-acting Preparations				
NSULIN ASPART WITH INSULIN ASPART PROTAMINE  Inj 100 iu per ml, 3 ml prefilled pen	52.15	5	✓ N	ovoMix 30 FlexPen

	Subsidy		Fully Brand or
	(Manufacturer's F		idised Generic
	\$	Per	✓ Manufacturer
NSULIN ISOPHANE			
Inj human 100 u per ml		10 ml OP	<ul><li>✓ Humulin NPH</li><li>✓ Protaphane</li></ul>
Inj human 100 u per ml, 3 ml	29.86	5	<ul><li>✓ Humulin NPH</li><li>✓ Protaphane Penfill</li></ul>
NSULIN ISOPHANE WITH INSULIN NEUTRAL			
Inj human with neutral insulin 100 u per ml	25.26	10 ml OP	<ul><li>✓ Humulin 30/70</li><li>✓ Mixtard 30</li></ul>
Inj human with neutral insulin 100 u per ml, 3 ml	42.66	5	<ul><li>✓ Humulin 30/70</li><li>✓ PenMix 30</li><li>✓ PenMix 50</li></ul>
Mixtard 30 Inj human with neutral insulin 100 u per ml to be delis PenMix 50 Inj human with neutral insulin 100 u per ml, 3 ml to be			
NSULIN LISPRO WITH INSULIN LISPRO PROTAMINE			
▲ Inj lispro 25% with insulin lispro protamine 75% 100 u per ml, 3 ml		5	✓ Humalog Mix 25
▲ Inj lispro 50% with insulin lispro protamine 50% 100 u per ml, 3 ml		5	✓ Humalog Mix 50
Insulin - Long-acting Preparations			
NSULIN GLARGINE			
Inj 100 u per ml, 10 ml	63.00	1	✓ Lantus
Inj 100 u per ml, 3 ml		5	✓ Lantus
Inj 100 u per ml, 3 ml disposable pen		5	✓ Lantus SoloStar
Insulin - Rapid Acting Preparations			
NSULIN ASPART			
Inj 100 u per ml, 10 ml	30.03	1	✓ NovoRapid
Inj 100 u per ml, 3 ml	51.19	5	✓ NovoRapid Penfill
Inj 100 u per ml, 3 ml syringe	51.19	5	NovoRapid FlexPen
NSULIN GLULISINE			
Inj 100 u per ml, 10 ml	27.03	1	✓ Apidra
Inj 100 u per ml, 3 ml	46.07	5	✓ Apidra
Inj 100 u per ml, 3 ml disposable pen	46.07	5	✓ Apidra SoloStar
NSULIN LISPRO			
Inj 100 u per ml, 10 ml	34.92	10 ml OP	✓ Humalog
Inj 100 u per ml, 3 ml		5	✓ Humalog
Alpha Glucosidase Inhibitors			
ACARBOSE			
★ Tab 50 mg Accarb to be Principal Supply on 1 February 2025	11.20	90	✓ Accarb
* Tab 100 mgAccarb to be Principal Supply on 1 February 2025	17.38	90	✓ Accarb
Oral Hypoglycaemic Agents			
GLIBENCLAMIDE			
<b>≮</b> Tab 5 mg	7.50	100	✓ Daonil

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	
GLICLAZIDE			_	
* Tab 80 mg	20.10	500	•	Glizide
GLIPIZIDE				
* Tab 5 mg	6.86	100	•	Minidiab
Minidiab to be Principal Supply on 1 March 2025				
METFORMIN HYDROCHLORIDE			_	
* Tab immediate-release 500 mg		1,000	_	Metformin Viatris
* Tab immediate-release 850 mg	11.28	500	•	Metformin Viatris
PIOGLITAZONE				
* Tab 15 mg		90	_	<u>Vexazone</u>
* Tab 30 mg		90	/	Vexazone
* Tab 45 mg	12.00	90	•	<u>Vexazone</u>
VILDAGLIPTIN				
Tab 50 mg	35.00	60	•	Galvus
VILDAGLIPTIN WITH METFORMIN HYDROCHLORIDE				
Tab 50 mg with 1,000 mg metformin hydrochloride	35.00	60	•	Galvumet
Tab 50 mg with 850 mg metformin hydrochloride	35.00	60	/	Galvumet

## **GLP-1 Agonists**

DULAGLUTIDE - Special Authority see SA2338 below - Retail pharmacy

Note: Not to be given in combination with a funded SGLT-2 inhibitor or other GLP-1 agonist.

## ⇒SA2338 Special Authority for Subsidy

Note: Subsidy for patients with existing approvals prior to 1 May 2024. Approvals valid without further renewal unless notified. No new patients will be granted from 1 May 2024 until further notice.

LIRAGLUTIDE - Special Authority see SA2339 below - Retail pharmacy

- a) Maximum of 9 inj per prescription
- b)
- a) Not to be given in combination with a funded SGLT-2 inhibitor or other GLP-1 agonist.
- b) Maximum of 1 pack of 3 (6 mg per ml, 3 ml) prefilled pens will be funded per month.

### ⇒SA2339 Special Authority for Subsidy

Note: Subsidy for patients with existing approvals prior to 1 May 2024. Approvals valid without further renewal unless notified. No new patients will be granted from 1 May 2024 until further notice.

### **SGLT2 Inhibitors**

### ⇒SA2408 Special Authority for Subsidy

**Initial application** — (heart failure reduced ejection fraction) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

### All of the following:

- 1 Patient has heart failure: and
- 2 Patient is in NYHA functional class II or III or IV; and
- 3 Either:
  - 3.1 Patient has a documented left ventricular ejection fraction (LVEF) of less than or equal to 40%; or

Subsidy (Manufacturer's Price)		Fully	Brand or Generic
 \$	Per	1	Manufacturer

continued...

- 3.2 An ECHO is not reasonably practicable, and in the opinion of the treating practitioner the patient would benefit from treatment; and
- 4 Patient is receiving concomitant optimal standard funded chronic heart failure treatment.

Initial application — (Type 2 Diabetes) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

- Either:
  - 1 Patient has previously received an initial approval for a GLP-1 agonist; or
  - 2 All of the following:
    - 2.1 Patient has type 2 diabetes; and
    - 2.2 Any of the following:
      - 2.2.1 Patient is Maori or any Pacific ethnicity\*; or
      - 2.2.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)\*; or
      - 2.2.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator\*; or
      - 2.2.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult\*; or
      - 2.2.5 Patient has diabetic kidney disease (see note b)\*; and

EMPAGLIFLOZIN - Special Authority see SA2408 on the previous page - Retail pharmacy

2.3 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of at least one blood-glucose lowering agent (e.g. metformin, vildagliptin, or insulin) for at least 3 months.

Notes: \* Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

- a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack, ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.
- b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m2 in the presence of diabetes, without alternative cause.
- c) Funded [empagliflozin / empagliflozin with metformin hydrochloride] treatment is not to be given in combination with a funded GLP-1 unless receiving (empagliflozin / empagliflozin with metformin hydrochloride] for the treatment of heart failure.

	Tab 10 mg		30 <sup>°</sup> 30	<ul><li>✓ Jardiance</li><li>✓ Jardiance</li></ul>
	PAGLIFLOZIN WITH METFORMIN HYDROCHLORIDE rmacy	- Special Authority see	SA2408	on the previous page - Retail
	Tab 5 mg with 1,000 mg metformin hydrochloride	58.56	60	✓ Jardiamet
	Tab 5 mg with 500 mg metformin hydrochloride		60	✓ Jardiamet
	Tab 12.5 mg with 1,000 mg metformin hydrochloride		60	✓ Jardiamet
*	Tab 12.5 mg with 500 mg metformin hydrochloride	58.56	60	<ul><li>Jardiamet</li></ul>

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer

## **Diabetes Management**

## **Ketone Testing**

BLOOD KETONE DIAGNOSTIC TEST STRIP - Subsidy by endorsement

- a) Not on a BSO
- b) Maximum of 20 strip per prescription
- c) Up to 10 strip available on a PSO
- d) Patient has any of the following:
  - 1) type 1 diabetes; or
  - 2) permanent neonatal diabetes: or
  - 3) undergone a pancreatectomy; or
  - 4) cystic fibrosis-related diabetes; or
  - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly.

## **Dual Blood Glucose and Blood Ketone Testing**

DUAL BLOOD GLUCOSE AND BLOOD KETONE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A dual blood glucose and blood ketone diagnostic test meter is subsidised for a patient who has:
  - 1) type 1 diabetes; or
  - 2) permanent neonatal diabetes; or
  - 3) undergone a pancreatectomy; or
  - 4) cystic fibrosis-related diabetes; or
  - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly. Only 1 meter per patient will be subsidised (no repeat prescriptions). For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Meter with 50 lancets, a lancing device and 10 blood glucose

				_
Subsidy	F	ully	Brand or	
(Manufacturer's Price)	Subsid	sed	Generic	
\$	Per	•	Manufacturer	

## **Blood Glucose Testing**

BLOOD GLUCOSE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A diagnostic blood glucose test meter is subsidised for a patient who:
  - 1) is receiving insulin or sulphonylurea therapy; or
  - 2) is pregnant with diabetes: or
  - 3) is on home TPN at risk of hypoglycaemia or hyperglycaemia; or
  - 4) has a genetic or an acquired disorder of glucose homeostasis, excluding type 1 or type 2 diabetes and metabolic syndrome.

The prescription must be endorsed accordingly. Only one CareSens meter per patient will be subsidised (no repeat prescriptions). Patients already using the CareSens N POP meter and CareSens N meter are not eligible for a new meter, unless they have:

- 1) type 1 diabetes; or
- 2) permanent neonatal diabetes: or
- 3) undergone a pancreatectomy; or
- 4) cystic fibrosis-related diabetes.

For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Note: Only 1 meter available per PSO

BLOOD GLUCOSE DIAGNOSTIC TEST STRIP - Up to 50 test available on a PSO

The number of test strips available on a prescription is restricted to 50 unless:

- Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the
  prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed; or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly; or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

Test strips	50 test OP	✓ CareSens N
		✓ CareSens PRO

## BLOOD GLUCOSE TEST STRIPS (VISUALLY IMPAIRED)

The number of test strips available on a prescription is restricted to 50 unless:

- Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the
  prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed; or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly: or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

Blood glucose test strips	33.69	50 test OP	✓ SensoCard
---------------------------	-------	------------	-------------

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	Subsidised	Generic	
\$	Per	/	Manufacturer	

## **Insulin Syringes and Needles**

Subsidy is available for disposable insulin syringes, needles, and pen needles if prescribed on the same form as the one used for the supply of insulin or liraglutide or when prescribed for a patient and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of insulin or liraglutide.

IIV	ocini i chi necoleo – maximum di 200 dev per prescripti	1011		
*	29 g × 12.7 mm	10.95	100	✓ B-D Micro-Fine
*	31 g × 5 mm		100	✓ B-D Micro-Fine
*	31 g × 6 mm	9.50	100	✓ Berpu
*	31 g × 8 mm		100	✓ B-D Micro-Fine
*	32 g × 4 mm		100	✓ B-D Micro-Fine
INS	SULIN SYRINGES, DISPOSABLE WITH ATTACHED NEED	LE - Maximum of 2	200 dev per p	orescription
*	Syringe 0.3 ml with 29 g x 12.7 mm needle	13.56	100	B-D Ultra Fine
		1.36	10	
		(1.99)		B-D Ultra Fine
*	Syringe 0.3 ml with 31 g x 8 mm needle	13.56	100	✓ B-D Ultra Fine II
	, ,	1.30	10	
		(1.99)		B-D Ultra Fine II
*	Syringe 0.5 ml with 29 g x 12.7 mm needle	13.56	100	✓ B-D Ultra Fine
	, ,	1.36	10	
		(1.99)		B-D Ultra Fine
*	Syringe 0.5 ml with 31 g x 8 mm needle	13.56	100	✓ B-D Ultra Fine II
	, ,	1.36	10	
		(1.99)		B-D Ultra Fine II
*	Syringe 1 ml with 29 g x 12.7 mm needle	13.56	100	✓ B-D Ultra Fine
	, ,	1.36	10	
		(1.99)		B-D Ultra Fine
*	Syringe 1 ml with 31 g × 8 mm needle	13.56	100	✓ B-D Ultra Fine II
		1.36	10	
		(1.99)		B-D Ultra Fine II
		` '		

## **Insulin Pumps**

INSULIN PUMP WITH ALGORITHM - Special Authority see SA2367 below - Retail pharmacy

- a) Maximum of 1 dev per prescription
- b) Only on a prescription

c) Maximum of 1 insulin pump per patient each four y	ear period.
Min basal rate 0.02 U/h	8,970.00
Min boool rate 0.1 LI/b	7 652 00

✓ mylife YpsoPump with CamAPS FX

✓ Tandem t:slim

X2 with Basal-IQ

✓ Tandem t:slim
X2 with Control-IQ

## **⇒SA2367** Special Authority for Subsidy

**Initial application — (type 1 diabetes)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

1 Any of the following:

continued...

1

(Manufacturer's Price) Subsidised Generic \$ Per ✓ Manufacturer	Subsidy	Full	y Brand or
\$ Per ✓ Manufacturer	(Manufacturer's	s Price) Subsidise	d Generic
	\$	Per <b>✓</b>	Manufacturer

continued...

- 1.1 The patient has type 1 diabetes; or
- 1.2 The patient has permanent neonatal diabetes or specific monogenic diabetes subtypes with insulin deficiency, considered by the treating endocrinologist as likely to benefit; or
- 1.3 The patient has Type 3c diabetes considered by the treating endocrinologist as likely to benefit (Type 3c diabetes includes insulin deficiency due to pancreatectomy, insulin deficiency secondary to cystic fibrosis or pancreatitis); or
- 1.4 The patient has atypical inherited forms of diabetes; and
- 2 Patient has been evaluated by a diabetes multidisciplinary team for their suitability for insulin pump therapy; and
- 3 In the opinion of the treating relevant practitioner the patient would benefit from an Automated Insulin Delivery (AID) system.

Renewal — (type 1 diabetes) from any relevant practitioner. Approvals valid for 6 months where the patient is continuing to derive benefit according to the treatment plan agreed at induction.

## **Insulin Pump Consumables**

## ⇒SA2380 Special Authority for Subsidy

**Initial application — (type 1 diabetes)** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 The patient has type 1 diabetes; or
  - 1.2 The patient has permanent neonatal diabetes or specific monogenic diabetes subtypes with insulin deficiency, considered by the treating endocrinologist as likely to benefit; or
  - 1.3 The patient has Type 3c diabetes considered by the treating endocrinologist as likely to benefit (Type 3c diabetes includes insulin deficiency due to pancreatectomy, insulin deficiency secondary to cystic fibrosis or pancreatitis); or
  - 1.4 The patient has atypical inherited forms of diabetes; and
- 2 Patient has been evaluated by a diabetes multidisciplinary team for their suitability for insulin pump therapy; and
- 3 In the opinion of the treating relevant practitioner the patient would benefit from an Automated Insulin Delivery (AID) system.

**Renewal — (type 1 diabetes)** from any relevant practitioner. Approvals valid for 2 years where the patient is continuing to derive benefit according to the treatment plan agreed at induction.

INSULIN PUMP CARTRIDGE - Special Authority see SA2380 above - Retail pharmacy

- a) Maximum of 5 sets per prescription
- b) Only on a prescription
- c) Maximum of 19 packs of cartridge sets will be funded per year.
- **★** Cartridge 300 U, t:lock × 10......86.00 1 OP **✓ Tandem Cartridge**

		Subsidy (Manufacturer's Price)	Per	Fully Subsidised	Brand or Generic Manufacturer
INS	SULIN PUMP INFUSION SET (STEEL CANNULA) - Special	•		ne previous	
	a) Maximum of 5 set per prescription	·		·	
	b) Only on a prescription				
*	c) Maximum of 19 infusion sets will be funded per year. 6 mm steel needle; 60 cm tubing × 10	120.00	1 OP	1	MiniMed Sure-T
~	o min steer needie, oo cin tubing x 10	150.00	1 01	•	MMT-864A
*	6 mm steel needle; 80 cm tubing × 10	130.00	1 OP	1	MiniMed Sure-T
	•				MMT-866A
*	8 mm steel needle; 60 cm tubing × 10	130.00	1 OP	/	MiniMed Sure-T MMT-874A
*	8 mm steel needle; 80 cm tubing × 10	130.00	1 OP	•	MiniMed Sure-T MMT-876A
(Mi	niMed Sure-T MMT-866A 6 mm steel needle; 80 cm tubing x niMed Sure-T MMT-874A 8 mm steel needle; 60 cm tubing x niMed Sure-T MMT-876A 8 mm steel needle; 80 cm tubing x	10 to be delisted 1 O 10 to be delisted 1 O	ctobe ctobe	r 2026) r 2026)	a CACCO on the provious
	SULIN PUMP INFUSION SET (STEEL CANNULA, STRAIGHT le – Retail pharmacy	INSERTION) – Spe	ecial A	uthority se	e SA2380 on the previous
μαί	a) Maximum of 5 sets per prescription				
	b) Only on a prescription				
	c) Maximum of 19 infusion sets will be funded per year.				
	5.5 mm steel cannula; straight insertion; 45 cm line × 10 with		4 00		
	10 needles	136.00	1 OP	•	mylife Orbit micro
	5.5 mm steel needle; straight insertion; 60 cm line × 10 with 10 needles	136 00	1 OP	1	mylife Orbit micro
*	5.5 mm steel needle; straight insertion; 80 cm line × 10 with		. 0.	-	mymo orbit mioro
	10 needles	136.00	1 OP	1	mylife Orbit micro
*	8.5 mm steel needle; straight insertion; 60 cm line $\times$ 10 with			_	
	10 needles	136.00	1 OP	•	mylife Orbit micro
*	8.5 mm steel needle; straight insertion; 80 cm line × 10 with 10 needles	136.00	1 OP	1	mylife Orbit micro

\* 6 mm steel cannula; straight insertion; 80 cm line x 10 with

\* 8 mm steel cannula; straight insertion; 80 cm line × 10 with

\* 6 mm steel cannula; straight insertion; 60 cm line x 10 with

\* 8 mm steel cannula; straight insertion; 60 cm line x 10 with

10 needles......182.00

1 OP

1 OP

1 OP

1 OP

✓ TruSteel

✓ TruSteel

✓ TruSteel

✓ TruSteel

		Subsidy		Fully	Brand or					
		(Manufacturer's Price) \$	Per	Subsidised	Generic Manufacturer					
_		T								
INS	INSULIN PUMP INFUSION SET (TEFLON CANNULA) – Special Authority see SA2380 on page 17 – Retail pharmacy									
	a) Maximum of 5 set per prescription									
	b) Only on a prescription									
*	c) Maximum of 19 infusion sets will be funded per year.  13 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	1	MiniMed Silhouette					
~	To minitenon needle, oo em tabing x to	100.00	1 01	•	MMT-381A					
*	17 mm teflon needle, 110 cm tubing × 10	130.00	1 OP	1	MiniMed Silhouette					
•	, , , , , , , , , , , , , , , , , , ,				MMT-377A					
*	17 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	1	MiniMed Silhouette					
	-				MMT-378A					
*	6 mm teflon needle, 110 cm tubing × 10	130.00	1 OP	✓	MiniMed Quick-Set					
					MMT-398A					
*	6 mm teflon needle, 45 cm blue tubing $\times$ 10	130.00	1 OP	•	MiniMed Mio					
					MMT-941A					
*	6 mm teflon needle, 45 cm pink tubing × 10	130.00	1 OP	•	MiniMed Mio MMT-921A					
*	6 mm toflan needle 60 em blue tubing 10	100.00	1 OP	./	MiniMed Mio					
不	6 mm teflon needle, 60 cm blue tubing x 10	130.00	I OF	•	MMT-943A					
*	6 mm teflon needle, 60 cm pink tubing × 10	130.00	1 OP	1	MiniMed Mio					
	o him tollon noodio, oo on piin tabiilg x ro		. 0.		MMT-923A					
*	6 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	1	MiniMed Quick-Set					
	·				MMT-399A					
*	6 mm teflon needle, 80 cm blue tubing	130.00	1 OP	✓	MiniMed Mio					
					MMT-945A					
*	6 mm teflon needle, 80 cm clear tubing × 10	130.00	1 OP	✓	MiniMed Mio					
					MMT-965A					
*	6 mm teflon needle, 80 cm pink tubing x 10	130.00	1 OP	•	MiniMed Mio					
*	0 mm toflan needle 110 em tubing 10	100.00	1 OP	./	MMT-925A MiniMed Quick-Set					
不	9 mm teflon needle, 110 cm tubing × 10	130.00	I OF	•	MMT-396A					
*	9 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	1	MiniMed Quick-Set					
-,-	o min tonon noodio, oo om tabing x 10		. 0.	•	MMT-397A					
*	9 mm teflon needle, 80 cm clear tubing × 10	130.00	1 OP	1	MiniMed Mio					
	,				MMT-975A					

(MiniMed Silhouette MMT-381A 13 mm teflon needle, 60 cm tubing × 10 to be delisted 1 October 2026) (MiniMed Silhouette MMT-377A 17 mm teflon needle, 110 cm tubing × 10 to be delisted 1 October 2026) (MiniMed Silhouette MMT-378A 17 mm teflon needle, 60 cm tubing × 10 to be delisted 1 October 2026) (MiniMed Quick-Set MMT-398A 6 mm teflon needle, 110 cm tubing × 10 to be delisted 1 October 2026) (MiniMed Mio MMT-941A 6 mm teflon needle, 45 cm blue tubing × 10 to be delisted 1 October 2026) (MiniMed Mio MMT-921A 6 mm teflon needle, 45 cm pink tubing × 10 to be delisted 1 October 2026) (MiniMed Mio MMT-943A 6 mm teflon needle, 60 cm blue tubing × 10 to be delisted 1 October 2026) (MiniMed Mio MMT-933A 6 mm teflon needle, 60 cm pink tubing × 10 to be delisted 1 October 2026) (MiniMed Mio MMT-945A 6 mm teflon needle, 60 cm blue tubing × 10 to be delisted 1 October 2026) (MiniMed Mio MMT-945A 6 mm teflon needle, 80 cm blue tubing to be delisted 1 October 2026) (MiniMed Mio MMT-965A 6 mm teflon needle, 80 cm clear tubing × 10 to be delisted 1 October 2026) (MiniMed Mio MMT-925A 6 mm teflon needle, 80 cm clear tubing × 10 to be delisted 1 October 2026) (MiniMed Mio MMT-936A 9 mm teflon needle, 110 cm tubing × 10 to be delisted 1 October 2026) (MiniMed Quick-Set MMT-396A 9 mm teflon needle, 60 cm tubing × 10 to be delisted 1 October 2026) (MiniMed Quick-Set MMT-397A 9 mm teflon needle, 60 cm tubing × 10 to be delisted 1 October 2026) (MiniMed Mio MMT-975A 9 mm teflon needle, 80 cm clear tubing × 10 to be delisted 1 October 2026)

		Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
INS	SULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE IN	ISERTION WITH INS	SERT	ON DEVICE	Special Authority see
	2380 on page 17 – Retail pharmacy				, ,
	a) Maximum of 5 sets per prescription				
	b) Only on a prescription				
	c) Maximum of 19 infusion sets will be funded per year.				
*	13 mm teflon cannula; angle insertion; insertion device; 110 c	m			
	line × 10 with 10 needles	182.00	1 OP	✓ A	utoSoft 30
*	13 mm teflon cannula; angle insertion; insertion device; 60 cn line × 10 with 10 needles	n	1 OP	✓ A	utoSoft 30
INIS	SULIN PUMP INFUSION SET (TEFLON CANNULA, FLEXIBLE	INSERTION WITH	INSFI	RTION DEV	ICE) - Special Authority
	SA2380 on page 17 – Retail pharmacy	INOLITION WITH	IIVOLI	THONDEV	opecial Authority
	a) Maximum of 5 set per prescription				
	b) Only on a prescription				
	c) Maximum of 19 infusion sets will be funded per year.				
*	6 mm teflon cannula; flexible insertion; insertion device; 46 cr	n			
•	line × 10 with 10 needles		1 OP	✓ m	ylife Inset soft
*	6 mm teflon cannula; flexible insertion; insertion device; 60 cr	n			•
•	line with integrated inserter × 10 with 10 needles		1 OP	✓ m	ylife Inset soft
*	6 mm teflon cannula; flexible insertion; insertion device; 80 cr				,
•	line × 10 with 10 needles		1 OP	✓ m	ylife Inset soft
*	9 mm teflon cannula; flexible insertion; insertion device; 60 cr				,
•	line × 10 with 10 needles		1 OP	✓ m	ylife Inset soft
*	9 mm teflon cannula; flexible insertion; insertion device; 80 cr				,
•	line × 10 with 10 needles		1 OP	✓ m	ylife Inset soft
INI	SULIN PUMP INFUSION SET (TEFLON CANNULA, STRAIGH		INICE		•
	e SA2380 on page 17 – Retail pharmacy	I INSLITTION WITH	IIVOL	ITTION DE	riot) - Special Authority
500	a) Maximum of 5 sets per prescription				
	b) Only on a prescription				
	c) Maximum of 19 infusion sets will be funded per year.				
*	6 mm teflon cannula; straight insertion; insertion device;				
•••	110 cm line × 10 with 10 needles	182.00	1 OP	✓ A	utoSoft 90
*	6 mm teflon cannula; straight insertion; insertion device; 60 ci				
-,-	line × 10 with 10 needles		1 OP	<b>✓</b> Δ	utoSoft 90
*	9 mm teflon cannula; straight insertion; insertion device;			- 70	4.000.1.00
-,-	110 cm line × 10 with 10 needles	182 00	1 OP	✓ A	utoSoft 90
*	9 mm teflon cannula; straight insertion; insertion device; 60 cr			- 70	4.0001.00
-,-	line × 10 with 10 needles		1 OP	<b>✓</b> Δ	utoSoft 90
INIC			. •.		
	SULIN PUMP INFUSION SET (TEFLON CANNULA, VARIABLI tail pharmacy	E INSERTION) - SP	eciai	Authority See	e SA2360 on page 17 -
ne	a) Maximum of 5 set per prescription				
	b) Only on a prescription				
	c) Maximum of 19 infusion sets will be funded per year.				
*	13 mm teflon cannula; variable insertion; 60 cm line × 10 with				
*	10 needles		1 OP	✓ V	ariSoft
	TO TICCUICS	102.00	1 01	- V	anoon

MMT-332A

	Subsidy (Manufacturer's Price) \$	Su Per	Fully Brand or ibsidised Generic Manufacture	r
INSULIN PUMP RESERVOIR - Special Authority see SA2380	on page 17 – Retail ph	armacy		
a) Maximum of 9 sets per prescription				
b) Only on a prescription				
<ul> <li>c) Maximum of 36 packs of resevoir sets will be funded per</li> </ul>	,			
* 10 × 1.6 ml glass reservoir for YpsoPump	50.00	1 OP	✓ mylife YpsoP  Reservoir	ump
* 10 × luer lock conversion cartridges 1.8 ml for Paradigm p	umps50.00	1 OP	✓ ADR Cartridg	e 1.8
* Cartridge for 7 series pump; 3.0 ml x 10	50.00	1 OP	✓ MiniMed	
			3.0 Reservo	oir

(ADR Cartridge 1.8 10 × luer lock conversion cartridges 1.8 ml for Paradigm pumps to be delisted 1 October 2026) (MiniMed 3.0 Reservoir MMT-332A Cartridge for 7 series pump; 3.0 ml × 10 to be delisted 1 October 2026)

### Continuous Glucose Monitor

CONTINUOUS GLUCOSE MONITOR (INTEROPERABLE) – Special Authority see SA2371 below – Retail pharmacy Only on a prescription

Maximum of 28 dev will be funded per year.

\* Sensor (9) and transmitter (Dexcom G6) – Maximum of 1 dev

## ⇒SA2371 Special Authority for Subsidy

Initial application — (type 1 diabetes) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

#### Both:

- 1 Any of the following:
  - 1.1 The patient has type 1 diabetes; or
  - 1.2 The patient has permanent neonatal diabetes or specific monogenic diabetes subtypes with insulin deficiency, considered by the treating endocrinologist or relevant secondary health care professional as practicable, as likely to benefit; or
  - 1.3 The patient has Type 3c diabetes considered by the treating endocrinologist or relevant secondary health care professional as practicable, as likely to benefit (Type 3c diabetes includes insulin deficiency due to pancreatectomy, insulin deficiency secondary to cystic fibrosis or pancreatitis); or
  - 1.4 The patient has atypical inherited forms of diabetes; and
- 2 In the opinion of the treating relevant practitioner the patient would benefit from an Automated Insulin Delivery (AID) system.

**Renewal** — (type 1 diabetes) from any relevant practitioner. Approvals valid for 2 years where the patient is continuing to derive benefit according to the treatment plan agreed at induction.

CONTINUOUS GLUCOSE MONITOR (STANDALONE) - Special Authority see SA2370 on the next page - Retail pharmacy Only on a prescription

★ Sensor (Dexcom ONE+) - Maximum of 9 dev per prescription ......81.00
 Maximum of 40 dev will be funded per year.

★ Sensor (Freestyle Libre 2) – Maximum of 7 dev per prescription.....92.83
1
✓ Freestyle Libre 2
Maximum of 29 dev will be funded per year.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

## ⇒SA2370 Special Authority for Subsidy

Initial application — (type 1 diabetes) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 The patient has type 1 diabetes; or
- 2 The patient has permanent neonatal diabetes or specific monogenic diabetes subtypes with insulin deficiency, considered by the treating endocrinologist or relevant secondary health care professional as practicable, as likely to benefit; or
- 3 The patient has Type 3c diabetes considered by the treating endocrinologist or relevant secondary health care professional as practicable, as likely to benefit (Type 3c diabetes includes insulin deficiency due to pancreatectomy, insulin deficiency secondary to cystic fibrosis or pancreatitis); or
- 4 The patient has atypical inherited forms of diabetes.

Renewal — (type 1 diabetes) from any relevant practitioner. Approvals valid for 2 years where the patient is continuing to derive benefit according to the treatment plan agreed at induction.

## **Digestives Including Enzymes**

#### PANCREATIC ENZYME

Cap pancreatin 150 mg (amylase 8,000 Ph Eur U, lipase			
10,000 Ph Eur U, total protease 600 Ph Eur U)3	34.93	100	✓ Creon 10000
Cap pancreatin 300 mg (amylase 18,000 Ph Eur U, lipase			
25,000 Ph Eur U, total protease 1,000 Ph Eur U)9	94.38	100	✓ Creon 25000
Modified release granules pancreatin 60.12 mg (amylase			
3,600 Ph Eur U, lipase 5,000 Ph Eur U, protease 200 Ph			
Eur U)3	34.93	20 g OP	<ul><li>Creon Micro</li></ul>
URSODEOXYCHOLIC ACID - Special Authority see SA1739 below - R	letail pharma	cv	
Cap 250 mg	33.95	100	✓ Ursosan

## ⇒SA1739 Special Authority for Subsidy

Initial application — (Alagille syndrome or progressive familial intrahepatic cholestasis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Either:

- 1 Patient has been diagnosed with Alagille syndrome: or
- 2 Patient has progressive familial intrahepatic cholestasis.

Initial application — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

#### All of the following:

- 1 Patient has chronic severe drug induced cholestatic liver injury; and
- 2 Cholestatic liver injury not due to Total Parenteral Nutrition (TPN) use in adults; and
- 3 Treatment with ursodeoxycholic acid may prevent hospital admission or reduce duration of stay.

Initial application — (Primary biliary cholangitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

#### Both:

- 1 Primary biliary cholangitis confirmed by antimitochondrial antibody titre (AMA) > 1:80, and raised cholestatic liver enzymes with or without raised serum IgM or, if AMA is negative, by liver biopsy; and
- 2 Patient not requiring a liver transplant (bilirubin > 100 umol/l; decompensated cirrhosis).

Initial application — (Pregnancy) from any relevant practitioner. Approvals valid for 6 months where the patient diagnosed with cholestasis of pregnancy.

Initial application — (Haematological Transplant) from any relevant practitioner. Approvals valid for 6 months for applications

Subsidy (Manufacturer's Price	) Per	Fully Subsidised	Brand or Generic Manufacturer	
•	rei		Manuacturer	

continued...

meeting the following criteria:

#### Both:

- 1 Patient at risk of veno-occlusive disease or has hepatic impairment and is undergoing conditioning treatment prior to allogenic stem cell or bone marrow transplantation; and
- 2 Treatment for up to 13 weeks.

Initial application — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Paediatric patient has developed abnormal liver function as indicated on testing which is likely to be induced by Total Parenteral Nutrition (TPN); and
- 2 Liver function has not improved with modifying the TPN composition.

Renewal — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 6 months where the patient continues to benefit from treatment.

Renewal — (Pregnancy/Primary biliary cholangitis) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months where the paediatric patient continues to require TPN and who is benefiting from treatment, defined as a sustained improvement in bilirubin levels

## Laxatives

## **Bulk-forming Agents**

ISPAGHULA (PSYLLIUM) HUSK - Only on a prescription			
M. Daviday fay aval asla	00.00	F00 = OD	

*	Powder for oral soln20.0	00	500 g OP	/	Konsyl-D
---	--------------------------	----	----------	---	----------

## **Faecal Softeners**

DC	COSATE SOCION - Only on a prescription			
*	Tab 50 mg	3.20	100	✓ Coloxyl
¥	Tob 100 mg	4.00	100	./ Calavul

★ Tab 120 mg .......4.98 100 ✓ Coloxy

DOCUSATE SODIUM WITH SENNOSIDES

Tab 50 mg with sennosides 8 mg.......3.50

Z00

Laxsol

POLOXAMER - Only on a prescription

## **Opioid Receptor Antagonists - Peripheral**

METHYLNALTREXONE BROMIDE - Special Authority see SA1691 below - Retail pharmacy

## ⇒SA1691 Special Authority for Subsidy

DOCURATE SODILIM Only on a procedintion

Initial application — (Opioid induced constipation) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient is receiving palliative care; and
- 2 Either:
  - 2.1 Oral and rectal treatments for opioid induced constipation are ineffective; or
  - 2.2 Oral and rectal treatments for opioid induced constipation are unable to be tolerated.

	Subsidy (Manufacturer's Price) \$	) Si Per	Fully ubsidised	Brand or Generic Manufacturer
Osmotic Laxatives				
GLYCEROL  * Suppos 2.8/4.0 g - Only on a prescription	10.39	20	<b>√</b> <u>L</u>	ax-suppositories Glycerol
LACTULOSE – Only on a prescription  * Oral liq 10 g per 15 ml  MACROGOL 3350 WITH POTASSIUM CHLORIDE, SODIUM BIO		500 ml SODIUN	_	aevolac DE
Powder for oral soln 13.125 g with potassium chloride 46.6 m sodium bicarbonate 178.5 mg and sodium chloride 350.7 SODIUM ACID PHOSPHATE — Only on a prescription Enema 16% with sodium phosphate 8%	7 mg8.50	30 1	_	lolaxole leet Phosphate
SODIUM CITRATE WITH SODIUM LAURYL SULPHOACETATE Enema 90 mg with sodium lauryl sulphoacetate 9 mg per ml, 5 ml		iption 50	✓ <u>N</u>	Enema licolette
Stimulant Laxatives				
BISACODYL – Only on a prescription  * Tab 5 mg  * Suppos 10 mg  Lax-Suppositories to be Principal Supply on 1 February 2	4.14	200 10	_	sisacodyl Viatris ax-Suppositories
SENNA – Only on a prescription  * Tab, standardised	2.17 (8.21) 0.43 (2.06)	100 20		enokot enokot
SODIUM PICOSULFATE - Special Authority see SA2053 below Oral soln 7.5 mg per ml	7.40 3	0 ml OP	_	Pulcolax SP Drop

Both:

- 1 The patient is a child with problematic constipation despite an adequate trial of other oral pharmacotherapies including macrogol where practicable; and
- 2 The patient would otherwise require a high-volume bowel cleansing preparation or hospital admission.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

## Metabolic Disorder Agents

ALGLUCOSIDASE ALFA - Special Authority see SA1986 below - Retail pharmacy ✓ Myozyme 1

## **⇒SA1986** Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

1 The patient is aged up to 24 months at the time of initial application and has been diagnosed with infantile Pompe disease;

continued...

and

- 2 Any of the following:
  - 2.1 Diagnosis confirmed by documented deficiency of acid alpha-glucosidase by prenatal diagnosis using chorionic villus biopsies and/or cultured amniotic cells; or
  - 2.2 Documented deficiency of acid alpha-glucosidase, and urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides; or
  - 2.3 Documented deficiency of acid alpha-glucosidase, and documented molecular genetic testing indicating a disease-causing mutation in the acid alpha-glucosidase gene (GAA gene); or
  - 2.4 Documented urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides, and molecular genetic testing indicating a disease-causing mutation in the GAA gene; and
- 3 Patient has not required long-term invasive ventilation for respiratory failure prior to starting enzyme replacement therapy (ERT); and
- 4 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by ERT or might be reasonably expected to compromise a response to ERT; and
- 5 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks.

**Renewal** only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks; and
- 3 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 4 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by ERT; and
- 5 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT: and
- 6 There is no evidence of life threatening progression of respiratory disease as evidenced by the needed for > 14 days of invasive ventilation; and
- 7 There is no evidence of new or progressive cardiomyopathy.

ARGININE - Special Authority see SA2042 below - Retail pharmacy			
Tab 1,000 mg	CBS	90	<ul><li>Clinicians</li></ul>
Cap 500 mg	CBS	50	✓ Solgar
Powder	CBS	400 g	✓ Biomed

### ⇒SA2042 Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to arginine supplementation.

**Renewal** only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to arginine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

### **⇒SA1987** Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has a confirmed diagnosis of homocystinuria; and
- 2 Any of the following:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 2.1 A cystathionine beta-synthase (CBS) deficiency; or
- 2.2 A 5,10-methylene-tetrahydrofolate reductase (MTHFR) deficiency; or
- 2.3 A disorder of intracellular cobalamin metabolism; and
- 3 An appropriate homocysteine level has not been achieved despite a sufficient trial of appropriate vitamin supplementation.

**Renewal** only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

## ⇒SA2039 Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to coenzyme Q10 supplementation.

**Renewal** only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to coenzyme Q10 supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

GALSULFASE – Special Authority see SA1988 below – Retail pharmacy
Inj 1 mg per ml, 5 ml vial.......2,234.00 1 ✓ Naglazyme

## ⇒SA1988 Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has been diagnosed with mucopolysaccharidosis VI; and
- 2 Either:
  - 2.1 Diagnosis confirmed by demonstration of N-acetyl-galactosamine-4-sulfatase (arylsulfatase B) deficiency by either enzyme activity assay in leukocytes or skin fibroblasts; or
  - 2.2 Detection of two disease causing mutations and patient has a sibling who is known to have mucopolysaccharidosis VI.

**Renewal** only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 3 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by Enzyme Replacement Therapy (ERT); and
- 4 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT.

### ⇒SA1623 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hunter Syndrome (mucopolysaccharidosis II); and
- 2 Fither:
  - 2.1 Diagnosis confirmed by demonstration of iduronate 2-sulfatase deficiency in white blood cells by either enzyme

(Manufacturer's Price) Subsidised Generic \$ Per ✓ Manufacturer	Subsidy	Fully	Brand or
\$ Per  Manufacturer	(Manufacturer's Price)		
	<u> </u>	Per 🗸	Manutacturer

continued...

assav in cultured skin fibroblasts: or

- 2.2 Detection of a disease causing mutation in the iduronate 2-sulfatase gene; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with idursulfase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT): and
- 5 Idursulfase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 weeks post-HSCT) at doses no greater than 0.5 mg/kg every week.

### ⇒SA1695 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hurler Syndrome (mucopolysacchardosis I-H); and
- 2 Either:
  - 2.1 Diagnosis confirmed by demonstration of alpha-L-iduronidase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts; or
  - 2.2 Detection of two disease causing mutations in the alpha-L-iduronidase gene and patient has a sibling who is known to have Hurler syndrome; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with laronidase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT): and
- 5 Laronidase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 post-HSCT) at doses no greater than 100 units/kg every week.

LEVOCARNITINE - Special Authority see SA2040 below - Retail pharmacy		
Tab 500 mgCBS	30	✓ Solgar
Cap 250 mgCBS	30	✓ Solgar
Cap 500 mgCBS	60	✓ Balance
	300	<ul><li>Metabolics</li></ul>
Oral liq 1 g per 10 mlCBS	118 ml	✓ Carnitor S29
, •,		✓ Novitium Sugar
		Free S29
Oral liq 500 mg per 10 mlCBS	300 ml	✓ Balance

## ⇒SA2040 Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to carnitine supplementation.

**Renewal** only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to carnitine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

RIBOFLAVIN - Special Authority see SA2041 on the nex	t page – Retail pharmacy		
Tab 100 mg	CBS	100	Country Life
Ü			✓ Puritan's Pride Vitamin B-2 100 mg \$29
Cap 100 mg	CBS	100	✓ Solgar

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

## ⇒SA2041 Special Authority for Subsidy

**Initial application** only from a metabolic physician or neurologist. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to riboflavin supplementation.

Renewal only from a metabolic physician or neurologist. Approvals valid for 24 months for applications meeting the following criteria:

#### Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to riboflavin supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

SAPROPTERIN DIHYDROCHLORIDE - Special Authority see SA1989 below - Retail pharmacy

## **⇒SA1989** Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 1 month for applications meeting the following criteria: All of the following:

- 1 Patient has phenylketonuria (PKU) and is pregnant or actively planning to become pregnant; and
- 2 Treatment with sapropterin is required to support management of PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

**Renewal** only from a metabolic physician or any relevant practitioner on the recommendation of a metabolic physician.

Approvals valid for 12 months for applications meeting the following criteria:

### All of the following:

- 1 Either:
  - 1.1 Following the initial one-month approval, the patient has demonstrated an adequate response to a 2 to 4 week trial of sapropterin with a clinically appropriate reduction in phenylalanine levels to support management of PKU during pregnancy; or
  - 1.2 On subsequent renewal applications, the patient has previously demonstrated response to treatment with sapropterin and maintained adequate phenylalanine levels to support management of PKU during pregnancy; and
- 2 Any of the following:
  - 2.1 Patient continues to be pregnant and treatment with sapropterin will not continue after delivery; or
  - 2.2 Patient is actively planning a pregnancy and this is the first renewal for treatment with sapropterin; or
  - 2.3 Treatment with sapropterin is required for a second or subsequent pregnancy to support management of their PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

SODIUM BENZOATE – Special Authority see SA1599 below – Retail pharmacy
Soln 100 mg per ml ......CBS

SODIUM BENZOATE – Special Authority see SA1599 below – Retail pharmacy

#### ⇒SA1599 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder.

**Renewal** only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

SODIUM PHENYLBUTYRATE - Special Authority see SA1990 on the next page - Retail pharmacy

Grans 483 mg per g......2,016.00 174 g OP ✓ Pheburane

100 ml

✓ Amzoate S29

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

## ⇒SA1990 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder involving a deficiency of carbamylphosphate synthetase, ornithine transcarbamylase or argininosuccinate synthetase.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

TAURINE - Special Authority see SA2043 below - Retail pharmacy

Cap 500 mg	CBS	50	✓ Solgar
Cap 1,000 mg	CBS	90	✓ Life Extension
Powder	CBS	300 g	<ul> <li>Life Extension</li> </ul>

### ⇒SA2043 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected specific mitochondrial disorder that may respond taurine supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has confirmed diagnosis of a specific mitochondrial disorder which responds to taurine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

TRIENTINE - Special Authority see SA2324 below - Retail pharmacy Cap 250 mg......2,022.00 100 **Trientine Waymade** 

## ⇒SA2324 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Wilson disease: and
- 2 Treatment with D-penicillamine has been trialled and discontinued because the person has experienced intolerable side effects or has not received sufficient benefit; and
- 3 Treatment with zinc has been trailled and discontinued because the person has experienced intolerable side effects or has not received sufficient benefit, or zinc is considered clinically inappropriate as the person has symptomatic liver disease and requires copper chelation.

### Gaucher's Disease

TALIGLUCERASE ALFA - Special Authority see SA2137 below - Retail pharmacy ✓ Elelvso

### ⇒SA2137 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has a diagnosis of symptomatic type 1 or type 3\* Gaucher disease confirmed by the demonstration of specific deficiency of glucocerebrosidase in leukocytes or cultured skin fibroblasts, and genotypic analysis; and
- 2 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by enzyme replacement therapy (ERT) or the disease might be reasonably expected to compromise a response to ERT; and
- 3 Any of the following:
  - 3.1 Patient has haematological complications of Gaucher disease; or
  - 3.2 Patient has skeletal complications of Gaucher disease; or
  - 3.3 Patient has significant liver dysfunction or hepatomegaly attributable to Gaucher disease; or
  - 3.4 Patient has reduced vital capacity from clinically significant or progressive pulmonary disease due to Gaucher

	Subsidy (Manufacturer's Prio \$	ce) Subs	Fully sidised	Brand or Generic Manufacturer
continued				
disease; or  3.5 Patient is a child and has experienced growt 6-12 month period; and	h failure with significant	decrease in p	ercentil	le linear growth over a
4 Taliglucerase alfa is to be administered at a dose no whole vial (200 units).	greater than 30 unit/kg	every other	week ro	unded to the nearest
Note: Indication marked with * is an unapproved indication Renewal only from a metabolic physician or any relevant p Approvals valid for 3 years for applications meeting the folk All of the following:	ractitioner on the recomi	mendation of	a meta	bolic physician.
<ol> <li>Patient has demonstrated a symptomatic improvem symptoms for which therapy was started; and</li> <li>Patient has demonstrated a clinically objective improvement.</li> </ol>		•		
liver and spleen size; and Radiological (MRI) signs of bone activity performed demonstrate no deterioration shown by the MRI, color adjusted dose; and	at two years since initiat	tion of treatm	ent, and	I five yearly thereafter,
4 Patient has not developed another medical condition				
ERT; and  5 Patient is adherent with regular treatment and taliglu every other week rounded to the nearest whole vial	ucerase alfa is to be adm	·		•
ERT; and 5 Patient is adherent with regular treatment and taliglu	ucerase alfa is to be adm	·		•
<ul><li>ERT; and</li><li>Patient is adherent with regular treatment and taligluevery other week rounded to the nearest whole vial</li></ul>	ucerase alfa is to be adm	·		•
ERT; and 5 Patient is adherent with regular treatment and taliglu every other week rounded to the nearest whole vial  Mouth and Throat	ucerase alfa is to be adm	·		•
ERT; and 5 Patient is adherent with regular treatment and taligluevery other week rounded to the nearest whole vial  Mouth and Throat  Agents Used in Mouth Ulceration  BENZYDAMINE HYDROCHLORIDE  Soln 0.15% – Higher subsidy of \$22.60 per 500 ml with	ucerase alfa is to be adm (200 units).	ninistered at a		•
ERT; and 5 Patient is adherent with regular treatment and taligluevery other week rounded to the nearest whole vial  Mouth and Throat  Agents Used in Mouth Ulceration  BENZYDAMINE HYDROCHLORIDE	ucerase alfa is to be adm (200 units).	·	a dose r	no greater than 30 unit/kç
ERT; and 5 Patient is adherent with regular treatment and taligluevery other week rounded to the nearest whole vial  Mouth and Throat  Agents Used in Mouth Ulceration  BENZYDAMINE HYDROCHLORIDE  Soln 0.15% - Higher subsidy of \$22.60 per 500 ml with Endorsement	cucerase alfa is to be adm (200 units).	ninistered at a	a dose r	no greater than 30 unit/kç
ERT; and 5 Patient is adherent with regular treatment and talight every other week rounded to the nearest whole vial  Mouth and Throat  Agents Used in Mouth Ulceration  BENZYDAMINE HYDROCHLORIDE  Soln 0.15% – Higher subsidy of \$22.60 per 500 ml with Endorsement	cucerase alfa is to be adm (200 units).	ninistered at a	a dose r	no greater than 30 unit/kç
ERT; and 5 Patient is adherent with regular treatment and talight every other week rounded to the nearest whole vial  Mouth and Throat  Agents Used in Mouth Ulceration  BENZYDAMINE HYDROCHLORIDE  Soln 0.15% – Higher subsidy of \$22.60 per 500 ml with Endorsement	cucerase alfa is to be adm (200 units).  h	500 ml	a dose r	no greater than 30 unit/kç
ERT; and 5 Patient is adherent with regular treatment and taligluevery other week rounded to the nearest whole vial  Mouth and Throat  Agents Used in Mouth Ulceration  BENZYDAMINE HYDROCHLORIDE  Soln 0.15% – Higher subsidy of \$22.60 per 500 ml with Endorsement	cucerase alfa is to be adm (200 units).  h	ninistered at a	a dose r	no greater than 30 unit/kg
ERT; and 5 Patient is adherent with regular treatment and talight every other week rounded to the nearest whole vial  Mouth and Throat  Agents Used in Mouth Ulceration  BENZYDAMINE HYDROCHLORIDE  Soln 0.15% – Higher subsidy of \$22.60 per 500 ml with Endorsement	h	500 ml a result of tre	Deatment	no greater than 30 unit/kg
ERT; and 5 Patient is adherent with regular treatment and talight every other week rounded to the nearest whole vial  Mouth and Throat  Agents Used in Mouth Ulceration  BENZYDAMINE HYDROCHLORIDE  Soln 0.15% – Higher subsidy of \$22.60 per 500 ml with Endorsement	h	500 ml a result of tre	Deatment	no greater than 30 unit/kg bifflam t for cancer, and the bitomahesive
ERT; and 5 Patient is adherent with regular treatment and talight every other week rounded to the nearest whole vial  Mouth and Throat  Agents Used in Mouth Ulceration  BENZYDAMINE HYDROCHLORIDE  Soln 0.15% — Higher subsidy of \$22.60 per 500 ml wit Endorsement	h	500 ml a result of tro 56 g OP 15 g OP	Deatment	no greater than 30 unit/kg bifflam t for cancer, and the
ERT; and 5 Patient is adherent with regular treatment and talight every other week rounded to the nearest whole vial  Mouth and Throat  Agents Used in Mouth Ulceration  BENZYDAMINE HYDROCHLORIDE  Soln 0.15% – Higher subsidy of \$22.60 per 500 ml with Endorsement	h	500 ml a result of tre	Deatment  S C	no greater than 30 unit/kg bifflam t for cancer, and the ctomahesive brabase
ERT; and 5 Patient is adherent with regular treatment and taligluevery other week rounded to the nearest whole vial  Mouth and Throat  Agents Used in Mouth Ulceration  BENZYDAMINE HYDROCHLORIDE  Soln 0.15% — Higher subsidy of \$22.60 per 500 ml wite Endorsement	h	500 ml a result of tro 56 g OP 15 g OP	Deatment  S C	no greater than 30 unit/kg bifflam t for cancer, and the bitomahesive

Oral gel 20 mg per g......5.19
Decozol to be Principal Supply on 1 February 2025

✓ Fungilin

✓ Decozol

20

40 g OP

AMPHOTERICIN B

**MICONAZOLE** 

	Subsidy (Manufacturer's F	Price) Subs	Fully Brand or sidised Generic
	\$	Per	✓ Manufacturer
NYSTATIN			_
Oral liq 100,000 u per ml	2.22	24 ml OP	✓ <u>Nilstat</u>
Vitamins			
Vitamin B			
HYDROXOCOBALAMIN			
* Inj 1 mg per ml, 1 ml ampoule – Up to 6 inj available on a PS	SO2.46	3	✓ Cobal-B12 \$29 ✓ Hydroxocobalamin Panpharma
	4.10	5	✓ Vita-B12 ✓ Cobalin-H \$29
	4.10	5	✓ Neo-Cytamen S29 829
	8.20	10	✓ Vitarubin Depot
(Vita-B12 Inj 1 mg per ml, 1 ml ampoule to be delisted 1 July 202	25)		Injection S29
PYRIDOXINE HYDROCHLORIDE			
a) No more than 100 mg per dose			
b) Only on a prescription		0.5	4 Mil. 1 70 00
* Tab 25 mg - No patient co-payment payable  * Tab 50 mg		90 500	✓ <u>Vitamin B6 25</u> ✓ Pyridoxine
* Tab 50 Hig	23.43	300	multichem
THIAMINE HYDROCHLORIDE - Only on a prescription			
* Tab 50 mg	4.65	100	✓ Thiamine multichem
VITAMIN B COMPLEX			
* Tab, strong, BPC	11.25	500	✓ Bplex
Vitamin C			
ASCORBIC ACID			
a) No more than 100 mg per dose			
b) Only on a prescription			_
* Tab 100 mg	12.50	500	✓ <u>Cvite</u>
Vitamin D			
ALFACALCIDOL			
* Cap 0.25 mcg	26.32	100	One-Alpha
atr. O I	07.00	400	✓ One-Alpha S29 S29
* Cap 1 mcg*  Oral drops 2 mcg per ml		100 20 ml OP	✓ One-Alpha ✓ One-Alpha
	00.00	20 ml OP	✓ One-Alpha
CALCITRIOL  * Cap 0.25 mcg	7.89	100	✓ <u>Calcitriol-AFT</u> ✓ Calcitriol-AFT
* Cap 0.5 mcg	13.68	100	S29 S29 ✓ Calcitriol-AFT
·		.50	✓ Calcitriol-AFT S29 S29
			OLJ VIII

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

	ALIMENTARY TRACT AND METABOLISM				
		Subsidy (Manufacturer's Pric	ce) Subs	Fully sidised	Brand or Generic Manufacturer
*	OLECALCIFEROL Cap 1.25 mg (50,000 iu) – Maximum of 12 cap per prescripti Coral liq 188 mcg per ml (7,500 iu per ml)		12 5 ml OP		Vit.D3 Clinicians
N	Multivitamin Preparations				
	IULTIVITAMIN RENAL - Special Authority see SA1546 below - € Cap		30	•	Clinicians Renal Vit
Ini the	SA1546 Special Authority for Subsidy litial application from any relevant practitioner. Approvals valid the following criteria: ither:	l without further re	newal unless	s notifie	ed for applications meeting
	<ol> <li>The patient has chronic kidney disease and is receiving eit</li> <li>The patient has chronic kidney disease grade 5, defined as 15 ml/min/1.73 m² body surface area (BSA).</li> </ol>				
	IULTIVITAMINS – Special Authority see SA1036 below – Retail F Powder		200 g OP	•	Paediatric Seravit
	»SA1036 Special Authority for Subsidy litial application from any relevant practitioner. Approvals valid	I without further re	newal unless	s notifie	ed where the patient has
inb	born errors of metabolism.				·
	enewal from any relevant practitioner. Approvals valid without f pproval for multivitamins.	urther renewal un	less notified	where	patient has had a previous
	ITAMINS	40.50	4 000		
	<ul> <li>₹ Tab (BPC cap strength)</li> <li>₹ Cap (fat soluble vitamins A, D, E, K) – Special Authority see</li> </ul>	18.50	1,000	•	<u>Mvite</u>
•	SA1720 below – Retail pharmacy	23.40	60	✓	Vitabdeck
Init	SA1720 Special Authority for Subsidy nitial application from any relevant practitioner. Approvals valid the following criteria: ny of the following:	I without further re	newal unless	s notific	ed for applications meeting
	Patient has cystic fibrosis with pancreatic insufficiency; or     Patient is an infant or child with liver disease or short gut s     Patient has severe malabsorption syndrome.	yndrome; or			
N	Minerals				
C	Calcium				
CA	ALCIUM CARBONATE				
	Tab 1.25 g (500 mg elemental) — Subsidy by endorsemental Subsidy by endorsemental Tab eff 1.25 g (500 mg elemental) — Subsidy by endorsemental Subsidiary Endorse		250 100		<u>Calci-Tab 500</u> Calcium 500 mg Hexal <sup>S29</sup>
	Subsidy by endorsement - Only when prescribed for pae	ediatric patients (<	5 years) whe	ere cal	cium carbonate oral liquid is

considered unsuitable.

\* Inj 10%, 10 ml ampoule......32.00

CALCIUM GLUCONATE

✓ Max Health - HameIn \$29

10

	Subsidy (Manufacturer's Price)	S Per	Fully Subsidised	Brand or Generic Manufacturer
lodine				
POTASSIUM IODATE  * Tab 253 mcg (150 mcg elemental iodine)	5.99	90	<b>√</b> <u>V</u>	NeuroTabs
Iron				
FERROUS FUMARATE  * Tab 200 mg (65 mg elemental)  Ferro-tab to be Principal Supply on 1 February 2025	3.49	100	<b>√</b> F	Ferro-tab
FERROUS FUMARATE WITH FOLIC ACID  * Tab 310 mg (100 mg elemental) with folic acid 350 mcg	5.98	100	<b>√</b> <u>F</u>	Ferro-F-Tabs
# Tab long-acting 325 mg (105 mg elemental)  * Oral liq 30 mg (6 mg elemental) per 1 ml	9.25	30 250 ml 500 ml	<b>✓</b> F	<del>Ferrograd</del> Ferro-Liquid Ferodan
IRON (AS FERRIC CARBOXYMALTOSE) – Special Authority so		etail ph		Ferinject

## SA2394 Special Authority for Subsidy

**Initial application — (Anaemia)** from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with anaemia; and
- 2 Any of the following:
  - 2.1 Serum ferritin level is 20 mcg/L or less; or
  - 2.2 Both:
    - 2.2.1 Serum ferritin is between 20 and 50 mcg/L: and
    - 2.2.2 C-Reactive Protein (CRP) is at least 5 mg/L; or
  - 2.3 Patient has chronic inflammatory disease with symptoms of anaemia despite normal iron levels; and
- 3 Any of the following:
  - 3.1 Oral iron treatment has proven ineffective; or
  - 3.2 Oral iron treatment has resulted in dose-limiting intolerance; or
  - 3.3 Rapid correction of anaemia is required.

Renewal — (Anaemia) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient continues to have anaemia with a serum ferritin level of 20 mcg/L, or less or between 20 and 50 mcg/L with CRP of at least 5 mg/L, or has chronic inflammatory disease with symptoms of anaemia despite normal iron levels; and
- 2 A trial (or re-trial) with oral iron is clinically inappropriate.

Initial application — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with iron-deficiency anaemia; and
- 2 Any of the following:
  - 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
  - 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or

Subsit		,
\$	Per	Manufacturer

continued...

- 2.3 Patient has symptomatic heart failure, chronic kidney disease stage 3 or more or active inflammatory bowel disease and a trial of oral iron is unlikely to be effective; or
- 2.4 Rapid correction of anaemia is required.

Renewal — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

1 Patient continues to have iron-deficiency anaemia; and

2 A re-trial with oral iron is clinically inappropriate.  IRON POLYMALTOSE  * Inj 50 mg per ml, 2 ml ampoule	5	✓ Ferrosig
Magnesium		
MAGNESIUM HYDROXIDE Suspension 8%	355 ml	✓ Phillips Milk of Magnesia \$29
MAGNESIUM SULPHATE  * Inj 2 mmol per ml, 5 ml ampoule	10	✓ Martindale
* Inj 2 minol per ml, 10 ml ampoule	10	✓ Inresa \$29
Zinc		
ZINC SULPHATE  * Cap 137.4 mg (50 mg elemental)11.00	100	✓ Zincaps

## **BLOOD AND BLOOD FORMING ORGANS**

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

## **Antianaemics**

## Hypoplastic and Haemolytic

## ⇒SA2266 Special Authority for Subsidy

**Initial application — (chronic renal failure)** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient in chronic renal failure: and
- 2 Haemoglobin is less than or equal to 100g/L; and
- 3 Any of the following:
  - 3.1 Both:
    - 3.1.1 Patient does not have diabetes mellitus: and
    - 3.1.2 Glomerular filtration rate is less than or equal to 30ml/min; or
  - 3.2 Both:
    - 3.2.1 Patient has diabetes mellitus; and
    - 3.2.2 Glomerular filtration rate is less than or equal to 45ml/min; or
  - 3.3 Patient is on haemodialysis or peritoneal dialysis.

Initial application — (myelodysplasia) from any specialist. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of myelodysplasia (MDS)\*; and
- 2 Has had symptomatic anaemia with haemoglobin < 100g/L and is red cell transfusion-dependent; and
- 3 Patient has very low, low or intermediate risk MDS based on the WHO classification based prognostic scoring system for myelodysplastic syndrome (WPSS); and
- 4 Other causes of anaemia such as B12 and folate deficiency have been excluded; and
- 5 Patient has a serum epoetin level of < 500 IU/L; and
- 6 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Note: Indication marked with \* is an unapproved indication

**Renewal** — **(chronic renal failure)** from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

**Renewal — (myelodysplasia)** from any specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient's transfusion requirement continues to be reduced with erythropoietin treatment; and
- 2 Transformation to acute myeloid leukaemia has not occurred; and
- 3 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Note: Indication marked with \* is an unapproved indication

### EPOETIN ALFA - Special Authority see SA2266 above - Retail pharmacy

Wastage claimable			
Inj 1,000 iu in 0.5 ml, syringe	250.00	6	Binocrit
Inj 2,000 iu in 1 ml, syringe		6	Binocrit
Inj 3,000 iu in 0.3 ml, syringe		6	Binocrit
Inj 4,000 iu in 0.4 ml, syringe		6	Binocrit
Inj 5,000 iu in 0.5 ml, syringe		6	<ul><li>Binocrit</li></ul>
Inj 6,000 iu in 0.6 ml, syringe	145.00	6	Binocrit
Inj 8,000 iu in 0.8 ml, syringe		6	Binocrit
Inj 10,000 iu in 1 ml, syringe		6	Binocrit
Inj 40,000 iu in 1 ml, syringe		1	<ul><li>Binocrit</li></ul>

## **BLOOD AND BLOOD FORMING ORGANS**

	Subsidy (Manufacturer's Price) \$		Fully dised	Brand or Generic Manufacturer
Megaloblastic				
FOLIC ACID  * Tab 0.8 mg	26.60	1,000		olic Acid multichem
* Tab 5 mg Oral liq 50 mcg per ml		100 5 ml OP	_	olic Acid Viatris iomed

## **Antifibrinolytics, Haemostatics and Local Sclerosants**

### EFTRENONACOG ALFA [RECOMBINANT FACTOR IX] - [Xpharm]

For patients with haemophilia B receiving prophylaxis treatment. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management group.

reaters Group in conjunction with the National Haemophili	ia ivianagement gro	up.	
Inj 250 iu vial	612.50	1	Alprolix
Inj 500 iu vial	1,225.00	1	✓ Alprolix
Inj 1,000 iu vial		1	✓ Alprolix
Inj 2,000 iu vial	4,900.00	1	✓ Alprolix
Inj 3,000 iu vial	7,350.00	1	✓ Alprolix
Inj 4,000 iu vial	9,800.00	1	✓ Alprolix
ELTROMBOPAG – Special Authority see SA1743 below – Ret	ail pharmacy		
Wastage claimable			
Tab 25 mg	1,550.00	28	Revolade
Tah 50 mg	3 100 00	28	✓ Revolade

### ⇒SA1743 Special Authority for Subsidy

**Initial application** — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has had a splenectomy; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Any of the following:
  - 3.1 Patient has a platelet count of 20,000 to 30,000 platelets per microlitre and has evidence of significant mucocutaneous bleeding; or
  - 3.2 Patient has a platelet count of less than or equal to 20,000 platelets per microlitre and has evidence of active bleeding; or
  - 3.3 Patient has a platelet count of less than or equal to 10,000 platelets per microlitre.

**Initial application** — (**idiopathic thrombocytopenic purpura - preparation for splenectomy**) only from a haematologist. Approvals valid for 6 weeks where the patient requires eltrombopage treatment as preparation for splenectomy.

Initial application — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a significant and well-documented contraindication to splenectomy for clinical reasons; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Either:
  - 3.1 Patient has immune thrombocytopenic purpura\* with a platelet count of less than or equal to 20,000 platelets per microliter: or
  - 3.2 Patient has immune thrombocytopenic purpura\* with a platelet count of 20,000 to 30,000 platelets per microlitre

Sub	sidy Fu	ully Brand or
(Manufactu		sed Generic
	Per Per	✓ Manufacturer

continued...

and significant mucocutaneous bleeding.

**Initial application — (severe aplastic anaemia)** only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Two immunosuppressive therapies have been trialled and failed after therapy of at least 3 months duration; and
- 2 Either:
  - 2.1 Patient has severe aplastic anaemia with a platelet count of less than or equal to 20,000 platelets per microliter; or
  - 2.2 Patient has severe aplastic anaemia with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

Renewal — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 12 months where the patient has obtained a response (see Note) from treatment during the initial approval or subsequent renewal periods and further treatment is required.

Note: Response to treatment is defined as a platelet count of > 30,000 platelets per microlitre.

Renewal — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

#### All of the following:

- 1 The patient's significant contraindication to splenectomy remains; and
- 2 The patient has obtained a response from treatment during the initial approval period; and
- 3 Patient has maintained a platelet count of at least 50,000 platelets per microlitre on treatment; and
- 4 Further treatment with eltrombopag is required to maintain response.

Renewal — (severe aplastic anaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

#### Both:

- 1 The patient has obtained a response from treatment of at least 20,000 platelets per microlitre above baseline during the initial approval period; and
- 2 Platelet transfusion independence for a minimum of 8 weeks during the initial approval period.

#### EMICIZUMAB - [Xpharm] - Special Authority see SA2272 below

Inj 30 mg in 1 ml vial	3,570.00	1	✓ Hemlibra
Inj 60 mg in 0.4 ml vial	· ·	1	✓ Hemlibra
Inj 105 mg in 0.7 ml vial	12,492.00	1	✓ Hemlibra
Inj 150 mg in 1 ml vial	17,846.00	1	✓ Hemlibra

# ⇒SA2272 Special Authority for Subsidy

Initial application — (Severe Haemophilia A with or without FVIII inhibitors) only from a haematologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has severe congenital haemophilia A with a severe bleeding phenotype (endogenous factor VIII activity less than or equal to 2%); and
- 2 Emicizumab is to be administered at a dose of no greater than 3 mg/kg weekly for 4 weeks followed by the equivalent of 1.5 mg/kg weekly.

# EPTACOG ALFA [RECOMBINANT FACTOR VIIA] - [Xpharm]

For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group. Rare Clinical Circumstances Brand of bypassing agent for > 14 days predicted use. Access to funded treatment for > 14 days predicted use is by named patient application to the Haemophilia Treaters Group, subject to access criteria.

Inj 1 mg syringe	1,178.30	1	✓ NovoSeven RT
Inj 2 mg syringe	2,356.60	1	✓ NovoSeven RT
Inj 5 mg syringe	5,891.50	1	✓ NovoSeven RT
Inj 8 mg syringe	9,426.40	1	✓ NovoSeven RT

Subsidy (Manufacturer's Price)	Fu Subsidise	,	
 \$	Per	<ul> <li>Manufacturer</li> </ul>	

#### FACTOR EIGHT INHIBITOR BYPASSING FRACTION - [Xpharm]

For patients with haemophilia. Preferred Brand of bypassing agent for > 14 days predicted use. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

	managed by the reasonsprima reduces droup in conjunction that the reasons	aaoop	aaagoo o
lr	j 500 U1,315.00	1	FEIBA NF
Ir	j 1,000 U2,630.00	1	✓ FEIBA NF
Ir	i 2.500 U	1	✓ FEIBA NF

#### MOROCTOCOG ALFA [RECOMBINANT FACTOR VIII] - [Xpharm]

For patients with haemophilia. Rare Clinical Circumstances Brand of short half-life recombinant factor VIII. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group, subject to criteria.

287.50	1	Xyntha
	1	Xyntha
1,150.00	1	Xyntha
	1	Xyntha
3,450.00	1	Xyntha
	575.00	575.00 1 1,150.00 1 2,300.00 1

#### NONACOG GAMMA, [RECOMBINANT FACTOR IX] - [Xpharm]

For patients with naemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

Inj 500 iu vial	435.00	1	✓ RIXUBIS
Inj 1,000 iu vial		1	✓ RIXUBIS
Inj 2,000 iu vial		1	✓ RIXUBIS
Ini 3.000 iu vial	·	1	✓ RIXUBIS

(RIXUBIS Inj 500 iu vial to be delisted 1 February 2025)

# OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (ADVATE) - [Xpharm]

For patients with haemophilia. Preferred Brand of short half-life recombinant factor VIII. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

Inj 250 iu vial	210.00	1	Advate
Inj 500 iu vial		1	✓ Advate
Inj 1,000 iu vial		1	✓ Advate
Inj 1,500 iu vial		1	✓ Advate
Inj 2,000 iu vial	•	1	✓ Advate
Ini 3.000 iu vial	•	1	✓ Advate

(Advate Inj 250 iu vial to be delisted 1 February 2025)

(Advate Inj 1,500 iu vial to be delisted 1 February 2025)

#### OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (KOGENATE FS) - [Xpharm]

For patients with haemophilia. Rare Clinical Circumstances Brand of short half-life recombinant factor VIII. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group, subject to criteria.

Inj 250 iu vial	237.50	1	✓ Kogenate FS
Inj 500 iu vial	475.00	1	✓ Kogenate FS
Inj 1,000 iu vial	950.00	1	✓ Kogenate FS
Inj 2,000 iu vial	1,900.00	1	✓ Kogenate FS
Inj 3,000 iu vial	2,850.00	1	✓ Kogenate FS

Fully

Brand or

Pytazen SR

✓ Ticagrelor Sandoz

Subsidy

	Subsidy (Manufactured Drice)		ully Brand or
	(Manufacturer's Price) \$	Subsidi Per	sed Generic  Manufacturer
RURIOCTOCOG ALFA PEGOL [RECOMBINANT FACTOR VIII]	- [Xpharm]		
For patients with haemophilia A receiving prophylaxis treatm		d treatment	is managed by the Haemophilia
Treaters Group in conjunction with the National Haemophilia Inj 250 iu vial		1	✓ Adynovate
Inj 500 iu vial		1	✓ Adynovate
Inj 1,000 iu vial		1	✓ Adynovate
Inj 2,000 iu vial	,	1	✓ Adynovate
(Adynovate Inj 250 iu vial to be delisted 1 February 2025)	2,400.00	•	Adynovate
(Adynovate Inj 500 iu vial to be delisted 1 February 2025)			
, , , , , , , , , , , , , , , , , , , ,			
SODIUM TETRADECYL SULPHATE  * Inj 3% 2 ml	20 50	5	
* Inj 3% 2 ml	(73.00)	5	Fibro-vein
TRANSVANIO ACID	(73.00)		Fibro-veiii
TRANEXAMIC ACID	40.45	00	/ Marray Diagram
Tab 500 mg		60	Mercury Pharma
	45.68	100	✓ Cyklokapron
Vitamin K			
PHYTOMENADIONE	0.00	-	/ Kanaldan MM
Inj 2 mg per 0.2 ml — Up to 5 inj available on a PSO		5	✓ Konakion MM
Inj 10 mg per ml, 1 ml - Up to 5 inj available on a PSO	9.21	5	✓ Konakion MM
Antithrombotic Agents			
Antiplatelet Agents			
ASPIRIN			
* Tab 100 mg	12.65	990	✓ Ethics Aspirin EC
CLOPIDOGREL			
* Tab 75 mg	5.07	84	✓ Arrow - Clopid
100 / 0 mg		U-T	- Allow - Olopiu

⇒SA1955 Special Authority for Subsidy

Initial application — (acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Initial application — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

1 Fither

DIPYRIDAMOI F

- 1.1 Patient has had a neurological stenting procedure\* in the last 60 days; or
- 1.2 Patient is about to have a neurological stenting procedure performed\*; and

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer	
----------------------------------------------------------------------------------------------	--

continued...

- 2 Either:
  - 2.1 Patient has demonstrated clopidogrel resistance using the P2Y12 (VerifyNow) assay or another appropriate platelet function assay and requires antiplatelet treatment with ticagrelor; or
  - 2.2 Either:
    - 2.2.1 Clopidogrel resistance has been demonstrated by the occurrence of a new cerebral ischemic event; or
    - 2.2.2 Clopidogrel resistance has been demonstrated by the occurrence of transient ischemic attack symptoms referable to the stent.

**Initial application** — (Percutaneous coronary intervention with stent deployment) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic\*\*.

**Initial application** — (Stent thrombosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has experienced cardiac stent thrombosis whilst on clopidogrel.

**Renewal — (subsequent acute coronary syndrome)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Renewal — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient is continuing to benefit from treatment; and
- 2 Treatment continues to be clinically appropriate.

**Renewal** — (**Percutaneous coronary intervention with stent deployment**) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic\*\*.

Notes: indications marked with \* are unapproved indications.

Note: \*\* Clopidogrel allergy is defined as a history of anaphylaxis, urticaria, generalised rash or asthma (in non-asthmatic patients) developing soon after clopidogrel is started and is considered unlikely to be caused by any other treatment.

Subsidised

Per

Fully

Brand or

Generic

Manufacturer

		manadatato	
Heparin and Antagonist Preparations			
ENOXAPARIN SODIUM - Special Authority see SA2152 below - Retail pharma	acv		
Inj 20 mg in 0.2 ml syringe	10	✓ Clexane	
Inj 40 mg in 0.4 ml syringe29.74	10	✓ Clexane	
Clexane to be Principal Supply on 1 February 2025			
Inj 60 mg in 0.6 ml syringe42.47	10	Clexane	
Clexane to be Principal Supply on 1 February 2025			
Inj 80 mg in 0.8 ml syringe56.62	10	Clexane	
Clexane to be Principal Supply on 1 February 2025			
Inj 100 mg in 1 ml syringe70.91	10	Clexane	
Clexane to be Principal Supply on 1 February 2025			
Inj 120 mg in 0.8 ml syringe88.11	10	Clexane Forte	
Clexane Forte to be Principal Supply on 1 February 2025			
Inj 150 mg in 1 ml syringe100.70	10	✓ Clexane Forte	
Clexane Forte to be Principal Supply on 1 February 2025			

Subsidy

(Manufacturer's Price)

# ⇒SA2152 Special Authority for Subsidy

**Initial application** — (**Pregnancy, Malignancy or Haemodialysis**) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patients pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Initial application — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 For the short-term treatment of venous thromboembolism prior to establishing a therapeutic level of oral anti-coagulant treatment: or
- 2 For the prophylaxis and treatment of venous thromboembolism in high risk surgery: or
- 3 To enable cessation/re-establishment of existing oral anticoagulant treatment pre/post surgery; or
- 4 For the prophylaxis and treatment of venous thromboembolism in Acute Coronary Syndrome surgical intervention; or
- 5 To be used in association with cardioversion of atrial fibrillation.

Initial application — (Short-term use during treatment of COVID-19 with nirmatrelvir with ritonavir) from any relevant practitioner. Approvals valid for 2 weeks for applications meeting the following criteria:

- All of the following:
  - 1 Patient is receiving an anticoagulation treatment that has drug/drug interactions with ritonavir that increases risk of bleeding; and
  - 2 Patient meets the Access Criteria for COVID-19 antivirals published on the Pharmac website\*; and
  - 3 Other antiviral treatments for COVID-19 have been considered and are not clinically suitable options.

Renewal — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patient's pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Renewal — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month where low molecular weight heparin treatment or prophylaxis is required for a second or subsequent event (surgery, ACS, cardioversion, or prior to oral anti-coagulation).

	Subsidy		Fully	
	(Manufacturer's Price)	_	Subsidised	
	\$	Per		Manufacturer
HEPARIN SODIUM				
Inj 1,000 iu per ml, 10 ml vial	127.44	25	1	Pfizer S29
Inj 1,000 iu per ml, 5 ml ampoule	25.49	10	✓	Wockhardt S29
	103.70		1	Wockhardt PSF S29
	127.44	50	✓	Pfizer
Inj 5,000 iu per ml, 5 ml vial	83.00	10	1	Heparin Sodium
				<u>Panpharma</u>
Inj 5,000 iu per ml, 1 ml	70.33	5	✓	Hospira
Inj 25,000 iu per ml, 0.2 ml	22.42	5	✓	Hospira
	42.40		1	Heparin DBL S29
	482.20	50	✓	Heparin DBL \$29
HEPARINISED SALINE				
Inj 10 iu per ml, 5 ml	96.91	50	✓	Pfizer
• • •				
Oral Anticoagulants				
DABIGATRAN				
Cap 75 mg - No more than 2 cap per day	27.99	60	✓	Pradaxa
Cap 110 mg	27.99	60	✓	Pradaxa
Cap 150 mg	27.99	60	1	Pradaxa
RIVAROXABAN				
Tab 10 mg - No more than 1 tab per day	15.60	30	/	Xarelto
Tab 15 mg - Up to 14 tab available on a PSO		28		Xarelto
Tab 20 mg		28		Xarelto
WARFARIN SODIUM				
Note: Marevan and Coumadin are not interchangeable.				
* Tab 1 mg	3.46	50	/	Coumadin
•	7.50	100	✓	Marevan
* Tab 2 mg		50	1	Coumadin
* Tab 3 mg		100	1	Marevan
* Tab 5 mg	5.93	50	✓	Coumadin
-	13.50	100	✓	Marevan

FILGRASTIM - Special Authority see SA1259 below - Retail pha	ırmacy		
Inj 300 mcg per 0.5 ml prefilled syringe	86.60	10	✓ <u>Nivestim</u>
Inj 480 mcg per 0.5 ml prefilled syringe	133.72	10	✓ <u>Nivestim</u>

# ⇒SA1259 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 20%\*); or
- 2 Peripheral blood stem cell mobilisation in patients undergoing haematological transplantation; or
- 3 Peripheral blood stem cell mobilisation or bone marrow donation from healthy donors for transplantation; or
- 4 Treatment of severe chronic neutropenia (ANC < 0.5 ×10<sup>9</sup>/L); or
- 5 Treatment of drug-induced prolonged neutropenia (ANC < 0.5 ×10<sup>9</sup>/L).

Note: \*Febrile neutropenia risk greater than or equal to 20% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer	
PEGFILGRASTIM – Special Authority see SA1912 below – Retain j 6 mg per 0.6 ml syringe	'	1	_	i <u>extenzo</u> iextenzo AU	

#### ⇒SA1912 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where used for prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 5%\*). Note: \*Febrile neutropenia risk greater than or equal to 5% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

# Fluids and Electrolytes

#### Intravenous Administration

GLUCOSE [DEXTROSE]  * Inj 50%, 10 ml ampoule – Up to 5 inj available on a PSO  * Inj 50%, 90 ml bottle – Up to 5 inj available on a PSO		5 1	✓ <u>Biomed</u> ✓ Biomed
POTASSIUM CHLORIDE			
* Inj 75 mg per ml, 10 ml	65.00	50	✓ Juno ✓ LumaCina ✓ Pfizer \$29
SODIUM BICARBONATE Inj 8.4%, 50 ml a) Up to 5 inj available on a PSO b) Not in combination	24.70	1	✓ Biomed
Inj 8.4%, 100 ml	25.31	1	✓ Biomed

- a) Up to 5 inj available on a PSO
- b) Not in combination

#### SODIUM CHLORIDE

Not funded for use as a nasal drop. Not funded for nebuliser use except when used in conjunction with an antibiotic intended for nebuliser use.

Inj 0.9%, bag - Up to 2000 ml available on a PSO	500 ml	✓ Baxter
1.58	1 000 ml	✓ Raytor

Only if prescribed on a prescription for renal dialysis, maternity or post-natal care in the home of the patient, or on a PSO for emergency use. (500 ml and 1.000 ml packs)

for emergency use. (500 mi and 1,000 mi packs)			
Inj 23.4% (4 mmol/ml), 20 ml ampoule	40.15	5	✓ Biomed
For Sodium chloride oral liquid formulation refer Standa	ard Formulae, page	274	
Inj 0.9%, 5 ml ampoule - Up to 5 inj available on a PSO	4.00	20	✓ Fresenius Kabi
Inj 0.9%, 10 ml ampoule - Up to 5 inj available on a PSO	5.25	50	✓ Fresenius Kabi
Inj 0.9%, 20 ml ampoule	5.00	20	✓ Fresenius Kabi
TAL PARENTERAL MILITRITION (TDM)			

# TOTAL PARENTERAL NUTRITION (TPN) Infusion .......CBS

### WATER

 On a prescription or Practitioner's Supply Order only when on the same form as an injection listed in the Pharmaceutical Schedule requiring a solvent or diluent; or

1 OP

✓ TPN

- 2) On a bulk supply order; or
- 3) When used in the extemporaneous compounding of eye drops; or
- 4) When used for the dilution of sodium chloride soln 7% for cystic fibrosis patients only.

Inj 10 ml ampoule - Up to 5 inj available on a PSO	7.60	50	✓ Multichem
Inj 20 ml ampoule - Up to 5 inj available on a PSO	5.00	20	✓ Fresenius Kabi

	Subsidy (Manufacturer's	,	Fully Brand or idised Generic  Manufacturer
Oral Administration	\$	Per	Manuacturer
CALCIUM POLYSTYRENE SULPHONATE Powder	169.85	300 g OP	✓ Calcium Resonium
COMPOUND ELECTROLYTES  Powder for oral soln — Up to 5 sach available on a PSO		50	✓ <u>Electral</u>
COMPOUND ELECTROLYTES WITH GLUCOSE [DEXTROSE Soln with electrolytes	•	1,000 ml OP	✓ <u>Hydralyte -</u> Lemonade
PHOSPHORUS Tab eff 500 mg (16 mmol)	82.50	100	✓ Phosphate Phebra
POTASSIUM CHLORIDE  * Tab eff 548 mg (14 m eq) with chloride 285 mg (8 m eq)	5.26	60	Chlorvescent
* Tab long-acting 600 mg (8 mmol)	, ,	200	✓ Span-K
Cap 840 mg	8.52	100	<ul><li>✓ Sodibic</li><li>✓ Sodibic</li></ul>
SODIUM POLYSTYRENE SULPHONATE	04.05	454 00	
Powder	84.65	454 a OP	✓ Resonium-A

Subsidy		Fully	Brand or
(Manufacturer's Price)	Su	bsidised	Generic
` <b>\$</b>	Por	1	Manufacturer

# **Alpha-Adrenoceptor Blockers**

# **Alpha Adrenoceptor Blockers**

DOXAZO			
* Tab 2	<sup>17.35</sup>	500	<ul><li>Doxazosin Clinect</li></ul>
* Tab 4	mg20.94	500	<ul><li>Doxazosin Clinect</li></ul>
PHENOX	YBENZAMINE HYDROCHLORIDE		
* Cap 1	0 mg65.00	30	✓ BNM \$29
	216.67	100	✓ Dibenzyline S29
PRAZOSI	N		
* Tab 1	mg5.53	100	✓ Arrotex-Prazosin
			<b>S29</b> S29
	9.98		✓ Minipress S29
* Tab 2	<sup>1</sup> mg7.00	100	✓ Arrotex-Prazosin
			<b>S29</b> S29
	13.29		✓ Minipress S29
* Tab 5	mg11.70	100	✓ Arrotex-Prazosin
			<b>S29</b> S29
	22.00		✓ Minipress S29
* Cap 1	mg15.40	100	✓ Prazosin Mylan S29
* Cap 2	2 mg15.58	100	✓ Prazosin Mylan S29
	5 mg23.32	100	✓ Prazosin Mylan S29

# Agents Affecting the Renin-Angiotensin System

#### **ACE Inhibitors**

#### **CAPTOPRIL**

*	Oral liq 5 mg per ml86.00	100 ml OP	DP-Captopril
	Oral liquid restricted to children under 12 years of age		

#### CILAZAPRIL - Subsidy by endorsement

Subsidy by endorsement - Subsidised for patients who were taking cilazapril prior to 1 May 2021 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of cilazapril.

alopolioning of onal Laprin			
* Tab 0.5 mg	2.69	90	✓ Zapril
* Tab 2.5 mg	5.79	90	✓ Zapril
Tab 5 mg		90	✓ Zapril
ENALAPRIL MALEATE			
* Tab 5 mg	1.75	90	✓ Acetec
* Tab 10 mg	1.97	90	✓ Acetec
* Tab 20 mg		90	✓ Acetec
LISINOPRIL			
* Tab 5 mg	11.07	90	<ul> <li>Ethics Lisinopril</li> </ul>
· ·			✓ Teva Lisinopril
* Tab 10 mg	11.67	90	✓ Ethics Lisinopril
Ç			✓ Teva Lisinopril
* Tab 20 mg	14.69	90	✓ Ethics Lisinopril
•			✓ Teva Lisinopril

	Subsidy		Fully	
	(Manufacturer's Price)	Per	Subsidised	Generic Manufacturer
PERINDOPRIL	Ψ	1 01		
* Tab 2 mg	1 70	30	1	Coversyl
★ Tab 4 mg		30		Coversyl
★ Tab 8 mg		30		Coversyl
•		00	•	<u>oorcioyi</u>
QUINAPRIL	10.04	00	./	Arran Oninandi F
* Tab 5 mg		90	V	Arrow-Quinapril 5
Arrow-Quinapril 5 to be Principal Supply on 1 March 202  * Tab 10 mg		90		Arrow-Quinapril 10
Arrow-Quinapril 10 to be Principal Supply on 1 March 20		30	•	Allow-Quillapili io
* Tab 20 mg		90	1	Arrow-Quinapril 20
Arrow-Quinapril 20 to be Principal Supply on 1 March 20		50	•	Allow-Quillapili 20
	<i>J</i> 23			
RAMIPRIL	17.05	00	.,	Truzon
* Cap 1.25 mg	17.25	90	•	Tryzan
Tryzan to be Principal Supply on 1 February 2025	10.50	00		T
* Cap 2.5 mg	16.50	90	•	Tryzan
Tryzan to be Principal Supply on 1 February 2025	16 00	90	./	Truzon
* Cap 5 mg	10.00	90	•	Tryzan
Tryzan to be Principal Supply on 1 February 2025  Cap 10 mg	17.62	90		Tryzan
Tryzan to be Principal Supply on 1 February 2025	17.00	90	•	11 y Zaii
Tryzan to be i imolpai oupply on i i obraally 2020				
Angiotensin II Antagonists				
CANDESARTAN CILEXETIL				
<b>米</b> Tab 4 mg	2.68	90	✓	Candestar
Candestar to be Principal Supply on 1 February 2025				
* Tab 8 mg	2.67	90	•	Candestar
Candestar to be Principal Supply on 1 February 2025			_	
* Tab 16 mg	4.22	90	/	Candestar
Candestar to be Principal Supply on 1 February 2025			_	
* Tab 32 mg	5.24	90	•	Candestar
Candestar to be Principal Supply on 1 February 2025				
LOSARTAN POTASSIUM				
* Tab 12.5 mg	2.00	84	✓	Losartan Actavis
<b>★</b> Tab 25 mg	2.29	84	✓	Losartan Actavis
<b>米</b> Tab 50 mg	2.86	84	✓	Losartan Actavis
* Tab 100 mg	4.57	84	1	Losartan Actavis
Angiotensin II Antagonists with Diuretics				
CANDESARTAN CILEXETIL WITH HYDROCHLOROTHIAZIDE				
		30	,	APO-Candesartan
* Tab 16 mg with hydrochlorothiazide 12.5 mg	4. IU	30	•	HCTZ 16/12.5
* Tob 30 mg with hydrophlorothic-ide 10 5 mg	E OF	20	.,	
* Tab 32 mg with hydrochlorothiazide 12.5 mg	5.25	30	•	APO-Candesartan
				HCTZ 32/12.5
OSARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE				
* Tab 50 mg with hydrochlorothiazide 12.5 mg	4.00	30	1	Arrow-Losartan &
				<u>Hydrochlorothiazi</u>

Subsidy		Fully	Brand or
(Manufacturer's Price)	S	ubsidised	Generic
\$	Per	✓	Manufacturer

# **Angiotensin II Antagonists with Neprilysin Inhibitors**

SACUBITRIL WITH VALSARTAN - Special Authority see SA	A2302 below - Retail	pharmacy	
Tab 24.3 mg with valsartan 25.7 mg	190.00	56	✓ Entresto 24/26
Tab 48.6 mg with valsartan 51.4 mg	190.00	56	✓ Entresto 49/51
Tab 97.2 mg with valsartan 102.8 mg	190.00	56	✓ Entresto 97/103

# ⇒SA2302 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

## All of the following:

- 1 Patient has heart failure; and
- 2 Any of the following:
  - 2.1 Patient is in NYHA/WHO functional class II; or
  - 2.2 Patient is in NYHA/WHO functional class III; or
  - 2.3 Patient is in NYHA/WHO functional class IV; and
- 3 Either:
  - 3.1 Patient has a documented left ventricular ejection fraction (LVEF) of less than or equal to 35%; or

ablarida refer to NEDVOLIC CVCTEM. Appendibation I and page 100

- 3.2 An ECHO is not reasonably practical, and in the opinion of the treating practitioner the patient would benefit from treatment; and
- 4 Patient is receiving concomitant optimal standard chronic heart failure treatments.

# **Antiarrhythmics**

For lignocaine hydrochloride refer to NERVOUS SYSTEM, Anaesthetics, Local,	page 123	
AMIODARONE HYDROCHLORIDE		
▲ Tab 100 mg	30	✓ Aratac
▲ Tab 200 mg	30	✓ Aratac
Inj 50 mg per ml, 3 ml ampoule - Up to 10 inj available on a PSO 9.12	6	✓ Cordarone-X
15.22	10	✓ Max Health
ATROPINE SULPHATE		
* Inj 600 mcg per ml, 1 ml ampoule - Up to 5 inj available on a		
PSO16.10	10	✓ Hikma S29
		✓ Juno S29
		✓ Martindale
Martindale to be Principal Supply on 1 February 2025		
DIGOXIN		
* Tab 62.5 mcg – Up to 30 tab available on a PSO	240	✓ Lanoxin PG
* Tab 250 mcg - Up to 30 tab available on a PSO	240	✓ Lanoxin
* Oral lig 50 mcg per ml	60 ml	✓ Lanoxin
		✓ Lanoxin Paediatric
		Elixir
		✓ Lanoxin S29 S29
DISOPYRAMIDE PHOSPHATE		
▲ Cap 100 mg23.87	100	✓ Rythmodan
55.90	84	✓ Rythmodan -
		Cheplafarm S29

(N	Subsidy lanufacturer's Price)	Suh	Fully	Brand or Generic
(IV	\$	Per	√	Manufacturer
LECAINIDE ACETATE				
▲ Tab 50 mg	19.95	60	✓	Flecainide BNM
Cap long-acting 100 mg	35.78	90	✓ ]	Flecainide
				Controlled
				Release Teva
Cap long-acting 200 mg	54.28	90	<b>√</b> <u>[</u>	<u>Flecainide</u>
				Controlled
				Release Teva
Inj 10 mg per ml, 15 ml ampoule		5		Almarytm S29
	108.16			Tambocor
			•	Tambocor
				German S29
MEXILETINE HYDROCHLORIDE				
▲ Cap 150 mg	162.00	100	✓.	Teva S29
▲ Cap 250 mg	202.00	100	✓.	Teva S29
ROPAFENONE HYDROCHLORIDE				
▲ Tab 150 mg	40.90	50	<b>✓</b>	Rytmonorm
·				•
Antihypotensives				
MIDODRINE - Special Authority see SA1474 below - Retail pharm	acv			
Tab 2.5 mg	,	100	<b>√</b> 1	MAR-Midodrine S29
1 db 2.5 mg	00.00	100	-	Midodrine
				Medsurge
Midodrine Medsurge to be Principal Supply on 1 February 2	2025			3-
Tab 5 mg		100	<b>✓</b> I	MAR-Midodrine S29
	50.00	.00	-	Midodrine
			-	Medsurge

Midodrine Medsurge to be Principal Supply on 1 February 2025

# **⇒SA1474** Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years where patient has disabling orthostatic hypotension not due to drugs.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

# **Beta-Adrenoceptor Blockers**

# **Beta Adrenoceptor Blockers**

ATENOLOL		
* Tab 50 mg11.00 Viatris to be Principal Supply on 1 February 2025	500	✓ Viatris
* Tab 100 mg	500	✓ Atenolol Viatris
* Oral liq 25 mg per 5 ml49.85  Restricted to children under 12 years of age.	300 ml OP	✓ Atenolol AFT
BISOPROLOL FUMARATE		
* Tab 2.5 mg	90	✓ Ipca-Bisoprolol
* Tab 5 mg1.91	90	✓ Ipca-Bisoprolol
* Tab 10 mg2.71	90	✓ Ipca-Bisoprolol

	Subsidy		Fully	y Brand or
	(Manufacturer's Price)		Subsidised	
	\$	Per	•	Manufacturer
CARVEDILOL				
* Tab 6.25 mg	2.24	60	1	Carvedilol Sandoz
* Tab 12.5 mg	2.30	60	•	Carvedilol Sandoz
* Tab 25 mg	2.95	60	•	Carvedilol Sandoz
LABETALOL				
* Tab 100 mg	14.50	100	/	Trandate
* Tab 200 mg		100	1	Trandate
* Inj 5 mg per ml, 20 ml ampoule		5		
, , , , , , , , , , , , , , , , , , , ,	(88.60)			Trandate
METOPROLOL SUCCINATE	, ,			
* Tab long-acting 23.75 mg	4 20	90	/	Myloc CR
* Tab long-acting 47.5 mg		90		Myloc CR
* Tab long-acting 95 mg		90		Myloc CR
* Tab long-acting 190 mg		90		Myloc CR
METOPROLOL TARTRATE				<u>,</u>
* Tab 50 mg	5.66	100		IPCA-Metoprolol
* Tab 30 mg		60		IPCA-Metoprolol
* Tab long-acting 200 mg		28		Slow-Lopresor
* Inj 1 mg per ml, 5 ml vial		5		Metoprolol IV Mylan
* III I IIIg pei IIII, 5 IIII viai	20.50	5		Metoprolol IV Viatris
NAPOLOI			•	wetoprolor iv viatris
NADOLOL National Communication of the Communication	10.10	400	,	Al-J-I-I DAM
* Tab 40 mg		100		Nadolol BNM
* Tab 80 mg	30.39	100	•	Nadolol BNM
PROPRANOLOL				
* Tab 10 mg	7.04	100		<u>Drofate</u>
* Tab 40 mg	8.75	100		IPCA-Propranolol
* Cap long-acting 160 mg	18.17	100	•	Cardinol LA
* Oral liq 4 mg per ml - Special Authority see SA1327 below	_			
Retail pharmacy	CBS	500 m	· •	Roxane-
				Propranolol \$29
				•

# **⇒SA1327** Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

**Renewal** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

S	U	ΙA	LC	)L

*	Tab 80 mg37.50	500	Mylan
	Tab 160 mg14.00		✓ Mylan

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per Manufacturer

# **Calcium Channel Blockers**

D	ihydropyridine Calcium Channel Blockers			
ΑM	LODIPINE			
	Tab 2.5 mg	1.45	90	✓ Vasorex
*	Tab 5 mg	1.21	90	✓ Vasorex
*	Tab 10 mg	1.31	90	✓ Vasorex
FE	LODIPINE			
*	Tab long-acting 2.5 mg	2.18	30	✓ Plendil ER
	Plendil ER to be Principal Supply on 1 February 2025			
*	Tab long-acting 5 mg	6.57	90	✓ Felo 5 ER
	Felo 5 ER to be Principal Supply on 1 February 2025			
*	Tab long-acting 10 mg	6.95	90	✓ Felo 10 ER
	Felo 10 ER to be Principal Supply on 1 February 2025			
	EDIPINE			
*	Tab long-acting 10 mg — Subsidy by endorsement	19.42	56	✓ Tensipine MR10 S29
	Subsidised for patients who were taking nifedipine tab lor endorsed accordingly. Pharmacists may annotate the prodispensing of nifedipine tab long-acting 10 mg.	escription as end	orsed where	there exists a record of prior
*	Tab long-acting 20 mg		100	✓ Nyefax Retard
*	Tab long-acting 30 mg	4.78	14	Mylan Italy (24 hr
				release) S29
		34.10	100	✓ Mylan (24 hr
				release) S29
*	Tab long-acting 60 mg	52.81	100	✓ Mylan (24 hr
				release) S29
0	ther Calcium Channel Blockers			
DIL	TIAZEM HYDROCHLORIDE			
	Cap long-acting 120 mg	65.35	500	✓ <u>Diltiazem CD Clinect</u>
*	Cap long-acting 180 mg	7.00	30	✓ Cardizem CD
*	Cap long-acting 240 mg	9.30	30	✓ Cardizem CD
PΕ	RHEXILINE MALEATE			
*	Tab 100 mg	62.90	100	✓ Pexsig
۷E	RAPAMIL HYDROCHLORIDE			
*	Tab 40 mg	7.01	100	✓ Isoptin
*	Tab 80 mg	11.74	100	✓ Isoptin
*	Tab long-acting 120 mg	36.02	100	✓ Isoptin Retard S29
				✓ Isoptin SR
*	Tab long-acting 240 mg	15.12	30	✓ Isoptin SR
*	Inj 2.5 mg per ml, 2 ml ampoule - Up to 5 inj available on a			

✓ Isoptin

			_
	Subsidy		Fully Brand or
	(Manufacturer's F		sidised Generic
	\$	Per	✓ Manufacturer
Centrally-Acting Agents			
CLONIDINE			
CLONIDINE  ** Potch 2.5 mg, 100 mag, new day. Only on a precediation.	11 70	4	./ Mulan
* Patch 2.5 mg, 100 mcg per day — Only on a prescription		4	✓ Mylan
* Patch 5 mg, 200 mcg per day – Only on a prescription		4 4	✓ <u>Mylan</u>
* Patch 7.5 mg, 300 mcg per day – Only on a prescription	17.90	4	✓ <u>Mylan</u>
CLONIDINE HYDROCHLORIDE			<b>.</b>
* Tab 25 mcg		112	✓ Clonidine Teva
* Tab 150 mcg	40.41	100	✓ Catapres
Catapres to be Principal Supply on 1 February 2025	4440	-	
* Inj 150 mcg per ml, 1 ml ampoule	14.10	5	✓ Catapres
METHYLDOPA			
* Tab 250 mg	15.10	100	Methyldopa Viatris
Discouling			
Diuretics			
Loop Diuretics			
BUMETANIDE			
* Tab 1 mg	16.36	100	✓ Burinex
* Inj 500 mcg per ml, 4 ml vial		5	✓ Burinex
FUROSEMIDE [FRUSEMIDE]			
Tab 40 mg – Up to 30 tab available on a PSO	12.80	1,000	✓ IPCA-Frusemide
IPCA-Frusemide to be Principal Supply on 1 February		.,000	
* Tab 500 mg		50	✓ Urex Forte
* Oral lig 10 mg per ml		30 ml OP	✓ Lasix
* Inj 10 mg per ml, 25 ml ampoule		6	✓ Lasix
* Inj 10 mg per ml, 2 ml ampoule - Up to 5 inj available on a		5	✓ Furosemide-Baxter
Potassium Sparing Diuretics			
Potassium Sparing Didiences			
AMILORIDE HYDROCHLORIDE			
Tab 5 mg	81.07	100	✓ Padagis ©29
•	171.41	28	✓ Wockhardt S29
Oral liq 1 mg per ml	35.40	25 ml OP	✓ Biomed
EPLERENONE – Special Authority see SA1728 below – Retail			
Tab 25 mg	15.84	30	✓ Inspra
Tab 50 mg		30	✓ Inspra
⇒SA1728 Special Authority for Subsidy			<u></u>
Initial application from any relevant practitioner. Approvals va	lid without further	ranawal unlass	s notified for applications meeting
the following criteria:	and without further	Teriewai uriiese	s notified for applications meeting
Both:			
1 Patient has heart failure with ejection fraction less than 4	10%: and		
2 Either:	10 /0, and		
<ul><li>2.1 Patient is intolerant to optimal dosing of spironola</li><li>2.2 Patient has experienced a clinically significant ad</li></ul>		on optimal dos	sing of spironolactone.
SPIRONOLACTONE		•	
* Tab 25 mg	3 68	100	✓ Spiractin
* Tab 100 mg		100	✓ Spiractin
Oral liq 5 mg per ml		25 ml OP	✓ Biomed
1 - 31 -			

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price) \$	F Subsidi Per	ully ised •	Brand or Generic Manufacturer
Potassium Sparing Combination Diuretics				
AMILORIDE HYDROCHLORIDE WITH FUROSEMIDE  * Tab 5 mg with furosemide 40 mg		28	/	Frumil
* Tab 5 mg with hydrochlorothiazide 50 mg	-	50	1	Moduretic
Thiazide and Related Diuretics				
BENDROFLUMETHIAZIDE [BENDROFLUAZIDE]  * Tab 2.5 mg – Up to 150 tab available on a PSO	51.50	500	/	Arrow- Bendrofluazide
May be supplied on a PSO for reasons other than emerge * Tab 5 mg	,	500	•	Arrow- Bendrofluazide
CHLOROTHIAZIDE Oral liq 50 mg per ml	30.67 25	5 ml OP	•	Biomed
* Tab 25 mg	6.95	50	•	Hygroton
INDAPAMIDE  * Tab 2.5 mg  METOLAZONE	16.00	90	•	Dapa-Tabs
Tab 5 mg	CBS	1 50		Metolazone S29 Zaroxolyn S29
Vasopressin receptor antagonists				
TOLVAPTAN – Special Authority see SA2166 below – Retail pha Tab 15 mg Tab 30 mg Tab 45 mg + 15 mg Tab 60 mg + 30 mg	873.50	28 OP 28 OP 56 OP 56 OP	<b>/</b>	Jinarc Jinarc Jinarc Jinarc

# ⇒SA2166 Special Authority for Subsidy

Initial application — (autosomal dominant polycystic kidney disease) only from a renal physician or any relevant practitioner on the recommendation of a renal physician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 Patient has a confirmed diagnosis of autosomal dominant polycystic kidney disease; and

- 2 Patient has an estimated glomerular filtration rate (eGFR) of greater than or equal to 25 ml/min/1.73 m<sup>2</sup> at treatment initiation; and
- 3 Either:
  - 3.1 Patient's disease is rapidly progressing, with a decline in eGFR of greater than or equal to 5 mL/min/1.73 m<sup>2</sup> within one-year; or
  - 3.2 Patient's disease is rapidly progressing, with an average decline in eGFR of greater than or equal to 2.5 mL/min/1.73 m² per year over a five-year period.

Renewal — (autosomal dominant polycystic kidney disease) only from a renal physician or any relevant practitioner on the

continued...

56 OP

✓ Jinarc

continued...

recommendation of a renal physician. Approvals valid for 12 months for applications meeting the following criteria:

- 1 Patient has not developed end-stage renal disease, defined as an eGFR of less than 15 mL/min/1.73 m²; and

2 Patient has not undergone a kidney transplant.		
Lipid-Modifying Agents		
Fibrates		
# Tab 200 mg	90 30	<ul><li>✓ Bezalip</li><li>✓ Bezalip Retard</li></ul>
Bezalip Retard to be Principal Supply on 1 March 2025		
Other Lipid-Modifying Agents		
ACIPIMOX * Cap 250 mg38.19	30	✓ Olbetam
Resins		
COLESTYRAMINE Powder for oral suspension 4 g sachet	50	✓ Colestyramine - Mylan \$29 ✓ Quantalan sugar free \$29
HMG CoA Reductase Inhibitors (Statins)		
ATORVASTATIN  Tab 10 mg	500 500 500 500	✓ Lorstat ✓ Lorstat ✓ Lorstat ✓ Lorstat
THAVAOTATIN		

~	Tab to mg						
RO	SUVASTATIN	- Special	Authority	see SA	A2093 be	low –	Retai

⇒SA2093 Special Authority for Subsidy

*	Tab 40 mg	100
	SUVASTATIN - Special Authority see SA2093 below - Retail pharmacy	
	Tab 5 mg	30
	Tab 10 mg1.69	30
	Tab 20 mg2.71	30
*	Tab 40 mg	30

✓ Rosuvastatin Viatris

✓ Rosuvastatin Viatris

✓ Clinect

✓ Clinect

100

✓ Rosuvastatin Viatris

✓ Rosuvastatin Viatris

Initial application — (cardiovascular disease risk) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

Subsidy (Manufacturer's Price)	Subsi Per	Fully dised	Brand or Generic Manufacturer
Ψ	1 01		- Iviarialactaroi

continued...

- 1 Both:
  - 1.1 Patient is considered to be at risk of cardiovascular disease; and
  - 1.2 Patient is Māori or any Pacific ethnicity: or
  - 2 Both:
    - 2.1 Patient has a calculated risk of cardiovascular disease of at least 15% over 5 years; and
    - 2.2 LDL cholesterol has not reduced to less than 1.8 mmol/litre with treatment with the maximum tolerated dose of atoryastatin and/or simyastatin.

Initial application — (familial hypercholesterolemia) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has familial hypercholesterolemia (defined as a Dutch Lipid Criteria score greater than or equal to 6); and
- 2 LDL cholesterol has not reduced to less than 1.8 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initial application — (established cardiovascular disease) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

1 Any of the following:

SIMVASTATIN

- 1.1 Patient has proven coronary artery disease (CAD); or
- 1.2 Patient has proven peripheral artery disease (PAD); or
- 1.3 Patient has experienced an ischaemic stroke; and
- 2 LDL cholesterol has not reduced to less than 1.4 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initial application — (recurrent major cardiovascular events) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has experienced a recurrent major cardiovascular event (defined as myocardial infarction, ischaemic stroke, coronary revascularisation, hospitalisation for unstable angina) in the last 2 years; and
- 2 LDL cholesterol has not reduced to less than 1.0 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

* Tab 10 mg1.68	90	<ul> <li>✓ Simvastatin Mylan</li> <li>✓ Simvastatin Viatris</li> </ul>
* Tab 20 mg2.54	90	✓ Simvastatin Viatris
* Tab 40 mg4.11	90	✓ Simvastatin Viatris
* Tab 80 mg8.81	90	✓ Simvastatin Viatris
Selective Cholesterol Absorption Inhibitors		
EZETIMIBE		
Tab 10 mg1.76	30	<ul><li>✓ Ezemibe Viatris</li><li>✓ Ezetimibe Sandoz</li></ul>
(Ezemibe Viatris Tab 10 mg to be delisted 1 July 2025)		
EZETIMIBE WITH SIMVASTATIN		
Tab 10 mg with simvastatin 10 mg5.15	30	✓ Zimybe
Tab 10 mg with simvastatin 20 mg6.15	30	✓ Zimybe
Tab 10 mg with simvastatin 40 mg7.15	30	✓ Zimybe
Tab 10 mg with simvastatin 80 mg8.15	30	✓ Zimybe

	Subsidy (Manufacturer's \$	Price) Subsi Per	Fully Brand or dised Generic ✓ Manufacturer
Nitrates			
GLYCERYL TRINITRATE			
* Oral pump spray, 400 mcg per dose – Up to 250 dose available on a PSO	7.48	250 dose OP	✓ Nitrolingual Pump Spray
* Patch 25 mg, 5 mg per day	15.73	30	✓ Nitroderm TTS
* Patch 50 mg, 10 mg per day	18.62	30	✓ Nitroderm TTS
ISOSORBIDE MONONITRATE			
* Tab 20 mg		100	✓ <u>Ismo 20</u>
* Tab long-acting 40 mg		30	✓ Ismo 40 Retard
* Tab long-acting 60 mg	13.50	90	✓ <u>Duride</u>
Sympathomimetics			
ADRENALINE Inj 1 in 1,000, 1 ml ampoule – Up to 5 inj available on a PSO	1 08	5	✓ Aspen Adrenaline
ing 1 in 1,000, 1 in ampoule – op to 3 ing available on a 1 30	13.27	3	✓ DBL Adrenaline
	25.30	10	✓ Hameln S29
Inj 1 in 10,000, 10 ml ampoule - Up to 5 inj available on a PS	SO27.00	5	✓ Hospira
	49.00	10	✓ Aspen Adrenaline
Vasodilators			
HYDRALAZINE HYDROCHLORIDE			
* Tab 25 mg - Special Authority see SA1321 below - Retail			
pharmacy	CBS	1	✓ Hydralazine
		56	✓ Onelink S29
		84	✓ AMDIPHARM \$29
		100	✓ Camber S29
* Inj 20 mg ampoule	25.90	5	✓ Apresoline
■ SA1321 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid the following criteria:  Either:	d without furthe	r renewal unless	notified for applications meeting
<ul><li>1 For the treatment of refractory hypertension; or</li><li>2 For the treatment of heart failure in combination with a nitr inhibitors and/or angiotensin receptor blockers.</li></ul>	ate, in patients	who are intolera	nt or have not responded to ACE

60

100

60

60

5

50

78.40

✓ Minoxidil Roma S29

✓ Loniten

✓ Max Health

✓ Max Health

✓ Trental 400

✓ Hospira

# ▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

▲ Tab 10 mg .......21.73

Tab 400 mg ......44.37

MINOXIDIL

NICORANDII

PAPAVERINE HYDROCHLORIDE

PENTOXIFYLLINE [OXPENTIFYLLINE]

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

# **Endothelin Receptor Antagonists**

AMBRISENTAN - Special Authority see SA2253 below - Retail	pharmacy		
Tab 5 mg	200.00	30	✓ Ambrisentan Viatris
Tab 10 mg	200.00	30	✓ Ambrisentan Viatris

⇒SA2253 Special Authority for Subsidy

Initial application — (PAH monotherapy) only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH has been confirmed by right heart catheterisation; and
    - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
    - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
    - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm<sup>-5</sup>); and
    - 4.1.5 Any of the following:
      - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH 2022 (see note below for link to these guidelines) †: or
      - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool\*\*: or
      - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
  - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
  - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Roth:
  - 5.1 Ambrisentan is to be used as PAH monotherapy; and
  - 5.2 Any of the following:
    - 5.2.1 Patient has experienced intolerable side effects with both sildenafil and bosentan; or
    - 5.2.2 Patient has an absolute contraindication to sildenafil and an absolute or relative contraindication to bosentan (e.g. due to current use of a combined oral contraceptive or liver disease); or
    - 5.2.3 Patient is a child with idiopathic PAH or PAH secondary to congenital heart disease.

Initial application — (PAH dual therapy) only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH has been confirmed by right heart catheterisation; and

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

- 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
- 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
- 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm<sup>-5</sup>); and
- 4.1.5 Any of the following:
  - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH 2022 (see note below for link to these guidelines) †; or
  - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool\*\*; or
  - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
- 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 All of the following:
  - 5.1 Ambrisentan is to be used as PAH dual therapy; and
  - 5.2 Either:
    - 5.2.1 Patient has tried a PAH monotherapy (sildenafil or bosentan) for at least three months and has not experienced an acceptable response to treatment according to a validated risk stratification tool\*\*; or
    - 5.2.2 Patient has tried PAH dual therapy including bosentan and has experienced intolerable side effects on bosentan; and
  - 5.3 Both:
    - 5.3.1 Patient is presenting in NYHA/WHO functional class III or IV, and in the opinion of the treating clinician would benefit from initial dual therapy; and
    - 5.3.2 Patient has an absolute or relative contraindication to bosentan (e.g. due to current use of a combined oral contraceptive or liver disease).

**Initial application** — **(PAH triple therapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH has been confirmed by right heart catheterisation; and
    - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
    - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
    - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm<sup>-5</sup>); and
    - 4.1.5 Any of the following:
      - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH 2022 (see note below for link to these guidelines) †; or
      - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool\*\*; or

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsid	dised	Generic	
\$	Per	/	Manufacturer	

continued...

- 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
- 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
  - 5.1 Ambrisentan is to be used as PAH triple therapy; and
  - 5.2 Any of the following:
    - 5.2.1 Patient is on the lung transplant list; or
    - 5.2.2 Both:
      - 5.2.2.1 Patient is presenting in NYHA/WHO functional class IV: and
      - 5.2.2.2 Patient has an absolute or relative contraindication to bosentan (e.g. due to current use of a combined oral contraceptive or liver disease); or
    - 5.2.3 Both:
      - 5.2.3.1 Patient has tried PAH dual therapy for at least three months and remains in an unacceptable risk category according to a validated risk stratification tool\*\*; and
      - 5.2.3.2 Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario.

**Renewal** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 2 years where the patient is continuing to derive benefit from ambrisentan treatment according to a validated PAH risk stratification tool\*\*.

Notes: † The European Respiratory Journal Guidelines can be found here: 2022 ECS/ERS Guidelines for the diagnosis and treatment of pulmonary hypertension PAH

\*\* the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

Tab 62.5 mg	100.00	60	✓ Bosentan Dr
			Reddy's
Tab 125 mg	100.00	60	✓ Bosentan Dr
			Reddy's

#### **⇒SA2254** Special Authority for Subsidy

**Initial application** — **(PAH monotherapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)\*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH has been confirmed by right heart catheterisation; and
    - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
    - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
    - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm<sup>5</sup>); and

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer
----------------------------------------------------------------------------------------------

continued...

- 4.1.5 Any of the following:
  - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these quidelines) †; or
  - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool\*\*; or
  - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
- 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or
- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
  - 5.1 Bosentan is to be used as PAH monotherapy; and
  - 5.2 Any of the following:
    - 5.2.1 Patient has experienced intolerable side effects on sildenafil: or
    - 5.2.2 Patient has an absolute contraindication to sildenafil: or
    - 5.2.3 Patient is a child with idiopathic PAH or PAH secondary to congenital heart disease.

**Initial application** — **(PAH dual therapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

#### All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)\*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II. III or IV: and
- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH has been confirmed by right heart catheterisation; and
    - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
    - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
    - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm<sup>-5</sup>); and
    - 4.1.5 Any of the following:
      - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these quidelines) †; or
      - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool\*\*: or
      - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
  - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or
  - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Bosentan is to be used as part of PAH dual therapy; and
- 6 Either:
  - 6.1 Patient has tried a PAH monotherapy (sildenafil) for at least three months and has experienced an inadequate therapeutic response to treatment according to a validated risk stratification tool\*\*; or
  - 6.2 Patient is presenting in NYHA/WHO functional class III or IV, and in the opinion of the treating clinician would likely

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

benefit from initial dual therapy.

**Initial application** — **(PAH triple therapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)\*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH has been confirmed by right heart catheterisation; and
    - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
    - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
    - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm<sup>5</sup>); and
    - 4.1.5 Any of the following:
      - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these quidelines) †: or
      - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool\*\*: or
      - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
  - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or
  - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
  - 5.1 Bosentan is to be used as part of PAH triple therapy; and
  - 5.2 Any of the following:
    - 5.2.1 Patient is on the lung transplant list; or
    - 5.2.2 Patient is presenting in NYHA/WHO functional class IV; or
    - 5.2.3 Both:
      - 5.2.3.1 Patient has tried PAH dual therapy for at least three months and has not experienced an acceptable response to treatment according to a validated risk stratification tool\*\*; and
      - 5.2.3.2 Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario.

**Renewal** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 2 years where patient is continuing to derive benefit from bosentan treatment according to a validated PAH risk stratification tool\*\*.

Notes: † The European Respiratory Journal Guidelines can be found here: 2022 ECS/ERS Guidelines for the diagnosis and treatment of pulmonary hypertension PAH

\*\* the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

# **Phosphodiesterase Type 5 Inhibitors**

SILDENAFIL - Special Authority see SA2255 on the next page - Retail	pharmacy		
Tab 25 mg	0.72	4	✓ Vedafil
Tab 50 mg	1.45	4	✓ Vedafil
Tab 100 mg	11.22	12	✓ Vedafil

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsi	dised	Generic	
\$	Per	✓	Manufacturer	

# ⇒SA2255 Special Authority for Subsidy

**Initial application** — (Raynaud's Phenomenon\*) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has Raynaud's Phenomenon\*; and
- 2 Patient has severe digital ischaemia (defined as severe pain requiring hospital admission or with a high likelihood of digital ulceration; digital ulcers; or gangrene); and
- 3 Patient is following lifestyle management (avoidance of cold exposure, sufficient protection, smoking cessation support, avoidance of sympathomimetic drugs); and
- 4 Patient is being treated with calcium channel blockers and nitrates (or these are contraindicated/not tolerated).

Initial application — (Pulmonary arterial hypertension\*) only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)\*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II. III or IV: and
- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH is confirmed by right heart catheterisation; and
    - 4.1.2 A mean pulmonary artery pressure (PAPm) of greater than 20 mmHg; and
    - 4.1.3 A pulmonary capillary wedge pressure (PCWP) that is less than or equal to 15 mmHg; and
    - 4.1.4 Pulmonary vascular resistance (PVR) of at least 2 Wood Units or greater than 160 International Units (dyn s cm<sup>-5</sup>); and
    - 4.1.5 Any of the following:
      - 4.1.5.1 PAH is non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) †; or
      - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool\*\*; or
      - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
  - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or
  - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures.

Initial application — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has a documented history of traumatic or non-traumatic spinal cord injury; and
- 2 Patient has erectile dysfunction secondary to spinal cord injury requiring pharmacological treatment.

Renewal — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Notes: Note: Indications marked with \* are Unapproved Indications.

† The European Respiratory Journal Guidelines can be found here: 2022 ECS/ERS Guidelines for the diagnosis and treatment of pulmonary hypertension PAH

\*\* the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

# **Prostacyclin Analogues**

EPOPROSTENOL - Special Authority see SA2256 below - Retail pharmacy 1 ✓ Veletri ✓ Veletri 

⇒SA2256 Special Authority for Subsidy

Initial application — (PAH dual therapy) only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class III or IV; and
- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH has been confirmed by right heart catheterisation; and
    - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
    - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
    - 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm<sup>-5</sup>); and
    - 4.1.5 Any of the following:
      - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these quidelines) † ; or
      - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool\*\*; or
      - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
  - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
  - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 All of the following:
  - 5.1 Epoprostenol is to be used as part of PAH dual therapy with either sildenafil or an endothelin receptor antagonist;
  - 5.2 Patient is presenting in NYHA/WHO functional class IV; and
  - 5.3 Patient has tried a PAH monotherapy for at least three months and remains in an unacceptable risk category according to a validated risk stratification tool.

Initial application — (PAH triple therapy) only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class III or IV; and
- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH has been confirmed by right heart catheterisation; and
    - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and

continued...

- 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
- 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm<sup>-5</sup>); and
- 4.1.5 Any of the following:
  - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these quidelines) †: or
  - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool\*\*; or
  - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
- 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
  - 5.1 Epoprostenol is to be used as PAH triple therapy; and
  - 5.2 Any of the following:
    - 5.2.1 Patient is on the lung transplant list; or
      - 5.2.2 Patient is presenting in NYHA/WHO functional class IV; or
      - 5.2.3 Both:
        - 5.2.3.1 Patient has tried PAH dual therapy for at least three months and has not experienced an acceptable response to treatment according to a validated risk stratification tool; and
        - 5.2.3.2 Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario.

**Renewal** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 2 years where patient is continuing to derive benefit from epoprostenol treatment according to a validated PAH risk stratification tool\*\*.

Notes: † The European Respiratory Journal Guidelines can be found here: 2022 ECS/ERS Guidelines for the diagnosis and treatment of pulmonary hypertension PAH

\*\* the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

ILOPROST - Special Authority see SA2257 below - Retail pharmacy

### ⇒SA2257 Special Authority for Subsidy

**Initial application** — **(PAH monotherapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH has been confirmed by right heart catheterisation; and
    - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
    - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	Subsidised	Generic	
\$	Per	/	Manufacturer	

continued...

- 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm<sup>-5</sup>); and
- 4.1.5 Any of the following:
  - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these quidelines) †: or
  - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool\*\*: or
  - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
- 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
  - 5.1 Iloprost is to be used as PAH monotherapy; and
  - 5.2 Either:
    - 5.2.1 Patient has experienced intolerable side effects on sildenafil and both the funded endothelin receptor antagonists (i.e. both bosentan and ambrisentan); or
    - 5.2.2 Patient has an absolute contraindication to sildenafil and an absolute or relative contraindication to endothelin receptor antagonists.

**Initial application** — **(PAH dual therapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH has been confirmed by right heart catheterisation; and
    - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
    - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
    - 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm<sup>-5</sup>); and
    - 4.1.5 Any of the following:
      - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) † : or
      - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool\*\*: or
      - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
  - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
  - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 All of the following:
  - 5.1 Iloprost is to be used as PAH dual therapy with either sildenafil or an endothelin receptor antagonist; and

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer	
----------------------------------------------------------------------------------------	--

continued...

- 5.2 Either:
  - 5.2.1 Patient has an absolute contraindication to or has experienced intolerable side effects on sildenafil; or
  - 5.2.2 Patient has an absolute or relative contraindication to or experienced intolerable side effects with a funded endothelin receptor antagonist; and
- 5.3 Either:
  - 5.3.1 Patient has tried a PAH monotherapy for at least three months and remains in an unacceptable risk category according to a validated risk stratification tool\*\*; or
  - 5.3.2 Patient is presenting in NYHA/WHO functional class III or IV, and in the opinion of the treating clinician would benefit from initial dual therapy.

**Initial application** — **(PAH triple therapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH has been confirmed by right heart catheterisation; and
    - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
    - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
    - 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm<sup>-5</sup>); and
    - 4.1.5 Any of the following:
      - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) †; or
      - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool\*\*; or
      - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
  - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
  - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
  - 5.1 Iloprost is to be used as PAH triple therapy; and
  - 5.2 Any of the following:
    - 5.2.1 Patient is on the lung transplant list: or
    - 5.2.2 Patient is presenting in NYHA/WHO functional class IV; or
    - 5.2.3 Both:
      - 5.2.3.1 Patient has tried PAH dual therapy for at least three months and has not experienced an acceptable response to treatment according to a validated risk stratification tool\*\*; and
      - 5.2.3.2 Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario.

**Renewal** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 2 years where patient is continuing to derive benefit from iloprost treatment according to a validated PAH risk stratification tool\*\*.

Subsidy	Fu	ly Brand or	
(Manufacturer's Price)	Subsidise	ed Generic	
\$	Per	Manufacturer	

continued...

Notes: † The European Respiratory Journal Guidelines can be found here: 2022 ECS/ERS Guidelines for the diagnosis and treatment of pulmonary hypertension PAH

<sup>\*\*</sup> the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

# **DERMATOLOGICALS**

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

# **Antiacne Preparations**

For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 95

#### **ADAPALENE**

- a) Maximum of 30 g per prescription
- b) Only on a prescription

22.89	30 g OP	<ul><li>Differin</li></ul>
harmacy		
11.26	60	<ul><li>Oratane</li></ul>
18.75	120	✓ Oratane
	120	✓ Oratane
	22.89 harmacy11.2618.7526.73	harmacy 6011.26 6018.75 120

# ⇒SA2023 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around isotretinoin and is competent to prescribe isotretinoin; and
- 3 Either:
  - 3.1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and that they must not become pregnant during treatment and for a period of one month after the completion of treatment; or
  - 3.2 Patient is not of child bearing potential.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and that they must not become pregnant during treatment and for a period of one month after the completion of treatment; or
- 2 Patient is not of child bearing potential.

#### **TRETINOIN**

Crm 0.5 mg per g - Maximum of 50 g per prescription	16.82	50 g OP	✓ ReTrieve
ReTrieve to be Principal Supply on 1 February 2025			

# **Antibacterials Topical**

For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 95

# HYDROGEN PEROXIDE

* Crm 1%	8.56	10 g OP	✓ Crystaderm
MUPIROCIN		· ·	
Oint 2%	6.60	15 g OP	
	(13.00)	-	Bactroban

- a) Only on a prescription
- b) Not in combination

# **DERMATOLOGICALS**

	Subsidy (Manufacturer's F \$	Price) Subs	Fully sidised	Brand or Generic Manufacturer
SODIUM FUSIDATE [FUSIDIC ACID]				
Crm 2%	1.69	5 g OP	<b>√</b> F	oban
a) Maximum of 5 g per prescription     b) Only on a prescription     c) Not in combination     d) Foban to be Principal Supply on 1 February 2025	4.00	5 OD	4.	•••••
Oint 2%	1.69	5 g OP	<b>V</b> F	Foban
SULFADIAZINE SILVER Crm 1%	10.80 15.44	50 g OP	-	Flamazine Ascend S29
<ul><li>a) Up to 250 g available on a PSO</li><li>b) Not in combination</li></ul>				
Antifungals Topical				
	100			
For systemic antifungals, refer to INFECTIONS, Antifungals, pa	ge 102			
AMOROLFINE  a) Only on a prescription b) Not in combination				
Nail soln 5%	21.87	5 ml OP	✓ N	/lycoNail
CLOTRIMAZOLE				
* Crm 1%	1.10	20 g OP	<b>√</b> <u>C</u>	Clomazol
<ul><li>a) Only on a prescription</li><li>b) Not in combination</li></ul>				
* Soln 1%	4.36	20 ml OP		
	(7.55)	20 01	C	Canesten
<ul><li>a) Only on a prescription</li><li>b) Not in combination</li></ul>				
ECONAZOLE NITRATE				
Crm 1%a) Only on a prescription	8.04	20 g OP	<b>√</b> F	Pevaryl
b) Not in combination Foaming soln 1%, 10 ml sachets	0.80	3		
1 January 2011 1 /0, 10 111 Sacricis	(18.64)	J	F	Pevaryl
	(10.04)			J J.

a) Only on a prescriptionb) Not in combination

	Subsidy (Manufacturer's P		Fully Brand or sidised Generic	
	\$	Per	✓ Manufacturer	
MICONAZOLE NITRATE				
* Crm 2%	0.90	15 g OP	✓ Multichem	
<ul> <li>a) Only on a prescription</li> </ul>		ŭ	<del></del>	
b) Not in combination				
* Lotn 2%	4.36	30 ml OP		
	(10.03)		Daktarin	
a) Only on a prescription	,			
b) Not in combination				
* Tinct 2%	4.36	30 ml OP		
	(12.10)		Daktarin	
a) Only on a prescription	( -/			
b) Not in combination				
b) Hot in combination				

CA	۱ ۸	NΛI	N	

a) Only on a prescription

b) Not in combination

100 g **✓ healthE Calamine**Aqueous

healthE Calamine Aqueous to be Principal Supply on 1 April 2025

#### CROTAMITON

a) Only on a prescription

b) Not in combination

Crm 10%......3.49

20 g OP ✓

✓ Itch-Soothe

#### MENTHOL - Only in combination

1) Only in combination with a dermatological base or proprietary Topical Corticosteriod - Plain

2) With or without other dermatological galenicals.

Itch-Soothe to be Principal Supply on 1 February 2025

# **Corticosteroids Topical**

For systemic corticosteroids, refer to CORTICOSTEROIDS AND RELATED AGENTS, page 85

### Corticosteroids - Plain

BE	TAMETHASONE DIPROPIONATE			
	Crm 0.05%	2.96	15 g OP	✓ Diprosone
	30	6.00	50 g OP	✓ Diprosone
	Oint 0.05%	2.96	15 g OP	✓ Diprosone
	30	6.00	50 g OP	✓ Diprosone
	Oint 0.05% in propylene glycol base	4.33	30 g OP	✓ Diprosone OV
BE	TAMETHASONE VALERATE			
*	Crm 0.1%	5.85	50 g OP	✓ Beta Cream
	Beta Cream to be Principal Supply on 1 February 2025		-	
*	Oint 0.1%	7.90	50 g OP	✓ Beta Ointment
	Beta Ointment to be Principal Supply on 1 February 2025			
*	Lotn 0.1%3	0.00	50 ml OP	✓ Betnovate

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy		Fully	Brand or
	(Manufacturer's F	Price) Subs Per	idised •	Generic Manufacturer
	Ψ	rei		Manufacturer
CLOBETASOL PROPIONATE	0.40	00 - 00	, ,	N 1
* Crm 0.05% * Oint 0.05%		30 g OP	-	<u>Dermol</u>
	2.33	30 g OP	V į	<u>Dermol</u>
CLOBETASONE BUTYRATE	T 00	00 = 00		
Crm 0.05%	5.38	30 g OP		Eumovate
LIVERGOOFFICONE	(10.00)			Lumovale
HYDROCORTISONE  ** Crm 19/ Only on a prescription	1 70	20 ~ OD	./ 1	-thia
* Crm 1% – Only on a prescription	20.40	30 g OP 500 g	-	<u>Ethics</u> Noumed
* Powder – Only in combination		25 g	_	ABM
Up to 5% in a dermatological base (not proprietary Topic				
galenicals				
HYDROCORTISONE AND PARAFFIN LIQUID AND LANOLIN				
Lotn 1% with paraffin liquid 15.9% and lanolin 0.6% - Only o			_	
a prescription	12.83	250 ml	<b>√</b> [	DP Lotn HC
HYDROCORTISONE BUTYRATE			_	
Lipocream 0.1%		100 g OP		Locoid Lipocream
Oint 0.1%		100 g OP	_	Locoid
Milky emul 0.1%	12.33	100 ml OP	<b>✓</b> [	Locoid Crelo
METHYLPREDNISOLONE ACEPONATE			_	
Crm 0.1%		15 g OP	-	Advantan
Oint 0.1%	4.95	15 g OP	• 1	<u>Advantan</u>
MOMETASONE FUROATE				
Crm 0.1%		15 g OP		Elocon Alcohol Free
Elocon Alcohol Free to be Principal Supply on 1 February	3.50	50 g OP	• [	Elocon Alcohol Free
Oint 0.1%		15 g OP	<b>✓</b> 1	Elocon
GIII G.1 / S.	3.50	50 g OP		Elocon
Elocon to be Principal Supply on 1 February 2025				
Lotn 0.1%	4.99	30 ml OP	<b>✓</b> [	Elocon
Elocon to be Principal Supply on 1 February 2025				
TRIAMCINOLONE ACETONIDE				
Crm 0.02%	6.49	100 g OP	1	<u>Aristocort</u>
Oint 0.02%	6.54	100 g OP	<b>✓</b> <u>I</u>	<u>Aristocort</u>
Corticosteroids - Combination				
BETAMETHASONE VALERATE WITH SODIUM FUSIDATE [FUS	SIDIC ACIDI			
Crm 0.1% with sodium fusidate (fusidic acid) 2%	•	15 g OP		
Onn o. 1 /o man occuran racidate (racidie acid) 2 /o	(10.45)	10 9 01	F	-ucicort
a) Maximum of 15 g per prescription	( /			
b) Only on a prescription				
HYDROCORTISONE WITH MICONAZOLE - Only on a prescript	tion			
* Crm 1% with miconazole nitrate 2%		15 g OP	<b>√</b> [	Micreme H
Micreme H to be Principal Supply on 1 February 2025		•		
HYDROCORTISONE WITH NATAMYCIN AND NEOMYCIN - O	nly on a prescrip	otion		
Oint 1% with natamycin 1% and neomycin sulphate 0.5%		15 g OP	<b>✓</b> [	Pimafucort
		•		

			LINIMATOLOGICALO
	Subsidy (Manufacturer's I \$		Fully Brand or dised Generic  Manufacturer
TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYC	ng		
and gramicidin 250 mcg per g - Only on a prescription	(9.28)	15 g OP	Viaderm KC
Barrier Creams and Emollients			
<b>Barrier Creams</b>			
DIMETHICONE  * Crm 5% pump bottle	4.30	500 ml OP	✓ <u>healthE</u> Dimethicone 5%
* Crm 10% pump bottle	4.52	500 ml OP	✓ healthE  Dimethicone 10%
ZINC AND CASTOR OIL  * Oint	4.25	500 g	✓ <u>Evara</u>
Emollients			
AQUEOUS CREAM Crm	1.30	100 g	✓ healthE Aqueous Cream SLS Free
	1.65 1.73	500 g	✓ <u>Evara</u> ✓ GEM Aqueous Cream
(healthE Aqueous Cream SLS Free Crm to be delisted 1 March (GEM Aqueous Cream Crm to be delisted 1 March 2025)	2025)		
CETOMACROGOL  * Crm BP  Cetomacrogol-AFT to be Principal Supply on 1 Februar		500 g	✓ Cetomacrogol-AFT
CETOMACROGOL WITH GLYCEROL Crm 90% with glycerol 10%	2.13 3.50	500 ml OP 1,000 ml OP	✓ <u>Evara</u> ✓ <u>Evara</u>
* Oint BP	3.13	500 g	✓ Emulsifying Ointment ADE
OIL IN WATER EMULSION  * Crm	2.04 2.10	500 g	<ul> <li>✓ Fatty Cream AFT</li> <li>✓ Fatty Emulsion</li> <li>Cream (Evara)</li> </ul>
Fatty Emulsion Cream (Evara) to be Principal Supply of (Fatty Cream AFT Crm to be delisted 1 April 2025)	n 1 April 2025		,/
PARAFFIN Oint liquid paraffin 50% with white soft paraffin 50%	4.94	500 g OP	✓ White Soft Liquid Paraffin AFT
UREA			<i>4</i> =

71

✓ healthE Urea Cream

100 g OP

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

# **DERMATOLOGICALS**

	Subsidy (Manufacturer's I \$	Price) Subsi	Fully idised	Brand or Generic Manufacturer
WOOL FAT WITH MINERAL OIL - Only on a prescription				
* Lotn hydrous 3% with mineral oil	5.60	1,000 ml		
	(14.96)			P Lotion
	(20.53)		Д	Alpha-Keri Lotion
	1.40	250 ml OP		
	(5.87)			P Lotion
	5.60	1,000 ml		
	(23.91)		В	BK Lotion
	1.40	250 ml OP		
	(7.73)		В	BK Lotion

# **Other Dermatological Bases**

PA	RA	FF	IN
----	----	----	----

White soft - Only in combination	4.74	450 g	✓ EVARA White Soft
	19.00	2,500 g	Paraffin ✓ EVARA White Soft Paraffin

Only in combination with a dermatological galenical or as a diluent for a proprietary Topical Corticosteroid – Plain.

# **Minor Skin Infections**

POVIDONE IODINE			
Oint 10%	7.40	65 g OP	✓ Betadine
a) Maximum of 130 g per prescription			
b) Only on a prescription			
Antiseptic Solution 10%	4.99	100 ml	✓ Riodine
Antiseptic soln 10%	3.83	15 ml	✓ Riodine
·	6.99	500 ml	✓ Riodine
Skin preparation, povidone iodine 10% with 30% alcohol	1.63	100 ml	
	(3.48)		Betadine Skin Prep

# **Parasiticidal Preparations**

DI	ΝЛΕ	=T	ш	$\sim$	$\sim$	NI	_
וט	IVII	= 1	ПΙ	U	U	IV	ᆮ

* Lotn 4%	200 ml OP	✓ healthE  Dimethicone 4%  Lotion
IVERMECTIN - Special Authority see SA2294 below - Retail pharmacy		

- Ivermectin available on BSO provided the BSO includes a valid Special Authority for a patient of the institution.
- For the purposes of subsidy of ivermectin, institution means age related residential care facilities, disability care facilities or prisons.

# **⇒SA2294** Special Authority for Subsidy

Initial application — (Scabies) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

### DERMATOLOGICALS

Subsidy (Manufacturer's Price)			Brand or Generic
 \$	Per	•	Manufacturer

continued...

Either:

- 1 The person has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
- 2 Both:
  - 2.1 The person has a confirmed diagnosis of scabies or is a close contact of a scabies case; and
  - 2.2 Fither:
    - 2.2.1 The person is unable to complete topical therapy; or
    - 2.2.2 Previous treatment with topical therapy has been tried and not cleared the infestation.

**Initial application — (Other parasitic infections)** from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 filariasis: or
- 2 cutaneous larva migrans (creeping eruption); or
- 3 strongyloidiasis.

Renewal — (Scables) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria: Either:

- 1 The person has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
- 2 Both:
  - 2.1 The person has a confirmed diagnosis of scabies or is a close contact of a scabies case; and
  - 2.2 Fither:
    - 2.2.1 The person is unable to complete topical therapy; or
    - 2.2.2 Previous treatment with topical therapy has been tried and not cleared the infestation.

Renewal — (Other parasitic infections) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 filariasis: or
- 2 cutaneous larva migrans (creeping eruption); or
- 3 strongyloidiasis.

#### **PERMETHRIN**

## **Psoriasis and Eczema Preparations**

ACITRETIN - Special Authority see SA2024 below - Retail ph	narmacy		
Cap 10 mg	26.20	60	Novatretin
Cap 25 mg	57.37	60	✓ Novatretin

#### ⇒SA2024 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around acitretin and is competent to prescribe acitretin; and
- 3 Either:
  - 3.1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and that they must not become pregnant during treatment and for a period of three years after the completion of treatment; or



Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✓ Manufacturer	
----------------------------------------------------------------------------------------	--

continued...

3.2 Patient is not of child bearing potential.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

- Fither:
  - 1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and that they must not become pregnant during treatment and for a period of three years after the completion of treatment:
  - 2 Patient is not of child bearing potential.

BETAMETHASONE DIPROPIONATE WITH CALCIPOTRIOL		
Foam spray 500 mcg with calcipotriol 50 mcg per g59.95	60 g OP	<ul><li>Enstilar</li></ul>
Gel 500 mcg with calcipotriol 50 mcg per g40.92	60 g OP	Daivobet
Oint 500 mcg with calcipotriol 50 mcg per g14.31		✓ Daivobet
CALCIPOTRIOL		
Oint 50 mcg per g40.00	120 g OP	Daivonex
COAL TAR		
Soln BP - Only in combination36.25	200 ml	✓ Midwest

- 1) Up to 10% only in combination with a dermatological base or proprietary Topical Corticosteriod Plain
- 2) With or without other dermatological galenicals.

#### COAL TAR WITH ALLANTOIN, MENTHOL, PHENOL AND SULPHUR Cala CO/ with a dalah w O CO/ manufical O 700/ mbanal O CO/ and

COAL TAR WITH SALICYLIC ACID AND SULPHUR Soln 12% with salicylic acid 2% and sulphur 4% oint	4.97	25 g OP	✓ Coco-Scalp
COAL TAR WITH CALICYLIC ACID AND CHILDHIRD	, ,		01 7
	(4.35)		Egopsoryl TA
	3.43	30 g OP	
	(8.00)		Egopsoryl TA
allantoin crm 2.5%	6.59	75 g OP	
Soin 5% with sulphur 0.5%, menthol 0.75%, phenol 0.5% and			

PIMECROLIMUS - Special Authority see SA1970 below - Retail pharmacy

- a) Maximum of 15 g per prescription
- b) Note: a maximum of 15 g per prescription and no more than one prescription per 12 weeks.

Cream 1%......33.00 15 g OP ✓ Elidel

### ⇒SA1970 Special Authority for Subsidy

Initial application only from a dermatologist, paediatrician, ophthalmologist or any relevant practitioner on the recommendation of a dermatologist, paediatrician or ophthalmologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

7.95

40 a OP

✓ Coco-Scalp

#### Both:

- 1 Patient has atopic dermatitis on the eyelid; and
- 2 Patient has at least one of the following contraindications to topical corticosteroids: periorificial dermatitis, rosacea, documented epidermal atrophy, documented allergy to topical corticosteroids, cataracts, glaucoma, or raised intraocular

PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORESCEIN - Only on a prescription

\* Soln 2.3% with trolamine laurilsulfate and fluorescein sodium...........5.41 500 ml **Pinetarsol** 

		I	DERM	ATOLOGICALS
	Subsidy (Manufacturer's Pr	rice) Sub:	Fully sidised	Brand or Generic Manufacturer
SALICYLIC ACID				
Powder – Only in combination		250 g al Corticostero		lidwest ain or collodion flexible
SULPHUR Precipitated - Only in combination		100 g al Corticostero		<b>lidwest</b> ain
TACROLIMUS				
Oint 0.1% – Special Authority see SA2074 below – Retail pharmacy	33.00	30 g OP	<b>√</b> <u>Z</u>	<u>ematop</u>
b) Note: a maximum of 30 g per prescription and no n	nore than one pres	cription per 1	2 weeks	i.
SA2074   Special Authority for Subsidy   Initial application only from a dermatologist, paediatrician or an paediatrician, . Approvals valid without further renewal unless r Both:  1    Patient has atopic dermatitis on the face; and 2    Patient has at least one of the following contraindications documented epidermal atrophy or documented allergy to	notified for applications to topical corticos	ions meeting to	the follo	wing criteria:
	topical corticoster	oldo.		
Scalp Preparations  BETAMETHASONE VALERATE  * Scalp app 0.1%  Beta Scalp to be Principal Supply on 1 February 2025	12.95	100 ml OP	<b>✓</b> E	leta Scalp
CLOBETASOL PROPIONATE	2.22	00 100		
* Scalp app 0.05%	6.26	30 ml OP	<b>✓</b> <u>L</u>	<u>lermol</u>
Scalp lotn 0.1%	6.57	100 ml OP	<b>✓</b> L	ocoid
KETOCONAZOLE Shampoo 2%	3.23 4.09	100 ml OP	_	ebizole ebizole
<ul><li>a) Maximum of 100 ml per prescription</li><li>b) Only on a prescription</li></ul>				
Sunscreens				
SUNSCREENS, PROPRIETARY – Subsidy by endorsement				
Only if prescribed for a patient with severe photosensitivity	secondary to a def	ined clinical c	ondition	and the prescription is

endorsed accordingly. 200 g OP

Lotn,......6.50

✓ Marine Blue Lotion SPF 50+

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

### **DERMATOLOGICALS**

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

✓ Efudix

### **Wart Preparations**

For salicylic acid preparations refer to PSORIASIS AND ECZEMA PREPARATIONS, page 73

**PODOPHYLLOTOXIN** 

- a) Maximum of 3.5 ml per prescription
- b) Only on a prescription

### **Other Skin Preparations**

### **Antineoplastics**

IMIQUIMOD

Crm 5%, 250 mg sachet......21.72 24 **✓ Perrigo** 

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

		Subsidy (Manufacturer's Price)		Fully Subsidised	
		(Manufacturer's Price) \$	Per	Subsidised	
C	ontraceptives - Non-hormonal				
C	ondoms				
-	NDOMS				
	49 mm - Up to 144 dev available on a PSO		144	•	Moments
K	53 mm	1.15	10	•	Moments
		14.25	144	•	Moments
	<ul> <li>a) Maximum of 60 dev per prescription</li> </ul>				
	b) Up to 60 dev available on a PSO				
K	53 mm, 0.05 mm thickness	1.15	10	✓	Moments
		14.25	144	✓	Moments
	a) Up to 60 dev available on a PSO				
	b) Maximum of 60 dev per prescription				
6	53 mm, chocolate, brown	1.15	10	/	Moments
		14.25	144		Moments
	a) Up to 60 dev available on a PSO	11.20		•	
	b) Maximum of 60 dev per prescription				
K	53 mm, strawberry, red	1 15	10		Moments
•	oo min, shawbony, rou	14.25	144		Moments
	a) Un to 60 day available as a DCO	14.20	144	•	MOHIEHIS
	a) Up to 60 dev available on a PSO				
	b) Maximum of 60 dev per prescription	4.45	40	,	
÷	56 mm		10		Moments
		14.50	144	•	Moments
	a) Maximum of 60 dev per prescription				
	b) Up to 60 dev available on a PSO				
6	56 mm, 0.05 mm thickness		12		Gold Knight
		24.10	144	•	Gold Knight
	a) Up to 60 dev available on a PSO				
	b) Maximum of 60 dev per prescription				
6	56 mm, 0.05mm thickness (bulk pack)	20.17	144	•	Gold Knight
	a) Maximum of 60 dev per prescription				-
	b) Up to 60 dev available on a PSO				
6	56 mm, 0.08 mm thickness	1.15	10	/	Moments
		14.25	144	/	Moments
	a) Up to 60 dev available on a PSO				
	b) Maximum of 60 dev per prescription				
6	56 mm, 0.08 mm thickness, red	1.15	10	/	Moments
	, , , , , , , , , , , , , , , , , , , ,	14.25	144		Moments
	a) Up to 60 dev available on a PSO	0		-	<del>-</del>
	b) Maximum of 60 dev per prescription				
6	56 mm, chocolate	1 70	12	1	Gold Knight
•	oo mm, onocolate	21.45	144		Gold Knight
	a) Unito 60 day available and PCO	21.40	1-1-4	•	Gold Killylit
	a) Up to 60 dev available on a PSO				
	b) Maximum of 60 dev per prescription 56 mm, strawberry	1 70	10		Cold Knieht
÷	oo miin, strawberry		12		Gold Knight
	\	21.45	144	•	Gold Knight
	a) Up to 60 dev available on a PSO				
	b) Maximum of 60 dev per prescription				
+	60 mm		12		Gold Knight XL
		21.89	144	•	Gold Knight XL
	a) Maximum of 60 dev per prescription				
_	b) 🎜 prita p Sude varse ilable on a PSO	S29 Unapprove	d med	icine suppli	ed under Section 29
ŏ	60 mm/hHttparetipply	cdiz Z8haidiaad	c144	lv 📝	Gold Knight XL

Subsidy	ce) Su	Fully	Brand or	
(Manufacturer's Pric		bsidised	Generic	
\$	Per	1	Manufacturer	

- a) Maximum of 60 dev per prescription
- b) Up to 60 dev available on a PSO

### **Contraceptive Devices**

#### INTRA-UTERINE DEVICE

- a) Up to 40 dev available on a PSO
- b) Only on a PSO

*	IUD 29.1 mm length × 23.2 mm width	29.80	1	✓ Choice 380 7med  Nsha Silver/ copper Short
*	IUD 33.6 mm length × 29.9 mm width	26.80	1	✓ TCu 380 Plus
*	IUD 35.5 mm length × 19.6 mm width	33.00	1	Normal  ✓ Cu 375 Standard

### **Contraceptives - Hormonal**

### **Combined Oral Contraceptives**

### ⇒SA0500 Special Authority for Alternate Subsidy

**Initial application** from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Either:
  - 1.1 Patient is on a Social Welfare benefit; or
  - 1.2 Patient has an income no greater than the benefit; and
- 2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient is on a Social Welfare benefit; or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- · on a Social Welfare benefit; or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

#### ETHINYLOESTRADIOL WITH DESOGESTREL

*	Tab 20 mcg with desogestrel 150 mcg and 7 inert tab - Up to			
	84 tab available on a PSO	10.00	84	✓ Mercilon 28

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
ETHINYLOESTRADIOL WITH LEVONORGESTREL				
* Tab 20 mcg with levonorgestrel 100 mcg and 7 inert tablets	_			
Up to 84 tab available on a PSO	1.50	84	<b>√</b> <u>L</u>	o-Oralcon 20 ED
* Tab 30 mcg with levonorgestrel 150 mcg	6.62	63		
	(16.50)		N	/licrogynon 30
<ul><li>a) Higher subsidy of \$15.00 per 63 tab with Special Aut</li><li>b) Up to 63 tab available on a PSO</li></ul>	hority see SA0500 on	the	previous pag	ge
* Tab 30 mcg with levonorgestrel 150 mcg and 7 inert tablets Up to 84 tab available on a PSO		84	<b>√</b> <u>c</u>	Oralcon 30 ED
ETHINYLOESTRADIOL WITH NORETHISTERONE				
Tab 35 mcg with norethisterone 1 mg and 7 inert tab - Up to 84 tab available on a PSO		84		Alyacen Brevinor 1/28
Tab 35 mcg with norethisterone 500 mcg and 7 inert tab - L	Jp			
to 84 tab available on a PSO	21.99	84	<b>√</b> N	lorimin

### **Progestogen-only Contraceptives**

### ⇒SA0500 Special Authority for Alternate Subsidy

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria:

- 1 Either:
  - 1.1 Patient is on a Social Welfare benefit; or
  - 1.2 Patient has an income no greater than the benefit; and
- 2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient is on a Social Welfare benefit; or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- on a Social Welfare benefit; or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

* Tab 30 mcg - Up to 84 tab available on a PSO	16.50	84 112	✓ Microlut ✓ Microlut
	22.00	112	• Wilci Olut
* Subdermal implant (2 x 75 mg rods) – Up to 3 pack available			
on a PSO	106.92	1	✓ <u>Jadelle</u>
MEDROXYPROGESTERONE ACETATE			
Inj 150 mg per ml, 1 ml syringe – Up to 5 inj available on a PSO	9.18	1	Depo-Provera

		GENIII	U-UKI	NARY SYSTEM
	Subsidy (Manufacturer's Pric	ee) Subs	Fully sidised	Brand or Generic Manufacturer
NORETHISTERONE Tab 350 mcg – Up to 84 tab available on a PSO	12.25	84	✓ N	lorethinderone - CDC loriday loriday 28
<b>Emergency Contraceptives</b>				
# Tab 1.5 mg	1.75	1	<b>√</b> <u>L</u>	evonorgestrel BNM
b) Up to 5 tab available on a PSO c) Note: Direct Provision by a pharmacist permitted u	nder the provisions	n Part I of S	ection A	١.
Antiandrogen Oral Contraceptives				
Prescribers may code prescriptions "contraceptive" (code "O") v and prescription charge will be as per other contraceptives, as f		ed for contra	aceptior	n. The period of supply
non-contraceptive period of supply. ie. Prescriptions may be w CYPROTERONE ACETATE WITH ETHINYLOESTRADIOL	nt) may apply.  Intraceptive prescript viritten for up to three		pply.	oly, and the
<ul> <li>prescription may be written for up to six months supply.</li> <li>Prescriptions coded in any other way are subject to any non connon-contraceptive period of supply. ie. Prescriptions may be w</li> <li>CYPROTERONE ACETATE WITH ETHINYLOESTRADIOL</li> <li>Tab 2 mg with ethinyloestradiol 35 mcg and 7 inert tabs</li> </ul>	nt) may apply.  Intraceptive prescript viritten for up to three	months sup	pply.	
prescription may be written for up to six months supply.  Prescriptions coded in any other way are subject to any non con non-contraceptive period of supply. ie. Prescriptions may be w CYPROTERONE ACETATE WITH ETHINYLOESTRADIOL      * Tab 2 mg with ethinyloestradiol 35 mcg and 7 inert tabs — to 168 tab available on a PSO	nt) may apply.  Intraceptive prescript written for up to three  Up  CACID  ate	months sup	<b>√</b> <u>G</u>	
prescription may be written for up to six months supply.  Prescriptions coded in any other way are subject to any non connon-contraceptive period of supply. ie. Prescriptions may be we CYPROTERONE ACETATE WITH ETHINYLOESTRADIOL      Tab 2 mg with ethinyloestradiol 35 mcg and 7 inert tabs — to 168 tab available on a PSO	ntraceptive prescript rritten for up to three up to the up to three up to the up t	months sup	<b>√</b> <u>G</u> A	iinet
<ul> <li>prescription may be written for up to six months supply.</li> <li>Prescriptions coded in any other way are subject to any non connon-contraceptive period of supply. ie. Prescriptions may be w</li> <li>CYPROTERONE ACETATE WITH ETHINYLOESTRADIOL</li> <li>* Tab 2 mg with ethinyloestradiol 35 mcg and 7 inert tabs — to 168 tab available on a PSO</li> <li>Gynaecological Anti-infectives</li> <li>ACETIC ACID WITH HYDROXYQUINOLINE AND RICINOLEIC Jelly with glacial acetic acid 0.94%, hydroxyquinoline sulph</li> </ul>	ntraceptive prescript rritten for up to three up to the up to th	168 100 g OP 35 g OP	Pply. ✓ G	inet ci-Jel

# ERGOMETRINE MALEATE

Inj 500 mcg per ml, 1 ml ampoule - Up to 5 inj available on a	100.00	-	A DDI. Europeatrino
PSO	. 160.00	5	DBL Ergometrine
OESTRIOL			
* Crm 1 mg per g with applicator	6.95	15 g OP	<ul><li>Ovestin</li></ul>
* Pessaries 500 mcg	7.55	15	✓ Ovestin

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price) \$	S Per	Fully ubsidised	Brand or Generic Manufacturer
OXYTOCIN – Up to 5 inj available on a PSO				
Inj 5 iu per ml, 1 ml ampoule	4.98	5	✓ (	Dxytocin BNM
Inj 10 iu per ml, 1 ml ampoule	5.98	5	✓ (	Dxytocin BNM
	11.96	10	<b>√</b> (	Oxytocin Panpharma
OXYTOCIN WITH ERGOMETRINE MALEATE – Up to 5 inj availa Inj 5 iu with ergometrine maleate 500 mcg per ml, 1 ml ampoul		5	<b>√</b> <u>9</u>	Syntometrine

### **Pregnancy Tests - hCG Urine**

David One Step Cassette Pregnancy Test to be Principal Supply on 1 March 2025 (Smith BioMed Rapid Pregnancy Test Cassette to be delisted 1 March 2025)

### **Urinary Agents**

For urinary tract Infections refer to INFECTIONS, Antibacterials, page 114

### 5-Alpha Reductase Inhibitors

FINASTERIDE - Special Authority see SA0928 below - Retail pharmacy

★ Tab 5 mg .......4.79 100 ✓ Ricit

#### ⇒SA0928 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

### Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 Either:
  - 2.1 The patient is intolerant of non-selective alpha blockers or these are contraindicated; or
  - 2.2 Symptoms are not adequately controlled with non-selective alpha blockers.

### Alpha-1A Adrenoreceptor Blockers

TAMSULOSIN HYDROCHLORIDE − Special Authority see SA1032 below − Retail pharmacy

\* Cap 400 mcg .......22.31 100 ✓ Tamsulosin-Rex

### ⇒SA1032 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 The patient is intolerant of non-selective alpha blockers or these are contraindicated.

**Pregnancy Test** 

		G.E.I.I.I.	J 0	
	Subsidy (Manufacturer's P \$	rice) Subs Per	Fully idised	Brand or Generic Manufacturer
Other Urinary Agents				
OXYBUTYNIN			_	
* Tab 5 mg	5.42	100	<b>✓</b> A	llchemy Oxybutynin
POTASSIUM CITRATE				Oxybutyiiii
Oral liq 3 mmol per ml - Special Authority see SA1083 below	N —			
Retail pharmacy	37.49	200 ml OP	<b>✓</b> E	Biomed
<ul> <li>SA1083 Special Authority for Subsidy</li> <li>Initial application from any relevant practitioner. Approvals valid</li> <li>Both:</li> <li>The patient has recurrent calcium oxalate urolithiasis; and</li> </ul>		or applications	meetin	g the following criteria:
2 The patient has had more than two renal calculi in the two				
Renewal from any relevant practitioner. Approvals valid for 2 yes benefitting from the treatment.	ars where the tre	atment remain	s appro	ppriate and the patient is
SODIUM CITRO-TARTRATE  * Grans eff 4 g sachets	3.50	28	<b>√</b> U	Iral
SOLIFENACIN SUCCINATE		20	· <u>·</u>	nui
Tab 5 mg	1.95	30	<b>√</b> S	olifenacin succinate Max Health
Tab 10 mg	2.05 3.53	30	-	olifenacin Viatris olifenacin succinate Max
				Health
	3.72		<b>√</b> S	olifenacin Viatris
(Solifenacin Viatris Tab 5 mg to be delisted 1 June 2025) (Solifenacin Viatris Tab 10 mg to be delisted 1 June 2025)				
Detection of Substances in Urine				
ORTHO-TOLIDINE				
* Compound diagnostic sticks		50 test OP	_	lemastix
TETRABROMOPHENOL	(8.25)		Г	ισιπαδιίλ
* Blue diagnostic strips	13.92	100 test OP	<b>✓</b> A	lbustix
Obstetric Preparations				
Antiprogesterones				
MIFEPRISTONE	00.00	,		116 an an a
Tab 200 mg - Up to 15 tab available on a PSO	83.90 180.00	1 3		lifegyne lifegyne

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	osidised	Generic	
\$	Per	✓	Manufacturer	

### Calcium Homeostasis

CALCITONIN			
* Inj 100 iu per ml, 1 ml ampoule	121.00	5	✓ Miacalcic
CINACALCET - Special Authority see SA2170 below - Retail	pharmacy		
Tab 30 mg - Wastage claimable	25.24	28	<ul> <li>Cinacalet Devatis</li> </ul>
Tab 60 mg - Wastage claimable	50.47	28	✓ Cinacalet Devatis

### ⇒SA2170 Special Authority for Subsidy

Initial application — (parathyroid carcinoma or calciphylaxis) only from a nephrologist or endocrinologist. Approvals valid for 6 months for applications meeting the following criteria:

- Fither: 1 All of the following:
  - 1.1 The patient has been diagnosed with a parathyroid carcinoma (see Note); and
  - 1.2 The patient has persistent hypercalcaemia (serum calcium greater than or equal to 3 mmol/L) despite previous first-line treatments including sodium thiosulfate (where appropriate) and bisphosphonates; and
  - 1.3 The patient is symptomatic; or
  - 2 All of the following:
    - 2.1 The patient has been diagnosed with calciphylaxis (calcific uraemic arteriolopathy); and
    - 2.2 The patient has symptomatic (e.g. painful skin ulcers) hypercalcaemia (serum calcium greater than or equal to 3 mmol/L): and
    - 2.3 The patient's condition has not responded to previous first-line treatments including bisphosphonates and sodium

Renewal — (parathyroid carcinoma or calciphylaxis) only from a nephrologist or endocrinologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

- Both:
  - 1 The patient's serum calcium level has fallen to < 3mmol/L; and
  - 2 The patient has experienced clinically significant symptom improvement.

Note: This does not include parathyroid adenomas unless these have become malignant.

Initial application — (primary hyperparathyroidism) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### All of the following:

- 1 Patient has primary hyperparathyroidism; and
- 2 Either:
  - 2.1 Patient has hypercalcaemia of more than 3 mmol/L with or without symptoms; or
  - 2.2 Patient has hypercalcaemia of more than 2.85 mmol/L with symptoms; and
- 3 Surgery is not feasible or has failed; and
- 4 Patient has other comorbidities, severe bone pain, or calciphylaxis.

Initial application — (secondary or tertiary hyperparathyroidism) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 Patient has tertiary hyperparathyroidism and markedly elevated parathyroid hormone (PTH) with hypercalcaemia:
  - 1.2 Patient has symptomatic secondary hyperparathyroidism and elevated PTH; and
- 2 Patient is on renal replacement therapy: and
- 3 Any of the following:
  - 3.1 Residual parathyroid tissue has not been localised despite repeat unsuccessful parathyroid explorations; or

S	Subsidy	Fully	Brand or
(Manufa	cturer's Price) Su	ıbsidised	Generic
	\$ Per	✓	Manufacturer

continued...

- 3.2 Parathyroid tissue is surgically inaccessible; or
- 3.3 Parathyroid surgery is not feasible.

**Renewal — (secondary or tertiary hyperparathyroidism)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

#### Fither:

- 1 The patient has had a kidney transplant, and following a treatment free interval of at least 12 weeks a clinically acceptable parathyroid hormone (PTH) level to support ongoing cessation of treatment has not been reached; or
- 2 The patient has not received a kidney transplant and trial of withdrawal of cinacalcet is clinically inappropriate.

#### **ZOLEDRONIC ACID**

Inj 4 mg per 5 ml, vial	15.65	1	1	Zoledronic acid
				Viatris

### Corticosteroids and Related Agents for Systemic Use

BETAMETHASONE SODIUM PHOSPHATE WITH BETAMETHASONE ACETATE		
* Inj 3.9 mg with betamethasone acetate 3 mg per ml, 1 ml	5	
(36.96)		Celestone
		Chronodose
DEXAMETHASONE		
* Tab 0.5 mg - Up to 60 tab available on a PSO1.80	30	<ul><li>Dexmethsone</li></ul>
Dexmethsone to be Principal Supply on 1 February 2025		
* Tab 4 mg - Up to 30 tab available on a PSO	30	<ul><li>Dexmethsone</li></ul>
Dexmethsone to be Principal Supply on 1 February 2025		
Oral liq 1 mg per ml53.86	25 ml OP	✓ Biomed
DEXAMETHASONE PHOSPHATE		
Dexamethasone phosphate injection will not be funded for oral use.		
* Inj 4 mg per ml, 1 ml ampoule – Up to 5 inj available on a PSO7.86	10	✓ <u>Hameln</u>
* Inj 4 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO13.10	10	✓ <u>Hameln</u>
FLUDROCORTISONE ACETATE		
* Tab 100 mcg11.46	100	✓ Florinef
HYDROCORTISONE		
* Tab 5 mg8.10	100	✓ Douglas
* Tab 20 mg20.32	100	Douglas
* Inj 100 mg vial	1	✓ Solu-Cortef
a) Not on a BSO		
b) Up to 5 inj available on a PSO		
METHYLPREDNISOLONE		
* Tab 4 mg112.00	100	✓ Medrol
* Tab 100 mg223.10	20	✓ Medrol
METHYLPREDNISOLONE (AS SODIUM SUCCINATE)		
Inj 40 mg vial22.30	1	✓ Solu-Medrol-Act-
		O-Vial
Inj 125 mg vial34.10	1	✓ Solu-Medrol-Act-
		O-Vial
Inj 500 mg vial26.88	1	✓ Solu-Medrol-Act-
, 3	•	O-Vial
Inj 1 g vial32.84	1	✓ Solu-Medrol

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

,	Subsidy		Fully Brand or
(	Manufacturer's Price \$	e) Subsid Per	ised Generic  Manufacturer
METHYLPREDNISOLONE ACETATE			
Inj 40 mg per ml, 1 ml vial	47.06	5	✓ Depo-Medrol
REDNISOLONE			
Oral liq 5 mg per ml – Up to 30 ml available on a PSO	6.00	30 ml OP	✓ Redipred
REDNISONE			
F Tab 1 mg		500	✓ Prednisone Clinect
: Tab 2.5 mg		500	✓ Prednisone Clinect
Tab 5 mg - Up to 30 tab available on a PSO	19.30	500	✓ Prednisone Clinect
Tab 20 mg - Up to 30 tab available on a PSO	50.51	500	✓ Prednisone Clinect
ETRACOSACTRIN			
Inj 250 mcg per ml, 1 ml ampoule	86.25	1	✓ Synacthen
,			✓ UK Synacthen
Inj 1 mg per ml, 1 ml ampoule	690.00	1	✓ Synacthen Depot
, .,			✓ Synacthene
			Retard S29
RIAMCINOLONE ACETONIDE			
Inj 10 mg per ml, 1 ml ampoule	21.42	5	✓ Kenacort-A 10
Inj 40 mg per ml, 1 ml ampoule		5	✓ Kenacort-A 40
Sex Hormones Non Contraceptive			
·			
Androgen Agonists and Antagonists			
YPROTERONE ACETATE			
Tab 50 mg		50	✓ Siterone
Tab 100 mg	28.03	50	✓ Siterone
ESTOSTERONE			
Gel (transdermal) 16.2 mg per g	52.00	88 g OP	✓ Testogel
ESTOSTERONE CIPIONATE			
Inj 100 mg per ml, 10 ml vial	85.00	1	✓ Depo-Testosterone
ESTOSTERONE ESTERS			
Inj 250 mg per ml, 1 ml	12.00	1	✓ Sustanon Ampoules
	12.30	1	- Justanon Ampudies
ESTOSTERONE UNDECANOATE			
Cap 40 mg - Subsidy by endorsement		100	✓ Steril-Gene S29
Subsidy by endorsement – subsidised for patients who we			
1 November 2021 and the prescription is endorsed accord			
where there exists a record of prior dispensing of testoster	rone undecanoate	cap 40 mg in	the preceding 12 months.

Inj 250 mg per ml, 4 ml vial......86.00

✓ Reandron 1000

Subsidy Fully Brand or
(Manufacturer's Price) Subsidised Generic
\$ Per ✓ Manufacturer

### Hormone Replacement Therapy - Systemic

UES	u	υy	CI	ı

	•			
-	STRADIOL Tab 1 mg	A 12	28 OP	
~	Tab Ting	(11.10)	20 01	Estrofem
*	Tab 2 mg	, ,	28 OP	LStrotein
*	rab 2 mg		28 UP	Catuatana
	0-1/h1\ 0.000/ (750/1\	(11.10)	00 · OD	Estrofem
	Gel (transdermal) 0.06% (750 mcg/actuation)		80 g OP	✓ Estrogel
	Patch 25 mcg per day	9.85	8	Estradiol TDP Mylan
		13.50		✓ Estraderm MX S29
		14.50		✓ Estradot
		21.35		✓ Lyllana
	a) No more than 2 patch per week			-
	b) Only on a prescription			
	Patch 50 mcg per day	10.75	8	✓ Estradiol TDP Mylan
	. a.o. ooog po. aa,		ū	✓ Estradiol Viatris
		14.50		✓ Estraderm MX S29
		14.50		
				✓ Estradiol Sandoz
		04.55		✓ Estradot
		21.55		✓ Lyllana
	<ul> <li>a) No more than 2 patch per week</li> </ul>			
	b) Only on a prescription			
	Patch 75 mcg per day	11.88	8	<ul><li>Estradiol TDP Mylan</li></ul>
				<ul><li>Estradiol Viatris</li></ul>
		14.50		✓ Estradiol Sandoz
				✓ Estradot
		22.37		✓ Lyllana
	a) No more than 2 patch per week			•
	b) Only on a prescription			
	Patch 100 mcg per day	12.05	8	✓ Estradiol TDP Mylan
	Tater 100 meg per day	12.33	U	✓ Estradiol Viatris
		14.50		✓ Estradiol Sandoz
		14.50		✓ Estradioi Sandoz
		15.50		✓ Estraderm MX S29
		22.77		✓ Lyllana
	<ul> <li>a) No more than 2 patch per week</li> </ul>			
	b) Only on a prescription			
OF	STRADIOL VALERATE			
	Tab 1 mg	12.36	84	✓ Progynova
	Tab 2 mg		84	✓ Progynova
	-	12.00	04	- i logyilova
	STROGENS			
*	Conjugated, equine tab 300 mcg	3.01	28	
		(19.25)		Premarin
*	Conjugated, equine tab 625 mcg	4.12	28	
		(19.25)		Premarin

	Subsidy (Manufacturer's Price) \$	Per	Subsidised G	rand or eneric anufacturer
Progestogens				
EDROXYPROGESTERONE ACETATE				
Tab 2.5 mg	6.56	30	✓ Prov	era
	8.75	56	✓ Prov	
Tab 5 mg		56	✓ Prov	
	20.13	100	✓ Prov	
Tab 10 mg		30	✓ Prov	
Progestogen and Oestrogen Combined Prepara	itions			
ESTRADIOL WITH NORETHISTERONE				
Tab 1 mg with 0.5 mg norethisterone acetate	5.40	28 OF	1	
3 0 111 1 111111111	(18.10)		Kliov	ance
Tab 2 mg with 1 mg norethisterone acetate	` '	28 OF		-
J	(18.10)		Kliog	est
Tab 2 mg with 1 mg norethisterone acetate (10), and 2 mg	( /		09	
oestradiol tab (12) and 1 mg oestradiol tab (6)	5.40	28 OF	1	
ossitudioi tab (12) and 1 mg ossitudioi tab (0)	(18.10)	_0 01		quens
	(10.10)		11130	quens
Other Oestrogen Preparations				
ESTRIOL				
Tab 2 mg	7.70	30	✓ Oves	tin
•	-			
Other Progestogen Preparations				
EVONORGESTREL				
Intra-uterine device 52 mg	269.50	1	✓ Mire	na
Intra-uterine device 13.5 mg	215.60	1	✓ Jayd	ess
EDROXYPROGESTERONE ACETATE			-	
Tab 100 mg	133 57	100	✓ Prov	era HD
	100.07	100	5 110V	oiu IID
ORETHISTERONE	F 15			
Tab 5 mg - Up to 30 tab available on a PSO	5.49	30	✓ Prim	olut N
ROGESTERONE	44.05	00	<b>4</b>	
Cap 100 mg	14.85	30	✓ <u>Utro</u>	gestan
Thyroid and Antithyroid Agents				
ARBIMAZOLE				
Tab 5 mg	7.56	100	✓ Neo-	Mercazole
· ·				
EVOTHYROXINE Tab 05 mag	E	00	./ 0	امامسا
Tab 25 mcg		90	✓ Synt	
Tab 50 mcg		28		ury Pharma
	5.79	90	✓ Synt	
T 11 150	64.28	1,000	✓ Eltro	
Tablet 50 mcg		200	✓ Eltro	
Tab 100 mcg		28		ury Pharma
	6.01	90	✓ Synt	
	66.78	1,000		
Tablet 100 mcg	13.36	200	✓ Eltro	vin

	Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
	\$	Per	✓	Manufacturer
PROPYLTHIOURACIL - Special Authority see SA1199 below -	Retail pharmacy			
Tab 50 mg	35.00	100	<b>✓</b> P	TU \$29
<b>⇒SA1199</b> Special Authority for Subsidy				

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

- 1 The patient has hyperthyroidism; and
- 2 The patient is intolerant of carbimazole or carbimazole is contraindicated.

**Renewal** from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

### **Trophic Hormones**

#### **Growth Hormones**

SOMATROPIN (OMNITROPE) – Special Authority see SA2032 below – Reta	ail pharmacy	
* Inj 5 mg cartridge80.21	1	<ul><li>Omnitrope</li></ul>
		✓ Omnitrope S29 S29
Omnitrope to be Principal Supply on 1 February 2025		
* Inj 10 mg cartridge80.21	1	✓ Omnitrope
		✓ Omnitrope S29 S29
Omnitrope to be Principal Supply on 1 February 2025		
* Inj 15 mg cartridge	) 1	✓ Omnitrope
		✓ Omnitrope S29 S29
Omnitrope to be Principal Supply on 1 February 2025		
(Omnitrope S29 S29 Inj 5 mg cartridge to be delisted 1 February 2025)		

(Omnitrope S29 S29 Inj 15 mg cartridge to be delisted 1 February 2025)

(Omnitrope S29 S29 Inj 10 mg cartridge to be delisted 1 February 2025)

■ SA2032 Special Authority for Subsidy
Initial application — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist.

Approvals valid for 9 months for applications meeting the following criteria:

Fither:

- 1 Growth hormone deficiency causing symptomatic hypoglycaemia, or with other significant growth hormone deficient sequelae (e.g. cardiomyopathy, hepatic dysfunction) and diagnosed with GH < 5 mcg/l on at least two random blood samples in the first 2 weeks of life, or from samples during established hypoglycaemia (whole blood glucose < 2 mmol/l using a laboratory device); or</p>
- 2 All of the following:
  - 2.1 Height velocity < 25th percentile for age adjusted for bone age/pubertal status if appropriate over 6 or 12 months using the standards of Tanner and Davies (1985); and
  - 2.2 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
  - 2.3 Peak growth hormone value of < 5.0 mcg per litre in response to two different growth hormone stimulation tests. In children who are 5 years or older, GH testing with sex steroid priming is required; and</p>
  - 2.4 If the patient has been treated for a malignancy, they should be disease free for at least one year based upon follow-up laboratory and radiological imaging appropriate for the malignancy, unless there are strong medical reasons why this is either not necessary or appropriate; and
  - 2.5 Appropriate imaging of the pituitary gland has been obtained.

**Renewal — (growth hormone deficiency in children)** only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Price	,	Subsidised	Generic	
\$	Per		Manutacturer	

continued...

- 1 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 2 Height velocity is greater than or equal to 25th percentile for age (adjusted for bone age/pubertal status if appropriate) while on growth hormone treatment, as calculated over six months using the standards of Tanner and Davis (1985); and
- 3 Height velocity is greater than or equal to 2.0 cm per year, as calculated over 6 months; and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

**Initial application** — (**Turner syndrome**) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a post-natal genotype confirming Turner Syndrome; and
- 2 Height velocity is < 25th percentile over 6-12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is < 14 years.

Renewal — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile for age (while on growth hormone calculated over 6 to 12 months using the Ranke's Turner Syndrome growth velocity charts); and
- 2 Height velocity is greater than or equal to 2 cm per year, calculated over six months; and
- 3 A current bone age is 14 years or under : and
- 4 No serious adverse effect that the specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 3 standard deviations below the mean for age or for bone age if there is marked growth acceleration or delay; and
- 2 Height velocity is < 25th percentile for age (adjusted for bone age/pubertal status if appropriate), as calculated over 6 to 12 months using the standards of Tanner and Davies(1985); and
- 3 A current bone age is < 14 years or under (female patients) or < 16 years (male patients); and
- 4 The patient does not have severe chronic disease (including malignancy or recognized severe skeletal dysplasia) and is not receiving medications known to impair height velocity.

**Renewal** — **(short stature without growth hormone deficiency)** only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred

Initial application — (short stature due to chronic renal insufficiency) only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

1 The patient's height is more than 2 standard deviations below the mean; and

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic	
 (Manufacturer's Price)	Per	• dipsidised ✓	Manufacturer	

continued.

- 2 Height velocity is < 25th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and</p>
- 3 A current bone age is to 14 years or under (female patients) or to 16 years or under (male patients); and
- 4 The patient is metabolically stable, has no evidence of metabolic bone disease and absence of any other severe chronic disease; and
- 5 The patient is under the supervision of a specialist with expertise in renal medicine; and
- 6 Either:
  - 6.1 The patient has a GFR less than or equal to 30 ml/min/1.73m<sup>2</sup> as measured by the Schwartz method (Height(cm)/plasma creatinine (umol/l)) × 40 = corrected GFR (ml/min/1.73m<sup>2</sup>) in a child who may or may not be receiving dialysis; or
  - 6.2 The patient has received a renal transplant and has received < 5mg/ m²/day of prednisone or equivalent for at least 6 months..

Renewal — (short stature due to chronic renal insufficiency) only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not experienced significant biochemical or metabolic deterioration confirmed by diagnostic results; and
- 7 The patient has not received renal transplantation since starting growth hormone treatment; and
- 8 If the patient requires transplantation, growth hormone prescription should cease before transplantation and a new application should be made after transplantation based on the above criteria.

**Initial application — (Prader-Willi syndrome)** only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of Prader-Willi syndrome that has been confirmed by genetic testing or clinical scoring criteria;
- 2 The patient is aged six months or older; and
- 3 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
- 4 Sleep studies or overnight oximetry have been performed and there is no obstructive sleep disorder requiring treatment, or if an obstructive sleep disorder is found, it has been adequately treated under the care of a paediatric respiratory physician and/or ENT surgeon; and
- 5 Either:
  - 5.1 Both:
    - 5.1.1 The patient is aged two years or older; and
    - 5.1.2 There is no evidence of type II diabetes or uncontrolled obesity defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months; or
  - 5.2 The patient is aged between six months and two years and a thorough upper airway assessment is planned to be undertaken prior to treatment commencement and at six to 12 weeks following treatment initiation.

Renewal — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

(manufacturer's Price) Subsidised Generic  \$ Per ✓ Manufacturer		Subsidy (Manufacturer's Price) \$		Fully bsidised	Brand or Generic Manufacturer	
------------------------------------------------------------------	--	-----------------------------------------	--	----------------	-------------------------------------	--

continued...

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred: and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not developed type II diabetes or uncontrolled obesity as defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months.

Initial application — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a medical condition that is known to cause growth hormone deficiency (e.g. surgical removal of the pituitary for treatment of a pituitary tumour); and
- 2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
- 3 The patient has severe growth hormone deficiency (see notes); and
- 4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
- 5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

Renewal — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 All of the following:
  - 1.1 The patient has been treated with somatropin for < 12 months; and
  - 1.2 There has been an improvement in Quality of Life defined as a reduction of at least 8 points on the Quality of Life Assessment of Growth Hormone Deficiency in Adults (QoL-AGHDA®) score from baseline; and
  - 1.3 Serum IGF-I levels have been increased within ±1SD of the mean of the normal range for age and sex; and
  - 1.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients, or 1 mg per day for female patients; or
- 2 All of the following:
  - 2.1 The patient has been treated with somatropin for more than 12 months; and
  - 2.2 The patient has not had a deterioration in Quality of Life defined as a 6 point or greater increase from their lowest QoL-AGHDA® score on treatment (other than due to obvious external factors such as external stressors); and
  - 2.3 Serum IGF-I levels have continued to be maintained within ±1SD of the mean of the normal range for age and sex (other than for obvious external factors); and
  - 2.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients or 1 mg per day for female patients; or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$ Pe	er	Manufacturer

continued...

- 3 All of the following:
  - 3.1 The patient has had a Special Authority approval for somatropin for childhood deficiency in children and no longer meets the renewal criteria under this indication; and
  - 3.2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
  - 3.3 The patient has severe growth hormone deficiency (see notes); and
  - 3.4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
  - 3.5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

GnRH	Ana	logues
------	-----	--------

GOSERELIN			
Implant 3.6 mg, syringe	66.48	1	✓ Zoladex
Implant 10.8 mg, syringe	138.23	1	✓ Zoladex
LEUPRORELIN			
Additional subsidy by endorsement where the patient is a c goserelin and the prescription is endorsed accordingly.	hild or adolescent a	nd is unable	e to tolerate administration of
Inj 3.75 mg prefilled dual chamber syringe - Higher subsid	y of		
\$221.60 per 1 inj with Endorsement	66.48	1	
	(221.60)		Lucrin Depot 1-month
Inj 11.25 mg prefilled dual chamber syringe - Higher subsi	dy		
of \$591.68 per 1 inj with Endorsement	177.50	1	
	(591.68)		Lucrin Depot 3-month

### **Vasopressin Agonists**

DESMOPRESSIN			
Wafer 120 mcg	47.00	30	Minirin Melt
DESMOPRESSIN ACETATE			
Tab 100 mcg	25.00	30	✓ Minirin
Tab 200 mcg	54.45	30	✓ Minirin
▲ Nasal spray 10 mcg per dose		6 ml OP	✓ <u>Desmopressin-</u> <u>PH&amp;T</u>
Ini 4 mcg ner ml. 1 ml	67 18	10	✓ Minirin

93

_			
	Subsidy	Fully	Brand or
	(Manufacturer's Price)	Subsidised	Generic
	\$	Per 🗸	Manufacturer

### **Other Endocrine Agents**

#### **CABERGOLINE**

### ⇒SA2070 Special Authority for Waiver of Rule

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Hyperprolactinemia; or
- 2 Acromegaly\*: or
- 3 Inhibition of lactation.

Renewal — (for patients who have previously been funded under Special Authority form SA1031) from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has previously held a valid Special Authority which has expired and the treatment remains appropriate and the patient is benefiting from treatment.

Note: Indication marked with \* is an unapproved indication.

### **CLOMIFENE CITRATE**

Tab 50 mg	29.84	10	✓ Mylan Clomiphen S29
METYRAPONE Cap 250 mg	558.00	50	✓ Metopirone

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer **Anthelmintics** ALBENDAZOLE - Special Authority see SA1318 below - Retail pharmacy 60 ✓ Eskazole S29 **⇒SA1318** Special Authority for Subsidy Initial application only from an infectious disease specialist or clinical microbiologist. Approvals valid for 6 months where the patient has hydatids. Renewal only from an infectious disease specialist or clinical microbiologist. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefitting from the treatment. MEBENDAZOLE - Only on a prescription 6 Vermox 15 ml (7.83)Vermox **PRAZIQUANTEL**  Biltricide **Antibacterials** a) For topical antibacterials, refer to DERMATOLOGICALS, page 67 b) For anti-infective eve preparations, refer to SENSORY ORGANS, page 267 Cephalosporins and Cephamycins

CEFACLOR MONOHYDRATE			
Cap 250 mg		100	✓ Ranbaxy-Cefactor
Grans for oral liq 125 mg per 5 ml - Wastage claimable	3.75	100 ml	✓ Ranbaxy-Cefactor
CEFALEXIN			
Cap 250 mg	3.85	20	Cephalexin ABM
Cap 500 mg		20	✓ Cephalexin ABM
Grans for oral liq 25 mg per ml - Wastage claimable	7.88	100 ml	✓ Flynn
Grans for oral liq 50 mg per ml - Wastage claimable	10.38	100 ml	✓ Flynn
	11.75		✓ Cefalexin Sandoz
CEFAZOLIN - Subsidy by endorsement			
Only if prescribed for dialysis or cellulitis in accordance with a	a Health NZ Hos	pital approved	protocol and the prescription is
endorsed accordingly.		P	p
Inj 500 mg vial	3.39	5	✓ Cefazolin-AFT
Inj 1 g vial		5	✓ Cefazolin-AFT
Inj 2 g vial	7.09	5	✓ Cefazolin-AFT
CEFTRIAXONE - Subsidy by endorsement			
a) Up to 10 inj available on a PSO			
b) Subsidised only if prescribed for a dialysis or cystic fibros	is patient, or the	treatment of g	onorrhoea, or the treatment of
pelvic inflammatory disease, or the treatment of suspecte	d meningococca	ıl disease, and	I the prescription or PSO is
endorsed accordingly.			
Inj 500 mg vial	0.79	1	✓ Ceftriaxone-AFT
Inj 1 g vial	3.59	5	✓ <u>Ceftriaxone-AFT</u>
CEFUROXIME AXETIL - Subsidy by endorsement			
Only if prescribed for prophylaxis of endocarditis and the pre-	scription is endo	rsed according	alv.
Tab 250 mg		20	✓ Ascend-
•			Cefuroxime S29

95

Subsidy		Fully	Brand or	
(Manufacturer's Price)	9	Subsidised	Generic	
\$	Per	✓	Manufacturer	

#### **Macrolides**

AZITHROMYCIN – Maximum of 5 days treatment per prescription; can be waived by Special Authority see SA1683 below A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised on Special Authority.

Tab 250 mg	8.19	30	✓ Apo-Azithromycin
Tab 500 mg - Up to 8 tab available on a PSO		2	✓ Zithromax
Grans for oral liq 200 mg per 5 ml (40 mg per ml) - Wastage			
claimable	16.97	15 ml	✓ Zithromax

#### **⇒SA1683** Special Authority for Waiver of Rule

Initial application — (bronchiolitis obliterans syndrome, cystic fibrosis and atypical Mycobacterium infections) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following:

- 1 Patient has received a lung transplant, stem cell transplant, or bone marrow transplant and requires treatment for bronchiolitis obliterans syndrome\*; or
- 2 Patient has received a lung transplant and requires prophylaxis for bronchiolitis obliterans syndrome\*; or
- 3 Patient has cystic fibrosis and has chronic infection with Pseudomonas aeruginosa or Pseudomonas-related gram negative organisms\*; or
- 4 Patient has an atypical Mycobacterium infection.

Note: Indications marked with \* are unapproved indications.

Initial application — (non-cystic fibrosis bronchiectasis\*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 For prophylaxis of exacerbations of non-cystic fibrosis bronchiectasis\*; and
- 2 Patient is aged 18 and under; and
- 3 Fither:
  - 3.1 Patient has had 3 or more exacerbations of their bronchiectasis, within a 12 month period; or
  - 3.2 Patient has had 3 acute admissions to hospital for treatment of infective respiratory exacerbations within a 12 month period.

Note: Indications marked with \* are unapproved indications.

Renewal — (non-cystic fibrosis bronchiectasis\*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has completed 12 months of azithromycin treatment for non-cystic fibrosis bronchiectasis; and
- 2 Following initial 12 months of treatment, the patient has not received any further azithromycin treatment for non-cystic fibrosis bronchiectasis for a further 12 months, unless considered clinically inappropriate to stop treatment; and
- 3 The patient will not receive more than a total of 24 months' azithromycin cumulative treatment (see note).

The patient must not have had more than 1 prior approval.

Note: No further renewals will be subsidised. A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised. Indications marked with \* are unapproved indications

CLARITHROMYCIN - Maximum of 500 mg per prescription; can be waived by Special Authority see SA1857 below

**⇒SA1857** Special Authority for Waiver of Rule

Initial application — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years for applications meeting the following criteria:

Either:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1 Atypical mycobacterial infection; or
  - 2 Mycobacterium tuberculosis infection where there is drug-resistance or intolerance to standard pharmaceutical agents.

**Initial application — (Helicobacter pylori eradication)** from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 For the eradication of helicobacter pylori in a patient unable to swallow tablets; and
- 2 For use only in combination with omeprazole and amoxicillin as part of a triple therapy regimen.

Initial application — (Prophylaxis of infective endocarditis) from any relevant practitioner. Approvals valid for 3 months where prophylaxis of infective endocarditis associated with surgical or dental procedures if amoxicillin is contra-indicated. Renewal — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

ERYTHROMYCIN (AS LACTOBIONATE) Inj 1 g vial10.0	00 1	✓ Erythrocin IV
ERYTHROMYCIN ETHYL SUCCINATE		
Tab 400 mg35.8	32 100	E-Mycin
a) Up to 20 tab available on a PSO		
b) Up to 2 x the maximum PSO quantity for RFPP		
Grans for oral liq 200 mg per 5 ml	3 100 n	nl 🗸 E-Mycin
a) Up to 300 ml available on a PSO		
b) Up to 2 x the maximum PSO quantity for RFPP		
c) Wastage claimable		_
Grans for oral liq 400 mg per 5 ml9.4	l1 100 n	nl <b>✓ E-Mycin</b>
a) Up to 200 ml available on a PSO		
b) Wastage claimable		
ROXITHROMYCIN		
Tab 150 mg13.1	9 50	✓ Arrow-
		<u>Roxithromycin</u>
T 1 000		
Tab 300 mg25.0	00 50	✓ Arrow-
		<u>Roxithromycin</u>

	Subsidy (Manufacturer's Pr \$	rice) Subs Per	Fully idised	Brand or Generic Manufacturer
Penicillins				
AMOXICILLIN Cap 250 mg a) Up to 30 cap available on a PSO	27.50	500	•	Miro-Amoxicillin
b) Up to 10 x the maximum PSO quantity for RFPP Cap 500 mg      a) Up to 30 cap available on a PSO	41.00	500	1	Miro-Amoxicillin
b) Up to 10 x the maximum PSO quantity for RFPP Grans for oral liq 125 mg per 5 ml	2.22	100 ml	✓	Alphamox 125
Grans for oral liq 250 mg per 5 ml	2.81	100 ml	•	Alphamox 250
Inj 250 mg vial	15.97	10	1	Ibiamox
Inj 500 mg vial		10	1	lbiamox
Inj 1 g vial - Up to 5 inj available on a PSO		10	1	lbiamox
AMOXICILLIN WITH CLAVULANIC ACID  Tab 500 mg with clavulanic acid 125 mg — Up to 30 tab available on a PSO		10	•	Curam Duo 500/125
per ml		100 ml	./	Augmentin
<ul><li>a) Up to 200 ml available on a PSO</li><li>b) Wastage claimable</li></ul>		100 1111	•	Augmenum
Grans for oral liq amoxicillin 50 mg with clavulanic acid 12.5 per ml – Up to 200 ml available on a PSO		100 ml OP		Curam Amoxiclav Devatis Forte
(Curam Grans for oral liq amoxicillin 50 mg with clavulanic acid 1.	2.5 ma per ml to l	be delisted 1 J	lune 2	025)
BENZATHINE BENZYLPENICILLIN	37.			,
Inj 900 mg (1.2 million units) in 2.3 ml syringe – Up to 5 inj available on a PSO	375.97	10	•	Bicillin LA
BENZYLPENICILLIN SODIUM [PENICILLIN G] Inj 600 mg (1 million units) vial – Up to 5 inj available on a Po	SO 16.50	10	1	<u>Sandoz</u>
FLUCLOXACILLIN	45.70	050	,	Flore Leave - 1111 AFT
Cap 250 mg — Up to 30 cap available on a PSO		250		Flucloxacillin-AFT Flucloxacillin-AFT
Cap 500 mg – Up to 30 cap available on a PSO		500 100 ml		AFT
Grans for oral liq 25 mg per ml				
Grans for oral liq 50 mg per ml	5.89	100 ml		AFT
Inj 250 mg vial	42.60	10	1	Flucloxin
Inj 500 mg vial		10		Flucloxin
Inj 1 g vial – Up to 5 inj available on a PSO		5		Flucil

	Subsidy (Manufacturer's Price)		Fully		
	\$	Per	<b>√</b>	Manufacturer	
PHENOXYMETHYLPENICILLIN (PENICILLIN V)					
Cap 250 mg - Up to 30 cap available on a PSO	7.68	50	•	Cilicaine VK	
Cilicaine VK to be Principal Supply on 1 February 2025			_		
Cap 500 mg	13.72	50	/	Cilicaine VK	
a) Up to 20 cap available on a PSO					
<ul><li>b) Up to 2 x the maximum PSO quantity for RFPP</li><li>c) Cilicaine VK to be Principal Supply on 1 February 202</li></ul>	)E				
Grans for oral lig 125 mg per 5 ml		100 ml	/	AFT	
a) Up to 200 ml available on a PSO		100 1111		<u>Al 1</u>	
b) Wastage claimable					
Grans for oral liq 250 mg per 5 ml	4.24	100 ml	1	<u>AFT</u>	
a) Up to 300 ml available on a PSO					
b) Up to 2 x the maximum PSO quantity for RFPP					
c) Wastage claimable					

### **Tetracyclines**

DO:	XYCYCLINE			
*	Tab 100 mg - Up to 30 tab available on a PSO	64.43	500	Doxine
MIN	IOCYCLINE HYDROCHLORIDE			
*	Tab 50 mg - Additional subsidy by Special Authority see			
	SA1355 below – Retail pharmacy	5.79	60	
		(12.05)		Mino-tabs
*	Cap 100 mg	19.32	100	
		(52.04)		Minomycin

### **⇒SA1355** Special Authority for Manufacturers Price

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has rosacea.

TETRACYCLINE - Special Authority see SA1332 below - Retail pharmacy
Tab 250 mg .......58.20 28 ✓ Accord S29

### ⇒SA1332 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 For the eradication of helicobacter pylori following unsuccessful treatment with appropriate first-line therapy; and
- 2 For use only in combination with bismuth as part of a quadruple therapy regimen.

### **Other Antibiotics**

For topical antibiotics, refer to DERMATOLOGICALS, page 67

#### **CIPROFLOXACIN**

Recommended for patients with any of the following:

- i) microbiologically confirmed and clinically significant pseudomonas infection; or
- ii) prostatitis; or
- iii) pyelonephritis; or
- iv) gonorrhoea.

Tab 250 mg - Up to 5 tab available on a PSO	1.95	28	✓ Ipca-Ciprofloxacin
Tab 500 mg - Up to 5 tab available on a PSO	3.10	28	✓ Ipca-Ciprofloxacin
Tab 750 mg	4.80	28	✓ Ipca-Ciprofloxacin

	Subsidy		Fully Brand or	
	(Manufacturer's Price)		Subsidised Generic	
	\$	Per	✓ Manufacti	urer
CLINDAMYCIN				
Cap hydrochloride 150 mg		24	✓ Dalacin C	
Inj 150 mg per ml, 4 ml ampoule	35.10	10	✓ <u>Hameln</u>	
COLISTIN SULPHOMETHATE - Retail pharmacy-Specialist -	Subsidy by endorseme	nt		
Only if prescribed for dialysis or cystic fibrosis patient and the			accordingly.	
Inj 150 mg	65.00	1	✓ Colistin-Lir	nk
GENTAMICIN SULPHATE				
Inj 40 mg per ml, 2 ml vial - Subsidy by endorsement	36.70	5	✓ Cidomycin	
			P/Free S2	9
Only if prescribed for a dialysis or cystic fibrosis patient endorsed accordingly.	or complicated urinary	trac	t infection and the pres	cription is
Inj 10 mg per ml, 1 ml ampoule - Subsidy by endorsement		5	DBL Genta	
Only if prescribed for a dialysis or cystic fibrosis patient endorsed accordingly.	or complicated urinary	trac	t infection and the pres	cription is
Inj 10 mg per ml, 2 ml ampoule - Subsidy by endorsement	91.00	5	✓ Wockhardt	S29
Only if prescribed for a dialysis or cystic fibrosis patient endorsed accordingly.	or complicated urinary	trac	t infection and the pres	cription is
Inj 40 mg per ml, 2 ml ampoule - Subsidy by endorsement	18.38	10	✓ Pfizer	
	91.90	50	<ul><li>Gentamicir</li></ul>	1
			Noridem	S29
Only if prescribed for a dialysis or cystic fibrosis patient endorsed accordingly.	or complicated urinary	trac	t infection and the pres	cription is
MOXIFLOXACIN - Special Authority see SA1740 below - Reta	ail pharmacy			
No patient co-payment payable	. ,			
Tab 400 mg	42.00	5	✓ Avelox	

### ⇒SA1740 Special Authority for Subsidy

Initial application — (Tuberculosis) only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Both:
  - 1.1 Active tuberculosis\*: and
  - 1.2 Any of the following:
    - 1.2.1 Documented resistance to one or more first-line medications; or
    - 1.2.2 Suspected resistance to one or more first-line medications (tuberculosis assumed to be contracted in an area with known resistance), as part of regimen containing other second-line agents; or
    - 1.2.3 Impaired visual acuity (considered to preclude ethambutol use); or
    - 1.2.4 Significant pre-existing liver disease or hepatotoxicity from tuberculosis medications; or
    - 1.2.5 Significant documented intolerance and/or side effects following a reasonable trial of first-line medications; or
- 2 Mycobacterium avium-intracellulare complex not responding to other therapy or where such therapy is contraindicated.\*; or
- 3 Patient is under five years of age and has had close contact with a confirmed multi-drug resistant tuberculosis case.

Note: Indications marked with \* are unapproved indications.

**Renewal** only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Mycoplasma genitalium) only from a sexual health specialist or Practitioner on the recommendation of a sexual health specialist. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Su	bsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1 Has nucleic acid amplification test (NAAT) confirmed Mycoplasma genitalium\* and is symptomatic; and
- 2 Either
  - 2.1 Has tried and failed to clear infection using azithromycin; or
  - 2.2 Has laboratory confirmed azithromycin resistance; and
- 3 Treatment is only for 7 days.

Initial application — (Penetrating eye injury) only from an ophthalmologist. Approvals valid for 1 month where the patient requires prophylaxis following a penetrating eye injury and treatment is for 5 days only.

Note: Indications marked with \* are unapproved indications.

PAROMOMYCIN - Special Authority see SA1689 below - Retail pharmacy

### ⇒SA1689 Special Authority for Subsidy

Initial application only from an infectious disease specialist, clinical microbiologist or gastroenterologist. Approvals valid for 1 month for applications meeting the following criteria:

### Either:

- 1 Patient has confirmed cryptosporidium infection; or
- 2 For the eradication of Entamoeba histolyica carriage.

**Renewal** only from an infectious disease specialist, clinical microbiologist or gastroenterologist. Approvals valid for 1 month for applications meeting the following criteria:

### Either:

- 1 Patient has confirmed cryptosporidium infection; or
- 2 For the eradication of Entamoeba histolyica carriage.

PYRIMETHAMINE - Special Authority see SA1328 below - Retail pharmacy

### ⇒SA1328 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 For the treatment of toxoplasmosis in patients with HIV for a period of 3 months; or
- 2 For pregnant patients for the term of the pregnancy; or
- 3 For infants with congenital toxoplasmosis until 12 months of age.

### SODIUM FUSIDATE [FUSIDIC ACID]

Tab 250 mg	135.70	36	✓ Fucidin
SULFADIAZINE SODIUM - Special Authority see SA13	331 below – Retail pharmacy		
Tab 500 mg	150.70	100	✓ Sulfadiazin-Heyl  S29
	543.20	56	✓ Wockhardt S29

#### ⇒SA1331 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 For the treatment of toxoplasmosis in patients with HIV for a period of 3 months; or
- 2 For pregnant patients for the term of the pregnancy; or
- 3 For infants with congenital toxoplasmosis until 12 months of age.

	Subsidy (Manufacturer's Prio \$	ce) Subs Per	Fully idised	Brand or Generic Manufacturer
TOBRAMYCIN				
Inj 40 mg per ml, 2 ml vial - Subsidy by endorsement Only if prescribed for dialysis or cystic fibrosis patient ar		5 is endorsed a		<b>obramycin (Viatris)</b> gly.
Solution for inhalation 60 mg per ml, 5 ml — Subsidy by endorsement		56 dose		obramycin BNM
TRIMETHOPRIM  * Tab 300 mg – Up to 30 tab available on a PSO TMP to be Principal Supply on 1 February 2025	27.83	50	<b>✓</b> TI	МР
TRIMETHOPRIM WITH SULPHAMETHOXAZOLE [CO-TRIMOX  * Tab trimethoprim 80 mg and sulphamethoxazole 400 mg — to 30 tab available on a PSO  Trisul to be Principal Supply on 1 February 2025  * Oral liq 8 mg sulphamethoxazole 40 mg per ml — Up to 200 available on a PSO	Up 115.74 ml	500 100 ml	✓ Tr	isul eprim
VANCOMYCIN – Subsidy by endorsement Only if prescribed for a dialysis or cystic fibrosis patient or fo difficile following metronidazole failure and the prescription is Inj 500 mg vial	s endorsed accordi			ment of Clostridium ylan
Antifungals				

- a) For topical antifungals refer to DERMATOLOGICALS, page 68
- b) For topical antifungals refer to GENITO URINARY, page 81

### **FLUCONAZOLE**

Cap 50 mg4.10	28	✓ Mylan
Cap 150 mg	1	✓ Mylan
Cap 200 mg8.90	28	✓ Mylan
Powder for oral suspension 10 mg per ml - Special Authority		
see SA1359 below – Retail pharmacy129.02	35 ml	<ul><li>Diflucan</li></ul>
Wastage claimable		

### **⇒SA1359** Special Authority for Subsidy

**Initial application — (Systemic candidiasis)** from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

#### Both:

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

**Initial application — (Immunocompromised)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Patient is at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

**Renewal — (Systemic candidiasis)** from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

✓ Burel \$29

✓ Strides Shasun S29

30 100

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
\$	Per	✓	Manufacturer

#### continued...

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

**Renewal — (Immunocompromised)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

#### All of the following:

- 1 Patient remains immunocompromised; and
- 2 Patient remains at moderate to high risk of invasive fungal infection; and

Tab 200 mg - PCT......CBS

3 Patient is unable to swallow capsules.

#### ITRACONAZOLE

Cap 100 mg6.83	15	✓ Itrazole
Oral liq 10 mg per ml - Special Authority see SA1322 below -		
Retail pharmacy141.80	150 ml OP	✓ Itraconazole
		Kent S29
		✓ Sporanox

#### ⇒SA1322 Special Authority for Subsidy

**Initial application** only from an infectious disease specialist, clinical microbiologist, clinical immunologist or any relevant practitioner on the recommendation of a infectious disease physician, clinical microbiologist or clinical immunologist. Approvals valid for 6 months where the patient has a congenital immune deficiency.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefitting from the treatment.

#### **KETOCONAZOLE**

		100	✓ Taro \$29 ✓ Teva- Ketoconazole \$29
NYSTATIN			
Tab 500,000 u	14.16	50	
	(17.09)		Nilstat
Cap 500,000 u	12.81	50	
	(15.47)		Nilstat
POSACONAZOLE - Special Authority see SA2383 below - Reta	il pharmacy		
Tab modified-release 100 mg	206.00	24	✓ Posaconazole Juno
Oral liq 40 mg per ml	342.51	105 ml OP	✓ Devatis

#### **⇒SA2383** Special Authority for Subsidy

**Initial application** only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

#### Either:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation chemotherapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppressive therapy\*.

**Renewal** only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

Subsidy		Fully	Brand or
(Manufacturer's Price)	S	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

Fither:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation therapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppression\* and requires on going posaconazole treatment.

Note: \* Graft versus host disease (GVHD) on significant immunosuppression is defined as acute GVHD, grade II to IV, or extensive chronic GVHD, or if they were being treated with intensive immunosuppressive therapy consisting of either high-dose corticosteroids (1 mg or greater per kilogram of body weight per day for patients with acute GVHD or 0.8 mg or greater per kilogram every other day for patients with chronic GVHD), antithymocyte globulin, or a combination of two or more immunosuppressive agents or types of treatment.

Initial application — (Invasive fungal infection prophylaxis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient is at risk of invasive fungal infection; and
- 2 Either:
  - 2.1 Posaconazole is prescribed by, or recommended by a haematologist, transplant physician, infectious disease specialist, paediatric haematologist or paediatric oncologist; or
  - 2.2 Prescribing posaconazole is in accordance with a protocol or guideline that has been endorsed by the Health New Zealand - Te Whatu Ora Hospital in the specific settings where there is a greater than 10% risk of invasive fungal infection (IFI).

Renewal — (Invasive fungal infection prophylaxis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient is at risk of invasive fungal infection; and
- 2 Either:
  - 2.1 Posaconazole is prescribed by, or recommended by a haematologist, transplant physician, infectious disease specialist, paediatric haematologist or paediatric oncologist; or
  - 2.2 Prescribing posaconazole is in accordance with a protocol or guideline that has been endorsed by the Health New Zealand - Te Whatu Ora Hospital in the specific settings where there is a greater than 10% risk of invasive fungal infection (IFI).

### **TERBINAFINE**

* Tab 250 mg8.97	84	✓ Deolate
VORICONAZOLE - Special Authority see SA2384 below - Retail pharmacy		
Tab 50 mg91.00	56	✓ Vttack
Tab 200 mg350.00	56	✓ Vttack
Powder for oral suspension 40 mg per ml - Wastage		
claimable1,523.22	70 ml	✓ Vfend

### ⇒SA2384 Special Authority for Subsidy

Initial application — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
  - 3.1 Patient has proven or probable invasive aspergillus infection; or
  - 3.2 Patient has possible invasive aspergillus infection; or

Subsidy (Manufacturer's Price)	Sub	Fully	Brand or Generic	
\$	Per	✓	Manufacturer	

continued...

- 3.3 Patient has fluconazole resistant candidiasis; or
- 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

**Renewal — (invasive fungal infection)** only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
  - 3.1 Patient continues to require treatment for proven or probable invasive aspergillus infection; or
  - 3.2 Patient continues to require treatment for possible invasive aspergillus infection; or
  - 3.3 Patient has fluconazole resistant candidiasis: or
  - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Initial application — (Invasive fungal infection prophylaxis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient is at risk of invasive fungal infection; and
- 2 Either:
  - 2.1 Voriconazole is prescribed by, or recommended by a haematologist, transplant physician, infectious disease specialist, paediatric haematologist or paediatric oncologist; or
  - 2.2 Prescribing voriconazole is in accordance with a protocol or guideline that has been endorsed by the Health New Zealand - Te Whatu Ora Hospital in the specific settings where there is a greater than 10% risk of invasive fungal infection (IFI).

Renewal — (Invasive fungal infection prophylaxis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient is at risk of invasive fungal infection; and
- 2 Either:
  - 2.1 Voriconazole is prescribed by, or recommended by a haematologist, transplant physician, infectious disease specialist, paediatric haematologist or paediatric oncologist; or
  - 2.2 Prescribing voriconazole is in accordance with a protocol or guideline that has been endorsed by the Health New Zealand - Te Whatu Ora Hospital in the specific settings where there is a greater than 10% risk of invasive fungal infection (IFI).

### **Antimalarials**

#### SA1684 Special Authority for Subsidy

**Initial application** only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 The patient has vivax or ovale malaria; and
- 2 Primaguine is to be given for a maximum of 21 days.

**Renewal** only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 The patient has relapsed vivax or ovale malaria; and
- 2 Primaguine is to be given for a maximum of 21 days.

	Subsidy		Fully Bra	nd or
	(Manufacturer's P	rice) Si Per		neric nufacturer
	\$	Per	₹ IVIa	nuiacturer
Antitrichomonal Agents				
METRONIDAZOLE				
Tab 200 mg – Up to 30 tab available on a PSO	25.86	250	✓ Metro	nidamed
J J	33.15		✓ Metro	gyl
Metronidamed to be Principal Supply on 1 March 2025				•
Tab 400 mg - Up to 15 tab available on a PSO		21	✓ Metro	
	5.23		✓ Metro	gyl
Metronidamed to be Principal Supply on 1 March 2025	05.00	4001	<b>4</b> Element	
Oral liq benzoate 200 mg per 5 ml		100 ml	✓ Flagy	
Suppos 500 mg	24.48	10	✓ Flagy	
(Metrogyl Tab 200 mg to be delisted 1 March 2025) (Metrogyl Tab 400 mg to be delisted 1 March 2025)				
ORNIDAZOLE				
Tab 500 mg	36 52	10	✓ Arrow	-Ornidazole
Arrow-Ornidazole to be Principal Supply on 1 March 202		10	V Allow	-Offiidazoie
7 mon of mazzolo to 50 minospar cappiy on 1 maron 202				
Antituberculotics and Antileprotics				
·				
Note: There is no co-payment charge for all pharmaceuticals list	ted in the Antitube	erculotics ar	nd Antileprotics	group regardless of
immigration status.				
BEDAQUILINE – Special Authority see SA2244 below – Retail p	harmacy			
No patient co-payment payable	0.004.51	24 OP	✓ Sirtur	_
Tab 100mg	3,064.51	24 UP	♥ Sirtur	U
⇒SA2244 Special Authority for Subsidy		-1:1: A.		C
Initial application — (multi-drug resistant tuberculosis) from applications meeting the following criteria:	any relevant prac	cutioner. A	pprovais valid i	or 6 months for
applications meeting the following chieffa: Both:				
The person has multi-drug resistant tuberculosis (MDR-TI	B)· and			
Ministry of Health's Tuberculosis Clinical Network has rev		ual case an	d recommends	bedaquiline as part
of the treatment regimen.	icwca tric iriaivia	uai oaoo aii	a recommends	bodaquiii io do part
•				
CLOFAZIMINE – Retail pharmacy-Specialist				
<ul><li>a) No patient co-payment payable</li><li>b) Prescriptions must be written by, or on the recommendat</li></ul>	ion of an infaction	ue diegaeg	nhveician clinic	cal microhiologiet or
dermatologist.	ion oi, an imedio	us uiscasc	priyolciari, ciiriic	cal microbiologist of
* Cap 50 mg	442.00	100	✓ Lamp	rene S29
CYCLOSERINE – Retail pharmacy-Specialist		100	- <b>-</b> amp	.0
a) No patient co-payment payable				
b) Prescriptions must be written by, or on the recommendat	ion of an infection	us disease i	nhysician clinic	eal microhiologist or
respiratory physician.	ion oi, an inicolo	us discase	priyololari, olirik	al microbiologist of
Cap 250 mg	344.00	60	✓ Cyclo	rin S29
DAPSONE - Retail pharmacy-Specialist			-,	
a) No patient co-payment payable				
b) Prescriptions must be written by, or on the recommendat	ion of an infection	us dispass	nhysician clinic	cal microhiologist or
dermatologist	ion oi, an inicctio	us discase	priyololari, olirik	al microbiologist of
Tab 25 mg	268.50	100	✓ Dapso	one
Tab 100 mg		100	✓ Dapso	
ů				

•	NFECTIONS - A	GLI	131011	OTOTEINIO GGE
	Subsidy		Fully	Brand or
	(Manufacturer's Price)	Per	Subsidised <	Generic Manufacturer
ETHAMBUTOL HYDROCHLORIDE - Retail pharmacy-Speciali	st			
<ul> <li>a) No patient co-payment payable</li> <li>b) Prescriptions must be written by, or on the recommendar respiratory physician</li> </ul>	tion of, an infectious	disease	physician	, clinical microbiologist or
Tab 100 mg	85.73	100	<b>✓</b> E	EMB Fatol S29
Tab 400 mg	49.34	56	<b>✓</b> I	Myambutol S29
ISONIAZID - Retail pharmacy-Specialist				
<ul> <li>a) No patient co-payment payable</li> <li>b) Prescriptions must be written by, or on the recommendal microbiologist, dermatologist or public health physician</li> </ul>	tion of, an internal me	edicine	physician,	paediatrician, clinical
* Tab 100 mg	23.00	100	<b>√</b> F	PSM
	94.50			soniazid Teva S29
(DOM T. ). 400	327.41		<b>✓</b> 1	Noumed Isoniazid
(PSM Tab 100 mg to be delisted 1 May 2025)				
ISONIAZID WITH RIFAMPICIN – Retail pharmacy-Specialist				
<ul> <li>a) No patient co-payment payable</li> <li>b) Prescriptions must be written by, or on the recommendar microbiologist, dermatologist or public health physician</li> </ul>				
* Tab 100 mg with rifampicin 150 mg Rifinah to be Principal Supply on 1 February 2025		100	<b>√</b> F	Rifinah
* Tab 150 mg with rifampicin 300 mg Rifinah to be Principal Supply on 1 February 2025	179.13	100	<b>✓</b> F	Rifinah
LINEZOLID - Special Authority see SA2234 below - Retail pha	rmacy			
No patient co-payment payable				_
Tab 600 mg		10 150 ml		<u>Zyvox</u>
Oral liq 20 mg per ml	1,079.00	130 1111	• 2	Zyvox
■ SA2234 Special Authority for Subsidy Initial application — (multi-drug resistant tuberculosis) from applications meeting the following criteria: Both:  1 The person has multi-drug resistant tuberculosis (MDR-T 2 Ministry of Health's Tuberculosis Clinical Network has revenue to regime.	B); and			
the treatment regimen.				
PARA-AMINO SALICYLIC ACID – Retail pharmacy-Specialist				
<ul> <li>a) No patient co-payment payable</li> <li>b) Prescriptions must be written by, or on the recommendar respiratory physician</li> </ul>	tion of, an infectious	disease	specialist,	, clinical microbiologist or
Grans for oral liq 4 g sachet	280.00	30	<b>✓</b> F	Paser S29
PROTIONAMIDE - Retail pharmacy-Specialist				
<ul> <li>a) No patient co-payment payable</li> <li>b) Prescriptions must be written by, or on the recommendar respiratory physician</li> </ul>		disease	specialist,	, clinical microbiologist or
Tab 250 mg	305.00	100	<b>✓</b> F	Peteha S29
PYRAZINAMIDE - Retail pharmacy-Specialist				
<ul> <li>a) No patient co-payment payable</li> <li>b) Prescriptions must be written by, or on the recommendar respiratory physician</li> </ul>	tion of, an infectious	disease	physician	, clinical microbiologist or
* Tab 500 mg	64.95	100	<b>✓</b>	AFT-Pyrazinamide
•				•

INFECTIONS - AGENTS FOR SYSTEMIC USE						
		Subsidy (Manufacturer's Pric	ce) Si	Fully ubsidised	Brand or Generic Manufacturer	
RIF	ABUTIN – Retail pharmacy-Specialist a) No patient co-payment payable b) Processinting must be written by a real the recommendation	on of an infactious	o diogga	nhusisian r	acciratory physician or	
	b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, respiratory physician or gastroenterologist					
	Cap 150 mgAMPICIN – Subsidy by endorsement	353.71	30	✓ My	cobutin	
	<ul> <li>a) No patient co-payment payable</li> <li>b) For confirmed recurrent Staphylococcus aureus infection in combination with other effective anti-staphylococcal antimicrobial based on susceptibilities and the prescription is endorsed accordingly; can be waived by endorsement - Retail pharmacy - Specialist. Specialist must be an internal medicine physician, clinical microbiologist, dermatologist, paediatrician, or public health physician.</li> </ul>					
	Cap 150 mg		100 100	✓ <u>Ri</u> ✓ Ri		
	, ,			✓ Ri	fadin Sanofi	
*	Oral liq 100 mg per 5 ml	12.60	60 ml	✓ <u>Ri</u>	<u>fadin</u>	
Aı	ntivirals					
For	eye preparations refer to Eye Preparations, Anti-Infective Pre	parations, page 26	67			
Н	epatitis B Treatment					
	ECAVIR	10.04	00		stanavin (Dav)	
	Tab 0.5 mg  IIVUDINE – Special Authority see SA1685 below – Retail pha		30	▼ <u>E</u>	tecavir (Rex)	
LAIN	Tab 100 mg		28	<b>✓</b> <u>Ze</u>	tlam	
_	Oral liq 5 mg per ml	270.00	240 ml OF	✓ Ze	ffix	
<b>Initi</b> App	A1685 Special Authority for Subsidy al application only from a relevant specialist or medical pract rovals valid for 1 year where used for the treatment or preven ewal from any relevant practitioner. Approvals valid for 2 year	tion of hepatitis B.			·	
TEN	IOFOVIR DISOPROXIL Tenofovir disoproxil prescribed under endorsement for the tre		ncluded in	the count of	of up to 4 subsidised	
æ.	antiretrovirals for the purposes of Special Authority SA2139., Tab 245 mg (300 mg as a maleate)		30	<b>√</b> To	nofovir Disoproxil	
*	Tab 245 mg (500 mg as a maleate)	13.00	30		<u>Viatris</u>	
Н	erpesvirus Treatments					
ACI	CLOVIR					
	Tab dispersible 200 mg		25	✓ <u>Lo</u>		
	Tab dispersible 400 mg Tab dispersible 800 mg		56 35	✓ <u>Lo</u> ✓ <u>Lo</u>		
	ACICLOVIR		00	· <u>Lo</u>	<u>, , , , , , , , , , , , , , , , , , , </u>	
• / 12	Tab 500 mg	9.64	30	✓ Va	clovir	
	Vaclovir to be Principal Supply on 1 February 2025 Tab 1,000 mg	17 70	20	√ Vo	clovir	
	Vaclovir to be Principal Supply on 1 February 2025	11./0	30	▼ va	ICIOVII	
VAL	GANCICLOVIR - Special Authority see SA1993 on the next		rmacy			
	Tab 450 mg		60		ılganciclovir Viatris	

Valganciclovir Viatris to be Principal Supply on 1 February 2025

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

## **⇒SA1993** Special Authority for Subsidy

**Initial application** — **(transplant cytomegalovirus prophylaxis)** only from a relevant specialist. Approvals valid for 3 months where the patient has undergone a solid organ transplant and requires valganciclovir for CMV prophylaxis.

**Renewal — (transplant cytomegalovirus prophylaxis)** only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

#### Either:

- 1 Both:
  - 1.1 Patient has undergone a solid organ transplant and received anti-thymocyte globulin and requires valganciclovir therapy for CMV prophylaxis; and
  - 1.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following anti-thymocyte globulin; or
- 2 Both:
  - 2.1 Patient has received pulse methylprednisolone for acute rejection and requires further valganciclovir therapy for CMV prophylaxis; and
  - 2.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following pulse methylprednisolone.

Initial application — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has undergone a solid organ transplant and received valganciclovir under Special Authority more than 2 years ago (27 months); and
- 2 Patient has received anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Renewal — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months where the patient has received a further course of anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Initial application — (Lung transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone a lung transplant; and
- 2 Either:
  - 2.1 The donor was cytomegalovirus positive and the patient is cytomegalovirus negative; or
  - 2.2 The recipient is cytomegalovirus positive; and
- 3 Patient has a high risk of CMV disease.

Initial application — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

#### Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
  - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
  - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
  - 2.3 Patient has cytomegalovirus retinitis.

**Renewal — (Cytomegalovirus in immunocompromised patients)** only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

#### Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
  - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
  - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
  - 2.3 Patient has cytomegalovirus retinitis.

Note: for the purpose of this Special Authority "immunocompromised" includes transplant recipients, patients with immunosuppressive diseases (e.g. HIV) or those receiving immunosuppressive treatment for other conditions.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic S Per ✔ Manufacturer

# **Hepatitis C Treatment**

GLECAPREVIR WITH PIBRENTASVIR - [Xpharm]

Note the supply of treatment is via Pharmac's approved direct distribution supply. Further details can be found on Pharmac's website https://pharmac.govt.nz/maviret

Tab 100 mg with pibrentasvir 40 mg .......24,750.00 84 OP ✓ Maviret

LEDIPASVIR WITH SOFOSBUVIR - [Xpharm] - Special Authority see SA1605 below

No patient co-payment payable

Tab 90 mg with sofosbuvir 400 mg......24,363.46 28 **✓ Harvoni** 

#### ⇒SA1605 Special Authority for Subsidy

Special Authority approved by the Hepatitis C Treatment Panel (HepCTP)

Notes: By application to the Hepatitis C Treatment Panel (HepCTP).

Applications will be considered by HepCTP and approved subject to confirmation of eligibility.

Application details may be obtained from Pharmac's website <a href="http://www.pharmac.govt.nz/maviret">http://www.pharmac.govt.nz/maviret</a> or:

The Coordinator, Hepatitis C Treatment Panel

Pharmac, PO Box 10-254, WELLINGTON Tel: (04) 460 4990.

Email: hepcpanel@pharmac.govt.nz

# **HIV Prophylaxis and Treatment**

EMTRICITABINE WITH TENOFOVIR DISOPROXIL - Subsidy by endorsement; can be waived by Special Authority see SA2138 below

- a) Funding for emtricitabine with tenofovir disporoxil for use as PrEP, should be applied using Special Authority SA2138.
- b) Endorsement for treatment of conditions approved via Special Authority SA2139 (antiretrovirals for confirmed HIV, prevention of maternal transmission, post-exposure prophylaxis following exposure to HIV and percutaneous exposure): Prescription is deemed to be endorsed if emtricitabine with tenofovir disoproxil is co-prescribed with another antiretroviral subsidised under Special Authority SA2139 and the prescription is annotated accordingly by the Pharmacist or endorsed by the prescriber.

Note: Emtricitabine with tenofovir disoproxil prescribed under endorsement, for treatment of conditions approved via Special Authority SA2139 (antiretrovirals for confirmed HIV, prevention of maternal transmission, post-exposure prophylaxis following exposure to HIV and percutaneous exposure), is included in the count of up to 4 subsidised antiretrovirals, and counts as two antiretroviral medications, for the purposes of Special Authority SA2139, page 111 There is an approval process to become a named specialist to prescribe antiretroviral therapy in New Zealand. Further information is available on the Pharmac website.

* Tab 200 mg with tenofovir disoproxil 245 mg (300 mg	as a
-------------------------------------------------------	------

## ⇒SA2138 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 Patient has tested HIV negative, does not have signs or symptoms of acute HIV infection and has been assessed for HIV seroconversion; and
- 2 The Practitioner considers the patient is at elevated risk of HIV exposure and use of PrEP is clinically appropriate.

Notes: Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines:

https://ashm.org.au/HIV/PrEP/

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per		Manutacturer

continued...

**Renewal** from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 Patient has tested HIV negative, does not have signs or symptoms of acute HIV infection and has been assessed for HIV seroconversion; and
- 2 The Practitioner considers the patient is at elevated risk of HIV exposure and use of PrEP is clinically appropriate.

Notes: Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines:

https://ashm.org.au/HIV/PrEP/

#### **COVID-19 Treatments**

MOLNUPIRAVIR - [Xpharm] - Subsidy by endorsement

- a) No patient co-payment payable
- b) Treatment is funded only if patient meets access criteria for oral antiviral COVID-19 treatments (as on <u>Pharmac's website</u>) and has been endorsed accordingly by the prescriber. The supply of treatment is via Pharmac's approved distribution process. Refer to the Pharmac website for more information about this and stock availability.

NIRMATRELVIR WITH RITONAVIR - [Xpharm] - Subsidy by endorsement

- a) No patient co-payment payable
- b) Treatment is funded only if patient meets access criteria for oral antiviral COVID-19 treatments (as on <u>Pharmac's website</u>) and has been endorsed accordingly by the prescriber. The supply of treatment is via Pharmac's approved distribution process. Refer to the Pharmac website for more information about this and stock availability.

#### **Antiretrovirals**

#### ⇒SA2139 Special Authority for Subsidy

Initial application — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the patient has confirmed HIV infection.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Prevention of maternal transmission) only from a named specialist. Approvals valid for 1 year for applications meeting the following criteria:

Either:

- 1 Prevention of maternal foetal transmission: or
- 2 Treatment of the newborn for up to eight weeks.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Some antiretrovirals are unapproved or contraindicated for this indication. Practitioners prescribing these medications should exercise their own skill, judgement, expertise and discretion, and make their own prescribing decisions with respect to the use of a Pharmaceutical for an indication for which it is not approved or contraindicated.

Subsidy		Fully	Brand or	
(Manufacturer's Pr	ice) S	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Initial application — (post-exposure prophylaxis following exposure to HIV) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
  - 2.1 Patient has had condomless anal intercourse or receptive vaginal intercourse with a known HIV positive person with an unknown or detectable viral load greater than 200 copies per ml; or
  - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
  - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required; or
  - 2.4 Patient has had condomless anal intercourse with a person from a high HIV prevalence country or risk group whose HIV status is unknown.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines for PEP (https://www.ashm.org.au/hiv/hiv-management/pep/).

Renewal — (second or subsequent post-exposure prophylaxis) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
  - 2.1 Patient has had condomless anal intercourse or receptive vaginal intercourse with a known HIV positive person with an unknown or detectable viral load greater than 200 copies per ml; or
  - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
  - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required; or
  - 2.4 Patient has had condomless anal intercourse with a person from a high HIV prevalence country or risk group whose HIV status is unknown.

**Initial application — (Percutaneous exposure)** only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Second or subsequent percutaneous exposure) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

# Non-nucleosides Reverse Transcriptase Inhibitors

EFAVIRENZ – Special Authority see SA2139 on the previous page – Retail phart Tab 600 mg65.38	macy 30	✓ Efavirenz  Milpharm ©29
ETRAVIRINE - Special Authority see SA2139 on the previous page - Retail pha	rmacy	
Tab 200 mg770.00	60	✓ Intelence
NEVIRAPINE - Special Authority see SA2139 on the previous page - Retail pha	rmacy	
Tab 200 mg198.25	60	✓ Nevirapine Viatris
Nevirapine Viatris to be Principal Supply on 1 February 2025		
Oral suspension 10 mg per ml203.55	240 ml OP	✓ Viramune Suspension

	Subsidy (Manufacturer's Price) \$	Sub Per	Fully sidised	Brand or Generic Manufacturer
Nucleosides Reverse Transcriptase Inhibitors				
ABACAVIR SULPHATE – Special Authority see SA2139 on pag Tab 300 mg		acy 60	✓ Z	iagen
ABACAVIR SULPHATE WITH LAMIVUDINE – Special Authority Note: abacavir with lamivudine (combination tablets) counts anti-retroviral Special Authority.	see SA2139 on pag			
Tab 600 mg with lamivudine 300 mg	29.50	30	✓ <u>A</u>	<u>bacavir/</u> <u>Lamivudine</u> <u>Viatris</u>
EFAVIRENZ WITH EMTRICITABINE AND TENOFOVIR DISOPP pharmacy Note: Efavirenz with emtricitabine and tenofovir disoproxil co anti-retroviral Special Authority		-		
Tab 600 mg with emtricitabine 200 mg and tenofovir disoproz		30	<b>✓</b> V	iatris
EMTRICITABINE - Special Authority see SA2139 on page 111 - Cap 200 mg		30	<b>√</b> E	mtriva
LAMIVUDINE – Special Authority see SA2139 on page 111 – Re Tab 150 mgOral liq 10 mg per ml	98.00	60 0 ml OP	✓ <u>L</u>	amivudine Viatris TC
ZIDOVUDINE [AZT] – Special Authority see SA2139 on page 11 Cap 100 mg Oral liq 10 mg per ml	152.25	100 0 ml OP		etrovir etrovir
ZIDOVUDINE [AZT] WITH LAMIVUDINE – Special Authority see Note: zidovudine [AZT] with lamivudine (combination tablets the anti-retroviral Special Authority.	1 0			,
Tab 300 mg with lamivudine 150 mg	92.40	60	<b>√</b> L	amivudine/ Zidovudine Viatris
Protease Inhibitors				
ATAZANAVIR SULPHATE – Special Authority see SA2139 on p Cap 150 mg Cap 200 mg	85.00	rmacy 60 60		tazanavir Mylan tazanavir Viatris
DARUNAVIR - Special Authority see SA2139 on page 111 - Re Tab 400 mg Tab 600 mg	150.00	60 60		arunavir Viatris arunavir Viatris
LOPINAVIR WITH RITONAVIR – Special Authority see SA2139 Tab 100 mg with ritonavir 25 mg	on page 111 - Retail	l pharmad 60		opinavir/Ritonavir

Lopinavir/Ritonavir Mylan to be Principal Supply on 1 February 2025

RITONAVIR – Special Authority see SA2139 on page 111 – Retail pharmacy
Tab 100 mg .......43.31

Mylan

✓ Norvir

✓ Lopinavir/Ritonavir Mylan

120

30

_			
	Subsidy	Fully	Brand or
	(Manufacturer's Price)	Subsidised	Generic
	\$	Per 🗸	Manufacturer

#### Strand Transfer Inhibitors

DOLUTEGRAVIR – Special Authority see SA2139 on page	je 111 – Retail pharmacy	1	
Tab 50 mg	1,090.00	30	Tivicay
DOLUTEGRAVIR WITH LAMIVUDINE - Special Authority	y see SA2139 on page 1	11 – Retail p	harmacy
Tab 50 mg with lamivudine 300 mg	1,090.00	30	✓ Dovato
RALTEGRAVIR POTASSIUM - Special Authority see SA	2139 on page 111 – Reta	ail pharmacy	1
Tab 400 mg	1,090.00	60	✓ Isentress
Tah 600 mg	1 090 00	60	✓ Isentress HD

#### **Immune Modulators**

PEGYLATED INTERFERON ALFA-2A - Special Authority see SA2034 below - Retail pharmacy

Note: Pharmac will consider funding ribavirin for the small group of patients who have a clinical need for ribavirin and meet Special Authority criteria. Please contact the Hepatitis C Coordinator at Pharmac on 0800-023-588 option 4.

Inj 180 mcg prefilled syringe......748.50

**⇒SA2034** Special Authority for Subsidy

Initial application — (chronic hepatitis C - genotype 1, 4, 5 or 6 infection or co-infection with HIV or genotype 2 or 3 post liver transplant) from any specialist. Approvals valid for 18 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 Patient has chronic hepatitis C, genotype 1, 4, 5 or 6 infection; or
  - 1.2 Patient has chronic hepatitis C and is co-infected with HIV; or
  - 1.3 Patient has chronic hepatitis C genotype 2 or 3 and has received a liver transplant; and
- 2 Maximum of 48 weeks therapy.

Renewal — (Chronic hepatitis C - genotype 1 infection) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

- All of the following:
  - 1 Patient has chronic hepatitis C, genotype 1; and
  - 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
  - 3 Fither:
    - 3.1 Patient has responder relapsed; or
    - 3.2 Patient was a partial responder; and
  - 4 Patient is to be treated in combination with boceprevir; and
  - 5 Maximum of 48 weeks therapy.

Initial application — (Chronic Hepatitis C - genotype 1 infection treatment more than 4 years prior) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Any of the following:
  - 3.1 Patient has responder relapsed; or
  - 3.2 Patient was a partial responder; or
  - 3.3 Patient received interferon treatment prior to 2004; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sub	osidised	Generic
\$	Per	1	Manufacturer

continued...

Initial application — (chronic hepatitis C - genotype 2 or 3 infection without co-infection with HIV) from any specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has chronic hepatitis C, genotype 2 or 3 infection; and
- 2 Maximum of 6 months therapy.

Initial application — (Hepatitis B) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Hepatitis B infection (HBsAg positive for more than 6 months); and
- 2 Patient is Hepatitis B treatment-naive; and
- 3 ALT > 2 times Upper Limit of Normal; and
- 4 HBV DNA < 10 log10 IU/ml; and
- 5 Either:
  - 5.1 HBeAg positive; or
  - 5.2 serum HBV DNA greater than or equal to 2,000 units/ml and significant fibrosis (Metavir Stage F2 or greater or moderate fibrosis); and
- 6 Compensated liver disease: and
- 7 No continuing alcohol abuse or intravenous drug use; and
- 8 Not co-infected with HCV, HIV or HDV; and
- 9 Neither ALT nor AST > 10 times upper limit of normal; and
- 10 No history of hypersensitivity or contraindications to pegylated interferon; and
- 11 Maximum of 48 weeks therapy.

Initial application — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 Patient has a cutaneous T cell lymphoma\*; or
- 2 All of the following:
  - 2.1 Patient has a myeloproliferative disorder\*; and
  - 2.2 Patient is intolerant of hydroxyurea; and
  - 2.3 Treatment with an agrelide and busulfan is not clinically appropriate; or
- 3 Both:
  - 3.1 Patient has a myeloproliferative disorder; and
  - 3.2 Patient is pregnant, planning pregnancy or lactating.

Renewal — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment; and
- 3 Either:
  - 3.1 Patient has a cutaneous T cell lymphoma\*; or
  - 3.2 Both:
    - 3.2.1 Patient has a myeloproliferative disorder\*; and
    - 3.2.2 Either:
      - 3.2.2.1 Remains intolerant of hydroxyurea and treatment with anagrelide and busulfan remains clinically inappropriate; or
      - 3.2.2.2 Patient is pregnant, planning pregnancy or lactating.

Note: Indications marked with \* are unapproved indications.

	Subsidy	Fully	Brand or
(Manufa	acturer's Price)	Subsidised	Generic
	\$ Per	✓	Manufacturer

continued...

**Initial application** — **(post-allogenic bone marrow transplant)** from any relevant practitioner. Approvals valid for 3 months where patient has received an allogeneic bone marrow transplant\* and has evidence of disease relapse.

Renewal — (post-allogenic bone marrow transplant) from any relevant practitioner. Approvals valid for 3 months where patient is responding and ongoing treatment remains appropriate.

Note: Indications marked with \* are unapproved indications.

# **Urinary Tract Infections**

### ⇒SA2406 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 2 months for applications meeting the following criteria: Both:

- 1 Patient has an acute, symptomatic, bacteriologically-proven uncomplicated urinary tract infection (UTI)/cystitis with Escherichia Coli: and
- 2 Either:
  - 2.1 Microbiological testing confirms the pathogen is resistant to all of: trimethoprim, nitrofurantoin, amoxicillin, cefaclor, cefalexin, amoxicillin with clavulanic acid, and norfloxacin; or
  - 2.2 The patient has a contraindication or intolerance to all of: trimethoprim, nitrofurantoin, amoxicillin, cefaclor, cefalexin, amoxicillin with clavulanic acid, and norfloxacin that the pathogen is susceptible to.

Renewal from any relevant practitioner. Approvals valid for 2 months for applications meeting the following criteria: Both:

- 1 Patient has an acute, symptomatic, bacteriologically-proven uncomplicated urinary tract infection (UTI)/cystitis with Escherichia Coli; and
- 2 Either:
  - 2.1 Microbiological testing confirms the pathogen is resistant to all of: trimethoprim, nitrofurantoin, amoxicillin, cefaclor, cefalexin, amoxicillin with clavulanic acid, and norfloxacin; or
  - 2.2 The patient has a contraindication or intolerance to all of: trimethoprim, nitrofurantoin, amoxicillin, cefaclor, cefalexin, amoxicillin with clavulanic acid, and norfloxacin that the pathogen is susceptible to.

#### METHENAMINE (HEXAMINE) HIPPURATE

* Tab 1 g	19.95	100	✓ <u>Hiprex</u>
NITROFURANTOIN			
* Tab 50 mg - Up to 30 tab available on a PSO	22.20	100	✓ Nifuran
* Tab 100 mg	37.50	100	✓ Nifuran
* Cap modified-release 100 mg - Up to 15 cap available of PSO		100	✓ Macrobid
NORFLOXACIN			
Tab 400 mg — Subsidy by endorsement			✓ Arrow-Norfloxacin

Only if prescribed for a patient with an uncomplicated urinary tract infection that is unresponsive to a first line agent or with proven resistance to first line agents and the prescription is endorsed accordingly.

	Subsidy		Fully Brand or
	(Manufacturer's Price)		Subsidised Generic
	\$	Per	✓ Manufacturer
	Ψ	1 01	- Warialactarci
Anticholinesterases			
NEOCTIONINE METH CHILEATE			
NEOSTIGMINE METILSULFATE	40.05	40	<b>4.88</b> 11 101
Inj 2.5 mg per ml, 1 ml ampoule	48.25	10	✓ Max Health
Max Health to be Principal Supply on 1 February 2025			
PYRIDOSTIGMINE BROMIDE			
▲ Tab 60 mg	50.28	100	✓ Mestinon
_ Tub 00 mg		100	· modifien
Non-Steroidal Anti-Inflammatory Drugs			
DICLOFENAC SODIUM			
* Tab EC 25 mg	2.19	50	<ul><li>Diclofenac Sandoz</li></ul>
Diclofenac Sandoz to be Principal Supply on 1 February	2025		
* Tab 50 mg dispersible	1.50	20	✓ Voltaren D
* Tab EC 50 mg		50	✓ Diclofenac Sandoz
Diclofenac Sandoz to be Principal Supply on 1 February		00	5 Bioloicina Ganage
		100	/ Voltavan CD
* Tab long-acting 75 mg		100	✓ Voltaren SR
* Inj 25 mg per ml, 3 ml ampoule – Up to 5 inj available on a P		5	✓ Voltaren
* Suppos 12.5 mg		10	✓ Voltaren
* Suppos 25 mg	2.44	10	✓ Voltaren
* Suppos 50 mg - Up to 10 supp available on a PSO	4.22	10	✓ Voltaren
* Suppos 100 mg	7.00	10	✓ Voltaren
IBUPROFEN			4 - "
* Tab 200 mg		1,000	
* Tab long-acting 800 mg	3.05	30	✓ Brufen SR
	3.65		✓ Ibuprofen SR BNM
Ibuprofen SR BNM to be Principal Supply on 1 April 2029	5		•
* Oral lig 20 mg per ml		200 m	✓ Ethics
Ethics to be Principal Supply on 1 April 2025			
(Brufen SR Tab long-acting 800 mg to be delisted 1 April 2025)			
KETOPROFEN			
* Cap long-acting 200 mg	12.07	28	Oruvail SR
MEFENAMIC ACID			
* Cap 250 mg	1.05	50	
ж Сар 200 mg		50	Demotors
	(10.82)		Ponstan
	0.50	20	
	(7.50)		Ponstan
NAPROXEN			
* Tab 250 mg	20.22	500	✓ Noflam 250
•	39.23	300	• Nonain 250
Noflam 250 to be Principal Supply on 1 February 2025	04.45	050	4 N. W. 500
* Tab 500 mg	34.45	250	✓ Noflam 500
Noflam 500 to be Principal Supply on 1 February 2025			
* Tab long-acting 750 mg		28	Naprosyn SR 750
Naprosyn SR 750 to be Principal Supply on 1 February 2	2025		
* Tab long-acting 1 g		28	✓ Naprosyn SR 1000
Naprosyn SR 1000 to be Principal Supply on 1 February			. ,
	- <del></del>		
TENOXICAM			<b>4 -</b> 111 .11
* Tab 20 mg		100	✓ <u>Tilcotil</u>
* Inj 20 mg vial	9.95	1	✓ AFT

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	d Generic
NSAIDs Other				
ELECOXIB				
Cap 100 mg	3.45	60		Celebrex
Cap 200 mg	3 20	30		Celecoxib Pfizer Celebrex
Сцр 200 mg		00		Celecoxib Pfizer
Topical Products for Joint and Muscula	r Pain			
APSAICIN				
Crm 0.025% - Special Authority see SA1289 belo	ow – Retail			
pharmacy		15 g O		Zo-Rub Osteo S29
	13.00	60 g O		Zostrix Rugby Capsaicin
	13.00	50 g O		Topical
SA1289 Special Authority for Subsidy				Cream S29
itial application from any relevant practitioner. App steoarthritis that is not responsive to paracetamol and				
Antirheumatoid Agents				
YDROXYCHLOROQUINE SULPHATE	7.80	100		Inca-
-	7.80	100		lpca- Hydroxychloroquin
YDROXYCHLOROQUINE SULPHATE	7.80 8.78	100	<b>✓</b>	•
YDROXYCHLOROQUINE SULPHATE		100	<b>✓</b>	Hydroxychloroquin
YDROXYCHLOROQUINE SULPHATE  Tab 200 mg  Plaquenil Tab 200 mg to be delisted 1 May 2025)  EFLUNOMIDE	8.78		/	Hydroxychloroquin
YDROXYCHLOROQUINE SULPHATE  Tab 200 mg  Plaquenil Tab 200 mg to be delisted 1 May 2025)  EFLUNOMIDE  Tab 10 mg	8.78	30	<i>,</i>	Hydroxychloroquin Plaquenil Arava
YDROXYCHLOROQUINE SULPHATE  Tab 200 mg  Plaquenil Tab 200 mg to be delisted 1 May 2025)  EFLUNOMIDE  Tab 10 mg	8.78		<i>,</i>	Hydroxychloroquin
YDROXYCHLOROQUINE SULPHATE  Tab 200 mg  Plaquenil Tab 200 mg to be delisted 1 May 2025)  EFLUNOMIDE  Tab 10 mg  Tab 20 mg  ENICILLAMINE	8.78 6.00 6.00	30	,	Hydroxychloroquin Plaquenil Arava
YDROXYCHLOROQUINE SULPHATE  Tab 200 mg  Plaquenil Tab 200 mg to be delisted 1 May 2025)  EFLUNOMIDE  Tab 10 mg	8.78 6.00 6.00	30 30	<i>y</i>	Hydroxychloroquin Plaquenil  Arava Arava
YDROXYCHLOROQUINE SULPHATE  Tab 200 mg  Plaquenil Tab 200 mg to be delisted 1 May 2025)  EFLUNOMIDE  Tab 10 mg  Tab 20 mg  ENICILLAMINE  Tab 125 mg	8.78 6.00 6.00	30 30	<i>y</i>	Hydroxychloroquin Plaquenil  Arava Arava D-Penamine
YDROXYCHLOROQUINE SULPHATE  Tab 200 mg  Plaquenil Tab 200 mg to be delisted 1 May 2025)  EFLUNOMIDE  Tab 10 mg  Tab 20 mg  ENICILLAMINE  Tab 125 mg  Tab 250 mg	8.78 6.00 6.00	30 30	<i>y</i>	Hydroxychloroquin Plaquenil  Arava Arava D-Penamine
YDROXYCHLOROQUINE SULPHATE  Tab 200 mg	8.78 6.00 6.00	30 30	<i>y</i>	Hydroxychloroquin Plaquenil  Arava Arava D-Penamine
YDROXYCHLOROQUINE SULPHATE  Tab 200 mg	8.78 	30 30		Hydroxychloroquin Plaquenil  Arava Arava D-Penamine
YDROXYCHLOROQUINE SULPHATE  Tab 200 mg	8.78 	30 30 100 100		Hydroxychloroquin Plaquenil Arava Arava D-Penamine D-Penamine
YDROXYCHLOROQUINE SULPHATE  Tab 200 mg	8.78 	30 30 100 100		Hydroxychloroquin Plaquenil Arava Arava D-Penamine D-Penamine
YDROXYCHLOROQUINE SULPHATE  Tab 200 mg	8.78 	30 30 100 100		Hydroxychloroquin Plaquenil Arava Arava D-Penamine D-Penamine
Plaquenil Tab 200 mg to be delisted 1 May 2025)  EFLUNOMIDE Tab 10 mg Tab 20 mg Tab 20 mg Tab 25 mg Tab 27 mg  LENDRONATE SODIUM Tab 70 mg  LENDRONATE SODIUM WITH COLECALCIFEROL Tab 70 mg with colecalciferol 5,600 iu	8.78 	30 30 100 100		Hydroxychloroquir Plaquenil Arava Arava D-Penamine D-Penamine

Subsidy (Manufacturer's Price)

Fully Subsidised Per

Brand or Generic Manufacturer

#### ⇒SA1777 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 Fither:
  - 2.1 The patient is female and postmenopausal; or
  - 2.2 The patient is male or non-binary; and
- 3 Any of the following:
  - 3.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
  - 3.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons; or
  - 3.3 History of two significant osteoporotic fractures demonstrated radiologically: or
  - 3.4 Documented T-Score less than or equal to -3.0 (see Note); or
  - 3.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
  - 3.6 Patient has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 4 Zoledronic acid is contraindicated because the patient's creatinine clearance is less than 35 mL/min; and
- 5 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes); and
- 6 The patient must not receive concomitant treatment with any other funded antiresorptive agent for this condition or teriparatide.

#### Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with denosumab
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body
- e) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: risedronate sodium tab 35 mg once weekly; alendronate sodium tab 70 mg or tab 70 mg with cholecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy

### PAMIDRONATE DISODIUM

Inj 3 mg per ml, 10 ml vial	32.49	1	Pamisol
Inj 6 mg per ml, 10 ml vial	88.11	1	✓ Pamisol
Inj 9 mg per ml, 10 ml vial	94.34	1	✓ Pamisol

RALOXIFENE HYDROCHLORIDE - Special Authority see SA1779 on the next page - Retail pharmacy

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
· · · · · · ·	Por 🖋	Manufacturor

#### ⇒SA1779 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Notes); or
- 2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
- 3 History of two significant osteoporotic fractures demonstrated radiologically; or
- 4 Documented T-Score less than or equal to -3.0 (see Notes); or
- 5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Notes); or
- 6 Patient has had a Special Authority approval for zoledronic acid (Underlying cause Osteoporosis) or has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) prior to 1 February 2019.

#### Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
   Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for raloxifene funding.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis, and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

#### RISEDRONATE SODIUM

Tab 35 mg2.50	4	✓ Risedronate Sandoz
TERIPARATIDE - Special Authority see SA1139 below - Retail pharmacy		
Inj 250 mcg per ml, 2.4 ml	1	✓ Teriparatide - Teva

#### ⇒SA1139 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 18 months for applications meeting the following criteria: All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 The patient has a documented T-score less than or equal to -3.0 (see Notes); and
- 3 The patient has had two or more fractures due to minimal trauma; and
- 4 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes).

#### Notes:

- a) The bone mineral density (BMD) measurement used to derive the T-score must be made using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: alendronate sodium tab 70 mg or tab 70 mg with colecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily; zoledronic acid 5 mg per year. If an intolerance of a severity necessitating permanent treatment withdrawal develops

Subsidy	Fi	ully	Brand or
(Manufacturer's Price)	Subsidis	sed	Generic
<b>`</b> \$	Per	✓	Manufacturer

#### continued...

during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy.

- c) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.
- d) A maximum of 18 months of treatment (18 cartridges) will be subsidised.

#### **ZOLEDRONIC ACID**

Inj 0.05 mg per ml, 100 ml, bag22.53	100 ml OP	✓ Zoledronic Acid
		Viatric

# **Hyperuricaemia and Antigout**

ALI	LOPURINOL			
*	Tab 100 mg	17.99	1,000	✓ <u>Ipca-Allopurinol</u>
*	Tab 300 mg	22.50	500	✓ Ipca-Allopurinol
BE	NZBROMARONE – Special Authority see SA1963 below –	Retail pharmacy		
	Tab 50 mg	32.00	100	✓ Narcaricin mite S29
_	<del></del>			

#### ⇒SA1963 Special Authority for Subsidy

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefitting from the treatment; and
- 2 There is no evidence of liver toxicity and patient is continuing to receive regular (at least every three months) liver function tests

#### COLCHICINE

002002			
* Tab 500 mcg	6.00	100	✓ Colgout
FEBUXOSTAT - Special Authority see SA2054 below -			
Tab 80 mg	, ,	28	✓ Febuxostat (Teva)
Tab 120 mg		28	✓ Febuxostat (Teva)

#### SA2054 Special Authority for Subsidy

**Initial application — (Gout)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

#### Both:

- 1 Patient has been diagnosed with gout; and
- 2 Any of the following:
  - 2.1 The patient has a serum urate level greater than 0.36 mmol/l despite treatment with allopurinol at doses of at least 600 mg/day and addition of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
  - 2.2 The patient has experienced intolerable side effects from allopurinol such that treatment discontinuation is required and serum urate remains greater than 0.36 mmol/l despite use of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
  - 2.3 The patient has renal impairment such that probenecid is contraindicated or likely to be ineffective and serum urate remains greater than 0.36 mmol/l despite optimal treatment with allopurinol (see Note); or
  - 2.4 The patient has previously had an initial Special Authority approval for benzbromarone for treatment of gout...

Initial application — (Tumour lysis syndrome) only from a haematologist or oncologist. Approvals valid for 6 weeks for applications meeting the following criteria:

#### Both:

1 Patient is scheduled to receive cancer therapy carrying an intermediate or high risk of tumour lysis syndrome; and

	MUSCULOSKELETAL SYSTEM					
		Subsidy		Fully	Brand or	Т
		(Manufacturer's Price)		Subsidised	Generic	
		\$	Per	<u> </u>	Manufacturer	
cor	ntinued					
	2 Patient has a documented history of allopurinol intolerance	).				
	newal — (Gout) from any relevant practitioner. Approvals val ient is benefitting from treatment.	id for 2 years where	the tre	eatment ren	nains appropriate and the	е
•	newal — (Tumour lysis syndrome) only from a haematologis	st or oncologist. App	rovals	s valid for 6	weeks where the	
	atment remains appropriate and the patient is benefitting from t					
PR	OBENECID					
*	Tab 500 mg	66.95	100	<b>√</b> P	robenecid-AFT	
W	luscle Relaxants					
BA	CLOFEN					_
*	Tab 10 mg	3.70	100	<b>√</b> P	acifen	
	Inj 0.05 mg per ml, 1 ml ampoule - Subsidy by endorsement.		1	_	ioresal Intrathecal	
	Subsidised only for use in a programmable pump in patie		oastic	agents have	e been ineffective or hav	e
	caused intolerable side effects and the prescription is end			ŭ		

a) Subsidised only for use in a programmable pump in patients where oral antispastic agents have been ineffective or have caused intolerable side effects and the prescription is endorsed accordingly.

5

10

✓ Medsurge

✓ Sintetica Baclofen Intrathecal

b) Sintetica Baclofen Intrathecal to be Principal Supply on 1 March 2025

Inj 2 mg per ml, 5 ml ampoule - Subsidy by endorsement......306.82

(Medsurge Inj 2 mg per ml, 5 ml ampoule to be delisted 1 l	March 2025)		
DANTROLENE			
Cap 25 mg	112.13	100	✓ Dantrium
			✓ Dantrium S29 S29
Cap 50 mg	77.00	100	✓ Dantrium
(Dantrium Cap 25 mg to be delisted 1 April 2025)			
ORPHENADRINE CITRATE			
Tab 100 mg	23.25	100	✓ Norflex
Norflex to be Principal Supply on 1 February 2025			

Subsidy (Manufacturer's Price)

Fully Subsidised Per

Brand or Generic Manufacturer

# **Agents for Parkinsonism and Related Disorders**

<b>Dopamine</b>	<b>Agonists</b>	and	Related	Agents
-----------------	-----------------	-----	---------	--------

AMANTADINE HYDROCHLORIDE			
▲ Cap 100 mg	38.24	60	✓ Symmetrel
	63.73	100	✓ Symmetrel
APOMORPHINE HYDROCHLORIDE			
▲ Inj 10 mg per ml, 2 ml ampoule	59.50	5	✓ Movapo
▲ Inj 10 mg per ml, 5 ml ampoule		5	✓ Movapo
ENTACAPONE			•
▲ Tab 200 mg	18.04	100	✓ Comtan
LEVODOPA WITH BENSERAZIDE			
* Tab dispersible 50 mg with benserazide 12.5 mg	12.25	100	✓ Madopar Rapid
* Cap 50 mg with benserazide 12.5 mg		100	✓ Madopar 62.5
Cap 30 mg with benserazide 12.5 mg      Cap 100 mg with benserazide 25 mg		100	✓ Madopar 125
* Cap long-acting 100 mg with benserazide 25 mg		100	✓ Madopar HBS
* Cap 200 mg with benserazide 50 mg		100	✓ Madopar 250
LEVODOPA WITH CARBIDOPA		100	inaaopai 200
* Tab 100 mg with carbidopa 25 mg	26.40	100	✓ Sinemet
Sinemet to be Principal Supply on 1 February 2025	20.49	100	▼ Sillelliet
* Tab long-acting 200 mg with carbidopa 50 mg	44 99	100	✓ Sinemet CR
Sinemet CR to be Principal Supply on 1 February 2025		100	5 Gillemet Git
* Tab 250 mg with carbidopa 25 mg	39.49	100	✓ Sinemet
Sinemet to be Principal Supply on 1 February 2025			
PRAMIPEXOLE HYDROCHLORIDE			
▲ Tab 0.25 mg	5 51	100	✓ Ramipex
▲ Tab 1 mg		100	✓ Ramipex
<u> </u>		100	- <u>Hampox</u>
RASAGILINE	50.50	00	/ A=111 000
* Tab 1 mg	53.50	30	✓ Azilect S29
ROPINIROLE HYDROCHLORIDE			
▲ Tab 0.25 mg		84	Ropin
▲ Tab 1 mg		84	Ropin
▲ Tab 2 mg		84	Ropin
▲ Tab 5 mg	14.50	84	✓ Ropin
TOLCAPONE			
▲ Tab 100 mg	152.38	100	✓ Tasmar
A 1			
Anticholinergics			
BENZATROPINE MESYLATE			
Tab 2 mg		60	✓ Benztrop
Inj 1 mg per ml, 2 ml	95.00	5	✓ Phebra
a). Un to 10 ini available on a PSO			

a) Up to 10 inj available on a PSCb) Only on a PSO

b) Only on a 1 00

PROCYCLIDINE HYDROCHLORIDE

Tab 5 mg .......7.40 100 ✓ Kemadrin



Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

# Agents for Essential Tremor, Chorea and Related Disorders

#### ⇒SA1403 Special Authority for Subsidy

**Initial application** only from a neurologist or respiratory specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has amyotrophic lateral sclerosis with disease duration of 5 years or less; and
- 2 The patient has at least 60 percent of predicted forced vital capacity within 2 months prior to the initial application; and
- 3 The patient has not undergone a tracheostomy; and
- 4 The patient has not experienced respiratory failure; and
- 5 Any of the following:
  - 5.1 The patient is ambulatory; or
  - 5.2 The patient is able to use upper limbs; or
  - 5.3 The patient is able to swallow.

Renewal from any relevant practitioner. Approvals valid for 18 months for applications meeting the following criteria: All of the following:

- 1 The patient has not undergone a tracheostomy; and
- 2 The patient has not experienced respiratory failure; and
- 3 Any of the following:
  - 3.1 The patient is ambulatory; or
  - 3.2 The patient is able to use upper limbs; or
  - 3.3 The patient is able to swallow.

#### **TETRABENAZINE**

### Anaesthetics

#### Local

#### LIDOCAINE [LIGNOCAINE]

Gel 2%, tube − Subsidy by endorsement .......14.50 30 ml **✓ Xylocaine 2% Jelly** 

a) Up to 150 ml available on a PSO

b) Subsidised only if prescribed for urethral or cervical administration and the prescription is endorsed accordingly.

Gel 2%, 11 ml urethral syringe − Subsidy by endorsement...............59.50 10 ✓ Instillagel Lido

- a) Up to 5 each available on a PSO
- Subsidised only if prescribed for urethral, cervical or rectal administration and the prescription is endorsed accordingly.

cturer's Price) \$	Per	sidised •	Generic
			Manufacturer
4.00	200 ml	✓ N	<b>Aucosoothe</b>
5.00	25	<b>✓</b> L	idocaine-Baxter.
7.50	50		
5.00)		Х	(ylocaine
7.50 <sup>°</sup>	25	<b>✓</b> L	idocaine-Baxter
2.00	5		
0.00)		Х	(ylocaine
6.85	5	✓ L	idocaine-Baxter
7.15	5	<b>√</b> L	idocaine-Baxter
DC.	10	✓ X	(ylocard 500 S29
	0.00) 6.85 7.15 CBS	6.85 5 7.15 5 CBS 10	6.85 5 ✓ L 7.15 5 ✓ L

# **Topical Local Anaesthetics**

## **⇒SA0906** Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 2 years where the patient is a child with a chronic medical condition requiring frequent injections or venepuncture.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

LIDOCAINE [LIGNOCAINE] - Special Authority se	ee SA0906 above – Retail pl	narmacy	
Crm 4%	5.40	5 g OP	✓ LMX4
	27.00	30 g OP	✓ LMX4
LIDOCAINE [LIGNOCAINE] WITH PRILOCAINE -	- Special Authority see SAOS	906 above – Reta	il pharmacy
Crm 2.5% with prilocaine 2.5%	45.00	30 g OP	✓ EMLA
Crm 2.5% with prilocaine 2.5% (5 g tubes)	45.00	5	✓ EMLA

# **Analgesics**

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 117

# **Non-opioid Analgesics**

ASPIRIN			
* Tab dispersible 300 mg - Up to 30 tab available on a PSO	5.65	100	✓ Ethics Aspirin
CAPSAICIN – Subsidy by endorsement Subsidised only if prescribed for post-herpetic neuralgia or diabetic accordingly.	peripheral n	europathy ar	nd the prescription is endorsed
Crm 0.075%	11.95	45 g OP	✓ Zo-Rub HP S29 ✓ Zostrix HP
	15.14	57 g OP	✓ Rugby Capsaicin Topical Cream S29
NEFOPAM HYDROCHLORIDE Tab 30 mg	23 40	90	✓ Acunan

				Subsidy		Fully	Brand or
			(M	anufacturer's Price)	S Per	ubsidised	Generic Manufacturer
=				Ψ	rei		Ivialiulaciulei
PAH	RACET.	_		10.75	1 000		Dasimal
			- blister packimum of 300 tab per prescription; can be waived by		1,000	•	Pacimol Pacimol
			o 30 tab available on a PSO	endorsement			
	c)		0 30 tab available off a F30				
	,	1)	Subsidy by endorsement for higher quantities is avergular daily dosing for one month or greater, and annotate the prescription as endorsed where dispersumment of 100 tab per dispensing for non-endorsed patients), then dispense in repeat-bottle pack — Maximum of 300 tab per	the prescription is ensing history suppeed patients. If qu	annota ports a uantities	ted accor long-term prescrib	rdingly. Pharmacists may a condition. ed for more than 100 tabs
			otion; can be waived by endorsement	17 92	1,000	1	Noumed
	рі	escrip	nion, can be waived by endorsement	17.32	1,000	•	Paracetamol
		da pre 2) Ma	ubsidy by endorsement for higher quantities is availa aily dosing for one month or greater, and the prescrip escription as endorsed where dispensing history sup- aximum of 100 tab per dispensing for non-endorsed on-endorsed patients), then dispense in repeat dispe	otion is annotated oports a long-term patients. If quant	accordi conditi tities pre	ngly. Ph on. escribed t	ditions who require regular armacists may annotate the for more than 100 tabs (for
	Oral lic	120	mg per 5 ml	3.98	200 ml	✓	Paracetamol (Ethics)
	b)	Up to	imum of 600 ml per prescription; can be waived by 6 o 200 ml available on a PSO in combination	endorsement			(ETHOS)
	u)	•	Maximum of 200 ml per dispensing for non-endors non-endorsed patients), then dispense in repeat di Subsidy by endorsement for higher quantities is av regular daily dosing for one month or greater and the Pharmacists may annotate the prescription as endocondition.	spensing not excer allable for patient he prescription is	eeding 2 s with lo endorse	200 ml pe ong term ed or ann	er dispensing. conditions who require otated accordingly.
		3)	Note: 200 ml presentations of paracetamol oral liq Pharmacist) under the provisions in Part I of Section	, , , , ,	ied on E	SSO to a	Vaccinator (other than a
		4)	Note: Direct Provision by a pharmacist of up to 20	0 ml permitted un			
	Oral lic	2050	conjunction with immunisation of a child under 2 years per 5 ml		ieningo 200 ml		multicomponent vaccine.  Pamol
			imum of 600 ml per prescription; can be waived by e		200 1111	•	<u>ramoi</u>
	,		o 200 ml available on a PSO	Huorsement			
			in combination				
	,	1)	Maximum of 200 ml per dispensing for non-endors non-endorsed patients), then dispense in repeat di				
		•	Subsidy by endorsement for higher quantities is av regular daily dosing for one month or greater and t Pharmacists may annotate the prescription as end condition.	ailable for patient he prescription is orsed where dispe	s with lo endorse ensing h	ong term ed or ann nistory su	conditions who require otated accordingly. pports a long-term
		•	Note: 200 ml presentations of paracetamol oral liq Pharmacist) under the provisions in Part I of Section Note: Direct Provision by a pharmacist of up to 20	on A			•
*	Sunno		conjunction with immunisation of a child under 2 ye	ears of age with m	eningo	coccal B	multicomponent vaccine.

				IVL	HVOOS STSTEW
		Subsidy (Manufacturer's Price)	Per	Fully Subsidised	
			10	_	Gacet
木	Suppos 500 mg	10.00	50	•	Gacet
C	pioid Analgesics				
CC	DEINE PHOSPHATE - Safety medicine; prescriber may dete	ermine dispensing fre	quen	су	
	Tab 15 mg	5.92	100	✓	Noumed
	Tab 30 mg	6.98	100	✓	Aspen
	•			✓	Noumed
	Tab 60 mg	13.89	100	•	Noumed
DII	HYDROCODEINE TARTRATE				
	Tab long-acting 60 mg	8.60	60	✓	<b>DHC Continus</b>
FF	NTANYL				
٠-	a) Only on a controlled drug form				
	b) No patient co-payment payable				
		allonov			
	c) Safety medicine; prescriber may determine dispensing fre		10	./	Davishay and Muly
	Inj 50 mcg per ml, 2 ml ampoule		10		Boucher and Muir
	Inj 50 mcg per ml, 10 ml ampoule		10	_	Boucher and Muir
	Patch 12.5 mcg per hour		5		Fentanyl Sandoz
	Patch 25 mcg per hour		5		Fentanyl Sandoz
	Patch 50 mcg per hour		5		Fentanyl Sandoz
	Patch 75 mcg per hour		5		Fentanyl Sandoz
	Patch 100 mcg per hour	16.37	5	•	Fentanyl Sandoz
ME	THADONE HYDROCHLORIDE				
	a) Only on a controlled drug form				
	b) No patient co-payment payable				
	c) Safety medicine; prescriber may determine dispensing fre				
	Tab 5 mg	1.45	10	✓	Methadone BNM
	Oral liq 2 mg per ml	7.80	200 m	nl 🗸	Biodone
	Biodone to be Principal Supply on 1 February 2025				
	Oral liq 5 mg per ml		200 m	nl 🗸	Biodone Forte
	Biodone Forte to be Principal Supply on 1 February 2025	5			
	Oral liq 10 mg per ml	9.65	200 m	nl 🗸	Biodone Extra Forte
	Biodone Extra Forte to be Principal Supply on 1 February				
	Inj 10 mg per ml, 1 ml	68.90	10	•	AFT
MC	DRPHINE HYDROCHLORIDE				
	a) Only on a controlled drug form				
	b) No patient co-payment payable				
	c) Safety medicine; prescriber may determine dispensing fre	equency			
	Oral liq 1 mg per ml		200 m	nl 🗸	RA-Morph
	Oral liq 2 mg per ml		200 m		RA-Morph
	Oral lig 5 mg per ml		200 m		RA-Morph
	Oral lig 10 mg par ml		200 11		DA Morph

200 ml

✓ RA-Morph

Oral liq 10 mg per ml .......40.25

	Subsidy (Manufacturer's Price	١	Fully Subsidised	
	(Manufacturer's Price \$	) Per	Subsidised	Manufacturer
DRPHINE SULPHATE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing free			_	
Tab immediate-release 10 mg		10		Sevredol
Tab immediate-release 20 mg		10		Sevredol
Cap long-acting 10 mg		10		m-Eslon
Cap long acting 60 mg		10 10		m-Eslon m-Eslon
Cap long-acting 60 mg  Cap long-acting 100 mg		10		m-Esion
, , ,		100 m		Wockhardt S29
Oral liq 2 mg per ml	29.80	100 111		Oramorph
	29.00			Oramorph CDC
			•	S29 S29
Inj 5 mg per ml, 1 ml ampoule - Up to 5 inj available on a PS	O 539	5	1	Medsurge
Inj 10 mg per ml, 1 ml ampoule – Up to 5 inj available on a PS		5		Medsurge Medsurge
Inj 15 mg per ml, 1 ml ampoule – Up to 5 inj available on a PS		5		Medsurge Medsurge
Inj 30 mg per ml, 1 ml ampoule – Up to 5 inj available on a PS		5		Medsurge
(YCODONE HYDROCHLORIDE	00	Ŭ		ouour go
a) Only on a controlled drug form				
<ul><li>b) No patient co-payment payable</li><li>c) Safety medicine; prescriber may determine dispensing free</li></ul>	auonov			
Tab controlled-release 5 mg		20	1	Oxycodone Sando
Tab controlled follower of mg	3.77	28		Oxycodone Sando
	0	0		S29 S29
	4.04	30	1	OxyContin S29
Tab immediate-release 5 mg		100		Oxycodone Amnea
Tab controlled-release 10 mg		20		Oxycodone Sando
	3.77	28		Oxycodone Sando
				<b>S29</b> S29
Tab immediate-release 10 mg	18 77	100	1	Oxycodone Amne
Tab controlled-release 20 mg		20		Oxycodone Sando
Tab immediate-release 20 mg		100		Oxycodone Amne
Tab controlled-release 40 mg		20		Oxycodone Sando
Tab controlled-release 80 mg	12.99	20	1	Oxycodone Sando
Cap immediate-release 20 mg	5.23	20	1	OxyNorm
Oral liq 1 mg per ml	37.08	250 m	<b>✓</b>	Oxycodone Lucis
				S29 S29
Inj 10 mg per ml, 1 ml ampoule	4.37	5	1	Hameln
Inj 10 mg per ml, 2 ml ampoule	8.62	5	1	Hameln
Inj 50 mg per ml, 1 ml ampoule		5	1	Hameln
xyNorm Cap immediate-release 20 mg to be delisted 1 March 2	2025)			
RACETAMOL WITH CODEINE - Safety medicine; prescriber	may determine disp	ensino	frequenc	y
Tab paracetamol 500 mg with codeine phosphate 8 mg		1,000		Paracetamol +
				Codeine (Relieve

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	I Generic
PETHIDINE HYDROCHLORIDE	· · · · · · · · · · · · · · · · · · ·			
a) Only on a controlled drug form				
b) No patient co-payment payable				
<ul> <li>Safety medicine; prescriber may determine dispensing free</li> </ul>	equency			
Tab 50 mg		10		Noumed Pethidine
Inj 50 mg per ml, 1 ml ampoule – Up to 5 inj available on a F	'SO29.88	5	•	DBL Pethidine
lai 50 man namani O mil amanania. Illa ta 5 ini amailabla an a 5	000 00 70	_		Hydrochloride DBL Pethidine
Inj 50 mg per ml, 2 ml ampoule – Up to 5 inj available on a F	3030.72	5	•	Hydrochloride
TRAMADOL HYDROCHLORIDE				Trydrocilloride
Tab sustained-release 100 mg	1 05	20	1	Tramal SR 100
Tab sustained-release 150 mg		20		Tramal SR 150
Tab sustained-release 200 mg		20		Tramal SR 200
Cap 50 mg		100		Arrow-Tramadol
Antidepressants				
Cyclic and Related Agents				
AMITRIPTYLINE - Safety medicine; prescriber may determine d	ispensing frequency			
Tab 10 mg		100	✓	Arrow-Amitriptyline
Tab 25 mg	1.99	100		Arrow-Amitriptyline
Tab 50 mg	3.14	100	•	Arrow-Amitriptyline
CLOMIPRAMINE HYDROCHLORIDE - Safety medicine; prescri		ispen		
Tab 10 mg		30		Clomipramine Teva
Tab 25 mg		30		Clomipramine Teva
	39.97	100		Anafranil S29
Cap 10 mg		28		Clomipramine Teva
Cap 25 mg		28	•	Clomipramine Teva
DOSULEPIN [DOTHIEPIN] HYDROCHLORIDE – Subsidy by en				
Safety medicine; prescriber may determine dispensing freb Subsidy by endorsement – Subsidised for patients who we 2019 and the prescription is endorsed accordingly. Phare exists a record of prior dispensing of dosulepin [dothiepin]	ere taking dosulepin macists may annotate ] hydrochloride.	the p	orescriptio	n as endorsed where there
Tab 75 mg		30		Dosulepin Viatris
Cap 25 mg	7.03	50	•	Dosulepin Viatris S29
IMIPRAMINE HYDROCHLORIDE - Safety medicine; prescriber	may determine dispe	nsina	frequency	V
Tab 10 mg		50		Tofranil
-	10.96	100	/	Tofranil
Tab 25 mg	4.93	28	/	Imipramine
				Crescent S29
	8.80	50	•	Tofranil
NORTRIPTYLINE HYDROCHLORIDE - Safety medicine; prescr	•			
Tab 10 mg		100		Norpress
Tab 25 mg		180	/	Norpress
Monoamine-Oxidase Inhibitors (MAOIs) - Non S	elective			
TRANYLCYPROMINE SULPHATE				
* Tab 10 mg	22.94	50	•	Parnate

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

NERVOUS SYSTEM				
	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
Monoamine-Oxidase Type A Inhibitors				
MOCLOBEMIDE  * Tab 150 mg	23.60	60	J 1	urorix
Aurorix to be Principal Supply on 1 February 2025			_	
* Tab 300 mg	38.50	60	<b>✓</b>	urorix
Selective Serotonin Reuptake Inhibitors				
CITALOPRAM HYDROBROMIDE			_	
* Tab 20 mg	2.86	84	<b>√</b> <u>c</u>	<u>Celapram</u>
ESCITALOPRAM * Tab 10 mg	0.79	28	✓ li	oca-Escitalopram
	1.07		_	scitalopram
* Tab 20 mg	1.49	28	<b>√</b>  ı	(Ethics) oca-Escitalopram
FLUOXETINE HYDROCHLORIDE				<u> </u>
* Tab dispersible 20 mg, scored – Subsidy by endorsement Subsidised by endorsement		28		iluox
<ol> <li>When prescribed for a patient who cannot swallow accordingly; or</li> </ol>	whole tablets of caps	ules	ind the pre	scription is endorsed
<ol><li>When prescribed in a daily dose that is not a multipendorsed. Note: Tablets should be combined with</li></ol>	•			
* Cap 20 mg	3.13	90	<b>√</b> µ	rrow-Fluoxetine
PAROXETINE				
* Tab 20 mg	4.11	90	<b>✓</b> <u>L</u>	<u>oxamine</u>
SERTRALINE * Tab 50 mg	0.99	30	<b>√</b> 9	Setrona
* Tab 100 mg		30		Setrona
Other Antidepressants				
MIRTAZAPINE				
Tab 30 mg	2.60	28		loumed
Tob 45 mg	0.45	30 28		loumed loumed
Tab 45 mg	3.45	28 30		loumed
VENLAFAXINE				
Cap 37.5 mg	8.29	84	<b>✓</b> E	inlafax XR
Cap 75 mg		84		nlafax XR
Cap 150 mg	13.95	84	<b>✓</b> E	inlafax XR

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

# **Antiepilepsy Drugs**

Agents for	Control of	Status E	pilepticus
------------	------------	----------	------------

DIAZEPAM – Safety medicine; prescriber may determine dispensing frequency Inj 5 mg per ml, 2 ml ampoule – Subsidy by endorsement27.92	5	✓ Hospira
a) Up to 5 inj available on a PSO		-
b) Only on a PSO		
<ul> <li>c) PSO must be endorsed "not for anaesthetic procedures".</li> </ul>		
Rectal tubes 5 mg - Up to 5 tube available on a PSO54.58	5	✓ Stesolid
PHENYTOIN SODIUM		
* Inj 50 mg per ml, 2 ml ampoule - Up to 5 inj available on a		
PSO104.58	5	<ul><li>Hospira</li></ul>
* Inj 50 mg per ml, 5 ml ampoule - Up to 5 inj available on a		
PSO154.01	5	<ul><li>Hospira</li></ul>
		-

# **Control of Epilepsy**

**CARBAMAZEPINE** 

*	Tab 200 mg	14.53	100	<ul><li>✓ Tegretol</li><li>✓ Tegretol AU</li></ul>
*	Tab long-acting 200 mg	16.98	100	✓ Tegretol CR
		33.96	200	✓ Tegretol CR
*	Tab 400 mg	34.58	100	✓ Tegretol
*	Tab long-acting 400 mg	39.17	100	✓ Tegretol CR
*	Oral liq 20 mg per ml	26.37	250 ml	✓ Tegretol
CL	OBAZAM - Safety medicine; prescriber may determine dispe	ensing frequency		
	Tab 10 mg		50	✓ Frisium
CL	ONAZEPAM - Safety medicine; prescriber may determine di	spensing frequen	су	
	Oral drops 2.5 mg per ml	7.38	10 ml OP	✓ Rivotril
ΕT	HOSUXIMIDE			
	Cap 250 mg	78.89	56	✓ Essential  Ethosuximide \$29
		140.88	100	✓ Zarontin
	Oral liq 250 mg per 5 ml	56.35	200 ml	✓ Zarontin
GA	BAPENTIN			
	Note: Not subsidised in combination with subsidised pregal	oalin		
*	Cap 100 mg	6.45	100	✓ Nupentin
*	Cap 300 mg	8.45	100	✓ Nupentin
*	Cap 400 mg	10.26	100	✓ Nupentin
LA	COSAMIDE - Special Authority see SA2267 on the next page	e – Retail pharma	acy	
$\blacktriangle$	Tab 50 mg		14	✓ Vimpat
$\blacktriangle$	Tab 100 mg	50.06	14	✓ Vimpat
		200.24	56	✓ Vimpat
•	Tab 150 mg	75.10	14	✓ Vimpat
_	140 100 mg			

56

56

300.40

✓ Vimpat

✓ Vimpat

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sı	ubsidised	Generic
\$	Per	✓	Manufacturer

#### **⇒SA2267** Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Patient has focal epilepsy; and
- 2 Seizures are not adequately controlled by, or patient has experienced unacceptable side effects from, optimal treatment with all of the following: sodium valproate, topiramate, levetiracetam and any two of carbamazepine, lamotrigine and phenytoin sodium (see Note).

Note: Those of childbearing potential are not required to trial phenytoin sodium, sodium valproate, or topiramate. Those who can father children are not required to trial sodium valproate.

Renewal from any relevant practitioner. Approvals valid for 24 months where the patient has demonstrated a significant and sustained improvement in seizure rate or severity and/or quality of life compared with that prior to starting lacosamide treatment.

LAMOTRIGINE			
▲ Tab dispersible 2 mg	55.00	30	✓ Lamictal
▲ Tab dispersible 5 mg	50.00	30	✓ Lamictal
* Tab dispersible 25 mg	4.20	56	✓ Logem
* Tab dispersible 50 mg	5.11	56	✓ Logem
* Tab dispersible 100 mg	6.75	56	✓ Logem
LEVETIRACETAM			
Tab 250 mg	5.84	60	✓ Everet
Tab 500 mg		60	✓ Everet
Tab 750 mg	16.71	60	✓ Everet
Tab 1,000 mg	21.82	60	✓ Everet
Oral liq 100 mg per ml	44.78	300 ml OP	✓ Levetiracetam-AFT
Inj 100 mg per ml, 5 ml vial	38.95	10	✓ Levetiracetam-AFT
PHENOBARBITONE			
For phenobarbitone oral liquid refer Standard	Formulae, page 274		
Tab 15 mg		500	✓ Noumed
•			Phenobarbitone
Tab 30 mg	398.50	500	✓ Noumed
3			Phenobarbitone
PHENYTOIN SODIUM			
* Tab 50 mg	75.00	200	✓ Dilantin Infatab
Cap 30 mg		200	✓ Dilantin
Cap 100 mg		200	✓ Dilantin
* Oral liq 30 mg per 5 ml		500 ml	✓ Dilantin Paediatric
PREGABALIN			
Note: Not subsidised in combination with sub	osidised gabanentin		
* Cap 25 mg	• •	56	✓ Pregabalin Pfizer
- Cap 20 mg	7.80	00	✓ Milpharm \$29
* Cap 75 mg		56	✓ Pregabalin Pfizer
- Cup 70 mg	8.10	00	✓ Milpharm \$29
* Cap 150 mg	****	56	✓ Lyrica
ж Оар 130 mg	4.01	30	✓ Pregabalin Pfizer
* Cap 300 mg	7 38	56	✓ Pregabalin Pfizer
, ,	7.50	50	- i regulariti ilzer
PRIMIDONE	07.05	100	✓ Drimidono Clinest
* Tab 250 mg	37.35	100	Primidone Clinect

Subsidy		Fully	Brand or
(Manufacturer's Price	e) Su	bsidised	Generic
\$	Per	/	Manufacturer
13.65	100	✓ E	pilim Crushable
27.44	100	✓ E	pilim
52.24	100	✓ E	pilim
20.48	300 ml	✓ E	pilim S/F Liquid
		✓ E	pilim Syrup
41.50	1	<b>√</b> E	pilim IV
- Retail pharmacy			
509.29	60	✓ D	iacomit
509.29	60	✓ D	iacomit
	(Manufacturer's Price \$	(Manufacturer's Price) Su Per Su Per Su Su Per Su Su Per Su Su Per Su	(Manufacturer's Price) Subsidised Per ✓ 13.65 100 ✓ E27.44 100 ✓ E52.24 100 ✓ E20.48 300 ml ✓ E41.50 1 ✓ E - Retail pharmacy509.29 60 ✓ D

#### ⇒SA2268 Special Authority for Subsidy

**Initial application** only from a paediatric neurologist or Practitioner on the recommendation of a paediatric neurologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has confirmed diagnosis of Dravet syndrome; and
- 2 Seizures have been inadequately controlled by appropriate courses of sodium valproate, clobazam and at least two of the following: topiramate, levetiracetam, ketogenic diet.

Note: Those of childbearing potential are not required to trial sodium valproate or topiramate. Those who can father children are not required to trial sodium valproate.

**Renewal** from any relevant practitioner. Approvals valid without further renewal unless notified where the patient continues to benefit from treatment as measured by reduced seizure frequency from baseline.

#### TOPIRAMATE

▲ Tab 25 mg	11.07	60	✓ Arrow-Topiramate
Ÿ			✓ Topiramate Actavis
	26.04		✓ Topamax
▲ Tab 50 mg	18.81	60	✓ Arrow-Topiramate
·			✓ Topiramate Actavis
	44.26		✓ Topamax
▲ Tab 100 mg	31.99	60	✓ Arrow-Topiramate
· ·			✓ Topiramate Actavis
	75.25		✓ Topamax
▲ Tab 200 mg	55.19	60	✓ Arrow-Topiramate
•			✓ Topiramate Actavis
	129.85		✓ Topamax
▲ Sprinkle cap 15 mg	20.84	60	✓ Topamax
▲ Sprinkle cap 25 mg		60	✓ Topamax
VIGABATRIN - Special Authority see SA2088 below - Re	tail pharmacy		
▲ Tab 500 mg	'	100	✓ Sabril
Powder for oral soln 500 mg per sachet		60	✓ Sabril

#### **⇒SA2088** Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Any of the following:
  - 1.1 Patient has infantile spasms; or
  - 1.2 Both:
    - 1.2.1 Patient has epilepsy: and
    - 1.2.2 Either:



continued...

- 1.2.2.1 Seizures are not adequately controlled with optimal treatment with other antiepilepsy agents; or
- 1.2.2.2 Seizures are controlled adequately but the patient has experienced unacceptable side effects from optimal treatment with other antiepilepsy agents; or
- 1.3 Patient has tuberous sclerosis complex; and
- 2 Either:
  - 2.1 Patient is, or will be, receiving regular automated visual field testing (ideally before starting therapy and on a 6-monthly basis thereafter): or
  - 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields...

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient has demonstrated a significant and sustained improvement in seizure rate or severity and or quality of life; and
- 2 Fither
  - 2.1 Patient is receiving regular automated visual field testing (ideally every 6 months) on an ongoing basis for duration of treatment with vigabatrin; or
  - 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields...

# **Antimigraine Preparations**

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 117

#### **Acute Migraine Treatment**

melt
agran
agran
tran

### **Prophylaxis of Migraine**

For Beta Adrenoceptor Blockers refer to CARDIOVASCULAR SYSTEM, page 48

PIZOTIFFN

**★** Tab 500 mcg......23.21 100 **✓ Sandomigran** 

# **Antinausea and Vertigo Agents**

For Antispasmodics refer to ALIMENTARY TRACT, page 8

APREPITANT – Special Authority see SA0987 below – Retail pharmacy

Cap 2 × 80 mg and 1 × 125 mg.......21.90 3 OP ✓ <u>Emend Tri-Pack</u>

#### SA0987 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 12 months where the patient is undergoing highly emetogenic chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.

**Renewal** from any relevant practitioner. Approvals valid for 12 months where the patient is undergoing highly emetogenic chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.

BETAHISTINE DIHYDROCHLORIDE

				_
	Subsidy (Manufacturer's Price) \$	S Per	Fully Subsidised	Brand or Generic Manufacturer
CYCLIZINE HYDROCHLORIDE Tab 50 mg Nausicalm to be Principal Supply on 1 February 2025	0.66	10	<b>✓</b> N	ausicalm
CYCLIZINE LACTATE Inj 50 mg per ml, 1 ml ampoule – Up to 10 inj available on a PSO		10	<b>√</b> <u>H</u>	<u>ameln</u>
DOMPERIDONE  * Tab 10 mg	4.00	100	<b>✓</b> <u>D</u>	omperidone <u>Viatris</u>
HYOSCINE HYDROBROMIDE				
* Inj 400 mcg per ml, 1 ml ampoule	93.00	10	✓ M	lartindale S29
below – Retail pharmacy	88.50	10		copolamine - Mylan copolamine - Mylan S29 (29)

(Scopolamine - Mylan S29 S29 Patch 1 mg per 72 hours to be delisted 1 February 2025)

# **⇒SA1998** Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Control of intractable nausea, vomiting, or inability to swallow saliva in the treatment of malignancy or chronic disease where the patient cannot tolerate or does not adequately respond to oral anti-nausea agents; or
- 2 Control of clozapine-induced hypersalivation where trials of at least two other alternative treatments have proven ineffective.

Renewal from any relevant practitioner. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

METOCLOPRAMIDE HYDROCHLORIDE		
* Tab 10 mg - Up to 30 tab available on a PSO1.57	100	✓ <u>Metoclopramide</u> <u>Actavis 10</u>
* Inj 5 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO7.00	10	✓ Baxter
ONDANSETRON		
* Tab 4 mg2.27	50	✓ Periset
Tab disp 4 mg - Up to 10 tab available on a PSO	10	✓ Periset ODT
* Tab 8 mg4.10	50	✓ Periset
Tab disp 8 mg - Up to 10 tab available on a PSO0.90	10	✓ Periset ODT
PROCHLORPERAZINE		
* Tab 3 mg buccal5.97	50	
(30.00)		Buccastem
(30.00)		Max Health \$29
(30.00)		Prochlorperazine
		Brown & Burk S29
* Tab 5 mg - Up to 30 tab available on a PSO25.00	250	✓ Nausafix
* Inj 12.5 mg per ml, 1 ml - Up to 5 inj available on a PSO25.81	10	✓ Stemetil

Subsidy (Manufacturer's Price) \$ Fully Subsidised Per

Brand or Generic Manufacturer

# Antipsychotics

# General

AMISULPRIDE - Safety medicine; prescriber may determine disp	ensing frequen	су	
Tab 100 mg	5.84	30	✓ Sulprix
Tab 200 mg	14.47	60	✓ Sulprix
Tab 400 mg	35.06	60	✓ Sulprix
ARIPIPRAZOLE - Safety medicine; prescriber may determine dis		ncv	<del></del>
Tab 5 mg		30	✓ Aripiprazole Sandoz
1 ab 5 mg	10.00	00	✓ Ampiprazoic dandoz
T 1 40	40.50		Aripiprazole \$29
Tab 10 mg		30	✓ Aripiprazole Sandoz
Tab 15 mg		30	✓ Aripiprazole Sandoz
Tab 20 mg		30	✓ Aripiprazole Sandoz
Tab 30 mg	10.50	30	Aripiprazole Sandoz
CHLORPROMAZINE HYDROCHLORIDE - Safety medicine; pre	scriber may det	ermine dispen	sing frequency
Tab 25 mg - Up to 30 tab available on a PSO	15.62	100	✓ Largactil
Tab 100 mg - Up to 30 tab available on a PSO	36.73	100	✓ Largactil
Inj 25 mg per ml, 2 ml - Up to 5 inj available on a PSO	30.79	10	✓ Largactil
CLOZAPINE – Hospital pharmacy [HP4]			•
Safety medicine; prescriber may determine dispensing freque	nev		
Tab 25 mg	•	50	✓ Clopine
1 ab 25 mg	0.03	30	✓ Clozaril
	13.37	100	✓ Clopine
	13.37	100	✓ Clopine ✓ Clozaril
Tab 50 mg	9.67	50	✓ Clopine
Tab 50 filg	17.33	100	✓ Clopine
Tab 100 mg		50	✓ Clopine
Tab 100 Hig	17.33	50	✓ Clopine ✓ Clozaril
	34.65	100	✓ Clopine
	34.03	100	•
Tob 000 mg	04.65	50	✓ Clozaril
Tab 200 mg		50	✓ Clopine
Commencian FO man man mi	69.30	100	✓ Clopine
Suspension 50 mg per ml		100 ml	✓ Versacloz
HALOPERIDOL - Safety medicine; prescriber may determine dis		ncy	
Tab 500 mcg - Up to 30 tab available on a PSO	6.23	100	Serenace
Tab 1.5 mg - Up to 30 tab available on a PSO		100	Serenace
Tab 5 mg - Up to 30 tab available on a PSO	14.86	50	Serenace
	29.72	100	Serenace
Oral liq 2 mg per ml - Up to 200 ml available on a PSO	23.84	100 ml	<ul><li>Serenace</li></ul>
Inj 5 mg per ml, 1 ml ampoule - Up to 5 inj available on a PS	O21.55	10	✓ Serenace
LEVOMEPROMAZINE - Safety medicine; prescriber may determ	nine dispensing t	frequency	
Tab 25 mg (33.8 mg as a maleate)		100	✓ Nozinan (Swiss)
Tab 25 mg as a maleate		100	✓ Nozinan
Tab 100 mg (135 mg as a maleate)		100	✓ Nozinan (Swiss)
Tab 100 mg as a maleate		100	✓ Nozinan
-			
LEVOMEPROMAZINE HYDROCHLORIDE – Safety medicine; pr	•		
Inj 25 mg per ml, 1 ml ampoule	24.48	10	✓ Nozinan S29 S29
			✓ Wockhardt

	Subsidy (Manufactureria Price)		Fully	
	(Manufacturer's Price) \$	Per	Subsidised •	I Generic Manufacturer
LITHIUM CARBONATE - Safety medicine; prescriber may dete	rmine dispensing fred	wency		
Tab long-acting 400 mg		100		Priadel
Priadel to be Principal Supply on 1 February 2025				
Cap 250 mg	22.36	100	1	Douglas
DLANZAPINE - Safety medicine; prescriber may determine dis				3
Tab 2.5 mg		30	1	Zypine
Tab 5 mg		30		Zypine
Tab orodispersible 5 mg		28		Zypine ODT
Tab 10 mg		30		Zypine
Tab orodispersible 10 mg		28		Zypine ODT
PERICYAZINE - Safety medicine; prescriber may determine dis				=71
Tab 2.5 mg		100	1	Neulactil
Tab 10 mg		100		Neulactil
· ·		100	•	Neulactii
QUETIAPINE – Safety medicine; prescriber may determine disp	. ,	00	,	0
Tab 25 mg		90		Quetapel
Tab 100 mg		90		Quetapel
Tab 200 mg		90		Quetapel
Tab 300 mg		90	•	Quetapel
ISPERIDONE - Safety medicine; prescriber may determine di	spensing frequency			
Tab 0.5 mg	0.72	20		Risperdal
	2.17	60		Risperidone (Teva)
	4.01		•	Risperidone
				Sandoz S29
Tab 1 mg	2.44	60		Risperdal
			1	Risperidone (Teva)
	3.68		•	Risperidone
				Sandoz S29
Tab 2 mg	2.72	60	1	Risperdal
•			1	Risperidone (Teva)
	5.38		/	Risperidone
				Sandoz S29
Tab 3 mg	4.50	60	1	Risperdal
· • • • • • • • • • • • • • • • • • • •				Risperidone (Teva)
	8.57			Risperidone
				Sandoz S29
Tab 4 mg	6.25	60	/	Risperdal
1 ab + mg	0.20	00		Risperidone (Teva)
Oral liq 1 mg per ml	10.29	30 ml		Risperon
		JU 1111		
IPRASIDONE – Safety medicine; prescriber may determine dis	, , ,	60	.1	7uodono
Cap 40 mg		60 60		Zusdone Zusdone
Cap 40 mg Cap 60 mg		60		Zusdone
Cap 80 mg		60		Zusdone
' '				
CUCLOPENTHIXOL HYDROCHLORIDE – Safety medicine; pre	•		•	
Tab 10 mg	31.45	100	/	Clopixol



Subsid (Manufacturer		ılly Brand or ed Generic	
	Per	✓ Manufacturer	

# **Depot Injections**

ARIPIPRAZOLE – Special Authority see SA2395 below – Retail pharmacy Safety medicine; prescriber may determine dispensing frequency		
Inj 300 mg vial273.56	1	<ul> <li>✓ Abilify Maintena</li> <li>✓ Abilify Maintena</li> <li>S29 S29</li> </ul>
Inj 400 mg vial341.96	1	<ul> <li>✓ Abilify Maintena</li> <li>✓ Abilify Maintena</li> <li>S29 S29</li> </ul>

### **⇒SA2395** Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Either:

- 1 Either:
  - 1.1 The patient has had an initial Special Authority approval for risperidone depot injection, paliperidone depot injection or olanzapine depot injection; or
  - 1.2 All of the following:
    - 1.2.1 The patient has schizophrenia or other psychotic disorder; and
    - 1.2.2 The patient has received treatment with oral atypical antipsychotic agents but has been unable to adhere; and
    - 1.2.3 The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months; or
- 2 Patient has been unable to access olanzapine depot injection due to supply issues with olanzapine depot injection, or otherwise would have been started on olanzapine depot injection but has been unable to due to supply issues with olanzapine depot injection.

Notes: The Olanzapine depot injection Special Authority criteria that apply to criterion 2 in this Aripiprazole Special Authority application are as follows:

- The patient has had an initial Special Authority approval for paliperidone depot injection or risperidone depot injection; or
- All of the following:
  - The patient has schizophrenia; and
  - The patient has not been able to adhere with treatment using oral atypical antipsychotic agents; and
  - The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months.

FLUPENTHIXOL DECANOATE - Safety medicine; prescriber may	determine disp	ensing freq	uency
Inj 20 mg per ml, 1 ml - Up to 5 inj available on a PSO	13.14	5	✓ Fluanxol
Inj 20 mg per ml, 2 ml - Up to 5 inj available on a PSO	20.90	5	✓ Fluanxol
Inj 100 mg per ml, 1 ml - Up to 5 inj available on a PSO	40.87	5	✓ Fluanxol
HALOPERIDOL DECANOATE - Safety medicine; prescriber may	determine dispe	nsing frequ	ency
Inj 50 mg per ml, 1 ml - Up to 5 inj available on a PSO	28.39	5	✓ Haldol
Inj 100 mg per ml, 1 ml - Up to 5 inj available on a PSO	55.90	5	✓ Haldol Concentrate
			✓ Haldol
			Decanoas \$29
OLANZAPINE - Special Authority see SA2313 on the next page -	Retail pharmacy	y	
a) Safety medicine; prescriber may determine dispensing freq	uency		
b) Note – no new patients to be initiated on olanzapine.	•		
Inj 210 mg vial	252.00	1	✓ Zyprexa Relprevv
Inj 300 mg vial	414.00	1	✓ Zyprexa Relprevv
Ini 405 mg vial		1	✓ Zyprexa Relprevy

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per ✓	Manufacturer

#### ⇒SA2313 Special Authority for Subsidy

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of olanzapine depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

PALIPERIDONE - Special Authority see SA2396 below - Retail pharmacy

Safety medicine; prescriber may determine dispensing frequency

Inj 25 mg syringe	1	✓ Invega Sustenna
Inj 50 mg syringe271.95	1	✓ Invega Sustenna
Inj 75 mg syringe	1	✓ Invega Sustenna
Inj 100 mg syringe435.12	1	✓ Invega Sustenna
Inj 150 mg syringe435.12	1	✓ Invega Sustenna

#### ⇒SA2396 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for risperidone depot injection or olanzapine depot injection or aripiprazole depot injection; or
- 2 All of the following:
  - 2.1 The patient has schizophrenia or other psychotic disorder; and
  - 2.2 Has been unable to adhere to treatment using oral atypical antipsychotic agents; and
  - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

**Renewal** from any relevant practitioner. Approvals valid for 12 months where the initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

#### PALIPERIDONE PALMITATE - Special Authority see SA2167 below - Retail pharmacy

Inj 175 mg syringe	815.85	1 •	Invega Trinza
Inj 263 mg syringe1,		1 •	/ Invega Trinza
Inj 350 mg syringe1,	305.36	1 •	/ Invega Trinza
Inj 525 mg syringe1,	305.36	1 •	Invega Trinza

#### ⇒SA2167 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has schizophrenia; and
- 2 The patient has had an initial Special Authority approval for paliperidone once-monthly depot injection.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

RISPERIDONE - Special Authority see SA2397 below - Retail pharmacy

Safety medicine; prescriber may determine dispensing	g frequency		
Inj 25 mg vial	135.98	1	✓ Risperdal Consta
Inj 37.5 mg vial	178.71	1	✓ Risperdal Consta
Inj 50 mg vial	217.56	1	✓ Risperdal Consta
_ · _ ·			•

#### ⇒SA2397 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Fither:

1 The patient has had an initial Special Authority approval for paliperidone depot injection or olanzapine depot injection or aripiprazole depot injection; or



Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 2 All of the following:
  - 2.1 The patient has schizophrenia or other psychotic disorder; and
  - 2.2 Has not been able to adhere with treatment using oral atypical antipsychotic agents; and
  - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of risperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

ZUCLOPENTHIXOL DECANOATE - Safety medicine; prescriber may determine dispensing frequency

✓ Clopixol

# **Anxiolytics**

BUSPIRONE HYDROCHLORIDE			
* Tab 5 mg	13.95	100	✓ Buspirone Viatris
* Tab 10 mg	12.50	100	<ul> <li>Buspirone Viatris</li> </ul>
CLONAZEPAM - Safety medicine; prescriber may determine d	ispensing frequency	1	
Tab 500 mcg	5.64	100	✓ Paxam
Tab 2 mg	10.78	100	✓ Paxam
DIAZEPAM - Safety medicine; prescriber may determine dispe	nsing frequency		
Tab 2 mg	95.00	500	✓ Arrow-Diazepam
Tab 5 mg		500	✓ Arrow-Diazepam
LORAZEPAM - Safety medicine; prescriber may determine dis	pensing frequency		
Tab 1 mg	10.20	250	✓ Ativan
Ativan to be Principal Supply on 1 February 2025			
Tab 2.5 mg	13.13	100	✓ Ativan
Ativan to be Principal Supply on 1 February 2025			

# **Multiple Sclerosis Treatments**

### ⇒SA2274 Special Authority for Subsidy

Initial application — (Multiple Sclerosis - dimethyl fumarate, fingolimod, glatiramer acetate, interferon beta-1-alpha, interferon beta-1-beta, natalizumab and teriflunomide) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 All of the following:
  - 1.1 Diagnosis of multiple sclerosis (MS) meets the McDonald 2017 diagnostic criteria for MS and has been confirmed by a neurologist; and
  - 1.2 Patient has an EDSS score between 0 6.0; and
  - 1.3 Patient has had at least one significant attack of MS in the previous 12 months or two significant attacks in the past 24 months: and
  - 1.4 All of the following:
    - 1.4.1 Each significant attack must be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the attack, but the neurologist/physician must be satisfied that the clinical features were characteristic); and
    - 1.4.2 Each significant attack is associated with characteristic new symptom(s)/sign(s) or substantially worsening

			NER	VOUS SYSTEM
Subs (Manufactur \$	er's Price)	Sul Per	Fully bsidised	Brand or Generic Manufacturer
continued				
of previously experienced symptoms(s)/sign(s); and 1.4.3 Each significant attack has lasted at least one week an previous attack (where relevant); and 1.4.4 Each significant attack can be distinguished from the e				
fever (T> 37.5°C); and 1.4.5 Either:				
1.4.5.1 Each significant attack is severe enough to chan Functional System scores by at least 1 point; or	•			
1.4.5.2 Each significant attack is a recurrent paroxysma seizures/spasms, trigeminal neuralgia, Lhermitte			ple sclero	osis (tonic
<ul><li>1.5 Evidence of new inflammatory activity on an MRI scan within t</li><li>1.6 Any of the following:</li></ul>	he past 24	months	s; and	
1.6.1 A sign of that new inflammatory activity on MRI scanning enhancing lesion; or	ng (in criter	rion 5 im	nmediatel	y above) is a gadolinium
<ul><li>1.6.2 A sign of that new inflammatory activity is a lesion show</li><li>1.6.3 A sign of that new inflammatory is a T2 lesion with asso</li></ul>	ociated loca	al swelli	ing; or	
1.6.4 A sign of that new inflammatory activity is a prominent features of a recent attack that occurred within the last	2 years; or	r		
1.6.5 A sign of that new inflammatory activity is new T2 lesio				is MRI scan; or
2 Patient has an active approval for ocrelizumab and does not have prin Note: Treatment on two or more funded multiple sclerosis treatments simultance.	,, ,			
Renewal — (Multiple Sclerosis - dimethyl fumarate, fingolimod, glatiram				a-1-alpha. interferon
beta-1-beta, natalizumab and teriflunomide) from any relevant practitione had an EDSS score of 0 to 6.0 (inclusive) with or without the use of unilatera the patient has walked 100 metres or more with or without aids in the last six Note: Treatment on two or more funded multiple sclerosis treatments simultations.	er. Approva I or bilatera months).	als valid al aids a	for 12 m It any time	onths where patient has
DIMETHYL FUMARATE - Special Authority see SA2274 on the previous pa	ı <mark>ge</mark> – Retai	il pharm	асу	
a) Wastage claimable				
b) Note: Treatment on two or more funded multiple sclerosis treatment Cap 120 mg520.0		ously is		nitted. <b>ecfidera</b>
Cap 240 mg2,000.0		56		ecfidera
FINGOLIMOD - Special Authority see SA2274 on the previous page - Reta	il pharmac	у		
a) Wastage claimable     b) Note: Treatment on two or more funded multiple sclerosis treatment Cap 0.5 mg2,200.0		eously is		nitted. <b>ilenya</b>
GLATIRAMER ACETATE – Special Authority see SA2274 on the previous properties.  Note: Treatment on two or more funded multiple sclerosis treatments singled may be prefilled syringe	multaneous		ot permitte	ed. opaxone
INTERFERON BETA-1-ALPHA — Special Authority see SA2274 on the prev			_	
Note: Treatment on two or more funded multiple sclerosis treatments si				
Inj 6 million iu prefilled syringe	0	4	` <b>✓</b> A	vonex vonex Pen

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

INTERFERON BETA-1-BETA - Special Authority see SA2274 on the previous page - Retail pharmacy Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

NATALIZUMAB - Special Authority see SA2274 on the previous page - Retail pharmacy

✓ Betaferon

✓ Tysabri

15



Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer

TERIFLUNOMIDE - Special Authority see SA2274 on page 140 - Retail pharmacy

a) Wastage claimable

b) Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

✓ Teriflunomide Sandoz

659.90

✓ Aubagio

Teriflunomide Sandoz to be Principal Supply on 1 April 2025 (Aubagio Tab 14 mg to be delisted 1 April 2025)

# **Multiple Sclerosis Treatments - Other**

OCRELIZUMAB - Special Authority see SA2273 below - Retail pharmacy

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

✓ Ocrevus

⇒SA2273 Special Authority for Subsidy

Initial application — (Multiple Sclerosis - ocrelizumab) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 All of the following:
  - 1.1 Diagnosis of multiple sclerosis (MS) meets the McDonald 2017 diagnostic criteria for MS and has been confirmed by a neurologist; and
  - 1.2 Patient has an EDSS score between 0 6.0; and
  - 1.3 Patient has had at least one significant attack of MS in the previous 12 months or two significant attacks in the past 24 months: and
  - 1.4 All of the following:
    - 1.4.1 Each significant attack must be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the attack, but the neurologist/physician must be satisfied that the clinical features were characteristic): and
    - 1.4.2 Each significant attack is associated with characteristic new symptom(s)/sign(s) or substantially worsening of previously experienced symptoms(s)/sign(s); and
    - 1.4.3 Each significant attack has lasted at least one week and has started at least one month after the onset of a previous attack (where relevant); and
    - 1.4.4 Each significant attack can be distinguished from the effects of general fatigue; and is not associated with a fever (T> 37.5°C); and
    - 1.4.5 Fither:
      - 1.4.5.1 Each significant attack is severe enough to change either the EDSS or at least one of the Kurtze Functional System scores by at least 1 point; or
      - 1.4.5.2 Each significant attack is a recurrent paroxysmal symptom of multiple sclerosis (tonic seizures/spasms, trigeminal neuralgia, Lhermitte's symptom); and
  - 1.5 Evidence of new inflammatory activity on an MRI scan within the past 24 months; and
  - 1.6 Any of the following:
    - 1.6.1 A sign of that new inflammatory activity on MRI scanning (in criterion 5 immediately above) is a gadolinium enhancing lesion; or
    - 1.6.2 A sign of that new inflammatory activity is a lesion showing diffusion restriction; or
    - 1.6.3 A sign of that new inflammatory is a T2 lesion with associated local swelling; or
    - 1.6.4 A sign of that new inflammatory activity is a prominent T2 lesion that clearly is responsible for the clinical features of a recent attack that occurred within the last 2 years; or
    - 1.6.5 A sign of that new inflammatory activity is new T2 lesions compared with a previous MRI scan; or
  - 2 Patient has an active Special Authority approval for either dimethyl fumarate, fingolimod, glatiramer acetate, interferon

#### **NERVOUS SYSTEM**

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

beta-1-alpha, interferon beta-1-beta, natalizumab or teriflunomide.

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

Renewal — (Multiple Sclerosis - ocrelizumab) from any relevant practitioner. Approvals valid for 12 months where patient has had an EDSS score of 0 to 6.0 (inclusive) with or without the use of unilateral or bilateral aids at any time in the last six months (ie the patient has walked 100 metres or more with or without aids in the last six months).

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

Initial application — (Primary Progressive Multiple Sclerosis) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Diagnosis of primary progressive multiple sclerosis (PPMS) meets the 2017 McDonald criteria and has been confirmed by a neurologist; and
- 2 Patient has an EDSS 2.0 (score equal to or greater than 2 on pyramidal functions) to EDSS 6.5; and
- 3 Patient has no history of relapsing remitting multiple sclerosis.

Renewal — (Primary Progressive Multiple Sclerosis) from any relevant practitioner. Approvals valid for 12 months where patient has had an EDSS score of less than or equal to 6.5 at any time in the last six months (ie patient has walked 20 metres with bilateral assistance/aids, without rest in the last six months).

# Sedatives and Hypnotics

MELATONIN - Special Authority see SA1666 below - Retail pharmacy

#### **⇒SA1666** Special Authority for Subsidy

**Initial application** only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with persistent and distressing insomnia secondary to a neurodevelopmental disorder (including, but not limited to, autism spectrum disorder or attention deficit hyperactivity disorder)\*; and
- 2 Behavioural and environmental approaches have been tried and were unsuccessful, or are inappropriate; and
- 3 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day; and
- 4 Patient is aged 18 years or under\*.

**Renewal** only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is aged 18 years or under\*; and
- 2 Patient has demonstrated clinically meaningful benefit from funded modified-release melatonin (clinician determined); and
- 3 Patient has had a trial of funded modified-release melatonin discontinuation within the past 12 months and has had a recurrence of persistent and distressing insomnia; and
- 4 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day.

Note: Indications marked with \* are unapproved indications.

#### **NERVOUS SYSTEM**

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Generic
MIDAZOLAM - Safety medicine; prescriber may determine dispe	ensing frequency			
Inj 1 mg per ml, 5 ml ampoule	7.80 16.75	10		Midazolam-Baxter Midazolam Viatris
Inj 1 mg per ml, 5 ml plastic ampoule – Up to 10 inj available on a PSO		10	/	Pfizer
On a PSO for status epilepticus use only. PSO must be	endorsed for status e	epilep	oticus use	only.
Inj 5 mg per ml, 1 ml plastic ampoule – Up to 10 inj available on a PSOOn a PSO for status epilepticus use only. PSO must be	22.50	10 epilep		Midazolam-Pfizer
Inj 5 mg per ml, 3 ml ampoule — Brand switch fee payable (Pharmacode 2695863) - see page 272 for details	4.75 5.50	5		Midazolam-Baxter Midazolam Viatris
Inj 5 mg per ml, 3 ml plastic ampoule — Up to 5 inj available a PSO		5	/	Pfizer
On a PSO for status epilepticus use only. PSO must be (Midazolam Viatris Inj 1 mg per ml, 5 ml ampoule to be delisted 1 (Midazolam Viatris Inj 5 mg per ml, 3 ml ampoule to be delisted :	May 2025)	epilep	oticus use	only.
PHENOBARBITONE SODIUM - Special Authority see SA1386 b	pelow – Retail pharm	acy		
Inj 200 mg per ml, 1 ml ampoule		10	1	Max Health S29
■ SA1386 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid the following criteria:  Both:	d without further rene	wal u	inless notif	ied for applications meeting
For the treatment of terminal agitation that is unresponsive     The applicant is part of a multidisciplinary team working in	<b>0</b> ,	t		
TEMAZEPAM – Safety medicine; prescriber may determine disp Tab 10 mg		25	•	Normison
ZOPICLONE - Safety medicine; prescriber may determine dispe	nsing frequency			

# **Spinal Muscular Atrophy**

### ⇒SA2174 Special Authority for Subsidy

Initial application — (spinal muscular atrophy (SMA)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has genetic documentation of homozygous SMN1 gene deletion, homozygous SMN1 point mutation, or compound heterozygous mutation; and
- 2 Patient is 18 years of age or under; and
- 3 Either
  - 3.1 Patient has experienced the defined signs and symptoms of SMA type I, II or IIIa prior to three years of age; or
  - 3.2 Both:
    - 3.2.1 Patient is pre-symptomatic; and
    - 3.2.2 Patient has three or less copies of SMN2.

continued...

500

✓ Zopiclone Actavis

#### **NERVOUS SYSTEM**

Subsidy	Fully		Brand or	
(Manufacturer's Price)	5	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Renewal — (spinal muscular atrophy (SMA)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 There has been demonstrated maintenance of motor milestone function since treatment initiation; and
- 2 Patient does not require invasive permanent ventilation (at least 16 hours per day) in the absence of a potentially reversible cause while being treated with nusinersen; and
- 3 Nusinersen not to be administered in combination other SMA disease modifying treatments or gene therapy.

# RISDIPLAM - [Xpharm] - Special Authority see SA2203 below

Note: the supply of risdiplam is via Pharmac's approved direct distribution supply. Further details can be found on Pharmac's website https://pharmac.govt.nz/risdiplam

Powder for oral soln 750 mcg per ml, 60 mg per bottle......14,100.00 80 ml OP ✓ Evrysdi

#### ⇒SA2203 Special Authority for Subsidy

**Initial application — (spinal muscular atrophy (SMA))** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has genetic documentation of homozygous SMN1 gene deletion, homozygous SMN1 point mutation, or compound heterozygous mutation; and
- 2 Patient is 18 years of age or under; and
- 3 Either:
  - 3.1 Patient has experienced the defined signs and symptoms of SMA type I, II or IIIa prior to three years of age; or
  - 3.2 Both:
    - 3.2.1 Patient is pre-symptomatic; and
    - 3.2.2 Patient has three or less copies of SMN2.

Renewal — (spinal muscular atrophy (SMA)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 There has been demonstrated maintenance of motor milestone function since treatment initiation: and
- 2 Patient does not require invasive permanent ventilation (at least 16 hours per day) in the absence of a potentially reversible cause while being treated with risdiplam; and
- 3 Risdiplam not to be administered in combination other SMA disease modifying treatments or gene therapy.

# Stimulants/ADHD Treatments

ATOMOXETINE			
Cap 10 mg	43.02	28	✓ APO-Atomoxetine
Cap 18 mg	45.57	28	✓ APO-Atomoxetine
Cap 25 mg	44.30	28	✓ APO-Atomoxetine
Cap 40 mg	46.21	28	✓ APO-Atomoxetine
Cap 60 mg		28	✓ APO-Atomoxetine
Cap 80 mg	65.20	28	✓ APO-Atomoxetine
Cap 100 mg	65.71	28	✓ APO-Atomoxetine

DEXAMFETAMINE SULFATE - Special Authority see SA2410 on the next page - Retail pharmacy

- a) Only on a controlled drug form
- b) Safety medicine; prescriber may determine dispensing frequency

Tab 5	mg	 	···········	 29.80	100	1	Noumed	
							Devamfetan	nine



Subsidy		Fully	Brand or
(Manufacturer's Price)	Sı	ubsidised	Generic
\$	Per	✓	Manufacturer

# ⇒SA2410 Special Authority for Subsidy

Initial application — (ADHD in patients aged 5 years or over) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid without further renewal unless notified for applications meeting the following criteria: All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) in patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
  - 3.1 Applicant is a paediatrician or psychiatrist; or
  - 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients aged under 5 years) only from a paediatrician or psychiatrist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) in patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

**Initial application** — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid without further renewal unless notified where the patient suffers from narcolepsy.

LISDEXAMFETAMINE DIMESILATE - Special Authority see SA2415 below - Retail pharmacy

- a) Only on a controlled drug form
- b) Safety medicine; prescriber may determine dispensing frequency

Cap 30 mg - No more than 1 cap per day60.	00 30	✓ Vyvanse
Cap 50 mg60.	00 30	✓ Vyvanse
Cap 70 mg60.	00 30	✓ Vyvanse

#### ⇒SA2415 Special Authority for Subsidy

**Initial application** only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 Patient is currently on treatment with lisdexamfetamine dimesilate and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
  - 2.1 ADHD (Attention Deficit and Hyperactivity Disorder); and
  - 2.2 Diagnosed according to DSM-V or ICD 11 criteria; and
  - 2.3 Either:
    - 2.3.1 Applicant is a paediatrician or psychiatrist; or
    - 2.3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing; and
  - 2.4 Any of the following:
    - 2.4.1 Patient is taking a currently subsidised formulation of atomoxetine or methylphenidate hydrochloride (extended-release) and has not received sufficient benefit or has experienced intolerable side effects; or
    - 2.4.2 Patient is taking a currently subsidised formulation of dexamfetamine sulfate (immediate-release) which has not been effective due to significant administration and/or treatment adherence difficulties; or
    - 2.4.3 There is significant concern regarding the risk of diversion or abuse of immediate release dexamfetamine sulfate; or
    - 2.4.4 Patient is taking a currently subsidised formulation of methylphenidate hydrochloride (immediate-release or sustained release) which has not been effective due to significant administration and/or treatment

#### **NERVOUS SYSTEM**

(Ma	Subsidy anufacturer's Price)	Subs	Fully idised	Brand or Generic
	\$	Per	•	Manufacturer

continued...

adherence difficulties: or

- 2.4.5 There is significant concern regarding the risk of diversion or abuse of immediate release methylphenidate hydrochloride; or
- 2.4.6 Both:
  - 2.4.6.1 Patient would have been prescribed a subsidised formulation of methylphenidate hydrochloride (extended-release) but has been unable to access due to supply issues with methylphenidate hydrochloride (extended-release); and
  - 2.4.6.2 Other alternative stimulant presentations (methylphenidate or dexamfetamine) are not appropriate; and
- 2.5 Lisdexamfetamine dimesilate is not to be used in combination with another funded methylphenidate presentation.

#### METHYLPHENIDATE HYDROCHLORIDE - Special Authority see SA2411 below - Retail pharmacy

- a) Only on a controlled drug form
- b) Safety medicine: prescriber may determine dispensing frequency

Tab immediate-release 5 mg	3.20	30	✓ Rubifen
Tab immediate-release 10 mg		30	✓ Rubifen
· ·	4.00		✓ Ritalin
Tab extended-release 18 mg	7.75	30	<ul><li>Methylphenidate ER</li><li>Teva</li></ul>
Tab immediate-release 20 mg	7.85	30	✓ Rubifen
Tab sustained-release 20 mg	10.95	30	Rubifen SR
Tab extended-release 27 mg	11.45	30	<ul><li>Methylphenidate ER</li><li>Teva</li></ul>
Tab extended-release 36 mg	15.50	30	<ul><li>Methylphenidate ER</li><li>Teva</li></ul>
Tab extended-release 54 mg	22.25	30	✓ Methylphenidate ER - Teva

#### ⇒SA2411 Special Authority for Subsidy

Initial application — (ADHD in patients aged 5 years or over) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid without further renewal unless notified for applications meeting the following criteria: All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) in patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Fither:
  - 3.1 Applicant is a paediatrician or psychiatrist; or
  - 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients aged under 5 years) only from a paediatrician or psychiatrist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) in patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy\*) only from a neurologist or respiratory specialist. Approvals valid without further renewal unless notified where the patient suffers from narcolepsy.

Note: \*narcolepsy is not a registered indication for Methylphenidate ER – Teva.

# **NERVOUS SYSTEM**

Subsidy		Fully	Brand or
(Manufacturer's Price)	S	Subsidised	Generic
 \$	Per	1	Manufacturer

METHYLPHENIDATE HYDROCHLORIDE EXTENDED-RELEASE - Special Authority see SA2412 below - Retail pharmacy

- a) Only on a controlled drug form
- b) Safety medicine; prescriber may determine dispensing frequency

b) dalety inedictie, prescriber may determine dispensing	requeries		
Tab extended-release 18 mg	58.96	30	<ul><li>Concerta</li></ul>
Tab extended-release 27 mg	65.44	30	<ul><li>Concerta</li></ul>
Tab extended-release 36 mg		30	<ul><li>Concerta</li></ul>
Tab extended-release 54 mg		30	<ul><li>Concerta</li></ul>
Cap modified-release 10 mg		30	Ritalin LA
Cap modified-release 20 mg	27.72	30	Ritalin LA
Cap modified-release 30 mg		30	Ritalin LA
Cap modified-release 40 mg	38.67	30	Ritalin LA

#### ⇒SA2412 Special Authority for Subsidy

Initial application — (ADHD) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder); and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Fither:
  - 3.1 Applicant is a paediatrician or psychiatrist; or
  - 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing; and
- 4 Either:
  - 4.1 Patient is taking a currently subsidised formulation of methylphenidate hydrochloride (immediate-release or sustained-release) which has not been effective due to significant administration and/or difficulties with adherence; or
  - 4.2 There is significant concern regarding the risk of diversion or abuse of immediate-release methylphenidate hydrochloride.

# MODAFINIL – Special Authority see SA2413 below – Retail pharmacy Tab 100 mg .......14.27 30 ✓ Modafinil Max Health 29.13 60 ✓ Modavigil

(Modavigil Tab 100 mg to be delisted 1 May 2025)

#### ⇒SA2413 Special Authority for Subsidy

**Initial application** only from a neurologist or respiratory specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### All of the following:

- 1 The patient has a diagnosis of narcolepsy and has excessive daytime sleepiness associated with narcolepsy occurring almost daily for three months or more; and
- almost dai 2 Either:
  - 2.1 The patient has a multiple sleep latency test with a mean sleep latency of less than or equal to 10 minutes and 2 or more sleep onset rapid eye movement periods; or
  - 2.2 The patient has at least one of: cataplexy, sleep paralysis or hypnagogic hallucinations; and
- 3 Either:
  - 3.1 An effective dose of a subsidised formulation of methylphenidate or dexamfetamine has been trialled and discontinued because of intolerable side effects; or
  - 3.2 Methylphenidate and dexamfetamine are contraindicated.

	Subsidy (Manufacturer's Price)		Fully sidised	Brand or Generic
	\$	Per		Manufacturer
Treatments for Dementia				
DONEPEZIL HYDROCHLORIDE				
* Tab 5 mg	3.70	84	<b>√</b> <u>l</u> j	pca-Donepezil
* Tab 10 mg		84	<b>✓</b> <u>l</u>	pca-Donepezil
RIVASTIGMINE - Special Authority see SA1488 below - Retail I	pharmacy			
Patch 4.6 mg per 24 hour	49.40	30	<b>√</b> F	Rivastigmine Patch BNM 5
	90.00		<b>√</b> E	xelon Patch 5
Rivastigmine Patch BNM 5 to be Principal Supply on 1 N	March 2025			
Patch 9.5 mg per 24 hour	49.40	30	<b>√</b> F	Rivastigmine Patch BNM 10
	90.00		<b>√</b> E	xelon Patch 10
Rivastigmine Patch BNM 10 to be Principal Supply on 1	March 2025			
(Exelon Patch 5 Patch 4.6 mg per 24 hour to be delisted 1 March	2025)			

## ⇒SA1488 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

1 The patient has been diagnosed with dementia; and

(Exelon Patch 10 Patch 9.5 mg per 24 hour to be delisted 1 March 2025)

2 The patient has experienced intolerable nausea and/or vomiting from donepezil tablets.

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate; and
- 2 The patient has demonstrated a significant and sustained benefit from treatment.

# Treatments for Substance Dependence

BUPRENORPHINE WITH NALOXONE - Special Authority see SA1203 below - Retail pharmacy

- a) No patient co-payment payable
- b) Safety medicine; prescriber may determine dispensing frequency

Tab sublingual 2 mg with naloxone 0.5 mg .......11.76

28

Tab sublingual 8 mg with naloxone 2 mg ......34.00

✓ Buprenorphine
 Naloxone BNM
 ✓ Buprenorphine

Naloxone BNM

# ⇒SA1203 Special Authority for Subsidy

**Initial application — (Detoxification)** from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health...

**Initial application — (Maintenance treatment)** from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient will not be receiving methadone; and



bsidy urer's Price) Subs	Fully	Brand or Generic
 \$ Per	•	Manufacturer

continued...

- 3 Patient is currently enrolled in an opioid substitution treatment program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient has previously trialled but failed detoxification with buprenorphine with naloxone with relapse back to opioid use and another attempt is planned; and
- 3 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is or has been receiving maintenance therapy with buprenorphine with naloxone (and is not receiving methadone); and
- 2 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health or is a medical practitioner authorised by the service to manage treatment in this patient.

Renewal — (Maintenance treatment where the patient has previously had an initial application for detoxification) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient received but failed detoxification with buprenorphine with naloxone; and
- 2 Maintenance therapy with buprenorphine with naloxone is planned (and patient will not be receiving methadone); and
- 3 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Tab modified-release 150 mg	15.00	30	✓ Zyban
DISULFIRAM			
Tab 200 mg	236.40	100	✓ Antabuse
NALTREXONE HYDROCHLORIDE - Special Authority see S	A1408 below – Reta	il pharmacy	
Tab 50 mg	77.77	28	✓ Naltrexone AOP \$29
	83.33	30	✓ Naltraccord
	102.60		✓ Naltrexone Max
			Health S29
	138.88	50	✓ Revia S29

#### ⇒SA1408 Special Authority for Subsidy

DI IDDODIONI UVDDOCUI ODIDE

**Initial application** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient is currently enrolled in a recognised comprehensive treatment programme for alcohol dependence; and
- 2 Applicant works in or with a community Alcohol and Drug Service contracted to Health NZ or accredited against the New Zealand Alcohol and Other Drug Sector Standard or the National Mental Health Sector Standard.

Renewal from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

1 Compliance with the medication (prescriber determined); and

Subsidy (Manufacturer's Pric	e)	Fully Subsidised	Brand or Generic	
(manusculor o 1 no	Per		Manufacturer	

#### continued...

- 2 Any of the following:
  - 2.1 Patient is still unstable and requires further treatment; or
  - 2.2 Patient achieved significant improvement but requires further treatment; or
  - 2.3 Patient is well controlled but requires maintenance therapy.

#### NICOTINE

- a) Nicotine will not be funded in amounts less than 4 weeks of treatment.
- b) Note: Direct Provision by a pharmacist permitted under the provisions in Part I of Section A. Patch 7 mg - Up to 28 patch available on a PSO ......19.62 ✓ Habitrol Patch 14 mg - Up to 28 patch available on a PSO ......21.57 28 ✓ Habitrol Patch 14 mg for direct distribution only - [Xpharm]......12.49 7 ✓ Habitrol Patch 21 mg - Up to 28 patch available on a PSO ......24.72 28 ✓ Habitrol Patch 21 mg for direct distribution only - [Xpharm]......13.19 7 ✓ Habitrol Lozenge 1 mg - Up to 216 loz available on a PSO......22.53 216 ✓ Habitrol 36 ✓ Habitrol ✓ Habitrol Lozenge 2 mg - Up to 216 loz available on a PSO......24.68 216 36 ✓ Habitrol ✓ Habitrol Gum 2 mg (Fruit) - Up to 204 piece available on a PSO ......23.02 204 Gum 2 mg (Fruit) for direct distribution only - [Xpharm].......17.57 96 ✓ Habitrol Gum 2 mg (Mint) - Up to 204 piece available on a PSO......23.02 204 ✓ Habitrol Gum 2 mg (Mint) for direct distribution only - [Xpharm]......17.57 96 ✓ Habitrol Gum 4 mg (Fruit) - Up to 204 piece available on a PSO ......25.98 204 ✓ Habitrol Gum 4 mg (Fruit) for direct distribution only - [Xpharm]...........23.87 ✓ Habitrol 96 ✓ Habitrol Gum 4 mg (Mint) - Up to 204 piece available on a PSO......25.98 204 Gum 4 mg (Mint) for direct distribution only - [Xpharm]......23.87 96 ✓ Habitrol

#### VARENICLINE TARTRATE - Special Authority see SA1845 below - Retail pharmacy

- a) A maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval, including the starter pack
- b) Varenicline will not be funded in amounts less than 4 weeks of treatment.
- c) The 6-month time period in which a patient can receive a funded 12-week course of varenicline tartrate starts from the date the Special Authority is approved.

Tab 0.5 mg × 11 and 1 mg × 42	16.67	53 OP	✓ Varenicline Pfizer
Tab 1 mg	17.62	56	✓ Varenicline Pfizer

#### ⇒SA1845 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 Either
  - 3.1 The patient has tried but failed to quit smoking after at least two separate trials of nicotine replacement therapy, at least one of which included the patient receiving comprehensive advice on the optimal use of nicotine replacement therapy; or
  - 3.2 The patient has tried but failed to guit smoking using bupropion or nortriptyline; and
- 4 The patient has not had a Special Authority for varenicline approved in the last 6 months; and
- 5 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and

# NERVOUS SYSTEM

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
 \$	Per	•	Manufacturer

continued...

- 6 The patient is not pregnant; and
- 7 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

Renewal from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 It has been 6 months since the patient's previous Special Authority was approved; and
- 4 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 5 The patient is not pregnant; and
- 6 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

The patient must not have had an approval in the past 6 months.

Notes: a maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval.

This includes the 4-week 'starter' pack.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

# **Chemotherapeutic Agents**

## Alkylating Agents

BENDAMUSTINE HYDROCHLORIDE - PCT only -	- Specialist - Special Authority	see SA2398	3 below
Inj 25 mg vial	50.05	1	✓ Bendamustine Sandoz
	77.00		✓ Ribomustin
Inj 100 mg vial	200.20	1	✓ Bendamustine Sandoz
	308.00		✓ Ribomustin
Inj 1 mg for ECP	3.23	1 mg	✓ Baxter

#### ⇒SA2398 Special Authority for Subsidy

Initial application — (CLL\*) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has chronic lymphocytic leukaemia requiring treatment; and
- 2 Patient has ECOG performance status of 0-2; and
- 3 Bendamustine is to be administered at a maximum dose of 100 mg/m<sup>2</sup> on days 1 and 2 every 4 weeks for a maximum of 6 cycles.

Note: Indication marked with a \* includes indications that are unapproved. 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL).

Initial application — (Indolent, Low-grade lymphomas) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has indolent low grade NHL requiring treatment; and
- 2 The patient has ECOG performance status of 0-2; and
- 3 Any of the following:
  - 3.1 Both:
    - 3.1.1 Patient is treatment naive; and
    - 3.1.2 Bendamustine is to be administered for a maximum of 6 cycles (in combination with rituximab when CD20+); or
  - 3.2 Both:
    - 3.2.1 Patient is refractory to or has relapsed within 12 months of a rituximab containing combined chemo-immunotherapy regimen; and
    - 3.2.2 Bendamustine is to be administered in combination with obinutuzumab for a maximum of 6 cycles; or
  - 3.3 All of the following:
    - 3.3.1 The patient has not received prior bendamustine therapy; and
    - 3.3.2 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
    - 3.3.3 Patient has had a rituximab treatment-free interval of 12 months or more; or
  - 3.4 Bendamustine is to be administered as monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Renewal — (Indolent, Low-grade lymphomas) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: Either:

- 1 Both:
  - 1.1 Patient is refractory to or has relapsed within 12 months of rituximab in combination with bendamustine; and

Subsidy (Manufacturer's Price)	Subs	Fully	Brand or Generic
\$	Per	1	Manufacturer

continued...

- 1.2 Bendamustine is to be administered in combination with obinutuzumab for a maximum of 6 cycles; or
- 2 Both:
  - 2.1 Patients have not received a bendamustine regimen within the last 12 months; and
  - 2.2 Either:
    - 2.2.1 Both:
      - 2.2.1.1 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
      - 2.2.1.2 Patient has had a rituximab treatment-free interval of 12 months or more: or
    - 2.2.2 Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Note: 'indolent, low-grade lymphomas' includes follicular, mantle cell, marginal zone and lymphoplasmacytic/ Waldenstrom's macroglobulinaemia.

Initial application — (Hodgkin's lymphoma\*) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has Hodgkin's lymphoma requiring treatment; and
- 2 Patient has a ECOG performance status of 0-2; and
- 3 Patient has received one prior line of chemotherapy; and
- 4 Patient's disease relapsed or was refractory following prior chemotherapy; and
- 5 Bendamustine is to be administered in combination with gemcitabine and vinorelbine (BeGeV) at a maximum dose of no greater than 90 mg/m2 twice per cycle, for a maximum of four cycles.

Note: Indications marked with \* are unapproved indications.

BUSINE FAN - PCT - Botail pharmacy-Specialist

BUSULFAN – PUT – Retail pharmacy-Specialist	00.05	100	. Mulanan
Tab 2 mg	89.25	100	✓ Myleran
CARBOPLATIN - PCT only - Specialist			
Inj 10 mg per ml, 45 ml vial	25.73	1	<ul> <li>Carboplatin Accord</li> </ul>
	32.59		✓ DBL Carboplatin
	48.50		✓ Carbaccord
Inj 1 mg for ECP	0.06	1 mg	✓ Baxter
CARMUSTINE - PCT only - Specialist			
Inj 100 mg vial	710.00	1	✓ BiCNU
			✓ BiCNU S29 S29
			✓ Novadoz S29
Inj 100 mg for ECP	710.00	100 mg OP	✓ Baxter
CHLORAMBUCIL - PCT - Retail pharmacy-Specialist			
Tab 2 mg	29.06	25	✓ Leukeran FC
CISPLATIN - PCT only - Specialist			
Inj 1 mg per ml, 50 ml vial	9.45	1	✓ Cisplatin Accord
, ,	15.00		✓ Cisplatin Ebewe
Inj 1 mg per ml, 100 ml vial	18.90	1	✓ Cisplatin Accord
, ,	21.00		✓ Cisplatin Ebewe
	29.66		✓ DBL Cisplatin
Inj 1 mg for ECP	0.31	1 mg	✓ Baxter

	Subsidy		Fully	
	Manufacturer's Price)	Per	Subsidised	Generic Manufacturer
CYCLOPHOSPHAMIDE	*			
Tab 50 mg - PCT - Retail pharmacy-Specialist	145.00	50	1	Cyclonex
Inj 1 g vial – PCT – Retail pharmacy-Specialist		1		Endoxan
, 3 , , . , . , . , .	127.80	6	1	Cytoxan
Endoxan to be Principal Supply on 1 February 2025				•
Inj 2 g vial - PCT only - Specialist	95.06	1	1	Endoxan
Endoxan to be Principal Supply on 1 February 2025				
Inj 1 mg for ECP - PCT only - Specialist	0.05	1 mg	<b>√</b>	Baxter
FOSFAMIDE - PCT only - Specialist				
Inj 1 g	96.00	1	1	Holoxan
lnj 2 g		1	1	Holoxan
Inj 1 mg for ECP	0.10	1 mg	· •	Baxter
OMUSTINE - PCT - Retail pharmacy-Specialist				
Cap 40 mg	880.00	20	1	Medac S29
MELPHALAN				
Tab 2 mg - PCT - Retail pharmacy-Specialist	40.70	25	1	Alkeran
Inj 50 mg - PCT only - Specialist		1		Megval S29
inj 30 mg = 1 01 omy = Specialist	40.23	'		Melpha
	67.80			Alkeran
DXALIPLATIN - PCT only - Specialist	0.100			
Inj 100 mg vial	25.01	1	1	Oxaliplatin Actavis
ing 100 mg viai	25.01	'	•	100
	110.00		1	Oxaliplatin Ebewe
Inj 5 mg per ml, 20 ml vial		1		Alchemy Oxaliplatin
, 0, 5, 20	46.32	•		Oxaliplatin Accord
Inj 1 mg for ECP	0.35	1 mg		Baxter
THIOTEPA - PCT only - Specialist		•		
Inj 15 mg vial	CBS	1	1	Bedford \$29
IIIJ 13 IIIg viai		'		Max Health S29
				THIO-TEPA \$29
	398.00			Tepadina
lei 400 eservial				Max Health S29
Inj 100 mg vial		1		
	1,800.00			Tepadina
Antimetabolites				
AZACITIDINE - PCT only - Specialist - Special Authority see SA	2141 below			
Inj 100 mg vial		1	1	Azacitidine Dr
, ,				Reddy's
Inj 1 mg for ECP	0.54	1 mg	· •	Baxter
⇒SA2141 Special Authority for Subsidy		·		

#### ⇒SA2141 Special Authority for Subsidy

Initial application only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 The patient has International Prognostic Scoring System (IPSS) intermediate-2 or high risk myelodysplastic syndrome; or

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	1	Manufacturer	

continued...

- 1.2 The patient has chronic myelomonocytic leukaemia (10%-29% marrow blasts without myeloproliferative disorder); or
- 1.3 The patient has acute myeloid leukaemia with 20-30% blasts and multi-lineage dysplasia, according to World Health Organisation Classification (WHO); and
- 2 The patient has performance status (WHO/ECOG) grade 0-2; and
- 3 The patient has an estimated life expectancy of at least 3 months.

**Renewal** only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

CALCIUM FOLINATE		
Tab 15 mg – PCT – Retail pharmacy-Specialist	10	✓ DBL Leucovorin Calcium
Inj 3 mg per ml, 1 ml - PCT - Retail pharmacy-Specialist	5	✓ Hospira
Inj 10 mg per ml, 5 ml vial - PCT - Retail pharmacy-Specialist7.28	1	✓ Calcium Folinate Sandoz
		✓ Calcium Folinate
		Sandoz S29 S29
36.48	5	✓ Eurofolic S29
Inj 50 mg - PCT - Retail pharmacy-Specialist72.80	10	✓ Leucovorin Pharmacia S29
Inj 10 mg per ml, 10 ml vial - PCT only - Specialist9.49	1	✓ Calcium Folinate Sandoz
47.45	5	✓ Eurofolic S29
Inj 100 mg - PCT only - Specialist7.33	1	<ul><li>Calcium Folinate Ebewe</li></ul>
94.90	10	✓ Leucovorin  Pharmacia \$29
Inj 300 mg - PCT only - Specialist21.55	1	✓ Leucovorin DBL ©29
22.51		<ul><li>Calcium Folinate Ebewe</li></ul>
Inj 10 mg per ml, 35 ml vial - PCT only - Specialist25.14	1	<ul><li>Calcium Folinate Sandoz</li></ul>
		✓ Calcium Folinate Sandoz S29 S29
Inj 1 g - PCT only - Specialist67.51	1	✓ Calcium Folinate Ebewe
Inj 10 mg per ml, 100 ml vial - PCT only - Specialist72.00	1	✓ Calcium Folinate Sandoz
		✓ Eurofolic S29
Inj 1 mg for ECP - PCT only - Specialist	1 mg	✓ Baxter
	9	<b>S</b> unto:
CAPECITABINE – Retail pharmacy-Specialist Tab 150 mg9.80	60	✓ Capecitabine Viatris
Tab 500 mg	120	✓ Capecitabine Viatris
1 ab 300 mg46.30	120	- capecitabilie viatris

	Subsidy		Fully Brand or
	(Manufacturer's F	Price) Subsid	dised Generic ✓ Manufacturer
CLADRIBINE - PCT only - Specialist			
Inj 1 mg per ml, 10 ml	749.96	1	✓ Leustatin
Inj 10 mg for ECP	749.96	10 mg OP	✓ Baxter
CYTARABINE			
Inj 20 mg per ml, 5 ml vial - PCT - Retail pharmacy-Specia	alist472.00	5	✓ Pfizer
Inj 100 mg per ml, 20 ml vial - PCT - Retail			
pharmacy-Specialist	48.80	1	<ul> <li>Cytarabine DBL</li> </ul>
			✓ Pfizer
			✓ Pfizer S29 S29
Inj 1 mg for ECP - PCT only - Specialist	0.29	10 mg	✓ Baxter
Inj 100 mg intrathecal syringe for ECP - PCT only - Specia	alist94.40	100 mg OP	✓ Baxter
FLUDARABINE PHOSPHATE			
Tab 10 mg - PCT - Retail pharmacy-Specialist	412.00	20	✓ Fludara Oral
Inj 50 mg vial - PCT only - Specialist		1	✓ Fludarabine
, , ,			Sagent S29
	634.00	5	✓ Fludarabine Ebewe
Inj 50 mg for ECP - PCT only - Specialist		50 mg OP	✓ Baxter
FLUOROURACIL		J	
Inj 50 mg per ml, 20 ml vial – PCT only – Specialist	10.51	1	✓ Fluorouracil Accord
Inj 50 mg per ml, 50 ml vial – PCT only – Specialist		1	✓ Fluorouracil Accord
Inj 50 mg per ml, 100 ml vial – PCT only – Specialist		1	✓ Fluorouracil Accord
Inj 1 mg for ECP - PCT only - Specialist		100 mg	✓ Baxter
GEMCITABINE HYDROCHLORIDE - PCT only - Specialist			
Inj 43.3 mg per ml (equivalent to 38 mg per ml gemcitabine	١		
26.3 ml vial		1	✓ DBL Gemcitabine
Inj 1 g		1	✓ Gemcitabine Ebewe
Inj 1 mg for ECP		1 mg	✓ Baxter
IRINOTECAN HYDROCHLORIDE - PCT only - Specialist		J	
Inj 20 mg per ml, 5 ml vial	52 57	1	✓ Accord
11) 20 11g por 1111, 0 1111 Val	71.44	•	✓ Irinotecan Actavis
			100
	100.00		✓ Irinotecan-Rex
Inj 1 mg for ECP		1 mg	✓ Baxter
MERCAPTOPURINE		3	
Tab 50 mg - PCT - Retail pharmacy-Specialist	25 90	25	✓ Puri-nethol
Oral suspension 20 mg per ml – Retail pharmacy-Specialis		20	- I difficulor
Special Authority see SA1725 below		100 ml OP	✓ Allmercap
oposici nationly soo ortifico bolon		.00 1111 01	✓ Xaluprine S29
			- Adiupillic de

# ⇒SA1725 Special Authority for Subsidy

Initial application only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where the patient requires a total dose of less than one full 50 mg tablet per day.

Renewal only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where patient still requires a total dose of less than one full 50 mg tablet per day.

(M	Subsidy anufacturer's Price) \$	Per	Fully Subsidised	Generic
ETHOTREXATE	•			
Tab 2.5 mg - PCT - Retail pharmacy-Specialist	7.80	90	1	Trexate
Tab 10 mg - PCT - Retail pharmacy-Specialist		90		Trexate
Inj 2.5 mg per ml, 2 ml - PCT - Retail pharmacy-Specialist		5		Methotrexate DBL
			✓	Methotrexate DBL S29 S29
Inj 7.5 mg prefilled syringe	29.17	1	✓	Methotrexate Sandoz
Methotrexate Sandoz to be Principal Supply on 1 February			_	
Inj 10 mg prefilled syringe	19.09	1	•	Methotrexate Sandoz
Methotrexate Sandoz to be Principal Supply on 1 February				
Inj 15 mg prefilled syringe		1	•	Methotrexate Sandoz
Methotrexate Sandoz to be Principal Supply on 1 February				
Inj 20 mg prefilled syringe		1	•	Methotrexate Sandoz
Methotrexate Sandoz to be Principal Supply on 1 February				Mathatuarrata
Inj 25 mg prefilled syringe		1	•	Methotrexate Sandoz
Methotrexate Sandoz to be Principal Supply on 1 February 2 Inj 30 mg prefilled syringe		1	1	Methotrexate
Methotrexate Sandoz to be Principal Supply on 1 February		'	•	Sandoz
Inj 25 mg per ml, 2 ml vial – PCT – Retail pharmacy-Specialist.		5	✓	Methotrexate DBL Onco-Vial
Inj 25 mg per ml, 20 ml vial - PCT - Retail pharmacy-Specialist	45.00	1	✓	DBL Methotrexate Onco-Vial
Inj 100 mg per ml, 10 ml - PCT - Retail pharmacy-Specialist Inj 100 mg per ml, 50 ml vial - PCT - Retail	25.00	1	•	Methotrexate Ebewe
pharmacy-Specialist		1		Methotrexate Ebewe
Inj 1 mg for ECP - PCT only - Specialist		1 mg		Baxter
Inj 5 mg intrathecal syringe for ECP - PCT only - Specialist EMETREXED - PCT only - Specialist		mg C	P 🗸	Baxter
Inj 100 mg vial		1		Pemetrexed-AFT
	60.89			Juno Pemetrexed
Inj 500 mg vial		1		Pemetrexed-AFT
Ini 1 mg for FCD	217.77	4		Juno Pemetrexed
Inj 1 mg for ECP HOGUANINE – PCT – Retail pharmacy-Specialist	0.11	1 mg	•	Baxter
Tab 40 mg	126.31	25	•	Lanvis
Other Cytotoxic Agents				
MSACRINE - PCT only - Specialist				
Inj 50 mg per ml, 1.5 ml ampoule	1,500.00	6	✓	Amsidine S29
	4,736.00		1	Amsidine S29
Inj 75 mg	1,250.00	5	✓	AmsaLyo S29
"ij 70 mg				

	Subsidy		Fully	
	(Manufacturer's P	Price) Subs Per	idised	
ANACREURE HVDDOCHLORIDE DOT Batail abovemon Co	•	101		Mandadarer
ANAGRELIDE HYDROCHLORIDE – PCT – Retail pharmacy-Sp Cap 0.5 mg		100	ſ	Agrylin
	1,175.07	100	٠	Agryiiii
ARSENIC TRIOXIDE - PCT only - Specialist Inj 1 mg per ml, 10 ml vial	4 917 00	10		Phenasen
Inj 10 mg for ECP		10 mg OP	_	Baxter
BLEOMYCIN SULPHATE – PCT only – Specialist		10 mg Oi	•	Duxiei
Inj 15,000 iu, vial	185 16	1	/	DBL Bleomycin
11j 10,000 id, vidi	100.10	•	٠	Sulfate
Inj 1,000 iu for ECP	14.32	1,000 iu	1	Baxter
BORTEZOMIB - PCT only - Specialist - Special Authority see S		1,000		
Inj 3.5 mg vial		1	1	DBL Bortezomib
Inj 1 mg for ECP		1 mg	_	Baxter
<b>⇒SA2355</b> Special Authority for Subsidy		· ·		
Initial application — (plasma cell dyscrasia) from any relevant	t practitioner. Au	oprovals valid v	vithou	ut further renewal unless
notified where the patient has plasma cell dyscrasia, not including				
DACARBAZINE - PCT only - Specialist	,	Ü		, ,
Inj 200 mg vial	72.11	1	1	DBL Dacarbazine
Inj 200 mg for ECP		200 mg OP	1	Baxter
DACTINOMYCIN [ACTINOMYCIN D] - PCT only - Specialist				
Inj 0.5 mg vial	255.00	1	1	Cosmegen
Inj 0.5 mg for ECP		0.5 mg OP		Baxter
DAUNORUBICIN - PCT only - Specialist				
Inj 2 mg per ml, 10 ml	171.93	1	1	Pfizer
Inj 20 mg for ECP	171.93	20 mg OP	1	Baxter
DOCETAXEL - PCT only - Specialist				
Inj 20 mg	48.75	1	✓	<b>Docetaxel Sandoz</b>
Inj 10 mg per ml, 8 ml vial	24.91	1	✓	DBL Docetaxel
Inj 20 mg per ml, 4 ml vial	26.95	1	•	Docetaxel
				Accord S29
Inj 80 mg		1	_	Docetaxel Sandoz
Inj 1 mg for ECP	0.35	1 mg		Baxter
DOXORUBICIN HYDROCHLORIDE - PCT only - Specialist				
Inj 2 mg per ml, 5 ml vial		1	_	Doxorubicin Ebewe
Inj 2 mg per ml, 25 ml vial		1	_	Doxorubicin Ebewe
Inj 2 mg per ml, 50 ml vial	17.00	1	_	Arrow-Doxorubicin
Inj 2 mg per ml, 100 ml vial		1		Doxorubicin Ebewe Arrow-Doxorubicin
inj z mg per mi, 100 mi viai	69.99	'		Doxorubicin Ebewe
Inj 1 mg for ECP		1 mg		Baxter
EPIRUBICIN HYDROCHLORIDE - PCT only - Specialist		3		
Inj 2 mg per ml, 5 ml vial	25.00	1	1	Epirubicin Ebewe
Inj 2 mg per ml, 25 ml vial	30.00	i		Epirubicin Ebewe
Inj 2 mg per ml, 100 ml vial		1		Epirubicin Ebewe
Inj 1 mg for ECP		1 mg		Baxter
ETOPOSIDE				
Cap 50 mg - PCT - Retail pharmacy-Specialist	340.73	20	1	Vepesid
Cap 100 mg - PCT - Retail pharmacy-Specialist		10		Vepesid
Inj 20 mg per ml, 5 ml vial - PCT - Retail pharmacy-Speciali	st7.90	1		Rex Medical
Inj 1 mg for ECP - PCT only - Specialist	0.09	1 mg	1	Baxter

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. 

★Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price \$	) Per	Fully Subsidised	Brand or Generic Manufacturer	
ETOPOSIDE PHOSPHATE – PCT only – Specialist Inj 100 mg (of etoposide base) Inj 1 mg (of etoposide base) for ECP		1 1 mg		Etopophos Baxter	
HYDROXYUREA [HYDROXYCARBAMIDE] – PCT – Retail phar Cap 500 mg	, ,	100	<b>√</b> <u>j</u>	<u>Devatis</u>	
IBRUTINIB – Special Authority see SA2168 below – Retail pharm Tab 140 mg Tab 420 mg	3,217.00	30 30	-	Imbruvica Imbruvica	

#### ⇒SA2168 Special Authority for Subsidy

Initial application — (chronic lymphocytic leukaemia (CLL)) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic lymphocytic leukaemia (CLL) requiring therapy; and
- 2 Patient has not previously received funded ibrutinib; and
- 3 Ibrutinib is to be used as monotherapy; and
- 4 Any of the following:
  - 4.1 Both:
    - 4.1.1 There is documentation confirming that patient has 17p deletion or TP53 mutation; and
    - 4.1.2 Patient has experienced intolerable side effects with venetoclax monotherapy; or
  - 4.2 All of the following:
    - 4.2.1 Patient has received at least one prior immunochemotherapy for CLL; and
    - 4.2.2 Patient's CLL has relapsed within 36 months of previous treatment; and
    - 4.2.3 Patient has experienced intolerable side effects with venetoclax in combination with rituximab regimen; or
  - 4.3 Patient's CLL is refractory to or has relapsed within 36 months of a venetoclax regimen.

**Renewal — (chronic lymphocytic leukaemia (CLL))** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 No evidence of clinical disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL) and B-cell prolymphocytic leukaemia (B-PLL)\*. Indications marked with \* are Unapproved indications.

#### IDARUBICIN HYDROCHI ORIDE

Inj 5 mg vial - PCT only - Specialist	109.74 1	✓ Zavedos
Inj 10 mg vial - PCT only - Specialist	233.64 1	✓ Zavedos
Inj 1 mg for ECP - PCT only - Specialist	25.77 1 mg	✓ Baxter

# LENALIDOMIDE (REVLIMID) – Retail pharmacy-Specialist – Special Authority see SA2047 on the next page Wastage claimable

Cap 5 mg	5,122.76	28	Revlimid
Cap 10 mg		28	✓ Revlimid
Cap 15 mg	7,239.18	28	✓ Revlimid
Can 25 mg	7 627 00	21	✓ Revlimid

(Revlimid Cap 5 mg to be delisted 1 February 2025)

(Revlimid Cap 10 mg to be delisted 1 February 2025)

(Revlimid Cap 15 mg to be delisted 1 February 2025)

(Revlimid Cap 25 mg to be delisted 1 February 2025)

Subsidy		Fully	Brand or	
(Manufacturer's Price)	;	Subsidised	Generic	
\$	Per	✓	Manufacturer	

#### ⇒SA2047 Special Authority for Subsidy

Initial application — (Relapsed/refractory disease) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has relapsed or refractory multiple myeloma with progressive disease; and
- 2 Patient has not previously been treated with lenalidomide; and
- 3 Either
  - 3.1 Lenalidomide to be used as third line\* treatment for multiple myeloma; or
  - 3.2 Both:
    - 3.2.1 Lenalidomide to be used as second line treatment for multiple myeloma; and
    - 3.2.2 The patient has experienced severe (grade 3 or higher), dose limiting, peripheral neuropathy with either bortezomib or thalidomide that precludes further treatment with either of these treatments; and
- 4 Lenalidomide to be administered at a maximum dose of 25 mg/day in combination with dexamethasone.

**Initial application** — (Maintenance following first-line autologous stem cell transplant (SCT)) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has newly diagnosed symptomatic multiple myeloma and has undergone first-line treatment that included an autologous stem cell transplantation; and
- 2 Patient has at least a stable disease response in the first 100 days after transplantation; and
- 3 Lenalidomide maintenance is to be commenced within 6 months of transplantation; and
- 4 Lenalidomide to be administered at a maximum dose of 15 mg/day.

Renewal — (Relapsed/refractory disease) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

Renewal — (Maintenance following first line autologous SCT) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

Note: Indication marked with \* is an unapproved indication. A line of treatment is considered to comprise either: a) a known therapeutic chemotherapy regimen and supportive treatments or b) a transplant induction chemotherapy regimen, stem cell transplantation and supportive treatments. Prescriptions must be written by a registered prescriber in the lenalidomide risk management programme operated by the supplier.

LENALIDOMIDE (VIATRIS) - Special Authority see SA2353 o	n the next page - Re	tail pharmad	СУ
Cap 5 mg	76.92	21	✓ Lenalidomide Viatris
Lenalidomide Viatris to be Principal Supply on 1 Febru	ary 2025		
Cap 10 mg	50.30	21	✓ Lenalidomide Viatris
Lenalidomide Viatris to be Principal Supply on 1 Febru	ary 2025		
Cap 15 mg	62.13	21	<ul><li>Lenalidomide Viatris</li></ul>
Lenalidomide Viatris to be Principal Supply on 1 Febru	ary 2025		
Cap 25 mg	65.09	21	✓ Lenalidomide Viatris
Lenalidomide Viatris to be Principal Supply on 1 Febru	ary 2025		

Subsidy (Manufacturer's Pri	ice)	Fu Subsidis	,	Brand or Generic
\$	P	er	/	Manufacturer

#### ⇒SA2353 Special Authority for Subsidy

Initial application — (Plasma cell dyscrasia) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Both:

- 1 Patient has plasma cell dyscrasia, not including Waldenström macroglobulinaemia, requiring treatment; and
- 2 Patient is not refractory to prior lenalidomide use.

**Initial application — (Myelodysplastic syndrome)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

#### Both:

- 1 Patient has low or intermediate-1 risk myelodysplastic syndrome (based on IPSS or an IPSS-R score of less than 3.5) associated with a deletion 5q cytogenetic abnormality; and
- 2 Patient has transfusion-dependent anaemia.

**Renewal — (Myelodysplastic syndrome)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

## Both:

- 1 Patient has not needed a transfusion in the last 4 months; and
- 2 No evidence of disease progression.

#### **MESNA**

Tab 400 mg - PCT - Retail pharmacy-Specialist	314.00	50	✓ Uromitexan
Tab 600 mg - PCT - Retail pharmacy-Specialist	448.50	50	✓ Uromitexan
Inj 100 mg per ml, 4 ml ampoule - PCT only - Specialist	177.45	15	✓ Uromitexan
Inj 100 mg per ml, 10 ml ampoule - PCT only - Specialist		15	✓ Uromitexan
Inj 1 mg for ECP - PCT only - Specialist		100 mg	✓ Baxter
MITOMYCIN C - PCT only - Specialist			
Inj 5 mg vial	517.65	1	✓ Mitomycin (Fresenius Kabi) ©29
	526.00		✓ Mitomycin (Sagent) S29
	641.70		✓ Accord S29
Inj 20 mg vial	1,250.00	1	<ul> <li>✓ Omegapharm S29</li> <li>✓ Teva</li> </ul>
Inj 1 mg for ECP	269.85	1 mg	✓ Baxter
MITOZANTRONE - PCT only - Specialist			
Inj 2 mg per ml, 10 ml vial	97.50	1	Mitozantrone Ebewe
Inj 1 mg for ECP		1 mg	✓ Baxter
NIRAPARIB – Special Authority see SA2325 below – Retail pharm Wastage claimable	nacy		
Tab 100 mg	13,393.50	84	✓ Zejula
Cap 100 mg	8,929.84	56	✓ Zejula
	13,393.50	84	✓ Zejula

## ⇒SA2325 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has advanced high-grade serous\* epithelial ovarian, fallopian tube, or primary peritoneal cancer; and
- 2 Patient has received at least one line\*\* of treatment with platinum-based chemotherapy; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	;	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 3 Patient has experienced a partial or complete response to the preceding treatment with platinum-based chemotherapy; and
- 4 Patient has not previously received funded treatment with a PARP inhibitor; and
- 5 Fither:
  - 5.1 Treatment will be commenced within 12 weeks of the patient's last dose of the preceding platinum-based regimen; or
- 5.2 Patient commenced treatment with niraparib prior to 1 May 2024; and
- 6 Treatment to be administered as maintenance treatment; and
- 7 Treatment not to be administered in combination with other chemotherapy.

**Renewal** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 No evidence of progressive disease; and
- 2 Treatment to be administered as maintenance treatment; and
- 3 Treatment not to be administered in combination with other chemotherapy; and
- 4 Either
  - 4.1 Treatment with niraparib to cease after a total duration of 36 months from commencement; or
  - 4.2 Treatment with niraparib is being used in the second-line or later maintenance setting.

Notes: \* "high-grade serous" includes tumours with high-grade serous features or a high-grade serous component.

\*\*A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments

pnarmacy-Specialist – Special Authority see SA2163 b	elow	
3,701.0	00 56	<ul><li>Lynparza</li></ul>
3,701.0	00 56	<ul><li>Lynparza</li></ul>

#### ⇒SA2163 Special Authority for Subsidy

Initial application — (Ovarian cancer) only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a high-grade serous\* epithelial ovarian, fallopian tube, or primary peritoneal cancer; and
- 2 There is documentation confirming pathogenic germline BRCA1 or BRCA2 gene mutation; and
- 3 Fither:
  - 3.1 All of the following:
    - 3.1.1 Patient has newly diagnosed, advanced disease; and
    - 3.1.2 Patient has received one line\*\* of previous treatment with platinum-based chemotherapy; and
    - 3.1.3 Patient's disease must have experienced a partial or complete response to the first-line platinum-based regimen; or
  - 3.2 All of the following:
    - 3.2.1 Patient has received at least two lines\*\* of previous treatment with platinum-based chemotherapy; and
    - 3.2.2 Patient has platinum sensitive disease defined as disease progression occurring at least 6 months after the last dose of the penultimate line\*\* of platinum-based chemotherapy; and
    - 3.2.3 Patient's disease must have experienced a partial or complete response to treatment with the immediately preceding platinum-based regimen; and
    - 3.2.4 Patient has not previously received funded olaparib treatment; and
- 4 Treatment will be commenced within 12 weeks of the patient's last dose of the immediately preceding platinum-based regimen; and
- 5 Treatment to be administered as maintenance treatment; and
- 6 Treatment not to be administered in combination with other chemotherapy.

Subsidy	y Full	Brand or
(Manufacturer	's Price) Subsidise	I Generic
\$	Per 🗸	Manufacturer

continued...

**Renewal — (Ovarian cancer)** only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from treatment; and
- 2 Either:
  - 2.1 No evidence of progressive disease; or
  - 2.2 Evidence of residual (not progressive) disease and the patient would continue to benefit from treatment in the clinician's opinion; and
- 3 Treatment to be administered as maintenance treatment; and
- 4 Treatment not to be administered in combination with other chemotherapy; and
- 5 Either:
  - 5.1 Both:
    - 5.1.1 Patient has received one line\*\* of previous treatment with platinum-based chemotherapy; and
    - 5.1.2 Documentation confirming that the patient has been informed and acknowledges that the funded treatment period of olaparib will not be continued beyond 2 years if the patient experiences a complete response to treatment and there is no radiological evidence of disease at 2 years; or
  - 5.2 Patient has received at least two lines\*\* of previous treatment with platinum-based chemotherapy.

Notes: \*Note "high-grade serous" includes tumours with high-grade serous features or a high-grade serous component.
\*\*A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

PACLITAXEL - PCT only - Specialist		
Inj 30 mg47.30	5	✓ Paclitaxel Ebewe
Inj 6 mg per ml, 16.7 ml vial19.59	1	✓ Anzatax
24.00		✓ Paclitaxel Ebewe
91.67		✓ Paclitaxel Actavis
Inj 150 mg26.69	1	✓ Paclitaxel Ebewe
137.50		✓ Anzatax
		✓ Paclitaxel Actavis
Inj 6 mg per ml, 50 ml vial	1	✓ Anzatax
44.00		✓ Paclitaxel Ebewe
275.00		✓ Paclitaxel Actavis
Inj 1 mg for ECP0.17	1 mg	✓ Baxter
PEGASPARGASE - PCT only - Special Authority see SA1979 below		
Inj 750 iu per ml, 5 ml vial	1	<ul><li>Oncaspar LYO</li></ul>

#### ⇒SA1979 Special Authority for Subsidy

**Initial application** — (Acute lymphoblastic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has newly diagnosed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

Initial application — (Lymphoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months where the patient has lymphoma requiring L-asparaginase containing protocols (e.g. SMILE).

Renewal — (Acute lymphoblastic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has relapsed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

	Subsidy (Manufacturer's Price)	Su Per	Fully bsidised	Brand or Generic Manufacturer
PENTOSTATIN [DEOXYCOFORMYCIN] - PCT only - Specialis	t			
Inj 10 mg	CBS	1	<b>✓</b> N	Nipent S29
POMALIDOMIDE - Special Authority see SA2354 below - Retai	l pharmacy			
Cap 1 mg	47.45	14	<b>✓</b> <u>F</u>	Pomolide Pomolide
•	71.18	21	<b>✓</b> F	Pomolide
Cap 2 mg	94.90	14	<b>✓</b> <u>F</u>	Pomolide Pomolide
	142.35	21	<b>✓</b> <u>F</u>	Pomolide Pomolide
Cap 3 mg	142.35	14	<b>✓</b> <u>F</u>	Pomolide Pomolide
	213.53	21	<b>✓</b> <u>F</u>	Pomolide Pomolide
Cap 4 mg	189.81	14	<b>✓</b> F	Pomolide
	284.71	21	<b>✓</b> F	Pomolide

#### ⇒SA2354 Special Authority for Subsidy

Initial application — (Relapsed/refractory plasma cell dyscrasia) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has relapsed or refractory plasma cell dyscrasia, not including Waldenström macroglobulinaemia, requiring treatment; and
- 2 Patient has not received prior funded pomalidomide.

Renewal — (Relapsed/refractory plasma cell dyscrasia) from any relevant practitioner. Approvals valid for 12 months where there is no evidence of disease progression.

PROCARBAZINE HYDROCHLORIDE – PCT – Retail phail Cap 50 mg	, ,	50	✓ Natulan S29
TEMOZOLOMIDE - Special Authority see SA2275 below -	- Retail pharmacy		
Cap 5 mg		5	<ul><li>✓ Temaccord</li><li>✓ Temozolomide- Taro \$29</li></ul>
Cap 20 mg		5	✓ Temaccord
•	18.30	_	✓ Apo-Temozolomide
Cap 100 mg	35.98	5	<ul><li>Temaccord</li></ul>
	40.20		✓ Apo-Temozolomide
Cap 140 mg	50.12	5	✓ Temaccord
Cap 250 mg	86.34	5	✓ Temaccord

#### ⇒SA2275 Special Authority for Subsidy

Initial application — (gliomas) only from a relevant specialist. Approvals valid for 12 months where the patient has a glioma. Renewal — (gliomas) only from a relevant specialist. Approvals valid for 12 months where treatment remains appropriate and patient is benefitting from treatment.

**Initial application** — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with metastatic or unresectable well-differentiated neuroendocrine tumour\*; and
- 2 Temozolomide is to be given in combination with capecitabine; and
- 3 Temozolomide is to be used in 28 day treatment cycles for a maximum of 5 days treatment per cycle at a maximum dose of 200 mg/m² per day; and
- 4 Temozolomide to be discontinued at disease progression.

**Renewal — (neuroendocrine tumours)** only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Subsidy	y Full	/ Brand or
(Manufacturer's	s Price) Subsidise	d Generic
\$	Per 🗸	Manufacturer

continued...

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

**Initial application** — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 9 months where the patient has relapsed/refractory Ewing's sarcoma.

Renewal — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indication marked with a \* is an unapproved indication. Temozolomide is not subsidised for the treatment of relapsed high grade glioma.

THALIDOMIDE - Retail pharmacy-Specialist - Special A	Authority see SA2356 below		
Cap 50 mg	378.00	28	Thalomid
Cap 100 mg	756.00	28	Thalomid

#### ⇒SA2356 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 12 months where the patient has plasma cell dyscrasia, not including Waldenström macroglobulinaemia, requiring treatment.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has obtained a response from treatment during the initial approval period.

Notes: Prescription must be written by a registered prescriber in the thalidomide risk management programme operated by the supplier.

Maximum dose of 400 mg daily as monotherapy or in a combination therapy regimen.

#### **TRETINOIN**

Cap 10 mg - PCT - Retail pharmacy-Specialist	479.50	100	Vesanoid
VENETOCLAX - Retail pharmacy-Specialist - Special Authority s	ee SA1868 belo	W	
Tab 14 × 10 mg, 7 × 50 mg, 21 × 100 mg	1,771.86	42 OP	✓ Venclexta
Tab 10 mg		2 OP	✓ Venclexta
Tab 50 mg	239.44	7 OP	✓ Venclexta
Tab 100 mg - Wastage claimable	8,209.41	120	✓ Venclexta

#### ⇒SA1868 Special Authority for Subsidy

Initial application — (relapsed/refractory chronic lymphocytic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic lymphocytic leukaemia requiring treatment; and
- 2 Patient has received at least one prior therapy for chronic lymphocytic leukaemia; and
- 3 Patient has not previously received funded venetoclax; and
- 4 The patient's disease has relapsed within 36 months of previous treatment; and
- 5 Venetoclax to be used in combination with six 28-day cycles of rituximab commencing after the 5-week dose titration schedule with venetoclax; and
- 6 Patient has an ECOG performance status of 0-2.

Renewal — (relapsed/refractory chronic lymphocytic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: Both:

1 Treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

2 Venetoclax is to be discontinued after a maximum of 24 months of treatment following the titration schedule unless earlier discontinuation is required due to disease progression or unacceptable toxicity.

Initial application — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation\*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has previously untreated chronic lymphocytic leukaemia; and
- 2 There is documentation confirming that patient has 17p deletion by FISH testing or TP53 mutation by sequencing; and
- 3 Patient has an ECOG performance status of 0-2.

Renewal — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation\*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where the treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL)\* and B-cell prolymphocytic leukaemia (B-PLL)\*. Indications marked with \* are Unapproved indications.

#### VINBLASTINE SULPHATE

Inj 1 mg per ml, 10 ml vial – PCT – Retail pharmacy-Specialist270.37	5	✓ Hospira
Inj 1 mg for ECP - PCT only - Specialist	1 mg	✓ Baxter
VINCRISTINE SULPHATE		
Inj 1 mg per ml, 1 ml vial - PCT - Retail pharmacy-Specialist74.52	5	<ul><li>DBL Vincristine Sulfate</li></ul>
Inj 1 mg per ml, 2 ml vial - PCT - Retail pharmacy-Specialist102.73	5	<ul><li>DBL Vincristine Sulfate</li></ul>
Inj 1 mg for ECP - PCT only - Specialist12.60	1 mg	✓ Baxter
VINORELBINE		
Cap 20 mg30.00	1	✓ Vinorelbine Te Arai
Cap 30 mg40.00	1	✓ Vinorelbine Te Arai
Cap 80 mg60.00	1	✓ Vinorelbine Te Arai
Inj 10 mg per ml, 1 ml vial - PCT only - Specialist42.00	1	✓ Vinorelbine Ebewe
Inj 10 mg per ml, 5 ml vial - PCT only - Specialist168.00	1	✓ Navelbine S29 S29
210.00		✓ Vinorelbine Ebewe
Inj 1 mg for ECP - PCT only - Specialist	1 mg	✓ Baxter

## **Protein-tyrosine Kinase Inhibitors**

ALECTINIB – Retail pharmacy-Specialist – Special Auth	ority see SA1870 below		
Wastage claimable			
Cap 150 mg	7,935.00	224	Alecensa

#### ⇒SA1870 Special Authority for Subsidy

**Initial application** only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-small cell lung cancer; and
- 2 There is documentation confirming that the patient has an ALK tyrosine kinase gene rearrangement using an appropriate ALK test; and
- 3 Patient has an ECOG performance score of 0-2.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid

1 No evidence of progressive disease according to RECIST criteria; and

	Subsidy	Fully	Brand or
	(Manufacturer's Price)	Subsidised	Generic
	\$	Per 🗸	Manufacturer
continued for 6 months for applications meeting the following criteria: Both:			

The patient is benefitting from and tolerating treatment.

DASATINIB – Special Authority see SA2385 below – Retail pharmacy

Wastage	claimable
---------	-----------

Tab 20 mg	132.88	60	✓ Dasatinib-Teva
	3,774.06		✓ Sprycel
Dasatinib-Teva to be Principal Supply on 1 March	2025		• •
Tab 50 mg	304.13	60	✓ Dasatinib-Teva
	6,214.20		✓ Sprycel
Dasatinib-Teva to be Principal Supply on 1 March	2025		
Tab 70 mg	415.75	60	✓ Dasatinib-Teva
-	7,692.58		✓ Sprycel

Dasatinib-Teva to be Principal Supply on 1 March 2025

(Sprycel Tab 20 mg to be delisted 1 March 2025)

(Sprycel Tab 50 mg to be delisted 1 March 2025)

(Sprycel Tab 70 mg to be delisted 1 March 2025)

# ⇒SA2385 Special Authority for Subsidy

Initial application only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis or accelerated phase; or
- 2 The patient has a diagnosis of Philadelphia chromosome-positive acute lymphoid leukaemia (Ph+ ALL); or
- 3 Both:
  - 3.1 The patient has a diagnosis of CML in chronic phase; and
  - 3.2 Any of the following:
    - 3.2.1 Patient has documented treatment failure\* with imatinib; or
    - 3.2.2 Patient has experienced treatment-limiting toxicity with imatinib precluding further treatment with imatinib; or
    - 3.2.3 Patient has high-risk chronic-phase CML defined by the Sokal or EURO scoring system.

**Renewal** only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

#### Both:

- 1 Lack of treatment failure while on dasatinib\*; and
- 2 Dasatinib treatment remains appropriate and the patient is benefiting from treatment.

Note: \*treatment failure for CML as defined by Leukaemia Net Guidelines.

ERLOTINIB - Retail pharmacy-Specialist - Special Authority see SA2422 below

Tab 100 mg	280.84	30	Alchemy
Tab 150 mg	484.24	30	✓ Alchemy

#### ⇒SA2422 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 There is documentation confirming that the disease expresses activating mutations of EGFR; and
- 3 Any of the following:
  - 3.1 Patient is treatment naive: or

Subsidy	Ful	y Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per •	Manufacturer	

continued...

- 3.2 Patient has received prior treatment in the adjuvant setting and/or while awaiting EGFR results; or
- 3.3 Both:
  - 3.3.1 The patient has discontinued osimertinib or gefitinib due to intolerance; and
  - 3.3.2 The cancer did not progress while on osimertinib or gefitinib.

Renewal from any relevant practitioner. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

GEFITINIB - Retail pharmacy-Specialist - Special Authority see SA2423 below

Tab 250 mg .......918.00 30 ✓ Iressa

#### ⇒SA2423 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 Any of the following:
  - 2.1 Patient is treatment naive; or
  - 2.2 Patient has received prior treatment in the adjuvant setting and/or while awaiting EGFR results; or
  - 2.3 Roth
    - 2.3.1 The patient has discontinued osimertinib or erlotinib due to intolerance; and
    - 2.3.2 The cancer did not progress whilst on osimertinib or erlotinib; and
- 3 There is documentation confirming that disease expresses activating mutations of EGFR.

Renewal from any relevant practitioner. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

#### **IMATINIB MESILATE**

* Cap 100 mg * Cap 400 mg		60 30	✓ <u>Imatinib-Rex</u> ✓ <u>Imatinib-Rex</u>
LENVATINIB – Special Authority see SA2407 below Wastage claimable	/ - Retail pharmacy		
Cap 4 mg	3,407.40	30	<ul><li>Lenvima</li></ul>
Cap 10 mg	3,407.40	30	<ul><li>Lenvima</li></ul>

#### ⇒SA2407 Special Authority for Subsidy

**Initial application** — **(thyroid cancer)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

#### Fither:

- 1 Patient is currently on treatment with lenvatinib and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
  - 2.1 The patient has locally advanced or metastatic differentiated thyroid cancer; and
  - 2.2 Either:
    - 2.2.1 Patient must have symptomatic progressive disease prior to treatment; or
    - 2.2.2 Patient must progressive disease at critical anatomical sites with a high risk of morbidity or mortality where local control cannot be achieved by other measures: and
  - 2.3 Any of the following:
    - 2.3.1 A lesion without iodine uptake in a RAI scan; or
    - 2.3.2 Receiving cumulative RAI greater than or equal to 600 mCi; or
    - 2.3.3 Experiencing disease progression after a RAI treatment within 12 months; or
    - 2.3.4 Experiencing disease progression after two RAI treatments administered within 12 months of each other; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 2.4 Patient has thyroid stimulating hormone (TSH) adequately supressed; and
- 2.5 Patient is not a candidate for radiotherapy with curative intent; and
- 2.6 Surgery is clinically inappropriate; and
- 2.7 Patient has an ECOG performance status of 0-2.

Renewal — (thyroid cancer) from any relevant practitioner. Approvals valid for 6 months where there is no evidence of disease progression.

Initial application — (unresectable hepatocellular carcinoma) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has unresectable hepatocellular carcinoma; and
- 2 Patient has preserved liver function (Childs-Pugh A); and
- 3 Transarterial chemoembolisation (TACE) is unsuitable; and
- 4 Patient has an ECOG performance status of 0-2; and
- 5 Patient has not received prior systemic therapy for their disease in the palliative setting.

Renewal — (unresectable hepatocellular carcinoma) from any relevant practitioner. Approvals valid for 6 months where there is no evidence of disease progression.

**Initial application — (renal cell carcinoma)** from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 All of the following:
  - 1.1 The patient has metastatic renal cell carcinoma; and
  - 1.2 The disease is of predominant clear-cell histology; and
  - 1.3 The patient has documented disease progression following one previous line of treatment; and
  - 1.4 The patient has an ECOG performance status of 0-2; and
  - 1.5 Lenvatinib is to be used in combination with everolimus; or
- 2 All of the following:
  - 2.1 Patient has received funded treatment with nivolumab for the second line treatment of metastatic renal cell carcinoma; and
  - 2.2 Patient has experienced treatment limiting toxicity from treatment with nivolumab; and
  - 2.3 Lenvatinib is to be used in combination with everolimus; and
  - 2.4 There is no evidence of disease progression.

**Renewal** — **(renal cell carcinoma)** from any relevant practitioner. Approvals valid for 4 months where there is no evidence of disease progression.

# ⇒SA2342 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria: All of the following:

- 1 Patient has a diagnosis of acute myeloid leukaemia; and
- 2 Condition must be FMS tyrosine kinase 3 (FLT3) mutation positive; and
- 3 Patient must not have received a prior line of intensive chemotherapy for acute myeloid leukaemia; and
- 4 Patient is to receive standard intensive chemotherapy in combination with midostaurin only; and
- 5 Midostaurin to be funded for a maximum of 4 cycles.

NILOTINIB - Special Authority see SA2301 on the next page - Retail pharmacy

Wastage claimable

 Cap 150 mg
 4,680.00
 120
 ✓ Tasigna

 Cap 200 mg
 6,532.00
 120
 ✓ Tasigna

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

#### ⇒SA2301 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis, high risk chronic phase, or in chronic phase; and
- 2 Fither:
  - 2.1 Patient has documented CML treatment failure\* with a tyrosine kinase inhibitor (TKI); or
  - 2.2 Patient has experienced treatment limiting toxicity with a tyrosine kinase inhibitor (TKI) precluding further treatment; and
- 3 Maximum nilotinib dose of 800 mg/day: and
- 4 Subsidised for use as monotherapy only.

Note: \*treatment failure as defined by Leukaemia Net Guidelines.

**Renewal** only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Lack of treatment failure while on nilotinib as defined by Leukaemia Net Guidelines; and
- 2 Nilotinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

OSIMERTINIB - Special Authority see SA2418 below - Retail pharmacy

Tab 40 mg	9,310.00	30	✓ Tagrisso
Tab 80 mg	9,310.00	30	✓ Tagrisso

#### ⇒SA2418 Special Authority for Subsidy

Initial application — (NSCLC - first line) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

#### Either:

- 1 Patient is currently on treatment with osimertinib and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
  - 2.1 Patient has locally advanced or metastatic, incurable, non-squamous non-small cell lung cancer (NSCLC); and
  - 2.2 Any of the following:
    - 2.2.1 Patient is treatment naïve; or
    - 2.2.2 Patient has received prior chemotherapy in the adjuvant setting and/or while awaiting EGFR results; or
    - 2.2.3 Both:
      - 2.2.3.1 The patient has discontinued gefitinib or erlotinib due to intolerance; and
      - 2.2.3.2 The cancer did not progress while on gefitinib or erlotinib; and
  - 2.3 There is documentation confirming that the cancer expresses activating mutations of EGFR; and
  - 2.4 Patient has an ECOG performance status 0-3; and
  - 2.5 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Renewal — (NSCLC – first line) from any relevant practitioner. Approvals valid for 6 months where response to or stable disease with treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period.

**Initial application — (NSCLC – second line)** from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

#### Either:

- 1 Patient is currently on treatment with osimertinib and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
  - 2.1 Patient has locally advanced or metastatic, incurable, non-squamous non-small cell lung cancer (NSCLC); and

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
	Per	•	Manufacturer

continued...

- 2.2 Patient has an ECOG performance status 0-3; and
- 2.3 The patient must have received previous treatment with erlotinib or gefitinib; and
- 2.4 There is documentation confirming that the cancer expresses T790M mutation of EGFR following progression on or after erlotinib or gefitinib: and
- 2.5 The treatment must be given as monotherapy; and
- 2.6 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Renewal — (NSCLC – second line) from any relevant practitioner. Approvals valid for 6 months where response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period.

PALBOCICLIB - Special Authority see SA2345 below - Retail pharmacy

Wastage claimable			
Tab 75 mg	4,000.00	21	Ibrance
Tab 100 mg	4,000.00	21	✓ Ibrance
Tab 125 mg	· · · · · · · · · · · · · · · · · · ·	21	✓ Ibrance
	,		

#### ⇒SA2345 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Either:

- 1 All of the following:
  - 1.1 Patient has unresectable locally advanced or metastatic breast cancer; and
  - 1.2 There is documentation confirming disease is hormone-receptor positive and HER2-negative; and
  - 1.3 Patient has an ECOG performance score of 0-2; and
  - 1.4 Either:
    - 1.4.1 Disease has relapsed or progressed during prior endocrine therapy; or
    - 1.4.2 Both:
      - 1.4.2.1 Patient is amenorrhoeic, either naturally or induced, with endocrine levels consistent with a postmenopausal or without menstrual-potential state; and
      - 1.4.2.2 Patient has not received prior systemic treatment for metastatic disease; and
  - 1.5 Treatment must be used in combination with an endocrine partner; and
  - 1.6 Patient has not received prior funded treatment with a CDK4/6 inhibitor; or
- 2 All of the following:
  - 2.1 Patient has an active Special Authority approval for ribociclib; and
  - 2.2 Patient has experienced a grade 3 or 4 adverse reaction to ribociclib that cannot be managed by dose reductions and requires treatment discontinuation; and
  - 2.3 Treatment must be used in combination with an endocrine partner; and
  - 2.4 There is no evidence of progressive disease since initiation of ribociclib.

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 Treatment must be used in combination with an endocrine partner; and
- 2 There is no evidence of progressive disease since initiation of palbociclib.

PAZOPANIB – Special Authority see SA1190 on the	e next page – Retail pharmacy		
Tab 200 mg	172.88	30	Pazopanib Teva
•	1,334.70		✓ Votrient
Tab 400 mg	464.00	30	Pazopanib Teva
-	2,669.40		✓ Votrient

(Votrient Tab 200 mg to be delisted 1 May 2025) (Votrient Tab 400 mg to be delisted 1 May 2025)

Subsidy	Fu	lly Brand or	
(Manufacturer's Price)	Subsidis	ed Generic	Generic
\$	Per	<ul> <li>Manufacturer</li> </ul>	

#### ⇒SA1190 Special Authority for Subsidy

**Initial application** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
  - 2.1 The patient is treatment naive; or
  - 2.2 The patient has only received prior cytokine treatment; or
  - 2.3 Both:
    - 2.3.1 The patient has discontinued sunitinib within 3 months of starting treatment due to intolerance; and
    - 2.3.2 The cancer did not progress whilst on sunitinib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and The patient has intermediate or poor prognosis defined as:
- 5 Any of the following:
  - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
  - 5.2 Haemoglobin level < lower limit of normal; or
  - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
  - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
  - 5.5 Karnofsky performance score of less than or equal to 70; or
  - 5.6 2 or more sites of organ metastasis; and
- 6 Pazopanib to be used for a maximum of 3 months.

**Renewal** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

#### Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Pazopanib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6.

RIBOCICLIB - Special Authority see SA2343 below - Retail pharmacy

Tractage ciairiable			
Tab 200 mg	1,883.00	21	<ul><li>Kisqali</li></ul>
•	3,767.00	42	✓ Kisqali
	5.650.00	63	✓ Kisqali

#### ⇒SA2343 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Either:

1 All of the following:

Wastage claimable

- 1.1 Patient has unresectable locally advanced or metastatic breast cancer; and
- 1.2 There is documentation confirming disease is hormone-receptor positive and HER2-negative; and
- 1.3 Patient has an ECOG performance score of 0-2; and
- 1.4 Any of the following:
  - 1.4.1 Disease has relapsed or progressed during prior endocrine therapy; or
  - 1.4.2 Both:
    - 1.4.2.1 Patient is amenorrhoeic, either naturally or induced, with endocrine levels consistent with a postmenopausal or without menstrual-potential state; and

Subsidy		Fully	Brand or
(Manufacturer's Price)	Su	bsidised	Generic
\$	Per	1	Manufacturer

continued...

- 1.4.2.2 Patient has not received prior systemic endocrine treatment for metastatic disease; or
- 1.4.3 Both:
  - 1.4.3.1 Patient commenced treatment with ribociclib in combination with an endocrine partner prior to 1 July 2024; and
  - 1.4.3.2 There is no evidence of progressive disease; and
- 1.5 Treatment to be used in combination with an endocrine partner; and
- 1.6 Patient has not received prior funded treatment with a CDK4/6 inhibitor; or
- 2 All of the following:
  - 2.1 Patient has an active Special Authority approval for palbociclib; and
  - 2.2 Patient has experienced a grade 3 or 4 adverse reaction to palbociclib that cannot be managed by dose reductions and requires treatment discontinuation; and
  - 2.3 Treatment must be used in combination with an endocrine partner; and
  - 2.4 There is no evidence of progressive disease since initiation of palbociclib.

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 Treatment must be used in combination with an endocrine partner; and
- 2 There is no evidence of progressive disease since initiation of ribociclib.

# RUXOLITINIB – Special Authority see SA1890 below – Retail pharmacy

wastage cialmable			
Tab 5 mg	2,500.00	56	Jakavi
Tab 10mg	5,000.00	56	<ul><li>Jakavi</li></ul>
Tab 15 mg	5,000.00	56	Jakavi
Tab 20 mg	5,000.00	56	<ul><li>Jakavi</li></ul>

#### ⇒SA1890 Special Authority for Subsidy

**Initial application** only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has primary myelofibrosis or post-polycythemia vera myelofibrosis or post-essential thrombocythemia myelofibrosis; and
- 2 Either:
  - 2.1 A classification of risk of intermediate-2 or high-risk myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS; or
  - 2.2 Both:
    - 2.2.1 A classification of risk of intermediate-1 myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS; and
    - 2.2.2 Patient has severe disease-related symptoms that are resistant, refractory or intolerant to available therapy; and
- 3 A maximum dose of 20 mg twice daily is to be given.

**Renewal** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 A maximum dose of 20 mg twice daily is to be given.

SUNITINIB - Special Authorit	v see SA2117 (	on the next page -	<ul> <li>Retail pharmacy</li> </ul>

Cap 12.5 mg	208.38	28	<ul> <li>Sunitinib Pfizer</li> </ul>
Cap 25 mg	416.77	28	<ul> <li>Sunitinib Pfizer</li> </ul>
Cap 50 mg	694.62	28	<ul> <li>Sunitinib Pfizer</li> </ul>

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

#### ⇒SA2117 Special Authority for Subsidy

Initial application — (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
  - 2.1 The patient is treatment naive; or
  - 2.2 The patient has only received prior cytokine treatment; or
  - 2.3 The patient has only received prior treatment with an investigational agent within the confines of a bona fide clinical trial which has Ethics Committee approval; or
  - 2.4 Both:
    - 2.4.1 The patient has discontinued pazopanib within 3 months of starting treatment due to intolerance; and
    - 2.4.2 The cancer did not progress whilst on pazopanib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and The patient has intermediate or poor prognosis defined as:
- 5 Any of the following:
  - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
  - 5.2 Haemoglobin level < lower limit of normal; or
  - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
  - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
  - 5.5 Karnofsky performance score of less than or equal to 70; or
  - 5.6 2 or more sites of organ metastasis; and
- 6 Sunitinib to be used for a maximum of 2 cycles.

Initial application — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:
Both:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal tumour (GIST); and
- 2 Either:
  - 2.1 The patient's disease has progressed following treatment with imatinib; or
  - 2.2 The patient has documented treatment-limiting intolerance, or toxicity to, imatinib.

Renewal — (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Sunitinib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6

**Renewal — (GIST)** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

The patient has responded to treatment or has stable disease as determined by Choi's modified CT response evaluation criteria as follows:

- 1 Any of the following:
  - 1.1 The patient has had a complete response (disappearance of all lesions and no new lesions); or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1.2 The patient has had a partial response (a decrease in size of 10% or more or decrease in tumour density in Hounsfield Units (HU) of 15% or more on CT and no new lesions and no obvious progression of non measurable disease): or
- 1.3 The patient has stable disease (does not meet criteria the two above) and does not have progressive disease and no symptomatic deterioration attributed to tumour progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: It is recommended that response to treatment be assessed using Choi's modified CT response evaluation criteria (J Clin Oncol, 2007, 25:1753-1759). Progressive disease is defined as either: an increase in tumour size of 10% or more and not meeting criteria of partial response (PR) by tumour density (HU) on CT; or: new lesions, or new intratumoral nodules, or increase in the size of the existing intratumoral nodules.

Renewal — (GIST pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal (GIST); and
- 2 The patient is clinically benifiting from treatment and continued treatment remains appropriate; and
- 3 Sunitinib is to be discontinued at progression; and
- 4 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

# **Endocrine Therapy**

For GnRH ANALOGUES - refer to HORMONE PREPARATIONS, Trophic Hormones, page 89

ABIRATERONE ACETATE - Retail pharmacy-Specialist - Special Authority see SA2118 below

Wastage claimable

Tab 250 mg .......4,276.19 120 **✓ Zytiga** 

## **⇒SA2118** Special Authority for Subsidy

**Initial application** only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has prostate cancer; and
- 2 Patient has metastases; and
- 3 Patient's disease is castration resistant; and
- 4 Fither:
  - 4.1 All of the following:
    - 4.1.1 Patient is symptomatic; and
    - 4.1.2 Patient has disease progression (rising serum PSA) after second line anti-androgen therapy; and
    - 4.1.3 Patient has ECOG performance score of 0-1; and
    - 4.1.4 Patient has not had prior treatment with taxane chemotherapy; or
  - 4.2 All of the following:
    - 4.2.1 Patient's disease has progressed following prior chemotherapy containing a taxane; and
    - 4.2.2 Patient has ECOG performance score of 0-2; and
    - 4.2.3 Patient has not had prior treatment with abiraterone.

Renewal — (abiraterone acetate) only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

1 Significant decrease in serum PSA from baseline; and

Subsidy (Manufacturer's Pric			Brand or Generic	
\$	Per	Subsidised	Manufacturer	

#### continued...

- 2 No evidence of clinical disease progression; and
- 3 No initiation of taxane chemotherapy with abiraterone; and
- 4 The treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

#### All of the following:

- 1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 2 Abiraterone acetate to be discontinued at progression; and
- 3 No initiation of taxane chemotherapy with abiraterone; and
- 4 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

# BICALUTAMIDE

Tab 50 mg4.18	28	✓ Binarex
FLUTAMIDE		
Tab 250 mg107.55	90	✓ Prostacur S29
119.50	100	✓ Flutamin
FULVESTRANT - Retail pharmacy-Specialist - Special Authority see SA1895	5 below	
Inj 50 mg per ml, 5 ml prefilled syringe1,068.00		✓ Faslodex

#### **⇒SA1895** Special Authority for Subsidy

**Initial application** only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

#### All of the following:

- 1 Patient has oestrogen-receptor positive locally advanced or metastatic breast cancer; and
- 2 Patient has disease progression following prior treatment with an aromatase inhibitor or tamoxifen for their locally advanced or metastatic disease; and
- 3 Treatment to be given at a dose of 500 mg monthly following loading doses; and
- 4 Treatment to be discontinued at disease progression.

**Renewal** only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

#### All of the following:

- 1 Treatment remains appropriate and patient is benefitting from treatment; and
- 2 Treatment to be given at a dose of 500 mg monthly; and
- 3 There is no evidence of disease progression.

## OCTREOTIDE

OCTRECTIDE			
Inj 50 mcg per ml, 1 ml vial	27.58	5	✓ Omega S29
Inj 100 mcg per ml, 1 ml vial	48.50	5	✓ Omega S29
Inj 500 mcg per ml, 1 ml vial	113.10	5	✓ Omega S29
Inj 50 mcg per ml, 1 ml ampoule		5	✓ Max Health
			✓ Octreotide GH S29
Inj 100 mcg per ml, 1 ml ampoule	32.71	5	✓ Max Health
			✓ Octreotide GH S29
			✓ Sun Pharma S29
Inj 500 mcg per ml, 1 ml ampoule	113.10	5	✓ Max Health
			✓ Octreotide GH S29
			✓ Sun Pharma S29
OCTREOTIDE LONG-ACTING - Special Authority see SA211	19 on the next page	- Retail pha	rmacy
Inj depot 10 mg prefilled syringe		1 '	Sandostatin LAR
Inj depot 20 mg prefilled syringe	583.70	1	✓ Sandostatin LAR

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Inj depot 30 mg prefilled syringe.......670.80

Sandostatin LAR

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
•	Por 🗸	Manufacturor

## ⇒SA2119 Special Authority for Subsidy

**Initial application** — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 The patient has nausea\* and vomiting\* due to malignant bowel obstruction\*; and
- 2 Treatment with antiemetics, rehydration, antimuscarinic agents, corticosteroids and analgesics for at least 48 hours has failed; and
- 3 Octreotide to be given at a maximum dose 1500 mcg daily for up to 4 weeks.

Note: Indications marked with \* are unapproved indications.

**Renewal — (Malignant Bowel Obstruction)** from any relevant practitioner. Approvals valid for 3 months where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:
Both:

- 1 The patient has acromegaly; and
- 2 Any of the following:
  - 2.1 Treatment with surgery, radiotherapy and a dopamine agonist has failed; or
  - 2.2 Treatment with octreotide is for an interim period while awaiting the effects of radiotherapy and a dopamine agonist has failed; or
  - 2.3 The patient is unwilling, or unable, to undergo surgery and/or radiotherapy.

Renewal — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 IGF1 levels have decreased since starting octreotide; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: In patients with Acromegaly octreotide treatment should be discontinued if IGF1 levels have not decreased after 3 months treatment. In patients treated with radiotherapy octreotide treatment should be withdrawn every 2 years, for 1 month, for assessment of remission. Octreotide treatment should be stopped where there is biochemical evidence of remission (normal IGF1 levels) following octreotide treatment withdrawal for at least 4 weeks

**Renewal — (Acromegaly - pandemic circumstances)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has acromedaly; and
- 2 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 3 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

Initial application — (Other Indications) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 VIPomas and Glucagonomas for patients who are seriously ill in order to improve their clinical state prior to definitive surgery; or
- 2 Both:
  - 2.1 Gastrinoma: and
  - 2.2 Either:
    - 2.2.1 Patient has failed surgery; or
    - 2.2.2 Patient in metastatic disease after H2 antagonists (or proton pump inhibitors) have failed; or
- 3 Both:
  - 3.1 Insulinomas: and

60

✓ <u>Tamoxifen Sandoz</u>
✓ <u>Tamoxifen Sandoz</u>

✓ Letrole

Subsidy		Fully	Brand or	
(Manufacturer's Price)	5	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 3.2 Surgery is contraindicated or has failed; or
- 4 For pre-operative control of hypoglycaemia and for maintenance therapy; or
- 5 Both:
  - 5.1 Carcinoid syndrome (diagnosed by tissue pathology and/or urinary 5HIAA analysis); and
  - 5.2 Disabling symptoms not controlled by maximal medical therapy.

Note: The use of octreotide in patients with fistulae, oesophageal varices, miscellaneous diarrhoea and hypotension will not be funded as a Special Authority item

Renewal — (Other Indications) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment. Initial application — (pre-operative acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has acromegaly; and
- 2 Patient has a large pituitary tumour, greater than 10 mm at its widest; and
- 3 Patient is scheduled to undergo pituitary surgery in the next six months.

ΙA	MOXIFEN CITRATE	
*	Tab 10 mg	.00

Tah 20 ma

* Tab 20 Hig	00	Tallioxileti Salidoz
Aromatase Inhibitors		
ANASTROZOLE	30	✓ Anatrole
EXEMESTANE	30	✓ Pfizer Exemestane
LETROZOLE		

#### **Immunosuppressants**

# Cytotoxic Immunosuppressants

AZATHIOPRINE	
* Tab 25 mg7.36	0 <b>✓ Azamun</b>
* Tab 50 mg8.10	00 <b>Azamun</b>
MYCOPHENOLATE MOFETIL	
Tab 500 mg35.90	0 ✓ Cellcept
Cap 250 mg35.90	00 Cellcept
Powder for oral liq 1 g per 5 ml - Subsidy by endorsement 187.25	nl OP   Cellcept

Mycophenolate powder for oral liquid is subsidised only for patients unable to swallow tablets and capsules, and when the prescription is endorsed accordingly.

#### **Fusion Proteins**

ETANERCEPT - Special Authority see SA2399 on the next page	- Retail pharmad	су	
Inj 25 mg	690.00	4	<ul><li>Enbrel</li></ul>
Inj 25 mg autoinjector	690.00	4	<ul><li>Enbrel</li></ul>
Inj 50 mg autoinjector	1,050.00	4	<ul><li>Enbrel</li></ul>
Inj 50 mg prefilled syringe		4	<ul><li>Enbrel</li></ul>

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic

\$ Per ✔ Manufacturer

⇒SA2399 Special Authority for Subsidy

Initial application — (adult-onset Still's disease) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

#### Fither:

- 1 Both:
  - 1.1 Either:
    - 1.1.1 The patient has had an initial Special Authority approval for adalimumab for adult-onset Still's disease (AOSD); or
    - 1.1.2 The patient has been started on tocilizumab for AOSD in a Health NZ Hospital; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or tocilizumab; or
    - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or tocilizumab such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
  - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
  - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal anti-inflammatory drugs (NSAIDs) and methotrexate; and
- 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
  - 1.1 Applicant is a rheumatologist; or
  - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 The patient has a sustained improvement in inflammatory markers and functional status.

**Initial application** — **(ankylosing spondylitis)** only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

#### Fither:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for ankylosing spondylitis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for ankylosing spondylitis; or
- 2 All of the following:
  - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis present for more than six months; and
  - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
  - 2.3 Patient has bilateral sacroiliitis demonstrated by plain radiographs, CT or MRI scan; and
  - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
  - 2.5 Either:
    - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender (see Notes); and
- 2.6 A Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of initial application.

Average normal chest expansion corrected for age and gender:

18-24 years - Male: 7.0 cm; Female: 5.5 cm 25-34 years - Male: 7.5 cm; Female: 5.5 cm 35-44 years - Male: 6.5 cm; Female: 4.5 cm 45-54 years - Male: 6.0 cm; Female: 5.0 cm 55-64 years - Male: 5.5 cm; Female: 4.0 cm 65-74 years - Male: 4.0 cm; Female: 4.0 cm

75+ years - Male: 3.0 cm; Female: 2.5 cm

**Renewal — (ankylosing spondylitis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

#### All of the following:

- 1 Fither:
  - 1.1 Applicant is a rheumatologist; or
  - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 3 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 4 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (polyarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for polyarticular course JIA; or
- 2 All of the following:
  - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
  - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
  - 2.3 Any of the following:
    - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
    - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
    - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Renewal — (polyarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidi	sed	Generic
\$	Per	1	

continued...

criteria:

#### Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
  - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

## Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for oligoarticular course juvenile idiopathic arthritis (JIA); and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for oligoarticular course JIA; or
- 2 All of the following:
  - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
  - 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
  - 2.3 Any of the following:
    - 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
    - 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose): or
    - 2.3.3 High disease activity (cJADAS10 score greater than 4) after a 6-month trial of methotrexate.

Renewal — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

#### Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Fither:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
  - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

## Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab or secukinumab for psoriatic arthritis; and
  - 1.2 Either
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab or secukinumab; or

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsidised		Generic
\$	Per	✓	Manufacturer

continued...

- 1.2.2 The patient has received insufficient benefit from adalimumab or secukinumab to meet the renewal criteria for adalimumab or secukinumab for psoriatic arthritis; or
- 2 All of the following:
  - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
  - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
  - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
  - 2.4 Fither:
    - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
    - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
  - 2.5 Any of the following:
    - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
    - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
    - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

**Renewal — (psoriatic arthritis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Either:
  - 1.1 Applicant is a rheumatologist; or
  - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

**Initial application — (pyoderma gangrenosum)** only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum\*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Indications marked with \* are unapproved indications.

**Renewal — (pyoderma gangrenosum)** only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

Initial application — (Arthritis - rheumatoid) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

#### Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for rheumatoid arthritis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects; or
    - 1.2.2 The patient has received insufficient benefit to meet the renewal criteria for rheumatoid arthritis; or
- 2 All of the following:
  - 2.1 Patient has had rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
  - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
  - 2.3 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
  - 2.4 Patient has tried and not responded to at least three months of methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses unless contraindicated); and
  - 2.5 Either:
    - 2.5.1 Patient has tried and not responded to at least three months of methotrexate in combination with the maximum tolerated dose of ciclosporin; or
    - 2.5.2 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with methotrexate: and
  - 2.6 Either:
    - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
    - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip.

Renewal — (Arthritis - rheumatoid) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
  - 2.1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

**Initial application** — (severe chronic plaque psoriasis) only from a dermatologist or any relevant practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria: Fither:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe chronic plague psoriasis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe chronic plaque psoriasis; or
- 2 All of the following:

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
<u> </u>	Per 🗸	Manufacturer	

continued...

- 2.1 Any of the following:
  - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
  - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; or
  - 2.1.3 Patient has severe chronic localised genital or flexural plaque psoriasis where the plaques or lesions have been present for at least 6 months from the time of initial diagnosis, and with a Dermatology Life Quality Index (DLQI) score greater than 10; and
- 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
- 2.3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 2.4 The most recent PASI or DLQI assessment is no more than 1 month old at the time of application.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand, foot, genital or flexural areas at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and for the face, palm of a hand or sole of a foot the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 Both:
    - 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
    - 1.1.2 Either:
      - 1.1.2.1 Following each prior etanercept treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-treatment baseline value; or
      - 1.1.2.2 Following each prior etanercept treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline value; or
  - 1.2 Both:
    - 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
    - 1.2.2 Either:
      - 1.2.2.1 Following each prior etanercept treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
      - 1.2.2.2 Following each prior etanercept treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; or
  - 1.3 Both:
    - 1.3.1 Patient had severe chronic localised genital or flexural plaque psoriasis at the start of treatment; and
    - 1.3.2 Either:
      - 1.3.2.1 The patient has experienced a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1.3.2.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing etanercept; and
- 2 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Note: A treatment course is defined as a minimum of 12 weeks of etanercept treatment

Initial application — (undifferentiated spondyloarthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undifferentiated peripheral spondyloarthritis\* with active peripheral joint arthritis in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day (or maximum tolerated dose); and
- 4 Patient has tried and not responded to at least three months of leflunomide at a dose of up to 20 mg daily (or maximum tolerated dose); and
- 5 Any of the following:
  - 5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
  - 5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour measured no more than one month prior to the date of this application; or
  - 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Note: Indications marked with \* are unapproved indications.

Renewal — (undifferentiated spondyloarthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 Applicant is a rheumatologist; or
  - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg dose every 7 days.

### Immune Modulators

ANTITHYMOCYTE GLOBULIN (EQUINE) - PCT only - Spec	cialist		
Inj 50 mg per ml, 5 ml	2,774.48	5	✓ ATGAM
BACILLUS CALMETTE-GUERIN (BCG) VACCINE - PCT on	ly – Specialist		
Subsidised only for bladder cancer.			
Inj 2-8 × 100 million CFU	149.37	1	✓ OncoTICE
Inj 40 mg per ml, vial	176.90	3	✓ SII-Onco-BCG S29

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	Subsidised	Generic	
\$	Per	✓	Manufacturer	

#### Monoclonal Antibodies

	асу	ee SA2400 below – Retail pharm	ADALIMUMAB (AMGEVITA) – Special Authority see S
✓ Amgevita	1	190.00	Inj 20 mg per 0.4 ml prefilled syringe
✓ Amgevita	2	375.00	Inj 40 mg per 0.8 ml prefilled pen
✓ Amgevita	2	375.00	Ini 40 mg per 0.8 ml prefilled syringe

#### ⇒SA2400 Special Authority for Subsidy

Initial application — (Behcet's disease - severe) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient has severe Behcet's disease\* that is significantly impacting the patient's quality of life; and
- 2 Either:
  - 2.1 The patient has severe ocular, neurological, and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s); or
  - 2.2 The patient has severe gastrointestinal, rheumatological, and/or mucocutaneous symptoms and has not responded adequately to two or more treatments appropriate for the particular symptom(s).

Note: Indications marked with \* are unapproved indications.

Initial application — (Hidradenitis suppurativa) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has hidradenitis suppurativa Hurley Stage II or Hurley Stage III lesions in distinct anatomic areas; and
- 2 Patient has tried, but had an inadequate response to at least a 90 day trial of systemic antibiotics or has demonstrated intolerance to or has contraindications for systemic antibiotics; and
- 3 Patient has 3 or more active lesions; and
- 4 The patient has a DLQI of 10 or more and the assessment is no more than 1 month old at time of application.

**Renewal — (Hidradenitis suppurativa)** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline; and
- 2 The patient has a DLQI improvement of 4 or more from baseline.

Initial application — (Plaque psoriasis - severe chronic) only from a dermatologist or any relevant practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria: Either:

- 1 Both:
  - 1.1 Patient has had an initial Special Authority approval for etanercept for severe chronic plaque psoriasis; and
  - 1.2 Fither:
    - 1.2.1 Patient has experienced intolerable side effects; or
    - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for etanercept for severe chronic plaque psoriasis; or
- 2 All of the following:
  - 2.1 Any of the following:
    - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a PASI score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
    - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; or
    - 2.1.3 Patient has severe chronic localised genital or flexural plaque psoriasis where the plaques or lesions have been present for at least 6 months from the time of initial diagnosis, and with a Dermatology Life Quality

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic	
\$	Per	✓	Manufacturer	

continued...

Index (DLQI) score greater than 10; and

- 2.2 Patient has tried, but had an inadequate response to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
- 2.3 A PASI assessment or DLQI assessment has been completed for at least the most recent prior treatment course but no longer than 1 month following cessation of each prior treatment course and is no more than 1 month old at the time of application.

Renewal — (Plaque psoriasis - severe chronic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 Both:
  - 1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced a 75% or more reduction in PASI score, or is sustained at this level, when compared with the pre-treatment baseline value; or
    - 1.2.2 The patient has a DLQI improvement of 5 or more, when compared with the pre-treatment baseline value; or

#### 2 Both:

- 2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
- 2.2 Either:
  - 2.2.1 The patient has experienced reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
  - 2.2.2 The patient has experienced reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre treatment baseline value; or
- 3 Both:
  - 3.1 Patient had severe chronic localised genital or flexural plaque psoriasis at the start of treatment; and
  - 3.2 Fither:
    - 3.2.1 The patient has experienced a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; or
    - 3.2.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing adalimumab.

**Initial application — (pyoderma gangrenosum)** only from a dermatologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has pyoderma gangrenosum\*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporin, azathioprine, or methotrexate) and has not received an adequate response.

Note: Indications marked with \* are unapproved indications.

Initial application — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active Crohn's disease: and
- 2 Any of the following:
  - 2.1 Patient has a CDAI score of greater than or equal to 300, or HBI score of greater than or equal to 10; or
  - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or

Subsidy (Manufacturer's Price)	Fully Subsidised		
\$	Per 🗸	Manufacturer	

continued...

- 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
- 2.4 Patient has an ileostomy or colostomy and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Renewal — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 CDAI score has reduced by 100 points from the CDAI score, or HBI score has reduced by 3 points, from when the patient was initiated on adalimumab; or
- 2 CDAI score is 150 or less, or HBI is 4 or less; or
- 3 The patient has demonstrated an adequate response to treatment, but CDAI score and/or HBI score cannot be assessed.

**Initial application — (Crohn's disease - children)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has active Crohn's disease; and
- 2 Either:
  - 2.1 Patient has a PCDAI score of greater than or equal to 30; or
  - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Renewal — (Crohn's disease - children) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab; or
- 2 PCDAI score is 15 or less: or
- 3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed.

**Initial application** — **(Crohn's disease - fistulising)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Crohn's disease: and
- 2 Any of the following:
  - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
  - 2.2 Patient has one or more rectovaginal fistula(e); or
  - 2.3 Patient has complex peri-anal fistula; and
- 3 A Baseline Fistula Assessment has been completed and is no more than 1 month old at the time of application.

Renewal — (Crohn's disease - fistulising) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 The number of open draining fistulae have decreased from baseline by at least 50%; or
- 2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain.

**Initial application — (Ocular inflammation - chronic)** from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 The patient has had an initial Special Authority approval for infliximab for chronic ocular inflammation; or
- 2 Both:

Subsidy (Manufacturer's Price)	Subsi	Fully idised	Brand or Generic
\$	Per	/	Manufacturer

continued...

- 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
- 2.2 Any of the following:
  - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective; or
  - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
  - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

**Renewal — (Ocular inflammation - chronic)** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 12 weeks' initial treatment; or
- 2 Following each 2 year treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 2 year treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

**Initial application — (Ocular inflammation - severe)** from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria: Fither:

- 1 Patient has had an initial Special Authority approval for infliximab for severe ocular inflammation; or
- 2 Both:
  - 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
  - 2.2 Any of the following:
    - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
    - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
    - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Renewal — (Ocular inflammation - severe) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 2 year treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
- 3 Following each 2 year treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

**Initial application — (ankylosing spondylitis)** only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
  - 1.1 Patient has had an initial Special Authority approval for etanercept for ankylosing spondylitis; and
  - 12 Fither
    - 1.2.1 The patient has experienced intolerable side effects; or

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	Subsidised	Generic	
\$	Per	1	Manufacturer	

continued...

- 1.2.2 The patient has received insufficient benefit to meet the renewal criteria for ankylosing spondylitis; or
- 2 All of the following:
  - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis for more than six months; and
  - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
  - 2.3 Patient has bilateral sacroiliitis demonstrated by radiology imaging; and
  - 2.4 Patient has not responded adequately to treatment with two or more NSAIDs, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
  - 2.5 Fither:
    - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following BASMI measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
    - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender; and
  - 2.6 A BASDAI of at least 6 on a 0-10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment and is no more than 1 month old at the time of application.

Renewal — (ankylosing spondylitis) from any relevant practitioner. Approvals valid for 2 years where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less.

Initial application — (Arthritis - oligoarticular course juvenile idiopathic) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for etanercept for oligoarticular course juvenile idiopathic arthritis (JIA); and
  - 1.2 Either:
    - 1.2.1 Patient has experienced intolerable side effects; or
    - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for oligoarticular course JIA; or
- 2 All of the following:
  - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
  - 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
  - 2.3 Either:
    - 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
    - 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose).

**Renewal — (Arthritis - oligoarticular course juvenile idiopathic)** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - polyarticular course juvenile idiopathic) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

1 Both:

_			
	Subsidy	Fully	Brand or
	(Manufacturer's Price)	Subsidised	Generic
	\$	Per 🗸	Manufacturer

continued...

- 1.1 Patient has had an initial Special Authority approval for etanercept for polyarticular course juvenile idiopathic arthritis (JIA): and
- 1.2 Either:
  - 1.2.1 Patient has experienced intolerable side effects; or
  - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for polyarticular course JIA: or
- 2 All of the following:
  - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance: and
  - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
  - 2.3 Any of the following:
    - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
    - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
    - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

**Renewal — (Arthritis - polyarticular course juvenile idiopathic)** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

- Either:
  - 1 Following initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
  - 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

**Initial application — (Arthritis - psoriatic)** only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

#### Either:

- 1 Both:
  - 1.1 Patient has had an initial Special Authority approval for etanercept or secukinumab for psoriatic arthritis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects; or
    - 1.2.2 The patient has received insufficient benefit from to meet the renewal criteria for psoriatic arthritis; or
- 2 All of the following:
  - 2.1 Patient has had active psoriatic arthritis for six months duration or longer; and
  - 2.2 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
  - 2.3 Patient has tried and not responded to at least three months of sulfasalazine or leflunomide at maximum tolerated doses (unless contraindicated); and
  - 2.4 Either:
    - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
    - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
  - 2.5 Any of the following:
    - 2.5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application: or
    - 2.5.2 Patient has an ESR greater than 25 mm per hour; or
    - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

Renewal — (Arthritis - psoriatic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in swollen joint count from baseline and a clinically significant response in the opinion of the physician; or
- 2 Patient demonstrates at least a continuing 30% improvement in swollen joint count from baseline and a clinically significant response in the opinion of the treating physician.

Initial application — (Arthritis - rheumatoid) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for etanercept for rheumatoid arthritis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects; or
    - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for rheumatoid arthritis: or
- 2 All of the following:
  - 2.1 Patient has had rheumatoid arthritis (either confirmed by radiology imaging, or the patient is CCP antibody positive) for six months duration or longer; and
  - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance: and
  - 2.3 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated): and
  - 2.4 Patient has tried and not responded to at least three months of methotrexate in combination with sulfasalazine and hydroxychloroguine sulphate at maximum tolerated doses (unless contraindicated); and
  - 2.5 Either:
    - 2.5.1 Patient has tried and not responded to at least three months of methotrexate in combination with the maximum tolerated dose of ciclosporin; or
    - 2.5.2 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with methotrexate: and
  - 2.6 Either:
    - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
    - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip.

Renewal — (Arthritis - rheumatoid) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician: or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Initial application — (Still's disease - adult-onset (AOSD)) only from a rheumatologist. Approvals valid without further renewal unless notified for applications meeting the following criteria: Fither:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for etanercept and/or tocilizumab for AOSD: and
  - 12 Fither

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subs	sidised	Generic	
\$	Per	•	Manufacturer	

continued...

- 1.2.1 Patient has experienced intolerable side effects from etanercept and/or tocilizumab; or
- 1.2.2 Patient has received insufficient benefit from at least a three-month trial of etanercept and/or tocilizumab; or
- 2 All of the following:
  - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria; and
  - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, NSAIDs and methotrexate: and
  - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

**Initial application — (ulcerative colitis)** from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active ulcerative colitis: and
- 2 Either:
  - 2.1 Patient's SCCAI score is greater than or equal to 4; or
  - 2.2 Patient's PUCAI score is greater than or equal to 20; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from prior therapy with immunomodulators and systemic corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 The SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on biologic therapy; or
- 2 The PUCAI score has reduced by 10 points or more from the PUCAI score when the patient was initiation on biologic therapy.

Initial application — (undifferentiated spondyloarthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undifferentiated peripheral spondyloarthritis\* with active peripheral joint arthritis in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2 Patient has tried and not responded to at least three months of each of methotrexate, sulfasalazine and leflunomide, at maximum tolerated doses (unless contraindicated); and
- 3 Any of the following:
  - 3.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
  - 3.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application; or
  - 3.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Note: Indications marked with \* are unapproved indications

Renewal — (undifferentiated spondyloarthritis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Fither:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response in the opinion of the treating physician.

Initial application — (inflammatory bowel arthritis – axial) only from a rheumatologist. Approvals valid for 6 months for

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
 \$	Per 🗸	Manufacturer

continued...

applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has axial inflammatory pain for six months or more; and
- 3 Patient is unable to take NSAIDs; and
- 4 Patient has unequivocal sacroiliitis demonstrated by radiological imaging or MRI; and
- 5 Patient has not responded adequately to prior treatment consisting of at least 3 months of an exercise regime supervised by a physiotherapist; and
- 6 A BASDAI of at least 6 on a 0-10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment.

Renewal — (inflammatory bowel arthritis – axial) from any relevant practitioner. Approvals valid for 2 years where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less.

Initial application — (inflammatory bowel arthritis – peripheral) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has active arthritis in at least four joints from the following: hip, knee, ankle, subtalar, tarsus, forefoot, wrist, elbow, shoulder, sternoclavicular; and
- 3 Patient has tried and not experienced a response to at least three months of methotrexate, or azathioprine at a maximum tolerated dose (unless contraindicated); and
- 4 Patient has tried and not experienced a response to at least three months of sulfasalazine at a maximum tolerated dose (unless contraindicated); and
- 5 Any of the following:
  - 5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
  - 5.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application: or
  - 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

**Renewal — (inflammatory bowel arthritis – peripheral)** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 Patient has experienced at least a continuing 30% improvement in active joint count from baseline in the opinion of the treating physician.

ADALIMUMAB (HUMIRA - ALTERNATIVE BRAND) - Special Authority see SA2157 below - Retail pharmacy

Inj 20 mg per 0.2 ml prefilled syringe	1,599.96	2	✓ Humira
Inj 40 mg per 0.4 ml prefilled pen	1,599.96	2	✓ HumiraPen
Inj 40 mg per 0.4 ml prefilled syringe	1,599.96	2	<ul><li>Humira</li></ul>

⇒SA2157 Special Authority for Subsidy

**Initial application** — (Behcet's disease – severe) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks

Subsidy		Fully	Brand or	_
(Manufacturer's Price)	)	Subsidised	Generic	
\$	Per	1	Manufacturer	

continued...

treatment: or

- 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Behcet's disease – severe) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient has had a good clinical response to treatment with measurably improved quality of life; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Hidradenitis suppurativa) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 7 days. Fortnightly dosing has been considered.

**Renewal — (Hidradenitis suppurativa)** only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline; and
- 2 The patient has a Dermatology Quality of Life Index improvement of 4 or more from baseline; and
- 3 Adalimumab is to be administered at doses no greater than 40mg every 7 days. Fortnightly dosing has been considered.

Initial application — (Psoriasis - severe chronic plaque) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:
All of the following:

- 1 Either:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Psoriasis - severe chronic plaque) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 Both:
    - 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
    - 1.1.2 Fither:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1.1.2.1 Following each prior adalimumab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-adalimumab treatment baseline value; or
- 1.1.2.2 Following each prior adalimumab treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline value; or
- 1.2 Both:
  - 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
  - 1.2.2 Either:
    - 1.2.2.1 Following each prior adalimumab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
    - 1.2.2.2 Following each prior adalimumab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-adalimumab treatment baseline value: and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

**Initial application — (Pyoderma gangrenosum)** only from a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 A maximum of 8 doses.

**Renewal — (Pyoderma gangrenosum)** only from a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient has demonstrated clinical improvement and continues to require treatment; and
- 2 A maximum of 8 doses.

Initial application — (Crohn's disease - adult) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevitat; or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
  - 1.3 Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Crohn's disease - adult) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

continued...

Both:

- 1 Any of the following:
  - 1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on adalimumab; or
  - 1.2 CDAI score is 150 or less; or
  - 1.3 The patient has demonstrated an adequate response to treatment, but CDAI score cannot be assessed; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Crohn's disease - children) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
  - 1.3 Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Crohn's disease - children) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab; or
  - 1.2 PCDAI score is 15 or less: or
  - 1.3 The patient has demonstrated an adequate response to treatment, but PCDAI score cannot be assessed; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Crohn's disease - fistulising) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
  - 1.3 Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Crohn's disease - fistulising) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
  - 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
  - 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain; and

Subsidy		Fully	Brand or	
(Manufacturer's Price	Sı	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Ocular inflammation – chronic) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with Amgevita, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
  - 1.3 Patient has uveitis and is considered to be at risk of vision loss if they were to change treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Ocular inflammation – chronic) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 The patient has had a good clinical response following 12 weeks' initial treatment; or
  - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < 1/2+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
  - 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Ocular inflammation – severe) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with Amgevita, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
  - 1.3 Patient has uveitis and is considered to be at risk of vision loss if they were to change treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Ocular inflammation – severe) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 The patient has had a good clinical response following 3 initial doses; or
  - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < 1/2+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
  - 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Initial application — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita); and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

**Renewal — (ankylosing spondylitis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Arthritis – oligoarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

Renewal — (Arthritis – oligoarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months where the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - polyarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

Renewal — (Arthritis - polyarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months where the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

Initial application — (Arthritis - psoriatic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Arthritis - psoriatic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Arthritis – rheumatoid) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Either:
  - 4.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or
  - 4.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

**Renewal — (Arthritis – rheumatoid)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Roth:

- 1 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 2 Fither
  - 2.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or
  - 2.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

Initial application — (Still's disease – adult-onset (AOSD)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or

Subsidy	Full	y Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per •	<ul> <li>Manufacturer</li> </ul>	

continued...

- 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

Renewal — (Still's disease – adult-onset (AOSD)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months where the patient has demonstrated a sustained improvement in inflammatory markers and functional status.

AFLIBERCEPT - Special Authority see SA1772 below - Retail pharmacy

## ⇒SA1772 Special Authority for Subsidy

**Initial application — (wet age related macular degeneration)** only from an ophthalmologist. Approvals valid for 3 months for applications meeting the following criteria:

#### Either:

- 1 All of the following:
  - 1.1 Any of the following:
    - 1.1.1 Wet age-related macular degeneration (wet AMD); or
    - 1.1.2 Polypoidal choroidal vasculopathy; or
    - 1.1.3 Choroidal neovascular membrane from causes other than wet AMD; and
  - 1.2 Either:
    - 1.2.1 The patient has developed severe endophthalmitis or severe posterior uveitis following treatment with bevacizumab; or
    - 1.2.2 There is worsening of vision or failure of retina to dry despite three intraocular injections of bevacizumab four weeks apart; and
  - 1.3 There is no structural damage to the central fovea of the treated eye; and
  - 1.4 Patient has not previously been treated with ranibizumab for longer than 3 months; or
- 2 Either:
  - 2.1 Patient has current approval to use ranibizumab for treatment of wAMD and was found to be intolerant to ranibizumab within 3 months; or
  - 2.2 Patient has previously\* (\*before June 2018) received treatment with ranibizumab for wAMD and disease was stable while on treatment.

Initial application — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has centre involving diabetic macular oedema (DMO); and
- 2 Patient's disease is non responsive to 4 doses of intravitreal bevacizumab when administered 4-6 weekly; and
- 3 Patient has reduced visual acuity between 6/9 6/36 with functional awareness of reduction in vision; and
- 4 Patient has DMO within central OCT (ocular coherence tomography) subfield > 350 micrometers; and
- 5 There is no centre-involving sub-retinal fibrosis or foveal atrophy.

Renewal — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Documented benefit must be demonstrated to continue; and
- 2 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 3 There is no structural damage to the central fovea of the treated eye.

Renewal — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

All of the following:

- 1 There is stability or two lines of Snellen visual acuity gain; and
- 2 There is structural improvement on OCT scan (with reduction in intra-retinal cysts, central retinal thickness, and sub-retinal fluid); and
- 3 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 4 There is no centre-involving sub-retinal fibrosis or foveal atrophy; and
- 5 After each consecutive 12 months treatment with (2nd line anti-VEGF agent), patient has retrialled with at least one injection of bevacizumab and had no response.

BENRALIZUMAB - Special Authority see SA2151 below - Retail pharmacy

## **⇒SA2151** Special Authority for Subsidy

Initial application — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Patient must have a diagnosis of severe eosinophilic asthma documented by a respiratory physician or clinical immunologist; and
- 3 Conditions that mimic asthma eg. vocal cord dysfunction, central airway obstruction, bronchiolitis etc. have been excluded: and
- 4 Patient has a blood eosinophil count of greater than 0.5 x 10^9 cells/L in the last 12 months; and
- 5 Patient must be adherent to optimised asthma therapy including inhaled corticosteroids (equivalent to at least 1000 mcg per day of fluticasone propionate) plus long-acting beta-2 agonist, or budesonide/formoterol as part of the anti-inflammatory reliever therapy plus maintenance regimen, unless contraindicated or not tolerated; and
- 6 Fither:
  - 6.1 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral corticosteroids; or
  - 6.2 Patient has received continuous oral corticosteroids of at least the equivalent of 10 mg per day over the previous 3 months: and
- 7 Treatment is not to be used in combination with subsidised mepolizumab; and
- 8 Patient has an Asthma Control Test (ACT) score of 10 or less. Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 52 weeks after the first dose to assess response to treatment; and
- 9 Either:
  - 9.1 Patient has not previously received an anti-IL5 biological therapy for their severe eosinophilic asthma; or
  - 9.2 Both:
    - 9.2.1 Patient was refractory or intolerant to previous anti-IL5 biological therapy; and
    - 9.2.2 Patient was not eligible to continue treatment with previous anti-IL5 biological therapy and discontinued within 12 months of commencing treatment.

**Renewal — (Severe eosinophilic asthma)** only from a respiratory physician or clinical immunologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 Either:
  - 2.1 Exacerbations have been reduced from baseline by 50% as a result of treatment with benralizumab; or
  - 2.2 Reduction in continuous oral corticosteroid use by 50% or by 10 mg/day while maintaining or improving asthma control.

BRENTUXIMAB VEDOTIN - PCT only - Special Authority see SA2289 on the next page

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

#### ⇒SA2289 Special Authority for Subsidy

**Initial application** — (relapsed/refractory Hodgkin lymphoma) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 Both:
    - 1.1.1 Patient has relapsed/refractory CD30-positive Hodgkin lymphoma after two or more lines of chemotherapy; and
    - 1.1.2 Patient is ineligible for autologous stem cell transplant; or
  - 1.2 Both:
    - 1.2.1 Patient has relapsed/refractory CD30-positive Hodgkin lymphoma; and
    - 1.2.2 Patient has previously undergone autologous stem cell transplant; and
- 2 Patient has not previously received funded brentuximab vedotin; and
- 3 Response to brentuximab vedotin treatment is to be reviewed after a maximum of 6 treatment cycles; and
- 4 Brentuximab vedotin to be administered at doses no greater than 1.8 mg/kg every 3 weeks.

**Renewal — (relapsed/refractory Hodgkin lymphoma)** from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has achieved a partial or complete response to brentuximab vedotin after 6 treatment cycles; and
- 2 Treatment remains clinically appropriate and the patient is benefitting from treatment and treatment is being tolerated; and
- 3 Patient is to receive a maximum of 16 total cycles of brentuximab vedotin treatment.

**Initial application — (anaplastic large cell lymphoma)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has relapsed/refractory CD30-positive systemic anaplastic large cell lymphoma; and
- 2 Patient has an ECOG performance status of 0-1; and
- 3 Patient has not previously received brentuximab vedotin; and
- 4 Response to brentuximab vedotin treatment is to be reviewed after a maximum of 6 treatment cycles; and
- 5 Brentuximab vedotin to be administered at doses no greater than 1.8 mg/kg every 3 weeks.

Renewal — (anaplastic large cell lymphoma) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has achieved a partial or complete response to brentuximab vedotin after 6 treatment cycles; and
- 2 Treatment remains clinically appropriate and the patient is benefitting from treatment and treatment is being tolerated; and
- 3 Patient is to receive a maximum of 16 total cycles of brentuximab vedotin treatment.

CASIRIVIMAB AND IMDEVIMAB - [Xpharm] - Special Authority see \$A2096 below

⇒SA2096 Special Authority for Subsidy

**Initial application — (Treatment of profoundly immunocompromised patients)** from any relevant practitioner. Approvals valid for 2 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed (or probable) COVID-19; and
- 2 The patient is in the community with mild to moderate disease severity\*; and
- 3 Patient is profoundly immunocompromised\*\* and is at risk of not having mounted an adequate response to vaccination against COVID-19 or is unvaccinated; and
- 4 Patient's symptoms started within the last 10 days; and

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

- 5 Patient is not receiving high flow oxygen or assisted/mechanical ventilation; and
- 6 Casirivimab and imdevimab is to be administered at a maximum dose of no greater than 2,400 mg.

Notes: \* Mild to moderate disease severity as described on the Ministry of Health Website

\*\* Examples include B-cell depletive illnesses or patients receiving treatment that is B-Cell depleting.

CETUXIMAB - PCT only - Specialist - Special Authority see SA2401 below

Inj 5 mg per ml, 20 ml vial	364.00	1	Erbitux
Inj 5 mg per ml, 100 ml vial	1,820.00	1	✓ Erbitux
Inj 1 mg for ECP	3.82	1 mg	✓ Baxter

## ⇒SA2401 Special Authority for Subsidy

Initial application — (head and neck cancer, locally advanced) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, non-metastatic, squamous cell cancer of the head and neck; and
- 2 Cisplatin is contraindicated or has resulted in intolerable side effects; and
- 3 Patient has an ECOG performance score of 0-2; and
- 4 To be administered in combination with radiation therapy.

Initial application — (colorectal cancer, metastatic) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has metastatic colorectal cancer located on the left side of the colon (see Note); and
- 2 There is documentation confirming disease is RAS and BRAF wild-type; and
- 3 Patient has an ECOG performance score of 0-2; and
- 4 Patient has not received prior funded treatment with cetuximab; and
- 5 Either:
  - 5.1 Cetuximab is to be used in combination with chemotherapy; or
  - 5.2 Chemotherapy is determined to not be in the best interest of the patient based on clinician assessment.

**Renewal — (colorectal cancer, metastatic)** only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where there is no evidence of disease progression.

Note: Left-sided colorectal cancer comprises of the distal one-third of the transverse colon, the splenic flexure, the descending colon, the sigmoid colon, or the rectum.

## ⇒SA2269 Special Authority for Subsidy

**Initial application** only from a haematologist, paediatric haematologist or paediatric oncologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has not received prior chemotherapy for this condition; and
- 2 Patient has de novo CD33-positive acute myeloid leukaemia; and
- 3 Patient does not have acute promyelocytic leukaemia; and
- 4 Gemtuzumab ozogamicin will be used in combination with standard anthracycline and cytarabine (AraC); and
- 5 Patient is being treated with curative intent; and
- 6 Patient's disease risk has been assessed by cytogenetic testing to be good or intermediate; and
- 7 Patient must be considered eligible for standard intensive remission induction chemotherapy with standard anthracycline and cytarabine (AraC); and
- 8 Gemtuzumab ozogamicin to be funded for one course only (one dose at 3 mg per m² body surface area or up to 2 vials of 5 mg as separate doses).

Note: Acute myeloid leukaemia excludes acute promyelocytic leukaemia and acute myeloid leukaemia that is secondary to another haematological disorder (eg myelodysplasia or myeloproliferative disorder).

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer	
INFLIXIMAB - PCT only - Special Authority see SA2402 below					
Inj 100 mg	428.00	1	<b>✓</b> F	Remicade	
Inj 1 mg for ECP	4.40	1 mg	<b>√</b> E	Baxter	
Inj 100 mg	428.00	Per 1	✓ F	Remicade	

⇒SA2402 Special Authority for Subsidy

**Initial application — (Crohn's disease (adults))** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active Crohn's disease; and
- 2 Any of the following:
  - 2.1 Patient has a CDAI score of greater than or equal to 300 or HBI score of greater than or equal to 10; or
  - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
  - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
  - 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but has experienced an inadequate response to, or intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Renewal — (Crohn's disease (adults)) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 CDAI score has reduced by 100 points from the CDAI score, or HBI score has reduced by 3 points, from when the patient was initiated on infliximab; or
  - 1.2 CDAI score is 150 or less, or HBI is 4 or less; or
  - 1.3 The patient has demonstrated an adequate response to treatment but CDAI score and/or HBI score cannot be assessed: and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

**Initial application** — (**Crohn's disease (children)**) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has active Crohn's disease; and
- 2 Either:
  - 2.1 Patient has a PCDAI score of greater than or equal to 30; or
  - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but experienced an inadequate response to, or intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Renewal — (Crohn's disease (children)) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on infliximab; or
  - 1.2 PCDAI score is 15 or less; or
  - 1.3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

**Initial application** — **(Graft vs host disease)** from any relevant practitioner. Approvals valid without further renewal unless notified where patient has steroid-refractory acute graft vs. host disease of the gut.

**Initial application — (Pulmonary sarcoidosis)** from any relevant practitioner. Approvals valid without further renewal unless notified where patient has life-threatening pulmonary sarcoidosis diagnosed by a multidisciplinary team that is refractory to other treatments.

Initial application — (acute fulminant ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 weeks for applications meeting the following criteria:
Both:

- 1 Patient has acute, fulminant ulcerative colitis; and
- 2 Treatment with intravenous or high dose oral corticosteroids has not been successful.

Initial application — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and
- 2 Either:
  - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
  - 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

**Renewal — (ankylosing spondylitis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Following 12 weeks of infliximab treatment, BASDAI has improved by 4 or more points from pre-infliximab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 3 Infliximab to be administered at doses no greater than 5 mg/kg every 6-8 weeks.

**Initial application — (chronic ocular inflammation)** from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

#### Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for chronic ocular inflammation; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for chronic ocular inflammation; or

#### 2 Both:

- 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
- 2.2 Any of the following:
  - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective: or
  - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
  - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

**Renewal — (chronic ocular inflammation)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.</p>

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (fistulising Crohn's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has confirmed Crohn's disease; and
- 2 Any of the following:
  - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
  - 2.2 Patient has one or more rectovaginal fistula(e); or
  - 2.3 Patent has complex peri-anal fistula.

Renewal — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
  - 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline (in the case of adult patients, as demonstrated by a reduction in the Fistula Assessment score), together with less induration and patient reported pain; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (neurosarcoidosis) only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with neurosarcoiosis by a multidisciplinary team; and
- 2 Patient has CNS involvement; and
- 3 Patient has steroid-refractory disease; and
- 4 Either:
  - 4.1 IV cyclophosphamide has been tried; or
  - 4.2 Treatment with IV cyclophosphamide is clinically inappropriate.

Renewal — (neurosarcoidosis) only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria:

Either:

- 1 A withdrawal period has been tried and the patient has relapsed; or
- 2 All of the following:
  - 2.1 A withdrawal period has been considered but would not be clinically appropriate; and
  - 2.2 There has been a marked reduction in prednisone dose; and

Subsidy		Fully	Brand or
(Manufacturer's Price)	9	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 2.3 Either:
  - 2.3.1 There has been an improvement in MRI appearances; or
  - 2.3.2 Marked improvement in other symptomology.

Initial application — (plaque psoriasis) only from a dermatologist or any relevant practitioner on the recommendation of a dermatologist. Approvals valid for 3 months for applications meeting the following criteria:

Fither:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab, etanercept or secukinumab for severe chronic plaque psoriasis; and
  - 1.2 Either:
    - 1.2.1 Patient has experienced intolerable side effects from adalimumab, etanercept or secukinumab; or
    - 1.2.2 Patient has received insufficient benefit from adalimumab, etanercept or secukinumab to meet the renewal criteria for adalimumab, etanercept or secukinumab for severe chronic plaque psoriasis; or
- 2 All of the following:
  - 2.1 Any of the following:
    - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
    - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; or
    - 2.1.3 Patient has severe chronic localised genital or flexural plaque psoriasis where the plaques or lesions have been present for at least 6 months from the time of initial diagnosis, and with a Dermatology Life Quality Index (DLQI) score greater than 10; and
  - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
  - 2.3 A PASI assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course: and
  - 2.4 The most recent PASI assessment is no more than 1 month old at the time of initiation.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand, foot, genital or flexural areas at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and for the face, palm of a hand or sole of a foot the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (plaque psoriasis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 Both:
    - 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
    - 1.1.2 Following each prior infliximab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-infliximab treatment baseline value; or
  - 1.2 Both:
    - 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and

Subsidy (Manufacturer's Price)	s	Fully	Brand or Generic
 \$	Per	1	Manufacturer

continued...

- 1.2.2 Either:
  - 1.2.2.1 Following each prior infliximab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
  - 1.2.2.2 Following each prior infliximab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-infliximab treatment baseline value: or
- 1.3 Both:
  - 1.3.1 Patient had severe chronic localised genital or flexural plague psoriasis at the start of treatment; and
  - 1.3.2 Fither:
    - 1.3.2.1 The patient has experienced a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; or
    - 1.3.2.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing infliximab; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient was being treated with infliximab prior to 1 February 2019; and
- 2 Any of the following:
  - 2.1 Rheumatoid arthritis; or
  - 2.2 Ankylosing spondylitis: or
  - 2.3 Psoriatic arthritis; or
  - 2.4 Severe ocular inflammation: or
  - 2.5 Chronic ocular inflammation: or
  - 2.6 Crohn's disease (adults): or
  - 2.7 Crohn's disease (children); or
  - 2.8 Fistulising Crohn's disease; or
  - 2.9 Severe fulminant ulcerative colitis: or
  - 2.10 Severe ulcerative colitis; or
  - 2.11 Plaque psoriasis: or
  - 2.12 Neurosarcoidosis; or
  - 2.13 Severe Behcet's disease.

Initial application — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis: and
- 2 Either:
  - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept and/or secukinumab; or
  - 2.2 Following 3-4 months' initial treatment with adalimumab and/or etanercept and/or secukinumab, the patient did not meet the renewal criteria for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis.

**Renewal** — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

1 Fither

Subsidy	5	Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	1	Manufacturer

continued...

- 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior infliximab treatment in the opinion of the treating physician; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
  - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
  - 2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept; and
- 3 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance.

**Renewal — (rheumatoid arthritis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Infliximab to be administered at doses no greater than 3 mg/kg every 8 weeks.

**Initial application — (severe Behcet's disease)** from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe Behcet's disease which is significantly impacting the patient's quality of life (see Notes); and
- 2 Either:
  - 2.1 The patient has severe ocular, neurological and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s) (see Notes); or
  - 2.2 The patient has severe gastrointestinal, rheumatologic and/or mucocutaneous symptoms and has not responded adequately to two or more treatment appropriate for the particular symptom(s) (see Notes); and
- 3 The patient is experiencing significant loss of quality of life.

Notes: Behcet's disease diagnosed according to the International Study Group for Behcet's Disease. Lancet 1990;335(8697):1078-80. Quality of life measured using an appropriate quality of life scale such as that published in Gilworth et al J Rheumatol. 2004;31:931-7.

Treatments appropriate for the particular symptoms are those that are considered standard conventional treatments for these symptoms, for example intravenous/oral steroids and other immunosuppressants for ocular symptoms; azathioprine, steroids, thalidomide, interferon alpha and ciclosporin for mucocutaneous symptoms; and colchicine, steroids and methotrexate for rheumatological symptoms.

Renewal — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 Patient has had a good clinical response to initial treatment with measurably improved quality of life; and

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic	
<u> </u>	Per	✓	Manufacturer	

continued...

2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Renewal — (fulminant ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Where maintenance treatment is considered appropriate, infliximab should be used in combination with immunomodulators and reassessed every 6 months; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

**Initial application** — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe ocular inflammation; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe ocular inflammation; or
- 2 Both:
  - 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
  - 2.2 Any of the following:
    - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
    - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
    - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

**Renewal — (severe ocular inflammation)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

**Initial application — (ulcerative colitis)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active ulcerative colitis; and
- 2 Either:
  - 2.1 Patients SCCAI is greater than or equal to 4; or
  - 2.2 Patients PUCAI score is greater than or equal to 20; and
- 3 Patient has tried but has experienced an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and systemic corticosteroids.

Subsidy	Full	y Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per 🗸	<ul> <li>Manufacturer</li> </ul>	

continued...

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 The SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on infliximab: or
  - 1.2 The PUCAI score has reduced by 10 points or more from the PUCAI score when the patient was initiated on infliximab: and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

**Initial application — (pyoderma gangrenosum)** only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum\*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Note: Indications marked with \* are unapproved indications.

**Renewal — (pyoderma gangrenosum)** only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

Initial application — (inflammatory bowel arthritis – axial) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has had axial inflammatory pain for six months or more; and
- 3 Patient is unable to take NSAIDs; and
- 4 Patient has unequivocal sacroiliitis demonstrated by radiological imaging or MRI; and
- 5 Patient's disease has not responded adequately to prior treatment consisting of at least 3 months of an exercise regime supervised by a physiotherapist; and
- 6 Patient has a BASDAI of at least 6 on a 0 10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment.

Renewal — (inflammatory bowel arthritis – axial) from any relevant practitioner. Approvals valid for 2 years where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10-point scale, or an improvement in BASDAI of 50%, whichever is less.

Initial application — (inflammatory bowel arthritis – peripheral) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has active arthritis in at least four joints from the following: hip, knee, ankle, subtalar, tarsus, forefoot, wrist, elbow, shoulder, sternoclavicular; and
- 3 Patient has tried and not experienced a response to at least three months of methotrexate or azathioprine at a maximum

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per Manufacturer

continued...

tolerated dose (unless contraindicated): and

- 4 Patient has tried and not experienced a response to at least three months of sulfasalazine at a maximum tolerated dose (unless contraindicated); and
- 5 Any of the following:
  - 5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
  - 5.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application: or
  - 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (inflammatory bowel arthritis – peripheral) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, patient has experienced at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 Patient has experienced at least a continuing 30% improvement in active joint count from baseline in the opinion of the treating physician.

MEPOLIZUMAB - Special Authority see SA2331 below - Retail pharmacy

✓ Nucala

⇒SA2331 Special Authority for Subsidy

Initial application — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Patient must have a diagnosis of severe eosinophilic asthma documented by a respiratory physician or clinical immunologist: and
- 3 Conditions that mimic asthma eq. vocal cord dysfunction, central airway obstruction, bronchiolitis etc. have been
- 4 Patient has a blood eosinophil count of greater than 0.5 × 10<sup>9</sup> cells/L in the last 12 months; and
- 5 Patient must be adherent to optimised asthma therapy including inhaled corticosteroids (equivalent to at least 1000 mcg per day of fluticasone propionate) plus long acting beta-2 agonist, or budesonide/formoterol as part of the single maintenance and reliever therapy regimen, unless contraindicated or not tolerated; and
- - 6.1 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral corticosteroids: or
  - 6.2 Patient has received continuous oral corticosteroids of at least the equivalent of 10 mg per day over the previous
- 7 Treatment is not to be used in combination with subsidised benralizumab; and
- 8 Patient has an Asthma Control Test (ACT) score of 10 or less. Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 52 weeks after the first dose to assess response to treatment; and
- - 9.1 Patient has not previously received an anti-IL5 biological therapy for their severe eosinophilic asthma; or
  - 9.2 Both:
    - 9.2.1 Patient was refractory or intolerant to previous anti-IL5 biological therapy; and
    - 9.2.2 Patient was not eligible to continue treatment with previous anti-IL5 biological therapy and discontinued

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
 \$	Per	1	Manufacturer

continued...

within 12 months of commencing treatment.

Renewal — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 Either:
  - 2.1 Exacerbations have been reduced from baseline by 50% as a result of treatment with mepolizumab; or
  - 2.2 Reduction in continuous oral corticosteroid use by 50% or by 10 mg/day while maintaining or improving asthma control.

Initial application — (eosinophilic granulomatosis with polyangiitis) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has eosinophilic granulomatosis with polyangiitis; and
- 2 The patient has trialled and not received adequate benefit from at least one of the following for at least three months (unless contraindicated to all): azathioprine, cyclophosphamide, leflunomide, methotrexate, mycophenolate, or rituximab; and
- 3 Fither:
  - 3.1 The patient has trialled prednisone for a minimum of three months and is unable to maintain disease control at doses below 7.5 mg per day; or
  - 3.2 Corticosteroids are contraindicated.

**Renewal** — (eosinophilic granulomatosis with polyangiitis) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months where patient has no evidence of clinical disease progression.

OBINUTUZUMAB - PCT only - Specialist - Special Authori	ty see SA2155 below		
Inj 25 mg per ml, 40 ml vial	5,910.00	1	Gazyva
Inj 1 mg for ECP	6.21	1 mg	✓ Baxter

⇒SA2155 Special Authority for Subsidy

Initial application — (chronic lymphocytic leukaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C CD20+ chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is obinutuzumab treatment naive; and
- 3 The patient is not eligible for full dose FCR due to comorbidities with a score > 6 on the Cumulative Illness Rating Scale (CIRS) or reduced renal function (creatinine clearance < 70mL/min); and</p>
- 4 Patient has adequate neutrophil and platelet counts\* unless the cytopenias are a consequence of marrow infiltration by CLL; and
- 5 Patient has good performance status; and
- 6 Obinutuzumab to be administered at a maximum cumulative dose of 8,000 mg and in combination with chlorambucil for a maximum of 6 cycles.

Notes: Chronic lymphocytic leukaemia includes small lymphocytic lymphoma. Comorbidity refers only to illness/impairment other than CLL induced illness/impairment in the patient. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with obinutuzumab is expected to improve symptoms and improve ECOG score to < 2.

\* Neutrophil greater than or equal to  $1.5 \times 10^9$ /L and platelets greater than or equal to  $75 \times 10^9$ /L.

**Initial application — (follicular / marginal zone lymphoma)** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

continued...

All of the following:

- 1 Either:
  - 1.1 Patient has follicular lymphoma; or
  - 1.2 Patient has marginal zone lymphoma; and
- 2 Patient is refractory to or has relapsed within 12 months of a rituximab containing combined chemo-immunotherapy regimen\*: and
- 3 Patient has an ECOG performance status of 0-2; and
- 4 Patient has been previously treated with no more than four chemotherapy regimens; and
- 5 Obinutuzumab to be administered at a maximum dose of 1000 mg for a maximum of 6 cycles in combination with chemotherapy\*.

Note: \* includes unapproved indications

Renewal — (follicular / marginal zone lymphoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 24 months for applications meeting the following criteria: All of the following:

- 1 Patient has no evidence of disease progression following obinutuzumab induction therapy; and
- 2 Obinutuzumab to be administered at a maximum of 1000 mg every 2 months for a maximum of 2 years; and
- 3 Obinutuzumab to be discontinued at disease progression.

		OMALIZUMAB - Special Authority see SA1744 below - Retail pharmacy
✓ Xolair	1	Inj 150 mg prefilled syringe450.00
✓ Xolair AU ✓ Xolair	1	Inj 150 mg vial450.00

## ⇒SA1744 Special Authority for Subsidy

**Initial application** — (severe asthma) only from a respiratory specialist or clinical immunologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 6 years or older; and
- 2 Patient has a diagnosis of severe asthma; and
- 3 Past or current evidence of atopy, documented by skin prick testing or RAST; and
- 4 Total serum human immunoglobulin E (IgE) between 76 IU/mL and 1300 IU/ml at baseline; and
- 5 Proven adherence with optimal inhaled therapy including high dose inhaled corticosteroid (budesonide 1,600 mcg per day or fluticasone propionate 1,000 mcg per day or equivalent), plus long-acting beta-2 agonist therapy (at least salmeterol 50 mcg bd or eformoterol 12 mcg bd) for at least 12 months, unless contraindicated or not tolerated; and
- 6 Either:
  - 6.1 Patient has received courses of systemic corticosteroids equivalent to at least 28 days treatment in the past 12 months, unless contraindicated or not tolerated; or
  - 6.2 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral steroids; and
- 7 Patient has an Asthma Control Test (ACT) score of 10 or less; and
- 8 Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 26 weeks after the first dose to assess response to treatment.

Initial application — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Fither:

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

- 2.1 Both:
  - 2.1.1 Patient is symptomatic with Urticaria Activity Score 7 (UAS7) of 20 or above; and
  - 2.1.2 Patient has a Dermatology life quality index (DLQI) of 10 or greater; or
- 2.2 Patient has a Urticaria Control Test (UCT) of 8 or less; and
- 3 Any of the following:
  - 3.1 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and ciclosporin (> 3 mg/kg day) for at least 6 weeks; or
  - 3.2 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and at least 3 courses of systemic corticosteroids (> 20 mg prednisone per day for at least 5 days) in the previous 6 months; or
  - 3.3 Patient has developed significant adverse effects whilst on corticosteroids or ciclosporin; and
- 4 Fither:
  - 4.1 Treatment to be stopped if inadequate response\* following 4 doses; or
  - 4.2 Complete response\* to 6 doses of omalizumab.

**Renewal — (severe asthma)** only from a clinical immunologist or respiratory specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 A reduction in the maintenance oral corticosteroid dose or number of exacerbations of at least 50% from baseline.

Renewal — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Patient has previously adequately responded\* to 6 doses of omalizumab; or
- 2 Both:
  - 2.1 Patient has previously had a complete response\* to 6 doses of omalizumab; and
  - 2.2 Patient has relapsed after cessation of omalizumab therapy.

Note: \*Inadequate response defined as less than 50% reduction in baseline UAS7 and DLQI score, or an increase in Urticaria Control Test (UCT) score of less than 4 from baseline. Patient is to be reassessed for response after 4 doses of omalizumab. Complete response is defined as UAS7 less than or equal to 6 and DLQI less than or equal to 5; or UCT of 16. Relapse of chronic urticaria on stopping prednisone/ciclosporin does not justify the funding of omalizumab.

PALIVIZUMAB - PCT only - Special Authority see SA2419 below

#### ⇒SA2419 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

- 1 Palivizumab to be administered during the annual RSV season; and
- 2 Either:
  - 2.1 Both:
    - 2.1.1 Infant was born in the last 12 months; and
    - 2.1.2 Infant was born at less than 32 weeks zero days' gestation; or
  - 2.2 Both:
    - 2.2.1 Child was born in the last 24 months; and
    - 2.2.2 Any of the following:
      - 2.2.2.1 Child has severe lung, airway, neurological or neuromuscular disease that requires ongoing ventilatory/respiratory support (see Note A) in the community; or
      - 2.2.2.2 Both:
        - 2.2.2.2.1 Child has haemodynamically significant heart disease; and

(Man	Subsidy ufacturer's Price)	Fu Subsidis	ılly	Brand or Generic
(wich	. ,	Per	<b>✓</b>	Manufacturer

continued...

- 2.2.2.2.2 Any of the following:
  - 2.2.2.2.2.1 Child has unoperated simple congenital heart disease with significant left to right shunt (see Note B): or
  - 2.2.2.2.2.2 Child has unoperated or surgically palliated complex congenital heart disease; or
  - 2.2.2.2.3 Child has severe pulmonary hypertension (see Note C); or
  - 2.2.2.2.2.4 Child has moderate or severe left ventricular (LV) failure (see Note D); or
- 2.2.2.3 Child has severe combined immune deficiency, confirmed by an immunologist, but has not received a stem cell transplant: or
- 2.2.2.4 Child has inborn errors of immunity (see Note E) that increase susceptibility to life-threatening viral respiratory infections, confirmed by an immunologist.

Renewal from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Palivizumab to be administered during the annual RSV season; and
- 2 Child was born in the last 24 months; and
- 3 Any of the following:
  - 3.1 Child has severe lung, airway, neurological or neuromuscular disease that requires ongoing ventilatory/respiratory support (see Note A) in the community; or
  - 3.2 Both:
    - 3.2.1 Child has haemodynamically significant heart disease; and
    - 3.2.2 Any of the following:
      - 3.2.2.1 Child has unoperated simple congenital heart disease with significant left to right shunt (see Note B);
      - 3.2.2.2 Child has unoperated or surgically palliated complex congenital heart disease; or
      - 3.2.2.3 Child has severe pulmonary hypertension (see Note C); or
      - 3.2.2.4 Child has moderate or severe left ventricular (LV) failure (see Note D); or
  - 3.3 Child has severe combined immune deficiency, confirmed by an immunologist, but has not received a stem cell transplant: or
  - 3.4 Child has inborn errors of immunity (see Note E) that increase susceptibility to life-threatening viral respiratory infections, confirmed by an immunologist.

#### Notes:

- a) Ventilatory/respiratory support includes those on home oxygen, CPAP/VPAP and those with tracheostomies in situ managed at home
- b) Child requires/will require heart failure medication, and/or child has significant pulmonary hypertension, and/or infant will require surgical palliation/definitive repair within the next 3 months
- c) Mean pulmonary artery pressure more than 25 mmHg
- d) LV Ejection Fraction less than 40%
- e) Inborn errors of immunity include, but are not limited to, IFNAR deficiencies

# PERTUZUMAB - PCT only - Specialist - Special Authority see SA2276 below

Perieta 420 mg OP ✓ Baxter

### ⇒SA2276 Special Authority for Subsidy

Initial application — (metastatic breast cancer) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer	
----------------------------------------------------------------------------------------------	--

#### continued...

- 2 Either:
  - 2.1 Patient is chemotherapy treatment naïve; or
  - 2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
  - 3 The patient has good performance status (ECOG grade 0-1); and
  - 4 Pertuzumab to be administered in combination with trastuzumab; and
  - 5 Pertuzumab maximum first dose of 840 mg, followed by maximum of 420 mg every 3 weeks; and
  - 6 Pertuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

#### Either:

- 1 Both:
  - 1.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
  - 1.2 The cancer has not progressed at any time point during the previous 12 months whilst on pertuzumab and trastuzumab; or
- 2 All of the following:
  - 2.1 Patient has previously discontinued treatment with pertuzumab and trastuzumab for reasons other than severe toxicity or disease progression; and
  - 2.2 Patient has signs of disease progression; and
  - 2.3 Disease has not progressed during previous treatment with pertuzumab and trastuzumab.

RITUXIMAB (MABTHERA) - PCT only - Specialist	cial Authority see SA197	76 below	
Inj 100 mg per 10 ml vial	1,075.50	2	✓ Mabthera
Inj 500 mg per 50 ml vial	2,688.30	1	✓ Mabthera
Inj 1 mg for ECP	5.64	1 mg	✓ Baxter (Mabthera)

#### ⇒SA1976 Special Authority for Subsidy

Initial application — (rheumatoid arthritis - TNF inhibitors contraindicated) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Treatment with a Tumour Necrosis Factor alpha inhibitor is contraindicated; and
- 2 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
- 5 Any of the following:
  - 5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or
  - 5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
  - 5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
- 6 Either:
  - 6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
  - 6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	;	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

wrist, elbow, knee, ankle, and either shoulder or hip; and

- 7 Either:
  - 7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application: or
  - 7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months; and
- 8 Either:
  - 8.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
  - 8.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 9 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Initial application — (rheumatoid arthritis - prior TNF inhibitor use) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Both:
  - 1.1 The patient has had an initial community Special Authority approval for at least one of etanercept and/or adalimumab for rheumatoid arthritis: and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
    - 1.2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
  - 2.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
  - 2.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 3 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Renewal — (rheumatoid arthritis - re-treatment in 'partial responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 At 4 months following the initial course of rituximab infusions the patient had between a 30% and 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 1.2 At 4 months following the second course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 1.3 At 4 months following the third and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Either:
  - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
  - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Renewal — (rheumatoid arthritis - re-treatment in 'responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

1 Fither:

continued...

- 1.1 At 4 months following the initial course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 1.2 At 4 months following the second and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Fither:
  - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
  - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

RITUXIMAB (RIXIMYO) - PC	Γonly – Specialist – Special Authority	see SA2233 below	٧
1-1-400		075.00	_

Inj 100 mg per 10 ml vial	2/5.33	2	✓ Riximyo
Inj 500 mg per 50 ml vial	688.20	1	✓ Riximyo
Inj 1 mg for ECP	1.38	1 mg	✓ Baxter (Riximyo)

#### ⇒SA2233 Special Authority for Subsidy

Initial application — (ABO-incompatible organ transplant) from any relevant practitioner. Approvals valid without further renewal unless notified where patient is to undergo an ABO-incompatible solid organ transplant\*.

Note: Indications marked with \* are unapproved indications.

Initial application — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis\*; and
- 2 The total rituximab dose would not exceed the equivalent of 375 mg/m<sup>2</sup> of body-surface area per week for a total of 4 weeks; and
- 3 Any of the following:
  - 3.1 Induction therapy with daily oral or pulse intravenous cyclophosphamide has failed to achieve significant improvement of disease after at least 3 months; or
  - 3.2 Patient has previously had a cumulative dose of cyclophosphamide > 15 g or a further repeat 3 month induction course of cyclophosphamide would result in a cumulative dose > 15 g; or
  - 3.3 Cyclophosphamide and methotrexate are contraindicated; or
  - 3.4 Patient is a female of child-bearing potential; or
  - 3.5 Patient has a previous history of haemorrhagic cystitis, urological malignancy or haematological malignancy.

Note: Indications marked with \* are unapproved indications.

**Renewal — (ANCA associated vasculitis)** from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis\*; and
- 2 Patient has previously responded to treatment with rituximab but is now experiencing an acute flare of vasculitis; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m<sup>2</sup> of body-surface area per week for a total of 4 weeks.

Note: Indications marked with \* are unapproved indications.

**Initial application** — (Antibody-mediated organ transplant rejection) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has been diagnosed with antibody-mediated organ transplant rejection\*.

Note: Indications marked with \* are unapproved indications.

Initial application — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	sidised	Generic	
\$	Per	1	Manufacturer	

continued...

- 1 The patient has progressive Binet stage A, B or C chronic lymphocytic leukaemia (CLL) requiring treatment; and
- 2 Any of the following:
  - 2.1 The patient is rituximab treatment naive; or
  - 2.2 Fither:
    - 2.2.1 The patient is chemotherapy treatment naive; or
    - 2.2.2 Both:
      - 2.2.2.1 The patient's disease has relapsed following no more than three prior lines of chemotherapy treatment: and
      - 2.2.2.2 The patient has had a treatment-free interval of 12 months or more if previously treated with fludarabine and cyclophosphamide chemotherapy; or
  - 2.3 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax; and
- 3 The patient has good performance status; and
- 4 Either:
  - 4.1 The patient does not have chromosome 17p deletion CLL; or
  - 4.2 Rituximab treatment is to be used in combination with funded venetoclax for relapsed/refractory chronic lymphocytic leukaemia; and
- 5 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles; and
- 6 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration), bendamustine or venetoclax.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with rituximab is expected to improve symptoms and improve ECOG score to < 2.

Renewal — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

#### Both:

- 1 Either:
  - 1.1 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax; or
  - 1.2 All of the following:
    - 1.2.1 The patient's disease has relapsed following no more than one prior line of treatment with rituximab for CLL; and
    - 1.2.2 The patient has had an interval of 36 months or more since commencement of initial rituximab treatment;
    - 1.2.3 The patient does not have chromosome 17p deletion CLL; and
    - 1.2.4 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration) or bendamustine; and
- 2 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initial application — (Neuromyelitis Optica Spectrum Disorder(NMOSD)) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks; and
- 2 Either:
  - 2.1 The patient has experienced a severe episode or attack of NMOSD (rapidly progressing symptoms and clinical investigations supportive of a severe attack of NMOSD); or
  - 2.2 All of the following:
    - 2.2.1 The patient has experienced a breakthrough attack of NMOSD; and
    - 2.2.2 The patient is receiving treatment with mycophenolate; and
    - 2.2.3 The patients is receiving treatment with corticosteroids.

Renewal — (Neuromyelitis Optica Spectrum Disorder) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks; and
- 2 The patients has responded to the most recent course of rituximab; and
- 3 The patient has not received rituximab in the previous 6 months.

Initial application — (Post-transplant) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has B-cell post-transplant lymphoproliferative disorder\*; and
- 2 To be used for a maximum of 8 treatment cycles.

Note: Indications marked with \* are unapproved indications.

Renewal — (Post-transplant) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has B-cell post-transplant lymphoproliferative disorder\*; and
- 3 To be used for no more than 6 treatment cycles.

Note: Indications marked with \* are unapproved indications.

Initial application — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
- 2 Either:
  - 2.1 Treatment with corticosteroids and at least one other immunosuppressant for at least a period of 12 months has been ineffective; or
  - 2.2 Both:
    - 2.2.1 Treatment with at least one other immunosuppressant for a period of at least 12 months; and
    - 2.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

Renewal — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
- 2 An initial response lasting at least 12 months was demonstrated; and

Subsidy (Manufacturer's Price)	Sı	Fully ubsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

- 3 Either:
  - 3.1 The patient has relapsed despite treatment with corticosteroids and at least one other immunosuppressant for a period of at least 12 months; or
  - 3.2 Both:
    - 3.2.1 The patient's myasthenia gravis has relapsed despite treatment with at least one immunosuppressant for a period of at least 12 months; and
    - 3.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

Initial application — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is a child with SDNS\* or FRNS\*; and
- 2 Treatment with steroids for at least a period of 3 months has been ineffective or associated with evidence of steroid toxicity; and
- 3 Treatment with ciclosporin for at least a period of 3 months has been ineffective and/or discontinued due to unacceptable side effects; and
- 4 Treatment with mycophenolate for at least a period of 3 months with no reduction in disease relapses; and
- 5 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with \* are unapproved indications.

Renewal — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome\*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with \* are unapproved indications.

Initial application — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is a child with SRNS\* where treatment with steroids and ciclosporin for at least 3 months have been ineffective; and
- 2 Treatment with tacrolimus for at least 3 months has been ineffective; and
- 3 Genetic causes of nephrotic syndrome have been excluded; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with \* are unapproved indications.

**Renewal** — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome\*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks

Note: Indications marked with \* are unapproved indications.

Initial application — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 All of the following:
  - 1.1 The patient has treatment naive aggressive CD20 positive NHL; and
  - 1.2 To be used with a multi-agent chemotherapy regimen given with curative intent; and
  - 1.3 To be used for a maximum of 8 treatment cycles; or
- 2 Both:
  - 2.1 The patient has aggressive CD20 positive NHL with relapsed disease following prior chemotherapy; and
  - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Renewal — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has relapsed refractory/aggressive CD20 positive NHL; and
- 3 To be used with a multi-agent chemotherapy regimen given with curative intent; and
- 4 To be used for a maximum of 4 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

**Initial application — (haemophilia with inhibitors)** only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria:

Any of the following:

- 1 Patient has mild congenital haemophilia complicated by inhibitors; or
- 2 Patient has severe congenital haemophilia complicated by inhibitors and has failed immune tolerance therapy; or
- 3 Patient has acquired haemophilia.

**Renewal** — (haemophilia with inhibitors) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for haemophilia with inhibitors; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment.

Initial application — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Fither:
  - 1.1 Patient has immune thrombocytopenic purpura\* with a platelet count of less than or equal to 20,000 platelets per microlitre: or
  - 1.2 Patient has immune thrombocytopenic purpura\* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding; and
- 2 Any of the following:
  - 2.1 Treatment with steroids and splenectomy have been ineffective; or
  - 2.2 Treatment with steroids has been ineffective and splenectomy is an absolute contraindication; or
  - 2.3 Other treatments including steroids have been ineffective and patient is being prepared for elective surgery (e.g. splenectomy); and

Subsidy	Fu	lly Brand or	
(Manufacturer's Price)	Subsidis	ed Generic	
\$	Per	<ul> <li>Manufacturer</li> </ul>	

continued...

3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of

Note: Indications marked with \* are unapproved indications.

**Renewal** — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
  - 2.1 Patient was previously treated with rituximab for immune thrombocytopenic purpura\*; and
  - 2.2 An initial response lasting at least 12 months was demonstrated; and
  - 2.3 Patient now requires repeat treatment.

Note: Indications marked with \* are unapproved indications.

Initial application — (indolent, low-grade lymphomas or hairy cell leukaemia\*) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has indolent low grade NHL or hairy cell leukaemia\* with relapsed disease following prior chemotherapy; and
  - 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
  - 2.1 The patient has indolent, low grade lymphoma or hairy cell leukaemia\* requiring first-line systemic chemotherapy; and
  - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. \*Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Renewal — (indolent, low-grade lymphomas or hairy cell leukaemia\*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has indolent, low-grade NHL or hairy cell leukaemia\* with relapsed disease following prior chemotherapy; and
- 3 To be used for no more than 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. \*Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Initial application — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient has autoimmune pure red cell aplasia\* associated with a demonstrable B-cell lymphoproliferative disorder.

Note: Indications marked with \* are unapproved indications.

Renewal — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient was previously treated with rituximab for pure red cell aplasia\* associated with a demonstrable B-cell lymphoproliferative disorder and demonstrated an initial response lasting at least 12 months.

Note: Indications marked with \* are unapproved indications.

**Initial application** — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

1 Patient has cold haemagglutinin disease\*; and

Subsidy	Full	/ Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per 🗸	Manufacturer Manufacturer	

continued...

- 2 Patient has severe disease which is characterized by symptomatic anaemia, transfusion dependence or disabling circulatory symptoms; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks

Note: Indications marked with \* are unapproved indications.

Renewal — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Fither:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
  - 2.1 Patient was previously treated with rituximab for severe cold haemagglutinin disease\*; and
  - 2.2 An initial response lasting at least 12 months was demonstrated; and
  - 2.3 Patient now requires repeat treatment.

Note: Indications marked with \* are unapproved indications.

Initial application — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Both:

- 1 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks; and
- 2 Either:
  - 2.1 Patient has thrombotic thrombocytopenic purpura\* and has experienced progression of clinical symptoms or persistent thrombocytopenia despite plasma exchange; or
  - 2.2 Patient has acute idiopathic thrombotic thrombocytopenic purpura\* with neurological or cardiovascular pathology.

Note: Indications marked with \* are unapproved indications.

Renewal — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for thrombotic thrombocytopenic purpura\*; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with \* are unapproved indications.

Initial application — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe, immediately life- or organ-threatening SLE\*; and
- 2 The disease has proved refractory to treatment with steroids at a dose of at least 1 mg/kg; and
- 3 The disease has relapsed following prior treatment for at least 6 months with maximal tolerated doses of azathioprine, mycophenolate mofetil and high dose cyclophosphamide, or cyclophosphamide is contraindicated; and
- 4 Maximum of four 1000 mg infusions of rituximab.

Note: Indications marked with \* are unapproved indications.

Renewal — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 6 months for applications meeting the following criteria:

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidis	sed	Generic
	Per	1	Manufacturer

continued...

All of the following:

- 1 Patient's SLE\* achieved at least a partial response to the previous round of prior rituximab treatment; and
- 2 The disease has subsequently relapsed; and
- 3 Maximum of two 1000 mg infusions of rituximab.

Note: Indications marked with \* are unapproved indications.

Initial application — (warm autoimmune haemolytic anaemia (warm AlHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has warm autoimmune haemolytic anaemia\*; and
- 2 One of the following treatments has been ineffective: steroids (including if patient requires ongoing steroids at doses equivalent to > 5 mg prednisone daily), cytotoxic agents (e.g. cyclophosphamide monotherapy or in combination), intravenous immunoglobulin; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with \* are unapproved indications.

Renewal — (warm autoimmune haemolytic anaemia (warm AIHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Fither:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
  - 2.1 Patient was previously treated with rituximab for warm autoimmune haemolytic anaemia\*; and
  - 2.2 An initial response lasting at least 12 months was demonstrated; and
  - 2.3 Patient now requires repeat treatment.

Note: Indications marked with \* are unapproved indications.

**Initial application — (severe antisynthetase syndrome)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed antisynthetase syndrome; and
- 2 Patient has severe, immediately life or organ threatening disease, including interstitial lung disease; and
- 3 Either:
  - 3.1 Treatment with at least 3 immunosuppressants (oral steroids, cyclophosphamide, methotrexate, mycophenolate, ciclosporin, azathioprine) has not be effective at controlling active disease; or
  - 3.2 Rapid treatment is required due to life threatening complications; and
- 4 Maximum of four 1,000mg infusions of rituximab.

Renewal — (severe antisynthetase syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in inflammatory markers, muscle strength and pulmonary function; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 Maximum of two cycles of 2 x 1,000mg infusions of rituximab given two weeks apart.

Initial application — (graft versus host disease) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

1 Patient has refractory graft versus host disease following transplant; and

Subsidy	Fı	ılly	Brand or	
(Manufacturer's Price)	Subsidis	ed	Generic	
\$	Per	✓	Manufacturer	

continued...

- 2 Treatment with at least 3 immunosuppressants (oral steroids, ciclosporin, tacrolimus, mycophenolate, sirolimus) has not be effective at controlling active disease; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks

Initial application — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has severe chronic inflammatory demyelinating polyneuropathy (CIPD); and
- 2 Either:
  - 2.1 Both:
    - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
    - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
  - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function compared to baseline; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Initial application — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe anti-NMDA receptor autoimmune encephalitis; and
- 2 Either:
  - 2.1 Both:
    - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
    - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
  - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 The patient has experienced a relapse and now requires further treatment; and
- 4 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1.000 mg doses given two weeks apart.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

Initial application — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has CD20+ low grade or follicular B-cell NHL with relapsed disease following prior chemotherapy; and
  - 1.2 To be used for a maximum of 6 treatment cycles: or
- 2 Both
  - 2.1 The patient has CD20+ low grade or follicular B-cell NHL requiring first-line systemic chemotherapy; and
  - 2.2 To be used for a maximum of 6 treatment cycles.

Renewal — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 Rituximab is to be used for maintenance in CD20+ low grade or follicular B-cell NHL following induction with first-line systemic chemotherapy; and
- 2 Patient is intended to receive rituximab maintenance therapy for 2 years at a dose of 375 mg/m2 every 8 weeks (maximum of 12 cycles).

**Initial application** — (Membranous nephropathy) only from a nephrologist or any relevant practitioner on the recommendation of a nephrologist. Approvals valid for 6 weeks for applications meeting the following criteria: All of the following:

- 1 Either:
  - 1.1 Patient has biopsy-proven primary/idiopathic membranous nephropathy\*; or
  - 1.2 Patient has PLA2 antibodies with no evidence of secondary cause, and an eGFR of > 60ml/min/1.73m2; and
- 2 Patient remains at high risk of progression to end-stage kidney disease despite more than 3 months of treatment with conservative measures (see Note); and
- 3 The total rituximab dose would not exceed the equivalent of 375mg/m2 of body surface area per week for a total of 4 weeks.

Renewal — (Membranous nephropathy) only from a nephrologist or any relevant practitioner on the recommendation of a nephrologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for membranous nephropathy\*; and
- 2 Either:
  - 2.1 Treatment with rituximab was previously successful, but the condition has relapsed, and the patient now requires repeat treatment: or
  - 2.2 Patient achieved partial response to treatment and requires repeat treatment (see Note); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks

#### Notes:

- a) Indications marked with \* are unapproved indications.
- b) High risk of progression to end-stage kidney disease defined as > 5g/day proteinuria.
- c) Conservative measures include renin-angiotensin system blockade, blood-pressure management, dietary sodium and protein restriction, treatment of dyslipidaemia, and anticoagulation agents unless contraindicated or the patient has experienced intolerable side effects.
- d) Partial response defined as a reduction of proteinuria of at least 50% from baseline, and between 0.3 grams and 3.5 grams per 24 hours.

Initial application — (B-cell acute lymphoblastic leukaemia/lymphoma\*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Subsidy (Manufacturer's Price)	Fully Subsidised		
(ivialitulacture) 5 Filoe)	Per 🗸	Manufacturer	
Φ	rei •	Manuacturei	

continued

All of the following:

- 1 Patient has newly diagnosed B-cell acute lymphoblastic leukaemia/lymphoma\*; and
- 2 Treatment must be in combination with an intensive chemotherapy protocol with curative intent; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m<sup>2</sup> per dose for a maximum of 18 doses.

Note: Indications marked with \* are unapproved indications.

Initial application — (desensisation prior to transplant) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires desensitisation prior to mismatched allogenic stem cell transplant\*; and
- 2 Patient would receive no more than two doses at 375 mg/m2 of body-surface area.

Note: Indications marked with \* are unapproved indications.

Initial application — (pemiphigus\*) only from a dermatologist or relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 All of the following:
  - 1.1 Patient has severe rapidly progressive pemphigus; and
  - 1.2 Is used in combination with systemic corticosteroids (20 mg/day); and
  - 1.3 Any of the following:
    - 1.3.1 Skin involvement is at least 5% body surface area; or
    - 1.3.2 Significant mucosal involvement (10 or more mucosal erosions) or diffuse gingivitis or confluent large erosions: or
    - 1.3.3 Involvement of two or more mucosal sites: or
- 2 Both:
  - 2.1 Patient has pemphigus; and
  - 2.2 Patient has not experienced adequate clinical benefit from systemic corticosteroids (20 mg/day) in combination with a steroid sparing agent, unless contraindicated.

Note: Indications marked with \* are unapproved indications.

Renewal — (pemiphigus\*) only from a dermatologist or relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has experienced adequate clinical benefit from rituximab treatment, with improvement in symptoms and healing of skin ulceration and reduction in corticosteroid requirement; and
- 2 Patient has not received rituximab in the previous 6 months.

Note: Indications marked with \* are unapproved indications.

Initial application — (immunoglobulin G4-related disease (IgG4-RD\*)) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed diagnosis of IgG4-RD\*; and
- 2 Either:
  - 2.1 Treatment with corticosteroids and/or disease modifying anti-rheumatic drugs for at least 3 months has been ineffective in lowering corticosteroid dose below 5 mg per day (prednisone equivalent) without relapse; or
  - 2.2 Treatment with corticosteroids and/or disease modifying anti-rheumatic drugs is contraindicated or associated with evidence of toxicity or intolerance; and
- 3 Total rituximab dose used should not exceed a maximum of two 1000 mg infusions of rituximab given two weeks apart. Note: Indications marked with \* are unapproved indications.

Renewal — (immunoglobulin G4-related disease (IgG4-RD\*)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✓ Manufacturer	
----------------------------------------------------------------------------------------	--

continued...

All of the following:

- 1 Either:
  - 1.1 Treatment with rituximab for IgG4-RD\* was previously successful and patient's disease has demonstrated sustained response, but the condition has relapsed; or
  - 1.2 Patient is receiving maintenance treatment for IgG4-RD\*; and
- 2 Rituximab re-treatment not to be given within 6 months of previous course of treatment; and
- 3 Maximum of two 1000 mg infusions of rituximab given two weeks apart.

Note: Indications marked with \* are unapproved indications.

SECUKINUMAB - Special Authority see SA2403 below - Retail pharmacy

### ⇒SA2403 Special Authority for Subsidy

Initial application — (severe chronic plaque psoriasis – second-line biologic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab or etanercept, or has trialled infliximab in a Health NZ Hospital, for severe chronic plaque psoriasis; and
- 2 Either:
  - 2.1 The patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
  - 2.2 The patient has received insufficient benefit from adalimumab, etanercept or infliximab; and
- 3 A Psoriasis Area and Severity Index (PASI) assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Initial application — (severe chronic plaque psoriasis – first-line biologic) only from a dermatologist or any relevant practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
  - 1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; or
  - 1.3 Patient has severe chronic localised genital or flexural plaque psoriasis where the plaques or lesions have been present for at least 6 months from the time of initial diagnosis, and with a Dermatology Life Quality Index (DLQI) score greater than 10; and
- 2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin: and
- 3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course: and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Note: A treatment course is defined as a minimum of 12 weeks of treatment. "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand. foot, genital or flexural areas, at least 2 of the 3 PASI symptom sub scores for erythema, thickness and scaling are rated as

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

severe or very severe, and for the face, palm of a hand or sole of a foot the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis – first and second-line biologic) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Roth:

- 1 Either:
  - 1.1 Either:
    - 1.1.1 Patient's PASI score has reduced by 75% or more (PASI 75) as compared to baseline PASI prior to commencing secukinumab; or
    - 1.1.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing secukinumab; or
  - 1.2 Both:
    - 1.2.1 Patient had severe chronic localised genital or flexural plague psoriasis at the start of treatment; and
    - 1.2.2 Fither:
      - 1.2.2.1 The patient has experienced a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; or
      - 1.2.2.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing secukinumab; and
- 2 Secukinumab to be administered at a maximum dose of 300 mg monthly.

Initial application — (ankylosing spondylitis – second-line biologic) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and
- 2 Either:
  - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
  - 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

Renewal — (ankylosing spondylitis – second-line biologic) only from a rheumatologist or medical practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Following 12 weeks initial treatment of secukinumab treatment, BASDAI has improved by 4 or more points from pre-secukinumab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefitted from treatment and that continued treatment is appropriate; and
- 3 Secukinumab to be administered at doses no greater than 150 mg monthly.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 Patient has had an initial Special Authority approval for adalimumab, etanercept or infliximab for psoriatic arthritis; and
  - 1.2 Either:
    - 1.2.1 Patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
    - 1.2.2 Patient has received insufficient benefit from adalimumab, etanercept or infliximab to meet the renewal criteria for adalimumab, etanercept or infliximab for psoriatic arthritis; or
- 2 All of the following:
  - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and

continued...

- 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
- 2.4 Either:
  - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
  - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2.5 Any of the following:
  - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
  - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
  - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

**Renewal — (psoriatic arthritis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Either:
  - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior secukinumab treatment in the opinion of the treating physician; and
- 2 Secukinumab to be administered at doses no greater than 300 mg monthly.

SILTUXIM	IAB - Special Authority see SA1596 below - Retail pharmacy	
Note:	Siltuximab is to be administered at doses no greater than 11 mg/kg every 3	weeks.

 Inj 100 mg vial
 770.57
 1
 ✓ Sylvant

 Inj 400 mg vial
 3,082.33
 1
 ✓ Svlvant

### ⇒SA1596 Special Authority for Subsidy

Initial application only from a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's Disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Siltuximab is to be administered at doses no greater than 11 mg/kg every 3 weeks.

**Renewal** only from a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has sustained improvement in inflammatory markers and functional status.

TIXAGEVIMAB WITH CILGAVIMAB - [Xpharm] - Subsidy by endorsement

- a) No patient co-payment payable
- b) Treatment is funded only if patient meets access criteria for tixagevimab with cilgavimab (as per https://pharmac.govt.nz/Evusheld) and has been endorsed accordingly by the prescriber. The supply of treatment is via Pharmac's approved distribution process. Refer to the Pharmac website for more information about this and stock availability.

inj 100 mg per mi, 1.5	mi viai with ciigavimad 100 mg per			
ml 1.5 ml vial		0.00	1	✓ Evusheld

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	
TOCILIZUMAB - PCT only - Special Authority see SA2404 belo	W			
Inj 20 mg per ml, 4 ml vial	220.00	1	✓	Actemra
Inj 20 mg per ml, 10 ml vial		1	✓	Actemra
Inj 20 mg per ml, 20 ml vial	1,100.00	1	✓	Actemra
Inj 1 mg for ECP		1 mg	✓	Baxter

### ⇒SA2404 Special Authority for Subsidy

Initial application — (cytokine release syndrome) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

### Either:

- 1 Both:
  - 1.1 The patient has developed grade 3 or 4 cytokine release syndrome associated with the administration of blinatumomab for the treatment of acute lymphoblastic leukaemia; and
  - 1.2 Tocilizumab is to be administered at doses no greater than 8 mg/kg IV for a maximum of 3 doses (if less than 30kg, maximum of 12 mg/kg); or
- 2 All of the following:
  - 2.1 The patient is enrolled in the Malaghan Institute of Medical Research ENABLE trial programme; and
  - 2.2 The patient has developed CRS or Immune Effector Cell-Associated Neurotoxicity Syndrome (ICANS) following CAR T-cell therapy for the treatment of relapsed or refractory B-cell non-Hodgkin lymphoma; and
  - 2.3 Tocilizumab is to be administered according to the consensus guidelines for CRS or ICANS for CAR T-cell therapy at doses no greater than 8 mg/kg IV for a maximum of 3 doses.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

#### Both:

- 1 Patient was being treated with tocilizumab prior to 1 February 2019; and
- 2 Any of the following:
  - 2.1 rheumatoid arthritis: or
  - 2.2 systemic juvenile idiopathic arthritis; or
  - 2.3 adult-onset Still's disease: or
  - 2.4 polyarticular juvenile idiopathic arthritis; or
  - 2.5 idiopathic multicentric Castleman's disease.

Initial application — (Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

#### All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Fither:
  - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
  - 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
- 3 Either:
  - 3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or
  - 3.2 Both:
    - 3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a Health NZ Hospital; and
    - 3.2.2 Either:
      - 3.2.2.1 The patient has experienced intolerable side effects from rituximab; or
      - 3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Subsidy	F	ully	Brand or	
(Manufacturer's Pr	rice) Subsid	ised	Generic	
\$	Per	✓	Manufacturer	

continued...

Initial application — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 2 Tocilizumab is to be used as monotherapy; and
- 3 Either:
  - 3.1 Treatment with methotrexate is contraindicated; or
  - 3.2 Patient has tried and did not tolerate oral and/or parenteral methotrexate; and
- 4 Fither:
  - 4.1 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of ciclosporin alone or in combination with another agent; or
  - 4.2 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of leflunomide alone or in combination with another agent; and
- 5 Either:
  - 5.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 active, swollen, tender joints;
  - 5.2 Patient has persistent symptoms of poorly controlled and active disease in at least four active joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 6 Fither:
  - 6.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
  - 6.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Initial application — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient diagnosed with systemic juvenile idiopathic arthritis: and
- 2 Patient has tried and not responded to a reasonable trial of all of the following, either alone or in combination: oral or parenteral methotrexate; non-steroidal anti-inflammatory drugs (NSAIDs); and systemic corticosteroids.

Initial application — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 Fither:
    - 1.1.1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for adult-onset Still's disease (AOSD); or
    - 1.1.2 The patient has been started on tocilizumab for AOSD in a Health NZ Hospital; and
  - 1.2 Fither:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
    - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
  - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
  - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal antiinflammatory drugs (NSAIDs) and methotrexate; and
  - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Initial application — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for both etanercept and adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
  - 1.2 The patient has experienced intolerable side effects, or has received insufficient benefit from, both etanercept and adalimumab; or
- 2 All of the following:
  - 2.1 Treatment with a tumour necrosis factor alpha inhibitor is contraindicated; and
  - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
  - 2.3 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
  - 2.4 Any of the following:
    - 2.4.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
    - 2.4.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose): or
    - 2.4.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Initial application — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Tocilizumab to be administered at doses no greater than 8 mg/kg IV every 3-4 weeks.

Initial application — (moderate to severe COVID-19) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed (or probable) COVID-19; and
- 2 Oxygen saturation of < 92% on room air, or requiring supplemental oxygen; and
- 3 Patient is receiving adjunct systemic corticosteroids, or systemic corticosteroids are contraindicated; and
- 4 Tocilizumab is to be administered at doses no greater than 8mg/kg IV for a maximum of one dose; and
- 5 Tocilizumab is not to be administered in combination with barcitinib.

Renewal — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Renewal — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Following up to 6 months' initial treatment, the patient has achieved at least an American College of Rheumatology paediatric 30% improvement criteria (ACR Pedi 30) response from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing ACR Pedi 30 response from baseline.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

**Renewal — (adult-onset Still's disease)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months where the patient has a sustained improvement in inflammatory markers and functional status.

Renewal — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
  - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Renewal — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has a sustained improvement in inflammatory markers and functional status.

TRASTUZUMAB (HERZUMA) - PCT only - Special Authority see SA2293 below

Inj 150 mg vial100.00	1	✓ Herzuma
Inj 440 mg vial293.35	1	✓ Herzuma
Inj 1 mg for ECP	1 mg	✓ Baxter

#### ⇒SA2293 Special Authority for Subsidy

**Initial application** — (early breast cancer) from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria:

Both:

- 1 The patient has early breast cancer expressing HER-2 IHC 3+ or ISH + (including FISH or other current technology); and
- 2 Maximum cumulative dose of 106 mg/kg (12 months' treatment).

Renewal — (early breast cancer\*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Fither:

- 1 All of the following:
  - 1.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
  - 1.2 The patient received prior adjuvant trastuzumab treatment for early breast cancer; and
  - 1.3 Any of the following:
    - 1.3.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
    - 1.3.2 The patient discontinued lapatinib within 3 months due to intolerable side effects and the cancer did not progress whilst on lapatinib; or
    - 1.3.3 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
  - 1.4 Fither:
    - 1.4.1 Trastuzumab will not be given in combination with pertuzumab; or
    - 1.4.2 All of the following:
      - 1.4.2.1 Trastuzumab to be administered in combination with pertuzumab; and
      - 1.4.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
      - 1.4.2.3 The patient has good performance status (ECOG grade 0-1); and
  - 1.5 Trastuzumab to be discontinued at disease progression; or

Subsidy	Full	/ Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per 🗸	Manufacturer Manufacturer	

continued...

- 2 All of the following:
  - 2.1 Patient has previously discontinued treatment with trastuzumab in the metastatic setting for reasons other than severe toxicity or disease progression; and
  - 2.2 Patient has signs of disease progression; and
  - 2.3 Disease has not progressed during previous treatment with trastuzumab.

Note: \* For patients with relapsed HER-2 positive disease who have previously received adjuvant trastuzumab for early breast cancer

Initial application — (metastatic breast cancer) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Fither:
  - 2.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
  - 2.2 The patient discontinued lapatinib within 3 months due to intolerable side effects and the cancer did not progress whilst on lapatinib; and
- 3 Either:
  - 3.1 Trastuzumab will not be given in combination with pertuzumab; or
  - 3.2 All of the following:
    - 3.2.1 Trastuzumab to be administered in combination with pertuzumab; and
    - 3.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer: and
    - 3.2.3 The patient has good performance status (ECOG grade 0-1); and
- 4 Trastuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 All of the following:
  - 1.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
  - 1.2 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
  - 1.3 Trastuzumab to be discontinued at disease progression; or
- 2 All of the following:
  - 2.1 Patient has previously discontinued treatment with trastuzumab for reasons other than severe toxicity or disease progression; and
  - 2.2 Patient has signs of disease progression; and
  - 2.3 Disease has not progressed during previous treatment with trastuzumab.

**Initial application** — (gastric, gastro-oesophageal junction and oesophageal cancer) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has locally advanced or metastatic gastric, gastro-oesophageal junction or oesophageal cancer expressing HER-2 IHC 2+ FISH+ or IHC3+ (or other current technology); and
- 2 Patient has an ECOG score of 0-2.

Renewal — (gastric, gastro-oesophageal junction and oesophageal cancer) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

Subsidy (Manufacturer's Price)	Suk	Fully	Brand or Generic
 \$	Per	√	Manufacturer

continued...

- 1 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 2 Trastuzumab to be discontinued at disease progression.

### TRASTUZUMAB DERUXTECAN - PCT only - Special Authority see SA2420 below

Inj 100 mg per ml, 1 ml vial	2,550.00	1	<ul><li>Enhertu</li></ul>
Inj 1 mg for ECP	27.05	1 mg	Baxter

#### ⇒SA2420 Special Authority for Subsidy

**Initial application** only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

#### Either:

- 1 Patient is currently on treatment with trastuzumab deruxtecan and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
  - Patient has metastatic breast cancer expressing HER-2 IHC3+ or ISH+ (including FISH or other current technology); and
  - 2.2 Patient has previously received trastuzumab and chemotherapy, separately or in combination; and
  - 2.3 Fither:
    - 2.3.1 The patient has received prior therapy for metastatic disease; or
    - 2.3.2 The patient developed disease recurrence during, or within six months of completing adjuvant therapy; and
  - 2.4 Patient has a good performance status (ECOG 0-1); and
  - 2.5 Patient has not received prior funded trastuzumab deruxtecan treatment; and
  - 2.6 Treatment to be discontinued at disease progression.

**Renewal** only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The cancer has not progressed at any time point during the previous approval period whilst on trastuzumab deruxtecan; and
- 2 Treatment to be discontinued at disease progression.

Note: Prior or adjuvant therapy includes anthracycline, other chemotherapy, biological drugs, or endocrine therapy.

#### TRASTUZUMAB FMTANSINF - PCT only - Specialist - Special Authority see SA2424 below

TO TO TO E CHITTAIN TO LINE	i o i oilly opoolaliot	opoolar riatifority ood or	IL IL I DOIO!	
Inj 100 mg vial		2,320.00	1	✓ Kadcyla
Inj 160 mg vial		3,712.00	1	✓ Kadcyla
Ini 1 mg for ECP		24.52	1 ma	✓ Baxter

#### ⇒SA2424 Special Authority for Subsidy

Initial application — (early breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has early breast cancer expressing HER2 IHC3+ or ISH+; and
- 2 Documentation of pathological invasive residual disease in the breast and/or auxiliary lymph nodes following completion of surgery; and
- 3 Patient has completed systemic neoadjuvant therapy with trastuzumab and chemotherapy prior to surgery; and
- 4 Disease has not progressed during neoadjuvant therapy; and
- 5 Patient has left ventricular ejection fraction of 45% or greater; and
- 6 Adjuvant treatment with trastuzumab emtansine to be commenced within 12 weeks of surgery; and
- 7 Trastuzumab emtansine to be discontinued at disease progression; and
- 8 Total adjuvant treatment duration must not exceed 42 weeks (14 cycles).

Initial application — (metastatic breast cancer) only from a relevant specialist or any relevant practitioner on the

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

continued...

recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Patient has previously received trastuzumab and chemotherapy, separately or in combination; and
- 3 Either:
  - 3.1 The patient has received prior therapy for metastatic disease\*; or
  - 3.2 The patient developed disease recurrence during, or within six months of completing adjuvant therapy\*; and
- 4 Patient has a good performance status (ECOG 0-1); and
- 5 Either:
  - 5.1 Patient does not have symptomatic brain metastases; or
  - 5.2 Patient has brain metastases and has received prior local CNS therapy, and
- 6 Fither:
  - 6.1 Patient has not received prior funded trastuzumab emtansine or trastuzumab deruxtecan treatment; or
  - 6.2 Both:
    - 6.2.1 Patient has discontinued trastuzumab deruxtecan due to intolerance; and
    - 6.2.2 The cancer did not progress while on trastuzumab deruxtecan; and
- 7 Treatment to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The cancer has not progressed at any time point during the previous approval period whilst on trastuzumab emtansine:
- 2 Treatment to be discontinued at disease progression.

Note: Prior or adjuvant therapy includes anthracycline, other chemotherapy, biological drugs, or endocrine therapy,

USTEKINUMAB - Special Authority see SA2182 below - Retail pharmacy

Inj 90 mg per ml, 1 ml pre-filled syringe......4,162.00 ✓ Stelara

#### ⇒SA2182 Special Authority for Subsidy

Initial application — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Fither:

- 1 Patient is currently on treatment with ustekinumab commenced prior to 1 February 2023 and met all remaining criteria (criterion 2) below at the time of commencing treatment: or
  - 2 Both:
    - 2.1 Patient has active Crohn's disease; and
    - 2.2 Either:
      - 2.2.1 Patient has had an initial approval for prior biologic therapy for Crohn's disease and has experienced intolerable side effects or insufficient benefit to meet renewal criteria: or
      - 2.2.2 Both:
        - 2.2.2.1 Patient meets the initiation criteria for prior biologic therapies for Crohn's disease; and
        - 2.2.2.2 Other biologics for Crohn's disease are contraindicated.

Renewal — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 CDAI score has reduced by 100 points, or HBI score has reduced by 3 points, from when the patient was initiated on biologic therapy; or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1.2 CDAI score is 150 or less, or HBI is 4 or less; or
- 1.3 The patient has experienced an adequate response to treatment, but CDAI score and/or HBI score cannot be assessed: and
- 2 Ustekinumab to be administered at a dose no greater than 90 mg every 8 weeks.

Initial application — (Crohn's disease - children\*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Patient is currently on treatment with ustekinumab commenced prior to 1 February 2023 and met all remaining criteria (criterion 2) below at the time of commencing treatment; or
- 2 Both:
  - 2.1 Patient has active Crohn's disease: and
  - 2.2 Either:
    - 2.2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria; or
    - 2.2.2 Both:
      - 2.2.2.1 Patient meets the initiation criteria for prior biologic therapies for Crohn's disease; and
      - 2.2.2.2 Other biologics for Crohn's disease are contraindicated.

Note: Indication marked with \* is an unapproved indication.

Renewal — (Crohn's disease - children\*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 PCDAI score has reduced by 10 points from when the patient was initiated on biologic therapy; or
  - 1.2 PCDAI score is 15 or less; or
  - 1.3 The patient has experienced an adequate response to treatment, but CDAI score cannot be assessed; and
- 2 Ustekinumab to administered at a dose no greater than 90 mg every 8 weeks.

Note: Indication marked with \* is an unapproved indication.

Initial application — (ulcerative colitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Patient is currently on treatment with ustekinumab commenced prior to 1 February 2023 and met all remaining criteria (criterion 2) below at the time of commencing treatment; or
- 2 Both:
  - 2.1 Patient has active ulcerative colitis: and
  - 2.2 Either:
    - 2.2.1 Patient has had an initial approval for prior biologic therapy for ulcerative colitis and has experienced intolerable side effects or insufficient benefit to meet renewal criteria; or
    - 2.2.2 Both:
      - 2.2.2.1 Patient meets the initiation criteria for prior biologic therapies for ulcerative colitis; and
      - 2.2.2.2 Other biologics for ulcerative colitis are contraindicated.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Fither:
  - 1.1 The SCCAI score has reduced by 2 points or more from the SCCAI score since initiation on biologic therapy; or
  - 1.2 PUCAI score has reduced by 10 points or more from the PUCAI score since initiation on biologic therapy\*; and
- 2 Ustekinumab will be used at a dose no greater than 90 mg intravenously every 8 weeks.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

Note: Criterion marked with \* is for an unapproved indication.

VEDOLIZUMAB - PCT only - Special Authority see SA2183 below

⇒SA2183 Special Authority for Subsidy

Initial application — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active Crohn's disease; and
- 2 Any of the following:
  - 2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria (unless contraindicated); or
  - 2.2 Patient has a CDAI score of greater than or equal to 300, or HBI score of greater than or equal to 10; or
  - 2.3 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
  - 2.4 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
  - 2.5 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Any of the following:
  - 3.1 Patient has tried but experienced an inadequate response to (including lack of initial response and/or loss of initial response) from prior therapy with immunomodulators and corticosteroids; or
  - 3.2 Patient has experienced intolerable side effects from immunomodulators and corticosteroids; or
  - 3.3 Immunomodulators and corticosteroids are contraindicated.

Renewal — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 CDAI score has reduced by 100 points, or HBI score has reduced by 3 points, from when the patient was initiated on biologic therapy; or
  - 1.2 CDAI score is 150 or less, or HBI is 4 or less; or
  - 1.3 The patient has experienced an adequate response to treatment, but CDAI score and/or HBI score cannot be assessed; and
- 2 Vedolizumab to administered at a dose no greater than 300 mg every 8 weeks.

Initial application — (Crohn's disease - children\*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has active Crohn's disease; and
- 2 Any of the following:
  - 2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria (unless contraindicated); or
  - 2.2 Patient has a Paediatric Crohn's Disease Activity Index (PCDAI) score of greater than or equal to 30; or
  - 2.3 Patient has extensive small intestine disease; and
- 3 Any of the following:
  - 3.1 Patient has tried but experienced an inadequate response to (including lack of initial response and/or loss of initial response) from prior therapy with immunomodulators and corticosteroids; or
  - 3.2 Patient has experienced intolerable side effects from immunomodulators and corticosteroids; or
  - 3.3 Immunomodulators and corticosteroids are contraindicated.

Note: Indication marked with \* is an unapproved indication.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

Renewal — (Crohn's disease - children\*) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 PCDAI score has reduced by 10 points from when the patient was initiated on biologic therapy; or
  - 1.2 PCDAI score is 15 or less; or
  - 1.3 The patient has experienced an adequate response to treatment, but CDAI score cannot be assessed; and
- 2 Vedolizumab to administered at a dose no greater than 300mg every 8 weeks.

Note: Indication marked with \* is an unapproved indication.

Initial application — (ulcerative colitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active ulcerative colitis; and
- 2 Any of the following:
  - 2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria (unless contraindicated); or
  - 2.2 Patient has a SCCAI score is greater than or equal to 4; or
  - 2.3 Patient's PUCAI score is greater than or equal to 20\*; and
- 3 Any of the following:
  - 3.1 Patient has tried but experienced an inadequate response to (including lack of initial response and/or loss of initial response) from prior therapy with immunomodulators and corticosteroids; or
  - 3.2 Patient has experienced intolerable side effects from immunomodulators and corticosteroids; or
  - 3.3 Immunomodulators and corticosteroids are contraindicated.

Note: Indication marked with \* is an unapproved indication.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 The SCCAI score has reduced by 2 points or more from the SCCAI score since initiation on biologic therapy; or
  - 1.2 The PUCAI score has reduced by 10 points or more from the PUCAI score since initiation on biologic therapy \*; and
- 2 Vedolizumab will be used at a dose no greater than 300 mg intravenously every 8 weeks.

Note: Indication marked with \* is an unapproved indication.

# Programmed Cell Death-1 (PD-1) Inhibitors

## ⇒SA2264 Special Authority for Subsidy

**Initial application** — (non-small cell lung cancer second line monotherapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced or metastatic non-small cell lung cancer; and
- 2 Patient has not received prior funded treatment with an immune checkpoint inhibitor for NSCLC; and
- 3 For patients with non-squamous histology there is documentation confirming that the disease does not express activating mutations of EGFR or ALK tyrosine kinase unless not possible to ascertain; and

Subsidy (Manufacturer's Price)	Subsi	Fully	Brand or Generic
 \$	Per	1	Manufacturer

continued...

- 4 Patient has an ECOG 0-2; and
- 5 Patient has documented disease progression following treatment with at least two cycles of platinum-based chemotherapy; and
- 6 Atezolizumab is to be used as monotherapy at a dose of 1200 mg every three weeks (or equivalent) for a maximum of 16 weeks; and
- 7 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Renewal — (non-small cell lung cancer second line monotherapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Any of the following:
  - 1.1 Patient's disease has had a complete response to treatment; or
  - 1.2 Patient's disease has had a partial response to treatment; or
  - 1.3 Patient has stable disease; and
- 2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period; and
- 3 No evidence of disease progression; and
- 4 The treatment remains clinically appropriate and patient is benefitting from treatment; and
- 5 Atezolizumab to be used at a maximum dose of 1200 mg every three weeks (or equivalent); and
- 6 Treatment with atezolizumab to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

## DURVALUMAB - PCT only - Specialist - Special Authority see SA2425 below

Inj 50 mg per ml, 10 ml vial	4,700.00	1	Imfinzi
Inj 50 mg per ml, 2.4 ml vial	1,128.00	1	Imfinzi
Inj 1 mg for ECP	9.59	1 mg	<ul><li>Baxter</li></ul>

#### ⇒SA2425 Special Authority for Subsidy

Initial application — (Non-small cell lung cancer) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Either:
  - 1.1 Patient has histologically or cytologically documented stage III, locally advanced, unresectable non-small cell lung cancer (NSCLC); or
  - 1.2 Patient has histologically or cytologically documented stage IIb (T1N2a only), locally advanced, unresectable non-small cell lung cancer (NSCLC); and
- 2 Patient has received two or more cycles of platinum-based chemotherapy concurrently with definitive radiation therapy; and
- 3 Patient has no disease progression following the second or subsequent cycle of platinum-based chemotherapy with definitive radiation therapy treatment; and
- 4 Patient has a ECOG performance status of 0 or 1; and
- 5 Patient has completed last radiation dose within 8 weeks of starting treatment with duryalumab; and
- 6 Patient must not have received prior PD-1 or PD-L1 inhibitor therapy for this condition; and
- 7 Fither
  - 7.1 Durvalumab is to be used at a maximum dose of no greater than 10 mg/kg every 2 weeks; or
  - 7.2 Durvalumab is to be used at a flat dose of 1500 mg every 4 weeks; and
- 8 Treatment with durvalumab to cease upon signs of disease progression.

Renewal — (Non-small cell lung cancer) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

	Subsidy	Fully	Brand or
(Manufa	acturer's Price)	Subsidised	Generic
	\$ Per	✓	Manufacturer

continued...

- 1 The treatment remains clinically appropriate and the patient is benefitting from treatment; and
- 2 Fither
  - 2.1 Durvalumab is to be used at a maximum dose of no greater than 10 mg/kg every 2 weeks; or
  - 2.2 Durvalumab is to be used at a flat dose of 1500 mg every 4 weeks; and
- 3 Treatment with durvalumab to cease upon signs of disease progression; and
- 4 Total continuous treatment duration must not exceed 12 months.

NIVOLUMAB - PCT only - Specialist - Special Authority see SA2405 below

Inj 10 mg per ml, 4 ml vial1,051.98	1	✓ Opdivo
Inj 10 mg per ml, 10 ml vial2,629.96	1	✓ Opdivo
Inj 1 mg for ECP27.62	1 mg	✓ Baxter

### ⇒SA2405 Special Authority for Subsidy

**Initial application** only from a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Baseline measurement of overall tumour burden is documented clinically and radiologically; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Either:
  - 4.1 Patient has not received funded pembrolizumab; or
  - 4.2 Both:
    - 4.2.1 Patient has received an initial Special Authority approval for pembrolizumab and has discontinued pembrolizumab within 12 weeks of starting treatment due to intolerance; and
    - 4.2.2 The cancer did not progress while the patient was on pembrolizumab; and
- 5 Documentation confirming that the patient has been informed and acknowledges that funded treatment with nivolumab will not be continued if their disease progresses.

Renewal — (less than 24 months on treatment) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 All of the following:
  - 1.1 Any of the following:
    - 1.1.1 Patient's disease has had a complete response to treatment; or
    - 1.1.2 Patient's disease has had a partial response to treatment; or
    - 1.1.3 Patient has stable disease; and
  - 1.2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period; and
  - 1.3 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
- 2 All of the following:
  - 2.1 Patient has previously discontinued treatment with nivolumab for reasons other than severe toxicity or disease progression; and
  - 2.2 Patient has signs of disease progression; and
  - 2.3 Disease has not progressed during previous treatment with nivolumab.

Renewal — (more than 24 months on treatment) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 Patient has been on treatment for more than 24 months; and
- 2 Either:
  - 2.1 All of the following:

|--|

continued...

- 2.1.1 Any of the following:
  - 2.1.1.1 Patient's disease has had a complete response to treatment; or
  - 2.1.1.2 Patient's disease has had a partial response to treatment: or
  - 2.1.1.3 Patient has stable disease; and
- 2.1.2 Response to treatment in target lesions has been determined by comparable radiologic or clinical assessment following the most recent treatment period: and
- 2.1.3 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
- 2.2 All of the following:
  - 2.2.1 Patient has previously discontinued treatment with nivolumab for reasons other than severe toxicity or disease progression; and
  - 2.2.2 Patient has signs of disease progression; and
  - 2.2.3 Disease has not progressed during previous treatment with nivolumab.

Initial application — (Renal cell carcinoma) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria: Either:

- 1 Patient is currently on treatment with nivolumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
  - 2.1 Patient has metastatic renal-cell carcinoma; and
  - 2.2 The disease is of predominant clear-cell histology; and
  - 2.3 Patient has an ECOG performance score of 0-2; and
  - 2.4 Patient has documented disease progression following one or two previous regimens of antiangiogenic therapy; and
  - 2.5 Nivolumab is to be used as monotherapy at a maximum dose of 240 mg every 2 weeks (or equivalent) and discontinued at disease progression.

Renewal — (Renal cell carcinoma) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 Patient's disease has had a complete response to treatment; or
  - 1.2 Patient's disease has had a partial response to treatment; or
  - 1.3 Patient has stable disease; and
- 2 No evidence of disease progression; and
- 3 Nivolumab is to be used as monotherapy at a maximum dose of 240 mg every 2 weeks (or equivalent) and discontinued at disease progression.

PEMBROLIZUMAB - PCT only - Specialist - Special Au	thority see SA2386 below		
Inj 25 mg per ml, 4 ml vial	4,680.00	1	✓ Keytruda
Inj 1 mg for ECP	47.74	1 mg	✓ Baxter

#### ⇒SA2386 Special Authority for Subsidy

Initial application — (unresectable or metastatic melanoma) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Baseline measurement of overall tumour burden is documented clinically and radiologically; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Either:
  - 4.1 Patient has not received funded nivolumab; or

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic	
\$	Per	✓	Manufacturer	

continued...

- 4.2 Both:
  - 4.2.1 Patient has received an initial Special Authority approval for nivolumab and has discontinued nivolumab within 12 weeks of starting treatment due to intolerance; and
  - 4.2.2 The cancer did not progress while the patient was on nivolumab; and
- 5 Documentation confirming that the patient has been informed and acknowledges that funded treatment with pembrolizumab will not be continued if their disease progresses.

Renewal — (unresectable or metastatic melanoma, less than 24 months on treatment) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 All of the following:
  - 1.1 Any of the following:
    - 1.1.1 Patient's disease has had a complete response to treatment; or
    - 1.1.2 Patient's disease has had a partial response to treatment; or
    - 1.1.3 Patient has stable disease; and
  - 1.2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period; and
  - 1.3 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
- 2 All of the following:
  - 2.1 Patient has previously discontinued treatment with pembrolizumab for reasons other than severe toxicity or disease progression; and
  - 2.2 Patient has signs of disease progression; and
  - 2.3 Disease has not progressed during previous treatment with pembrolizumab.

Renewal — (unresectable or metastatic melanoma, more than 24 months on treatment) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 Patient has been on treatment for more than 24 months; and
- 2 Either:
  - 2.1 All of the following:
    - 2.1.1 Any of the following:
      - 2.1.1.1 Patient's disease has had a complete response to treatment; or
      - 2.1.1.2 Patient's disease has had a partial response to treatment; or
      - 2.1.1.3 Patient has stable disease; and
    - 2.1.2 Response to treatment in target lesions has been determined by comparable radiologic or clinical assessment following the most recent treatment period; and
    - 2.1.3 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
  - 2.2 All of the following:
    - 2.2.1 Patient has previously discontinued treatment with pembrolizumab for reasons other than severe toxicity or disease progression; and
    - 2.2.2 Patient has signs of disease progression; and
    - 2.2.3 Disease has not progressed during previous treatment with pembrolizumab.

**Initial application — (non-small cell lung cancer first-line monotherapy)** only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

1 Patient has locally advanced or metastatic, unresectable, non-small cell lung cancer; and

Subsidy (Manufacturer's Price)	,	Fully Subsidised	Brand or Generic	
\$	Per	✓	Manufacturer	

#### continued...

- 2 Patient has not had chemotherapy for their disease in the palliative setting; and
- 3 Patient has not received prior funded treatment with an immune checkpoint inhibitor for NSCLC; and
- 4 For patients with non-squamous histology there is documentation confirming that the disease does not express activating mutations of EGFR or ALK tyrosine kinase unless not possible to ascertain; and
- 5 Pembrolizumab to be used as monotherapy; and
- 6 Fither:
  - 6.1 There is documentation confirming the disease expresses PD-L1 at a level greater than or equal to 50% as determined by a validated test unless not possible to ascertain; or
  - 6.2 Both:
    - 6.2.1 There is documentation confirming the disease expresses PD-L1 at a level greater than or equal to 1% as determined by a validated test unless not possible to ascertain; and
    - 6.2.2 Chemotherapy is determined to be not in the best interest of the patient based on clinician assessment; and
- 7 Patient has an ECOG 0-2: and
- 8 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks; and
- 9 Baseline measurement of overall tumour burden is documented clinically and radiologically.

**Renewal — (non-small cell lung cancer first line monotherapy)** only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Any of the following:
  - 1.1 Patient's disease has had a complete response to treatment; or
  - 1.2 Patient's disease has had a partial response to treatment; or
  - 1.3 Patient has stable disease; and
- 2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period; and
- 3 No evidence of disease progression; and
- 4 The treatment remains clinically appropriate and patient is benefitting from treatment; and
- 5 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 6 Treatment with pembrolizumab to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

**Initial application — (non-small cell lung cancer first-line combination therapy)** only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

#### All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-small cell lung cancer; and
- 2 The patient has not had chemotherapy for their disease in the palliative setting; and
- 3 Patient has not received prior funded treatment with an immune checkpoint inhibitor for NSCLC; and
- 4 For patients with non-squamous histology there is documentation confirming that the disease does not express activating mutations of EGFR or ALK tyrosine kinase unless not possible to ascertain; and
- 5 Pembrolizumab to be used in combination with platinum-based chemotherapy; and
- 6 Patient has an ECOG 0-2: and
- 7 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks; and
- 8 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Renewal — (non-small cell lung cancer first line combination therapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

Subsidy (Manufacturer's F	Price) Sub	Fully	Brand or Generic
\$	Per	•	Manufacturer

continued...

All of the following:

- 1 Any of the following:
  - 1.1 Patient's disease has had a complete response to treatment; or
  - 1.2 Patient's disease has had a partial response to treatment; or
  - 1.3 Patient has stable disease; and
- 2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period: and
- 3 No evidence of disease progression; and
- 4 The treatment remains clinically appropriate and patient is benefitting from treatment; and
- 5 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 6 Treatment with pembrolizumab to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

**Initial application** — (breast cancer, advanced) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: Fither:

- 1 Patient is currently on treatment with pembrolizumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
  - 2.1 Either:
    - 2.1.1 Patient has recurrent or de novo unresectable, inoperable locally advanced triple-negative breast cancer (that does not express ER, PR or HER2 IHC3+ or ISH+ [including FISH or other technology]); or
    - 2.1.2 Patient has recurrent or de novo metastatic triple-negative breast cancer (that does not express ER, PR or HER2 IHC3+ or ISH+ [including FISH or other technology]); and
  - 2.2 Patient is treated with palliative intent; and
  - 2.3 Patient's cancer has confirmed PD-L1 Combined Positive Score (CPS) is greater than or equal to 10; and
  - 2.4 Patient has received no prior systemic therapy in the palliative setting; and
  - 2.5 Patient has an ECOG score of 0-2; and
  - 2.6 Pembrolizumab is to be used in combination with chemotherapy; and
  - 2.7 Baseline measurement of overall tumour burden is documented clinically and radiologically; and
  - 2.8 Pembrolizumab is to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks.

Renewal — (breast cancer, advanced) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 Patient's disease has had a complete response to treatment; or
  - 1.2 Patient's disease has had a partial response to treatment; or
  - 1.3 Patient has stable disease; and
- 2 No evidence of disease progression; and
- 3 Response to treatment in target lesions has been determined by a comparable radiologic assessment following the most recent treatment period: and
- 4 Pembrolizumab is to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 5 Treatment with pembrolizumab is to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initial application — (head and neck squamous cell carcinoma) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria: Either:

1 Patient is currently on treatment with pembrolizumab and met all remaining criteria prior to commencing treatment; or

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sul	bsidised	Generic
\$	Per	1	Manufacturer

continued...

- 2 All of the following:
  - 2.1 Patient has recurrent or metastatic head and neck squamous cell carcinoma of mucosal origin (excluding nasopharyngeal carcinoma) that is incurable by local therapies; and
  - 2.2 Patient has not received prior systemic therapy in the recurrent or metastatic setting; and
  - 2.3 Patient has a positive PD-L1 combined positive score (CPS) of greater than or equal to 1; and
  - 2.4 Patient has an ECOG performance score of 0-2; and
  - 2.5 Fither:
    - 2.5.1 Pembrolizumab to be used in combination with platinum-based chemotherapy; or
    - 2.5.2 Pembrolizumab to be used as monotherapy; and
  - 2.6 Pembrolizumab is to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks.

Renewal — (head and neck squamous cell carcinoma) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 Patient's disease has had a complete response to treatment; or
  - 1.2 Patient's disease has had a partial response to treatment: or
  - 1.3 Patient has stable disease; and
- 2 No evidence of disease progression; and
- 3 Pembrolizumab is to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 4 Treatment with pembrolizumab is to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initial application — (MSI-H/dMMR advanced colorectal cancer) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria: Fither:

- 1 Patient is currently on treatment with pembrolizumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
  - 2.1 Either:
    - 2.1.1 Patient has deficient mismatch repair (dMMR) or microsatellite instability-high (MSI-H) metastatic colorectal
    - 2.1.2 Patient has deficient mismatch repair (dMMR) or microsatellite instability-high (MSI-H) unresectable colorectal cancer; and
  - 2.2 Patient is treated with palliative intent: and
  - 2.3 Patient has not previously received funded treatment with pembrolizumab; and
  - 2.4 Patient has an ECOG performance score of 0-2; and
  - 2.5 Baseline measurement of overall tumour burden is documented clinically and radiologically; and
  - 2.6 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks.

**Renewal — (MSI-H/dMMR advanced colorectal cancer)** from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 3 Treatment with pembrolizumab is to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

**Initial application** — **(Urothelial carcinoma)** only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

Subs (Manufactur		
\$	Per	Manufacturer

continued...

Fither:

- 1 Patient is currently on treatment with pembrolizumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
  - 2.1 Patient has inoperable locally advanced (T4) or metastatic urothelial carcinoma; and
  - 2.2 Patient has an ECOG performance score of 0-2; and
  - 2.3 Patient has documented disease progression following treatment with chemotherapy; and
  - 2.4 Pembrolizumab to be used as monotherapy at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks

Renewal — (Urothelial carcinoma) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Any of the following:
  - 1.1 Patient's disease has had a complete response to treatment; or
  - 1.2 Patient's disease has had a partial response to treatment; or
  - 1.3 Patient has stable disease; and
- 2 No evidence of disease progression; and
- 3 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 4 Treatment with pembrolizumab is to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initial application — (relapsed/refractory Hodgkin lymphoma) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria: Fither:

- 1 Patient is currently on treatment with pembrolizumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
  - 2.1 Either:
    - 2.1.1 Both:
      - 2.1.1.1 Patient has relapsed/refractory Hodgkin lymphoma after two or more lines of chemotherapy; and
      - 2.1.1.2 Patient is ineligible for autologous stem cell transplant; or
    - 2.1.2 Patient has relapsed/refractory Hodgkin lymphoma and has previously undergone an autologous stem cell transplant: and
  - 2.2 Patient has not previously received funded pembrolizumab; and
  - 2.3 Pembrolizumab to be administered at doses no greater than 200 mg once every 3 weeks.

Renewal — (relapsed/refractory Hodgkin lymphoma) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient has received a partial or complete response to pembrolizumab; and
- 2 Treatment with pembrolizumab is to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

# Other Immunosuppressants

CICLOSPORIN			
Cap 25 mg	44.63	50	✓ Neoral
Cap 50 mg		50	✓ Neoral
Cap 100 mg		50	✓ Neoral
Oral liq 100 mg per ml	198.13	50 ml OP	✓ Neoral
EVEROLIMUS – Special Authority see SA2414 on the next page Wastage claimable	– Retail pharm	acy	
Tab 10 mg	6,512.29	30	✓ Afinitor
Tab 5 mg	4,555.76	30	Afinitor

				_
Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsid	lised	Generic	
\$	Per	✓	Manufacturer	

#### ⇒SA2414 Special Authority for Subsidy

Initial application only from a neurologist or oncologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has tuberous sclerosis: and
- 2 Patient has progressively enlarging sub-ependymal giant cell astrocytomas (SEGAs) that require treatment.

**Renewal** only from a neurologist or oncologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Documented evidence of SEGA reduction or stabilisation by MRI within the last 3 months; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Everolimus to be discontinued at progression of SEGAs.

**Initial application — (renal cell carcinoma)** from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 All of the following:
  - 1.1 The patient has metastatic renal cell carcinoma; and
  - 1.2 The disease is of predominant clear-cell histology; and
  - 1.3 The patient has documented disease progression following one previous line of treatment; and
  - 1.4 The patient has an ECOG performance status of 0-2; and
  - 1.5 Everolimus is to be used in combination with lenvatinib; or
- 2 All of the following:
  - 2.1 Patient has received funded treatment with nivolumab for the second line treatment of metastatic renal cell carcinoma; and
  - 2.2 Patient has experienced treatment limiting toxicity from treatment with nivolumab; and
  - 2.3 Everolimus is to be used in combination with lenvatinib; and
  - 2.4 There is no evidence of disease progression.

**Renewal — (renal cell carcinoma)** from any relevant practitioner. Approvals valid for 4 months where there is no evidence of disease progression.

CIDOLIMITIC	Special	Authority coo	CA2270 bolow	- Retail pharmacy
ZIROLIMOS .	– Special	Allinoriiv see	SAZZZU DEIOW -	- Refall bharmacy

Tab 1 mg	749.99	100	Rapamune
Tab 2 mg		100	✓ Rapamune
Oral liq 1 mg per ml	449.99	60 ml OP	Rapamune

#### ⇒SA2270 Special Authority for Subsidy

**Initial application** from any medical practitioner. Approvals valid without further renewal unless notified where the drug is to be used for rescue therapy for an organ transplant recipient.

Notes: Rescue therapy defined as unresponsive to calcineurin inhibitor treatment as defined by refractory rejection; or intolerant to calcineurin inhibitor treatment due to any of the following:

- GFR< 30 ml/min: or
- Rapidly progressive transplant vasculopathy; or
- Rapidly progressive obstructive bronchiolitis: or
- . HUS or TTP: or
- · Leukoencepthalopathy: or
- · Significant malignant disease

Initial application — (severe non-malignant lymphovascular malformations\*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

1 Patient has severe non-malignant lymphovascular malformation\*; and

Subsidy (Manufacturer's Price)	Su	Fully bsidised	Brand or Generic
\$	Per	1	Manufacturer

continued...

- 2 Any of the following:
  - 2.1 Malformations are not adequately controlled by sclerotherapy and surgery; or
  - 2.2 Malformations are widespread/extensive and sclerotherapy and surgery are not considered clinically appropriate; or
  - 2.3 Sirolimus is to be used to reduce malformation prior to consideration of surgery; and
- 3 Patient is being treated by a specialist lymphovascular malformation multi-disciplinary team; and
- 4 Patient has measurable disease as defined by RECIST version 1.1 (see Note).

Renewal — (severe non-malignant lymphovascular malformations\*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 Patient's disease has had either a complete response or a partial response to treatment, or patient has stable disease according to RECIST version 1.1 (see Note); or
  - 1.2 Patient's disease has stabilised or responded clinically and disease response to treatment has been clearly documents in patient notes; and
- 2 No evidence of progressive disease: and
- 3 The treatment remains clinically appropriate and the patient is benefitting from the treatment.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer et al. Eur J Cancer 2009;45:228-47)

Indications marked with \* are unapproved indications

Initial application — (renal angiomyolipoma(s) associated with tuberous sclerosis complex\*) only from a nephrologist or urologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has tuberous sclerosis complex\*; and
- 2 Evidence of renal angiomyolipoma(s) measuring 3 cm or greater and that have shown interval growth.

Renewal — (renal angiomyolipoma(s) associated with tuberous sclerosis complex\*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Documented evidence of renal angiomyolipoma reduction or stability by magnetic resonance imaging (MRI) or ultrasound; and
- 2 Demonstrated stabilisation or improvement in renal function; and
- 3 The patient has not experienced angiomyolipoma haemorrhage or significant adverse effects to sirolimus treatment; and
- 4 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indications marked with \* are unapproved indications

Initial application — (refractory seizures associated with tuberous sclerosis complex\*) only from a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has epilepsy with a background of documented tuberous sclerosis complex; and
- 2 Either:
  - 2.1 Both:
    - 2.1.1 Vigabatrin has been trialled and has not adequately controlled seizures; and
    - 2.1.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least two of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); or
  - 2.2 Both:
    - 2.2.1 Vigabatrin is contraindicated; and
    - 2.2.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least three of the following: sodium valproate, topiramate, levetiracetam,

Subsidy (Manufacturer's Price)	Fu Subsidise	,	
 \$	Per	<ul> <li>Manufacturer</li> </ul>	

continued...

carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); and

- 3 Seizures have a significant impact on quality of life; and
- 4 Patient has been assessed and surgery is considered inappropriate for this patient, or the patient has been assessed and would benefit from mTOR inhibitor treatment prior to surgery.

Note: Those of childbearing age potential are not required to trial phenytoin sodium, sodium valproate, or topiramate. Those who can father children are not required to trial sodium valproate.

Renewal — (refractory seizures associated with tuberous sclerosis complex\*) only from a neurologist. Approvals valid for 12 months where demonstrated significant and sustained improvement in seizure rate (e.g. 50% reduction in seizure frequency) or severity and/or patient quality of life compared with baseline prior to starting sirolimus treatment.

Note: Indications marked with \* are unapproved indications

TACROLIMUS - Special Authority see SA2271 below - Retail pharmacy

Cap 0.5 mg	49.60	100	✓ Tacrolimus Sandoz
Cap 0.75 mg	99.30	100	✓ Tacrolimus Sandoz
Cap 1 mg	84.30	100	✓ Tacrolimus Sandoz
Cap 5 mg	248.20	50	✓ Tacrolimus Sandoz

## ⇒SA2271 Special Authority for Subsidy

Initial application — (organ transplant) only from a relevant specialist. Approvals valid without further renewal unless notified where the patient is an organ transplant recipient.

Note: Subsidy applies for either primary or rescue therapy.

Initial application — (non-transplant indications\*) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient requires long-term systemic immunosuppression; and
- 2 Fither:
  - 2.1 Ciclosporin has been trialled and discontinued treatment because of unacceptable side effects or inadequate clinical response; or
  - 2.2 Patient is a child with nephrotic syndrome\*.

Note: Indications marked with \* are unapproved indications

#### JAK inhibitors

UPADACITINIB — Special Authority see SA2079 below — Retail pharmacy
Tab 15 mg .......1,271.00 28 ✓ RINVOQ

#### ⇒SA2079 Special Authority for Subsidy

Initial application — (Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
  - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
  - 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
- 3 Either:
  - 3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or
  - 3.2 Both:

Subsidy		Fully	Brand or
(Manufacturer's Price)	S Per	ubsidised	Generic Manufacturer
Ψ	1 61		Waltulacturei

continued...

- 3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a Health NZ Hospital; and 3.2.2 Either:
  - 3.2.2.1 The patient has experienced intolerable side effects from rituximab; or
    - 3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Renewal — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Either:

- 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic 
\$ Per ✓ Manufacturer

# **Antiallergy Preparations**

## Allergic Emergencies

ADRENALINE - Special Authority see SA2185 below - Retail pharmacy

- a) Maximum of 2 ini per prescription
- Additional prescriptions limited to replacement of up to two devices prior to expiry, or replacement of used device for treatment of anaphylaxis.

Inj 0.15 mg per 0.3 ml auto-injector	90.00	1 OP	Epipen Jr
Inj 0.3 mg per 0.3 ml auto-injector	90.00	1 OP	Epipen

## ⇒SA2185 Special Authority for Subsidy

Initial application — (anaphylaxis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 Patient has experienced an anaphylactic reaction which has resulted in presentation to a hospital or emergency department; or
  - 1.2 Patient has been assessed to be at significant risk of anaphylaxis by a relevant practitioner; and
- 2 Patient is not to be prescribed more than two devices in initial prescription.

#### ⇒SA1558 Special Authority for Subsidy

Initial application only from a clinical immunologist or relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Supply for anticipated emergency treatment of laryngeal/oro-pharyngeal or severe abdominal attacks of acute hereditary angioedema (HAE) for patients with confirmed diagnosis of C1-esterase inhibitor deficiency; and
- 2 The patient has undergone product training and has agreed upon an action plan for self-administration.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

# **Allergy Desensitisation**

#### ⇒SA1367 Special Authority for Subsidy

Initial application only from a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 RAST or skin test positive; and
- 2 Patient has had severe generalised reaction to the sensitising agent.

**Renewal** only from a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

BEE VENOM ALLERGY TREATMENT - Special Authority see SA1367 above - Retail pharmacy Initiation kit - 5 yials freeze dried venom with diluent 305.00 1 OP ✓ VENOX S29 1 OP ✓ VENOX S29 Maintenance kit - 6 vials 120 mcg freeze dried venom, with 1 OP ✓ Venomil S29 Treatment kit - 1 vial 550 mcg freeze dried venom, 1 diluent 1 OP ✓ Albev Treatment kit - 1 vial 550 mcg freeze dried venom, with diluent ..... 305.00 1 OP ✓ Hymenoptera S29

	Subsidy		Fully Brand or	
	(Manufacturer's Pr		idised Generic	
/AOD VENOMALLEDOV TREATMENT Or side Authority	\$	Per	✓ Manufacturer	
VASP VENOM ALLERGY TREATMENT — Special Authority see	e SA1367 on the	previous page	– Hetaii pnarmacy	
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze dried polistes venom, 1 diluent 9 ml, 3 diluent 1.8 ml	382.23	1 OP	✓ Albey	
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze dried venom, with diluent	305.00	1 OP	✓ Hymenoptera S2S	9
Treatment kit (Paper wasp venom) - 6 vials 120 mcg freeze dried venom, with diluent	305.00	1 OP	✓ Venomil S29	
Treatment kit (Yellow Jacket venom) - 1 vial 550 mcg freeze dried venom, with diluent	305.00	1 OP	✓ Hymenoptera S29	9)
Treatment kit (Yellow jacket venom) - 1 vial 550 mcg freeze dried vespula venom, 1 diluent 9 ml, 3 diluent 1.8 ml	431.24	1 OP	✓ Albey	
Treatment kit (Yellow jacket venom) - 6 vials 120 mcg freeze dried venom, with diluent		1 OP	✓ Venomil S29	
Antihistamines				
ETIRIZINE HYDROCHLORIDE				
₭ Tab 10 mg	1.71	100	✓ Zista	
Oral liq 1 mg per ml		200 ml	✓ Histaclear	
EXTROCHLORPHENIRAMINE MALEATE				
← Tab 2 mg	2.02	40		
	(8.40)		Polaramine	
	1.01	20		
	(5.99)		Polaramine	
Oral liq 2 mg per 5 ml		100 ml		
	(10.29)		Polaramine	
EXOFENADINE HYDROCHLORIDE				
Fab 60 mg	4.34	20		
1 /00	(8.23)		Telfast	
← Tab 120 mg		10	T-164	
	(8.23)	20	Telfast	
	14.22	30	Telfast	
	(26.44)		renasi	
ORATADINE				
* Tab 10 mg	1.78	100	✓ <u>Lorafix</u>	
Gral liq 1 mg per ml	1.43	100 ml	Haylor syrup	
ROMETHAZINE HYDROCHLORIDE				
€ Tab 10 mg		50	Allersoothe	
€ Tab 25 mg		50	✓ <u>Allersoothe</u>	
Oral liq 1 mg per 1 ml		100 ml	✓ Allersoothe	
. In OF many and Outlemands. The tellinian like and F	10.47	-	✓ Phenergan Elixir	r
	<sup>2</sup> SO21.09	5	✓ Hospira	
Inhaled Corticosteroids				
BECLOMETHASONE DIPROPIONATE				
Aerosol inhaler, 50 mcg per dose		200 dose OP	✓ Qvar	
Aerosol inhaler, 50 mcg per dose CFC-free		200 dose OP	✓ Beclazone 50	
Aerosol inhaler, 100 mcg per dose	17.52	200 dose OP	✓ Qvar	
Aerosol inhaler, 100 mcg per dose CFC-free		200 dose OP	✓ Beclazone 100	
Aerosol inhaler, 250 mcg per dose CFC-free	22.67	200 dose OP	✓ Beclazone 250	

	Subsidy (Manufacturer's	Price) Sub	Fully Brand or osidised Generic
	` \$	Per	✓ Manufacturer
BUDESONIDE			
Powder for inhalation, 100 mcg per dose	17.00	200 dose OF	Pulmicort Turbuhaler
Powder for inhalation, 200 mcg per dose	19.00	200 dose OF	Pulmicort Turbuhaler
Powder for inhalation, 400 mcg per dose	32.00	200 dose OF	Pulmicort Turbuhaler
LUTICASONE			
Aerosol inhaler, 50 mcg per dose	7.19	120 dose OF	✓ Flixotide
Powder for inhalation, 50 mcg per dose		60 dose OP	✓ Flixotide Accuhaler
Powder for inhalation, 100 mcg per dose		60 dose OP	✓ Flixotide Accuhaler
Aerosol inhaler, 125 mcg per dose		120 dose OF	✓ Flixotide
Aerosol inhaler, 250 mcg per dose		120 dose OF	✓ Flixotide
Powder for inhalation, 250 mcg per dose	11.93	60 dose OP	✓ Flixotide Accuhaler
Inhaled Long-acting Beta-adrenoceptor Agonis	ts		
FORMOTEROL FUMARATE DIHYDRATE			
Powder for inhalation 4.5 mcg per dose, breath activated			
(equivalent to eformoterol fumarate 6 mcg metered dose	10.32	60 dose OP	
(equivalent to clothoteror furnarate of fleg frictered dose	(16.90)	00 0030 01	Oxis Turbuhaler
VD 4 C 4 T T D C 4	(10.50)		Oxio Turburialer
NDACATEROL			
Powder for inhalation 150 mcg		30 dose OP	
Powder for inhalation 300 mcg	61.00	30 dose OP	Onbrez Breezhaler
ALMETEROL			
Aerosol inhaler CFC-free, 25 mcg per dose	26.25	120 dose OF	✓ Serevent
Powder for inhalation, 50 mcg per dose, breath activated	26.25	60 dose OP	Serevent Accuhaler
Inhaled Corticosteroids with Long-Acting Beta-	Adrenocept	or Agonists	s
SUDESONIDE WITH EFORMOTEROL			
Powder for inhalation 160 mcg with 4.5 mcg eformoterol			
fumarate per dose (equivalent to 200 mcg budesonide w	vith		
6 mcg eformoterol fumarate metered dose)		120 dose OF	✓ DuoResp Spiromax
Powder for inhalation 320 mcg with 9 mcg eformoterol fumar		120 0030 01	5 Buorlesp opiromax
per dose (equivalent to 400 mcg budesonide with 12 mc			
eformoterol fumarate metered dose) – No more than 2	g		
·	90.50	120 dose OF	A DuoBoon Spiromov
dose per day			
Aerosol inhaler 100 mcg with eformaterol fumarate 6 mcg		120 dose OF	
Powder for inhalation 100 mcg with eformoterol fumarate 6 n	ncg33.74	120 dose OF	O Symbicort Turbuhaler 100/6
A 111 1 200 111 1 1 1 1 1 1 1	24.42	400 1 05	
Aerosol inhaler 200 mcg with eformoterol fumarate 6 mcg		120 dose OF	
Powder for inhalation 200 mcg with eformoterol fumarate 6 n	ncg33.74	120 dose OF	
			Turbuhaler 200/6
Powder for inhalation 400 mcg with eformoterol fumarate			
12 mcg - No more than 2 dose per day	33.74	60 dose OP	•
			Turbuhaler 400/12
LUTICASONE FUROATE WITH VILANTEROL			
Powder for inhalation 100 mcg with vilanterol 25 mcg	44.08	30 dose OP	✓ Breo Ellipta
ů ů			•

	Subsidy		Fully	
	(Manufacturer's I \$	Price) Sub Per	sidised •	
LUTICA CONE WITH CALMETEROL	Ψ	101		Mariaracturer
LUTICASONE WITH SALMETEROL  Aerosol inhaler 50 mcg with salmeterol 25 mcg	25.70	120 dose OP	1	Seretide
Aerosol inhaler 30 mcg with salmeterol 25 mcg		120 dose OF		Seretide
Powder for inhalation 100 mcg with salmeterol 50 mcg – No	02.00	120 0000 01	•	Colonido
more than 2 dose per day	33.74	60 dose OP	1	Seretide Accuhaler
Powder for inhalation 250 mcg with salmeterol 50 mcg - No				
more than 2 dose per day	44.08	60 dose OP	•	Seretide Accuhaler
Beta-Adrenoceptor Agonists				
ALBUTAMOL				
Oral lig 400 mcg per ml	50.00	150 ml	1	Ventolin
Infusion 1 mg per ml, 5 ml		10		Ventolin
Inj 500 mcg per ml, 1 ml - Up to 5 inj available on a PSO		5	•	Ventolin
Inhaled Beta-Adrenoceptor Agonists				
ALBUTAMOL				
Aerosol inhaler, 100 mcg per dose CFC free - Up to 1000				
dose available on a PSO	3.80	200 dose OP	1	SalAir
	(6.80)			Ventolin
Nebuliser soln, 1 mg per ml, 2.5 ml ampoule - Up to 30 neb				
available on a PSO	8.96	20	1	Asthalin
Nebuliser soln, 2 mg per ml, 2.5 ml ampoule – Up to 30 neb	0.40	00	,	A - Al III-
available on a PSO	9.43	20	•	Asthalin
ERBUTALINE SULPHATE				
Powder for inhalation, 200 mcg per dose (equivalent to	00.00	400 de e OD	,	Bullian and Treatment along
250 mcg metered dose), breath activated	22.20	120 dose OP	•	Bricanyl Turbuhaler
Anticholinergic Agents				
PRATROPIUM BROMIDE				
Aerosol inhaler, 20 mcg per dose CFC-free - Up to 400 dose				
available on a PSO	16.20	200 dose OP	1	Atrovent
Nebuliser soln, 250 mcg per ml, 2 ml ampoule - Up to 40 neb	)			
available on a PSO	5.86	10		Pharmascience S29
	11.73	20	1	Ipratropium
				IVAX S29
			1	Univent
Pharmascience S29 Nebuliser soln, 250 mcg per ml, 2 ml ampo	ule to be delist	ed 1 May 2025	)	
pratropium IVAX S29 Nebuliser soln, 250 mcg per ml, 2 ml amp	oule to be deli	sted 1 Februar	/ 2025	5)
Inhaled Beta-Adrenoceptor Agonists with Antich	olinergic A	gents		
ALBUTAMOL WITH IPRATROPIUM BROMIDE				
Aerosol inhaler, 100 mcg with ipratropium bromide, 20 mcg pe	er			
dose CFC-free		200 dose OP	1	Duolin HFA
Nebuliser soln, 2.5 mg with ipratropium bromide 0.5 mg per				
vial, 2.5 ml ampoule – Up to 20 neb available on a PSO.	11.04	20	1	Duolin
			./	Duralin Cinta on
			v	Duolin Cipla S29

260

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic 
\$ Per ✔ Manufacturer

## **Long-Acting Muscarinic Antagonists**

GLYCOPYRRONIUM - Subsidy by endorsement

- a) Inhaled glycopyrronium treatment will not be subsidised if patient is also receiving treatment with subsidised tiotropium or umeclidinium
- b) Glycopyrronium powder for inhalation 50 mcg per dose is subsidised only for patients who have been diagnosed as having COPD using spirometry if spirometry is possible, and the prescription is endorsed accordingly.

TIOTROPIUM BROMIDE - Subsidy by endorsement

- a) Tiotropium treatment will not be subsidised if patient is also receiving treatment with subsidised inhaled glycopyrronium or umeclidinium.
- b) Tiotropium bromide is subsidised only for patients who have been diagnosed as having COPD using spirometry if spirometry is possible, and the prescription is endorsed accordingly. Patients who had tiotropium dispensed before 1 October 2018 with a valid Special Authority are deemed endorsed.

#### UMECLIDINIUM - Subsidy by endorsement

- a) Umeclidinium will not be subsidised if patient is also receiving treatment with subsidised inhaled glycopyrronium or tiotropium bromide.
- b) Umeclidinium powder for inhalation 62.5 mcg per dose is subsidised only for patients who have been diagnosed as having COPD using spirometry if spirometry is possible, and the prescription is endorsed accordingly.

# Long-Acting Muscarinic Antagonists with Long-Acting Beta-Adrenoceptor Agonists

Combination long acting muscarinic antagonist and long acting beta-2 agonist will not be subsidised if patient is also receiving treatment with a combination inhaled corticosteroid and long acting beta-2 agonist.

#### ⇒SA1584 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Patient has been stabilised on a long acting muscarinic antagonist; and
- 2 The prescriber considers that the patient would receive additional benefit from switching to a combination product.

**Renewal** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Patient is compliant with the medication; and
- 2 Patient has experienced improved COPD symptom control (prescriber determined).

GLYCOPYRRONIUM WITH INDACATEROL — Special Authority see SA1584 above — Retail pharmacy
Powder for Inhalation 50 mcg with indacaterol 110 mcg......81.00 30 dose OP ✓ Ultibro Breezhaler

TIOTROPIUM BROMIDE WITH OLODATEROL - Special Authority see SA1584 above - Retail pharmacy

UMECLIDINIUM WITH VILANTEROL - Special Authority see SA1584 above - Retail pharmacy

# Inhaled Corticosteroid with Long-Acting Muscarinic Antagonist and Beta Agonist

BUDESONIDE WITH GLYCOPYRRONIUM AND EFORMOTEROL – Special Authority see SA2421 on the next page – Retail pharmacy

Aerosol inhaler budesonide 160 mcg with glycopyrronium

7.2 mcg and formoterol 5 mcg per dose......79.15 120 dose OP ✓ Breztri Aerosphere

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

#### ⇒SA2421 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

# Both:

- 1 Patient has a diagnosis of COPD confirmed by spirometry or spirometry has been attempted and technically acceptable results are not possible; and
- 2 Either:
  - 2.1 Both:
    - 2.1.1 Patient is currently receiving an inhaled corticosteroid with long acting beta-2 agonist (ICS/LABA) or a long acting muscarinic antagonist with long acting beta-2 agonist (LAMA/LABA); and
    - 2.1.2 Any of the following:

Clinical criteria:

- 2.1.2.1 Patient has a COPD Assessment Test (CAT) score greater than 10; or
- 2.1.2.2 Patient has had 2 or more exacerbations in the previous 12 months; or
- 2.1.2.3 Patient has had one exacerbation requiring hospitalisation in the previous 12 months; or
- 2.1.2.4 Patient has had an eosinophil count greater than or equal to  $0.3 \times 10^9$  cells/L in the previous 12 months; or
- 2.2 Patient is currently receiving multiple inhaler triple therapy (inhaled corticosteroid with long-acting muscarinic antagonist and long-acting beta-2 agonist ICS/LAMA/LABA) and met at least one of the clinical criteria above prior to commencing multiple inhaler therapy.

FLUTICASONE FUROATE WITH UMECLIDINIUM AND VILANTEROL – Special Authority see SA2326 below – Retail pharmacy

Powder for inhalation fluticasone furoate 100 mcg with umeclidinium 62.5 mcg and vilanterol 25 mcg......104.24

104.24 30 dose OP

✓ Trelegy Ellipta

## **⇒SA2326** Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has a diagnosis of COPD confirmed by spirometry or spirometry has been attempted and technically acceptable results are not possible; and
- 2 Either:
  - 2.1 Both:
    - 2.1.1 Patient is currently receiving an inhaled corticosteroid with long acting beta-2 agonist (ICS/LABA) or a long acting muscarinic antagonist with long acting beta-2 agonist (LAMA/LABA); and
    - 2.1.2 Any of the following:

Clinical criteria:

- 2.1.2.1 Patient has a COPD Assessment Test (CAT) score greater than 10; or
- 2.1.2.2 Patient has had 2 or more exacerbations in the previous 12 months; or
- 2.1.2.3 Patient has had one exacerbation requiring hospitalisation in the previous 12 months; or
- 2.1.2.4 Patient has had an eosinophil count greater than or equal to  $0.3 \times 10^{\circ}9$  cells/L in the previous 12 months; or
- 2.2 Patient is currently receiving multiple inhaler triple therapy (inhaled corticosteroid with long acting muscarinic antagonist and long acting beta-2 agonist ICS/LAMA/LABA) and met at least one of the clinical criteria above prior to commencing multiple inhaler triple therapy.

## **Antifibrotics**

NINTEDANIB - Special Authority see SA2012 on the next page - Retail pharmacy

Note: Nintedanib not subsidised in combination with subsidised pirfenidone.

Cap 100 mg	2,554.00	60 OP	<ul><li>Ofev</li></ul>
Cap 150 mg	3,870.00	60 OP	Ofev

Subsidy		Fully	Brand or	
(Manufacturer's Price)	:	Subsidised	Generic	
\$	Per	✓	Manufacturer	

#### ⇒SA2012 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Nintedanib is to be discontinued at disease progression (See Note); and
- 4 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 5 Any of the following:
  - 5.1 The patient has not previously received treatment with pirfenidone; or
  - 5.2 Patient has previously received pirfenidone, but discontinued pirfenidone within 12 weeks due to intolerance; or
  - 5.3 Patient has previously received pirfenidone, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with pirfenidone).

**Renewal — (idiopathic pulmonary fibrosis)** only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 3 Nintedanib is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

PIRFENIDONE – Retail pharmacy-Specialist – Special Authority see SA2013 below Note: Pirfenidone is not subsidised in combination with subsidised nintedanib.

Tab 801 mg	3,645.00	90 OP	✓ Esbriet
Tab 267 mg	1,215.00	90	<ul><li>Esbriet</li></ul>

#### ⇒SA2013 Special Authority for Subsidy

**Initial application — (idiopathic pulmonary fibrosis)** only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note); and
- 4 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 5 Any of the following:
  - 5.1 The patient has not previously received treatment with nintedanib; or
  - 5.2 Patient has previously received nintedanib, but discontinued nintedanib within 12 weeks due to intolerance; or
  - 5.3 Patient has previously received nintedanib, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with nintedanib).

**Renewal — (idiopathic pulmonary fibrosis)** only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	
Leukotriene Receptor Antagonists				
MONTELUKAST  * Tab 4 mg  * Tab 5 mg  * Tab 10 mg	3.10	28 28 28	/	Montelukast Viatris Montelukast Viatris Montelukast Viatris
Methylxanthines				
AMINOPHYLLINE  * Inj 25 mg per ml, 10 ml ampoule – Up to 5 inj available on a PSO		5	•	DBL Aminophylline
THEOPHYLLINE  * Tab long-acting 250 mg  * Oral liq 80 mg per 15 ml		100 00 m		Nuelin-SR Nuelin
Mucolytics				
DORNASE ALFA – Special Authority see SA1978 below – Retai Nebuliser soln, 2.5 mg per 2.5 ml ampoule		6	1	Pulmozyme

#### ⇒SA1978 Special Authority for Subsidy

Initial application — (cystic fibrosis) only from a respiratory physician or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of cystic fibrosis: and
- 2 Patient has previously undergone a trial with, or is currently being treated with, hypertonic saline; and
- 3 Any of the following:
  - 3.1 Patient has required one or more hospital inpatient respiratory admissions in the previous 12 month period; or
  - 3.2 Patient has had 3 exacerbations due to CF, requiring oral or intravenous (IV) antibiotics in the previous 12 month period: or
  - 3.3 Patient has had 1 exacerbation due to CF, requiring oral or IV antibiotics in the previous 12 month period and a Brasfield score of < 22/25; or</p>
  - 3.4 Patient has a diagnosis of allergic bronchopulmonary aspergillosis (ABPA).

**Renewal** — **(cystic fibrosis)** only from a respiratory physician or paediatrician. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient continues to benefit from treatment.

ELEXACAFTOR WITH TEZACAFTOR, IVACAFTOR AND IVACAFTOR - PCT only - Special Authority see SA2196 below

Tab elexacaftor 50 mg with tezacaftor 25 mg, ivacaftor 37.5 mg (56) and ivacaftor 75 mg (28) .......27,647.39

84 OP ✓ Trikafta

Tab elexacaftor 100 mg with tezacaftor 50 mg, ivacaftor 75 mg

(56) and ivacaftor 150 mg (28) ......27,647.39

84 OP ✓ Trikafta

#### ⇒SA2196 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with cystic fibrosis; and
- 2 Patient is 6 years of age or older; and
- 3 Fither:

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsi	dised	Generic
 \$	Per	1	Manufacturer

#### continued...

- 3.1 Patient has two cystic fibrosis-causing mutations in the cystic fibrosis transmembrane regulator (CFTR) gene (one from each parental allele); or
- 3.2 Patient has a sweat chloride value of at least 60 mmol/L by quantitative pilocarpine iontophoresis or by Macroduct sweat collection system; and
- 4 Fither:
  - 4.1 Patient has a heterozygous or homozygous F508del mutation; or
  - 4.2 Patient has a G551D mutation or other mutation responsive in vitro to elexacaftor/tezacaftor/ivacaftor (see note a); and
- 5 The treatment must be the sole funded CFTR modulator therapy for this condition; and
- 6 Treatment with elexacaftor/tezacaftor/ivacaftor must be given concomitantly with standard therapy for this condition.

#### Notes:

 a) Eligible mutations are listed in the Food and Drug Administration (FDA) Trikafta prescribing information https://www.accessdata.fda.gov/drugsatfda\_docs/label/2021/212273s004lbl.pdf

#### IVACAFTOR - PCT only - Specialist - Special Authority see SA2017 below

Tab 150 mg	29,386.00	56	✓ Kalydeco
Oral granules 50 mg, sachet	29,386.00	56	✓ Kalydeco
Oral granules 75 mg, sachet	29,386.00	56	✓ Kalydeco

#### **⇒SA2017** Special Authority for Subsidy

Initial application only from a respiratory specialist or paediatrician. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### All of the following:

- 1 Patient has been diagnosed with cystic fibrosis; and
- 2 Either:
  - 2.1 Patient must have G551D mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene on at least 1 allele; or
  - 2.2 Patient must have other gating (class III) mutation (G1244E, G1349D, G178R, G551S, S1251N, S1255P, S549N and S549R) in the CFTR gene on at least 1 allele; and
- 3 Patients must have a sweat chloride value of at least 60 mmol/L by quantitative pilocarpine iontophoresis or by Macroduct sweat collection system; and
- 4 Treatment with ivacaftor must be given concomitantly with standard therapy for this condition; and
- 5 Patient must not have an acute upper or lower respiratory infection, pulmonary exacerbation, or changes in therapy (including antibiotics) for pulmonary disease in the last 4 weeks prior to commencing treatment with ivacaftor; and
- 6 The dose of ivacaftor will not exceed one tablet or one sachet twice daily; and
- 7 Applicant has experience and expertise in the management of cystic fibrosis.

#### SODIUM CHLORIDE

Not funded for use as a nasal drop.

## **Nasal Preparations**

## **Allergy Prophylactics**

#### BUDESONIDE

Metered aqueous nasal spray, 50 mcg per dose2.59	200 dose OP	✓ SteroClear
SteroClear to be Principal Supply on 1 February 2025		
Metered aqueous nasal spray, 100 mcg per dose2.89	200 dose OP	✓ SteroClear
SteroClear to be Principal Supply on 1 February 2025		

	Subsidy (Manufacturer's \$	Price) Subsi	Fully Brand or idised Generic Manufacturer
LUTICASONE PROPIONATE			
Metered aqueous nasal spray, 50 mcg per dose	1.98	120 dose OP	✓ Flixonase Hayfever & Allergy
PRATROPIUM BROMIDE			
Aqueous nasal spray, 0.03%	5.23	15 ml OP	✓ Univent
Respiratory Devices			
ASK FOR SPACER DEVICE			
a) Up to 50 dev available on a PSO			
b) Only on a PSO			
<ul> <li>c) Only for children aged six years and under</li> </ul>			_
Small	2.70	1	<ul><li>e-chamber Mask</li></ul>
EAK FLOW METER			
<ul> <li>a) Up to 25 dev available on a PSO</li> </ul>			
b) Only on a PSO			<b>4</b>
Low range	9.54	1	✓ Mini-Wright AFS
Normal range	0.54	1	Low Range  ✓ Mini-Wright
Norma range	9.04	'	Standard
PACER DEVICE			Otanidard
a) Up to 50 dev available on a PSO     b) Only on a PSO			
220 ml (single patient)	3.65	1	✓ e-chamber Turbo
510 ml (single patient)		i	✓ e-chamber La
, ,			Grande
800 ml	6.50	1	✓ Volumatic

Oral liq 20 mg per ml (10 mg base per ml)......16.91 25 ml OP

✓ Biomed

CAFFEINE CITRATE

	Subsidy (Manufacturer's Pr \$	rice) Subs	Fully Brand or sidised Generic ✓ Manufacturer
Ear Preparations			
FLUMETASONE PIVALATE Ear drops 0.02% with clioquinol 1%	4.46	7.5 ml OP	✓ Locacorten-Viaform ED's ✓ Locorten-Vioform
TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYC Ear drops 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 mg and gramicidin 250 mcg per g		7.5 ml OP	✓ Kenacomb
Ear/Eye Preparations			
DEXAMETHASONE WITH FRAMYCETIN AND GRAMICIDIN Ear/Eye drops 500 mcg with framycetin sulphate 5 mg and gramicidin 50 mcg per ml	4.50 (9.27) (9.27)	8 ml OP	Otodex \$29 Sofradex
FRAMYCETIN SULPHATE Ear/Eye drops 0.5%	4.13 (8.65)	8 ml OP	Soframycin
Eye Preparations			
Eye preparations are only funded for use in the eye, unless expl	icitly stated otherv	vise.	
Anti-Infective Preparations			
* Eye oint 3%  ViruPOS to be Principal Supply on 1 February 2025	15.89	4.5 g OP	✓ ViruPOS
CHLORAMPHENICOL  Eye oint 1%  Eye drops 0.5%  Funded for use in the ear*. Indications marked with * a	1.45	5 g OP 10 ml OP dications.	✓ <u>Devatis</u> ✓ <u>Chlorsig</u>
CIPROFLOXACIN  Eye drops 0.3% — Subsidy by endorsement	itis or severe bactorive otitis media (CS unapproved indica	SOM)*; and th	
SODIUM FUSIDATE [FUSIDIC ACID] Eye drops 1%	5.29	5 g OP	✓ Fucithalmic ✓ Fucithalmic S29 S29
TOBRAMYCIN	40.45	0.5 0.0	/ Tabasa

3.5 g OP

5 ml OP

✓ Tobrex

✓ Tobrex



Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per		Manufacturer

## **Corticosteroids and Other Anti-Inflammatory Preparations**

*	Eye oint 0.1%	3.5 g OP	Maxidex
*	Eye drops 0.1%	5 ml OP	<ul><li>Maxidex</li></ul>
	Ocular implant 700 mcg - Special Authority see SA1680 below		
	- Retail pharmacy	1	<ul><li>Ozurdex</li></ul>

#### **⇒SA1680** Special Authority for Subsidy

Initial application — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has diabetic macular oedema with pseudophakic lens; and
- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- 3 Fither
  - 3.1 Patient's disease has progressed despite 3 injections with bevacizumab; or
  - 3.2 Patient is unsuitable or contraindicated to treatment with anti-VEGF agents; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Initial application — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has diabetic macular oedema; and
- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- 3 Patient is of child bearing potential and has not yet completed a family; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Patient is of child bearing potential and has not yet completed a family; and
- 3 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

#### DEXAMETHASONE WITH NEOMYCIN SULPHATE AND POLYMYXIN B SULPHATE

* Eye oint 0.1% with neomycin sulphate 0.35% and polymyxin b			
sulphate 6,000 u per g	5.39	3.5 g OP	✓ Maxitrol
* Eye drops 0.1% with neomycin sulphate 0.35% and polymyxin		•	
b sulphate 6,000 u per ml	4.50	5 ml OP	✓ Maxitrol
FLUOROMETHOLONE			
* Eye drops 0.1%	3.09	5 ml OP	✓ FML
,	5.20		✓ Flucon
LEVOCABASTINE			
Eye drops 0.5 mg per ml	8.71	4 ml OP	
, , ,	(10.34)		Livostin

			SEN	ISORY ORGANS
	Subsidy (Manufacturer's P \$	rice) Subsi Per	Fully idised	Brand or Generic Manufacturer
LODOXAMIDE				
Eye drops 0.1%	8.71	10 ml OP	•	Lomide
NEPAFENAC				
Eye drops 0.3%	8.80	3 ml OP	1	llevro
PREDNISOLONE ACETATE				
Eye drops 1%		10 ml OP		Prednisolone-AFT
	7.00	5 ml OP		Pred Forte
PREDNISOLONE SODIUM PHOSPHATE - Special Authority s	ee SA1715 below	– Retail pharm		
Eye drops 0.5%, single dose (preservative free)	43.26	20 dose	1	Minims Prednisolone
SA1715   Special Authority for Subsidy	n eye drops. onths where the t		ins ap	
Eye drops 2%	2.02	10 1111 OP	•	Alleriix
Glaucoma Preparations - Beta Blockers				
BETAXOLOL  * Eye drops 0.25%  * Eye drops 0.5%	7.50	5 ml OP 5 ml OP	•	Betoptic S Betoptic
* Eye drops 0.25%  * Eye drops 0.5%		5 ml OP 5 ml OP		Arrow-Timolol Arrow-Timolol
Glaucoma Preparations - Carbonic Anhydrase	Inhibitors			
ACETAZOLAMIDE  * Tab 250 mg  BRINZOLAMIDE  * Eye drops 1%		100 5 ml OP		Diamox <u>Azopt</u>
DORZOLAMIDE WITH TIMOLOL  * Eye drops 2% with timolol 0.5%  Dortimopt to be Principal Supply on 1 February 2025	3.58	5 ml OP	1	Dortimopt

# Glaucoma Preparations - Prostaglandin Analogues

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy		Fully	Brand or
	(Manufacturer's Pri	ice) Sub: Per	sidised •	Generic Manufacturer
TRAVOPROST	· ·			
* Eye drops 0.004%	6.80	2.5 ml OP	1	<u>Travatan</u>
Glaucoma Preparations - Other				
BRIMONIDINE TARTRATE				
* Eye drops 0.2%		5 ml OP	•	Arrow-Brimonidine
BRIMONIDINE TARTRATE WITH TIMOLOL MALEATE				
* Eye drops 0.2% with timolol maleate 0.5%	7.13	5 ml OP	1	<u>Combigan</u>
LATANOPROST WITH TIMOLOL				
* Eye drops 0.005% with timolol 0.5%	4.95	2.5 ml OP	✓ .	Arrow - Lattim
PILOCARPINE HYDROCHLORIDE				
* Eye drops 1%	4.26	15 ml OP	✓	Isopto Carpine
* Eye drops 2%		15 ml OP	✓	Isopto Carpine
* Eye drops 4%		15 ml OP	✓	Isopto Carpine
Subsidised for oral use pursuant to the Standard Formula	ae.			
PILOCARPINE NITRATE				
* Eye drops 2% single dose - Special Authority see SA0895				
below – Retail pharmacy	35.90	20 dose	1	Minims Pilocarpine
TO CARROLL Crossed Authority for Cubaidy				•

## SA0895 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient has to use an unpreserved solution due to an allergy to the preservative; or
- 2 Patient wears soft contact lenses.

**Mydriatics and Cycloplegics** 

Note: Minims for a general practice are considered to be "tools of trade" and are not approved as special authority items. Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

ATROPINE SULPHATE	
* Eye drops 1%18.27	15 ml OP
CYCLOPENTOLATE HYDROCHLORIDE	
* Evo drope 19/	15 ml OD

✓ Atropt

Cyclogyl 15 ml OF Eye drops 1%, single dose (preservative free) - Only on a

20 dose ✓ Minims prescription......84.85 Cyclopentolate

(Minims Cyclopentolate Eye drops 1%, single dose (preservative free) to be delisted 1 February 2025) **TROPICAMIDE** 

*	Eye drops 0.5%	15 ml OP	✓ Mydriacyl
*	Eye drops 1%24.82	15 ml OP	Mydriacyl

# **Preparations for Tear Deficiency**

For acetylcysteine eye drops refer Standard Formulae, page 274

HY	PROMELLOSE
*	Eye drops 0.5%

**Principal Supply** 

*	Eye drops 0.5%	19.50	15 ml OP	Methopt
HY	PROMELLOSE WITH DEXTRAN			
*	Eye drops 0.3% with dextran 0.1%	2.30	15 ml OP	✓ Poly-Tears

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

#### **Preservative Free Ocular Lubricants**

#### ⇒SA2134 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Confirmed diagnosis by slit lamp or Schirmer test of severe secretory dry eye; and
- 2 Either:
  - 2.1 Patient is using eye drops more than four times daily on a regular basis; or
  - 2.2 Patient has had a confirmed allergic reaction to preservative in eye drop.

**Renewal** from any relevant practitioner. Approvals valid for 24 months where the patient continues to require lubricating eye drops and has benefited from treatment.

CARBOMER – Special Authority see SA2134 above – Retail pharmac Ophthalmic gel 0.3%, 0.5 g(Poly-Gel Ophthalmic gel 0.3%, 0.5 g to be delisted 1 July 2025)	,	30	✓ Poly-Gel
POLYETHYLENE GLYCOL 400 AND PROPYLENE GLYCOL - Speci Eye drops 0.4% and propylene glycol 0.3%, 0.8 ml			ove – Retail pharmacy  ✓ Systane Unit Dose
SODIUM HYALURONATE [HYALURONIC ACID] - Special Authority s	see SA2134 abo	ve – Retail	pharmacy
Eye drops 1 mg per ml			
Hylo-Fresh has a 6 month expiry after opening. The Pharmac month is not relevant and therefore only the prescribed dosage			

Other Ev	e Preparations
----------	----------------

NAPHAZOLINE HYDROCHLORIDE  * Eye drops 0.1%5.65	15 ml OP	✓ <u>Albalon</u>
OLOPATADINE  Eye drops 0.1%2.17	5 ml OP	✓ Olopatadine Teva
PARAFFIN LIQUID WITH WOOL FAT  * Eye oint 3% with wool fat 3%	3.5 g OP	✓ Poly-Visc
RETINOL PALMITATE  Eve girt 138 mcg per g 3 80	5 a OP	✓ VitA-POS



VARIOUS				
	Subsidy (Manufacturer's Prio \$	ce) Per	Fully Subsidised	
	<b>3</b>	Per		Manufacturer
Various				
PHARMACY SERVICES				
* Brand switch fee	4.50	1 fee	•	BSF Midazolam-Baxter
a) May only be claimed once per patient.				
b) The Pharmacode for BSF Midazolam-Baxter is 26958	863 - see also <mark>pag</mark>	e 144		
* Immunisation administration fee - flu		1 fee		Immunisation - Flu
* Immunisation administration fee - other		1 fee		Immunisation Other
* Immunisation co-administration fee - flu and shingles	0.00	1 fee	•	Immunisation Flu and Shingles
(BSF Midazolam-Baxter Brand switch fee to be delisted 1 February	ary 2025)			·
Agents Used in the Treatment of Poisonings				
Antidotes				
ACETYLCYSTEINE				
Inj 200 mg per ml, 10 ml ampoule	42.99 52.88	10		DBL Acetylcysteine Martindale Pharma
DBL Acetylcysteine to be Principal Supply on 1 April 202 (Martindale Pharma Inj 200 mg per ml, 10 ml ampoule to be delis	25			
NALOXONE HYDROCHLORIDE	, ,			
a) Up to 10 inj available on a PSO				
b) Only on a PSO				
# Inj 400 mcg per ml, 1 ml ampoule	13.29	5	/	DBL Naloxone
,				Hydrochloride
	35.26	10	1	Hameln
DBL Naloxone Hydrochloride to be Principal Supply on 1 (Hameln Inj 400 mcg per ml, 1 ml ampoule to be delisted 1 April 2				
Removal and Elimination				
CHARCOAL				
* Oral liq 50 g per 250 ml	43.50	250 ml (	OP 🗸	Carbosorb-X
a) Up to 250 ml available on a PSO b) Only on a PSO				
DEFERASIROX - Special Authority see SA1492 below - Retail Wastage claimable	pharmacy			
Tab 125 mg dispersible	276.00	28	1	Exjade
Tab 250 mg dispersible		28	_	Exjade
Tab 500 mg dispersible		28		Exjade

#### ⇒SA1492 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; and
- 2 Deferasirox is to be given at a daily dose not exceeding 40 mg/kg/day; and
- 3 Any of the following:

Subsidy (Manufacturer's Price)	Subsi	Fully dised	Brand or Generic
 \$	Per	✓	Manufacturer

- 3.1 Treatment with maximum tolerated doses of deferiprone monotherapy or deferiprone and desferrioxamine combination therapy have proven ineffective as measured by serum ferritin levels, liver or cardiac MRI T2\*; or
- 3.2 Treatment with deferiprone has resulted in severe persistent vomiting or diarrhoea; or
- 3.3 Treatment with deferiprone has resulted in arthritis; or
- 3.4 Treatment with deferiprone is contraindicated due to a history of agranulocytosis (defined as an absolute neutrophil count (ANC) of < 0.5 cells per μL) or recurrent episodes (greater than 2 episodes) of moderate neutropenia (ANC 0.5 1.0 cells per μL).</p>

**Renewal** only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the first renewal following 2 years of therapy, the treatment has been tolerated and has resulted in clinical improvement in all three parameters namely serum ferritin, cardiac MRI T2\* and liver MRI T2\* levels; or
- 2 For subsequent renewals, the treatment has been tolerated and has resulted in clinical stability or continued improvement in all three parameters namely serum ferritin, cardiac MRI T2\* and liver MRI T2\* levels.

DEFERIPRONE - Special Authority see SA1480 below - Retail	il pharmacy		
Tab 500 mg	533.17	100	✓ Ferriprox
Oral liq 100 mg per 1 ml	266.59	250 ml OP	✓ Ferriprox

## ⇒SA1480 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Either:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; or
- 2 The patient has been diagnosed with chronic iron overload due to acquired red cell aplasia.

# Inj 500 mg vial	151.31	10	✓ DBL  Desferrioxamine  Mesylate for Inj
			BP  ✓ Deferoxamine Pfizer S29 S29
SODIUM CALCIUM EDETATE			
* Inj 200 mg per ml, 5 ml	53.31	6	
	(156.71)		Calcium Disodium Versenate



# **Standard Formulae**

Otaliaala i Olillalac			
ACETYLCYSTEINE EYE DROPS Acetylcysteine inj 200 mg per ml, 10 ml	qs	PHENOBARBITONE SODIUM PAEDIATRIC ORAL mg per ml)	LIQUID (10
Suitable eye drop base	qs qs	Phenobarbitone Sodium	400 mg
, .	•	Glycerol BP	4 ml
CODEINE LINCTUS (3 mg per 5 ml)		Water	to 40 ml
Codeine phosphate	60 mg		
Glycerol	40 ml	PILOCARPINE ORAL LIQUID	
Preservative	qs	Pilocarpine 4% eye drops	qs
Water	to 100 ml	Preservative	qs
CODEINE LINCTUS (15 mg per 5 ml)		Water	to 500 ml
Codeine phosphate	300 mg	(Preservative should be used if quantity supplied is	for more
Glycerol	40 ml	than 5 days.)	
Preservative	qs	SALIVA SUBSTITUTE FORMULA	
Water	to 100 ml	Methylcellulose	5 g
water	10 100 1111	Preservative	qs
FOLINIC MOUTHWASH		Water	to 500 ml
Calcium folinate 15 mg tab	1 tab	(Preservative should be used if quantity supplied is	
Preservative	qs	than 5 days. Maximum 500 ml per prescription.)	
Water	to 500 ml		
(Preservative should be used if quantity supplied is	for more	SODIUM CHLORIDE ORAL LIQUID	
than 5 days. Maximum 500 ml per prescription.)		Sodium chloride inj 23.4%, 20 ml	qs
METHYL HYDROXYBENZOATE 10% SOLUTION		Water	ds
Methyl hydroxybenzoate	10 g	(Only funded if prescribed for treatment of hyponatr	aemia)
Propylene glycol	to 100 ml	VANCOMYCIN ORAL SOLUTION (25 mg per ml)	
(Use 1 ml of the 10% solution per 100 ml of oral liqu		Vancomycin 500 mg injection	5 vials
(OSC 1 IIII OF THE TO /O SOLUTION PER TOO THE OF ORAL HIGH	ia illixiale)	Glycerin with sucrose suspension	37.5 ml
OMEPRAZOLE SUSPENSION		Water	to 100 ml
Omeprazole capsules or powder	qs	(Only funded if prescribed for treatment of Clostridia	ım difficile
Sodium bicarbonate powder BP	8.4 g	following metronidazole failure)	
Water	to 100 ml		
PHENOBARBITONE ORAL LIQUID			
Phenobarbitone Sodium	1 g		
Glycerol BP	70 ml		
G1,00101 D1	, 5 1111		

to 100 ml

Water

# EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic

Per Manufacturer

# **Extemporaneously Compounded Preparations and Galenicals**

CODEINE PHOSPHATE – Safety medicine; prescriber may determin	e dispensina f	requency	
Powder – Only in combination		25 g	
	(90.09)		Douglas
Only in extemporaneously compounded codeine linctus.			
COLLODION FLEXIBLE			
Note: This product is no longer being manufactured by the suppli determined.	er and will be	delisted from 1	ine Schedule at a date to be
Collodion flexible	19.30	100 ml	✓ PSM
COMPOUND HYDROXYBENZOATE – Only in combination	10.00	100 1111	
Only in extemporaneously compounded oral mixtures.			
Soln	30.00	100 ml	✓ Midwest
GLYCERIN WITH SODIUM SACCHARIN - Only in combination			
Only in combination with Ora-Plus or when used in the vancomyc	in oral Iquuid S	Standard Form	nulae.
Suspension	30.95	473 ml	✓ Ora-Sweet SF
GLYCERIN WITH SUCROSE - Only in combination			
Only in combination with Ora-Plus or when used in the vancomyc	in oral Iquuid S		
Suspension	30.95	473 ml	✓ Ora-Sweet
GLYCEROL			•
* Liquid – Only in combination		500 ml	✓ healthE Glycerol BP
Only in extemporaneously compounded oral liquid preparatio	ns.		
METHYL HYDROXYBENZOATE	0.00	05	/ Misharat
Powder	8.98	25 g	✓ Midwest
METHYLCELLULOSE Powder	26.05	100 ~	✓ MidWoot
Suspension – Only in combination		100 g 473 ml	<ul><li>✓ MidWest</li><li>✓ Ora-Plus</li></ul>
METHYLCELLULOSE WITH GLYCERIN AND SODIUM SACCHARIN			· Old-I lub
Suspension		473 ml	✓ Ora-Blend SF
METHYLCELLULOSE WITH GLYCERIN AND SUCROSE - Only in a		4701111	ora bicina or
Suspension		473 ml	✓ Ora-Blend
PHENOBARBITONE SODIUM			0.0 2.0
Powder – Only in combination	52.50	10 g	✓ MidWest
	325.00	100 g	✓ MidWest
Only in children up to 12 years		-	
PROPYLENE GLYCOL			
Only in extemporaneously compounded methyl hydroxybenzoate			_
Liq	11.25	500 ml	✓ Midwest
SODIUM BICARBONATE			•
Powder BP – Only in combination		500 g	✓ Midwest
Only in extemporaneously compounded omeprazole and lans	soprazoie susp	ension.	
SYRUP (PHARMACEUTICAL GRADE) – Only in combination			
Only in extemporaneously compounded oral liquid preparations.  Liq	14 95	500 ml	✓ Midwest
•	لق.⊷ا	JUU IIII	- WIIWWEST
WATER Tap – Only in combination	0.00	1 ml	✓ Tap water
rap Only in combination	0.00	1 1111	- Iup water

Subsidy (Manufacturer's Price) Fully Subsidised Per

Brand or Generic Manufacturer

# **Nutrient Modules**

## Carbohydrate

#### ⇒SA1930 Special Authority for Subsidy

Initial application — (Cystic fibrosis or kidney disease) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Fither:

- 1 cystic fibrosis; or
- 2 chronic kidney disease.

**Initial application — (Indications other than cystic fibrosis or renal failure)** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 cancer in children: or
- 2 cancers affecting alimentary tract where there are malabsorption problems in patients over the age of 20 years; or
- 3 faltering growth in an infant/child; or
- 4 bronchopulmonary dysplasia; or
- 5 premature and post premature infant; or
- 6 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified where the patient has inborn errors of metabolism. Renewal — (Cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

CARBOHYDRATE SUPPLEMENT – Special Authority see SA1930 above – Hospital pharmacy [HP3]

# Carbohydrate And Fat

## **⇒SA1376** Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subs	idised	Generic	
\$	Per	✓	Manufacturer	

- 1 Infant or child aged four years or under; and
- 2 cystic fibrosis.

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 infant or child aged four years or under; and
- 2 Any of the following:
  - 2.1 cancer in children; or
  - 2.2 faltering growth; or
  - 2.3 bronchopulmonary dysplasia; or
  - 2.4 premature and post premature infants.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

#### Fat

## ⇒SA2204 Special Authority for Subsidy

Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified where the patient has an inborn error of metabolism.

Initial application — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 faltering growth in an infant/child; or
- 2 bronchopulmonary dysplasia; or
- 3 fat malabsorption; or
- 4 lymphangiectasia; or
- 5 short bowel syndrome: or
- 6 infants with necrotising enterocolitis; or
- 7 biliary atresia; or
- 8 for use in a ketogenic diet; or
- 9 chyle leak; or

continued...

✓ fully subsidised 277

Subsidy		Fully	Brand or
(Manufacturer's Price)	_ 8	Subsidised	Generic
\$	Per		Manufacturer

- 10 ascites: or
- 11 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. Renewal — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT SUPPLEMENT	- Special Authority	y see SA2204 o	on the previous	page - Hos	pital pharmacy [HP3]	Ĺ

Emulsion (neutral)		
38.44	500 ml OP	✓ Calogen
Emulsion (strawberry)15.38	200 ml OP	✓ Calogen
Oil		✓ MCT oil (Nutricia)
MCT Emulsion, 250 ml143.65	4 OP	<ul><li>Liquigen</li></ul>

#### **Protein**

## ⇒SA1524 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 protein losing enteropathy; or
- 2 high protein needs; or
- 3 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

	armacy [HP3]	ROTEIN SUPPLEMENT - Special Authority see SA1524 above - Hospital pha	PRO
✓ Resource	227 g OP	Powder8.95	
Beneprotein			
✓ Protifar	225 a OP	13 82	

Subsidy (Manufacturer's Price) \$

Subsidised Per 🗸

Fully

Brand or Generic Manufacturer

## **Oral and Enteral Feeds**

#### **Diabetic Products**

#### ⇒SA1095 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is a type I or and II diabetic who is suffering weight loss and malnutrition that requires nutritional support. Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

DIABETIC ENTERAL FEED 1KCAL/ML - Special Authority see	SA1095 above -	Hospital pharm	acy [HP3]
Liquid	4.65	500 ml OP	Glucerna Select
DIABETIC ORAL FEED 1KCAL/ML - Special Authority see SA	1095 above – Hos	pital pharmacy	[HP3]
Liquid (strawberry)	2.25	200 ml OP	✓ Diasip
Liquid (vanilla)	2.10	200 ml OP	✓ Nutren Diabetes
, , ,	2.25		Diasip

## **Fat Modified Products**

## ⇒SA2205 Special Authority for Subsidy

Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified where the patient has an inborn error of metabolism. Initial application — (Indications other than errors of inborn metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Patient has a chyle leak; or
- 2 Modified as a modular feed, made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule, for adults.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

#### Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

✓ fully subsidised 279

Subsidy (Manufacturer's Price)

Fully Subsidised Brand or Generic Manufacturer

## Paediatric Products For Children Awaiting Liver Transplant

#### ⇒SA1098 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) who requires a liver transplant.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL FEED 1KCAL/ML - Special Authority see SA1098 above - Hospital pharmacy (HP3)

400 g OP ✓ Heparon Junior 

## Paediatric Products For Children With Chronic Renal Failure

# **⇒SA1099** Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 vears where the patient is a child (up to 18 years) with acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL FEED 1KCAL/ML - Special Authority see SA1099 above - Hospital pharmacy [HP3] Powder .......64.26 400 a OP ✓ Kindergen

#### **Paediatric Products**

## ⇒SA1379 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Child is aged one to ten years; and
- 2 Any of the following:
  - 2.1 the child is being fed via a tube or a tube is to be inserted for the purposes of feeding; or
  - 2.2 any condition causing malabsorption; or
  - 2.3 faltering growth in an infant/child; or
  - 2.4 increased nutritional requirements; or
  - 2.5 the child is being transitioned from TPN or tube feeding to oral feeding.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for

			SI	PECIAL FOODS
	Subsidy (Manufacturer's Price) \$	Subs Per	Fully sidised	Brand or Generic Manufacturer
continued applications meeting the following criteria: Both:  1 The treatment remains appropriate and the patient is 2 General Practitioners must include the name of the dipractitioner and date contacted.			nally re	egistered general
PAEDIATRIC ENTERAL FEED 1.5KCAL/ML - Special Auth Liquid	,	previous p 00 ml OP	<b>V</b> F	Hospital pharmacy [HP3] Frebini Energy Jutrini Energy RTH
PAEDIATRIC ENTERAL FEED 1KCAL/ML - Special Author Liquid	•	evious pag 00 ml OP	je – Ho ✓ F ✓ N	= -
PAEDIATRIC ENTERAL FEED WITH FIBRE 1.5KCAL/ML - pharmacy [HP3]	•			
Liquid	7.00 50	00 ml OP	<b>✓</b> F	rebini Energy Fibre

PAEDIATRIC ENTERAL FEED WITH FIBRE 1KCAL/ML	- Special Authority see SA1379 on the previous page - Hospital
nharmacy [HP3]	

7.14

Liquid	500 ml OP	Frebini Original
		Fibre

FALDIATRIC ORAL FEED T.SRCAL/IVIL	- Special Authority see SA 1379 of the p	nevious page -	- Hospilai phannacy	[LIE o
Liquid (strawberry)	1.90	200 ml OP	✓ Fortini	

Liquid (vanilla)	1.90	200 ml OP	✓ Fortini
	8.67	500 ml OP	Pediasure Plus

PAEDIATRIC ORAL F	EED INCAL/IVIL - 5	pecial Authority see	SA 1379 on the pro	evious page –	nospital pharmacy	יחחן
Liquid (chocolate)			1.33	200 ml OP	✓ Pediasure	
Liquid (strawberry	)		1.33	200 ml OP	Pediasure	

200 ml OP ✓ Pediasure 250 ml OP ✓ Pediasure

PAEDIATRIC ORAL FEED WITH FIBRE 1.5KCAL/ML - Special Authority see SA1379 on the previous page - Hospital pharmacy [HP3]

Liquid (unflavoured)	1.90 200 ml OP	✓ Fortini Multi Fibre
Liquid (chocolate)		✓ Fortini Multi Fibre
Liquid (strawberry)		✓ Fortini Multi Fibre
Liquid (vanilla)		✓ Fortini Multi Fibre

PEPTIDE-BASED ORAL FEED - Special Authority see SA1379 on the previous page - Hospital pharmacy [HP3] ✓ Peptamen Junior

Powder .......43.60 400 g OP

## **Renal Products**

#### ⇒SA1101 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the

continued...

✓ Nutrini Energy Multi Fibre

✓ fully subsidised 281

Subsidy	)	Fully	Brand or
(Manufacturer's Price		Subsidised	Generic
	Per	✓	Manufacturer

recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

#### Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

RENAL ORAL FEED 1.8 KCAL/ML - Special Authority see SA1101 on the previous page - Hospital pharmacy [HP3] ✓ Nepro HP 220 ml OP (strawberry) ✓ Nepro HP (vanilla) RENAL ORAL FEED 2 KCAL/ML - Special Authority see SA1101 on the previous page - Hospital pharmacy [HP3] 4 OP ✓ NovaSource Renal 4 OP ✓ Renilon 7.5 4 OP ✓ Renilon 7.5 

## **Specialised And Elemental Products**

#### **⇒SA1377** Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 malabsorption: or
- 2 short bowel syndrome; or
- 3 enterocutaneous fistulas: or
- 4 eosinophilic oesophagitis; or
- 5 inflammatory bowel disease: or
- 6 patients with multiple food allergies requiring enteral feeding.

Notes: Each of these products is highly specialised and would be prescribed only by an expert for a specific disorder. The alternative is hospitalisation.

Elemental 028 Extra is more expensive than other products listed in this section and should only be used where the alternatives have been tried first and/or are unsuitable.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

#### Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL SEMI-ELEMENTAL FEED 1.5KCAL/ML - Special Authority see \$A1377 above - Hospital pharmacy [HP3] 1.000 ml OP ORAL ELEMENTAL FEED 0.8KCAL/ML - Special Authority see SA1377 above - Hospital pharmacy [HP3] 18 OP ✓ Elemental 028 Extra Liquid (pineapple & orange), 250 ml carton......179.46 18 OP ✓ Elemental 028 Extra 18 OP ✓ Elemental 028 Extra ORAL ELEMENTAL FEED 1KCAL/ML - Special Authority see SA1377 above - Hospital pharmacy [HP3] ✓ Vivonex TEN 80 a OP

	Subsidy (Manufacturer's Pr \$	rice) Subs Per	Fully sidised	Brand or Generic Manufacturer
SEMI-ELEMENTAL ENTERAL FEED 1KCAL/ML - Special Autl	,			
Liquid	9.60	500 ml OP	-	lutrison Advanced Peptisorb Survimed OPD

## Paediatric Products For Children With Low Energy Requirements

#### ⇒SA1196 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 Child aged one to eight years; and
- 2 The child has a low energy requirement but normal protein and micronutrient requirements.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

- Both:
  - 1 The treatment remains appropriate and the patient is benefiting from treatment; and
  - 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

PAEDIATRIC ENTERAL FEED WITH FIBRE 0.76 KCAL/ML - Special Authority see SA1196 above - Hospital pharmacy [HP3] 500 ml OP ✓ Nutrini Low Energy Multi Fibre

# Standard Supplements

#### ⇒SA1859 Special Authority for Subsidy

Initial application — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- - 1 The patient is under 18 years of age; and
  - 2 Any of the following:
    - 2.1 The patient has a condition causing malabsorption; or
    - 2.2 The patient has failure to thrive; or
    - 2.3 The patient has increased nutritional requirements; and
  - 3 Nutrition goal has been set (eg reach a specific weight or BMI).

Renewal — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 A nutrition goal has been set (eg reach a specific weight or BMI).

Initial application — (Children - exclusive enteral nutrition for Crohn's disease) only from a gastroenterologist or dietitian on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

continued...

✓ fully subsidised 283

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
 \$	Per 🗸	Manufacturer

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 Dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Renewal — (Children - exclusive enteral nutrition for Crohn's disease) from any relevant practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 General Practitioners and dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Initial application — (Adults) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

1 Any of the following:

Patient is Malnourished

- 1.1 Patient has a body mass index (BMI) of less than 18.5 kg/m<sup>2</sup>; or
- 1.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 1.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months; and
- 2 Any of the following:

Patient has not responded to first-line dietary measures over a 4 week period by:

- 2.1 Increasing their food intake frequency (eg snacks between meals); or
- 2.2 Using high-energy foods (e.g. milkshakes, full fat milk, butter, cream, cheese, sugar etc); or
- 2.3 Using over the counter supplements (e.g. Complan); and
- 3 A nutrition goal has been set (e.g. to reach a specific weight or BMI).

Renewal — (Adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 A nutrition goal has been set (eg reach a specific weight or BMI); and
- 2 Any of the following:

Patient is Malnourished

- 2.1 Patient has a body mass index (BMI) of less than 18.5 kg/m<sup>2</sup>; or
- 2.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 2.3 Patient has a BMI of less than 20 kg/m<sup>2</sup> and unintentional weight loss greater than 5% within the last 3-6 months.

**Initial application — (Short-term medical condition)** from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube or a nasogastric tube is to be inserted for feeding; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Is undergoing a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
  - 5.1 Pregnant; and
  - 5.2 Any of the following:
    - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or

Subsidy	Fı	ılly	Brand or
(Manufacturer's Price)	Subsidis	ed	Generic
\$	Per	1	Manufacturer

- 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
- 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

Renewal — (Short-term medical condition) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Has undergone a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
  - 5.1 Pregnant: and
  - 5.2 Any of the following:
    - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis
    - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
    - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

Initial application — (Long-term medical condition) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis: or
- 3 Liver disease: or
- 4 Chronic Renal failure: or
- 5 Inflammatory bowel disease; or
- 6 Chronic obstructive pulmonary disease with hypercapnia: or
- 7 Short bowel syndrome; or
- 8 Bowel fistula: or
- 9 Severe chronic neurological conditions; or
- 10 Epidermolysis bullosa: or
- 11 AIDS (CD4 count < 200 cells/mm3): or
- 12 Chronic pancreatitis.

Renewal — (Chronic disease OR tube feeding for patients who have previously been funded under Special Authority forms SA0702 or SA0583) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis: or
- 3 Liver disease: or

continued...

285 ✓ fully subsidised

	Subsidy		Fully	Brand or
(	Manufacturer's Price)	Sub	sidised	Generic
	\$	Per	1	Manufacturer

- 4 Chronic Renal failure; or
- 5 Inflammatory bowel disease; or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome; or
- 8 Bowel fistula; or

9 Severe chronic neurological conditions.			
ENTERAL FEED 1.5KCAL/ML - Special Authority see SA1859 on p Liquid		Hospital pharmac 250 ml OP 1,000 ml OP	y [HP3]  ✓ Ensure Plus HN ✓ Ensure Plus HN RTH
	9.00 9.60		<ul><li>✓ Nutrison Energy</li><li>✓ Fresubin HP Energy</li></ul>
ENTERAL FEED 1KCAL/ML – Special Authority see SA1859 on pa	•	spital pharmacy [ 250 ml OP 1,000 ml OP	HP3]  Isosource Standard Fresubin Original Osmolite RTH Nutrison RTH
ENTERAL FEED WITH FIBRE 0.83 KCAL/ML – Special Authority s Liquid		on page 283 – Ho 1,000 ml OP	ospital pharmacy [HP3]  Nutrison  800 Complete  Multi Fibre
ENTERAL FEED WITH FIBRE 1 KCAL/ML - Special Authority see Liquid		page 283 – Hosp 1,000 ml OP	ital pharmacy [HP3]  Jevity RTH Fresubin Original Fibre
	7.21		✓ Nutrison Multi Fibre
ENTERAL FEED WITH FIBRE 1.2KCAL/ML - Special Authority see Liquid		page 283 – Hos 1,000 ml OP	pital pharmacy [HP3]  ✓ Jevity Plus RTH
ENTERAL FEED WITH FIBRE 1.5KCAL/ML - Special Authority see Liquid		1,000 ml OP	pital pharmacy [HP3]  ✓ Jevity HiCal RTH  ✓ Nutrison Energy  Multi Fibre
	9.80		✓ Fresubin HP Energy Fibre
ENTERAL FEED WITH PROTEIN 1.2KCAL/ML - Special Authority Liquid		on page 283 – F 500 ml OP	Hospital pharmacy [HP3]  ✓ Fresubin Intensive
ORAL FEED (POWDER) - Special Authority see SA1859 on page 2	283 – Hospit	al pharmacy [HP	3]
Powder (chocolate)		840 g OP	✓ Sustagen Hospital Formula
	26.00	850 g OP	✓ Ensure
Powder (vanilla)	14.00	840 g OP	<ul> <li>Sustagen Hospital Formula Active</li> </ul>
	26.00	850 g OP	✓ Ensure

	Subsidy	Fully	Brand or
(1)	Manufacturer's Price)	Subsidised	Generic
	\$	Per 🗸	Manufacturer

#### ORAL FEED 1.5KCAL/ML - Special Authority see SA1859 on page 283 - Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, who have severe epidermolysis bullosa, or as exclusive enteral nutrition in children under the age of 18 years for the treatment of Crohn's disease, or for patients with COPD and hypercapnia, defined as CO2 value exceeding 55mmHg. The prescription must be endorsed accordingly.

)P
Ensure Plus Fortisip
)P
Ensure Plus Fortisip
)P
Ensure Plus
)P
Fortisip
_
P Ensure Plus
P Erisure Plus
Ensure Plus Fortisip

ORAL FEED WITH FIBRE 1.5 KCAL/ML — Special Authority see SA1859 on page 283 — Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, or who have severe epidermolysis bullosa. The prescription must be endorsed accordingly.

Liquid (chocolate) – Higher subsidy of \$1.76 per 200 ml with	0.70	000 100	
Endorsement	0./2	200 ml OP	
	(1.76)		Fortisip Multi Fibre
Liquid (strawberry) - Higher subsidy of \$1.76 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.76)		Fortisip Multi Fibre
Liquid (vanilla) - Higher subsidy of \$1.76 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.76)		Fortisip Multi Fibre

# **High Calorie Products**

## ⇒SA1195 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

All of the following:

- 1 Cystic fibrosis; and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements.

continued...

✓ fully subsidised 287

# **SPECIAL FOODS**

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	1	Manufacturer

continued...

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 any condition causing malabsorption; or
  - 1.2 faltering growth in an infant/child; or
  - 1.3 increased nutritional requirements; or
  - 1.4 fluid restricted: and

practitioner and date contacted.

- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements or is fluid restricted.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL FEED 2 KCAL/ML - Special Authority see SA1195 on the previous	oage – Hospital p	harmacy [HP3]
Liquid	500 ml OP	✓ Fresubin 2kcal HP
6.82		✓ Nutrison Concentrated
13.64	1,000 ml OP	✓ Ensure Two Cal HN

ORAL FEED 2 KCAL/ML - Special Authority see SA1195 on the previous page - Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, or who have severe epidermolysis bullosa. The prescription must be endorsed accordingly.

Liquid (vanilla) - Higher subsidy of \$2.34 per 200 ml with

(2.34) Two Cal HN

## **Food Thickeners**

# ⇒SA1106 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient has motor neurone disease with swallowing disorder.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

Subsidy		Fully	Brand or
(Manufacturer's Price)	5	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

#### **Gluten Free Foods**

The funding of gluten free foods is no longer being actively managed by Pharmac from 1 April 2011. This means that we are no longer considering the listing of new products, or making subsidy, or other changes to the existing listings. As a result we anticipate that the range of funded items will reduce over time. Management of Coeliac disease with a gluten free diet is necessary for good outcomes. A range of gluten free options are available through retail outlets.

#### ⇒SA1729 Special Authority for Subsidy

Initial application — (all patients) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Either:

- 1 Gluten enteropathy has been diagnosed by biopsy; or
- 2 Patient suffers from dermatitis herpetiformis.

Initial application — (paediatric patients diagnosed by ESPGHAN criteria) only from a paediatric gastroenterologist. Approvals valid without further renewal unless notified where the paediatric patient fulfils ESPGHAN criteria for biopsy free diagnosis of coeliac disease.

GLUTEN FREE BAKING MIX – Special Authority see SA1729 Powder		pharmacy [HP3] 1,000 g OP	
	(5.15)		Healtheries Simple Baking Mix
GLUTEN FREE BREAD MIX - Special Authority see SA1729	above – Hospital r	oharmacy [HP3]	
Powder		1,000 g OP	
	(7.32)		NZB Low Gluten Bread Mix
	3.51		
	(10.87)		Horleys Bread Mix
GLUTEN FREE FLOUR - Special Authority see SA1729 abov	e – Hospital pharr	nacy [HP3]	
Powder		2,000 g OP	
	(18.10)	. 3	Horleys Flour

✓ fully subsidised 289

	Subsidy		Fully Brand or	
	(Manufacturer's F			
	\$	Per	✓ Manufacturer	
GLUTEN FREE PASTA – Special Authority see SA1729 on the	previous page -	Hospital pharma	cy [HP3]	
Buckwheat Spirals	2.00	250 g OP		
	(3.11)	-	Orgran	
Corn and Vegetable Shells	2.00	250 g OP		
	(2.92)		Orgran	
Corn and Vegetable Spirals	2.00	250 g OP		
	(2.92)		Orgran	
Rice and Corn Lasagne Sheets	1.60	200 g OP		
	(3.82)		Orgran	
Rice and Corn Macaroni	2.00	250 g OP		
	(2.92)		Orgran	
Rice and Corn Penne		250 g OP		
	(2.92)		Orgran	
Rice and Maize Pasta Spirals		250 g OP		
	(2.92)		Orgran	
Rice and Millet Spirals		250 g OP		
	(3.11)		Orgran	
Rice and corn spaghetti noodles		375 g OP	_	
	(2.92)		Orgran	
Vegetable and Rice Spirals		250 g OP	_	
	(2.92)		Orgran	
Italian long style spaghetti		220 g OP		
	(3.11)		Orgran	
(Orgran Buckwheat Spirals to be delisted 1 July 2025)				
(Orgran Corn and Vegetable Shells to be delisted 1 July 2025)				
(Orgran Corn and Vegetable Spirals to be delisted 1 July 2025)				
(Orgran Rice and Corn Lasagne Sheets to be delisted 1 July 202	?5)			
(Orgran Rice and Corn Macaroni to be delisted 1 July 2025)				
(Orgran Rice and Corn Penne to be delisted 1 July 2025)				
(Orgran Rice and Maize Pasta Spirals to be delisted 1 July 2025,	)			
(Orgran Rice and Millet Spirals to be delisted 1 July 2025)				
(Orgran Rice and corn spaghetti noodles to be delisted 1 July 20	25)			

# **Foods And Supplements For Inherited Metabolic Disease**

(Orgran Vegetable and Rice Spirals to be delisted 1 July 2025) (Orgran Italian long style spaghetti to be delisted 1 July 2025)

#### ⇒SA2357 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid without further renewal unless notified where patient requires dietary management of inherited metabolic disorders.

# **Supplements For Homocystinuria**

AMINOACID FORMULA WITHOUT METHIONINE	- Special Authority see SA2357	above -	Hospital pharmacy [HP3]
Powder (neutral), 36 g sachets	750.30	30	✓ HCU Anamix Junior
Powder, 12.5 g sachets	349.65	30	✓ HCU Explore 5
Powder, 25 g sachets	1,048.95	30	✓ HCU Express 15
Powder (neutral), can	480.42	500 g Ol	P <b>XMET Maxamum</b>
Powder (unflavoured), can		400 g Ol	P <b>V HCU Anamix Infant</b>
Liquid (juicy berries), 125 ml bottle		30	✓ HCU Lophlex LQ
Liquid (orange), 125 ml bottle		36	✓ HCU Anamix Junior
			LQ

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	/	Manufacturer

# Supplements For MSUD and short chain enoyl coA hydratase deficiency

AMINOACID FORMULA WITHOUT VALINE, LEUCINE AND ISOLEUCINE - Special Authority see \$A2357 on the previous page - Hospital pharmacy [HP3]

Powder (neutral) 36 g sachets7	50.00 30	<ul><li>MSUD Anamix Junior</li></ul>
Powder, 12.5 g sachets3	49.65 30	✓ MSUD Explore 5
Powder, 25 g sachets		✓ MSUD Express 15
Powder (neutral), can4		✓ MSUD Maxamum
Powder (orange), can4		✓ MSUD Maxamum
Powder (unflavoured), can2		MSUD Anamix Infant
Liquid (orange) 125 ml bottles9	41.40 36	<ul><li>MSUD Anamix Junior LQ</li></ul>
Liquid (juicy berries) 125 ml pouches	84.80 30	✓ MSUD Lophlex LQ 20

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

# **Supplements For PKU**

••			
AMINOACID FORMULA WITHOUT PHENYLALANINE	- Special Authority see S		290 – Hospital pharmacy [HP3]
Tabs	99.00	75 OP	✓ Phlexy 10
Powder (Lemon), 34 g sachets	883.50	30	✓ PKU Express 20
Powder (Neutral), 12.5 g sachets	220.88	30	✓ PKU Explore 5
Powder (Neutral), 34 g sachets	883.50	30	✓ PKU Express 20
Powder (Orange), 25 g sachets	441.75	30	✓ PKU Explore 10
Powder (Orange), 34 g sachets	883.50	30	✓ PKU Express 20
Powder (Raspberry), 25 g sachets	441.75	30	✓ PKU Explore 10
Powder (Tropical), 34 g sachets		30	✓ PKU Express 20
Powder (berry) 28 g sachets	936.00	30	✓ PKU Lophlex Powder
Powder (chocolate) 36 g sachet	393.00	30	✓ PKU Anamix Junior Chocolate
Powder (neutral) 28 g sachets	936.00	30	✓ PKU Lophlex Powder
Powder (neutral) 36 g sachets	393.00	30	✓ PKU Anamix Junior
Powder (orange) 28 g sachets	936.00	30	✓ PKU Lophlex
, , ,			Powder
Powder (orange) 36 g sachet	393.00	30	✓ PKU Anamix Junior Orange
Powder (unflavoured) 12.5 g sachets	234.00	30	✓ PKU First Spoon
Powder (vanilla) 36 g sachet	393.00	30	✓ PKU Anamix Junior Vanilla
Infant formula	174.72	400 g OP	✓ PKU Anamix Infant
Powder (orange)	320.00	500 g OP	✓ XP Maxamum
Powder (unflavoured)		500 g OP	✓ XP Maxamum
Liquid (berry)		125 ml OP	✓ PKU Anamix Junior LQ
Liquid (orange)	13.10	125 ml OP	✓ PKU Anamix Junior LQ
Liquid (forest berries), 250 ml carton	540.00	18 OP	✓ Easiphen Liquid
Liquid (juicy tropical) 125 ml	936.00	30 OP	✓ PKU Lophlex LQ 20
Oral semi-solid (berries) 109 g		36 OP	✓ PKU Lophlex Sensation 20
Powder (neutral), 400 g can	715.16	4 OP	✓ PKU Start
Liquid (juicy berries) 62.5 ml		60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy berries) 125 ml		30 OP	✓ PKU Lophlex LQ 20
Liquid (juicy orange) 125 ml		30 OP	✓ PKU Lophlex LQ 20
1 0 -7 0-7 -			

	Subsidy (Manufacturer's Price) \$	S Per	Fully Subsidised	Brand or Generic Manufacturer
OLYGONA OR OR FRIDE AND ANNUA AND CONTAINS CONT	*			
GLYCOMACROPEPTIDE AND AMINO ACID CONTAINS SOME	PHENYLALANINE	– Spec	iai Autho	rity see SA2357 on
page 290 – Hospital pharmacy [HP3] Powder (Banana) 35 g sachets	030.00	30	1	PKU
Fowder (Bariana) 55 g sacriets	930.00	30	•	sphere20 Banana
Powder (Berry), 20 g sachets	440.00	60	1	PKU Restore
Fowder (Derry), 20 g Sacriets	443.20	00	•	Powder
Powder (Chocolate) 32 g sachets	808 56	30	1	PKU Build
1 owder (Orlocolate) 32 g sacriets	030.30	30	•	20 Chocolate
Powder (Chocolate) 35 g sachets	930.00	30	1	PKU
1 owder (Griocolate) oo g cacricie		00	•	sphere20 Chocolate
				opilorozo ollocolato
Powder (Lemon) 35 g sachets	930.00	30	1	PKU
				sphere20 Lemon
Powder (Lemonade) 33.4 g sachets	936.00	30	1	PKU GMPro Ultra
				Lemonade
Powder (Neutral), 15 g sachets	449.28	30	1	PKU Build 10
Powder (Orange), 20 g sachets	449.28	60	1	PKU Restore
				Powder
Powder (Raspberry Lemonade) 31 g sachets	898.56	30	•	PKU Build
				20 Raspberry
			_	Lemonade
Powder (Smooth) 31 g sachets	898.56	30	•	PKU Build
				20 Smooth
Powder (Vanilla) 33 g sachets		30		PKU Build 20 Vanilla
Powder (neutral), 40 g sachets		30		Glytactin Bettermilk
Powder (unflavoured) 12.5 g sachets		30 30		PKU GMPro Mix-In PKU GMPro Ultra
Powder (vanilla) 33.4 g sachets	936.00	30	•	Vanilla
Powder (Red Berry) 35 g sachets	030 NO	30	1	PKU sphere20 Red
Fowder (Hed Berry) 35 g sacriets	930.00	30	•	Berry
Powder (Vanilla) 35 g sachets	030 NO	30	1	PKU
1 Owder (Variilla) 65 g Sacricis		00	•	sphere20 Vanilla
Liquid (neutral), 250 ml carton	280.80	18	1	PKU GMPro LQ
Liquid (original), 250 ml carton		30 OP		PKU Glytactin RTD
Liquid (original), 200 m out or		00 01	-	15
Liquid (Coffee Mocha), 250 ml carton	684.45	30 OP	1	PKU Glytactin RTD
				15 Lite
Liquid (chocolate), 250 ml carton	684.45	30 OP	1	PKU Glytactin RTD
1 (*	· ·			15
Liquid (vanilla), 250 ml carton	684.45	30 OP	1	PKU Glytactin RTD
, , , , , , , , , , , , , , , , , , , ,				15 Lite

# Foods

LOW PROTEIN BAKING MIX - Special Authority see SA2357 on p	page 290 -	Hospital pharmacy	y [HP3]
Powder	8.55	500 g OP	✓ Loprofin Mix

✓ fully subsidised 293

	Subsidy		Fully Brand or
	(Manufacturer's F	Price) Subs Per	idised Generic  Manufacturer
LOW DOCTON DAOTA Constitution of the CASSES CONTRACTOR OF THE CONT	. 200 Harrital		
LOW PROTEIN PASTA – Special Authority see SA2357 on pag	•		
Animal shapes		500 g OP	✓ Loprofin
Lasagne		250 g OP	✓ Loprofin
Low protein rice pasta		500 g OP	✓ Loprofin
Macaroni		250 g OP	✓ Loprofin
Penne		500 g OP	✓ Loprofin
Spaghetti		500 g OP	✓ Loprofin
Spirals	12.39	500 g OP	✓ Loprofin
Supplements for Tyrosinaemia			
AMINOACID FORMULA WITHOUT PHENYLALANINE AND TY pharmacy [HP3]	ROSINE - Spec	ial Authority see	e SA2357 on page 290 – Hospi
Powder (Neutral), 12.5 g sachets	349.65	30	✓ TYR Explore 5
Powder (neutral) 36 g sachets		30	✓ TYR Anamix Junior
Powder, can		400 g OP	✓ TYR Anamix Infant
Liquid (juicy berries) 125 ml pouches	1,684.80	30	✓ TYR Lophlex LQ 20
Liquid (orange) 125 ml bottle		36	TYR Anamix Junior
			LQ
GLYCOMACROPEPTIDE AND AMINO ACID CONTAINS SOME	TYROSINE AN	D PHENYLALA	ANINE - Special Authority see
SA2357 on page 290 – Hospital pharmacy [HP3]			
Powder (Red Berry), 35 g sachets	1,398.60	30	✓ TYR Sphere 20
Powder (Vanilla), 35 g sachets		30	✓ TYR Sphere 20
Supplements for Organic Acidaemias  AMINOACID FORMULA WITHOUT ISOLEUCINE, METHIONIN on page 290 – Hospital pharmacy [HP3]	E, THREONINE	AND VALINE -	- Special Authority see SA2357
Powder, can	260.00	400 g OP	<ul><li>MMA/PA Anamix Infant</li></ul>
AMINOACID FORMULA WITHOUT METHIONINE, THREONINE Hospital pharmacy [HP3]	E AND VALINE -	- Special Autho	rity see SA2357 on page 290 -
Powder (neutral), 18 g sachets	750.30	30	<ul><li>MMA/PA Anamix Junior</li></ul>
Powder (neutral), 18 g sachets		30 30	
	349.65		Junior
Powder, 12.5 g sachets	349.65	30	Junior ✓ MMA/PA Explore 5
Powder, 12.5 g sachets	349.65 1,048.95	30 30	Junior  ✓ MMA/PA Explore 5  ✓ MMA/PA Express 15
Powder, 12.5 g sachets  Powder, 25 g sachets  Supplements for Glutaric Aciduria type 1  AMINOACID FORMULA WITHOUT LYSINE – Special Authority	349.65 1,048.95 see SA2357 on	30 30 page 290 – Ho	Junior  MMA/PA Explore 5  MMA/PA Express 15  spital pharmacy [HP3]
Powder, 12.5 g sachets  Powder, 25 g sachets  Supplements for Glutaric Aciduria type 1  AMINOACID FORMULA WITHOUT LYSINE – Special Authority Powder (neutral), 18 g sachets	349.65 1,048.95 see SA2357 on 750.30	30 30 page 290 – Hos	Junior  MMA/PA Explore 5  MMA/PA Express 15  spital pharmacy [HP3]  GA1 Anamix Junior
Powder, 12.5 g sachets  Powder, 25 g sachets  Supplements for Glutaric Aciduria type 1  AMINOACID FORMULA WITHOUT LYSINE – Special Authority Powder (neutral), 18 g sachets	349.65 1,048.95 see SA2357 on 750.30 349.65	30 30 page 290 – Hos 30 30	Junior  MMA/PA Explore 5  MMA/PA Express 15  spital pharmacy [HP3]  GA1 Anamix Junior  GA Explore 5
Powder, 12.5 g sachets	349.65 1,048.95 see SA2357 on 750.30 349.65	30 30 page 290 – Hos	Junior  MMA/PA Explore 5  MMA/PA Express 15  spital pharmacy [HP3]  GA1 Anamix Junior
Powder, 12.5 g sachets	349.65 1,048.95 see SA2357 on 750.30 349.65 260.00	30 30 page 290 – Hos 30 30 400 g OP	Junior  MMA/PA Explore 5  MMA/PA Express 15  spital pharmacy [HP3]  GA1 Anamix Junior  GA Explore 5  GA1 Anamix Infant
Powder, 12.5 g sachets	349.65 1,048.95 see SA2357 on 750.30 349.65 260.00	30 30 page 290 – Hos 30 30 400 g OP	Junior  MMA/PA Explore 5  MMA/PA Express 15  spital pharmacy [HP3]  GA1 Anamix Junior  GA Explore 5  GA1 Anamix Infant
Powder, 12.5 g sachets	349.65 1,048.95 see SA2357 on 750.30 349.65 260.00	30 30 page 290 – Hos 30 30 400 g OP	Junior  MMA/PA Explore 5  MMA/PA Express 15  spital pharmacy [HP3]  GA1 Anamix Junior  GA Explore 5  GA1 Anamix Infant

Subsidy (Manufacturer's Price) \$	S Per	Fully Subsidised	
CITRULLINE – Special Authority see SA2357 on page 290 – Hospital pharmacy [HP3] Powder, 4 g sachets211.45	30	1	Citrulline1000
ISOLEUCINE - Special Authority see SA2357 on page 290 - Hospital pharmacy [HP3]   Powder, 4 g sachets141.05	] 30	•	Isoleucine50
LEUCINE - Special Authority see SA2357 on page 290 - Hospital pharmacy [HP3] Powder, 4 g sachets141.05	30	1	Leucine100
PHENYLALANINE – Special Authority see SA2357 on page 290 – Hospital pharmacy [ Powder, 4 g sachets141.05	[HP3] 30		Phenylalanine50
TYROSINE - Special Authority see SA2357 on page 290 - Hospital pharmacy [HP3] Powder, 4 g sachets211.45	30	1	Tyrosine1000
VALINE - Special Authority see SA2357 on page 290 - Hospital pharmacy [HP3] Powder, 4 g sachets141.05	30	1	Valine50
Other Fat Modified Products			
ELEMENTAL FEED WITH HIGH MEDIUM CHAIN TRIGLYCERIDES – Special Authori pharmacy [HP3]	ity see	e SA2357	on page 290 – Hospital
, ,, ,	10	•	Emsogen
Carbohydrate and Fat with added vitamins and minerals			
PROTEIN FREE SUPPLEMENT CONTAINING CARBOHYDRATE, FAT WITH ADDED Authority see SA2357 on page 290 – Hospital pharmacy [HP3]	VITA	AMINS AN	ND MINERALS - Special
	g OF	• 🗸	Energivit
Essential Amino Acids			
ESSENTIAL AMINOACID FORMULA - Special Authority see SA2357 on page 290 - H Powder (neutral), can313.73 200	Hospit ) g OF	•	acy [HP3] Essential Amino Acid Mix
Infant Formulae			

#### Infant Formulae

## For Williams Syndrome

# ⇒SA1110 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is an infant suffering from Williams Syndrome and associated hypercalcaemia.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

✓ fully subsidised 295

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

## **Gastrointestinal and Other Malabsorptive Problems**

AMINO ACID FORMULA - Special Authority see SA209	2 below – Hospital pharr	nacy [HP3]	
Powder	43.60	400 g OP	<ul><li>✓ Alfamino</li><li>✓ Alfamino Junior</li></ul>
Powder (unflavoured)	55.61	400 g OP	<ul><li>✓ Neocate Gold</li><li>✓ Neocate Junior</li><li>Unflavoured</li></ul>
			✓ Neocate SYNEO
	65.72		✓ Elecare ✓ Elecare LCP
Powder (vanilla)	55.61	400 g OP	✓ Neocate Junior Vanilla
	65.72		✓ Elecare

#### ⇒SA2092 Special Authority for Subsidy

**Initial application** — (Infants under 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 History of anaphylaxis to cow's milk protein formula or dairy products; or
- 2 Eosinophilic oesophagitis; or
- 3 Ultra-short gut; or
- 4 Severe Immune deficiency; or
- 5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
- 6 Both
  - 6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
  - 6.2 Either:
    - 6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or 6.2.2 Patient has IgE mediated allergy.

Initial application — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or
  - 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
  - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products; or
  - 2.2 Eosinophilic oesophagitis; or
  - 2.3 Ultra-short gut; or
  - 2.4 Severe Immune deficiency: or
  - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
  - 2.6 Both:
    - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
    - 2.6.2 Either:
      - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval

continued...

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

number: or

2.6.2.2 Patient has IgE mediated allergy.

Renewal — (Infants up to 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 Patient has IgE mediated allergy; and
  - 1.2 All of the following:
    - 1.2.1 Patient remains allergic to cow's milk; and
    - 1.2.2 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy or extensively hydrolysed infant formula has been undertaken; and
    - 1.2.3 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
    - 1.2.4 Amino acid formula is required for a nutritional deficit; and
    - 1.2.5 It has been more than three months from the previous approval; or

#### 2 Both:

- 2.1 Patient has non IgE mediated severe gastrointestinal intolerance (including eosinophilic oesophagitis, ultra-short gut and severe immune deficiency); and
- 2.2 All of the following:
  - 2.2.1 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy, or extensively hydrolysed infant formula has been undertaken; and
    - 2.2.2 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
    - 2.2.3 Amino acid formula is required for a nutritional deficit; and
    - 2.2.4 It has been more than three months from the previous approval.

Renewal — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or
  - 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
  - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products; or
  - 2.2 Eosinophilic oesophagitis; or
  - 2.3 Ultra-short gut; or
  - 2.4 Severe Immune deficiency; or
  - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
  - 2.6 Both:
    - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or alleroy or malabsorotion; and
    - 2.6.2 Either:
      - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or
      - 2.6.2.2 Patient has IgE mediated allergy.

Initial application — (for patients who have a current funding under Special Authority form SA1557) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 months for applications meeting the following criteria:

continued...

✓ fully subsidised 297

ubsidy cturer's Price) Subs	Fully	Brand or Generic
 \$ Per	•	Manufacturer

continued...

All of the following:

- 1 Patient has a valid Special Authority approval for extensively hydrolysed formula (SA1557); and
- 2 Extensively hydrolysed formula (Aptamil Gold+ Pepti Junior, AllerPro SYNEO 1 and 2) is unable to be supplied at this time; and
- 3 The approval only applies to funded dispensings of Neocate Gold and Neocate Syneo.

Note: This criteria is short term funding to cover an out-of-stock situation on some extensively hydrolysed formula powder funded under Special Authority form SA1557. There is no renewal criteria under this restriction.

ENTERAL LIQUID PEPTIDE FORMULA - Special Authority see SA1953 below - Hospital pharmacy [HP3]

Liquid 1 kcal/ml12.44	500 ml OP	✓ Nutrini Peptisorb
Liquid 1.5 kcal/ml18.66	500 ml OP	✓ Nutrini Peptisorb
		Energy

### ⇒SA1953 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has impaired gastrointestinal function and either cannot tolerate polymeric feeds, or polymeric feeds are unsuitable; and
- 2 Any of the following:
  - 2.1 Severe malabsorption; or
  - 2.2 Short bowel syndrome: or
  - 2.3 Intractable diarrhoea; or
  - 2.4 Biliary atresia; or
  - 2.5 Cholestatic liver diseases causing malabsorption; or
  - 2.6 Cystic fibrosis; or
  - 2.7 Proven fat malabsorption; or
  - 2.8 Severe intestinal motility disorders causing significant malabsorption; or
  - 2.9 Intestinal failure: or
  - 2.10 Both:
    - 2.10.1 The patient is currently receiving funded amino acid formula; and
    - 2.10.2 The patient is to be trialled on, or transitioned to, an enteral liquid peptide formula; and
- 3 Either:
  - 3.1 A semi-elemental or partially hydrolysed powdered feed has been reasonably trialled and considered unsuitable; or
  - 3.2 For step down from intravenous nutrition.

Note: A reasonable trial is defined as a 2-4 week trial.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

#### All of the following:

- 1 An assessment as to whether the patient can be transitioned to a cows milk protein or soy infant formula or extensively hydrolysed formula has been undertaken; and
- 2 The outcome of the assessment is that the patient continues to require an enteral liquid peptide formula; and
- 3 General practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

EXTENSIVELY HYDROLYSED FORMULA -	<ul> <li>Special Authority see SA1557 on</li> </ul>	the next page - I	Hospital pharmacy [HP3]
Powder	18.10	450 g OP	✓ Pepti-Junior
	36.20	900 g OP	✓ Allerpro Syneo 1
			✓ Allerpro Syneo 2

Subsidy (Manufacturer's Price) \$

Fully Subsidised Per

Brand or Generic Manufacturer

#### ⇒SA1557 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 Both:
  - 1.1 Cows milk formula is inappropriate due to severe intolerance or allergy to its protein content; and
  - 12 Fither
    - 1.2.1 Soy milk formula has been reasonably trialled without resolution of symptoms; or
    - 1.2.2 Soy milk formula is considered clinically inappropriate or contraindicated; or
- 2 Severe malabsorption; or
- 3 Short bowel syndrome; or
- 4 Intractable diarrhoea; or
- 5 Biliary atresia; or
- 6 Cholestatic liver diseases causing malsorption; or
- 7 Cystic fibrosis: or
- 8 Proven fat malabsorption; or
- 9 Severe intestinal motility disorders causing significant malabsorption; or
- 10 Intestinal failure; or
- 11 All of the following:
  - 11.1 For step down from Amino Acid Formula: and
  - 11.2 The infant is currently receiving funded amino acid formula; and
  - 11.3 The infant is to be trialled on, or transitioned to, an extensively hydrolysed formula; and
  - 11.4 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

Note: A reasonable trial is defined as a 2-4 week trial, or signs of an immediate IgE mediated allergic reaction.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 An assessment as to whether the infant can be transitioned to a cows milk protein or soy infant formula has been undertaken; and
- 2 The outcome of the assessment is that the infant continues to require an extensively hydrolysed infant formula; and
- 3 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

#### Fluid Restricted

PAEDIATRIC ORAL/ENTERAL FEED 1 KCAL/ML - Special Authority see SA1698 below - Hospital pharmacy [HP3] Liquid.......2.80 125 ml OP ✓ Infatrini

#### ⇒SA1698 Special Authority for Subsidy

**Initial application** only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is fluid restricted or volume intolerant and has been diagnosed with faltering growth; and
- 2 Patient is under the care of a paediatrician or dietitian who has recommended treatment with a high energy infant formula; and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

continued...

✓ fully subsidised 299



Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
<u> </u>	Per		Manufacturer

continued...

Note: "Volume intolerant" patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

Renewal only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient continues to be fluid restricted or volume intolerant and has faltering growth; and
- 2 Patient is under the care of a hospital paediatrician or dietitian who has recommended treatment with a high energy infant formula; and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: 'Volume intolerant' patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

# **Ketogenic Diet**

## ⇒SA1197 Special Authority for Subsidy

**Initial application** only from a metabolic physician or paediatric neurologist. Approvals valid for 3 months where the patient has intractable epilepsy, pyruvate dehydrogenase deficiency or glucose transported type-1 deficiency and other conditions requiring a ketogenic diet.

**Renewal** only from a metabolic physician or paediatric neurologist. Approvals valid for 2 years where the patient is on a ketogenic diet and the patient is benefiting from the diet.

HIGH FAT LOW CARBOHYDRATE FORMULA - Special Authority see SA1197 above - Retail pharmacy

Powder (unflavoured)36.92	300 g OP	✓ KetoCal 4:1
		✓ Ketocal 3:1
Powder (vanilla)36.92	300 g OP	✓ KetoCal 4:1

# **SECTION I: NATIONAL IMMUNISATION SCHEDULE**

Subsidy (Manufacturer's Price) \$ P

Subsidised Per 🗸

Fully

Brand or Generic Manufacturer

# **Vaccinations**

#### BACILLUS CALMETTE-GUERIN VACCINE - [Xpharm]

For infants at increased risk of tuberculosis. Increased risk is defined as:

- 1) living in a house or family with a person with current or past history of TB; or
- 2) having one or more household members or carers who within the last 5 years lived in a country with a rate of TB > or equal to 40 per 100,000 for 6 months or longer; or
- 3) during their first 5 years will be living 3 months or longer in a country with a rate of TB > or equal to 40 per 100,000 Note a list of countries with high rates of TB are available at www.health.govt.nz/tuberculosis (search for downloads) or www.bcgatlas.org/index.php.
- Inj Mycobacterium bovis BCG (Bacillus Calmette-Guerin),

Danish strain 1331, live attenuated, vial with diluent................0.00

10

✓ BCG Vaccine AJV

	Subsidy		Fully	
	(Manufacturer's Price) \$	Per	Subsidised	I Generic Manufacturer
VID-19 VACCINE - [Xpharm]				
Inj 10 mcg raxtozinameran per 0.3 ml, 0.48 ml vial; paediatric vaccine, light blue cap	0.00	10	•	Comirnaty Omicron (XBB.1.5)
Either:				(ADD:1.0)
One dose for previously unvaccinated children aged     Up to three doses for immunocompromised children	•	l.		
Inj 3 mcg raxtozinameran per 0.2 ml, 0.4 ml vial; infant vaccin maroon cap		10	,	Comirnaty Omicron
παισστισαμ			•	(XBB.1.5)
Up to three doses for previously unvaccinated children ag	ged 6 months - 4 yea	rs at	high risk o	of severe illness.
Inj 30 mcg raxtozinameran per 0.3 ml, 0.48 ml vial; adult				
vaccine, light grey cap	0.00	10	✓	<b>Comirnaty Omicron</b>
Any of the following:				(XBB.1.5)
<ol> <li>Up to three doses for immunocompromised people</li> <li>Up to two doses for previously unvaccinated people</li> <li>Up to four doses for people aged 16-29 at high risk</li> <li>One dose for previously unvaccinated people aged</li> <li>One additional dose every 6 months for previously given at least 6 months after last dose.</li> </ol>	16-29 years old; or of severe illness; or 30 and older; or		0 years ar	nd over – additional dose i
Inj 30 mcg raxtozinameran per 0.3 ml, 2.25 ml vial; adult				
vaccine, dark grey cap	0.00	10	✓	Comirnaty Omicron (XBB.1.5)
Any of the following:				
<ol> <li>One dose for previously unvaccinated people aged</li> <li>Up to three doses for immunocompromised people</li> <li>Up to two doses for previously unvaccinated people</li> <li>Up to four doses for people aged 16-29 at high risk</li> <li>One dose for previously unvaccinated people aged</li> <li>One additional dose every 6 months for previously vision at least 6 months after last dose.</li> </ol>	aged 12-15 years old 16-29 years old; or of severe illness; or 30 and older; or		0 years ar	nd over – additional dose i
Inj 3 mcg bretovameran per 0.3 ml, 0.48 ml vial; infant vaccing yellow cap		10	•	Comirnaty Omicron
Up to three doses for previously unvaccinated children ag	ged 6 months - 4 yea	rs at	high risk o	(JN.1) of severe illness.
Inj 10 mcg bretovameran per 0.3 ml, 0.48 ml vial; paediatric				
vaccine, light blue cap	0.00	10	•	Comirnaty Omicron (JN.1)
Either:				(0)
<ol> <li>One dose for previously unvaccinated children aged</li> <li>Up to three doses for immunocompromised children</li> </ol>		l.		

	Subsidy (Manufacturer's Price) \$	Sub Per	Fully sidised	Brand or Generic Manufacturer
Inj 30 mcg bretovameran per 0.3 ml, 0.48 ml vial; adult vaccii light grey cap	,	10		omirnaty Omicron (JN.1)

Any of the following:

- 1) One dose for previously unvaccinated people aged 12-15 years old; or
- 2) Up to three doses for immunocompromised people aged 12-15 years old: or
- 3) Up to two doses for previously unvaccinated people 16-29 years old; or
- 4) Up to four doses for people aged 16-29 at high risk of severe illness; or
- 5) One dose for previously unvaccinated people aged 30 and older; or
- 6) One additional dose every 6 months for previously vaccinated people aged 30 years and over additional dose is given at least 6 months after last dose.

(Comirnaty Omicron (XBB.1.5) Inj 10 mcg raxtozinameran per 0.3 ml, 0.48 ml vial; paediatric vaccine, light blue cap to be delisted 1 February 2025)

(Comirnaty Omicron (XBB.1.5) Inj 3 mcg raxtozinameran per 0.2 ml, 0.4 ml vial; infant vaccine, maroon cap to be delisted 1 February 2025)

(Comirnaty Omicron (XBB.1.5) Inj 30 mcg raxtozinameran per 0.3 ml, 0.48 ml vial; adult vaccine, light grey cap to be delisted 1 February 2025)

(Comirnaty Omicron (XBB.1.5) Ini 30 mcg raxtozinameran per 0.3 ml. 2.25 ml vial; adult vaccine, dark grey cap to be delisted 1 February 2025)

#### DIPHTHERIA. TETANUS AND PERTUSSIS VACCINE

- a) Only on a prescription
- b) No patient co-payment payable

c)

- A) Funded for any of the following criteria:
  - 1) A single dose for pregnant women in the second or third trimester of each pregnancy; or
  - 2) A single dose for parents or primary caregivers of infants admitted to a Neonatal Intensive Care Unit or Specialist Care Baby Unit for more than 3 days, who had not been exposed to maternal vaccination at least 14 days prior to birth: or
  - 3) A course of up to four doses is funded for children from age 7 up to the age of 18 years inclusive to complete full primary immunisation; or
  - 4) An additional four doses (as appropriate) are funded for (re-)immunisation for patients post haematopoietic stem cell transplantation or chemotherapy; pre or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or
  - 5) A single dose for vaccination of patients aged from 65 years old; or
  - 6) A single dose for vaccination of patients aged from 45 years old who have not had 4 previous tetanus doses; or
  - 7) For vaccination of previously unimmunised or partially immunised patients; or
  - 8) For revaccination following immunosuppression; or
  - 9) For boosting of patients with tetanus-prone wounds.

Notes: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.

- B) Contractors will be entitled to claim payment from the Funder for the supply of diphtheria, tetanus and pertussis vaccine to patients eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the diphtheria, tetanus and pertussis vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraphs 1-9 above.

Inj 2 10 diphtheria toxold with 20 10 tetanus toxolo	, 8 mcg
pertussis toxoid, 8 mcg pertussis filamentous	

haemagglutinin and 2.5 mcg pertactin in 0.5 ml prefilled

**Boostrix** 

Subsidy		Fully	Brand or	
(Manufacturer's Price)	5	Subsidised	Generic	
\$	Per	✓	Manufacturer	

#### DIPHTHERIA, TETANUS, PERTUSSIS AND POLIO VACCINE

- a) Only on a prescription
- b) No patient co-payment payable

c)

- A) Funded for any of the following:
  - 1) A single dose for children up to the age of 7 who have completed primary immunisation; or
  - A course of four vaccines is funded for catch up programmes for children (to the age of 10 years) to complete full primary immunisation; or
  - 3) An additional four doses (as appropriate) are funded for (re-)immunisation for people post HSCT, or chemotherapy; pre- or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or
  - 4) Five doses will be funded for children requiring solid organ transplantation.
- B) Contractors will be entitled to claim payment from the Funder for the supply of Diphtheria, tetanus, pertussis and polio vaccine to people eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Diphtheria, tetanus, pertussis and polio vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.

Inj 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg

pertussis toxoid, 25 mcg pertussis filamentous

haemagglutinin, 8 mcg pertactin and 80 D-antigen units

10 ✓ Infanrix IPV

### DIPHTHERIA, TETANUS, PERTUSSIS, POLIO, HEPATITIS B AND HAEMOPHILUS INFLUENZAE TYPE B VACCINE

- a) Only on a prescription
- b) No patient co-payment payable

c)

- A) Funded for children meeting any of the following criteria
  - 1) Up to four doses for children under the age of 10 years for primary immunisation; or
  - An additional four doses (as appropriate) for (re-)immunisation of children under the age of 18 years post haematopoietic stem cell transplantation; or
  - 3) An additional four doses (as appropriate) for (re-)immunisation of children under the age of 10 years who are post chemotherapy; pre or post splenectomy; undergoing renal dialysis and other severely immunosuppressive regimens; or
  - 4) Up to five doses for children under the age of 10 years receiving solid organ transplantation.
- B) Contractors will be entitled to claim payment from the Funder for the supply of Diphtheria, tetanus, pertussis, polio, hepatitis B and haemophilus influenzae type b vaccine to people eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Diphtheria, tetanus, pertussis, polio, hepatitis B and haemophilus influenzae type b vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Note: A course of up-to four vaccines is funded for catch up programmes for children (up to and under the age of 10 years) to complete full primary immunisation. Please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes.

Inj 30IU diphtheria with 40IU tetanus and 25mcg pertussis toxoids, 25mcg pertussis filamentous haemagglutinin, 8mcg pertactin, 80D-AgU polio virus, 10mcg hepatitis B antigen, 10mcg H. influenzae type b with tetanus toxoid

10

✓ Infanrix-hexa

✓ Havrix Junior

			NATIONAL	IIVIIVI	JNISATI	ON SCHEDOLL
			Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
HAEM	OPHII	LUS INFLUENZAE TYPE B VACCINE				
a)	Only	on a prescription				
b)	No p	atient co-payment payable				
c)						
	A)	One dose for people meeting any of the following:				
		<ol> <li>For primary vaccination in children; or</li> <li>An additional dose (as appropriate) is funded</li> </ol>	for (re-)immunication f	or no	onle noet he	ematonojetic stem cell
		transplantation, or chemotherapy; functional a				
		transplant, pre or post cochlear implants, rena				
		3) For use in testing for primary immunodeficien	cy diseases, on the re	comm	endation of	an internal medicine
		physician or paediatrician.				
	B)					
		vaccine to people eligible under the above criteria programme for subsidised immunisation, and they may only do				,
		in the Pharmaceutical Schedule.	so in respect of the ric	aemo <sub>k</sub>	Jillius IIIIlue	lizae type b vaccine listeu
	C)	Contractors may only claim for populations within the	ne criteria that are cove	ered b	v their cont	ract, which may be a
	,	sub-set of the population described in paragraph A			,	
lnj	10 m	cg vial with diluent syringe	0.00	1	✓ A	ct-HIB
HEPAT	TITIS .	A VACCINE - [Xpharm]				
Fu	nded	for patients meeting any of the following criteria:				
		o vaccinations for use in transplant patients; or				
	,	o vaccinations for use in children with chronic liver of	·			
(	3) Or	e dose of vaccine for close contacts of known hepa	titis A cases.			
lnj	1440	ELISA units in 1 ml syringe	0.00	1	<b>✓</b> <u>H</u>	avrix 1440

Inj 720 ELISA units in 0.5 ml syringe.......0.00

	Subsidy		Fully	Brand or	
	(Manufacturer's Price)	Sı	bsidised	Generic	
	\$	Per	<b>✓</b>	Manufacturer	
HEPATITIS B RECOMBINANT VACCINE - [Xpharm]					

✓ Engerix-B

Funded for patients meeting any of the following criteria:

- 1) for household or sexual contacts of known acute hepatitis B patients or hepatitis B carriers; or
- 2) for children born to mothers who are hepatitis B surface antigen (HBsAq) positive; or
- 3) for children up to and under the age of 18 years inclusive who are considered not to have achieved a positive serology and require additional vaccination or require a primary course of vaccination; or
- 4) for HIV positive patients: or
- 5) for hepatitis C positive patients; or
- 6) for patients following non-consensual sexual intercourse; or
- 7) for patients prior to planned immunosuppression for greater than 28 days; or
- 8) for patients following immunosuppression; or
- 9) for solid organ transplant patients; or
- 10) for post-haematopoietic stem cell transplant (HSCT) patients; or
- 11) following needle stick injury.

Inj 20 mcg per 1 ml prefilled syringe......0.00 ✓ Engerix-B

Funded for patients meeting any of the following criteria:

- 1) for household or sexual contacts of known acute hepatitis B patients or hepatitis B carriers; or
- 2) for children born to mothers who are hepatitis B surface antigen (HBsAg) positive; or
- 3) for children up to and under the age of 18 years inclusive who are considered not to have achieved a positive serology and require additional vaccination or require a primary course of vaccination; or
- 4) for HIV positive patients; or
- 5) for hepatitis C positive patients; or
- 6) for patients following non-consensual sexual intercourse; or
- 7) for patients prior to planned immunosuppression for greater than 28 days; or
- 8) for patients following immunosuppression; or
- 9) for solid organ transplant patients; or
- 10) for post-haematopoietic stem cell transplant (HSCT) patients; or
- 11) following needle stick injury; or
- 12) for dialysis patients; or
- 13) for liver or kidney transplant patients.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

HUMAN PAPILLOMAVIRUS (6, 11, 16, 18, 31, 33, 45, 52 AND 58) VACCINE [HPV]

- a) Maximum of 1 inj per prescription
- b) Only on a prescription
- c) No patient co-payment payable
- d
- a) A) Any of the following:
  - 1) Maximum of two doses for children aged 14 years and under; or
  - 2) Maximum of three doses for people meeting any of the following criteria:
    - 1) People aged 15 to 26 years inclusive; or
    - 2) Either:

People aged 9 to 26 years inclusive who have

- 1) Confirmed HIV infection; or
- 2) Received a transplant (including stem cell): or
- 3) Maximum of four doses for people aged 9 to 26 years inclusive post chemotherapy
- B) Contractors will be entitled to claim payment from the Funder for the supply of Human papillomavirus vaccine to people eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Human papillomavirus vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraphs A above.

nj 270 mcg in 0.5 ml syringe	0.00	10	Gardasil 9
------------------------------	------	----	------------

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
INFLUENZA VACCINE Inj 60 mcg in 0.5 ml syringe (quadrivalent vaccine)	120.00	10		nfluvac Tetra (2024 formulation)

Subsidy		Fully	Brand or	
(Manufacturer's Price)	;	Subsidised	Generic	
\$	Per	✓	Manufacturer	

- a) Maximum of 1 inj per prescription
- b) Only on a prescription
- c) No patient co-payment payable
- d

#### A) INFLUENZA VACCINE

is available each year for patients who meet the following criteria, as set by Pharmac:

- a) all people 65 years of age and over; or
- b) people under 65 years of age who:
  - i) have any of the following cardiovascular diseases:
    - a) ischaemic heart disease, or
    - b) congestive heart failure, or
    - c) rheumatic heart disease, or
    - d) congenital heart disease, or
    - e) cerebo-vascular disease; or
  - ii) have either of the following chronic respiratory diseases:
    - a) asthma, if on a regular preventative therapy, or
    - b) other chronic respiratory disease with impaired lung function; or
  - iii) have diabetes; or
  - iv) have chronic renal disease; or
  - v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
  - vi) have any of the following other conditions:
    - a) autoimmune disease, or
    - b) immune suppression or immune deficiency, or
    - c) HIV, or
    - d) transplant recipients, or
    - e) neuromuscular and CNS diseases/disorders, or
    - f) haemoglobinopathies, or
    - g) are children on long term aspirin, or
    - h) have a cochlear implant, or
    - i) errors of metabolism at risk of major metabolic decompensation, or
    - i) pre and post splenectomy, or
    - k) Down syndrome, or
  - vii) are pregnant; or
- c) children 4 years of age and under who have been hospitalised for respiratory illness or have a history of significant respiratory illness; or
- d) people under 65 years of age who:
  - i) have any of the following serious mental health conditions:
    - a) schizophrenia, or
    - b) major depressive disorder, or
    - c) bipolar disorder, or
    - d) schizoaffective disorder, or
  - ii) are currently accessing secondary or tertiary mental health and addiction services; or

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy,
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Contractors will be entitled to claim payment for the supply of influenza vaccine to patients eligible under the above criteria pursuant to their contract with Health NZ for subsidised immunisation, and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	✓	Manufacturer	

#### MEASLES. MUMPS AND RUBELLA VACCINE

- a) Only on a prescription
- b) No patient co-payment payable

c)

#### A) Measles, mumps and rubella vaccine

A maximum of two doses for any patient meeting the following criteria:

- 1) For primary vaccination in children; or
- 2) For revaccination following immunosuppression; or
- 3) For any individual susceptible to measles, mumps or rubella; or
- 4) A maximum of three doses for children who have had their first dose prior to 12 months.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes. Although a price is listed for the vaccine, doctors can still order measles mumps and rubella vaccine free of charge, as with other Schedule vaccines.

- B) Contractors will be entitled to claim payment for the supply of measles, mumps and rubella vaccine to patients eligible under the above criteria pursuant to their contract with Health NZ for subsidised immunisation, and they may only do so in respect of the measles, mumps and rubella vaccine listed in the Pharmaceutical Schedule.
- C) Contractors can only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Inj, measles virus 1,000 CCID50, mumps virus 5,012 CCID50,			
Rubella virus 1,000 CCID50; prefilled syringe/ampoule of			
diluent 0.5 ml	0.00	10	Priorix

Subsidy (Manufacturer's Price)	Sub	Fully	Brand or Generic
 \$	Per	✓	Manufacturer

#### MENINGOCOCCAL (GROUPS A. C. Y AND W-135) CONJUGATE VACCINE

Inj 10 mcg of each meningococcal polysaccharide conjugated to a total of approximately 55 mcg of tetanus toxoid carrier 

✓ MenQuadfi

- a) Only on a prescription
- b) No patient co-payment payable

- A) Any of the following:
  - 1) Up to three doses and a booster every five years for patients pre- and post splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre or post solid organ transplant; or
  - 2) One dose for close contacts of meningococcal cases of any group; or
  - 3) One dose for person who has previously had meningococcal disease of any group; or
  - 4) A maximum of two doses for bone marrow transplant patients; or
  - 5) A maximum of two doses for person pre- and post-immunosuppression\*: or
- B) Both:
  - 1) Person is aged between 13 and 25 years, inclusive; and
  - 2) Either:
    - 1) One dose for individuals who are entering within the next three months, or in their first year of living in boarding school hostels, tertiary education halls of residence, military barracks, Youth Justice residences, or prisons; or
    - 2) One dose for individuals who turn 13 years of age while living in boarding school hostels.
- C) Contractors will be entitled to claim payment from the Funder for the supply of Meningococcal A, C, Y and W-135 vaccine to patients eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Meningococcal A, C, Y and W-135 vaccine listed in the Pharmaceutical Schedule.
- D) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraphs A-B above.

Note: children under seven years of age require two doses 8 weeks apart, a booster dose three years after the primary series and then five yearly.

\*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than

Ini 5 mcg of each meningococcal polysaccharide conjugated to a total of approximately 44 mcg of tetanus toxoid carrier

✓ Nimenrix

- A) Both:
  - 1) The child is under 12 months of age; and
  - 2) Any of the following:
    - 1) A maximum of three doses (dependant on age at first dose) for patients pre- and post- splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre- or post- solid organ transplant; or
    - 2) A maximum of three doses (dependant on age at first dose) for close contacts of meningococcal cases
    - 3) A maximum of three doses (dependant on age at first dose) for child who has previously had meningococcal disease of any group; or
    - 4) A maximum of three doses (dependant on age at first dose) for bone marrow transplant patients; or
    - 5) A maximum of three doses (dependant on age at first dose) for child pre- and post-immunosuppression\*.

Note: infants from 6 weeks to less than 6 months of age require a 2+1 schedule, infants from 6 months to less than 12 months of age require a 1+1 schedule. Refer to the Immunisation Handbook for recommended booster schedules with meningococcal ACWY vaccine.

\*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days.

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
<b>\$</b>	Per	✓	Manufacturer	

#### MENINGOCOCCAL B MULTICOMPONENT VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- c) Any of the following:
  - A) Three doses for children up to 12 months of age (inclusive) for primary immunisation; or
  - B) Up to three doses (dependent on age at first dose) for a catch-up programme for children from 13 months to 59 months of age (inclusive) for primary immunisation, from 1 March 2023 to 31 August 2025; or
  - C) Both:
    - 1) Person is one year of age or over; and
    - 2) Any of the following:
      - i) up to two doses and a booster every five years for patients pre- and post-splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre- or post-solid organ transplant; or
      - ii) up to two doses for close contacts of meningococcal cases of any group; or
      - iii) up to two doses for person who has previously had meningococcal disease of any group; or
      - iv) up to two doses for bone marrow transplant patients; or
      - v) up to two doses for person pre- and post-immunosuppression\*; or
  - D) Both:
    - 1) Person is aged between 13 and 25 years (inclusive); and
    - 2) Either:
      - Two doses for individuals who are entering within the next three months, or in their first year of living in boarding school hostels, tertiary education halls of residence, military barracks, Youth Justice residences or prisons; or
      - ii) Two doses for individuals who turn 13 years of age while living in boarding school hostels.
  - E) Contractors will be entitled to claim payment from the Funder for the supply of Meningococcal B multicomponent vaccine to patients eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Meningococcal B multicomponent vaccine listed in the Pharmaceutical Schedule.
  - F) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraphs A-D above.

\*Immunosuppression due to corticosteroid or other immunosuppressive therapy must be for a period of greater than 28 days.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

#### PNEUMOCOCCAL (PCV13) CONJUGATE VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- c)
- A) Any of the following:
  - 1) A course of three doses for previously unvaccinated children up to the age of 59 months inclusive; or
  - Two doses are funded for high risk individuals (over the age of 12 months and under 18 years) who have previously received two doses of the primary course of PCV10: or
  - 3) Up to an additional four doses (as appropriate) are funded for the (re)immunisation of high risk children aged under 5 years with any of the following:
    - a) on immunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient immune response; or
    - b) primary immune deficiencies; or
    - c) HIV infection: or
    - d) renal failure, or nephrotic syndrome; or
    - e) who are immune-suppressed following organ transplantation (including haematopoietic stem cell transplant); or
    - f) cochlear implants or intracranial shunts; or
    - g) cerebrospinal fluid leaks; or
    - h) receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage of 20 mg or greater; or
    - i) chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or
    - j) pre term infants, born before 28 weeks gestation; or
    - k) cardiac disease, with cyanosis or failure; or
    - diabetes: or
    - m) Down syndrome; or
    - n) who are pre-or post-splenectomy, or with functional asplenia; or
  - 4) Up to an additional four doses (as appropriate) are funded for the (re-)immunisation of individuals 5 years and over with HIV, pre or post haematopoietic stem cell transplantation, or chemotherapy; pre- or post splenectomy; functional asplenia, pre- or post- solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, intracranial shunts, cerebrospinal fluid leaks or primary immunodeficiency; or
  - 5) For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.
- B) Contractors will be entitled to claim payment from the Funder for the supply of Pneumococcal (PCV13) conjugate vaccine to people eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Pneumococcal (PCV13) conjugate vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Note: please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes Inj 30.8 mcg of pneumococcal polysaccharide serotypes 1, 3, 4,

5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and 23F in 0.5ml		
syringe0.00	10	✓ Prevenar 13
	1	✓ Prevenar 13

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer PNEUMOCOCCAL (PPV23) POLYSACCHARIDE VACCINE - [Xpharm] Either: 1) Up to three doses (as appropriate) for patients with HIV, for patients post haematopoietic stem cell transplant, or chemotherapy; pre- or post-splenectomy or with functional asplenia, pre- or post-solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, or primary immunodeficiency; or 2) All of the following: a) Patient is a child under 18 years for (re-)immunisation; and b) Treatment is for a maximum of two doses; and c) Any of the following: i) on immunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient immune response; or ii) with primary immune deficiencies; or iii) with HIV infection; or iv) with renal failure, or nephrotic syndrome; or v) who are immune-suppressed following organ transplantation (including haematopoietic stem cell transplant); vi) with cochlear implants or intracranial shunts; or vii) with cerebrospinal fluid leaks; or viii) receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage of 20 mg or greater: or ix) with chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or x) pre term infants, born before 28 weeks gestation; or xi) with cardiac disease, with cyanosis or failure; or xii) with diabetes: or xiii) with Down syndrome; or xiv) who are pre-or post-splenectomy, or with functional asplenia. Inj 575 mcg in 0.5 ml prefilled syringe (25 mcg of each 1 ✓ Pneumovax 23 POLIOMYELITIS VACCINE - [Xpharm] Up to three doses for patients meeting either of the following: 1) For partially vaccinated or previously unvaccinated individuals: or

2) For revaccination following immunosuppression.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch-up programmes.

✓ IPOL

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

#### **ROTAVIRUS ORAL VACCINE**

- a) Only on a prescription
- b) No patient co-payment payable
- c)
- A) Maximum of two doses for people meeting the following:
  - 1) first dose to be administered in infants aged under 14 weeks of age: and
  - 2) no vaccination being administered to children aged 24 weeks or over.
- B) Contractors will be entitled to claim payment from the Funder for the supply of Rotavirus oral vaccine to people eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Rotavirus oral vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Oral susp live attenuated human rotavirus			
1,000,000 CCID50 per dose, squeezable tube0.	.00	10	✓ Rotarix
Oral susp live attenuated human rotavirus			
1,000,000 CCID50 per dose, prefilled oral applicator0.	.00	10	✓ Rotarix

(Manu	Subsidy	Fully	Brand or
	facturer's Price)	Subsidised	Generic
	\$ Per	✓	Manufacturer

#### VARICELLA VACCINE [CHICKENPOX VACCINE]

- a) Only on a prescription
- b) No patient co-payment payable
- c)
- A) Either:
  - 1) Maximum of one dose for primary vaccination for either:
    - a) Any infant born on or after 1 April 2016; or
    - For previously unvaccinated children turning 11 years old on or after 1 July 2017, who have not previously had a varicella infection (chickenpox), or
  - 2) Maximum of two doses for any of the following:
    - a) Any of the following for non-immune individuals:
      - i) with chronic liver disease who may in future be candidates for transplantation; or
      - ii) with deteriorating renal function before transplantation; or
      - iii) prior to solid organ transplant; or
      - iv) prior to any elective immunosuppression\*; or
      - v) for post exposure prophylaxis who are immune competent inpatients; or
    - b) For individuals at least 2 years after bone marrow transplantation, on advice of their specialist; or
    - c) For individuals at least 6 months after completion of chemotherapy, on advice of their specialist; or
    - d) For HIV positive non immune to varicella with mild or moderate immunosuppression on advice of HIV specialist; or
    - e) For individuals with inborn errors of metabolism at risk of major metabolic decompensation, with no clinical history of varicella; or
    - f) For household contacts of paediatric patients who are immunocompromised, or undergoing a procedure leading to immune compromise where the household contact has no clinical history of varicella; or
    - g) For household contacts of adult patients who have no clinical history of varicella and who are severely immunocompromised, or undergoing a procedure leading to immune compromise where the household contact has no clinical history of varicella.
- B) Contractors will be entitled to claim payment from the Funder for the supply of Varicella vaccine [Chickenpox vaccine] vaccine to people eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Varicella vaccine [Chickenpox vaccine] listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraphs A above.

\* immunosuppression due to steroid or other immunosuppressive therapy must be for a treatment period of greater than 28 days

# MATIONAL IMMUNICATION COLEDINE

	NATIONAL	IIVIIVIUNISATI	ON SCHEDULE
	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
VARICELLA ZOSTER VACCINE [SHINGLES VACCINE]			
a) Only on a prescription			
b) No patient co-payment payable			
c)			
A) Funded for patients meeting the following criteria:			
1) Either:			
1) Two doses for all people aged 65 years	, or		

- 2) Two doses for people 18 years of age or older with any of the following:
  - a) pre- and post-haematopoietic stem cell transplant or cellular therapy; or
  - b) pre- or post-solid organ transplant; or
  - c) haematological malignancies; or
  - d) people living with poorly controlled HIV infection; or
  - e) planned or receiving disease modifying anti-rheumatic drugs (DMARDs targeted synthetic, biologic, or conventional synthetic) for polymyalgia rheumatica, systemic lupus erythematosus or rheumatoid arthritis: or
  - f) end stage kidney disease (CKD 4 or 5); or
  - g) primary immunodeficiency
- B) Contractors will be entitled to claim payment from the Funder for the supply of Varicella zoster vaccine (Shingles vaccine) to patients eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Varicella zoster vaccine [Shingles vaccine] listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Inj 50 mcg per 0.5 ml vial plus vial	0.00	1	✓ Shingrix
		10	✓ Shingrix

# **Diagnostic Agents**

TUBERCULIN PPD [MANTOUX] TEST - [Xpharm]			
Inj 5 TU per 0.1 ml, 1 ml vial	0.00	1	✓ Tubersol

	Symbols -		Albendazole	95	Amzoate	28
		113	Albey		Anaesthetics	
	- A -		Albustix		Anafranil	129
A-Scabies		73	Alchemy Oxaliplatin	155	Anagrelide hydrochloride	
Abacavir sulpha	ate	113	Alchemy Oxybutynin		Analgesics	
Abacavir sulpha			Aldurazyme		Anastrozole	
		113	Alecensa		Anatrole	179
	udine Viatris		Alectinib		Anoro Ellipta	
Abilify Maintena	a	138	Alendronate sodium	118	Antabuse	150
	a S29		Alendronate sodium with		Antacids and Antiflatulents	
	etate		colecalciferol	118	Anthelmintics	9
			Alfacalcidol		Antiacne Preparations	
Accarb		11	Alfamino	296	Antiallergy Preparations	25
Acetazolamide		269	Alfamino Junior	296	Antianaemics	
Acetec		45	Alginic acid	6	Antiandrogen Oral	
Acetic acid with	n hydroxyquinoline	and	Alglucosidase alfa		Contraceptives	8 <sup>.</sup>
	id		Alkeran		Antiarrhythmics	
			Allerfix		Antibacterials	
			Allerpro Syneo 1	298	Antibacterials Topical	
Aciclovir			Allerpro Syneo 2		Anticholinergic Agents	
Infection		108	Allersoothe		Anticholinesterases	
			Allmercap		Antidepressants	
,			Allopurinol		Antidiarrhoeals	
Acipimox		53	Almarytm		Antiepilepsy Drugs	
•			Alpha-Adrenoceptor Blockers.		Antifibrinolytics, Haemostatics an	
			Alpha-Keri Lotion		Local Sclerosants	
			Alphamox 125		Antifibrotics	
			Alphamox 250		Antifungals	
•			Alprolix		Antifungals Topical	
			Alu-Tab		Antihistamines	
			Aluminium hydroxide	6	Antihypotensives	
•	mgevita)		Alyacen		Antimalarials	
,	łumira - Alternative		Amantadine hydrochloride		Antimigraine Preparations	134
,			Ambrisentan		Antinausea and Vertigo Agents	
Adapalene		67	Ambrisentan Viatris	<u>56</u>	Antipruritic Preparations	69
Adcetris		203	Amgevita	187	Antipsychotics	
ADR Cartridge	1.8	21	Amiloride hydrochloride	51	Antiretrovirals	11
Adrenaline			Amiloride hydrochloride with		Antirheumatoid Agents	118
Cardiovascu	ılar	55	furosemide	<mark>52</mark>	Antispasmodics and Other Agent	S
Respiratory.		257	Amiloride hydrochloride with		Altering Gut Motility	8
Advantan		70	hydrochlorothiazide	<mark>52</mark>	Antithrombotic Agents	39
Advate		38	Aminophylline	264	Antithymocyte globulin	
Adynovate		39	Amiodarone hydrochloride	47	(equine)	186
Afinitor		252	Amisulpride	136	Antitrichomonal Agents	106
Aflibercept		202	Amitriptyline	129	Antituberculotics and	
AFT-Pyrazinam	nide	107	Amlodipine	<mark>50</mark>	Antileprotics	
Agents Affectin	g the		Amorolfine	68	Antiulcerants	9
	tensin System		Amoxicillin		Antivirals	
	kinsonism and Rela		Amoxicillin with clavulanic acid		Anxiolytics	
Disorders		123	Amoxiclav Devatis Forte		Anzatax	
Agents Used in	the Treatment of		Amphotericin B		Apidra	
			Amsacrine		Apidra SoloStar	
			AmsaLyo		APO-Atomoxetine	
Albalon		271	Amsidine	158	Apo-Azithromycin	96

APO-Candesartan HCTZ		ATGAM	186	Benzylpenicillin sodium [Penicillin	
16/12.5	46	Ativan		G]	98
APO-Candesartan HCTZ		Atnahs Olsalazine		Beta Cream	
32/12.5	46	Atomoxetine		Beta Ointment	
Apo-Temozolomide		Atorvastatin		Beta Scalp	
Apomorphine hydrochloride		Atropine sulphate		Beta-Adrenoceptor Agonists	
Aprepitant		Cardiovascular	47	Beta-Adrenoceptor Blockers	
Apresoline		Sensory		Beta-hCG low sensitivity urine test	
Aptamil Feed Thickener		Atropt		kit	8
Aqueous cream		Atrovent		Betadine	
Aratac		Aubagio		Betadine Skin Prep	
Arava		Augmentin		Betaferon	
Arginine		Aurorix		Betahistine dihydrochloride	
Arginine2000		AutoSoft 30		Betaine	
Aripiprazole		AutoSoft 90		Betamethasone dipropionate	
Aripiprazole Sandoz		Avelox		Betamethasone dipropionate with	
Aristocort		Avonex		calcipotriol	7,
Arrotex-Prazosin S29		Avonex Pen		Betamethasone sodium phosphate	
Arrow - Clopid		Azacitidine		with betamethasone acetate	
Arrow - Lattim		Azacitidine Dr Reddy's		Betamethasone valerate	
Arrow-Amitriptyline		Azamun		Betamethasone valerate with sodiu	
Arrow-Bendrofluazide		Azathioprine		fusidate [fusidic acid]	
Arrow-Brimonidine		Azilect		Betaxolol	
Arrow-Diazepam		Azithromycin		Betnovate	
Arrow-Doxorubicin		Azopt		Betoptic	
Arrow-Fluoxetine		AZT		Betoptic S	
Arrow-Losartan &	100	-B-	110	Bexsero	
Hydrochlorothiazide	46	B-D Micro-Fine	16	Bezafibrate	
Arrow-Norfloxacin		B-D Ultra Fine		Bezalip	
Arrow-Ornidazole		B-D Ultra Fine II		Bezalip Retard	
Arrow-Quinapril 10		Bacillus Calmette-Guerin (BCG		Bicalutamide	
Arrow-Quinapril 20		vaccine	,	Bicillin LA	
		Bacillus Calmette-Guerin	100	BiCNU	
Arrow-Quinapril 5		vaccine	201	BiCNU S29	
Arrow-Roxithromycin		Baclofen		Bile and Liver Therapy	
Arrow-Timolol		Bactroban		Biltricide	
Arrow-Topiramate					
Arrow-Tramadol		Balance		Bimatoprost	
Arsenic trioxide		Barrier Creams and Emollients		Binarex Binocrit	
Asacol S29		BCG Vaccine AJV Beclazone 100		Biodone	
Ascend		Beclazone 250		Biodone Extra Forte	
Ascend Aripiprazole Ascend-Cefuroxime		Beclazone 50		Biodone Forte	
		Beclomethasone dipropionate.		Bisacodyl	
Ascorbic acid		Bedaquiline		Bisacodyl Viatris	24
Aspen Adrenaline		Bee venom allergy treatment		Bisoprolol fumarate	
Aspirin	00	Bendamustine hydrochloride		BK Lotion	
Blood		Bendamustine Sandoz		Bleomycin sulphate	15
Nervous		Bendrofluazide	52	Blood Colony-stimulating	
Asthalin		Bendroflumethiazide		Factors	42
Atazanavir Mylan		[Bendrofluazide]		Blood glucose diagnostic test	4
Atazanavir sulphate	113	Benralizumab		meter	18
Atazanavir Viatris		Benzathine benzylpenicillin		Blood glucose diagnostic test	4
Atenolol		Benzatropine mesylate		strip	1
Atenolol AFT		Benzbromarone		Blood glucose test strips (visually	4
Atenolol Viatris		Benztrop		impaired)	1
Atezolizumab	244	Benzydamine hydrochloride	30	Blood Ketone Diagnostic Test	

# **INDEX: Generic Chemicals and Brands**

Strip	14	Calcium polystyrene sulphona	te44	Cetomacrogol with glycerol	7
Boostrix		Calcium Resonium		Cetomacrogol-AFT	<mark>7</mark>
Bortezomib	159	Calogen	278	Cetuximab	20
Bosentan	58	Camber	55	Charcoal	27
Bosentan Dr Reddy's	58	Candesartan cilexetil	46	CheckTop	8
Bplex		Candesartan cilexetil with		Chemotherapeutic Agents	
Brentuximab Vedotin		hydrochlorothiazide	46	Chickenpox vaccine	
Breo Ellipta		Candestar		Chlorambucil	
Brevinor 1/28		Canesten		Chloramphenicol	
Breztri Aerosphere		Capecitabine		Chlorothiazide	
Bricanyl Turbuhaler		Capecitabine Viatris		Chlorpromazine hydrochloride	
Brimonidine tartrate		Capsaicin		Chlorsig	
Brimonidine tartrate with timolol		Musculoskeletal	118	Chlortalidone [Chlorthalidone]	
maleate	270	Nervous		Chlorthalidone	5
Brinzolamide		Captopril		Chlorvescent	
Brufen SR		Carafate		Choice 380 7med Nsha Silver/co	
BSF Midazolam-Baxter		Carbaccord		Short	
Buccastem		Carbamazepine		Ciclosporin	
Budesonide		Carbimazole		Cidomycin P/Free	
Alimentary	6	Carbomer		Cilazapril	10
Respiratory2		Carbonler		Cilicaine VK	a
Budesonide Te Arai		Carboplatin Accord		Cinacalcet	
Budesonide with eformoterol		Carbosorb-X		Cinacalet Devatis	
Budesonide with glycopyrronium		Cardinol LA		Ciprofloxacin	0
eformoterol		Cardizem CD		Infection	Q
Bumetanide		CareSens Dual		Sensory	
Buprenorphine Naloxone BNM		CareSens N		Ciprofloxacin Teva	
		CareSens N POP		Cisplatin	20 15
Buprenorphine with naloxone	150	CareSens N Premier		Cisplatin Accord	10 15
Bupropion hydrochloride					
		CareSens PRO		Cisplatin Ebewe Citalopram hydrobromide	روا ده
Burinex		Carmellose sodium with gelati		Citrulline1000	
Buscopan		pectin		Cladribine	
Buspirone hydrochloride		Carmustine			10
Buspirone Viatris		Carnitor		Clarithromycin	
Busulfan	154	Carvedilol		Alimentary	
	0.4	Carvedilol Sandoz		Infection	9
Cabergoline		Casirivimab and imdevimab		Clexane	
Caffeine citrate		Catapres		Clexane Forte	
Calamine		Cefaclor monohydrate		Clindamycin	
Calci-Tab 500		Cefalexin		Clinicians	
Calcipotriol		Cefalexin Sandoz		Clinicians Renal Vit	
Calcitonin		Cefazolin		Clobazam	
Calcitriol		Cefazolin-AFT		Clobetasol propionate	/0, /
Calcitriol-AFT		Ceftriaxone		Clobetasone butyrate	
Calcitriol-AFT S29		Ceftriaxone-AFT		Clofazimine	10
Calcium 500 mg Hexal		Cefuroxime axetil		Clomazol	
Calcium carbonate		Celapram		Dermatological	
Calcium carbonate PAI		Celebrex		Genito-Urinary	
Calcium Channel Blockers		Celecoxib		Clomifene citrate	
Calcium Disodium Versenate		Celecoxib Pfizer		Clomipramine hydrochloride	12
Calcium folinate		Celestone Chronodose		Clomipramine Teva	12
Calcium Folinate Ebewe		Cellcept	179	Clonazepam1	
Calcium Folinate Sandoz		Centrally-Acting Agents		Clonidine	
Calcium Folinate Sandoz S29		Cephalexin ABM		Clonidine hydrochloride	
Calcium gluconate		Cetirizine hydrochloride		Clonidine Teva	
Calcium Homeostasis		Cetomacrogol	71	Clopidogrel	3

Clopine	Cosentyx	232	DBL Aminophylline	26/
Clopixol	Cosmegen		DBL Bleomycin Sulfate	
Clotrimazole	Coumadin		DBL Bortezomib	150
Dermatological68	Country Life		DBL Carboplatin	
Genito-Urinary81	Coversyl		DBL Cisplatin	
Clozapine136	COVID-19 vaccine		DBL Dacarbazine	
Clozaril	Creon 10000		DBL Desferrioxamine Mesylate for	
Clustran	Creon 25000		BP	
Co-trimoxazole	Creon Micro		DBL Docetaxel	
Coal tar	Crotamiton		DBL Ergometrine	
Coal tar with allantoin, menthol,	Crystaderm		DBL Gemcitabine	
phenol and sulphur74	Cu 375 Standard		DBL Gentamicin	
Coal tar with salicylic acid and	Curam		DBL Leucovorin Calcium	
	Curam Duo 500/125		DBL Methotrexate Onco-Vial	
sulphur	Cvite		DBL Naloxone Hydrochloride	
Cobalin-H31	Cyclizine hydrochloride		DBL Pethidine Hydrochloride	
Coco-Scalp74			DBL Vincristine Sulfate	
•	Cyclizine lactate		Decozol	
Codeine phosphate	Cyclogyl		Deferasirox	
Extemporaneous	Cyclonex			
Nervous	Cyclopentolate hydrochloride		Deferiprone	
Collabining 26	Cyclophosphamide		Deferoxamine Pfizer S29	
Colchicine 121	Cyclorin		Denosumab	
Colectiferol	Cycloserine		Deolate	
Colestyramine 53	Cyklokapron		Deoxycoformycin	
Colestyramine - Mylan53	Cyproterone acetate	00	Depo-Medrol	
Colgout	Cyproterone acetate with	04	Depo-Provera	
Colistin culphomethate 100	ethinyloestradiol		Depo-Testosterone	
Colistin Link	Cystadane		Deprim	
Collection florible	Cytarabine		Dermol	
Collodion flexible275 Colloidal bismuth subcitrate	Cytotec		Desferrioxamine mesilate	
	Cytoxan	100	Desmopressin	
Colorac8	<del>-</del>	440	Desmopressin acetate	
Coloxyl23	D-Penamine		Desmopressin-PH&T	90
Combigan270	Dabigatran		Detection of Substances in	0.0
Comirnaty Omicron	Dacarbazine		Urine	80
(JN.1)	Dactinomycin [Actinomycin D]		Dexamethasone	0.0
Command all and the latest than 302	Daivobet		Hormone	
Compound electrolytes44	Daivonex		Sensory	
Compound electrolytes with glucose	Daktarin		Dexamethasone phosphate	
[Dextrose]	Dalacin C		Dexamethasone with framycetin a	na
Compound hydroxybenzoate275	Dantrium		gramicidin	267
Comtan	Dantrium S29		Dexamethasone with neomycin	
Concerta	Dantrolene		sulphate and polymyxin B	000
Condoms	Daonil		sulphate	
Condyline	Dapa-Tabs		Dexamfetamine sulfate	
Continuous glucose monitor	Dapsone		Dexcom G6	
(interoperable)21	Daraprim		Dexcom G7	
Continuous glucose monitor	Darunavir		Dexcom ONE+	
(standalone)21	Darunavir Viatris		Dexmethsone	85
Contraceptives - Hormonal79	Dasatinib		Dextrochlorpheniramine	0.5
Contraceptives - Non-hormonal77	Dasatinib-Teva		maleate	
Copaxone	Daunorubicin		Dextrose	
Cordarone-X47	David One Step Cassette Pregn		DHC Continus	
Corticosteroids and Related Agents	Test		Diabetes	
for Systemic Use85	DBL Acetylcysteine		Diabetes Management	
Corticosteroids Topical69	DBL Adrenaline	55	Diacomit	133

# **INDEX: Generic Chemicals and Brands**

Diagnostic Agents317	Doxazosin45	EMB Fatol	. 10
Diamide Relief6	Doxazosin Clinect45	Emend Tri-Pack	. 13
Diamox269	Doxine99	Emicizumab	3
Diasip279	Doxorubicin Ebewe159	EMLA	. 12
Diazepam131, 140	Doxorubicin hydrochloride159	Empagliflozin	1
Diazoxide10	Doxycycline99	Empagliflozin with metformin	
Dibenzyline45	DP Lotion72	hydrochloride	1
Diclofenac Sandoz117	DP Lotn HC70	Emsogen	
Diclofenac sodium117	DP-Captopril45	Emtricitabine	.11
Differin67	Dr Reddy's Omeprazole9	Emtricitabine with tenofovir	
Difflam30	Drofate49	disoproxil	. 11
Diflucan102	Drugs Affecting Bone	Emtriva	.11
Digestives Including Enzymes22	Metabolism 118	Emulsifying ointment	7
Digoxin47	Dual blood glucose and blood ketone	Emulsifying Ointment ADE	
Dihydrocodeine tartrate127	diagnostic test meter14	Enalapril maleate	
Dilantin132	Dulaglutide12	Enbrel	
Dilantin Infatab132	Dulcolax SP Drop24	Endocrine Therapy	
Dilantin Paediatric132	Duocal Super Soluble Powder277	Endoxan	. 15
Diltiazem CD Clinect50	Duolin260	Energivit	. 29
Diltiazem hydrochloride50	Duolin Cipla260	Engerix-B	.30
Dimethicone71–72	Duolin HFA260	Enhertu	. 24
Dimethyl fumarate141	DuoResp Spiromax259	Enlafax XR	.13
Dipentum8	Duride55	Enoxaparin sodium	4
Diphtheria, tetanus and pertussis	Durvalumab245	Enstilar	7
vaccine303	-E-	Ensure	. 28
Diphtheria, tetanus, pertussis and	e-chamber La Grande266	Ensure Plus	
polio vaccine304	e-chamber Mask266	Ensure Plus HN	. 28
Diphtheria, tetanus, pertussis, polio,	e-chamber Turbo266	Ensure Plus HN RTH	. 28
hepatitis B and haemophilus	E-Mycin97	Ensure Two Cal HN RTH	
influenzae type B vaccine304	e5 Pharma10	Entacapone	
Diprosone69	Ear Preparations267	Entecavir	
Diprosone OV69	Ear/Eye Preparations267	Entecavir (Rex)	. 10
Dipyridamole39	Easiphen Liquid292	Entresto 24/26	
Disopyramide phosphate47	Econazole nitrate68	Entresto 49/51	
Disulfiram150	Efavirenz112	Entresto 97/103	
Diuretics51	Efavirenz Milpharm112	Entyvio	
Docetaxel159	Efavirenz with emtricitabine and	Epilim	
Docetaxel Accord159	tenofovir disoproxil113	Epilim Crushable	
Docetaxel Sandoz159	Eformoterol fumarate dihydrate259	Epilim IV	
Docusate sodium23	Eftrenonacog alfa [Recombinant	Epilim S/F Liquid	
Docusate sodium with	factor IX]36	Epilim Syrup	
sennosides23	Efudix76	Epipen	
Dolutegravir114	Egopsoryl TA74	Epipen Jr	
Dolutegravir with lamivudine114	Elaprase26	Epirubicin Ebewe	
Domperidone	Elecare	Epirubicin hydrochloride	
Domperidone Viatris135	Elecare LCP296	Eplerenone	
Donepezil hydrochloride	Electral44	Epoetin alfa	
Dornase alfa	Elelyso29	Epoprostenol	6
Dortimopt269	Elemental 028 Extra282	Eptacog alfa [Recombinant factor	
Dorzolamide with timolol269	Elexacaftor with tezacaftor, ivacaftor	VIIa]	
Dostinex94	and ivacaftor	Erbitux	
Dosulepin [Dothiepin]	Elidel74	Ergometrine maleate	
hydrochloride	Elocon70	Erlotinib	
Dosulepin Viatris129	Elocon Alcohol Free70	Erythrocin IV	
Dothiepin	Eltrombopag36	Erythromycin (as lactobionate)	
Dovato114	Eltroxin88	Erythromycin ethyl succinate	9

Fabricat	000	Facenta	202	fluorantalana nivalata and	
Esbriet		Fasenra		fluocortolone pivalate and	
Escitalopram (Ethica)		Faslodex		cinchocaineFluorometholone	
Escitalopram (Ethics)		Fatty Cream AFT			
Eskazole		Fatty Emulsion Cream (Evara)		Fluorouracil	
Essential Amino Acid Mix		Febuxostat		Fluorouracil Accord	
Essential Ethosuximide		Febuxostat (Teva)		Fluorouracil sodium	
Estraderm MX		FEIBA NF		Fluox	
Estradiol Sandoz		Felo 10 ER	50	Fluoxetine hydrochloride	
Estradiol TDP Mylan	87	Felo 5 ER	50	Flupenthixol decanoate	13
Estradiol Viatris	87	Felodipine	50	Flutamide	17
Estradot	87	Fentanyl	127	Flutamin	17
Estrofem	87	Fentanyl Sandoz	127	Fluticasone	25
Estrogel	87	Ferinject	33	Fluticasone furoate with	
Etanercept		Ferodan		umeclidinium and vilanterol	26
Ethambutol hydrochloride		Ferriprox		Fluticasone furoate with	
Ethics Aspirin		Ferro-F-Tabs		vilanterol	259
Ethics Aspirin EC		Ferro-Liquid		Fluticasone propionate	
Ethics Lisinopril		Ferro-tab		Fluticasone with salmeterol	
	45				
Ethinyloestradiol with	70	Ferrograd		Flynn	
desogestrel	79	Ferrosig		FML	
Ethinyloestradiol with		Ferrous fumarate		Foban	
levonorgestrel	80	Ferrous fumarate with folic acid		Folic acid	
Ethinyloestradiol with		Ferrous sulfate		Folic Acid multichem	
norethisterone		Fexofenadine hydrochloride		Folic Acid Viatris	
Ethosuximide		Fibro-vein	39	Food Thickeners	28
Etopophos		Filgrastim	42	Foods And Supplements For	
Etoposide	159	Finasteride	82	Inherited Metabolic Disease	29
Etoposide phosphate		Fingolimod	141	Fortini	28
Etravirine		Firazyr	257	Fortini Multi Fibre	28
Eumovate	70	Flagyl	106	Fortisip	28
Eurofolic		Flagyl-S		Fortisip Multi Fibre	
Evara		Flamazine		Fosamax	
EVARA White Soft Paraffin		Flecainide acetate		Fosamax Plus	
Everet		Flecainide BNM		Fosfomycin	
Everolimus		Flecainide Controlled Release		Framycetin sulphate	
Evista		Teva	10	Frebini Energy	
Evrysdi		Fleet Phosphate Enema		Frebini Energy Fibre	
Evusheld		Flixonase Hayfever & Allergy		Frebini Original	
Exelon Patch 10		Flixotide		Frebini Original Fibre	
Exelon Patch 5		Flixotide Accuhaler		Freestyle Libre 2	2
Exemestane		Florinef		Freestyle Libre 3 Plus	
Exjade		Fluanxol		Fresubin 2kcal HP	
Extemporaneously Compound	ded	Flucil		Fresubin HP Energy	
Preparations and		Flucloxacillin		Fresubin HP Energy Fibre	28
Galenicals	275	Flucloxacillin-AFT	98	Fresubin Intensive	28
Eye Preparations	267	Flucloxin	98	Fresubin Original	28
Eylea	202	Flucon	268	Fresubin Original Fibre	28
Ezemibe Viatris		Fluconazole	102	Frisium	13 <sup>-</sup>
Ezetimibe		Fludara Oral		Frumil	
Ezetimibe Sandoz		Fludarabine Ebewe		Frusemide	
Ezetimibe with simvastatin		Fludarabine phosphate		Fucicort	
- F -		Fludarabine Sagent	157	Fucidin	
Factor eight inhibitor bypassin	na	Fludrocortisone acetate		Fucithalmic	
fraction		Fluids and Electrolytes		Fucithalmic S29	
Famotidine		Flumetasone pivalate		Fulvestrant	
Famotidine Hovid		Fluocortolone caproate with	201		
ramouume moviu	<del>y</del>	riuoconoione caproate with		Fungilin	ال

# **INDEX: Generic Chemicals and Brands**

Furosemide [Frusemide]	51	Goserelin	93	Humira	19
Furosemide-Baxter	51	Gynaecological Anti-infectives	81	HumiraPen	19
fusidic acid		- H -		Humulin 30/70	1
Dermatological	68, 70	Habitrol	151	Humulin NPH	
Infection		Haemophilus influenzae type B		Humulin R	
Sensory	267	vaccine		Hyaluronic acid	27
- G -		Haldol	138	Hydralazine	5
GA Explore 5	294	Haldol Concentrate	138	Hydralazine hydrochloride	
GA1 Anamix Infant	294	Haldol Decanoas		Hydralyte - Lemonade	4
GA1 Anamix Junior		Haloperidol	136	Hydrocortisone	
Gabapentin	131	Haloperidol decanoate		Dermatological	
Gacet	126–127	Harvoni	110	Hormone	8
Galsulfase	26	Havrix 1440	305	Hydrocortisone acetate	
Galvumet		Havrix Junior	305	Hydrocortisone acetate with	
Galvus	12	Haylor syrup	258	pramoxine hydrochloride	
Gardasil 9	307	HCU Anamix Infant	290	Hydrocortisone and paraffin liquid	i
Gastrodenol	10	HCU Anamix Junior	290	and lanolin	70
Gaviscon Extra Strength	6	HCU Anamix Junior LQ	290	Hydrocortisone butyrate	.70, 7
Gaviscon Infant	6	HCU Explore 5	290	Hydrocortisone with cinchocaine.	8
Gazyva	215	HCU Express 15	290	Hydrocortisone with miconazole	70
Gefitinib	169	HCU Lophlex LQ	290	Hydrocortisone with natamycin ar	nd
GEM Aqueous Cream		healthE Aqueous Cream SLS		neomycin	70
Gemcitabine Ebewe	157	Free	71	Hydrogen peroxide	6
Gemcitabine hydrochloride	157	healthE Calamine Aqueous	69	Hydroxocobalamin	3
Gemtuzumab ozogamicin		healthE Dimethicone 10%	71	Hydroxocobalamin Panpharma	3
Gentamicin Noridem	100	healthE Dimethicone 4% Lotion .	72	hydroxycarbamide	160
Gentamicin sulphate	100	healthE Dimethicone 5%		Hydroxychloroquine sulphate	118
Gilenya	141	healthE Glycerol BP	275	Hydroxyurea	
Ginet	81	healthE Urea Cream	71	[hydroxycarbamide]	160
Glatiramer acetate	141	Healtheries Simple Baking Mix	289	Hygroton	52
Glecaprevir with pibrentasvir	110	Hemastix	83	Hylo-Fresh	27
Glibenclamide	11	Hemlibra	37	Hymenoptera25	57-258
Gliclazide	12	Heparin sodium	42	Hyoscine butylbromide	
Glipizide	12	Heparin Sodium Panpharma	42	Hyoscine Butylbromide	
Glizide	12	Heparinised saline	42	(Adiramedica)	8
Glucagen Hypokit	10	Heparon Junior	280	Hyoscine hydrobromide	13
Glucagon hydrochloride	10	Hepatitis A vaccine	305	Hyperuricaemia and Antigout	
Glucerna Select		Hepatitis B recombinant		Hypromellose	270
Glucose [Dextrose]	43	vaccine	306	Hypromellose with dextran	270
Gluten Free Foods	289	Herzuma	238	-1-	
Glycerin with sodium sacchar	in2 <mark>75</mark>	Hikma	47	Ibiamox	98
Glycerin with sucrose	275	Hiprex	116	Ibrance	172
Glycerol		Histaclear	258	Ibrutinib	160
Alimentary	<mark>24</mark>	Holoxan	155	Ibuprofen	117
Extemporaneous	275	Horleys Bread Mix	289	Ibuprofen SR BNM	117
Glyceryl trinitrate		Horleys Flour	289	Icatibant	25
Alimentary	8	Hormone Replacement Therapy	-	Idarubicin hydrochloride	160
Cardiovascular	55	Systemic	87	Idursulfase	20
Glycopyrronium	261	HPV	307	Ifosfamide	15
Glycopyrronium bromide		Humalog		llevro	
Glycopyrronium with		Humalog Mix 25	11	lloprost	
indacaterol		Humalog Mix 50		Imatinib mesilate	
Glycosade	294	Human papillomavirus (6, 11, 16		Imatinib-Rex	
Glytactin Bettermilk		31, 33, 45, 52 and 58) vaccine		Imbruvica	
Gold Knight		[HPV]		Imfinzi	24
Gold Knight XL		Humatin	101	Imipramine Crescent	129

Imipramine hydrochloride12		16	Jevity Plus RTH	286
Imiquimod7			Jevity RTH	
Immune Modulators11	4 attached needle	16	Jinarc	
Immunisation - Flu27	2 Intelence	112	Juno Pemetrexed	158
Immunisation Flu and Shingles27	2 Interferon beta-1-alpha	141	- K -	
Immunisation Other27	2 Interferon beta-1-beta	141	Kadcyla	
Immunosuppressants17	9 Intra-uterine device	79	Kalydeco	265
Incruse Ellipta26	1 Invega Sustenna	139	Kemadrin	123
Indacaterol25		139	Kenacomb	267
Indapamide5			Kenacort-A 10	
Infanrix IPV30			Kenacort-A 40	86
Infanrix-hexa30	4 Ipca-Ciprofloxacin	99	Kenalog in Orabase	
Infant Formulae29		149	Ketocal 3:1	300
Infatrini29			KetoCal 4:1	300
Infliximab20	6 IPCA-Frusemide	51	Ketoconazole	
Influenza vaccine30	8 Ipca-Hydroxychloroquine	118	Dermatological	75
Influvac Tetra	IPCA-Metoprolol	49	Infection	103
(2024 formulation) 30	8 IPCA-Propranolol	49	Ketogenic Diet	300
Inhaled Corticosteroids25	8 IPOL	314	Ketoprofen	117
Inhaled Long-acting	Ipratropium bromide2	60, 266	KetoSens	14
Beta-adrenoceptor Agonists 25	9 Ipratropium IVAX	260	Keytruda	247
Inresa3			Kindergen	280
Inspra5	1 Irinotecan Actavis 100	157	Kisqali	173
Instillagel Lido12		157	Klacid	
Insulin aspart1	1 Irinotecan-Rex	157	Alimentary	9
Insulin aspart with insulin aspart	Iron (as ferric carboxymaltose)	33	Infection	96
protamine1	0 Iron polymaltose	34	Kliogest	8
Insulin glargine1			Kliovance	8
Insulin glulisine1		114	Kogenate FS	38
Insulin isophane1	1 Ismo 20	55	Konakion MM	
Insulin isophane with insulin	Ismo 40 Retard	55	Konsyl-D	23
neutral1	1 Isoleucine50	295	Kuvan	
Insulin lispro1	1 Isoniazid	107	-L-	
Insulin lispro with insulin lispro	Isoniazid Teva		Labetalol	49
protamine1	1 Isoniazid with rifampicin	107	Lacosamide	131
Insulin neutral1	·		Lactulose	24
Insulin pen needles1			Laevolac	
Insulin pump cartridge1			Lagevrio	111
Insulin pump infusion set (steel	Isopto Carpine		Lamictal	
cannula)1			Lamivudine1	
Insulin pump infusion set (steel	Isosource Standard	286	Lamivudine Viatris	113
cannula, straight insertion) 1	8 Isotretinoin	67	Lamivudine/Zidovudine Viatris	113
Insulin pump infusion set (teflon	Ispaghula (psyllium) husk	23	Lamotrigine	132
cannula)1			Lamprene	
Insulin pump infusion set (teflon	Itraconazole		Lanoxin	
cannula, angle insertion with	Itraconazole Kent		Lanoxin Paediatric Elixir	
insertion device)2	0 Itrazole	103	Lanoxin PG	47
Insulin pump infusion set (teflon	Ivacaftor		Lanoxin S29	47
cannula, flexible insertion with	Ivermectin		Lansoprazole	
insertion device)2			Lantus	
Insulin pump infusion set (teflon	Jadelle	80	Lantus SoloStar	
cannula, straight insertion with	Jakavi		Lanvis	
insertion device)2			Lanzol Relief	
Insulin pump infusion set (teflon	Jardiance		Largactil	
cannula, variable insertion) 2			Laronidase	
Insulin pump reservoir2			Lasix	
1 1	,			

Latanoprost	269	Rectal Disorders	8	MCT oil (Nutricia)	278
Latanoprost with timolol	270	Locasol	295	Measles, mumps and rubella	
Lax-Suppositories	<mark>24</mark>	Locoid	70, 75	vaccine	310
Lax-suppositories Glycerol	<mark>24</mark>	Locoid Crelo	70	Mebendazole	
Laxatives		Locoid Lipocream	70	Mebeverine hydrochloride	8
Laxsol	23	Locorten-Vioform		Medac	15
Ledipasvir with sofosbuvir	110	Lodoxamide	269	Medrol	8
Leflunomide	118	Logem	132	Medroxyprogesterone acetate	
Lenalidomide (Revlimid)	160	Lomide	269	Genito-Urinary	80
Lenalidomide (Viatris)	161	Lomustine	155	Hormone	8
Lenalidomide Viatris		Loniten	<u>55</u>	Mefenamic acid	117
Lenvatinib	169	Loperamide hydrochloride		Megval	15
Lenvima	169	Lopinavir with ritonavir	113	Melatonin	143
Letrole		Lopinavir/Ritonavir Mylan		Melpha	15
Letrozole	179	Loprofin		Melphalan	15
Leucine100	295	Loprofin Mix	293	Meningococcal (groups A, C, Y a	nd
Leukeran FC	154	Lorafix		W-135) conjugate vaccine	31 <sup>-</sup>
Leukotriene Receptor		Loratadine	258	Meningococcal B multicomponen	
Antagonists	264	Lorazepam	140	vaccine	312
Leuprorelin		Lorstat	53	MenQuadfi	31°
Leustatin	157	Losartan Actavis	46	Menthol	69
Levetiracetam		Losartan potassium	46	Mepolizumab	
Levetiracetam-AFT	132	Losartan potassium with		Mercaptopurine	157
Levocabastine	268	hydrochlorothiazide	46	Mercilon 28	79
Levocarnitine	27	Lovir		Mesalazine	
Levodopa with benserazide	123	Loxamine	130	Mesna	162
Levodopa with carbidopa	123	Lucrin Depot 1-month	93	Mestinon	117
Levomepromazine		Lucrin Depot 3-month		Metabolic Disorder Agents	2
Levomepromazine		LumaCina		Metabolics	
hydrochloride	136	Lumigan	269	Metformin hydrochloride	12
Levonorgestrel		Lyllana		Metformin Viatris	12
Genito-Urinary	80-81	Lynparza		Methadone BNM	
Hormone		Lyrica		Methadone hydrochloride	12
Levonorgestrel BNM		- M -		Methenamine (hexamine)	
Levothyroxine		m-Eslon	128	hippurate	116
Lidocaine [Lignocaine]		Mabthera	219	Methopt	
Lidocaine [Lignocaine]		Macrobid	116	Methotrexate	
hydrochloride	125	Macrogol 3350 with potassiu	m	Methotrexate DBL Onco-Vial	158
Lidocaine [Lignocaine] with		chloride, sodium bicarbon		Methotrexate DBL S29	
prilocaine	125	sodium chloride		Methotrexate Ebewe	158
Lidocaine-Baxter		Madopar 125		Methotrexate Sandoz	
Life Extension		Madopar 250		Methyl hydroxybenzoate	
Lignocaine		Madopar 62.5		Methylcellulose	
Linezolid		Madopar HBS		Methylcellulose with glycerin and	
Lioresal Intrathecal	122	Madopar Rapid		sodium saccharin	
Lipid-Modifying Agents		Magnesium hydroxide		Methylcellulose with glycerin and	
Liquigen		Magnesium sulphate		sucrose	
Liraglutide		Mantoux		Methyldopa	_
Lisdexamfetamine dimesilate		MAR-Midodrine		Methyldopa Viatris	
Lisinopril		Marevan		Methylnaltrexone bromide	2
Lithium carbonate		Marine Blue Lotion SPF 50+		Methylphenidate ER - Teva	
Livostin		Martindale Pharma		Methylphenidate hydrochloride	
LMX4		Mask for spacer device		Methylphenidate hydrochloride	
Lo-Oralcon 20 ED		Maviret		extended-release	148
Locacorten-Viaform ED's		Maxidex		Methylprednisolone	
Local preparations for Anal and		Maxitrol		Methylprednisolone (as sodium	
				• •	

succinate)	85	MiniMed Silhouette MMT-381A	19	MSUD Lophlex LQ 20	291
Methylprednisolone aceponate	70	MiniMed Sure-T MMT-864A	18	MSUD Maxamum	291
Methylprednisolone acetate	86	MiniMed Sure-T MMT-866A	18	Mucolytics	264
Methylxanthines		MiniMed Sure-T MMT-874A	18	Mucosoothe	125
Metoclopramide Actavis 10		MiniMed Sure-T MMT-876A	18	Multiple Sclerosis Treatments	
Metoclopramide hydrochloride	135	Minims Cyclopentolate	270	Multivitamin renal	
Metolazone	52	Minims Pilocarpine	270	Multivitamins	
Metopirone	94	Minims Prednisolone	269	Mupirocin	67
Metoprolol IV Mylan	49	Minipress	45	Muscle Relaxants	122
Metoprolol IV Viatris		Minirin	93	Mvite	32
Metoprolol succinate	49	Minirin Melt		Myambutol	107
Metoprolol tartrate	49	Mino-tabs	99	Mycobutin	
Metrogyl		Minocycline hydrochloride	99	MycoNail	68
Metronidamed		Minomycin		Mycophenolate mofetil	
Metronidazole	106	Minor Skin Infections		Mydriacyl	270
Metyrapone	94	Minoxidil	55	Mylan (24 hr release)	50
Mexiletine hydrochloride	48	Minoxidil Roma	55	Mylan Clomiphen	
Miacalcic	84	Mirena	88	Mylan Italy (24 hr release)	50
Micolette		Miro-Amoxicillin	98	Myleran	154
Miconazole	30	Mirtazapine	130	mylife Inset soft	
Miconazole nitrate		Misoprostol		mylife Orbit micro	
Dermatological	69	Mitomycin (Fresenius Kabi)		mylife YpsoPump Reservoir	
Genito-Urinary		Mitomycin (Sagent)	162	mylife YpsoPump with CamAPS	
Micreme		Mitomycin C		FX	16
Micreme H	70	Mitozantrone		Myloc CR	49
Microgynon 30	80	Mitozantrone Ebewe		Mylotarg	
Microlut		Mixtard 30		Myometrial and Vaginal Hormone	
Midazolam	144	MMA/PA Anamix Infant	294	Preparations	81
Midazolam Viatris	144	MMA/PA Anamix Junior	294	Myozyme	24
Midazolam Viatris Midazolam-Baxter				Myozyme N -	24
	144	MMA/PA Explore 5	294		
Midazolam-Baxter	144 144		294 294	- N - Nadolol	49
Midazolam-Baxter Midazolam-Pfizer	144 144 48	MMA/PA Explore 5 MMA/PA Express 15	294 294 130	- N - Nadolol Nadolol BNM	49
Midazolam-BaxterMidazolam-PfizerMidodrineMidostaurin	144 144 48 170	MMA/PA Explore 5 MMA/PA Express 15 Moclobemide	294 294 130 148	- N - Nadolol Nadolol BNM Naglazyme	49 49
Midazolam-Baxter	144 144 48 170 83	MMA/PA Explore 5MMA/PA Express 15MoclobemideModafinilModafinil Max Health	294 130 148	- N - Nadolol Nadolol BNM Naglazyme Naloxone hydrochloride	49 49 26
Midazolam-Baxter	144 48 170 83 83	MMA/PA Explore 5 MMA/PA Express 15 Moclobemide Modafinil	294 130 148 148	- N - Nadolol Nadolol BNM Naglazyme	49 26 272 150
Midazolam-Baxter	144 144 170 83 83 132	MMA/PA Explore 5	294 130 148 148 148	- N - Nadolol Nadolol BNM Naglazyme Naloxone hydrochloride Naltraccord Naltrexone AOP	49 49 27 272 150
Midazolam-Baxter	144 144 170 83 83 132 32	MMA/PA Explore 5 MMA/PA Express 15 Moclobemide Modafinil Modafinil Max Health Modavigil Moduretic Molaxole	294 130 148 148 148 148 52	- N - Nadolol Nadolol BNM Naglazyme Naloxone hydrochloride Naltraccord	49 26 272 150 150
Midazolam-Baxter	144 144 48 170 83 83 132 32 32	MMA/PA Explore 5	294 130 148 148 148 52 24	- N - Nadolol	49 26 272 150 150
Midazolam-Baxter	144 144 48 170 83 132 32 32 266 266	MMA/PA Explore 5 MMA/PA Express 15 Moclobemide Modafinil Modafinil Max Health Modavigil Moduretic Molaxole Molnupiravir	294 294 130 148 148 148 52 24 111	- N - Nadolol	49 26 272 150 150 150
Midazolam-Baxter	144 144 48 170 83 132 32 32 266 266	MMA/PA Explore 5 MMA/PA Express 15 Moclobemide Modafinil Modafinil Max Health Modavigil Moduretic Molaxole Molnupiravir Moments Mometasone furoate	294 294 130 148 148 52 24 111 78	- N - Nadolol	49 26 150 150 150 151
Midazolam-Baxter	144 144 48 170 83 83 32 32 266 266 12	MMA/PA Explore 5 MMA/PA Express 15 Moclobemide Modafinil Modafinil Max Health Modavigil Moduretic Molaxole Molnupiravir Moments Mometasone furoate Monogen		- N - Nadolol	49 26 150 150 150 151
Midazolam-Baxter	1444817083831323226626612	MMA/PA Explore 5 MMA/PA Express 15 Moclobemide Modafinil Modafinil Max Health Modavigil Moduretic Molaxole Molnupiravir Mometasone furoate Monogen Montelukast		Nadolol	49 272 150 150 150 177 117
Midazolam-Baxter	144481708383132322662661211	MMA/PA Explore 5 MMA/PA Express 15 Moclobemide Modafinil Modafinil Max Health Modavigil Moduretic Molaxole Molnupiravir Moments Mometasone furoate Monogen Montelukast Montelukast Montelukast Viatris		- N - Nadolol	492615015015017117117
Midazolam-Baxter	1444817083831322662661219	MMA/PA Explore 5 MMA/PA Express 15 Moclobemide Modafinil Modafinil Max Health Modavigil Moduretic Molaxole Molnupiravir Moments Mometasone furoate Monogen Montelukast Montelukast Viatris Moroctocog alfa [Recombinant		- N - Nadolol	49 26 150 150 150 150 117 117 117
Midazolam-Baxter	1444817083833226626612211919	MMA/PA Explore 5 MMA/PA Express 15 Moclobemide Modafinil Modafinil Max Health Modavigil Moduretic Molaxole Molnupiravir Moments Mometasone furoate Monogen Montelukast Montelukast Moroctocog alfa [Recombinant VIII]		- N - Nadolol	
Midazolam-Baxter	1444817083831322662661221191919	MMA/PA Explore 5 MMA/PA Express 15 Moclobemide Modafinil Modafinil Max Health Modavigil Moduretic Molaxole Molnupiravir Moments Mometasone furoate Monogen Montelukast Montelukast Moroctocog alfa [Recombinant VIII] Morphine hydrochloride		- N - Nadolol	
Midazolam-Baxter	144481708383132266266121919191919	MMA/PA Explore 5 MMA/PA Express 15 Moclobemide Modafinil Modafinil Max Health Modavigil Moduretic Molaxole Molnupiravir Moments Mometasone furoate Monogen Montelukast Montelukast Viatris Moroctocog alfa [Recombinant VIII] Morphine hydrochloride Morphine sulphate		- N - Nadolol	
Midazolam-Baxter	144144481708383132266266121919191919	MMA/PA Explore 5 MMA/PA Express 15 Moclobemide Modafinil Modafinil Max Health Modavigil Moduretic Molaxole Molnupiravir Moments Mometasone furoate Mondelukast Montelukast Viatris Moroctocog alfa [Recombinant VIII] Morphine hydrochloride Morphine sulphate Motetis	29429413014814852241117870279264264 factor38127128	- N - Nadolol	
Midazolam-Baxter	1441444817083322662661219191919191919	MMA/PA Explore 5 MMA/PA Express 15 Moclobemide Modafinil Modafinil Max Health Modavigil Moduretic Molaxole Molnupiravir Moments Mometasone furoate Monogen Montelukast Viatris Moroctocog alfa [Recombinant VIII] Morphine hydrochloride Motetis Motetis Motetis Motetis Motetis Motetis Motetis Mouth and Throat	29429413014814852241117870279264264 factor38127128	- N - Nadolol	
Midazolam-Baxter	144144481708332266266122119191919191919	MMA/PA Explore 5 MMA/PA Express 15 Moclobemide Modafinil Modafinil Max Health Modavigil Moduretic Molaxole Molnupiravir Moments Mometasone furoate Monogen Montelukast Viatris Moroctocog alfa [Recombinant VIII] Morphine hydrochloride Morphine sulphate Motetis Mouth and Throat Movapo		- N - Nadolol	
Midazolam-Baxter	1444817083833226626612191919191919191919	MMA/PA Explore 5 MMA/PA Express 15 Moclobemide Modafinil Modafinil Max Health Modavigil Moduretic Molaxole Molnupiravir Mometasone furoate Monogen Montelukast Viatris Moroctocog alfa [Recombinant VIII] Morphine hydrochloride Morphine sulphate Motetis Mouth and Throat Movapo Moxifloxacin		- N - Nadolol	
Midazolam-Baxter	144481708383132266266121919191919191919191919	MMA/PA Explore 5 MMA/PA Express 15 Moclobemide Modafinil Modafinil Max Health Modavigil Moduretic Molaxole Molnupiravir Mometasone furoate Monogen Montelukast Viatris Moroctocog alfa [Recombinant VIII] Morphine hydrochloride Morphine sulphate Motetis Mouth and Throat Movapo Moxifloxacin MSUD Anamix Infant		- N - Nadolol	
Midazolam-Baxter	14448170838313226626612191919191919191919191919	MMA/PA Explore 5 MMA/PA Express 15 Moclobemide Modafinil Modafinil Max Health Modavigil Moduretic Molaxole Molnupiravir Moments Mometasone furoate Monogen Montelukast Montelukast Viatris Moroctocog alfa [Recombinant VIII] Morphine hydrochloride Morphine sulphate Motetis Mouth and Throat Movapo Moxifloxacin MSUD Anamix Junior		- N - Nadolol	
Midazolam-Baxter	1444817083833226612211919191919191919191919191919	MMA/PA Explore 5 MMA/PA Express 15 Moclobemide Modafinil Modafinil Max Health Modavigil Moduretic Molaxole Molnupiravir Moments Mometasone furoate Monogen Montelukast Montelukast Viatris Moroctocog alfa [Recombinant VIII] Morphine hydrochloride Morphine sulphate Motetis Mouth and Throat Movapo Moxifloxacin MSUD Anamix Junior LQ		- N - Nadolol	
Midazolam-Baxter	14448170838332266122119191919191919191919191919191919	MMA/PA Explore 5 MMA/PA Express 15 Moclobemide Modafinil Modafinil Max Health Modavigil Moduretic Molaxole Molnupiravir Moments Mometasone furoate Monogen Montelukast Montelukast Viatris Moroctocog alfa [Recombinant VIII] Morphine hydrochloride Morphine sulphate Motetis Mouth and Throat Movapo Moxifloxacin MSUD Anamix Junior		- N - Nadolol	

Neoral	252	Novatretin	73	Hormone	8
Neostigmine metilsulfate	117	Novitium Sugar Free	27	Oestrogens	8
Nepafenac	269	NovoMix 30 FlexPen	10	Ofev	26
Nepro HP (strawberry)		NovoRapid	11	Oil in water emulsion	<mark>7</mark>
Nepro HP (vanilla)	282	NovoRapid FlexPen	11	Olanzapine	137-13
Neulactil		NovoRapid Penfill		Olaparib	
NeuroTabs	33	NovoSeven RT		Olbetam	
Nevirapine	112	Nozinan	136	Olopatadine	27
Nevirapine Viatris	112	Nozinan (Swiss)	136	Olopatadine Teva	
Nicorandil		Nozinan S29	136	Olsalazine	
Nicotine	151	Nucala	214	Omalizumab	21
Nifedipine	50	Nuelin	264	Omeprazole	
Nifuran	116	Nuelin-SR	264	Omeprazole actavis 10	
Nilotinib	170	Nupentin	131	Omeprazole actavis 20	
Nilstat		Nusinersen		Omeprazole actavis 40	
Alimentary	31	Nutilis	289	Omeprazole Teva	
Genito-Urinary		Nutren Diabetes	279	Omnitrope	8
Infection	103	Nutrient Modules	276	Omnitrope S29	8
Nimenrix	311	Nutrini Energy Multi Fibre	281	Onbrez Breezhaler	
Nintedanib		Nutrini Energy RTH		Oncaspar LYO	16
Nipent	165	Nutrini Low Energy Multi Fibre		OncoTICE	18
Niraparib		Nutrini Peptisorb		Ondansetron	
Nirmatrelvir with ritonavir		Nutrini Peptisorb Energy		One-Alpha	3
Nitrates		Nutrini RTH		One-Alpha S29	
Nitroderm TTS		Nutrison 800 Complete Multi		Opdivo	
Nitrofurantoin	116	Fibre	286	Ora-Blend	
Nitrolingual Pump Spray		Nutrison Advanced Peptisorb	283	Ora-Blend SF	27
Nivestim		Nutrison Concentrated		Ora-Plus	27
Nivolumab	246	Nutrison Energy	286	Ora-Sweet	
Nodia	6	Nutrison Energy Multi Fibre		Ora-Sweet SF	
Noflam 250	117	Nutrison Multi Fibre		Orabase	3
Noflam 500	117	Nutrison RTH		Oral and Enteral Feeds	
Non-Steroidal Anti-Inflammatory		Nyefax Retard		Oralcon 30 ED	
Drugs	117	Nystatin		Oramorph	
Nonacog gamma, [Recombinant		Alimentary	31	Oramorph CDC S29	
Factor IX]	38	Genito-Urinary		Oratane	
Norethinderone - CDC		Infection		Orgran	
Norethisterone		NZB Low Gluten Bread Mix	289	Ornidazole	
Genito-Urinary	81	-0-		Orphenadrine citrate	
Hormone		Obinutuzumab	215	Ortho-tolidine	
Norflex	122	Obstetric Preparations		Oruvail SR	
Norfloxacin	116	Ocicure		Osimertinib	17
Noriday	81	Ocrelizumab	142	Osmolite RTH	28
Noriday 28		Ocrevus		Other Endocrine Agents	
Norimin		Octocog alfa [Recombinant factor	or	Other Oestrogen Preparations	
Normison		VIII] (Advate)		Other Progestogen	
Norpress		Octocog alfa [Recombinant factor		Preparations	8
Nortriptyline hydrochloride		VIII] (Kogenate FS)		Other Skin Preparations	
Norvir		Octreotide		Otodex	26
Noumed Dexamfetamine	145	Octreotide GH	177	Ovestin	
Noumed Isoniazid		Octreotide long-acting		Genito-Urinary	8
Noumed Paracetamol		Oestradiol		Hormone	
Noumed Pethidine		Oestradiol valerate		Oxaliplatin	
Noumed Phenobarbitone		Oestradiol with norethisterone		Oxaliplatin Accord	
Novadoz		Oestriol		Oxaliplatin Actavis 100	
NovaSource Renal		Genito-Urinary	81	Oxaliplatin Ebewe	
		~~ ~			

Oxis Turbuhaler	259	Pegaspargase164		
Oxpentifylline	<u>55</u>	Pegasys114		7
Oxybutynin		Pegfilgrastim43		
Oxycodone Amneal	128	Pegylated interferon alfa-2a114	Pirfenidone	26
Oxycodone hydrochloride		Pembrolizumab247	Pizotifen	13
Oxycodone Lucis S29	128	Pemetrexed 158		29
Oxycodone Sandoz		Pemetrexed-AFT158	PKU Anamix Junior	29
Oxycodone Sandoz S29	128	Penicillamine118	PKU Anamix Junior Chocolate	29
OxyContin	128	Penicillin G98	PKU Anamix Junior LQ	29
OxyNorm	128	PenMix 3011	PKU Anamix Junior Orange	29
Oxytocin	82	PenMix 5011	PKU Anamix Junior Vanilla	29
Oxytocin BNM		Pentasa7	PKU Build 10	29
Oxytocin Panpharma	82	Pentostatin [Deoxycoformycin]165	PKU Build 20 Chocolate	29
Oxytocin with ergometrine		Pentoxifylline [Oxpentifylline]55	PKU Build 20 Raspberry	
maleate	82	Peptamen Junior281	Lemonade	29
Ozurdex	268	Pepti-Junior298	PKU Build 20 Smooth	29
- P -		Perhexiline maleate50	PKU Build 20 Vanilla	29
Pacifen	122	Pericyazine137	PKU Explore 10	29
Pacimol	126	Perindopril46	PKU Explore 5	
Paclitaxel	164	Periset135		29
Paclitaxel Actavis	164	Periset ODT135	PKU First Spoon	29
Paclitaxel Ebewe	164	Perjeta218	PKU Glytactin RTD 15	29
Padagis		Permethrin73		29
Paediatric Seravit		Perrigo76	PKU GMPro LQ	
Palbociclib	172	Pertuzumab218		29
Paliperidone	139	Peteha 107	PKU GMPro Ultra Lemonade	29
Paliperidone palmitate	139	Pethidine hydrochloride 129	PKU GMPro Ultra Vanilla	29
Palivizumab		Pevaryl68		
Pamidronate disodium	119	Pexsig50		
Pamisol	119	Pfizer Exemestane 179		
Pamol	126	Pfizer S29157		
Pancreatic enzyme	<mark>22</mark>	Pharmacy Services272		
Pantoprazole		Pharmascience260		29
Panzop Relief		Pheburane28		
Papaverine hydrochloride		Phenasen 159		
Para-amino salicylic acid		Phenergan Elixir258		
Paracetamol		Phenobarbitone		
Paracetamol (Ethics)		Phenobarbitone sodium	PKU Start	
Paracetamol + Codeine		Extemporaneous275		
(Relieve)	128	Nervous144	•	
Paracetamol with codeine		Phenoxybenzamine	Pneumococcal (PCV13) conjugate	
Paraffin		hydrochloride45		
Paraffin liquid with wool fat		Phenoxymethylpenicillin (Penicillin	Pneumococcal (PPV23)	
Parasiticidal Preparations		V)99		314
Parnate		Phenylalanine50295		
Paromomycin	101	Phenytoin sodium131–132		
Paroxetine		Phillips Milk of Magnesia34		
Paser		Phlexy 10292	<b>.</b>	
Paxam		Phosphate Phebra44		
Paxlovid		Phosphorus44		
Pazopanib		Phytomenadione39		
Pazopanib Teva		Pilocarpine hydrochloride270	Poly-Visc	27
Peak flow meter	266	Pilocarpine nitrate270	Polycal	27
Pediasure		Pimafucort70		
Pediasure Plus		Pimecrolimus		27
Pediasure RTH		Pine tar with trolamine laurilsulfate	Pomalidomide	

Pomolide	165	Provera	88	Rifabutin	10
Ponstan	117	Provera HD	88	Rifadin	10
Posaconazole	103	Psoriasis and Eczema		Rifadin Sanofi	10
Posaconazole Juno	103	Preparations	<del>73</del>	Rifampicin	10
Potassium chloride	43-44	PTU		Rifaximin	
Potassium citrate	83	Pulmicort Turbuhaler	259	Rifinah	10
Potassium iodate	33	Pulmozyme	264	Rilutek	
Povidone iodine	72	Puri-nethol		Riluzole	12
Pradaxa	42	Puritan's Pride Vitamin		RINVOQ	
Pramipexole hydrochloride		B-2 100 mg	27	Riodine	
Pravastatin		Pyrazinamide		Risdiplam	
Praziguantel		Pyridostigmine bromide		Risedronate Sandoz	
Prazosin		Pyridoxine hydrochloride		Risedronate sodium	
Prazosin Mylan		Pyridoxine multichem		Risperdal	
Pred Forte		Pyrimethamine		Risperdal Consta	
Prednisolone		Pytazen SR		Risperidone	
Prednisolone acetate		- Q -		Risperidone (Teva)	
Prednisolone sodium	200	Quantalan sugar free	53	Risperidone Sandoz	
phosphate	260	Quetapel		Risperon	
Prednisolone-AFT		Quetiapine		Ritalin	14
Prednisone		Quinapril		Ritalin LA	
Prednisone Clinect		Qvar		Ritonavir	
Pregabalin		- R -	200	Rituximab (Mabthera)	
		RA-Morph	107	Rituximab (Riximyo)	
Pregabalin Pfizer				Rivaroxaban	
Pregnancy Tests - hCG Urine		Ralicrom Raloxifene hydrochloride		Rivastigmine	
Premarin					
		Raltegravir potassium		Rivastigmine Patch BNM 10	14
Priadel		Ramipex		Rivastigmine Patch BNM 5	
Primaquine		Ramipril		Rivotril	
Primidone		Ranbaxy-Cefaclor		Riximyo	
Primidone Clinect		Rapamune	253	RIXUBIS	
Primolut N		Rasagiline		Rizamelt	
Priorix		Reandron 1000		Rizatriptan	
Probenecid		Recombinant factor IX		Robinul	
Probenecid-AFT		Recombinant factor VIIa		Ronapreve	
Procarbazine hydrochloride		Recombinant factor VIII		Ropin	
Prochlorperazine	135	Rectogesic		Ropinirole hydrochloride	12
Prochlorperazine Brown &		Redipred		Rosuvastatin	5
Burk		Relieve		Rosuvastatin Viatris	
Proctofoam		Relistor		Rotarix	
Proctosedyl		Remicade		Rotavirus oral vaccine	
Procyclidine hydrochloride		Renilon 7.5		Roxane-Propranolol	4
Progesterone		Resonium-A		Roxithromycin	
Proglicem		Resource Beneprotein		Rubifen	
Progynova		Respiratory Devices		Rubifen SR	
Prolia		Respiratory Stimulants	266	Rugby Capsaicin Topical Crean	n
Promethazine hydrochloride		Retinol palmitate	271	Musculoskeletal	
Propafenone hydrochloride	48	ReTrieve	67	Nervous	12
Propranolol	49	Retrovir	113	Rurioctocog alfa pegol [Recomb	oinant
Propylene glycol	275	Revia	150	factor VIII]	
Propylthiouracil		Revlimid	160	Ruxolitinib	17
Prostacur		Revolade		Rydapt	17
Protaphane	11	Ribociclib	173	Rythmodan	4
Protaphane Penfill	11	Riboflavin		Rythmodan - Cheplafarm	
Protifar		Ribomustin	153	Rytmonorm	
Protionamide	107	Ricit	82	´ - S -	

Sabril	133	Sodium bicarbonate		Strides Shasun	10
Sacubitril with valsartan		Blood	43–44	Stromectol	
SalAir		Extemporaneous		Sucralfate	
Salazopyrin		Sodium calcium edetate		Sulfadiazin-Heyl	
Salazopyrin EN		Sodium chloride		Sulfadiazine Silver	
Salbutamol		Blood	43	Sulfadiazine sodium	
Salbutamol with ipratropium	200	Respiratory		Sulfasalazine	
bromide	260	Sodium citrate with sodium laury		Sulphur	
Salicylic acid		sulphoacetate		Sulprix	
Salmeterol		Sodium citro-tartrate		Sumagran	
Sandomigran		Sodium cromoglicate		Sumatriptan	
Sandostatin LAR		Alimentary	Q	Sunitinib	
Sanofi Primaguine				Sunitinib Pfizer	
Sapropterin dihydrochloride		SensorySodium Fusidate [fusidic acid]	209	Sunscreens	
			60		
Scalp Preparations		Dermatological		Sunscreens, proprietary	
Scopolamine - Mylan	130	Infection		Survimed OPD	
Scopolamine - Mylan S29		Sensory		Sustagen Hospital Formula	201
Sebizole		Sodium hyaluronate [Hyaluronic		Sustagen Hospital Formula	00
Secukinumab		acid]		Active	
Sedatives and Hypnotics		Sodium phenylbutyrate		Sustanon Ampoules	
Seebri Breezhaler		Sodium picosulfate		Sylvant	
Senna		Sodium polystyrene sulphonate		Symbicort Turbuhaler 100/6	
Senokot		Sodium tetradecyl sulphate		Symbicort Turbuhaler 200/6	
SensoCard		Sodium valproate		Symbicort Turbuhaler 400/12	
Serc		Sofradex		Symmetrel	12
Serenace		Soframycin		Sympathomimetics	
Seretide		Solgar2		Synacthen	
Seretide Accuhaler		Solifenacin succinate	83	Synacthen Depot	80
Serevent		Solifenacin succinate Max		Synacthene Retard	
Serevent Accuhaler		Health		Synagis	
Sertraline		Solifenacin Viatris		Synthroid	
Setrona		Solu-Cortef		Syntometrine	
Sevredol	128	Solu-Medrol		Syrup (pharmaceutical grade)	
Sex Hormones Non		Solu-Medrol-Act-O-Vial		Systane Unit Dose	27
Contraceptive		Somatropin (Omnitrope)		-T-	
Shingles vaccine		Sotalol		Tacrolimus	
Shingrix	317	Spacer device		Dermatological	7
SII-Onco-BCG	186	Span-K		Oncology	
Sildenafil	60	Spazmol	8	Tacrolimus Sandoz	
Siltuximab		Spinal Muscular Atrophy	144	Tagrisso	17
Simvastatin		Spinraza	144	Taliglucerase alfa	
Simvastatin Mylan	54	Spiolto Respimat	261	Tambocor	
Simvastatin Viatris		Spiractin	51	Tambocor German	
Sinemet	123	Spiriva	261	Tamoxifen citrate	17
Sinemet CR	123	Spiriva Respimat	261	Tamoxifen Sandoz	17
Sintetica Baclofen Intrathecal	122	Spironolactone	51	Tamsulosin hydrochloride	8
Sirolimus		Sporanox		Tamsulosin-Rex	8
Sirturo	106	Sprycel	168	Tandem Cartridge	1
Siterone	86	Stelara		Tandem t:slim X2 with Basal-IQ	
Slow-Lopresor	49	Stemetil	135	Tandem t:slim X2 with	
Smith BioMed Rapid Pregnand	у	Steril-Gene		Control-IQ	
Test		SteroClear	265	Tap water	27
Sodibic	44	Stesolid		Taro	10
Sodium acid phosphate	24	Stimulants/ADHD Treatments	145	Tasigna	
Sodium alginate	6	Stiripentol	133	Tasmar	
Sodium benzoate		Stomahesive	30	Taurine	

TCu 380 Plus Normal	79	Sensory	267	TruSteel	18
Tecentriq	244	Tobramycin (Viatris)	102	Tryzan	46
Tecfidera		Tobramycin BNM	102	Tuberculin PPD [Mantoux] test	317
Tegretol	131	Tobrex	267	Tubersol	
Tegretol AU	131	Tocilizumab	235	Two Cal HN	288
Tegretol CR		Tofranil	129	TYR Anamix Infant	294
Telfast		Tolcapone	123	TYR Anamix Junior	294
Temaccord	165	Tolvaptan	52	TYR Anamix Junior LQ	294
Temazepam	144	Topamax	133	TYR Explore 5	294
Temozolomide		Topical Products for Joint and		TYR Lophlex LQ 20	294
Temozolomide-Taro	165	Muscular Pain	118	TYR Sphere 20	
Tenofovir disoproxil	108	Topiramate	133	Tyrosine1000	29
Tenofovir Disoproxil Emtricitabine	е	Topiramate Actavis		Tysabri	14
Viatr		Total parenteral nutrition (TPN).		. U -	
Tenofovir Disoproxil Viatris	108	TPN		UK Synacthen	8
Tenoxicam		Tramadol hydrochloride	129	Ultibro Breezhaler	26
Tensipine MR10		Tramal SR 100		Ultraproct	
Tepadina		Tramal SR 150		Umeclidinium	
Terbinafine		Tramal SR 200		Umeclidinium with vilanterol	
Terbutaline sulphate		Trandate		Univent26	
Teriflunomide		Tranexamic acid		Upadacitinib	
Teriflunomide Sandoz		Tranylcypromine sulphate		Ural	
Teriparatide		Trastuzumab (Herzuma)		Urea	
Teriparatide - Teva		Trastuzumab deruxtecan		Urex Forte	
Testogel		Trastuzumab emtansine		Urinary Agents	
Testosterone		Travatan		Urinary Tract Infections	
Testosterone cipionate		Travoprost		UroFos	116
Testosterone esters		Treatments for Dementia		Uromitexan	
Testosterone undecanoate		Treatments for Substance		Ursodeoxycholic acid	
Tetrabenazine		Dependence	149	Ursosan	
Tetrabromophenol		Trelegy Ellipta		Ustekinumab	
Tetracosactrin		Trental 400		Utrogestan	
Tetracycline		Tretinoin		- V -	
Teva Lisinopril		Dermatological	67	Vaccinations	30
Teva-Ketoconazole		Oncology		Vaclovir	
Thalidomide		Trexate		Valaciclovir	
Thalomid		Triamcinolone acetonide		Valganciclovir	
Theophylline		Alimentary	30	Valganciclovir Viatris	
Thiamine hydrochloride		Dermatological		Valine50	29!
Thiamine multichem		Hormone		Vancomycin	
THIO-TEPA		Triamcinolone acetonide with		Vannair	
Thioguanine		gramicidin, neomycin and nys	statin	Varenicline Pfizer	
Thiotepa		Dermatological		Varenicline tartrate	
Thyroid and Antithyroid Agents		Sensory		Varicella vaccine [Chickenpox	
Ticagrelor		Trientine		vaccine]	316
Ticagrelor Sandoz		Trientine Waymade		Varicella zoster vaccine [Shingles	
Tilcotil		Trikafta		vaccine]	
Timolol		Trimethoprim		Varilrix	
Tiotropium bromide		Trimethoprim with		Various	
Tiotropium bromide with		sulphamethoxazole		VariSoft	
olodaterol	261	[Co-trimoxazole]	102	Vasodilators	
Tivicay		Trisequens		Vasopressin Agonists	
Tixagevimab with cilgavimab		Trisul		Vasorex	5t
TMP		Trophic Hormones		Vebulis	
Tobramycin	102	Tropicamide		Vedafil	
Infection	102	Trulicity		Vedolizumab	
		uy		* 000=201100	41

Veletri	- X -
Venclexta166	Xaluprine157
Venetoclax166	Xarelto42
Venlafaxine130	Xifaxan10
Venomil257–258	XMET Maxamum290
VENOX257	Xolair216
Ventolin260	Xolair AU216
Vepesid159	XP Maxamum292
Verapamil hydrochloride50	Xylocaine125
Vermox95	Xylocaine 2% Jelly124
Versacloz	Xylocard 500125
Vesanoid	Xyntha38
Vexazone12	, - Z -
Vfend 104	Zapril45
Viaderm KC71	Zarontin131
Victoza12	Zaroxolyn52
Vigabatrin133	Zavedos160
Vigisom143	Zeffix108
Vildagliptin12	Zejula162
Vildagliptin with metformin	Zematop75
hydrochloride12	Zetlam108
Vimpat131	Ziagen113
Vinblastine sulphate167	Zidovudine [AZT]113
Vincristine sulphate167	Zidovudine [AZT] with
Vinorelbine167	lamivudine113
Vinorelbine Ebewe167	Ziextenzo 43
Vinorelbine Te Arai167	Ziextenzo AU43
Viramune Suspension112	Zimybe54
ViruPOS267	Zinc and castor oil71
Vit.D332	Zinc sulphate34
Vita-B1231	Zincaps34
VitA-POS271	Ziprasidone137
Vitabdeck32	Zista258
Vital282	Zithromax96
Vitamin B complex31	Zo-Rub HP125
Vitamin B6 2531	Zo-Rub Osteo118
Vitamins31–32	Zoladex93
Vitarubin Depot Injection31	Zoledronic acid
Vivonex TEN282	Hormone85
Voltaren 117	Musculoskeletal121
Voltaren D117	Zoledronic acid Viatris
Voltaren SR117	Hormone85
Volumatic266	Musculoskeletal121
Voriconazole104	Zopiclone144
Votrient	Zopiclone Actavis144
Vttack104	Zostrix118
Vyvanse146	Zostrix HP125
- W -	Zuclopenthixol decanoate140
Warfarin sodium42	Zuclopenthixol hydrochloride137
Wart Preparations	Zusdone137
Wasp venom allergy treatment258	Zyban
Water	Zypine137
Blood43	Zypine ODT137
Extemporaneous275	Zyprexa Relprevv138
White Soft Liquid Paraffin AFT71	Zytiga176
Wool fat with mineral oil	Zyvox107
	-, · - · · · · · · · · · · · · · · · · ·