SA2402 - Infliximab

Crohn's disease (adults) - Initial application
Crohn's disease (adults) - Renewal
Crohn's disease (children) - Initial application
Crohn's disease (children) - Renewal
Graft vs host disease - Initial application
Pulmonary sarcoidosis - Initial application
Acute fulminant ulcerative colitis - Initial application
Ankylosing spondylitis - Initial application
Ankylosing spondylitis - Renewal
Chronic ocular inflammation - Initial application
Chronic ocular inflammation - Renewal
Fistulising Crohn's disease - Initial application
Fistulising Crohn's disease - Renewal
Fulminant ulcerative colitis - Renewal
Inflammatory bowel arthritis – axial - Initial application
Inflammatory bowel arthritis – axial - Renewal
Inflammatory bowel arthritis – peripheral - Initial application
Inflammatory bowel arthritis - peripheral - Renewal
Neurosarcoidosis - Initial application
Neurosarcoidosis - Renewal
Plaque psoriasis - Initial application
Plaque psoriasis - Renewal
Previous use - Initial application
Psoriatic arthritis - Initial application
Psoriatic arthritis - Renewal
Pyoderma gangrenosum - Initial application
Pyoderma gangrenosum - Renewal
Rheumatoid arthritis - Initial application11
Rheumatoid arthritis - Renewal
Severe Behcet's disease - Initial application
Severe Behcet's disease - Renewal
Severe ocular inflammation - Initial application
Severe ocular inflammation - Renewal
Ulcerative colitis - Initial application
Ulcerative colitis - Renewal

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
	Address:	
Fax Number:		Fax Number:

Infliximab

Appli	catior	i on — Crohn's disease (adults) om any relevant practitioner. Approvals valid for 6 months. tick boxes where appropriate)	
	and	Patient has active Crohn's disease	
		Patient has a CDAI score of greater than or equal to 300 or HBI score of greater than or equal to 10	
		Patient has extensive small intestine disease affecting more than 50 cm of the small intestine	
		Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection	
		Patient has an ileostomy or colostomy, and has intestinal inflammation	
	and [Patient has tried but has experienced an inadequate response to, or intolerable side effects from, prior therapy with imm and corticosteroids	unomodulators
Rene	wal –	ohn's disease (adults)	
		al Number (if known):	
		m any relevant practitioner. Approvals valid for 2 years. tick boxes where appropriate)	
		CDAI score has reduced by 100 points from the CDAI score, or HBI score has reduced by 3 points, from when the initiated on infliximab	e patient was
		CDAI score is 150 or less, or HBI is 4 or less	
		The patient has demonstrated an adequate response to treatment but CDAI score and/or HBI score cannot be as	sessed
	and [Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treat prior to 1 February 2019	d sixteen weeks

Enquiries	to Ministry	of Health
0800 855	066	

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
	Address:	
Fax Number:		Fax Number:

Infliximab - continued

Initial application — Crohn's disease (children) Applications from any relevant practitioner. Approvals valid for 6 months. Prerequisites(tick boxes where appropriate)	
Paediatric patient has active Crohn's disease	
Patient has a PCDAI score of greater than or equal to 30	
Patient has extensive small intestine disease	
and Patient has tried but experienced an inadequate response to, or intolerable side effects from, prior therapy with immunomodulators and corticosteroids	
Renewal — Crohn's disease (children)	
Current approval Number (if known):	
Applications from any relevant practitioner. Approvals valid for 2 years. Prerequisites (tick boxes where appropriate)	
PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on infliximab	
PCDAI score is 15 or less	
or The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed	
and	1
Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up	
to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose	;
prior to 1 February 2019	
Initial application — Graft vs host disease Applications from any relevant practitioner. Approvals valid without further renewal unless notified. Prerequisites(tick box where appropriate)	
Patient has steroid-refractory acute graft vs. host disease of the gut	
Initial application — Pulmonary sarcoidosis Applications from any relevant practitioner. Approvals valid without further renewal unless notified. Prerequisites(tick box where appropriate)	
Patient has life-threatening pulmonary sarcoidosis diagnosed by a multidisciplinary team that is refractory to other treatments	

Enquiries	to Ministry	of Health
0800 855	066	

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
	Address:	
Fax Number:		Fax Number:

Infliximab - continued

Initial application — acute fulminant ulcerative colitis Applications only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 weeks. Prerequisites(tick boxes where appropriate)
Patient has acute, fulminant ulcerative colitis and Treatment with intravenous or high dose oral corticosteroids has not been successful
Initial application — ankylosing spondylitis Applications only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months. Prerequisites(tick boxes where appropriate)
The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis and
The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept or
Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis
Renewal — ankylosing spondylitis

Current approval Number (if known):.....

Applications only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months. **Prerequisites**(tick boxes where appropriate)

> Following 12 weeks of infliximab treatment, BASDAI has improved by 4 or more points from pre-infliximab baseline on a 10 point scale, or by 50%, whichever is less

Physician considers that the patient has benefited from treatment and that continued treatment is appropriate

Infliximab to be administered at doses no greater than 5 mg/kg every 6-8 weeks

I confirm the above details are correct and that in signing this form I understand I may be audited.

and

and

Enquiries	to Ministry	of Health
0800 855	066	

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
	Address:	
Fax Number:		Fax Number:

Infliximab - continued

Initial application — chronic ocular inflammation

Applications from any relevant practitioner. Approvals valid for 4 months.

Prerequisites(tick boxes where appropriate)

	and		The patient has had an initial Special Authority approval for adalimumab for chronic ocular inflammation
		or	The patient has experienced intolerable side effects from adalimumab
		U1	The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for chronic ocular inflammation
or			
	and		Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision lo
	and		Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision logonal patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective
	and	or	

Rene	ewal — chronic ocular inflammation
Curr	ent approval Number (if known):
	ications from any relevant practitioner. Approvals valid for 12 months.
	equisites(tick boxes where appropriate)
	 The patient has had a good clinical response following 3 initial doses Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema) Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old
	A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible n loss if infliximab is withdrawn.

Enquiries	to Ministry	of Health
0800 855	066	

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
	Address:	
Fax Number:		Fax Number:

Infliximab - continued

Initial application — fistulising Crohn's disease Applications from any relevant practitioner. Approvals valid for 6 months. Prerequisites(tick boxes where appropriate)		
Patient has confirmed Crohn's disease		
Patient has one or more complex externally draining enterocutaneous fistula(e) or		
Patient has one or more rectovaginal fistula(e) or		
Patent has complex peri-anal fistula		
Renewal — fistulising Crohn's disease		
Current approval Number (if known): Applications only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 2 years. Prerequisites (tick boxes where appropriate)		
or The number of open draining fistulae have decreased from baseline by at least 50% There has been a marked reduction in drainage of all fistula(e) from baseline (in the case of adult patients, as demonstrated by a reduction in the Fistula Assessment score), together with less induration and patient reported pain		
and Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019		
Initial application — neurosarcoidosis Applications only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months. Prerequisites(tick boxes where appropriate)		
Patient has been diagnosed with neurosarcoiosis by a multidisciplinary team and		
and Patient has CNS involvement		
and		
IV cyclophosphamide has been tried		
Treatment with IV cyclophosphamide is clinically inappropriate		

Enquiries to Ministry of Health 0800 855 066

APPLICATION FOR SUBSIDY BY SPECIAL AUTHORITY

APPLICANT (stamp or sti	cker acceptable)	PATIENT NHI:	REFERRER Reg No:	
Reg No:		First Names:	First Names:	
Name:		Surname:	Surname:	
Address:		DOB:	Address:	
		Address:		
Fax Number:			Fax Number:	
Infliximab - continued				
Renewal — neurosarco	idosis			
Current approval Number (if known):				
	0	on the recommendation of a neurologist. Approvals	valid for 18 months.	
Prerequisites(tick boxes	where appropriate)			
A withdra	wal period has been tried	and the patient has relapsed		
	vithdrawal period has bee	n considered but would not be clinically appropriate		
and The and The and	ere has been a marked re	duction in prednisone dose		
	There has been an im	provement in MRI appearances		
or	Marked improvement	in other symptomology		

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
	Address:	
Fax Number:		Fax Number:

Infliximab - continued

App	lication	ns only	on — plaque psoriasis y from a dermatologist or any relevant practitioner on the recommendation of a dermatologist. Approvals valid for 3 months. ck boxes where appropriate)	
		and	The patient has had an initial Special Authority approval for adalimumab, etanercept or secukinumab for severe chronic plaque psoriasis	
			Patient has experienced intolerable side effects from adalimumab, etanercept or secukinumab	
			Patient has received insufficient benefit from adalimumab, etanercept or secukinumab to meet the renewal criteria for adalimumab, etanercept or secukinumab for severe chronic plaque psoriasis	
	or			_
			 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis 	
			Patient has severe chronic localised genital or flexural plaque psoriasis where the plaques or lesions have been present for at least 6 months from the time of initial diagnosis, and with a Dermatology Life Quality Index (DLQI) score greater than 10	
		and		
		and	Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin	
		and	A PASI assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course	
		[The most recent PASI assessment is no more than 1 month old at the time of initiation	
while face seve	e still o , hand, re, and	n treat foot, g d for th	te response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably tment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the genital or flexural areas at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very ne face, palm of a hand or sole of a foot the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed still on treatment but no longer than 1 month following cessation of the most recent prior treatment.	b

Enquiries	to Ministry	of Health
0800 855	066	

APPLICANT (sta	mp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:	
Reg No:		First Names:	First Names:	
Name:		Surname:	Surname:	
Address:		DOB:	Address:	
		Address:		
Fax Number:			Fax Number:	
nfliximab - co	ontinued			
Renewal — plac	que psoriasis			
Current approval	l Number (if known):			
	n any relevant practitioner. Approv			
	ck boxes where appropriate)			
or	and Following each prior in	dy" severe chronic plaque psoriasis at the start of tre nfliximab treatment course the patient has a PASI sco , when compared with the pre-infliximab treatment ba	ore which is reduced by 75% or more, or is	
or	Patient had severe ch	ronic plaque psoriasis of the face, or palm of a hand	or sole of a foot at the start of treatment	
	Following each	prior infliximab treatment course the patient has a rec a, thickness and scaling, to slight or better, or sustain values		
		prior infliximab treatment course the patient has a rea tained at this level, as compared to the pre-infliximab		
or	Patient had severe ch	ronic localised genital or flexural plaque psoriasis at t	he start of treatment	
		experienced a reduction of 75% or more in the skin a pre-treatment baseline value	area affected, or sustained at this level, as	
		ermatology Quality of Life Index (DLQI) improvement ncing infliximab	of 5 or more, as compared to baseline DLQI	
and I	nfliximab to be administered at do	ses no greater than 5 mg/kg every 8 weeks		

Enquiries to Ministry of Health 0800 855 066

APPLICATION FOR SUBSIDY BY SPECIAL AUTHORITY

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
	Address:	
Fax Number:		Fax Number:

Infliximab - continued

Initial application — previous use Applications from any relevant practitioner. Approvals valid for 6 months. Prerequisites(tick boxes where appropriate)		
and	Patient was being treated with infliximab prior to 1 February 2019	
	Rheumatoid arthritis	
	Ankylosing spondylitis	
	Psoriatic arthritis or	
	or Severe ocular inflammation	
	Chronic ocular inflammation	
	Crohn's disease (adults)	
	Crohn's disease (children) or	
	Fistulising Crohn's disease or	
	Severe fulminant ulcerative colitis or	
	Severe ulcerative colitis or	
	Plaque psoriasis or	
	or	
	Severe Behcet's disease	
nitial appli	cation — psoriatic arthritis	
Application	s only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months. es(tick boxes where appropriate)	

and		The p	patient has had an initial Special Authority approval for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis
	or		The patient has experienced intolerable side effects from adalimumab and/or etanercept and/or secukinumab
	01		Following 3-4 months' initial treatment with adalimumab and/or etanercept and/or secukinumab, the patient did not meet the renewal criteria for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis

Enquiries	to Ministry	of Health
0800 855	066	

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
	Address:	
Fax Number:		Fax Number:

Renewal — psoriatic arthritis			
urrent ap	proval Number (if known):		
-	s only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months. tes (tick boxes where appropriate)		
	 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior infliximab treatment in the opinion of the treating physician 		
and [Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks		
pplicatior	lication — rheumatoid arthritis ns only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months. tes (tick boxes where appropriate)		
and	The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis		
	The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept or		
	Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumat and/or etanercept		
and [Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance		
enewal –	- rheumatoid arthritis		
urrent ap	proval Number (if known):		
-	s only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months. tes (tick boxes where appropriate)		
and	Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance		
	Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician		
	The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician		

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
	DOB:	
	Address:	
Fax Number:		Fax Number:

Infliximab - continued

Initial application — severe Behcet's disease Applications from any relevant practitioner. Approvals valid for 4 months.				
Prerequisites (tick boxes where appropriate)				
The patient has severe Behcet's disease which is significantly impacting the patient's quality of life (see Notes) and				
The patient has severe ocular, neurological and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s) (see Notes)				
The patient has severe gastrointestinal, rheumatologic and/or mucocutaneous symptoms and has not responded adequately to two or more treatment appropriate for the particular symptom(s) (see Notes)				
and The patient is experiencing significant loss of quality of life				
Note: Behcet's disease diagnosed according to the International Study Group for Behcet's Disease. Lancet 1990;335(8697):1078-80. Quality of life measured using an appropriate quality of life scale such as that published in Gilworth et al J Rheumatol. 2004;31:931-7. Treatments appropriate for the particular symptoms are those that are considered standard conventional treatments for these symptoms, for example intravenous/oral steroids and other immunosuppressants for ocular symptoms; azathioprine, steroids, thalidomide, interferon alpha and ciclosporin for mucocutaneous symptoms; and colchicine, steroids and methotrexate for rheumatological symptoms.				
Renewal — severe Behcet's disease				
Current approval Number (if known):				
Applications from any relevant practitioner. Approvals valid for 6 months.				
Prerequisites(tick boxes where appropriate)				
Patient has had a good clinical response to initial treatment with measurably improved quality of life and				
Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks				
Renewal — fulminant ulcerative colitis				
Current approval Number (if known):				
Applications from any relevant practitioner. Approvals valid for 2 years.				
Prerequisites(tick boxes where appropriate)				
Where maintenance treatment is considered appropriate, infliximab should be used in combination with immunomodulators and reassessed every 6 months and				
Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019				

Enquiries to Mir	nistry of Health
0800 855 066	-

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
	Address:	
Fax Number:		Fax Number:

Infliximab - continued

Initial application — severe ocular inflammation Applications from any relevant practitioner. Approvals valid for 4 months. Prerequisites(tick boxes where appropriate)					
	The patient has had an initial Special Authority approval for adalimumab for severe ocular inflammation and				
			or	The patient has experienced intolerable side effects from adalimumab	
				The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe ocular inflammation	
	or				
	Patient has severe, vision-threatening ocular inflammation requiring rapid control and				
	Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms				
			or	Patient developed new inflammatory symptoms while receiving high dose steroids	
			or	Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms	

Renewal — severe ocular inflammation				
Current approval Number (if known): Applications from any relevant practitioner. Approvals valid for 12 months. Prerequisites (tick boxes where appropriate)				
 or The patient has had a good clinical response following 3 initial doses Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema) or Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg 				
daily, or steroid drops less than twice daily if under 18 years old Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.				

Enquiries	to Ministry	of Health
0800 855	066	

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
	Address:	
Fax Number:		Fax Number:

Infliximab - continued

itial application — ulcerative colitis Applications from any relevant practitioner. Approvals valid for 6 months. rerequisites(tick boxes where appropriate)			
Patient has active ulcerative colitis			
Patients SCCAI is greater than or equal to 4			
Patients PUCAI score is greater than or equal to 20			
and Patient has tried but has experienced an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and systemic corticosteroids			
Renewal — ulcerative colitis			
Current approval Number (if known):			
Applications from any relevant practitioner. Approvals valid for 2 years.			
Prerequisites(tick boxes where appropriate)			
The SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on infliximab			
The PUCAI score has reduced by 10 points or more from the PUCAI score when the patient was initiated on infliximab			
and Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019			
Initial application — pyoderma gangrenosum Applications only from a dermatologist. Approvals valid for 4 months. Prerequisites(tick boxes where appropriate)			
Patient has pyoderma gangrenosum*			
Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response			
A maximum of 8 doses			

Note: Note: Indications marked with * are unapproved indications.

Enquiries	to Ministry	of Health
0800 855	066	

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:		
Reg No:	First Names:	First Names:		
Name:	Surname:	Surname:		
Address:	DOB:	Address:		
	Address:			
Fax Number:		Fax Number:		
Infliximab - continued				
Renewal — pyoderma gangrenosum				
Current approval Number (if known):				
Applications only from a dermatologist or Practition	ner on the recommendation of a dermatologist. Appre	ovals valid for 4 months.		
Prerequisites(tick boxes where appropriate)				
Patient has shown clinical improvement and Patient continues to require treatment and				
A maximum of 8 doses				
Initial application — inflammatory bowel arthritis – axial Applications from any relevant practitioner. Approvals valid for 6 months. Prerequisites(tick boxes where appropriate)				
Patient has a diagnosis of active u	lcerative colitis or active Crohn's disease			
Patient has had axial inflammatory	pain for six months or more			
and Patient is unable to take NSAIDs				
and Patient has unequivocal sacroiliitis demonstrated by radiological imaging or MRI and				
	ed adequately to prior treatment consisting of at least	3 months of an exercise regime supervised by a		
	on a 0 - 10 scale completed after the 3 month exer	rcise trial, but prior to ceasing any previous		
Renewal — inflammatory bowel arthritis – axial				
Current approval Number (if known): Applications from any relevant practitioner. Approvals valid for 2 years.				

Prerequisites(tick box where appropriate)

Treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10-point scale, or an improvement in BASDAI of 50%, whichever is less

Enquiries	to Ministry	of Health
0800 855	066	

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
	Address:	
Fax Number:		Fax Number:

Infliximab - continued

Initial application — inflammatory bowel arthritis – peripheral Applications from any relevant practitioner. Approvals valid for 6 months. Prerequisites(tick boxes where appropriate)				
	and	Patient has a diagnosis of active ulcerative colitis or active Crohn's disease		
	and	Patient has active arthritis in at least four joints from the following: hip, knee, ankle, subtalar, tarsus, forefoot, wrist, elbow, shoulder, sternoclavicular		
		Patient has tried and not experienced a response to at least three months of methotrexate or azathioprine at a maximum tolerated dose (unless contraindicated)		
	and	Patient has tried and not experienced a response to at least three months of sulfasalazine at a maximum tolerated dose (unless contraindicated)		
		Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application or		
		Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application		
		ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months		
Rene	Renewal — inflammatory bowel arthritis – peripheral			
Curre	Current approval Number (if known):			
Applications from any relevant practitioner. Approvals valid for 2 years. Prerequisites (tick boxes where appropriate)				

Following initial treatment, patient has experienced at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician

Patient has experienced at least a continuing 30% improvement in active joint count from baseline in the opinion of the treating physician

I confirm the above details are correct and that in signing this form I understand I may be audited.

or