February 2025

Editors:

Kave Wilson & Doris Chong email: enquiry@pharmac.govt.nz Telephone +64 4 460 4990 Level 9, 40 Mercer Street PO Box 10 254 Wellington

Freephone Information Line 0800 66 00 50 (9am - 5pm weekdays)

Circulation

You can register to have an electronic version of the Pharmaceutical Schedule (link to PDF copy) emailed to your nominated email address each month by subscribing at pharmac.govt.nz/subscribe.

Production

Typeset automatically from XML and T_FX. XML version of the Schedule available from schedule.pharmac.govt.nz/pub/schedule

Programmers

Anrik Drenth

email: texschedule@pharmac.govt.nz ©Pharmaceutical Management Agency



ISSN 1179-3686

This work is licensed under the Creative Commons Attribution 4.0 International licence. In essence, you are free to copy, distribute and adapt it, as long as you attribute the work to Pharmac and abide by the other licence terms. To view a copy of this licence, visit: creativecommons.org/licenses/by/4.0/. Attribution to Pharmac should be in written form and not by reproduction of the Pharmac logo. While care has been taken in compiling this Schedule, Pharmac takes no responsibility for any errors or omissions, and shall not be liable for any consequences arising there from.

| introducing File | illac Z |
|-------------------|-----------------|
| General | Rules 5 |
| ıry Tract & Metab | olism 6 |
| Blood Forming O | gans 35 |
| Cardiovascular S | stem 44 |
| Dermatolo | icals 66 |
| Genito Urinary S | stem 76 |
| eparations – Sys | emic 83 |
| ents For Systemi | Use 94 |
| usculoskeletal S | stem 116 |
| Nervous S | stem 122 |
| Immunosuppres | sants 152 |
| ory System & Alle | rgies 254 |
| Sensory O | gans 265 |
| Va | rious 270 |
| ıs Compounds (E | CPs) 272 |
| Special I | oods 274 |
| mmunisation Sch | edule 298 |
| | 7010V 914 |

Introducing Pharmac

Index

Introducing Pharmac

The Pharmaceutical Management Agency (Pharmac) makes decisions that help control Government spending on pharmaceuticals. This includes community pharmaceuticals, hospital pharmaceuticals, vaccines and increasingly, hospital medical devices. Pharmac negotiates prices, sets subsidy levels and conditions, and makes decisions on changes to the subsidised list.

Pharmac's role:

"to secure for eligible people in need of pharmaceuticals, the best health outcomes that are reasonably achievable from pharmaceutical treatment and from within the amount of funding provided."

Pae Ora (Healthy Futures) Act 2022

To ensure our decisions are as fair and robust as possible we use a decision-making process that incorporates clinical, economic and commercial issues. We also seek the views of users and the wider community through consultation. The processes we generally use are outlined in our Operating Policies and Procedures.

Further information about Pharmac and the way we make funding decisions can be found on the Pharmac website at https://pharmac.govt.nz/about.

Purpose of the Pharmaceutical Schedule

The purpose of the Schedule is to list:

- the Community Pharmaceuticals that are subsidised by the Government and to show the amount of the subsidy paid to contractors, as well as the manufacturer's price and any access conditions that may apply;
- the Hospital Pharmaceuticals that may be used in Health NZ Hospitals, as well as any access conditions that may apply; and
- the Pharmaceuticals, including Medical Devices, used in Health NZ Hospitals for which national prices have been negotiated by Pharmac.

The Schedule does not show the final cost to Government of subsidising each Community Pharmaceutical or to Health NZ Hospitals in purchasing each Pharmaceutical, since that will depend on any rebate and other arrangements Pharmac has with the supplier and, for Pharmaceuticals used in Health NZ Hospitals, on any logistics arrangements put in place.

This book contains sections A to D and Section I of the Pharmaceutical Schedule and lists the Pharmaceuticals funded for use in the community, including vaccines, as well as haemophilia and cancer treatments given in Health NZ Hospitals. Section H lists the Pharmaceuticals that that can be used in Health NZ Hospitals and is a separate publication.


The Pharmaceuticals in this book are listed by therapeutic group, which is based on the WHO Anatomical Therapeutic Chemical (ATC) system. The listings are displayed alphabetically under each heading.

The index lists both chemical entities and product brand names.

Explaining pharmaceutical entries

The Pharmaceutical Schedule lists pharmaceuticals subsidised by the Government, the subsidy, the supplier's price and the access conditions that may apply.

Example



Glossary

Units of Measure

| gramg kilogramkg international unitiu | mi mi mi |
|---------------------------------------|----------------|
| Abbreviations | |
| AmpouleAmp | Ge |
| CapsuleCap | Gr |
| Cream | Inf |
| DeviceDev | Ini |
| DispersibleDisp | Lic |
| EffervescentEff | Lo |
| EmulsionEmul | Oi |
| Enteric Coated EC | Sa |

| microgrammilligrammillilitre | mg |
|------------------------------|------|
| Gelatinous | |
| Granules | |
| Infusion | Inf |
| Injection | Inj |
| Liquid | Liq |
| Long Acting | LA |
| Ointment | Oint |
| Sachet | Sach |

| millimoleunit | |
|------------------------------|-------------|
| Solution | Supp Tab |
| Trans Dermal Delivery System | TDDS |

Read the General Rules: https://pharmac.govt.nz/section-a.

SECTION B: ALIMENTARY TRACT AND METABOLISM

| SECTION B: ALIMENTARY TRACT AND MET | TABOLISM | | | |
|---|---|------------------|-------------------|---|
| | Subsidy (Manufacturer's Price) \$ | Su Per | Fully bsidised | Brand or Generic Manufacturer |
| Antacids and Antiflatulents | | | | |
| Antacids and Reflux Barrier Agents | | | | |
| ALGINIC ACID Sodium alginate 225 mg and magnesium alginate 87.5 mg pe sachet | | 30 | √ G | aviscon Infant |
| * Tab 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg - peppermint flavour | 1.80 (14.39) | 60 | G | aviscon Extra Strength |
| Oral liq 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg per 10 ml | | 500 ml | A | cidex |
| Phosphate Binding Agents | | | | |
| ALUMINIUM HYDROXIDE * Tab 600 mg CALCIUM CARBONATE Oral liq 1,250 mg per 5 ml (500 mg elemental per 5 ml) – | | 100 | | lu-Tab |
| Subsidy by endorsement | 47.30 | 500 ml 473 ml | √ C | oxane alcium carbonate PAI ⁸²⁹ |
| Only when prescribed for patients unable to swallow calc inappropriate and the prescription is endorsed accordingl | | ts or whe | ere calciur | n carbonate tablets are |
| Antidiarrhoeals | | | | |
| Agents Which Reduce Motility | | | | |
| LOPERAMIDE HYDROCHLORIDE – Up to 30 cap available on a * Tab 2 mg * Cap 2 mg | 10.75 | 400 400 | ✓ N ✓ <u>D</u> | odia iamide Relief |
| Rectal and Colonic Anti-inflammatories | | | | |
| BUDESONIDE Cap modified-release 3 mg - Special Authority see SA1886 below - Retail pharmacy | 87.60 | 90 | ✓ <u>B</u> | udesonide Te Arai |

⇒SA1886 Special Authority for Subsidy

Initial application — (Crohn's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Mild to moderate ileal, ileocaecal or proximal Crohn's disease; and
- 2 Any of the following:
 - 2.1 Diabetes; or

| Subsidy | F | ully | Brand or |
|------------------------|--------|------|--------------|
| (Manufacturer's Price) | Subsid | ised | Generic |
| \$ | Per | 1 | Manufacturer |

continued...

- 2.2 Cushingoid habitus; or
- 2.3 Osteoporosis where there is significant risk of fracture; or
- 2.4 Severe acne following treatment with conventional corticosteroid therapy; or
- 2.5 History of severe psychiatric problems associated with corticosteroid treatment; or
- 2.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
- 2.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated).

Initial application — (collagenous and lymphocytic colitis (microscopic colitis)) from any relevant practitioner. Approvals valid for 6 months where patient has a diagnosis of microscopic colitis (collagenous or lymphocytic colitis) by colonoscopy with biopsies.

Initial application — (gut Graft versus Host disease) from any relevant practitioner. Approvals valid for 6 months where patient has a gut Graft versus Host disease following allogenic bone marrow transplantation*.

Note: Indication marked with * is an unapproved indication.

Initial application — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has autoimmune hepatitis*: and
- 2 Patient does not have cirrhosis; and
- 3 Any of the following:
 - 3.1 Diabetes: or
 - 3.2 Cushingoid habitus; or
 - 3.3 Osteoporosis where there is significant risk of fracture; or
 - 3.4 Severe acne following treatment with conventional corticosteroid therapy; or
 - 3.5 History of severe psychiatric problems associated with corticosteroid treatment; or
 - 3.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
 - 3.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated); or3.8 Adolescents with poor linear growth (where conventional corticosteroid use may limit further growth).

Note: Indication marked with * is an unapproved indication.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

HYDROCORTISONE ACETATE Rectal foam 10%, CFC-Free (14 applications)......57.09 15 g OP ✓ Colifoam HYDROCORTISONE ACETATE WITH PRAMOXINE HYDROCHLORIDE Topical aerosol foam, 1% with pramoxine hydrochloride 1%.......26.55 10 g OP ✓ Proctofoam S29 **MESALAZINE** 100 ✓ Asacol 100 ✓ Pentasa Tab 800 mg85.50 90 ✓ Asacol ✓ Asacol S29 S29 100 OP ✓ Pentasa ✓ Pentasa Enema 1 g per 100 ml41.30 7 20 ✓ Asacol

✓ Pentasa

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | |
|--------------------------------|---|-----|---------------------|----------------|
| OLSALAZINE | | | | |
| Tab 500 mg | 56.02 | 60 | • | Atnahs |
| | | | | Olsalazine S29 |
| | 93.37 | 100 | ✓ | Dipentum |
| Cap 250 mg | 53.00 | 100 | ✓ | Dipentum |
| SODIUM CROMOGLICATE Cap 100 mg | 113.35 | 100 | ✓ | Ralicrom |
| SULFASALAZINE * Tab 500 mg | 16.52 | 100 | 1 | Salazopyrin |
| * Tab EC 500 mg | | 100 | | Salazopyrin EN |

Local preparations for Anal and Rectal Disorders

Antihaemorrhoidal Preparations

| FLUOCORTOLONE CAPROATE WITH FLUOCORTOLONE PIVALATE AND CII | NCHOCAINE | |
|--|-----------|------------------------------|
| Oint 950 mcg, with fluocortolone pivalate 920 mcg, and | | |
| cinchocaine hydrochloride 5 mg per g13.05 | 30 g OP | Ultraproct |
| Suppos 630 mcg, with fluocortolone pivalate 610 mcg, and | | |
| cinchocaine hydrochloride 1 mg8.61 | 12 | Ultraproct |
| HYDROCORTISONE WITH CINCHOCAINE | | |
| Oint 5 mg with cinchocaine hydrochloride 5 mg per g15.00 | 30 g OP | ✓ Proctosedyl |
| Suppos 5 mg with cinchocaine hydrochloride 5 mg per g9.90 | 12 | ✓ Proctosedyl |

Management of Anal Fissures

GLYCERYL TRINITRATE − Special Authority see SA1329 below − Retail pharmacy

★ Oint 0.2%......22.00 30 g OP

✓ Rectogesic

⇒SA1329 Special Authority for Subsidy

CL VCODVDDONII IM BDOMIDE

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has a chronic anal fissure that has persisted for longer than three weeks.

Antispasmodics and Other Agents Altering Gut Motility

| GLYCOPYRRONIUM BROMIDE | | |
|---|-----|---|
| Inj 200 mcg per ml, 1 ml ampoule – Up to 10 inj available on a PSO19.00 | 5 | ✓ Robinul |
| HYOSCINE BUTYLBROMIDE | | |
| * Tab 10 mg2.25 | 20 | ✓ Hyoscine Butylbromide (Adiramedica) |
| 6.35 | 100 | ✓ Buscopan |
| Hyoscine Butylbromide (Adiramedica) to be Principal Supply on 1 April 2025 | | · |
| * Inj 20 mg, 1 ml – Up to 5 inj available on a PSO1.91 (Buscopan Tab 10 mg to be delisted 1 April 2025) | 5 | ✓ <u>Spazmol</u> |
| MEBEVERINE HYDROCHLORIDE | | |
| * Tab 135 mg8.50 | 90 | ✓ Colofac |

Subsidy Fully Brand or
(Manufacturer's Price) Subsidised Generic
\$ Per Manufacturer

Antiulcerants

Antisecretory and Cytoprotective

MISOPROSTOL - Wastage claimable

Helicobacter Pylori Eradication

CLARITHROMYCIN

Tab 500 mg − Subsidy by endorsement.......14.58 14 ✓ Klacid

- a) Maximum of 28 tab per prescription
- Subsidised only if prescribed for helicobacter pylori eradication and prescription is endorsed accordingly.
 Note: the prescription is considered endorsed if clarithromycin is prescribed in conjunction with a proton pump inhibitor and either amoxicillin or metronidazole.

H2 Antagonists

| FΑ | MOTIDINE - Only on a prescription | | | |
|----|---|----------------------|----------------|--------------|
| * | Tab 20 mg | 4.91 | 100 | ✓ Famotidine |
| | • | | | Hovid S29 |
| * | Tab 40 mg | 10.32 | 100 | ✓ Famotidine |
| | • | | | Hovid S29 |
| * | Inj 10 mg per ml, 4 ml - Subsidy by endorsement | CBS | 10 | ✓ Mylan S29 |
| | Subsidy by endorsement – Subsidised for natients rece | iving treatment as i | nart of nallia | tive care |

Proton Pump Inhibitors

| LANSOPRAZOLE | | |
|---|-----|-----------------------------------|
| * Cap 15 mg4.04 | 100 | ✓ Lanzol Relief |
| * Cap 30 mg5.43 | 100 | ✓ Lanzol Relief |
| OMEPRAZOLE | | |
| For omeprazole suspension refer Standard Formulae, page 272 | | |
| * Cap 10 mg2.06 | 90 | Omeprazole Teva |
| | | ✓ Omeprazole actavis 10 |
| * Cap 20 mg2.02 | 90 | Omeprazole Teva |
| · · | | ✓ Omeprazole actavis 20 |
| * Cap 40 mg3.18 | 90 | ✓ Omeprazole Teva |
| 3 | | ✓ Omeprazole actavis 40 |
| * Powder – Only in combination42.50 | 5 q | ✓ Midwest |
| Only in extemporaneously compounded omeprazole suspension. | · | |
| * Inj 40 mg ampoule with diluent | 5 | ✓ <u>Dr Reddy's</u> Omeprazole |
| | | ✓ Ocicure S29 |
| PANTOPRAZOLE | | |
| | 00 | A Bonzon Bolief |
| * Tab EC 20 mg | 90 | ✓ Panzop Relief |
| * Tab EC 40 mg2.74 | 90 | ✓ Panzop Relief |

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

| | Subsidy (Manufacturer's Price \$ | e) Per | Fully Subsidised | Brand or Generic Manufacturer |
|---|--|-----------------------|--------------------------------|---|
| Site Protective Agents | | | | |
| COLLOIDAL BISMUTH SUBCITRATE Tab 120 mgSUCRALFATE | 14.51 | 50 | ✓ G | astrodenol \$29 |
| Tab 1 g | 35.50 (48.28) | 120 | С | arafate |
| Bile and Liver Therapy | | | | |
| RIFAXIMIN – Special Authority see SA1461 below – Retail pl Tab 550 mg SA1461 Special Authority for Subsidy | , | 56 | √ <u>X</u> | <u>ifaxan</u> |
| Initial application only from a gastroenterologist, hepatologist nepatologist. Approvals valid for 6 months where the patient tolerated doses of lactulose. Renewal only from a gastroenterologist, hepatologist or Pract nepatologist. Approvals valid without further renewal unless repending from treatment. | has hepatic encephalo | oathy d | lespite an ad n of a gastro | lequate trial of maximum enterologist or |
| Diabetes | | | | |
| | | | | |
| Hyperglycaemic Agents | la a mara a su | | | |
| DIAZOXIDE - Special Authority see SA1320 below - Retail p Cap 25 mg Cap 100 mg Oral liq 50 mg per ml → SA1320 Special Authority for Subsidy | 110.00 | 100 100 30 ml C | ✓ P | roglicem \$29 roglicem \$29 5 Pharma \$29 |
| nitial application from any relevant practitioner. Approvals hypoglycaemia caused by hyperinsulinism. | valid for 12 months whe | ere use | d for the trea | atment of confirmed |
| Renewal from any relevant practitioner. Approvals valid with appropriate and the patient is benefiting from treatment. GLUCAGON HYDROCHLORIDE | out further renewal unle | ess noti | ified where th | he treatment remains |
| Inj 1 mg syringe kit – Up to 5 kit available on a PSO | 32.00 | 1 | ✓ G | lucagen Hypokit |
| Insulin - Short-acting Preparations | | | | |
| NSULIN NEUTRAL ▲ Inj human 100 u per ml | 25.26 | 10 ml C | | ctrapid lumulin R |
| ▲ Inj human 100 u per ml, 3 ml | 42.66 | 5 | ✓ A | ctrapid Penfill umulin R |
| Insulin - Intermediate-acting Preparations | | | | |
| NSULIN ASPART WITH INSULIN ASPART PROTAMINE Inj 100 iu per ml, 3 ml prefilled pen | 52.15 | 5 | ✓ N | ovoMix 30 FlexPen |

| | Subsidy (Manufacturer's P \$ | rice) Subs Per | Fully Brand or idised Generic Manufacturer |
|--|------------------------------------|-------------------|--|
| NSULIN ISOPHANE | | | |
| Inj human 100 u per ml | 17.68 | 10 ml OP | ✓ Humulin NPH✓ Protaphane |
| Inj human 100 u per ml, 3 ml | 29.86 | 5 | ✓ Humulin NPH✓ Protaphane Penfill |
| NSULIN ISOPHANE WITH INSULIN NEUTRAL | | | |
| Inj human with neutral insulin 100 u per ml | 25.26 | 10 ml OP | ✓ Humulin 30/70✓ Mixtard 30 |
| Inj human with neutral insulin 100 u per ml, 3 ml | 42.66 | 5 | ✓ Humulin 30/70 ✓ PenMix 30 ✓ PenMix 50 |
| Mixtard 30 Inj human with neutral insulin 100 u per ml to be deli PenMix 50 Inj human with neutral insulin 100 u per ml, 3 ml to b NSULIN LISPRO WITH INSULIN LISPRO PROTAMINE | e delisted 1 June | | Penwix 50 |
| Inj lispro 25% with insulin lispro protamine 75% 100 u per ml | 42.66 | 5 | ✓ Humalog Mix 25 |
| Inj lispro 50% with insulin lispro protamine 50% 100 u per ml 3 ml | | 5 | ✓ Humalog Mix 50 |
| Insulin - Long-acting Preparations | | | |
| NSULIN GLARGINE | | | |
| Inj 100 u per ml, 10 ml | 63.00 | 1 | ✓ Lantus |
| Inj 100 u per ml, 3 ml | 94.50 | 5 | ✓ Lantus |
| Inj 100 u per ml, 3 ml disposable pen | 94.50 | 5 | ✓ Lantus SoloStar |
| Insulin - Rapid Acting Preparations | | | |
| NSULIN ASPART | | | |
| Inj 100 u per ml, 10 ml | | 1_ | ✓ NovoRapid |
| Inj 100 u per ml, 3 ml | | 5 | ✓ NovoRapid Penfill |
| ▲ Inj 100 u per ml, 3 ml syringe | 51.19 | 5 | ✓ NovoRapid FlexPen |
| Inj 100 u per ml, 10 ml | 27.03 | 1 | ✓ Apidra |
| Inj 100 u per ml, 3 ml | | 5 | ✓ Apidra |
| Inj 100 u per ml, 3 ml disposable pen | 46.07 | 5 | ✓ Apidra SoloStar |
| | | 10 ml OP 5 | ✓ Humalog✓ Humalog |
| Alpha Glucosidase Inhibitors | | | |
| • | | | |
| CARBOSE ★ Tab 50 mg ★ Tab 100 mg | | 90 90 | ✓ <u>Accarb</u> ✓ <u>Accarb</u> |
| Oral Hypoglycaemic Agents | | | |
| GLIBENCLAMIDE | | | |
| k Tab 5 mg | 7 50 | 100 | ✓ Daonil |
| v | 7.50 | 100 | - Duyiiii |
| GLICLAZIDE 烙 Tab 80 mg | 20.10 | 500 | ✓ Clizido |
| r 1 au 00 mg | ∠0.10 | 500 | ✓ <u>Glizide</u> |

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Brand or Subsidised Generic Manufacturer |
|---|---|----------------|---|
| # Tab 5 mg Minidiab to be Principal Supply on 1 March 2025 | 6.86 | 100 | ✓ Minidiab |
| METFORMIN HYDROCHLORIDE * Tab immediate-release 500 mg * Tab immediate-release 850 mg | | 1,000 500 | ✓ <u>Metformin Viatris</u> ✓ <u>Metformin Viatris</u> |
| PIOGLITAZONE * Tab 15 mg * Tab 30 mg * Tab 45 mg | 7.25 | 90 90 90 | ✓ <u>Vexazone</u> ✓ <u>Vexazone</u> ✓ <u>Vexazone</u> |
| VILDAGLIPTIN Tab 50 mg | | 60 | ✓ Galvus |
| VILDAGLIPTIN WITH METFORMIN HYDROCHLORIDE Tab 50 mg with 1,000 mg metformin hydrochloride Tab 50 mg with 850 mg metformin hydrochloride | | 60 60 | ✓ Galvumet✓ Galvumet |

GLP-1 Agonists

DULAGLUTIDE - Special Authority see SA2338 below - Retail pharmacy

Note: Not to be given in combination with a funded SGLT-2 inhibitor or other GLP-1 agonist.

⇒SA2338 Special Authority for Subsidy

Note: Subsidy for patients with existing approvals prior to 1 May 2024. Approvals valid without further renewal unless notified. No new patients will be granted from 1 May 2024 until further notice.

LIRAGLUTIDE - Special Authority see SA2339 below - Retail pharmacy

- a) Maximum of 9 inj per prescription
- b)
- a) Not to be given in combination with a funded SGLT-2 inhibitor or other GLP-1 agonist.
- b) Maximum of 1 pack of 3 (6 mg per ml, 3 ml) prefilled pens will be funded per month.

⇒SA2339 Special Authority for Subsidy

Note: Subsidy for patients with existing approvals prior to 1 May 2024. Approvals valid without further renewal unless notified. No new patients will be granted from 1 May 2024 until further notice.

SGLT2 Inhibitors

⇒SA2408 Special Authority for Subsidy

Initial application — (heart failure reduced ejection fraction) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has heart failure: and
- 2 Patient is in NYHA functional class II or III or IV; and
- 3 Either:
 - 3.1 Patient has a documented left ventricular ejection fraction (LVEF) of less than or equal to 40%; or
 - 3.2 An ECHO is not reasonably practicable, and in the opinion of the treating practitioner the patient would benefit from treatment; and

| Subsidy | | Fully | Brand or | |
|---------------------|-----|------------|--------------|--|
| (Manufacturer's Pri | ce) | Subsidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

continued...

4 Patient is receiving concomitant optimal standard funded chronic heart failure treatment.

Initial application — (Type 2 Diabetes) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 Patient has previously received an initial approval for a GLP-1 agonist; or
- 2 All of the following:
 - 2.1 Patient has type 2 diabetes; and
 - 2.2 Any of the following:
 - 2.2.1 Patient is Māori or any Pacific ethnicity*; or
 - 2.2.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)*; or
 - 2.2.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator*; or
 - 2.2.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult*; or
 - 2.2.5 Patient has diabetic kidney disease (see note b)*; and

FMPAGLIFI OZIN - Special Authority see SA2408 on the previous page - Retail pharmacy

2.3 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of at least one blood-glucose lowering agent (e.g. metformin, vildagliptin, or insulin) for at least 3 months.

Notes: * Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

- a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack, ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.
- b) Diabetic kidney disease defined as: persistent albuminuria (albumin: creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m2 in the presence of diabetes, without alternative cause.
- c) Funded [empagliflozin / empagliflozin with metformin hydrochloride] treatment is not to be given in combination with a funded GLP-1 unless receiving (empagliflozin / empagliflozin with metformin hydrochloride] for the treatment of heart failure.

| * Tab 10 mg | | 30 | ✓ Jardiance |
|---|-------------------------|----------|------------------------------|
| * Tab 25 mg | | 30 | ✓ Jardiance |
| EMPAGLIFLOZIN WITH METFORMIN HYDROCHLORIDE - | - Special Authority see | SA2408 o | n the previous page - Retail |
| pharmacy | | | |
| * Tab 5 mg with 1,000 mg metformin hydrochloride | 58.56 | 60 | Jardiamet |
| * Tab 5 mg with 500 mg metformin hydrochloride | 58.56 | 60 | Jardiamet |
| * Tab 12.5 mg with 1,000 mg metformin hydrochloride | 58.56 | 60 | ✓ Jardiamet |
| * Tab 12.5 mg with 500 mg metformin hydrochloride | 58.56 | 60 | ✓ Jardiamet |

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer

Diabetes Management

Ketone Testing

BLOOD KETONE DIAGNOSTIC TEST STRIP - Subsidy by endorsement

- a) Not on a BSO
- b) Maximum of 20 strip per prescription
- c) Up to 10 strip available on a PSO
- d) Patient has any of the following:
 - 1) type 1 diabetes; or
 - 2) permanent neonatal diabetes: or
 - 3) undergone a pancreatectomy; or
 - 4) cystic fibrosis-related diabetes; or
 - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly.

Dual Blood Glucose and Blood Ketone Testing

DUAL BLOOD GLUCOSE AND BLOOD KETONE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A dual blood glucose and blood ketone diagnostic test meter is subsidised for a patient who has:
 - 1) type 1 diabetes; or
 - 2) permanent neonatal diabetes; or
 - 3) undergone a pancreatectomy; or
 - 4) cystic fibrosis-related diabetes; or
 - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly. Only 1 meter per patient will be subsidised (no repeat prescriptions). For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Meter with 50 lancets, a lancing device and 10 blood glucose

| | | | | _ |
|------------------------|-----|-----------|--------------|---|
| Subsidy | | Fully | Brand or | |
| (Manufacturer's Price) | S | ubsidised | Generic | |
| \$ | Per | 1 | Manufacturer | |

Blood Glucose Testing

BLOOD GLUCOSE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A diagnostic blood glucose test meter is subsidised for a patient who:
 - 1) is receiving insulin or sulphonylurea therapy; or
 - 2) is pregnant with diabetes: or
 - 3) is on home TPN at risk of hypoglycaemia or hyperglycaemia; or
 - 4) has a genetic or an acquired disorder of glucose homeostasis, excluding type 1 or type 2 diabetes and metabolic syndrome.

The prescription must be endorsed accordingly. Only one CareSens meter per patient will be subsidised (no repeat prescriptions). Patients already using the CareSens N POP meter and CareSens N meter are not eligible for a new meter, unless they have:

- 1) type 1 diabetes; or
- 2) permanent neonatal diabetes: or
- 3) undergone a pancreatectomy; or
- 4) cystic fibrosis-related diabetes.

For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Note: Only 1 meter available per PSO

BLOOD GLUCOSE DIAGNOSTIC TEST STRIP - Up to 50 test available on a PSO

The number of test strips available on a prescription is restricted to 50 unless:

- Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the
 prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed; or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly; or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

| Test strips | 50 test OP | ✓ CareSens N |
|-------------|------------|----------------|
| | | ✓ CareSens PRO |

BLOOD GLUCOSE TEST STRIPS (VISUALLY IMPAIRED)

The number of test strips available on a prescription is restricted to 50 unless:

- Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the
 prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed; or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly: or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

| Blood glucose test strips | 33.69 | 50 test OP | ✓ SensoCard |
|---------------------------|-------|------------|-------------|
|---------------------------|-------|------------|-------------|

| Subsidy | Fu | ly B | rand or |
|------------------------|-----------|------|--------------|
| (Manufacturer's Price) | Subsidise | ed G | Generic |
| \$ | Per | / N | Manufacturer |

Insulin Syringes and Needles

Subsidy is available for disposable insulin syringes, needles, and pen needles if prescribed on the same form as the one used for the supply of insulin or liraglutide or when prescribed for a patient and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of insulin or liraglutide.

| IIV | ocini i chi necocco – maximum di 200 dev pei prescripti | 1011 | | |
|-----|---|-------------------|---------------|---------------------|
| * | 29 g × 12.7 mm | 10.95 | 100 | ✓ B-D Micro-Fine |
| * | 31 g × 5 mm | | 100 | ✓ B-D Micro-Fine |
| * | 31 g × 6 mm | 9.50 | 100 | ✓ Berpu |
| * | 31 g × 8 mm | | 100 | ✓ B-D Micro-Fine |
| * | 32 g × 4 mm | | 100 | ✓ B-D Micro-Fine |
| INS | SULIN SYRINGES, DISPOSABLE WITH ATTACHED NEED | LE - Maximum of 2 | 200 dev per p | orescription |
| * | Syringe 0.3 ml with 29 g x 12.7 mm needle | 13.56 | 100 | B-D Ultra Fine |
| | | 1.36 | 10 | |
| | | (1.99) | | B-D Ultra Fine |
| * | Syringe 0.3 ml with 31 g x 8 mm needle | 13.56 | 100 | ✓ B-D Ultra Fine II |
| | , , | 1.30 | 10 | |
| | | (1.99) | | B-D Ultra Fine II |
| * | Syringe 0.5 ml with 29 g x 12.7 mm needle | 13.56 | 100 | ✓ B-D Ultra Fine |
| | , , | 1.36 | 10 | |
| | | (1.99) | | B-D Ultra Fine |
| * | Syringe 0.5 ml with 31 g x 8 mm needle | 13.56 | 100 | ✓ B-D Ultra Fine II |
| | , , | 1.36 | 10 | |
| | | (1.99) | | B-D Ultra Fine II |
| * | Syringe 1 ml with 29 g x 12.7 mm needle | 13.56 | 100 | ✓ B-D Ultra Fine |
| | , , | 1.36 | 10 | |
| | | (1.99) | | B-D Ultra Fine |
| * | Syringe 1 ml with 31 g × 8 mm needle | 13.56 | 100 | ✓ B-D Ultra Fine II |
| | | 1.36 | 10 | |
| | | (1.99) | | B-D Ultra Fine II |
| | | ` ' | | |

Insulin Pumps

INSULIN PUMP WITH ALGORITHM - Special Authority see SA2367 below - Retail pharmacy

- a) Maximum of 1 dev per prescription
- b) Only on a prescription

| c) Maximum of 1 insulin pump per patient each four year period. | | |
|---|----------|--|
| Min basal rate 0.02 U/h8,970.00 | | |
| Min boool rate 0.1 LI/b | 7 652 00 | |

✓ mylife YpsoPump with CamAPS FX

✓ Tandem t:slim

X2 with Basal-IQ

✓ Tandem t:slim
X2 with Control-IQ

⇒SA2367 Special Authority for Subsidy

Initial application — (type 1 diabetes) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

1 Any of the following:

continued...

1

| (Manufacturer's Price) Subsidised Generic \$ Per ✓ Manufacturer | Subsidy | Full | y Brand or |
|--|-----------------|--------------------|--------------|
| \$ Per ✓ Manufacturer | (Manufacturer's | s Price) Subsidise | d Generic |
| | \$ | Per ✓ | Manufacturer |

continued...

- 1.1 The patient has type 1 diabetes; or
- 1.2 The patient has permanent neonatal diabetes or specific monogenic diabetes subtypes with insulin deficiency, considered by the treating endocrinologist as likely to benefit; or
- 1.3 The patient has Type 3c diabetes considered by the treating endocrinologist as likely to benefit (Type 3c diabetes includes insulin deficiency due to pancreatectomy, insulin deficiency secondary to cystic fibrosis or pancreatitis); or
- 1.4 The patient has atypical inherited forms of diabetes; and
- 2 Patient has been evaluated by a diabetes multidisciplinary team for their suitability for insulin pump therapy; and
- 3 In the opinion of the treating relevant practitioner the patient would benefit from an Automated Insulin Delivery (AID) system.

Renewal — (type 1 diabetes) from any relevant practitioner. Approvals valid for 6 months where the patient is continuing to derive benefit according to the treatment plan agreed at induction.

Insulin Pump Consumables

⇒SA2380 Special Authority for Subsidy

Initial application — (type 1 diabetes) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has type 1 diabetes; or
 - 1.2 The patient has permanent neonatal diabetes or specific monogenic diabetes subtypes with insulin deficiency, considered by the treating endocrinologist as likely to benefit; or
 - 1.3 The patient has Type 3c diabetes considered by the treating endocrinologist as likely to benefit (Type 3c diabetes includes insulin deficiency due to pancreatectomy, insulin deficiency secondary to cystic fibrosis or pancreatitis); or
 - 1.4 The patient has atypical inherited forms of diabetes; and
- 2 Patient has been evaluated by a diabetes multidisciplinary team for their suitability for insulin pump therapy; and
- 3 In the opinion of the treating relevant practitioner the patient would benefit from an Automated Insulin Delivery (AID) system.

Renewal — (type 1 diabetes) from any relevant practitioner. Approvals valid for 2 years where the patient is continuing to derive benefit according to the treatment plan agreed at induction.

INSULIN PUMP CARTRIDGE - Special Authority see SA2380 above - Retail pharmacy

- a) Maximum of 5 sets per prescription
- b) Only on a prescription
- c) Maximum of 19 packs of cartridge sets will be funded per year.
- **★** Cartridge 300 U, t:lock × 10......86.00 1 OP **✓ Tandem Cartridge**

| | | Subsidy (Manufacturer's Price) | Per | Fully Subsidised | Brand or Generic Manufacturer |
|-----|---|--|----------------|---------------------|-------------------------------------|
| INS | SULIN PUMP INFUSION SET (STEEL CANNULA) - Special | • | | ne previous | |
| | a) Maximum of 5 set per prescription | · | | · | |
| | b) Only on a prescription | | | | |
| * | c) Maximum of 19 infusion sets will be funded per year. 6 mm steel needle; 60 cm tubing × 10 | 120.00 | 1 OP | 1 | MiniMed Sure-T |
| ~ | o min steer needie, oo cin tubing x 10 | 150.00 | 1 01 | • | MMT-864A |
| * | 6 mm steel needle; 80 cm tubing × 10 | 130.00 | 1 OP | 1 | MiniMed Sure-T |
| | • | | | | MMT-866A |
| * | 8 mm steel needle; 60 cm tubing × 10 | 130.00 | 1 OP | / | MiniMed Sure-T MMT-874A |
| * | 8 mm steel needle; 80 cm tubing × 10 | 130.00 | 1 OP | • | MiniMed Sure-T MMT-876A |
| (Mi | niMed Sure-T MMT-866A 6 mm steel needle; 80 cm tubing x niMed Sure-T MMT-874A 8 mm steel needle; 60 cm tubing x niMed Sure-T MMT-876A 8 mm steel needle; 80 cm tubing x | 10 to be delisted 1 O 10 to be delisted 1 O | ctobe ctobe | r 2026) r 2026) | a CACCO on the provious |
| | SULIN PUMP INFUSION SET (STEEL CANNULA, STRAIGHT le – Retail pharmacy | INSERTION) – Spe | ecial A | uthority se | e SA2380 on the previous |
| μαί | a) Maximum of 5 sets per prescription | | | | |
| | b) Only on a prescription | | | | |
| | c) Maximum of 19 infusion sets will be funded per year. | | | | |
| | 5.5 mm steel cannula; straight insertion; 45 cm line × 10 with | | 4 00 | | |
| | 10 needles | 136.00 | 1 OP | • | mylife Orbit micro |
| | 5.5 mm steel needle; straight insertion; 60 cm line × 10 with 10 needles | 136 00 | 1 OP | 1 | mylife Orbit micro |
| * | 5.5 mm steel needle; straight insertion; 80 cm line × 10 with | | . 0. | - | mymo orbit mioro |
| | 10 needles | 136.00 | 1 OP | 1 | mylife Orbit micro |
| * | 8.5 mm steel needle; straight insertion; 60 cm line \times 10 with | | | _ | |
| | 10 needles | 136.00 | 1 OP | • | mylife Orbit micro |
| * | 8.5 mm steel needle; straight insertion; 80 cm line × 10 with 10 needles | 136.00 | 1 OP | 1 | mylife Orbit micro |

* 6 mm steel cannula; straight insertion; 80 cm line x 10 with

* 8 mm steel cannula; straight insertion; 80 cm line × 10 with

* 6 mm steel cannula; straight insertion; 60 cm line x 10 with

* 8 mm steel cannula; straight insertion; 60 cm line x 10 with

1 OP

1 OP

1 OP

1 OP

✓ TruSteel

✓ TruSteel

✓ TruSteel

✓ TruSteel

| | | Subsidy | | Fully | Brand or |
|-----|---|------------------------------|-------|------------|-------------------------------|
| | | (Manufacturer's Price) \$ | Per | Subsidised | Generic Manufacturer |
| _ | | T | | | |
| INS | SULIN PUMP INFUSION SET (TEFLON CANNULA) – Specia | I Authority see SA23 | 80 on | page 17 – | Retail pharmacy |
| | a) Maximum of 5 set per prescription | | | | |
| | b) Only on a prescription | | | | |
| * | c) Maximum of 19 infusion sets will be funded per year. 13 mm teflon needle, 60 cm tubing × 10 | 130.00 | 1 OP | 1 | MiniMed Silhouette |
| ~ | To minitenon needle, oo em tabing x to | 100.00 | 1 01 | • | MMT-381A |
| * | 17 mm teflon needle, 110 cm tubing × 10 | 130.00 | 1 OP | 1 | MiniMed Silhouette |
| • | , , , , , , , , , , , , , , , , , , , | | | | MMT-377A |
| * | 17 mm teflon needle, 60 cm tubing × 10 | 130.00 | 1 OP | 1 | MiniMed Silhouette |
| | - | | | | MMT-378A |
| * | 6 mm teflon needle, 110 cm tubing × 10 | 130.00 | 1 OP | ✓ | MiniMed Quick-Set |
| | | | | | MMT-398A |
| * | 6 mm teflon needle, 45 cm blue tubing \times 10 | 130.00 | 1 OP | • | MiniMed Mio |
| | | | | | MMT-941A |
| * | 6 mm teflon needle, 45 cm pink tubing × 10 | 130.00 | 1 OP | • | MiniMed Mio MMT-921A |
| * | 6 mm toflan needle 60 em blue tubing 10 | 100.00 | 1 OP | ./ | MiniMed Mio |
| 不 | 6 mm teflon needle, 60 cm blue tubing x 10 | 130.00 | I OF | • | MMT-943A |
| * | 6 mm teflon needle, 60 cm pink tubing × 10 | 130.00 | 1 OP | 1 | MiniMed Mio |
| | o him tollon noodio, oo oni piim tabilig x ro | | . 0. | | MMT-923A |
| * | 6 mm teflon needle, 60 cm tubing × 10 | 130.00 | 1 OP | 1 | MiniMed Quick-Set |
| | · | | | | MMT-399A |
| * | 6 mm teflon needle, 80 cm blue tubing | 130.00 | 1 OP | ✓ | MiniMed Mio |
| | | | | | MMT-945A |
| * | 6 mm teflon needle, 80 cm clear tubing × 10 | 130.00 | 1 OP | ✓ | MiniMed Mio |
| | | | | | MMT-965A |
| * | 6 mm teflon needle, 80 cm pink tubing x 10 | 130.00 | 1 OP | • | MiniMed Mio |
| * | 0 mm toflan needle 110 em tubing 10 | 100.00 | 1 OP | ./ | MMT-925A MiniMed Quick-Set |
| 不 | 9 mm teflon needle, 110 cm tubing × 10 | 130.00 | I OF | • | MMT-396A |
| * | 9 mm teflon needle, 60 cm tubing × 10 | 130.00 | 1 OP | 1 | MiniMed Quick-Set |
| -,- | o min tonon noodio, oo om tabing x 10 | | . 0. | • | MMT-397A |
| * | 9 mm teflon needle, 80 cm clear tubing × 10 | 130.00 | 1 OP | 1 | MiniMed Mio |
| | , | | | | MMT-975A |
| | | | | | |

(MiniMed Silhouette MMT-381A 13 mm teflon needle, 60 cm tubing × 10 to be delisted 1 October 2026) (MiniMed Silhouette MMT-377A 17 mm teflon needle, 110 cm tubing × 10 to be delisted 1 October 2026) (MiniMed Silhouette MMT-378A 17 mm teflon needle, 60 cm tubing × 10 to be delisted 1 October 2026) (MiniMed Quick-Set MMT-398A 6 mm teflon needle, 110 cm tubing × 10 to be delisted 1 October 2026) (MiniMed Mio MMT-941A 6 mm teflon needle, 45 cm blue tubing × 10 to be delisted 1 October 2026) (MiniMed Mio MMT-921A 6 mm teflon needle, 45 cm pink tubing × 10 to be delisted 1 October 2026) (MiniMed Mio MMT-943A 6 mm teflon needle, 60 cm blue tubing × 10 to be delisted 1 October 2026) (MiniMed Mio MMT-933A 6 mm teflon needle, 60 cm pink tubing × 10 to be delisted 1 October 2026) (MiniMed Mio MMT-945A 6 mm teflon needle, 60 cm blue tubing × 10 to be delisted 1 October 2026) (MiniMed Mio MMT-945A 6 mm teflon needle, 80 cm blue tubing to be delisted 1 October 2026) (MiniMed Mio MMT-965A 6 mm teflon needle, 80 cm clear tubing × 10 to be delisted 1 October 2026) (MiniMed Mio MMT-925A 6 mm teflon needle, 80 cm clear tubing × 10 to be delisted 1 October 2026) (MiniMed Mio MMT-936A 9 mm teflon needle, 110 cm tubing × 10 to be delisted 1 October 2026) (MiniMed Quick-Set MMT-396A 9 mm teflon needle, 60 cm tubing × 10 to be delisted 1 October 2026) (MiniMed Quick-Set MMT-397A 9 mm teflon needle, 60 cm tubing × 10 to be delisted 1 October 2026) (MiniMed Mio MMT-975A 9 mm teflon needle, 80 cm clear tubing × 10 to be delisted 1 October 2026)

| | | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer |
|------|--|---|--------|---------------------|-------------------------------------|
| INS | SULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE IN | ISERTION WITH INS | SERT | ON DEVICE | Special Authority see |
| | 2380 on page 17 – Retail pharmacy | | | | , , |
| | a) Maximum of 5 sets per prescription | | | | |
| | b) Only on a prescription | | | | |
| | c) Maximum of 19 infusion sets will be funded per year. | | | | |
| * | 13 mm teflon cannula; angle insertion; insertion device; 110 c | m | | | |
| | line × 10 with 10 needles | 182.00 | 1 OP | ✓ A | utoSoft 30 |
| * | 13 mm teflon cannula; angle insertion; insertion device; 60 cn line × 10 with 10 needles | n | 1 OP | ✓ A | utoSoft 30 |
| INIS | SULIN PUMP INFUSION SET (TEFLON CANNULA, FLEXIBLE | INSERTION WITH | INSFI | RTION DEV | ICE) - Special Authority |
| | SA2380 on page 17 – Retail pharmacy | INOLITION WITH | IIVOLI | THONDEV | opecial Authority |
| | a) Maximum of 5 set per prescription | | | | |
| | b) Only on a prescription | | | | |
| | c) Maximum of 19 infusion sets will be funded per year. | | | | |
| * | 6 mm teflon cannula; flexible insertion; insertion device; 46 cr | n | | | |
| • | line × 10 with 10 needles | | 1 OP | ✓ m | ylife Inset soft |
| * | 6 mm teflon cannula; flexible insertion; insertion device; 60 cr | n | | | • |
| • | line with integrated inserter × 10 with 10 needles | | 1 OP | ✓ m | ylife Inset soft |
| * | 6 mm teflon cannula; flexible insertion; insertion device; 80 cr | | | | , |
| • | line × 10 with 10 needles | | 1 OP | ✓ m | ylife Inset soft |
| * | 9 mm teflon cannula; flexible insertion; insertion device; 60 cr | | | | , |
| • | line × 10 with 10 needles | | 1 OP | ✓ m | ylife Inset soft |
| * | 9 mm teflon cannula; flexible insertion; insertion device; 80 cr | | | | , |
| • | line × 10 with 10 needles | | 1 OP | ✓ m | ylife Inset soft |
| INI | SULIN PUMP INFUSION SET (TEFLON CANNULA, STRAIGH | | INICE | | • |
| | e SA2380 on page 17 – Retail pharmacy | I INSLITTION WITH | IIVOL | ITTION DE | riot) - Special Authority |
| 500 | a) Maximum of 5 sets per prescription | | | | |
| | b) Only on a prescription | | | | |
| | c) Maximum of 19 infusion sets will be funded per year. | | | | |
| * | 6 mm teflon cannula; straight insertion; insertion device; | | | | |
| ••• | 110 cm line × 10 with 10 needles | 182.00 | 1 OP | ✓ A | utoSoft 90 |
| * | 6 mm teflon cannula; straight insertion; insertion device; 60 ci | | | | |
| -,- | line × 10 with 10 needles | | 1 OP | ✓ Δ | utoSoft 90 |
| * | 9 mm teflon cannula; straight insertion; insertion device; | | | - 70 | 4.000.1.00 |
| -,- | 110 cm line × 10 with 10 needles | 182 00 | 1 OP | ✓ A | utoSoft 90 |
| * | 9 mm teflon cannula; straight insertion; insertion device; 60 cr | | | - 70 | 4.0001.00 |
| -,- | line × 10 with 10 needles | | 1 OP | ✓ Δ | utoSoft 90 |
| INIC | | | . •. | | |
| | SULIN PUMP INFUSION SET (TEFLON CANNULA, VARIABLI tail pharmacy | E INSERTION) - SP | eciai | Authority See | e SA2360 on page 17 - |
| ne | a) Maximum of 5 set per prescription | | | | |
| | b) Only on a prescription | | | | |
| | c) Maximum of 19 infusion sets will be funded per year. | | | | |
| * | 13 mm teflon cannula; variable insertion; 60 cm line × 10 with | | | | |
| * | 10 needles | | 1 OP | ✓ V | ariSoft |
| | TO TICCUICS | 102.00 | 1 01 | - V | anoon |
| | | | | | |

MMT-332A

| | Subsidy (Manufacturer's Price) \$ | Su Per | Fully Brand or ibsidised Generic Manufacture | r |
|--|---|-----------|--|-------|
| INSULIN PUMP RESERVOIR - Special Authority see SA2380 | on page 17 – Retail ph | armacy | | |
| a) Maximum of 9 sets per prescription | | | | |
| b) Only on a prescription | | | | |
| c) Maximum of 36 packs of resevoir sets will be funded per | , | | | |
| * 10 × 1.6 ml glass reservoir for YpsoPump | 50.00 | 1 OP | ✓ mylife YpsoP Reservoir | ump |
| * 10 × luer lock conversion cartridges 1.8 ml for Paradigm p | umps50.00 | 1 OP | ✓ ADR Cartridg | e 1.8 |
| * Cartridge for 7 series pump; 3.0 ml x 10 | 50.00 | 1 OP | ✓ MiniMed | |
| | | | 3.0 Reservo | oir |

(ADR Cartridge 1.8 10 × luer lock conversion cartridges 1.8 ml for Paradigm pumps to be delisted 1 October 2026) (MiniMed 3.0 Reservoir MMT-332A Cartridge for 7 series pump; 3.0 ml × 10 to be delisted 1 October 2026)

Continuous Glucose Monitor

CONTINUOUS GLUCOSE MONITOR (INTEROPERABLE) – Special Authority see SA2371 below – Retail pharmacy Only on a prescription

Maximum of 28 dev will be funded per year.

⇒SA2371 Special Authority for Subsidy

Initial application — (type 1 diabetes) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has type 1 diabetes; or
 - 1.2 The patient has permanent neonatal diabetes or specific monogenic diabetes subtypes with insulin deficiency, considered by the treating endocrinologist or relevant secondary health care professional as practicable, as likely to benefit; or
 - 1.3 The patient has Type 3c diabetes considered by the treating endocrinologist or relevant secondary health care professional as practicable, as likely to benefit (Type 3c diabetes includes insulin deficiency due to pancreatectomy, insulin deficiency secondary to cystic fibrosis or pancreatitis); or
 - 1.4 The patient has atypical inherited forms of diabetes; and
- 2 In the opinion of the treating relevant practitioner the patient would benefit from an Automated Insulin Delivery (AID) system.

Renewal — **(type 1 diabetes)** from any relevant practitioner. Approvals valid for 2 years where the patient is continuing to derive benefit according to the treatment plan agreed at induction.

CONTINUOUS GLUCOSE MONITOR (STANDALONE) - Special Authority see SA2370 on the next page - Retail pharmacy Only on a prescription

★ Sensor (Dexcom ONE+) - Maximum of 9 dev per prescription81.00
 Maximum of 40 dev will be funded per year.

| Subsid | idy Fully | Brand or |
|--------------|-----------|--------------|
| (Manufacture | | |
| \$ | Per 🗸 | Manufacturer |

⇒SA2370 Special Authority for Subsidy

Initial application — (type 1 diabetes) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 The patient has type 1 diabetes; or
- 2 The patient has permanent neonatal diabetes or specific monogenic diabetes subtypes with insulin deficiency, considered by the treating endocrinologist or relevant secondary health care professional as practicable, as likely to benefit; or
- 3 The patient has Type 3c diabetes considered by the treating endocrinologist or relevant secondary health care professional as practicable, as likely to benefit (Type 3c diabetes includes insulin deficiency due to pancreatectomy, insulin deficiency secondary to cystic fibrosis or pancreatitis); or
- 4 The patient has atypical inherited forms of diabetes.

Renewal — (type 1 diabetes) from any relevant practitioner. Approvals valid for 2 years where the patient is continuing to derive benefit according to the treatment plan agreed at induction.

Digestives Including Enzymes

PANCREATIC ENZYME

| Cap pancreatin 150 mg (amylase 8,000 Ph Eur U, lipase | | |
|--|---------|------------------|
| 10,000 Ph Eur U, total protease 600 Ph Eur U)34.93 | 100 | ✓ Creon 10000 |
| Cap pancreatin 300 mg (amylase 18,000 Ph Eur U, lipase | | |
| 25,000 Ph Eur U, total protease 1,000 Ph Eur U)94.38 | 100 | ✓ Creon 25000 |
| Modified release granules pancreatin 60.12 mg (amylase | | |
| 3,600 Ph Eur U, lipase 5,000 Ph Eur U, protease 200 Ph | | |
| Eur U) | 20 g OP | Creon Micro |
| URSODEOXYCHOLIC ACID - Special Authority see SA1739 below - Retail pharm | nacy | |
| Cap 250 mg33.95 | 100 | ✓ <u>Ursosan</u> |

⇒SA1739 Special Authority for Subsidy

Initial application — (Alagille syndrome or progressive familial intrahepatic cholestasis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 Patient has been diagnosed with Alagille syndrome: or
- 2 Patient has progressive familial intrahepatic cholestasis.

Initial application — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic severe drug induced cholestatic liver injury; and
- 2 Cholestatic liver injury not due to Total Parenteral Nutrition (TPN) use in adults; and
- 3 Treatment with ursodeoxycholic acid may prevent hospital admission or reduce duration of stay.

Initial application — (Primary biliary cholangitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Primary biliary cholangitis confirmed by antimitochondrial antibody titre (AMA) > 1:80, and raised cholestatic liver enzymes with or without raised serum IgM or, if AMA is negative, by liver biopsy; and
- 2 Patient not requiring a liver transplant (bilirubin > 100 umol/l; decompensated cirrhosis).

Initial application — (Pregnancy) from any relevant practitioner. Approvals valid for 6 months where the patient diagnosed with cholestasis of pregnancy.

Initial application — (Haematological Transplant) from any relevant practitioner. Approvals valid for 6 months for applications

ColoxvI

| Subsidy | e) | Fully | Brand or |
|----------------------|-----|------------|--------------|
| (Manufacturer's Pric | | Subsidised | Generic |
| \$ | Per | 1 | Manufacturer |

continued...

meeting the following criteria:

Both:

- 1 Patient at risk of veno-occlusive disease or has hepatic impairment and is undergoing conditioning treatment prior to allogenic stem cell or bone marrow transplantation; and
- 2 Treatment for up to 13 weeks.

Initial application — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Paediatric patient has developed abnormal liver function as indicated on testing which is likely to be induced by Total Parenteral Nutrition (TPN); and
- 2 Liver function has not improved with modifying the TPN composition.

Renewal — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 6 months where the patient continues to benefit from treatment.

Renewal — (Pregnancy/Primary biliary cholangitis) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months where the paediatric patient continues to require TPN and who is benefiting from treatment, defined as a sustained improvement in bilirubin levels

Laxatives

Bulk-forming Agents

| ISPAGHULA (PSYLLIUM) HUSK – Only on a prescription | | | |
|--|-------|----------|---------|
| W Douglas for oral calm | 20.00 | E00 ~ OD | ./ Vana |

| * Powder for oral soln | 20.00 | 500 g OP | ✓ Konsyl-D |
|------------------------|-------|----------|------------|
| Faecal Softeners | | | |

| DO | CUSATE SODIUM - Only on a prescription | | |
|----|--|-----|-----|
| * | Tab 50 mg | 100 | ✓ (|

Tab 50 mg with sennosides 8 mg.......3.50 200 ✓ Laxsol POLOXAMER – Only on a prescription

Opioid Receptor Antagonists - Peripheral

METHYLNALTREXONE BROMIDE - Special Authority see SA1691 below - Retail pharmacy

⇒SA1691 Special Authority for Subsidy

Initial application — (Opioid induced constipation) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient is receiving palliative care; and
- 2 Fither:
 - 2.1 Oral and rectal treatments for opioid induced constipation are ineffective; or
 - 2.2 Oral and rectal treatments for opioid induced constipation are unable to be tolerated.

| | Subsidy | | Fully | Brand or |
|---|------------------------|------------------------|----------------|---|
| | (Manufacturer's Price) |) S Per | ubsidised • | Generic Manufacturer |
| Osmotic Laxatives | | | | |
| GLYCEROL * Suppos 2.8/4.0 g - Only on a prescription | 10.39 | 20 | 1 | Lax-suppositories Glycerol |
| LACTULOSE – Only on a prescription * Oral liq 10 g per 15 ml | 3.61 | 500 ml | • | <u>Laevolac</u> |
| MACROGOL 3350 WITH POTASSIUM CHLORIDE, SODIUM BIO | | SODIUN | I CHLOF | RIDE |
| Powder for oral soln 13.125 g with potassium chloride 46.6 m sodium bicarbonate 178.5 mg and sodium chloride 350.7 | • | 30 | ✓ | <u>Molaxole</u> |
| SODIUM ACID PHOSPHATE – Only on a prescription Enema 16% with sodium phosphate 8% | 2.50 | 1 | • | Fleet Phosphate Enema |
| SODIUM CITRATE WITH SODIUM LAURYL SULPHOACETATE | | iption | | |
| Enema 90 mg with sodium lauryl sulphoacetate 9 mg per ml, 5 ml | | 50 | ✓ | Micolette |
| Stimulant Laxatives | | | | |
| BISACODYL – Only on a prescription * Tab 5 mg * Suppos 10 mg SENNA – Only on a prescription * Tab, standardised | | 200 10 100 20 | • | Bisacodyl Viatris Lax-Suppositories Senokot |
| SODIUM PICOSULFATE – Special Authority see SA2053 below Oral soln 7.5 mg per ml | | 0 ml OF | | Senokot Dulcolax SP Drop |
| | | · ···· • · | | |

⇒SA2053 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The patient is a child with problematic constipation despite an adequate trial of other oral pharmacotherapies including macrogol where practicable; and
- 2 The patient would otherwise require a high-volume bowel cleansing preparation or hospital admission.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Metabolic Disorder Agents

⇒SA1986 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

1 The patient is aged up to 24 months at the time of initial application and has been diagnosed with infantile Pompe disease; and

|--|

continued...

- 2 Any of the following:
 - 2.1 Diagnosis confirmed by documented deficiency of acid alpha-glucosidase by prenatal diagnosis using chorionic villus biopsies and/or cultured amniotic cells; or
 - 2.2 Documented deficiency of acid alpha-glucosidase, and urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides; or
 - 2.3 Documented deficiency of acid alpha-glucosidase, and documented molecular genetic testing indicating a disease-causing mutation in the acid alpha-glucosidase gene (GAA gene); or
 - 2.4 Documented urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides, and molecular genetic testing indicating a disease-causing mutation in the GAA gene; and
 - 3 Patient has not required long-term invasive ventilation for respiratory failure prior to starting enzyme replacement therapy (ERT); and
 - 4 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by ERT or might be reasonably expected to compromise a response to ERT; and
 - 5 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks.

Renewal only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks; and
- 3 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 4 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by ERT: and
- 5 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT: and
- 6 There is no evidence of life threatening progression of respiratory disease as evidenced by the needed for > 14 days of invasive ventilation; and
- 7 There is no evidence of new or progressive cardiomyopathy.

ARGININE - Special Authority see SA2042 below - Retail pharmacy

| Tab 1,000 mg | CBS | 90 | Clinicians |
|--------------|-----|-------|------------|
| Cap 500 mg | CBS | 50 | ✓ Solgar |
| Powder | | 400 g | ✓ Biomed |

⇒SA2042 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to arginine supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to arginine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

⇒SA1987 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has a confirmed diagnosis of homocystinuria; and
- 2 Any of the following:
 - 2.1 A cystathionine beta-synthase (CBS) deficiency; or

| | | | |
|-------------|------------------------|------------|--------------|
| | Subsidy | Fully | Brand or |
| | (Manufacturer's Price) | Subsidised | Generic |
| | \$ | Por 🗸 | Manufacturer |

continued...

- 2.2 A 5,10-methylene-tetrahydrofolate reductase (MTHFR) deficiency; or
- 2.3 A disorder of intracellular cobalamin metabolism; and
- 3 An appropriate homocysteine level has not been achieved despite a sufficient trial of appropriate vitamin supplementation.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

COENZYME Q10 - Special Authority see SA2039 below - Retail pharmacy

| Cap 120 mg | CBS | 30 | Solgar |
|------------|-----|----|------------|
| Cap 160 mg | CBS | 60 | Go Healthy |

⇒SA2039 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to coenzyme Q10 supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to coenzyme Q10 supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

GALSULFASE − Special Authority see SA1988 below − Retail pharmacy
Inj 1 mg per ml, 5 ml vial......2,234.00 1 ✓ Naglazyme

⇒SA1988 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has been diagnosed with mucopolysaccharidosis VI; and
- 2 Fither:
 - 2.1 Diagnosis confirmed by demonstration of N-acetyl-galactosamine-4-sulfatase (arylsulfatase B) deficiency by either enzyme activity assay in leukocytes or skin fibroblasts; or
 - 2.2 Detection of two disease causing mutations and patient has a sibling who is known to have mucopolysaccharidosis VI

Renewal only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 3 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by Enzyme Replacement Therapy (ERT); and
- 4 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT.

⇒SA1623 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hunter Syndrome (mucopolysaccharidosis II); and
- 2 Either:
 - 2.1 Diagnosis confirmed by demonstration of iduronate 2-sulfatase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts: or

| | Subsidy (Manufacturer's Price) \$ | F Subsidis Per | ully sed | Brand or Generic Manufacturer |
|-----------|---|----------------------|-------------|-------------------------------------|
| continued | | | | |

- 2.2 Detection of a disease causing mutation in the iduronate 2-sulfatase gene; and
 - 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with idursulfase would be bridging treatment to transplant; and
 - 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT); and
 - 5 Idursulfase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 weeks post-HSCT) at doses no greater than 0.5 mg/kg every week.

| LARONIDASE - Special Authority see SA1695 below | - Retail pharmacy | | |
|---|-------------------|---|------------|
| Inj 100 U per ml, 5 ml vial | 1,335.16 | 1 | Aldurazyme |

⇒SA1695 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hurler Syndrome (mucopolysacchardosis I-H); and
- 2 Fither:

L

- 2.1 Diagnosis confirmed by demonstration of alpha-L-iduronidase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts; or
- 2.2 Detection of two disease causing mutations in the alpha-L-iduronidase gene and patient has a sibling who is known to have Hurler syndrome; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with laronidase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT): and
- 5 Laronidase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 post-HSCT) at doses no greater than 100 units/kg every week.

| LEVOCARNITINE - Special Authority see SA204 | 10 below - Retail pharmacy | | |
|---|----------------------------|--------|------------------|
| Tab 500 mg | CBS | 30 | ✓ Solgar |
| Cap 250 mg | CBS | 30 | ✓ Solgar |
| Cap 500 mg | | 60 | ✓ Balance |
| | | 300 | ✓ Metabolics |
| Oral lig 1 g per 10 ml | CBS | 118 ml | ✓ Carnitor S29 |
| 1 31 | | | ✓ Novitium Sugar |
| | | | Free S29 |
| Oral lig 500 mg per 10 ml | CBS | 300 ml | ✓ Balance |

⇒SA2040 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to carnitine supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to carnitine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

| RIBOFLAVIN – Special Authority see SA2041 on the next Tab 100 mg | , , | 100 | ✓ Country Life ✓ Puritan's Pride |
|--|-----|-----|----------------------------------|
| | | | Vitamin B-2 100 mg S29 |
| Cap 100 mg | CBS | 100 | ✓ Solgar |

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA2041 Special Authority for Subsidy

Initial application only from a metabolic physician or neurologist. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to riboflavin supplementation.

Renewal only from a metabolic physician or neurologist. Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to riboflavin supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

SAPROPTERIN DIHYDROCHLORIDE - Special Authority see SA1989 below - Retail pharmacy

⇒SA1989 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 1 month for applications meeting the following criteria: All of the following:

- 1 Patient has phenylketonuria (PKU) and is pregnant or actively planning to become pregnant; and
- 2 Treatment with sapropterin is required to support management of PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

Renewal only from a metabolic physician or any relevant practitioner on the recommendation of a metabolic physician.

Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Following the initial one-month approval, the patient has demonstrated an adequate response to a 2 to 4 week trial of sapropterin with a clinically appropriate reduction in phenylalanine levels to support management of PKU during pregnancy; or
 - 1.2 On subsequent renewal applications, the patient has previously demonstrated response to treatment with sapropterin and maintained adequate phenylalanine levels to support management of PKU during pregnancy; and
- 2 Any of the following:
 - 2.1 Patient continues to be pregnant and treatment with sapropterin will not continue after delivery; or
 - 2.2 Patient is actively planning a pregnancy and this is the first renewal for treatment with sapropterin; or
 - 2.3 Treatment with sapropterin is required for a second or subsequent pregnancy to support management of their PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

SODIUM BENZOATE – Special Authority see SA1599 below – Retail pharmacy
Soln 100 mg per mlCBS

SODIUM BENZOATE – Special Authority see SA1599 below – Retail pharmacy

⇒SA1599 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

SODIUM PHENYLBUTYRATE - Special Authority see SA1990 on the next page - Retail pharmacy

Grans 483 mg per g......2,016.00 174 g OP ✓ Pheburane

100 ml

✓ Amzoate S29

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

⇒SA1990 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder involving a deficiency of carbamylphosphate synthetase, ornithine transcarbamylase or argininosuccinate synthetase.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

TAURINE - Special Authority see SA2043 below - Retail pharmacy

| Cap 500 mg | CBS | 50 | ✓ Solgar |
|--------------|-----|-------|------------------------------------|
| Cap 1,000 mg | CBS | 90 | ✓ Life Extension |
| Powder | CBS | 300 g | Life Extension |

⇒SA2043 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected specific mitochondrial disorder that may respond taurine supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has confirmed diagnosis of a specific mitochondrial disorder which responds to taurine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

TRIENTINE - Special Authority see SA2324 below - Retail pharmacy Cap 250 mg......2,022.00 100 **Trientine Waymade**

⇒SA2324 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Wilson disease: and
- 2 Treatment with D-penicillamine has been trialled and discontinued because the person has experienced intolerable side effects or has not received sufficient benefit; and
- 3 Treatment with zinc has been trailled and discontinued because the person has experienced intolerable side effects or has not received sufficient benefit, or zinc is considered clinically inappropriate as the person has symptomatic liver disease and requires copper chelation.

Gaucher's Disease

TALIGLUCERASE ALFA - Special Authority see SA2137 below - Retail pharmacy ✓ Elelvso

⇒SA2137 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has a diagnosis of symptomatic type 1 or type 3* Gaucher disease confirmed by the demonstration of specific deficiency of glucocerebrosidase in leukocytes or cultured skin fibroblasts, and genotypic analysis; and
- 2 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by enzyme replacement therapy (ERT) or the disease might be reasonably expected to compromise a response to ERT; and
- 3 Any of the following:
 - 3.1 Patient has haematological complications of Gaucher disease; or
 - 3.2 Patient has skeletal complications of Gaucher disease; or
 - 3.3 Patient has significant liver dysfunction or hepatomegaly attributable to Gaucher disease; or
 - 3.4 Patient has reduced vital capacity from clinically significant or progressive pulmonary disease due to Gaucher

| | Subsidy (Manufacturer's P \$ | rice) Sul Per | Fully osidised | Brand or Generic Manufacturer |
|---|------------------------------------|------------------|----------------|-------------------------------------|
| continued | | | | |
| disease; or | | | | |
| 3.5 Patient is a child and has experienced growth failt6-12 month period; and | ıre with significan | t decrease in | percenti | ile linear growth over a |
| 4 Taliglucerase alfa is to be administered at a dose no great whole vial (200 units). | ater than 30 unit/k | g every other | week ro | ounded to the nearest |
| Note: Indication marked with * is an unapproved indication | | | | |
| Renewal only from a metabolic physician or any relevant practiti | | nmendation of | of a meta | ıbolic physician. |
| Approvals valid for 3 years for applications meeting the following All of the following: | | | | |
| Patient has demonstrated a symptomatic improvement as symptoms for which therapy was started; and Patient has demonstrated a linically chiesting improvement. | | · | | • • |
| 2 Patient has demonstrated a clinically objective improvem liver and spleen size; and | ent or no detenor | alion in naen | logiobiri | ieveis, piateiet courits and |
| 3 Radiological (MRI) signs of bone activity performed at two demonstrate no deterioration shown by the MRI, compare | | | | |
| or adjusted dose; and 4 Patient has not developed another medical condition that | might reasonably | be expected | I to comp | promise a response to |
| ERT; and | | | | |
| 5 Patient is adherent with regular treatment and taliglucera: every other week rounded to the nearest whole vial (200 | | ministered at | a dose i | no greater than 30 unit/kg |
| Mouth and Throat | | | | |
| Agents Used in Mouth Ulceration | | | | |
| BENZYDAMINE HYDROCHLORIDE | | | | |
| Soln 0.15% - Higher subsidy of \$22.60 per 500 ml with | | | | |
| Endorsement | 9.00 (22.60) | 500 ml | | Difflam |
| Additional subsidy by endorsement for a patient who ha prescription is endorsed accordingly. | s oral mucositis a | s a result of t | reatmen | t for cancer, and the |
| CARMELLOSE SODIUM WITH GELATIN AND PECTIN | | | _ | |
| Paste | | 56 g OP | √ S | Stomahesive |
| | 4.55 | 15 g OP | _ | North and |
| | (7.90) | E a OB | C | Drabase |
| | 1.52 (3.60) | 5 g OP | (| Drabase |
| Powder | | 28 g OP | | 7140436 |
| | (10.95) | _0 g O | S | Stomahesive |
| TRIAMCINOLONE ACETONIDE | • | | | |
| Paste 0.1% | 5.49 | 5 g OP | ✓ <u>K</u> | Kenalog in Orabase |
| Oropharyngeal Anti-infectives | | | | |
| AMPHOTERICIN B | | | | |
| Lozenges 10 mg | | 20 | | ungilin |

| MICONAZOLE |
|----------------------|
| Oral gel 20 mg per g |
| NYSTATIN |

40 g OP

.....5.19

✓ Decozol

Oral liq 100,000 u per ml2.22

24 ml OP

✓ Nilstat

| | Subsidy (Manufacturer's Price \$ |) Su Per | Fully bsidised | Brand or Generic Manufacturer |
|---|--|-------------|-------------------|---|
| Vitamins | | | | |
| Vitamin B | | | | |
| HYDROXOCOBALAMIN | | | | |
| Inj 1 mg per ml, 1 ml ampoule – Up to 6 inj available on a F | PSO2.46 | 3 | | cobal-B12 ©29 Vita-B12 |
| | 3.95 | | √ } | lydroxocobalamin Panpharma |
| | 4.10 | 5 | _ | cobalin-H S29 leo-Cytamen S29 S29 |
| | 8.20 | 10 | ✓ \ | itarubin Depot Injection S29 |
| (Cobal-B12 S29 Inj 1 mg per ml, 1 ml ampoule to be delisted 1 (Vita-B12 Inj 1 mg per ml, 1 ml ampoule to be delisted 1 July 20 (Cobalin-H S29 Inj 1 mg per ml, 1 ml ampoule to be delisted 1 (Neo-Cytamen S29 S29 Inj 1 mg per ml, 1 ml ampoule to be delisted 1 (Vitarubin Depot Injection S29 Inj 1 mg per ml, 1 ml ampoule to be de (Vitarubin Depot Injection S29 Inj 1 mg per ml, 1 ml ampoule to PYRIDOXINE HYDROCHLORIDE a) No more than 100 mg per dose b) Only on a prescription * Tab 25 mg — No patient co-payment payable | 025) July 2025) elisted 1 July 2025) o be delisted 1 July 20 3.43 23.45 | 90 500 | √ <u>F</u> | ritamin B6 25 Pyridoxine multichem hiamine multichem |
| * Tab, strong, BPC | 11.25 | 500 | ✓ E | Splex |
| Vitamin C | | | | |
| ASCORBIC ACID a) No more than 100 mg per dose b) Only on a prescription * Tab 100 mg | 12.50 | 500 | √ <u>(</u> | evite evite |
| Vitamin D | | | | |
| ALFACALCIDOL * Cap 0.25 mcg * Cap 1 mcg | | 100 | ✓ (| One-Alpha One-Alpha S29 S29 One-Alpha |
| * Oral drops 2 mcg per ml | | 20 ml OP | | ne-Alpha |

| Subs (Manufactur | | | Fully Subsidised | Generic |
|---|-------|-------------|---------------------|---------------------------------------|
| \$ | | Per | | Manufacturer |
| CALCITRIOL | | | | |
| * Cap 0.25 mcg7.8 | 9 | 100 | | Calcitriol-AFT Calcitriol-AFT |
| * Cap 0.5 mcg13.6 | 8 | 100 | | S29 S29 Calcitriol-AFT Calcitriol-AFT |
| 00/ 50 // 0/5500/ | | | | S29 S29 |
| COLECALCIFEROL * Cap 1.25 mg (50,000 iu) – Maximum of 12 cap per prescription3.6 * Oral liq 188 mcg per ml (7,500 iu per ml)9.0 | | 12 ml Of | | Vit.D3 Clinicians |
| Multivitamin Preparations | | | | |
| MULTIVITAMIN RENAL - Special Authority see SA1546 below - Retail pha | rmacy | | | |

✓ Clinicians Renal Vit.

⇒SA1546 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 The patient has chronic kidney disease and is receiving either peritoneal dialysis or haemodialysis; or
- 2 The patient has chronic kidney disease grade 5, defined as patient with an estimated glomerular filtration rate of < 15 ml/min/1.73 m² body surface area (BSA).

MULTIVITAMINS - Special Authority see SA1036 below - Retail pharmacy

200 a OP ✓ Paediatric Seravit

⇒SA1036 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has inborn errors of metabolism.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where patient has had a previous approval for multivitamins.

VITAMINS

| * | Tab (BPC cap strength) | 1,000 | ✓ Mvite |
|---|---|-------|-------------|
| * | Cap (fat soluble vitamins A, D, E, K) - Special Authority see | | |
| | SA1720 below – Retail pharmacy23.40 | 60 | ✓ Vitabdeck |

⇒SA1720 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Patient has cystic fibrosis with pancreatic insufficiency: or
- 2 Patient is an infant or child with liver disease or short gut syndrome; or
- 3 Patient has severe malabsorption syndrome.

| | ALIMENTAR | Y TRAC | T AND | METABOLISM |
|---|--|------------|------------------|-------------------------------------|
| | Subsidy (Manufacturer's Price \$ |) Sub | Fully sidised | Brand or Generic Manufacturer |
| Minerals | | | | |
| Calcium | | | | |
| CALCIUM CARBONATE | | | | |
| * Tab 1.25 g (500 mg elemental) | 7.28 | 250 | √ 0 | Calci-Tab 500 |
| * Tab eff 1.25 g (500 mg elemental) – Subsidy by endorsem | | 100 | √ 0 | Calcium 500 mg Hexal S29 |
| Subsidy by endorsement – Only when prescribed for poconsidered unsuitable. | aediatric patients (< 5 | years) wh | ere calc | ium carbonate oral liquid is |
| CALCIUM GLUCONATE | | | | |
| * Inj 10%, 10 ml ampoule | 32.00 | 10 | ✓ N | Max Health - Hameln S29 |
| lodine | | | | |
| POTASSIUM IODATE * Tab 253 mcg (150 mcg elemental iodine) | 5.99 | 90 | ✓ V | leuroTabs |
| Iron | | | | |
| FERROUS FUMARATE | | | | |
| * Tab 200 mg (65 mg elemental) | 3.49 | 100 | √ F | erro-tab |
| FERROUS FUMARATE WITH FOLIC ACID | | | | |
| * Tab 310 mg (100 mg elemental) with folic acid 350 mcg | 5.98 | 100 | √ F | erro-F-Tabs |
| FERROUS SULFATE | | | _ | |
| * Tab long-acting 325 mg (105 mg elemental) | 2.55 | 30 | √ F | errograd |
| * Oral liq 30 mg (6 mg elemental) per 1 ml | | 250 ml | _ | erro-Liquid |
| 7 3 7 7 | 13.10 | 500 ml | | erodan ' |
| IRON (AS FERRIC CARBOXYMALTOSE) - Special Authority | see SA2394 below - | Retail pha | rmacv | |
| Inj 50 mg per ml, 10 ml vial | | 1 | | erinject |
| ⇒SA2394 Special Authority for Subsidy | | | | • |
| Initial application — (Anaemia) from any relevant practitioner following criteria: All of the following: | r. Approvals valid for | 3 months t | for applic | cations meeting the |
| 1 Patient has been diagnosed with anaemia: and | | | | |

- 1 Patient has been diagnosed with anaemia; and
- 2 Any of the following:
 - 2.1 Serum ferritin level is 20 mcg/L or less; or
 - 2.2 Both:
 - 2.2.1 Serum ferritin is between 20 and 50 mcg/L; and
 - 2.2.2 C-Reactive Protein (CRP) is at least 5 mg/L; or
 - 2.3 Patient has chronic inflammatory disease with symptoms of anaemia despite normal iron levels; and
- 3 Any of the following:
 - 3.1 Oral iron treatment has proven ineffective; or
 - 3.2 Oral iron treatment has resulted in dose-limiting intolerance; or
 - 3.3 Rapid correction of anaemia is required.

Renewal — (Anaemia) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following

| Subsidy | Fully | | Brand or |
|------------------------|------------|----------|----------|
| (Manufacturer's Price) | Subsidised | | Generic |
| \$ | Per | √ | |

continued...

Both:

- 1 Patient continues to have anaemia with a serum ferritin level of 20 mcg/L, or less or between 20 and 50 mcg/L with CRP of at least 5 mg/L, or has chronic inflammatory disease with symptoms of anaemia despite normal iron levels; and
- 2 A trial (or re-trial) with oral iron is clinically inappropriate.

Initial application — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with iron-deficiency anaemia; and
- 2 Any of the following:
 - 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
 - 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
 - 2.3 Patient has symptomatic heart failure, chronic kidney disease stage 3 or more or active inflammatory bowel disease and a trial of oral iron is unlikely to be effective; or
- 2.4 Rapid correction of anaemia is required.

Renewal — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient continues to have iron-deficiency anaemia; and
- 2 A re-trial with oral iron is clinically inappropriate.

| Magnesium | | |
|--|--------|---------------------------------|
| MAGNESIUM HYDROXIDE Suspension 8%33.60 | 355 ml | ✓ Phillips Milk of Magnesia ©29 |
| MAGNESIUM SULPHATE | | |
| * Inj 2 mmol per ml, 5 ml ampoule | 10 | ✓ Martindale |
| * Inj 2 mmol per ml, 10 ml ampoule | 10 | ✓ Inresa S29 |
| Zinc | | |
| ZINC SULPHATE * Cap 137.4 mg (50 mg elemental)11.00 | 100 | ✓ Zincaps |

✓ Ferrosia

5

BLOOD AND BLOOD FORMING ORGANS

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

Antianaemics

Hypoplastic and Haemolytic

⇒SA2266 Special Authority for Subsidy

Initial application — (chronic renal failure) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient in chronic renal failure: and
- 2 Haemoglobin is less than or equal to 100g/L; and
- 3 Any of the following:
 - 3.1 Both:
 - 3.1.1 Patient does not have diabetes mellitus; and
 - 3.1.2 Glomerular filtration rate is less than or equal to 30ml/min; or
 - 3.2 Both:
 - 3.2.1 Patient has diabetes mellitus; and
 - 3.2.2 Glomerular filtration rate is less than or equal to 45ml/min; or
 - 3.3 Patient is on haemodialysis or peritoneal dialysis.

Initial application — (myelodysplasia) from any specialist. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of myelodysplasia (MDS)*; and
- 2 Has had symptomatic anaemia with haemoglobin < 100g/L and is red cell transfusion-dependent; and
- 3 Patient has very low, low or intermediate risk MDS based on the WHO classification based prognostic scoring system for myelodysplastic syndrome (WPSS); and
- 4 Other causes of anaemia such as B12 and folate deficiency have been excluded; and
- 5 Patient has a serum epoetin level of < 500 IU/L; and
- 6 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Note: Indication marked with * is an unapproved indication

Renewal — **(chronic renal failure)** from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (myelodysplasia) from any specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient's transfusion requirement continues to be reduced with erythropoietin treatment; and
- 2 Transformation to acute myeloid leukaemia has not occurred; and
- 3 The minimum necessary dose of epoetin would be used and will not exceed 80,000 ju per week.

Note: Indication marked with * is an unapproved indication

EPOETIN ALFA - Special Authority see SA2266 above - Retail pharmacy

| Wastage claimable | | | |
|---------------------------------|--------|---|----------|
| Inj 1,000 iu in 0.5 ml, syringe | 250.00 | 6 | Binocrit |
| Inj 2,000 iu in 1 ml, syringe | 100.00 | 6 | Binocrit |
| Inj 3,000 iu in 0.3 ml, syringe | 150.00 | 6 | Binocrit |
| Inj 4,000 iu in 0.4 ml, syringe | 96.50 | 6 | Binocrit |
| Inj 5,000 iu in 0.5 ml, syringe | 125.00 | 6 | Binocrit |
| Inj 6,000 iu in 0.6 ml, syringe | 145.00 | 6 | Binocrit |
| Inj 8,000 iu in 0.8 ml, syringe | 175.00 | 6 | Binocrit |
| Inj 10,000 iu in 1 ml, syringe | 197.50 | 6 | Binocrit |
| Inj 40,000 iu in 1 ml, syringe | 250.00 | 1 | Binocrit |
| | | | |

BLOOD AND BLOOD FORMING ORGANS

| | Subsidy (Manufacturer's Price) \$ | | Fully ised ✓ | Brand or Generic Manufacturer |
|--------------------------------------|---|----------------|--------------------|-------------------------------------|
| Megaloblastic | | | | |
| FOLIC ACID * Tab 0.8 mg | 26.60 | 1,000 | | olic Acid multichem |
| * Tab 5 mg Oral liq 50 mcg per ml | | 100 5 ml OP | _ | olic Acid Viatris omed |

Antifibrinolytics, Haemostatics and Local Sclerosants

EFTRENONACOG ALFA [RECOMBINANT FACTOR IX] - [Xpharm]

For patients with haemophilia B receiving prophylaxis treatment. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management group.

| reaters Group in conjunction with the National | Haemophilia Management gro | up. | |
|---|----------------------------|-----|------------|
| Inj 250 iu vial | 612.50 | 1 | Alprolix |
| Inj 500 iu vial | 1,225.00 | 1 | ✓ Alprolix |
| Inj 1,000 iu vial | 2,450.00 | 1 | ✓ Alprolix |
| Inj 2,000 iu vial | 4,900.00 | 1 | ✓ Alprolix |
| Inj 3,000 iu vial | 7,350.00 | 1 | ✓ Alprolix |
| lnj 4,000 iu vial | 9,800.00 | 1 | Alprolix |
| ELTROMBOPAG - Special Authority see SA1743 b Wastage claimable | pelow - Retail pharmacy | | |
| Tab 25 mg | 1,550.00 | 28 | Revolade |
| Tab 50 mg | 3,100.00 | 28 | Revolade |
| | | | |

⇒SA1743 Special Authority for Subsidy

Initial application — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

Ε

- 1 Patient has had a splenectomy; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Any of the following:
 - 3.1 Patient has a platelet count of 20,000 to 30,000 platelets per microlitre and has evidence of significant mucocutaneous bleeding; or
 - 3.2 Patient has a platelet count of less than or equal to 20,000 platelets per microlitre and has evidence of active bleeding; or
 - 3.3 Patient has a platelet count of less than or equal to 10,000 platelets per microlitre.

Initial application — (**idiopathic thrombocytopenic purpura - preparation for splenectomy**) only from a haematologist. Approvals valid for 6 weeks where the patient requires eltrombopage treatment as preparation for splenectomy.

Initial application — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a significant and well-documented contraindication to splenectomy for clinical reasons; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Either:
 - 3.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per microliter: or
 - 3.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre

| Sub | sidy Fu | ully Brand or |
|------------|---------|----------------|
| (Manufactu | | sed Generic |
| | Per Per | ✓ Manufacturer |

continued...

and significant mucocutaneous bleeding.

Initial application — (severe aplastic anaemia) only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Two immunosuppressive therapies have been trialled and failed after therapy of at least 3 months duration; and
- 2 Either:
 - 2.1 Patient has severe aplastic anaemia with a platelet count of less than or equal to 20,000 platelets per microliter; or
 - 2.2 Patient has severe aplastic anaemia with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

Renewal — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 12 months where the patient has obtained a response (see Note) from treatment during the initial approval or subsequent renewal periods and further treatment is required.

Note: Response to treatment is defined as a platelet count of > 30,000 platelets per microlitre.

Renewal — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient's significant contraindication to splenectomy remains; and
- 2 The patient has obtained a response from treatment during the initial approval period; and
- 3 Patient has maintained a platelet count of at least 50,000 platelets per microlitre on treatment; and
- 4 Further treatment with eltrombopag is required to maintain response.

Renewal — (severe aplastic anaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has obtained a response from treatment of at least 20,000 platelets per microlitre above baseline during the initial approval period; and
- 2 Platelet transfusion independence for a minimum of 8 weeks during the initial approval period.

EMICIZUMAB - [Xpharm] - Special Authority see SA2272 below

| Inj 30 mg in 1 ml vial | 3,570.00 | 1 | ✓ Hemlibra |
|---------------------------|-----------|---|------------|
| Inj 60 mg in 0.4 ml vial | • | 1 | ✓ Hemlibra |
| Inj 105 mg in 0.7 ml vial | 12,492.00 | 1 | ✓ Hemlibra |
| Inj 150 mg in 1 ml vial | 17,846.00 | 1 | ✓ Hemlibra |

⇒SA2272 Special Authority for Subsidy

Initial application — (Severe Haemophilia A with or without FVIII inhibitors) only from a haematologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has severe congenital haemophilia A with a severe bleeding phenotype (endogenous factor VIII activity less than or equal to 2%); and
- 2 Emicizumab is to be administered at a dose of no greater than 3 mg/kg weekly for 4 weeks followed by the equivalent of 1.5 mg/kg weekly.

EPTACOG ALFA [RECOMBINANT FACTOR VIIA] - [Xpharm]

For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group. Rare Clinical Circumstances Brand of bypassing agent for > 14 days predicted use. Access to funded treatment for > 14 days predicted use is by named patient application to the Haemophilia Treaters Group, subject to access criteria.

| Inj 1 mg syringe | 1,178.30 | 1 | ✓ NovoSeven RT |
|------------------|----------|---|----------------|
| Inj 2 mg syringe | 2,356.60 | 1 | ✓ NovoSeven RT |
| Inj 5 mg syringe | 5,891.50 | 1 | ✓ NovoSeven RT |
| Inj 8 mg syringe | 9,426.40 | 1 | ✓ NovoSeven RT |

| | Subsidy | | Fully | Brand or |
|--|------------------------------|----------|---------------------------|-----------------------------|
| | (Manufacturer's Price) \$ | S Per | ubsidised • | Generic Manufacturer |
| FACTOR EIGHT INHIBITOR BYPASSING FRACTION - [Xph | narm] | | | |
| For patients with haemophilia. Preferred Brand of bypass | | | | |
| is managed by the Haemophilia Treaters Group in conjunc | ction with the National H | aemop | hilia Man | agement Group. |
| Inj 500 U | 1,315.00 | 1 | 1 | FEIBA NF |
| Inj 1,000 U | 2,630.00 | 1 | ✓ | FEIBA NF |
| Inj 2,500 U | 6,575.00 | 1 | ✓ | FEIBA NF |
| MOROCTOCOG ALFA [RECOMBINANT FACTOR VIII] - [Xp | harm] | | | |
| For patients with haemophilia. Rare Clinical Circumstance | | e recon | nbinant fa | ctor VIII. Access to funded |
| treatment is managed by the Haemophilia Treaters Group | in conjunction with the | Nationa | ıl Haemo | ohilia Management Group |
| subject to criteria. | • | | · | |
| Inj 250 iu prefilled syringe | 287.50 | 1 | 1 | Xyntha |
| Inj 500 iu prefilled syringe | 575.00 | 1 | 1 | Xyntha |
| Inj 1,000 iu prefilled syringe | 1,150.00 | 1 | 1 | Xyntha |
| Inj 2,000 iu prefilled syringe | 2,300.00 | 1 | ✓ : | Xyntha |
| Inj 3,000 iu prefilled syringe | 3,450.00 | 1 | 1 | Xyntha |
| NONACOG GAMMA, [RECOMBINANT FACTOR IX] - [Xphai | rm] | | | |
| For patients with haemophilia. Access to funded treatmer | | emophi | ilia Treate | ers Group in conjunction |
| with the National Haemophilia Management Group. | | | | , , , |
| Inj 1,000 iu vial | 870.00 | 1 | 1 | RIXUBIS |
| Ini 2.000 iu vial | | 1 | _ | RIXUBIS |
| Inj 3,000 iu vial | 2,610.00 | 1 | | RIXUBIS |
| OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (ADVATE) | _ [Ynharm] | | | |
| For patients with haemophilia. Preferred Brand of short h | | or VIII | Access to | funded treatment is |
| managed by the Haemophilia Treaters Group in conjunction | | | | |
| Inj 500 iu vial | | 1 | | Advate |
| Inj 1,000 iu vial | | 1 | | Advate |
| Inj 2,000 iu vial | | 1 | _ | Advate |
| Inj 3,000 iu vial | | 1 | | Advate |
| OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (KOGENA | • | | | |
| For patients with haemophilia. Rare Clinical Circumstance | | a racon | nhinant fa | ctor VIII Access to funder |
| treatment is managed by the Haemophilia Treaters Group | | | | |
| subject to criteria. | in conjunction with the | valione | ii i iaciiio _i | orilla mariagement Group |
| Inj 250 iu vial | 237 50 | 1 | 1 | Kogenate FS |
| Inj 500 iu vial | | i | | Kogenate FS |
| Inj 1,000 iu vial | | 1 | | Kogenate FS |
| Inj 2,000 iu vial | | i | | Kogenate FS |
| Inj 3,000 iu vial | | 1 | | Kogenate FS |
| • • | • | • | - ' | Acgoniato i o |
| RURIOCTOCOG ALFA PEGOL [RECOMBINANT FACTOR V For patients with haemophilia A receiving prophylaxis trea | | d +=aa+= | nantia m | anagad by the Heamanhill |
| Treaters Group in conjunction with the National Haemoph | | u ireair | nent is m | anaged by the Haemophii |
| | | 4 | ./ | A duma vata |
| Inj 1,000 iu vial | | 1 | | Adynovate |
| Inj 2,000 iu vial | ∠,400.00 | 1 | • | Adynovate |
| SODIUM TETRADECYL SULPHATE | | _ | | |
| * Inj 3% 2 ml | | 5 | | |
| | (73.00) | | | Fibro-vein |
| | | | | |
| TRANEXAMIC ACID | | | | |
| TRANEXAMIC ACID Tab 500 mg | 10.45 | 60 | • | Mercury Pharma |

| | Subsidy (Manufacturer's Price) \$ | Su Per | Fully bsidised | Brand or Generic Manufacturer | |
|---|---|-----------|-------------------|-------------------------------------|--|
| Vitamin K | | | | | |
| PHYTOMENADIONE Inj 2 mg per 0.2 ml - Up to 5 inj available on a PSO Inj 10 mg per ml, 1 ml - Up to 5 inj available on a PSO | | 5 5 | • • | onakion MM onakion MM | |

Antithrombotic Agents

Antiplatelet Agents

| • | | | |
|---|-------|-----|---------------------------------------|
| ASPIRIN | 12.65 | 990 | ✓ Ethics Aspirin EC |
| CLOPIDOGREL | 12.00 | 990 | Ettiles Aspiriii Ee |
| * Tab 75 mg | 5.07 | 84 | ✓ Arrow - Clopid |
| DIPYRIDAMOLE | 10.00 | 00 | 4 P. danser O.D. |
| * Tab long-acting 150 mg | | 60 | ✓ Pytazen SR |
| TICAGRELOR – Special Authority see SA1955 below | , , | | |
| * Tab 90 mg | 20.35 | 56 | Ticagrelor Sandoz |

⇒SA1955 Special Authority for Subsidy

Initial application — (acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Initial application — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

oun.

- 1 Fither:
 - 1.1 Patient has had a neurological stenting procedure* in the last 60 days; or
 - 1.2 Patient is about to have a neurological stenting procedure performed*; and
- 2 Either:
 - 2.1 Patient has demonstrated clopidogrel resistance using the P2Y12 (VerifyNow) assay or another appropriate platelet function assay and requires antiplatelet treatment with ticagrelor; or
 - 2.2 Either:
 - 2.2.1 Clopidogrel resistance has been demonstrated by the occurrence of a new cerebral ischemic event; or
 - 2.2.2 Clopidogrel resistance has been demonstrated by the occurrence of transient ischemic attack symptoms referable to the stent.

Initial application — (Percutaneous coronary intervention with stent deployment) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic**.

Initial application — (Stent thrombosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has experienced cardiac stent thrombosis whilst on clopidogrel.

Renewal — (subsequent acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for

| Subsidy | F | ully | Brand or |
|------------------------|---------|------|--------------|
| (Manufacturer's Price) | Subsidi | sed | Generic |
| \$ | Per | • | Manufacturer |

continued...

applications meeting the following criteria:

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Renewal — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient is continuing to benefit from treatment; and
- 2 Treatment continues to be clinically appropriate.

Renewal — (**Percutaneous coronary intervention with stent deployment**) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic**.

Notes: indications marked with * are unapproved indications.

Note: ** Clopidogrel allergy is defined as a history of anaphylaxis, urticaria, generalised rash or asthma (in non-asthmatic patients) developing soon after clopidogrel is started and is considered unlikely to be caused by any other treatment.

Heparin and Antagonist Preparations

| ENOXAPARIN SODIUM - Special Authority see SA215 | 2 below – Retail pharmacy | | |
|---|---------------------------|----|-----------------|
| Inj 20 mg in 0.2 ml syringe | 21.90 | 10 | ✓ Clexane |
| Inj 40 mg in 0.4 ml syringe | 29.74 | 10 | ✓ Clexane |
| Inj 60 mg in 0.6 ml syringe | | 10 | ✓ Clexane |
| Inj 80 mg in 0.8 ml syringe | | 10 | ✓ Clexane |
| Inj 100 mg in 1 ml syringe | 70.91 | 10 | ✓ Clexane |
| Inj 120 mg in 0.8 ml syringe | 88.11 | 10 | ✓ Clexane Forte |
| Ini 150 mg in 1 ml syringe | | 10 | ✓ Clexane Forte |

⇒SA2152 Special Authority for Subsidy

Initial application — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patients pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Initial application — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 For the short-term treatment of venous thromboembolism prior to establishing a therapeutic level of oral anti-coagulant treatment; or
- 2 For the prophylaxis and treatment of venous thromboembolism in high risk surgery; or
- 3 To enable cessation/re-establishment of existing oral anticoagulant treatment pre/post surgery; or
- 4 For the prophylaxis and treatment of venous thromboembolism in Acute Coronary Syndrome surgical intervention; or
- 5 To be used in association with cardioversion of atrial fibrillation.

Initial application — (Short-term use during treatment of COVID-19 with nirmatrelvir with ritonavir) from any relevant

| Subsidy | | Fully | Brand or |
|-----------------------|-----|------------|--------------|
| (Manufacturer's Price | | Subsidised | Generic |
| \$ | Per | ✓ | Manufacturer |

continued...

practitioner. Approvals valid for 2 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is receiving an anticoagulation treatment that has drug/drug interactions with ritonavir that increases risk of bleeding; and
- 2 Patient meets the Access Criteria for COVID-19 antivirals published on the Pharmac website*; and
- 3 Other antiviral treatments for COVID-19 have been considered and are not clinically suitable options.

Renewal — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

HEPARIN SODIUM

- 1 Low molecular weight heparin treatment is required during a patient's pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Renewal — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month where low molecular weight heparin treatment or prophylaxis is required for a second or subsequent event (surgery, ACS, cardioversion, or prior to oral anti-coagulation).

| Inj 1,000 iu per ml, 10 ml vial | 127.44 | 25 | ✓ Pfizer S29 |
|---|--------|-----|-----------------------------------|
| Inj 1,000 iu per ml, 5 ml ampoule | 25.49 | 10 | ✓ Wockhardt S29 |
| | 103.70 | | ✓ Wockhardt PSF S29 |
| | 127.44 | 50 | ✓ Pfizer |
| Inj 5,000 iu per ml, 5 ml vial | 83.00 | 10 | ✓ <u>Heparin Sodium</u> Panpharma |
| Inj 5,000 iu per ml, 1 ml | 70.33 | 5 | ✓ Hospira |
| Inj 25,000 iu per ml, 0.2 ml | | 5 | ✓ Hospira |
| | 42.40 | | ✓ Heparin DBL S29 |
| | 482.20 | 50 | ✓ Heparin DBL S29 |
| HEPARINISED SALINE | | | |
| Inj 10 iu per ml, 5 ml | 96.91 | 50 | ✓ Pfizer |
| Oral Anticoagulants | | | |
| DABIGATRAN | | | |
| Cap 75 mg - No more than 2 cap per day | 27.99 | 60 | ✓ Pradaxa |
| Cap 110 mg | 27.99 | 60 | ✓ Pradaxa |
| Cap 150 mg | 27.99 | 60 | ✓ Pradaxa |
| RIVAROXABAN | | | |
| Tab 10 mg - No more than 1 tab per day | 15.60 | 30 | ✓ Xarelto |
| Tab 15 mg - Up to 14 tab available on a PSO | 14.56 | 28 | ✓ Xarelto |
| Tab 20 mg | | 28 | ✓ Xarelto |
| WARFARIN SODIUM | | | |
| Note: Marevan and Coumadin are not interchangeable. | | | |
| * Tab 1 mg | 3.46 | 50 | ✓ Coumadin |
| | 7.50 | 100 | ✓ Marevan |
| * Tab 2 mg | 4.31 | 50 | ✓ Coumadin |
| * Tab 3 mg | 12.00 | 100 | ✓ Marevan |
| * Tab 5 mg | 5.93 | 50 | ✓ Coumadin |

13.50

100

✓ Marevan

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

| Subsidy | | Fully | Brand or | |
|------------------------|-----|---------|--------------|--|
| (Manufacturer's Price) | Sub | sidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

Blood Colony-stimulating Factors

| FILGRASTIM - Special Authority see SA1259 below - Retail pharm | acy | | |
|--|--------|----|------------|
| Inj 300 mcg per 0.5 ml prefilled syringe | 86.60 | 10 | ✓ Nivestim |
| Inj 480 mcg per 0.5 ml prefilled syringe | 133.72 | 10 | ✓ Nivestim |

⇒SA1259 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 20%*); or
- 2 Peripheral blood stem cell mobilisation in patients undergoing haematological transplantation; or
- 3 Peripheral blood stem cell mobilisation or bone marrow donation from healthy donors for transplantation; or
- 4 Treatment of severe chronic neutropenia (ANC < 0.5 ×109/L); or
- 5 Treatment of drug-induced prolonged neutropenia (ANC < 0.5 ×10⁹/L).

Note: *Febrile neutropenia risk greater than or equal to 20% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

⇒SA1912 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where used for prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 5%*). Note: *Febrile neutropenia risk greater than or equal to 5% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

Fluids and Electrolytes

Intravenous Administration

| GLUCOSE [DEXTROSE] * Inj 50%, 10 ml ampoule – Up to 5 inj available on a PSO * Inj 50%, 90 ml bottle – Up to 5 inj available on a PSO POTASSIUM CHLORIDE | | 5 1 | ✓ Biomed ✓ Biomed |
|---|-------|--------|--|
| * Inj 75 mg per ml, 10 ml | 65.00 | 50 | ✓ Juno✓ LumaCina✓ Pfizer S29 |
| SODIUM BICARBONATE | | | |
| Inj 8.4%, 50 ml | 24.70 | 1 | Biomed |
| a) Up to 5 inj available on a PSO b) Not in combination Inj 8.4%, 100 ml | 25.31 | 1 | ✓ Biomed |
| a) Up to 5 inj available on a PSO b) Not in combination | | | |

| | Subsidy | | Fully | Brand or |
|--|--------------------|-------------------|-------------------|-----------------------------|
| | (Manufacturer's Pr | rice) Subs Per | idised • | Generic Manufacturer |
| SODIUM CHLORIDE | | | | |
| Not funded for use as a nasal drop. Not funded for nebulise | r use except when | n used in conju | unction | with an antibiotic intended |
| for nebuliser use. | | | | |
| Inj 0.9%, bag - Up to 2000 ml available on a PSO | | 500 ml | | axter |
| | 1.58 | 1,000 ml | | axter |
| Only if prescribed on a prescription for renal dialysis, ma for emergency use. (500 ml and 1,000 ml packs) | | | | |
| Inj 23.4% (4 mmol/ml), 20 ml ampoule | | 5 | ✓ B | iomed |
| For Sodium chloride oral liquid formulation refer Standar | 71 0 | | , _ | |
| Inj 0.9%, 5 ml ampoule – Up to 5 inj available on a PSO | | 20 | _ | resenius Kabi |
| Inj 0.9%, 10 ml ampoule – Up to 5 inj available on a PSO | | 50 | _ | resenius Kabi |
| Inj 0.9%, 20 ml ampoule | 5.00 | 20 | V <u>r</u> | resenius Kabi |
| TOTAL PARENTERAL NUTRITION (TPN) | | | | |
| Infusion | CBS | 1 OP | ✓ T | PN |
| VATER | | | | |
| On a prescription or Practitioner's Supply Order only w Schedule requiring a solvent or diluent; or On a bulk supply order; or When used in the extemporaneous compounding of ey When used for the dilution of sodium chloride soln 7% | e drops; or | | ection ii | sieu III ille Filannaceulic |
| Inj 10 ml ampoule – Up to 5 inj available on a PSO | | 50 | _ | lultichem |
| Inj 20 ml ampoule – Up to 5 inj available on a PSO | 5.00 | 20 | ▼ <u>F</u> | resenius Kabi |
| Oral Administration | | | | |
| CALCIUM POLYSTYRENE SULPHONATE | | | | |
| Powder | 169.85 | 300 g OP | ✓ 0 | alcium Resonium |
| COMPOUND ELECTROLYTES | | 3 - | | |
| Powder for oral soln — Up to 5 sach available on a PSO | 9.53 | 50 | √ F | lectral |
| • | | 30 | · <u>-</u> | icottai |
| COMPOUND ELECTROLYTES WITH GLUCOSE [DEXTROSE] | | 1 000 00 | | la calma la aka |
| Soln with electrolytes | 6.53 | 1,000 ml OP | V <u>H</u> | lydralyte - |
| | | | | Lemonade |
| PHOSPHORUS | | | | |
| Tab eff 500 mg (16 mmol) | 82.50 | 100 | ✓ P | hosphate Phebra |
| POTASSIUM CHLORIDE | | | | |
| * Tab eff 548 mg (14 m eq) with chloride 285 mg (8 m eq) | 5.26 | 60 | | |
| | (17.10) | | | chlorvescent |
| * Tab long-acting 600 mg (8 mmol) | 15.35 | 200 | ✓ S | pan-K |
| SODIUM BICARBONATE | | | | |
| Cap 840 mg | 8.52 | 100 | ✓ S | odibic |
| - | | | ✓ S | odibic |
| SODIUM POLYSTYRENE SULPHONATE | | | | |
| Decides Of the Control of the Contro | 04.05 | 454 - 00 | | |

✓ Resonium-A

454 g OP

Powder84.65

| | (Manufacturer's Pr | ice) Subs | idised | Generic |
|---|--------------------|-----------|---------------------|------------------|
| | \$ | Per | ✓ | Manufacturer |
| Alpha-Adrenoceptor Blockers | | | | |
| Alpha Adrenoceptor Blockers | | | | |
| DOXAZOSIN | | | | |
| * Tab 2 mg | 17.35 | 500 | | xazosin Clinect |
| * Tab 4 mg | 20.94 | 500 | ✓ Do | xazosin Clinect |
| PHENOXYBENZAMINE HYDROCHLORIDE | | | | |
| * Cap 10 mg | 65.00 | 30 | ✓ BN | IM \$29 |
| | 216.67 | 100 | ✓ Dib | oenzyline \$29 |
| PRAZOSIN | | | | |
| * Tab 1 mg | 5.53 | 100 | ✓ Arı | rotex-Prazosin |
| | | | 5 | S29 S29 |
| | 9.98 | | 🗸 Mii | nipress S29 |
| * Tab 2 mg | 7.00 | 100 | ✓ Arı | rotex-Prazosin |
| | | | 5 | S29 S29 |
| | 13.29 | | | nipress S29 |
| * Tab 5 mg | 11.70 | 100 | | rotex-Prazosin |
| | | | 5 | S29 S29 |
| | 22.00 | | ✓ Mii | nipress S29 |
| * Cap 1 mg | 15.40 | 100 | ✓ Pra | azosin Mylan S29 |
| * Cap 2 mg | 15.58 | 100 | ✓ Pragram | azosin Mylan S29 |
| * Cap 5 mg | 23.32 | 100 | ✓ Pra | azosin Mylan S29 |
| Agents Affecting the Renin-Angiotensin System ACE Inhibitors | n | | | |
| CAPTOPRIL | | | | |
| Oral liq 5 mg per ml Oral liquid restricted to children under 12 years of age. | 86.00 | 100 ml OP | ✓ <u>DP</u> | -Captopril |
| CILAZAPRIL – Subsidy by endorsement | | | | |
| Subsidy by endorsement – Subsidised for patients who wer endorsed accordingly. Pharmacists may annotate the preso dispensing of cilazapril. | | | | |
| * Tab 0.5 mg | | 90 | ✓ Za | |
| * Tab 2.5 mg | | 90 | ✓ Za | |
| Tab 5 mg | 10.05 | 90 | ✓ Za | prii |
| ENALAPRIL MALEATE | | 0.5 | , . | |
| * Tab 10 mg | | 90 | ✓ <u>Ac</u> | |
| * Tab 10 mg* Tab 20 mg | | 90 90 | ✓ <u>Ac</u> ✓ Ac | |
| · · | 2.00 | 30 | ▼ AU | <u> </u> |
| LISINOPRIL * Tab 5 mg | 11.07 | 90 | ✓ Eth | nics Lisinopril |
| rab o my | | 00 | | va Lisinopril |
| * Tab 10 mg | 11.67 | 90 | | nics Lisinopril |
| • | - | | | va Lisinopril |
| * Tab 20 mg | 14.69 | 90 | ✓ Eth | nics Lisinopril |
| | | | ✓ <u>Te</u> | va Lisinopril |
| | | | | |

Subsidy

Fully

Brand or

| | Subsidy | | Fully | |
|--|------------------------|----------|------------|---|
| | (Manufacturer's Price) | Per | Subsidised | Generic Manufacturer |
| PERINDOPRIL | Ψ | | | a.iaiaaiai |
| * Tab 2 mg | 1 70 | 30 | 1 | Coversyl |
| * Tab 4 mg | | 30 | | Coversyl |
| • | | | | |
| | 3.94 | 30 | • | Coversyl |
| QUINAPRIL | | | | |
| * Tab 5 mg Arrow-Quinapril 5 to be Principal Supply on 1 March 2025 | | 90 | • | Arrow-Quinapril 5 |
| * Tab 10 mg | | 90 | ✓ | Arrow-Quinapril 10 |
| Arrow-Quinapril 10 to be Principal Supply on 1 March 202 * Tab 20 mg | | 90 | 1 | Arrow-Quinapril 20 |
| Arrow-Quinapril 20 to be Principal Supply on 1 March 202 | | 90 | • | Arrow-Quinaprii 20 |
| RAMIPRIL | | | | |
| * Cap 1.25 mg | 17.25 | 90 | 1 | Tryzan |
| * Cap 2.5 mg | 16.50 | 90 | 1 | Tryzan |
| * Cap 5 mg | 16.88 | 90 | 1 | Tryzan |
| ₭ Cap 10 mg | 17.63 | 90 | ✓ | Tryzan |
| Angiotensin II Antagonists | | | | |
| CANDESARTAN CILEXETIL | | | | |
| * Tab 4 mg | 2.68 | 90 | 1 | Candestar |
| ₭ Tab 8 mg | | 90 | | Candestar |
| ★ Tab 16 mg | | 90 | | Candestar |
| ₭ Tab 32 mg | | 90 | / | Candestar |
| OSARTAN POTASSIUM | | | | |
| | 2.00 | 84 | ./ | Locarton Actovia |
| ★ Tab 12.5 mg ★ Tab 25 mg | | 84 | | Losartan Actavis Losartan Actavis |
| | | | | |
| * Tab 50 mg | | 84 | | Losartan Actavis |
| ≰ Tab 100 mg | 4.57 | 84 | • | Losartan Actavis |
| Angiotensin II Antagonists with Diuretics | | | | |
| CANDESARTAN CILEXETIL WITH HYDROCHLOROTHIAZIDE | | | | |
| * Tab 16 mg with hydrochlorothiazide 12.5 mg | 4.10 | 30 | • | APO-Candesartan HCTZ 16/12.5 |
| * Tab 32 mg with hydrochlorothiazide 12.5 mg | 5.25 | 30 | ✓ | APO-Candesartan HCTZ 32/12.5 |
| OSARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE | | | | |
| * Tab 50 mg with hydrochlorothiazide 12.5 mg | 4.00 | 30 | • | Arrow-Losartan & Hydrochlorothiazide |
| Angiotensin II Antagonists with Neprilysin Inhibi | tors | | | |
| SACUBITRIL WITH VALSARTAN - Special Authority see SA230 | on the next nego | Rota | il nharma | 21/ |
| | | | | |
| Tab 24.3 mg with valsartan 25.7 mg | | 56 56 | | Entresto 24/26 |
| Tab 48.6 mg with valsartan 51.4 mg | | 56 | | Entresto 49/51 |

Tab 97.2 mg with valsartan 102.8 mg190.00

✓ Entresto 97/103

56

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

⇒SA2302 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has heart failure; and
- 2 Any of the following:
 - 2.1 Patient is in NYHA/WHO functional class II; or
 - 2.2 Patient is in NYHA/WHO functional class III; or
 - 2.3 Patient is in NYHA/WHO functional class IV; and
- 3 Either:
 - 3.1 Patient has a documented left ventricular ejection fraction (LVEF) of less than or equal to 35%; or
 - 3.2 An ECHO is not reasonably practical, and in the opinion of the treating practitioner the patient would benefit from treatment; and
- 4 Patient is receiving concomitant optimal standard chronic heart failure treatments.

Antiarrhythmics

| For lignocaine hydrochloride refer to NERVOUS SYSTEM, Anaesthetics, Local, p | age 122 | |
|--|---------|--|
| AMIODARONE HYDROCHLORIDE | | |
| ▲ Tab 100 mg3.49 | 30 | ✓ Aratac |
| ▲ Tab 200 mg4.49 | 30 | ✓ Aratac |
| Inj 50 mg per ml, 3 ml ampoule - Up to 10 inj available on a PSO 9.12 | 6 | ✓ Cordarone-X |
| 15.22 | 10 | ✓ Max Health |
| ATROPINE SULPHATE | | |
| * Inj 600 mcg per ml, 1 ml ampoule - Up to 5 inj available on a | | |
| PSO16.10 | 10 | ✓ Hikma S29 |
| | | ✓ Juno S29 |
| | | ✓ <u>Martindale</u> |
| DIGOXIN | | |
| * Tab 62.5 mcg - Up to 30 tab available on a PSO | 240 | ✓ Lanoxin PG |
| * Tab 250 mcg - Up to 30 tab available on a PSO16.90 | 240 | ✓ Lanoxin |
| * Oral liq 50 mcg per ml | 60 ml | ✓ Lanoxin |
| | | Lanoxin Paediatric |
| | | Elixir |
| | | ✓ Lanoxin S29 S29 |
| DISOPYRAMIDE PHOSPHATE | | |
| ▲ Cap 100 mg23.87 | 100 | ✓ Rythmodan |
| 55.90 | 84 | Rythmodan - |
| | | Cheplafarm \$29 |

| (1 | Subsidy Manufacturer's Price) \$ | Per | Fully Subsidised | |
|---|--|-----|---------------------|--|
| FLECAINIDE ACETATE | | | | |
| ▲ Tab 50 mg | | 60 | | Flecainide BNM |
| ▲ Cap long-acting 100 mg | 35.78 | 90 | • | Flecainide Controlled Release Teva |
| ▲ Cap long-acting 200 mg | 54.28 | 90 | ✓ | Flecainide Controlled Release Teva |
| Inj 10 mg per ml, 15 ml ampoule | 102.79 | 5 | 1 | Almarytm \$29 |
| | 108.16 | | | Tambocor Tambocor German S29 |
| MEXILETINE HYDROCHLORIDE | | | | |
| ▲ Cap 150 mg | 162.00 | 100 | 1 | Teva S29 |
| A Cap 250 mgPROPAFENONE HYDROCHLORIDE | 202.00 | 100 | ✓ | Teva S29 |
| ▲ Tab 150 mg | 40.90 | 50 | ✓ | Rytmonorm |
| Antihypotensives MIDODRINE – Special Authority see SA1474 below – Retail pharm | nacy | | | |
| Tab 2.5 mg | • | 100 | | MAR-Midodrine S29 <u>Midodrine</u> <u>Medsurge</u> |

■SA1474 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years where patient has disabling orthostatic hypotension not due to drugs.

100

✓ MAR-Midodrine S29
✓ Midodrine
Medsurge

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Beta-Adrenoceptor Blockers

Beta Adrenoceptor Blockers

| ATENOLOL * Tab 50 mg 11.00 * Tab 100 mg 18.50 * Oral liq 25 mg per 5 ml 49.85 Restricted to children under 12 years of age. | 500 500 300 ml OP | ✓ <u>Viatris</u> ✓ <u>Atenolol Viatris</u> ✓ Atenolol AFT |
|---|-------------------------|---|
| BISOPROLOL FUMARATE * Tab 2.5 mg | 90 90 90 | ✓ Ipca-Bisoprolol ✓ Ipca-Bisoprolol ✓ Ipca-Bisoprolol |
| CARVEDILOL * Tab 6.25 mg | 60 60 60 | ✓ Carvedilol Sandoz ✓ Carvedilol Sandoz ✓ Carvedilol Sandoz |

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

| | Subsidy | | Fully | Brand or |
|--|------------------------|-------|------------|-----------------------|
| | (Manufacturer's Price) | | Subsidised | |
| | \$ | Per | | Manufacturer |
| ABETALOL | | | | |
| * Tab 100 mg | 14.50 | 100 | ✓ | Trandate |
| * Tab 200 mg | 27.00 | 100 | ✓ | Trandate |
| * Inj 5 mg per ml, 20 ml ampoule | | 5 | | |
| | (88.60) | | | Trandate |
| METOPROLOL SUCCINATE | | | | |
| * Tab long-acting 23.75 mg | 4.20 | 90 | 1 | Myloc CR |
| * Tab long-acting 47.5 mg | | 90 | | Myloc CR |
| * Tab long-acting 95 mg | | 90 | 1 | Myloc CR |
| * Tab long-acting 190 mg | | 90 | 1 | Myloc CR |
| METOPROLOL TARTRATE | | | | |
| * Tab 50 mg | 5.66 | 100 | 1 | IPCA-Metoprolol |
| * Tab 100 mg | | 60 | | IPCA-Metoprolol |
| * Tab long-acting 200 mg | | 28 | | Slow-Lopresor |
| * Inj 1 mg per ml, 5 ml vial | | 5 | | Metoprolol IV Mylan |
| ,, ., ., ., ., ., ., ., ., ., ., | | - | | Metoprolol IV Viatris |
| NADOLOL | | | | |
| * Tab 40 mg | 19.19 | 100 | 1 | Nadolol BNM |
| * Tab 80 mg | | 100 | ✓ | Nadolol BNM |
| PROPRANOLOL | | | | |
| * Tab 10 mg | 7.04 | 100 | / | Drofate |
| * Tab 40 mg | | 100 | | IPCA-Propranolol |
| * Cap long-acting 160 mg | | 100 | | Cardinol LA |
| * Oral liq 4 mg per ml - Special Authority see SA1327 be | | | | - |
| Retail pharmacy | | 500 m | n 🗸 | Roxane- |
| • • | | | | Propranolol \$29 |

⇒SA1327 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Fither:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

SOTALOL

AMI ODIDINI

| * | Tab 80 mg37.50 | 500 | ✓ Mylan |
|---|-----------------|-----|---------|
| | Tab 160 mg14.00 | | ✓ Mylan |

Calcium Channel Blockers

Dihydropyridine Calcium Channel Blockers

| AIV | LODIPINE | | | |
|-----|------------|------|----|-----------|
| * | Tab 2.5 mg | 1.45 | 90 | ✓ Vasorex |
| | Tab 5 mg | | 90 | ✓ Vasorex |
| * | Tob 10 mg | 1 21 | 00 | / Vacaray |

| | Subsidy | | Fully | Brand or |
|--|------------------------|------|------------|------------------------------------|
| | (Manufacturer's Price) | Per | Subsidised | Generic Manufacturer |
| FLADIDINE | Ψ | 1 01 | • | Warranacturer |
| ELODIPINE | 0.10 | 30 | ./ | Plendil ER |
| ★ Tab long-acting 2.5 mg ★ Tab long-acting 5 mg | | 90 | | Felo 5 ER |
| ★ Tab long-acting 5 mg ★ Tab long-acting 10 mg | | 90 | | Felo 10 ER |
| | 0.93 | 90 | • | reio io En |
| IFEDIPINE | | | | |
| ← Tab long-acting 10 mg – Subsidy by endorsement | 19.42 | 56 | | Tensipine MR10 S29 |
| Subsidised for patients who were taking nifedipine to endorsed accordingly. Pharmacists may annotate the dispensing of nifedipine tab long-acting 10 mg. | 0 0 | | • | |
| Tab long-acting 20 mg | 17.72 | 100 | | Nyefax Retard |
| * Tab long-acting 30 mg | 4.78 | 14 | • | Mylan Italy (24 hr release) §29 |
| | 34.10 | 100 | 1 | Mylan (24 hr release) S29 |
| ₭ Tab long-acting 60 mg | 52.81 | 100 | ✓ | Mylan (24 hr |
| | | | | release) S29 |
| Other Calcium Channel Blockers | | | | |
| ILTIAZEM HYDROCHLORIDE | | | | |
| Cap long-acting 120 mg | 65.35 | 500 | ✓ | Diltiazem CD Clinect |
| ★ Cap long-acting 180 mg | 7.00 | 30 | ✓ | Cardizem CD |
| Cap long-acting 240 mg | 9.30 | 30 | ✓ | Cardizem CD |
| PERHEXILINE MALEATE | | | | |
| ★ Tab 100 mg | 62.90 | 100 | 1 | Pexsig |
| /ERAPAMIL HYDROCHLORIDE | | | | · · |
| ₹ Tab 40 mg | 7.01 | 100 | 1 | Isoptin |
| ₭ Tab 80 mg | | 100 | | Isoptin |
| ₹ Tab long-acting 120 mg | | 100 | | Isoptin Retard \$29 |
| Tab long doding 120 mg | | 100 | | Isoptin SR |
| Fab long-acting 240 mg | 15.12 | 30 | | Isoptin SR |
| Inj 2.5 mg per ml, 2 ml ampoule – Up to 5 inj available o | | | | |
| PSO | | 5 | ✓ | Isoptin |
| | | | | |
| Centrally-Acting Agents | | | | |
| | | | | |
| CLONIDINE | | | | |
| ★ Patch 2.5 mg, 100 mcg per day — Only on a prescription | | 4 | | <u>Mylan</u> |
| ★ Patch 5 mg, 200 mcg per day − Only on a prescription | | 4 | _ ` | <u>Mylan</u> |
| ▶ Patch 7.5 mg, 300 mcg per day — Only on a prescription | 17.90 | 4 | ✓. | <u>Mylan</u> |
| CLONIDINE HYDROCHLORIDE | | | | |
| ★ Tab 25 mcg | 29.32 | 112 | 1 | Clonidine Teva |
| ★ Tab 150 mcg | | 100 | 1 | <u>Catapres</u> |
| Inj 150 mcg per ml, 1 ml ampoule | 14.10 | 5 | ✓ | Catapres |
| METHYLDOPA | | | | |
| ⊁ Tab 250 mg | 15.10 | 100 | 1 | Methyldopa Viatris |
| · | | | - | |

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

| | | (Manufacturer's P | rice) Subs | sidised (| Brand or Generic Manufacturer |
|-----|---|---------------------|----------------|----------------|-------------------------------------|
| | | \$ | Per | V 1 | wanulacturer |
| ט | iuretics | | | | |
| L | pop Diuretics | | | | |
| | METANIDE | 40.00 | 100 | 4 D | |
| * | Tab 1 mg | | 100 5 | ✓ Bur ✓ Bur | |
| FU | ROSEMIDE [FRUSEMIDE] | | | _ | |
| * | Tab 40 mg — Up to 30 tab available on a PSO Tab 500 mg | | 1,000 50 | | <u>A-Frusemide</u> x Forte |
| * | Oral liq 10 mg per ml | 11.20 | 30 ml OP | ✓ Las | ix |
| | Inj 10 mg per ml, 25 ml ampoule | | 6 5 | ✓ Las | ix osemide-Baxter |
| | | 002.40 | | - <u>- u</u> | OSCIIIUC BUXICI |
| P | otassium Sparing Diuretics | | | | |
| AM | ILORIDE HYDROCHLORIDE | 04.07 | 100 | 45 | |
| | Tab 5 mg | 81.07 | 100 28 | | agis S29 ckhardt S29 |
| | Oral liq 1 mg per ml | | 25 ml OP | ✓ Bio | |
| EP | LERENONE - Special Authority see SA1728 below - Retail p | | 00 | . In a | |
| | Tab 25 mg | | 30 30 | ✓ <u>Ins</u> p | |
| | SA1728 Special Authority for Subsidy | | | | |
| | ial application from any relevant practitioner. Approvals valid following criteria: | d without further i | renewal unless | s notified f | or applications meeting |
| Bot | • | | | | |
| | 1 Patient has heart failure with ejection fraction less than 402 Either: | %; and | | | |
| | 2.1 Patient is intolerant to optimal dosing of spironolact | | | | |
| | 2.2 Patient has experienced a clinically significant adve | erse effect while | on optimal dos | sing of spir | onolactone. |
| | RONOLACTONE Tab 25 mg | 3.68 | 100 | ✓ Spir | ractin |
| | Tab 100 mg | 10.65 | 100 | ✓ Spir | ractin |
| | Oral liq 5 mg per ml | 35.70 | 25 ml OP | ✓ Bio | med |
| P | otassium Sparing Combination Diuretics | | | | |
| | ILORIDE HYDROCHLORIDE WITH FUROSEMIDE | 0.00 | | | |
| | Tab 5 mg with furosemide 40 mgILORIDE HYDROCHLORIDE WITH HYDROCHLOROTHIAZI | | 28 | ✓ Fru | mii |
| | Tab 5 mg with hydrochlorothiazide 50 mg | | 50 | ✓ Mod | duretic |

Subsidy

Fully

Brand or

✓ Jinarc

✓ Jinarc

56 OP

| | Subsidy (Manufacturer's Price \$ | e) Subs Per | Fully sidised | Brand or Generic Manufacturer |
|--|--|-------------------------|-------------------|-------------------------------------|
| Thiazide and Related Diuretics | | | | |
| BENDROFLUMETHIAZIDE [BENDROFLUAZIDE] * Tab 2.5 mg - Up to 150 tab available on a PSO | 51.50 | 500 | ✓ <u>A</u> | Arrow- Bendrofluazide |
| May be supplied on a PSO for reasons other than emerg * Tab 5 mg | , , | 500 | ✓ <u>A</u> | Arrow- Bendrofluazide |
| CHLOROTHIAZIDE Oral liq 50 mg per ml | 30.67 2 | 25 ml OP | ✓ B | Biomed |
| CHLORTALIDONE [CHLORTHALIDONE] * Tab 25 mg | 6.95 | 50 | ✓ <u>H</u> | lygroton |
| INDAPAMIDE * Tab 2.5 mg METOLAZONE | 16.00 | 90 | ✓ <u>D</u> | Dapa-Tabs |
| Tab 5 mg | CBS | 1 50 | | Metolazone S29 Zaroxolyn S29 |
| Vasopressin receptor antagonists | | | | |
| TOLVAPTAN – Special Authority see SA2166 below – Retail pha Tab 15 mg Tab 30 mg Tab 45 mg + 15 mg | 873.50 873.50 | 28 OP 28 OP 56 OP | √ J | linarc linarc linarc |

⇒SA2166 Special Authority for Subsidy

Initial application — (autosomal dominant polycystic kidney disease) only from a renal physician or any relevant practitioner on the recommendation of a renal physician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 Patient has a confirmed diagnosis of autosomal dominant polycystic kidney disease; and

- 2 Patient has an estimated glomerular filtration rate (eGFR) of greater than or equal to 25 ml/min/1.73 m² at treatment initiation; and
- 3 Fither:
 - 3.1 Patient's disease is rapidly progressing, with a decline in eGFR of greater than or equal to 5 mL/min/1.73 m² within one-year; or
 - 3.2 Patient's disease is rapidly progressing, with an average decline in eGFR of greater than or equal to 2.5 mL/min/1.73 m² per year over a five-year period.

Renewal — (autosomal dominant polycystic kidney disease) only from a renal physician or any relevant practitioner on the recommendation of a renal physician. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has not developed end-stage renal disease, defined as an eGFR of less than 15 mL/min/1.73 m²; and
- 2 Patient has not undergone a kidney transplant.

| | Subsidy | | Fully | Brand or |
|--|---|----------|----------------|---------------------------|
| | (Manufacturer's Price) | | ubsidised | Generic |
| | \$ | Per | ubsidised ✓ | Manufacturer |
| | * | | | |
| Lipid-Modifying Agents | | | | |
| Fibrates | | | | |
| BEZAFIBRATE | | | | |
| * Tab 200 mg | 22.65 | 90 | ✓ E | Bezalip |
| Bezalip to be Principal Supply on 1 March 2025 | | | | • |
| * Tab long-acting 400 mg | 21.54 | 30 | ✓ E | Bezalip Retard |
| Bezalip Retard to be Principal Supply on 1 March 2025 | | | | • |
| | | | | |
| Other Lipid-Modifying Agents | | | | |
| ACIPIMOX | | | | |
| * Cap 250 mg | 38.19 | 30 | ✓ (| Olbetam |
| Resins | | | | |
| 11001110 | | | | |
| COLESTYRAMINE | | | | |
| Powder for oral suspension 4 g sachet | 61.50 | 50 | ✓ (| Colestyramine - |
| | | | | Mylan S29 |
| | | | ✓ (| Quantalan sugar |
| | | | | free S29 |
| | | | | |
| HMG CoA Reductase Inhibitors (Statins) | | | | |
| ATODVACTATINI | | | | |
| ATORVASTATIN | 0.01 | 30 | ./ 1 | avatat |
| Tab 10 mg | 5.16 | 500 | _ | <u>-orstat</u> -orstat |
| Tab 20 mg | • | 500 | _ | orstat |
| Tab 40 mg | | 500 | _ | orstat |
| Tab 80 mg | | 30 | _ | orstat |
| 1 ab 00 mg | 25.39 | 500 | | orstat |
| DDAMACTATINI | 20.00 | 000 | - | <u> </u> |
| PRAVASTATIN * Tab 20 mg | 7.16 | 100 | | Clinect |
| * Tab 20 mg | | 100 | _ | <u>Simect</u> Clinect |
| • | | 100 | <u> </u> | <u>Jilliect</u> |
| ROSUVASTATIN – Special Authority see SA2093 below – Reta | ' | | | |
| * Tab 5 mg | | 30 | _ | Rosuvastatin Viatris |
| * Tab 10 mg | | 30 | _ | Rosuvastatin Viatris |
| * Tab 40 mg | | 30 30 | _ | Rosuvastatin Viatris |
| * Tab 40 mg | 4.55 | 30 | V I | Rosuvastatin Viatris |
| ⇒SA2093 Special Authority for Subsidy | | _ | | |
| Initial application — (cardiovascular disease risk) from any | relevant practitioner. | Approv | als valid w | vithout further renewal |
| unless notified for applications meeting the following criteria: | | | | |
| Either: | | | | |
| 1 Both: | | | | |
| 1.1 Patient is considered to be at risk of cardiovascula1.2 Patient is Māori or any Pacific ethnicity; or | ar disease; and | | | |
| 2 Both: | | | | |
| E Bouil. | | | | |

/ Olasson - Latin Madan

| Subsidy | | Fully | Brand or | |
|------------------------|------------|-------|--------------|--|
| (Manufacturer's Price) | Subsidised | | Generic | |
| \$ | Per | ✓ | Manufacturer | |

continued...

- 2.1 Patient has a calculated risk of cardiovascular disease of at least 15% over 5 years; and
- 2.2 LDL cholesterol has not reduced to less than 1.8 mmol/litre with treatment with the maximum tolerated dose of atoryastatin and/or simvastatin.

Initial application — (familial hypercholesterolemia) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has familial hypercholesterolemia (defined as a Dutch Lipid Criteria score greater than or equal to 6); and
- 2 LDL cholesterol has not reduced to less than 1.8 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initial application — (established cardiovascular disease) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:
Both:

- 1 Any of the following:
 - 1.1 Patient has proven coronary artery disease (CAD); or
 - 1.2 Patient has proven peripheral artery disease (PAD); or
 - 1.3 Patient has experienced an ischaemic stroke; and
- 2 LDL cholesterol has not reduced to less than 1.4 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initial application — (recurrent major cardiovascular events) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

SIMVASTATIN

- 1 Patient has experienced a recurrent major cardiovascular event (defined as myocardial infarction, ischaemic stroke, coronary revascularisation, hospitalisation for unstable angina) in the last 2 years; and
- 2 LDL cholesterol has not reduced to less than 1.0 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

| * | 1.68 10 mg | 90 | ✓ Simvastatin Mylan ✓ Simvastatin Viatris |
|-----|--|----|---|
| * | Tab 20 mg2.54 | 90 | Simvastatin Viatris |
| * | Tab 40 mg4.11 | 90 | ✓ Simvastatin Viatris |
| * | Tab 80 mg8.81 | 90 | ✓ Simvastatin Viatris |
| S | elective Cholesterol Absorption Inhibitors | | |
| ΕZ | ETIMIBE | | |
| * | Tab 10 mg1.76 | 30 | Ezemibe Viatris |
| | | | Ezetimibe Sandoz |
| (Ez | zemibe Viatris Tab 10 mg to be delisted 1 July 2025) | | |
| ΕZ | ETIMIBE WITH SIMVASTATIN | | |
| | Tab 10 mg with simvastatin 10 mg5.15 | 30 | ✓ Zimybe |
| | Tab 10 mg with simvastatin 20 mg6.15 | 30 | ✓ Zimybe |
| | Tab 10 mg with simvastatin 40 mg7.15 | 30 | ✓ Zimybe |

30

Zimvbe

Tab 10 mg with simvastatin 80 mg......8.15

| | Subsidy | | Fully Brand or |
|---|---|---|--|
| | (Manufacturer's | | idised Generic |
| | \$ | Per | ✓ Manufacturer |
| | | | |
| Nitrates | | | |
| OLVOEDVI TRINITRATE | | | |
| GLYCERYL TRINITRATE | | | |
| * Oral pump spray, 400 mcg per dose – Up to 250 dose | | | |
| available on a PSO | 7.48 | 250 dose OP | Nitrolingual Pump |
| | | | Spray |
| * Patch 25 mg, 5 mg per day | 15.73 | 30 | ✓ Nitroderm TTS |
| * Patch 50 mg, 10 mg per day | 18.62 | 30 | ✓ Nitroderm TTS |
| ISOSORBIDE MONONITRATE | | | |
| * Tab 20 mg | 22.40 | 100 | ✓ Ismo 20 |
| * Tab long-acting 40 mg | | 30 | ✓ Ismo 40 Retard |
| | | 90 | _ |
| * Tab long-acting 60 mg | 13.30 | 90 | ✓ <u>Duride</u> |
| Sympathomimetics | | | |
| Sympathonimetics | | | |
| ADRENALINE | | | |
| Inj 1 in 1,000, 1 ml ampoule – Up to 5 inj available on a PS | SO4.98 | 5 | ✓ Aspen Adrenaline |
| .,, | 13.27 | - | ✓ DBL Adrenaline |
| | 25.30 | 10 | ✓ Hameln S29 |
| Inj 1 in 10,000, 10 ml ampoule - Up to 5 inj available on a | | 5 | ✓ Hospira |
| ing this to,000, to the ampoule - op to 5 ing available on a | 49.00 | 10 | ✓ Aspen Adrenaline |
| | 49.00 | 10 | Aspen Autenanne |
| Vasodilators | | | |
| Vasouliators | | | |
| | | | |
| HYDRALAZINE HYDROCHLORIDE | | | |
| | 1 | | |
| * Tab 25 mg - Special Authority see SA1321 below - Retai | | 1 | ✓ Hvdralazine |
| | | 1 56 | ✓ Hydralazine |
| * Tab 25 mg - Special Authority see SA1321 below - Retai | | 56 | ✓ Onelink S29 |
| * Tab 25 mg - Special Authority see SA1321 below - Retai | | 56 84 | ✓ Onelink \$29 ✓ AMDIPHARM \$29 |
| * Tab 25 mg - Special Authority see SA1321 below - Retai pharmacy | CBS | 56 84 100 | ✓ Onelink \$29 ✓ AMDIPHARM \$29 ✓ Camber \$29 |
| ** Tab 25 mg – Special Authority see SA1321 below – Retai pharmacy ** Inj 20 mg ampoule | CBS | 56 84 | ✓ Onelink \$29 ✓ AMDIPHARM \$29 |
| * Tab 25 mg - Special Authority see SA1321 below - Retai pharmacy | CBS | 56 84 100 | ✓ Onelink \$29 ✓ AMDIPHARM \$29 ✓ Camber \$29 |
| ** Tab 25 mg – Special Authority see SA1321 below – Retai pharmacy ** Inj 20 mg ampoule ** SA1321 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals va | CBS | 56 84 100 5 | ✓ Onelink \$29 ✓ AMDIPHARM \$29 ✓ Camber \$29 ✓ Apresoline |
| ** Tab 25 mg - Special Authority see SA1321 below - Retai pharmacy ** Inj 20 mg ampoule SA1321 Special Authority for Subsidy | CBS | 56 84 100 5 | ✓ Onelink \$29 ✓ AMDIPHARM \$29 ✓ Camber \$29 ✓ Apresoline |
| ** Tab 25 mg – Special Authority see SA1321 below – Retai pharmacy ** Inj 20 mg ampoule ** SA1321 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals va | CBS | 56 84 100 5 | ✓ Onelink \$29 ✓ AMDIPHARM \$29 ✓ Camber \$29 ✓ Apresoline |
| Tab 25 mg - Special Authority see SA1321 below - Retain pharmacy Inj 20 mg ampoule | CBS | 56 84 100 5 | ✓ Onelink \$29 ✓ AMDIPHARM \$29 ✓ Camber \$29 ✓ Apresoline |
| ** Tab 25 mg – Special Authority see SA1321 below – Retai pharmacy ** Inj 20 mg ampoule ** SA1321 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals value following criteria: Either: 1 For the treatment of refractory hypertension; or | | 56 84 100 5 r renewal unless | ✓ Onelink \$29 ✓ AMDIPHARM \$29 ✓ Camber \$29 ✓ Apresoline snotified for applications meeting |
| ** Tab 25 mg – Special Authority see SA1321 below – Retai pharmacy ** Inj 20 mg ampoule ** SA1321 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals vathe following criteria: Either: 1 For the treatment of refractory hypertension; or 2 For the treatment of heart failure in combination with a result of the second sec | | 56 84 100 5 r renewal unless | ✓ Onelink \$29 ✓ AMDIPHARM \$29 ✓ Camber \$29 ✓ Apresoline snotified for applications meeting |
| * Tab 25 mg – Special Authority see SA1321 below – Retain pharmacy | | 56 84 100 5 r renewal unless | ✓ Onelink \$29 ✓ AMDIPHARM \$29 ✓ Camber \$29 ✓ Apresoline snotified for applications meeting |
| * Tab 25 mg – Special Authority see SA1321 below – Retain pharmacy | 25.90 alid without furthe | 56 84 100 5 r renewal unless who are intolera | ✓ Onelink \$29 ✓ AMDIPHARM \$29 ✓ Camber \$29 ✓ Apresoline s notified for applications meeting and or have not responded to AC |
| * Tab 25 mg – Special Authority see SA1321 below – Retain pharmacy | 25.90 alid without furthe hitrate, in patients | 56 84 100 5 r renewal unless who are intolera | ✓ Onelink \$29 ✓ AMDIPHARM \$29 ✓ Camber \$29 ✓ Apresoline s notified for applications meeting ant or have not responded to AC ✓ Minoxidil Roma \$29 |
| * Tab 25 mg – Special Authority see SA1321 below – Retain pharmacy | 25.90 alid without furthe | 56 84 100 5 r renewal unless who are intolera | ✓ Onelink \$29 ✓ AMDIPHARM \$29 ✓ Camber \$29 ✓ Apresoline s notified for applications meeting and or have not responded to AC |
| * Tab 25 mg – Special Authority see SA1321 below – Retain pharmacy | 25.90 alid without furthe hitrate, in patients | 56 84 100 5 r renewal unless who are intolera | ✓ Onelink \$29 ✓ AMDIPHARM \$29 ✓ Camber \$29 ✓ Apresoline s notified for applications meeting ant or have not responded to AC ✓ Minoxidil Roma \$29 ✓ Loniten |
| # Tab 25 mg – Special Authority see SA1321 below – Retain pharmacy # Inj 20 mg ampoule | | 56 84 100 5 r renewal unless who are intolera | ✓ Onelink \$29 ✓ AMDIPHARM \$29 ✓ Camber \$29 ✓ Apresoline s notified for applications meeting ant or have not responded to AC ✓ Minoxidil Roma \$29 ✓ Loniten ✓ Max Health |
| Tab 25 mg – Special Authority see SA1321 below – Retain pharmacy | | 56 84 100 5 r renewal unless who are intolera 60 100 | ✓ Onelink \$29 ✓ AMDIPHARM \$29 ✓ Camber \$29 ✓ Apresoline s notified for applications meeting ant or have not responded to AC ✓ Minoxidil Roma \$29 ✓ Loniten |
| * Inj 20 mg ampoule * Inj 20 mg ampoule * Inj 20 mg ampoule * SA1321 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals vathe following criteria: Either: 1 For the treatment of refractory hypertension; or 2 For the treatment of heart failure in combination with a rinhibitors and/or angiotensin receptor blockers. MINOXIDIL A Tab 10 mg | | 56 84 100 5 r renewal unless who are intolera 60 100 | ✓ Onelink \$29 ✓ AMDIPHARM \$29 ✓ Camber \$29 ✓ Apresoline s notified for applications meeting ant or have not responded to AC ✓ Minoxidil Roma \$29 ✓ Loniten ✓ Max Health |
| * Inj 20 mg ampoule | | 56 84 100 5 r renewal unless who are intolera 60 100 60 | ✓ Onelink \$29 ✓ AMDIPHARM \$29 ✓ Camber \$29 ✓ Apresoline snotified for applications meeting ant or have not responded to AC ✓ Minoxidil Roma \$29 ✓ Loniten ✓ Max Health ✓ Max Health |
| * Tab 25 mg - Special Authority see SA1321 below - Retain pharmacy | | 56 84 100 5 r renewal unless who are intolera 60 100 | ✓ Onelink \$29 ✓ AMDIPHARM \$29 ✓ Camber \$29 ✓ Apresoline s notified for applications meeting ant or have not responded to AC ✓ Minoxidil Roma \$29 ✓ Loniten ✓ Max Health |
| * Inj 20 mg ampoule | | 56 84 100 5 r renewal unless who are intolera 60 100 60 60 | ✓ Onelink \$29 ✓ AMDIPHARM \$29 ✓ Camber \$29 ✓ Apresoline snotified for applications meeting and or have not responded to AC ✓ Minoxidil Roma \$29 ✓ Loniten ✓ Max Health ✓ Max Health ✓ Hospira |
| * Tab 25 mg - Special Authority see SA1321 below - Retain pharmacy | | 56 84 100 5 r renewal unless who are intolera 60 100 60 | ✓ Onelink \$29 ✓ AMDIPHARM \$29 ✓ Camber \$29 ✓ Apresoline snotified for applications meeting ant or have not responded to AC ✓ Minoxidil Roma \$29 ✓ Loniten ✓ Max Health ✓ Max Health |

| Subsidy | | Fully | Brand or | |
|------------------------|-----|------------|--------------|--|
| (Manufacturer's Price) | 5 | Subsidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

Endothelin Receptor Antagonists

| AMBRISENTAN - Special Authority see SA2253 below - Retail | pharmacy | | |
|---|----------|----|-----------------------|
| Tab 5 mg | 200.00 | 30 | ✓ Ambrisentan Viatris |
| Tab 10 mg | 200.00 | 30 | ✓ Ambrisentan Viatris |

⇒SA2253 Special Authority for Subsidy

Initial application — **(PAH monotherapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁻⁵); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH 2022 (see note below for link to these guidelines) †; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
 - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease: or
 - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
 - 5.1 Ambrisentan is to be used as PAH monotherapy; and
 - 5.2 Any of the following:
 - 5.2.1 Patient has experienced intolerable side effects with both sildenafil and bosentan; or
 - 5.2.2 Patient has an absolute contraindication to sildenafil and an absolute or relative contraindication to bosentan (e.g. due to current use of a combined oral contraceptive or liver disease); or
 - 5.2.3 Patient is a child with idiopathic PAH or PAH secondary to congenital heart disease.

Initial application — **(PAH dual therapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and

| | | | | _ |
|------------------------|-----|------------|--------------|---|
| Subsidy | | Fully | Brand or | |
| (Manufacturer's Price) | S | Subsidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

continued...

- 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
- 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
- 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁵); and
- 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH 2022 (see note below for link to these guidelines) †; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
- 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 All of the following:
 - 5.1 Ambrisentan is to be used as PAH dual therapy; and
 - 5.2 Either:
 - 5.2.1 Patient has tried a PAH monotherapy (sildenafil or bosentan) for at least three months and has not experienced an acceptable response to treatment according to a validated risk stratification tool**; or
 - 5.2.2 Patient has tried PAH dual therapy including bosentan and has experienced intolerable side effects on bosentan; and
 - 5.3 Both:
 - 5.3.1 Patient is presenting in NYHA/WHO functional class III or IV, and in the opinion of the treating clinician would benefit from initial dual therapy; and
 - 5.3.2 Patient has an absolute or relative contraindication to bosentan (e.g. due to current use of a combined oral contraceptive or liver disease).

Initial application — **(PAH triple therapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁵); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH 2022 (see note below for link to these guidelines) †; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or

| Subsidy | | Fully | Brand or |
|------------------------|------------|-------|--------------|
| (Manufacturer's Price) | Subsidised | | Generic |
| \$ | Per | ✓ | Manufacturer |

continued...

- 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
- 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
 - 5.1 Ambrisentan is to be used as PAH triple therapy; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is on the lung transplant list; or
 - 5.2.2 Both:
 - 5.2.2.1 Patient is presenting in NYHA/WHO functional class IV: and
 - 5.2.2.2 Patient has an absolute or relative contraindication to bosentan (e.g. due to current use of a combined oral contraceptive or liver disease); or
 - 5.2.3 Both:
 - 5.2.3.1 Patient has tried PAH dual therapy for at least three months and remains in an unacceptable risk category according to a validated risk stratification tool**; and
 - 5.2.3.2 Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario.

Renewal only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 2 years where the patient is continuing to derive benefit from ambrisentan treatment according to a validated PAH risk stratification tool**.

Notes: † The European Respiratory Journal Guidelines can be found here: 2022 ECS/ERS Guidelines for the diagnosis and treatment of pulmonary hypertension PAH

** the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

BOSENTAN - Special Authority see SA2254 below - Retail pharmacy

| Tab 62.5 mg | 100.00 | 60 | ✓ Bosentan Dr |
|-------------|--------|----|---------------|
| | | | Reddy's |
| Tab 125 mg | 100.00 | 60 | ✓ Bosentan Dr |
| | | | Reddy's |

⇒SA2254 Special Authority for Subsidy

Initial application — **(PAH monotherapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁻⁵); and

| Subsidy (Manufacturer's Price) | , | | Brand or Generic |
|-----------------------------------|-----|---|---------------------|
| \$ | Per | ✓ | Manufacturer |

continued...

- 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) †; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
- 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease: or
- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
 - 5.1 Bosentan is to be used as PAH monotherapy; and
 - 5.2 Any of the following:
 - 5.2.1 Patient has experienced intolerable side effects on sildenafil: or
 - 5.2.2 Patient has an absolute contraindication to sildenafil: or
 - 5.2.3 Patient is a child with idiopathic PAH or PAH secondary to congenital heart disease.

Initial application — **(PAH dual therapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁻⁵); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these quidelines) †: or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**: or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
 - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or
 - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Bosentan is to be used as part of PAH dual therapy; and
- 6 Either:
 - 6.1 Patient has tried a PAH monotherapy (sildenafil) for at least three months and has experienced an inadequate therapeutic response to treatment according to a validated risk stratification tool**; or
 - 6.2 Patient is presenting in NYHA/WHO functional class III or IV, and in the opinion of the treating clinician would likely

| Subsidy | | Fully | Brand or | |
|----------------------|-----|------------|--------------|--|
| (Manufacturer's Pric | ce) | Subsidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

continued...

benefit from initial dual therapy.

Initial application — (PAH triple therapy) only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁻⁵); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these quidelines) † : or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**: or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
 - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or
 - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
 - 5.1 Bosentan is to be used as part of PAH triple therapy; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is on the lung transplant list; or
 - 5.2.2 Patient is presenting in NYHA/WHO functional class IV; or
 - 5.2.3 Both:
 - 5.2.3.1 Patient has tried PAH dual therapy for at least three months and has not experienced an acceptable response to treatment according to a validated risk stratification tool**; and
 - 5.2.3.2 Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario.

Renewal only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 2 years where patient is continuing to derive benefit from bosentan treatment according to a validated PAH risk stratification tool**.

Notes: † The European Respiratory Journal Guidelines can be found here: 2022 ECS/ERS Guidelines for the diagnosis and treatment of pulmonary hypertension PAH

** the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

Phosphodiesterase Type 5 Inhibitors

| SILDENAFIL – Special Authority see SA2255 on the next page – Retail p | oharmacy | | |
|---|----------|----|-----------|
| Tab 25 mg | 0.72 | 4 | ✓ Vedafil |
| Tab 50 mg | | 4 | ✓ Vedafil |
| Tab 100 mg1 | | 12 | ✓ Vedafil |
| | | | |

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| • | Por 🗸 | Manufacturer |

⇒SA2255 Special Authority for Subsidy

Initial application — (Raynaud's Phenomenon*) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has Raynaud's Phenomenon*; and
- 2 Patient has severe digital ischaemia (defined as severe pain requiring hospital admission or with a high likelihood of digital ulceration; digital ulcers; or gangrene); and
- 3 Patient is following lifestyle management (avoidance of cold exposure, sufficient protection, smoking cessation support, avoidance of sympathomimetic drugs); and
- 4 Patient is being treated with calcium channel blockers and nitrates (or these are contraindicated/not tolerated).

Initial application — (Pulmonary arterial hypertension*) only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II. III or IV: and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH is confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) of greater than 20 mmHg; and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) that is less than or equal to 15 mmHg; and
 - 4.1.4 Pulmonary vascular resistance (PVR) of at least 2 Wood Units or greater than 160 International Units (dyn s cm⁵); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH is non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) †; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
 - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or
 - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures.

Initial application — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has a documented history of traumatic or non-traumatic spinal cord injury; and
- 2 Patient has erectile dysfunction secondary to spinal cord injury requiring pharmacological treatment.

Renewal — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Notes: Note: Indications marked with * are Unapproved Indications.

† The European Respiratory Journal Guidelines can be found here: 2022 ECS/ERS Guidelines for the diagnosis and treatment of pulmonary hypertension PAH

** the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

| Subsidy | | Fully | Brand or | |
|------------------------|------|---------|--------------|--|
| (Manufacturer's Price) | Subs | sidised | Generic | |
| \$ | Per | 1 | Manufacturer | |

Prostacyclin Analogues

| EPOPROSTENOL - Special Authority see SA2256 below - Retail pharm | macy | | |
|--|-------|---|-----------|
| Inj 500 mcg vial | 36.61 | 1 | ✓ Veletri |
| Inj 1.5 mg vial | 73.21 | 1 | ✓ Veletri |

⇒SA2256 Special Authority for Subsidy

Initial application — **(PAH dual therapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁵); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these quidelines) † : or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
 - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
 - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 All of the following:
 - 5.1 Epoprostenol is to be used as part of PAH dual therapy with either sildenafil or an endothelin receptor antagonist;
 - 5.2 Patient is presenting in NYHA/WHO functional class IV; and
 - 5.3 Patient has tried a PAH monotherapy for at least three months and remains in an unacceptable risk category according to a validated risk stratification tool.

Initial application — **(PAH triple therapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and

| Subsidy | F | ully | Brand or |
|-----------------------|------------|------|--------------|
| Manufacturer's Price) | Subsidised | | Generic |
| \$ | Per | ✓ | Manufacturer |

continued...

- 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
- 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁻⁵); and
- 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**: or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
- 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
 - 5.1 Epoprostenol is to be used as PAH triple therapy; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is on the lung transplant list; or
 - 5.2.2 Patient is presenting in NYHA/WHO functional class IV; or
 - 5.2.3 Both:
 - 5.2.3.1 Patient has tried PAH dual therapy for at least three months and has not experienced an acceptable response to treatment according to a validated risk stratification tool; and
 - 5.2.3.2 Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario.

Renewal only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 2 years where patient is continuing to derive benefit from epoprostenol treatment according to a validated PAH risk stratification tool**.

Notes: † The European Respiratory Journal Guidelines can be found here: 2022 ECS/ERS Guidelines for the diagnosis and treatment of pulmonary hypertension PAH

** the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

ILOPROST – Special Authority see SA2257 below – Retail pharmacy

Nebuliser soln 10 mcg per ml, 2 ml185.03 Vebulis

⇒SA2257 Special Authority for Subsidy

Initial application — (PAH monotherapy) only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

continued...

- 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁻⁵); and
- 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these quidelines) † : or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**: or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
- 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
 - 5.1 Iloprost is to be used as PAH monotherapy; and
 - 5.2 Either:
 - 5.2.1 Patient has experienced intolerable side effects on sildenafil and both the funded endothelin receptor antagonists (i.e. both bosentan and ambrisentan); or
 - 5.2.2 Patient has an absolute contraindication to sildenafil and an absolute or relative contraindication to endothelin receptor antagonists.

Initial application — (PAH dual therapy) only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁵); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) † : or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**: or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
 - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
 - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 All of the following:
 - 5.1 Iloprost is to be used as PAH dual therapy with either sildenafil or an endothelin receptor antagonist; and

| Subsidy | | Fully | Brand or |
|------------------------|-----|------------|--------------|
| (Manufacturer's Price) | | Subsidised | Generic |
| \$ | Per | 1 | Manufacturer |

continued...

- 5.2 Either:
 - 5.2.1 Patient has an absolute contraindication to or has experienced intolerable side effects on sildenafil; or
 - 5.2.2 Patient has an absolute or relative contraindication to or experienced intolerable side effects with a funded endothelin receptor antagonist; and
- 5.3 Either:
 - 5.3.1 Patient has tried a PAH monotherapy for at least three months and remains in an unacceptable risk category according to a validated risk stratification tool**; or
 - 5.3.2 Patient is presenting in NYHA/WHO functional class III or IV, and in the opinion of the treating clinician would benefit from initial dual therapy.

Initial application — **(PAH triple therapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁻⁵); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) †; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
 - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease: or
 - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
 - 5.1 Iloprost is to be used as PAH triple therapy; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is on the lung transplant list; or
 - 5.2.2 Patient is presenting in NYHA/WHO functional class IV; or
 - 5.2.3 Both:
 - 5.2.3.1 Patient has tried PAH dual therapy for at least three months and has not experienced an acceptable response to treatment according to a validated risk stratification tool**; and
 - 5.2.3.2 Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario.

Renewal only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 2 years where patient is continuing to derive benefit from iloprost treatment according to a validated PAH risk stratification tool**.

| | Subsidy | F | ully | Brand or |
|----|-----------------------|------------|------|--------------|
| (N | lanufacturer's Price) | Subsidised | | Generic |
| | \$ | Per | 1 | Manufacturer |

continued...

Notes: † The European Respiratory Journal Guidelines can be found here: 2022 ECS/ERS Guidelines for the diagnosis and treatment of pulmonary hypertension PAH

^{**} the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

Subsidy Fully Brand or
(Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

Antiacne Preparations

For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 94

ADAPALENE

- a) Maximum of 30 g per prescription
- b) Only on a prescription

| b) Only on a proscription | | | |
|---|----------------|---------|----------------------------|
| Gel 0.1% | 22.89 | 30 g OP | Differin |
| ISOTRETINOIN - Special Authority see SA2023 below - R | etail pharmacy | | |
| Cap 5 mg | 11.26 | 60 | Oratane |
| Cap 10 mg | 18.75 | 120 | ✓ Oratane |
| Cap 20 mg | 26.73 | 120 | ✓ Oratane |

⇒SA2023 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around isotretinoin and is competent to prescribe isotretinoin; and
- 3 Either:
 - 3.1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and that they must not become pregnant during treatment and for a period of one month after the completion of treatment; or
 - 3.2 Patient is not of child bearing potential.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and that they must not become pregnant during treatment and for a period of one month after the completion of treatment; or
- 2 Patient is not of child bearing potential.

TRETINOIN

Crm 0.5 mg per g − Maximum of 50 g per prescription16.82 50 g OP ✓ ReTrieve

Antibacterials Topical

For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 94

HYDROGEN PEROXIDE

| * Crm 1% | 8.56 | 10 g OP | ✓ Crystaderm |
|-----------|---------|---------|--------------|
| MUPIROCIN | | | |
| Oint 2% | 6.60 | 15 g OP | |
| | (13.00) | - | Bactroban |

- a) Only on a prescription
- b) Not in combination

| | | | /ENIVI | ATOLOGICALS |
|---|------------------------------------|-------------|-------------------|-------------------------------------|
| | Subsidy (Manufacturer's F \$ | Price) Subs | Fully idised | Brand or Generic Manufacturer |
| SODIUM FUSIDATE [FUSIDIC ACID] Crm 2% | 1.69 | 5 g OP | √ <u>F</u> | oban |
| a) Maximum of 5 g per prescription b) Only on a prescription c) Not in combination Oint 2% a) Maximum of 5 g per prescription b) Only on a prescription | 1.69 | 5 g OP | ✓ <u>F</u> | 'oban |
| c) Not in combination SULFADIAZINE SILVER Crm 1% | 10.80 | 50 g OP | - | Flamazine Ascend 829 |
| a) Up to 250 g available on a PSOb) Not in combination | 10.44 | | . , | Nacciiu |
| Antifungals Topical | | | | |
| For systemic antifungals, refer to INFECTIONS, Antifungals, page AMOROLFINE a) Only on a prescription b) Not in combination | | | | |
| Nail soln 5% CLOTRIMAZOLE | 21.87 | 5 ml OP | ✓ <u>N</u> | <u>//ycoNail</u> |
| ** Crm 1% a) Only on a prescription b) Not in combination | 1.10 | 20 g OP | √ <u>(</u> | Clomazol |
| Soln 1% a) Only on a prescription b) Not in combination | 4.36 (7.55) | 20 ml OP | (| Canesten |
| ECONAZOLE NITRATE Crm 1% | 1.00 | 20 g OP | | |
| a) Only on a prescription b) Not in combination | (8.09) | - | F | Pevaryl |
| Foaming soln 1%, 10 ml sachets | 9.89 | 3 | - | Davie and |

Pevaryl

(18.64)

a) Only on a prescription b) Not in combination

| | Subsidy | O., 1 | Fully | Brand or |
|--|--|---|--|---|
| | (Manufacturer's F \$ | Per Subs | sidised • | Generic Manufacturer |
| ICONAZOLE NITRATE | | | | |
| F Crm 2% | 0.90 | 15 g OP | ✓ <u>N</u> | <u>//ultichem</u> |
| a) Only on a prescription | | | | |
| b) Not in combination | | | | |
| £ Lotn 2% | | 30 ml OP | | |
| | (10.03) | | | Daktarin |
| a) Only on a prescription | | | | |
| b) Not in combination Tinct 2% | 4.26 | 30 ml OP | | |
| 1 III Gt 2 /0 | (12.10) | 30 1111 OF | г | Daktarin |
| a) Only on a prescription | (12.10) | | _ | Jaktann |
| b) Not in combination | | | | |
| , | | | | |
| Antipruritic Preparations | | | | |
| ALAMINE | | | | |
| a) Only on a prescription | | | | |
| b) Not in combination | | | _ | |
| Crm, aqueous, BP | 3.45 | 100 g | ✓ t | nealthE Calamine |
| | 4.4. "1.0005 | | | Aqueous |
| healthE Calamine Aqueous to be Principal Supply of | on 1 April 2025 | | | |
| ROTAMITON | | | | |
| a) Only on a prescription b) Not in combination | | | | |
| | | | | |
| | 3.40 | 20 a OP | √ 1 | tch-Sootha |
| Ćrm 10% | 3.49 | 20 g OP | ✓ <u>I</u> | tch-Soothe |
| Crm 10%ENTHOL – Only in combination | | · · | _ | tch-Soothe |
| Crm 10% ENTHOL – Only in combination 1) Only in combination with a dermatological base or | | · · | _ | tch-Soothe |
| Crm 10%ENTHOL – Only in combination | | · · | _ | tch-Soothe |
| Crm 10% ENTHOL – Only in combination 1) Only in combination with a dermatological base or | proprietary Topical C | · · | Plain | tch-Soothe MidWest |
| Crm 10% ENTHOL – Only in combination 1) Only in combination with a dermatological base or 2) With or without other dermatological galenicals. | proprietary Topical C | Corticosteriod – | Plain | |
| Crm 10% ENTHOL – Only in combination 1) Only in combination with a dermatological base or 2) With or without other dermatological galenicals. Crystals | proprietary Topical C | Corticosteriod – 25 g | Plain | MidWest |
| Crm 10% | proprietary Topical C 6.92 29.60 | Corticosteriod – 25 g 100 g | Plain | MidWest |
| Crm 10% ENTHOL – Only in combination 1) Only in combination with a dermatological base or 2) With or without other dermatological galenicals. Crystals | proprietary Topical C 6.92 29.60 | Corticosteriod – 25 g 100 g | Plain | MidWest |
| Crm 10% | proprietary Topical C 6.92 29.60 | Corticosteriod – 25 g 100 g | Plain | MidWest |
| Crm 10% | proprietary Topical C 6.92 29.60 | Corticosteriod – 25 g 100 g | Plain | MidWest |
| Crm 10% | proprietary Topical C6.92 29.60 AND RELATED AGE | Corticosteriod – 25 g 100 g | Plain • N | MidWest |
| Crm 10% | proprietary Topical C | Corticosteriod – 25 g 100 g | Plain M M | MidWest MidWest |
| Crm 10% | proprietary Topical C | Corticosteriod – 25 g 100 g ENTS, page 84 15 g OP 50 g OP 15 g OP | Plain N N | MidWest MidWest Diprosone Diprosone Diprosone |
| Crm 10% | proprietary Topical C | Corticosteriod – 25 g 100 g ENTS, page 84 15 g OP 50 g OP 15 g OP 50 g OP | Plain | MidWest MidWest Diprosone Diprosone Diprosone Diprosone Diprosone |
| Crm 10% | proprietary Topical C | Corticosteriod – 25 g 100 g ENTS, page 84 15 g OP 50 g OP 15 g OP | Plain | MidWest MidWest Diprosone Diprosone Diprosone |
| Crm 10% | proprietary Topical C | 25 g 100 g 100 g ENTS, page 84 15 g OP 50 g OP 15 g OP 50 g OP 30 g OP | Plain A A C C C C C C C C C C C | MidWest MidWest Diprosone Diprosone Diprosone Diprosone Diprosone Diprosone OV |
| Crm 10% | proprietary Topical C | 25 g 100 g 100 g ENTS, page 84 15 g OP 50 g OP 15 g OP 50 g OP 30 g OP | Plain I M I M I M I M I M I M I M I | MidWest MidWest Diprosone Diprosone Diprosone Diprosone Diprosone Oiprosone OV Beta Cream |
| Crm 10% | proprietary Topical C | 25 g 100 g 25 g 100 g 25 T 100 g 25 T 20 T 20 T 20 T 20 T 20 T 20 T 20 T 20 | Plain N N L L L L L L L L L L L | AidWest AidWest Diprosone Diprosone Diprosone Diprosone Diprosone Oiprosone OV Beta Cream Beta Ointment |
| Crm 10% | proprietary Topical C | 25 g 100 g 100 g ENTS, page 84 15 g OP 50 g OP 15 g OP 50 g OP 30 g OP | Plain N N L L L L L L L L L L L | MidWest MidWest Diprosone Diprosone Diprosone Diprosone Diprosone Oiprosone OV Beta Cream |
| Crm 10% | proprietary Topical C | 25 g 100 g 25 g 100 g 25 T 100 g 25 T 20 T 20 T 20 T 20 T 20 T 20 T 20 T 20 | Plain N N L L L L L L L L L L L | AidWest AidWest Diprosone Diprosone Diprosone Diprosone Diprosone Oiprosone OV Beta Cream Beta Ointment |
| Crm 10% | 2.96 AND RELATED AGE 2.96 36.00 2.96 36.00 4.33 5.85 7.90 30.00 | 25 g 100 g 25 g 100 g 25 T 100 g 25 T 20 T 20 T 20 T 20 T 20 T 20 T 20 T 20 | Plain N L L L L L L L L L L L L | AidWest AidWest Diprosone Diprosone Diprosone Diprosone Diprosone Oiprosone OV Beta Cream Beta Ointment |

| | Subsidy | | Fully | Brand or |
|---|--------------------|-----------------------|--------------|-------------------------|
| | (Manufacturer's Pi | rice) Sub: Per | sidised • | Generic Manufacturer |
| OLODETA COLIE DUTA (DATE | Ψ | 1 61 | | Wallulacturer |
| CLOBETASONE BUTYRATE | T 00 | 00 = OD | | |
| Crm 0.05% | | 30 g OP | г., | |
| | (10.00) | | EU | ımovate |
| HYDROCORTISONE | | | | |
| * Crm 1% – Only on a prescription | | 30 g OP | ✓ <u>Et</u> | |
| W Daviday Only in combination | 20.40 | 500 g | | <u>oumed</u> |
| * Powder – Only in combination | 49.95 | 25 g | ✓ AE | |
| galenicals | ai Corticosteriou | - Flaili) Willi | Ji Williou | t other dermatological |
| • | | | | |
| HYDROCORTISONE AND PARAFFIN LIQUID AND LANOLIN | | | | |
| Lotn 1% with paraffin liquid 15.9% and lanolin 0.6% – Only o | | 0E0 ml | ./ DI | I atm UC |
| a prescription | 12.83 | 250 ml | ₩ <u>DF</u> | Lotn HC |
| HYDROCORTISONE BUTYRATE | | 05 | ٠. | |
| Lipocream 0.1% | | 100 g OP | | coid Lipocream |
| Oint 0.1% | | 100 g OP 100 ml OP | ✓ Lo | ocola ocold Crelo |
| Milky emul 0.1% | 12.33 | 100 mi OP | ♥ Lo | ocola Creio |
| METHYLPREDNISOLONE ACEPONATE | 4.05 | 45 · OD | | Lt |
| Crm 0.1% | | 15 g OP | _ | <u>lvantan</u> |
| Oint 0.1% | 4.95 | 15 g OP | ♥ AC | <u>lvantan</u> |
| MOMETASONE FUROATE | | | <i>-</i> | |
| Crm 0.1% | | 15 g OP | _ | ocon Alcohol Free |
| Oint 0.1% | 3.50 | 50 g OP | ✓ Ele | ocon Alcohol Free |
| OIIIL 0.1% | 3.50 | 15 g OP 50 g OP | ✓ Ele | |
| Lotn 0.1% | | 30 ml OP | ✓ Ele | |
| TRIAMCINOLONE ACETONIDE | | 00 1111 01 | | <u> </u> |
| Crm 0.02% | 6.40 | 100 g OP | ✓ Ar | istocort |
| Oint 0.02% | | 100 g OF | _ | istocort |
| Ont 0.02 / 0 | | 100 g O1 | · <u>Ai</u> | 15100011 |
| Corticosteroids - Combination | | | | |
| | | | | |
| BETAMETHASONE VALERATE WITH SODIUM FUSIDATE [FU | | 45 · OD | | |
| Crm 0.1% with sodium fusidate (fusidic acid) 2% | | 15 g OP | г., | anian wt |
| a) Marian was of 45 are as a security to a | (10.45) | | Fu | cicort |
| a) Maximum of 15 g per prescriptionb) Only on a prescription | | | | |
| , , . | | | | |
| HYDROCORTISONE WITH MICONAZOLE – Only on a prescrip | | 15 = OD | Z M: | |
| * Crm 1% with miconazole nitrate 2% | | 15 g OP | V IVII | creme H |
| HYDROCORTISONE WITH NATAMYCIN AND NEOMYCIN - O | , , , | | | |
| Oint 1% with natamycin 1% and neomycin sulphate 0.5% | 3.35 | 15 g OP | ✓ Pii | mafucort |
| TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYCI | N AND NYSTAT | ΊN | | |
| Crm 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 mg | | | | |
| and gramicidin 250 mcg per g - Only on a prescription | | 15 g OP | | |
| | (9.28) | | Via | aderm KC |

Subsidy (Manufacturer's Price) Fully Subsidised Per

Brand or Generic Manufacturer

Barrier Creams and Emollients

| _ | | _ | |
|-----|------|------|-----|
| Rar | riar | (re | ams |
| | | | |

| DIMETHICONE | | |
|---|-------------|-------------------------------|
| * Crm 5% pump bottle4.30 | 500 ml OP | ✓ <u>healthE</u> |
| | | Dimethicone 5% |
| * Crm 10% pump bottle4.52 | 500 ml OP | ✓ healthE |
| | | Dimethicone 10% |
| ZINC AND CASTOR OIL | | |
| * Oint | 500 g | ✓ Evara |
| - | | |
| Emollients | | |
| AQUEOUS CREAM | | |
| Crm | 100 g | ✓ healthE Aqueous |
| | ŭ | Cream SLS Free |
| 1.65 | 500 g | ✓ Evara |
| 1.73 | • | ✓ GEM Aqueous |
| | | Cream |
| (healthE Aqueous Cream SLS Free Crm to be delisted 1 March 2025) | | |
| (GEM Aqueous Cream Crm to be delisted 1 March 2025) | | |
| CETOMACROGOL | | |
| * Crm BP | 500 g | ✓ Cetomacrogol-AFT |
| CETOMACROGOL WITH GLYCEROL | | |
| Crm 90% with glycerol 10%2.13 | 500 ml OP | ✓ Evara |
| 3.50 | 1,000 ml OP | ✓ <u>Evara</u> |
| EMULSIFYING OINTMENT | | |
| * Oint BP | 500 g | Emulsifying |
| | | Ointment ADE |
| OIL IN WATER EMULSION | | |
| * Crm | 500 g | ✓ Fatty Cream AFT |
| 2.10 | | ✓ Fatty Emulsion |
| | | Cream (Evara) |
| Fatty Emulsion Cream (Evara) to be Principal Supply on 1 April 2025 | | |
| (Fatty Cream AFT Crm to be delisted 1 April 2025) | | |
| PARAFFIN | | |
| Oint liquid paraffin 50% with white soft paraffin 50%4.94 | 500 g OP | ✓ White Soft Liquid |

100 g OP

Paraffin AFT

✓ healthE Urea Cream

UREA

✓ EVARA White Soft Paraffin

| | | _ | |
|--|---------------------|--------------|--------------------------------|
| | Subsidy | | Fully Brand or |
| | (Manufacturer's I | Price) Subsi | dised Generic |
| | \$ | Per | ✓ Manufacturer |
| WOOL FAT WITH MINERAL OIL - Only on a prescription | | | |
| * Lotn hydrous 3% with mineral oil | 5.60 | 1,000 ml | |
| • | (14.96) | | DP Lotion |
| | (20.53) | | Alpha-Keri Lotion |
| | ` 1.40 [′] | 250 ml OP | , |
| | (5.87) | | DP Lotion |
| | `5.60 [°] | 1,000 ml | |
| | (23.91) | | BK Lotion |
| | ` 1.40 [′] | 250 ml OP | |
| | (7.73) | | BK Lotion |
| Other Dermatological Bases | | | |
| PARAFFIN | | | |
| White soft - Only in combination | 4.74 | 450 g | ✓ EVARA White Soft Paraffin |

Only in combination with a dermatological galenical or as a diluent for a proprietary Topical Corticosteroid – Plain.

19.00

2.500 a

| Minor Skin Infections | | | |
|--|--------|---------|--------------------|
| POVIDONE IODINE | | | |
| Oint 10% | 7.40 | 65 g OP | ✓ Betadine |
| a) Maximum of 130 g per prescription | | | |
| b) Only on a prescription | | | |
| Antiseptic Solution 10% | 4.99 | 100 ml | ✓ Riodine |
| Antiseptic soln 10% | 3.83 | 15 ml | ✓ Riodine |
| | 6.99 | 500 ml | ✓ Riodine |
| Skin preparation, povidone iodine 10% with 30% alcohol | 1.63 | 100 ml | |
| | (3.48) | | Betadine Skin Prep |

Parasiticidal Preparations

DIMETHICONE

| * Lotn 4% | 200 ml OP | ✓ <u>healthE</u> <u>Dimethicone 4%</u> <u>Lotion</u> |
|---|-----------|--|
| IVERMECTIN - Special Authority see SA2294 below - Retail pharmacy | | |
| Tab 3 mg - Up to 100 tab available on a PSO | 4 | ✓ Stromectol |

- 1) PSO for institutional use only. Must be endorsed with the name of the institution for which the PSO is required and a valid Special Authority for patient of that institution.
- 2) Ivermectin available on BSO provided the BSO includes a valid Special Authority for a patient of the institution.
- For the purposes of subsidy of ivermectin, institution means age related residential care facilities, disability care facilities or prisons.

⇒SA2294 Special Authority for Subsidy

Initial application — (Scables) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

DERMATOLOGICALS

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

continued...

Fither:

- 1 The person has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
- 2 Both:
 - 2.1 The person has a confirmed diagnosis of scabies or is a close contact of a scabies case; and
 - - 2.2.1 The person is unable to complete topical therapy: or
 - 2.2.2 Previous treatment with topical therapy has been tried and not cleared the infestation.

Initial application — (Other parasitic infections) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 filariasis: or
- 2 cutaneous larva migrans (creeping eruption); or
- 3 strongyloidiasis.

Renewal — (Scabies) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria: Either:

- 1 The person has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
- 2 Both:
 - 2.1 The person has a confirmed diagnosis of scabies or is a close contact of a scabies case; and
 - 2.2 Fither:
 - 2.2.1 The person is unable to complete topical therapy; or
 - 2.2.2 Previous treatment with topical therapy has been tried and not cleared the infestation.

Renewal — (Other parasitic infections) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 filariasis: or
- 2 cutaneous larva migrans (creeping eruption); or
- 3 strongyloidiasis.

PERMETHRIN

Lotn 5% 30 ml OP ✓ A-Scabies

Psoriasis and Eczema Preparations

| ACTIRETIN — Special Authority see SA2024 below — Ret | ali pharmacy | | |
|--|--------------|----|--------------|
| Cap 10 mg | 26.20 | 60 | Novatretin |
| Cap 25 mg | 57.37 | 60 | ✓ Novatretin |
| | | | |

⇒SA2024 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around acitretin and is competent to prescribe acitretin; and
- 3 Either:
 - 3.1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and that they must not become pregnant during treatment and for a period of three years after the completion of treatment: or

DERMATOLOGICALS

| Subsidy (Manufacturer's Price) | S | Fully ubsidised | Brand or Generic |
|-----------------------------------|-----|--------------------|---------------------|
| \$ | Per | 1 | Manufacturer |

continued...

3.2 Patient is not of child bearing potential.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Fither:

- 1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and that they must not become pregnant during treatment and for a period of three years after the completion of treatment; or
- 2 Patient is not of child bearing potential.

BETAMETHASONE DIPROPIONATE WITH CALCIPOTRIOL

| Foam spray 500 mcg with calcipotriol 50 mcg per g | 59.95 | 60 g OP | Enstilar |
|--|------------------|------------------|----------------------------|
| Gel 500 mcg with calcipotriol 50 mcg per g | 40.92 | 60 g OP | Daivobet |
| Oint 500 mcg with calcipotriol 50 mcg per g | | 30 g OP | ✓ Daivobet |
| CALCIPOTRIOL | | | |
| Oint 50 mcg per g | 40.00 | 120 g OP | Daivonex |
| COAL TAR | | | |
| Soln BP - Only in combination | 36.25 | 200 ml | ✓ Midwest |
| 1) I In to 10% only in combination with a dermatologic | al hace or propr | iotany Tonical (| orticostariod - Pl |

- 1) Up to 10% only in combination with a dermatological base or proprietary Topical Corticosteriod Plain
- 2) With or without other dermatological galenicals.

COAL TAR WITH ALLANTOIN, MENTHOL, PHENOL AND SULPHUR

| Soin 5% with sulphur 0.5%, menthol 0.75%, phenol 0.5% and | | | |
|---|--------|---------|--------------|
| allantoin crm 2.5% | 6.59 | 75 g OP | |
| | (8.00) | J | Egopsoryl TA |
| | 3.43 | 30 g OP | • • • |
| | (4.35) | • | Egopsoryl TA |
| COAL TAR WITH SALICYLIC ACID AND SULPHUR | | | |
| Soln 12% with salicylic acid 2% and sulphur 4% oint | 4.97 | 25 g OP | ✓ Coco-Scalp |
| · | 7.95 | 40 g OP | ✓ Coco-Scalp |

PIMECROLIMUS - Special Authority see SA1970 below - Retail pharmacy

- a) Maximum of 15 g per prescription
- b) Note: a maximum of 15 g per prescription and no more than one prescription per 12 weeks.

⇒SA1970 Special Authority for Subsidy

Initial application only from a dermatologist, paediatrician, ophthalmologist or any relevant practitioner on the recommendation of a dermatologist, paediatrician or ophthalmologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has atopic dermatitis on the eyelid; and
- 2 Patient has at least one of the following contraindications to topical corticosteroids: periorificial dermatitis, rosacea, documented epidermal atrophy, documented allergy to topical corticosteroids, cataracts, glaucoma, or raised intraocular pressure.

PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORESCEIN - Only on a prescription

* Soln 2.3% with trolamine laurilsulfate and fluorescein sodium..........5.41 500 ml / Pinetarsol

DERMATOLOGICALS

| | Subsidy (Manufacturer's Pr \$ | ice) Subs Per | Fully sidised | Brand or Generic Manufacturer |
|---|---|--------------------------|-------------------|--|
| SALICYLIC ACID Powder – Only in combination | | 250 g al Corticostero | | flidwest ain or collodion flexible |
| SULPHUR Precipitated – Only in combination | 6.35 | 100 g | ✓ N | /lidwest |
| Only in combination with a dermatological base With or without other dermatological galenicals. | or proprietary Topica | al Corticostero | oid – Pla | ain |
| TACROLIMUS Oint 0.1% – Special Authority see SA2074 below – Retail pharmacy | 33.00 | 30 g OP | _ | <u>Cematop</u> |
| ➤ SA2074 Special Authority for Subsidy Initial application only from a dermatologist, paediatrician or a paediatrician, . Approvals valid without further renewal unless Both: 1 Patient has atopic dermatitis on the face; and 2 Patient has at least one of the following contraindication documented epidermal atrophy or documented allergy to | notified for applications to topical corticos | ons meeting t | the follo | wing criteria: |
| Scalp Preparations | | | | |
| BETAMETHASONE VALERATE * Scalp app 0.1% CLOBETASOL PROPIONATE | 12.95 | 100 ml OP | ✓ <u>B</u> | Beta Scalp |
| Scalp app 0.05% HYDROCORTISONE BUTYRATE | 6.26 | 30 ml OP | ✓ 0 | <u>Dermol</u> |
| Scalp lotn 0.1%KETOCONAZOLE | 6.57 | 100 ml OP | √ L | ocoid |
| Shampoo 2% | 3.23 4.09 | 100 ml OP | _ | Sebizole Sebizole |
| a) Maximum of 100 ml per prescriptionb) Only on a prescription | | | | _ |
| Sunscreens | | | | |
| SUNSCREENS, PROPRIETARY – Subsidy by endorsement Only if prescribed for a patient with severe photosensitivity | secondary to a defi | ned clinical co | ondition | and the prescription is |
| endorsed accordingly. Lotn, | 6.50 | 200 g OP | ✓ <u>N</u> | Marine Blue Lotion |

SPF 50+

DERMATOLOGICALS

Subsidy Fully Brand or
(Manufacturer's Price) Subsidised Generic
\$ Per ✓ Manufacturer

Wart Preparations

For salicylic acid preparations refer to PSORIASIS AND ECZEMA PREPARATIONS, page 72

PODOPHYLLOTOXIN

- a) Maximum of 3.5 ml per prescription
- b) Only on a prescription

Other Skin Preparations

Antineoplastics

| ı | | - 1 | \sim | п | \sim | ш | п | ١٨ | \sim | 11 | S | $^{\sim}$ | п | ш | IN | |
|---|--|-----|--------|---|--------|---|---|----|--------|----|---|-----------|---|---|----|--|
| | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | |

IMIQUIMOD

GENITO-URINARY SYSTEM

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

Contraceptives - Non-hormonal

Condoms

| COND | | 14.05 | 144 | ✓ Moments |
|------|--|---------------|-----------|-----------------------------------|
| | mm – Up to 144 dev available on a PSOmm | | 10 | ✓ Moments |
| • 55 | | 14.25 | 144 | ✓ Moments |
| | a) Maximum of 60 dev per prescription | 14.20 | 177 | - momento |
| | b) Up to 60 dev available on a PSO | | | |
| ÷ 53 | mm, 0.05 mm thickness | 1.15 | 10 | ✓ Moments |
| | , | 14.25 | 144 | ✓ Moments |
| | a) Up to 60 dev available on a PSO | | | |
| | b) Maximum of 60 dev per prescription | | | |
| 53 | mm, chocolate, brown | 1.15 | 10 | ✓ Moments |
| | | 14.25 | 144 | ✓ Moments |
| | a) Up to 60 dev available on a PSO | | | |
| | b) Maximum of 60 dev per prescription | | | |
| 53 | mm, strawberry, red | 1.15 | 10 | ✓ Moments |
| | | 14.25 | 144 | ✓ Moments |
| | a) Up to 60 dev available on a PSO | | | |
| | b) Maximum of 60 dev per prescription | | | |
| 56 | mm | | 10 | ✓ Moments |
| | | 14.50 | 144 | ✓ Moments |
| | a) Maximum of 60 dev per prescription | | | |
| | b) Up to 60 dev available on a PSO | | | |
| 56 | mm, 0.05 mm thickness | | 12 | ✓ Gold Knight |
| | | 24.10 | 144 | Gold Knight |
| | a) Up to 60 dev available on a PSO | | | |
| | b) Maximum of 60 dev per prescription | | | |
| 56 | mm, 0.05mm thickness (bulk pack) | 20.17 | 144 | Gold Knight |
| | a) Maximum of 60 dev per prescription | | | |
| | b) Up to 60 dev available on a PSO | | | |
| 56 | mm, 0.08 mm thickness | | 10 | ✓ Moments |
| | | 14.25 | 144 | ✓ Moments |
| | a) Up to 60 dev available on a PSO | | | |
| | b) Maximum of 60 dev per prescription | | | |
| 56 | mm, 0.08 mm thickness, red | | 10 | ✓ Moments |
| | | 14.25 | 144 | ✓ Moments |
| | a) Up to 60 dev available on a PSO | | | |
| | b) Maximum of 60 dev per prescription | 4 70 | 40 | A Cald Madala |
| 56 | mm, chocolate | | 12 | ✓ Gold Knight |
| | a) Ha ta 00 day aya'labla ay a BOO | 21.45 | 144 | ✓ Gold Knight |
| | a) Up to 60 dev available on a PSO | | | |
| EC | b) Maximum of 60 dev per prescription | 1 70 | 10 | ✓ Cald Value |
| 56 | mm, strawberry | 1.79 21.45 | 12 144 | ✓ Gold Knight |
| | a) Ha ta CO day available an a DCO | ∠1.45 | 144 | ✓ Gold Knight |
| | a) Up to 60 dev available on a PSO | | | |
| 60 | b) Maximum of 60 dev per prescription mm | 1 00 | 12 | ✓ Gold Knight XL |
| 60 | | 21.89 | 144 | ✓ Gold Knight XL ✓ Gold Knight XL |
| | a) Maximum of 60 day now proporting | 21.09 | 144 | - Goid Killyill AL |
| | a) Maximum of 60 dev per prescription | | | |

GENITO-URINARY SYSTEM

| Subsidy (Manufacturer's Price) | | Fully Subsidised | Brand or Generic | |
|-----------------------------------|-----|---------------------|---------------------|--|
| \$ | Per | / | Manufacturer | |

- a) Maximum of 60 dev per prescription
- b) Up to 60 dev available on a PSO

Contraceptive Devices

INTRA-UTERINE DEVICE

- a) Up to 40 dev available on a PSO
- b) Only on a PSO

| * | IUD 29.1 mm length × 23.2 mm width29. | .80 1 | ✓ | Choice 380 7med |
|---|---------------------------------------|-------|---|-----------------|
| | | | | Nsha Silver/ |
| | | | | copper Short |
| * | IUD 33.6 mm length × 29.9 mm width | .80 1 | • | TCu 380 Plus |
| | • | | | Normal |
| * | IUD 35.5 mm length × 19.6 mm width | .00 1 | • | Cu 375 Standard |

Contraceptives - Hormonal

Combined Oral Contraceptives

⇒SA0500 Special Authority for Alternate Subsidy

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Either:
 - 1.1 Patient is on a Social Welfare benefit; or
 - 1.2 Patient has an income no greater than the benefit; and
- 2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient is on a Social Welfare benefit; or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- · on a Social Welfare benefit; or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

ETHINYLOESTRADIOL WITH DESOGESTREL

| * | Tab 20 mcg with desogestrel 150 mcg and 7 inert tab - Up to | | | |
|---|---|-------|----|---------------|
| | 84 tab available on a PSO | 10.00 | 84 | ✓ Mercilon 28 |

| | | | | - | |
|--|------------------------|-----|-------------------|------------------------|--|
| | Subsidy | | Fully | Brand or | |
| | (Manufacturer's Price) | | Subsidised | Generic | |
| | \$ | Per | ✓ | Manufacturer | |
| ETHINYLOESTRADIOL WITH LEVONORGESTREL | | | | | |
| * Tab 20 mcg with levonorgestrel 100 mcg and 7 inert tablets | _ | | | | |
| Up to 84 tab available on a PSO | 1.50 | 84 | ✓ L | o-Oralcon 20 ED | |
| * Tab 30 mcg with levonorgestrel 150 mcg | | 63 | _ | | |
| | (16.50) | | N | licrogynon 30 | |
| a) Higher subsidy of \$15.00 per 63 tab with Special Aut b) Up to 63 tab available on a PSO * Tab 30 mcg with levonorgestrel 150 mcg and 7 inert tablets | - | | | • | |
| Up to 84 tab available on a PSO | 1.50 | 84 | √ <u>0</u> | ralcon 30 ED | |
| ETHINYLOESTRADIOL WITH NORETHISTERONE | | | | | |
| Tab 35 mcg with norethisterone 1 mg and 7 inert tab - Up to 84 tab available on a PSO | | 84 | | lyacen revinor 1/28 | |
| Tab 35 mcg with norethisterone 500 mcg and 7 inert tab - U to 84 tab available on a PSO | • | 84 | ✓ N | lorimin | |
| | | | | | |

Progestogen-only Contraceptives

⇒SA0500 Special Authority for Alternate Subsidy

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Fither
 - 1.1 Patient is on a Social Welfare benefit; or
 - 1.2 Patient has an income no greater than the benefit; and
- 2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient is on a Social Welfare benefit: or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- on a Social Welfare benefit; or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

LEVONORGESTREL

| * | Tab 30 mcg - Up to 84 tab available on a PSO | .16.50 22.00 | 84 112 | ✓ Microlut✓ Microlut |
|----|--|-----------------|-----------|---|
| * | Subdermal implant (2 x 75 mg rods) – Up to 3 pack available on a PSO | 106.92 | 1 | ✓ <u>Jadelle</u> |
| ME | DROXYPROGESTERONE ACETATE Inj 150 mg per ml, 1 ml syringe – Up to 5 inj available on a PSO | 9.18 | 1 | ✓ Depo-Provera |

GENITO-URINARY SYSTEM

| | Subsidy (Manufacturer's Price) | Per | Fully Subsidised | Brand or Generic Manufacturer |
|---|-----------------------------------|-----|---------------------|--|
| NORETHISTERONE Tab 350 mcg - Up to 84 tab available on a PSO | 12.25 | 84 | ✓ N | Norethinderone - CDC Noriday Noriday 28 |

Emergency Contraceptives

LEVONORGESTREL

- - a) Maximum of 2 tab per prescription
 - b) Up to 5 tab available on a PSO
 - c) Note: Direct Provision by a pharmacist permitted under the provisions in Part I of Section A.

Antiandrogen Oral Contraceptives

Prescribers may code prescriptions "contraceptive" (code "O") when used as indicated for contraception. The period of supply and prescription charge will be as per other contraceptives, as follows:

- A maximum \$5.00 prescription charge (patient co-payment) may apply.
- prescription may be written for up to six months supply.

Prescriptions coded in any other way are subject to any non contraceptive prescription charges that apply, and the non-contraceptive period of supply. ie. Prescriptions may be written for up to three months supply.

CYPROTERONE ACETATE WITH ETHINYLOESTRADIOL

Gynaecological Anti-infectives

| ACETIC ACID WITH HYDROXYQUINOLINE AND RICINOLEIC ACID Jelly with glacial acetic acid 0.94%, hydroxyquinoline sulphate | | | |
|--|----------|------------|--|
| 0.025%, glycerol 5% and ricinoleic acid 0.75% with applicator8.43 (24.87) | 100 g OP | Aci-Jel | |
| CLOTRIMAZOLE | | | |
| * Vaginal crm 1% with applicators | 35 g OP | ✓ Clomazol | |
| * Vaginal crm 2% with applicators | 20 g OP | ✓ Clomazol | |
| MICONAZOLE NITRATE | | | |
| * Vaginal crm 2% with applicator6.89 | 40 g OP | ✓ Micreme | |
| NYSTATIN | | | |
| Vaginal crm 100,000 u per 5 g with applicator(s)5.70 | 75 g OP | ✓ Nilstat | |

Myometrial and Vaginal Hormone Preparations

| | PSOPSO | 160.00 | 5 | ✓ DBL Ergometrine |
|----|--------------------------------|--------|---------|-------------------|
| 0E | STRIOL | | | |
| * | Crm 1 mg per g with applicator | 6.95 | 15 g OP | ✓ Ovestin |
| | Pessaries 500 mcg | | 15 | ✓ Ovestin |

FRGOMETRINE MAI FATE

✓ David One Step Cassette Pregnancy Test

| | Subsidy | | Fully | Brand or |
|---|-----------------------|-----------|-------------------|--------------------------------------|
| | (Manufacturer's Price | e) Sub | sidised | Generic |
| | \$ | Per | • | Manufacturer |
| OXYTOCIN – Up to 5 inj available on a PSO | | | | |
| Inj 5 iu per ml, 1 ml ampoule | 4.98 | 5 | 1 | Oxytocin BNM |
| Inj 10 iu per ml, 1 ml ampoule | | 5 | _ | Dxytocin BNM |
| , apoulo | 11.96 | 10 | _ | Oxytocin |
| | 11.00 | 10 | • | Panpharma |
| OXYTOCIN WITH ERGOMETRINE MALEATE – Up to 5 inj avai Inj 5 iu with ergometrine maleate 500 mcg per ml, 1 ml ampo Pregnancy Tests - hCG Urine | | 5 | √ <u>9</u> | Syntometrine |
| BETA-HCG LOW SENSITIVITY URINE TEST KIT – Up to 15 tes Note: For use in abortion services only. | st available on a PS | 0 | | |
| Midstream | 16.28 | 1 test OP | ✓ (| CheckTop |
| PREGNANCY TESTS - HCG URINE | | | | |
| a) Up to 200 test available on a PSO | | | | |
| b) Only on a PSO | | | | |
| Cassette | 12.00 4 | 0 test OP | √ 9 | Smith BioMed Rapid Pregnancy Test |

16.00

David One Step Cassette Pregnancy Test to be Principal Supply on 1 March 2025 (Smith BioMed Rapid Pregnancy Test Cassette to be delisted 1 March 2025)

Urinary Agents

For urinary tract Infections refer to INFECTIONS, Antibacterials, page 113

5-Alpha Reductase Inhibitors

FINASTERIDE – Special Authority see SA0928 below – Retail pharmacy

★ Tab 5 mg4.79 100 ✓ Ricit

⇒SA0928 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 Either:
 - 2.1 The patient is intolerant of non-selective alpha blockers or these are contraindicated; or
 - 2.2 Symptoms are not adequately controlled with non-selective alpha blockers.

Alpha-1A Adrenoreceptor Blockers

TAMSULOSIN HYDROCHLORIDE − Special Authority see SA1032 below − Retail pharmacy

* Cap 400 mcg22.31 100

✓ Tamsulosin-Rex

⇒SA1032 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has symptomatic benian prostatic hyperplasia; and
- 2 The patient is intolerant of non-selective alpha blockers or these are contraindicated.

GENITO-URINARY SYSTEM

| | Subsidy (Manufacturer's I \$ | Price) Subs Per | Fully sidised | Brand or Generic Manufacturer |
|--|------------------------------------|--------------------|-------------------|--|
| Other Urinary Agents | | | | |
| DXYBUTYNIN | | | | |
| * Tab 5 mg | 5.42 | 100 | ✓ A | llchemy Oxybutynin |
| POTASSIUM CITRATE | | | | ,, |
| Oral liq 3 mmol per ml - Special Authority see SA1083 b | | | | |
| Retail pharmacy | 37.49 | 200 ml OP | ✓ E | Biomed |
| ⇒SA1083 Special Authority for Subsidy nitial application from any relevant practitioner. Approvals 3oth: | valid for 12 months | for applications | meetir | ng the following criteria: |
| 1 The patient has recurrent calcium oxalate urolithiasis;2 The patient has had more than two renal calculi in the | | ne application. | | |
| Renewal from any relevant practitioner. Approvals valid for 2 penefitting from the treatment. | 2 years where the tro | eatment remain | s appro | priate and the patient is |
| SODIUM CITRO-TARTRATE | | | | |
| F Grans eff 4 g sachets | 3.50 | 28 | √ <u>L</u> | <u>Iral</u> |
| SOLIFENACIN SUCCINATE | 1.05 | 00 | | alifamasin |
| Tab 5 mg | 1.95 | 30 | • 5 | olifenacin succinate Max Health |
| | 2.05 | | | olifenacin Viatris |
| Tab 10 mg | 3.53 | 30 | √ S | solifenacin succinate Max Health |
| | 3.72 | | √ S | Solifenacin Viatris |
| Solifenacin Viatris Tab 5 mg to be delisted 1 June 2025) Solifenacin Viatris Tab 10 mg to be delisted 1 June 2025) | | | | |
| Detection of Substances in Urine | | | | |
| DRTHO-TOLIDINE | | | | |
| Compound diagnostic sticks | 7.50 | 50 test OP | | |
| | (8.25) | | H | lemastix |
| ETRABROMOPHENOL | 40.00 | 1001-105 | | |
| Blue diagnostic strips | 13.92 | 100 test OP | ✓ A | Albustix |
| Obstetric Preparations | | | | |
| Antiprogesterones | | | | |
| MIFEPRISTONE | | | | |
| Tab 200 mg - Up to 15 tab available on a PSO | 83.90 180.00 | 1 3 | | lifegyne lifegyne |

| Subsidy | | Fully | Brand or | |
|----------------------|-----|------------|--------------|--|
| (Manufacturer's Pric | ce) | Subsidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |
| | | | | |

Calcium Homeostasis

| CALCITONIN | | |
|---|----|---------------------|
| * Inj 100 iu per ml, 1 ml ampoule | 5 | ✓ Miacalcic |
| CINACALCET – Special Authority see SA2170 below – Retail pharmacy | | |
| Tab 30 mg - Wastage claimable25.24 | 28 | ✓ Cinacalet Devatis |

⇒SA2170 Special Authority for Subsidy

Initial application — (parathyroid carcinoma or calciphylaxis) only from a nephrologist or endocrinologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 All of the following:
 - 1.1 The patient has been diagnosed with a parathyroid carcinoma (see Note); and

- 1.2 The patient has persistent hypercalcaemia (serum calcium greater than or equal to 3 mmol/L) despite previous first-line treatments including sodium thiosulfate (where appropriate) and bisphosphonates; and
- 1.3 The patient is symptomatic; or
- 2 All of the following:
 - 2.1 The patient has been diagnosed with calciphylaxis (calcific uraemic arteriolopathy); and
 - 2.2 The patient has symptomatic (e.g. painful skin ulcers) hypercalcaemia (serum calcium greater than or equal to 3 mmol/L); and
 - 2.3 The patient's condition has not responded to previous first-line treatments including bisphosphonates and sodium thiosulfate.

Renewal — (parathyroid carcinoma or calciphylaxis) only from a nephrologist or endocrinologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient's serum calcium level has fallen to < 3mmol/L; and
- 2 The patient has experienced clinically significant symptom improvement.

Note: This does not include parathyroid adenomas unless these have become malignant.

Initial application — (primary hyperparathyroidism) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has primary hyperparathyroidism; and
- 2 Either:
 - 2.1 Patient has hypercalcaemia of more than 3 mmol/L with or without symptoms; or
 - 2.2 Patient has hypercalcaemia of more than 2.85 mmol/L with symptoms; and
- 3 Surgery is not feasible or has failed; and
- 4 Patient has other comorbidities, severe bone pain, or calciphylaxis.

Initial application — (secondary or tertiary hyperparathyroidism) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient has tertiary hyperparathyroidism and markedly elevated parathyroid hormone (PTH) with hypercalcaemia; or
 - 1.2 Patient has symptomatic secondary hyperparathyroidism and elevated PTH; and
- 2 Patient is on renal replacement therapy; and
- 3 Any of the following:
 - 3.1 Residual parathyroid tissue has not been localised despite repeat unsuccessful parathyroid explorations; or

continued...

Cinacalet Devatis

| Subsidy | | Fully | Brand or |
|------------------------|-----|------------|--------------|
| (Manufacturer's Price) | | Subsidised | Generic |
| \$ | Per | ✓ | Manufacturer |

continued...

- 3.2 Parathyroid tissue is surgically inaccessible; or
- 3.3 Parathyroid surgery is not feasible.

Renewal — (secondary or tertiary hyperparathyroidism) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 The patient has had a kidney transplant, and following a treatment free interval of at least 12 weeks a clinically acceptable parathyroid hormone (PTH) level to support ongoing cessation of treatment has not been reached; or
- 2 The patient has not received a kidney transplant and trial of withdrawal of cinacalcet is clinically inappropriate.

ZOLEDRONIC ACID

| Inj 4 mg per 5 ml, vial | 15.65 | 1 | ✓ Zoledronic acid |
|-------------------------|-------|---|-------------------|
| | | | Viatris |

Corticosteroids and Related Agents for Systemic Use

BETAMETHASONE SODIUM PHOSPHATE WITH BETAMETHASONE ACETATE

| * Inj 3.9 mg with betamethasone acetate 3 mg per ml, 1 ml | 5 | Celestone Chronodose |
|---|----------|------------------------------|
| DEXAMETHASONE | | |
| * Tab 0.5 mg - Up to 60 tab available on a PSO1.80 | 30 | ✓ Dexmethsone |
| * Tab 4 mg - Up to 30 tab available on a PSO | 30 | ✓ <u>Dexmethsone</u> |
| Oral liq 1 mg per ml53.86 | 25 ml OP | ✓ Biomed |
| DEXAMETHASONE PHOSPHATE | | |
| Dexamethasone phosphate injection will not be funded for oral use. | | |
| * Inj 4 mg per ml, 1 ml ampoule – Up to 5 inj available on a PSO7.86 | 10 | ✓ <u>Hameln</u> |
| * Inj 4 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO13.10 | 10 | ✓ <u>Hameln</u> |
| FLUDROCORTISONE ACETATE | | |
| * Tab 100 mcg11.46 | 100 | ✓ Florinef |
| HYDROCORTISONE | | |
| * Tab 5 mg8.10 | 100 | Douglas |
| * Tab 20 mg20.32 | 100 | ✓ Douglas |
| * Inj 100 mg vial | 1 | ✓ Solu-Cortef |
| a) Not on a BSO | | |
| b) Up to 5 inj available on a PSO | | |
| METHYLPREDNISOLONE | | |
| * Tab 4 mg112.00 | 100 | ✓ Medrol |
| * Tab 100 mg223.10 | 20 | ✓ Medrol |
| METHYLPREDNISOLONE (AS SODIUM SUCCINATE) | | |
| Inj 40 mg vial22.30 | 1 | ✓ Solu-Medrol-Act- O-Vial |
| Inj 125 mg vial34.10 | 1 | ✓ Solu-Medrol-Act- |
| ., 1=0g va | • | O-Vial |
| Inj 500 mg vial26.88 | 1 | ✓ Solu-Medrol-Act- O-Vial |
| Inj 1 g vial32.84 | 1 | ✓ Solu-Medrol |

| | Subsidy | | Fully Brand or |
|---|-----------------------|-------------|----------------------------------|
| | (Manufacturer's Price | e) S Per | Subsidised Generic Manufacturer |
| AETUNA BEEDMISSU ONE VOETATE | \$ | rei | ▼ ividilulacturer |
| IETHYLPREDNISOLONE ACETATE | 47.00 | _ | ✓ Dama Madual |
| Inj 40 mg per ml, 1 ml vial | 47.06 | 5 | ✓ Depo-Medrol |
| REDNISOLONE | | | _ |
| Oral liq 5 mg per ml – Up to 30 ml available on a PSO Restricted to children under 12 years of age. | 6.00 | 30 ml OP | P ✓ Redipred |
| REDNISONE | | | |
| F Tab 1 mg | 18.58 | 500 | ✓ Prednisone Clinect |
| Fab 2.5 mg | 21.04 | 500 | ✓ Prednisone Clinect |
| Tab 5 mg - Up to 30 tab available on a PSO | 19.30 | 500 | Prednisone Clinect |
| Tab 20 mg - Up to 30 tab available on a PSO | 50.51 | 500 | ✓ Prednisone Clinect |
| ETRACOSACTRIN | | | |
| Inj 250 mcg per ml, 1 ml ampoule | 86.25 | 1 | ✓ Synacthen |
| , <u>-</u> 559 pc, apcas | | • | ✓ UK Synacthen |
| Inj 1 mg per ml, 1 ml ampoule | 690.00 | 1 | ✓ Synacthen Depot |
| ., | | | ✓ Synacthene |
| | | | Retard \$29 |
| RIAMCINOLONE ACETONIDE | | | |
| Inj 10 mg per ml, 1 ml ampoule | 21.42 | 5 | ✓ Kenacort-A 10 |
| Inj 40 mg per ml, 1 ml ampoule | | 5 5 | ✓ Kenacort-A 10 ✓ Kenacort-A 40 |
| inj 40 mg per mi, i mi ampoule | 52.03 | 5 | Nenacon-A 40 |
| Sex Hormones Non Contraceptive | | | |
| | | | |
| Androgen Agonists and Antagonists | | | |
| YPROTERONE ACETATE | | | |
| Tab 50 mg | 17.05 | 50 | ✓ Siterone |
| Tab 100 mg | | 50 | ✓ Siterone |
| ESTOSTERONE | | | |
| Gel (transdermal) 16.2 mg per g | 52.00 | 88 g OP | ✓ Testogel |
| | | 55 g 51 | 10010901 |
| ESTOSTERONE CIPIONATE | 0F 00 | 1 | / Dono Testestaus |
| Inj 100 mg per ml, 10 ml vial | 05.00 | ı | ✓ Depo-Testosterone |
| ESTOSTERONE ESTERS | | | |
| Inj 250 mg per ml, 1 ml | 12.98 | 1 | Sustanon Ampoules |
| ESTOSTERONE UNDECANOATE | | | |
| Cap 40 mg - Subsidy by endorsement | 36.00 | 100 | ✓ Steril-Gene S29 |
| Subsidy by endorsement – subsidised for patients who | | rone und | lecanoate cap 40mg prior to |
| 1 November 2021 and the prescription is endorsed ac | | | |
| where there exists a record of prior dispensing of testo | | | |
| | 00.00 | | / D |

Inj 250 mg per ml, 4 ml vial......86.00

✓ Reandron 1000

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

Hormone Replacement Therapy - Systemic

Oestrogens

| - | STRADIOL | | | |
|----|--|-------------------|---------|---------------------------------------|
| * | Tab 1 mg | 4.12 | 28 OP | |
| | | (11.10) | | Estrofem |
| * | Tab 2 mg | 4.12 | 28 OP | |
| | | (11.10) | | Estrofem |
| | Gel (transdermal) 0.06% (750 mcg/actuation) | 14.25 | 80 g OP | ✓ Estrogel |
| | Patch 25 mcg per day | | 8 | ✓ Estradiol TDP Mylan |
| | | 13.50 | | ✓ Estraderm MX S29 |
| | | 14.50 | | ✓ Estradot |
| | | | | |
| | | 21.35 | | ✓ Lyllana |
| | a) No more than 2 patch per week | | | |
| | b) Only on a prescription | | | |
| | Patch 50 mcg per day | 10.75 | 8 | Estradiol TDP Mylan |
| | | | | Estradiol Viatris |
| | | 14.50 | | ✓ Estraderm MX S29 |
| | | | | ✓ Estradiol Sandoz |
| | | | | ✓ Estradot |
| | | 21.55 | | ✓ Lyllana |
| | Alamana than Omatah manasala | 21.00 | | Lylialia |
| | a) No more than 2 patch per week | | | |
| | b) Only on a prescription | | | |
| | Patch 75 mcg per day | 11.88 | 8 | Estradiol TDP Mylan |
| | | | | Estradiol Viatris |
| | | 14.50 | | Estradiol Sandoz |
| | | | | ✓ Estradot |
| | | 22.37 | | ✓ Lyllana |
| | a) No more than 2 patch per week | | | • |
| | b) Only on a prescription | | | |
| | Patch 100 mcg per day | 12 95 | 8 | ✓ Estradiol TDP Mylan |
| | Tatori 100 mag par day | 12.00 | Ü | ✓ Estradiol Viatris |
| | | 14.50 | | ✓ Estradiol Sandoz |
| | | 14.50 | | ✓ Estradot |
| | | | | |
| | | 15.50 | | ✓ Estraderm MX S29 |
| | | 22.77 | | ✓ Lyllana |
| | a) No more than 2 patch per week | | | |
| | b) Only on a prescription | | | |
| ΩF | STRADIOL VALERATE | | | |
| | Tab 1 mg | 10.26 | 84 | ✓ Progynova |
| | • | | 84 | ✓ Progynova |
| | Tab 2 mg | 12.30 | 04 | ♥ Progynova |
| | STROGENS | | | |
| * | Conjugated, equine tab 300 mcg | 3.01 | 28 | |
| | - | (19.25) | | Premarin |
| * | Conjugated, equine tab 625 mcg | 4.12 [′] | 28 | |
| | | (19.25) | | Premarin |

| | Subsidy | | Fully Brand or |
|---|------------------------------|-------|----------------------------------|
| | (Manufacturer's Price) \$ | Per | Subsidised Generic Manufacturer |
| Progestogens | | | |
| MEDROXYPROGESTERONE ACETATE | | | |
| ₹ Tab 2.5 mg | 6.56 | 30 | ✓ Provera |
| - | 8.75 | 56 | ✓ Provera |
| ₹ Tab 5 mg | 9.80 | 56 | ✓ Provera |
| | 20.13 | 100 | |
| Tab 10 mg | 10.28 | 30 | ✓ Provera |
| Progestogen and Oestrogen Combined Prepara | tions | | |
| ESTRADIOL WITH NORETHISTERONE | | | |
| Tab 1 mg with 0.5 mg norethisterone acetate | 5.40 | 28 OF | |
| | (18.10) | | Kliovance |
| Tab 2 mg with 1 mg norethisterone acetate | | 28 OF | |
| | (18.10) | | Kliogest |
| Tab 2 mg with 1 mg norethisterone acetate (10), and 2 mg | F 40 | 00.0- | n |
| oestradiol tab (12) and 1 mg oestradiol tab (6) | | 28 OF | |
| | (18.10) | | Trisequens |
| Other Oestrogen Preparations | | | |
| ESTRIOL | | | |
| F Tab 2 mg | 7.70 | 30 | ✓ Ovestin |
| Other Progestogen Preparations | | | |
| EVONORGESTREL | | | |
| Intra-uterine device 52 mg | 269.50 | 1 | ✓ Mirena |
| Intra-uterine device 13.5 mg | | 1 | ✓ Jaydess |
| EDROXYPROGESTERONE ACETATE | | | • |
| Tab 100 mg | 133.57 | 100 | ✓ Provera HD |
| | | 100 | 1101014112 |
| ORETHISTERONE · Tab 5 mg – Up to 30 tab available on a PSO | 5.40 | 30 | ✓ Primolut N |
| | | 30 | ▼ FINNOIULIN |
| ROGESTERONE : Cap 100 mg | 1/1 05 | 30 | ✓ Utropostan |
| Cap 100 mg | 14.00 | 30 | ✓ <u>Utrogestan</u> |
| Thyroid and Antithyroid Agents | | | |
| ARBIMAZOLE | | | |
| ₹ Tab 5 mg | 7.56 | 100 | ✓ Neo-Mercazole |
| EVOTHYROXINE | | | |
| EVOTHTHOXINE ₹ Tab 25 mcg | 5 55 | 90 | ✓ Synthroid |
| Tab 50 mcg | | 28 | ✓ Mercury Pharma |
| 55 mg | 5.79 | 90 | ✓ Synthroid |
| | 64.28 | 1,000 | _ * |
| Tablet 50 mcg | | 200 | |
| : Tab 100 mcg | | 28 | ✓ Mercury Pharma |
| | 6.01 | 90 | ✓ Synthroid |
| | 66.78 | 1,000 | |
| Tablet 100 mcg | 13.36 | 200 | ✓ Eltroxin |

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

| | Subsidy (Manufacturer's Price) | | Fully Subsidised | Brand or Generic |
|---|-----------------------------------|-----|---------------------|---------------------|
| | \$ | Per | 1 | Manufacturer |
| PROPYLTHIOURACIL - Special Authority see SA1199 below - | Retail pharmacy | | | |
| Tab 50 mg | 35.00 | 100 | √ P | TU \$29 |
| ⇒SA1199 Special Authority for Subsidy | | | | |

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

- 1 The patient has hyperthyroidism; and
- 2 The patient is intolerant of carbimazole or carbimazole is contraindicated.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

Trophic Hormones

Growth Hormones

| SO | MATROPIN (OMNITROPE) - Special Authority see SA2032 belo | ow – Retail ph | armacy | |
|----|--|----------------|--------|-------------|
| * | Inj 5 mg cartridge | 80.21 | 1 | ✓ Omnitrope |
| * | Inj 10 mg cartridge | 80.21 | 1 | ✓ Omnitrope |
| * | Inj 15 mg cartridge | 139.50 | 1 | ✓ Omnitrope |

⇒SA2032 Special Authority for Subsidy

Initial application — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

Fither:

- 1 Growth hormone deficiency causing symptomatic hypoglycaemia, or with other significant growth hormone deficient sequelae (e.g. cardiomyopathy, hepatic dysfunction) and diagnosed with GH < 5 mcg/l on at least two random blood samples in the first 2 weeks of life, or from samples during established hypoglycaemia (whole blood glucose < 2 mmol/l using a laboratory device); or</p>
- 2 All of the following:
 - 2.1 Height velocity < 25th percentile for age adjusted for bone age/pubertal status if appropriate over 6 or 12 months using the standards of Tanner and Davies (1985); and</p>
 - 2.2 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
 - 2.3 Peak growth hormone value of < 5.0 mcg per litre in response to two different growth hormone stimulation tests. In children who are 5 years or older, GH testing with sex steroid priming is required; and</p>
 - 2.4 If the patient has been treated for a malignancy, they should be disease free for at least one year based upon follow-up laboratory and radiological imaging appropriate for the malignancy, unless there are strong medical reasons why this is either not necessary or appropriate; and
 - 2.5 Appropriate imaging of the pituitary gland has been obtained.

Renewal — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 2 Height velocity is greater than or equal to 25th percentile for age (adjusted for bone age/pubertal status if appropriate) while on growth hormone treatment, as calculated over six months using the standards of Tanner and Davis (1985); and
- 3 Height velocity is greater than or equal to 2.0 cm per year, as calculated over 6 months; and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (**Turner syndrome**) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

| | Fully bsidised | Brand or Generic |
|------------|-------------------|---------------------|
| \$ Per | | Manufacturer |

continued...

All of the following:

- 1 The patient has a post-natal genotype confirming Turner Syndrome; and
- 2 Height velocity is < 25th percentile over 6-12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is < 14 years.

Renewal — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile for age (while on growth hormone calculated over 6 to 12 months using the Ranke's Turner Syndrome growth velocity charts); and
- 2 Height velocity is greater than or equal to 2 cm per year, calculated over six months; and
- 3 A current bone age is 14 years or under; and
- 4 No serious adverse effect that the specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 3 standard deviations below the mean for age or for bone age if there is marked growth acceleration or delay; and
- 2 Height velocity is < 25th percentile for age (adjusted for bone age/pubertal status if appropriate), as calculated over 6 to 12 months using the standards of Tanner and Davies(1985); and
- 3 A current bone age is < 14 years or under (female patients) or < 16 years (male patients); and
- 4 The patient does not have severe chronic disease (including malignancy or recognized severe skeletal dysplasia) and is not receiving medications known to impair height velocity.

Renewal — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred.

Initial application — (short stature due to chronic renal insufficiency) only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 2 standard deviations below the mean; and
- 2 Height velocity is < 25th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is to 14 years or under (female patients) or to 16 years or under (male patients); and
- 4 The patient is metabolically stable, has no evidence of metabolic bone disease and absence of any other severe chronic disease; and
- 5 The patient is under the supervision of a specialist with expertise in renal medicine; and
- 6 Either:
 - 6.1 The patient has a GFR less than or equal to 30 ml/min/1.73m² as measured by the Schwartz method (Height(cm)/plasma creatinine (umol/l)) × 40 = corrected GFR (ml/min/1.73m²) in a child who may or may not be receiving dialysis; or

| Subsidy (Manufacturer's Price) | Sub | Fully sidised | Brand or Generic |
|-----------------------------------|-----|---------------|---------------------|
| \$ | Per | 1 | Manufacturer |

continued...

6.2 The patient has received a renal transplant and has received < 5mg/ m²/day of prednisone or equivalent for at least 6 months..

Renewal — (short stature due to chronic renal insufficiency) only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone has occurred;
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not experienced significant biochemical or metabolic deterioration confirmed by diagnostic results; and
- 7 The patient has not received renal transplantation since starting growth hormone treatment; and
- 8 If the patient requires transplantation, growth hormone prescription should cease before transplantation and a new application should be made after transplantation based on the above criteria.

Initial application — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria: All of the following:

- 1 The patient has a diagnosis of Prader-Willi syndrome that has been confirmed by genetic testing or clinical scoring criteria:
- 2 The patient is aged six months or older; and
- 3 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
- 4 Sleep studies or overnight oximetry have been performed and there is no obstructive sleep disorder requiring treatment, or if an obstructive sleep disorder is found, it has been adequately treated under the care of a paediatric respiratory physician and/or ENT surgeon; and
- 5 Either:
 - 5.1 Both:
 - 5.1.1 The patient is aged two years or older; and
 - 5.1.2 There is no evidence of type II diabetes or uncontrolled obesity defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months; or
 - 5.2 The patient is aged between six months and two years and a thorough upper airway assessment is planned to be undertaken prior to treatment commencement and at six to 12 weeks following treatment initiation.

Renewal — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred: and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not developed type II diabetes or uncontrolled obesity as defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months.

Initial application — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9

| Subsidy (Manufacturer's Price) | | Fully Subsidised | Brand or Generic | |
|-----------------------------------|-----|---------------------|---------------------|--|
| \$ | Per | | Manufacturer | |

continued...

months for applications meeting the following criteria:

All of the following:

- 1 The patient has a medical condition that is known to cause growth hormone deficiency (e.g. surgical removal of the pituitary for treatment of a pituitary tumour); and
- 2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
- 3 The patient has severe growth hormone deficiency (see notes); and
- 4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
- 5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

Renewal — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 All of the following:
 - 1.1 The patient has been treated with somatropin for < 12 months; and
 - 1.2 There has been an improvement in Quality of Life defined as a reduction of at least 8 points on the Quality of Life Assessment of Growth Hormone Deficiency in Adults (QoL-AGHDA®) score from baseline; and
 - 1.3 Serum IGF-I levels have been increased within ±1SD of the mean of the normal range for age and sex; and
 - 1.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients, or 1 mg per day for female patients; or

2 All of the following:

- 2.1 The patient has been treated with somatropin for more than 12 months; and
- 2.2 The patient has not had a deterioration in Quality of Life defined as a 6 point or greater increase from their lowest QoL-AGHDA® score on treatment (other than due to obvious external factors such as external stressors); and
- 2.3 Serum IGF-I levels have continued to be maintained within ±1SD of the mean of the normal range for age and sex (other than for obvious external factors); and
- 2.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients or 1 mg per day for female patients; or

3 All of the following:

- 3.1 The patient has had a Special Authority approval for somatropin for childhood deficiency in children and no longer meets the renewal criteria under this indication; and
- 3.2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
- 3.3 The patient has severe growth hormone deficiency (see notes); and
- 3.4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
- 3.5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

| Subsidy | | Fully | Brand or |
|------------------------|-----|-----------|--------------|
| (Manufacturer's Price) | Su | ubsidised | Generic |
| \$ | Per | ✓ | Manufacturer |

continued...

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

GnRH Analogues

| GOSENELIN | | | |
|--------------------------|-------|---|-----------|
| Implant 3.6 mg, syringe | 66.48 | 1 | ✓ Zoladex |
| Implant 10.8 mg, syringe | | 1 | ✓ Zoladex |

I FUPRORFI IN

COCEDELIN

Additional subsidy by endorsement where the patient is a child or adolescent and is unable to tolerate administration of goserelin and the prescription is endorsed accordingly.

| inj 3.75 mg premied duai chamber syninge – | nigher subsidy of | | |
|---|-------------------|---|----------------------|
| \$221.60 per 1 inj with Endorsement | 66.48 | 1 | |
| | (221.60) | | Lucrin Depot 1-month |
| Ini 11 OF ma profilled dual abambar auringa | Higher subsidy | | |

Inj 11.25 mg prefilled dual chamber syringe — Higher subsidy of \$591.68 per 1 inj with Endorsement......177.50

(591.68) Lucrin Depot 3-month

Vasopressin Agonists

DECMODDECCIN

| Wafer 120 mcg | 47.00 | 30 | ✓ Minirin Melt |
|-------------------------------|-------|---------|-----------------|
| DESMOPRESSIN ACETATE | | | |
| Tab 100 mcg | 25.00 | 30 | ✓ Minirin |
| Tab 200 mcg | | 30 | ✓ Minirin |
| ▲ Nasal spray 10 mcg per dose | 34.95 | 6 ml OP | ✓ Desmopressin- |
| , , , | | | PH&T |
| Inj 4 mcg per ml, 1 ml | 67.18 | 10 | ✓ Minirin |

Other Endocrine Agents

CABERGOLINE

| | | Tab 0.5 mg - Maximum of 2 tab per prescription; can be |
|----------------------------|---|--|
| Dostinex | 2 | waived by Special Authority see SA2070 below4.43 |
| Dostinex | 8 | 17.94 |

⇒SA2070 Special Authority for Waiver of Rule

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

1 Hyperprolactinemia; or

| Subsidy (Manufacturer's Price) | , | Fully Subsidised | Brand or Generic |
|-----------------------------------|-----|---------------------|---------------------|
| \$ | Per | ✓ | Manufacturer |

continued...

- 2 Acromegaly*; or
- 3 Inhibition of lactation.

Renewal — (for patients who have previously been funded under Special Authority form SA1031) from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has previously held a valid Special Authority which has expired and the treatment remains appropriate and the patient is benefiting from treatment.

Note: Indication marked with * is an unapproved indication.

| CLOMIFENE (| CITRATE |
|-------------|---------|
|-------------|---------|

| Tab 50 mg | 29.84 | 10 | ✓ Mylan Clomiphen S29 |
|--------------------------|--------|----|--------------------------|
| METYRAPONE Cap 250 mg | 558.00 | 50 | ✓ Metopirone |

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| ` ¢ ′ | Por ./ | Manufacturar |

Anthelmintics

| ALBENDAZOLE – Special Authority see SA1318 bek | ow – Retail pharmacy | | |
|--|----------------------|----|----------------|
| Tab 400 mg | 469.20 | 60 | ✓ Eskazole S29 |

⇒SA1318 Special Authority for Subsidy

Initial application only from an infectious disease specialist or clinical microbiologist. Approvals valid for 6 months where the patient has hydatids.

Renewal only from an infectious disease specialist or clinical microbiologist. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefitting from the treatment.

MEBENDAZOLE - Only on a prescription

| Tab 100 mg | 5.18 | 6 | ✓ Vermox |
|--------------------------|--------|-------|--------------|
| Oral lig 100 mg per 5 ml | | 15 ml | |
| 31. | (7.83) | | Vermox |
| PRAZIQUANTEL | | | |
| Tab 600 mg | 68.00 | 8 | ✓ Biltricide |

Antibacterials

- a) For topical antibacterials, refer to DERMATOLOGICALS, page 66
- b) For anti-infective eye preparations, refer to SENSORY ORGANS, page 265

Cephalosporins and Cephamycins

| Cap 250 mg | 25.85 | 100 | ✓ Ranbaxy-Cefaclor |
|--|-------|--------|--------------------------------------|
| Grans for oral liq 125 mg per 5 ml - Wastage claimable | | 100 ml | ✓ Ranbaxy-Cefaclor |
| CEFALEXIN | | | |
| Cap 250 mg | 3.85 | 20 | ✓ Cephalexin ABM |
| Cap 500 mg | 5.85 | 20 | ✓ Cephalexin ABM |
| Grans for oral liq 25 mg per ml - Wastage claimable | 7.88 | 100 ml | ✓ Flynn |
| Grans for oral liq 50 mg per ml - Wastage claimable | 10.38 | 100 ml | ✓ Flynn |
| | 11.75 | | Cefalexin Sandoz |
| CFFAZOLIN - Subsidy by endorsement | | | |

Only if prescribed for dialysis or cellulitis in accordance with a Health NZ Hospital approved protocol and the prescription is endorsed accordingly.

| Inj 500 mg vial3.39 | 5 | ✓ Cefazolin-AFT |
|---------------------|---|-----------------|
| Inj 1 g vial | 5 | ✓ Cefazolin-AFT |
| Inj 2 g vial | 5 | ✓ Cefazolin-AFT |

CEFTRIAXONE - Subsidy by endorsement

- a) Up to 10 inj available on a PSO
- b) Subsidised only if prescribed for a dialysis or cystic fibrosis patient, or the treatment of gonorrhoea, or the treatment of pelvic inflammatory disease, or the treatment of suspected meningococcal disease, and the prescription or PSO is endorsed accordingly.

| Inj 500 mg vial | 0.79 | 1 | ✓ Ceftriaxone-AFT |
|-----------------|------|---|-------------------|
| lnj 1 g vial | 3.59 | 5 | ✓ Ceftriaxone-AFT |

CEFUROXIME AXETIL - Subsidy by endorsement

Only if prescribed for prophylaxis of endocarditis and the prescription is endorsed accordingly

| Tab 250 mg | 20 | ✓ Ascend- |
|------------|----|----------------|
| | | Cefuroxime S29 |

| Subsidy | | Fully | Brand or |
|------------------------|-----|-----------|--------------|
| (Manufacturer's Price) | S | ubsidised | Generic |
| \$ | Per | ✓ | Manufacturer |

Macrolides

AZITHROMYCIN – Maximum of 5 days treatment per prescription; can be waived by Special Authority see SA1683 below A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised on Special Authority.

| Tab 250 mg | .8.19 | 30 | ✓ Apo-Azithromycin |
|---|-------|-------|--------------------|
| Tab 500 mg - Up to 8 tab available on a PSO | | 2 | ✓ Zithromax |
| Grans for oral liq 200 mg per 5 ml (40 mg per ml) - Wastage | | | |
| claimable | 16.97 | 15 ml | ✓ Zithromax |

⇒SA1683 Special Authority for Waiver of Rule

Initial application — (bronchiolitis obliterans syndrome, cystic fibrosis and atypical Mycobacterium infections) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following:

- 1 Patient has received a lung transplant, stem cell transplant, or bone marrow transplant and requires treatment for bronchiolitis obliterans syndrome*; or
- 2 Patient has received a lung transplant and requires prophylaxis for bronchiolitis obliterans syndrome*; or
- 3 Patient has cystic fibrosis and has chronic infection with Pseudomonas aeruginosa or Pseudomonas-related gram negative organisms*; or
- 4 Patient has an atypical Mycobacterium infection.

Note: Indications marked with * are unapproved indications.

Initial application — (non-cystic fibrosis bronchiectasis*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 For prophylaxis of exacerbations of non-cystic fibrosis bronchiectasis*; and
- 2 Patient is aged 18 and under; and
- 3 Either:
 - 3.1 Patient has had 3 or more exacerbations of their bronchiectasis, within a 12 month period; or
 - 3.2 Patient has had 3 acute admissions to hospital for treatment of infective respiratory exacerbations within a 12 month period.

Note: Indications marked with * are unapproved indications.

Renewal — (non-cystic fibrosis bronchiectasis*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has completed 12 months of azithromycin treatment for non-cystic fibrosis bronchiectasis; and
- 2 Following initial 12 months of treatment, the patient has not received any further azithromycin treatment for non-cystic fibrosis bronchiectasis for a further 12 months, unless considered clinically inappropriate to stop treatment; and
- 3 The patient will not receive more than a total of 24 months' azithromycin cumulative treatment (see note).

The patient must not have had more than 1 prior approval.

Note: No further renewals will be subsidised. A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised. Indications marked with * are unapproved indications

CLARITHROMYCIN - Maximum of 500 mg per prescription; can be waived by Special Authority see SA1857 below

⇒SA1857 Special Authority for Waiver of Rule

Initial application — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years for applications meeting the following criteria:

Fither:

| Subsidy | Fi | ully | Brand or |
|------------------------|----------|------|--------------|
| (Manufacturer's Price) | Subsidis | sed | Generic |
| \$ | Per | ✓ | Manufacturer |

continued...

- 1 Atypical mycobacterial infection; or
- 2 Mycobacterium tuberculosis infection where there is drug-resistance or intolerance to standard pharmaceutical agents.

Initial application — (Helicobacter pylori eradication) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 For the eradication of helicobacter pylori in a patient unable to swallow tablets; and
- 2 For use only in combination with omeprazole and amoxicillin as part of a triple therapy regimen.

Initial application — (Prophylaxis of infective endocarditis) from any relevant practitioner. Approvals valid for 3 months where prophylaxis of infective endocarditis associated with surgical or dental procedures if amoxicillin is contra-indicated. Renewal — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

| ERYTHROMYCIN ETHYL SUCCINATE |
|--|
| Tab 400 mg35.82 100 ✓ E-Mycin |
| a) Up to 20 tab available on a PSO |
| b) Up to 2 x the maximum PSO quantity for RFPP |
| Grans for oral liq 200 mg per 5 ml |
| a) Up to 300 ml available on a PSO |
| b) Up to 2 x the maximum PSO quantity for RFPP |
| c) Wastage claimable |
| Grans for oral liq 400 mg per 5 ml |
| a) Up to 200 ml available on a PSO |
| b) Wastage claimable |
| ROXITHROMYCIN |
| Tab 150 mg13.19 50 ✓ <u>Arrow-</u> |
| Roxithromycin |
| Tab 300 mg25.00 50 ✓ Arrow - |
| Roxithromycin |

| | 0.1.11 | | | - |
|--|------------------------------|---------------|-------------|----------------------------|
| | Subsidy (Manufacturer's I | Drine) Cuba | Fully | Brand or |
| | (Manufacturer S) | Price) Subs | idised • | Generic Manufacturer |
| | | | | That is a said to |
| Penicillins | | | | |
| AMOXICILLIN | | | | |
| Cap 250 mg | 27.50 | 500 | 1 | Miro-Amoxicillin |
| a) Up to 30 cap available on a PSO | | | | |
| b) Up to 10 x the maximum PSO quantity for RFPP | | | | |
| Cap 500 mg | 41.00 | 500 | 1 | Miro-Amoxicillin |
| a) Up to 30 cap available on a PSO | | | | |
| b) Up to 10 x the maximum PSO quantity for RFPP | | | | |
| Grans for oral liq 125 mg per 5 ml | 2.22 | 100 ml | 1 | Alphamox 125 |
| a) Up to 200 ml available on a PSO | | | | |
| b) Wastage claimable | | | | |
| Grans for oral liq 250 mg per 5 ml | 2.81 | 100 ml | 1 | Alphamox 250 |
| a) Up to 300 ml available on a PSO | | | | |
| b) Up to 10 x the maximum PSO quantity for RFPP | | | | |
| c) Wastage claimable | | | | |
| Inj 250 mg vial | 15.97 | 10 | 1 | lbiamox |
| Inj 500 mg vial | | 10 | ✓ | lbiamox |
| Inj 1 g vial - Up to 5 inj available on a PSO | 21.64 | 10 | 1 | lbiamox |
| AMOXICILLIN WITH CLAVULANIC ACID | | | | |
| Tab 500 mg with clavulanic acid 125 mg - Up to 30 tab | | | | |
| available on a PSO | 1 59 | 10 | 1 | Curam Duo 500/125 |
| Grans for oral lig amoxicillin 25 mg with clavulanic acid 6.25 | | 10 | - | <u> </u> |
| per ml | 0 | 100 ml | 1 | Augmentin |
| a) Up to 200 ml available on a PSO | | | | |
| b) Wastage claimable | | | | |
| c) Augmentin to be Principal Supply on 1 May 2025 | | | | |
| Grans for oral lig amoxicillin 50 mg with clavulanic acid 12.5 | ma | | | |
| per ml – Up to 200 ml available on a PSO | | 100 ml OP | 1 | Curam |
| por mili op to 200 mil avallable on a 1 00 | 5.61 | 100 1111 01 | | Amoxiclay Devatis |
| | | | | Forte |
| (Curam Grans for oral liq amoxicillin 50 mg with clavulanic acid 1 | 2.5 ma per ml to | be delisted 1 | lune 2 | 025) |
| BENZATHINE BENZYLPENICILLIN | 37 | | | / |
| Inj 900 mg (1.2 million units) in 2.3 ml syringe – Up to 5 inj | | | | |
| available on a PSO | 375.07 | 10 | 1 | Bicillin LA |
| | | 10 | • | DICIIIII LA |
| BENZYLPENICILLIN SODIUM [PENICILLIN G] | 100 | 40 | , | 0 |
| Inj 600 mg (1 million units) vial – Up to 5 inj available on a P | 30 16.50 | 10 | • | <u>Sandoz</u> |
| FLUCLOXACILLIN | | | _ | |
| Cap 250 mg – Up to 30 cap available on a PSO | | 250 | | Flucloxacillin-AFT |
| Cap 500 mg – Up to 30 cap available on a PSO | | 500 | _ | Flucloxacillin-AFT |
| Grans for oral liq 25 mg per ml | 4.89 | 100 ml | • | <u>AFT</u> |
| a) Up to 200 ml available on a PSO | | | | |
| b) Wastage claimable | 5.00 | 4001 | , | A === |
| Grans for oral liq 50 mg per ml | 5.89 | 100 ml | • | <u>AFT</u> |
| a) Up to 200 ml available on a PSO | | | | |
| b) Wastage claimable | 40.60 | 10 | ./ | Eluciovin |
| Inj 250 mg vialInj 500 mg vial | 42.0U 45.60 | 10 10 | | Flucloxin Flucloxin |
| Inj 1 g vial – Up to 5 inj available on a PSO | 6 00 6 00 | 5 | | <u>Flucioxin</u> Flucil |
| 111 1 9 VIGI OP 10 3 111 AVAIIADIE 011 A 1 30 | 0.00 | J | • | i iuvii |

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

| | Subsidy (Manufacturer's Price) | Per | Fully Subsidised | |
|--|-----------------------------------|----------|---------------------|------------------------------|
| PHENOXYMETHYLPENICILLIN (PENICILLIN V) Cap 250 mg — Up to 30 cap available on a PSO Cap 500 mg | | 50 50 | | Cilicaine VK Cilicaine VK |
| Grans for oral liq 125 mg per 5 ml | 3.40 | 100 m | · • | <u>AFT</u> |
| Grans for oral liq 250 mg per 5 ml | 4.24 | 100 m | l / | <u>AFT</u> |

Tetracyclines

| DO | XYCYCLINE | | | |
|-----|---|---------|-----|-----------|
| * | Tab 100 mg - Up to 30 tab available on a PSO | 64.43 | 500 | Doxine |
| MIN | NOCYCLINE HYDROCHLORIDE | | | |
| * | Tab 50 mg - Additional subsidy by Special Authority see | | | |
| | SA1355 below – Retail pharmacy | 5.79 | 60 | |
| | | (12.05) | | Mino-tabs |
| * | Cap 100 mg | 19.32 | 100 | |
| | • | (52.04) | | Minomycin |

⇒SA1355 Special Authority for Manufacturers Price

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has rosacea.

 ${\sf TETRACYCLINE\ - Special\ Authority\ see\ SA1332\ below\ - \ Retail\ pharmacy}$

⇒SA1332 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 For the eradication of helicobacter pylori following unsuccessful treatment with appropriate first-line therapy; and
- 2 For use only in combination with bismuth as part of a quadruple therapy regimen.

Other Antibiotics

For topical antibiotics, refer to DERMATOLOGICALS, page 66

CIPROFLOXACIN

Recommended for patients with any of the following:

- i) microbiologically confirmed and clinically significant pseudomonas infection; or
- ii) prostatitis; or
- iii) pyelonephritis; or
- iv) gonorrhoea.

| Tab 250 mg — Up to 5 tab available on a PSO | 28 28 | ✓ <u>Ipca-Ciprofloxacin</u> ✓ Ipca-Ciprofloxacin |
|---|----------|--|
| Tab 750 mg4.80 | 28 | ✓ Ipca-Ciprofloxacin |
| CLINDAMYCIN | | |
| Cap hydrochloride 150 mg4.94 | 24 | ✓ Dalacin C |
| Inj 150 mg per ml, 4 ml ampoule35.10 | 10 | ✓ HameIn |

| | Subsidy (Manufacturer's Price) | | Fully Subsidised | Generic |
|--|-----------------------------------|-------|---------------------|-------------------------|
| | \$ | Per | · • | Manufacturer |
| COLISTIN SULPHOMETHATE - Retail pharmacy-Specialist - S | Subsidy by endorseme | ent | | |
| Only if prescribed for dialysis or cystic fibrosis patient and th | | | accordingl | ٧. |
| Inj 150 mg | | 1 | | Colistin-Link |
| GENTAMICIN SULPHATE | | | | |
| Inj 40 mg per ml, 2 ml vial – Subsidy by endorsement | 36.70 | 5 | 1 | Cidomycin |
| ing 40 mg per mi, 2 mi viai – Subsidy by endorsement | | J | • | P/Free S29 |
| Only if preservined for a distrain or quatic fibracia nations | ar aamaliaatad urinan | . + | t infaction | |
| Only if prescribed for a dialysis or cystic fibrosis patient endorsed accordingly. | or complicated unitary | rirac | i mection | and the prescription is |
| Inj 10 mg per ml, 1 ml ampoule - Subsidy by endorsement. | 95.00 | 5 | 1 | DBL Gentamicin |
| Only if prescribed for a dialysis or cystic fibrosis patient endorsed accordingly. | | | t infection | and the prescription is |
| Inj 10 mg per ml, 2 ml ampoule - Subsidy by endorsement. | 91.00 | 5 | 1 | Wockhardt S29 |
| Only if prescribed for a dialysis or cystic fibrosis patient endorsed accordingly. | | trac | t infection | and the prescription is |
| Inj 40 mg per ml, 2 ml ampoule – Subsidy by endorsement. | 18.38 | 10 | 1 | Pfizer |
| | 91.90 | 50 | 1 | Gentamicin |
| | | | | Noridem \$29 |
| Only if prescribed for a dialysis or cystic fibrosis patient endorsed accordingly. | or complicated urinary | trac | t infection | and the prescription is |
| MOXIFLOXACIN - Special Authority see SA1740 below - Retail | I pharmacy | | | |
| No patient co-payment payable | | | | |
| Tab 400 mg | 42.00 | 5 | / | Avelox |
| ⇒SA1740 Special Authority for Subsidy | | - | | |
| Initial and lighting (Tubersulation) and from a required man | | | | |

>>

Initial application — (Tuberculosis) only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 Active tuberculosis*: and
 - 1.2 Any of the following:
 - 1.2.1 Documented resistance to one or more first-line medications; or
 - 1.2.2 Suspected resistance to one or more first-line medications (tuberculosis assumed to be contracted in an area with known resistance), as part of regimen containing other second-line agents; or
 - 1.2.3 Impaired visual acuity (considered to preclude ethambutol use); or
 - 1.2.4 Significant pre-existing liver disease or hepatotoxicity from tuberculosis medications; or
 - 1.2.5 Significant documented intolerance and/or side effects following a reasonable trial of first-line medications;
- 2 Mycobacterium avium-intracellulare complex not responding to other therapy or where such therapy is contraindicated.*; or
- 3 Patient is under five years of age and has had close contact with a confirmed multi-drug resistant tuberculosis case.

Note: Indications marked with * are unapproved indications.

Renewal only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Mycoplasma genitalium) only from a sexual health specialist or Practitioner on the recommendation of a sexual health specialist. Approvals valid for 1 month for applications meeting the following criteria: All of the following:

- 1 Has nucleic acid amplification test (NAAT) confirmed Mycoplasma genitalium* and is symptomatic; and
- 2 Either:
 - 2.1 Has tried and failed to clear infection using azithromycin; or
 - 2.2 Has laboratory confirmed azithromycin resistance; and

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

continued...

3 Treatment is only for 7 days.

Initial application — (Penetrating eye injury) only from an ophthalmologist. Approvals valid for 1 month where the patient requires prophylaxis following a penetrating eve injury and treatment is for 5 days only.

Note: Indications marked with * are unapproved indications.

PAROMOMYCIN - Special Authority see SA1689 below - Retail pharmacy

✓ Humatin S29 16

⇒SA1689 Special Authority for Subsidy

Initial application only from an infectious disease specialist, clinical microbiologist or gastroenterologist. Approvals valid for 1 month for applications meeting the following criteria:

Either:

- 1 Patient has confirmed cryptosporidium infection; or
- 2 For the eradication of Entamoeba histolyica carriage.

Renewal only from an infectious disease specialist, clinical microbiologist or gastroenterologist. Approvals valid for 1 month for applications meeting the following criteria:

Fither:

- 1 Patient has confirmed cryptosporidium infection; or
- 2 For the eradication of Entamoeba histolyica carriage.

PYRIMETHAMINE - Special Authority see SA1328 below - Retail pharmacy

30 ✓ Daraprim S29

⇒SA1328 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 For the treatment of toxoplasmosis in patients with HIV for a period of 3 months; or
- 2 For pregnant patients for the term of the pregnancy; or
- 3 For infants with congenital toxoplasmosis until 12 months of age.

SODIUM FUSIDATE [FUSIDIC ACID]

36 ✓ Fucidin Tab 250 mg135.70 SULFADIAZINE SODIUM - Special Authority see SA1331 below - Retail pharmacy 100 ✓ Sulfadiazin-Heyl S29 ✓ Wockhardt S29 543.20 56

⇒SA1331 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 For the treatment of toxoplasmosis in patients with HIV for a period of 3 months; or
- 2 For pregnant patients for the term of the pregnancy; or
- 3 For infants with congenital toxoplasmosis until 12 months of age.

TOBRAMYCIN

Inj 40 mg per ml, 2 ml vial - Subsidy by endorsement.......15.50 ✓ Tobramvcin (Viatris) Only if prescribed for dialysis or cystic fibrosis patient and the prescription is endorsed accordingly. Solution for inhalation 60 mg per ml, 5 ml - Subsidy by ✓ Tobramvcin BNM endorsement......395.00 56 dose

- a) Wastage claimable
- b) Only if prescribed for a cystic fibrosis patient and the prescription is endorsed accordingly.

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer |
|---|---|-------|---------------------|-------------------------------------|
| TRIMETHOPRIM | | | | |
| * Tab 300 mg - Up to 30 tab available on a PSO | 27.83 | 50 | √ <u>T</u> | <u>MP</u> |
| TRIMETHOPRIM WITH SULPHAMETHOXAZOLE [CO-TRIMOX | AZOLE] | | | |
| * Tab trimethoprim 80 mg and sulphamethoxazole 400 mg - l | | | | |
| to 30 tab available on a PSO | 115.74 | 500 | √ <u>T</u> | <u>'risul</u> |
| * Oral liq 8 mg sulphamethoxazole 40 mg per ml - Up to 200 | | | | |
| available on a PSO | 5.00 | 100 m | · |)eprim |
| VANCOMYCIN - Subsidy by endorsement | | | | |
| Only if prescribed for a dialysis or cystic fibrosis patient or for | | | s or for trea | tment of Clostridium |
| difficile following metronidazole failure and the prescription is | | ly. | | |
| Inj 500 mg vial | 3.38 | 1 | ✓ <u>N</u> | <u>lylan</u> |
| | | | | |

Antifungals

- a) For topical antifungals refer to DERMATOLOGICALS, page 67
- b) For topical antifungals refer to GENITO URINARY, page 80

FLUCONAZOLE

| OCONAZOLL | | |
|---|-------|----------------------------|
| Cap 50 mg4.10 | 28 | ✓ Mylan |
| Cap 150 mg0.45 | 1 | ✓ Mylan |
| Cap 200 mg8.90 | 28 | ✓ Mylan |
| Powder for oral suspension 10 mg per ml - Special Authority | | |
| see SA1359 below – Retail pharmacy129.02 | 35 ml | Diflucan |
| Wastage claimable | | |

⇒SA1359 Special Authority for Subsidy

Initial application — (Systemic candidiasis) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

Initial application — (Immunocompromised) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Patient is at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

Renewal — (Systemic candidiasis) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

Renewal — (Immunocompromised) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient remains immunocompromised; and
- 2 Patient remains at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

| | (| | (Manufacturer's Price) Subsidis | | er's Price) Subsidised Generic | | |
|--|--------|-----------|---------------------------------|-------------------------|--------------------------------|--|--|
| | \$ | Per | ✓ | Manufacturer | | | |
| ITRACONAZOLE | | | | | | | |
| Cap 100 mg | 6.83 | 15 | ✓ It | trazole | | | |
| | 27.32 | 60 | ✓ It | tracap \$29 | | | |
| Oral liq 10 mg per ml - Special Authority see SA1322 below | · — | | | | | | |
| Retail pharmacy | 141.80 | 150 ml OP | ✓ It | traconazole Kent S29 | | | |
| | | | √ S | Sporanox | | | |

⇒SA1322 Special Authority for Subsidy

Initial application only from an infectious disease specialist, clinical microbiologist, clinical immunologist or any relevant practitioner on the recommendation of a infectious disease physician, clinical microbiologist or clinical immunologist. Approvals valid for 6 months where the patient has a congenital immune deficiency.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefitting from the treatment.

KETOCONAZOLE

| Tab 200 mg - PCT | CBS | 30 100 | ✓ Burel \$29 ✓ Strides Shasun \$29 ✓ Taro \$29 ✓ Teva- Ketoconazole \$29 |
|---|---------------------|-----------|--|
| NYSTATIN | | | |
| Tab 500,000 u | 14.16 | 50 | |
| | (17.09) | | Nilstat |
| Cap 500,000 u | 12.81 | 50 | |
| | (15.47) | | Nilstat |
| POSACONAZOLE - Special Authority see SA2383 below | v – Retail pharmacy | | |
| Tab modified-release 100 mg | , , | 24 | Posaconazole Juno |
| Oral liq 40 mg per ml | | 105 ml OP | ✓ Devatis |
| 0.0000 | | | |

⇒SA2383 Special Authority for Subsidy

Initial application only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

Either:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation chemotherapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppressive therapy*.

Renewal only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

Either:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation therapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppression* and requires on going posaconazole treatment.

Note: * Graft versus host disease (GVHD) on significant immunosuppression is defined as acute GVHD, grade II to IV, or extensive chronic GVHD, or if they were being treated with intensive immunosuppressive therapy consisting of either high-dose corticosteroids (1 mg or greater per kilogram of body weight per day for patients with acute GVHD or 0.8 mg or greater per kilogram every other day for patients with chronic GVHD), antithymocyte globulin, or a combination of two or more

| Subsidy | Fully | | Brand or |
|------------------------|------------|---|----------|
| (Manufacturer's Price) | Subsidised | | Generic |
| \$ | Per | 1 | |

continued...

immunosuppressive agents or types of treatment.

Initial application — (Invasive fungal infection prophylaxis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient is at risk of invasive fungal infection; and
- 2 Fither:
 - 2.1 Posaconazole is prescribed by, or recommended by a haematologist, transplant physician, infectious disease specialist, paediatric haematologist or paediatric oncologist; or
 - 2.2 Prescribing posaconazole is in accordance with a protocol or guideline that has been endorsed by the Health New Zealand - Te Whatu Ora Hospital in the specific settings where there is a greater than 10% risk of invasive fungal infection (IFI).

Renewal — (Invasive fungal infection prophylaxis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- - 1 The patient is at risk of invasive fungal infection; and
 - 2 Either:
 - 2.1 Posaconazole is prescribed by, or recommended by a haematologist, transplant physician, infectious disease specialist, paediatric haematologist or paediatric oncologist; or
 - 2.2 Prescribing posaconazole is in accordance with a protocol or guideline that has been endorsed by the Health New Zealand - Te Whatu Ora Hospital in the specific settings where there is a greater than 10% risk of invasive fungal infection (IFI).

TERBINAFINE

| * Tab 250 mg8.97 | 84 | ✓ Deolate |
|---|-------|-----------|
| VORICONAZOLE - Special Authority see SA2384 below - Retail pharmacy | | |
| Tab 50 mg91.00 | 56 | ✓ Vttack |
| Tab 200 mg350.00 | 56 | ✓ Vttack |
| Powder for oral suspension 40 mg per ml - Wastage | | |
| claimable | 70 ml | ✓ Vfend |

⇒SA2384 Special Authority for Subsidy

Initial application — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
 - 3.1 Patient has proven or probable invasive aspergillus infection; or
 - 3.2 Patient has possible invasive aspergillus infection; or
 - 3.3 Patient has fluconazole resistant candidiasis: or
 - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Renewal — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
 - 3.1 Patient continues to require treatment for proven or probable invasive aspergillus infection; or

| ubsidy cturer's Price) Subs | Fully | Brand or Generic |
|--------------------------------|-------|---------------------|
| \$ Per | • | Manufacturer |

continued...

- 3.2 Patient continues to require treatment for possible invasive aspergillus infection; or
- 3.3 Patient has fluconazole resistant candidiasis; or
- 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Initial application — (Invasive fungal infection prophylaxis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient is at risk of invasive fungal infection; and
- 2 Either:
 - 2.1 Voriconazole is prescribed by, or recommended by a haematologist, transplant physician, infectious disease specialist, paediatric haematologist or paediatric oncologist; or
 - 2.2 Prescribing voriconazole is in accordance with a protocol or guideline that has been endorsed by the Health New Zealand - Te Whatu Ora Hospital in the specific settings where there is a greater than 10% risk of invasive fungal infection (IFI).

Renewal — (Invasive fungal infection prophylaxis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient is at risk of invasive fungal infection; and
- 2 Either:
 - 2.1 Voriconazole is prescribed by, or recommended by a haematologist, transplant physician, infectious disease specialist, paediatric haematologist or paediatric oncologist; or
 - 2.2 Prescribing voriconazole is in accordance with a protocol or guideline that has been endorsed by the Health New Zealand - Te Whatu Ora Hospital in the specific settings where there is a greater than 10% risk of invasive fungal infection (IFI).

Antimalarials

⇒SA1684 Special Authority for Subsidy

Initial application only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 The patient has vivax or ovale malaria; and
- 2 Primaguine is to be given for a maximum of 21 days.

Renewal only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 The patient has relapsed vivax or ovale malaria; and
- 2 Primaguine is to be given for a maximum of 21 days.

| | Subsidy | | Fully | Brand or |
|---|---------------------|----------------------|---------------------------------------|-------------------------------|
| | (Manufacturer's F | Price) | Subsidised | Generic |
| | \$ | Per | • | Manufacturer |
| Antitrichomonal Agents | | | | |
| METRONIDAZOLE | | | | |
| Tab 200 mg - Up to 30 tab available on a PSO | 25.86 | 250 | 1 | Metronidamed |
| | 33.15 | | | Metrogyl |
| Metronidamed to be Principal Supply on 1 March 2025 | | | | |
| Tab 400 mg - Up to 15 tab available on a PSO | 4.29 | 21 | 1 | Metronidamed |
| | 5.23 | | ✓ | Metrogyl |
| Metronidamed to be Principal Supply on 1 March 2025 | | | | 5 , |
| Oral lig benzoate 200 mg per 5 ml | 25.00 | 100 ml | 1 | Flagyl-S |
| Suppos 500 mg | 24.48 | 10 | | Flagyl |
| (Metrogyl Tab 200 mg to be delisted 1 March 2025) | | | | • |
| (Metrogyl Tab 400 mg to be delisted 1 March 2025) | | | | |
| ORNIDAZOLE | | | | |
| Tab 500 mg | 36.52 | 10 | 1 | Arrow-Ornidazole |
| Arrow-Ornidazole to be Principal Supply on 1 March 202 | | 10 | • | ATTOW-OTTIIUazoie |
| Altow-Offiidazole to be i filicipal Supply off i March 202 | 3 | | | |
| Antituberculotics and Antileprotics | | | | |
| Antituber culotics and Antileprotics | | | | |
| Note: There is no co-payment charge for all pharmaceuticals liste | ed in the Antitub | erculotics a | nd Antile | protics group regardless |
| immigration status. | | | | 3 3 |
| BEDAQUILINE - Special Authority see SA2244 below - Retail p | harmacy | | | |
| No patient co-payment payable | паппасу | | | |
| Tab 100mg | 3 084 51 | 24 OP | 1 | Sirturo |
| | 0,004.01 | 24 01 | - | Ontaro |
| ⇒SA2244 Special Authority for Subsidy | | . 1212 | · · · · · · · · · · · · · · · · · · · | |
| Initial application — (multi-drug resistant tuberculosis) from | any relevant pra | ictitioner. <i>F</i> | Approvais | valid for 6 months for |
| applications meeting the following criteria: | | | | |
| Both: | | | | |
| 1 The person has multi-drug resistant tuberculosis (MDR-TE | , : | | | |
| 2 Ministry of Health's Tuberculosis Clinical Network has revi | ewed the individ | lual case ai | nd recomi | mends bedaquiline as pai |
| of the treatment regimen. | | | | |
| CLOFAZIMINE - Retail pharmacy-Specialist | | | | |
| a) No patient co-payment payable | | | | |
| b) Prescriptions must be written by, or on the recommendation | on of, an infection | ous disease | physiciar | n, clinical microbiologist o |
| dermatologist. | | | . , | |
| * Cap 50 mg | 442.00 | 100 | 1 | Lamprene S29 |
| CYCLOSERINE - Retail pharmacy-Specialist | | | | • |
| a) No patient co-payment payable | | | | |
| b) Prescriptions must be written by, or on the recommendati | on of an infectio | nie diepaed | nhveicia | n clinical microhiologist o |
| respiratory physician. | on or, an imedic | ius uiscasc | priyololal | i, ciiriicai microbiologist o |
| Cap 250 mg | 344.00 | 60 | 1 | Cyclorin \$29 |
| | | 00 | • | Oycioiii |
| DAPSONE - Retail pharmacy-Specialist | | | | |
| a) No patient co-payment payable | | | | |
| b) Prescriptions must be written by, or on the recommendati | on of, an infection | ous disease | physiciar | n, clinical microbiologist o |
| dermatologist | | | _ | _ |
| Tab 25 mg | | 100 | | Dapsone |
| Tab 100 mg | 329.50 | 100 | ✓ | Dapsone |
| | | | | |
| | | | | |

| | Subsidy | | Fully | Brand or |
|---|--|--------------------------|----------------------------|---|
| | (Manufacturer's Price) \$ | Per | Subsidised < | Generic Manufacturer |
| THAMBUTOL HYDROCHLORIDE - Retail pharmacy-Specia | list | | | |
| a) No patient co-payment payable | | | | |
| b) Prescriptions must be written by, or on the recommend | ation of, an infectious o | lisease | e physiciar | n, clinical microbiologist o |
| respiratory physician Tab 100 mg | 0F 70 | 100 | ./ | EMB Fatol \$29 |
| Tab 400 mg | | 100 56 | _ | Myambutol \$29 |
| - | 49.34 | 50 | • | wyanibulor 323 |
| SONIAZID – Retail pharmacy-Specialist a) No patient co-payment payable | | | | |
| b) Prescriptions must be written by, or on the recommendation microbiologist, dermatologist or public health physician | ation of, an internal me | dicine | physician | paediatrician, clinical |
| * Tab 100 mg | 23.00 | 100 | 1 | PSM |
| Tub 100 mg | 94.50 | 100 | | Isoniazid Teva S29 |
| | 327.41 | | | Noumed Isoniazid |
| Noumed Isoniazid to be Principal Supply on 1 May 202 (PSM Tab 100 mg to be delisted 1 May 2025) | 25 | | | |
| SONIAZID WITH RIFAMPICIN - Retail pharmacy-Specialist | | | | |
| a) No patient co-payment payable | | | | |
| Prescriptions must be written by, or on the recommendamicrobiologist, dermatologist or public health physician | ation of, an internal me | dicine | physician | paediatrician, clinical |
| * Tab 100 mg with rifampicin 150 mg | | 100 | | Rifinah |
| * Tab 150 mg with rifampicin 300 mg | | 100 | • | <u>Rifinah</u> |
| .INEZOLID – Special Authority see SA2234 below – Retail ph No patient co-payment payable | | | | |
| Tab 600 mg | | 10 | | <u>Zyvox</u> |
| Oral liq 20 mg per ml | 1,879.00 | 150 m | ı • | Zyvox |
| ⇒SA2234 Special Authority for Subsidy nitial application — (multi-drug resistant tuberculosis) fro | m any relevant practition | oner. | Approvals | valid for 18 months for |
| applications meeting the following criteria: | | | | |
| applications meeting the following criteria: Both: | TR): and | | | |
| applications meeting the following criteria: Both: 1 The person has multi-drug resistant tuberculosis (MDR- 2 Ministry of Health's Tuberculosis Clinical Network has re | | case a | ınd recomr | nends linezolid as part of |
| applications meeting the following criteria: Both: 1 The person has multi-drug resistant tuberculosis (MDR-2 Ministry of Health's Tuberculosis Clinical Network has rethe treatment regimen. | | case a | and recomr | nends linezolid as part of |
| applications meeting the following criteria: Both: 1 The person has multi-drug resistant tuberculosis (MDR-2 Ministry of Health's Tuberculosis Clinical Network has rethe treatment regimen. | eviewed the individual | | | · |
| applications meeting the following criteria: Both: 1 The person has multi-drug resistant tuberculosis (MDR-2 Ministry of Health's Tuberculosis Clinical Network has rethe treatment regimen. PARA-AMINO SALICYLIC ACID — Retail pharmacy-Specialist a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendarespiratory physician | eviewed the individual | lisease | e specialis | t, clinical microbiologist o |
| applications meeting the following criteria: Both: 1 The person has multi-drug resistant tuberculosis (MDR-2 Ministry of Health's Tuberculosis Clinical Network has rethe treatment regimen. PARA-AMINO SALICYLIC ACID — Retail pharmacy-Specialist a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendations. | eviewed the individual | | e specialis | · |
| pplications meeting the following criteria: Both: 1 The person has multi-drug resistant tuberculosis (MDR-2 Ministry of Health's Tuberculosis Clinical Network has rethe treatment regimen. PARA-AMINO SALICYLIC ACID — Retail pharmacy-Specialist a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendarespiratory physician Grans for oral liq 4 g sachet | eviewed the individual | lisease | e specialis | t, clinical microbiologist c |
| applications meeting the following criteria: Both: 1 The person has multi-drug resistant tuberculosis (MDR-2 Ministry of Health's Tuberculosis Clinical Network has rethe treatment regimen. PARA-AMINO SALICYLIC ACID — Retail pharmacy-Specialist a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendarespiratory physician Grans for oral liq 4 g sachet | eviewed the individual of the entire of the | lisease 30 | e specialis | t, clinical microbiologist o |
| applications meeting the following criteria: Both: 1 The person has multi-drug resistant tuberculosis (MDR-2 Ministry of Health's Tuberculosis Clinical Network has rethe treatment regimen. PARA-AMINO SALICYLIC ACID — Retail pharmacy-Specialist a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendarespiratory physician Grans for oral liq 4 g sachet | ation of, an infectious of the control of the contr | lisease 30 lisease | e specialis • e specialis | t, clinical microbiologist o Paser \$29 t, clinical microbiologist o |
| applications meeting the following criteria: Both: 1 The person has multi-drug resistant tuberculosis (MDR-2 Ministry of Health's Tuberculosis Clinical Network has rethe treatment regimen. PARA-AMINO SALICYLIC ACID — Retail pharmacy-Specialist a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendarespiratory physician Grans for oral liq 4 g sachet | ation of, an infectious of the control of the contr | lisease 30 | e specialis • e specialis | t, clinical microbiologist o |
| applications meeting the following criteria: Both: 1 The person has multi-drug resistant tuberculosis (MDR-2 Ministry of Health's Tuberculosis Clinical Network has rethe treatment regimen. PARA-AMINO SALICYLIC ACID — Retail pharmacy-Specialist a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendarespiratory physician Grans for oral liq 4 g sachet | ation of, an infectious of the control of the contr | lisease 30 lisease | e specialis • e specialis | t, clinical microbiologist o Paser \$29 t, clinical microbiologist o |
| applications meeting the following criteria: Both: 1 The person has multi-drug resistant tuberculosis (MDR-2 Ministry of Health's Tuberculosis Clinical Network has rethe treatment regimen. PARA-AMINO SALICYLIC ACID — Retail pharmacy-Specialist a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendarespiratory physician Grans for oral liq 4 g sachet | ation of, an infectious of ation of ati | 30 lisease 100 | e specialis | t, clinical microbiologist o Paser \$29 t, clinical microbiologist o Peteha \$29 |

| Subsidy | Fully | | Brand or |
|------------------------|------------|---|----------|
| (Manufacturer's Price) | Subsidised | | Generic |
| \$ | Per | 1 | |

RIFABUTIN - Retail pharmacy-Specialist

- a) No patient co-payment payable
- b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, respiratory physician or gastroenterologist
- ✓ Mycobutin

RIFAMPICIN - Subsidy by endorsement

- a) No patient co-payment payable
- b) For confirmed recurrent Staphylococcus aureus infection in combination with other effective anti-staphylococcal antimicrobial based on susceptibilities and the prescription is endorsed accordingly; can be waived by endorsement -Retail pharmacy - Specialist. Specialist must be an internal medicine physician, clinical microbiologist, dermatologist, paediatrician, or public health physician

| Cap 150 mg | 100 100 | ✓ <u>Rifadin</u> ✓ Rifadin |
|------------|------------|-------------------------------|
| Cap 300 mg | 100 | ✓ Rifadin Sanofi ✓ Rifadin |

Antivirals

For eye preparations refer to Eye Preparations, Anti-Infective Preparations, page 265

Oral liq 5 mg per ml270.00

Hepatitis B Treatment

| • | |
|-----------|--|
| ENTECAVIR | |

| * Tab 0.5 mg | .04 30 | ✓ Entecavir (Rex) |
|---|--------------|-------------------|
| LAMIVUDINE - Special Authority see SA1685 below - Retail pharmacy | | |
| Tab 100 mg12 | .06 28 | ✓ Zetlam |
| Oral lig 5 mg per ml270 | .00 240 ml (| OP ✓ Zeffix |

⇒SA1685 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 1 year where used for the treatment or prevention of hepatitis B.

Renewal from any relevant practitioner. Approvals valid for 2 years where used for the treatment or prevention of hepatitis B. TENOFOVIR DISOPROXIL

Tenofovir disoproxil prescribed under endorsement for the treatment of HIV is included in the count of up to 4 subsidised antiretrovirals for the nurnoses of Special Authority SA2139 page 110

| | antifetrovitals for the purposes of opecial Authority GAZT | oo., page 110 | | | |
|---|--|---------------|----|-------------------|---------------------|
| * | Tab 245 mg (300 mg as a maleate) | 15.00 | 30 | ✓ <u>I</u> | enofovir Disoproxil |
| | | | | | Viatris |

Herpesvirus Treatments

| ACICLOVIR | | | |
|--|-----------------------|--------|------------------------------------|
| * Tab dispersible 200 mg | 1.78 | 25 | ✓ Lovir |
| * Tab dispersible 400 mg | 5.81 | 56 | ✓ Lovir |
| * Tab dispersible 800 mg | 6.46 | 35 | ✓ Lovir |
| VALACICLOVIR | | | |
| Tab 500 mg | 9.64 | 30 | ✓ Vaclovir |
| Tab 1,000 mg | 17.78 | 30 | ✓ Vaclovir |
| VALGANCICLOVIR - Special Authority see SA1993 on the n | ext page – Retail pha | ırmacy | |
| Tab 450 mg | 140.89 | 60 | ✓ <u>Valganciclovir</u> Viatris |

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA1993 Special Authority for Subsidy

Initial application — **(transplant cytomegalovirus prophylaxis)** only from a relevant specialist. Approvals valid for 3 months where the patient has undergone a solid organ transplant and requires valganciclovir for CMV prophylaxis.

Renewal — (transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has undergone a solid organ transplant and received anti-thymocyte globulin and requires valganciclovir therapy for CMV prophylaxis: and
 - 1.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following anti-thymocyte globulin; or
- 2 Both:
 - 2.1 Patient has received pulse methylprednisolone for acute rejection and requires further valganciclovir therapy for CMV prophylaxis; and
 - 2.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following pulse methylprednisolone.

Initial application — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has undergone a solid organ transplant and received valganciclovir under Special Authority more than 2 years ago (27 months); and
- 2 Patient has received anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Renewal — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months where the patient has received a further course of anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Initial application — (Lung transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone a lung transplant; and
- 2 Either:
 - 2.1 The donor was cytomegalovirus positive and the patient is cytomegalovirus negative; or
 - 2.2 The recipient is cytomegalovirus positive; and
- 3 Patient has a high risk of CMV disease.

Initial application — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
 - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
 - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
 - 2.3 Patient has cytomegalovirus retinitis.

Renewal — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
 - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
 - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
 - 2.3 Patient has cytomegalovirus retinitis.

Note: for the purpose of this Special Authority "immunocompromised" includes transplant recipients, patients with immunosuppressive diseases (e.g. HIV) or those receiving immunosuppressive treatment for other conditions.

Subsidy (Manufacturer's Price) \$

Subsidised Per 🗸

Fully

Brand or Generic Manufacturer

Hepatitis C Treatment

GLECAPREVIR WITH PIBRENTASVIR - [Xpharm]

Note the supply of treatment is via Pharmac's approved direct distribution supply. Further details can be found on Pharmac's website https://pharmac.govt.nz/maviret

Tab 100 mg with pibrentasvir 40 mg24,750.00 84 OP ✓ Maviret

LEDIPASVIR WITH SOFOSBUVIR – [Xpharm] – Special Authority see SA1605 below

No patient co-payment payable

Tab 90 mg with sofosbuvir 400 mg......24,363.46 28 **✓ Harvoni**

⇒SA1605 Special Authority for Subsidy

Special Authority approved by the Hepatitis C Treatment Panel (HepCTP)

Notes: By application to the Hepatitis C Treatment Panel (HepCTP).

Applications will be considered by HepCTP and approved subject to confirmation of eligibility.

Application details may be obtained from Pharmac's website http://www.pharmac.govt.nz/mayiret or:

The Coordinator, Hepatitis C Treatment Panel

Pharmac, PO Box 10-254, WELLINGTON Tel: (04) 460 4990.

Email: hepcpanel@pharmac.govt.nz

HIV Prophylaxis and Treatment

EMTRICITABINE WITH TENOFOVIR DISOPROXIL - Subsidy by endorsement; can be waived by Special Authority see SA2138 below

- a) Funding for emtricitabine with tenofovir disoproxil for use as PrEP, should be applied using Special Authority SA2138.
- b) Endorsement for treatment of conditions approved via Special Authority SA2139 (antiretrovirals for confirmed HIV, prevention of maternal transmission, post-exposure prophylaxis following exposure to HIV and percutaneous exposure): Prescription is deemed to be endorsed if emtricitabine with tenofovir disoproxil is co-prescribed with another antiretroviral subsidised under Special Authority SA2139 and the prescription is annotated accordingly by the Pharmacist or endorsed by the prescriber.

Note: Emtricitabine with tenofovir disoproxil prescribed under endorsement, for treatment of conditions approved via Special Authority SA2139 (antiretrovirals for confirmed HIV, prevention of maternal transmission, post-exposure prophylaxis following exposure to HIV and percutaneous exposure), is included in the count of up to 4 subsidised antiretrovirals, and counts as two antiretroviral medications, for the purposes of Special Authority SA2139, page 110 There is an approval process to become a named specialist to prescribe antiretroviral therapy in New Zealand. Further information is available on the Pharmac website.

* Tab 200 mg with tenofovir disoproxil 245 mg (300 mg as a

⇒SA2138 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria:

Both:

1 Patient has tested HIV negative, does not have signs or symptoms of acute HIV infection and has been assessed for HIV seroconversion; and

30

✓ Teva

2 The Practitioner considers the patient is at elevated risk of HIV exposure and use of PrEP is clinically appropriate.

Notes: Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines:

https://ashm.org.au/HIV/PrEP/

| Subsidy | | Fully | Brand or |
|------------------------|-----|------------|--------------|
| (Manufacturer's Price) | | Subsidised | Generic |
| \$ | Per | | Manufacturer |

continued...

Renewal from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 Patient has tested HIV negative, does not have signs or symptoms of acute HIV infection and has been assessed for HIV seroconversion; and
- 2 The Practitioner considers the patient is at elevated risk of HIV exposure and use of PrEP is clinically appropriate.

Notes: Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines:

https://ashm.org.au/HIV/PrEP/

COVID-19 Treatments

MOLNUPIRAVIR - [Xpharm] - Subsidy by endorsement

- a) No patient co-payment payable
- b) Treatment is funded only if patient meets access criteria for oral antiviral COVID-19 treatments (as on <u>Pharmac's website</u>) and has been endorsed accordingly by the prescriber. The supply of treatment is via Pharmac's approved distribution process. Refer to the Pharmac website for more information about this and stock availability.

NIRMATRELVIR WITH RITONAVIR - [Xpharm] - Subsidy by endorsement

- a) No patient co-payment payable
- b) Treatment is funded only if patient meets access criteria for oral antiviral COVID-19 treatments (as on <u>Pharmac's website</u>) and has been endorsed accordingly by the prescriber. The supply of treatment is via Pharmac's approved distribution process. Refer to the Pharmac website for more information about this and stock availability.

Antiretrovirals

⇒SA2139 Special Authority for Subsidy

Initial application — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the patient has confirmed HIV infection.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Prevention of maternal transmission) only from a named specialist. Approvals valid for 1 year for applications meeting the following criteria:

Fither:

- 1 Prevention of maternal foetal transmission: or
- 2 Treatment of the newborn for up to eight weeks.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Some antiretrovirals are unapproved or contraindicated for this indication. Practitioners prescribing these medications should exercise their own skill, judgement, expertise and discretion, and make their own prescribing decisions with respect to the use of a Pharmaceutical for an indication for which it is not approved or contraindicated.

| Subsidy | | Fully | Brand or |
|-----------------------|-----|------------|--------------|
| (Manufacturer's Price |) | Subsidised | Generic |
| \$ | Per | ✓ | Manufacturer |

continued...

Initial application — (post-exposure prophylaxis following exposure to HIV) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

- Both:
 - 1 Treatment course to be initiated within 72 hours post exposure; and
 - 2 Any of the following:
 - 2.1 Patient has had condomless anal intercourse or receptive vaginal intercourse with a known HIV positive person with an unknown or detectable viral load greater than 200 copies per ml; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required; or
 - 2.4 Patient has had condomless anal intercourse with a person from a high HIV prevalence country or risk group whose HIV status is unknown.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines for PEP (https://www.ashm.org.au/hiv/hiv-management/pep/).

Renewal — (second or subsequent post-exposure prophylaxis) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had condomless anal intercourse or receptive vaginal intercourse with a known HIV positive person with an unknown or detectable viral load greater than 200 copies per ml; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required; or
 - 2.4 Patient has had condomless anal intercourse with a person from a high HIV prevalence country or risk group whose HIV status is unknown.

Initial application — (**Percutaneous exposure**) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Second or subsequent percutaneous exposure) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Non-nucleosides Reverse Transcriptase Inhibitors

| EFAVIRENZ - Special Authority see SA2139 on the previous pag | je – Retail pha | rmacy | |
|--|-----------------|-----------|--|
| Tab 600 mg | 65.38 | 30 | ✓ Efavirenz |
| | | | Milpharm \$29 |
| ETRAVIRINE - Special Authority see SA2139 on the previous pa | ge – Retail ph | armacy | |
| Tab 200 mg | 770.00 | 60 | ✓ Intelence |
| NEVIRAPINE - Special Authority see SA2139 on the previous pa | ge – Retail ph | armacy | |
| Tab 200 mg | 198.25 | 60 | Nevirapine Viatris |
| Oral suspension 10 mg per ml | 203.55 | 240 ml OP | ✓ Viramune |
| | | | Suspension |

| | Subsidy | | Fully | Brand or |
|---|------------------------|----------------|-------------------|---|
| | (Manufacturer's Price) | Suk | sidised | Generic |
| | \$ | Per | ✓ | Manufacturer |
| Nucleosides Reverse Transcriptase Inhibitors | | | | |
| ABACAVIR SULPHATE - Special Authority see SA2139 on pag | e 110 – Retail pharma | асу | | |
| Tab 300 mg | | 60 | √ Z | Ziagen |
| ABACAVIR SULPHATE WITH LAMIVUDINE - Special Authority | see SA2139 on page | 110 – F | Retail ph | armacy |
| Note: abacavir with lamivudine (combination tablets) counts anti-retroviral Special Authority. | | medicati | ons for t | he purposes of the |
| Tab 600 mg with lamivudine 300 mg | 29.50 | 30 | √ <u>I</u> | <u>Abacavir/</u> <u>Lamivudine</u> <u>Viatris</u> |
| EFAVIRENZ WITH EMTRICITABINE AND TENOFOVIR DISOPI | ROXIL - Special Auth | nority see | SA213 | 9 on page 110 – Retail |
| pharmacy | | | | |
| Note: Efavirenz with emtricitabine and tenofovir disoproxil or anti-retroviral Special Authority | | rovirai m | iedicatio | ns for the purposes of the |
| Tab 600 mg with emtricitabine 200 mg and tenofovir disopro: 245 mg (300 mg as a fumarate) | | 30 | / 1 | riovir \$29 |
| Tab 600 mg with emtricitabine 200 mg and tenofovir disopro | | 30 | • ' | IIIOVII 623 |
| 245 mg (300 mg as a maleate) | | 30 | ✓ \ | /iatris |
| EMTRICITABINE - Special Authority see SA2139 on page 110 - | | | | |
| Cap 200 mg | | 30 | ✓ E | Emtriva |
| LAMIVUDINE - Special Authority see SA2139 on page 110 - Re | etail pharmacy | | | |
| Tab 150 mg | | 60 | _ | amivudine Viatris |
| Oral liq 10 mg per ml | | 0 ml OP | √ 3 | TC |
| ZIDOVUDINE [AZT] - Special Authority see SA2139 on page 11 | | | | |
| Cap 100 mg | | 100 0 ml OP | | Retrovir Retrovir |
| Oral liq 10 mg per ml | | - | | |
| ZIDOVUDINE [AZT] WITH LAMIVUDINE – Special Authority see Note: zidovudine [AZT] with lamivudine (combination tablets the anti-retroviral Special Authority. | | | | |
| Tab 300 mg with lamivudine 150 mg | 92.40 | 60 | ✓ L | amivudine/ |
| | | | | Zidovudine Viatris |
| Protease Inhibitors | | | | |
| ATAZANAVIR SULPHATE - Special Authority see SA2139 on p | | rmacy | | |
| Cap 150 mg | 85.00 | 60 | | Atazanavir Mylan |
| Cap 200 mg | 110.00 | 60 | √ <u>I</u> | Atazanavir Viatris |
| DARUNAVIR - Special Authority see SA2139 on page 110 - Re | ' ' | | | |
| Tab 400 mg | | 60 | _ | Darunavir Viatris |
| Tab 600 mg | | 60 | _ | Darunavir Viatris |
| LOPINAVIR WITH RITONAVIR – Special Authority see SA2139 Tab 100 mg with ritonavir 25 mg | | pharmad 60 | • | .opinavir/Ritonavir |
| Tab 200 mg with ritonavir 50 mg | 875 00 | 120 | √ 1 | Mylan .opinavir/Ritonavir |
| | | 120 | • <u>1</u> | <u>Mylan</u> |
| RITONAVIR - Special Authority see SA2139 on page 110 - Ret | | 00 | | La caradia |
| Tab 100 mg | 43.31 | 30 | • | lorvir |

| Subsidy | | Fully | Brand or | |
|------------------------|-----|------------|--------------|--|
| (Manufacturer's Price) | ; | Subsidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

Strand Transfer Inhibitors

| DOLUTEGRAVIR - Special Authority see SA2139 on page 1 | 10 – Retail pharmacy | | |
|---|--------------------------|--------------|----------------|
| Tab 50 mg | 1,090.00 | 30 | ✓ Tivicay |
| DOLUTEGRAVIR WITH LAMIVUDINE - Special Authority se | e SA2139 on page 110 | - Retail pha | rmacy |
| Tab 50 mg with lamivudine 300 mg | | | ✓ Dovato |
| RALTEGRAVIR POTASSIUM - Special Authority see SA213 | 9 on page 110 – Retail p | harmacy | |
| Tab 400 mg | 1,090.00 | 60 | ✓ Isentress |
| Tah 600 mg | 1 090 00 | 60 | ✓ Isentress HD |

Immune Modulators

PEGYLATED INTERFERON ALFA-2A - Special Authority see SA2034 below - Retail pharmacy

⇒SA2034 Special Authority for Subsidy

Initial application — (chronic hepatitis C - genotype 1, 4, 5 or 6 infection or co-infection with HIV or genotype 2 or 3 post liver transplant) from any specialist. Approvals valid for 18 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 Patient has chronic hepatitis C, genotype 1, 4, 5 or 6 infection; or
 - 1.2 Patient has chronic hepatitis C and is co-infected with HIV; or
 - 1.3 Patient has chronic hepatitis C genotype 2 or 3 and has received a liver transplant; and
- 2 Maximum of 48 weeks therapy.

Renewal — (Chronic hepatitis C - genotype 1 infection) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Fither:
 - 3.1 Patient has responder relapsed; or
 - 3.2 Patient was a partial responder; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application — (Chronic Hepatitis C - genotype 1 infection treatment more than 4 years prior) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Any of the following:
 - 3.1 Patient has responder relapsed; or
 - 3.2 Patient was a partial responder; or
 - 3.3 Patient received interferon treatment prior to 2004; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

| Subsidy (Manufacturer's Pr | ice) | Fully Subsidised | Brand or Generic | |
|-------------------------------|------|---------------------|---------------------|--|
| \$ | Per | • | Manufacturer | |

continued...

Initial application — (chronic hepatitis C - genotype 2 or 3 infection without co-infection with HIV) from any specialist.

Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has chronic hepatitis C, genotype 2 or 3 infection; and
- 2 Maximum of 6 months therapy.

Initial application — (Hepatitis B) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Hepatitis B infection (HBsAq positive for more than 6 months); and
- 2 Patient is Hepatitis B treatment-naive; and
- 3 ALT > 2 times Upper Limit of Normal: and
- 4 HBV DNA < 10 log10 IU/ml; and
- 5 Either:
 - 5.1 HBeAg positive: or
 - 5.2 serum HBV DNA greater than or equal to 2,000 units/ml and significant fibrosis (Metavir Stage F2 or greater or moderate fibrosis); and
- 6 Compensated liver disease: and
- 7 No continuing alcohol abuse or intravenous drug use; and
- 8 Not co-infected with HCV, HIV or HDV; and
- 9 Neither ALT nor AST > 10 times upper limit of normal; and
- 10 No history of hypersensitivity or contraindications to pegylated interferon; and
- 11 Maximum of 48 weeks therapy.

Initial application — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 Patient has a cutaneous T cell lymphoma*; or
- 2 All of the following:
 - 2.1 Patient has a myeloproliferative disorder*; and
 - 2.2 Patient is intolerant of hydroxyurea; and
 - 2.3 Treatment with an agrelide and busulfan is not clinically appropriate; or
- 3 Roth:
 - 3.1 Patient has a myeloproliferative disorder; and
 - 3.2 Patient is pregnant, planning pregnancy or lactating.

Renewal — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment; and
- 3 Either:
 - 3.1 Patient has a cutaneous T cell lymphoma*; or
 - 3.2 Both:
 - 3.2.1 Patient has a myeloproliferative disorder*; and
 - 3.2.2 Either:
 - 3.2.2.1 Remains intolerant of hydroxyurea and treatment with anagrelide and busulfan remains clinically inappropriate; or
 - 3.2.2.2 Patient is pregnant, planning pregnancy or lactating.

Note: Indications marked with * are unapproved indications.

| Subsidy | | Fully | Brand or | |
|------------------------|-----|------------|--------------|--|
| (Manufacturer's Price) | 9 | Subsidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

continued

Initial application — **(post-allogenic bone marrow transplant)** from any relevant practitioner. Approvals valid for 3 months where patient has received an allogeneic bone marrow transplant* and has evidence of disease relapse.

Renewal — (post-allogenic bone marrow transplant) from any relevant practitioner. Approvals valid for 3 months where patient is responding and ongoing treatment remains appropriate.

Note: Indications marked with * are unapproved indications.

Urinary Tract Infections

⇒SA2406 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 months for applications meeting the following criteria: Both:

- 1 Patient has an acute, symptomatic, bacteriologically-proven uncomplicated urinary tract infection (UTI)/cystitis with Escherichia Coli: and
- 2 Fither:
 - 2.1 Microbiological testing confirms the pathogen is resistant to all of: trimethoprim, nitrofurantoin, amoxicillin, cefaclor, cefalexin, amoxicillin with clavulanic acid, and norfloxacin; or
 - 2.2 The patient has a contraindication or intolerance to all of: trimethoprim, nitrofurantoin, amoxicillin, cefaclor, cefalexin, amoxicillin with clavulanic acid, and norfloxacin that the pathogen is susceptible to.

Renewal from any relevant practitioner. Approvals valid for 2 months for applications meeting the following criteria: Both:

- 1 Patient has an acute, symptomatic, bacteriologically-proven uncomplicated urinary tract infection (UTI)/cystitis with Escherichia Coli; and
- 2 Either:
 - 2.1 Microbiological testing confirms the pathogen is resistant to all of: trimethoprim, nitrofurantoin, amoxicillin, cefaclor, cefalexin, amoxicillin with clavulanic acid, and norfloxacin; or
 - 2.2 The patient has a contraindication or intolerance to all of: trimethoprim, nitrofurantoin, amoxicillin, cefaclor, cefalexin, amoxicillin with clavulanic acid, and norfloxacin that the pathogen is susceptible to.

METHENAMINE (HEXAMINE) HIPPURATE

| * Tab 1 g | 19.95 | 100 | ✓ Hiprex |
|---|--------|-----|---------------------|
| NITROFURANTOIN | | | |
| * Tab 50 mg - Up to 30 tab available on a PSO | 22.20 | 100 | ✓ Nifuran |
| * Tab 100 mg | 37.50 | 100 | ✓ Nifuran |
| * Cap modified-release 100 mg - Up to 15 cap available on a | | | |
| PSO | 81.20 | 100 | ✓ Macrobid |
| NORFLOXACIN | | | |
| Tab 400 mg - Subsidy by endorsement | 245.00 | 100 | Arrow-Norfloxacin |
| | | | ✓ Arrow-Norfloxacin |

Only if prescribed for a patient with an uncomplicated urinary tract infection that is unresponsive to a first line agent or with proven resistance to first line agents and the prescription is endorsed accordingly.

| ly 's Price) Sul | Fully Brand or bsidised Generic |
|---------------------|------------------------------------|
| Per | ✓ Manufacturer |
| | |
| | |
| | |
| 10 | ✓ Max Health |
| | |
| 100 | ✓ Mestinon |
| | |
| | |
| 50 | ✓ Diclofenac Sandoz |
| 20 | ✓ Voltaren D |
| 50 | ✓ Diclofenac Sandoz |
| 100 | ✓ Voltaren SR |
| 5 | ✓ Voltaren |
| 10 | ✓ Voltaren |
| | |
| 1,000 | ✓ Relieve |
| 30 | ✓ Brufen SR |
| | ✓ Ibuprofen SR BNM |
| | |
| 200 ml | ✓ Ethics |
| | |
| | |
| | |
| 28 | Oruvail SR |
| | |
| 50 | |
| 30 | Ponstan |
| 20 | ronotari |
| 20 | Ponstan |
| | |
| 500 | ✓ Noflam 250 |
| 250 | ✓ Noflam 500 |
| 28 | ✓ Naprosyn SR 750 |
| 28 | ✓ Naprosyn SR 1000 |
| 20 | Maprosyn Sh 1000 |
| 400 | / = 11 111 |
| 100 | ✓ <u>Tilcotil</u> |
| 1 | ✓ AFT |
| | |
| | |
| 60 | ✓ Celebrex |
| δU | ✓ Celecoxib Pfizer |
| 30 | ✓ Celecoxib Plizer ✓ Celebrex |
| 30 | ✓ Celecoxib Pfizer |
| | 60 30 |

| | Subsidy (Manufacturer's Price | | Fully sidised | Brand or Generic |
|--|----------------------------------|---------|------------------|--|
| | \$ | Per | | Manufacturer |
| Topical Products for Joint and Muscular Pain | | | | |
| CAPSAICIN | | | | |
| Crm 0.025% - Special Authority see SA1289 below - Retai | I | | | |
| pharmacy | 9.75 | 45 g OP | | Co-Rub Osteo S29 Costrix |
| | 13.00 | 60 g OP | ✓ R | Rugby Capsaicin Topical Cream ^{©29} |
| ■ SA1289 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valiosteoarthritis that is not responsive to paracetamol and oral non- | | | | |
| Antirheumatoid Agents | | | | |
| HYDROXYCHLOROQUINE SULPHATE | 7.00 | 100 | ./ 1 | |

| DROXYCHLOROQUINE SULPHATE | | |
|--|----------------|--|
| Tab 200 mg7.80 |) 100 | ✓ Ipca- Hydroxychloroquine |
| 8.78 | 3 | ✓ Plaquenil |
| Ipca-Hydroxychloroquine to be Principal Supply on 1 May 2025 | | |
| , , , | | |
| | | |
| | | ✓ Arava |
| Tab 20 mg | 30 | ✓ Arava |
| VICILLAMINE | | |
| Tab 125 mg | 3 100 | ✓ D-Penamine |
| Tab 250 mg | | ✓ D-Penamine |
| Ipca-Hydroxychloroquine to be Principal Supply on 1 May 2025 Iquenil Tab 200 mg to be delisted 1 May 2025) ILUNOMIDE Tab 10 mg | 30 30 30 | ✓ <u>Arava</u> ✓ <u>Arava</u> ✓ D-Penamine |

| Drugs Affecting Bone Metabolism | | |
|--|---|----------------|
| Alendronate for Osteoporosis | | |
| ALENDRONATE SODIUM * Tab 70 mg3.10 | 4 | ✓ Fosamax |
| ALENDRONATE SODIUM WITH COLECALCIFEROL * Tab 70 mg with colecalciferol 5,600 iu | 4 | ✓ Fosamax Plus |
| | | |

Other Treatments

DENOSUMAB - Special Authority see SA2427 below - Retail pharmacy

Note: Denosumab inj 60 mg per 1 ml pre-filled syringe is Medsafe approved for use in osteoporosis. Denosumab inj 120 mg per 1.7 ml vial is Medsafe approved for use in hypercalcaemia of malignancy.

 Inj 120 mg per 1.7 ml vial
 500.00
 1
 ✓ Xgeva

 Inj 60 mg prefilled syringe
 326.00
 1
 ✓ Prolia

⇒SA2427 Special Authority for Subsidy

Initial application — (Osteoporosis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 Either:
 - 2.1 The patient is female and postmenopausal; or
 - 2.2 The patient is male or non-binary; and
- 3 Any of the following:
 - 3.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
 - 3.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons; or
 - 3.3 History of two significant osteoporotic fractures demonstrated radiologically; or
 - 3.4 Documented T-Score less than or equal to -3.0 (see Note); or
 - 3.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
 - 3.6 Patient has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 4 Zoledronic acid is contraindicated because the patient's creatinine clearance is less than 35 mL/min; and
- 5 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes); and
- 6 The patient must not receive concomitant treatment with any other funded antiresorptive agent for this condition or teriparatide.

Initial application — (Hypercalcaemia) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has hypercalcaemia of malignancy; and
- 2 Patient has severe renal impairment.

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
 Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with denosumab
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body
- e) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: risedronate sodium tab 35 mg once weekly; alendronate sodium tab 70 mg or tab 70 mg with cholecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy

PAMIDRONATE DISODIUM

| Inj 3 mg per ml, 10 ml vial | 32.49 | 1 | Pamisol |
|-----------------------------|-------|---|-----------|
| Inj 6 mg per ml, 10 ml vial | 88.11 | 1 | ✓ Pamisol |
| Inj 9 mg per ml, 10 ml vial | 94.34 | 1 | ✓ Pamisol |

| | Subsidy (Manufacturer's Price) | Sub | Fully sidised | Brand or Generic | |
|--|-----------------------------------|--------|------------------|---------------------|--|
| | \$ | Per | • | Manufacturer | |
| RALOXIFENE HYDROCHLORIDE - Special Authority see SA1 | 779 below – Retail ph | armacy | | | |
| * Tab 60 mg | 53.76 | 28 | ✓ E | vista | |

⇒SA1779 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Notes); or
- 2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
- 3 History of two significant osteoporotic fractures demonstrated radiologically; or
- 4 Documented T-Score less than or equal to -3.0 (see Notes); or
- 5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Notes); or
- 6 Patient has had a Special Authority approval for zoledronic acid (Underlying cause Osteoporosis) or has had a Special Authority approval for alendronate (Underlying cause - Osteoporosis) prior to 1 February 2019.

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
 Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for raloxifene funding.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis, and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

RISEDRONATE SODIUM

| Tab 35 mg2.50 | 4 | ✓ Risedronate Sandoz |
|---|---|----------------------|
| TERIPARATIDE - Special Authority see SA1139 below - Retail pharmacy | | |
| Inj 250 mcg per ml, 2.4 ml | 1 | Teriparatide - Teva |

⇒SA1139 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 18 months for applications meeting the following criteria: All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 The patient has a documented T-score less than or equal to -3.0 (see Notes); and
- 3 The patient has had two or more fractures due to minimal trauma; and
- 4 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes).

Notes:

a) The bone mineral density (BMD) measurement used to derive the T-score must be made using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable

| Subsidy | | Fully | Brand or | |
|----------------------|-----|------------|--------------|--|
| (Manufacturer's Pric | e) | Subsidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

continued...

- b) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: alendronate sodium tab 70 mg or tab 70 mg with colecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily; zoledronic acid 5 mg per year. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy.
- c) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.
- d) A maximum of 18 months of treatment (18 cartridges) will be subsidised.

ZOLEDRONIC ACID

| Inj 0.05 mg per ml, 100 ml, bag | 22.53 | 100 ml OP | ✓ Zoledronic Acid |
|---------------------------------|-------|-----------|-------------------|
| | | | Viatris |

Hyperuricaemia and Antigout

| ٨ | | \sim | ш | INI | വ | |
|---|--|--------|---|-----|---|--|
| | | | | | | |
| | | | | | | |

| ALLOI OTIINOL | | | |
|--|---------------------|-------|-----------------------|
| * Tab 100 mg | 17.99 | 1,000 | ✓ Ipca-Allopurinol |
| * Tab 300 mg | 22.50 | 500 | ✓ Ipca-Allopurinol |
| BENZBROMARONE - Special Authority see SA1963 below | / – Retail pharmacy | | |
| Tab 50 mg | 32.00 | 100 | ✓ Narcaricin mite S29 |

⇒SA1963 Special Authority for Subsidy

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefitting from the treatment; and
- 2 There is no evidence of liver toxicity and patient is continuing to receive regular (at least every three months) liver function tests

COLCHICINE

| COLUNIONE | | | |
|---|------|-----|---------------------|
| * Tab 500 mcg | 6.00 | 100 | ✓ Colgout |
| FEBUXOSTAT - Special Authority see SA2054 below - F | | | |
| Tab 80 mg | ' | 28 | ✓ Febuxostat (Teva) |
| Tab 120 mg | | 28 | ✓ Febuxostat (Teva) |

⇒SA2054 Special Authority for Subsidy

Initial application — (Gout) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with gout; and
- 2 Any of the following:
 - 2.1 The patient has a serum urate level greater than 0.36 mmol/l despite treatment with allopurinol at doses of at least 600 mg/day and addition of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.2 The patient has experienced intolerable side effects from allopurinol such that treatment discontinuation is required and serum urate remains greater than 0.36 mmol/l despite use of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.3 The patient has renal impairment such that probenecid is contraindicated or likely to be ineffective and serum urate remains greater than 0.36 mmol/l despite optimal treatment with allopurinol (see Note); or
 - 2.4 The patient has previously had an initial Special Authority approval for benzbromarone for treatment of gout...

Initial application — (Tumour lysis syndrome) only from a haematologist or oncologist. Approvals valid for 6 weeks for

100

100

100

✓ Dantrium ✓ Dantrium S29 S29

✓ Dantrium

✓ Norflex

| | MU | ISCULOSKEI | LETAL SYSTEM |
|---|---|----------------------------|---|
| | Subsidy (Manufacturer's Price) \$ | Fully Subsidised Per | Brand or Generic Manufacturer |
| continued applications meeting the following criteria: Both: 1 Patient is scheduled to receive cancer therapy carrying ar 2 Patient has a documented history of allopurinol intoleranc Renewal — (Gout) from any relevant practitioner. Approvals va | e. | - | |
| patient is benefitting from treatment. Renewal — (Tumour lysis syndrome) only from a haematolog treatment remains appropriate and the patient is benefitting from PROBENECID * Tab 500 mg | treatment. | | weeks where the |
| Muscle Relaxants | | | |
| BACLOFEN * Tab 10 mg | t11.55 ents where oral antispedorsed accordingly. | 1 | Medsurge Sintetica Baclofen |
| a) Subsidised only for use in a programmable pump in have caused intolerable side effects and the prescrip b) Sintetica Baclofen Intrathecal to be Principal Supply (Medsurge Inj 2 mg per ml, 5 ml ampoule to be delisted 1 March DANTROLENE | tion is endorsed acco | | Intrathecal have been ineffective or |

(Dantrium Cap 25 mg to be delisted 1 April 2025)

ORPHENADRINE CITRATE

Subsidy (Manufacturer's Price) Fully Subsidised

Brand or Generic Manufacturer

Agents for Parkinsonism and Related Disorders

| Dopamine Agonists and Related Agents |
|---|
|---|

| AMANTADINE HYDROCHLORIDE | | | |
|---|--------|--------|-----------------|
| ▲ Cap 100 mg | 38 24 | 60 | ✓ Symmetrel |
| _ Oap 100 mg | 63.73 | 100 | ✓ Symmetrel |
| APOMORPHINE HYDROCHLORIDE | 00.70 | 100 | - Cymmon or |
| ▲ Inj 10 mg per ml, 2 ml ampoule | E0 E0 | 5 | ✓ Movapo |
| ▲ Inj 10 mg per ml, 2 ml ampoule | | 5 5 | ✓ Movapo |
| , , , | 121.04 | 3 | • wovapo |
| ENTACAPONE | 40.70 | 400 | 4 - . |
| ▲ Tab 200 mg | | 100 | ✓ Entapone |
| (Comton Tob 200 mg to be delicted 1 July 2025) | 18.04 | | ✓ Comtan |
| (Comtan Tab 200 mg to be delisted 1 July 2025) | | | |
| LEVODOPA WITH BENSERAZIDE | | | |
| * Tab dispersible 50 mg with benserazide 12.5 mg | | 100 | ✓ Madopar Rapid |
| * Cap 50 mg with benserazide 12.5 mg | | 100 | ✓ Madopar 62.5 |
| * Cap 100 mg with benserazide 25 mg | | 100 | ✓ Madopar 125 |
| * Cap long-acting 100 mg with benserazide 25 mg | | 100 | ✓ Madopar HBS |
| * Cap 200 mg with benserazide 50 mg | 26.25 | 100 | ✓ Madopar 250 |
| LEVODOPA WITH CARBIDOPA | | | |
| * Tab 100 mg with carbidopa 25 mg | | 100 | Sinemet |
| * Tab long-acting 200 mg with carbidopa 50 mg | | 100 | ✓ Sinemet CR |
| * Tab 250 mg with carbidopa 25 mg | 39.49 | 100 | ✓ Sinemet |
| LEVODOPA WITH CARBIDOPA AND ENTACAPONE | | | |
| * Tab 50 mg with carbidopa 12.5 mg and entacapone 200 mg | | 100 | ✓ Stalevo |
| * Tab 100 mg with carbidopa 25 mg and entacapone 200 mg | | 100 | ✓ Stalevo |
| * Tab 150 mg with carbidopa 37.5 mg and entacapone 200 mg | | 100 | ✓ Stalevo |
| * Tab 200 mg with carbidopa 50 mg and entacapone 200 mg | 51.23 | 100 | ✓ Stalevo |
| PRAMIPEXOLE HYDROCHLORIDE | | | |
| ▲ Tab 0.25 mg | 5.51 | 100 | ✓ Ramipex |
| ▲ Tab 1 mg | 18.66 | 100 | ✓ Ramipex |
| RASAGILINE | | | |
| * Tab 1 mg | 53.50 | 30 | ✓ Azilect S29 |
| ROPINIROLE HYDROCHLORIDE | | | |
| ▲ Tab 0.25 mg | 4 05 | 84 | ✓ Ropin |
| ▲ Tab 1 mg | | 84 | ✓ Ropin |
| ▲ Tab 2 mg | | 84 | ✓ Ropin |
| ▲ Tab 5 mg | | 84 | ✓ Ropin |
| TOLCAPONE | | | |
| ▲ Tab 100 mg | 152 38 | 100 | ✓ Tasmar |
| _ 140 100 mg | | 100 | - Iugiliui |

Anticholinergics

| BENZATROPINE MESYLAT | TE | Ξ |
|----------------------|----|---|
|----------------------|----|---|

| LIVE/TITIOT IIVE WILCOTE/TIE | | | |
|------------------------------|-------|----|------------|
| Tab 2 mg | 9.59 | 60 | ✓ Benztrop |
| Inj 1 mg per ml, 2 ml | 95.00 | 5 | ✓ Phebra |
| | | | |

a) Up to 10 inj available on a PSO

b) Only on a PSO

| | | | INE | NVOUS STSTEIN |
|--|---|-------|---------------------|--|
| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | |
| PROCYCLIDINE HYDROCHLORIDE Tab 5 mg | 7.40 | 100 | ✓ | Kemadrin |
| Agents for Essential Tremor, Chorea and Relate | ed Disorders | | | |
| RILUZOLE – Special Authority see SA1403 below – Retail phan Wastage claimable Tab 50 mg | • | 56 | ✓ | Rilutek |
| ■ SA1403 Special Authority for Subsidy Initial application only from a neurologist or respiratory specialis following criteria: All of the following: | | | | |
| 1 The patient has amyotrophic lateral sclerosis with disease 2 The patient has at least 60 percent of predicted forced vit 3 The patient has not undergone a tracheostomy; and 4 The patient has not experienced respiratory failure; and 5 Any of the following: 5.1 The patient is ambulatory; or 5.2 The patient is able to use upper limbs; or 5.3 The patient is able to swallow. | | | | e initial application; and |
| Renewal from any relevant practitioner. Approvals valid for 18 manual All of the following: 1 The patient has not undergone a tracheostomy; and 2 The patient has not experienced respiratory failure; and 3 Any of the following: 3.1 The patient is ambulatory; or 3.2 The patient is able to use upper limbs; or 3.3 The patient is able to swallow. | nonths for application | s mee | eting the fo | llowing criteria: |
| TETRABENAZINE Tab 25 mg | 106.59 | 112 | 1 | <u>Motetis</u> |
| Anaesthetics | | | | |
| Local | | | | |
| LIDOCAINE [LIGNOCAINE] Gel 2%, tube – Subsidy by endorsement | | 30 m | | Xylocaine 2% Jelly endorsed accordingly. |

b) Subsidised only if prescribed for urethral, cervical or rectal administration and the prescription is endorsed

10

✓ Instillagel Lido

Gel 2%, 11 ml urethral syringe – Subsidy by endorsement...........59.50

a) Up to 5 each available on a PSO

accordingly.

| | Subsidy | | Fully | Brand or |
|--|-----------------------|-----------|------------|-----------------------|
| | (Manufacturer's Price | e) : | Subsidised | Generic |
| | \$ | Per | • | Manufacturer |
| LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE | | | | |
| Oral (gel) soln 2% | 44.00 | 200 ml | ✓ | Mucosoothe |
| Inj 1%, 5 ml ampoule - Up to 25 inj available on a PSO | 15.00 | 25 | ✓ | Lidocaine-Baxter |
| | 17.50 | 50 | | |
| | (35.00) | | | Xylocaine |
| Inj 2%, 5 ml ampoule - Up to 5 inj available on a PSO | 27.50 | 25 | 1 | Lidocaine-Baxter |
| Inj 1%, 20 ml ampoule - Up to 5 inj available on a PSO | 12.00 | 5 | | |
| | (20.00) | | | Xylocaine |
| Inj 1%, 20 ml vial - Up to 5 inj available on a PSO | 6.85 | 5 | 1 | Lidocaine-Baxter |
| Inj 2%, 20 ml vial - Up to 5 inj available on a PSO | 7.15 | 5 | 1 | Lidocaine-Baxter |
| Inj 10%, 5 ml ampoule - Subsidy by endorsement | CBS | 10 | ✓ | Xylocard 500 S29 |
| Subsidised only for people receiving palliative care serving | ices where other and | algesic a | agents ha | ven't been effective. |

Topical Local Anaesthetics

⇒SA0906 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years where the patient is a child with a chronic medical condition requiring frequent injections or venepuncture.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

| LIDOCAINE [LIGNOCAINE] - Special Authority : | see SA0906 above – Retail ph | narmacy | |
|--|------------------------------|------------------|-------------|
| Crm 4% | 5.40 | 5 g OP | ✓ LMX4 |
| | 27.00 | 30 g OP | ✓ LMX4 |
| LIDOCAINE [LIGNOCAINE] WITH PRILOCAINE | - Special Authority see SA09 | 906 above – Reta | il pharmacy |
| Crm 2.5% with prilocaine 2.5% | 45.00 | 30 g OP | ✓ EMLA |
| Crm 2.5% with prilocaine 2.5% (5 g tubes) | 45.00 | 5 | EMLA |

Analgesics

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 116

Non-opioid Analgesics

| ASPIRIN * Tab dispersible 300 mg - Up to 30 tab available on a PSO | 5.65 | 100 | ✓ Ethics Aspirin |
|---|---------------|---------------|-------------------------------------|
| CAPSAICIN - Subsidy by endorsement | | | |
| Subsidised only if prescribed for post-herpetic neuralgia or diabetic paccordingly. | peripheral ne | europathy and | the prescription is endorsed |
| Crm 0.075% | 11.95 | 45 g OP | ✓ Zo-Rub HP S29 ✓ Zostrix HP |
| • | 15.14 | 57 g OP | ✓ Rugby Capsaicin Topical Cream S29 |
| NEFOPAM HYDROCHLORIDE Tab 30 mg | 23.40 | 90 | ✓ Acupan |

NERVOUS SYSTEM

| | Subsidy (Manufacturer's Price) \$ | S Per | Fully ubsidised | Brand or Generic Manufacturer |
|--|---|----------------------------------|--------------------------------------|--|
| PARACETAMOL | | | | |
| Tab 500 mg - blister pack | | 1,000 | ✓ <u>I</u> | Pacimol Pacimol |
| a) Maximum of 300 tab per prescription; can be wait b) Up to 30 tab available on a PSO c) | ved by endorsement | | | |
| Subsidy by endorsement for higher quantities regular daily dosing for one month or greated annotate the prescription as endorsed wher Maximum of 100 tab per dispensing for non (for non-endorsed patients), then dispense Tab 500 mg - bottle pack — Maximum of 300 tab per | er, and the prescription is e dispensing history sup -endorsed patients. If qu | annota ports a uantities | ted accor ong-term prescrib | dingly. Pharmacists may condition. ed for more than 100 tabs |
| prescription; can be waived by endorsement | 17.92 | 1,000 | √ <u>i</u> | Noumed Paracetamol |
| Subsidy by endorsement for higher quantities is daily dosing for one month or greater, and the p prescription as endorsed where dispensing hist Maximum of 100 tab per dispensing for non-en- non-endorsed patients), then dispense in repeat | orescription is annotated cory supports a long-term dorsed patients. If quant | accordi conditi tities pre | ngly. Pha on. escribed f | armacists may annotate the or more than 100 tabs (for |
| Oral liq 120 mg per 5 ml | 3.98 | 200 ml | √ <u>i</u> | Paracetamol (Ethics) |
| a) Maximum of 600 ml per prescription; can be waiv b) Up to 200 ml available on a PSO c) Not in combination d) | ed by endorsement | | | |
| 1) Maximum of 200 ml per dispensing for non- non-endorsed patients), then dispense in re 2) Subsidy by endorsement for higher quantitic regular daily dosing for one month or greate Pharmacists may annotate the prescription condition. | peat dispensing not exce es is available for patient er and the prescription is | eeding 2 s with lo endorse | 200 ml pe ong term o ed or ann | r dispensing. conditions who require otated accordingly. |
| 3) Note: 200 ml presentations of paracetamol Pharmacist) under the provisions in Part I o 4) Note: Direct Provision by a pharmacist of u | f Section A | | | ` |
| conjunction with immunisation of a child und | der 2 years of age with m | | coccal B r | |
| a) Maximum of 600 ml per prescription; can be waivb) Up to 200 ml available on a PSOc) Not in combination | | 200 1111 | | unoi |
| d) Maximum of 200 ml per dispensing for non-non-endorsed patients), then dispense in re Subsidy by endorsement for higher quantitie | peat dispensing not exce | eeding 2 | 200 ml pe | r dispensing. |

regular daily dosing for one month or greater and the prescription is endorsed or annotated accordingly.

Pharmacists may annotate the prescription as endorsed where dispensing history supports a long-term

condition.

3) Note: 200 ml presentations of paracetamol oral liquid may be supplied on BSO to a Vaccinator (other than a Pharmacist) under the provisions in Part I of Section A

4) Note: Direct Provision by a pharmacist of up to 200 ml permitted under the provisions in Part I of Section A in conjunction with immunisation of a child under 2 years of age with meningococcal B multicomponent vaccine.

Suppos 125 mg4.29 10 ✓ Gacet

| | NERVOUS SYSTEM | | | | |
|------|---|---|-----------------|---------------------|--------------------------|
| | | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | |
| * | Suppos 250 mg | | 10 50 | | Gacet Gacet |
| 0 | pioid Analgesics | | | | |
| CC | DEINE PHOSPHATE - Safety medicine; prescriber may dete | rmine dispensina fre | auen | CV | |
| • | Tab 15 mg | , , | 100 | | Noumed |
| | Tab 30 mg | | 100 | | Aspen |
| | | | | | Noumed |
| | Tab 60 mg | 13.89 | 100 | _ | Noumed |
| DIF | HYDROCODEINE TARTRATE | | | | |
| ווט | Tab long-acting 60 mg | 8 60 | 60 | 1 | DHC Continus |
| | | | 00 | • | Dirio Continua |
| FE | NTANYL | | | | |
| | a) Only on a controlled drug form | | | | |
| | b) No patient co-payment payable | | | | |
| | c) Safety medicine; prescriber may determine dispensing fre | , , | 40 | | Davishan and Mule |
| | Inj 50 mcg per ml, 2 ml ampoule | 4.25 | 10 | • | Boucher and Muir |
| | Boucher and Muir to be Principal Supply on 1 May 2025 | 0.44 | 10 | ./ | Boucher and Muir |
| | Inj 50 mcg per ml, 10 ml ampoule | 9.41 | 10 | • | boucher and wuir |
| | Boucher and Muir to be Principal Supply on 1 May 2025 Patch 12.5 mcg per hour | 6.00 | 5 | ./ | Fentanyl Sandoz |
| | Patch 25 mcg per hour | | 5 | | Fentanyl Sandoz |
| | Patch 50 mcg per hour | | 5 | | Fentanyl Sandoz |
| | Patch 75 mcg per hour | | 5 | | Fentanyl Sandoz |
| | Patch 100 mcg per hour | | 5 | | Fentanyl Sandoz |
| | • . | 10.07 | J | • | r chianyi dandoz |
| IVIE | THADONE HYDROCHLORIDE | | | | |
| | a) Only on a controlled drug form | | | | |
| | b) No patient co-payment payable | | | | |
| | c) Safety medicine; prescriber may determine dispensing fre | , , | 10 | ./ | Mathadana DNM |
| | Tab 5 mg | | 10 200 n | | Methadone BNM Biodone |
| | Oral lig 5 mg per ml | | 200 II 200 n | | Biodone Forte |
| | Oral liq 5 mg per ml Oral liq 10 mg per ml | | 200 ii 200 n | | Biodone Extra Forte |
| | Inj 10 mg per ml, 1 ml | | 10 | | AFT |
| | , 01 | 00.30 | 10 | • | ALI |
| MC | RPHINE HYDROCHLORIDE | | | | |
| | a) Only on a controlled drug form | | | | |
| | b) No patient co-payment payable | | | | |
| | c) Safety medicine; prescriber may determine dispensing fre | | 000 | | DA 14 b |
| | Oral liq 1 mg per ml | | 200 n | | RA-Morph |
| | Oral lig 5 mg per ml | | 200 n | | RA-Morph |
| | Oral lig 10 mg per ml | | 200 n | _ | RA-Morph |
| | Oral liq 10 mg per ml | 40.25 | 200 n | II • | RA-Morph |

| | Subsidy | | Fully | Brand or |
|---|------------------------|--------------|------------|-----------------------------------|
| | (Manufacturer's Price) | | Subsidised | |
| | \$ | Per | | Manufacturer |
| MORPHINE SULPHATE | | | | |
| a) Only on a controlled drug form | | | | |
| b) No patient co-payment payable | | | | |
| c) Safety medicine; prescriber may determine dispensing free | | | | |
| Tab immediate-release 10 mg | | 10 | | Sevredol |
| Tab immediate-release 20 mg | | 10 | | Sevredol |
| Cap long-acting 10 mg | | 10 | | m-Eslon |
| Cap long-acting 30 mg | | 10 | | m-Eslon |
| Cap long-acting 60 mg | | 10 | | m-Eslon |
| Cap long-acting 100 mg | 10.50 | 10 | | m-Eslon |
| Oral liq 2 mg per ml | 16.31 | 100 m | | Wockhardt S29 |
| | 29.80 | | | Oramorph |
| | | | ✓ | Oramorph CDC |
| | | | | S29 S29 |
| Inj 5 mg per ml, 1 ml ampoule - Up to 5 inj available on a PS | SO5.38 | 5 | ✓ | Medsurge |
| Inj 10 mg per ml, 1 ml ampoule - Up to 5 inj available on a P | SO 4.68 | 5 | ✓ | Medsurge |
| Inj 15 mg per ml, 1 ml ampoule - Up to 5 inj available on a P | SO5.53 | 5 | | Medsurge |
| Inj 30 mg per ml, 1 ml ampoule - Up to 5 inj available on a P | SO6.28 | 5 | ✓ | Medsurge |
| OXYCODONE HYDROCHLORIDE | | | | |
| a) Only on a controlled drug form | | | | |
| b) No patient co-payment payable | | | | |
| c) Safety medicine; prescriber may determine dispensing fre | equency | | | |
| Tab controlled-release 5 mg | | 20 | 1 | Oxycodone Sandoz |
| 145 6011101104 1010400 0 119 | 3.77 | 28 | | Oxycodone Sandoz |
| | 0.17 | | | S29 S29 |
| | 4.04 | 30 | ./ | OxyContin S29 |
| Tab immediate-release 5 mg | | 100 | | Oxycodone Amneal |
| Tab controlled-release 10 mg | | 20 | | Oxycodone Sandoz |
| Tab controlled-release to mg | 3.77 | 28 | | Oxycodone Sandoz |
| | 3.77 | 20 | • | S29 S29 |
| Tab immediate values 40 mm | 10.77 | 100 | | |
| Tab immediate-release 10 mg | | 100 | | Oxycodone Amneal |
| Tab controlled-release 20 mg | | 20 | | Oxycodone Sandoz |
| Tab immediate-release 20 mg | | 100 | | Oxycodone Amneal |
| Tab controlled release 40 mg | | 20 | | Oxycodone Sandoz Oxycodone Sandoz |
| Tab controlled-release 80 mg | | 20 20 | | OxyCodone Sandoz OxyNorm |
| Cap immediate-release 20 mg | | ∠u 250 m | | Oxycodone Lucis |
| Oral liq 1 mg per ml | | 250 III 5 | | Hameln |
| Inj 10 mg per ml, 2 ml ampoule | | 5 | | Hameln |
| | | 5 | | Hameln |
| Inj 50 mg per ml, 1 ml ampoule(OxyNorm Cap immediate-release 20 mg to be delisted 1 March 2 | | J | • | <u>Hamelli</u> |
| | , | | | |
| PARACETAMOL WITH CODEINE – Safety medicine; prescriber | | | | |
| * Tab paracetamol 500 mg with codeine phosphate 8 mg | 27.50 | 1,000 | • | Paracetamol + |
| | | | | |

Codeine (Relieve)

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Brand or Subsidised Generic Manufacturer |
|--|---|---------------------------------|---|
| PETHIDINE HYDROCHLORIDE | | | |
| a) Only on a controlled drug form | | | |
| b) No patient co-payment payable | | | |
| c) Safety medicine; prescriber may determine dispensing | | | |
| Tab 50 mg | | 10 | ✓ Noumed Pethidine |
| Inj 50 mg per ml, 1 ml ampoule - Up to 5 inj available on a | 1 PSU 29.88 | 5 | ✓ DBL Pethidine Hydrochloride |
| Inj 50 mg per ml, 2 ml ampoule – Up to 5 inj available on a | DSO 20.72 | 5 | ✓ DBL Pethidine |
| Inj 50 mg per mi, 2 mi ampoule – Op to 5 inj available on a | 173030.72 | 5 | Hydrochloride |
| TRAMADOL HYDROCHLORIDE | | | Tiyaroomonac |
| Tab sustained-release 100 mg | 1 95 | 20 | ✓ Tramal SR 100 |
| Tab sustained-release 150 mg | | 20 | ✓ Tramal SR 150 |
| Tab sustained-release 200 mg | | 20 | ✓ Tramal SR 200 |
| Cap 50 mg | | 100 | |
| | | | |
| Antidepressants | | | |
| Cyclic and Related Agents | | | |
| AMITRIPTYLINE - Safety medicine; prescriber may determine | dispensing frequency | | |
| Tab 10 mg | | 100 | ✓ Arrow-Amitriptyline |
| Tab 25 mg | 1.99 | 100 | ✓ Arrow-Amitriptyline |
| Tab 50 mg | 3.14 | 100 | Arrow-Amitriptyline |
| CLOMIPRAMINE HYDROCHLORIDE - Safety medicine; pres | criber may determine d | isper | nsing frequency |
| Tab 10 mg | | 30 | Clomipramine Teva |
| Tab 25 mg | 11.99 | 30 | Clomipramine Teva |
| | 16.99 | 50 | APO Clomipramine |
| | 39.97 | 100 | |
| Cap 10 mg | | 28 | ✓ Clomipramine Teva |
| Cap 25 mg | 35.50 | 28 | Clomipramine Teva |
| (Clomipromine Teva Tab 10 mg to be delisted 1 July 2025) | | | |
| (Clomipramine Teva Tab 25 mg to be delisted 1 July 2025) | | | |
| (Anafranil S29 Tab 25 mg to be delisted 1 July 2025) | | | |
| (Clomipramine Teva Cap 10 mg to be delisted 1 July 2025) (Clomipramine Teva Cap 25 mg to be delisted 1 July 2025) | | | |
| DOSULEPIN [DOTHIEPIN] HYDROCHLORIDE - Subsidy by | andaraamant | | |
| . , , | | | |
| | | | |
| Safety medicine; prescriber may determine dispensing Subsidy by endorsement – Subsidised for nationts who | wara taking docularin | [doth | ienin) hydrochloride prior to 1 lune |
| b) Subsidy by endorsement - Subsidised for patients who | were taking dosulepin | | |
| Subsidy by endorsement – Subsidised for patients who 2019 and the prescription is endorsed accordingly. Pha | were taking dosulepin armacists may annotate | | |
| Subsidy by endorsement – Subsidised for patients who 2019 and the prescription is endorsed accordingly. Pha exists a record of prior dispensing of dosulepin [dothiep | were taking dosulepin armacists may annotate in] hydrochloride. | | prescription as endorsed where the |
| Subsidy by endorsement – Subsidised for patients who 2019 and the prescription is endorsed accordingly. Pha | were taking dosulepin armacists may annotate in] hydrochloride. 3.85 | e the | |
| b) Subsidy by endorsement – Subsidised for patients who 2019 and the prescription is endorsed accordingly. Pha exists a record of prior dispensing of dosulepin [dothiep Tab 75 mg | were taking dosulepin armacists may annotate in] hydrochloride. 3.85 | the 30 | prescription as endorsed where the ✓ Dosulepin Viatris |
| b) Subsidy by endorsement – Subsidised for patients who 2019 and the prescription is endorsed accordingly. Pha exists a record of prior dispensing of dosulepin [dothiep Tab 75 mg | were taking dosulepin armacists may annotate in] hydrochloride3.85 | 30 50 | prescription as endorsed where the Dosulepin Viatris Dosulepin Viatris \$29 |
| b) Subsidy by endorsement – Subsidised for patients who 2019 and the prescription is endorsed accordingly. Phaexists a record of prior dispensing of dosulepin [dothiep Tab 75 mg | were taking dosulepin armacists may annotate in] hydrochloride3.857.83 | 30 50 | prescription as endorsed where the Dosulepin Viatris Dosulepin Viatris \$29 |
| b) Subsidy by endorsement – Subsidised for patients who 2019 and the prescription is endorsed accordingly. Phaexists a record of prior dispensing of dosulepin [dothiep Tab 75 mg | were taking dosulepin armacists may annotate in] hydrochloride3.857.83 | 30 50 ensing | prescription as endorsed where the Dosulepin Viatris Dosulepin Viatris \$29 g frequency Tofranil |
| b) Subsidy by endorsement – Subsidised for patients who 2019 and the prescription is endorsed accordingly. Phaexists a record of prior dispensing of dosulepin [dothiep Tab 75 mg | were taking dosulepin armacists may annotate in] hydrochloride | 30 50 ensing | prescription as endorsed where the Dosulepin Viatris Dosulepin Viatris \$29 g frequency Tofranil |
| b) Subsidy by endorsement – Subsidised for patients who 2019 and the prescription is endorsed accordingly. Phaexists a record of prior dispensing of dosulepin [dothiep Tab 75 mg | were taking dosulepin armacists may annotate in] hydrochloride | 30 50 ensing 50 100 | prescription as endorsed where the Dosulepin Viatris Dosulepin Viatris 229 g frequency Tofranil Tofranil |

| | | | NE | RVOUS SYSTEM |
|--|-----------------------------------|--------------------------|------------------|---|
| | Subsidy (Manufacturer's Price) | Subs Per | Fully sidised | Generic |
| NORTRIPTYLINE HYDROCHLORIDE – Safety medicine; prescr Tab 10 mg Tab 25 mg | 2.46 | lispensing 100 180 | 1 | uency <u>Norpress</u> <u>Norpress</u> |
| Monoamine-Oxidase Inhibitors (MAOIs) - Non So | elective | | | |
| TRANYLCYPROMINE SULPHATE * Tab 10 mg | 22.94 | 50 | / | Parnate |
| Monoamine-Oxidase Type A Inhibitors | | | | |
| MOCLOBEMIDE * Tab 150 mg * Tab 300 mg | | 60 60 | _ | <u>Aurorix</u> <u>Aurorix</u> |
| Selective Serotonin Reuptake Inhibitors | | | | |
| CITALOPRAM HYDROBROMIDE * Tab 20 mg | 2.86 | 84 | / | <u>Celapram</u> |
| * Tab 10 mg | 0.79 1.07 | 28 | | Ipca-Escitalopram Escitalopram (Ethics) |
| * Tab 20 mg | 1.49 | 28 | 1 | Ipca-Escitalopram |
| FLUOXETINE HYDROCHLORIDE * Tab dispersible 20 mg, scored – Subsidy by endorsement Subsidised by endorsement | 2.50 | 28 | 1 | Fluox |
| When prescribed for a patient who cannot swallow accordingly; or When prescribed in a daily dose that is not a multip endorsed. Note: Tablets should be combined with | le of 20 mg in which | case the p | rescr | ription is deemed to be |
| * Cap 20 mg | 3.13 | 90 | 1 | Arrow-Fluoxetine |
| PAROXETINE * Tab 20 mg SERTRALINE | 4.11 | 90 | / | Loxamine |
| * Tab 50 mg* * Tab 100 mg | | 30 30 | | Setrona Setrona |
| Other Antidepressants | | | | |
| MIRTAZAPINE Tab 30 mg | 2.60 | 28 30 | | Noumed Noumed |
| Tab 45 mg | 3.45 | 28 30 | 1 | Noumed Noumed |
| VENLAFAXINE | | | | |
| Cap 37.5 mg Cap 75 mg Cap 150 mg | 10.32 | 84 84 84 | 1 | Enlafax XR Enlafax XR Enlafax XR |

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

Antiepilepsy Drugs

Agents for Control of Status Epilepticus

| DIAZEPAM – Safety medicine; prescriber may determine dispensing frequency Inj 5 mg per ml, 2 ml ampoule – Subsidy by endorsement27.92 | 5 | ✓ Hospira |
|---|---|---------------------------|
| a) Up to 5 inj available on a PSO | | · |
| b) Only on a PSO | | |
| c) PSO must be endorsed "not for anaesthetic procedures". | | |
| Rectal tubes 5 mg - Up to 5 tube available on a PSO54.58 | 5 | ✓ <u>Stesolid</u> |
| PHENYTOIN SODIUM | | |
| * Inj 50 mg per ml, 2 ml ampoule - Up to 5 inj available on a | | |
| PSO104.58 | 5 | Hospira |
| * Inj 50 mg per ml, 5 ml ampoule - Up to 5 inj available on a | | |
| PSO154.01 | 5 | Hospira |
| | | |

Control of Epilepsy

| ★ Tab 200 mg 14.53 100 ✓ Tegretol ★ Tab long-acting 200 mg 16.98 100 ✓ Tegretol CR ★ Tab 400 mg 33.96 200 ✓ Tegretol CR ★ Tab long-acting 400 mg 34.58 100 ✓ Tegretol ★ Tab long-acting 400 mg 39.17 100 ✓ Tegretol ★ Oral liq 20 mg per ml 26.37 250 ml ✓ Tegretol CLOBAZAM - Safety medicine; prescriber may determine dispensing frequency Tab 10 mg 9.12 50 ✓ Frisium CLONAZEPAM - Safety medicine; prescriber may determine dispensing frequency Oral drops 2.5 mg per ml 7.38 10 ml OP ✓ Rivotril ETHOSUXIMIDE Cap 250 mg 78.89 56 ✓ Essential Ethosuximide Ethosuximide \$239 Tab 100 mg per 5 ml 56.35 200 ml ✓ Zarontin Oral liq 250 mg per 5 ml 56.35 200 ml ✓ Zarontin GABAPENTIN Note: Not subsidised in combination with subsidised pregabalin ★ Cap 100 mg 6.45 100 ✓ Nupentin ★ Cap 400 mg 8.45 100 ✓ Nupentin ★ Cap 300 mg | CARBAMAZEPINE | | | |
|--|--|------------------------|----------|-------------------|
| ** Tab long-acting 200 mg 16.98 100 ✓ Tegretol CR ** Tab 400 mg 33.96 200 ✓ Tegretol CR ** Tab long-acting 400 mg 34.58 100 ✓ Tegretol CR ** Tab long-acting 400 mg 39.17 100 ✓ Tegretol CR ** Oral liq 20 mg per ml 26.37 250 ml ✓ Tegretol CR ** CLOBAZAM – Safety medicine; prescriber may determine dispensing frequency Oral drops 2.5 mg per ml 9.12 50 ✓ Frisium CLONAZEPAM – Safety medicine; prescriber may determine dispensing frequency Oral drops 2.5 mg per ml 7.38 10 ml OP ✓ Rivotril ETHOSUXIMIDE Cap 250 mg 78.89 56 ✓ Essential Ethosuximide \$29 Toral liq 250 mg per 5 ml 56.35 200 ml ✓ Zarontin GABAPENTIN Note: Not subsidised in combination with subsidised pregabalin * Cap 300 mg 6.45 100 ✓ Nupentin * Cap 300 mg 8.45 100 ✓ Nupentin * Cap 400 mg 10.26 100 ✓ Nupentin * Cap 400 mg 10.26 100 ✓ Nupentin * Cap 400 mg 10.26 100 ✓ Nupentin * Cap 400 mg 25.04 | * Tab 200 mg | 14.53 | 100 | ✓ Tegretol |
| # Tab 400 mg | | | | • |
| ** Tab 400 mg 34.58 100 ✓ Tegretol ** Tab long-acting 400 mg 39.17 100 ✓ Tegretol CR ** Oral liq 20 mg per ml 26.37 250 ml ✓ Tegretol CLOBAZAM – Safety medicine; prescriber may determine dispensing frequency Tab 10 mg 9.12 50 ✓ Frisium CLONAZEPAM – Safety medicine; prescriber may determine dispensing frequency Oral drops 2.5 mg per ml 7.38 10 ml OP ✓ Rivotril ETHOSUXIMIDE 78.89 56 ✓ Essential Ethosuximide S29 Cap 250 mg 78.89 56 ✓ Essential Ethosuximide S29 Total liq 250 mg per 5 ml 56.35 200 ml ✓ Zarontin Oral liq 250 mg per 5 ml 56.35 200 ml ✓ Zarontin GABAPENTIN Note: Not subsidised in combination with subsidised pregabalin X Cap 100 mg X Nupentin * Cap 300 mg 8.45 100 Nupentin * Cap 300 mg 8.45 100 Nupentin * Cap 400 mg 10.26 100 Nupentin * LACOSAMIDE - Special Authority see SA2267 on the next page - Retail pharmacy A Tab 50 mg 4 ✓ Vimpat A Tab 100 mg 50.06 14 ✓ | * Tab long-acting 200 mg | | | • |
| ** Tab long-acting 400 mg 39.17 100 ✓ Tegretol CR ** Oral liq 20 mg per ml 26.37 250 ml ✓ Tegretol CLOBAZAM – Safety medicine; prescriber may determine dispensing frequency Tab 10 mg 9.12 50 ✓ Frisium CLONAZEPAM – Safety medicine; prescriber may determine dispensing frequency Oral drops 2.5 mg per ml 7.38 10 ml OP ✓ Rivotril ETHOSUXIMIDE 78.89 56 ✓ Essential Ethosuximide S29 Cap 250 mg 78.89 56 ✓ Zarontin Oral liq 250 mg per 5 ml 56.35 200 ml ✓ Zarontin GABAPENTIN Note: Not subsidised in combination with subsidised pregabalin A.45 100 ✓ Nupentin ★ Cap 100 mg 8.45 100 ✓ Nupentin ★ Cap 300 mg 8.45 100 ✓ Nupentin ★ Cap 400 mg 10.26 100 ✓ Nupentin ★ Cap 400 mg 10.26 100 ✓ Nupentin ★ Tab 50 mg 25.04 14 ✓ Vimpat ▲ Tab 50 mg 50.06 14 ✓ Vimpat ▲ Tab 150 mg 75.10 14 ✓ Vimpat ▲ Tab 150 mg 75.10 14< | | | | • |
| ★ Oral liq 20 mg per ml 26.37 250 ml ✓ Tegretol CLOBAZAM - Safety medicine; prescriber may determine dispensing frequency Tab 10 mg 9.12 50 ✓ Frisium CLONAZEPAM - Safety medicine; prescriber may determine dispensing frequency Oral drops 2.5 mg per ml 7.38 10 ml OP ✓ Rivotril ETHOSUXIMIDE 78.89 56 ✓ Essential Ethosuximide S29 Cap 250 mg 78.89 56 ✓ Zarontin Oral liq 250 mg per 5 ml 56.35 200 ml ✓ Zarontin GABAPENTIN Note: Not subsidised in combination with subsidised pregabalin Xarontin Xarontin ★ Cap 100 mg 6.45 100 Xarontin ★ Cap 300 mg 8.45 100 Xarontin ★ Cap 400 mg 10.26 100 Xarontin LACOSAMIDE - Special Authority see SA2267 on the next page - Retail pharmacy Xarontin Xarontin Xarontin A Tab 50 mg 25.04 14 Yimpat A Tab 150 mg 75.10 14 Yimpat A Tab 150 mg 75.10 14 Yimpat | | | | • |
| CLOBAZAM - Safety medicine; prescriber may determine dispensing frequency 9.12 50 ✓ Frisium CLONAZEPAM - Safety medicine; prescriber may determine dispensing frequency 7.38 10 ml OP ✓ Rivotril ETHOSUXIMIDE Cap 250 mg 78.89 56 ✓ Essential Ethosuximide \$29 140.88 100 ✓ Zarontin Oral liq 250 mg per 5 ml 56.35 200 ml ✓ Zarontin GABAPENTIN Note: Not subsidised in combination with subsidised pregabalin * Cap 100 mg 6.45 100 ✓ Nupentin * Cap 300 mg 8.45 100 ✓ Nupentin * Cap 400 mg 10.26 100 ✓ Nupentin * Cap 400 mg 10.26 100 ✓ Nupentin * A Tab 50 mg 25.04 14 ✓ Vimpat A Tab 100 mg 50.06 14 ✓ Vimpat A Tab 150 mg 75.10 14 ✓ Vimpat A Tab 150 mg 75.10 14 ✓ Vimpat | 0 0 0 | | | • |
| Tab 10 mg | * Oral liq 20 mg per ml | 26.37 | 250 ml | ✓ Tegretol |
| CLONAZEPAM — Safety medicine; prescriber may determine dispensing frequency Oral drops 2.5 mg per ml | CLOBAZAM - Safety medicine; prescriber may determine | dispensing frequency | | |
| Oral drops 2.5 mg per ml | Tab 10 mg | 9.12 | 50 | ✓ Frisium |
| ETHOSUXIMIDE Cap 250 mg | CLONAZEPAM - Safety medicine; prescriber may determine | ine dispensing frequer | ncy | |
| ETHOSUXIMIDE Cap 250 mg | Oral drops 2.5 mg per ml | 7.38 | 10 ml OP | ✓ Rivotril |
| Tab 150 mg | | | | |
| Tab 150 mg | | 78.89 | 56 | ✓ Essential |
| Oral liq 250 mg per 5 ml .56.35 200 ml ✓ Zarontin GABAPENTIN | 3 | | | Ethosuximide \$29 |
| GABAPENTIN Note: Not subsidised in combination with subsidised pregabalin * Cap 100 mg | | 140.88 | 100 | ✓ Zarontin |
| Note: Not subsidised in combination with subsidised pregabalin * Cap 100 mg | Oral liq 250 mg per 5 ml | 56.35 | 200 ml | ✓ Zarontin |
| ** Cap 100 mg 6.45 100 ✓ Nupentin ** Cap 300 mg 8.45 100 ✓ Nupentin ** Cap 400 mg 10.26 100 ✓ Nupentin LACOSAMIDE – Special Authority see SA2267 on the next page – Retail pharmacy ▲ Tab 50 mg 25.04 14 ✓ Vimpat ▲ Tab 100 mg 50.06 14 ✓ Vimpat ▲ Tab 150 mg 75.10 14 ✓ Vimpat ▲ Tab 150 mg 300.40 56 ✓ Vimpat | GABAPENTIN | | | |
| ** Cap 300 mg 8.45 100 ✓ Nupentin ** Cap 400 mg 10.26 100 ✓ Nupentin LACOSAMIDE – Special Authority see SA2267 on the next page – Retail pharmacy ▲ Tab 50 mg 25.04 14 ✓ Vimpat ▲ Tab 100 mg 50.06 14 ✓ Vimpat ▲ 200.24 56 ✓ Vimpat ▲ Tab 150 mg 75.10 14 ✓ Vimpat 300.40 56 ✓ Vimpat | Note: Not subsidised in combination with subsidised p | regabalin | | |
| ** Cap 400 mg 10.26 100 Nupentin LACOSAMIDE − Special Authority see SA2267 on the next page − Retail pharmacy ▲ Tab 50 mg .25.04 14 ✓ Vimpat ▲ Tab 100 mg .50.06 14 ✓ Vimpat 200.24 56 ✓ Vimpat ▲ Tab 150 mg .75.10 14 ✓ Vimpat 300.40 56 ✓ Vimpat | * Cap 100 mg | 6.45 | 100 | ✓ Nupentin |
| LACOSAMIDE – Special Authority see SA2267 on the next page – Retail pharmacy ▲ Tab 50 mg | * Cap 300 mg | 8.45 | 100 | ✓ Nupentin |
| ▲ Tab 50 mg .25.04 14 ✓ Vimpat ▲ Tab 100 mg .50.06 14 ✓ Vimpat 200.24 56 ✓ Vimpat ▲ Tab 150 mg .75.10 14 ✓ Vimpat 300.40 56 ✓ Vimpat | * Cap 400 mg | 10.26 | 100 | ✓ Nupentin |
| ▲ Tab 100 mg .50.06 14 ✓ Vimpat 200.24 56 ✓ Vimpat ▲ Tab 150 mg .75.10 14 ✓ Vimpat 300.40 56 ✓ Vimpat | LACOSAMIDE - Special Authority see SA2267 on the nex | t page – Retail pharm | acy | |
| 200.24 56 ✓ Vimpat Tab 150 mg | ▲ Tab 50 mg | 25.04 | 14 | ✓ Vimpat |
| ▲ Tab 150 mg75.10 14 ✓ Vimpat 300.40 56 ✓ Vimpat | ▲ Tab 100 mg | 50.06 | 14 | ✓ Vimpat |
| 300.40 56 ✓ Vimpat | • | 200.24 | 56 | ✓ Vimpat |
| 300.40 56 ✓ Vimpat | ▲ Tab 150 mg | 75.10 | 14 | ✓ Vimpat |
| ▲ Tab 200 mg | - | 300.40 | 56 | |
| | ▲ Tab 200 mg | 400.55 | 56 | ✓ Vimpat |

| Subsidy | | Fully | Brand or |
|------------------------|-----|------------|--------------|
| (Manufacturer's Price) | | Subsidised | Generic |
| \$ | Per | ✓ | Manufacturer |

⇒SA2267 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Patient has focal epilepsy; and
- 2 Seizures are not adequately controlled by, or patient has experienced unacceptable side effects from, optimal treatment with all of the following: sodium valproate, topiramate, levetiracetam and any two of carbamazepine, lamotrigine and phenytoin sodium (see Note).

Note: Those of childbearing potential are not required to trial phenytoin sodium, sodium valproate, or topiramate. Those who can father children are not required to trial sodium valproate.

Renewal from any relevant practitioner. Approvals valid for 24 months where the patient has demonstrated a significant and sustained improvement in seizure rate or severity and/or quality of life compared with that prior to starting lacosamide treatment.

| LAN | MOTRIGINE | | | |
|------------------|--|-----------|---|---------------------|
| \blacktriangle | Tab dispersible 2 mg55.00 | 30 | 1 | Lamictal |
| \blacktriangle | Tab dispersible 5 mg50.00 | 30 | 1 | Lamictal |
| * | Tab dispersible 25 mg4.20 | | 1 | Logem |
| * | Tab dispersible 50 mg5.11 | | / | Logem |
| * | Tab dispersible 100 mg6.75 | 5 56 | 1 | Logem |
| LEV | 'ETIRACETAM | | | |
| | Tab 250 mg5.84 | 60 | 1 | Everet |
| | Tab 500 mg10.51 | | 1 | Everet |
| | Tab 750 mg16.71 | 60 | 1 | Everet |
| | Tab 1,000 mg21.82 | 2 60 | 1 | Everet |
| | Oral liq 100 mg per ml44.78 | 300 ml OP | 1 | Levetiracetam-AFT |
| | Inj 100 mg per ml, 5 ml vial | 5 10 | 1 | Levetiracetam-AFT |
| PHE | ENOBARBITONE | | | |
| | For phenobarbitone oral liquid refer Standard Formulae, page 272 | | | |
| | Tab 15 mg248.50 | 500 | 1 | Noumed |
| | · | | | Phenobarbitone |
| | Tab 30 mg | 500 | 1 | Noumed |
| | · · | | | Phenobarbitone |
| РН | ENYTOIN SODIUM | | | |
| | Tab 50 mg75.00 | 200 | / | Dilantin Infatab |
| • | Cap 30 mg | | | Dilantin |
| | Cap 100 mg | | | Dilantin |
| * | Oral liq 30 mg per 5 ml22.03 | | 1 | Dilantin Paediatric |
| | EGABALIN | | | |
| 1 111 | Note: Not subsidised in combination with subsidised gabapentin | | | |
| * | Cap 25 mg | 5 56 | / | Pregabalin Pfizer |
| ••• | 7.80 | | | Milpharm S29 |
| * | Cap 75 mg | | | Pregabalin Pfizer |
| 71 | 8.10 | | | Milpharm S29 |
| * | Cap 150 mg | | | Lyrica |
| * | Cap 150 mg4.01 | 30 | | Pregabalin Pfizer |
| * | Cap 300 mg | 3 56 | | Pregabalin Pfizer |
| | | , 30 | • | i regavann r nzer |
| | MIDONE | . 100 | | Duineldone Olineet |
| * | Tab 250 mg37.35 | 5 100 | • | Primidone Clinect |

NERVOUS SYSTEM

| | Subsidy | | Fully | Brand or |
|--|-----------------------|-------|------------|-------------------|
| | (Manufacturer's Price | e) | Subsidised | I Generic |
| | \$ | Per | • | Manufacturer |
| SODIUM VALPROATE | | | | |
| Tab 100 mg | 13.65 | 100 | ✓ | Epilim Crushable |
| Tab 200 mg EC | 27.44 | 100 | ✓ | Epilim |
| Tab 500 mg EC | 52.24 | 100 | ✓ | Epilim |
| * Oral liq 200 mg per 5 ml | 20.48 | 300 n | nl 🗸 | Epilim S/F Liquid |
| | | | ✓ | Epilim Syrup |
| * Inj 100 mg per ml, 4 ml | 41.50 | 1 | ✓ | Epilim IV |
| STIRIPENTOL - Special Authority see SA2268 below - Retail ph | narmacy | | | |
| Cap 250 mg | 509.29 | 60 | ✓ | Diacomit |
| Powder for oral liq 250 mg sachet | 509.29 | 60 | 1 | Diacomit |
| | | | | |

⇒SA2268 Special Authority for Subsidy

Initial application only from a paediatric neurologist or Practitioner on the recommendation of a paediatric neurologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient has confirmed diagnosis of Dravet syndrome; and
- 2 Seizures have been inadequately controlled by appropriate courses of sodium valproate, clobazam and at least two of the following: topiramate, levetiracetam, ketogenic diet.

Note: Those of childbearing potential are not required to trial sodium valproate or topiramate. Those who can father children are not required to trial sodium valproate.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where the patient continues to benefit from treatment as measured by reduced seizure frequency from baseline.

| . • | | |
|----------|-----------|--|
| A | Tab 25 mg | |
| | | |

TOPIRAMATE

| ▲ Tab 25 mg | 11.07 | 60 | ✓ Arrow-Topiramate |
|--|------------|-----|----------------------|
| - | | | ✓ Topiramate Actavis |
| | 26.04 | | ✓ Topamax |
| ▲ Tab 50 mg | 18.81 | 60 | ✓ Arrow-Topiramate |
| · · | | | ✓ Topiramate Actavis |
| | 44.26 | | ✓ Topamax |
| ▲ Tab 100 mg | 31.99 | 60 | Arrow-Topiramate |
| · · | | | ✓ Topiramate Actavis |
| | 75.25 | | ✓ Topamax |
| ▲ Tab 200 mg | 55.19 | 60 | Arrow-Topiramate |
| · · | | | ✓ Topiramate Actavis |
| | 129.85 | | ✓ Topamax |
| ▲ Sprinkle cap 15 mg | 20.84 | 60 | ✓ Topamax |
| ▲ Sprinkle cap 25 mg | | 60 | ✓ Topamax |
| VIGABATRIN - Special Authority see SA2088 below - Retain | l pharmacy | | |
| ▲ Tab 500 mg | | 100 | ✓ Sabril |
| Powder for oral soln 500 mg per sachet | | 60 | ✓ Sabril |
| | | | |

⇒SA2088 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Any of the following:
 - 1.1 Patient has infantile spasms; or
 - 1.2 Both:
 - 1.2.1 Patient has epilepsy; and
 - 1.2.2 Either:

| | | | NER | VOUS SYSTEM |
|--|--|--------------|--------------------|-------------------------------------|
| (A) | Subsidy flanufacturer's Price \$ | e) Su Per | Fully ubsidised | Brand or Generic Manufacturer |
| continued | | | | |
| 1.2.2.1 Seizures are not adequately controlled 1.2.2.2 Seizures are controlled adequately but 1 optimal treatment with other antiepileps 1.3 Patient has tuberous sclerosis complex; and 2 Either: | the patient has e | | | |
| Patient is, or will be, receiving regular automated visu 6-monthly basis thereafter); or It is impractical or impossible (due to comorbid condit | • | • | | |
| Renewal from any relevant practitioner. Approvals valid without fur following criteria: Both: | , | | | |
| The patient has demonstrated a significant and sustained im Either: | provement in sei | zure rate | or severit | y and or quality of life; and |
| 2.1 Patient is receiving regular automated visual field test of treatment with vigabatrin; or2.2 It is impractical or impossible (due to comorbid condit | | | • | |
| Antimigraine Preparations | | | | |
| For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page | 116 | | | |
| Acute Migraine Treatment | | | | |
| RIZATRIPTAN Tab orodispersible 10 mg | 4.84 | 30 | ✓ <u>F</u> | Rizamelt |
| SUMATRIPTAN Tab 50 mg Tab 100 mg | | 90 90 | _ | Sumagran Sumagran |
| Inj 12 mg per ml, 0.5 ml prefilled pen – Maximum of 10 inj per prescription | | 2 OP | √ <u>c</u> | Clustran |
| Prophylaxis of Migraine | | | | |
| For Beta Adrenoceptor Blockers refer to CARDIOVASCULAR SYST | EM, page 47 | | | |
| PIZOTIFEN * Tab 500 mcg | 23.21 | 100 | √ 9 | Sandomigran |
| Antinausea and Vertigo Agents | | | | |
| For Antispasmodics refer to ALIMENTARY TRACT, page 8 | | | | |
| APREPITANT – Special Authority see SA0987 below – Retail phan Cap 2 × 80 mg and 1 × 125 mg | | 3 OP | √ <u>E</u> | Emend Tri-Pack |
| ■ SA0987 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid for | or 12 months whe | ere the pa | tient is ur | ndergoing highly |

Renewal from any relevant practitioner. Approvals valid for 12 months where the patient is undergoing highly emetogenic

emetogenic chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.

chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.

BETAHISTINE DIHYDROCHLORIDE

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

NERVOUS SYSTEM

| | Subsidy (Manufacturer's Price) | Per | Fully Subsidised | Brand or Generic Manufacturer |
|--|-----------------------------------|-----|---------------------|--------------------------------------|
| CYCLIZINE HYDROCHLORIDE Tab 50 mg CYCLIZINE LACTATE | 0.66 | 10 | ✓ | <u>Nausicalm</u> |
| Inj 50 mg per ml, 1 ml ampoule - Up to 10 inj available on a PSO | | 10 | ✓ | <u>Hameln</u> |
| DOMPERIDONE * Tab 10 mg | 4.00 | 100 | • | <u>Domperidone</u> <u>Viatris</u> |
| HYOSCINE HYDROBROMIDE | | | | |
| * Inj 400 mcg per ml, 1 ml ampoule | 93.00 | 10 | • | Martindale S29 |
| below – Retail pharmacy | 88.50 | 10 | ✓ | Scopolamine - Mylan |

⇒SA1998 Special Authority for Subsidy

METOCI OPRAMIDE HYDROCHI ORIDE

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Control of intractable nausea, vomiting, or inability to swallow saliva in the treatment of malignancy or chronic disease where the patient cannot tolerate or does not adequately respond to oral anti-nausea agents; or
- 2 Control of clozapine-induced hypersalivation where trials of at least two other alternative treatments have proven ineffective.

Renewal from any relevant practitioner. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

| METOCLOPRAMIDE HYDROCHLORIDE | | |
|--|-----|--------------------------------|
| * Tab 10 mg - Up to 30 tab available on a PSO1.57 | 100 | ✓ Metoclopramide Actavis 10 |
| * Inj 5 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO7.00 | 10 | ✓ Baxter |
| ONDANSETRON | | |
| * Tab 4 mg2.27 | 50 | ✓ Periset |
| Tab disp 4 mg - Up to 10 tab available on a PSO | 10 | ✓ Periset ODT |
| * Tab 8 mg4.10 | 50 | ✓ Periset |
| Tab disp 8 mg - Up to 10 tab available on a PSO0.90 | 10 | ✓ Periset ODT |
| PROCHLORPERAZINE | | |
| * Tab 3 mg buccal5.97 | 50 | |
| (30.00) | | Buccastem |
| (30.00) | | Max Health \$29 |
| (30.00) | | Prochlorperazine |
| , | | Brown & Burk \$29 |
| * Tab 5 mg - Up to 30 tab available on a PSO25.00 | 250 | ✓ Nausafix |
| * Inj 12.5 mg per ml, 1 ml - Up to 5 inj available on a PSO25.81 | 10 | ✓ Stemetil |

Antipsychotics

General

| AMISULPRIDE - Safety medicine; prescriber may determine | dispensing frequency | | |
|---|----------------------|----|---------------------------|
| Tab 100 mg | 5.84 | 30 | Sulprix |
| Tab 200 mg | 14.47 | 60 | ✓ Sulprix |
| Tab 400 mg | 35.06 | 60 | ✓ Sulprix |

| | Subsidy (Manufacturer's Price) | Per | Fully Subsidised | I Generic |
|--|-----------------------------------|-------------|---------------------|-------------------------------|
| | \$ | rei | | Manuacturer |
| ARIPIPRAZOLE – Safety medicine; prescriber may determine of Tab 5 mg | | 30 | | Aripiprazole Sandoz Ascend |
| | | | | Aripiprazole S29 |
| Tab 10 mg | 10.50 | 30 | 1 | Aripiprazole Sandoz |
| Tab 15 mg | | 30 | | Aripiprazole Sandoz |
| Tab 20 mg | | 30 | | Aripiprazole Sandoz |
| Tab 30 mg | 10.50 | 30 | | Aripiprazole Sandoz |
| CHLORPROMAZINE HYDROCHLORIDE - Safety medicine; pr | escriber may determine | ne dis | spensing f | requency |
| Tab 25 mg - Up to 30 tab available on a PSO | | 100 | | Largactil |
| Tab 100 mg - Up to 30 tab available on a PSO | 36.73 | 100 | ✓ | Largactil |
| Inj 25 mg per ml, 2 ml - Up to 5 inj available on a PSO | | 10 | • | Largactil |
| CLOZAPINE – Hospital pharmacy [HP4] | IODO! | | | |
| Safety medicine; prescriber may determine dispensing frequency Tab 25 mg | • | 50 | _ | Clopine |
| Tab 25 mg | 0.09 | 50 | | Clozaril |
| | 13.37 | 100 | | Clopine |
| | 10.07 | | | Clozaril |
| Tab 50 mg | 8.67 | 50 | | Clopine |
| | 17.33 | 100 | | Clopine |
| Tab 100 mg | 17.33 | 50 | ✓ | Clopine |
| • | | | • | Clozaril |
| | 34.65 | 100 | • | Clopine |
| | | | | Clozaril |
| Tab 200 mg | | 50 | | Clopine |
| | 69.30 | 100 | | Clopine |
| Suspension 50 mg per ml | | 100 m | nl 🗸 | Versacloz |
| HALOPERIDOL - Safety medicine; prescriber may determine d | | | _ | |
| Tab 500 mcg - Up to 30 tab available on a PSO | | 100 | | Serenace |
| Tab 1.5 mg – Up to 30 tab available on a PSO | | 100 | | Serenace |
| Tab 5 mg - Up to 30 tab available on a PSO | | 50 | | Serenace |
| Oral lia 0 ma nor ml . Un to 000 ml available on a DCO | 29.72 | 100 | | Serenace |
| Oral liq 2 mg per ml - Up to 200 ml available on a PSO Inj 5 mg per ml, 1 ml ampoule - Up to 5 inj available on a P | | 100 m 10 | | Serenace Serenace |
| | | | | Serenace |
| LEVOMEPROMAZINE – Safety medicine; prescriber may deter | 1 0 1 | , | _ | |
| Tab 25 mg (33.8 mg as a maleate) | | 100 | | Nozinan (Swiss) |
| Tab 25 mg as a maleate | | 100 | | Nozinan |
| Tab 100 mg (135 mg as a maleate) | | 100 100 | | Nozinan (Swiss) Nozinan |
| Tab 100 mg as a maleate | | | | |
| LEVOMEPROMAZINE HYDROCHLORIDE - Safety medicine; | • | | | |
| Inj 25 mg per ml, 1 ml ampoule | 24.48 | 10 | | Nozinan S29 S29 |
| | | | • | <u>Wockhardt</u> |
| LITHIUM CARBONATE - Safety medicine; prescriber may dete | | | • | |
| Tab long-acting 400 mg | | 100 | | <u>Priadel</u> |
| Cap 250 mg | 22.36 | 100 | • | Douglas |
| | | | | |

NERVOUS SYSTEM

| | Subsidy | | Fully | Brand or |
|---|------------------------|---------|-------------|---------------------------|
| | (Manufacturer's Price) | Per | Subsidised | I Generic Manufacturer |
| | \$ | Per | | Manufacturer |
| DLANZAPINE – Safety medicine; prescriber may determine dis | | | | |
| Tab 2.5 mg | | 30 | | Zypine |
| Tab 5 mg | | 30 | | Zypine |
| Tab orodispersible 5 mg | | 28 | | Zypine ODT |
| Tab 10 mg | | 30 | | Zypine |
| Tab orodispersible 10 mg | 2.89 | 28 | • | Zypine ODT |
| PERICYAZINE - Safety medicine; prescriber may determine di | spensing frequency | | | |
| Tab 2.5 mg | 13.61 | 100 | ✓ | Neulactil |
| Tab 10 mg | 48.45 | 100 | ✓ | Neulactil |
| QUETIAPINE - Safety medicine; prescriber may determine disp | pensing frequency | | | |
| Tab 25 mg | 2.36 | 90 | ✓ | Quetapel |
| Tab 100 mg | | 90 | ✓ | Quetapel |
| Tab 200 mg | | 90 | ✓ | Quetapel |
| Tab 300 mg | | 90 | 1 | Quetapel |
| RISPERIDONE – Safety medicine; prescriber may determine d | | | | |
| Tab 0.5 mg | | 20 | 1 | Risperdal |
| 1 ab 0.0 mg | 2.17 | 60 | | Risperidone (Teva) |
| | 4.01 | 00 | | Risperidone |
| | 1.01 | | | Sandoz S29 |
| Tab 1 mg | 2.44 | 60 | 1 | Risperdal |
| rab ring | 2.44 | 00 | | Risperidone (Teva) |
| | 3.68 | | | Risperidone |
| | 3.00 | | • | • |
| Tale O area | 0.70 | 00 | , | Sandoz S29 |
| Tab 2 mg | 2.72 | 60 | | Risperdal |
| | 5.00 | | | Risperidone (Teva) |
| | 5.38 | | • | Risperidone |
| | | | _ | Sandoz S29 |
| Tab 3 mg | 4.50 | 60 | | Risperdal |
| | | | _ | Risperidone (Teva) |
| | 8.57 | | • | Risperidone |
| | | | | Sandoz S29 |
| Tab 4 mg | 6.25 | 60 | | Risperdal |
| | | | | Risperidone (Teva) |
| Oral liq 1 mg per ml | 10.29 | 30 m | · • | Risperon |
| IPRASIDONE - Safety medicine; prescriber may determine di | ispensing frequency | | | |
| Cap 20 mg | | 60 | 1 | Zusdone |
| Cap 40 mg | | 60 | | Zusdone |
| Cap 60 mg | | 60 | _ | Zusdone |
| Cap 80 mg | | 60 | / | Zusdone |
| UCLOPENTHIXOL HYDROCHLORIDE - Safety medicine; pro | | ne diei | nensina fra | equency |
| Tab 10 mg | | 100 | | Clopixol |

| Subsidy | | Fully | Brand or | |
|------------------------|-----|----------|--------------|--|
| (Manufacturer's Price) | Sul | bsidised | Generic | |
| \$ | Per | | Manufacturer | |

Depot Injections

| ARIPIPRAZOLE – Special Authority see SA2395 below – Retail pharmacy Safety medicine; prescriber may determine dispensing frequency | | |
|--|---|---|
| Inj 300 mg vial | 1 | ✓ Abilify Maintena✓ Abilify MaintenaS29 S29 |
| Inj 400 mg vial341.96 | 1 | ✓ Abilify Maintena ✓ Abilify Maintena S29 S29 |

⇒SA2395 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 Either:
 - 1.1 The patient has had an initial Special Authority approval for risperidone depot injection, paliperidone depot injection or olanzapine depot injection; or
 - 1.2 All of the following:
 - 1.2.1 The patient has schizophrenia or other psychotic disorder; and

ELLIDENTHIVOL DECANOATE Cofety medicine: procesiber may determine dispension fractions

- 1.2.2 The patient has received treatment with oral atypical antipsychotic agents but has been unable to adhere; and
- 1.2.3 The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months; or
- 2 Patient has been unable to access olanzapine depot injection due to supply issues with olanzapine depot injection, or otherwise would have been started on olanzapine depot injection but has been unable to due to supply issues with olanzapine depot injection.

Notes: The Olanzapine depot injection Special Authority criteria that apply to criterion 2 in this Aripiprazole Special Authority application are as follows:

- The patient has had an initial Special Authority approval for paliperidone depot injection or risperidone depot injection; or
- All of the following:
 - The patient has schizophrenia; and
 - The patient has not been able to adhere with treatment using oral atypical antipsychotic agents; and
 - The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months.

| FLUPENTHIXOL DECANOATE - Safety medicine; presci | iber may determine dispensing t | requency |
|---|----------------------------------|--------------------|
| Inj 20 mg per ml, 1 ml - Up to 5 inj available on a PS0 | D13.14 5 | ✓ Fluanxol |
| Inj 20 mg per ml, 2 ml - Up to 5 inj available on a PS0 | D20.90 5 | ✓ Fluanxol |
| Inj 100 mg per ml, 1 ml - Up to 5 inj available on a PS | SO40.87 5 | ✓ Fluanxol |
| HALOPERIDOL DECANOATE - Safety medicine; prescri | ber may determine dispensing fro | equency |
| Inj 50 mg per ml, 1 ml - Up to 5 inj available on a PS0 | D28.39 5 | ✓ Haldol |
| Inj 100 mg per ml, 1 ml - Up to 5 inj available on a PS | SO55.90 5 | Haldol Concentrate |
| | | ✓ Haldol |
| | | Decanoas \$29 |
| OLANZAPINE - Special Authority see SA2313 on the nex | t page - Retail pharmacy | |
| a) Safety medicine; prescriber may determine dispen- | sing frequency | |
| b) Note - no new patients to be initiated on olanzapin | e. | |
| Inj 210 mg vial | 252.00 1 | ✓ Zyprexa Relprevv |
| Inj 300 mg vial | 414.00 1 | ✓ Zyprexa Relprevv |
| Inj 405 mg vial | 504.00 1 | ✓ Zyprexa Relprevv |
| | | |

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.



| Subsidy | F | ully | Brand or | |
|------------------------|----------|------|--------------|--|
| (Manufacturer's Price) | Subsidis | sed | Generic | |
| \$ | Per | ✓ | Manufacturer | |

⇒SA2313 Special Authority for Subsidy

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of olanzapine depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

PALIPERIDONE - Special Authority see SA2396 below - Retail pharmacy

Safety medicine; prescriber may determine dispensing frequency

| Inj 25 mg syringe | 194.25 | 1 | ✓ Invega Sustenna |
|--------------------|--------|---|-------------------|
| Inj 50 mg syringe | 271.95 | 1 | ✓ Invega Sustenna |
| Inj 75 mg syringe | 357.42 | 1 | ✓ Invega Sustenna |
| Inj 100 mg syringe | 435.12 | 1 | ✓ Invega Sustenna |
| Inj 150 mg syringe | | 1 | ✓ Invega Sustenna |

⇒SA2396 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for risperidone depot injection or olanzapine depot injection or aripiprazole depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 Has been unable to adhere to treatment using oral atypical antipsychotic agents; and
 - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

PALIPERIDONE PALMITATE - Special Authority see SA2167 below - Retail pharmacy

| Inj 175 mg syringe | 815.85 | 1 | Invega Trinza |
|--------------------|----------|---|-----------------|
| | 1,072.26 | 1 | ✓ Invega Trinza |
| Inj 350 mg syringe | 1,305.36 | 1 | ✓ Invega Trinza |
| Inj 525 mg syringe | 1,305.36 | 1 | ✓ Invega Trinza |

⇒SA2167 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has schizophrenia; and
- 2 The patient has had an initial Special Authority approval for paliperidone once-monthly depot injection.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

RISPERIDONE - Special Authority see SA2397 below - Retail pharmacy

⇒SA2397 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Fither:

1 The patient has had an initial Special Authority approval for paliperidone depot injection or olanzapine depot injection or aripiprazole depot injection; or

| Subsidy | F | ully | Brand or |
|------------------------|---------|------|--------------|
| (Manufacturer's Price) | Subsidi | sed | Generic |
| \$ | Per | • | Manufacturer |

continued...

- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 Has not been able to adhere with treatment using oral atypical antipsychotic agents; and
 - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of risperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

ZUCLOPENTHIXOL DECANOATE - Safety medicine; prescriber may determine dispensing frequency
Inj 200 mg per ml, 1 ml - Up to 5 inj available on a PSO......19.80 5

✓ Clopixol

Anxiolytics

| BUS | SPIRONE HYDROCHLORIDE | | | |
|-----|--|-----------------|-----|---------------------------------------|
| * | Tab 5 mg | 13.95 | 100 | Buspirone Viatris |
| * | Tab 10 mg | 12.50 | 100 | ✓ Buspirone Viatris |
| CLC | DNAZEPAM - Safety medicine; prescriber may determine disper | nsing frequency | | |
| | Tab 500 mcg | 5.64 | 100 | ✓ Paxam |
| | Tab 2 mg | 10.78 | 100 | ✓ Paxam |
| DIA | ZEPAM - Safety medicine; prescriber may determine dispensing | g frequency | | |
| | Tab 2 mg | 95.00 | 500 | ✓ Arrow-Diazepam |
| | Tab 5 mg | 115.00 | 500 | ✓ Arrow-Diazepam |
| LOF | RAZEPAM - Safety medicine; prescriber may determine dispens | sing frequency | | |
| | Tab 1 mg | 10.20 | 250 | ✓ Ativan |
| | Tab 2.5 mg | 13.13 | 100 | ✓ Ativan |
| | | | | |

Multiple Sclerosis Treatments

⇒SA2274 Special Authority for Subsidy

Initial application — (Multiple Sclerosis - dimethyl fumarate, fingolimod, glatiramer acetate, interferon beta-1-alpha, interferon beta-1-beta, natalizumab and teriflunomide) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

1 All of the following:

- 1.1 Diagnosis of multiple sclerosis (MS) meets the McDonald 2017 diagnostic criteria for MS and has been confirmed by a neurologist; and
- 1.2 Patient has an EDSS score between 0 6.0; and
- 1.3 Patient has had at least one significant attack of MS in the previous 12 months or two significant attacks in the past 24 months; and
- 1.4 All of the following:
 - 1.4.1 Each significant attack must be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the attack, but the neurologist/physician must be satisfied that the clinical features were characteristic); and
 - 1.4.2 Each significant attack is associated with characteristic new symptom(s)/sign(s) or substantially worsening of previously experienced symptoms(s)/sign(s); and
 - 1.4.3 Each significant attack has lasted at least one week and has started at least one month after the onset of a



Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

previous attack (where relevant); and

- 1.4.4 Each significant attack can be distinguished from the effects of general fatigue; and is not associated with a fever (T> 37.5°C); and
- 1.4.5 Either:
 - 1.4.5.1 Each significant attack is severe enough to change either the EDSS or at least one of the Kurtze Functional System scores by at least 1 point; or
 - 1.4.5.2 Each significant attack is a recurrent paroxysmal symptom of multiple sclerosis (tonic seizures/spasms, trigeminal neuralgia, Lhermitte's symptom); and
- 1.5 Evidence of new inflammatory activity on an MRI scan within the past 24 months; and
- 1.6 Any of the following:
 - 1.6.1 A sign of that new inflammatory activity on MRI scanning (in criterion 5 immediately above) is a gadolinium enhancing lesion; or
 - 1.6.2 A sign of that new inflammatory activity is a lesion showing diffusion restriction; or
 - 1.6.3 A sign of that new inflammatory is a T2 lesion with associated local swelling; or
 - 1.6.4 A sign of that new inflammatory activity is a prominent T2 lesion that clearly is responsible for the clinical features of a recent attack that occurred within the last 2 years; or
 - 1.6.5 A sign of that new inflammatory activity is new T2 lesions compared with a previous MRI scan; or
- 2 Patient has an active approval for ocrelizumab and does not have primary progressive MS.

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

Renewal — (Multiple Sclerosis - dimethyl fumarate, fingolimod, glatiramer acetate, interferon beta-1-alpha, interferon beta-1-beta, natalizumab and teriflunomide) from any relevant practitioner. Approvals valid for 12 months where patient has had an EDSS score of 0 to 6.0 (inclusive) with or without the use of unilateral or bilateral aids at any time in the last six months (ie the patient has walked 100 metres or more with or without aids in the last six months).

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

DIMETHYL FUMARATE - Special Authority see SA2274 on the previous page - Retail pharmacy

- a) Wastage claimable
- b) Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

| Cap 120 mg | 520.00 | 14 | Tecfidera |
|------------|----------|----|-------------|
| Cap 240 mg | 2,000.00 | 56 | ✓ Tecfidera |

FINGOLIMOD - Special Authority see SA2274 on the previous page - Retail pharmacy

- a) Wastage claimable
- b) Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

GLATIRAMER ACETATE - Special Authority see SA2274 on the previous page - Retail pharmacy

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

INTERFERON BETA-1-ALPHA - Special Authority see SA2274 on the previous page - Retail pharmacy

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

Treation of the policy of the

INTERFERON BETA-1-BETA – Special Authority see SA2274 on the previous page – Retail pharmacy Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

NATALIZUMAB - Special Authority see SA2274 on the previous page - Retail pharmacy

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

NERVOUS SYSTEM

Aubagio

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

TERIFLUNOMIDE - Special Authority see SA2274 on page 139 - Retail pharmacy

a) Wastage claimable

b) Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

659.90

Teriflunomide Sandoz to be Principal Supply on 1 April 2025 (Aubagio Tab 14 mg to be delisted 1 April 2025)

Multiple Sclerosis Treatments - Other

OCRELIZUMAB - Special Authority see SA2273 below - Retail pharmacy

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

⇒SA2273 Special Authority for Subsidy

Initial application — (Multiple Sclerosis - ocrelizumab) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 Diagnosis of multiple sclerosis (MS) meets the McDonald 2017 diagnostic criteria for MS and has been confirmed by a neurologist; and
 - 1.2 Patient has an EDSS score between 0 6.0; and
 - 1.3 Patient has had at least one significant attack of MS in the previous 12 months or two significant attacks in the past 24 months; and
 - 1.4 All of the following:
 - 1.4.1 Each significant attack must be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the attack, but the neurologist/physician must be satisfied that the clinical features were characteristic); and
 - 1.4.2 Each significant attack is associated with characteristic new symptom(s)/sign(s) or substantially worsening of previously experienced symptoms(s)/sign(s); and
 - 1.4.3 Each significant attack has lasted at least one week and has started at least one month after the onset of a previous attack (where relevant); and
 - 1.4.4 Each significant attack can be distinguished from the effects of general fatigue; and is not associated with a fever (T> 37.5°C); and
 - 1.4.5 Fither:
 - 1.4.5.1 Each significant attack is severe enough to change either the EDSS or at least one of the Kurtze Functional System scores by at least 1 point; or
 - 1.4.5.2 Each significant attack is a recurrent paroxysmal symptom of multiple sclerosis (tonic seizures/spasms, trigeminal neuralgia, Lhermitte's symptom); and
 - 1.5 Evidence of new inflammatory activity on an MRI scan within the past 24 months; and
 - 1.6 Any of the following:
 - 1.6.1 A sign of that new inflammatory activity on MRI scanning (in criterion 5 immediately above) is a gadolinium enhancing lesion; or
 - 1.6.2 A sign of that new inflammatory activity is a lesion showing diffusion restriction; or
 - 1.6.3 A sign of that new inflammatory is a T2 lesion with associated local swelling; or
 - 1.6.4 A sign of that new inflammatory activity is a prominent T2 lesion that clearly is responsible for the clinical features of a recent attack that occurred within the last 2 years; or
 - 1.6.5 A sign of that new inflammatory activity is new T2 lesions compared with a previous MRI scan; or
- 2 Patient has an active Special Authority approval for either dimethyl fumarate, fingolimod, glatiramer acetate, interferon



Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

beta-1-alpha, interferon beta-1-beta, natalizumab or teriflunomide.

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

Renewal — (Multiple Sclerosis - ocrelizumab) from any relevant practitioner. Approvals valid for 12 months where patient has had an EDSS score of 0 to 6.0 (inclusive) with or without the use of unilateral or bilateral aids at any time in the last six months (ie the patient has walked 100 metres or more with or without aids in the last six months).

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

Initial application — (Primary Progressive Multiple Sclerosis) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Diagnosis of primary progressive multiple sclerosis (PPMS) meets the 2017 McDonald criteria and has been confirmed by a neurologist; and
- 2 Patient has an EDSS 2.0 (score equal to or greater than 2 on pyramidal functions) to EDSS 6.5; and
- 3 Patient has no history of relapsing remitting multiple sclerosis.

Renewal — (Primary Progressive Multiple Sclerosis) from any relevant practitioner. Approvals valid for 12 months where patient has had an EDSS score of less than or equal to 6.5 at any time in the last six months (ie patient has walked 20 metres with bilateral assistance/aids, without rest in the last six months).

Sedatives and Hypnotics

⇒SA1666 Special Authority for Subsidy

Initial application only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with persistent and distressing insomnia secondary to a neurodevelopmental disorder (including, but not limited to, autism spectrum disorder or attention deficit hyperactivity disorder)*; and
- 2 Behavioural and environmental approaches have been tried and were unsuccessful, or are inappropriate; and
- 3 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day; and
- 4 Patient is aged 18 years or under*.

Renewal only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is aged 18 years or under*; and
- 2 Patient has demonstrated clinically meaningful benefit from funded modified-release melatonin (clinician determined); and
- 3 Patient has had a trial of funded modified-release melatonin discontinuation within the past 12 months and has had a recurrence of persistent and distressing insomnia; and
- 4 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day.

Note: Indications marked with * are unapproved indications.

| | | | | VOOCOTOTEM |
|---|---|----------------|--------------------|--|
| | Subsidy (Manufacturer's Price) \$ | Sı Per | Fully ubsidised | Brand or Generic Manufacturer |
| MIDAZOLAM - Safety medicine; prescriber may determine dispe | nsing frequency | | | |
| Inj 1 mg per ml, 5 ml ampoule | 7.80 16.75 | 10 | | /lidazolam-Baxter /lidazolam Viatris |
| Inj 1 mg per ml, 5 ml plastic ampoule – Up to 10 inj available on a PSO | | 10 | | Pfizer |
| On a PSO for status epilepticus use only. PSO must be | endorsed for status e | epileptic | us use or | nly. |
| Inj 5 mg per ml, 1 ml plastic ampoule – Up to 10 inj available on a PSO | | 10 | - | /lidazolam-Pfizer |
| On a PSO for status epilepticus use only. PSO must be Inj 5 mg per ml, 3 ml ampoule | | epileptic 5 | ✓ I | ^{nly.} Midazolam-Baxter Midazolam Viatris |
| Inj 5 mg per ml, 3 ml plastic ampoule — Up to 5 inj available a PSO | 22.50 | 5 epileptic | | Pfizer |
| (Midazolam Viatris Inj 1 mg per ml, 5 ml ampoule to be delisted 1 (Midazolam Viatris Inj 5 mg per ml, 3 ml ampoule to be delisted 1 | May 2025) | | | ,. |
| PHENOBARBITONE SODIUM - Special Authority see SA1386 b | elow – Retail pharma | асу | | |
| Inj 200 mg per ml, 1 ml ampoule | 113.37 | 10 | ✓ I | Max Health S29 |
| ■ SA1386 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid the following criteria: Both: | without further rene | wal unle | ess notifie | d for applications meeting |
| 1 For the treatment of terminal agitation that is unresponsive2 The applicant is part of a multidisciplinary team working in | | l | | |
| TEMAZEPAM – Safety medicine; prescriber may determine dispertant 10 mg | . , | 25 | ✓ <u>I</u> | <u>lormison</u> |
| ZOPICLONE – Safety medicine; prescriber may determine disper Tab 7.5 mg | | 500 | √ <u>Z</u> | Copiclone Actavis |
| Spinal Muscular Atrophy | | | | |
| | | | | |

⇒SA2174 Special Authority for Subsidy

Initial application — (spinal muscular atrophy (SMA)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has genetic documentation of homozygous SMN1 gene deletion, homozygous SMN1 point mutation, or compound heterozygous mutation; and
- 2 Patient is 18 years of age or under; and
- 3 Either:
 - 3.1 Patient has experienced the defined signs and symptoms of SMA type I, II or IIIa prior to three years of age; or
 - 3.2 Both:
 - 3.2.1 Patient is pre-symptomatic; and
 - 3.2.2 Patient has three or less copies of SMN2.

Renewal — (spinal muscular atrophy (SMA)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:



| Subsidy | | Fully | Brand or |
|------------------------|-----|------------|--------------|
| (Manufacturer's Price) | | Subsidised | Generic |
| \$ | Per | ✓ | Manufacturer |

continued...

All of the following:

- 1 There has been demonstrated maintenance of motor milestone function since treatment initiation; and
- 2 Patient does not require invasive permanent ventilation (at least 16 hours per day) in the absence of a potentially reversible cause while being treated with nusinersen; and
- 3 Nusinersen not to be administered in combination other SMA disease modifying treatments or gene therapy.

RISDIPLAM - [Xpharm] - Special Authority see SA2203 below

Note: the supply of risdiplam is via Pharmac's approved direct distribution supply. Further details can be found on Pharmac's website https://pharmac.govt.nz/risdiplam

Powder for oral soln 750 mcg per ml, 60 mg per bottle......14,100.00 80 ml OP ✓ Evrysdi

⇒SA2203 Special Authority for Subsidy

Initial application — (spinal muscular atrophy (SMA)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has genetic documentation of homozygous SMN1 gene deletion, homozygous SMN1 point mutation, or compound heterozygous mutation; and
- 2 Patient is 18 years of age or under; and
- 3 Either:
 - 3.1 Patient has experienced the defined signs and symptoms of SMA type I, II or IIIa prior to three years of age; or
 - 3.2 Both:
 - 3.2.1 Patient is pre-symptomatic; and
 - 3.2.2 Patient has three or less copies of SMN2.

Renewal — (spinal muscular atrophy (SMA)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 There has been demonstrated maintenance of motor milestone function since treatment initiation; and
- 2 Patient does not require invasive permanent ventilation (at least 16 hours per day) in the absence of a potentially reversible cause while being treated with risdiplam; and
- 3 Risdiplam not to be administered in combination other SMA disease modifying treatments or gene therapy.

Stimulants/ADHD Treatments

| ATOMOXETINE | | | |
|---|-------|---------------|-------------------------------|
| Cap 10 mg | 43.02 | 28 | ✓ APO-Atomoxetine |
| Cap 18 mg | 45.57 | 28 | ✓ APO-Atomoxetine |
| Cap 25 mg | 44.30 | 28 | ✓ APO-Atomoxetine |
| Cap 40 mg | 46.21 | 28 | ✓ APO-Atomoxetine |
| Cap 60 mg | 51.31 | 28 | ✓ APO-Atomoxetine |
| Cap 80 mg | 65.20 | 28 | ✓ APO-Atomoxetine |
| Cap 100 mg | 65.71 | 28 | ✓ APO-Atomoxetine |
| DEXAMFETAMINE SULFATE – Special Authority see SA2410 a) Only on a controlled drug form b) Safety medicine; prescriber may determine dispensing to | , , | - Retail phar | macy |
| Tab 5 mg | | 100 | ✓ <u>Noumed</u> Dexamfetamine |

| Subsidy | | |
|------------------------|-----|------|
| (Manufacturer's Price) | | Subs |
| \$ | Per | |

Fully Subsidised

Brand or Generic Manufacturer

⇒SA2410 Special Authority for Subsidy

Initial application — (ADHD in patients aged 5 years or over) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid without further renewal unless notified for applications meeting the following criteria: All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) in patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients aged under 5 years) only from a paediatrician or psychiatrist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) in patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid without further renewal unless notified where the patient suffers from narcolepsy.

LISDEXAMFETAMINE DIMESILATE - Special Authority see SA2415 below - Retail pharmacy

- a) Only on a controlled drug form
- b) Safety medicine; prescriber may determine dispensing frequency

| Cap 30 mg - No more than 1 cap per day | 60.00 | 30 | Vyvanse |
|--|-------|----|-----------|
| Cap 50 mg | 60.00 | 30 | Vyvanse |
| Cap 70 mg | 60.00 | 30 | ✓ Vyvanse |

⇒SA2415 Special Authority for Subsidy

Initial application only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 Patient is currently on treatment with lisdexamfetamine dimesilate and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 ADHD (Attention Deficit and Hyperactivity Disorder); and
 - 2.2 Diagnosed according to DSM-V or ICD 11 criteria; and
 - 2.3 Either:
 - 2.3.1 Applicant is a paediatrician or psychiatrist; or
 - 2.3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing; and
 - 2.4 Any of the following:
 - 2.4.1 Patient is taking a currently subsidised formulation of atomoxetine or methylphenidate hydrochloride (extended-release) and has not received sufficient benefit or has experienced intolerable side effects; or
 - 2.4.2 Patient is taking a currently subsidised formulation of dexamfetamine sulfate (immediate-release) which has not been effective due to significant administration and/or treatment adherence difficulties; or
 - 2.4.3 There is significant concern regarding the risk of diversion or abuse of immediate release dexamfetamine sulfate; or
 - 2.4.4 Patient is taking a currently subsidised formulation of methylphenidate hydrochloride (immediate-release or sustained release) which has not been effective due to significant administration and/or treatment



| Subsidy | Fully | Brand or | |
|------------------------|------------|--------------|--|
| (Manufacturer's Price) | Subsidised | Generic | |
| \$ | Per 🗸 | Manufacturer | |

continued...

adherence difficulties: or

- 2.4.5 There is significant concern regarding the risk of diversion or abuse of immediate release methylphenidate hydrochloride; or
- 2.4.6 Both:
 - 2.4.6.1 Patient would have been prescribed a subsidised formulation of methylphenidate hydrochloride (extended-release) but has been unable to access due to supply issues with methylphenidate hydrochloride (extended-release); and
 - 2.4.6.2 Other alternative stimulant presentations (methylphenidate or dexamfetamine) are not appropriate;
- 2.5 Lisdexamfetamine dimesilate is not to be used in combination with another funded methylphenidate presentation.

METHYLPHENIDATE HYDROCHLORIDE - Special Authority see SA2411 below - Retail pharmacy

- a) Only on a controlled drug form
- b) Safety medicine; prescriber may determine dispensing frequency 30 ✓ Rubifen ✓ Rubifen Tab immediate-release 10 mg......3.00 30 ✓ Ritalin ✓ Methylphenidate ER 30 - Teva ✓ Rubifen 30 30 ✓ Rubifen SR 30 ✓ Methylphenidate ER - Teva ✓ Methylphenidate ER 30 - Teva Tab extended-release 54 mg......22.25 30 Methylphenidate ER

⇒SA2411 Special Authority for Subsidy

Initial application — (ADHD in patients aged 5 years or over) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid without further renewal unless notified for applications meeting the following criteria: All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) in patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Fither:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients aged under 5 years) only from a paediatrician or psychiatrist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) in patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy*) only from a neurologist or respiratory specialist. Approvals valid without further renewal unless notified where the patient suffers from narcolepsy.

Note: *narcolepsy is not a registered indication for Methylphenidate ER – Teva.

- Teva

NERVOUS SYSTEM

| Subsidy | | Fully | Brand or |
|------------------------|-----|------------|--------------|
| (Manufacturer's Price) | S | Subsidised | Generic |
| \$ | Per | 1 | Manufacturer |

METHYLPHENIDATE HYDROCHLORIDE EXTENDED-RELEASE - Special Authority see SA2412 below - Retail pharmacy

- a) Only on a controlled drug form
- b) Safety medicine: prescriber may determine dispensing frequency

| b) Salety medicine, prescriber may determine dispensing i | requericy | | |
|---|-----------|----|----------------------------|
| Tab extended-release 18 mg | 58.96 | 30 | Concerta |
| Tab extended-release 27 mg | 65.44 | 30 | Concerta |
| Tab extended-release 36 mg | 71.93 | 30 | Concerta |
| Tab extended-release 54 mg | 86.24 | 30 | Concerta |
| Cap modified-release 10 mg | | 30 | Ritalin LA |
| Cap modified-release 20 mg | | 30 | Ritalin LA |
| Cap modified-release 30 mg | | 30 | Ritalin LA |
| Cap modified-release 40 mg | | 30 | Ritalin LA |
| | | | |

⇒SA2412 Special Authority for Subsidy

Initial application — (ADHD) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder); and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Fither:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing; and
- 4 Either:
 - 4.1 Patient is taking a currently subsidised formulation of methylphenidate hydrochloride (immediate-release or sustained-release) which has not been effective due to significant administration and/or difficulties with adherence; or
 - 4.2 There is significant concern regarding the risk of diversion or abuse of immediate-release methylphenidate hydrochloride.

| MODAFINIL – Special Authority see SA2413 below – Retail pharma | acy | | |
|--|-------|----|---------------|
| Tab 100 mg | 14.27 | 30 | Modafinil Max |
| • | | | Health |
| | 29.13 | 60 | ✓ Modavigil |

Modafinil Max Health to be Principal Supply on 1 May 2025 (Modavigil Tab 100 mg to be delisted 1 May 2025)

⇒SA2413 Special Authority for Subsidy

Initial application only from a neurologist or respiratory specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of narcolepsy and has excessive daytime sleepiness associated with narcolepsy occurring almost daily for three months or more; and
- 2 Either:
 - 2.1 The patient has a multiple sleep latency test with a mean sleep latency of less than or equal to 10 minutes and 2 or more sleep onset rapid eye movement periods; or
 - 2.2 The patient has at least one of: cataplexy, sleep paralysis or hypnagogic hallucinations; and
- 3 Either:
 - 3.1 An effective dose of a subsidised formulation of methylphenidate or dexamfetamine has been trialled and discontinued because of intolerable side effects; or
 - 3.2 Methylphenidate and dexamfetamine are contraindicated.



| | (Manufacturer's Price) \$ | Per | Subsidised 🗸 | Generic Manufacturer |
|--|------------------------------|-----|--------------|------------------------------|
| Treatments for Dementia | | | | |
| DONEPEZIL HYDROCHLORIDE | | | | |
| * Tab 5 mg | 3.70 | 84 | ✓ | Ipca-Donepezil |
| * Tab 10 mg | 5.50 | 84 | ✓ | Ipca-Donepezil |
| RIVASTIGMINE - Special Authority see SA1488 below - Retail p | oharmacv | | | |
| Patch 4.6 mg per 24 hour | • | 30 | 1 | Rivastigmine Patch BNM 5 |
| | 90.00 | | ✓ | Exelon Patch 5 |
| Rivastigmine Patch BNM 5 to be Principal Supply on 1 M | larch 2025 | | | |
| Patch 9.5 mg per 24 hour | 49.40 | 30 | ✓ | Rivastigmine Patch BNM 10 |
| | 90.00 | | ✓ | Exelon Patch 10 |
| Rivastigmine Patch BNM 10 to be Principal Supply on 1 l | March 2025 | | | |
| (Exelon Patch 5 Patch 4.6 mg per 24 hour to be delisted 1 March | 2025) | | | |
| (Exelon Patch 10 Patch 9.5 mg per 24 hour to be delisted 1 March | h 2025) | | | |
| OA4400 Our alst Authority for Out alst | | | | |

Subsidy

Fully

Brand or

⇒SA1488 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The patient has been diagnosed with dementia; and
- 2 The patient has experienced intolerable nausea and/or vomiting from donepezil tablets.

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate; and
- 2 The patient has demonstrated a significant and sustained benefit from treatment.

Treatments for Substance Dependence

BUPRENORPHINE WITH NALOXONE - Special Authority see SA1203 below - Retail pharmacy

- a) No patient co-payment payable
- b) Safety medicine: prescriber may determine dispensing frequency

| Tab sublingual 2 mg with naloxone 0.5 mg | | 28 | ✓ Buprenorphine Naloxone BNM |
|--|-------|----|---------------------------------|
| Tab cublingual 9 mg with palayona 2 mg | 24.00 | 20 | ✓ Punyanayahina |

⇒SA1203 Special Authority for Subsidy

Initial application — (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health..

Initial application — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient will not be receiving methadone: and

| Sub | osidy Fu | ly Brand or |
|------------|-------------------------|--------------|
| (Manufactu | urer's Price) Subsidise | d Generic |
| | \$ Per | Manufacturer |

continued...

- 3 Patient is currently enrolled in an opioid substitution treatment program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient has previously trialled but failed detoxification with buprenorphine with naloxone with relapse back to opioid use and another attempt is planned; and
- 3 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is or has been receiving maintenance therapy with buprenorphine with naloxone (and is not receiving methadone); and
- 2 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health or is a medical practitioner authorised by the service to manage treatment in this patient.

Renewal — (Maintenance treatment where the patient has previously had an initial application for detoxification) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient received but failed detoxification with buprenorphine with naloxone; and
- 2 Maintenance therapy with buprenorphine with naloxone is planned (and patient will not be receiving methadone); and
- 3 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

| Tab modified-release 150 mg | 15.00 | 30 | ✓ <u>Zyban</u> |
|---|---------------------|-------------|----------------------|
| DISULFIRAM | | | |
| Tab 200 mg | 236.40 | 100 | ✓ Antabuse |
| NALTREXONE HYDROCHLORIDE - Special Authority see \$ | SA1408 below – Reta | il pharmacy | |
| Tab 50 mg | 77.77 | 28 | ✓ Naltrexone AOP S29 |
| | 83.33 | 30 | ✓ Naltraccord |
| | 102.60 | | ✓ Naltrexone Max |
| | | | Health S29 |
| | 138.88 | 50 | ✓ Revia S29 |

⇒SA1408 Special Authority for Subsidy

BLIBBODION HADDOCHI ODIDE

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient is currently enrolled in a recognised comprehensive treatment programme for alcohol dependence; and
- 2 Applicant works in or with a community Alcohol and Drug Service contracted to Health NZ or accredited against the New Zealand Alcohol and Other Drug Sector Standard or the National Mental Health Sector Standard.

Renewal from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

1 Compliance with the medication (prescriber determined); and



| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

continued...

- 2 Any of the following:
 - 2.1 Patient is still unstable and requires further treatment; or
 - 2.2 Patient achieved significant improvement but requires further treatment; or
 - 2.3 Patient is well controlled but requires maintenance therapy.

NICOTINE

- a) Nicotine will not be funded in amounts less than 4 weeks of treatment.
- b) Note: Direct Provision by a pharmacist permitted under the provisions in Part I of Section A. Patch 7 mg - Up to 28 patch available on a PSO19.62 ✓ Habitrol Patch 14 mg - Up to 28 patch available on a PSO21.57 28 ✓ Habitrol Patch 14 mg for direct distribution only - [Xpharm]......12.49 7 ✓ Habitrol Patch 21 mg - Up to 28 patch available on a PSO24.72 28 ✓ Habitrol Patch 21 mg for direct distribution only - [Xpharm]......13.19 7 ✓ Habitrol Lozenge 1 mg - Up to 216 loz available on a PSO......22.53 216 ✓ Habitrol 36 ✓ Habitrol ✓ Habitrol Lozenge 2 mg - Up to 216 loz available on a PSO......24.68 216 36 ✓ Habitrol Gum 2 mg (Fruit) - Up to 204 piece available on a PSO23.02 204 ✓ Habitrol Gum 2 mg (Fruit) for direct distribution only - [Xpharm].......17.57 96 ✓ Habitrol Gum 2 mg (Mint) - Up to 204 piece available on a PSO......23.02 204 ✓ Habitrol Gum 2 mg (Mint) for direct distribution only - [Xpharm]......17.57 96 ✓ Habitrol Gum 4 mg (Fruit) - Up to 204 piece available on a PSO25.98 204 ✓ Habitrol Gum 4 mg (Fruit) for direct distribution only - [Xpharm]......23.87 ✓ Habitrol 96 Gum 4 mg (Mint) – Up to 204 piece available on a PSO......25.98 204 ✓ Habitrol Gum 4 mg (Mint) for direct distribution only - [Xpharm]......23.87 96 ✓ Habitrol

VARENICLINE TARTRATE - Special Authority see SA1845 below - Retail pharmacy

- a) A maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval, including the starter pack
- b) Varenicline will not be funded in amounts less than 4 weeks of treatment.
- c) The 6-month time period in which a patient can receive a funded 12-week course of varenicline tartrate starts from the date the Special Authority is approved.

| Tab 0.5 mg × 11 and 1 mg × 42 | 16.67 | 53 OP | ✓ Varenicline Pfizer |
|-------------------------------|-------|-------|----------------------|
| Tab 1 mg | 17.62 | 56 | ✓ Varenicline Pfizer |

⇒SA1845 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 Either:
 - 3.1 The patient has tried but failed to quit smoking after at least two separate trials of nicotine replacement therapy, at least one of which included the patient receiving comprehensive advice on the optimal use of nicotine replacement therapy; or
 - 3.2 The patient has tried but failed to guit smoking using bupropion or nortriptyline; and
- 4 The patient has not had a Special Authority for varenicline approved in the last 6 months; and
- 5 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and

NERVOUS SYSTEM

| Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer | (Manufacturer's Price) Subsidised Generic |
|--|---|
|--|---|

continued...

- 6 The patient is not pregnant; and
- 7 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

Renewal from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 It has been 6 months since the patient's previous Special Authority was approved; and
- 4 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 5 The patient is not pregnant; and
- 6 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

The patient must not have had an approval in the past 6 months.

Notes: a maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval.

This includes the 4-week 'starter' pack.

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

Chemotherapeutic Agents

Alkylating Agents

| BENDAMUSTINE HYDROCHLORIDE - PCT only - Specialist - Special Authorit | y see SA2398 | |
|---|--------------|---|
| Inj 25 mg vial50.05 | 1 | Bendamustine Sandoz |
| 77.00 | | ✓ Ribomustin |
| Inj 100 mg vial200.20 | 1 | ✓ Bendamustine Sandoz |
| 308.00 | | ✓ Ribomustin |
| Inj 1 mg for ECP | 1 mg | ✓ Baxter |

⇒SA2398 Special Authority for Subsidy

Initial application — (CLL*) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has chronic lymphocytic leukaemia requiring treatment; and
- 2 Patient has ECOG performance status of 0-2; and
- 3 Bendamustine is to be administered at a maximum dose of 100 mg/m² on days 1 and 2 every 4 weeks for a maximum of 6 cycles.

Note: Indication marked with a * includes indications that are unapproved. 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL).

Initial application — (Indolent, Low-grade lymphomas) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has indolent low grade NHL requiring treatment; and
- 2 The patient has ECOG performance status of 0-2; and
- 3 Any of the following:
 - 3.1 Both:
 - 3.1.1 Patient is treatment naive; and
 - 3.1.2 Bendamustine is to be administered for a maximum of 6 cycles (in combination with rituximab when CD20+); or
 - 3.2 Both:
 - 3.2.1 Patient is refractory to or has relapsed within 12 months of a rituximab containing combined chemo-immunotherapy regimen; and
 - 3.2.2 Bendamustine is to be administered in combination with obinutuzumab for a maximum of 6 cycles; or
 - 3.3 All of the following:
 - 3.3.1 The patient has not received prior bendamustine therapy; and
 - 3.3.2 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
 - 3.3.3 Patient has had a rituximab treatment-free interval of 12 months or more; or
 - 3.4 Bendamustine is to be administered as monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Renewal — (Indolent, Low-grade lymphomas) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: Either:

- 1 Both:
 - 1.1 Patient is refractory to or has relapsed within 12 months of rituximab in combination with bendamustine; and

| | | - :: | | |
|------------------------|-----|-----------|--------------|--|
| Subsidy | | Fully | Brand or | |
| (Manufacturer's Price) | Sı | ubsidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

continued...

- 1.2 Bendamustine is to be administered in combination with obinutuzumab for a maximum of 6 cycles; or
- 2 Both
 - 2.1 Patients have not received a bendamustine regimen within the last 12 months; and
 - 2.2 Fither:
 - 2.2.1 Both:
 - 2.2.1.1 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
 - 2.2.1.2 Patient has had a rituximab treatment-free interval of 12 months or more; or
 - 2.2.2 Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Note: 'indolent, low-grade lymphomas' includes follicular, mantle cell, marginal zone and lymphoplasmacytic/ Waldenstrom's macroglobulinaemia.

Initial application — (**Hodgkin's lymphoma***) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has Hodgkin's lymphoma requiring treatment; and
- 2 Patient has a ECOG performance status of 0-2; and
- 3 Patient has received one prior line of chemotherapy; and
- 4 Patient's disease relapsed or was refractory following prior chemotherapy; and
- 5 Bendamustine is to be administered in combination with gemcitabine and vinorelbine (BeGeV) at a maximum dose of no greater than 90 mg/m2 twice per cycle, for a maximum of four cycles.

Note: Indications marked with * are unapproved indications.

| BUSULFAN – PCT – Retail pharmacy-Specialist Tab 2 mg | 89.25 | 100 | ✓ Myleran |
|--|--------|-----------|--|
| CARBOPLATIN - PCT only - Specialist | | | • |
| Inj 10 mg per ml, 45 ml vial | 25.73 | 1 | ✓ Carboplatin Accord ✓ DBL Carboplatin S29 S29 |
| | 32.59 | | ✓ DBL Carboplatin |
| | 48.50 | | ✓ Carbaccord |
| Inj 1 mg for ECP | 0.06 | 1 mg | ✓ Baxter |
| CARMUSTINE - PCT only - Specialist | | | |
| Inj 100 mg vial | 710.00 | 1 | ✓ BiCNU |
| , , | | | ✓ BiCNU S29 S29 |
| | | | ✓ Novadoz S29 |
| Inj 100 mg for ECP | 710.00 | 100 mg OP | ✓ Baxter |
| CHLORAMBUCIL – PCT – Retail pharmacy-Specialist | | • | |
| Tab 2 mg | 29.06 | 25 | ✓ Leukeran FC |
| CISPLATIN - PCT only - Specialist | | | |
| Inj 1 mg per ml, 50 ml vial | 9.45 | 1 | Cisplatin Accord |
| , -, | 15.00 | | ✓ Cisplatin Ebewe |
| Inj 1 mg per ml, 100 ml vial | 18.90 | 1 | ✓ Cisplatin Accord |
| | 21.00 | | Cisplatin Ebewe |
| | 29.66 | | DBL Cisplatin |
| Ini 1 ma for ECP | 0.19 | 1 ma | ✓ Baxter |

| | Subsidy | Fu | | |
|---|-----------------------------|--------------|-------------|---------------------|
| (N | lanufacturer's Price) \$ | Subsi Per | idised • | |
| YCLOPHOSPHAMIDE | - | | | |
| Tab 50 mg - PCT - Retail pharmacy-Specialist | 145.00 | 50 | 1 | Cyclonex |
| Inj 1 g vial – PCT – Retail pharmacy-Specialist | | 1 | | Endoxan |
| , · g · · · · · · · · · · · · · · · · · | 127.80 | 6 | | Cytoxan |
| Inj 2 g vial - PCT only - Specialist | 95.06 | 1 | | Endoxan |
| Inj 1 mg for ECP - PCT only - Specialist | | 1 mg | 1 | Baxter |
| FOSFAMIDE - PCT only - Specialist | | • | | |
| Inj 1 g | 96.00 | 1 | / | Holoxan |
| Inj 2 g | | 1 | | Holoxan |
| Inj 1 mg for ECP | | 1 mg | | Baxter |
| OMUSTINE - PCT - Retail pharmacy-Specialist | | 9 | | |
| | 990 00 | 20 | ./ | Medac S29 |
| Cap 40 mg | 000.00 | 20 | • | Wedac 329 |
| MELPHALAN | | | | |
| Tab 2 mg - PCT - Retail pharmacy-Specialist | | 25 | | Alkeran |
| Inj 50 mg - PCT only - Specialist | 48.25 | 1 | | Megval S29 |
| | | | | Melpha |
| | 67.80 | | • | Alkeran |
| DXALIPLATIN - PCT only - Specialist | | | | |
| Inj 100 mg vial | 25.01 | 1 | / | Oxaliplatin Actavis |
| | | | | 100 |
| | 110.00 | | | Oxaliplatin Ebewe |
| Inj 5 mg per ml, 20 ml vial | 33.35 | 1 | | Alchemy Oxaliplatin |
| | 46.32 | | | Oxaliplatin Accord |
| Inj 1 mg for ECP | 0.35 | 1 mg | / | Baxter |
| HIOTEPA - PCT only - Specialist | | | | |
| Inj 15 mg vial | CBS | 1 | 1 | Bedford S29 |
| . • | | | 1 | Max Health \$29 |
| | | | / | THIO-TEPA \$29 |
| | 398.00 | | | Tepadina |
| Inj 100 mg vial | | 1 | | Max Health S29 |
| iiij 100 iiig va | 1,800.00 | • | | Tepadina |
| | 1,000.00 | | | Topuumu |
| Antimetabolites | | | | |
| AZACITIDINE - PCT only - Specialist - Special Authority see SA2 | 141 below | | | |
| Inj 100 mg vial | | 1 | • | Azacitidine Dr |
| In: 4 may few FOD | 0.54 | 4 | , | Reddy's |
| Inj 1 mg for ECP | 0.54 | 1 mg | • | Baxter |

⇒SA2141 Special Authority for Subsidy

Initial application only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has International Prognostic Scoring System (IPSS) intermediate-2 or high risk myelodysplastic syndrome; or
 - 1.2 The patient has chronic myelomonocytic leukaemia (10%-29% marrow blasts without myeloproliferative disorder); or

| Subsidy | | Fully | Brand or | |
|------------------------|-----|---------|--------------|--|
| (Manufacturer's Price) | Sub | sidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

continued...

- 1.3 The patient has acute myeloid leukaemia with 20-30% blasts and multi-lineage dysplasia, according to World Health Organisation Classification (WHO); and
- 2 The patient has performance status (WHO/ECOG) grade 0-2; and
- 3 The patient has an estimated life expectancy of at least 3 months.

Renewal only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

| CALCIUM FOLINATE | | |
|--|----------|---|
| Tab 15 mg - PCT - Retail pharmacy-Specialist | 10 | ✓ DBL Leucovorin Calcium |
| Inj 3 mg per ml, 1 ml - PCT - Retail pharmacy-Specialist | 5 | ✓ Hospira |
| Inj 10 mg per ml, 5 ml vial - PCT - Retail pharmacy-Specialist7.28 | 1 | ✓ Calcium Folinate Sandoz |
| | | ✓ Calcium Folinate Sandoz S29 S29 |
| 36.48 | 5 | ✓ Eurofolic S29 |
| Inj 50 mg - PCT - Retail pharmacy-Specialist72.80 | 10 | ✓ Leucovorin Pharmacia S29 |
| Inj 10 mg per ml, 10 ml vial - PCT only - Specialist9.49 | 1 | Calcium Folinate Sandoz |
| 47.45 | 5 | ✓ Eurofolic S29 |
| Inj 100 mg - PCT only - Specialist7.33 | 1 | Calcium Folinate Ebewe |
| 94.90 | 10 | ✓ Leucovorin Pharmacia S29 |
| Inj 300 mg - PCT only - Specialist21.55 | 1 | ✓ Leucovorin DBL S29 |
| 22.51 | | ✓ Calcium Folinate Ebewe |
| Inj 10 mg per ml, 35 ml vial - PCT only - Specialist25.14 | 1 | ✓ Calcium Folinate Sandoz |
| | | ✓ Calcium Folinate Sandoz S29 S29 |
| Inj 1 g - PCT only - Specialist67.51 | 1 | Calcium Folinate Ebewe |
| Inj 10 mg per ml, 100 ml vial - PCT only - Specialist72.00 | 1 | ✓ Calcium Folinate Sandoz |
| | | ✓ Eurofolic S29 |
| Inj 1 mg for ECP - PCT only - Specialist | 1 mg | ✓ Baxter |
| CAPECITABINE - Retail pharmacy-Specialist | | |
| Tab 150 mg9.80 | 60 | ✓ Capecitabine Viatris |
| Tab 500 mg46.50 | 120 | ✓ Capecitabine Viatris |
| CLADRIBINE - PCT only - Specialist | | |
| Inj 1 mg per ml, 10 ml749.96 | 1 | ✓ Leustatin |
| Inj 10 mg for ECP749.96 | 10 mg OP | ✓ Baxter |

| | Subsidy | | Fully Brand or |
|--|-------------------|--------------------|--------------------------------------|
| | (Manufacturer's F | Price) Subs Per | idised Generic Manufacturer |
| CYTARABINE | · · | | |
| Inj 20 mg per ml, 5 ml vial – PCT – Retail pharmacy-Specia | list472.00 | 5 | ✓ Pfizer |
| Inj 100 mg per ml, 20 ml vial – PCT – Retail | | · · | |
| pharmacy-Specialist | 48.80 | 1 | ✓ Cytarabine DBL |
| priarriacy openialist | | ' | ✓ Pfizer |
| | | | ✓ Pfizer S29 S29 |
| Inj 1 mg for ECP - PCT only - Specialist | 0.20 | 10 mg | ✓ Baxter |
| Inj 100 mg intrathecal syringe for ECP - PCT only - Specia | liet 0/ /0 | 100 mg OP | ✓ Baxter |
| | 113134.40 | 100 mg Oi | Daxiei |
| FLUDARABINE PHOSPHATE | | | |
| Tab 10 mg – PCT – Retail pharmacy-Specialist | | 20 | ✓ Fludara Oral |
| Inj 50 mg vial - PCT only - Specialist | 126.80 | 1 | ✓ Fludarabine Sagent \$29 |
| | 634.00 | 5 | ✓ Fludarabine Ebewe |
| Inj 50 mg for ECP - PCT only - Specialist | 126.80 | 50 mg OP | ✓ Baxter |
| LUOROURACIL | | Ü | |
| Inj 50 mg per ml, 20 ml vial – PCT only – Specialist | 10.51 | 1 | ✓ Fluorouracil Accor |
| Inj 50 mg per ml, 50 ml vial – PCT only – Specialist | | 1 | ✓ Fluorouracii Accor |
| Inj 50 mg per ml, 30 ml vial – PCT only – Specialist | | 1 | ✓ Fluorouracii Accor |
| Inj 1 mg for ECP – PCT only – Specialist | | 100 mg | ✓ Baxter |
| , , | 0.41 | 100 mg | Daxiei |
| SEMCITABINE HYDROCHLORIDE - PCT only - Specialist | | | |
| Inj 43.3 mg per ml (equivalent to 38 mg per ml gemcitabine). | | | |
| 26.3 ml vial | | 1 | ✓ DBL Gemcitabine |
| Inj 1 g | | 1 | Gemcitabine Ebew |
| Inj 1 mg for ECP | 0.02 | 1 mg | ✓ Baxter |
| RINOTECAN HYDROCHLORIDE - PCT only - Specialist | | | |
| Inj 20 mg per ml, 5 ml vial | 52.57 | 1 | ✓ Accord |
| , , , | 71.44 | | ✓ Irinotecan Actavis |
| | | | 100 |
| | 100.00 | | ✓ Irinotecan-Rex |
| Inj 1 mg for ECP | | 1 mg | ✓ Baxter |
| MERCAPTOPURINE | | 9 | |
| | 25.00 | 25 | ✓ Puri-nethol |
| Tab 50 mg - PCT - Retail pharmacy-Specialist | | 25 | ▼ Puri-nethol |
| Oral suspension 20 mg per ml – Retail pharmacy-Specialist | | 100 105 | 4 A11 |
| Special Authority see SA1725 below | 428.00 | 100 ml OP | ✓ Allmercap |
| | | | ✓ Xaluprine S29 |

⇒SA1725 Special Authority for Subsidy

Initial application only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where the patient requires a total dose of less than one full 50 mg tablet per day.

Renewal only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where patient still requires a total dose of less than one full 50 mg tablet per day.

| METHOTREXATE | _ | | 0.1.11 | | | | |
|---|--|---|--|----------|-----|--------------------|--|
| ### Tab 2.5 mg - PCT - Retail pharmacy-Specialist | | | | ۵) | , | | |
| ### Tab 2.5 mg — PCT — Retail pharmacy-Specialist | | | | | | | |
| # Tab 2.5 mg — PCT — Retail pharmacy-Specialist | | THOTDEVATE | · · | | | | |
| # Tab 10 mg − PCT − Retail pharmacy-Specialist | | | 7.00 | 00 | | Tuessate | |
| # Inj 2.5 mg per ml, 2 ml − PCT − Retail pharmacy-Specialist | | , , , | | | | | |
| # Inj 7.5 mg prefilled syringe | | | | | | | |
| ★ Inj 7.5 mg prefilled syringe | 不 | inj 2.5 mg per mi, 2 mi – PCT – netali pharmacy-Specialist | 30.03 | 5 | | | |
| # Inj 7.5 mg prefilled syringe | | | | | • | | |
| ★ Inj 10 mg prefilled syringe | | Ini 7 F and an City of a series of | 00.47 | | , | | |
| # Inj 10 mg prefilled syringe | * | inj 7.5 mg pretilled syringe | 29.17 | ı | • | | |
| X Inj 15 mg prefilled syringe | | let 40 mm and filled and an | 40.00 | | , | | |
| ★ Inj 15 mg prefilled syringe | * | Inj 10 mg pretilled syringe | 19.09 | 1 | • | | |
| ★ Inj 20 mg prefilled syringe. 16.64 1 ✓ Methotrexate Sandoz ★ Inj 25 mg prefilled syringe. 20.72 1 ✓ Methotrexate Sandoz ★ Inj 30 mg prefilled syringe. 55.00 1 ✓ Methotrexate Sandoz ★ Inj 25 mg per ml, 2 ml vial – PCT – Retail pharmacy-Specialist. 30.00 5 ✓ Methotrexate DBL Onco-Vial ★ Inj 25 mg per ml, 20 ml vial – PCT – Retail pharmacy-Specialist. 45.00 1 ✓ DBL Methotrexate DBL Onco-Vial ★ Inj 100 mg per ml, 10 ml – PCT – Retail pharmacy-Specialist. 45.00 1 ✓ BMETHOTEXALE DBL Onco-Vial ★ Inj 100 mg per ml, 50 ml vial – PCT – Retail pharmacy-Specialist. 25.00 1 ✓ Methotrexate Ebewe ★ Inj 1 mg for ECP – PCT only – Specialist. 67.99 1 ✓ Methotrexate Ebewe ★ Inj 1 mg for ECP – PCT only – Specialist. 4.73 5 mg OP ✓ Baxter PEMETREXED – PCT only – Specialist. 4.73 5 mg OP ✓ Baxter PEMETREXED – PCT only – Specialist. 8.99 1 ✓ Pemetrexed-AFT Inj 10 mg vial 2.99 1 ✓ Pemetrexed-AFT Juno Pemetrexed 2.00 Methotrexate Ebewe | | 1.45 | 0.4.50 | | , | | |
| # Inj 20 mg prefilled syringe | * | Inj 15 mg pretilled syringe | 24.53 | 1 | • | | |
| ★ Inj 25 mg prefilled syringe | | | | | _ | | |
| ★ Inj 30 mg prefilled syringe | * | Inj 20 mg prefilled syringe | 16.64 | 1 | • | | |
| * Inj 30 mg prefilled syringe | | | | | _ | | |
| ★ Inj 30 mg prefilled syringe. .55.00 1 ★ Methotrexate Sandoz ★ Inj 25 mg per ml, 2 ml vial − PCT − Retail pharmacy-Specialist30.00 5 ★ Methotrexate DBL Onco-Vial ★ Inj 100 mg per ml, 20 ml vial − PCT − Retail pharmacy-Specialist | * | Inj 25 mg prefilled syringe | 20.72 | 1 | • | | |
| # Inj 25 mg per ml, 2 ml vial — PCT — Retail pharmacy-Specialist | | | | | _ | | |
| * Inj 25 mg per ml, 2 ml vial — PCT — Retail pharmacy-Specialist30.00 5 Methotrexate DBL Onco-Vial * Inj 25 mg per ml, 20 ml vial — PCT — Retail pharmacy-Specialist45.00 1 ✓ DBL Methotrexate Onco-Vial * Inj 100 mg per ml, 10 ml — PCT — Retail pharmacy-Specialist | * | Inj 30 mg prefilled syringe | 55.00 | 1 | • | | |
| Nonco-Vial | | | | | _ | | |
| ★ Inj 25 mg per ml, 20 ml vial - PCT - Retail pharmacy-Specialist | * | Inj 25 mg per ml, 2 ml vial - PCT - Retail pharmacy-Specialis | st30.00 | 5 | / | | |
| Methotrexate Ebewe | | | | | | | |
| ★ Inj 100 mg per ml, 10 ml − PCT − Retail pharmacy-Specialist .25.00 1 ✓ Methotrexate Ebewe ★ Inj 100 mg per ml, 50 ml vial − PCT − Retail pharmacy-Specialist .67.99 1 ✓ Methotrexate Ebewe ★ Inj 1 mg for ECP − PCT only − Specialist .0.06 1 mg ✓ Baxter ★ Inj 5 mg intrathecal syringe for ECP − PCT only − Specialist .4.73 5 mg OP ✓ Baxter PEMETREXED − PCT only − Specialist .8.99 1 ✓ Pemetrexed-AFT Inj 100 mg vial .8.99 1 ✓ Pemetrexed-AFT Juno Pemetrexed .9.99 1 ✓ Pemetrexed-AFT Juni 1 mg for ECP .0.11 1 mg ✓ Baxter THIOGUANINE − PCT − Retail pharmacy-Specialist Tab 40 mg .126.31 .25 ✓ Lanvis Other Cytotoxic Agents AMSACRINE − PCT only − Specialist Inj 50 mg per ml, 1.5 ml ampoule .1,500.00 6 ✓ Amsidine \$29 Inj 75 mg .1,250.00 5 ✓ AmsaLyo \$29 ANAGRELIDE HYDROCHLORIDE − PCT − Retail pharmacy-Specialist .1,175.87 100 ✓ Agrylin <td c<="" td=""><td>*</td><td>Inj 25 mg per ml, 20 ml vial - PCT - Retail pharmacy-Special</td><td>ist45.00</td><td>1</td><td>•</td><td></td></td> | <td>*</td> <td>Inj 25 mg per ml, 20 ml vial - PCT - Retail pharmacy-Special</td> <td>ist45.00</td> <td>1</td> <td>•</td> <td></td> | * | Inj 25 mg per ml, 20 ml vial - PCT - Retail pharmacy-Special | ist45.00 | 1 | • | |
| ★ Inj 100 mg per ml, 50 ml vial − PCT − Retail pharmacy-Specialist 67.99 1 ✓ Methotrexate Ebewe ★ Inj 1 mg for ECP − PCT only − Specialist 0.06 1 mg ✓ Baxter ★ Inj 5 mg intrathecal syringe for ECP − PCT only − Specialist 5 mg OP ✓ Baxter PEMETREXED − PCT only − Specialist 4.73 5 mg OP ✓ Baxter PEMETREXED − PCT only − Specialist 8.99 1 ✓ Pemetrexed-AFT Inj 100 mg vial 8.99 1 ✓ Pemetrexed-AFT Juno Pemetrexed 9.99 1 ✓ Pemetrexed-AFT Juny 1 mg for ECP 0.11 1 mg ✓ Baxter THIOGUANINE − PCT − Retail pharmacy-Specialist 126.31 25 ✓ Lanvis Other Cytotoxic Agents AMSACRINE − PCT only − Specialist Inj 50 mg per ml, 1.5 ml ampoule 1,500.00 6 ✓ Amsidine \$29 Inj 75 mg 1,250.00 5 ✓ AmsaLyo \$29 ANAGRELIDE HYDROCHLORIDE − PCT − Retail pharmacy-Specialist 0,218.00 ✓ Agrylin ARSENIC TRIOXIDE − PCT only − Specialist 1,175.87 100 ✓ Phenasen | | | | | | Onco-Vial | |
| Pharmacy-Specialist Fig. Fig. Post of the state Post of | * | Inj 100 mg per ml, 10 ml - PCT - Retail pharmacy-Specialist | 25.00 | 1 | • | Methotrexate Ebewe | |
| ★ Inj 1 mg for ECP - PCT only - Specialist .0.06 1 mg ✔ Baxter ★ Inj 5 mg intrathecal syringe for ECP - PCT only - Specialist .4.73 5 mg OP ✔ Baxter PEMETREXED - PCT only - Specialist .8.99 1 ✔ Pemetrexed-AFT Inj 100 mg vial .8.99 1 ✔ Pemetrexed-AFT Juno Pemetrexed .29.99 1 ✔ Pemetrexed-AFT ✓ Juno Pemetrexed .0.11 1 mg ✔ Baxter THIOGUANINE - PCT - Retail pharmacy-Specialist .0.11 1 mg ✔ Baxter THIOGUANINE - PCT only - Specialist Inj 50 mg per ml, 1.5 ml ampoule .1,500.00 6 ✔ Amsidine 529 Inj 50 mg per ml, 1.5 ml ampoule .1,250.00 5 ✔ AmsaLyo 529 Inj 75 mg .1,250.00 5 ✔ AmsaLyo 529 ANAGRELIDE HYDROCHLORIDE - PCT - Retail pharmacy-Specialist .1,175.87 100 ✔ Agrylin ARSENIC TRIOXIDE - PCT only - Specialist .1,175.87 100 ✔ Phenasen | * | , 01 | | | | | |
| ★ Inj 5 mg intrathecal syringe for ECP − PCT only − Specialist | | | | | | | |
| PEMETREXED − PCT only − Specialist Inj 100 mg vial | | | | Ū | _ | | |
| Inj 100 mg vial | * | Inj 5 mg intrathecal syringe for ECP – PCT only – Specialist | 4.73 | 5 mg O | P 🗸 | Baxter | |
| Solution Solution | PE | METREXED - PCT only - Specialist | | | | | |
| Inj 500 mg vial | | Inj 100 mg vial | 8.99 | 1 | | | |
| 217.77 | | | | | | | |
| Inj 1 mg for ECP | | Inj 500 mg vial | 29.99 | 1 | | | |
| THIOGUANINE − PCT − Retail pharmacy-Specialist | | | | | | | |
| Tab 40 mg | | Inj 1 mg for ECP | 0.11 | 1 mg | / | Baxter | |
| Other Cytotoxic Agents AMSACRINE – PCT only – Specialist | THI | OGUANINE - PCT - Retail pharmacy-Specialist | | | | | |
| AMSACRINE – PCT only – Specialist Inj 50 mg per ml, 1.5 ml ampoule | | Tab 40 mg | 126.31 | 25 | ✓ | Lanvis | |
| AMSACRINE – PCT only – Specialist Inj 50 mg per ml, 1.5 ml ampoule | _ | | | | | | |
| Inj 50 mg per ml, 1.5 ml ampoule | 0 | ther Cytotoxic Agents | | | | | |
| Inj 50 mg per ml, 1.5 ml ampoule | Δ Ι. Λ | SACRINE - PCT only - Specialist | | | | | |
| 4,736.00 | \(\tau\) | | 1 500 00 | c | | Ameiding con | |
| Inj 75 mg 1,250.00 5 ✓ AmsaLyo \$29 6,218.00 ✓ AmsaLyo \$29 ANAGRELIDE HYDROCHLORIDE – PCT – Retail pharmacy-Specialist 1,175.87 100 ✓ Agrylin ARSENIC TRIOXIDE – PCT only – Specialist 4,817.00 10 ✓ Phenasen | | inj 50 mg per mi, 1.5 mi ampoule | | Ö | | | |
| 6,218.00 | | 1.5 | | _ | | | |
| ANAGRELIDE HYDROCHLORIDE – PCT – Retail pharmacy-Specialist Cap 0.5 mg | | Inj /5 mg | | 5 | | • | |
| Cap 0.5 mg | | | 6,218.00 | | / | AmsaLyo S29 | |
| ARSENIC TRIOXIDE − PCT only − Specialist Inj 1 mg per ml, 10 ml vial4,817.00 10 ✓ Phenasen | AN | AGRELIDE HYDROCHLORIDE - PCT - Retail pharmacy-Spe | ecialist | | | | |
| Inj 1 mg per ml, 10 ml vial | | Cap 0.5 mg | 1,175.87 | 100 | 1 | Agrylin | |
| Inj 1 mg per ml, 10 ml vial | AR | SENIC TRIOXIDE - PCT only - Specialist | | | | | |
| | | | 4,817.00 | 10 | 1 | Phenasen | |
| | | , , , | | | | | |
| | | • • | | 0 | | | |

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

| | Subsidy | , | Fully | |
|--|---------------------------|----------------|--------------|--------------------------|
| | (Manufacturer's Pri \$ | ce) Sub Per | sidised • | Generic Manufacturer |
| FOLIVOIN OUR BUATE DOT L. O | Ψ | FEI | | Manuacturei |
| LEOMYCIN SULPHATE – PCT only – Specialist | 105.10 | | | DDI Disamusia |
| Inj 15,000 iu, vial | 185.16 | 1 | • | DBL Bleomycin Sulfate |
| Inj 1,000 iu for ECP | | 1,000 iu | ✓ | Baxter |
| DRTEZOMIB - PCT only - Specialist - Special Authority see | SA2355 below | | | |
| Inj 3.5 mg vial | | 1 | 1 | DBL Bortezomib |
| Inj 1 mg for ECP | 22.26 | 1 mg | ✓ | Baxter |
| SA2355 Special Authority for Subsidy | | | | |
| tial application — (plasma cell dyscrasia) from any relevar | nt practitioner. App | orovals valid | withou | it further renewal unle |
| tified where the patient has plasma cell dyscrasia, not includin | g Waldenström ma | acroglobulina | aemia, | requiring treatment. |
| ACARBAZINE - PCT only - Specialist | • | • | | |
| Inj 200 mg vial | 72.11 | 1 | 1 | DBL Dacarbazine |
| Inj 200 mg for ECP | | 200 mg OP | _ | Baxter |
| ACTINOMYCIN [ACTINOMYCIN D] - PCT only - Specialist | | | | |
| Inj 0.5 mg vial | 255.00 | 1 | / | Cosmegen |
| Inj 0.5 mg for ECP | | 0.5 mg OP | | Baxter |
| | 200.00 | o.o my OF | • | Davici |
| AUNORUBICIN – PCT only – Specialist | 474.00 | | _ | D6 |
| Inj 2 mg per ml, 10 ml | | 1 | | Pfizer |
| Inj 20 mg for ECP | 1/1.93 | 20 mg OP | • | Baxter |
| OCETAXEL - PCT only - Specialist | | | | |
| Inj 20 mg | 48.75 | 1 | | Docetaxel Sandoz |
| Inj 10 mg per ml, 8 ml vial | 24.91 | 1 | 1 | DBL Docetaxel |
| Inj 20 mg per ml, 4 ml vial | 26.95 | 1 | • | Docetaxel |
| | | | | Accord S29 |
| Inj 80 mg | 195.00 | 1 | 1 | Docetaxel Sandoz |
| Inj 1 mg for ECP | 0.35 | 1 mg | 1 | Baxter |
| OXORUBICIN HYDROCHLORIDE - PCT only - Specialist | | | | |
| Inj 2 mg per ml, 5 ml vial | 10.00 | 1 | 1 | Doxorubicin Ebewe |
| Inj 2 mg per ml, 25 ml vial | | 1 | | Doxorubicin Ebewe |
| , 3, , , , | 17.00 | | 1 | Arrow-Doxorubicin |
| Inj 2 mg per ml, 50 ml vial | 23.00 | 1 | 1 | Doxorubicin Ebewe |
| Inj 2 mg per ml, 100 ml vial | | 1 | 1 | Arrow-Doxorubicin |
| , • | 69.99 | | 1 | Doxorubicin Ebewe |
| Inj 1 mg for ECP | 0.35 | 1 mg | ✓ | Baxter |
| PIRUBICIN HYDROCHLORIDE - PCT only - Specialist | | - | | |
| Inj 2 mg per ml, 5 ml vial | 25.00 | 1 | 1 | Epirubicin Ebewe |
| Inj 2 mg per ml, 25 ml vial | | 1 | | Epirubicin Ebewe |
| Inj 2 mg per ml, 100 ml vial | | i | | Epirubicin Ebewe |
| Inj 1 mg for ECP | | 1 mg | | Baxter |
| ΓOPOSIDE | | 3 | | |
| Cap 50 mg - PCT - Retail pharmacy-Specialist | 340.73 | 20 | 1 | Vepesid |
| Cap 100 mg - PCT - Retail pharmacy-Specialist | | 10 | | Vepesid |
| Inj 20 mg per ml, 5 ml vial – PCT – Retail pharmacy-Special st | | 1 | | Rex Medical |
| Inj 1 mg for ECP – PCT only – Specialist | | 1 mg | _ | Baxter |
| | | illy | • | Dunter |
| TOPOSIDE PHOSPHATE – PCT only – Specialist | 40.00 | | , | Flamanha - |
| Inj 100 mg (of etoposide base) | | 1 | / | Etopophos |
| Inj 1 mg (of etoposide base) for ECP | | 1 mg | • | Baxter |
| /DROXYUREA [HYDROXYCARBAMIDE] - PCT - Retail pha | rmacy-Specialist | | | |
| Cap 500 mg | | 100 | | Devatis |

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | |
|---|---|-----|---------------------|-----------|
| IBRUTINIB - Special Authority see SA2168 below - Retail phare | nacy | | | |
| Tab 140 mg | 3,217.00 | 30 | ✓ | Imbruvica |
| Tab 420 mg | 9,652.00 | 30 | • | Imbruvica |

⇒SA2168 Special Authority for Subsidy

Initial application — (chronic lymphocytic leukaemia (CLL)) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic lymphocytic leukaemia (CLL) requiring therapy; and
- 2 Patient has not previously received funded ibrutinib: and
- 3 Ibrutinib is to be used as monotherapy; and
- 4 Any of the following:
 - 4.1 Both:
 - 4.1.1 There is documentation confirming that patient has 17p deletion or TP53 mutation; and
 - 4.1.2 Patient has experienced intolerable side effects with venetoclax monotherapy; or
 - 4.2 All of the following:
 - 4.2.1 Patient has received at least one prior immunochemotherapy for CLL; and
 - 4.2.2 Patient's CLL has relapsed within 36 months of previous treatment; and
 - 4.2.3 Patient has experienced intolerable side effects with venetoclax in combination with rituximab regimen; or

4.3 Patient's CLL is refractory to or has relapsed within 36 months of a venetoclax regimen.

Renewal — (chronic lymphocytic leukaemia (CLL)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 No evidence of clinical disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL) and B-cell prolymphocytic leukaemia (B-PLL)*. Indications marked with * are Unapproved indications.

IDARUBICIN HYDROCHLORIDE

| Inj 5 mg vial - PCT only - Specialist | 109.74 | 1 | ✓ Zavedos |
|--|----------------------|------|-----------------------|
| Inj 10 mg vial - PCT only - Specialist | | 1 | ✓ Zavedos |
| Inj 1 mg for ECP - PCT only - Specialist | | 1 mg | ✓ Baxter |
| LENALIDOMIDE (VIATRIS) - Special Authority see \$A2353 | below - Retail pharm | nacy | |
| Cap 5 mg | | 21 | ✓ Lenalidomide |
| | | | <u>Viatris</u> |
| Cap 10 mg | 50.30 | 21 | ✓ Lenalidomide |
| | | | <u>Viatris</u> |
| Cap 15 mg | 62.13 | 21 | ✓ <u>Lenalidomide</u> |
| | | | <u>Viatris</u> |
| Cap 25 mg | 65.09 | 21 | ✓ <u>Lenalidomide</u> |
| | | | Viatris |

⇒SA2353 Special Authority for Subsidy

Initial application — (Plasma cell dyscrasia) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Both:

- 1 Patient has plasma cell dyscrasia, not including Waldenström macroglobulinaemia, requiring treatment; and
- 2 Patient is not refractory to prior lenalidomide use.

Initial application — (Myelodysplastic syndrome) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

| Subsidy | | Fully | Brand or |
|------------------------|-----|-----------|--------------|
| (Manufacturer's Price) | S | ubsidised | Generic |
| \$ | Per | 1 | Manufacturer |

continued...

Both:

- 1 Patient has low or intermediate-1 risk myelodysplastic syndrome (based on IPSS or an IPSS-R score of less than 3.5) associated with a deletion 5g cytogenetic abnormality; and
- 2 Patient has transfusion-dependent anaemia.

Renewal — (Myelodysplastic syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has not needed a transfusion in the last 4 months: and
- 2 No evidence of disease progression.

MESNA

| MEGIVI | | |
|---|--------|---|
| Tab 400 mg - PCT - Retail pharmacy-Specialist314.00 | 50 | ✓ Uromitexan |
| Tab 600 mg - PCT - Retail pharmacy-Specialist448.50 | 50 | ✓ Uromitexan |
| Inj 100 mg per ml, 4 ml ampoule - PCT only - Specialist177.45 | 15 | ✓ Uromitexan |
| Inj 100 mg per ml, 10 ml ampoule - PCT only - Specialist407.40 | 15 | ✓ Uromitexan |
| Inj 1 mg for ECP - PCT only - Specialist2.96 | 100 mg | ✓ Baxter |
| MITOMYCIN C - PCT only - Specialist | | |
| Inj 5 mg vial517.65 | 1 | ✓ Mitomycin (Fresenius Kabi) ©29 |
| 526.00 | | ✓ Mitomycin (Sagent) S29 |
| 641.70 | | ✓ Accord S29 |
| Inj 20 mg vial1,250.00 | 1 | ✓ Omegapharm (\$29) ✓ Teva |
| Inj 1 mg for ECP269.85 | 1 mg | ✓ Baxter |
| MITOZANTRONE - PCT only - Specialist | | |
| Inj 2 mg per ml, 10 ml vial97.50 | 1 | ✓ Mitozantrone Ebewe |
| Inj 1 mg for ECP5.51 | 1 mg | ✓ Baxter |
| NIRAPARIB – Special Authority see SA2325 below – Retail pharmacy Wastage claimable | | |
| Tab 100 mg13,393.50 | 84 | ✓ Zejula |
| Cap 100 mg8,929.84 | 56 | ✓ Zejula |
| 13,393.50 | 84 | ✓ Zejula |
| | | • |

⇒SA2325 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has advanced high-grade serous* epithelial ovarian, fallopian tube, or primary peritoneal cancer; and
- 2 Patient has received at least one line** of treatment with platinum-based chemotherapy; and
- 3 Patient has experienced a partial or complete response to the preceding treatment with platinum-based chemotherapy; and
- 4 Patient has not previously received funded treatment with a PARP inhibitor; and
- 5 Either:
 - 5.1 Treatment will be commenced within 12 weeks of the patient's last dose of the preceding platinum-based regimen;
 or
 - 5.2 Patient commenced treatment with niraparib prior to 1 May 2024; and

Lynparza

| Subsidy | Fully | Brand or | |
|------------------------|------------|--------------|--|
| (Manufacturer's Price) | Subsidised | Generic | |
| \$ | Per 🗸 | Manufacturer | |

continued...

- 6 Treatment to be administered as maintenance treatment; and
- 7 Treatment not to be administered in combination with other chemotherapy.

Renewal from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 No evidence of progressive disease; and
- 2 Treatment to be administered as maintenance treatment; and
- 3 Treatment not to be administered in combination with other chemotherapy; and
- 4 Fither:
 - 4.1 Treatment with niraparib to cease after a total duration of 36 months from commencement; or
 - 4.2 Treatment with niraparib is being used in the second-line or later maintenance setting.

Notes: * "high-grade serous" includes tumours with high-grade serous features or a high-grade serous component.

**A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments

| OLAPARIB - Retail pharmacy-Specialist - Special A | uthority see SA2163 below | | |
|---|---------------------------|----|----------------------------|
| Tab 100 mg | 3,701.00 | 56 | Lynparza |
| Tab 150 mg | 3.701.00 | 56 | ✓ Lynparza |

⇒SA2163 Special Authority for Subsidy

Initial application — (Ovarian cancer) only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Patient has a high-grade serous* epithelial ovarian, fallopian tube, or primary peritoneal cancer; and
- 2 There is documentation confirming pathogenic germline BRCA1 or BRCA2 gene mutation; and
- 3 Fither:
 - 3.1 All of the following:
 - 3.1.1 Patient has newly diagnosed, advanced disease; and
 - 3.1.2 Patient has received one line** of previous treatment with platinum-based chemotherapy; and
 - 3.1.3 Patient's disease must have experienced a partial or complete response to the first-line platinum-based regimen; or
 - 3.2 All of the following:
 - 3.2.1 Patient has received at least two lines** of previous treatment with platinum-based chemotherapy; and
 - 3.2.2 Patient has platinum sensitive disease defined as disease progression occurring at least 6 months after the last dose of the penultimate line** of platinum-based chemotherapy; and
 - 3.2.3 Patient's disease must have experienced a partial or complete response to treatment with the immediately preceding platinum-based regimen; and
 - 3.2.4 Patient has not previously received funded olaparib treatment; and
- 4 Treatment will be commenced within 12 weeks of the patient's last dose of the immediately preceding platinum-based regimen: and
- 5 Treatment to be administered as maintenance treatment; and
- 6 Treatment not to be administered in combination with other chemotherapy.

Renewal — (Ovarian cancer) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from treatment; and
- 2 Fither:
 - 2.1 No evidence of progressive disease: or
 - 2.2 Evidence of residual (not progressive) disease and the patient would continue to benefit from treatment in the

| Subsidy (Manufacturer's Price) | Sub | Fully | Brand or Generic | |
|-----------------------------------|-----|-------|---------------------|--|
| \$ | Per | 1 | Manufacturer | |

continued...

clinician's opinion: and

- 3 Treatment to be administered as maintenance treatment; and
- 4 Treatment not to be administered in combination with other chemotherapy; and
- 5 Either:
 - 5.1 Both:
 - 5.1.1 Patient has received one line** of previous treatment with platinum-based chemotherapy; and
 - 5.1.2 Documentation confirming that the patient has been informed and acknowledges that the funded treatment period of olaparib will not be continued beyond 2 years if the patient experiences a complete response to treatment and there is no radiological evidence of disease at 2 years; or
 - 5.2 Patient has received at least two lines** of previous treatment with platinum-based chemotherapy.

Notes: *Note "high-grade serous" includes tumours with high-grade serous features or a high-grade serous component.
**A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

| PACLITAXEL - PCT only - Specialist | | | |
|--|-----------------|------|--------------------------------|
| Inj 30 mg | 47.30 | 5 | ✓ Paclitaxel Ebewe |
| Inj 6 mg per ml, 16.7 ml vial | | 1 | ✓ Anzatax |
| | 24.00 | | ✓ Paclitaxel Ebewe |
| | 91.67 | | ✓ Paclitaxel Actavis |
| Inj 150 mg | 26.69 | 1 | ✓ Paclitaxel Ebewe |
| | 137.50 | | ✓ Anzatax |
| | | | ✓ Paclitaxel Actavis |
| Inj 6 mg per ml, 50 ml vial | 37.89 | 1 | ✓ Anzatax |
| | 44.00 | | ✓ Paclitaxel Ebewe |
| | 275.00 | | ✓ Paclitaxel Actavis |
| Inj 1 mg for ECP | 0.17 | 1 mg | ✓ Baxter |
| PEGASPARGASE - PCT only - Special Authority se | ee SA1979 below | | |
| Inj 750 iu per ml, 5 ml vial | 3,973.25 | 1 | Oncaspar LYO |
| | | | |

⇒SA1979 Special Authority for Subsidy

Initial application — (Acute lymphoblastic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has newly diagnosed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

Initial application — (Lymphoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months where the patient has lymphoma requiring L-asparaginase containing protocols (e.g. SMILE).

Renewal — (Acute lymphoblastic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has relapsed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

PENTOSTATIN [DEOXYCOFORMYCIN] - PCT only - Specialist

| | Subsidy (Manufacturer's Price) \$ | Sı Per | Fully ubsidised | Brand or Generic Manufacturer | |
|---|---|-----------|--------------------|-------------------------------------|--|
| POMALIDOMIDE - Special Authority see SA2354 below - Retai | l pharmacy | | | | |
| Cap 1 mg | 47.45 | 14 | ✓ P | omolide | |
| , , | 71.18 | 21 | ✓ P | omolide | |
| Cap 2 mg | 94.90 | 14 | ✓ P | omolide | |
| , , | 142.35 | 21 | ✓ P | omolide | |
| Cap 3 mg | 142.35 | 14 | ✓ P | omolide | |
| , , | 213.53 | 21 | ✓ P | omolide | |
| Cap 4 mg | 189.81 | 14 | ✓ P | omolide | |
| , , | 284.71 | 21 | ✓ P | omolide | |

⇒SA2354 Special Authority for Subsidy

Initial application — (Relapsed/refractory plasma cell dyscrasia) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has relapsed or refractory plasma cell dyscrasia, not including Waldenström macroglobulinaemia, requiring treatment; and
- 2 Patient has not received prior funded pomalidomide.

Renewal — (Relapsed/refractory plasma cell dyscrasia) from any relevant practitioner. Approvals valid for 12 months where there is no evidence of disease progression.

PROCARBAZINE HYDROCHLORIDE - PCT - Retail pharmacy-Specialist

| Cap 50 mg | 980.00 | 50 | ✓ Natulan S29 |
|--------------------------------------|----------------------------------|----|--------------------|
| TEMOZOLOMIDE - Special Authority see | e SA2275 below - Retail pharmacy | | |
| Cap 5 mg | | 5 | ✓ Temaccord |
| | | | ✓ Temozolomide- |
| | | | Taro S29 |
| Cap 20 mg | 16.38 | 5 | ✓ Temaccord |
| | 18.30 | | ✓ Apo-Temozolomide |
| Cap 100 mg | 35.98 | 5 | ✓ Temaccord |
| | 40.20 | | ✓ Apo-Temozolomide |
| Cap 140 mg | 50.12 | 5 | ✓ Temaccord |
| Cap 250 mg | 86.34 | 5 | ✓ Temaccord |

⇒SA2275 Special Authority for Subsidy

Initial application — (gliomas) only from a relevant specialist. Approvals valid for 12 months where the patient has a glioma. Renewal — (gliomas) only from a relevant specialist. Approvals valid for 12 months where treatment remains appropriate and patient is benefitting from treatment.

Initial application — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with metastatic or unresectable well-differentiated neuroendocrine tumour*: and
- 2 Temozolomide is to be given in combination with capecitabine; and
- 3 Temozolomide is to be used in 28 day treatment cycles for a maximum of 5 days treatment per cycle at a maximum dose of 200 mg/m² per day; and
- 4 Temozolomide to be discontinued at disease progression.

Renewal — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

continued...

Initial application — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 9 months where the patient has relapsed/refractory Ewing's sarcoma.

Renewal — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indication marked with a * is an unapproved indication. Temozolomide is not subsidised for the treatment of relapsed high grade glioma.

| THALIDOMIDE - Retail pharmacy-Specialist - Special Au | thority see SA2356 below | | |
|---|--------------------------|----|----------|
| Cap 50 mg | 378.00 | 28 | Thalomid |
| Cap 100 mg | 756.00 | 28 | Thalomid |

⇒SA2356 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months where the patient has plasma cell dyscrasia, not including Waldenström macroglobulinaemia, requiring treatment.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has obtained a response from treatment during the initial approval period.

Notes: Prescription must be written by a registered prescriber in the thalidomide risk management programme operated by the supplier.

Maximum dose of 400 mg daily as monotherapy or in a combination therapy regimen.

TRETINOIN

| Cap 10 mg - PCT - Retail pharmacy-Specialist | 479.50 | 100 | ✓ Vesanoid |
|--|----------------|-------|-------------|
| VENETOCLAX - Retail pharmacy-Specialist - Special Authority se | e SA1868 below | ı | |
| Tab 14 × 10 mg, 7 × 50 mg, 21 × 100 mg | .1,771.86 | 42 OP | ✓ Venclexta |
| Tab 10 mg | | 2 OP | ✓ Venclexta |
| Tab 50 mg | 239.44 | 7 OP | ✓ Venclexta |
| Tab 100 mg - Wastage claimable | .8,209.41 | 120 | ✓ Venclexta |

⇒SA1868 Special Authority for Subsidy

Initial application — (relapsed/refractory chronic lymphocytic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic lymphocytic leukaemia requiring treatment; and
- 2 Patient has received at least one prior therapy for chronic lymphocytic leukaemia; and
- 3 Patient has not previously received funded venetoclax; and
- 4 The patient's disease has relapsed within 36 months of previous treatment; and
- 5 Venetoclax to be used in combination with six 28-day cycles of rituximab commencing after the 5-week dose titration schedule with venetoclax; and
- 6 Patient has an ECOG performance status of 0-2.

Renewal — (relapsed/refractory chronic lymphocytic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment; and
- 2 Venetoclax is to be discontinued after a maximum of 24 months of treatment following the titration schedule unless earlier discontinuation is required due to disease progression or unacceptable toxicity.

Initial application — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation*) only from

| Subsidy | | Fully | Brand or | |
|--------------------|-------|------------|--------------|--|
| (Manufacturer's Pr | rice) | Subsidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

continued...

a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has previously untreated chronic lymphocytic leukaemia; and
- 2 There is documentation confirming that patient has 17p deletion by FISH testing or TP53 mutation by sequencing; and
- 3 Patient has an ECOG performance status of 0-2.

Renewal — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where the treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL)* and B-cell prolymphocytic leukaemia (B-PLL)*. Indications marked with * are Unapproved indications.

VINBLASTINE SULPHATE

| Inj 1 mg per ml, 10 ml vial – PCT – Retail pharmacy-Specialist270.37 Inj 1 mg for ECP – PCT only – Specialist6.00 | 5 1 mg | ✓ Hospira✓ Baxter |
|---|-----------|---|
| VINCRISTINE SULPHATE | | |
| Inj 1 mg per ml, 1 ml vial - PCT - Retail pharmacy-Specialist74.52 | 5 | DBL Vincristine Sulfate |
| Inj 1 mg per ml, 2 ml vial - PCT - Retail pharmacy-Specialist 102.73 | 5 | DBL Vincristine Sulfate |
| Inj 1 mg for ECP - PCT only - Specialist12.60 | 1 mg | ✓ Baxter |
| VINORELBINE | | |
| Cap 20 mg30.00 | 1 | ✓ Vinorelbine Te Arai |
| Cap 30 mg40.00 | 1 | ✓ Vinorelbine Te Arai |
| Cap 80 mg60.00 | 1 | ✓ Vinorelbine Te Arai |
| Inj 10 mg per ml, 1 ml vial - PCT only - Specialist42.00 | 1 | ✓ Vinorelbine Ebewe |
| Inj 10 mg per ml, 5 ml vial - PCT only - Specialist168.00 | 1 | ✓ Navelbine S29 S29 |
| 210.00 | | ✓ Vinorelbine Ebewe |
| Inj 1 mg for ECP - PCT only - Specialist | 1 mg | ✓ Baxter |
| | | |

Protein-tyrosine Kinase Inhibitors

| ALECTINIB – Retail pharmacy-Specialist – Special Authority see SA1870 below | | |
|---|-----|------------|
| Wastage claimable | | |
| Cap 150 mg7,935.00 | 224 | ✓ Alecensa |

⇒SA1870 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-small cell lung cancer; and
- 2 There is documentation confirming that the patient has an ALK tyrosine kinase gene rearrangement using an appropriate ALK test: and
- 3 Patient has an ECOG performance score of 0-2.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of progressive disease according to RECIST criteria; and
- 2 The patient is benefitting from and tolerating treatment.

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer | |
|--|---|-----|---------------------|-------------------------------------|---|
| DASATINIB – Special Authority see SA2385 below – Retail phar | macy | | | | _ |
| Wastage claimable | | | | | |
| Tab 20 mg | 132.88 | 60 | ✓ [| Dasatinib-Teva | |
| · | 3,774.06 | | ✓ 9 | Sprycel | |
| Dasatinib-Teva to be Principal Supply on 1 March 2025 | • | | | . , | |
| Tab 50 mg | 304.13 | 60 | ✓ [| Dasatinib-Teva | |
| 3 | 6,214.20 | | √ 9 | Sprycel | |
| Dasatinib-Teva to be Principal Supply on 1 March 2025 | | | | | |
| Tab 70 mg | 415.75 | 60 | ✓ [| Dasatinib-Teva | |
| · · | 7,692.58 | | ✓ 9 | Sprycel | |
| Dasatinib-Teva to be Principal Supply on 1 March 2025 | | | | | |
| (Spryool Tab 20 mg to be delicted 1 March 2025) | | | | | |

(Sprycel Tab 20 mg to be delisted 1 March 2025)

(Sprycel Tab 50 mg to be delisted 1 March 2025)

(Sprycel Tab 70 mg to be delisted 1 March 2025)

⇒SA2385 Special Authority for Subsidy

Initial application only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis or accelerated phase; or
- 2 The patient has a diagnosis of Philadelphia chromosome-positive acute lymphoid leukaemia (Ph+ ALL); or
- 3 Roth
 - 3.1 The patient has a diagnosis of CML in chronic phase; and
 - 3.2 Any of the following:
 - 3.2.1 Patient has documented treatment failure* with imatinib; or
 - 3.2.2 Patient has experienced treatment-limiting toxicity with imatinib precluding further treatment with imatinib; or
 - 3.2.3 Patient has high-risk chronic-phase CML defined by the Sokal or EURO scoring system.

Renewal only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Lack of treatment failure while on dasatinib*: and
- 2 Dasatinib treatment remains appropriate and the patient is benefiting from treatment.

Note: *treatment failure for CML as defined by Leukaemia Net Guidelines.

| ERLOTINIB – Retail pharmacy-Specialist – Special Authority | y see SA2422 below | | |
|--|--------------------|----|-----------|
| Tab 100 mg | 280.84 | 30 | ✓ Alchemy |
| Tab 150 mg | 484.24 | 30 | ✓ Alchemy |

⇒SA2422 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 There is documentation confirming that the disease expresses activating mutations of EGFR; and
- 3 Any of the following:
 - 3.1 Patient is treatment naive: or
 - 3.2 Patient has received prior treatment in the adjuvant setting and/or while awaiting EGFR results; or
 - 3.3 Both:
 - 3.3.1 The patient has discontinued osimertinib or gefitinib due to intolerance; and
 - 3.3.2 The cancer did not progress while on osimertinib or gefitinib.

Renewal from any relevant practitioner. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

GEFITINIB − Retail pharmacy-Specialist − Special Authority see SA2423 on the next page
Tab 250 mg918.00 30 ✓ Iressa

| Subsidy | | Fully | Brand or | |
|------------------------|-----|------------|--------------|--|
| (Manufacturer's Price) | | Subsidised | Generic | |
| \$ | Per | / | Manufacturer | |

⇒SA2423 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 Any of the following:
 - 2.1 Patient is treatment naive; or
 - 2.2 Patient has received prior treatment in the adjuvant setting and/or while awaiting EGFR results; or
 - 2.3 Both:
 - 2.3.1 The patient has discontinued osimertinib or erlotinib due to intolerance; and
 - 2.3.2 The cancer did not progress whilst on osimertinib or erlotinib; and
- 3 There is documentation confirming that disease expresses activating mutations of EGFR.

Renewal from any relevant practitioner. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

IMATINIR MESILATE

| * Cap 100 mg * Cap 400 mg | | 60 30 | ✓ <u>Imatinib-Rex</u> ✓ <u>Imatinib-Rex</u> |
|---|---------------------|----------|--|
| LENVATINIB – Special Authority see SA2407 below Wastage claimable | / - Retail pharmacy | | |
| Cap 4 mg | 3,407.40 | 30 | ✓ Lenvima |
| Cap 10 mg | 3,407.40 | 30 | ✓ Lenvima |

⇒SA2407 Special Authority for Subsidy

Initial application — (thyroid cancer) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Patient is currently on treatment with lenvatinib and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 The patient has locally advanced or metastatic differentiated thyroid cancer; and
 - 2.2 Either
 - 2.2.1 Patient must have symptomatic progressive disease prior to treatment; or
 - 2.2.2 Patient must progressive disease at critical anatomical sites with a high risk of morbidity or mortality where local control cannot be achieved by other measures; and
 - 2.3 Any of the following:
 - 2.3.1 A lesion without iodine uptake in a RAI scan; or
 - 2.3.2 Receiving cumulative RAI greater than or equal to 600 mCi; or
 - 2.3.3 Experiencing disease progression after a RAI treatment within 12 months; or
 - 2.3.4 Experiencing disease progression after two RAI treatments administered within 12 months of each other; and
 - 2.4 Patient has thyroid stimulating hormone (TSH) adequately supressed; and
 - 2.5 Patient is not a candidate for radiotherapy with curative intent; and
 - 2.6 Surgery is clinically inappropriate; and
 - 2.7 Patient has an ECOG performance status of 0-2.

Renewal — (thyroid cancer) from any relevant practitioner. Approvals valid for 6 months where there is no evidence of disease progression.

Initial application — (unresectable hepatocellular carcinoma) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

1 Patient has unresectable hepatocellular carcinoma; and

| Subsidy | Ful | y Brand or | |
|------------------------|-----------|----------------------------------|--|
| (Manufacturer's Price) | Subsidise | d Generic | |
| \$ | Per • | Manufacturer | |

continued...

- 2 Patient has preserved liver function (Childs-Pugh A); and
- 3 Transarterial chemoembolisation (TACE) is unsuitable; and
- 4 Patient has an ECOG performance status of 0-2; and
- 5 Patient has not received prior systemic therapy for their disease in the palliative setting.

Renewal — (unresectable hepatocellular carcinoma) from any relevant practitioner. Approvals valid for 6 months where there is no evidence of disease progression.

Initial application — (renal cell carcinoma) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient has metastatic renal cell carcinoma; and
 - 1.2 The disease is of predominant clear-cell histology; and
 - 1.3 The patient has documented disease progression following one previous line of treatment; and
 - 1.4 The patient has an ECOG performance status of 0-2; and
 - 1.5 Lenvatinib is to be used in combination with everolimus; or
- 2 All of the following:
 - 2.1 Patient has received funded treatment with nivolumab for the second line treatment of metastatic renal cell carcinoma; and
 - 2.2 Patient has experienced treatment limiting toxicity from treatment with nivolumab; and
 - 2.3 Lenvatinib is to be used in combination with everolimus; and
 - 2.4 There is no evidence of disease progression.

Renewal — (renal cell carcinoma) from any relevant practitioner. Approvals valid for 4 months where there is no evidence of disease progression.

MIDOSTAURIN - PCT only - Special Authority see SA2342 below

Cap 25 mg.......10,981.00 56 **✓ Rydapt**

⇒SA2342 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria: All of the following:

- 1 Patient has a diagnosis of acute myeloid leukaemia; and
- 2 Condition must be FMS tyrosine kinase 3 (FLT3) mutation positive; and
- 3 Patient must not have received a prior line of intensive chemotherapy for acute myeloid leukaemia; and
- 4 Patient is to receive standard intensive chemotherapy in combination with midostaurin only; and
- 5 Midostaurin to be funded for a maximum of 4 cycles.

NILOTINIB - Special Authority see SA2301 below - Retail pharmacy

Wastage claimable

| Cap 150 mg | 4,680.00 | 120 | Tasigna |
|------------|----------|-----|-----------|
| Cap 200 mg | 6,532.00 | 120 | ✓ Tasigna |

⇒SA2301 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis, high risk chronic phase, or in chronic phase; and
- 2 Either:
 - 2.1 Patient has documented CML treatment failure* with a tyrosine kinase inhibitor (TKI); or
 - 2.2 Patient has experienced treatment limiting toxicity with a tyrosine kinase inhibitor (TKI) precluding further treatment; and

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

continued...

- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

Note: *treatment failure as defined by Leukaemia Net Guidelines.

Renewal only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Lack of treatment failure while on nilotinib as defined by Leukaemia Net Guidelines; and
- 2 Nilotinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

| OSIMERTINIB - Special Authority see SA2418 below - | Retail pharmacy | | |
|--|-----------------|----|------------|
| Tab 40 mg | 9,310.00 | 30 | ✓ Tagrisso |
| Tab 80 mg | 9,310.00 | 30 | ✓ Tagrisso |

⇒SA2418 Special Authority for Subsidy

Initial application — (NSCLC – first line) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 Patient is currently on treatment with osimertinib and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 Patient has locally advanced or metastatic, incurable, non-squamous non-small cell lung cancer (NSCLC); and
 - 2.2 Any of the following:
 - 2.2.1 Patient is treatment naïve: or
 - 2.2.2 Patient has received prior chemotherapy in the adjuvant setting and/or while awaiting EGFR results; or
 - 2.2.3 Both:
 - 2.2.3.1 The patient has discontinued gefitinib or erlotinib due to intolerance; and
 - 2.2.3.2 The cancer did not progress while on gefitinib or erlotinib; and
 - 2.3 There is documentation confirming that the cancer expresses activating mutations of EGFR; and
 - 2.4 Patient has an ECOG performance status 0-3; and
 - 2.5 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Renewal — (NSCLC – first line) from any relevant practitioner. Approvals valid for 6 months where response to or stable disease with treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period.

Initial application — (NSCLC – second line) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 Patient is currently on treatment with osimertinib and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 Patient has locally advanced or metastatic, incurable, non-squamous non-small cell lung cancer (NSCLC); and
 - 2.2 Patient has an ECOG performance status 0-3; and
 - 2.3 The patient must have received previous treatment with erlotinib or gefitinib; and
 - 2.4 There is documentation confirming that the cancer expresses T790M mutation of EGFR following progression on or after erlotinib or gefitinib; and
 - 2.5 The treatment must be given as monotherapy; and
 - 2.6 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Renewal — (NSCLC – second line) from any relevant practitioner. Approvals valid for 6 months where response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period.

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer |
|---|---|-----|---------------------|-------------------------------------|
| PALBOCICLIB – Special Authority see SA2345 below – Retail pl Wastage claimable | harmacy | | | |
| Tab 75 mg | 4,000.00 | 21 | ✓ | Ibrance |
| Tab 100 mg | 4,000.00 | 21 | ✓ | Ibrance |
| Tab 125 mg | 4,000.00 | 21 | ✓ | Ibrance |

⇒SA2345 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Fither:

- 1 All of the following:
 - 1.1 Patient has unresectable locally advanced or metastatic breast cancer; and
 - 1.2 There is documentation confirming disease is hormone-receptor positive and HER2-negative; and
 - 1.3 Patient has an ECOG performance score of 0-2; and
 - 1.4 Either:
 - 1.4.1 Disease has relapsed or progressed during prior endocrine therapy; or
 - 1.4.2 Both:
 - 1.4.2.1 Patient is amenorrhoeic, either naturally or induced, with endocrine levels consistent with a postmenopausal or without menstrual-potential state; and
 - 1.4.2.2 Patient has not received prior systemic treatment for metastatic disease; and
 - 1.5 Treatment must be used in combination with an endocrine partner; and
 - 1.6 Patient has not received prior funded treatment with a CDK4/6 inhibitor; or
- 2 All of the following:
 - 2.1 Patient has an active Special Authority approval for ribociclib; and
 - 2.2 Patient has experienced a grade 3 or 4 adverse reaction to ribociclib that cannot be managed by dose reductions and requires treatment discontinuation; and
 - 2.3 Treatment must be used in combination with an endocrine partner; and
 - 2.4 There is no evidence of progressive disease since initiation of ribociclib.

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 Treatment must be used in combination with an endocrine partner; and
- 2 There is no evidence of progressive disease since initiation of palbociclib.

| PAZOPANIB - Special Authority see SA2429 below - F | Retail pharmacy | | |
|--|-----------------|----|----------------|
| Tab 200 mg | 172.88 | 30 | Pazopanib Teva |
| · | 1,334.70 | | ✓ Votrient |
| Pazopanib Teva to be Principal Supply on 1 Ma | ay 2025 | | |
| Tab 400 mg | 464.00 | 30 | Pazopanib Teva |
| ř | 2,669.40 | | ✓ Votrient |

Pazopanib Teva to be Principal Supply on 1 May 2025

(Votrient Tab 200 mg to be delisted 1 May 2025)

(Votrient Tab 400 mg to be delisted 1 May 2025)

⇒SA2429 Special Authority for Subsidy

Initial application only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

- Either:
 - 1 All of the following:
 - 1.1 The patient has metastatic renal cell carcinoma of predominantly clear cell histology; and
 - - 1.2.1 The patient is treatment naive; or

| Subsidy (Manufacturer's Price) | S | Fully Subsidised | Brand or Generic |
|-----------------------------------|-----|---------------------|---------------------|
| \$ | Per | ✓ | Manufacturer |

continued...

- 1.2.2 The patient has only received prior cytokine treatment; and
- 1.3 The patient has an ECOG performance score of 0-2; and The patient has intermediate or poor prognosis defined as:
- 1.4 Any of the following:
 - 1.4.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
 - 1.4.2 Haemoglobin level < lower limit of normal; or
 - 1.4.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
 - 1.4.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
 - 1.4.5 Karnofsky performance score of less than or equal to 70; or
 - 1.4.6 2 or more sites of organ metastasis; and
- 1.5 Pazopanib to be used for a maximum of 3 months; or
- 2 All of the following:
 - 2.1 The patient has metastatic renal cell carcinoma; and
 - 2.2 The patient has discontinued sunitinib within 3 months of starting treatment due to intolerance; and
 - 2.3 The cancer did not progress whilst on sunitinib; and
 - 2.4 Pazopanib to be used for a maximum of 3 months.

Renewal only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months where there is no evidence of disease progression.

RIBOCICLIB - Special Authority see SA2343 below - Retail pharmacy

| Wastage claimable | · | | |
|-------------------|----------|----|---------------------------|
| Tab 200 mg | 1,883.00 | 21 | Kisqali |
| | 3,767.00 | 42 | ✓ Kisqali |
| | 5,650.00 | 63 | ✓ Kisqali |

⇒SA2343 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Either:

- 1 All of the following:
 - 1.1 Patient has unresectable locally advanced or metastatic breast cancer; and
 - 1.2 There is documentation confirming disease is hormone-receptor positive and HER2-negative; and
 - 1.3 Patient has an ECOG performance score of 0-2; and
 - 1.4 Any of the following:
 - 1.4.1 Disease has relapsed or progressed during prior endocrine therapy; or
 - 1.4.2 Both:
 - 1.4.2.1 Patient is amenorrhoeic, either naturally or induced, with endocrine levels consistent with a postmenopausal or without menstrual-potential state; and
 - 1.4.2.2 Patient has not received prior systemic endocrine treatment for metastatic disease; or
 - 1.4.3 Both:
 - 1.4.3.1 Patient commenced treatment with ribociclib in combination with an endocrine partner prior to 1 July 2024; and
 - 1.4.3.2 There is no evidence of progressive disease; and
 - 1.5 Treatment to be used in combination with an endocrine partner; and
 - 1.6 Patient has not received prior funded treatment with a CDK4/6 inhibitor; or
- 2 All of the following:
 - 2.1 Patient has an active Special Authority approval for palbociclib; and
 - 2.2 Patient has experienced a grade 3 or 4 adverse reaction to palbociclib that cannot be managed by dose reductions and requires treatment discontinuation; and

| Subsidy | Fully | | Brand or |
|------------------------|------------|---|----------|
| (Manufacturer's Price) | Subsidised | | Generic |
| \$ | Per | 1 | |

continued...

- 2.3 Treatment must be used in combination with an endocrine partner; and
- 2.4 There is no evidence of progressive disease since initiation of palbociclib.

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 Treatment must be used in combination with an endocrine partner; and
- 2 There is no evidence of progressive disease since initiation of ribociclib.

RUXOLITINIB - Special Authority see SA1890 below - Retail pharmacy

| Wastage | claimable |
|----------|-----------|
| Tab 5 mg | |

| Tractage claimable | | | |
|--------------------|----------|----|--------------------------|
| Tab 5 mg | 2,500.00 | 56 | Jakavi |
| Tab 10mg | 5,000.00 | 56 | Jakavi |
| Tab 15 mg | 5,000.00 | 56 | Jakavi |
| Tab 20 mg | 5,000.00 | 56 | Jakavi |
| - | | | |

⇒SA1890 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has primary myelofibrosis or post-polycythemia vera myelofibrosis or post-essential thrombocythemia myelofibrosis; and
- 2 Either:
 - 2.1 A classification of risk of intermediate-2 or high-risk myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS; or
 - - 2.2.1 A classification of risk of intermediate-1 myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS; and
 - 2.2.2 Patient has severe disease-related symptoms that are resistant, refractory or intolerant to available therapy:
- 3 A maximum dose of 20 mg twice daily is to be given.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 A maximum dose of 20 mg twice daily is to be given.

SUNITINIB - Special Authority see SA2430 below - Retail pharmacy

| Cap 12.5 mg | 208.38 | 28 | Sunitinib Pfizer |
|-------------|--------|----|--------------------------------------|
| , , | 416.77 | 28 | Sunitinib Pfizer |
| , , | 694.62 | 28 | Sunitinib Pfizer |

⇒SA2430 Special Authority for Subsidy

Initial application — (RCC) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic renal cell carcinoma of predominantly clear cell histology; and
- 2 Any of the following:
 - 2.1 The patient is treatment naive; or
 - 2.2 The patient has only received prior cytokine treatment; or
 - 2.3 The patient has only received prior treatment with an investigational agent within the confines of a bona fide clinical

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

trial which has Ethics Committee approval; or

- 2.4 Both:
 - 2.4.1 The patient has discontinued pazopanib within 3 months of starting treatment due to intolerance; and
 - 2.4.2 The cancer did not progress whilst on pazopanib; and
- 3 The patient has an ECOG performance score of 0-2; and
- 4 Sunitinib to be used for a maximum of 2 cycles.

Initial application — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal tumour (GIST); and
- 2 Fither:
 - 2.1 The patient's disease has progressed following treatment with imatinib; or
 - 2.2 The patient has documented treatment-limiting intolerance, or toxicity to, imatinib.

Renewal — (RCC) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months where there is no evidence of disease progression.

Renewal — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

The patient has responded to treatment or has stable disease as determined by Choi's modified CT response evaluation criteria as follows:

- 1 Any of the following:
 - 1.1 The patient has had a complete response (disappearance of all lesions and no new lesions); or
 - 1.2 The patient has had a partial response (a decrease in size of 10% or more or decrease in tumour density in Hounsfield Units (HU) of 15% or more on CT and no new lesions and no obvious progression of non measurable disease); or
 - 1.3 The patient has stable disease (does not meet criteria the two above) and does not have progressive disease and no symptomatic deterioration attributed to tumour progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: It is recommended that response to treatment be assessed using Choi's modified CT response evaluation criteria (J Clin Oncol, 2007, 25:1753-1759). Progressive disease is defined as either: an increase in tumour size of 10% or more and not meeting criteria of partial response (PR) by tumour density (HU) on CT; or: new lesions, or new intratumoral nodules, or increase in the size of the existing intratumoral nodules.

Renewal — (GIST pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal (GIST); and
- 2 The patient is clinically benifiting from treatment and continued treatment remains appropriate; and
- 3 Sunitinib is to be discontinued at progression; and
- 4 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

Endocrine Therapy

For GnRH ANALOGUES - refer to HORMONE PREPARATIONS, Trophic Hormones, page 88

ABIRATERONE ACETATE - Retail pharmacy-Specialist - Special Authority see SA2118 on the next page

Wastage claimable

| Subsidy (Manufacturer's Price) | Subsic | Fully dised | Brand or Generic |
|-----------------------------------|--------|----------------|---------------------|
| \$ | Per | 1 | Manufacturer |

⇒SA2118 Special Authority for Subsidy

Initial application only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has prostate cancer; and
- 2 Patient has metastases; and
- 3 Patient's disease is castration resistant; and
- 4 Fither:
 - 4.1 All of the following:
 - 4.1.1 Patient is symptomatic; and
 - 4.1.2 Patient has disease progression (rising serum PSA) after second line anti-androgen therapy; and
 - 4.1.3 Patient has ECOG performance score of 0-1; and
 - 4.1.4 Patient has not had prior treatment with taxane chemotherapy; or
 - 4.2 All of the following:
 - 4.2.1 Patient's disease has progressed following prior chemotherapy containing a taxane; and
 - 4.2.2 Patient has ECOG performance score of 0-2; and
 - 4.2.3 Patient has not had prior treatment with abiraterone.

Renewal — (abiraterone acetate) only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Significant decrease in serum PSA from baseline; and
- 2 No evidence of clinical disease progression; and
- 3 No initiation of taxane chemotherapy with abiraterone: and
- 4 The treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 2 Abiraterone acetate to be discontinued at progression; and
- 3 No initiation of taxane chemotherapy with abiraterone; and
- 4 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

| | - | | | _ |
|---|-----|---------|------|-----|
| н | ICA | IΔN | /111 | ۱۱– |
| | | | | |

| Tab 50 mg | 4.18 | 28 | ✓ Binarex |
|--|-----------------------|-----|-----------------|
| FLUTAMIDE | | | |
| Tab 250 mg | 107.55 | 90 | ✓ Prostacur S29 |
| · · | 119.50 | 100 | ✓ Flutamin |
| FULVESTRANT - Retail pharmacy-Specialist - Special Aut | hority see SA1895 bel | ow | |
| Inj 50 mg per ml, 5 ml prefilled syringe | 1,068.00 | 2 | ✓ Faslodex |

⇒SA1895 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has oestrogen-receptor positive locally advanced or metastatic breast cancer; and
- 2 Patient has disease progression following prior treatment with an aromatase inhibitor or tamoxifen for their locally advanced or metastatic disease; and

| Subsidy (Manufacturer | | Fully idised | Brand or Generic | |
|--------------------------|-----|--------------|---------------------|--|
| \$ | Per | ✓ | Manufacturer | |

continued...

- 3 Treatment to be given at a dose of 500 mg monthly following loading doses; and
- 4 Treatment to be discontinued at disease progression.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains appropriate and patient is benefitting from treatment; and
- 2 Treatment to be given at a dose of 500 mg monthly; and
- 3 There is no evidence of disease progression.

OCTREOTIDE

| OCTRECTIDE | | | |
|---|------------------------|--------|---------------------|
| Inj 50 mcg per ml, 1 ml vial | 27.58 | 5 | ✓ Omega S29 |
| Inj 100 mcg per ml, 1 ml vial | 48.50 | 5 | ✓ Omega S29 |
| Inj 500 mcg per ml, 1 ml vial | | 5 | ✓ Omega S29 |
| Inj 50 mcg per ml, 1 ml ampoule | 27.58 | 5 | ✓ Max Health |
| | | | ✓ Octreotide GH S29 |
| Inj 100 mcg per ml, 1 ml ampoule | 32.71 | 5 | Max Health |
| | | | ✓ Octreotide GH S29 |
| | | | ✓ Sun Pharma S29 |
| Inj 500 mcg per ml, 1 ml ampoule | 113.10 | 5 | ✓ Max Health |
| | | | ✓ Octreotide GH S29 |
| | | | ✓ Sun Pharma S29 |
| OCTREOTIDE LONG-ACTING - Special Authority see SA | 2119 below – Retail ph | armacy | |
| Inj depot 10 mg prefilled syringe | | í | ✓ Sandostatin LAR |
| Inj depot 20 mg prefilled syringe | | 1 | ✓ Sandostatin LAR |
| Inj depot 30 mg prefilled syringe | | 1 | ✓ Sandostatin LAR |

⇒SA2119 Special Authority for Subsidy

Initial application — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 The patient has nausea* and vomiting* due to malignant bowel obstruction*; and
- 2 Treatment with antiemetics, rehydration, antimuscarinic agents, corticosteroids and analgesics for at least 48 hours has failed; and
- 3 Octreotide to be given at a maximum dose 1500 mcg daily for up to 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 3 months where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has acromegaly; and
- 2 Any of the following:
 - 2.1 Treatment with surgery, radiotherapy and a dopamine agonist has failed; or
 - 2.2 Treatment with octreotide is for an interim period while awaiting the effects of radiotherapy and a dopamine agonist has failed; or
 - 2.3 The patient is unwilling, or unable, to undergo surgery and/or radiotherapy.

Renewal — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist.

| Subsidy | Fully | | Brand or |
|------------------------|------------|---|----------|
| (Manufacturer's Price) | Subsidised | | Generic |
| \$ | Per | 1 | |

Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 IGF1 levels have decreased since starting octreotide; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: In patients with Acromegaly octreotide treatment should be discontinued if IGF1 levels have not decreased after 3 months treatment. In patients treated with radiotherapy octreotide treatment should be withdrawn every 2 years, for 1 month, for assessment of remission. Octreotide treatment should be stopped where there is biochemical evidence of remission (normal IGF1 levels) following octreotide treatment withdrawal for at least 4 weeks

Renewal — (Acromegaly - pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has acromegaly: and
- 2 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 3 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

Initial application — (Other Indications) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 VIPomas and Glucagonomas for patients who are seriously ill in order to improve their clinical state prior to definitive surgery; or
- 2 Both:
 - 2.1 Gastrinoma: and
 - 2.2 Either:
 - 2.2.1 Patient has failed surgery; or
 - 2.2.2 Patient in metastatic disease after H2 antagonists (or proton pump inhibitors) have failed; or
- 3 Both:
 - 3.1 Insulinomas: and
 - 3.2 Surgery is contraindicated or has failed; or
- 4 For pre-operative control of hypoglycaemia and for maintenance therapy; or
- 5 Both:
 - 5.1 Carcinoid syndrome (diagnosed by tissue pathology and/or urinary 5HIAA analysis); and
 - 5.2 Disabling symptoms not controlled by maximal medical therapy.

Note: The use of octreotide in patients with fistulae, oesophageal varices, miscellaneous diarrhoea and hypotension will not be funded as a Special Authority item

Renewal — (Other Indications) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment. Initial application — (pre-operative acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has acromegaly; and
- 2 Patient has a large pituitary tumour, greater than 10 mm at its widest; and
- 3 Patient is scheduled to undergo pituitary surgery in the next six months.

TAMOXIFEN CITRATE

| * | Tab 10 mg15.00 | 60 | ✓ <u>Tamoxifen Sandoz</u> |
|---|----------------|----|---------------------------|
| | Tab 20 mg5.32 | 60 | ✓ <u>Tamoxifen Sandoz</u> |

Aromatase Inhibitors

| AN. | AS ⁻ | ΓR | ΟZ | OLE |
|-----|-----------------|----|----|-----|
|-----|-----------------|----|----|-----|

30 ✓ Anatrole

| | Subsidy (Manufacturer's Price) | | Fully Subsidised | Brand or Generic Manufacturer |
|-------------------------|-----------------------------------|----|---------------------|-------------------------------------|
| EXEMESTANE * Tab 25 mg | 9.86 | 30 | √ <u>P</u> | fizer Exemestane |
| LETROZOLE * Tab 2.5 mg | 4.67 | 30 | √ <u>L</u> | <u>etrole</u> |

Immunosuppressants

Cytotoxic Immunosuppressants

| AZATHIOPRINE | | |
|---|-----------|----------------------------|
| * Tab 25 mg7.36 | 60 | ✓ Azamun |
| * Tab 50 mg8.10 | 100 | ✓ Azamun |
| MYCOPHENOLATE MOFETIL | | |
| Tab 500 mg35.90 | 50 | Cellcept |
| Cap 250 mg35.90 | 100 | ✓ Cellcept |
| Powder for oral liq 1 g per 5 ml - Subsidy by endorsement187.25 | 165 ml OP | Cellcept |

Mycophenolate powder for oral liquid is subsidised only for patients unable to swallow tablets and capsules, and when the prescription is endorsed accordingly.

Fusion Proteins

| ETANERCEPT - Special Authority see SA2399 below - Ref | ail pharmacy | | |
|---|--------------|---|--------------------------|
| Inj 25 mg | 690.00 | 4 | Enbrel |
| Inj 25 mg autoinjector | 690.00 | 4 | Enbrel |
| Inj 50 mg autoinjector | | 4 | Enbrel |
| Inj 50 mg prefilled syringe | | 4 | Enbrel |

⇒SA2399 Special Authority for Subsidy

Initial application — (adult-onset Still's disease) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Fither:
 - 1.1.1 The patient has had an initial Special Authority approval for adalimumab for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a Health NZ Hospital; and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or tocilizumab; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or tocilizumab such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal anti-inflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 Fither:

| Subsidy (Manufacturer's Price) | Sub | Fully | Brand or Generic |
|-----------------------------------|-----|-------|---------------------|
| ` \$ | Per | ✓ | Manufacturer |

continued...

- 1.1 Applicant is a rheumatologist; or
- 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 The patient has a sustained improvement in inflammatory markers and functional status.

Initial application — **(ankylosing spondylitis)** only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for ankylosing spondylitis; and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for ankylosing spondylitis; or
- 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis present for more than six months; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.3 Patient has bilateral sacroillitis demonstrated by plain radiographs, CT or MRI scan; and
 - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
 - 2.5 Either:
 - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
 - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender (see Notes); and
 - 2.6 A Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of initial application.

Average normal chest expansion corrected for age and gender:

18-24 years - Male: 7.0 cm; Female: 5.5 cm 25-34 years - Male: 7.5 cm; Female: 5.5 cm 35-44 years - Male: 6.5 cm; Female: 4.5 cm 45-54 years - Male: 6.0 cm; Female: 5.0 cm 55-64 years - Male: 5.5 cm; Female: 4.0 cm 65-74 years - Male: 4.0 cm; Female: 4.0 cm

75+ years - Male: 3.0 cm; Female: 2.5 cm

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less;

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

continued...

and

- 3 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 4 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (polyarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for polyarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
 - 2.3 Any of the following:
 - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose): or
 - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Renewal — (polyarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Fither:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for oligoarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for oligoarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and

| Subsidy | | Fully | Brand or | |
|------------------------|-----|------------|--------------|--|
| (Manufacturer's Price) | S | Subsidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

continued...

- 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
- 2.3 Any of the following:
 - 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.3 High disease activity (cJADAS10 score greater than 4) after a 6-month trial of methotrexate.

Renewal — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab or secukinumab for psoriatic arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab or secukinumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab or secukinumab to meet the renewal criteria for adalimumab or secukinumab for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
 - 2.4 Either:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints;
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

| Subsidy | | Fully | Brand or | |
|------------------------|-----|-----------|--------------|--|
| (Manufacturer's Price) | Sı | ıbsidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

continued...

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
 - 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
 - 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Indications marked with * are unapproved indications.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement: and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

Initial application — (Arthritis - rheumatoid) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for rheumatoid arthritis; and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects; or
 - 1.2.2 The patient has received insufficient benefit to meet the renewal criteria for rheumatoid arthritis; or
 - 2 All of the following:
 - 2.1 Patient has had rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
 - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.3 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
 - 2.4 Patient has tried and not responded to at least three months of methotrexate in combination with sulfasalazine and hydroxychloroguine sulphate (at maximum tolerated doses unless contraindicated); and
 - 2.5 Either:
 - 2.5.1 Patient has tried and not responded to at least three months of methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 2.5.2 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with methotrexate; and
 - 2.6 Either:

| | Subsidy | Fully | Brand or |
|-------|-------------------|------------|--------------|
| (Manu | facturer's Price) | Subsidised | Generic |
| | \$ Pe | er 🗸 | Manufacturer |

continued...

- 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
- 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip.

Renewal — (Arthritis - rheumatoid) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (severe chronic plaque psoriasis) only from a dermatologist or any relevant practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria: Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe chronic plaque psoriasis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe chronic plaque psoriasis; or
- 2 All of the following:
 - 2.1 Any of the following:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.3 Patient has severe chronic localised genital or flexural plaque psoriasis where the plaques or lesions have been present for at least 6 months from the time of initial diagnosis, and with a Dermatology Life Quality Index (DLQI) score greater than 10; and
 - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
 - 2.3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
 - 2.4 The most recent PASI or DLQI assessment is no more than 1 month old at the time of application.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand, foot, genital or flexural areas at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and for the face, palm of a hand or sole of a foot the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis) from any relevant practitioner. Approvals valid for 6 months for applications

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

continued...

meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 Both:
 - 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 1.1.2 Either:
 - 1.1.2.1 Following each prior etanercept treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-treatment baseline value; or
 - 1.1.2.2 Following each prior etanercept treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline value; or
 - 1.2 Both:
 - 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 1.2.2 Either:
 - 1.2.2.1 Following each prior etanercept treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 1.2.2.2 Following each prior etanercept treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; or
 - 1.3 Both:
 - 1.3.1 Patient had severe chronic localised genital or flexural plague psoriasis at the start of treatment; and
 - 1.3.2 Either:
 - 1.3.2.1 The patient has experienced a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; or
 - 1.3.2.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing etanercept; and
- 2 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Note: A treatment course is defined as a minimum of 12 weeks of etanercept treatment

Initial application — (undifferentiated spondyloarthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undifferentiated peripheral spondyloarthritis* with active peripheral joint arthritis in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day (or maximum tolerated dose); and
- 4 Patient has tried and not responded to at least three months of leflunomide at a dose of up to 20 mg daily (or maximum tolerated dose); and
- 5 Any of the following:
 - 5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application: or
 - 5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour measured no more than one month prior to the date of this application; or
 - 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Note: Indications marked with * are unapproved indications.

| Subsidy | | Fully | Brand or | |
|------------------------|-----|-----------|--------------|--|
| (Manufacturer's Price) | Sı | ubsidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

continued...

Renewal — (undifferentiated spondyloarthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Fither:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Fither:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg dose every 7 days.

Immune Modulators

| ANTITHYMOCYTE GLOBULIN (EQUINE) - PCT only - Special | ist | | |
|--|--------------|---|--------------------|
| Inj 50 mg per ml, 5 ml | 2,774.48 | 5 | ✓ ATGAM |
| BACILLUS CALMETTE-GUERIN (BCG) VACCINE - PCT only - | - Specialist | | |
| Subsidised only for bladder cancer. | | | |
| Inj 2-8 × 100 million CFU | 149.37 | 1 | ✓ OncoTICE |
| Inj 40 mg per ml, vial | 176.90 | 3 | ✓ SII-Onco-BCG S29 |

Monoclonal Antibodies

| | | i pnarmacy | ai Authority see SA2400 delow - | ADALII |
|------------------------------|-----|------------|---------------------------------|--------|
| ✓ Amgevita | 1 🗸 | 0 | ge | Inj |
| Amgevita | 2 | 0 | - | Inj |
| ✓ Amgevita | 2 | 0 | ge | lnj |

⇒SA2400 Special Authority for Subsidy

Initial application — (Behcet's disease - severe) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Both:

- 1 The patient has severe Behcet's disease* that is significantly impacting the patient's quality of life; and
 - 2 Fither:
 - 2.1 The patient has severe ocular, neurological, and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s); or
 - 2.2 The patient has severe gastrointestinal, rheumatological, and/or mucocutaneous symptoms and has not responded adequately to two or more treatments appropriate for the particular symptom(s).

Note: Indications marked with * are unapproved indications.

Initial application — (Hidradenitis suppurativa) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has hidradenitis suppurativa Hurley Stage II or Hurley Stage III lesions in distinct anatomic areas; and
- 2 Patient has tried, but had an inadequate response to at least a 90 day trial of systemic antibiotics or has demonstrated intolerance to or has contraindications for systemic antibiotics; and
- 3 Patient has 3 or more active lesions; and

| Subsidy | | Fully | Brand or | |
|------------------------|-----|-----------|--------------|--|
| (Manufacturer's Price) | S | ubsidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

continued...

4 The patient has a DLQI of 10 or more and the assessment is no more than 1 month old at time of application.

Renewal — (Hidradenitis suppurativa) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline; and
- 2 The patient has a DLQI improvement of 4 or more from baseline.

Initial application — (Plaque psoriasis - severe chronic) only from a dermatologist or any relevant practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria: Either:

- 1 Both:
 - 1.1 Patient has had an initial Special Authority approval for etanercept for severe chronic plague psoriasis; and
 - 12 Fither
 - 1.2.1 Patient has experienced intolerable side effects; or
 - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for etanercept for severe chronic plaque psoriasis; or
- 2 All of the following:
 - 2.1 Any of the following:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a PASI score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.3 Patient has severe chronic localised genital or flexural plaque psoriasis where the plaques or lesions have been present for at least 6 months from the time of initial diagnosis, and with a Dermatology Life Quality Index (DLQI) score greater than 10; and
 - 2.2 Patient has tried, but had an inadequate response to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
 - 2.3 A PASI assessment or DLQI assessment has been completed for at least the most recent prior treatment course but no longer than 1 month following cessation of each prior treatment course and is no more than 1 month old at the time of application.

Renewal — (Plaque psoriasis - severe chronic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 Patient had "whole body" severe chronic plague psoriasis at the start of treatment; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced a 75% or more reduction in PASI score, or is sustained at this level, when compared with the pre-treatment baseline value; or
 - 1.2.2 The patient has a DLQI improvement of 5 or more, when compared with the pre-treatment baseline value; or
- 2 Both:
 - 2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 2.2 Either:
 - 2.2.1 The patient has experienced reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or

| Subsidy (Manufacturer's Price) | F Subsidi | ully | Brand or Generic |
|-----------------------------------|--------------|----------|---------------------|
| \$ | Per | √ | Manufacturer |

continued...

2.2.2 The patient has experienced reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre treatment baseline value; or

- 3 Both:
 - 3.1 Patient had severe chronic localised genital or flexural plague psoriasis at the start of treatment; and
 - - 3.2.1 The patient has experienced a reduction of 75% or more in the skin area affected, or sustained at this level. as compared to the pre-treatment baseline value; or
 - 3.2.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing adalimumab.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid without further renewal unless notified for applications meeting the following criteria: Both:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporin, azathioprine, or methotrexate) and has not received an adequate response.

Note: Indications marked with * are unapproved indications.

Initial application — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active Crohn's disease: and
- 2 Any of the following:
 - 2.1 Patient has a CDAI score of greater than or equal to 300, or HBI score of greater than or equal to 10; or
 - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection;
 - 2.4 Patient has an ileostomy or colostomy and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Renewal — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 CDAI score has reduced by 100 points from the CDAI score, or HBI score has reduced by 3 points, from when the patient was initiated on adalimumab; or
- 2 CDAI score is 150 or less, or HBI is 4 or less; or
- 3 The patient has demonstrated an adequate response to treatment, but CDAI score and/or HBI score cannot be assessed.

Initial application — (Crohn's disease - children) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has active Crohn's disease; and
- 2 Fither:
 - 2.1 Patient has a PCDAI score of greater than or equal to 30; or
 - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Renewal — (Crohn's disease - children) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

| Subsidy | | Fully | Brand or |
|------------------------|-----|-----------|--------------|
| (Manufacturer's Price) | Sı | ubsidised | Generic |
| \$ | Per | ✓ | Manufacturer |

continued...

- 1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab; or
- 2 PCDAI score is 15 or less: or
- 3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed.

Initial application — (Crohn's disease - fistulising) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
 - 2.2 Patient has one or more rectovaginal fistula(e); or
 - 2.3 Patient has complex peri-anal fistula; and
- 3 A Baseline Fistula Assessment has been completed and is no more than 1 month old at the time of application.

Renewal — (Crohn's disease - fistulising) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 The number of open draining fistulae have decreased from baseline by at least 50%; or
- 2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain.

Initial application — (Ocular inflammation - chronic) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 The patient has had an initial Special Authority approval for infliximab for chronic ocular inflammation; or
- 2 Both:
 - 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
 - 2.2 Any of the following:
 - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective; or
 - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
 - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Renewal — (Ocular inflammation - chronic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 12 weeks' initial treatment; or
- 2 Following each 2 year treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 2 year treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Initial application — **(Ocular inflammation - severe)** from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 Patient has had an initial Special Authority approval for infliximab for severe ocular inflammation; or
- 2 Both:

| Subsidy | | Fully | Brand or | |
|------------------------|-----|----------|--------------|--|
| (Manufacturer's Price) | Su | bsidised | Generic | |
| \$ | Per | / | Manufacturer | |

continued...

- 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
- 2.2 Any of the following:
 - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
 - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
 - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Renewal — (Ocular inflammation - severe) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 2 year treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
- 3 Following each 2 year treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Initial application — (ankylosing spondylitis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has had an initial Special Authority approval for etanercept for ankylosing spondylitis; and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects; or
 - 1.2.2 The patient has received insufficient benefit to meet the renewal criteria for ankylosing spondylitis; or
- 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis for more than six months; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.3 Patient has bilateral sacroiliitis demonstrated by radiology imaging; and
 - 2.4 Patient has not responded adequately to treatment with two or more NSAIDs, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
 - 2.5 Either:
 - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following BASMI measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
 - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender; and
 - 2.6 A BASDAI of at least 6 on a 0-10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment and is no more than 1 month old at the time of application.

Renewal — (ankylosing spondylitis) from any relevant practitioner. Approvals valid for 2 years where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less.

Initial application — (Arthritis - oligoarticular course juvenile idiopathic) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for oligoarticular course juvenile idiopathic arthritis (JIA): and

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

- 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects; or
 - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for oligoarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
 - 2.3 Either:
 - 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose).

Renewal — (Arthritis - oligoarticular course juvenile idiopathic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - polyarticular course juvenile idiopathic) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has had an initial Special Authority approval for etanercept for polyarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects; or
 - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for polyarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
 - 2.3 Any of the following:
 - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Renewal — (Arthritis - polyarticular course juvenile idiopathic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Fither:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - psoriatic) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

| Subsidy | F | ully | Brand or | |
|------------------------|----------|------|--------------|--|
| (Manufacturer's Price) | Subsidis | sed | Generic | |
| \$ | Per | ✓ | Manufacturer | |

continued...

Either:

- 1 Both:
 - 1.1 Patient has had an initial Special Authority approval for etanercept or secukinumab for psoriatic arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects; or
 - 1.2.2 The patient has received insufficient benefit from to meet the renewal criteria for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine or leflunomide at maximum tolerated doses (unless contraindicated); and
 - 2.4 Fither:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.5 Any of the following:
 - 2.5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an ESR greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (Arthritis - psoriatic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Fither:

- 1 Following initial treatment, the patient has at least a 50% decrease in swollen joint count from baseline and a clinically significant response in the opinion of the physician; or
- 2 Patient demonstrates at least a continuing 30% improvement in swollen joint count from baseline and a clinically significant response in the opinion of the treating physician.

Initial application — (Arthritis - rheumatoid) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for rheumatoid arthritis; and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for rheumatoid arthritis: or
- 2 All of the following:
 - 2.1 Patient has had rheumatoid arthritis (either confirmed by radiology imaging, or the patient is CCP antibody positive) for six months duration or longer; and
 - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.3 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
 - 2.4 Patient has tried and not responded to at least three months of methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate at maximum tolerated doses (unless contraindicated); and

| Subsidy (Manufacturer's Price) | Ful Subsidise | d Generic | |
|-----------------------------------|------------------|--------------|--|
| • | Per • | Manufacturer | |

continued...

- 2.5 Either:
 - 2.5.1 Patient has tried and not responded to at least three months of methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 2.5.2 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with methotrexate; and
- 2.6 Fither:
 - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
 - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip.

Renewal — (Arthritis - rheumatoid) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Initial application — (Still's disease - adult-onset (AOSD)) only from a rheumatologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept and/or tocilizumab for AOSD; and
 - 1.2 Fither:
 - 1.2.1 Patient has experienced intolerable side effects from etanercept and/or tocilizumab; or
 - 1.2.2 Patient has received insufficient benefit from at least a three-month trial of etanercept and/or tocilizumab; or
 - 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria; and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, NSAIDs and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Initial application — (ulcerative colitis) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active ulcerative colitis; and
- 2 Either:
 - 2.1 Patient's SCCAI score is greater than or equal to 4; or
 - 2.2 Patient's PUCAI score is greater than or equal to 20; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from prior therapy with immunomodulators and systemic corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Fither:

- 1 The SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on biologic therapy: or
- 2 The PUCAI score has reduced by 10 points or more from the PUCAI score when the patient was initiation on biologic therapy.

| Subsidy | | Fully | Brand or |
|------------------------|-----|---------|--------------|
| (Manufacturer's Price) | Sub | sidised | Generic |
| \$ | Per | • | Manufacturer |

continued...

Initial application — (undifferentiated spondyloarthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undifferentiated peripheral spondyloarthritis* with active peripheral joint arthritis in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2 Patient has tried and not responded to at least three months of each of methotrexate, sulfasalazine and leflunomide, at maximum tolerated doses (unless contraindicated); and
- 3 Any of the following:
 - 3.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application: or
 - 3.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application; or
 - 3.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Note: Indications marked with * are unapproved indications

Renewal — (undifferentiated spondyloarthritis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response in the opinion of the treating physician.

Initial application — (inflammatory bowel arthritis – axial) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has axial inflammatory pain for six months or more; and
- 3 Patient is unable to take NSAIDs: and
- 4 Patient has unequivocal sacroiliitis demonstrated by radiological imaging or MRI; and
- 5 Patient has not responded adequately to prior treatment consisting of at least 3 months of an exercise regime supervised by a physiotherapist; and
- 6 A BASDAI of at least 6 on a 0-10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment.

Renewal — (inflammatory bowel arthritis – axial) from any relevant practitioner. Approvals valid for 2 years where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less.

Initial application — (inflammatory bowel arthritis – peripheral) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has active arthritis in at least four joints from the following: hip, knee, ankle, subtalar, tarsus, forefoot, wrist, elbow, shoulder, sternoclavicular; and
- 3 Patient has tried and not experienced a response to at least three months of methotrexate, or azathioprine at a maximum tolerated dose (unless contraindicated); and
- 4 Patient has tried and not experienced a response to at least three months of sulfasalazine at a maximum tolerated dose (unless contraindicated); and
- 5 Any of the following:
 - 5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this

| (Manufacturer 5 Files) Substitused General General Substitution Substitution Substitution General General Control of the Substitution | | Subsidy Manufacturer's Price) | Fully Subsidise | |
|--|----|----------------------------------|--------------------|--|
| | Ţ, | \$ | | |

continued...

application: or

- 5.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application; or
- 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (inflammatory bowel arthritis – peripheral) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 Patient has experienced at least a continuing 30% improvement in active joint count from baseline in the opinion of the treating physician.

ADALIMUMAB (HUMIRA - ALTERNATIVE BRAND) - Special Authority see SA2157 below - Retail pharmacy

| Inj 20 mg per 0.2 ml prefilled syringe | 1,599.96 | 2 | Humira |
|--|----------|---|--------------------------|
| Inj 40 mg per 0.4 ml prefilled pen | 1,599.96 | 2 | ✓ HumiraPen |
| Inj 40 mg per 0.4 ml prefilled syringe | 1,599.96 | 2 | Humira |

⇒SA2157 Special Authority for Subsidy

Initial application — (Behcet's disease – severe) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Behcet's disease – severe) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient has had a good clinical response to treatment with measurably improved quality of life; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Hidradenitis suppurativa) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 7 days. Fortnightly dosing has been considered.

Renewal — (Hidradenitis suppurativa) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

| Subsidy (Manufacturer's Price) | Subs | Fully | Brand or Generic | |
|-----------------------------------|------|-------|---------------------|--|
| \$ | Per | 1 | Manufacturer | |

continued...

- 1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline; and
- 2 The patient has a Dermatology Quality of Life Index improvement of 4 or more from baseline; and
- 3 Adalimumab is to be administered at doses no greater than 40mg every 7 days. Fortnightly dosing has been considered.

Initial application — (Psoriasis - severe chronic plaque) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Psoriasis - severe chronic plaque) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Both:
 - 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 1.1.2 Either:
 - 1.1.2.1 Following each prior adalimumab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-adalimumab treatment baseline value; or
 - 1.1.2.2 Following each prior adalimumab treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline value; or
 - 1.2 Both:
 - 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 1.2.2 Either:
 - 1.2.2.1 Following each prior adalimumab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 1.2.2.2 Following each prior adalimumab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-adalimumab treatment baseline value: and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Pyoderma gangrenosum) only from a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and

| Subsidy | Fully | Brand or | |
|------------------------|------------|--------------|--|
| (Manufacturer's Price) | Subsidised | Generic | |
| \$ | Per 🗸 | Manufacturer | |

continued...

- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 A maximum of 8 doses.

Renewal — (Pyoderma gangrenosum) only from a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient has demonstrated clinical improvement and continues to require treatment; and
- 2 A maximum of 8 doses.

Initial application — (Crohn's disease - adult) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevitat; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
 - 1.3 Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Crohn's disease - adult) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on adalimumab; or
 - 1.2 CDAI score is 150 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment, but CDAI score cannot be assessed; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Crohn's disease - children) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
 - 1.3 Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Crohn's disease - children) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab; or
 - 1.2 PCDAI score is 15 or less: or

| Subsidy (Manufacturer's Price) | F Subsidi | ully | Brand or Generic |
|-----------------------------------|--------------|----------|---------------------|
| \$ | Per | √ | Manufacturer |

continued...

- 1.3 The patient has demonstrated an adequate response to treatment, but PCDAI score cannot be assessed; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Crohn's disease - fistulising) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
 - 1.3 Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Crohn's disease - fistulising) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
 - 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Ocular inflammation – chronic) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with Amgevita, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
 - 1.3 Patient has uveitis and is considered to be at risk of vision loss if they were to change treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Ocular inflammation – chronic) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has had a good clinical response following 12 weeks' initial treatment; or
 - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
 - 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and</p>
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

| Subsidy | | Fully | Brand or | |
|------------------------|-----|------------|--------------|--|
| (Manufacturer's Price) | S | Subsidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

continued...

Initial application — (Ocular inflammation – severe) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with Amgevita, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
 - 1.3 Patient has uveitis and is considered to be at risk of vision loss if they were to change treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Ocular inflammation – severe) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has had a good clinical response following 3 initial doses: or
 - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < 1/2+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
 - 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita): and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale. or an improvement in BASDAI of 50%, whichever is less; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Arthritis - oligoarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or

| Subsidy | | Fully | Brand or | |
|------------------------|-----|------------|--------------|--|
| (Manufacturer's Price) | | Subsidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

continued...

- 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

Renewal — (Arthritis – oligoarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months where the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - polyarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

Renewal — (Arthritis - polyarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months where the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from

Initial application — (Arthritis - psoriatic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Fither:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Arthritis - psoriatic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Arthritis - rheumatoid) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Fither:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

- 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Fither:
 - 4.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or
 - 4.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

Renewal — (Arthritis – rheumatoid) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 2 Fither:
 - 2.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or
 - 2.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

Initial application — (Still's disease – adult-onset (AOSD)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

Renewal — (Still's disease – adult-onset (AOSD)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months where the patient has demonstrated a sustained improvement in inflammatory markers and functional status

AFLIBERCEPT - Special Authority see SA1772 below - Retail pharmacy

⇒SA1772 Special Authority for Subsidy

Initial application — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 3 months for applications meeting the following criteria:

Fither:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Wet age-related macular degeneration (wet AMD); or
 - 1.1.2 Polypoidal choroidal vasculopathy; or
 - 1.1.3 Choroidal neovascular membrane from causes other than wet AMD; and
 - 1.2 Either:
 - 1.2.1 The patient has developed severe endophthalmitis or severe posterior uveitis following treatment with bevacizumab: or
 - 1.2.2 There is worsening of vision or failure of retina to dry despite three intraocular injections of bevacizumab four weeks apart; and

| Subsidy | F | ılly | Brand or | |
|------------------------|----------|------|--------------|--|
| (Manufacturer's Price) | Subsidis | sed | Generic | |
| \$ | Per | ✓ | Manufacturer | |

continued...

- 1.3 There is no structural damage to the central fovea of the treated eye; and
- 1.4 Patient has not previously been treated with ranibizumab for longer than 3 months; or
- 2 Fither:
 - 2.1 Patient has current approval to use ranibizumab for treatment of wAMD and was found to be intolerant to ranibizumab within 3 months; or
 - 2.2 Patient has previously* (*before June 2018) received treatment with ranibizumab for wAMD and disease was stable while on treatment.

Initial application — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has centre involving diabetic macular oedema (DMO); and
- 2 Patient's disease is non responsive to 4 doses of intravitreal bevacizumab when administered 4-6 weekly; and
- 3 Patient has reduced visual acuity between 6/9 6/36 with functional awareness of reduction in vision; and
- 4 Patient has DMO within central OCT (ocular coherence tomography) subfield > 350 micrometers; and
- 5 There is no centre-involving sub-retinal fibrosis or foveal atrophy.

Renewal — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Documented benefit must be demonstrated to continue; and
- 2 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 3 There is no structural damage to the central fovea of the treated eye.

Renewal — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 There is stability or two lines of Snellen visual acuity gain; and
- 2 There is structural improvement on OCT scan (with reduction in intra-retinal cysts, central retinal thickness, and sub-retinal fluid): and
- 3 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 4 There is no centre-involving sub-retinal fibrosis or foveal atrophy; and
- 5 After each consecutive 12 months treatment with (2nd line anti-VEGF agent), patient has retrialled with at least one injection of bevacizumab and had no response.

BENRALIZUMAB - Special Authority see SA2151 below - Retail pharmacy

⇒SA2151 Special Authority for Subsidy

Initial application — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Patient must have a diagnosis of severe eosinophilic asthma documented by a respiratory physician or clinical immunologist; and
- 3 Conditions that mimic asthma eg. vocal cord dysfunction, central airway obstruction, bronchiolitis etc. have been excluded; and
- 4 Patient has a blood eosinophil count of greater than 0.5 x 10^9 cells/L in the last 12 months; and
- 5 Patient must be adherent to optimised asthma therapy including inhaled corticosteroids (equivalent to at least 1000 mcg per day of fluticasone propionate) plus long-acting beta-2 agonist, or budesonide/formoterol as part of the anti-inflammatory reliever therapy plus maintenance regimen, unless contraindicated or not tolerated; and

| Subsidy | | Fully | Brand or | |
|------------------------|-----|---------|--------------|--|
| (Manufacturer's Price) | Sub | sidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

continued...

- 6 Either:
 - 6.1 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral corticosteroids; or
 - 6.2 Patient has received continuous oral corticosteroids of at least the equivalent of 10 mg per day over the previous 3 months: and
- 7 Treatment is not to be used in combination with subsidised mepolizumab; and
- 8 Patient has an Asthma Control Test (ACT) score of 10 or less. Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 52 weeks after the first dose to assess response to treatment; and
- 9 Either:
 - 9.1 Patient has not previously received an anti-IL5 biological therapy for their severe eosinophilic asthma; or
 - 9.2 Both:
 - 9.2.1 Patient was refractory or intolerant to previous anti-IL5 biological therapy; and
 - 9.2.2 Patient was not eligible to continue treatment with previous anti-IL5 biological therapy and discontinued within 12 months of commencing treatment.

Renewal — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 Either:
 - 2.1 Exacerbations have been reduced from baseline by 50% as a result of treatment with benralizumab; or
 - 2.2 Reduction in continuous oral corticosteroid use by 50% or by 10 mg/day while maintaining or improving asthma control.

1 Adcetris

⇒SA2289 Special Authority for Subsidy

Initial application — (relapsed/refractory Hodgkin lymphoma) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Both:
 - 1.1.1 Patient has relapsed/refractory CD30-positive Hodgkin lymphoma after two or more lines of chemotherapy; and
 - 1.1.2 Patient is ineligible for autologous stem cell transplant; or
 - 1.2 Both:
 - 1.2.1 Patient has relapsed/refractory CD30-positive Hodgkin lymphoma; and
 - 1.2.2 Patient has previously undergone autologous stem cell transplant; and
- 2 Patient has not previously received funded brentuximab vedotin; and
- 3 Response to brentuximab vedotin treatment is to be reviewed after a maximum of 6 treatment cycles; and
- 4 Brentuximab vedotin to be administered at doses no greater than 1.8 mg/kg every 3 weeks.

Renewal — (relapsed/refractory Hodgkin lymphoma) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has achieved a partial or complete response to brentuximab vedotin after 6 treatment cycles; and
- 2 Treatment remains clinically appropriate and the patient is benefitting from treatment and treatment is being tolerated; and

| Subsidy (Manufacturer's Price) | | Fully Subsidised | Brand or Generic |
|-----------------------------------|-----|---------------------|---------------------|
| \$ | Per | ✓ | Manufacturer |

continued...

3 Patient is to receive a maximum of 16 total cycles of brentuximab vedotin treatment.

Initial application — (anaplastic large cell lymphoma) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has relapsed/refractory CD30-positive systemic anaplastic large cell lymphoma; and
- 2 Patient has an ECOG performance status of 0-1; and
- 3 Patient has not previously received brentuximab vedotin; and
- 4 Response to brentuximab vedotin treatment is to be reviewed after a maximum of 6 treatment cycles; and
- 5 Brentuximab vedotin to be administered at doses no greater than 1.8 mg/kg every 3 weeks.

Renewal — (anaplastic large cell lymphoma) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has achieved a partial or complete response to brentuximab vedotin after 6 treatment cycles; and
- 2 Treatment remains clinically appropriate and the patient is benefitting from treatment and treatment is being tolerated; and
- 3 Patient is to receive a maximum of 16 total cycles of brentuximab vedotin treatment.

CASIRIVIMAB AND IMDEVIMAB - [Xpharm] - Special Authority see \$A2096 below

⇒SA2096 Special Authority for Subsidy

Initial application — (Treatment of profoundly immunocompromised patients) from any relevant practitioner. Approvals valid for 2 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed (or probable) COVID-19; and
- 2 The patient is in the community with mild to moderate disease severity*; and
- 3 Patient is profoundly immunocompromised** and is at risk of not having mounted an adequate response to vaccination against COVID-19 or is unvaccinated; and
- 4 Patient's symptoms started within the last 10 days; and
- 5 Patient is not receiving high flow oxygen or assisted/mechanical ventilation; and
- 6 Casirivimab and imdevimab is to be administered at a maximum dose of no greater than 2,400 mg.

Notes: * Mild to moderate disease severity as described on the Ministry of Health Website

** Examples include B-cell depletive illnesses or patients receiving treatment that is B-Cell depleting.

CETUXIMAB - PCT only - Specialist - Special Authority see SA2401 below

| Inj 5 mg per ml, 20 ml vial | 364.00 | 1 | Erbitux |
|------------------------------|----------|------|-----------|
| Inj 5 mg per ml, 100 ml vial | 1,820.00 | 1 | ✓ Erbitux |
| Inj 1 mg for ECP | 3.82 | 1 mg | ✓ Baxter |

⇒SA2401 Special Authority for Subsidy

Initial application — (head and neck cancer, locally advanced) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, non-metastatic, squamous cell cancer of the head and neck; and
- 2 Cisplatin is contraindicated or has resulted in intolerable side effects; and
- 3 Patient has an ECOG performance score of 0-2; and
- 4 To be administered in combination with radiation therapy.

Initial application — (colorectal cancer, metastatic) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

| Subsidy | F | ılly | Brand or | |
|------------------------|----------|------|--------------|--|
| (Manufacturer's Price) | Subsidis | ed | Generic | |
| \$ | Per | ✓ | Manufacturer | |

continued...

- 1 Patient has metastatic colorectal cancer located on the left side of the colon (see Note); and
- 2 There is documentation confirming disease is RAS and BRAF wild-type; and
- 3 Patient has an ECOG performance score of 0-2; and
- 4 Patient has not received prior funded treatment with cetuximab; and
- 5 Either:
 - 5.1 Cetuximab is to be used in combination with chemotherapy; or
 - 5.2 Chemotherapy is determined to not be in the best interest of the patient based on clinician assessment.

Renewal — (colorectal cancer, metastatic) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where there is no evidence of disease progression.

Note: Left-sided colorectal cancer comprises of the distal one-third of the transverse colon, the splenic flexure, the descending colon, the sigmoid colon, or the rectum.

GEMTUZUMAB OZOGAMICIN − PCT only − Specialist − Special Authority see SA2269 below Inj 5 mg vial12,973.00 1 ✓ Mylotarg

⇒SA2269 Special Authority for Subsidy

Initial application only from a haematologist, paediatric haematologist or paediatric oncologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has not received prior chemotherapy for this condition; and
- 2 Patient has de novo CD33-positive acute myeloid leukaemia; and
- 3 Patient does not have acute promyelocytic leukaemia; and
- 4 Gemtuzumab ozogamicin will be used in combination with standard anthracycline and cytarabine (AraC); and
- 5 Patient is being treated with curative intent; and
- 6 Patient's disease risk has been assessed by cytogenetic testing to be good or intermediate; and
- 7 Patient must be considered eligible for standard intensive remission induction chemotherapy with standard anthracycline and cytarabine (AraC); and
- 8 Gemtuzumab ozogamicin to be funded for one course only (one dose at 3 mg per m² body surface area or up to 2 vials of 5 mg as separate doses).

Note: Acute myeloid leukaemia excludes acute promyelocytic leukaemia and acute myeloid leukaemia that is secondary to another haematological disorder (eg myelodysplasia or myeloproliferative disorder).

⇒SA2402 Special Authority for Subsidy

Initial application — (Crohn's disease (adults)) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has a CDAI score of greater than or equal to 300 or HBI score of greater than or equal to 10; or
 - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
 - 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but has experienced an inadequate response to, or intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Renewal — (Crohn's disease (adults)) from any relevant practitioner. Approvals valid for 2 years for applications meeting the

| Subsidy | Fully Subsidised | | Brand or |
|----------------------------------|---------------------|------------------|-------------------------|
| (Manufacturer's Price) \$ | Per | isea ✓ | Generic Manufacturer |

continued...

following criteria:

Both:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points from the CDAI score, or HBI score has reduced by 3 points, from when the patient was initiated on infliximab; or
 - 1.2 CDAI score is 150 or less, or HBI is 4 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment but CDAI score and/or HBI score cannot be assessed: and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (Crohn's disease (children)) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has active Crohn's disease; and
- 2 Either:
 - 2.1 Patient has a PCDAI score of greater than or equal to 30; or
 - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but experienced an inadequate response to, or intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Renewal — (Crohn's disease (children)) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on infliximab; or
 - 1.2 PCDAI score is 15 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (**Graft vs host disease**) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has steroid-refractory acute graft vs. host disease of the qut.

Initial application — (Pulmonary sarcoidosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has life-threatening pulmonary sarcoidosis diagnosed by a multidisciplinary team that is refractory to other treatments.

Initial application — (acute fulminant ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient has acute, fulminant ulcerative colitis; and
- 2 Treatment with intravenous or high dose oral corticosteroids has not been successful.

Initial application — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and
- 2 Fither

| Subsidy | Full | / Brand or | |
|------------------------|-----------|---------------------------|--|
| (Manufacturer's Price) | Subsidise | d Generic | |
| \$ | Per 🗸 | Manufacturer Manufacturer | |

continued...

- 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Following 12 weeks of infliximab treatment, BASDAI has improved by 4 or more points from pre-infliximab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 3 Infliximab to be administered at doses no greater than 5 mg/kg every 6-8 weeks.

Initial application — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for chronic ocular inflammation; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for chronic ocular inflammation; or
- 2 Both:
 - 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
 - 2.2 Any of the following:
 - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective; or
 - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
 - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Renewal — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (fistulising Crohn's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has confirmed Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or

| Subsidy (Manufacturer's Pri | ice) | Fully Subsidised | Brand or Generic | |
|--------------------------------|------|---------------------|---------------------|--|
| \$ | Per | ✓ | Manufacturer | |

continued...

- 2.2 Patient has one or more rectovaginal fistula(e); or
- 2.3 Patent has complex peri-anal fistula.

Renewal — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
 - 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline (in the case of adult patients, as demonstrated by a reduction in the Fistula Assessment score), together with less induration and patient reported pain; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (neurosarcoidosis) only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria:

- All of the following:
 - 1 Patient has been diagnosed with neurosarcojosis by a multidisciplinary team; and
 - 2 Patient has CNS involvement; and
 - 3 Patient has steroid-refractory disease; and
 - 4 Either:
 - 4.1 IV cyclophosphamide has been tried; or
 - 4.2 Treatment with IV cyclophosphamide is clinically inappropriate.

Renewal — (neurosarcoidosis) only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria:

Either:

- 1 A withdrawal period has been tried and the patient has relapsed; or
- 2 All of the following:
 - 2.1 A withdrawal period has been considered but would not be clinically appropriate; and
 - 2.2 There has been a marked reduction in prednisone dose; and
 - 2.3 Either:
 - 2.3.1 There has been an improvement in MRI appearances; or
 - 2.3.2 Marked improvement in other symptomology.

Initial application — (plaque psoriasis) only from a dermatologist or any relevant practitioner on the recommendation of a dermatologist. Approvals valid for 3 months for applications meeting the following criteria:

Fither:

Eitner:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab, etanercept or secukinumab for severe chronic plaque psoriasis; and
 - 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects from adalimumab, etanercept or secukinumab; or
 - 1.2.2 Patient has received insufficient benefit from adalimumab, etanercept or secukinumab to meet the renewal criteria for adalimumab, etanercept or secukinumab for severe chronic plaque psoriasis; or
- 2 All of the following:
 - 2.1 Any of the following:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial

| \$ Per ✔ Manufacturer | | Subsidy (Manufacturer's Price) \$ | | Fully Subsidised | Brand or Generic Manufacturer |
|-----------------------|--|---|--|---------------------|-------------------------------------|
|-----------------------|--|---|--|---------------------|-------------------------------------|

continued...

- diagnosis; or
- 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; or
- 2.1.3 Patient has severe chronic localised genital or flexural plaque psoriasis where the plaques or lesions have been present for at least 6 months from the time of initial diagnosis, and with a Dermatology Life Quality Index (DLQI) score greater than 10; and
- 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
- 2.3 A PASI assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course: and
- 2.4 The most recent PASI assessment is no more than 1 month old at the time of initiation.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand, foot, genital or flexural areas at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and for the face, palm of a hand or sole of a foot the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (plaque psoriasis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 Both:
 - 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 1.1.2 Following each prior infliximab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-infliximab treatment baseline value; or
 - 1.2 Both:
 - 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 1.2.2 Either:
 - 1.2.2.1 Following each prior infliximab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 1.2.2.2 Following each prior infliximab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-infliximab treatment baseline value: or
 - 1.3 Both:
 - 1.3.1 Patient had severe chronic localised genital or flexural plaque psoriasis at the start of treatment; and
 - 1.3.2 Either:
 - 1.3.2.1 The patient has experienced a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; or
 - 1.3.2.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing infliximab; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

continued...

Both:

- 1 Patient was being treated with infliximab prior to 1 February 2019; and
- 2 Any of the following:
 - 2.1 Rheumatoid arthritis: or
 - 2.2 Ankylosing spondylitis; or
 - 2.3 Psoriatic arthritis: or
 - 2.4 Severe ocular inflammation; or
 - 2.5 Chronic ocular inflammation: or
 - 2.6 Crohn's disease (adults); or
 - 2.7 Crohn's disease (children): or
 - 2.8 Fistulising Crohn's disease; or
 - 2.9 Severe fulminant ulcerative colitis; or
 - 2.10 Severe ulcerative colitis: or
 - 2.11 Plague psoriasis; or
 - 2.12 Neurosarcoidosis: or
 - 2.13 Severe Behcet's disease.

Initial application — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis; and
- 2 Fither:
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept and/or secukinumab; or
 - 2.2 Following 3-4 months' initial treatment with adalimumab and/or etanercept and/or secukinumab, the patient did not meet the renewal criteria for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Either:
 - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior infliximab treatment in the opinion of the treating physician; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept; and
- 3 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance.

Renewal — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

| Subsidy | | Fully | Brand or |
|------------------------|-----|-----------|--------------|
| (Manufacturer's Price) | Sı | ubsidised | Generic |
| \$ | Per | ✓ | Manufacturer |

continued...

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Infliximab to be administered at doses no greater than 3 mg/kg every 8 weeks.

Initial application — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe Behcet's disease which is significantly impacting the patient's quality of life (see Notes); and
- 2 Fither
 - 2.1 The patient has severe ocular, neurological and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s) (see Notes); or
 - 2.2 The patient has severe gastrointestinal, rheumatologic and/or mucocutaneous symptoms and has not responded adequately to two or more treatment appropriate for the particular symptom(s) (see Notes); and
- 3 The patient is experiencing significant loss of quality of life.

Notes: Behcet's disease diagnosed according to the International Study Group for Behcet's Disease. Lancet 1990;335(8697):1078-80. Quality of life measured using an appropriate quality of life scale such as that published in Gilworth et al J Rheumatol. 2004;31:931-7.

Treatments appropriate for the particular symptoms are those that are considered standard conventional treatments for these symptoms, for example intravenous/oral steroids and other immunosuppressants for ocular symptoms; azathioprine, steroids, thalidomide, interferon alpha and ciclosporin for mucocutaneous symptoms; and colchicine, steroids and methotrexate for rheumatological symptoms.

Renewal — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has had a good clinical response to initial treatment with measurably improved quality of life; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Renewal — (fulminant ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Where maintenance treatment is considered appropriate, infliximab should be used in combination with immunomodulators and reassessed every 6 months; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe ocular inflammation; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or

| Subsidy | y Full | / Brand or |
|-----------------|--------------------|--------------|
| (Manufacturer's | s Price) Subsidise | d Generic |
| \$ | Per 🗸 | Manufacturer |

continued...

1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe ocular inflammation; or

- 2 Both:
 - 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
 - 2.2 Any of the following:
 - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
 - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
 - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Renewal — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < \(\frac{1}{2} + \) anterior chamber or vitreous cells, absence of active vitreous or retinal lesions. or resolution of uveitic cystoid macular oedema); or
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (ulcerative colitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active ulcerative colitis: and
- - 2.1 Patients SCCAI is greater than or equal to 4: or
 - 2.2 Patients PUCAI score is greater than or equal to 20; and
- 3 Patient has tried but has experienced an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and systemic corticosteroids.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 The SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on infliximab: or
 - 1.2 The PUCAI score has reduced by 10 points or more from the PUCAI score when the patient was initiated on infliximab: and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

1 Patient has pvoderma gangrenosum*; and

| Subsidy | Fully | Brand or | |
|------------------------|------------|--------------|--|
| (Manufacturer's Price) | Subsidised | Generic | |
| \$ | Per 🗸 | Manufacturer | |

continued...

- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Note: Indications marked with * are unapproved indications.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

Initial application — (inflammatory bowel arthritis – axial) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has had axial inflammatory pain for six months or more; and
- 3 Patient is unable to take NSAIDs; and
- 4 Patient has unequivocal sacroiliitis demonstrated by radiological imaging or MRI; and
- 5 Patient's disease has not responded adequately to prior treatment consisting of at least 3 months of an exercise regime supervised by a physiotherapist; and
- 6 Patient has a BASDAI of at least 6 on a 0 10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment.

Renewal — (inflammatory bowel arthritis – axial) from any relevant practitioner. Approvals valid for 2 years where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10-point scale, or an improvement in BASDAI of 50%, whichever is less.

Initial application — (inflammatory bowel arthritis – peripheral) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has active arthritis in at least four joints from the following: hip, knee, ankle, subtalar, tarsus, forefoot, wrist, elbow, shoulder, sternoclavicular; and
- 3 Patient has tried and not experienced a response to at least three months of methotrexate or azathioprine at a maximum tolerated dose (unless contraindicated); and
- 4 Patient has tried and not experienced a response to at least three months of sulfasalazine at a maximum tolerated dose (unless contraindicated); and
- 5 Any of the following:
 - 5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 5.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application: or
 - 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (inflammatory bowel arthritis – peripheral) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Fither:

- 1 Following initial treatment, patient has experienced at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 Patient has experienced at least a continuing 30% improvement in active joint count from baseline in the opinion of the

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

continued...

treating physician.

MEPOLIZUMAB - Special Authority see SA2331 below - Retail pharmacy

Inj 100 mg prefilled pen1,638.00 1 Nucala

⇒SA2331 Special Authority for Subsidy

Initial application — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Patient must have a diagnosis of severe eosinophilic asthma documented by a respiratory physician or clinical immunologist: and
- 3 Conditions that mimic asthma eq. vocal cord dysfunction, central airway obstruction, bronchiolitis etc. have been excluded: and
- 4 Patient has a blood eosinophil count of greater than 0.5 × 10°9 cells/L in the last 12 months; and
- 5 Patient must be adherent to optimised asthma therapy including inhaled corticosteroids (equivalent to at least 1000 mcg per day of fluticasone propionate) plus long acting beta-2 agonist, or budesonide/formoterol as part of the single maintenance and reliever therapy regimen, unless contraindicated or not tolerated; and
- 6 Either:
 - 6.1 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral corticosteroids: or
 - 6.2 Patient has received continuous oral corticosteroids of at least the equivalent of 10 mg per day over the previous 3 months: and
- 7 Treatment is not to be used in combination with subsidised benralizumab; and
- 8 Patient has an Asthma Control Test (ACT) score of 10 or less. Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 52 weeks after the first dose to assess response to treatment; and
- 9 Either:
 - 9.1 Patient has not previously received an anti-IL5 biological therapy for their severe eosinophilic asthma; or
 - 9.2 Both:
 - 9.2.1 Patient was refractory or intolerant to previous anti-IL5 biological therapy; and
 - 9.2.2 Patient was not eligible to continue treatment with previous anti-IL5 biological therapy and discontinued within 12 months of commencing treatment.

Renewal — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 Either:
 - 2.1 Exacerbations have been reduced from baseline by 50% as a result of treatment with mepolizumab; or
 - 2.2 Reduction in continuous oral corticosteroid use by 50% or by 10 mg/day while maintaining or improving asthma control.

Initial application — (eosinophilic granulomatosis with polyangiitis) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has eosinophilic granulomatosis with polyangiitis; and
- 2 The patient has trialled and not received adequate benefit from at least one of the following for at least three months (unless contraindicated to all): azathioprine, cyclophosphamide, leflunomide, methotrexate, mycophenolate, or rituximab;

| Subsidy | Fully | Brand or | |
|------------------------|------------|--------------|--|
| (Manufacturer's Price) | Subsidised | Generic | |
| <u> </u> | Per 🗸 | Manufacturer | |

continued...

and

- 3 Either:
 - 3.1 The patient has trialled prednisone for a minimum of three months and is unable to maintain disease control at doses below 7.5 mg per day; or
 - 3.2 Corticosteroids are contraindicated.

Renewal — (eosinophilic granulomatosis with polyangiitis) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months where patient has no evidence of clinical disease progression.

| OBINUTUZUMAB - PCT only - Specialist - Special Author | ity see SA2155 below | | |
|---|----------------------|------|----------|
| Inj 25 mg per ml, 40 ml vial | 5,910.00 | 1 | ✓ Gazyva |
| Inj 1 mg for ECP | 6.21 | 1 mg | ✓ Baxter |

⇒SA2155 Special Authority for Subsidy

Initial application — (chronic lymphocytic leukaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C CD20+ chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is obinutuzumab treatment naive: and
- 3 The patient is not eligible for full dose FCR due to comorbidities with a score > 6 on the Cumulative Illness Rating Scale (CIRS) or reduced renal function (creatinine clearance < 70mL/min); and
- 4 Patient has adequate neutrophil and platelet counts* unless the cytopenias are a consequence of marrow infiltration by CLL: and
- 5 Patient has good performance status; and
- 6 Obinutuzumab to be administered at a maximum cumulative dose of 8,000 mg and in combination with chlorambucil for a maximum of 6 cycles.

Notes: Chronic lymphocytic leukaemia includes small lymphocytic lymphoma. Comorbidity refers only to illness/impairment other than CLL induced illness/impairment in the patient. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with obinutuzumab is expected to improve symptoms and improve ECOG score to < 2.

* Neutrophil greater than or equal to 1.5×10^9 /L and platelets greater than or equal to 75×10^9 /L.

Initial application — (follicular / marginal zone lymphoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 Patient has follicular lymphoma; or
 - 1.2 Patient has marginal zone lymphoma; and
- 2 Patient is refractory to or has relapsed within 12 months of a rituximab containing combined chemo-immunotherapy regimen*; and
- 3 Patient has an ECOG performance status of 0-2; and
- 4 Patient has been previously treated with no more than four chemotherapy regimens; and
- 5 Obinutuzumab to be administered at a maximum dose of 1000 mg for a maximum of 6 cycles in combination with chemotherapy*.

Note: * includes unapproved indications

Renewal — (follicular / marginal zone lymphoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 24 months for applications meeting the following criteria: All of the following:

- 1 Patient has no evidence of disease progression following obinutuzumab induction therapy; and
- 2 Obinutuzumab to be administered at a maximum of 1000 mg every 2 months for a maximum of 2 years; and
- 3 Obinutuzumab to be discontinued at disease progression.

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer |
|--|---|-----|---------------------|-------------------------------------|
| OMALIZUMAB – Special Authority see SA1744 below – Retail phonon 150 mg prefilled syringe | , | 1 | | Colair Colair AU |
| Inj 150 mg vial | 450.00 | 1 | | Colair |

⇒SA1744 Special Authority for Subsidy

Initial application — (severe asthma) only from a respiratory specialist or clinical immunologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 6 years or older; and
- 2 Patient has a diagnosis of severe asthma; and
- 3 Past or current evidence of atopy, documented by skin prick testing or RAST; and
- 4 Total serum human immunoglobulin E (IgE) between 76 IU/mL and 1300 IU/ml at baseline; and
- 5 Proven adherence with optimal inhaled therapy including high dose inhaled corticosteroid (budesonide 1,600 mcg per day or fluticasone propionate 1,000 mcg per day or equivalent), plus long-acting beta-2 agonist therapy (at least salmeterol 50 mcg bd or eformoterol 12 mcg bd) for at least 12 months, unless contraindicated or not tolerated; and
- 6 Fither:
 - 6.1 Patient has received courses of systemic corticosteroids equivalent to at least 28 days treatment in the past 12 months. unless contraindicated or not tolerated; or
 - 6.2 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral steroids; and
- 7 Patient has an Asthma Control Test (ACT) score of 10 or less; and
- 8 Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 26 weeks after the first dose to assess response to treatment.

Initial application — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Fither:
 - 2.1 Both:
 - 2.1.1 Patient is symptomatic with Urticaria Activity Score 7 (UAS7) of 20 or above; and
 - 2.1.2 Patient has a Dermatology life quality index (DLQI) of 10 or greater; or
 - 2.2 Patient has a Urticaria Control Test (UCT) of 8 or less; and
- 3 Any of the following:
 - 3.1 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and ciclosporin (> 3 mg/kg day) for at least 6 weeks; or
 - 3.2 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and at least 3 courses of systemic corticosteroids (> 20 mg prednisone per day for at least 5 days) in the previous 6 months; or
 - 3.3 Patient has developed significant adverse effects whilst on corticosteroids or ciclosporin; and
- 4 Fither:
 - 4.1 Treatment to be stopped if inadequate response* following 4 doses; or
 - 4.2 Complete response* to 6 doses of omalizumab.

Renewal — (severe asthma) only from a clinical immunologist or respiratory specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 A reduction in the maintenance oral corticosteroid dose or number of exacerbations of at least 50% from baseline.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

Renewal — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Patient has previously adequately responded* to 6 doses of omalizumab; or
- 2 Both:
 - 2.1 Patient has previously had a complete response* to 6 doses of omalizumab; and
 - 2.2 Patient has relapsed after cessation of omalizumab therapy.

Note: *Inadequate response defined as less than 50% reduction in baseline UAS7 and DLQI score, or an increase in Urticaria Control Test (UCT) score of less than 4 from baseline. Patient is to be reassessed for response after 4 doses of omalizumab. Complete response is defined as UAS7 less than or equal to 6 and DLQI less than or equal to 5; or UCT of 16. Relapse of chronic urticaria on stopping prednisone/ciclosporin does not justify the funding of omalizumab.

PALIVIZUMAB - PCT only - Special Authority see SA2419 below

Inj 100 mg per ml, 1 ml vial.......1,700.00

1 ✓ Synagis

⇒SA2419 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Palivizumab to be administered during the annual RSV season; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Infant was born in the last 12 months; and
 - 2.1.2 Infant was born at less than 32 weeks zero days' gestation; or
 - 2.2 Both:
 - 2.2.1 Child was born in the last 24 months; and
 - 2.2.2 Any of the following:
 - 2.2.2.1 Child has severe lung, airway, neurological or neuromuscular disease that requires ongoing ventilatory/respiratory support (see Note A) in the community: or
 - 2.2.2.2 Both:
 - 2.2.2.2.1 Child has haemodynamically significant heart disease; and
 - 2.2.2.2.2 Any of the following:
 - 2.2.2.2.2.1 Child has unoperated simple congenital heart disease with significant left to right shunt (see Note B); or
 - 2.2.2.2.2.2 Child has unoperated or surgically palliated complex congenital heart disease; or
 - 2.2.2.2.3 Child has severe pulmonary hypertension (see Note C); or
 - 2.2.2.2.2.4 Child has moderate or severe left ventricular (LV) failure (see Note D); or
 - 2.2.2.3 Child has severe combined immune deficiency, confirmed by an immunologist, but has not received a stem cell transplant: or
 - 2.2.2.4 Child has inborn errors of immunity (see Note E) that increase susceptibility to life-threatening viral respiratory infections, confirmed by an immunologist.

Renewal from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Palivizumab to be administered during the annual RSV season; and
- 2 Child was born in the last 24 months; and
- 3 Any of the following:
 - 3.1 Child has severe lung, airway, neurological or neuromuscular disease that requires ongoing ventilatory/respiratory support (see Note A) in the community; or
 - 3.2 Both:

| Subsidy (Manufacturer's Price) | Subsi | Fully dised | Brand or Generic |
|-----------------------------------|-------|----------------|---------------------|
| \$ | Per | ✓ | Manufacturer |

continued...

- 3.2.1 Child has haemodynamically significant heart disease; and
- 3.2.2 Any of the following:
 - 3.2.2.1 Child has unoperated simple congenital heart disease with significant left to right shunt (see Note B); or
 - 3.2.2.2 Child has unoperated or surgically palliated complex congenital heart disease; or
 - 3.2.2.3 Child has severe pulmonary hypertension (see Note C); or
 - 3.2.2.4 Child has moderate or severe left ventricular (LV) failure (see Note D); or
- 3.3 Child has severe combined immune deficiency, confirmed by an immunologist, but has not received a stem cell transplant; or
- 3.4 Child has inborn errors of immunity (see Note E) that increase susceptibility to life-threatening viral respiratory infections, confirmed by an immunologist.

Notes:

- a) Ventilatory/respiratory support includes those on home oxygen, CPAP/VPAP and those with tracheostomies in situ managed at home
- b) Child requires/will require heart failure medication, and/or child has significant pulmonary hypertension, and/or infant will require surgical palliation/definitive repair within the next 3 months
- c) Mean pulmonary artery pressure more than 25 mmHg
- d) LV Eiection Fraction less than 40%
- e) Inborn errors of immunity include, but are not limited to, IFNAR deficiencies

PERTUZUMAB – PCT only – Specialist – Special Authority see SA2276 below

| Inj 30 mg per ml, 14 ml vial | 3,927.00 | 1 | Perjeta |
|------------------------------|----------|-----------|----------|
| Inj 420 mg for ECP | 3,927.00 | 420 mg OP | ✓ Baxter |

⇒SA2276 Special Authority for Subsidy

Initial application — (metastatic breast cancer) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Either:
 - 2.1 Patient is chemotherapy treatment naïve: or
 - 2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
- 3 The patient has good performance status (ECOG grade 0-1); and
- 4 Pertuzumab to be administered in combination with trastuzumab; and
- 5 Pertuzumab maximum first dose of 840 mg, followed by maximum of 420 mg every 3 weeks; and
- 6 Pertuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
 - 1.2 The cancer has not progressed at any time point during the previous 12 months whilst on pertuzumab and trastuzumab; or
- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with pertuzumab and trastuzumab for reasons other than severe toxicity or disease progression; and
 - 2.2 Patient has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with pertuzumab and trastuzumab.

Fully

Brand or

Subsidy

| | (Manufacturer's Price) | Per | Subsidised Generic Manufacturer |
|--|------------------------|------|----------------------------------|
| | Ψ | | Wallulacturei |
| RITUXIMAB (MABTHERA) - PCT only - Specialist - Special A | uthority see SA1976 b | elow | |
| Inj 100 mg per 10 ml vial | | 2 | Mabthera |
| Inj 500 mg per 50 ml vial | 2,688.30 | 1 | ✓ Mabthera |
| Inj 1 mg for ECP | 5.64 | 1 mg | ✓ Baxter (Mabthera) |

⇒SA1976 Special Authority for Subsidy

Initial application — (rheumatoid arthritis - TNF inhibitors contraindicated) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Treatment with a Tumour Necrosis Factor alpha inhibitor is contraindicated; and
- 2 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroguine sulphate (at maximum tolerated doses); and
- 5 Any of the following:
 - 5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin: or
 - 5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
 - 5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and

6 Either:

- 6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
- 6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and

7 Either:

- 7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
- 7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months; and

8 Either:

- 8.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
- 8.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 9 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Initial application — (rheumatoid arthritis - prior TNF inhibitor use) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

1 Both:

- 1.1 The patient has had an initial community Special Authority approval for at least one of etanercept and/or adalimumab for rheumatoid arthritis; and
- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 1.2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept for rheumatoid arthritis: and

2 Fither:

2.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or

| bsidy urer's Price) Subs | Fully | Brand or Generic |
|-----------------------------|-------|---------------------|
| \$ Per | • | Manufacturer |

continued...

- 2.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 3 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Renewal — (rheumatoid arthritis - re-treatment in 'partial responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 At 4 months following the initial course of rituximab infusions the patient had between a 30% and 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 At 4 months following the second course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.3 At 4 months following the third and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Either:
 - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Renewal — (rheumatoid arthritis - re-treatment in 'responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 At 4 months following the initial course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 At 4 months following the second and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Either:
 - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

RITUXIMAB (RIXIMYO) - PCT only - Specialist - Special Authority see \$A2233 below

| inj 100 mg per 10 mi viai | 2/5.33 | 2 | ✓ Riximyo |
|---------------------------|--------|------|--------------------|
| Inj 500 mg per 50 ml vial | 688.20 | 1 | ✓ Riximyo |
| Ini 1 mg for ECP | 1.38 | 1 ma | ✓ Baxter (Riximvo) |

⇒SA2233 Special Authority for Subsidy

Initial application — (ABO-incompatible organ transplant) from any relevant practitioner. Approvals valid without further renewal unless notified where patient is to undergo an ABO-incompatible solid organ transplant*.

Note: Indications marked with * are unapproved indications.

Initial application — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*: and
- 2 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per ✓ | Manufacturer |

continued...

- 4 weeks: and
- 3 Any of the following:
 - 3.1 Induction therapy with daily oral or pulse intravenous cyclophosphamide has failed to achieve significant improvement of disease after at least 3 months; or
 - 3.2 Patient has previously had a cumulative dose of cyclophosphamide > 15 g or a further repeat 3 month induction course of cyclophosphamide would result in a cumulative dose > 15 g; or
 - 3.3 Cyclophosphamide and methotrexate are contraindicated; or
 - 3.4 Patient is a female of child-bearing potential; or
 - 3.5 Patient has a previous history of haemorrhagic cystitis, urological malignancy or haematological malignancy.

Note: Indications marked with * are unapproved indications.

Renewal — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 Patient has previously responded to treatment with rituximab but is now experiencing an acute flare of vasculitis; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (Antibody-mediated organ transplant rejection) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has been diagnosed with antibody-mediated organ transplant rejection*.

Note: Indications marked with * are unapproved indications.

Initial application — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C chronic lymphocytic leukaemia (CLL) requiring treatment; and
- 2 Any of the following:
 - 2.1 The patient is rituximab treatment naive; or
 - 2.2 Fither:
 - 2.2.1 The patient is chemotherapy treatment naive; or
 - 2.2.2 Both:
 - 2.2.2.1 The patient's disease has relapsed following no more than three prior lines of chemotherapy treatment: and
 - 2.2.2.2 The patient has had a treatment-free interval of 12 months or more if previously treated with fludarabine and cyclophosphamide chemotherapy; or
 - 2.3 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax; and
- 3 The patient has good performance status; and
- 4 Either:
 - 4.1 The patient does not have chromosome 17p deletion CLL; or
 - 4.2 Rituximab treatment is to be used in combination with funded venetoclax for relapsed/refractory chronic lymphocytic leukaemia: and
- 5 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles; and
- 6 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration), bendamustine or venetoclax.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments. 'Good performance

| Subsidy | | Fully | Brand or | |
|------------------------|-----|------------|--------------|--|
| (Manufacturer's Price) | 5 | Subsidised | Generic | |
| \$ | Per | 1 | Manufacturer | |

continued...

status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with rituximab is expected to improve symptoms and improve ECOG score to < 2. **Renewal — (Chronic lymphocytic leukaemia)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax; or
 - 1.2 All of the following:
 - 1.2.1 The patient's disease has relapsed following no more than one prior line of treatment with rituximab for CLL; and
 - 1.2.2 The patient has had an interval of 36 months or more since commencement of initial rituximab treatment; and
 - 1.2.3 The patient does not have chromosome 17p deletion CLL; and
 - 1.2.4 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration) or bendamustine; and
- 2 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initial application — (Neuromyelitis Optica Spectrum Disorder(NMOSD)) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks; and
- 2 Either:
 - 2.1 The patient has experienced a severe episode or attack of NMOSD (rapidly progressing symptoms and clinical investigations supportive of a severe attack of NMOSD); or
 - 2.2 All of the following:
 - 2.2.1 The patient has experienced a breakthrough attack of NMOSD; and
 - 2.2.2 The patient is receiving treatment with mycophenolate; and
 - 2.2.3 The patients is receiving treatment with corticosteroids.

Renewal — (Neuromyelitis Optica Spectrum Disorder) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks; and
- 2 The patients has responded to the most recent course of rituximab; and
- 3 The patient has not received rituximab in the previous 6 months.

Initial application — (Post-transplant) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 2 To be used for a maximum of 8 treatment cycles.

Note: Indications marked with * are unapproved indications.

Renewal — (Post-transplant) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 3 To be used for no more than 6 treatment cycles.

Note: Indications marked with * are unapproved indications.

Initial application — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
- 2 Either:
 - 2.1 Treatment with corticosteroids and at least one other immunosuppressant for at least a period of 12 months has been ineffective; or
 - 2.2 Both:
 - 2.2.1 Treatment with at least one other immunosuppressant for a period of at least 12 months; and
 - 2.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects

Renewal — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Either:
 - 3.1 The patient has relapsed despite treatment with corticosteroids and at least one other immunosuppressant for a period of at least 12 months; or
 - 3.2 Both:
 - 3.2.1 The patient's myasthenia gravis has relapsed despite treatment with at least one immunosuppressant for a period of at least 12 months; and
 - 3.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

Initial application — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is a child with SDNS* or FRNS*; and
- 2 Treatment with steroids for at least a period of 3 months has been ineffective or associated with evidence of steroid toxicity; and
- 3 Treatment with ciclosporin for at least a period of 3 months has been ineffective and/or discontinued due to unacceptable side effects; and
- 4 Treatment with mycophenolate for at least a period of 3 months with no reduction in disease relapses; and
- 5 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting

| Subsidy | F | ully | Brand or |
|------------------------|----------|------|--------------|
| (Manufacturer's Price) | Subsidis | sed | Generic |
| | Per | 1 | Manufacturer |

continued...

the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of

Note: Indications marked with * are unapproved indications.

Initial application — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Patient is a child with SRNS* where treatment with steroids and ciclosporin for at least 3 months have been ineffective; and
- 2 Treatment with tacrolimus for at least 3 months has been ineffective; and
- 3 Genetic causes of nephrotic syndrome have been excluded; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient has treatment naive aggressive CD20 positive NHL; and
 - 1.2 To be used with a multi-agent chemotherapy regimen given with curative intent; and
 - 1.3 To be used for a maximum of 8 treatment cycles: or
- 2 Both:
 - 2.1 The patient has aggressive CD20 positive NHL with relapsed disease following prior chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Renewal — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has relapsed refractory/aggressive CD20 positive NHL; and
- 3 To be used with a multi-agent chemotherapy regimen given with curative intent; and
- 4 To be used for a maximum of 4 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Initial application — (haemophilia with inhibitors) only from a haematologist or Practitioner on the recommendation of a

| Subsidy | Full | y Brand or | |
|------------------------|-----------|----------------------------------|--|
| (Manufacturer's Price) | Subsidise | d Generic | |
| \$ | Per 🗸 | Manufacturer | |

continued...

haematologist. Approvals valid for 4 months for applications meeting the following criteria:

Any of the following:

- 1 Patient has mild congenital haemophilia complicated by inhibitors; or
- 2 Patient has severe congenital haemophilia complicated by inhibitors and has failed immune tolerance therapy; or
- 3 Patient has acquired haemophilia.

Renewal — **(haemophilia with inhibitors)** only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for haemophilia with inhibitors; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment.

Initial application — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Fither:
 - 1.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per microlitre: or
 - 1.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding; and
- 2 Any of the following:
 - 2.1 Treatment with steroids and splenectomy have been ineffective; or
 - 2.2 Treatment with steroids has been ineffective and splenectomy is an absolute contraindication; or
 - 2.3 Other treatments including steroids have been ineffective and patient is being prepared for elective surgery (e.g. splenectomy); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks

Note: Indications marked with * are unapproved indications.

Renewal — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for immune thrombocytopenic purpura*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (indolent, low-grade lymphomas or hairy cell leukaemia*) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has indolent low grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
 - 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
 - 2.1 The patient has indolent, low grade lymphoma or hairy cell leukaemia* requiring first-line systemic chemotherapy; and

| bsidy | Fully | Brand or |
|-----------------------|-------|----------|
| turer's Price) Subsid | dised | Generic |
| \$ Per | ✓ | |

continued...

2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Renewal — (indolent, low-grade lymphomas or hairy cell leukaemia*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has indolent, low-grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
- 3 To be used for no more than 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Initial application — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient has autoimmune pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder.

Note: Indications marked with * are unapproved indications.

Renewal — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient was previously treated with rituximab for pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder and demonstrated an initial response lasting at least 12 months.

Note: Indications marked with * are unapproved indications.

Initial application — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Patient has cold haemagglutinin disease*; and
- 2 Patient has severe disease which is characterized by symptomatic anaemia, transfusion dependence or disabling circulatory symptoms; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Fither:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for severe cold haemagglutinin disease*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Both:

- 1 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks; and
- 2 Either:
 - 2.1 Patient has thrombotic thrombocytopenic purpura* and has experienced progression of clinical symptoms or persistent thrombocytopenia despite plasma exchange; or

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

2.2 Patient has acute idiopathic thrombotic thrombocytopenic purpura* with neurological or cardiovascular pathology.

Note: Indications marked with * are unapproved indications.

Renewal — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for thrombotic thrombocytopenic purpura*; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe, immediately life- or organ-threatening SLE*; and
- 2 The disease has proved refractory to treatment with steroids at a dose of at least 1 mg/kg; and
- 3 The disease has relapsed following prior treatment for at least 6 months with maximal tolerated doses of azathioprine, mycophenolate mofetil and high dose cyclophosphamide, or cyclophosphamide is contraindicated; and
- 4 Maximum of four 1000 mg infusions of rituximab.

Note: Indications marked with * are unapproved indications.

Renewal — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient's SLE* achieved at least a partial response to the previous round of prior rituximab treatment; and
- 2 The disease has subsequently relapsed; and
- 3 Maximum of two 1000 mg infusions of rituximab.

Note: Indications marked with * are unapproved indications.

Initial application — (warm autoimmune haemolytic anaemia (warm AlHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has warm autoimmune haemolytic anaemia*; and
- 2 One of the following treatments has been ineffective: steroids (including if patient requires ongoing steroids at doses equivalent to > 5 mg prednisone daily), cytotoxic agents (e.g. cyclophosphamide monotherapy or in combination), intravenous immunoglobulin; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (warm autoimmune haemolytic anaemia (warm AIHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for warm autoimmune haemolytic anaemia*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and

| Subsidy | , | Fully | Brand or | |
|-----------------------|-----|------------|--------------|--|
| (Manufacturer's Price |) | Subsidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

continued...

2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (severe antisynthetase syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed antisynthetase syndrome; and
- 2 Patient has severe, immediately life or organ threatening disease, including interstitial lung disease; and
- 3 Either:
 - 3.1 Treatment with at least 3 immunosuppressants (oral steroids, cyclophosphamide, methotrexate, mycophenolate, ciclosporin, azathioprine) has not be effective at controlling active disease; or
 - 3.2 Rapid treatment is required due to life threatening complications; and
- 4 Maximum of four 1,000mg infusions of rituximab.

Renewal — (severe antisynthetase syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in inflammatory markers, muscle strength and pulmonary function; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 Maximum of two cycles of 2 x 1,000mg infusions of rituximab given two weeks apart.

Initial application — (graft versus host disease) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has refractory graft versus host disease following transplant; and
- 2 Treatment with at least 3 immunosuppressants (oral steroids, ciclosporin, tacrolimus, mycophenolate, sirolimus) has not be effective at controlling active disease; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Initial application — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has severe chronic inflammatory demyelinating polyneuropathy (CIPD); and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
 - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
 - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function compared to baseline; and
- 2 The patient has not received rituximab in the previous 6 months; and

| Subsidy | Fully | Brand or | Т |
|------------------------|------------|--------------|---|
| (Manufacturer's Price) | Subsidised | Generic | |
| \$ | Per 🗸 | Manufacturer | |

continued...

3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Initial application — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has severe anti-NMDA receptor autoimmune encephalitis: and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
 - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
 - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function: and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 The patient has experienced a relapse and now requires further treatment; and
- 4 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Initial application — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria: Fither:

- 1 Both:
 - 1.1 The patient has CD20+ low grade or follicular B-cell NHL with relapsed disease following prior chemotherapy; and
 - 1.2 To be used for a maximum of 6 treatment cycles; or
 - 2 Both:
 - 2.1 The patient has CD20+ low grade or follicular B-cell NHL requiring first-line systemic chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Renewal — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria: Both:

- - 1 Rituximab is to be used for maintenance in CD20+ low grade or follicular B-cell NHL following induction with first-line systemic chemotherapy; and
 - 2 Patient is intended to receive rituximab maintenance therapy for 2 years at a dose of 375 mg/m2 every 8 weeks (maximum of 12 cycles).

Initial application — (Membranous nephropathy) only from a nephrologist or any relevant practitioner on the recommendation of a nephrologist. Approvals valid for 6 weeks for applications meeting the following criteria: All of the following:

- 1 Fither:
 - 1.1 Patient has biopsy-proven primary/idiopathic membranous nephropathy*; or
 - 1.2 Patient has PLA2 antibodies with no evidence of secondary cause, and an eGFR of > 60ml/min/1,73m2; and

| Subsidy | | Fully | Brand or | |
|------------------------|-----|------------|--------------|--|
| (Manufacturer's Price) | | Subsidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

continued...

- 2 Patient remains at high risk of progression to end-stage kidney disease despite more than 3 months of treatment with conservative measures (see Note); and
- 3 The total rituximab dose would not exceed the equivalent of 375mg/m2 of body surface area per week for a total of 4 weeks.

Renewal — (Membranous nephropathy) only from a nephrologist or any relevant practitioner on the recommendation of a nephrologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for membranous nephropathy*; and
- 2 Either:
 - 2.1 Treatment with rituximab was previously successful, but the condition has relapsed, and the patient now requires repeat treatment; or
 - 2.2 Patient achieved partial response to treatment and requires repeat treatment (see Note); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Notes:

- a) Indications marked with * are unapproved indications.
- b) High risk of progression to end-stage kidney disease defined as > 5g/day proteinuria.
- c) Conservative measures include renin-angiotensin system blockade, blood-pressure management, dietary sodium and protein restriction, treatment of dyslipidaemia, and anticoagulation agents unless contraindicated or the patient has experienced intolerable side effects.
- d) Partial response defined as a reduction of proteinuria of at least 50% from baseline, and between 0.3 grams and 3.5 grams per 24 hours.

Initial application — (B-cell acute lymphoblastic leukaemia/lymphoma*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient has newly diagnosed B-cell acute lymphoblastic leukaemia/lymphoma*; and
- 2 Treatment must be in combination with an intensive chemotherapy protocol with curative intent; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m² per dose for a maximum of 18 doses.

Note: Indications marked with * are unapproved indications.

Initial application — (desensisation prior to transplant) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires desensitisation prior to mismatched allogenic stem cell transplant*; and
- 2 Patient would receive no more than two doses at 375 mg/m2 of body-surface area.

Note: Indications marked with * are unapproved indications.

Initial application — (pemiphigus*) only from a dermatologist or relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 Patient has severe rapidly progressive pemphigus; and
 - 1.2 Is used in combination with systemic corticosteroids (20 mg/day); and
 - 1.3 Any of the following:
 - 1.3.1 Skin involvement is at least 5% body surface area; or
 - 1.3.2 Significant mucosal involvement (10 or more mucosal erosions) or diffuse gingivitis or confluent large erosions: or
 - 1.3.3 Involvement of two or more mucosal sites; or

| Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer |
|--|
|--|

continued...

2 Both:

2.1 Patient has pemphigus; and

2.2 Patient has not experienced adequate clinical benefit from systemic corticosteroids (20 mg/day) in combination with a steroid sparing agent, unless contraindicated.

Note: Indications marked with * are unapproved indications.

Renewal — (pemiphigus*) only from a dermatologist or relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has experienced adequate clinical benefit from rituximab treatment, with improvement in symptoms and healing of skin ulceration and reduction in corticosteroid requirement; and
- 2 Patient has not received rituximab in the previous 6 months.

Note: Indications marked with * are unapproved indications.

Initial application — (immunoglobulin G4-related disease (IgG4-RD*)) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed diagnosis of IgG4-RD*; and
- 2 Fither:
 - 2.1 Treatment with corticosteroids and/or disease modifying anti-rheumatic drugs for at least 3 months has been ineffective in lowering corticosteroid dose below 5 mg per day (prednisone equivalent) without relapse; or
 - 2.2 Treatment with corticosteroids and/or disease modifying anti-rheumatic drugs is contraindicated or associated with evidence of toxicity or intolerance; and
- 3 Total rituximab dose used should not exceed a maximum of two 1000 mg infusions of rituximab given two weeks apart. Note: Indications marked with * are unapproved indications.

Renewal — (immunoglobulin G4-related disease (IgG4-RD*)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Treatment with rituximab for IgG4-RD* was previously successful and patient's disease has demonstrated sustained response, but the condition has relapsed; or
 - 1.2 Patient is receiving maintenance treatment for IgG4-RD*; and
- 2 Rituximab re-treatment not to be given within 6 months of previous course of treatment; and
- 3 Maximum of two 1000 mg infusions of rituximab given two weeks apart.

Note: Indications marked with * are unapproved indications.

SECUKINUMAB - Special Authority see SA2403 below - Retail pharmacy

⇒SA2403 Special Authority for Subsidy

Initial application — (severe chronic plaque psoriasis – second-line biologic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab or etanercept, or has trialled infliximab in a Health NZ Hospital, for severe chronic plaque psoriasis; and
- 2 Fither:
 - 2.1 The patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
 - 2.2 The patient has received insufficient benefit from adalimumab, etanercept or infliximab; and
- 3 A Psoriasis Area and Severity Index (PASI) assessment or Dermatology Quality of Life Index (DLQI) assessment has

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

continued...

been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and

4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Initial application — (severe chronic plaque psoriasis – first-line biologic) only from a dermatologist or any relevant practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; or
 - 1.3 Patient has severe chronic localised genital or flexural plaque psoriasis where the plaques or lesions have been present for at least 6 months from the time of initial diagnosis, and with a Dermatology Life Quality Index (DLQI) score greater than 10; and
- 2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
- 3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Note: A treatment course is defined as a minimum of 12 weeks of treatment. "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand. foot, genital or flexural areas, at least 2 of the 3 PASI symptom sub scores for erythema, thickness and scaling are rated as severe or very severe, and for the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis – first and second-line biologic) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Roth:

- 1 Either:
 - 1.1 Fither:
 - 1.1.1 Patient's PASI score has reduced by 75% or more (PASI 75) as compared to baseline PASI prior to commencing secukinumab; or
 - 1.1.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing secukinumab: or
 - 1.2 Both:
 - 1.2.1 Patient had severe chronic localised genital or flexural plaque psoriasis at the start of treatment; and
 - 1.2.2 Either:
 - 1.2.2.1 The patient has experienced a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; or
 - 1.2.2.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing secukinumab; and
- 2 Secukinumab to be administered at a maximum dose of 300 mg monthly.

Initial application — (ankylosing spondylitis – second-line biologic) only from a rheumatologist or Practitioner on the

| Subsidy | | Fully | Brand or |
|------------------------|-----|-----------|--------------|
| (Manufacturer's Price) | Sı | ubsidised | Generic |
| \$ | Per | ✓ | Manufacturer |

continued...

recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and
- 2 Fither:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

Renewal — (ankylosing spondylitis – second-line biologic) only from a rheumatologist or medical practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Following 12 weeks initial treatment of secukinumab treatment, BASDAI has improved by 4 or more points from pre-secukinumab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefitted from treatment and that continued treatment is appropriate; and
- 3 Secukinumab to be administered at doses no greater than 150 mg monthly.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has had an initial Special Authority approval for adalimumab, etanercept or infliximab for psoriatic arthritis; and
 - 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
 - 1.2.2 Patient has received insufficient benefit from adalimumab, etanercept or infliximab to meet the renewal criteria for adalimumab, etanercept or infliximab for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
 - 2.4 Either:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints;
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

continued...

- 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior secukinumab treatment in the opinion of the treating physician; and
- 2 Secukinumab to be administered at doses no greater than 300 mg monthly.

SILTUXIMAB - Special Authority see SA1596 below - Retail pharmacy

| Note: Siltuximab is to be administered at doses no greater th | • | ry 3 weeks. | |
|---|----------|-------------|-----------|
| Inj 100 mg vial | 770.57 | 1 | Sylvant |
| Inj 400 mg vial | 3,082.33 | 1 | ✓ Sylvant |

⇒SA1596 Special Authority for Subsidy

Initial application only from a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's Disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Siltuximab is to be administered at doses no greater than 11 mg/kg every 3 weeks.

Renewal only from a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has sustained improvement in inflammatory markers and functional status.

TIXAGEVIMAB WITH CILGAVIMAB - [Xpharm] - Subsidy by endorsement

- a) No patient co-payment payable
- b) Treatment is funded only if patient meets access criteria for tixagevimab with cilgavimab (as per https://pharmac.govt.nz/Evusheld) and has been endorsed accordingly by the prescriber. The supply of treatment is via Pharmac's approved distribution process. Refer to the Pharmac website for more information about this and stock availability.

| | | 00 mg per | Inj 100 mg per ml, 1.5 ml vial with cilgavimab 1 |
|----------------------------|------|----------------|--|
| Evusheld | 1 | 0.00 | ml,1.5 ml vial |
| | | e SA2404 below | TOCILIZUMAB - PCT only - Special Authority see |
| ✓ Actemra | 1 | 220.00 | Inj 20 mg per ml, 4 ml vial |
| ✓ Actemra | 1 | 550.00 | Inj 20 mg per ml, 10 ml vial |
| ✓ Actemra | 1 | | Inj 20 mg per ml, 20 ml vial |
| ✓ Baxter | 1 mg | 2.85 | Inj 1 mg for ECP |

⇒SA2404 Special Authority for Subsidy

Initial application — (cytokine release syndrome) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

1 Both:

- 1.1 The patient has developed grade 3 or 4 cytokine release syndrome associated with the administration of blinatumomab for the treatment of acute lymphoblastic leukaemia; and
- 1.2 Tocilizumab is to be administered at doses no greater than 8 mg/kg IV for a maximum of 3 doses (if less than 30kg, maximum of 12 mg/kg); or
- 2 All of the following:
 - 2.1 The patient is enrolled in the Malaghan Institute of Medical Research ENABLE trial programme; and
 - 2.2 The patient has developed CRS or Immune Effector Cell-Associated Neurotoxicity Syndrome (ICANS) following CAR T-cell therapy for the treatment of relapsed or refractory B-cell non-Hodgkin lymphoma; and
 - 2.3 Tocilizumab is to be administered according to the consensus guidelines for CRS or ICANS for CAR T-cell therapy at doses no greater than 8 mg/kg IV for a maximum of 3 doses.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

| Subsidy | | Fully | Brand or | |
|-----------------------|-----|-----------|--------------|--|
| (Manufacturer's Price | Sı | ubsidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

continued...

Both:

- 1 Patient was being treated with tocilizumab prior to 1 February 2019; and
- 2 Any of the following:
 - 2.1 rheumatoid arthritis; or
 - 2.2 systemic juvenile idiopathic arthritis; or
 - 2.3 adult-onset Still's disease; or
 - 2.4 polyarticular juvenile idiopathic arthritis; or
 - 2.5 idiopathic multicentric Castleman's disease.

Initial application — (Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Fither:
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
- 3 Either:
 - 3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or
 - 3.2 Both:
 - 3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a Health NZ Hospital; and
 - 3.2.2 Either:
 - 3.2.2.1 The patient has experienced intolerable side effects from rituximab; or
 - 3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Initial application — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 2 Tocilizumab is to be used as monotherapy; and
- 3 Either:
 - 3.1 Treatment with methotrexate is contraindicated: or
 - 3.2 Patient has tried and did not tolerate oral and/or parenteral methotrexate; and
- 4 Fither:
 - 4.1 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of ciclosporin alone or in combination with another agent; or
 - 4.2 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of leflunomide alone or in combination with another agent; and
- 5 Either:
 - 5.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 active, swollen, tender joints;
 - 5.2 Patient has persistent symptoms of poorly controlled and active disease in at least four active joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 6 Either:
 - 6.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or

| Subsidy | | Fully | Brand or | |
|-----------------------|-----|------------|--------------|--|
| (Manufacturer's Price |) | Subsidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

continued...

6.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Initial application — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient diagnosed with systemic juvenile idiopathic arthritis; and
- 2 Patient has tried and not responded to a reasonable trial of all of the following, either alone or in combination: oral or parenteral methotrexate; non-steroidal anti-inflammatory drugs (NSAIDs); and systemic corticosteroids.

Initial application — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 Fither:
 - 1.1.1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a Health NZ Hospital; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for AOSD: or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal antiinflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Initial application — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for both etanercept and adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 The patient has experienced intolerable side effects, or has received insufficient benefit from, both etanercept and adalimumab; or
- 2 All of the following:
 - 2.1 Treatment with a tumour necrosis factor alpha inhibitor is contraindicated; and
 - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
 - 2.3 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.4 Any of the following:
 - 2.4.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.4.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.4.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Initial application — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

continued...

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Tocilizumab to be administered at doses no greater than 8 mg/kg IV every 3-4 weeks.

Initial application — (moderate to severe COVID-19) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed (or probable) COVID-19; and
- 2 Oxygen saturation of < 92% on room air, or requiring supplemental oxygen; and
- 3 Patient is receiving adjunct systemic corticosteroids, or systemic corticosteroids are contraindicated; and
- 4 Tocilizumab is to be administered at doses no greater than 8mg/kg IV for a maximum of one dose; and
- 5 Tocilizumab is not to be administered in combination with barcitinib.

Renewal — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Renewal — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Following up to 6 months' initial treatment, the patient has achieved at least an American College of Rheumatology paediatric 30% improvement criteria (ACR Pedi 30) response from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing ACR Pedi 30 response from baseline.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months where the patient has a sustained improvement in inflammatory markers and functional status. Renewal — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Renewal — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has a sustained improvement in inflammatory markers and functional status.

 TRASTUZUMAB (HERZUMA) – PCT only – Special Authority see SA2293 below

 Inj 150 mg vial
 100.00
 1
 ✓ Herzuma

 Inj 440 mg vial
 293.35
 1
 ✓ Herzuma

 Inj 1 mg for ECP
 0.70
 1 mg
 ✓ Baxter

⇒SA2293 Special Authority for Subsidy

Initial application — (early breast cancer) from any relevant practitioner. Approvals valid for 15 months for applications

| Subsidy (Manufacturer's Price) | Sub | Fully sidised | Brand or Generic |
|-----------------------------------|-----|------------------|---------------------|
| \$ | Per | 1 | Manufacturer |

continued...

meeting the following criteria:

Both:

- 1 The patient has early breast cancer expressing HER-2 IHC 3+ or ISH + (including FISH or other current technology); and
- 2 Maximum cumulative dose of 106 mg/kg (12 months' treatment).

Renewal — (early breast cancer*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology): and
 - 1.2 The patient received prior adjuvant trastuzumab treatment for early breast cancer; and
 - 1.3 Any of the following:
 - 1.3.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
 - 1.3.2 The patient discontinued lapatinib within 3 months due to intolerable side effects and the cancer did not progress whilst on lapatinib; or
 - 1.3.3 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
 - 1.4 Fither:
 - 1.4.1 Trastuzumab will not be given in combination with pertuzumab; or
 - 1.4.2 All of the following:
 - 1.4.2.1 Trastuzumab to be administered in combination with pertuzumab; and
 - 1.4.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer: and
 - 1.4.2.3 The patient has good performance status (ECOG grade 0-1); and
 - 1.5 Trastuzumab to be discontinued at disease progression; or
- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with trastuzumab in the metastatic setting for reasons other than severe toxicity or disease progression; and
 - 2.2 Patient has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with trastuzumab.

Note: * For patients with relapsed HER-2 positive disease who have previously received adjuvant trastuzumab for early breast cancer

Initial application — (metastatic breast cancer) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Either:
 - 2.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
 - 2.2 The patient discontinued lapatinib within 3 months due to intolerable side effects and the cancer did not progress whilst on lapatinib; and
- 3 Either:
 - 3.1 Trastuzumab will not be given in combination with pertuzumab; or
 - 3.2 All of the following:
 - 3.2.1 Trastuzumab to be administered in combination with pertuzumab; and
 - 3.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

continued...

cancer: and

- 3.2.3 The patient has good performance status (ECOG grade 0-1); and
- 4 Trastuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
 - 1.2 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
 - 1.3 Trastuzumab to be discontinued at disease progression; or
- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with trastuzumab for reasons other than severe toxicity or disease progression; and
 - 2.2 Patient has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with trastuzumab.

Initial application — (gastric, gastro-oesophageal junction and oesophageal cancer) from any relevant practitioner.
Approvals valid for 12 months for applications meeting the following criteria:
Both:

- 1 The patient has locally advanced or metastatic gastric, gastro-oesophageal junction or oesophageal cancer expressing HER-2 IHC 2+ FISH+ or IHC3+ (or other current technology); and
- 2 Patient has an ECOG score of 0-2.

Renewal — (gastric, gastro-oesophageal junction and oesophageal cancer) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 2 Trastuzumab to be discontinued at disease progression.

TRASTUZUMAB DERUXTECAN - PCT only - Special Authority see SA2420 below

| Inj 100 mg per ml, 1 ml vial | 2,550.00 | 1 | Enhertu |
|------------------------------|----------|------|----------|
| Inj 1 mg for ECP | 27.05 | 1 mg | ✓ Baxter |

⇒SA2420 Special Authority for Subsidy

Initial application only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Patient is currently on treatment with trastuzumab deruxtecan and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 Patient has metastatic breast cancer expressing HER-2 IHC3+ or ISH+ (including FISH or other current technology); and
 - 2.2 Patient has previously received trastuzumab and chemotherapy, separately or in combination; and
 - 2.3 Fither:
 - 2.3.1 The patient has received prior therapy for metastatic disease; or
 - 2.3.2 The patient developed disease recurrence during, or within six months of completing adjuvant therapy; and
 - 2.4 Patient has a good performance status (ECOG 0-1); and
 - 2.5 Patient has not received prior funded trastuzumab deruxtecan treatment; and
 - 2.6 Treatment to be discontinued at disease progression.

| Subsidy | | Fully | Brand or |
|------------------------|-----|------------|--------------|
| (Manufacturer's Price) | | Subsidised | Generic |
| \$ | Per | ✓ | Manufacturer |

continued...

Renewal only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The cancer has not progressed at any time point during the previous approval period whilst on trastuzumab deruxtecan; and
- 2 Treatment to be discontinued at disease progression.

Note: Prior or adjuvant therapy includes anthracycline, other chemotherapy, biological drugs, or endocrine therapy.

TRASTUZUMAB EMTANSINE - PCT only - Specialist - Special Authority see SA2424 below

| Inj 100 mg vial | 2,320.00 | 1 | Kadcyla |
|------------------|----------|------|-----------|
| Inj 160 mg vial | 3,712.00 | 1 | ✓ Kadcyla |
| Inj 1 mg for ECP | 24.52 | 1 mg | ✓ Baxter |

⇒SA2424 Special Authority for Subsidy

Initial application — (early breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Patient has early breast cancer expressing HER2 IHC3+ or ISH+; and
- 2 Documentation of pathological invasive residual disease in the breast and/or axiliary lymph nodes following completion of
- 3 Patient has completed systemic neoadjuvant therapy with trastuzumab and chemotherapy prior to surgery; and
- 4 Disease has not progressed during neoadjuvant therapy; and
- 5 Patient has left ventricular ejection fraction of 45% or greater; and
- 6 Adjuvant treatment with trastuzumab emtansine to be commenced within 12 weeks of surgery; and
- 7 Trastuzumab emtansine to be discontinued at disease progression; and
- 8 Total adjuvant treatment duration must not exceed 42 weeks (14 cycles).

Initial application — (metastatic breast cancer) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Patient has previously received trastuzumab and chemotherapy, separately or in combination; and
- 3 Either:
 - 3.1 The patient has received prior therapy for metastatic disease*; or
 - 3.2 The patient developed disease recurrence during, or within six months of completing adjuvant therapy*; and
- 4 Patient has a good performance status (ECOG 0-1); and
- 5 Fither:
 - 5.1 Patient does not have symptomatic brain metastases; or
 - 5.2 Patient has brain metastases and has received prior local CNS therapy; and
- 6 Fither:
 - 6.1 Patient has not received prior funded trastuzumab emtansine or trastuzumab deruxtecan treatment; or

6.2 Both:

- 6.2.1 Patient has discontinued trastuzumab deruxtecan due to intolerance; and
- 6.2.2 The cancer did not progress while on trastuzumab deruxtecan; and
- 7 Treatment to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The cancer has not progressed at any time point during the previous approval period whilst on trastuzumab emtansine;
- 2 Treatment to be discontinued at disease progression.

Note: Prior or adjuvant therapy includes anthracycline, other chemotherapy, biological drugs, or endocrine therapy.

| (Man | Subsidy jufacturer's Price) | Fully Subsidised | Brand or Generic |
|------|--------------------------------|---------------------|---------------------|
| | \$ P | er 🗸 | Manufacturer |
| | | | |

USTEKINUMAB - Special Authority see SA2182 below - Retail pharmacy

Inj 90 mg per ml, 1 ml pre-filled syringe........................4,162.00 1 ✓ Stelara

⇒SA2182 Special Authority for Subsidy

Initial application — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Patient is currently on treatment with ustekinumab commenced prior to 1 February 2023 and met all remaining criteria (criterion 2) below at the time of commencing treatment; or
- 2 Both:
 - 2.1 Patient has active Crohn's disease; and
 - 2.2 Fither:
 - 2.2.1 Patient has had an initial approval for prior biologic therapy for Crohn's disease and has experienced intolerable side effects or insufficient benefit to meet renewal criteria; or
 - 2.2.2 Both:
 - 2.2.2.1 Patient meets the initiation criteria for prior biologic therapies for Crohn's disease; and
 - 2.2.2.2 Other biologics for Crohn's disease are contraindicated.

Renewal — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points, or HBI score has reduced by 3 points, from when the patient was initiated on biologic therapy; or
 - 1.2 CDAI score is 150 or less, or HBI is 4 or less; or
 - 1.3 The patient has experienced an adequate response to treatment, but CDAI score and/or HBI score cannot be assessed: and
- 2 Ustekinumab to be administered at a dose no greater than 90 mg every 8 weeks.

Initial application — (Crohn's disease - children*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Patient is currently on treatment with ustekinumab commenced prior to 1 February 2023 and met all remaining criteria (criterion 2) below at the time of commencing treatment; or
- 2 Both:
 - 2.1 Patient has active Crohn's disease; and
 - 2.2 Fither:
 - 2.2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria; or
 - 2.2.2 Both:
 - 2.2.2.1 Patient meets the initiation criteria for prior biologic therapies for Crohn's disease; and
 - 2.2.2.2 Other biologics for Crohn's disease are contraindicated.

Note: Indication marked with * is an unapproved indication.

Renewal — (Crohn's disease - children*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from when the patient was initiated on biologic therapy; or
 - 1.2 PCDAI score is 15 or less; or
 - 1.3 The patient has experienced an adequate response to treatment, but CDAI score cannot be assessed; and
- 2 Ustekinumab to administered at a dose no greater than 90 mg every 8 weeks.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

Note: Indication marked with * is an unapproved indication.

Initial application — (ulcerative colitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Patient is currently on treatment with ustekinumab commenced prior to 1 February 2023 and met all remaining criteria (criterion 2) below at the time of commencing treatment; or
- 2 Both
 - 2.1 Patient has active ulcerative colitis: and
 - 2.2 Either:
 - 2.2.1 Patient has had an initial approval for prior biologic therapy for ulcerative colitis and has experienced intolerable side effects or insufficient benefit to meet renewal criteria; or
 - 2.2.2 Both:
 - 2.2.2.1 Patient meets the initiation criteria for prior biologic therapies for ulcerative colitis; and
 - 2.2.2.2 Other biologics for ulcerative colitis are contraindicated.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 The SCCAI score has reduced by 2 points or more from the SCCAI score since initiation on biologic therapy; or
 - 1.2 PUCAI score has reduced by 10 points or more from the PUCAI score since initiation on biologic therapy*; and
- 2 Ustekinumab will be used at a dose no greater than 90 mg intravenously every 8 weeks.

Note: Criterion marked with * is for an unapproved indication.

VEDOLIZUMAB - PCT only - Special Authority see SA2183 below

⇒SA2183 Special Authority for Subsidy

Initial application — (**Crohn's disease - adults**) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria (unless contraindicated); or
 - 2.2 Patient has a CDAI score of greater than or equal to 300, or HBI score of greater than or equal to 10; or
 - 2.3 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.4 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
 - 2.5 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Any of the following:
 - 3.1 Patient has tried but experienced an inadequate response to (including lack of initial response and/or loss of initial response) from prior therapy with immunomodulators and corticosteroids; or
 - 3.2 Patient has experienced intolerable side effects from immunomodulators and corticosteroids; or
 - 3.3 Immunomodulators and corticosteroids are contraindicated.

Renewal — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

1 Any of the following:

| Subsidy | F | ully | Brand or | _ |
|------------------------|---------|------|--------------|---|
| (Manufacturer's Price) | Subsidi | sed | Generic | |
| \$ | Per | 1 | Manufacturer | |

continued...

- 1.1 CDAI score has reduced by 100 points, or HBI score has reduced by 3 points, from when the patient was initiated on biologic therapy; or
- 1.2 CDAI score is 150 or less, or HBI is 4 or less; or
- 1.3 The patient has experienced an adequate response to treatment, but CDAI score and/or HBI score cannot be assessed; and
- 2 Vedolizumab to administered at a dose no greater than 300 mg every 8 weeks.

Initial application — (Crohn's disease - children*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria (unless contraindicated); or
 - 2.2 Patient has a Paediatric Crohn's Disease Activity Index (PCDAI) score of greater than or equal to 30; or
 - 2.3 Patient has extensive small intestine disease; and
- 3 Any of the following:
 - 3.1 Patient has tried but experienced an inadequate response to (including lack of initial response and/or loss of initial response) from prior therapy with immunomodulators and corticosteroids; or
 - 3.2 Patient has experienced intolerable side effects from immunomodulators and corticosteroids; or
 - 3.3 Immunomodulators and corticosteroids are contraindicated.

Note: Indication marked with * is an unapproved indication.

Renewal — (Crohn's disease - children*) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from when the patient was initiated on biologic therapy; or
 - 1.2 PCDAI score is 15 or less: or
 - 1.3 The patient has experienced an adequate response to treatment, but CDAI score cannot be assessed; and
- 2 Vedolizumab to administered at a dose no greater than 300mg every 8 weeks.

Note: Indication marked with * is an unapproved indication.

Initial application — (ulcerative colitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active ulcerative colitis; and
- 2 Any of the following:
 - 2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria (unless contraindicated); or
 - 2.2 Patient has a SCCAI score is greater than or equal to 4; or
 - 2.3 Patient's PUCAI score is greater than or equal to 20*; and
- 3 Any of the following:
 - 3.1 Patient has tried but experienced an inadequate response to (including lack of initial response and/or loss of initial response) from prior therapy with immunomodulators and corticosteroids; or
 - 3.2 Patient has experienced intolerable side effects from immunomodulators and corticosteroids; or
 - 3.3 Immunomodulators and corticosteroids are contraindicated.

Note: Indication marked with * is an unapproved indication.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

| Subsidy | :) | Fully | Brand or |
|-----------------------|-----|------------|--------------|
| (Manufacturer's Price | | Subsidised | Generic |
| <u> </u> | Per | ✓ | Manufacturer |

continued...

Both:

- 1 Fither:
 - 1.1 The SCCAI score has reduced by 2 points or more from the SCCAI score since initiation on biologic therapy; or
 - 1.2 The PUCAI score has reduced by 10 points or more from the PUCAI score since initiation on biologic therapy *; and
- 2 Vedolizumab will be used at a dose no greater than 300 mg intravenously every 8 weeks.

Note: Indication marked with * is an unapproved indication.

Programmed Cell Death-1 (PD-1) Inhibitors

| ATEZOLIZUMAB - PCT only - Specialist - Special Authority see SA2264 bel | low | |
|---|------|-------------|
| Inj 60 mg per ml, 20 ml vial9,503.00 | 1 | ✓ Tecentriq |
| Inj 1 mg for ECP8.08 | 1 mg | ✓ Baxter |

⇒SA2264 Special Authority for Subsidy

Initial application — (non-small cell lung cancer second line monotherapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced or metastatic non-small cell lung cancer; and
- 2 Patient has not received prior funded treatment with an immune checkpoint inhibitor for NSCLC; and
- 3 For patients with non-squamous histology there is documentation confirming that the disease does not express activating mutations of EGFR or ALK tyrosine kinase unless not possible to ascertain; and
- 4 Patient has an ECOG 0-2; and
- 5 Patient has documented disease progression following treatment with at least two cycles of platinum-based chemotherapy; and
- 6 Atezolizumab is to be used as monotherapy at a dose of 1200 mg every three weeks (or equivalent) for a maximum of 16 weeks; and
- 7 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Renewal — (non-small cell lung cancer second line monotherapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Any of the following:
 - 1.1 Patient's disease has had a complete response to treatment; or
 - 1.2 Patient's disease has had a partial response to treatment; or
 - 1.3 Patient has stable disease; and
- 2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period; and
- 3 No evidence of disease progression; and
- 4 The treatment remains clinically appropriate and patient is benefitting from treatment; and
- 5 Atezolizumab to be used at a maximum dose of 1200 mg every three weeks (or equivalent); and
- 6 Treatment with atezolizumab to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

| DURVALUMAB – PCT only – Specialist – Special Author | rity see SA2425 on the ne | xt page | |
|---|---------------------------|---------|----------|
| Inj 50 mg per ml, 10 ml vial | 4,700.00 | 1 | Imfinzi |
| Inj 50 mg per ml, 2.4 ml vial | | 1 | Imfinzi |
| Inj 1 mg for ECP | | 1 mg | ✓ Baxter |

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA2425 Special Authority for Subsidy

Initial application — (Non-small cell lung cancer) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 Patient has histologically or cytologically documented stage III, locally advanced, unresectable non-small cell lung cancer (NSCLC): or
 - 1.2 Patient has histologically or cytologically documented stage IIb (T1N2a only), locally advanced, unresectable non-small cell lung cancer (NSCLC); and
- 2 Patient has received two or more cycles of platinum-based chemotherapy concurrently with definitive radiation therapy; and
- 3 Patient has no disease progression following the second or subsequent cycle of platinum-based chemotherapy with definitive radiation therapy treatment; and
- 4 Patient has a ECOG performance status of 0 or 1; and
- 5 Patient has completed last radiation dose within 8 weeks of starting treatment with durvalumab; and
- 6 Patient must not have received prior PD-1 or PD-L1 inhibitor therapy for this condition; and
- 7 Fither:
 - 7.1 Durvalumab is to be used at a maximum dose of no greater than 10 mg/kg every 2 weeks; or
 - 7.2 Durvalumab is to be used at a flat dose of 1500 mg every 4 weeks; and
- 8 Treatment with durvalumab to cease upon signs of disease progression.

Renewal — (Non-small cell lung cancer) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The treatment remains clinically appropriate and the patient is benefitting from treatment; and
- 2 Either:
 - 2.1 Durvalumab is to be used at a maximum dose of no greater than 10 mg/kg every 2 weeks; or
 - 2.2 Durvalumab is to be used at a flat dose of 1500 mg every 4 weeks; and
- 3 Treatment with durvalumab to cease upon signs of disease progression; and
- 4 Total continuous treatment duration must not exceed 12 months.

| NIVOLUMAB - PCT only - Specialist - Special Authority see SA2405 be | elow |
|---|-------------------------|
| Inj 10 mg per ml, 4 ml vial1,05 | 1.98 1 ✓ Opdivo |
| Inj 10 mg per ml, 10 ml vial2,62 | 9.96 1 ✓ Opdivo |
| Inj 1 mg for ECP2 | 7.62 1 mg Saxter |

⇒SA2405 Special Authority for Subsidy

Initial application only from a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Baseline measurement of overall tumour burden is documented clinically and radiologically; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Fither:
 - 4.1 Patient has not received funded pembrolizumab; or
 - 4.2 Both:
 - 4.2.1 Patient has received an initial Special Authority approval for pembrolizumab and has discontinued pembrolizumab within 12 weeks of starting treatment due to intolerance; and
 - 4.2.2 The cancer did not progress while the patient was on pembrolizumab; and
- 5 Documentation confirming that the patient has been informed and acknowledges that funded treatment with nivolumab will

| Subsidy | | Fully | Brand or | |
|------------------------|-----|------------|--------------|--|
| (Manufacturer's Price) | | Subsidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

continued...

not be continued if their disease progresses.

Renewal — (less than 24 months on treatment) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Patient's disease has had a complete response to treatment; or
 - 1.1.2 Patient's disease has had a partial response to treatment; or
 - 1.1.3 Patient has stable disease; and
 - 1.2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period: and
 - 1.3 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with nivolumab for reasons other than severe toxicity or disease progression; and
 - 2.2 Patient has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with nivolumab.

Renewal — (more than 24 months on treatment) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: Both:

- 1 Patient has been on treatment for more than 24 months; and
- 2 Either:
 - 2.1 All of the following:
 - 2.1.1 Any of the following:
 - 2.1.1.1 Patient's disease has had a complete response to treatment; or
 - 2.1.1.2 Patient's disease has had a partial response to treatment; or
 - 2.1.1.3 Patient has stable disease; and
 - 2.1.2 Response to treatment in target lesions has been determined by comparable radiologic or clinical assessment following the most recent treatment period; and
 - 2.1.3 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
 - 2.2 All of the following:
 - 2.2.1 Patient has previously discontinued treatment with nivolumab for reasons other than severe toxicity or disease progression; and
 - 2.2.2 Patient has signs of disease progression; and
 - 2.2.3 Disease has not progressed during previous treatment with nivolumab.

Initial application — (Renal cell carcinoma) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:
Either:

- 1 Patient is currently on treatment with nivolumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 Patient has metastatic renal-cell carcinoma; and
 - 2.2 The disease is of predominant clear-cell histology; and
 - 2.3 Patient has an ECOG performance score of 0-2; and
 - 2.4 Patient has documented disease progression following one or two previous regimens of antiangiogenic therapy; and
 - 2.5 Nivolumab is to be used as monotherapy at a maximum dose of 240 mg every 2 weeks (or equivalent) and discontinued at disease progression.

| Subsidy | Fully | Brand or | |
|------------------------|------------|--------------|--|
| (Manufacturer's Price) | Subsidised | Generic | |
| \$ | Per 🗸 | Manufacturer | |

continued...

Renewal — (Renal cell carcinoma) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 Patient's disease has had a complete response to treatment; or
 - 1.2 Patient's disease has had a partial response to treatment; or
 - 1.3 Patient has stable disease: and
- 2 No evidence of disease progression; and
- 3 Nivolumab is to be used as monotherapy at a maximum dose of 240 mg every 2 weeks (or equivalent) and discontinued at disease progression.

| PEMBROLIZUMAB - PCT only - Specialist - Special A | Authority see SA2386 below | | |
|---|----------------------------|----|----------|
| Inj 25 mg per ml, 4 ml vial | 4,680.00 | 1 | Keytruda |
| Inj 1 mg for ECP | 47.74 1 | mg | ✓ Baxter |

⇒SA2386 Special Authority for Subsidy

Initial application — (unresectable or metastatic melanoma) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Baseline measurement of overall tumour burden is documented clinically and radiologically; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Either:
 - 4.1 Patient has not received funded nivolumab; or
 - 4.2 Both:
 - 4.2.1 Patient has received an initial Special Authority approval for nivolumab and has discontinued nivolumab within 12 weeks of starting treatment due to intolerance; and
 - 4.2.2 The cancer did not progress while the patient was on nivolumab; and
- 5 Documentation confirming that the patient has been informed and acknowledges that funded treatment with pembrolizumab will not be continued if their disease progresses.

Renewal — (unresectable or metastatic melanoma, less than 24 months on treatment) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Patient's disease has had a complete response to treatment; or
 - 1.1.2 Patient's disease has had a partial response to treatment; or
 - 1.1.3 Patient has stable disease; and
 - 1.2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period; and
 - 1.3 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with pembrolizumab for reasons other than severe toxicity or disease progression; and
 - 2.2 Patient has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with pembrolizumab.

Renewal — (unresectable or metastatic melanoma, more than 24 months on treatment) only from a medical oncologist or

| Subsidy | | Fully | Brand or |
|------------------------|-----|------------|--------------|
| (Manufacturer's Price) | | Subsidised | Generic |
| \$ | Per | / | Manufacturer |

continued...

medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 Patient has been on treatment for more than 24 months; and
- 2 Either:
 - 2.1 All of the following:
 - 2.1.1 Any of the following:
 - 2.1.1.1 Patient's disease has had a complete response to treatment; or
 - 2.1.1.2 Patient's disease has had a partial response to treatment; or
 - 2.1.1.3 Patient has stable disease; and
 - 2.1.2 Response to treatment in target lesions has been determined by comparable radiologic or clinical assessment following the most recent treatment period; and
 - 2.1.3 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
 - 2.2 All of the following:
 - 2.2.1 Patient has previously discontinued treatment with pembrolizumab for reasons other than severe toxicity or disease progression; and
 - 2.2.2 Patient has signs of disease progression; and
 - 2.2.3 Disease has not progressed during previous treatment with pembrolizumab.

Initial application — (non-small cell lung cancer first-line monotherapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-small cell lung cancer; and
- 2 Patient has not had chemotherapy for their disease in the palliative setting; and
- 3 Patient has not received prior funded treatment with an immune checkpoint inhibitor for NSCLC; and
- 4 For patients with non-squamous histology there is documentation confirming that the disease does not express activating mutations of EGFR or ALK tyrosine kinase unless not possible to ascertain; and
- 5 Pembrolizumab to be used as monotherapy; and
- 6.1 There is documentation confirming the disease expresses PD-L1 at a level greater than or equal to 50% as determined by a validated test unless not possible to ascertain; or
 - 6.2 Both:
 - 6.2.1 There is documentation confirming the disease expresses PD-L1 at a level greater than or equal to 1% as determined by a validated test unless not possible to ascertain; and
 - 6.2.2 Chemotherapy is determined to be not in the best interest of the patient based on clinician assessment; and
- 7 Patient has an ECOG 0-2; and
- 8 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks; and
- 9 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Renewal — (non-small cell lung cancer first line monotherapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Any of the following:
 - 1.1 Patient's disease has had a complete response to treatment; or
 - 1.2 Patient's disease has had a partial response to treatment: or
 - 1.3 Patient has stable disease; and

| Subsidy | Fu | , | |
|------------------------|----------|----------------------------------|--|
| (Manufacturer's Price) | Subsidis | ed Generic | |
| \$ | Per | Manufacturer | |

continued...

- 2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period; and
- 3 No evidence of disease progression; and
- 4 The treatment remains clinically appropriate and patient is benefitting from treatment; and
- 5 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 6 Treatment with pembrolizumab to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initial application — (non-small cell lung cancer first-line combination therapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-small cell lung cancer; and
- 2 The patient has not had chemotherapy for their disease in the palliative setting; and
- 3 Patient has not received prior funded treatment with an immune checkpoint inhibitor for NSCLC; and
- 4 For patients with non-squamous histology there is documentation confirming that the disease does not express activating mutations of EGFR or ALK tyrosine kinase unless not possible to ascertain; and
- 5 Pembrolizumab to be used in combination with platinum-based chemotherapy; and
- 6 Patient has an ECOG 0-2; and
- 7 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks; and
- 8 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Renewal — (non-small cell lung cancer first line combination therapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 Patient's disease has had a complete response to treatment; or
 - 1.2 Patient's disease has had a partial response to treatment; or
 - 1.3 Patient has stable disease; and
- 2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period: and
- 3 No evidence of disease progression; and
- 4 The treatment remains clinically appropriate and patient is benefitting from treatment; and
- 5 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 6 Treatment with pembrolizumab to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initial application — (breast cancer, advanced) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: Fither:

- 1 Patient is currently on treatment with pembrolizumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has recurrent or de novo unresectable, inoperable locally advanced triple-negative breast cancer (that does not express ER, PR or HER2 IHC3+ or ISH+ [including FISH or other technology]); or
 - 2.1.2 Patient has recurrent or de novo metastatic triple-negative breast cancer (that does not express ER, PR or HER2 IHC3+ or ISH+ [including FISH or other technology]); and
 - 2.2 Patient is treated with palliative intent; and

| Subsidy | | Fully | Brand or | |
|------------------------|-----|------------|--------------|--|
| (Manufacturer's Price) | | Subsidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

continued...

- 2.3 Patient's cancer has confirmed PD-L1 Combined Positive Score (CPS) is greater than or equal to 10; and
- 2.4 Patient has received no prior systemic therapy in the palliative setting; and
- 2.5 Patient has an ECOG score of 0-2; and
- 2.6 Pembrolizumab is to be used in combination with chemotherapy; and
- 2.7 Baseline measurement of overall tumour burden is documented clinically and radiologically; and
- 2.8 Pembrolizumab is to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of

Renewal — (breast cancer, advanced) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 Patient's disease has had a complete response to treatment; or
 - 1.2 Patient's disease has had a partial response to treatment; or
 - 1.3 Patient has stable disease; and
- 2 No evidence of disease progression; and
- 3 Response to treatment in target lesions has been determined by a comparable radiologic assessment following the most recent treatment period; and
- 4 Pembrolizumab is to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 5 Treatment with pembrolizumab is to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initial application — (head and neck squamous cell carcinoma) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria: Either:

- 1 Patient is currently on treatment with pembrolizumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 Patient has recurrent or metastatic head and neck squamous cell carcinoma of mucosal origin (excluding nasopharyngeal carcinoma) that is incurable by local therapies; and
 - 2.2 Patient has not received prior systemic therapy in the recurrent or metastatic setting; and
 - 2.3 Patient has a positive PD-L1 combined positive score (CPS) of greater than or equal to 1; and
 - 2.4 Patient has an ECOG performance score of 0-2; and
 - 2.5 Either:
 - 2.5.1 Pembrolizumab to be used in combination with platinum-based chemotherapy; or
 - 2.5.2 Pembrolizumab to be used as monotherapy; and
- 2.6 Pembrolizumab is to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks.

Renewal — (head and neck squamous cell carcinoma) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 Patient's disease has had a complete response to treatment; or
 - 1.2 Patient's disease has had a partial response to treatment; or
 - 1.3 Patient has stable disease; and
- 2 No evidence of disease progression; and
- 3 Pembrolizumab is to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 4 Treatment with pembrolizumab is to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initial application — (MSI-H/dMMR advanced colorectal cancer) only from a relevant specialist or any relevant practitioner on

| Subsidy | F | ully | Brand or | _ |
|------------------------|--------|------|--------------|---|
| (Manufacturer's Price) | Subsid | sed | Generic | |
| \$ | Per | 1 | Manufacturer | |

continued

the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria: Fither:

- 1 Patient is currently on treatment with pembrolizumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 Fither:
 - 2.1.1 Patient has deficient mismatch repair (dMMR) or microsatellite instability-high (MSI-H) metastatic colorectal
 - 2.1.2 Patient has deficient mismatch repair (dMMR) or microsatellite instability-high (MSI-H) unresectable colorectal cancer; and
 - 2.2 Patient is treated with palliative intent: and
 - 2.3 Patient has not previously received funded treatment with pembrolizumab; and
 - 2.4 Patient has an ECOG performance score of 0-2; and
 - 2.5 Baseline measurement of overall tumour burden is documented clinically and radiologically; and
 - 2.6 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks.

Renewal — (MSI-H/dMMR advanced colorectal cancer) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 3 Treatment with pembrolizumab is to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initial application — (Urothelial carcinoma) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Patient is currently on treatment with pembrolizumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 Patient has inoperable locally advanced (T4) or metastatic urothelial carcinoma; and
 - 2.2 Patient has an ECOG performance score of 0-2; and
 - 2.3 Patient has documented disease progression following treatment with chemotherapy; and
 - 2.4 Pembrolizumab to be used as monotherapy at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks.

Renewal — (Urothelial carcinoma) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 Patient's disease has had a complete response to treatment; or
 - 1.2 Patient's disease has had a partial response to treatment; or
 - 1.3 Patient has stable disease; and
- 2 No evidence of disease progression; and
- 3 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 4 Treatment with pembrolizumab is to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initial application — (relapsed/refractory Hodgkin lymphoma) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

1 Patient is currently on treatment with pembrolizumab and met all remaining criteria prior to commencing treatment; or

| Subsidy | Fully | | Brand or |
|------------------------|------------|----------|----------|
| (Manufacturer's Price) | Subsidised | | Generic |
| \$ | Per | √ | |

continued...

- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Both:
 - 2.1.1.1 Patient has relapsed/refractory Hodgkin lymphoma after two or more lines of chemotherapy; and
 - 2.1.1.2 Patient is ineligible for autologous stem cell transplant; or
 - 2.1.2 Patient has relapsed/refractory Hodgkin lymphoma and has previously undergone an autologous stem cell transplant; and
 - 2.2 Patient has not previously received funded pembrolizumab; and
 - 2.3 Pembrolizumab to be administered at doses no greater than 200 mg once every 3 weeks.

Renewal — (relapsed/refractory Hodgkin lymphoma) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient has received a partial or complete response to pembrolizumab; and
- 2 Treatment with pembrolizumab is to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Other Immunosuppressants

| CICLOSPORIN | | | |
|--|---------------|----------|------------|
| Cap 25 mg | 44.63 | 50 | Neoral |
| Cap 50 mg | 88.91 | 50 | Neoral |
| Cap 100 mg | | 50 | ✓ Neoral |
| Oral liq 100 mg per ml | | 50 ml OP | ✓ Neoral |
| EVEROLIMUS - Special Authority see SA2414 below - Re | tail pharmacy | | |
| Wastage claimable | | | |
| Tab 10 mg | 6,512.29 | 30 | ✓ Afinitor |
| Tab 5 mg | 4,555.76 | 30 | ✓ Afinitor |
| | | | |

⇒SA2414 Special Authority for Subsidy

Initial application only from a neurologist or oncologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has tuberous sclerosis: and
- 2 Patient has progressively enlarging sub-ependymal giant cell astrocytomas (SEGAs) that require treatment.

Renewal only from a neurologist or oncologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Documented evidence of SEGA reduction or stabilisation by MRI within the last 3 months; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Everolimus to be discontinued at progression of SEGAs.

Initial application — (renal cell carcinoma) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 All of the following:
 - 1.1 The patient has metastatic renal cell carcinoma; and
 - 1.2 The disease is of predominant clear-cell histology; and
 - 1.3 The patient has documented disease progression following one previous line of treatment; and
 - 1.4 The patient has an ECOG performance status of 0-2; and

| Subsidy (Manufacturer's Price) | Subsid | Fully dised | Brand or Generic |
|-----------------------------------|--------|----------------|---------------------|
| \$ | Per | | Manufacturer |

continued...

- 1.5 Everolimus is to be used in combination with lenvatinib; or
- 2 All of the following:
 - 2.1 Patient has received funded treatment with nivolumab for the second line treatment of metastatic renal cell carcinoma; and
 - 2.2 Patient has experienced treatment limiting toxicity from treatment with nivolumab; and
 - 2.3 Everolimus is to be used in combination with lenvatinib; and
 - 2.4 There is no evidence of disease progression.

Renewal — (renal cell carcinoma) from any relevant practitioner. Approvals valid for 4 months where there is no evidence of disease progression.

| OLIMUS – Special Authority see SA2270 below – Retail pharmacy | | | |
|---|---------|-------------------|------|
| Tab 1 mg749. | 99 10 | 0 ✓ Rapa | mune |
| Tab 2 mg | 99 10 | 00 🗸 Rapa | mune |
| Oral liq 1 mg per ml449. | 99 60 m | IOP ✓ Rapa | mune |

⇒SA2270 Special Authority for Subsidy

Initial application from any medical practitioner. Approvals valid without further renewal unless notified where the drug is to be used for rescue therapy for an organ transplant recipient.

Notes: Rescue therapy defined as unresponsive to calcineurin inhibitor treatment as defined by refractory rejection; or intolerant to calcineurin inhibitor treatment due to any of the following:

- GFR< 30 ml/min; or
- · Rapidly progressive transplant vasculopathy; or
- Rapidly progressive obstructive bronchiolitis: or
- . HUS or TTP: or
- Leukoencepthalopathy: or
- Significant malignant disease

Initial application — (severe non-malignant lymphovascular malformations*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe non-malignant lymphovascular malformation*; and
- 2 Any of the following:
 - 2.1 Malformations are not adequately controlled by sclerotherapy and surgery; or
 - 2.2 Malformations are widespread/extensive and sclerotherapy and surgery are not considered clinically appropriate; or
 - 2.3 Sirolimus is to be used to reduce malformation prior to consideration of surgery; and
- 3 Patient is being treated by a specialist lymphovascular malformation multi-disciplinary team; and
- 4 Patient has measurable disease as defined by RECIST version 1.1 (see Note).

Renewal — (severe non-malignant lymphovascular malformations*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Patient's disease has had either a complete response or a partial response to treatment, or patient has stable disease according to RECIST version 1.1 (see Note); or
 - 1.2 Patient's disease has stabilised or responded clinically and disease response to treatment has been clearly documents in patient notes; and
- 2 No evidence of progressive disease: and
- 3 The treatment remains clinically appropriate and the patient is benefitting from the treatment.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer et al. Eur J Cancer 2009;45:228-47)

| Subsidy | | Fully | Brand or |
|------------------------|-----|------------|--------------|
| (Manufacturer's Price) | | Subsidised | Generic |
| <u> </u> | Per | | Manufacturer |

continued...

Indications marked with * are unapproved indications

Initial application — (renal angiomyolipoma(s) associated with tuberous sclerosis complex*) only from a nephrologist or urologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient has tuberous sclerosis complex*; and
- 2 Evidence of renal angiomyolipoma(s) measuring 3 cm or greater and that have shown interval growth.

Renewal — (renal angiomyolipoma(s) associated with tuberous sclerosis complex*) from any relevant practitioner.

Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Documented evidence of renal angiomyolipoma reduction or stability by magnetic resonance imaging (MRI) or ultrasound;
- 2 Demonstrated stabilisation or improvement in renal function; and
- 3 The patient has not experienced angiomyolipoma haemorrhage or significant adverse effects to sirolimus treatment; and
- 4 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indications marked with * are unapproved indications

Initial application — (refractory seizures associated with tuberous sclerosis complex*) only from a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has epilepsy with a background of documented tuberous sclerosis complex; and
- 2 Fither:
 - 2.1 Both:
 - 2.1.1 Vigabatrin has been trialled and has not adequately controlled seizures; and
 - 2.1.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least two of the following: sodium valproate, topiramate, levetiracetam. carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); or
 - 2.2 Both:
 - 2.2.1 Vigabatrin is contraindicated; and
 - 2.2.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from. optimal treatment with at least three of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); and
- 3 Seizures have a significant impact on quality of life; and
- 4 Patient has been assessed and surgery is considered inappropriate for this patient, or the patient has been assessed and would benefit from mTOR inhibitor treatment prior to surgery.

Note: Those of childbearing age potential are not required to trial phenytoin sodium, sodium valproate, or topiramate. Those who can father children are not required to trial sodium valproate.

Renewal — (refractory seizures associated with tuberous sclerosis complex*) only from a neurologist. Approvals valid for 12 months where demonstrated significant and sustained improvement in seizure rate (e.g. 50% reduction in seizure frequency) or severity and/or patient quality of life compared with baseline prior to starting sirolimus treatment.

Note: Indications marked with * are unapproved indications

TACROLIMUS - Special Authority see SA2271 below - Retail pharmacy

| ACTIOE Openial Authority Sec CALETT BOILD 11 | otali priarriacy | | |
|--|------------------|-----|---------------------|
| Cap 0.5 mg | 49.60 | 100 | ✓ Tacrolimus Sandoz |
| Cap 0.75 mg | 99.30 | 100 | ✓ Tacrolimus Sandoz |
| Cap 1 mg | 84.30 | 100 | ✓ Tacrolimus Sandoz |
| Cap 5 mg | 248.20 | 50 | ✓ Tacrolimus Sandoz |

⇒SA2271 Special Authority for Subsidy

Initial application — (organ transplant) only from a relevant specialist. Approvals valid without further renewal unless notified

ONCOLOGY AGENTS AND IMMUNOSUPPRESSANTS

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

continued...

where the patient is an organ transplant recipient.

Note: Subsidy applies for either primary or rescue therapy.

Initial application — (non-transplant indications*) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient requires long-term systemic immunosuppression; and
- - 2.1 Ciclosporin has been trialled and discontinued treatment because of unacceptable side effects or inadequate clinical response: or
 - 2.2 Patient is a child with nephrotic syndrome*.

Note: Indications marked with * are unapproved indications

JAK inhibitors

UPADACITINIB - Special Authority see SA2079 below - Retail pharmacy 28 ✓ RINVOQ

⇒SA2079 Special Authority for Subsidy

Initial application — (Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Fither:
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis: and
- 3 Either:
 - 3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or
 - 3.2 Both:
 - 3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a Health NZ Hospital; and

3.2.2 Either:

- 3.2.2.1 The patient has experienced intolerable side effects from rituximab; or
- 3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Renewal — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Either:

- - 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per

✓ Manufacturer

Antiallergy Preparations

Allergic Emergencies

ADRENALINE - Special Authority see SA2185 below - Retail pharmacy

- a) Maximum of 2 ini per prescription
- Additional prescriptions limited to replacement of up to two devices prior to expiry, or replacement of used device for treatment of anaphylaxis.

| Inj 0.15 mg per 0.3 ml auto-injector | 90.00 | 1 OP | Epipen Jr |
|--------------------------------------|-------|------|-----------|
| Inj 0.3 mg per 0.3 ml auto-injector | 90.00 | 1 OP | Epipen |

⇒SA2185 Special Authority for Subsidy

Initial application — (anaphylaxis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Patient has experienced an anaphylactic reaction which has resulted in presentation to a hospital or emergency department; or
 - 1.2 Patient has been assessed to be at significant risk of anaphylaxis by a relevant practitioner; and
- 2 Patient is not to be prescribed more than two devices in initial prescription.

⇒SA1558 Special Authority for Subsidy

Initial application only from a clinical immunologist or relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Supply for anticipated emergency treatment of laryngeal/oro-pharyngeal or severe abdominal attacks of acute hereditary angioedema (HAE) for patients with confirmed diagnosis of C1-esterase inhibitor deficiency; and
- 2 The patient has undergone product training and has agreed upon an action plan for self-administration.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Allergy Desensitisation

⇒SA1367 Special Authority for Subsidy

Initial application only from a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 RAST or skin test positive; and
- 2 Patient has had severe generalised reaction to the sensitising agent.

Renewal only from a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

| | Subsidy | , , | Fully Brand or |
|---|--------------------------|-----------------|-------------------------------------|
| | (Manufacturer's Price \$ | e) Subsi Per | idised Generic Manufacturer |
| BEE VENOM ALLERGY TREATMENT – Special Authority see S | A1367 on the previ | | |
| Initiation kit - 1 vial freeze dried venom with diluent | | 1 OP | ✓ VENOX \$29 |
| Initiation kit - 5 vials freeze dried venom with diluent | | 1 OP | ✓ VENOX S29 |
| Maintenance kit - 1 vial freeze dried venom with diluent | | 1 OP | ✓ VENOX \$29 |
| | 305.00 | TOP | A AEMOX 258 |
| Maintenance kit - 6 vials 120 mcg freeze dried venom, with diluent | 205.00 | 1 OP | ✓ Venomil S29 |
| | 205.00 | TOP | Venomin 329 |
| Treatment kit - 1 vial 550 mcg freeze dried venom, 1 diluent 9 ml, 3 diluent 1.8 ml | 334 30 | 1 OP | ✓ Albey |
| Treatment kit - 1 vial 550 mcg freeze dried venom, with diluer | | 1 OP | ✓ Hymenoptera S29 |
| (VENOX \$29) Initiation kit - 5 vials freeze dried venom with diluer | | - | • Hymenoptera 323 |
| • | | • / | |
| WASP VENOM ALLERGY TREATMENT - Special Authority see | SA1367 on the pre | evious page | Retail pharmacy |
| Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze | 000.00 | 4.00 | 4 A11 |
| dried polistes venom, 1 diluent 9 ml, 3 diluent 1.8 ml | 382.23 | 1 OP | ✓ Albey |
| Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze | 005.00 | 4.00 | / 11 |
| dried venom, with diluent | 305.00 | 1 OP | ✓ Hymenoptera S29 |
| Treatment kit (Paper wasp venom) - 6 vials 120 mcg freeze | 005.00 | 4 OD | / Vanamil oo |
| dried venom, with diluent | 305.00 | 1 OP | ✓ Venomil ©29 |
| Treatment kit (Yellow Jacket venom) - 1 vial 550 mcg freeze | 205.00 | 1 OD | ./ Uumanantara (200 |
| dried venom, with diluent | 305.00 | 1 OP | ✓ Hymenoptera S29 |
| Treatment kit (Yellow jacket venom) - 1 vial 550 mcg freeze | 421.04 | 1 OP | ✓ Albey |
| dried vespula venom, 1 diluent 9 ml, 3 diluent 1.8 ml Treatment kit (Yellow jacket venom) - 6 vials 120 mcg freeze | 431.24 | TOP | ▼ Albey |
| dried venom, with diluent 6 viais 120 mcg freeze | 305.00 | 1 OP | ✓ Venomil S29 |
| unca venom, with andert | 000.00 | 1 01 | · venonini es |
| Antihistamines | | | |
| | | | |
| CETIRIZINE HYDROCHLORIDE | | | 4 - · · |
| * Tab 10 mg | | 100 | ✓ Zista |
| * Oral liq 1 mg per ml | 3.99 | 200 ml | ✓ Histaclear |
| DEXTROCHLORPHENIRAMINE MALEATE | | | |
| * Tab 2 mg | | 40 | |
| | (8.40) | 00 | Polaramine |
| | 1.01 | 20 | Polaramine |
| * Oral liq 2 mg per 5 ml | (5.99) | 100 ml | Polaramine |
| 本 Oral liq 2 ftig per 5 ftil | (10.29) | 100 1111 | Polaramine |
| EEVOEEN A DINIE LIVEDOOUII ODIDE | (10.23) | | i diaramino |
| FEXOFENADINE HYDROCHLORIDE | 4.04 | 00 | |
| * Tab 60 mg | (8.23) | 20 | Telfast |
| * Tab 120 mg | ` ' | 30 | ✓ Fexaclear |
| - 100 120 mg | 14.22 | 00 | - I SAUGICUI |
| | (26.44) | | Telfast |
| * Tab 180 mg | ` ' | 30 | ✓ Fexaclear |
| (Telfast Tab 120 mg to be delisted 1 July 2025) | | | |
| LORATADINE | | | |
| * Tab 10 mg | 1.78 | 100 | ✓ Lorafix |
| * Oral liq 1 mg per ml | | 100 ml | ✓ Haylor syrup |
| 1 01 | | | , , , , , |

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

| | Subsidy | | Fully | Brand or |
|---|-----------------------|-------------|-------------|-------------------------|
| | (Manufacturer's \$ | Price) Subs | idised | Generic Manufacturer |
| ROMETHAZINE HYDROCHLORIDE | Ψ | 101 | | Warrandotarer |
| Tab 10 mg | 1 20 | 50 | √ ∧1 | lersoothe |
| · · · · · · · · · · · · · · · · · · · | | 50 50 | | lersoothe |
| Tab 25 mg Oral lig 1 mg per 1 ml | | 100 ml | | lersoothe |
| Oral liq 1 mg per 1 ml | 10.47 | 100 1111 | | energan Elixir |
| Inj 25 mg per ml, 2 ml ampoule - Up to 5 inj available on a F | | 5 | ✓ Ho | |
| nhaled Corticosteroids | | | | |
| | | | | |
| CLOMETHASONE DIPROPIONATE | 14.04 | 200 doos OD | ✓ Qı | ·0* |
| Aerosol inhaler, 50 mcg per dose | | 200 dose OP | | |
| Aerosol inhaler, 50 mcg per dose CFC-free | | 200 dose OP | | clazone 50 |
| Aerosol inhaler, 100 mcg per dose | | 200 dose OP | ✓ Q\ | |
| Aerosol inhaler, 100 mcg per dose CFC-free | | 200 dose OP | | clazone 100 |
| Aerosol inhaler, 250 mcg per dose CFC-free | 22.67 | 200 dose OP | ▼ B6 | clazone 250 |
| JDESONIDE | 17.00 | 000 daaa OD | / D. | lual a a ut |
| Powder for inhalation, 100 mcg per dose | 17.00 | 200 dose OP | | ılmicort Furbuhaler |
| Powder for inhalation, 200 mcg per dose | 10.00 | 200 dose OP | | Imicort |
| rowder for initialation, 200 mag per dose | 19.00 | 200 dose OF | | Turbuhaler |
| Powder for inhalation, 400 mcg per dose | 32.00 | 200 dose OP | | Imicort |
| | | | | Turbuhaler |
| UTICASONE | | | | |
| Aerosol inhaler, 50 mcg per dose | 7.19 | 120 dose OP | ✓ Fli | xotide |
| Powder for inhalation, 50 mcg per dose | | 60 dose OP | ✓ Fli | xotide Accuhaler |
| Powder for inhalation, 100 mcg per dose | | 60 dose OP | ✓ Fli | xotide Accuhaler |
| Aerosol inhaler, 125 mcg per dose | | 120 dose OP | ✓ Fli | xotide |
| Aerosol inhaler, 250 mcg per dose | | 120 dose OP | | xotide |
| Powder for inhalation, 250 mcg per dose | | 60 dose OP | | xotide Accuhaler |
| nhaled Long-acting Beta-adrenoceptor Agonisi | is | | | |
| FORMOTEROL FUMARATE DIHYDRATE | | | | |
| Powder for inhalation 4.5 mcg per dose, breath activated | | | | |
| (equivalent to eformoterol fumarate 6 mcg metered dose |) 10.32 | 60 dose OP | | |
| (equivalent to elorinoterol lumarate o mog metered dose | (16.90) | ou dose OF | 0, | ris Turbuhaler |
| DACATEROL | (10.50) | | 0) | io i uibuilalei |
| DACATEROL | 04.00 | 00 -1 05 | | . h D |
| Powder for inhalation 150 mcg | | 30 dose OP | | brez Breezhaler |
| Powder for inhalation 300 mcg | 61.00 | 30 dose OP | ✓ Or | brez Breezhaler |
| ALMETEROL | | | | |
| Aerosol inhaler CFC-free, 25 mcg per dose | | 120 dose OP | ✓ Se | revent |
| Powder for inhalation, 50 mcg per dose, breath activated | 26.25 | 60 dose OP | ✓ Se | revent Accuhaler |

| (I | Subsidy Manufacturer's Price) | | Fully idised | Brand or Generic | |
|----|----------------------------------|-----|--------------|---------------------|--|
| | \$ | Per | | Manufacturer | |
| | | | | | |

Inhaled Corticosteroids with Long-Acting Beta-Adrenoceptor Agonists

| minated obtained steroids with Long-Acting Deta-Adrenocopt | ioi Agoilisis | |
|---|----------------------------|--|
| BUDESONIDE WITH EFORMOTEROL | | |
| Powder for inhalation 160 mcg with 4.5 mcg eformoterol | | |
| fumarate per dose (equivalent to 200 mcg budesonide with | | 4 |
| 6 mcg eformoterol fumarate metered dose)41.50 | 120 dose OP | DuoResp Spiromax |
| Powder for inhalation 320 mcg with 9 mcg eformoterol fumarate | | |
| per dose (equivalent to 400 mcg budesonide with 12 mcg | | |
| eformoterol fumarate metered dose) – No more than 2 | 400 de OD | (D D O |
| dose per day | 120 dose OP 120 dose OP | ✓ DuoResp Spiromax✓ Vannair |
| Aerosol inhaler 100 mcg with eformoterol fumarate 6 mcg | 120 dose OP | ✓ vannair✓ Symbicort |
| Powder for illinatation 100 micg with elormoteror furnarate 6 micg55.74 | 120 dose OF | Turbuhaler 100/6 |
| Aerosol inhaler 200 mcg with eformoterol fumarate 6 mcg21.40 | 120 dose OP | ✓ Vannair |
| Powder for inhalation 200 mcg with eformoterol furnarate 6 mcg33.74 | 120 dose OP | ✓ Symbicort |
| 1 owder for illinatation 200 mag with elormoteror furnarate of mag55.74 | 120 dose O1 | Turbuhaler 200/6 |
| Powder for inhalation 400 mcg with eformoterol fumarate | | Turbunaler 200/0 |
| 12 mcg - No more than 2 dose per day | 60 dose OP | ✓ Symbicort |
| 12 mag | 00 0000 01 | Turbuhaler 400/12 |
| FLUTICASONE FUROATE WITH VILANTEROL | | |
| Powder for inhalation 100 mcg with vilanterol 25 mcg44.08 | 30 dose OP | ✓ Breo Ellipta |
| 5 | 30 dose Oi | • Breo Empta |
| FLUTICASONE WITH SALMETEROL | 100 dasa OD | / Carrellala |
| Aerosol inhaler 50 mcg with salmeterol 25 mcg | 120 dose OP | ✓ Seretide✓ Seretide |
| Aerosol inhaler 125 mcg with salmeterol 25 mcg32.60 | 120 dose OP | Serelide |
| Powder for inhalation 100 mcg with salmeterol 50 mcg – No | 60 dose OP | ✓ Seretide Accuhaler |
| more than 2 dose per day | 60 dose OP | Serelide Accunaler |
| Powder for inhalation 250 mcg with salmeterol 50 mcg - No more than 2 dose per day44.08 | 60 dose OP | ✓ Seretide Accuhaler |
| 11101e that12 dose per day44.06 | 60 dose OF | Serelide Accumaler |
| Beta-Adrenoceptor Agonists | | |
| SALBUTAMOL | | |
| Oral liq 400 mcg per ml50.00 | 150 ml | ✓ Ventolin |
| Ventolin to be Principal Supply on 1 May 2025 | 130 1111 | Ventonii |
| Infusion 1 mg per ml, 5 ml | 10 | ✓ Ventolin |
| Inj 500 mcg per ml, 1 ml – Up to 5 inj available on a PSO130.00 | 5 | ✓ Ventolin |
| ., | | |
| Inhaled Beta-Adrenoceptor Agonists | | |
| SALBUTAMOL | | |
| Aerosol inhaler, 100 mcg per dose CFC free - Up to 1000 | | |
| dose available on a PSO4.18 | 200 dose OP | ✓ SalAir |
| (6.80) | | Ventolin |
| Nebuliser soln, 1 mg per ml, 2.5 ml ampoule - Up to 30 neb | | |
| available on a PSO8.96 | 20 | ✓ Asthalin |
| Nebuliser soln, 2 mg per ml, 2.5 ml ampoule - Up to 30 neb | | |
| available on a PSO9.43 | 20 | ✓ Asthalin |
| TERBUTALINE SULPHATE | | |
| Powder for inhalation, 200 mcg per dose (equivalent to | | |
| 250 mcg metered dose), breath activated | 120 dose OP | ✓ Bricanyl Turbuhaler |
| | | |

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✓ Manufacturer

✓ Pharmascience S29

✓ Univent

Anticholinergic Agents

(Pharmascience S29 Nebuliser soln, 250 mcg per ml, 2 ml ampoule to be delisted 1 May 2025)

Inhaled Beta-Adrenoceptor Agonists with Anticholinergic Agents

SALBUTAMOL WITH IPRATROPIUM BROMIDE

✓ Duolin Cipla S29

Long-Acting Muscarinic Antagonists

GLYCOPYRRONIUM - Subsidy by endorsement

- a) Inhaled glycopyrronium treatment will not be subsidised if patient is also receiving treatment with subsidised tiotropium or umeclidinium.
- b) Glycopyrronium powder for inhalation 50 mcg per dose is subsidised only for patients who have been diagnosed as having COPD using spirometry if spirometry is possible, and the prescription is endorsed accordingly.

(Duolin Cipla \$29 Nebuliser soln, 2.5 mg with ipratropium bromide 0.5 mg per vial, 2.5 ml ampoule to be delisted 1 April 2025)

TIOTROPIUM BROMIDE - Subsidy by endorsement

- a) Tiotropium treatment will not be subsidised if patient is also receiving treatment with subsidised inhaled glycopyrronium or umeclidinium.
- b) Tiotropium bromide is subsidised only for patients who have been diagnosed as having COPD using spirometry if spirometry is possible, and the prescription is endorsed accordingly. Patients who had tiotropium dispensed before 1 October 2018 with a valid Special Authority are deemed endorsed.

UMECLIDINIUM - Subsidy by endorsement

- a) Umeclidinium will not be subsidised if patient is also receiving treatment with subsidised inhaled glycopyrronium or tiotropium bromide.
- b) Umeclidinium powder for inhalation 62.5 mcg per dose is subsidised only for patients who have been diagnosed as having COPD using spirometry if spirometry is possible, and the prescription is endorsed accordingly.

Subsidy (Manufacturer's Price) \$ Per

Subsidised

Fully

Brand or Generic Manufacturer

Long-Acting Muscarinic Antagonists with Long-Acting Beta-Adrenoceptor Agonists

Combination long acting muscarinic antagonist and long acting beta-2 agonist will not be subsidised if patient is also receiving treatment with a combination inhaled corticosteroid and long acting beta-2 agonist.

⇒SA1584 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Patient has been stabilised on a long acting muscarinic antagonist; and
- 2 The prescriber considers that the patient would receive additional benefit from switching to a combination product.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Patient is compliant with the medication; and
- 2 Patient has experienced improved COPD symptom control (prescriber determined).

GLYCOPYRRONIUM WITH INDACATEROL - Special Authority see SA1584 above - Retail pharmacy

Powder for Inhalation 50 mcg with indacaterol 110 mcg......81.00 30 dose OP ✓ Ultibro Breezhaler

TIOTROPIUM BROMIDE WITH OLODATEROL - Special Authority see SA1584 above - Retail pharmacy

UMECLIDINIUM WITH VILANTEROL - Special Authority see SA1584 above - Retail pharmacy

Powder for inhalation 62.5 mcg with vilanterol 25 mcg77.00 30 dose OP ✓ Anoro Ellipta

Inhaled Corticosteroid with Long-Acting Muscarinic Antagonist and Beta Agonist

 ${\tt BUDESONIDE\ WITH\ GLYCOPYRRONIUM\ AND\ EFORMOTEROL\ -Special\ Authority\ see\ SA2421\ below\ -\ Retail\ pharmacy}$

Aerosol inhaler budesonide 160 mcg with glycopyrronium

7.2 mcg and formoterol 5 mcg per dose......79.15 120 dose OP

✓ Breztri Aerosphere

⇒SA2421 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has a diagnosis of COPD confirmed by spirometry or spirometry has been attempted and technically acceptable results are not possible; and
- 2 Fither:
 - 2.1 Both:
 - 2.1.1 Patient is currently receiving an inhaled corticosteroid with long acting beta-2 agonist (ICS/LABA) or a long acting muscarinic antagonist with long acting beta-2 agonist (LAMA/LABA); and
 - 2.1.2 Any of the following:

Clinical criteria:

- 2.1.2.1 Patient has a COPD Assessment Test (CAT) score greater than 10; or
- 2.1.2.2 Patient has had 2 or more exacerbations in the previous 12 months; or
- 2.1.2.3 Patient has had one exacerbation requiring hospitalisation in the previous 12 months; or
- 2.1.2.4 Patient has had an eosinophil count greater than or equal to 0.3×10^9 cells/L in the previous 12 months; or
- 2.2 Patient is currently receiving multiple inhaler triple therapy (inhaled corticosteroid with long-acting muscarinic antagonist and long-acting beta-2 agonist ICS/LAMA/LABA) and met at least one of the clinical criteria above prior to commencing multiple inhaler therapy.

FLUTICASONE FUROATE WITH UMECLIDINIUM AND VILANTEROL – Special Authority see SA2326 below – Retail pharmacy Powder for inhalation fluticasone furoate 100 mcg with

umeclidinium 62.5 mcg and vilanterol 25 mcg.......104.24 30 dose OP ✓ Trelegy Ellipta

| Subsidy | | Fully | Brand or | |
|------------------------|-----|-----------|--------------|--|
| (Manufacturer's Price) | Su | ıbsidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

⇒SA2326 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has a diagnosis of COPD confirmed by spirometry or spirometry has been attempted and technically acceptable results are not possible; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient is currently receiving an inhaled corticosteroid with long acting beta-2 agonist (ICS/LABA) or a long acting muscarinic antagonist with long acting beta-2 agonist (LAMA/LABA); and
 - 2.1.2 Any of the following:

Clinical criteria:

- 2.1.2.1 Patient has a COPD Assessment Test (CAT) score greater than 10; or
- 2.1.2.2 Patient has had 2 or more exacerbations in the previous 12 months; or
- 2.1.2.3 Patient has had one exacerbation requiring hospitalisation in the previous 12 months; or
- 2.1.2.4 Patient has had an eosinophil count greater than or equal to 0.3×10^9 cells/L in the previous 12 months; or
- 2.2 Patient is currently receiving multiple inhaler triple therapy (inhaled corticosteroid with long acting muscarinic antagonist and long acting beta-2 agonist ICS/LAMA/LABA) and met at least one of the clinical criteria above prior to commencing multiple inhaler triple therapy.

Antifibrotics

NINTEDANIB - Special Authority see SA2012 below - Retail pharmacy

| Note: Nintedanib not subsidised in combination with subsidise | | | |
|---|---|-------|------------------------|
| Cap 100 mg | • | 60 OP | Ofev |

⇒SA2012 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Nintedanib is to be discontinued at disease progression (See Note); and
- 4 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 5 Any of the following:
 - 5.1 The patient has not previously received treatment with pirfenidone; or
 - 5.2 Patient has previously received pirfenidone, but discontinued pirfenidone within 12 weeks due to intolerance; or
 - 5.3 Patient has previously received pirfenidone, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with pirfenidone).

Renewal — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 3 Nintedanib is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

| | Subsidy (Manufacturer's Price) | | Fully sidised | Brand or Generic Manufacturer | |
|--|-----------------------------------|-------------|---------------|-------------------------------------|--|
| PIRFENIDONE – Retail pharmacy-Specialist – Special Authority Note: Pirfenidone is not subsidised in combination with subs | | | | | |
| Tab 801 mg | | 90 OP 90 | _ | sbriet sbriet | |

⇒SA2013 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note); and
- 4 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 5 Any of the following:
 - 5.1 The patient has not previously received treatment with nintedanib; or
 - 5.2 Patient has previously received nintedanib, but discontinued nintedanib within 12 weeks due to intolerance; or
 - 5.3 Patient has previously received nintedanib, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with nintedanib).

Renewal — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

Leukotriene Receptor Antagonists

| MC | NTELUKAST | | |
|----|-----------|----|-----------------------|
| * | Tab 4 mg | 28 | ✓ Montelukast Viatris |
| * | Tab 5 mg | 28 | ✓ Montelukast Viatris |
| * | Tab 10 mg | 28 | ✓ Montelukast Viatris |

Methylxanthines

AMINOPHYLLINE

| * | Inj 25 mg per ml, 10 ml ampoule — Up to 5 inj available on a PSO180.00 | 5 | ✓ DBL Aminophylline |
|----|--|--------|---------------------|
| TH | EOPHYLLINE | | |
| * | Tab long-acting 250 mg24.90 | 100 | ✓ Nuelin-SR |
| * | Oral liq 80 mg per 15 ml17.95 | 500 ml | ✓ Nuelin |

Mucolytics

| DORNASE ALFA – Special Authority see SA1978 on the next | page – Retail pharma | су | |
|---|----------------------|----|-------------|
| Nebuliser soln, 2.5 mg per 2.5 ml ampoule | 250.00 | 6 | ✓ Pulmozyme |

⇒SA1978 Special Authority for Subsidy

Initial application — (cystic fibrosis) only from a respiratory physician or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

| | Subsidy | F | ully | Brand or |
|-----|----------------------|---------|------|--------------|
| (Ma | anufacturer's Price) | Subsidi | sed | Generic |
| | \$ | Per | ✓ | Manufacturer |

continued...

All of the following:

- 1 Patient has a confirmed diagnosis of cystic fibrosis; and
- 2 Patient has previously undergone a trial with, or is currently being treated with, hypertonic saline; and
- 3 Any of the following:
 - 3.1 Patient has required one or more hospital inpatient respiratory admissions in the previous 12 month period; or
 - 3.2 Patient has had 3 exacerbations due to CF, requiring oral or intravenous (IV) antibiotics in the previous 12 month
 - 3.3 Patient has had 1 exacerbation due to CF, requiring oral or IV antibiotics in the previous 12 month period and a Brasfield score of < 22/25: or
 - 3.4 Patient has a diagnosis of allergic bronchopulmonary aspergillosis (ABPA).

Renewal — (cystic fibrosis) only from a respiratory physician or paediatrician. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient continues to benefit from treatment.

ELEXACAFTOR WITH TEZACAFTOR, IVACAFTOR AND IVACAFTOR - PCT only - Special Authority see SA2196 below

Tab elexacaftor 50 mg with tezacaftor 25 mg, ivacaftor 37.5 mg (56) and ivacaftor 75 mg (28)27,647.39 84 OP ✓ Trikafta Tab elexacaftor 100 mg with tezacaftor 50 mg, ivacaftor 75 mg ✓ Trikafta (56) and ivacaftor 150 mg (28)27,647.39 84 OP

⇒SA2196 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with cystic fibrosis; and
- 2 Patient is 6 years of age or older; and
- 3 Fither:
 - 3.1 Patient has two cystic fibrosis-causing mutations in the cystic fibrosis transmembrane regulator (CFTR) gene (one from each parental allele); or
 - 3.2 Patient has a sweat chloride value of at least 60 mmol/L by quantitative pilocarpine iontophoresis or by Macroduct sweat collection system; and
- 4 Fither
 - 4.1 Patient has a heterozygous or homozygous F508del mutation; or
 - 4.2 Patient has a G551D mutation or other mutation responsive in vitro to elexacaftor/tezacaftor/ivacaftor (see note a); and
- 5 The treatment must be the sole funded CFTR modulator therapy for this condition; and
- 6 Treatment with elexacaftor/tezacaftor/ivacaftor must be given concomitantly with standard therapy for this condition.

Notes:

a) Eligible mutations are listed in the Food and Drug Administration (FDA) Trikafta prescribing information https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/212273s004lbl.pdf

| IVACAFTOR - PCT only - Specialist - Special Author | rity see SA2017 below | | |
|--|-----------------------|----|----------------------------|
| Tab 150 mg | 29,386.00 | 56 | ✓ Kalydeco |
| Oral granules 50 mg, sachet | 29,386.00 | 56 | ✓ Kalydeco |
| Oral granules 75 mg, sachet | 29,386.00 | 56 | Kalydeco |

⇒SA2017 Special Authority for Subsidy

Initial application only from a respiratory specialist or paediatrician. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

| Subsidy (Manufacturer's Price) Subsidised Per ✓ Biomed Subsidity (Manufacturer's Price) Subsidised Per ✓ Biomed Subsidised Per ✓ Per Manufacturer Subsidity (Manufacturer Per Ver Ver Ver Ver Ver Ver Ver Ver Ver V |
|---|
| 1 Patient has been diagnosed with cystic fibrosis; and 2 Either: 2.1 Patient must have G551D mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene of least 1 allele; or 2.2 Patient must have other gating (class III) mutation (G1244E, G1349D, G178R, G551S, S1251N, S1255P, S548 and S549R) in the CFTR gene on at least 1 allele; and 3 Patients must have a sweat chloride value of at least 60 mmol/L by quantitative pilocarpine iontophoresis or by Macros sweat collection system; and 4 Treatment with vacaftor must be given concomitantly with standard therapy for this condition; and 5 Patient must not have an acute upper or lower respiratory infection, pulmonary exacerbation, or changes in therapy (including antibiotics) for pulmonary disease in the last 4 weeks prior to commencing treatment with ivacaftor; and 6 The dose of ivacaftor will not exceed one tablet or one sachet twice daily; and 7 Applicant has experience and expertise in the management of cystic fibrosis. SODIUM CHLORIDE Not funded for use as a nasal drop. Soln 7% |
| 2 Either: 2.1 Patient must have G551D mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene of least 1 allele; or 2.2 Patient must have other gating (class III) mutation (G1244E, G1349D, G178R, G551S, S1251N, S1255P, S548 and S549R) in the CFTR gene on at least 1 allele; and 3 Patients must have a sweat chloride value of at least 60 mmol/L by quantitative pilocarpine iontophoresis or by Macros sweat collection system; and 4 Treatment with ivacaftor must be given concomitantly with standard therapy for this condition; and 5 Patient must not have an acute upper or lower respiratory infection, pulmonary exacerbation, or changes in therapy (including antibiotics) for pulmonary disease in the last 4 weeks prior to commencing treatment with ivacaftor; and 6 The dose of ivacaftor will not exceed one tablet or one sachet twice daily; and 7 Applicant has experience and expertise in the management of cystic fibrosis. SODIUM CHLORIDE Not funded for use as a nasal drop. Soln 7% |
| 2.1 Patient must have G551D mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene of least 1 allele; or 2.2 Patient must have other gating (class III) mutation (G1244E, G1349D, G178R, G551S, S1251N, S1255P, S548 and S549R) in the CFTR gene on at least 1 allele; and 3 Patients must have a sweat chloride value of at least 60 mmol/L by quantitative pilocarpine iontophoresis or by Macros sweat collection system; and 4 Treatment with ivacaftor must be given concomitantly with standard therapy for this condition; and 5 Patient must not have an acute upper or lower respiratory infection, pulmonary exacerbation, or changes in therapy (including antibiotics) for pulmonary disease in the last 4 weeks prior to commencing treatment with ivacaftor; and 6 The dose of ivacaftor will not exceed one tablet or one sachet twice daily; and 7 Applicant has experience and expertise in the management of cystic fibrosis. SODIUM CHLORIDE Not funded for use as a nasal drop. Soln 7% |
| least 1 allele; or 2.2 Patient must have other gating (class III) mutation (G1244E, G1349D, G178R, G551S, S1251N, S1255P, S548 and S549R) in the CFTR gene on at least 1 allele; and 3 Patients must have a sweat chloride value of at least 60 mmol/L by quantitative pilocarpine iontophoresis or by Macros sweat collection system; and 4 Treatment with ivacaftor must be given concomitantly with standard therapy for this condition; and 5 Patient must not have an acute upper or lower respiratory infection, pulmonary exacerbation, or changes in therapy (including antibiotics) for pulmonary disease in the last 4 weeks prior to commencing treatment with ivacaftor; and 6 The dose of ivacaftor will not exceed one tablet or one sachet twice daily; and 7 Applicant has experience and expertise in the management of cystic fibrosis. SODIUM CHLORIDE Not funded for use as a nasal drop. Soln 7% |
| 2.2 Patient must have other gating (class III) mutation (G1244E, G1349D, G178R, G551S, S1251N, S1255P, S545 and S549R) in the CFTR gene on at least 1 allele; and 3 Patients must have a sweat chloride value of at least 60 mmol/L by quantitative pilocarpine iontophoresis or by Macrosweat collection system; and 4 Treatment with ivacaftor must be given concomitantly with standard therapy for this condition; and 5 Patient must not have an acute upper or lower respiratory infection, pulmonary exacerbation, or changes in therapy (including antibiotics) for pulmonary disease in the last 4 weeks prior to commencing treatment with ivacaftor; and 6 The dose of ivacaftor will not exceed one tablet or one sachet twice daily; and 7 Applicant has experience and expertise in the management of cystic fibrosis. SODIUM CHLORIDE Not funded for use as a nasal drop. Soln 7% |
| and S549R) in the CFTR gene on at least 1 allele; and 3 Patients must have a sweat chloride value of at least 60 mmol/L by quantitative pilocarpine iontophoresis or by Macrosweat collection system; and 4 Treatment with ivacaftor must be given concomitantly with standard therapy for this condition; and 5 Patient must not have an acute upper or lower respiratory infection, pulmonary exacerbation, or changes in therapy (including antibiotics) for pulmonary disease in the last 4 weeks prior to commencing treatment with ivacaftor; and 6 The dose of ivacaftor will not exceed one tablet or one sachet twice daily; and 7 Applicant has experience and expertise in the management of cystic fibrosis. SODIUM CHLORIDE Not funded for use as a nasal drop. Soln 7% |
| sweat collection system; and 4 Treatment with ivacaftor must be given concomitantly with standard therapy for this condition; and 5 Patient must not have an acute upper or lower respiratory infection, pulmonary exacerbation, or changes in therapy (including antibiotics) for pulmonary disease in the last 4 weeks prior to commencing treatment with ivacaftor; and 6 The dose of ivacaftor will not exceed one tablet or one sachet twice daily; and 7 Applicant has experience and expertise in the management of cystic fibrosis. SODIUM CHLORIDE Not funded for use as a nasal drop. Soln 7% |
| 4 Treatment with ivacaftor must be given concomitantly with standard therapy for this condition; and 5 Patient must not have an acute upper or lower respiratory infection, pulmonary exacerbation, or changes in therapy (including antibiotics) for pulmonary disease in the last 4 weeks prior to commencing treatment with ivacaftor; and 6 The dose of ivacaftor will not exceed one tablet or one sachet twice daily; and 7 Applicant has experience and expertise in the management of cystic fibrosis. SODIUM CHLORIDE Not funded for use as a nasal drop. Soln 7% |
| 5 Patient must not have an acute upper or lower respiratory infection, pulmonary exacerbation, or changes in therapy (including antibiotics) for pulmonary disease in the last 4 weeks prior to commencing treatment with ivacaftor; and 6 The dose of ivacaftor will not exceed one tablet or one sachet twice daily; and 7 Applicant has experience and expertise in the management of cystic fibrosis. SODIUM CHLORIDE Not funded for use as a nasal drop. Soln 7% |
| (including antibiotics) for pulmonary disease in the last 4 weeks prior to commencing treatment with ivacaftor; and 6 The dose of ivacaftor will not exceed one tablet or one sachet twice daily; and 7 Applicant has experience and expertise in the management of cystic fibrosis. SODIUM CHLORIDE Not funded for use as a nasal drop. Soln 7% |
| 7 Applicant has experience and expertise in the management of cystic fibrosis. SODIUM CHLORIDE Not funded for use as a nasal drop. Soln 7% |
| SODIUM CHLORIDE Not funded for use as a nasal drop. Soln 7% |
| Not funded for use as a nasal drop. Soln 7%25.73 90 ml OP ✓ Biomed |
| Soln 7% |
| |
| Nasal Preparations |
| |
| Allergy Prophylactics |
| BUDESONIDE |
| Metered aqueous nasal spray, 50 mcg per dose2.59 200 dose OP ✓ <u>SteroClear</u> |
| Metered aqueous nasal spray, 100 mcg per dose2.89 200 dose OP ✓ <u>SteroClear</u> |
| FLUTICASONE PROPIONATE |
| Metered aqueous nasal spray, 50 mcg per dose1.98 120 dose OP ✓ Flixonase Hayfever & Allergy |
| IPRATROPIUM BROMIDE |
| Aqueous nasal spray, 0.03% |
| |
| Respiratory Devices |
| MASK FOR SPACER DEVICE |
| a) Up to 50 dev available on a PSO |
| b) Only on a PSO |
| c) Only for children aged six years and under Small2.70 1 ✓ e-chamber Mask |

✓ Mini-Wright AFS Low Range

✓ Mini-Wright Standard

Normal range......9.54

PEAK FLOW METER

b) Only on a PSO

a) Up to 25 dev available on a PSO

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | d Generic |
|------------------------------------|---|-----|---------------------|------------------------|
| SPACER DEVICE | | | | |
| a) Up to 50 dev available on a PSO | | | | |
| b) Only on a PSO | | | | |
| 220 ml (single patient) | 3.65 | 1 | ✓ | e-chamber Turbo |
| 510 ml (single patient) | 5.95 | 1 | • | e-chamber La Grande |
| 800 ml | 6.50 | 1 | ✓ | Volumatic |

| CAFFEINE CITRATE | | | |
|---|-------|----------|--------------------------|
| Oral lig 20 mg per ml (10 mg base per ml) | 16.91 | 25 ml OP | Biomed |

| | Subsidy | | Fully Brand or |
|--|----------------------|-----------|---------------------------|
| | (Manufacturer's Pri | | |
| | \$ | Per | ✓ Manufacturer |
| Ear Preparations | | | |
| FLUMETASONE PIVALATE | | | |
| Ear drops 0.02% with clioquinol 1% | 4.46 | 7.5 ml OP | ✓ Locacorten-Viaform ED's |
| | | | ✓ Locorten-Vioform |
| | | | - Locoi ten-violonii |
| TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYCI | N AND NYSTATII | N | |
| Ear drops 1 mg with nystatin 100,000 u, neomycin sulphate | | | |
| 2.5 mg and gramicidin 250 mcg per g | 5.16 | 7.5 ml OP | ✓ Kenacomb |
| Ear/Eye Preparations | | | |
| DEXAMETHASONE WITH FRAMYCETIN AND GRAMICIDIN | | | |
| | | | |
| Ear/Eye drops 500 mcg with framycetin sulphate 5 mg and | 4.50 | 0 1 OD | |
| gramicidin 50 mcg per ml | | 8 ml OP | Otodov 520 |
| | (9.27) | | Otodex S29 Sofradex |
| | (9.27) | | Solradex |
| FRAMYCETIN SULPHATE | | | |
| Ear/Eye drops 0.5% | | 8 ml OP | |
| | (8.65) | | Soframycin |
| Eye Preparations | | | |
| Eye preparations are only funded for use in the eye, unless explicit | citly stated otherwi | ise. | |
| Anti-Infective Preparations | | | |
| ACICLOVIR | | | |
| * Eye oint 3% | 15.89 | 4.5 g OP | ✓ <u>ViruPOS</u> |
| CHLORAMPHENICOL | | | |
| Eye oint 1% | 1.09 | 5 g OP | ✓ Devatis |
| Eye drops 0.5% | 1.45 | 10 ml OP | ✓ Chlorsig |
| Funded for use in the ear*. Indications marked with * are | e unapproved indi | cations. | |
| CIPROFLOXACIN | | | |
| Eye drops 0.3% – Subsidy by endorsement | 10.85 | 5 ml OP | ✓ Ciprofloxacin Teva |
| a) When prescribed for the treatment of bacterial keratit | | | • |
| or for the second line treatment of chronic suppurativ | | | |
| accordingly. Note: Indication marked with a * is an u | | | F |
| b) Ciprofloxacin Teva to be Principal Supply on 1 March | | | |
| SODIUM FUSIDATE [FUSIDIC ACID] | | | |
| Eye drops 1% | 5 29 | 5 g OP | ✓ Fucithalmic |
| _, v alopo 1/v | | o y Oi | ✓ Fucithalmic S29 S29 |
| | | | - i-uciuiaiiiiiC 323 323 |
| TOBRAMYCIN | | | |
| Eye oint 0.3% | 10.45 | 3.5 g OP | ✓ Tobrex |
| Eye drops 0.3% | | 5 ml OP | ✓ Tobrex |
| =, 0 310p0 0.0 /0 | | 3 1111 31 | |



| Subsidy | | Fully | Brand or |
|------------------------|-----|-----------|--------------|
| (Manufacturer's Price) | S | ubsidised | Generic |
| \$ | Per | 1 | Manufacturer |

Corticosteroids and Other Anti-Inflammatory Preparations

| * | Eye oint 0.1% | 3.5 g OP | Maxidex |
|---|---|----------|---------------------------|
| * | Eye drops 0.1%4.50 | 5 ml OP | Maxidex |
| | Ocular implant 700 mcg - Special Authority see SA1680 below | | |
| | - Retail pharmacy | 1 | Ozurdex |

⇒SA1680 Special Authority for Subsidy

Initial application — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has diabetic macular oedema with pseudophakic lens: and
- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- 3 Fither
 - 3.1 Patient's disease has progressed despite 3 injections with bevacizumab; or
 - 3.2 Patient is unsuitable or contraindicated to treatment with anti-VEGF agents; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Initial application — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has diabetic macular oedema; and
- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- 3 Patient is of child bearing potential and has not yet completed a family; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Patient is of child bearing potential and has not yet completed a family; and
- 3 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

DEXAMETHASONE WITH NEOMYCIN SULPHATE AND POLYMYXIN B SULPHATE

| * Eye oint 0.1% with neomycin sulphate 0.35% and polymyxin b | | | |
|--|------|----------|----------------------|
| sulphate 6,000 u per g | 5.39 | 3.5 g OP | ✓ Maxitrol |
| * Eye drops 0.1% with neomycin sulphate 0.35% and polymyxin | | | |
| b sulphate 6,000 u per ml | 4.50 | 5 ml OP | ✓ Maxitrol |
| DICLOFENAC SODIUM | | | |
| Eye drops 0.1%, single dose | 1.85 | 10 dose | Diclofenac Devatis |
| , , , , , , | 5.54 | 30 dose | ✓ Diclofenac Devatis |
| FLUOROMETHOLONE | | | |
| * Eye drops 0.1% | 3.09 | 5 ml OP | ✓ FML |
| | 5.20 | | ✓ Flucon |

| | Subsidy (Manufacturer's Pr | ice) Subs | Fully | Brand or Generic |
|---|-------------------------------|----------------|------------|---------------------|
| | \$ | Per | ✓ | Manufacturer |
| EVOCABASTINE | | | | |
| Eye drops 0.5 mg per ml | 8.71 | 4 ml OP | | |
| | (10.34) | | L | ivostin |
| LODOXAMIDE | | | | |
| Eye drops 0.1% | 8.71 | 10 ml OP | √ L | omide |
| NEPAFENAC | | | | |
| Eye drops 0.3% | 8.80 | 3 ml OP | ✓ | evro |
| llevro Eye drops 0.3% to be delisted 1 July 2025) | | | | |
| PREDNISOLONE ACETATE | | | | |
| Eye drops 1% | 6.92 | 10 ml OP | ✓ P | rednisolone-AFT |
| , ' | 7.00 | 5 ml OP | ✓ P | red Forte |
| PREDNISOLONE SODIUM PHOSPHATE - Special Authority | see SA1715 below | - Retail pharr | nacv | |
| Eye drops 0.5%, single dose (preservative free) | | 20 dose | | linims |
| | | | | Prednisolone |

⇒SA1715 Special Authority for Subsidy

Initial application only from an ophthalmologist or optometrist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has severe inflammation; and
- 2 Patient has a confirmed allergic reaction to preservative in eye drops.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

| SODIUM CROMOGLICATE Eye drops 2% | 10 ml OP | ✓ <u>Allerfix</u> |
|--|--------------------|---|
| Glaucoma Preparations - Beta Blockers | | |
| # Eye drops 0.25% | 5 ml OP 5 ml OP | ✓ Betoptic S ✓ Betoptic |
| TIMOLOL * Eye drops 0.25% 2.42 * Eye drops 0.5% 2.50 | 5 ml OP 5 ml OP | ✓ <u>Arrow-Timolol</u> ✓ <u>Arrow-Timolol</u> |
| Glaucoma Preparations - Carbonic Anhydrase Inhibitors | | |
| ACETAZOLAMIDE * Tab 250 mg | 100 | ✓ Diamox |
| * Eye drops 1% | 5 ml OP | ✓ Azopt |
| * Eye drops 2% with timolol 0.5% | 5 ml OP | ✓ <u>Dortimopt</u> |
| | | |

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

| | Subsidy | | Fully Brand | or |
|---|-------------------|-----------|-------------------|---------------|
| | (Manufacturer's I | | idised Generi | |
| | \$ | Per | ✓ Manufa | acturer |
| LATANOPROST | | | | |
| * Eye drops 0.005% | 2.08 | 2.5 ml OP | ✓ Teva | |
| | | | | |
| TRAVOPROST | 0.00 | 0.5 | / Tuescates | |
| * Eye drops 0.004% | 6.80 | 2.5 ml OP | ✓ <u>Travatan</u> | |
| Glaucoma Preparations - Other | | | | |
| • | | | | |
| BRIMONIDINE TARTRATE | F 10 | 5 ml OP | ✓ Arrow-Bi | da a midina |
| Eye drops 0.2% | | 5 1111 OP | Allow-bi | imoniaine |
| | 25 | | | |
| BRIMONIDINE TARTRATE WITH TIMOLOL MALEATE | 7.40 | 5l OD | . | _ |
| * Eye drops 0.2% with timolol maleate 0.5% | 7.13 | 5 ml OP | ✓ Combiga | <u>ın</u> |
| LATANOPROST WITH TIMOLOL | | | _ | |
| * Eye drops 0.005% with timolol 0.5% | 4.95 | 2.5 ml OP | Arrow - L | <u>.attim</u> |
| PILOCARPINE HYDROCHLORIDE | | | | |
| * Eye drops 1% | 4.26 | 15 ml OP | ✓ Isopto Ca | arpine |
| * Eye drops 2% | 5.35 | 15 ml OP | ✓ Isopto Ca | |
| * Eye drops 4% | | 15 ml OP | ✓ Isopto Ca | arpine |
| Subsidised for oral use pursuant to the Standard Formul | ae. | | | |
| PILOCARPINE NITRATE | | | | |
| * Eye drops 2% single dose - Special Authority see SA0895 | | | | |
| below – Retail pharmacy | 35.90 | 20 dose | ✓ Minims F | Pilocarpine |
| ⇒SA0895 Special Authority for Subsidy | | | | - |
| | | | | |

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient has to use an unpreserved solution due to an allergy to the preservative; or
- 2 Patient wears soft contact lenses.

Note: Minims for a general practice are considered to be "tools of trade" and are not approved as special authority items. **Renewal** from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

| Mydriatics and Cycloplegics | | |
|---|----------------------|---|
| * Eye drops 1% | 15 ml OP | ✓ Atropt |
| CYCLOPENTOLATE HYDROCHLORIDE * Eye drops 1%25.16 TROPICAMIDE | 15 ml OP | ✓ Cyclogyl |
| * Eye drops 0.5% | 15 ml OP 15 ml OP | MydriacylMydriacyl |
| Preparations for Tear Deficiency | | |
| For acetylcysteine eye drops refer Standard Formulae, page 272 HYPROMELLOSE | | |
| * Eye drops 0.5% | 15 ml OP | ✓ Methopt |
| HYPROMELLOSE WITH DEXTRAN * Eye drops 0.3% with dextran 0.1%2.30 | 15 ml OP | ✓ Poly-Tears |

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

Preservative Free Ocular Lubricants

⇒SA2431 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Confirmed diagnosis by slit lamp or Schirmer test of severe secretory dry eye; and
- 2 Either:
 - 2.1 Patient is using eye drops more than four times daily on a regular basis; or
 - 2.2 Patient has had a confirmed allergic reaction to preservative in eye drop.

| CARBOMER - Special Authority see SA2431 above - Retail ph | armacy | | |
|--|---------------------|--------------|-----------------------------------|
| Ophthalmic gel 0.3%, 0.5 g | 8.25 | 30 | ✓ Poly-Gel |
| (Poly-Gel Ophthalmic gel 0.3%, 0.5 g to be delisted 1 July 2025) | | | · |
| POLYETHYLENE GLYCOL 400 AND PROPYLENE GLYCOL - | Special Authority s | ee SA2431 | above - Retail pharmacy |
| Eye drops 0.4% and propylene glycol 0.3%, 0.8 ml | 10.78 | 30 | Systane Unit Dose |
| SODIUM HYALURONATE [HYALURONIC ACID] - Special Auth | hority see SA2431 a | ibove – Reta | ail pharmacy |
| Eye drops 1 mg per ml | 13.58 | 10 ml OP | ✓ Hylo-Fresh |
| Hylo-Fresh has a 6 month expiry after opening. The Ph | armacy Procedures | Manual res | striction allowing one bottle per |
| month is not relevant and therefore only the prescribed | dosage to the neare | st OP may | be claimed. |

Other Eye Preparations

| NAPHAZOLINE HYDROCHLORIDE | | |
|--------------------------------|----------|--------------------|
| * Eye drops 0.1% | 15 ml OP | ✓ <u>Albalon</u> |
| OLOPATADINE | | |
| Eye drops 0.1%2.17 | 5 ml OP | ✓ Olopatadine Teva |
| PARAFFIN LIQUID WITH WOOL FAT | | |
| * Eye oint 3% with wool fat 3% | 3.5 g OP | ✓ Poly-Visc |
| RETINOL PALMITATE | | |
| Eve oint 138 mcg per g. 3.80 | 5 a OP | ✓ VitA-POS |



| | VARIOUS | | | | |
|-----|---|---------------------------------|-------------------------|---------------------|--|
| | | Subsidy (Manufacturer's Pric | ce) Per | Fully Subsidised | Brand or Generic Manufacturer |
| ٧ | arious | | | | |
| * * | ARMACY SERVICES Immunisation administration fee - flu Immunisation administration fee - other Immunisation co-administration fee - flu and shingles | 0.00 | 1 fee 1 fee 1 fee | • | Immunisation - Flu Immunisation Other Immunisation Flu and Shingles |
| A | gents Used in the Treatment of Poisonings | | | | |
| A | ntidotes | | | | |
| | ETYLCYSTEINE Inj 200 mg per ml, 10 ml ampoule DBL Acetylcysteine to be Principal Supply on 1 April 202 artindale Pharma Inj 200 mg per ml, 10 ml ampoule to be delis | 52.88 5 | 10 | | DBL Acetylcysteine Martindale Pharma |
| NA | LOXONE HYDROCHLORIDE a) Up to 10 inj available on a PSO b) Only on a PSO | | | | |
| * | Inj 400 mcg per ml, 1 ml ampoule | 13.29 | 5 | • | DBL Naloxone Hydrochloride |
| (На | DBL Naloxone Hydrochloride to be Principal Supply on 1 ameln Inj 400 mcg per ml, 1 ml ampoule to be delisted 1 April 2 | | 10 | • | Hameln |
| R | emoval and Elimination | | | | |
| | ARCOAL Oral liq 50 g per 250 ml a) Up to 250 ml available on a PSO b) Only on a PSO | 43.50 | 250 ml (| OP 🗸 | Carbosorb-X |
| DE | FERASIROX – Special Authority see SA1492 below – Retail p Wastage claimable | oharmacy | | | |

Wastage claimable

| Tab 125 mg dispersible | 276.00 | 28 | Exjade |
|------------------------|----------|----|--------------------------|
| Tab 250 mg dispersible | 552.00 | 28 | ✓ Exjade |
| Tab 500 mg dispersible | 1,105.00 | 28 | Exjade |

⇒SA1492 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; and
- 2 Deferasirox is to be given at a daily dose not exceeding 40 mg/kg/day; and
- 3 Any of the following:
 - 3.1 Treatment with maximum tolerated doses of deferiprone monotherapy or deferiprone and desferrioxamine combination therapy have proven ineffective as measured by serum ferritin levels, liver or cardiac MRI T2*; or
 - 3.2 Treatment with deferiprone has resulted in severe persistent vomiting or diarrhoea; or
 - 3.3 Treatment with deferiprone has resulted in arthritis; or
 - 3.4 Treatment with deferiprone is contraindicated due to a history of agranulocytosis (defined as an absolute neutrophil

| | Subsidy (Manufacturer's Pric \$ | ce) Per | Fully Subsidised | Brand or Generic Manufacturer |
|---|---------------------------------------|------------------------|---------------------------------|-------------------------------------|
| continued | | | | |
| count (ANC) of < 0.5 cells per μL) or recurrent episco. 5 - 1.0 cells per μL). | odes (greater than | n 2 episo | odes) of mod | derate neutropenia (ANC |
| Renewal only from a haematologist. Approvals valid for 2 years for Either: | or applications me | eeting th | e following | criteria: |
| For the first renewal following 2 years of therapy, the treatn improvement in all three parameters namely serum ferritin, For subsequent renewals, the treatment has been tolerated in all three parameters namely serum ferritin, cardiac MRI | cardiac MRI T2* d and has resulted | and live d in clini | r MRI T2* le cal stability o | vels; or |
| DEFERIPRONE – Special Authority see SA1480 below – Retail p Tab 500 mg Oral liq 100 mg per 1 ml | 533.17 | 100 250 ml | | erriprox erriprox |

⇒SA1480 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; or
- 2 The patient has been diagnosed with chronic iron overload due to acquired red cell aplasia.

| * Inj 500 mg vial | 151.31 | 10 | ✓ DBL Desferrioxamine Mesylate for Inj |
|---------------------------|----------|----|--|
| | | | BP ✓ Deferoxamine Pfizer S29 S29 |
| SODIUM CALCIUM EDETATE | | | |
| * Inj 200 mg per ml, 5 ml | 53.31 | 6 | |
| | (156.71) | | Calcium Disodium Versenate |



Standard Formulae

| ACETYLCYSTEINE EYE DROPS Acetylcysteine inj 200 mg per ml, 10 ml Suitable eye drop base Qs Phenobarbitone Sodium Qs Qlycerol BP Water Qs Qlycerol BP Water Qs Qs Qs Preservative should be used if quantity supplied is for more than 5 days. Maximum 500 ml per prescription.) METHYL HYDROXYBENZOATE 10% SOLUTION Methyl hydroxybenzoate Qs Propylene glycol Qs Propylene glycol Qs | Otaliaala i Olillalac | | | |
|--|---|-------------|--|--------------|
| Suitable eye drop base qs Phenobarbitone Sodium 400 mg Glycerol BP Water to 40 ml Vater to 100 ml Vater qs Pilocarpine 4% eye drops qs Vater to 100 ml Vater to 100 ml Vater to 500 ml Vater Vater to 100 ml Vater Vater to 100 ml Vater Vater Vater to 100 ml Vater Vat | | ae | | LIQUID (10 |
| CODEINE LINCTUS (3 mg per 5 ml) Codeine phosphate Glycerol Water CODEINE LINCTUS (15 mg per 5 ml) Codeine phosphate Glycerol Preservative Water CODEINE LINCTUS (15 mg per 5 ml) Codeine phosphate Glycerol Preservative Water CODEINE LINCTUS (15 mg per 5 ml) Codeine phosphate Glycerol Preservative Water Vater Codeine phosphate Glycerol Preservative Water Codeine phosphate Glycerol Preservative Water Codeine phosphate Glycerol A 0 ml Preservative Water Codeine phosphate Glycerol A 0 ml Preservative should be used if quantity supplied is for more than 5 days. Methyl cellulose Preservative should be used if quantity supplied is for more than 5 days. Maximum 500 ml per prescription.) Methyl hydroxybenzoate Propylene glycol (Use 1 ml of the 10% solution per 100 ml of oral liquid mixture) Comeprazole capsules or powder Sodium bicarbonate powder BP Water Conly funded if prescribed for treatment of Clostridium difficile following metronidazole failure) PHENOBARBITONE ORAL LIQUID Phenobarbitone Sodium 1 g | , | • | | 400 mg |
| Codeine phosphate Glycerol 40 ml Preservative qs Water to 100 ml CODEINE LINCTUS (15 mg per 5 ml) Codeine phosphate 300 mg Glycerol 40 ml Preservative qs Water to 100 ml Preservative water to 100 ml Preservative 40 ml Preservative 5 days. Water to 100 ml Preservative 40 ml Preservative 40 ml Preservative 40 ml Preservative 5 mg tab Calcium folinate 15 mg tab Preservative 40 ml Preservative 5000 ml Preservative 5 mg tab Preservative 40 ml Preservative 5000 ml Preservative 5 mg tab Preservative 5 mg tab Preservative 5 mg tab Preservative should be used if quantity supplied is for more than 5 days. Maximum 500 ml per prescription.) Methyl hydroxybenzoate 10 g Propylene glycol 5 mg to 100 ml (Use 1 ml of the 10% solution per 100 ml of oral liquid mixture) OMEPRAZOLE SUSPENSION 4 mg to 100 ml Omeprazole capsules or powder 5 mg water 5 mg to 100 ml Omeprazole capsules or powder 5 mg water 5 mg per ml PHENOBARBITONE ORAL LIQUID Phenobarbitone Sodium 1 g | , , | · | Glycerol BP | 4 ml |
| Glycerol Preservative qs Vater to 100 ml Vater to 100 ml CODEINE LINCTUS (15 mg per 5 ml) Codeine phosphate Glycerol 40 ml Preservative qs Glycerol 40 ml Preservative qs Vater to 100 ml Preservative 40 ml Preservative qs Vater (Preservative should be used if quantity supplied is for more than 5 days.) FOLINIC MOUTHWASH Calcium folinate 15 mg tab Preservative qs Water (Preservative should be used if quantity supplied is for more than 5 days. Maximum 500 ml per prescription.) METHYL HYDROXYBENZOATE 10% SOLUTION Methyl hydroxybenzoate 10 grouplene glycol (Use 1 ml of the 10% solution per 100 ml of oral liquid mixture) OMEPRAZOLE SUSPENSION Qmprazole capsules or powder Sodium bicarbonate powder BP Water to 100 ml Phenobarbitone Sodium 1 g g Vater (Conly funded if prescribed for treatment of Clostridium difficile following metronidazole failure) PHENOBARBITONE ORAL LIQUID PILOCARPINE ORAL LIQUID Pilocarpine 4% eye drops Preservative qs Water (Preservative should be used if quantity supplied is for more than 5 days.) Water (Preservative should be used if quantity supplied is for more than 5 days.) METHYL HYDROXYBENZOM 1 tab 1 tab Vater qs (Only funded if prescribed for treatment of hyponatraemia) Vancomycin 500 mg injection 5 vials Glycerin with sucrose suspension 37.5 ml Water (Only funded if prescribed for treatment of Clostridium difficile following metronidazole failure) PHENOBARBITONE ORAL LIQUID Phenobarbitone Sodium 1 g g | CODEINE LINCTUS (3 mg per 5 ml) | | Water | to 40 ml |
| Preservative Water to 100 ml Vocabine LINCTUS (15 mg per 5 ml) Codeine phosphate Glycerol Preservative 40 ml Preservative Should be used if quantity supplied is for more than 5 days.) FOLINIC MOUTHWASH Sater Says. Maximum 500 ml per prescription.) Methyl hydroxybenzoate Says. Maximum 500 ml per prescription.) Methyl hydroxybenzoate Propylene glycol (Use 1 ml of the 10% solution per 100 ml of oral liquid mixture) Omeprazole capsules or powder Sodium bicarbonate powder BP Water (bottom of the 100 ml Preservative solution per 100 ml of oral liquid mixture) Phenobarbitone Sodium 1 g | | | | |
| Water to 100 ml CODEINE LINCTUS (15 mg per 5 ml) Codeine phosphate 300 mg Glycerol 40 ml Preservative qs Water to 100 ml Preservative should be used if quantity supplied is for more than 5 days.) FOLINIC MOUTHWASH Calcium folinate 15 mg tab Preservative qs Water to 500 ml CPreservative should be used if quantity supplied is for more than 5 days. Maximum 500 ml per prescription.) Water qs Water to 500 ml CPreservative should be used if quantity supplied is for more than 5 days. Maximum 500 ml per prescription.) Water to 500 ml CPreservative should be used if quantity supplied is for more than 5 days. Maximum 500 ml per prescription.) Water to 500 ml CPreservative should be used if quantity supplied is for more than 5 days. Maximum 500 ml per prescription.) Water (Only funded if prescribed for treatment of hyponatraemia) WATERLY HYDROXYBENZOATE 10% SOLUTION Methyl hydroxybenzoate 10 g Propylene glycol to 100 ml (Use 1 ml of the 10% solution per 100 ml of oral liquid mixture) OMEPRAZOLE SUSPENSION VAICE TO METHORIAN SOLUTION (25 mg per ml) VANCOMYCIN ORAL SOLUTION (25 mg per ml) | | 40 ml | | |
| CODEINE LINCTUS (15 mg per 5 ml) Codeine phosphate Glycerol A0 ml Preservative Qs Water Volume Verservative should be used if quantity supplied is for more than 5 days. Maximum 500 ml per prescription.) Water Volume Water Volume Volume Vancomycin 500 mg injection Volume Volu | | | | qs |
| CODEINE LINCTUS (15 mg per 5 ml) Codeine phosphate Glycerol 40 ml Preservative Water to 100 ml Calcium folinate 15 mg tab Calcium folinate 15 mg tab Preservative Water to 500 ml Preservative should be used if quantity supplied is for more than 5 days. POLINIC MOUTHWASH Calcium folinate 15 mg tab Calcium folinate 15 mg tab Preservative Water to 500 ml Preservative should be used if quantity supplied is for more than 5 days. Maximum 500 ml per prescription.) Water | Water | to 100 ml | Preservative | qs |
| Codeine phosphate Glycerol 40 ml Preservative qs Water to 100 ml Calcium folinate 15 mg tab Preservative should be used if quantity supplied is for more than 5 days. Maximum 500 ml per prescription.) METHYL HYDROXYBENZOATE 10% SOLUTION Methyl hydroxybenzoate 10 g Propylene glycol to 100 ml (Use 1 ml of the 10% solution per 100 ml of oral liquid mixture) OMEPRAZOLE SUSPENSION Omeprazole capsules or powder Sodium bicarbonate powder BP Water to 100 ml PHENOBARBITONE ORAL LIQUID Phenobarbitone Sodium 1 g than 5 days.) SALIVA SUBSTITUTE FORMULA Methylcellulose 5 g Preservative water to 500 ml Methyl clylicolog 4 g Preservative should be used if quantity supplied is for more than 5 days. Maximum 500 ml per prescription.) SODIUM CHLORIDE ORAL LIQUID Sodium chloride inj 23.4%, 20 ml qs Water qs (Only funded if prescribed for treatment of hyponatraemia) VANCOMYCIN ORAL SOLUTION (25 mg per ml) Vancomycin 500 mg injection 5 vials Glycerin with sucrose suspension 37.5 ml Water to 100 ml One prazole capsules or powder qs Sodium bicarbonate powder BP Water to 100 ml PHENOBARBITONE ORAL LIQUID Phenobarbitone Sodium 1 g | CODEINE LINOTHO (45 | | | |
| Glycerol 40 ml Preservative qs Water to 100 ml Calcium folinate 15 mg tab Preservative should be used if quantity supplied is for more than 5 days. Maximum 500 ml per prescription.) Wethyl hydroxybenzoate 10% SOLUTION Methyl hydroxybenzoate 100 ml of oral liquid mixture) Water (Indicate 15 mg tab) Preservative should be used if quantity supplied is for more than 5 days. Maximum 500 ml per prescription.) METHYL HYDROXYBENZOATE 10% SOLUTION Methyl hydroxybenzoate 100 ml of oral liquid mixture) METHYL HYDROXYBENZOATE 10% SOLUTION Methyl hydroxybenzoate 100 ml of oral liquid mixture) METHYL HYDROXYBENZOATE 10% SOLUTION Methyl hydroxybenzoate 100 ml of oral liquid mixture) METHYL HYDROXYBENZOATE 10% SOLUTION Methyl hydroxybenzoate 100 ml of oral liquid mixture) METHYL HYDROXYBENZOATE 10% SOLUTION Methyl hydroxybenzoate 10% SOLUTION (25 mg per ml) VANCOMYCIN ORAL SOLUTION (25 mg per ml) Vancomycin 500 mg injection 5 vials Glycerin with sucrose suspension 37.5 ml Water to 100 ml Vancomycin 500 mg injection 5 vials Glycerin with sucrose suspension 37.5 ml Water to 100 ml Omeprazole capsules or powder 98 Water to 100 ml Omeprazole Capsules or powder 98 Water to 100 ml Omeprazole Capsules or powder 98 Water to 100 ml Omeprazole Capsules or powder 98 Water to 100 ml Omeprazole Capsules or powder 98 Water to 100 ml Omeprazole Capsules or powder 98 Water to 100 ml Omeprazole Capsules or powder 98 Water to 100 ml Omeprazole Capsules or powder 98 Water to 100 ml Omeprazole Capsules or powder 98 Water to 100 ml Omeprazole Capsules or powder 98 Water to 100 ml Omeprazole Capsules or powder 98 Water to 100 ml Omeprazole Capsules or powder 98 Water to 100 ml Omeprazole Capsules or powder 98 Water to 100 ml | (), | 000 | (Preservative should be used if quantity supplied is | for more |
| Preservative Water to 100 ml Wethylcellulose 5 g Preservative qs Water to 500 ml Calcium folinate 15 mg tab Calcium folinate 15 mg tab Preservative qs Water to 500 ml (Preservative should be used if quantity supplied is for more than 5 days. Maximum 500 ml per prescription.) METHYL HYDROXYBENZOATE 10% SOLUTION Methyl hydroxybenzoate Propylene glycol (Use 1 ml of the 10% solution per 100 ml of oral liquid mixture) OMEPRAZOLE SUSPENSION Omeprazole capsules or powder Sodium bicarbonate powder BP Water to 100 ml Water to 100 ml Phenobarbitone Sodium 1 g SA4 g Phenobarbitone Sodium 1 g SA4 g SALIVA SUBSTITUTE FORMULA Methyl prescribet of the 100 ml Methylcellulose 5 g g Preservative qqs Water (Preservative should be used if quantity supplied is for more than 5 days. Maximum 500 ml per prescription.) SODIUM CHLORIDE ORAL LIQUID Sodium chloride inj 23.4%, 20 ml qqs Water (Only funded if prescribed for treatment of hyponatraemia) VANCOMYCIN ORAL SOLUTION (25 mg per ml) Vancomycin 500 mg injection 5 vials Glycerin with sucrose suspension 37.5 ml Water (Only funded if prescribed for treatment of Clostridium difficile following metronidazole failure) | • • | J | than 5 days.) | |
| Water to 100 ml Calcium folinate 15 mg tab Preservative qs Water to 500 ml Calcium folinate 15 mg tab Preservative qs Water to 500 ml (Preservative should be used if quantity supplied is for more than 5 days. Maximum 500 ml per prescription.) METHYL HYDROXYBENZOATE 10% SOLUTION Methyl hydroxybenzoate 10 g Propylene glycol to 100 ml (Use 1 ml of the 10% solution per 100 ml of oral liquid mixture) OMEPRAZOLE SUSPENSION Omeprazole capsules or powder Sodium bicarbonate powder BP Water to 100 ml PHENOBARBITONE ORAL LIQUID Phenobarbitone Sodium 1 g Methylcellulose Preservative y Water to 500 ml (Preservative should be used if quantity supplied is for more than 5 days. Maximum 500 ml per prescription.) SODIUM CHLORIDE ORAL LIQUID Sodium chloride inj 23.4%, 20 ml qs Water (Only funded if prescribed for treatment of hyponatraemia) VANCOMYCIN ORAL SOLUTION (25 mg per ml) Vancomycin 500 mg injection 5 vials Glycerin with sucrose suspension 37.5 ml Water to 100 ml Oneprazole capsules or powder qs Sodium bicarbonate powder BP Water to 100 ml Oneprazole capsules or powder to 100 ml Oneprazole capsules or powder BP Water to 100 ml Oneprazole capsules or powder to 100 ml | | | CALIVA CUIDCTITUTE FORMULA | |
| FOLINIC MOUTHWASH Calcium folinate 15 mg tab Preservative Qs Water (Preservative should be used if quantity supplied is for more than 5 days. Maximum 500 ml per prescription.) Water (Preservative should be used if quantity supplied is for more than 5 days. Maximum 500 ml per prescription.) METHYL HYDROXYBENZOATE 10% SOLUTION Methyl hydroxybenzoate Propylene glycol (Use 1 ml of the 10% solution per 100 ml of oral liquid mixture) OMEPRAZOLE SUSPENSION Omeprazole capsules or powder Sodium bicarbonate powder BP Water ONLY | | | | r |
| FOLINIC MOUTHWASH Calcium folinate 15 mg tab Calcium folinate 15 mg tab Preservative Water (Preservative should be used if quantity supplied is for more than 5 days. Maximum 500 ml per prescription.) Wethyl hydroxybenzoate Propylene glycol (Use 1 ml of the 10% solution per 100 ml of oral liquid mixture) OMEPRAZOLE SUSPENSION Omeprazole capsules or powder Sodium bicarbonate powder BP Water Water Vancomycin 500 mg injection Solution per 100 ml (Only funded if prescribed for treatment of hyponatraemia) VANCOMYCIN ORAL SOLUTION (25 mg per ml) Vancomycin 500 mg injection Solium chloride inj 23.4%, 20 ml Water (Only funded if prescribed for treatment of hyponatraemia) VANCOMYCIN ORAL SOLUTION (25 mg per ml) Vancomycin 500 mg injection Glycerin with sucrose suspension Water Vancomycin 500 mg injection Glycerin with sucrose suspension Water Vancomycin 500 mg injection Glycerin with sucrose suspension Water Vancomycin 500 mg injection Glycerin with sucrose suspension Water Vancomycin 500 mg injection Glycerin with sucrose suspension Water Vancomycin 500 mg injection Glycerin with sucrose suspension Water Vancomycin 500 mg injection Glycerin with sucrose suspension Water Vancomycin 500 mg injection Glycerin with sucrose suspension Water Vancomycin 500 mg injection Glycerin with sucrose suspension Water Vancomycin 500 mg injection Glycerin with sucrose suspension Water Vancomycin 500 mg injection Glycerin with sucrose suspension Water Vancomycin 500 mg injection Glycerin with sucrose suspension Water Vancomycin 500 mg injection Glycerin with sucrose suspension Water Vancomycin 500 mg injection Glycerin with sucrose suspension Water Vancomycin 500 mg injection Glycerin with sucrose suspension Water Vancomycin 500 mg injection Glycerin with sucrose suspension Water Vancomycin 500 mg injection Glycerin with sucrose suspension Water Vancomycin 500 mg injection Glycerin with sucrose suspension Water Vancomycin 500 mg injection Glycerin with sucrose suspension Vancomycin 500 mg injection Glycerin with sucrose suspension Vanco | Water | to 100 mi | | • |
| Calcium folinate 15 mg tab Preservative should be used if quantity supplied is for more than 5 days. Maximum 500 ml per prescription.) Water (Preservative should be used if quantity supplied is for more than 5 days. Maximum 500 ml per prescription.) METHYL HYDROXYBENZOATE 10% SOLUTION Methyl hydroxybenzoate Propylene glycol (Use 1 ml of the 10% solution per 100 ml of oral liquid mixture) OMEPRAZOLE SUSPENSION Omeprazole capsules or powder Sodium bicarbonate powder BP Water Vancomycin 500 mg injection Glycerin with sucrose suspension Water (Only funded if prescribed for treatment of hyponatraemia) VANCOMYCIN ORAL SOLUTION (25 mg per ml) Vancomycin 500 mg injection Glycerin with sucrose suspension 37.5 ml Water (Only funded if prescribed for treatment of Clostridium difficile following metronidazole failure) PHENOBARBITONE ORAL LIQUID Phenobarbitone Sodium 1 g | FOLINIC MOLITHWASH | | | • |
| Preservative Water to 500 ml (Preservative should be used if quantity supplied is for more than 5 days. Maximum 500 ml per prescription.) METHYL HYDROXYBENZOATE 10% SOLUTION Methyl hydroxybenzoate Propylene glycol 10% solution per 100 ml of oral liquid mixture) OMEPRAZOLE SUSPENSION Omeprazole capsules or powder Sodium bicarbonate powder BP Water to 100 ml Phenobarbitone Sodium 1 g water than 5 days. Maximum 500 ml per prescription.) SODIUM CHLORIDE ORAL LIQUID Sodium chloride inj 23.4%, 20 ml qs Water (Only funded if prescribed for treatment of hyponatraemia) VANCOMYCIN ORAL SOLUTION (25 mg per ml) Vancomycin 500 mg injection 5 vials Glycerin with sucrose suspension 37.5 ml Water to 100 ml (Only funded if prescribed for treatment of Clostridium difficile following metronidazole failure) | | 1 tah | | |
| Water to 500 ml (Preservative should be used if quantity supplied is for more than 5 days. Maximum 500 ml per prescription.) METHYL HYDROXYBENZOATE 10% SOLUTION Methyl hydroxybenzoate Propylene glycol to 100 ml (Use 1 ml of the 10% solution per 100 ml of oral liquid mixture) OMEPRAZOLE SUSPENSION Omeprazole capsules or powder Sodium bicarbonate powder BP Water to 100 ml PHENOBARBITONE ORAL LIQUID Phenobarbitone Sodium 1 g | ŭ | | | for more |
| (Preservative should be used if quantity supplied is for more than 5 days. Maximum 500 ml per prescription.) METHYL HYDROXYBENZOATE 10% SOLUTION Methyl hydroxybenzoate 10 g Propylene glycol to 100 ml (Use 1 ml of the 10% solution per 100 ml of oral liquid mixture) OMEPRAZOLE SUSPENSION Omeprazole capsules or powder Sodium bicarbonate powder BP Water to 100 ml PHENOBARBITONE ORAL LIQUID Phenobarbitone Sodium 1 g SODIUM CHLORIDE ORAL LIQUID Sodium chloride inj 23.4%, 20 ml Water (Only funded if prescribed for treatment of hyponatraemia) VANCOMYCIN ORAL SOLUTION (25 mg per ml) Vancomycin 500 mg injection 5 vials Glycerin with sucrose suspension 37.5 ml Water (Only funded if prescribed for treatment of Clostridium difficile following metronidazole failure) | | | than 5 days. Maximum 500 mi per prescription.) | |
| than 5 days. Maximum 500 ml per prescription.) METHYL HYDROXYBENZOATE 10% SOLUTION Methyl hydroxybenzoate Propylene glycol (Use 1 ml of the 10% solution per 100 ml of oral liquid mixture) OMEPRAZOLE SUSPENSION Omeprazole capsules or powder Sodium bicarbonate powder BP Water PHENOBARBITONE ORAL LIQUID Phenobarbitone Sodium Sodium chloride inj 23.4%, 20 ml Water (Only funded if prescribed for treatment of hyponatraemia) VANCOMYCIN ORAL SOLUTION (25 mg per ml) Vancomycin 500 mg injection 5 vials Glycerin with sucrose suspension Water (Only funded if prescribed for treatment of Clostridium difficile following metronidazole failure) | | | SODIUM CHLORIDE ORAL LIQUID | |
| METHYL HYDROXYBENZOATE 10% SOLUTION Methyl hydroxybenzoate Propylene glycol (Use 1 ml of the 10% solution per 100 ml of oral liquid mixture) OMEPRAZOLE SUSPENSION Omeprazole capsules or powder Sodium bicarbonate powder BP Water Water Water Water (Only funded if prescribed for treatment of hyponatraemia) VANCOMYCIN ORAL SOLUTION (25 mg per ml) Vancomycin 500 mg injection Glycerin with sucrose suspension 37.5 ml Water (Only funded if prescribed for treatment of Clostridium difficile following metronidazole failure) PHENOBARBITONE ORAL LIQUID Phenobarbitone Sodium 1 g | | ioi ilioie | Sodium chloride ini 23.4%. 20 ml | as |
| METHYL HYDROXYBENZOATE 10% SOLUTION Methyl hydroxybenzoate Propylene glycol (Use 1 ml of the 10% solution per 100 ml of oral liquid mixture) OMEPRAZOLE SUSPENSION Omeprazole capsules or powder Sodium bicarbonate powder BP Water PHENOBARBITONE ORAL LIQUID Phenobarbitone Sodium Only funded if prescribed for treatment of hyponatraemia) VANCOMYCIN ORAL SOLUTION (25 mg per ml) Vancomycin 500 mg injection Glycerin with sucrose suspension 37.5 ml Water (Only funded if prescribed for treatment of Clostridium difficile following metronidazole failure) | than 5 days. Maximum 500 mi per prescription.) | | | • |
| Methyl hydroxybenzoate Propylene glycol (Use 1 ml of the 10% solution per 100 ml of oral liquid mixture) OMEPRAZOLE SUSPENSION Omeprazole capsules or powder Sodium bicarbonate powder BP Water OMEPRAZOLE SUSPENSION Omeprazole capsules or powder Sodium bicarbonate powder BP Water ONLY ORAL SOLUTION (25 mg per ml) Vancomycin 500 mg injection Glycerin with sucrose suspension 37.5 ml Water (Only funded if prescribed for treatment of Clostridium difficile following metronidazole failure) PHENOBARBITONE ORAL LIQUID Phenobarbitone Sodium 1 g | METHYL HYDROXYBENZOATE 10% SOLUTION | | (Only funded if prescribed for treatment of hyponatr | • |
| (Use 1 ml of the 10% solution per 100 ml of oral liquid mixture) OMEPRAZOLE SUSPENSION Omeprazole capsules or powder Sodium bicarbonate powder BP Water Vancomycin 500 mg injection Glycerin with sucrose suspension Water (Only funded if prescribed for treatment of Clostridium difficile following metronidazole failure) PHENOBARBITONE ORAL LIQUID Phenobarbitone Sodium 1 g | Methyl hydroxybenzoate | 10 g | | , |
| Glycerin with sucrose suspension 37.5 ml OMEPRAZOLE SUSPENSION Omeprazole capsules or powder Sodium bicarbonate powder BP Water Vo 100 ml PHENOBARBITONE ORAL LIQUID Phenobarbitone Sodium 1 g | Propylene glycol | to 100 ml | | |
| OMEPRAZOLE SUSPENSION Omeprazole capsules or powder Sodium bicarbonate powder BP Water Water Water Gollowing metronidazole failure) Water HENOBARBITONE ORAL LIQUID Phenobarbitone Sodium 1 g | (Use 1 ml of the 10% solution per 100 ml of oral liqu | id mixture) | | |
| Omeprazole capsules or powder Sodium bicarbonate powder BP Water PHENOBARBITONE ORAL LIQUID Phenobarbitone Sodium Only funded if prescribed for treatment of Clostridium difficile following metronidazole failure) to 100 ml Only funded if prescribed for treatment of Clostridium difficile following metronidazole failure) 1 g | OMEDBAZOLE CHORENGION | | , | |
| Sodium bicarbonate powder BP 8.4 g following metronidazole failure) Water to 100 ml PHENOBARBITONE ORAL LIQUID Phenobarbitone Sodium 1 g | | | | |
| Water to 100 ml PHENOBARBITONE ORAL LIQUID Phenobarbitone Sodium 1 g | · | | ` , | um difficile |
| PHENOBARBITONE ORAL LIQUID Phenobarbitone Sodium 1 g | • | | following metronidazole failure) | |
| Phenobarbitone Sodium 1 g | water | to 100 mi | | |
| Phenobarbitone Sodium 1 g | PHENORARRITONE ORAL LIQUID | | | |
| 3 | | 1 a | | |
| | Glycerol BP | 70 ml | | |

to 100 ml

Water

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic

\$ Per Manufacturer

| Extemporaneously | Compounded Pre | parations and Galenicals |
|------------------|----------------|--------------------------|
|------------------|----------------|--------------------------|

| CODEINE PHOSPHATE – Safety medicine; prescriber may determine | ne dispensina | frequency | |
|---|-----------------|---------------|------------------------------|
| Powder – Only in combination | | 25 g | |
| | (90.09) | | Douglas |
| Only in extemporaneously compounded codeine linctus. | | | |
| COLLODION FLEXIBLE | | | |
| Note: This product is no longer being manufactured by the suppl | ier and will be | delisted from | the Schedule at a date to be |
| determined. Collodion flexible | 10.00 | 100 ml | ✓ PSM |
| | 19.30 | 100 ml | ▼ PSIVI |
| COMPOUND HYDROXYBENZOATE – Only in combination | | | |
| Only in extemporaneously compounded oral mixtures. Soln | 30.00 | 100 ml | ✓ Midwest |
| | 50.00 | 100 1111 | • mawest |
| GLYCERIN WITH SODIUM SACCHARIN – Only in combination Only in combination with Ora-Plus or when used in the vancomyo | rin oral louuid | Standard Form | mulaa |
| Suspension | | 473 ml | ✓ Ora-Sweet SF |
| GLYCERIN WITH SUCROSE – Only in combination | 00.00 | ., 0 | 014 011001 01 |
| Only in combination with Ora-Plus or when used in the vancomy | rin oral louuid | Standard Form | mulae |
| Suspension | | 473 ml | ✓ Ora-Sweet |
| GLYCEROL | | | |
| * Liquid – Only in combination | 3.23 | 500 ml | ✓ healthE Glycerol BP |
| Only in extemporaneously compounded oral liquid preparation | | | |
| METHYL HYDROXYBENZOATE | | | |
| Powder | 8.98 | 25 g | ✓ Midwest |
| METHYLCELLULOSE | | J | |
| Powder | 36.95 | 100 g | ✓ MidWest |
| Suspension - Only in combination | 30.95 | 473 ml | ✓ Ora-Plus |
| METHYLCELLULOSE WITH GLYCERIN AND SODIUM SACCHARIN | N – Only in co | ombination | |
| Suspension | | 473 ml | ✓ Ora-Blend SF |
| METHYLCELLULOSE WITH GLYCERIN AND SUCROSE - Only in | combination | | |
| Suspension | | 473 ml | ✓ Ora-Blend |
| PHENOBARBITONE SODIUM | | | |
| Powder – Only in combination | 52.50 | 10 g | ✓ MidWest |
| | 325.00 | 100 g | ✓ MidWest |
| Only in children up to 12 years | | | |
| PROPYLENE GLYCOL | | | |
| Only in extemporaneously compounded methyl hydroxybenzoate | | | • |
| Liq | 11.25 | 500 ml | ✓ Midwest |
| SODIUM BICARBONATE | | | |
| Powder BP – Only in combination | | 500 g | ✓ Midwest |
| Only in extemporaneously compounded omeprazole and lan | soprazole sus | pension. | |
| SYRUP (PHARMACEUTICAL GRADE) – Only in combination | | | |
| Only in extemporaneously compounded oral liquid preparations. | 14.05 | E00! | ✓ Midwest |
| Liq | 14.95 | 500 ml | ▼ IVIIQWeSt |
| WATER | 0.00 | 1 ml | √ Ton water |
| Tap - Only in combination | 0.00 | 1 ml | ✓ Tap water |

Subsidy (Manufacturer's Price) \$ Fully Subsidised Per

Brand or Generic Manufacturer

Nutrient Modules

Carbohydrate

⇒SA1930 Special Authority for Subsidy

Initial application — (Cystic fibrosis or kidney disease) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Fither:

- 1 cystic fibrosis; or
- 2 chronic kidney disease.

Initial application — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 cancer in children: or
- 2 cancers affecting alimentary tract where there are malabsorption problems in patients over the age of 20 years; or
- 3 faltering growth in an infant/child; or
- 4 bronchopulmonary dysplasia; or
- 5 premature and post premature infant; or
- 6 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified where the patient has inborn errors of metabolism. Renewal — (Cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

CARBOHYDRATE SUPPLEMENT – Special Authority see SA1930 above – Hospital pharmacy [HP3]

Carbohydrate And Fat

⇒SA1376 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

| Subsidy | Fully | Brand or | |
|------------------------|------------|--------------|--|
| (Manufacturer's Price) | Subsidised | Generic | |
| \$ | Per 🗸 | Manufacturer | |

continued...

- 1 Infant or child aged four years or under; and
- 2 cystic fibrosis.

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 infant or child aged four years or under; and
- 2 Any of the following:
 - 2.1 cancer in children; or
 - 2.2 faltering growth; or
 - 2.3 bronchopulmonary dysplasia; or
 - 2.4 premature and post premature infants.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Fat

⇒SA2204 Special Authority for Subsidy

Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified where the patient has an inborn error of metabolism.

Initial application — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 faltering growth in an infant/child; or
- 2 bronchopulmonary dysplasia; or
- 3 fat malabsorption; or
- 4 lymphangiectasia; or
- 5 short bowel syndrome: or
- 6 infants with necrotising enterocolitis; or
- 7 biliary atresia; or
- 8 for use in a ketogenic diet; or
- 9 chyle leak; or

continued...

| Subsidy | | Fully | Brand or |
|------------------------|-----|------------|--------------|
| (Manufacturer's Price) | S | Subsidised | Generic |
| \$ | Per | ✓ | Manufacturer |

continued...

- 10 ascites: or
- 11 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. Renewal — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

| FAT SUPPLEMENT – Special Authority see SA2204 on the prev | vious page – I | Hospital pharmacy | [HP3] |
|---|----------------|-------------------|------------|
| Emulsion (noutral) | 15 20 | 200 ml OD | √ ∩ |

| Emulsion (neutral)1 | 5.38 20 | 00 ml OP | Calogen |
|------------------------|---------|------------|--------------------|
| 3 | 8.44 50 | 00 ml OP | Calogen |
| Emulsion (strawberry)1 | 5.38 20 | 00 ml OP | Calogen |
| Oil3 | 7.50 50 | 00 ml OP 🗸 | MCT oil (Nutricia) |
| MCT Emulsion, 250 ml14 | 3.65 | 4 OP ✓ | Liquigen |

Protein

⇒SA1524 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 protein losing enteropathy; or
- 2 high protein needs; or
- 3 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

| PROTEIN SUPPLEMENT – Special Authority see SA1524 above – Hospital pharmacy [HP3 |
|--|
|--|

| Powder | 8.95 | 227 g OP | ✓ Resource Beneprotein |
|--------|-------|----------|---------------------------|
| | 13.82 | 225 a OP | ✓ Protifar |

Subsidy (Manufacturer's Price)

Subsidised Per

Fully

Brand or Generic Manufacturer

Oral and Enteral Feeds

Diabetic Products

⇒SA1095 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is a type I or and II diabetic who is suffering weight loss and malnutrition that requires nutritional support. Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

- Both:
 - 1 The treatment remains appropriate and the patient is benefiting from treatment; and
 - 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

| DIABETIC ENTERAL FEED 1KCAL/ML - Special Authority see | SA1095 above - | Hospital pharm | acy [HP3] |
|--|------------------|----------------|-------------------|
| Liquid | 4.65 | 500 ml OP | Glucerna Select |
| DIABETIC ORAL FEED 1KCAL/ML - Special Authority see SA | 1095 above – Hos | pital pharmacy | [HP3] |
| Liquid (strawberry) | 2.25 | 200 ml OP | ✓ Diasip |
| Liquid (vanilla) | 2.10 | 200 ml OP | ✓ Nutren Diabetes |
| , , , | 2.25 | | Diasip |

Fat Modified Products

⇒SA2205 Special Authority for Subsidy

Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified where the patient has an inborn error of metabolism. Initial application — (Indications other than errors of inborn metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Patient has a chyle leak; or
- 2 Modified as a modular feed, made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule, for adults,

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

| FAT MODIFIED FEED - Special Authority see SA2205 above - Hospital phar | macy [HP3] | |
|--|------------|---------|
| Powder | 400 g OP | Monogen |

Subsidy (Manufacturer's Price) Fully Subsidised Per

Brand or Generic Manufacturer

Paediatric Products For Children Awaiting Liver Transplant

⇒SA1098 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) who requires a liver transplant.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL FEED 1KCAL/ML - Special Authority see SA1098 above - Hospital pharmacy [HP3]

Paediatric Products For Children With Chronic Renal Failure

⇒SA1099 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) with acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Paediatric Products

⇒SA1379 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Child is aged one to ten years; and
- 2 Any of the following:
 - 2.1 the child is being fed via a tube or a tube is to be inserted for the purposes of feeding; or
 - 2.2 any condition causing malabsorption; or
 - 2.3 faltering growth in an infant/child; or
 - 2.4 increased nutritional requirements; or
 - 2.5 the child is being transitioned from TPN or tube feeding to oral feeding.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for

| Subsidy | | Fully | Brand or | |
|--------------------|-------|------------|--------------|--|
| (Manufacturer's Pr | rice) | Subsidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

continued...

applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

| PAEDIATRIC ENTERAL FEED 1.5KCAL/ML - Special Authority see SA1379 on the previous placed in | |
|--|---|
| PAEDIATRIC ENTERAL FEED 1KCAL/ML - Special Authority see \$A1379 on the previous pa Liquid3.32 500 ml OP 4.69 | ge – Hospital pharmacy [HP3] ✓ Pediasure RTH ✓ Nutrini RTH |
| PAEDIATRIC ENTERAL FEED WITH FIBRE 1.5KCAL/ML - Special Authority see SA1379 on pharmacy [HP3] | the previous page – Hospital |
| Liquid | Nutrini Energy Multi Fibre |
| PAEDIATRIC ORAL FEED 1.5KCAL/ML - Special Authority see SA1379 on the previous page | - Hospital pharmacy [HP3] |
| Liquid (strawberry) | ✓ Fortini |
| Liquid (vanilla) | ✓ Fortini |
| 8.67 500 ml OP | ✓ Pediasure Plus |
| PAEDIATRIC ORAL FEED 1KCAL/ML - Special Authority see SA1379 on the previous page - | - Hospital pharmacy [HP3] |
| Liquid (chocolate) | ✓ Pediasure |
| Liquid (strawberry) | ✓ Pediasure |
| Liquid (vanilla) | ✓ Pediasure |
| 1.66 250 ml OP | ✓ Pediasure |
| PAEDIATRIC ORAL FEED WITH FIBRE 1.5KCAL/ML - Special Authority see SA1379 on the pharmacy [HP3] | previous page – Hospital |
| Liquid (unflavoured) | ✓ Fortini Multi Fibre |
| Liquid (chocolate) | ✓ Fortini Multi Fibre |
| Liquid (strawberry)1.90 200 ml OP | ✓ Fortini Multi Fibre |
| Liquid (vanilla) | ✓ Fortini Multi Fibre |
| PEPTIDE-BASED ORAL FEED - Special Authority see SA1379 on the previous page - Hospit | tal pharmacy [HP3] |
| Powder | ✓ Peptamen Junior |

Renal Products

⇒SA1101 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

| RENAL ORAL FEED 1.8 KCAL/ML – Special Authority see SA1101 above – He | ospital pharmacy | [HP3] |
|---|------------------|----------------------|
| Liquid3.31 | 220 ml OP | ✓ Nepro HP |
| | | (strawberry) |
| | | ✓ Nepro HP (vanilla) |

| | (Manufacturer's Price) \$ | Subsid Per | lised • | Generic Manufacturer |
|---|------------------------------|---------------|------------|-------------------------|
| RENAL ORAL FEED 2 KCAL/ML - Special Authority see SA110 | 1 on the previous page | ge – Hospita | | |
| Liquid, 200 ml bottle | 13.24 | 4 OP | ✓ No | ovaSource Renal |
| Liquid (apricot) 125 ml | 13.72 | 4 OP | ✓ Re | enilon 7.5 |
| Liquid (caramel) 125 ml | 13.72 | 4 OP | ✓ Re | enilon 7.5 |

Subsidy

Fully

Brand or

Specialised And Elemental Products

⇒SA1377 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 malabsorption; or
- 2 short bowel syndrome; or
- 3 enterocutaneous fistulas; or
- 4 eosinophilic oesophagitis; or
- 5 inflammatory bowel disease; or
- 6 patients with multiple food allergies requiring enteral feeding.

Notes: Each of these products is highly specialised and would be prescribed only by an expert for a specific disorder. The alternative is hospitalisation.

Elemental 028 Extra is more expensive than other products listed in this section and should only be used where the alternatives have been tried first and/or are unsuitable.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

| LiquidLiquid | , | 1,000 ml OP | |
|--|---------------------|-----------------|-----------------------|
| ORAL ELEMENTAL FEED 0.8KCAL/ML - Special Authority s | see SA1377 above – | Hospital pharr | nacy [HP3] |
| Liquid (grapefruit), 250 ml carton | 179.46 | 18 OP | ✓ Elemental 028 Extra |
| Liquid (pineapple & orange), 250 ml carton | 179.46 | 18 OP | ✓ Elemental 028 Extra |
| Liquid (summer fruits), 250 ml carton | 179.46 | 18 OP | ✓ Elemental 028 Extra |
| ORAL ELEMENTAL FEED 1KCAL/ML - Special Authority se | e SA1377 above – H | lospital pharma | acy [HP3] |
| Powder (unflavoured) | 4.50 | 80 g OP | ✓ Vivonex TEN |
| SEMI-ELEMENTAL ENTERAL FEED 1KCAL/ML - Special A | uthority see SA1377 | above – Hosp | ital pharmacy [HP3] |
| Liquid | 7.47 | 500 ml OP | ✓ Nutrison Advanced |
| | | | Pentisorb |

Paediatric Products For Children With Low Energy Requirements

⇒SA1196 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

continued...

- 1 Child aged one to eight years; and
 - 2 The child has a low energy requirement but normal protein and micronutrient requirements.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

PAEDIATRIC ENTERAL FEED WITH FIBRE 0.76 KCAL/ML - Special Authority see SA1196 on the previous page - Hospital pharmacy [HP3]

Standard Supplements

⇒SA1859 Special Authority for Subsidy

Initial application — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age; and
- 2 Any of the following:
 - 2.1 The patient has a condition causing malabsorption; or
 - 2.2 The patient has failure to thrive; or
 - 2.3 The patient has increased nutritional requirements; and
- 3 Nutrition goal has been set (eg reach a specific weight or BMI).

Renewal — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 A nutrition goal has been set (eg reach a specific weight or BMI).

Initial application — (Children - exclusive enteral nutrition for Crohn's disease) only from a gastroenterologist or dietitian on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 Dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Renewal — (Children - exclusive enteral nutrition for Crohn's disease) from any relevant practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 General Practitioners and dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

continued...

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

Initial application — (Adults) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

1 Any of the following:

Patient is Malnourished

- 1.1 Patient has a body mass index (BMI) of less than 18.5 kg/m²; or
- 1.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 1.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months; and
- 2 Any of the following:

Patient has not responded to first-line dietary measures over a 4 week period by:

- 2.1 Increasing their food intake frequency (eg snacks between meals); or
- 2.2 Using high-energy foods (e.g. milkshakes, full fat milk, butter, cream, cheese, sugar etc); or
- 2.3 Using over the counter supplements (e.g. Complan); and
- 3 A nutrition goal has been set (e.g. to reach a specific weight or BMI).

Renewal — (Adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 A nutrition goal has been set (eg reach a specific weight or BMI); and
- 2 Any of the following:

Patient is Malnourished

- 2.1 Patient has a body mass index (BMI) of less than 18.5 kg/m²; or
- 2.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 2.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months.

Initial application — (Short-term medical condition) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube or a nasogastric tube is to be inserted for feeding; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Is undergoing a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both
 - 5.1 Pregnant: and
 - 5.2 Any of the following:
 - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or
 - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
 - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

Renewal — (Short-term medical condition) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Has undergone a bone marrow transplant; or

| Subsidy | | Fully | Brand or | |
|------------------------|-----|------------|--------------|--|
| (Manufacturer's Price) | S | Subsidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

continued...

- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
 - 5.1 Pregnant: and
 - 5.2 Any of the following:
 - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or
 - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
 - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

Initial application — (Long-term medical condition) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis; or
- 3 Liver disease: or
- 4 Chronic Renal failure: or
- 5 Inflammatory bowel disease; or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome: or
- 8 Bowel fistula: or
- 9 Severe chronic neurological conditions: or
- 10 Epidermolysis bullosa; or
- 11 AIDS (CD4 count < 200 cells/mm3): or
- 12 Chronic pancreatitis.

Renewal — (Chronic disease OR tube feeding for patients who have previously been funded under Special Authority forms SA0702 or SA0583) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis; or
- 3 Liver disease; or
- 4 Chronic Renal failure; or
- 5 Inflammatory bowel disease; or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome; or
- 8 Bowel fistula: or
- 9 Severe chronic neurological conditions.

| ENTERAL FEED 1.5KCAL/ML - Special Authority see SA1859 on | page 281 - H | lospital pharmac | y [HP3] |
|---|--------------|------------------|----------------------------------|
| Liquid | 2.17 | 250 ml OP | ✓ Ensure Plus HN |
| | 8.68 | 1,000 ml OP | Ensure Plus HN |
| | | | RTH |
| | 9.00 | | ✓ Nutrison Energy |

| | Subsidy | | Fully Brand or |
|--|-------------------|-------------------|------------------------------|
| | (Manufacturer's I | Price) Subsid | dised Generic |
| | \$ | Per | ✓ Manufacturer |
| ENTERAL FEED 1KCAL/ML - Special Authority see SA1859 on | nage 281 – Ho | enital pharmacy l | THD31 |
| · | | | |
| Liquid | | 250 ml OP | ✓ Isosource Standard |
| | 6.56 | 1,000 ml OP | ✓ Osmolite RTH |
| | 6.90 | | ✓ Nutrison RTH |
| ENTERAL FEED WITH FIBRE 0.83 KCAL/ML - Special Authorit | v see SA1859 d | on page 281 – Ho | ospital pharmacy [HP3] |
| Liquid | • | 1,000 ml OP | ✓ Nutrison |
| Liquiu | | 1,000 1111 01 | |
| | | | 800 Complete |
| | | | Multi Fibre |
| ENTERAL FEED WITH FIBRE 1 KCAL/ML - Special Authority s | ee SA1859 on r | page 281 – Hosp | ital pharmacy [HP3] |
| Liquid | | 1,000 ml OP | ✓ Jevity RTH |
| Liquid | 7.21 | 1,000 1111 01 | ✓ Nutrison Multi Fibre |
| | | | |
| ENTERAL FEED WITH FIBRE 1.2KCAL/ML - Special Authority | see SA1859 on | page 281 - Hos | pital pharmacy [HP3] |
| Liquid | 7.87 | 1,000 ml OP | ✓ Jevity Plus RTH |
| ENTERAL FEED WITH FIBRE 1.5KCAL/ML - Special Authority | 000 CA10E0 on | nogo 201 Hoo | nital pharmany [UD2] |
| | | | |
| Liquid | 8.68 | 1,000 ml OP | Jevity HiCal RTH |
| | | | ✓ Nutrison Energy |
| | | | Multi Fibre |
| ORAL FEED (POWDER) - Special Authority see SA1859 on page | no 281 – Hoenit | al nharmacy [HD | 31 |
| , , , | • | , ,, | • |
| Powder (chocolate) | 14.00 | 840 g OP | ✓ Sustagen Hospital |
| | | | Formula |
| | 26.00 | 850 g OP | ✓ Ensure |
| Powder (vanilla) | 14.00 | 840 g OP | ✓ Sustagen Hospital |
| , | | J | Formula Active |
| | 26.00 | 850 g OP | ✓ Ensure |
| | | - | |
| ORAL FEED 1.5KCAL/ML - Special Authority see SA1859 on pa | | | |
| Additional subsidy by endorsement is available for patients b | eing bolus fed t | hrough a feeding | tube, who have severe |
| epidermolysis bullosa, or as exclusive enteral nutrition in chil | dren under the | age of 18 years f | or the treatment of Crohn's |
| disease, or for patients with COPD and hypercapnia, defined | | | |
| endorsed accordingly. | ao ooz valao (| oxocoding comm | rig. The precemption must be |
| · · · · · · · · · · · · · · · · · · · | | | |
| Liquid (banana) – Higher subsidy of up to \$1.76 per 200 ml | | | |
| with Endorsement | 0.72 | 200 ml OP | |
| | (1.56) | | Ensure Plus |
| | (1.76) | | Fortisip |
| Liquid (chocolate) - Higher subsidy of up to \$1.76 per 200 m | nl | | • |
| with Endorsement | | 200 ml OP | |
| WILL ENGOSOMOTIC | | 200 1111 01 | Ensure Plus |
| | (1.56) | | |
| | (1.76) | | Fortisip |
| Liquid (fruit of the forest) - Higher subsidy of \$1.56 per 200 | ml | | |
| with Endorsement | | 200 ml OP | |
| | (1.56) | | Ensure Plus |
| Liquid (strawberry) - Higher subsidy of \$1.76 per 200 ml with | , , | | |
| , | | 000 000 | |
| Endorsement | | 200 ml OP | |
| | (1.76) | | Fortisip |
| Liquid (vanilla) - Higher subsidy of up to \$1.76 per 200 ml w | ith | | |
| Endorsement | | 237 ml OP | |
| | (1.65) | ···· - · | Ensure Plus |
| | 0.72 | 200 ml OP | 211001011100 |
| | | ZOU IIII OF | Enguro Divo |
| | (1.56) | | Ensure Plus |
| | (1.76) | | Fortisip |
| | | | |

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

ORAL FEED WITH FIBRE 1.5 KCAL/ML - Special Authority see SA1859 on page 281 - Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, or who have severe epidermolysis bullosa. The prescription must be endorsed accordingly.

Liquid (chocolate) - Higher subsidy of \$1.76 per 200 ml with

| Elquid (onocoldic) Triginor Subsidy of \$1.70 per 200 mi with | | | |
|--|--------|-----------|----------------------|
| Endorsement | 0.72 | 200 ml OP | |
| | (1.76) | | Fortisip Multi Fibre |
| Liquid (strawberry) - Higher subsidy of \$1.76 per 200 ml with | | | |
| Endorsement | 0.72 | 200 ml OP | |
| | (1.76) | | Fortisip Multi Fibre |
| Liquid (vanilla) - Higher subsidy of \$1.76 per 200 ml with | | | |
| Endorsement | 0.72 | 200 ml OP | |
| | (1.76) | | Fortisip Multi Fibre |

High Calorie Products

⇒SA1195 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

All of the following:

- 1 Cystic fibrosis; and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements.

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Any of the following:
 - 1.1 any condition causing malabsorption; or
 - 1.2 faltering growth in an infant/child; or
 - 1.3 increased nutritional requirements; or
 - 1.4 fluid restricted: and
 - 2 other lower calorie products have been tried; and
 - 3 patient has substantially increased metabolic requirements or is fluid restricted.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
 - 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

| | Subsidy (Manufacturer's \$ | Price) Subs Per | idised | Brand or Generic Manufacturer |
|--|-----------------------------------|-------------------------------|------------|-------------------------------------|
| ENTERAL FEED 2 KCAL/ML – Special Authority see SA1195 (Liquid | | age – Hospital p 500 ml OP | ✓ Nut C | rison oncentrated |
| | 13.64 | 1,000 ml OP | | sure Two Cal HN TH |
| ORAL FEED 2 KCAL/ML – Special Authority see SA1195 on the Additional subsidy by endorsement is available for patients epidermolysis bullosa. The prescription must be endorsed Liquid (vanilla) – Higher subsidy of \$2.34 per 200 ml with Endorsement | being bolus fed t accordingly. | | g tube, or | • |

Food Thickeners

⇒SA1106 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient has motor neurone disease with swallowing disorder.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

| FOOD THICKENER - Special Authority see SA1106 abov | 'e – Hospital pharmacy | [HP3] | |
|--|-----------------------------------|----------|----------------|
| Powder | 8.29 | 300 g OP | ✓ Nutilis |
| | 24.00 | 380 g OP | ✓ Aptamil Feed |
| | | _ | Thickener |

Gluten Free Foods

The funding of gluten free foods is no longer being actively managed by Pharmac from 1 April 2011. This means that we are no longer considering the listing of new products, or making subsidy, or other changes to the existing listings. As a result we anticipate that the range of funded items will reduce over time. Management of Coeliac disease with a gluten free diet is necessary for good outcomes. A range of gluten free options are available through retail outlets.

⇒SA1729 Special Authority for Subsidy

Initial application — (all patients) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Either:

- 1 Gluten enteropathy has been diagnosed by biopsy; or
- 2 Patient suffers from dermatitis herpetiformis.

Initial application — (paediatric patients diagnosed by ESPGHAN criteria) only from a paediatric gastroenterologist. Approvals valid without further renewal unless notified where the paediatric patient fulfils ESPGHAN criteria for biopsy free diagnosis of coeliac disease.

| GLUTEN FREE BAKING MIX – Special Authority see SA1 | 729 above – Hospital p | harmacy [HP3] | |
|--|------------------------|---------------|--------------------|
| Powder | 2.81 | 1,000 g OP | |
| | (5.15) | | Healtheries Simple |
| | | | Baking Mix |

| | Subsidy | Fu | |
|---|-----------------------|-----------------|----------------------------|
| | (Manufacturer's Price | | ed Generic ✓ Manufacturer |
| | Ψ | - | |
| GLUTEN FREE BREAD MIX – Special Authority see SA1729 of | | | macy [HP3] |
| Powder | | ,000 g OP | |
| | (7.32) | | NZB Low Gluten |
| | | | Bread Mix |
| | 3.51 | | |
| | (10.87) | | Horleys Bread Mix |
| GLUTEN FREE FLOUR - Special Authority see SA1729 on the | previous page - Ho | spital pharmac | y [HP3] |
| Powder | 5.62 2 | ,000 g OP | |
| | (18.10) | | Horleys Flour |
| GLUTEN FREE PASTA - Special Authority see SA1729 on the | previous page – Ho | spital pharmacy | / [HP3] |
| Buckwheat Spirals | | 250 g OP | , [] |
| | (3.11) | 3 - | Orgran |
| Corn and Vegetable Shells | ` ' | 250 g OP | - 3 |
| | (2.92) | 3 - | Orgran |
| Corn and Vegetable Spirals | 2.00 [°] 2 | 250 g OP | Ŭ |
| , | (2.92) | · · | Orgran |
| Rice and Corn Lasagne Sheets | 1.60 | 200 g OP | |
| | (3.82) | | Orgran |
| Rice and Corn Macaroni | 2.00 | 250 g OP | |
| | (2.92) | | Orgran |
| Rice and Corn Penne | 2.00 | 250 g OP | |
| | (2.92) | | Orgran |
| Rice and Maize Pasta Spirals | 2.00 | 250 g OP | |
| | (2.92) | | Orgran |
| Rice and Millet Spirals | | 250 g OP | |
| | (3.11) | | Orgran |
| Rice and corn spaghetti noodles | | 375 g OP | |
| | (2.92) | | Orgran |
| Vegetable and Rice Spirals | | 250 g OP | _ |
| | (2.92) | | Orgran |
| Italian long style spaghetti | | 220 g OP | _ |
| | (3.11) | | Orgran |

(Orgran Buckwheat Spirals to be delisted 1 July 2025)

(Orgran Corn and Vegetable Shells to be delisted 1 July 2025)

(Orgran Corn and Vegetable Spirals to be delisted 1 July 2025)

(Orgran Rice and Corn Lasagne Sheets to be delisted 1 July 2025)

(Orgran Rice and Corn Macaroni to be delisted 1 July 2025)

(Orgran Rice and Corn Penne to be delisted 1 July 2025)

(Orgran Rice and Maize Pasta Spirals to be delisted 1 July 2025)

(Orgran Rice and Millet Spirals to be delisted 1 July 2025)

(Orgran Rice and corn spaghetti noodles to be delisted 1 July 2025)

(Orgran Vegetable and Rice Spirals to be delisted 1 July 2025)

(Orgran Italian long style spagnetti to be delisted 1 July 2025)

Foods And Supplements For Inherited Metabolic Disease

⇒SA2357 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where patient requires dietary management of inherited metabolic disorders.

| | Subsidy | Fully | Brand or |
|--------|------------------|------------|--------------|
| (Manuf | acturer's Price) | Subsidised | Generic |
| | \$ P6 | er 🗸 | Manufacturer |

Supplements For Homocystinuria

AMINOACID FORMULA WITHOUT METHIONINE – Special Authority see SA2357 on the previous page – Hospital pharmacy [HP3]

| Powder (neutral), 36 g sachets | 750.30 | 30 | HCU Anamix Junior |
|---------------------------------------|--------|----------|---------------------|
| Powder, 12.5 g sachets | 349.65 | 30 | ✓ HCU Explore 5 |
| Powder, 25 g sachets | | 30 | ✓ HCU Express 15 |
| Powder (neutral), can | | 500 g OP | ✓ XMET Maxamum |
| Powder (unflavoured), can | | 400 g OP | ✓ HCU Anamix Infant |
| Liquid (juicy berries), 125 ml bottle | | 30 | ✓ HCU Lophlex LQ |
| Liquid (orange), 125 ml bottle | | 36 | ✓ HCU Anamix Junior |
| 1 (0 // | | | 10 |

Supplements For MSUD and short chain enoyl coA hydratase deficiency

AMINOACID FORMULA WITHOUT VALINE, LEUCINE AND ISOLEUCINE - Special Authority see SA2357 on the previous page - Hospital pharmacy [HP3]

| Powder (neutral) 36 g sachets | 750.00 | 30 | MSUD Anamix Junior |
|---------------------------------------|----------|----------|---|
| Powder, 12.5 g sachets | 349.65 | 30 | ✓ MSUD Explore 5 |
| Powder, 25 g sachets | | 30 | ✓ MSUD Express 15 |
| Powder (neutral), can | | 500 g OP | ✓ MSUD Maxamum |
| Powder (orange), can | | 500 g OP | ✓ MSUD Maxamum |
| Powder (unflavoured), can | 260.00 | 400 g OP | MSUD Anamix Infant |
| Liquid (orange) 125 ml bottles | 941.40 | 36 | MSUD Anamix Junior LQ |
| Liquid (juicy berries) 125 ml pouches | 1,684.80 | 30 | ✓ MSUD Lophlex LQ 20 |

| Subsidy | | Fully | Brand or |
|------------------------|-----|------------|--------------|
| (Manufacturer's Price) | | Subsidised | Generic |
| \$ | Per | / | Manufacturer |

Supplements For PKU

| MINOACID FORMULA WITHOUT PHENYLALANINE | - Special Authority see | SA2357 on page | e 287 – Hospital pharmacy [H |
|--|-------------------------|----------------|-------------------------------------|
| Tabs | 99.00 | 75 OP | ✓ Phlexy 10 |
| Powder (Lemon), 34 g sachets | | 30 | ✓ PKU Express 20 |
| Powder (Neutral), 12.5 g sachets | 220.88 | 30 | ✓ PKU Explore 5 |
| Powder (Neutral), 34 g sachets | 883.50 | 30 | ✓ PKU Express 20 |
| Powder (Orange), 25 g sachets | 441.75 | 30 | ✓ PKU Explore 10 |
| Powder (Orange), 34 g sachets | 883.50 | 30 | ✓ PKU Express 20 |
| Powder (Raspberry), 25 g sachets | 441.75 | 30 | ✓ PKU Explore 10 |
| Powder (Tropical), 34 g sachets | 883.50 | 30 | ✓ PKU Express 20 |
| Powder (berry) 28 g sachets | 936.00 | 30 | ✓ PKU Lophlex Powder |
| Powder (chocolate) 36 g sachet | 393.00 | 30 | ✓ PKU Anamix Junior Chocolate |
| Powder (neutral) 28 g sachets | 936.00 | 30 | ✓ PKU Lophlex Powder |
| Powder (neutral) 36 g sachets | 393.00 | 30 | ✓ PKU Anamix Junior |
| Powder (orange) 28 g sachets | | 30 | ✓ PKU Lophlex Powder |
| Powder (orange) 36 g sachet | 393.00 | 30 | ✓ PKU Anamix Junior Orange |
| Powder (unflavoured) 12.5 g sachets | 234.00 | 30 | ✓ PKU First Spoon |
| Powder (vanilla) 36 g sachet | 393.00 | 30 | ✓ PKU Anamix Junior Vanilla |
| Infant formula | 174.72 | 400 g OP | ✓ PKU Anamix Infant |
| Powder (orange) | 320.00 | 500 g OP | ✓ XP Maxamum |
| Powder (unflavoured) | 320.00 | 500 g OP | ✓ XP Maxamum |
| Liquid (berry) | | 125 ml OP | ✓ PKU Anamix Junior LQ |
| Liquid (orange) | 13.10 | 125 ml OP | ✓ PKU Anamix Junior LQ |
| Liquid (forest berries), 250 ml carton | 540.00 | 18 OP | Easiphen Liquid |
| Liquid (juicy tropical) 125 ml | 936.00 | 30 OP | ✓ PKU Lophlex LQ 20 |
| Oral semi-solid (berries) 109 g | 1,123.20 | 36 OP | ✓ PKU Lophlex Sensation 20 |
| Powder (neutral), 400 g can | 715.16 | 4 OP | ✓ PKU Start |
| Liquid (juicy berries) 62.5 ml | | 60 OP | ✓ PKU Lophlex LQ 10 |
| Liquid (juicy berries) 125 ml | | 30 OP | ✓ PKU Lophlex LQ 20 |
| Liquid (juicy orange) 125 ml | | 30 OP | ✓ PKU Lophlex LQ 20 |

✓ fully subsidised 289

| | Cubaidu | | Fully Drand or |
|---|-----------------------------------|----------|--|
| | Subsidy (Manufacturer's Price) | Sı | Fully Brand or ubsidised Generic |
| | \$ | Per | ✓ Manufacturer |
| GLYCOMACROPEPTIDE AND AMINO ACID CONTAINS SOM | E PHENYLALANINE | – Specia | al Authority see SA2357 on |
| page 287 – Hospital pharmacy [HP3] | | • | , |
| Powder (Banana) 35 g sachets | 930.00 | 30 | ✓ PKU |
| , , , , , , , , , , , , , , , , , , , | | | sphere20 Banana |
| Powder (Berry), 20 g sachets | 449.28 | 60 | ✓ PKU Restore |
| (),, | | | Powder |
| Powder (Chocolate) 32 g sachets | 898.56 | 30 | ✓ PKU Build |
| 3 (* , 3 | | | 20 Chocolate |
| Powder (Chocolate) 35 g sachets | 930.00 | 30 | ✓ PKU |
| 3 · · · · (· · · · · · · · · · · · · · | | | sphere20 Chocolate |
| D 1 (1) 25 1 . | 000.00 | | |
| Powder (Lemon) 35 g sachets | 930.00 | 30 | ✓ PKU |
| | | | sphere20 Lemon |
| Powder (Lemonade) 33.4 g sachets | 936.00 | 30 | ✓ PKU GMPro Ultra |
| | | | Lemonade |
| Powder (Neutral), 15 g sachets | | 30 | PKU Build 10 |
| Powder (Orange), 20 g sachets | 449.28 | 60 | ✓ PKU Restore |
| | | | Powder |
| Powder (Raspberry Lemonade) 31 g sachets | 898.56 | 30 | ✓ PKU Build |
| | | | 20 Raspberry |
| | | | Lemonade |
| Powder (Smooth) 31 g sachets | 898.56 | 30 | ✓ PKU Build |
| | | | 20 Smooth |
| Powder (Vanilla) 33 g sachets | | 30 | PKU Build 20 Vanilla |
| Powder (neutral), 40 g sachets | | 30 | Glytactin Bettermilk |
| Powder (unflavoured) 12.5 g sachets | | 30 | ✓ PKU GMPro Mix-In |
| Powder (vanilla) 33.4 g sachets | 936.00 | 30 | PKU GMPro Ultra |
| | | | Vanilla |
| Powder (Red Berry) 35 g sachets | 930.00 | 30 | ✓ PKU sphere20 Red |
| | | | Berry |
| Powder (Vanilla) 35 g sachets | 930.00 | 30 | ✓ PKU |
| | | | sphere20 Vanilla |
| Liquid (neutral), 250 ml carton | 280.80 | 18 | ✓ PKU GMPro LQ |
| Liquid (original), 250 ml carton | 684.45 | 30 OP | PKU Glytactin RTD |
| | | | 15 |
| Liquid (Coffee Mocha), 250 ml carton | 684.45 | 30 OP | ✓ PKU Glytactin RTD |
| | | | 15 Lite |
| Liquid (chocolate), 250 ml carton | 684.45 | 30 OP | ✓ PKU Glytactin RTD |
| | | | 15 |
| Liquid (vanilla), 250 ml carton | 684.45 | 30 OP | ✓ PKU Glytactin RTD |
| | | | 15 Lite |
| | | | |

Foods

LOW PROTEIN BAKING MIX - Special Authority see SA2357 on page 287 - Hospital pharmacy [HP3]
Powder8.55 500 g OP ✓ Loprofin Mix

| | Subsidy | | Fully | Brand or |
|--|---------------------|-----------------|--------|-----------------------------------|
| | (Manufacturer's Pri | | | Generic |
| | \$ | Per | _ | Manufacturer |
| LOW PROTEIN PASTA - Special Authority see SA2357 on page | e 287 – Hospital p | harmacy [HP3] | | |
| Animal shapes | 12.39 | 500 g OP | | Loprofin |
| Lasagne | 6.19 | 250 g OP | | Loprofin |
| Low protein rice pasta | 12.39 | 500 g OP | 1 | Loprofin |
| Macaroni | 6.19 | 250 g OP | 1 | Loprofin |
| Penne | 12.39 | 500 g OP | 1 | Loprofin |
| Spaghetti | 12.39 | 500 g OP | 1 | Loprofin |
| Spirals | 12.39 | 500 g OP | 1 | Loprofin |
| Supplements for Tyrosinaemia | | | | |
| AMINOACID FORMULA WITHOUT PHENYLALANINE AND TYP | ROSINE - Special | Authority see | SA2 | 357 on page 287 – Hospita |
| pharmacy [HP3] | • | , | | |
| Powder (Neutral), 12.5 g sachets | 349.65 | 30 | 1 | TYR Explore 5 |
| Powder (neutral) 36 g sachets | | 30 | | TYR Anamix Junior |
| Powder, can | | 400 g OP | | TYR Anamix Infant |
| Liquid (juicy berries) 125 ml pouches | | 30 | _ | TYR Lophlex LQ 20 |
| Liquid (orange) 125 ml bottle | | 36 | | TYR Anamix Junior |
| Liquid (orange) 125 mi bottle | 341.40 | 50 | ٠ | LQ |
| GLYCOMACROPEPTIDE AND AMINO ACID CONTAINS SOME | TYROSINE AND | ΡΗΕΝΙΊ ΔΙ ΔΝ | MNE | - Special Authority see |
| SA2357 on page 287 – Hospital pharmacy [HP3] | TITIOONIL AND | IIILINILALAI | VII V. | opeolal Authority see |
| Powder (Red Berry), 35 g sachets | 1 308 60 | 30 | 1 | TYR Sphere 20 |
| Powder (Vanilla), 35 g sachets | | 30 | | TYR Sphere 20 |
| Fowder (varilla), 35 g sacriets | 1,390.00 | 30 | • | i in Spliele 20 |
| Supplements for Organic Acidaemias | | | | |
| AMINOACID FORMULA WITHOUT ISOLEUCINE, METHIONINE | . THREONINE AN | ND VALINE - | Spec | cial Authority see SA2357 |
| on page 287 – Hospital pharmacy [HP3] | ., | | Opoc | , a. / lat. only 500 0/ 12001 |
| Powder, can | 260.00 | 400 g OP | 1 | MMA/PA Anamix |
| 1 011001, 001 | | 100 g 01 | - | Infant |
| ANNINO A CID ECDANIII A MITUOLIT METUONINE TUDEONINE | AND VALINE (| D | | |
| AMINOACID FORMULA WITHOUT METHIONINE, THREONINE | AND VALINE - | Special Authori | ty se | e SA2357 on page 287 – |
| Hospital pharmacy [HP3] | | | | |
| Powder (neutral), 18 g sachets | 750.30 | 30 | • | MMA/PA Anamix |
| | | | | Junior |
| Powder, 12.5 g sachets | 349.65 | 30 | 1 | MMA/PA Explore 5 |
| Powder, 25 g sachets | 1,048.95 | 30 | 1 | MMA/PA Express 15 |
| Supplements for Glutaric Aciduria type 1 | | | | |
| AMINOACID FORMULA WITHOUT LYSINE - Special Authority | see SA2357 on n | ane 287 – Hoer | nital | nharmacy [HP3] |
| Powder (neutral), 18 g sachets | | • | | GA1 Anamix Junior |
| , , | | 30 | | |
| Powder, 12.5 g sachets | | 30 | | GA Explore 5 GA1 Anamix Infant |
| Powder, can | 200.00 | 400 g OP | • | GAT Anamix iniant |
| Supplements for Glycogen Storage Disease | | | | |
| HIGH AMYLOPECTIN CORN-STARCH - Special Authority see | SA2357 on page 2 | 287 – Hospital | phar | macv [HP3] |
| Powder, 60 g sachets | | 30 | | Glycosade |
| | | | | . , , |
| Single dose amino acids | | | | |
| ARGININE – Special Authority see SA2357 on page 287 – Hosp | ital nharmacy [HD | 31 | | |
| Powder, 4 g sachets | | 30 | 1 | Arginine2000 |
| 1 371301, 1 g 30011010 | | 00 | - | go. |

✓ fully subsidised 291

| | Subsidy (Manufacturer's Price \$ |) Per | Fully Subsidised | |
|---|---|---------------------------------|--------------------------------------|--|
| CITRULLINE – Special Authority see SA2357 on page 287 – H Powder, 4 g sachets | | ² 3] 30 | ✓ | Citrulline1000 |
| ISOLEUCINE - Special Authority see SA2357 on page 287 - F Powder, 4 g sachets | | ³ 3] | ✓ | Isoleucine50 |
| LEUCINE – Special Authority see SA2357 on page 287 – Hosp Powder, 4 g sachets | 141.05 | 30 | | Leucine100 |
| PHENYLALANINE – Special Authority see SA2357 on page 28 Powder, 4 g sachets | | y [HP3 30 | | Phenylalanine50 |
| TYROSINE – Special Authority see SA2357 on page 287 – Ho Powder, 4 g sachets | |] 30 | ✓ | Tyrosine1000 |
| VALINE – Special Authority see SA2357 on page 287 – Hospit Powder, 4 g sachets | | 30 | • | Valine50 |
| Other Fat Modified Products | | | | |
| ELEMENTAL FEED WITH HIGH MEDIUM CHAIN TRIGLYCEF pharmacy [HP3] | RIDES - Special Auth | ority se | ee SA2357 | 7 on page 287 – Hospital |
| Powder (neutral), 100 g sachets | 47.01 | 10 | ✓ | Emsogen |
| Carbohydrate and Fat with added vitamins and | l minerals | | | |
| PROTEIN FREE SUPPLEMENT CONTAINING CARBOHYDRA Authority see SA2357 on page 287 – Hospital pharmacy [HP3] Powder (neutral), can | | ED VIT | | ND MINERALS - Special Energivit |
| Essential Amino Acids | | | | |
| ESSENTIAL AMINOACID FORMULA - Special Authority see S Powder (neutral), can | | – Hosp 200 g O | | acy [HP3] Essential Amino Acid Mix |
| Infant Formulae | | | | |
| For Williams Syndrome | | | | |
| ■ SA1110 Special Authority for Subsidy Initial application only from a dietitian, relevant specialist or vo year where the patient is an infant suffering from Williams Synd Renewal only from a dietitian, relevant specialist, vocationally recommendation of a dietitian, relevant specialist or vocationally applications meeting the following criteria: Both: 1 The treatment remains appropriate and the patient is being 2 General Practitioners must include the name of the dietit practitioner and date contacted. | rome and associated egistered general pract registered general pract registered general practical promotes and treatment reatments. | hyperc ctitione ractition | alcaemia. r or gener ner. Appr | al practitioner on the ovals valid for 1 year for |

LOW CALCIUM INFANT FORMULA - Special Authority see SA1110 above - Hospital pharmacy [HP3]

Powder46.18

400 g OP

✓ Locasol

| | Subsidy | Fu | illy | Brand or |
|-----|---------------------|----------|------|--------------|
| (Ma | nufacturer's Price) | Subsidis | ed | Generic |
| | \$ | Per | ✓ | Manufacturer |

Gastrointestinal and Other Malabsorptive Problems

| AMINO ACID FORMULA - Special Authority see SA20 | 92 below – Hospital pharmad | y [HP3] | |
|---|-----------------------------|----------|---|
| Powder | 43.60 | 400 g OP | ✓ Alfamino✓ Alfamino Junior |
| Powder (unflavoured) | 55.61 | 400 g OP | ✓ Neocate Gold ✓ Neocate Junior Unflavoured |
| | 65.72 | | ✓ Neocate SYNEO✓ Elecare✓ Elecare LCP |
| Powder (vanilla) | 55.61 | 100 g OP | ✓ Neocate Junior Vanilla |
| | 65.72 | | ✓ Elecare |

⇒SA2092 Special Authority for Subsidy

Initial application — (Infants under 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 History of anaphylaxis to cow's milk protein formula or dairy products; or
- 2 Eosinophilic oesophagitis; or
- 3 Ultra-short gut; or
- 4 Severe Immune deficiency; or
- 5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
- 6 Both
 - 6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 6.2 Either:
 - 6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or 6.2.2 Patient has IoE mediated allerov.

Initial application — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or
 - 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
 - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products; or
 - 2.2 Eosinophilic oesophagitis; or
 - 2.3 Ultra-short gut; or
 - 2.4 Severe Immune deficiency: or
 - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
 - 2.6 Both:
 - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 2.6.2 Either
 - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval

continued...

✓ fully subsidised 293

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

continued...

number: or

2.6.2.2 Patient has IgE mediated allergy.

Renewal — (Infants up to 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has IgE mediated allergy; and
 - 1.2 All of the following:
 - 1.2.1 Patient remains allergic to cow's milk; and
 - 1.2.2 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy or extensively hydrolysed infant formula has been undertaken; and
 - 1.2.3 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
 - 1.2.4 Amino acid formula is required for a nutritional deficit; and
 - 1.2.5 It has been more than three months from the previous approval; or

2 Both:

- 2.1 Patient has non IgE mediated severe gastrointestinal intolerance (including eosinophilic oesophagitis, ultra-short gut and severe immune deficiency); and
- 2.2 All of the following:
 - 2.2.1 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy, or extensively hydrolysed infant formula has been undertaken; and
 - 2.2.2 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
 - 2.2.3 Amino acid formula is required for a nutritional deficit; and
 - 2.2.4 It has been more than three months from the previous approval.

Renewal — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 Either:

- 1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or
- 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
 - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products: or
 - 2.2 Eosinophilic oesophagitis; or
 - 2.3 Ultra-short gut; or
 - 2.4 Severe Immune deficiency; or
 - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
 - 2.6 Both:
 - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 2.6.2 Either:
 - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or
 - 2.6.2.2 Patient has IgE mediated allergy.

Initial application — (for patients who have a current funding under Special Authority form SA1557) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 months for applications meeting the following criteria:

continued...

| Sut | bsidy F | ully | Brand or |
|------------|-----------------------|------|--------------|
| (Manufacti | urer's Price) Subsidi | sed | Generic |
| | \$ Per | ✓ | Manufacturer |

continued...

All of the following:

- 1 Patient has a valid Special Authority approval for extensively hydrolysed formula (SA1557); and
- 2 Extensively hydrolysed formula (Aptamil Gold+ Pepti Junior, AllerPro SYNEO 1 and 2) is unable to be supplied at this time; and
- 3 The approval only applies to funded dispensings of Neocate Gold and Neocate Syneo.

Note: This criteria is short term funding to cover an out-of-stock situation on some extensively hydrolysed formula powder funded under Special Authority form SA1557. There is no renewal criteria under this restriction.

⇒SA1953 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has impaired gastrointestinal function and either cannot tolerate polymeric feeds, or polymeric feeds are unsuitable; and
- 2 Any of the following:
 - 2.1 Severe malabsorption; or
 - 2.2 Short bowel syndrome; or
 - 2.3 Intractable diarrhoea; or
 - 2.4 Biliary atresia; or
 - 2.5 Cholestatic liver diseases causing malabsorption; or
 - 2.6 Cystic fibrosis; or
 - 2.7 Proven fat malabsorption; or
 - 2.8 Severe intestinal motility disorders causing significant malabsorption; or
 - 2.9 Intestinal failure: or
 - 2.10 Both:
 - 2.10.1 The patient is currently receiving funded amino acid formula; and
 - 2.10.2 The patient is to be trialled on, or transitioned to, an enteral liquid peptide formula; and
- 3 Either:
 - 3.1 A semi-elemental or partially hydrolysed powdered feed has been reasonably trialled and considered unsuitable; or
 - $3.2\,$ For step down from intravenous nutrition.

Note: A reasonable trial is defined as a 2-4 week trial.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 An assessment as to whether the patient can be transitioned to a cows milk protein or soy infant formula or extensively hydrolysed formula has been undertaken; and
- 2 The outcome of the assessment is that the patient continues to require an enteral liquid peptide formula; and
- 3 General practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

| EXTENSIVELY HYDROLYSED FORMULA - Spec | cial Authority see SA1557 on the | e next page | Hospital pharmacy [HP3] |
|---------------------------------------|----------------------------------|-------------|---|
| Powder | 18.10 | 450 g OP | ✓ Pepti-Junior |
| | 36.20 | 900 g OP | ✓ Allerpro Syneo 1 |
| | | - | ✓ Allerpro Syneo 2 |

✓ fully subsidised

295

Subsidy (Manufacturer's Price)

Fully Subsidised Brand or Generic Manufacturer

⇒SA1557 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 Cows milk formula is inappropriate due to severe intolerance or allergy to its protein content; and
 - 1.2 Fither:
 - 1.2.1 Soy milk formula has been reasonably trialled without resolution of symptoms; or
 - 1.2.2 Soy milk formula is considered clinically inappropriate or contraindicated; or
- 2 Severe malabsorption: or
- 3 Short bowel syndrome; or
- 4 Intractable diarrhoea; or
- 5 Biliary atresia: or
- 6 Cholestatic liver diseases causing malsorption; or
- 7 Cystic fibrosis: or
- 8 Proven fat malabsorption; or
- 9 Severe intestinal motility disorders causing significant malabsorption; or
- 10 Intestinal failure; or
- 11 All of the following:
 - 11.1 For step down from Amino Acid Formula: and
 - 11.2 The infant is currently receiving funded amino acid formula; and
 - 11.3 The infant is to be trialled on, or transitioned to, an extensively hydrolysed formula; and
 - 11.4 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

Note: A reasonable trial is defined as a 2-4 week trial, or signs of an immediate IgE mediated allergic reaction.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 An assessment as to whether the infant can be transitioned to a cows milk protein or soy infant formula has been undertaken: and
- 2 The outcome of the assessment is that the infant continues to require an extensively hydrolysed infant formula; and
- 3 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

Fluid Restricted

PAEDIATRIC ORAL/ENTERAL FEED 1 KCAL/ML - Special Authority see SA1698 below - Hospital pharmacy [HP3] ✓ Infatrini 125 ml OP

⇒SA1698 Special Authority for Subsidy

Initial application only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is fluid restricted or volume intolerant and has been diagnosed with faltering growth; and
- 2 Patient is under the care of a paediatrician or dietitian who has recommended treatment with a high energy infant formula;
- 3 Patient is under 18 months of age or weighs less than 8 kg.

continued...

| Subsidy | Ful | y Brand or | Т |
|------------------------|-----------|--------------|---|
| (Manufacturer's Price) | Subsidise | d Generic | |
| \$ | Per • | Manufacturer | |

continued...

Note: "Volume intolerant" patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

Renewal only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient continues to be fluid restricted or volume intolerant and has faltering growth; and
- 2 Patient is under the care of a hospital paediatrician or dietitian who has recommended treatment with a high energy infant formula; and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: 'Volume intolerant' patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

Ketogenic Diet

⇒SA1197 Special Authority for Subsidy

Initial application only from a metabolic physician or paediatric neurologist. Approvals valid for 3 months where the patient has intractable epilepsy, pyruvate dehydrogenase deficiency or glucose transported type-1 deficiency and other conditions requiring a ketogenic diet.

Renewal only from a metabolic physician or paediatric neurologist. Approvals valid for 2 years where the patient is on a ketogenic diet and the patient is benefiting from the diet.

| HIGH FAT LOW CARBOHYDRATE FORMULA - Special Authority see SA11 | 97 above – Hospital pharmacy [HP3] |
|--|------------------------------------|
| Powder (unflavoured)36.92 | 300 g OP 	✓ KetoCal 4:1 |
| | ✓ Ketocal 3:1 |
| Powder (vanilla)36.92 | 300 g OP 		✓ KetoCal 4:1 |

✓ fully subsidised 297

SECTION I: NATIONAL IMMUNISATION SCHEDULE

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

Vaccinations

BACILLUS CALMETTE-GUERIN VACCINE - [Xpharm]

For infants at increased risk of tuberculosis. Increased risk is defined as:

- 1) living in a house or family with a person with current or past history of TB; or
- 2) having one or more household members or carers who within the last 5 years lived in a country with a rate of TB > or egual to 40 per 100,000 for 6 months or longer; or
- 3) during their first 5 years will be living 3 months or longer in a country with a rate of TB > or equal to 40 per 100,000 Note a list of countries with high rates of TB are available at www.health.govt.nz/tuberculosis (search for downloads) or www.bcgatlas.org/index.php.

| Inj Mycobacterium | bovis BCG (Bacillus | Calmette-Guerin), | |
|-------------------|-----------------------|----------------------|------|
| Danish strain | 1331, live attenuated | I, vial with diluent | 0.00 |

COVID-19 VACCINE - [Xpharm]

10

✓ BCG Vaccine AJV

Inj 3 mcg bretovameran per 0.3 ml, 0.48 ml vial; infant vaccine,

yellow cap......0.00

10

✓ Comirnaty Omicron (JN.1)

Up to three doses for previously unvaccinated children aged 6 months - 4 years at high risk of severe illness.

Ini 10 mcg bretovameran per 0.3 ml. 0.48 ml vial; paediatric vaccine, light blue cap0.00

10

✓ Comirnaty Omicron (JN.1)

Either:

- 1) One dose for previously unvaccinated children aged 5-11 years old; or
- 2) Up to three doses for immunocompromised children aged 5-11 years old.

Inj 30 mcg bretovameran per 0.3 ml, 0.48 ml vial; adult vaccine,

10

✓ Comirnaty Omicron (JN.1)

Any of the following:

- 1) One dose for previously unvaccinated people aged 12-15 years old; or
- 2) Up to three doses for immunocompromised people aged 12-15 years old; or
- 3) Up to two doses for previously unvaccinated people 16-29 years old; or
- 4) Up to four doses for people aged 16-29 at high risk of severe illness; or
- 5) One dose for previously unvaccinated people aged 30 and older; or
- 6) One additional dose every 6 months for previously vaccinated people aged 30 years and over additional dose is given at least 6 months after last dose.

| Subsidy (Manufacturer's Price) | Fully Subsidised | |
|-----------------------------------|---------------------|--------------|
| \$ | Per 🗸 | Manufacturer |

DIPHTHERIA. TETANUS AND PERTUSSIS VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- c)
- A) Funded for any of the following criteria:
 - 1) A single dose for pregnant women in the second or third trimester of each pregnancy; or
 - 2) A single dose for parents or primary caregivers of infants admitted to a Neonatal Intensive Care Unit or Specialist Care Baby Unit for more than 3 days, who had not been exposed to maternal vaccination at least 14 days prior to birth: or
 - A course of up to four doses is funded for children from age 7 up to the age of 18 years inclusive to complete full primary immunisation; or
 - 4) An additional four doses (as appropriate) are funded for (re-)immunisation for patients post haematopoietic stem cell transplantation or chemotherapy; pre or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or
 - 5) A single dose for vaccination of patients aged from 65 years old; or
 - 6) A single dose for vaccination of patients aged from 45 years old who have not had 4 previous tetanus doses; or
 - 7) For vaccination of previously unimmunised or partially immunised patients; or
 - 8) For revaccination following immunosuppression; or
 - 9) For boosting of patients with tetanus-prone wounds.

Notes: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.

- B) Contractors will be entitled to claim payment from the Funder for the supply of diphtheria, tetanus and pertussis vaccine to patients eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the diphtheria, tetanus and pertussis vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraphs 1 – 9 above.

| pertussis toxoid, 8 mcg pertussis filamentous | | |
|--|----|------------|
| haemagglutinin and 2.5 mcg pertactin in 0.5 ml prefilled | | |
| syringe | 10 | ✓ Boostrix |

| Subsidy (Manufacturer's Price) | Sı | Fully ubsidised | Brand or Generic | |
|-----------------------------------|-----|--------------------|---------------------|--|
| \$ | Per | √ | Manufacturer | |

DIPHTHERIA, TETANUS, PERTUSSIS AND POLIO VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- c)
- A) Funded for any of the following:
 - 1) A single dose for children up to the age of 7 who have completed primary immunisation; or
 - 2) A course of four vaccines is funded for catch up programmes for children (to the age of 10 years) to complete full primary immunisation: or
 - 3) An additional four doses (as appropriate) are funded for (re-)immunisation for people post HSCT, or chemotherapy; pre- or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or
 - 4) Five doses will be funded for children requiring solid organ transplantation.
- B) Contractors will be entitled to claim payment from the Funder for the supply of Diphtheria, tetanus, pertussis and polio vaccine to people eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Diphtheria, tetanus, pertussis and polio vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.

Inj 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg

pertussis toxoid, 25 mcg pertussis filamentous haemagglutinin, 8 mcg pertactin and 80 D-antigen units

10 ✓ Infanrix IPV

DIPHTHERIA, TETANUS, PERTUSSIS, POLIO, HEPATITIS B AND HAEMOPHILUS INFLUENZAE TYPE B VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- c)
- A) Funded for children meeting any of the following criteria
 - 1) Up to four doses for children under the age of 10 years for primary immunisation; or
 - 2) An additional four doses (as appropriate) for (re-)immunisation of children under the age of 18 years post haematopoietic stem cell transplantation; or
 - 3) An additional four doses (as appropriate) for (re-)immunisation of children under the age of 10 years who are post chemotherapy; pre or post splenectomy; undergoing renal dialysis and other severely immunosuppressive regimens: or
 - 4) Up to five doses for children under the age of 10 years receiving solid organ transplantation.
- B) Contractors will be entitled to claim payment from the Funder for the supply of Diphtheria, tetanus, pertussis, polio. hepatitis B and haemophilus influenzae type b vaccine to people eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Diphtheria, tetanus, pertussis, polio, hepatitis B and haemophilus influenzae type b vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Note: A course of up-to four vaccines is funded for catch up programmes for children (up to and under the age of 10 years) to complete full primary immunisation. Please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes.

Inj 30IU diphtheria with 40IU tetanus and 25mcg pertussis toxoids, 25mcg pertussis filamentous haemagglutinin, 8mcg pertactin, 80D-AgU polio virus, 10mcg hepatitis B antigen, 10mcg H. influenzae type b with tetanus toxoid

✓ Infanrix-hexa

10

✓ Havrix 1440

✓ Havrix Junior

1

| NATIONAL IMMUNISATION SCHEDULE | | | | | | |
|--|---|--------------------|--------------|-------------------------------------|--|--|
| | Subsidy (Manufacturer's Price) \$ | F Subsid Per | ully ised | Brand or Generic Manufacturer | | |
| HAEMOPHILUS INFLUENZAE TYPE B VACCINE | | | | _ | | |
| a) Only on a prescription | | | | | | |
| b) No patient co-payment payable c) | | | | | | |
| A) One dose for people meeting any of the followin | g: | | | | | |
| For primary vaccination in children; or | | | | | | |
| An additional dose (as appropriate) is fund transplantation, or chemotherapy; function | | | | | | |
| transplant, pre or post cochlear implants, r | | | | | | |
| For use in testing for primary immunodefic physician or paediatrician. | iency diseases, on the re- | commendat | ion of | an internal medicine | | |
| B) Contractors will be entitled to claim payment froit vaccine to people eligible under the above criter | ia pursuant to their contra | ct with Hea | ılth Ne | w Zealand (Health NZ) | | |
| for subsidised immunisation, and they may only in the Pharmaceutical Schedule. | do so in respect of the Ha | aemophilus | influer | nzae type b vaccine listed | | |
| C) Contractors may only claim for populations withing sub-set of the population described in paragraph | | ered by thei | r contr | act, which may be a | | |
| Inj 10 mcg vial with diluent syringe | 0.00 | 1 | ✓ <u>Ac</u> | ct-HIB | | |
| HEPATITIS A VACCINE - [Xpharm] | | | | | | |
| Funded for patients meeting any of the following criteria: | | | | | | |
| 1) Two vaccinations for use in transplant patients; or | | | | | | |
| Two vaccinations for use in children with chronic live | · | | | | | |
| One dose of vaccine for close contacts of known he | patitis A cases. | | | | | |

| | Subsidy (Manufacturer's Price) | S | Fully ubsidised | Brand or Generic | |
|--|-----------------------------------|-----|--------------------|---------------------|--|
| | \$ | Per | 1 | Manufacturer | |
| HEPATITIS B RECOMBINANT VACCINE - [Xnharm] | | | | | |

✓ Engerix-B

Funded for patients meeting any of the following criteria:

- 1) for household or sexual contacts of known acute hepatitis B patients or hepatitis B carriers; or
- 2) for children born to mothers who are hepatitis B surface antigen (HBsAq) positive; or
- 3) for children up to and under the age of 18 years inclusive who are considered not to have achieved a positive serology and require additional vaccination or require a primary course of vaccination; or
- 4) for HIV positive patients: or
- 5) for hepatitis C positive patients; or
- 6) for patients following non-consensual sexual intercourse; or
- 7) for patients prior to planned immunosuppression for greater than 28 days; or
- 8) for patients following immunosuppression; or
- 9) for solid organ transplant patients; or
- 10) for post-haematopoietic stem cell transplant (HSCT) patients; or
- 11) following needle stick injury.

Inj 20 mcg per 1 ml prefilled syringe......0.00 ✓ Engerix-B

Funded for patients meeting any of the following criteria:

- 1) for household or sexual contacts of known acute hepatitis B patients or hepatitis B carriers; or
- 2) for children born to mothers who are hepatitis B surface antigen (HBsAg) positive; or
- 3) for children up to and under the age of 18 years inclusive who are considered not to have achieved a positive serology and require additional vaccination or require a primary course of vaccination; or
- 4) for HIV positive patients; or
- 5) for hepatitis C positive patients; or
- 6) for patients following non-consensual sexual intercourse; or
- 7) for patients prior to planned immunosuppression for greater than 28 days; or
- 8) for patients following immunosuppression; or
- 9) for solid organ transplant patients; or
- 10) for post-haematopoietic stem cell transplant (HSCT) patients; or
- 11) following needle stick injury; or
- 12) for dialysis patients; or
- 13) for liver or kidney transplant patients.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

HUMAN PAPILLOMAVIRUS (6, 11, 16, 18, 31, 33, 45, 52 AND 58) VACCINE [HPV]

- a) Maximum of 1 inj per prescription
- b) Only on a prescription
- c) No patient co-payment payable
- d
- a) A) Any of the following:
 - 1) Maximum of two doses for children aged 14 years and under; or
 - 2) Maximum of three doses for people meeting any of the following criteria:
 - 1) People aged 15 to 26 years inclusive; or
 - 2) Either:

People aged 9 to 26 years inclusive who have

- 1) Confirmed HIV infection; or
- 2) Received a transplant (including stem cell): or
- 3) Maximum of four doses for people aged 9 to 26 years inclusive post chemotherapy
- B) Contractors will be entitled to claim payment from the Funder for the supply of Human papillomavirus vaccine to people eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Human papillomavirus vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraphs A above.

| Inj 270 mcg in 0.5 ml syringe | 0.00 | 10 | Gardasil 9 |
|-------------------------------|------|----|------------|
|-------------------------------|------|----|------------|

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer |
|---|---|-----|---------------------|-------------------------------------|
| INFLUENZA VACCINE Inj 60 mcg in 0.5 ml syringe (quadrivalent vaccine) | 120.00 | 10 | | nfluvac Tetra (2024 formulation) |

| Subsidy | | Fully | Brand or | |
|------------------------|-----|----------|--------------|--|
| (Manufacturer's Price) | Su | bsidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

- a) Maximum of 1 inj per prescription
- b) Only on a prescription
- c) No patient co-payment payable
- d

A) INFLUENZA VACCINE

is available each year for patients who meet the following criteria, as set by Pharmac:

- a) all people 65 years of age and over; or
- b) people under 65 years of age who:
 - i) have any of the following cardiovascular diseases:
 - a) ischaemic heart disease, or
 - b) congestive heart failure, or
 - c) rheumatic heart disease, or
 - d) congenital heart disease, or
 - e) cerebo-vascular disease; or
 - ii) have either of the following chronic respiratory diseases:
 - a) asthma, if on a regular preventative therapy, or
 - b) other chronic respiratory disease with impaired lung function; or
 - iii) have diabetes; or
 - iv) have chronic renal disease; or
 - v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
 - vi) have any of the following other conditions:
 - a) autoimmune disease, or
 - b) immune suppression or immune deficiency, or
 - c) HIV, or
 - d) transplant recipients, or
 - e) neuromuscular and CNS diseases/disorders, or
 - f) haemoglobinopathies, or
 - g) are children on long term aspirin, or
 - h) have a cochlear implant, or
 - i) errors of metabolism at risk of major metabolic decompensation, or
 - i) pre and post splenectomy, or
 - k) Down syndrome, or
 - vii) are pregnant; or
- c) children 4 years of age and under who have been hospitalised for respiratory illness or have a history of significant respiratory illness; or
- d) people under 65 years of age who:
 - i) have any of the following serious mental health conditions:
 - a) schizophrenia, or
 - b) major depressive disorder, or
 - c) bipolar disorder, or
 - d) schizoaffective disorder, or
 - ii) are currently accessing secondary or tertiary mental health and addiction services; or

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy,
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Contractors will be entitled to claim payment for the supply of influenza vaccine to patients eligible under the above criteria pursuant to their contract with Health NZ for subsidised immunisation, and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

| Subsidy | | Fully | Brand or | |
|-----------------------|-----|------------|--------------|--|
| (Manufacturer's Price | e) | Subsidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

MEASLES. MUMPS AND RUBELLA VACCINE

- a) Only on a prescription
- b) No patient co-payment payable

c)

A) Measles, mumps and rubella vaccine

A maximum of two doses for any patient meeting the following criteria:

- 1) For primary vaccination in children; or
- 2) For revaccination following immunosuppression; or
- 3) For any individual susceptible to measles, mumps or rubella; or
- 4) A maximum of three doses for children who have had their first dose prior to 12 months.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes. Although a price is listed for the vaccine, doctors can still order measles mumps and rubella vaccine free of charge, as with other Schedule vaccines.

- B) Contractors will be entitled to claim payment for the supply of measles, mumps and rubella vaccine to patients eligible under the above criteria pursuant to their contract with Health NZ for subsidised immunisation, and they may only do so in respect of the measles, mumps and rubella vaccine listed in the Pharmaceutical Schedule.
- C) Contractors can only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

| Inj, measles virus 1,000 CCID50, mumps virus 5,012 CCID50, | | |
|--|----|-----------|
| Rubella virus 1,000 CCID50; prefilled syringe/ampoule of | | |
| diluent 0.5 ml | 10 | ✓ Priorix |

| Subsidy (Manufacturer's Price) \$ | Full Subsidise Per | | |
|---|--------------------------|--|--|
|---|--------------------------|--|--|

MENINGOCOCCAL (GROUPS A, C, Y AND W-135) CONJUGATE VACCINE

1 U.5 IIII VIAI

✓ MenQuadfi

- a) Only on a prescription
- b) No patient co-payment payable

C)

- A) Any of the following:
 - Up to three doses and a booster every five years for patients pre- and post splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre or post solid organ transplant; or
 - 2) One dose for close contacts of meningococcal cases of any group; or
 - 3) One dose for person who has previously had meningococcal disease of any group; or
 - 4) A maximum of two doses for bone marrow transplant patients; or
 - 5) A maximum of two doses for person pre- and post-immunosuppression*; or
- B) Both:
 - 1) Person is aged between 13 and 25 years, inclusive; and
 - 2) Either:
 - One dose for individuals who are entering within the next three months, or in their first year of living in boarding school hostels, tertiary education halls of residence, military barracks, Youth Justice residences, or prisons; or
 - 2) One dose for individuals who turn 13 years of age while living in boarding school hostels.
- C) Contractors will be entitled to claim payment from the Funder for the supply of Meningococcal A, C, Y and W-135 vaccine to patients eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Meningococcal A, C, Y and W-135 vaccine listed in the Pharmaceutical Schedule.
- D) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraphs A-B above.

Note: children under seven years of age require two doses 8 weeks apart, a booster dose three years after the primary series and then five yearly.

*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days.

Inj 5 mcg of each meningococcal polysaccharide conjugated to a total of approximately 44 mcg of tetanus toxoid carrier

per 0.5 ml vial - [Xpharm]......0.00

✓ Nimenrix

A) Both:

- 1) The child is under 12 months of age; and
- 2) Any of the following:
 - A maximum of three doses (dependant on age at first dose) for patients pre- and post- splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre- or post- solid organ transplant; or
 - A maximum of three doses (dependant on age at first dose) for close contacts of meningococcal cases of any group; or
 - A maximum of three doses (dependant on age at first dose) for child who has previously had meningococcal disease of any group; or
 - 4) A maximum of three doses (dependant on age at first dose) for bone marrow transplant patients; or
 - A maximum of three doses (dependant on age at first dose) for child pre- and post-immunosuppression*.

Note: infants from 6 weeks to less than 6 months of age require a 2+1 schedule, infants from 6 months to less than 12 months of age require a 1+1 schedule. Refer to the Immunisation Handbook for recommended booster schedules with meningococcal ACWY vaccine.

*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days.

| Subsidy | | Fully | Brand or | |
|----------------------|-----|------------|--------------|--|
| (Manufacturer's Pric | e) | Subsidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

MENINGOCOCCAL B MULTICOMPONENT VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- c) Any of the following:
 - A) Three doses for children up to 12 months of age (inclusive) for primary immunisation; or
 - B) Up to three doses (dependent on age at first dose) for a catch-up programme for children from 13 months to 59 months of age (inclusive) for primary immunisation, from 1 March 2023 to 31 August 2025; or
 - C) Both:
 - 1) Person is one year of age or over; and
 - 2) Any of the following:
 - i) up to two doses and a booster every five years for patients pre- and post-splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre- or post-solid organ transplant; or
 - ii) up to two doses for close contacts of meningococcal cases of any group; or
 - iii) up to two doses for person who has previously had meningococcal disease of any group; or
 - iv) up to two doses for bone marrow transplant patients; or
 - v) up to two doses for person pre- and post-immunosuppression*; or
 - D) Both:
 - 1) Person is aged between 13 and 25 years (inclusive); and
 - 2) Either:
 - Two doses for individuals who are entering within the next three months, or in their first year of living in boarding school hostels, tertiary education halls of residence, military barracks, Youth Justice residences or prisons; or
 - ii) Two doses for individuals who turn 13 years of age while living in boarding school hostels.
 - E) Contractors will be entitled to claim payment from the Funder for the supply of Meningococcal B multicomponent vaccine to patients eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Meningococcal B multicomponent vaccine listed in the Pharmaceutical Schedule.
 - F) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraphs A-D above.

*Immunosuppression due to corticosteroid or other immunosuppressive therapy must be for a period of greater than 28 days.

| Subsidy | | Fully | Brand or | _ |
|------------------------|------------|-------|--------------|---|
| (Manufacturer's Price) | Subsidised | | Generic | |
| \$ | Per | ✓ | Manufacturer | |

PNEUMOCOCCAL (PCV13) CONJUGATE VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- c)
- A) Any of the following:
 - 1) A course of three doses for previously unvaccinated children up to the age of 59 months inclusive; or
 - Two doses are funded for high risk individuals (over the age of 12 months and under 18 years) who have previously received two doses of the primary course of PCV10: or
 - 3) Up to an additional four doses (as appropriate) are funded for the (re)immunisation of high risk children aged under 5 years with any of the following:
 - a) on immunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient immune response; or
 - b) primary immune deficiencies; or
 - c) HIV infection: or
 - d) renal failure, or nephrotic syndrome; or
 - e) who are immune-suppressed following organ transplantation (including haematopoietic stem cell transplant): or
 - f) cochlear implants or intracranial shunts; or
 - g) cerebrospinal fluid leaks; or
 - h) receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage of 20 mg or greater; or
 - i) chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or
 - j) pre term infants, born before 28 weeks gestation; or
 - k) cardiac disease, with cyanosis or failure; or
 - diabetes: or
 - m) Down syndrome; or
 - n) who are pre-or post-splenectomy, or with functional asplenia; or
 - 4) Up to an additional four doses (as appropriate) are funded for the (re-)immunisation of individuals 5 years and over with HIV, pre or post haematopoietic stem cell transplantation, or chemotherapy; pre- or post splenectomy; functional asplenia, pre- or post- solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, intracranial shunts, cerebrospinal fluid leaks or primary immunodeficiency; or
 - For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.
- B) Contractors will be entitled to claim payment from the Funder for the supply of Pneumococcal (PCV13) conjugate vaccine to people eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Pneumococcal (PCV13) conjugate vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Note: please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes Inj 30.8 mcg of pneumococcal polysaccharide serotypes 1, 3, 4,

| 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and 23F in 0.5ml | | |
|---|----|---------------|
| syringe0.00 | 10 | ✓ Prevenar 13 |
| , - | 1 | ✓ Prevenar 13 |

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer PNEUMOCOCCAL (PPV23) POLYSACCHARIDE VACCINE - [Xpharm] Either: 1) Up to three doses (as appropriate) for patients with HIV, for patients post haematopoietic stem cell transplant, or chemotherapy; pre- or post-splenectomy or with functional asplenia, pre- or post-solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, or primary immunodeficiency; or 2) All of the following: a) Patient is a child under 18 years for (re-)immunisation; and b) Treatment is for a maximum of two doses; and c) Any of the following: i) on immunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient immune response; or ii) with primary immune deficiencies; or iii) with HIV infection; or iv) with renal failure, or nephrotic syndrome; or v) who are immune-suppressed following organ transplantation (including haematopoietic stem cell transplant); vi) with cochlear implants or intracranial shunts; or vii) with cerebrospinal fluid leaks; or viii) receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage of 20 mg or greater: or ix) with chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or x) pre term infants, born before 28 weeks gestation; or xi) with cardiac disease, with cyanosis or failure; or xii) with diabetes: or xiii) with Down syndrome; or xiv) who are pre-or post-splenectomy, or with functional asplenia. Inj 575 mcg in 0.5 ml prefilled syringe (25 mcg of each 1 ✓ Pneumovax 23 POLIOMYELITIS VACCINE - [Xpharm] Up to three doses for patients meeting either of the following: 1) For partially vaccinated or previously unvaccinated individuals: or 2) For revaccination following immunosuppression.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch-up programmes.

✓ IPOL

| Subsidy | | Fully | Brand or | |
|------------------------|------------|-------|--------------|--|
| (Manufacturer's Price) | Subsidised | | Generic | |
| \$ | Per | 1 | Manufacturer | |

ROTAVIRUS ORAL VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- c)
- A) Maximum of two doses for people meeting the following:
 - 1) first dose to be administered in infants aged under 14 weeks of age: and
 - 2) no vaccination being administered to children aged 24 weeks or over.
- B) Contractors will be entitled to claim payment from the Funder for the supply of Rotavirus oral vaccine to people eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Rotavirus oral vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

| Oral susp live attenuated human rotavirus 1,000,000 CCID50 per dose, squeezable tube | 0.00 | 10 | ✓ Rotarix |
|--|------|----|------------------|
| Oral susp live attenuated human rotavirus 1,000,000 CCID50 per dose, prefilled oral applicator | 0.00 | 10 | ✓ <u>Rotarix</u> |

| Subsidy (Manufacturer's Price) | | Fully bsidised | Brand or |
|-----------------------------------|-----|-------------------|-------------------------|
| (Manufacturer's Price) \$ | Per | ⊅Sidised ✓ | Generic Manufacturer |

VARICELLA VACCINE [CHICKENPOX VACCINE]

- a) Only on a prescription
- b) No patient co-payment payable
- c)
- A) Either:
 - 1) Maximum of one dose for primary vaccination for either:
 - a) Any infant born on or after 1 April 2016; or
 - For previously unvaccinated children turning 11 years old on or after 1 July 2017, who have not previously had a varicella infection (chickenpox), or
 - 2) Maximum of two doses for any of the following:
 - a) Any of the following for non-immune individuals:
 - i) with chronic liver disease who may in future be candidates for transplantation; or
 - ii) with deteriorating renal function before transplantation; or
 - iii) prior to solid organ transplant; or
 - iv) prior to any elective immunosuppression*; or
 - v) for post exposure prophylaxis who are immune competent inpatients; or
 - b) For individuals at least 2 years after bone marrow transplantation, on advice of their specialist; or
 - c) For individuals at least 6 months after completion of chemotherapy, on advice of their specialist; or
 - d) For HIV positive non immune to varicella with mild or moderate immunosuppression on advice of HIV specialist; or
 - e) For individuals with inborn errors of metabolism at risk of major metabolic decompensation, with no clinical history of varicella; or
 - f) For household contacts of paediatric patients who are immunocompromised, or undergoing a procedure leading to immune compromise where the household contact has no clinical history of varicella; or
 - g) For household contacts of adult patients who have no clinical history of varicella and who are severely immunocompromised, or undergoing a procedure leading to immune compromise where the household contact has no clinical history of varicella.
- B) Contractors will be entitled to claim payment from the Funder for the supply of Varicella vaccine [Chickenpox vaccine] vaccine to people eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Varicella vaccine [Chickenpox vaccine] listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraphs A above.

* immunosuppression due to steroid or other immunosuppressive therapy must be for a treatment period of greater than 28 days

| | NATIONAL | IIMIMUNISATI | ON SCHEDULE |
|---|---|----------------------------|-------------------------------------|
| | Subsidy (Manufacturer's Price) \$ | Fully Subsidised Per | Brand or Generic Manufacturer |
| VARICELLA ZOSTER VACCINE [SHINGLES VACCINE] | | | |
| a) Only on a prescription | | | |
| b) No patient co-payment payable | | | |
| c) | | | |
| A) Funded for patients meeting the following criteria: 1) Either: | | | |

- 1) Two doses for all people aged 65 years, or
- 2) Two doses for people 18 years of age or older with any of the following:
 - a) pre- and post-haematopoietic stem cell transplant or cellular therapy; or
 - b) pre- or post-solid organ transplant; or
 - c) haematological malignancies; or
 - d) people living with poorly controlled HIV infection; or
 - e) planned or receiving disease modifying anti-rheumatic drugs (DMARDs targeted synthetic, biologic, or conventional synthetic) for polymyalgia rheumatica, systemic lupus erythematosus or rheumatoid arthritis: or
 - f) end stage kidney disease (CKD 4 or 5); or
 - g) primary immunodeficiency
- B) Contractors will be entitled to claim payment from the Funder for the supply of Varicella zoster vaccine (Shingles vaccine) to patients eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Varicella zoster vaccine [Shingles vaccine] listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

| Inj 50 mcg per 0.5 ml vial plus vial | 0.00 1 | ✓ Shingrix |
|--------------------------------------|--------|------------|
| | 10 | ✓ Shinarix |

Diagnostic Agents

| TUBERCULIN PPD [MANTOUX] TEST - [Xpharm] | | | |
|--|------|---|------------|
| Inj 5 TU per 0.1 ml, 1 ml vial | 0.00 | 1 | ✓ Tubersol |

| - Symbols - | Albendazole | 94 | Amzoate | 28 |
|---------------------------------------|-----------------------------------|-----|-------------------------------------|-----|
| 3TC112 | Albey | 255 | Anaesthetics | |
| - A - | Albustix | 82 | Anafranil | 128 |
| A-Scabies72 | Alchemy Oxaliplatin | 154 | Anagrelide hydrochloride | 15 |
| Abacavir sulphate112 | Alchemy Oxybutynin | | Analgesics | |
| Abacavir sulphate with | Aldurazyme | | Anastrozole | |
| lamivudine112 | Alecensa | | Anatrole | |
| Abacavir/Lamivudine Viatris112 | Alectinib | | Anoro Ellipta | |
| Abilify Maintena137 | Alendronate sodium | | Antabuse | |
| Abilify Maintena S29137 | Alendronate sodium with | | Antacids and Antiflatulents | |
| Abiraterone acetate173 | colecalciferol | 117 | Anthelmintics | |
| Acarbose11 | Alfacalcidol | | Antiacne Preparations | |
| Accarb11 | Alfamino | | Antiallergy Preparations | |
| Acetazolamide267 | Alfamino Junior | | Antianaemics | |
| Acetec44 | Alginic acid | | Antiandrogen Oral | |
| Acetic acid with hydroxyquinoline and | Alglucosidase alfa | | Contraceptives | 80 |
| ricinoleic acid80 | Alkeran | | Antiarrhythmics | |
| Acetylcysteine270 | Allerfix | | Antibacterials | |
| Aci-Jel80 | Allerpro Syneo 1 | | Antibacterials Topical | |
| Aciclovir | Allerpro Syneo 2 | | Anticholinergic Agents | |
| Infection107 | Allersoothe | | Anticholinesterases | |
| Sensory | Allmercap | | Antidepressants | |
| Acidex6 | Allopurinol | | Antidiarrhoeals | |
| Acipimox | Almarytm | | Antiepilepsy Drugs | |
| Acitretin72 | Alpha-Adrenoceptor Blockers | | Antifibrinolytics, Haemostatics and | |
| Act-HIB301 | Alpha-Keri Lotion | | Local Sclerosants | |
| Actemra232 | Alphamox 125 | | Antifibrotics | |
| Actinomycin D | Alphamox 250 | | Antifungals | |
| Actrapid10 | Alprolix | | Antifungals Topical | |
| Actrapid Penfill10 | Alu-Tab | | Antihistamines | |
| Acupan124 | Aluminium hydroxide | | Antihypotensives | |
| Adalimumab (Amgevita) 184 | Alyacen | | Antimalarials | |
| Adalimumab (Humira - Alternative | Amantadine hydrochloride | | Antimigraine Preparations | |
| brand) | Ambrisentan | | Antinausea and Vertigo Agents | |
| Adapalene66 | Ambrisentan Viatris | | Antipruritic Preparations | |
| Adcetris201 | Amgevita | | Antipsychotics | |
| ADR Cartridge 1.821 | Amiloride hydrochloride | | Antiretrovirals | |
| Adrenaline | Amiloride hydrochloride with | | Antirheumatoid Agents | |
| Cardiovascular54 | furosemide | 50 | Antispasmodics and Other Agents | |
| Respiratory254 | Amiloride hydrochloride with | | Altering Gut Motility | |
| Advantan69 | hydrochlorothiazide | 50 | Antithrombotic Agents | |
| Advate38 | Aminophylline | | Antithymocyte globulin | |
| Adynovate38 | Amiodarone hydrochloride | | (equine) | 184 |
| Afinitor250 | Amisulpride | | Antitrichomonal Agents | |
| Aflibercept | Amitriptyline | | Antituberculotics and | |
| AFT-Pyrazinamide106 | Amlodipine | | Antileprotics | 10 |
| Agents Affecting the | Amorolfine | | Antiulcerants | |
| Renin-Angiotensin System 44 | Amoxicillin | | Antivirals | |
| Agents for Parkinsonism and Related | Amoxicillin with clavulanic acid. | | Anxiolytics | |
| Disorders 122 | Amoxiclav Devatis Forte | | Anzatax | |
| Agents Used in the Treatment of | Amphotericin B | | Apidra | |
| Poisonings270 | Amsacrine | | Apidra SoloStar | |
| Agrylin157 | AmsaLyo | | APO Clomipramine | |
| Albalon269 | Amsidine | | APO-Atomoxetine | 144 |

| Apo-Azithromycin | 95 | Atezolizumab | 242 | Benzydamine hydrochloride | 30 |
|---------------------------|-----|--------------------------------|-----|-------------------------------------|---------|
| APO-Candesartan HCTZ | | ATGAM | | Benzylpenicillin sodium [Penicillin | |
| 16/12.5 | 45 | Ativan | 139 | G] | |
| APO-Candesartan HCTZ | | Atnahs Olsalazine | 8 | Beta Cream | 68 |
| 32/12.5 | 45 | Atomoxetine | 144 | Beta Ointment | |
| Apo-Temozolomide | 163 | Atorvastatin | 52 | Beta Scalp | 74 |
| Apomorphine hydrochloride | 122 | Atropine sulphate | | Beta-Adrenoceptor Agonists | |
| Aprepitant | | Cardiovascular | 46 | Beta-Adrenoceptor Blockers | |
| Apresoline | | Sensory | | Beta-hCG low sensitivity urine test | |
| Aptamil Feed Thickener | | Atropt | | kit | |
| Aqueous cream | | Atrovent | | Betadine | |
| Aratac | | Aubagio | | Betadine Skin Prep | |
| Arava | | Augmentin | | Betaferon | |
| Arginine | | Aurorix | | Betahistine dihydrochloride | |
| Arginine2000 | | AutoSoft 30 | | Betaine | |
| Aripiprazole | | AutoSoft 90 | | Betamethasone dipropionate | |
| | | Avelox | | • • | |
| Aripiprazole Sandoz | | Avonex | | Betamethasone dipropionate with | 70 |
| Aristocort | | | | calcipotriol | |
| Arrotex-Prazosin S29 | | Avonex Pen | | Betamethasone sodium phosphate | |
| Arrow - Clopid | | Azacitidine | | with betamethasone acetate | |
| Arrow - Lattim | | Azacitidine Dr Reddy's | | Betamethasone valerate | , |
| Arrow-Amitriptyline | | Azamun | | Betamethasone valerate with sodiu | |
| Arrow-Bendrofluazide | | Azathioprine | | fusidate [fusidic acid] | |
| Arrow-Brimonidine | | Azilect | | Betaxolol | |
| Arrow-Diazepam | | Azithromycin | | Betnovate | |
| Arrow-Doxorubicin | 158 | Azopt | 267 | Betoptic | |
| Arrow-Fluoxetine | 129 | AZT | 112 | Betoptic S | 267 |
| Arrow-Losartan & | | - B - | | Bexsero | 308 |
| Hydrochlorothiazide | 45 | B-D Micro-Fine | 16 | Bezafibrate | 52 |
| Arrow-Norfloxacin | 115 | B-D Ultra Fine | 16 | Bezalip | |
| Arrow-Ornidazole | 105 | B-D Ultra Fine II | 16 | Bezalip Retard | 52 |
| Arrow-Quinapril 10 | 45 | Bacillus Calmette-Guerin (BCG) | | Bicalutamide | |
| Arrow-Quinapril 20 | | vaccine | 184 | Bicillin LA | 97 |
| Arrow-Quinapril 5 | | Bacillus Calmette-Guerin | | BiCNU | |
| Arrow-Roxithromycin | | vaccine | 298 | BiCNU S29 | 153 |
| Arrow-Timolol | | Baclofen | | Bile and Liver Therapy | |
| Arrow-Topiramate | | Bactroban | | Biltricide | |
| Arrow-Tramadol | | Balance | | Bimatoprost | |
| Arsenic trioxide | | Barrier Creams and Emollients | | Binarex | |
| Asacol | | BCG Vaccine AJV | | Binocrit | |
| Asacol S29 | | Beclazone 100 | | Biodone | |
| Ascend | | Beclazone 250 | | Biodone Extra Forte | |
| Ascend Aripiprazole | | Beclazone 50 | | Biodone Forte | |
| Ascend-Cefuroxime | | Beclomethasone dipropionate | | Bisacodyl | |
| Ascorbic acid | | Bedaquiline | | Bisacodyl Viatris | ر ۲۰۰۰۰ |
| | | | | Disacouyi Viatris | -2 |
| Aspen Adrenaline | 54 | Bee venom allergy treatment | | Bisoprolol fumarate | |
| Aspirin | | Bendamustine hydrochloride | | BK Lotion | |
| Blood | | Bendamustine Sandoz | | Bleomycin sulphate | 158 |
| Nervous | | Bendrofluazide | 51 | Blood Colony-stimulating | |
| Asthalin | | Bendroflumethiazide | | Factors | 42 |
| Atazanavir Mylan | 112 | [Bendrofluazide] | | Blood glucose diagnostic test | |
| Atazanavir sulphate | | Benralizumab | | meter | 15 |
| Atazanavir Viatris | | Benzathine benzylpenicillin | | Blood glucose diagnostic test | |
| Atenolol | | Benzatropine mesylate | | strip | 15 |
| Atenolol AFT | | Benzbromarone | | Blood glucose test strips (visually | |
| Atenolol Viatris | 47 | Benztrop | 122 | impaired) | 15 |

| Blood Ketone Diagnostic Test | Calcium polystyrene sulphonate43 | Cetomacrogol with glycerol7 |
|------------------------------------|--|------------------------------------|
| Strip 14 | Calcium Resonium43 | Cetomacrogol-AFT7 |
| Boostrix299 | Calogen276 | Cetuximab20 |
| Bortezomib | Camber54 | Charcoal27 |
| Bosentan57 | Candesartan cilexetil45 | CheckTop8 |
| Bosentan Dr Reddy's57 | Candesartan cilexetil with | Chemotherapeutic Agents15 |
| Bplex31 | hydrochlorothiazide45 | Chickenpox vaccine31 |
| Brentuximab Vedotin201 | Candestar45 | Chlorambucil15 |
| Breo Ellipta257 | Canesten67 | Chloramphenicol26 |
| Brevinor 1/2879 | Capecitabine155 | Chlorothiazide5 |
| Breztri Aerosphere259 | Capecitabine Viatris155 | Chlorpromazine hydrochloride13 |
| Bricanyl Turbuhaler257 | Capsaicin | Chlorsig26 |
| Brimonidine tartrate268 | Musculoskeletal117 | Chlortalidone [Chlorthalidone]5 |
| Brimonidine tartrate with timolol | Nervous124 | Chlorthalidone5 |
| maleate268 | Captopril44 | Chlorvescent4 |
| Brinzolamide267 | Carafate10 | Choice 380 7med Nsha Silver/copper |
| Brufen SR | Carbaccord | Short |
| Buccastem 134 | Carbamazepine | Ciclosporin25 |
| Budesonide | Carbimazole87 | Cidomycin P/Free9 |
| Alimentary6 | Carbomer | Cilazapril4 |
| | | Cilicaine VK9 |
| Respiratory256, 263 | Carboplatin | Cinacalcet8 |
| Budesonide Te Arai | Carboplatin Accord | |
| Budesonide with eformoterol257 | Carbosorb-X | Cinacalet Devatis8 |
| Budesonide with glycopyrronium and | Cardinol LA48 | Ciprofloxacin |
| eformoterol | Cardizem CD49 | Infection9 |
| Bumetanide50 | CareSens Dual14 | Sensory26 |
| Buprenorphine Naloxone BNM 148 | CareSens N15 | Ciprofloxacin Teva26 |
| Buprenorphine with naloxone 148 | CareSens N POP15 | Cisplatin15 |
| Bupropion hydrochloride149 | CareSens N Premier15 | Cisplatin Accord15 |
| Burel102 | CareSens PRO15 | Cisplatin Ebewe15 |
| Burinex50 | Carmellose sodium with gelatin and | Citalopram hydrobromide12 |
| Buscopan8 | pectin30 | Citrulline100029 |
| Buspirone hydrochloride139 | Carmustine 153 | Cladribine15 |
| Buspirone Viatris139 | Carnitor27 | Clarithromycin |
| Busulfan153 | Carvedilol47 | Alimentary |
| - C - | Carvedilol Sandoz47 | Infection9 |
| Cabergoline92 | Casirivimab and imdevimab202 | Clexane4 |
| Caffeine citrate264 | Catapres49 | Clexane Forte4 |
| Calamine68 | Cefaclor monohydrate94 | Clindamycin9 |
| Calci-Tab 50033 | Cefalexin94 | Clinicians25, 3 |
| Calcipotriol73 | Cefalexin Sandoz94 | Clinicians Renal Vit3 |
| Calcitonin83 | Cefazolin94 | Clobazam13 |
| Calcitriol32 | Cefazolin-AFT94 | Clobetasol propionate68, 74 |
| Calcitriol-AFT32 | Ceftriaxone94 | Clobetasone butyrate6 |
| Calcitriol-AFT S2932 | Ceftriaxone-AFT94 | Clofazimine10 |
| Calcium 500 mg Hexal33 | Cefuroxime axetil94 | Clomazol |
| Calcium carbonate | Celapram129 | Dermatological6 |
| Calcium carbonate PAI6 | Celebrex | Genito-Urinary8 |
| Calcium Channel Blockers48 | Celecoxib | Clomifene citrate9 |
| Calcium Disodium Versenate271 | Celecoxib Pfizer | Clomipramine hydrochloride12 |
| Calcium folinate | Celestone Chronodose84 | Clomipramine Teva12 |
| Calcium Folinate Ebewe | Cellcept177 | Clonazepam130, 13 |
| Calcium Folinate Ebewe | Centrally-Acting Agents49 | Clonidine4 |
| Calcium Folinate Sandoz S29 155 | | Clonidine hydrochloride4 |
| | Cephalexin ABM94 Cetirizine hydrochloride255 | |
| Calcium gluconate | | Clonidine Teva4 |
| Calcium Homeostasis83 | Cetomacrogol70 | Clopidogrel3 |

| | | | | | _ |
|----------------------------------|-----------------|---|-----|----------------------------------|-------|
| Clopine | 135 | Coumadin | 41 | DBL Bortezomib | |
| Clopixol | 136, 139 | Country Life | 27 | DBL Carboplatin | |
| Clotrimazole | | Coversyl | 45 | DBL Carboplatin S29 | 153 |
| Dermatological | 67 | COVID-19 vaccine | 298 | DBL Cisplatin | 153 |
| Genito-Urinary | 80 | Creon 10000 | | DBL Dacarbazine | 158 |
| Clozapine | 135 | Creon 25000 | 22 | DBL Desferrioxamine Mesylate for | r Inj |
| Clozaril | 135 | Creon Micro | 22 | BP | 271 |
| Clustran | 133 | Crotamiton | 68 | DBL Docetaxel | 158 |
| Co-trimoxazole | 101 | Crystaderm | 66 | DBL Ergometrine | 80 |
| Coal tar | 73 | Cu 375 Standard | 78 | DBL Gemcitabine | |
| Coal tar with allantoin, mentho | ol, | Curam | 97 | DBL Gentamicin | 99 |
| phenol and sulphur | <mark>73</mark> | Curam Duo 500/125 | 97 | DBL Leucovorin Calcium | 155 |
| Coal tar with salicylic acid and | t | Cvite | 31 | DBL Methotrexate Onco-Vial | 157 |
| sulphur | | Cyclizine hydrochloride | 134 | DBL Naloxone Hydrochloride | 270 |
| Cobal-B12 | 31 | Cyclizine lactate | 134 | DBL Pethidine Hydrochloride | 128 |
| Cobalin-H | 31 | Cyclogyl | 268 | DBL Vincristine Sulfate | 165 |
| Coco-Scalp | 73 | Cyclonex | 154 | Decozol | 30 |
| Codeine phosphate | | Cyclopentolate hydrochloride | 268 | Deferasirox | 270 |
| Extemporaneous | 273 | Cyclophosphamide | 154 | Deferiprone | 271 |
| Nervous | | Cyclorin | | Deferoxamine Pfizer S29 | |
| Coenzyme Q10 | | Cycloserine | | Denosumab | 117 |
| Colchicine | | Cyklokapron | | Deolate | 103 |
| Colecalciferol | 32 | Cyproterone acetate | | Deoxycoformycin | 162 |
| Colestyramine | 52 | Cyproterone acetate with | | Depo-Medrol | |
| Colestyramine - Mylan | 52 | ethinyloestradiol | 80 | Depo-Provera | |
| Colgout | | Cystadane | | Depo-Testosterone | 85 |
| Colifoam | | Cytarabine | | Deprim | |
| Colistin sulphomethate | 99 | Cytotec | | Dermol | |
| Colistin-Link | | Cytoxan | | Desferrioxamine mesilate | |
| Collodion flexible | | _ D - | | Desmopressin | 92 |
| Colloidal bismuth subcitrate | | D-Penamine | 117 | Desmopressin acetate | |
| Colofac | | Dabigatran | | Desmopressin-PH&T | |
| Coloxyl | <mark>23</mark> | Dacarbazine | | Detection of Substances in | |
| Combigan | | Dactinomycin [Actinomycin D] | | Urine | 82 |
| Comirnaty Omicron (JN.1) | 298 | Daivobet | | Dexamethasone | |
| Compound electrolytes | | Daivonex | | Hormone | 84 |
| Compound electrolytes with g | | Daktarin | | Sensory | |
| [Dextrose] | | Dalacin C | | Dexamethasone phosphate | |
| Compound hydroxybenzoate. | | Dantrium | | Dexamethasone with framycetin a | |
| Comtan | | Dantrium S29 | 121 | gramicidin | |
| Concerta | 147 | Dantrolene | 121 | Dexamethasone with neomycin | |
| Condoms | <mark>77</mark> | Daonil | | sulphate and polymyxin B | |
| Condyline | 75 | Dapa-Tabs | 51 | sulphate | 266 |
| Continuous glucose monitor | | Dapsone | | Dexamfetamine sulfate | 144 |
| (interoperable) | <mark>21</mark> | Daraprim | | Dexcom G6 | |
| Continuous glucose monitor | | Darunavir | 112 | Dexcom G7 | |
| (standalone) | 21 | Darunavir Viatris | 112 | Dexcom ONE+ | 21 |
| Contraceptives - Hormonal | | Dasatinib | 166 | Dexmethsone | 84 |
| Contraceptives - Non-hormon | | Dasatinib-Teva | | Dextrochlorpheniramine | |
| Copaxone | | Daunorubicin | | maleate | 255 |
| Cordarone-X | | David One Step Cassette Pregna | | Dextrose | |
| Corticosteroids and Related A | | Test | | DHC Continus | |
| for Systemic Use | | DBL Acetylcysteine | | Diabetes | |
| Corticosteroids Topical | | DBL Adrenaline | | Diabetes Management | |
| Cosentyx | | | | | |
| | 229 | DDL AMMODNYIIINE | 201 | Diacomil | 102 |
| Cosmegen | | DBL Aminophylline DBL Bleomycin Sulfate | | Diacomit Diagnostic Agents | |

| Diamide Relief | 6 | Dothiepin | 128 | Eltrombopag | 3 |
|-------------------------------------|---------|------------------------------------|------|----------------------------------|----|
| Diamox | 267 | Dovato | 113 | Eltroxin | 8 |
| Diasip | 277 | Doxazosin | 44 | EMB Fatol | 10 |
| Diazepam1 | 30, 139 | Doxazosin Clinect | 44 | Emend Tri-Pack | 13 |
| Diazoxide | | Doxine | 98 | Emicizumab | 3 |
| Dibenzyline | 44 | Doxorubicin Ebewe | 158 | EMLA | |
| Diclofenac Devatis | 266 | Doxorubicin hydrochloride | 158 | Empagliflozin | 1 |
| Diclofenac Sandoz | 116 | Doxycycline | 98 | Empagliflozin with metformin | |
| Diclofenac sodium | | DP Lotion | 71 | hydrochloride | 1 |
| Musculoskeletal | 116 | DP Lotn HC | 69 | Emsogen | |
| Sensory | 266 | DP-Captopril | 44 | Emtricitabine | 11 |
| Differin | | Dr Reddy's Omeprazole | 9 | Emtricitabine with tenofovir | |
| Difflam | 30 | Drofate | 48 | disoproxil | 10 |
| Diflucan | 101 | Drugs Affecting Bone | | Emtriva | |
| Digestives Including Enzymes | 22 | Metabolism | 117 | Emulsifying ointment | 7 |
| Digoxin | | Dual blood glucose and blood ke | tone | Emulsifying Ointment ADE | |
| Dihydrocodeine tartrate | | diagnostic test meter | | Enalapril maleate | |
| Dilantin | | Dulaglutide | | Enbrel | |
| Dilantin Infatab | | Dulcolax SP Drop | | Endocrine Therapy | |
| Dilantin Paediatric | | Duocal Super Soluble Powder | | Endoxan | |
| Diltiazem CD Clinect | | Duolin | | Energivit | |
| Diltiazem hydrochloride | | Duolin Cipla | | Engerix-B | |
| Dimethicone | | Duolin HFA | | Enhertu | |
| Dimethyl fumarate | | DuoResp Spiromax | | Enlafax XR | |
| Dipentum | | Duride | | Enoxaparin sodium | |
| Diphtheria, tetanus and pertussis | | Durvalumab | | Enstilar | |
| vaccine | | - E - | 272 | Ensure | |
| Diphtheria, tetanus, pertussis and | | e-chamber La Grande | 264 | Ensure Plus | |
| polio vaccine | | e-chamber Mask | | Ensure Plus HN | |
| Diphtheria, tetanus, pertussis, pol | | e-chamber Turbo | | Ensure Plus HN RTH | |
| hepatitis B and haemophilus | 110, | E-Mycin | | Ensure Two Cal HN RTH | |
| influenzae type B vaccine | 200 | e5 Pharma | | Entacapone | |
| • • | | | | | |
| Diprosone | | Ear Preparations | | Entapone Entecavir | |
| Diprosone OV | | Ear/Eye Preparations | | | |
| Dipyridamole | | Easiphen Liquid | | Entecavir (Rex) | |
| Disopyramide phosphate | | Econazole nitrate | | Entresto 24/26 | |
| Disulfiram | | Efavirenz | | Entresto 49/51 | |
| Diuretics | | Efavirenz Milpharm | 111 | Entresto 97/103 | |
| Docetaxel | | Efavirenz with emtricitabine and | 440 | Entyvio | |
| Docetaxel Accord | | tenofovir disoproxil | | Epilim | 13 |
| Docetaxel Sandoz | | Eformoterol fumarate dihydrate | 256 | Epilim Crushable | 13 |
| Docusate sodium | 23 | Eftrenonacog alfa [Recombinant | | Epilim IV | |
| Docusate sodium with | | factor IX] | | Epilim S/F Liquid | |
| sennosides | | Efudix | | Epilim Syrup | |
| Dolutegravir | | Egopsoryl TA | | Epipen | |
| Dolutegravir with lamivudine | | Elaprase | | Epipen Jr | 25 |
| Domperidone | | Elecare | | Epirubicin Ebewe | 15 |
| Domperidone Viatris | | Elecare LCP | | Epirubicin hydrochloride | |
| Donepezil hydrochloride | | Electral | | Eplerenone | 5 |
| Dornase alfa | | Elelyso | | Epoetin alfa | |
| Dortimopt | | Elemental 028 Extra | | Epoprostenol | 6 |
| Dorzolamide with timolol | | Elexacaftor with tezacaftor, ivaca | ftor | Eptacog alfa [Recombinant factor | |
| Dostinex | 92 | and ivacaftor | 262 | VIIa] | |
| Dosulepin [Dothiepin] | | Elidel | 73 | Erbitux | |
| hydrochloride | 128 | Elocon | | Ergometrine maleate | 8 |
| Dosulepin Viatris | | Elocon Alcohol Free | 69 | Erlotinib | 16 |
| | | | | | |

| Erythrocin IV | 96 | fraction | 38 | Fludrocortisone acetate | 84 |
|----------------------------------|-----|----------------------------------|-----|-----------------------------|----------|
| Erythromycin (as lactobionate) | 96 | Famotidine | 9 | Fluids and Electrolytes | 42 |
| Erythromycin ethyl succinate | 96 | Famotidine Hovid | 9 | Flumetasone pivalate | |
| Esbriet | | Fasenra | 200 | Fluocortolone caproate with | |
| Escitalopram | 129 | Faslodex | 174 | fluocortolone pivalate and | |
| Escitalopram (Ethics) | 129 | Fatty Cream AFT | 70 | cinchocaine | 8 |
| Eskazole | 94 | Fatty Emulsion Cream (Evara). | 70 | Fluorometholone | 266 |
| Essential Amino Acid Mix | 292 | Febuxostat | 120 | Fluorouracil | 156 |
| Essential Ethosuximide | 130 | Febuxostat (Teva) | 120 | Fluorouracil Accord | 156 |
| Estraderm MX | 86 | FEIBA NF | 38 | Fluorouracil sodium | 75 |
| Estradiol Sandoz | 86 | Felo 10 ER | 49 | Fluox | |
| Estradiol TDP Mylan | 86 | Felo 5 ER | 49 | Fluoxetine hydrochloride | 129 |
| Estradiol Viatris | 86 | Felodipine | 49 | Flupenthixol decanoate | 137 |
| Estradot | 86 | Fentanyl | 126 | Flutamide | |
| Estrofem | 86 | Fentanyl Sandoz | 126 | Flutamin | 174 |
| Estrogel | 86 | Ferinject | 33 | Fluticasone | 256 |
| Etanercept | 177 | Ferodan | | Fluticasone furoate with | |
| Ethambutol hydrochloride | 106 | Ferriprox | 271 | umeclidinium and vilanterol | 259 |
| Ethics Aspirin | 124 | Ferro-F-Tabs | | Fluticasone furoate with | |
| Ethics Aspirin EC | | Ferro-Liquid | 33 | vilanterol | 257 |
| Ethics Lisinopril | | Ferro-tab | 33 | Fluticasone propionate | 263 |
| Ethinyloestradiol with | | Ferrograd | 33 | Fluticasone with salmeterol | 25 |
| desogestrel | 78 | Ferrosig | 34 | Flynn | 94 |
| Ethinyloestradiol with | | Ferrous fumarate | | FML | |
| levonorgestrel | 79 | Ferrous fumarate with folic acid | 33 | Foban | 67 |
| Ethinyloestradiol with | | Ferrous sulfate | 33 | Folic acid | 36 |
| norethisterone | 79 | Fexaclear | 255 | Folic Acid multichem | 36 |
| Ethosuximide | 130 | Fexofenadine hydrochloride | 255 | Folic Acid Viatris | |
| Etopophos | 158 | Fibro-vein | | Food Thickeners | 286 |
| Etoposide | | Filgrastim | 42 | Foods And Supplements For | |
| Etoposide phosphate | | Finasteride | 81 | Inherited Metabolic Disease | 287 |
| Etravirine | | Fingolimod | 140 | Fortini | 279 |
| Eumovate | 69 | Firazyr | | Fortini Multi Fibre | 279 |
| Eurofolic | 155 | Flagyl | 105 | Fortisip | 284 |
| Evara | 70 | Flagyl-S | | Fortisip Multi Fibre | |
| EVARA White Soft Paraffin | 71 | Flamazine | | Fosamax | 117 |
| Everet | 131 | Flecainide acetate | 47 | Fosamax Plus | 117 |
| Everolimus | 250 | Flecainide BNM | 47 | Fosfomycin | 118 |
| Evista | 119 | Flecainide Controlled Release | | Framycetin sulphate | 265 |
| Evrysdi | 144 | Teva | 47 | Freestyle Libre 2 | 21 |
| Evusheld | | Fleet Phosphate Enema | 24 | Freestyle Libre 3 Plus | 21 |
| Exelon Patch 10 | 148 | Flixonase Hayfever & Allergy | 263 | Frisium | |
| Exelon Patch 5 | 148 | Flixotide | | Frumil | 50 |
| Exemestane | 177 | Flixotide Accuhaler | 256 | Frusemide | 50 |
| Exjade | 270 | Florinef | 84 | Fucicort | 69 |
| Extemporaneously Compounded | l | Fluanxol | 137 | Fucidin | 100 |
| Preparations and | | Flucil | 97 | Fucithalmic | 265 |
| Galenicals | 273 | Flucloxacillin | 97 | Fucithalmic S29 | 265 |
| Eye Preparations | 265 | Flucloxacillin-AFT | 97 | Fulvestrant | |
| Eylea | | Flucloxin | | Fungilin | |
| Ezemibe Viatris | | Flucon | | Furosemide [Frusemide] | |
| Ezetimibe | | Fluconazole | | Furosemide-Baxter | |
| Ezetimibe Sandoz | | Fludara Oral | | fusidic acid | |
| Ezetimibe with simvastatin | | Fludarabine Ebewe | | Dermatological | . 67, 69 |
| -F- | | Fludarabine phosphate | | Infection | |
| Factor eight inhibitor bypassing | | Fludarabine Sagent | | Sensory | |
| | | | | | |

| - G - | Haldol | 137 | Hydralazine | 54 |
|---------------------------------|--------------------------------|-------------------|------------------------------------|----------------|
| GA Explore 52 | 91 Haldol Concentrate | 137 | Hydralazine hydrochloride | 54 |
| GA1 Anamix Infant2 | | | Hydralyte - Lemonade | 43 |
| GA1 Anamix Junior2 | 91 Haloperidol | 135 | Hydrocortisone | |
| Gabapentin1 | 30 Haloperidol decanoate | 137 | Dermatological | 69 |
| Gacet125-1 | 26 Harvoni | 109 | Hormone | 84 |
| Galsulfase | 26 Havrix 1440 | 301 | Hydrocortisone acetate | <mark>7</mark> |
| Galvumet | 12 Havrix Junior | 301 | Hydrocortisone acetate with | |
| Galvus | 12 Haylor syrup | 2 <u>55</u> | pramoxine hydrochloride | <mark>7</mark> |
| Gardasil 93 | 03 HCU Anamix Infant | 288 | Hydrocortisone and paraffin liquid | |
| Gastrodenol | 10 HCU Anamix Junior | 288 | and lanolin | 69 |
| Gaviscon Extra Strength | .6 HCU Anamix Junior LQ | 288 | Hydrocortisone butyrate | 69, 74 |
| Gaviscon Infant | .6 HCU Explore 5 | 288 | Hydrocortisone with cinchocaine | 8 |
| Gazyva2 | 13 HCU Express 15 | 288 | Hydrocortisone with miconazole | 69 |
| Gefitinib1 | | | Hydrocortisone with natamycin an | d |
| GEM Aqueous Cream | | | neomycin | 69 |
| Gemcitabine Ebewe1 | 56 Free | 70 | Hydrogen peroxide | 66 |
| Gemcitabine hydrochloride1 | | 68 | Hydroxocobalamin | |
| Gemtuzumab ozogamicin2 | 03 healthE Dimethicone 10% | 70 | Hydroxocobalamin Panpharma | 31 |
| Gentamicin Noridem | 99 healthE Dimethicone 4% Loti | on71 | hydroxycarbamide | 158 |
| Gentamicin sulphate | 99 healthE Dimethicone 5% | 70 | Hydroxychloroquine sulphate | |
| Gilenya1 | 40 healthE Glycerol BP | 273 | Hydroxyurea | |
| Ginet | | | [hydroxycarbamide] | 158 |
| Glatiramer acetate1 | 40 Healtheries Simple Baking M | ix286 | Hygroton | 51 |
| Glecaprevir with pibrentasvir1 | | | Hylo-Fresh | 269 |
| Glibenclamide | 11 Hemlibra | 37 | Hymenoptera | 255 |
| Gliclazide | 11 Heparin sodium | 41 | Hyoscine butylbromide | |
| Glipizide | 12 Heparin Sodium Panpharma | 41 | Hyoscine Butylbromide | |
| Glizide | 11 Heparinised saline | 41 | (Adiramedica) | 8 |
| Glucagen Hypokit | 10 Heparon Junior | 278 | Hyoscine hydrobromide | |
| Glucagon hydrochloride | 10 Hepatitis A vaccine | 301 | Hyperuricaemia and Antigout | 120 |
| Glucerna Select2 | | | Hypromellose | |
| Glucose [Dextrose] | 42 vaccine | 302 | Hypromellose with dextran | |
| Gluten Free Foods2 | | 2 <mark>35</mark> | -1- | |
| Glycerin with sodium saccharin2 | | 46 | Ibiamox | 97 |
| Glycerin with sucrose2 | | 115 | Ibrance | 170 |
| Glycerol | Histaclear | | Ibrutinib | 159 |
| Alimentary | | | Ibuprofen | |
| Extemporaneous2 | | | Ibuprofen SR BNM | |
| Glyceryl trinitrate | Horleys Flour | | lcatibant | |
| Alimentary | | | Idarubicin hydrochloride | |
| Cardiovascular | | | Idursulfase | |
| Glycopyrronium2 | | | Ifosfamide | |
| Glycopyrronium bromide | | 11 | llevro | 267 |
| Glycopyrronium with | Humalog Mix 25 | | lloprost | |
| indacaterol2 | | | Imatinib mesilate | |
| Glycosade2 | | | Imatinib-Rex | |
| Glytactin Bettermilk2 | | | Imbruvica | 159 |
| Gold Knight | | | Imfinzi | |
| Gold Knight XL | | | Imipramine Crescent | |
| Goserelin | | | Imipramine hydrochloride | |
| Gynaecological Anti-infectives | | | Imiquimod | |
| - H - | Humulin 30/70 | | Immune Modulators | |
| Habitrol1 | | | Immunisation - Flu | |
| Haemophilus influenzae type B | Humulin R | | Immunisation Flu and Shingles | |
| vaccine3 | | | Immunisation Other | |
| | | | | |

| | | | 17.1.1 | |
|------------------------------------|---------------------------------|----------|-------------------------------|-------|
| Immunosuppressants 177 | Intra-uterine device | | Kalydeco | |
| Incruse Ellipta | Invega Sustenna | | Kemadrin | |
| Indacaterol256 | Invega Trinza | | Kenacomb | |
| Indapamide51 | Ipca-Allopurinol | | Kenacort-A 10 | |
| Infanrix IPV300 | Ipca-Bisoprolol | | Kenacort-A 40 | |
| Infanrix-hexa300 | Ipca-Ciprofloxacin | | Kenalog in Orabase | 30 |
| Infant Formulae292 | Ipca-Donepezil | 148 | Ketocal 3:1 | |
| Infatrini296 | Ipca-Escitalopram | 129 | KetoCal 4:1 | 29 |
| Infliximab203 | IPCA-Frusemide | 50 | Ketoconazole | |
| Influenza vaccine304 | Ipca-Hydroxychloroquine | 117 | Dermatological | 74 |
| Influvac Tetra | IPCA-Metoprolol | 48 | Infection | 10 |
| (2024 formulation) 304 | IPCA-Propranolol | 48 | Ketogenic Diet | 29 |
| Inhaled Corticosteroids256 | IPOL | | Ketoprofen | |
| Inhaled Long-acting | Ipratropium bromide | 258. 263 | KetoSens | 14 |
| Beta-adrenoceptor Agonists 256 | Iressa | | Keytruda | |
| Inresa34 | Irinotecan Actavis 100 | | Kindergen | |
| Inspra50 | Irinotecan hydrochloride | | Kisqali | |
| Instillagel Lido123 | Irinotecan-Rex | | Klacid | |
| Insulin aspart11 | Iron (as ferric carboxymaltose) | | Alimentary | |
| Insulin aspart with insulin aspart | Iron polymaltose | | Infection | |
| | Isentress | | Kliogest | |
| protamine | | | Kliovance | |
| Insulin glargine | Isentress HD | | | |
| Insulin glulisine | Ismo 20 | | Kogenate FS | 30 |
| Insulin isophane11 | Ismo 40 Retard | | Konakion MM | |
| Insulin isophane with insulin | Isoleucine50 | | Konsyl-D | |
| neutral11 | Isoniazid | | Kuvan | 2 |
| Insulin lispro11 | Isoniazid Teva | | -L- | |
| Insulin lispro with insulin lispro | Isoniazid with rifampicin | | Labetalol | |
| protamine11 | Isoptin | | Lacosamide | |
| Insulin neutral10 | Isoptin Retard | | Lactulose | |
| Insulin pen needles16 | Isoptin SR | | Laevolac | |
| Insulin pump cartridge17 | Isopto Carpine | | Lagevrio | 110 |
| Insulin pump infusion set (steel | Isosorbide mononitrate | 54 | Lamictal | 13 |
| cannula)18 | Isosource Standard | 284 | Lamivudine10 | 7, 11 |
| Insulin pump infusion set (steel | Isotretinoin | 66 | Lamivudine Viatris | 112 |
| cannula, straight insertion) 18 | Ispaghula (psyllium) husk | 23 | Lamivudine/Zidovudine Viatris | 112 |
| Insulin pump infusion set (teflon | Itch-Soothe | 68 | Lamotrigine | 13 |
| cannula)19 | Itracap | 102 | Lamprene | |
| Insulin pump infusion set (teflon | Itraconazole | | Lanoxin | |
| cannula, angle insertion with | Itraconazole Kent | | Lanoxin Paediatric Elixir | |
| insertion device)20 | Itrazole | | Lanoxin PG | |
| Insulin pump infusion set (teflon | lvacaftor | | Lanoxin S29 | |
| cannula, flexible insertion with | lvermectin | | Lansoprazole | |
| insertion device)20 | - J - | | Lantus | |
| Insulin pump infusion set (teflon | Jadelle | 70 | Lantus SoloStar | |
| cannula, straight insertion with | Jakavi | | Lanvis | |
| insertion device)20 | Jardiamet | | Lanzol Relief | |
| | | | | |
| Insulin pump infusion set (teflon | Jardiance | | Largactil | |
| cannula, variable insertion)20 | Jaydess | | Laronidase | |
| Insulin pump reservoir | Jevity HiCal RTH | | Lasix | |
| Insulin pump with algorithm16 | Jevity Plus RTH | | Latanoprost | |
| Insulin syringes, disposable with | Jevity RTH | | Latanoprost with timolol | 26 |
| attached needle16 | Jinarc | | Lax-Suppositories | 24 |
| Intelence111 | Juno Pemetrexed | 157 | Lax-suppositories Glycerol | |
| Interferon beta-1-alpha140 | - K - | | Laxatives | |
| Interferon beta-1-beta140 | Kadcyla | 238 | Laxsol | 2 |

| Ledipasvir with sofosbuvir | 109 | Locorten-Vioform | 265 | Medac | 15 |
|--|---|---|---|--|--|
| Leflunomide | 117 | Lodoxamide | | Medrol | |
| Lenalidomide (Viatris) | 159 | Logem | 131 | Medroxyprogesterone acetate | |
| Lenalidomide Viatris | 159 | Lomide | | Genito-Urinary | 7 |
| Lenvatinib | | Lomustine | | Hormone | 8 |
| Lenvima | 167 | Loniten | 54 | Mefenamic acid | |
| Letrole | 177 | Loperamide hydrochloride | 6 | Megval | 15 |
| Letrozole | 177 | Lopinavir with ritonavir | 112 | Melatonin | |
| Leucine100 | 292 | Lopinavir/Ritonavir Mylan | | Melpha | 15 |
| Leukeran FC | 153 | Loprofin | | Melphalan | 15 |
| Leukotriene Receptor | | Loprofin Mix | | Meningococcal (groups A, C, Y an | |
| Antagonists | 261 | Lorafix | 255 | W-135) conjugate vaccine | |
| Leuprorelin | | Loratadine | 255 | Meningococcal B multicomponent | |
| Leustatin | | Lorazepam | 139 | vaccine | 30 |
| Levetiracetam | 131 | Lorstat | | MenQuadfi | 30 |
| Levetiracetam-AFT | | Losartan Actavis | | Menthol | 6 |
| Levocabastine | 267 | Losartan potassium | | Mepolizumab | |
| Levocarnitine | | Losartan potassium with | | Mercaptopurine | |
| Levodopa with benserazide | | hydrochlorothiazide | 45 | Mercilon 28 | |
| Levodopa with carbidopa | | Lovir | | Mesalazine | |
| Levodopa with carbidopa and | | Loxamine | | Mesna | |
| entacapone | 122 | Lucrin Depot 1-month | | Mestinon | |
| Levomepromazine | | Lucrin Depot 3-month | | Metabolic Disorder Agents | |
| Levomepromazine | | LumaCina | | Metabolics | |
| hydrochloride | 135 | Lumigan | | Metformin hydrochloride | |
| Levonorgestrel | | Lyllana | | Metformin Viatris | |
| Genito-Urinary | 79-80 | Lynparza | | Methadone BNM | |
| Hormone | | Lyrica | | Methadone hydrochloride | |
| | | | | | |
| Levonorgestrel BNM | 80 | - M - | | Methenamine (hexamine) | |
| Levonorgestrel BNM | | - M - m-Eslon | 127 | Methenamine (hexamine) hippurate | 11 |
| LevothyroxineLidocaine [Lignocaine] | 87 | ••• | | hippurate | |
| LevothyroxineLidocaine [Lignocaine] | 87 | m-Eslon | 217 | | 26 |
| LevothyroxineLidocaine [Lignocaine]Lidocaine [Lignocaine] | 87 123–124 | m-Eslon Mabthera Macrobid | 217 115 | hippurate Methopt Methotrexate | 26 15 |
| Levothyroxine | 87 123–124 | m-Eslon | 217 115 m | hippurate Methopt Methotrexate Methotrexate DBL Onco-Vial | 26 15 15 |
| Levothyroxine | 87 123–124 124 | m-Eslon | 217 115 m ate and | hippurate Methopt Methotrexate | 26 15 15 |
| Levothyroxine | 87 123–124 124 | m-Eslon Mabthera Macrobid Macrogol 3350 with potassiur chloride, sodium bicarbona sodium chloride | 217 115 m ate and 24 | hippurate Methopt Methotrexate Methotrexate DBL Onco-Vial Methotrexate DBL S29 | 26 15 15 15 |
| Levothyroxine | 87 123–124 124 124 | m-Eslon | 217115 m ate and2424 | hippurate | 26 15 15 15 |
| Levothyroxine | 87 123–124 124 124 124 29 | m-Eslon | 217115 m ate and2424122 | hippurate | 26 15 15 15 15 |
| Levothyroxine | 87 123–124 124 124 124 29 123–124 | m-Eslon | 217115 m ate and24122122 | hippurate | 26 15 15 15 15 |
| Levothyroxine | 87 123–124 124 124 124 29 123–124 106 | m-Eslon | 217115 m ate and24122122122 | hippurate | 26 15 15 15 15 27 |
| Levothyroxine | 87 123–124 124 124 124 29 123–124 106 121 | m-Eslon | 217115 m ate and24122122122122122 | hippurate | 26 15 15 15 15 27 |
| Levothyroxine Lidocaine [Lignocaine] Lidocaine [Lignocaine] hydrochloride Lidocaine [Lignocaine] with prilocaine Lidocaine-Baxter | 87 123–12412412429 123–124106121 | m-Eslon | 217115 m ate and2412212212212212234 | hippurate | 26 15 15 15 15 27 27 |
| Levothyroxine | 87 123–12412412429 123–12410612152276 | m-Eslon | 217115 m ate and241221221221221223434 | hippurate | 26 15 15 15 15 27 27 |
| Levothyroxine Lidocaine [Lignocaine] Lidocaine [Lignocaine] hydrochloride Lidocaine [Lignocaine] with prilocaine Lidocaine-Baxter | 87 123–12412412412412412412412412529 123–1241061215227612 | m-Eslon | 217115 m ate and2412212212212212234313 | hippurate | 26 15 15 15 15 27 27 27 |
| Levothyroxine Lidocaine [Lignocaine] Lidocaine [Lignocaine] hydrochloride Lidocaine [Lignocaine] with prilocaine Lidocaine-Baxter Life Extension Lignocaine Linezolid Lioresal Intrathecal Lipid-Modifying Agents Liquigen Liraglutide Lisdexamfetamine dimesilate | 87 123–124124124124291291201215227612145 | m-Eslon | 217115 m ate and241221221221221243431347 | hippurate | 26 15 15 15 15 27 27 27 |
| Levothyroxine Lidocaine [Lignocaine] Lidocaine [Lignocaine] hydrochloride Lidocaine [Lignocaine] with prilocaine Lidocaine-Baxter Life Extension Lignocaine Linezolid Lioresal Intrathecal Lipid-Modifying Agents Liquigen Liraglutide | 87 123–1241241241241241241261271261215212125145 | m-Eslon Mabthera Macrobid Macrogol 3350 with potassiur chloride, sodium bicarbona sodium chloride Madopar 125 Madopar 250 Madopar 62.5 Madopar HBS Madopar Rapid Magnesium hydroxide Magnesium sulphate Mantoux MAR-Midodrine Marevan | 217115 m ste and241221221221221223431347 | hippurate | 26 |
| Levothyroxine Lidocaine [Lignocaine] Lidocaine [Lignocaine] hydrochloride Lidocaine [Lignocaine] with prilocaine Lidocaine-Baxter Life Extension Lignocaine Linezolid Lioresal Intrathecal Lipid-Modifying Agents Liquigen Liraglutide Lisdexamfetamine dimesilate Lisinopril | 87 123–12412412412429 123–1241061215227612145 | m-Eslon Mabthera Macrobid | 217115 m ate and2412212212212234313474174 | hippurate | 26 15 15 15 27 27 27 27 27 27 27 |
| Levothyroxine Lidocaine [Lignocaine] Lidocaine [Lignocaine] hydrochloride Lidocaine [Lignocaine] with prilocaine Lidocaine-Baxter Life Extension Lignocaine Linezolid Lioresal Intrathecal Lipid-Modifying Agents Liquigen Liraglutide Lisdexamfetamine dimesilate Lisinopril Lithium carbonate | 87 123–12412412412429 123–124106121522761214544135267 | m-Eslon Mabthera Macrobid | 217115 m ste and2412212212212234313474174 | hippurate | 26 15 15 15 27 27 27 27 27 27 27 |
| Levothyroxine Lidocaine [Lignocaine] hydrochloride Lidocaine [Lignocaine] with prilocaine Lidocaine-Baxter Life Extension Lignocaine Linezolid Lioresal Intrathecal Lipid-Modifying Agents Liquigen Liraglutide Lisdexamfetamine dimesilate Lisinopril Lithium carbonate Livostin | 87 123–12412412429 123–124106121522761214544135267124 | m-Eslon Mabthera Macrobid Macrogol 3350 with potassiur chloride, sodium bicarbona sodium chloride Madopar 125 Madopar 250 Madopar 62.5 Madopar HBS Madopar Rapid Magnesium hydroxide Magnesium sulphate Mantoux MAR-Midodrine Marevan Marine Blue Lotion SPF 50+ Martindale Pharma | 217115 m site and24122122122122124314313474174270263 | hippurate | 26 15 15 15 15 15 27 27 27 27 21 24 44 14 |
| Levothyroxine Lidocaine [Lignocaine] hydrochloride Lidocaine [Lignocaine] with prilocaine Lidocaine-Baxter Life Extension Lignocaine Linezolid Lioresal Intrathecal Lipid-Modifying Agents Liquigen Liraglutide Lisdexamfetamine dimesilate Lisinopril Lithium carbonate Livostin LMX4 | 87 123–12412412429 123–1241061215227612135267124 | m-Eslon Mabthera Macrobid Macrogol 3350 with potassiur chloride, sodium bicarbona sodium chloride Madopar 125 Madopar 250 Madopar 62.5 Madopar HBS Madopar Rapid Magnesium hydroxide Magnesium sulphate Mantoux MAR-Midodrine Marevan Marine Blue Lotion SPF 50+ Martindale Pharma Mask for spacer device | 217115 m ste and2412212212212212234313474174270263109 | hippurate | 2615151515 |
| Levothyroxine Lidocaine [Lignocaine] hydrochloride Lidocaine [Lignocaine] hydrochloride Lidocaine [Lignocaine] with prilocaine Lidocaine-Baxter. Life Extension Lignocaine Linezolid Lioresal Intrathecal Lipid-Modifying Agents Liquigen Liraglutide Lisdexamfetamine dimesilate Lisinopril Lithium carbonate Livostin LMX4 Lo-Oralcon 20 ED Locacorten-Viaform ED's. | 87 123–12412412429 123–1241061215227612135267124 | m-Eslon Mabthera Macrobid | 217115 m ste and24122122122122122 | hippurate | 2615151515 |
| Levothyroxine Lidocaine [Lignocaine] hydrochloride Lidocaine [Lignocaine] with prilocaine Lidocaine-Baxter Life Extension Lignocaine Linezolid Lioresal Intrathecal Liquigen Liraglutide Lisdexamfetamine dimesilate Lisinopril Lithium carbonate Livostin LMX4 Lo-Oralcon 20 ED | 87 123–12412412412429 123–1241061215227612145135267124 | m-Eslon Mabthera Macrobid | | hippurate | |
| Levothyroxine Lidocaine [Lignocaine] hydrochloride Lidocaine [Lignocaine] hydrochloride Lidocaine [Lignocaine] with prilocaine Lidocaine-Baxter Lidocaine-Baxter Life Extension Lignocaine Linezolid Lioresal Intrathecal Lipid-Modifying Agents Liquigen Liraglutide Lisdexamfetamine dimesilate Lisinopril Lithium carbonate Livostin LMX4 Lo-Oralcon 20 ED Locacorten-Viaform ED's Local preparations for Anal and | 87 123–12412412412429 123–124106121522761214544135267267265 | m-Eslon | | hippurate | |
| Levothyroxine Lidocaine [Lignocaine] hydrochloride Lidocaine [Lignocaine] with prilocaine Lidocaine [Lignocaine] with prilocaine Lidocaine-Baxter Life Extension Lignocaine Linezolid Lioresal Intrathecal Lipid-Modifying Agents Liquigen Liraglutide Lisdexamfetamine dimesilate Lisinopril Lithium carbonate Livostin LMX4 Lo-Oralcon 20 ED Locacorten-Viaform ED's Local preparations for Anal and Rectal Disorders | 87 123–12412412412429 123–12410612152276121454413526712479265 | m-Eslon Mabthera Macrobid | | hippurate | |
| Levothyroxine Lidocaine [Lignocaine] Lidocaine [Lignocaine] hydrochloride Lidocaine [Lignocaine] with prilocaine Lidocaine-Baxter Life Extension Lignocaine Linezolid Lioresal Intrathecal Lipid-Modifying Agents Liquigen Liraglutide Lisdexamfetamine dimesilate Lisinopril Lithium carbonate Livostin LMX4 Lo-Oralcon 20 ED Locacorten-Viaform ED's Local preparations for Anal and Rectal Disorders Locaid Locoid Locoid | 87 123–12412412412429 123–124106121522761254413526712479265 | m-Eslon | | hippurate | |
| Levothyroxine Lidocaine [Lignocaine] Lidocaine [Lignocaine] hydrochloride Lidocaine [Lignocaine] with prilocaine Lidocaine-Baxter Life Extension Lignocaine Linezolid Lioresal Intrathecal Lipid-Modifying Agents Liquigen Liraglutide Lisdexamfetamine dimesilate Lisinopril Lithium carbonate Livostin LMX4 Lo-Oralcon 20 ED Locacorten-Viaform ED's Local preparations for Anal and Rectal Disorders Locasol Locoid | 87 123–12412412412429 123–124106121522761254413526712479265 | m-Eslon | | hippurate | |

| Metoclopramide hydrochloride . | 134 | Minims Pilocarpine | 268 | Multivitamins | 3 |
|--------------------------------|-----------------|-------------------------------|------------------|--------------------------------|-----|
| Metolazone | 51 | Minims Prednisolone | | Mupirocin | 6 |
| Metopirone | 93 | Minipress | 44 | Muscle Relaxants | 12 |
| Metoprolol IV Mylan | 48 | Minirin | 92 | Mvite | 3 |
| Metoprolol IV Viatris | 48 | Minirin Melt | 92 | Myambutol | 10 |
| Metoprolol succinate | | Mino-tabs | 98 | Mycobutin | 10 |
| Metoprolol tartrate | 48 | Minocycline hydrochloride | 98 | MycoNail | |
| Metrogyl | 105 | Minomycin | 98 | Mycophenolate mofetil | 17 |
| Metronidamed | | Minor Skin Infections | | Mydriacyl | 26 |
| Metronidazole | 105 | Minoxidil | 54 | Mylan (24 hr release) | |
| Metyrapone | 93 | Minoxidil Roma | 54 | Mylan Clomiphen | 9 |
| Mexiletine hydrochloride | 47 | Mirena | 87 | Mylan Italy (24 hr release) | 4 |
| Miacalcic | | Miro-Amoxicillin | 97 | Myleran | 15 |
| Micolette | <mark>24</mark> | Mirtazapine | 129 | mylife Inset soft | |
| Miconazole | 30 | Misoprostol | 9 | mylife Orbit micro | 18 |
| Miconazole nitrate | | Mitomycin (Fresenius Kabi) | 160 | mylife YpsoPump Reservoir | 2 |
| Dermatological | 68 | Mitomycin (Sagent) | 160 | mylife YpsoPump with CamAPS | |
| Genito-Urinary | 80 | Mitomycin C | 160 | FX | |
| Micreme | | Mitozantrone | 160 | Myloc CR | 4 |
| Micreme H | 69 | Mitozantrone Ebewe | 160 | Mylotarg | |
| Microgynon 30 | 79 | Mixtard 30 | 11 | Myometrial and Vaginal Hormone | Э |
| Microlut | 79 | MMA/PA Anamix Infant | 291 | Preparations | 8 |
| Midazolam | 143 | MMA/PA Anamix Junior | 291 | Myozyme | 2 |
| Midazolam Viatris | 143 | MMA/PA Explore 5 | 291 | - N - | |
| Midazolam-Baxter | 143 | MMA/PA Express 15 | 291 | Nadolol | 4 |
| Midazolam-Pfizer | 143 | Moclobemide | 129 | Nadolol BNM | |
| Midodrine | 47 | Modafinil | 147 | Naglazyme | |
| Midostaurin | | Modafinil Max Health | 147 | Naloxone hydrochloride | |
| Mifegyne | 82 | Modavigil | 147 | Naltraccord | 149 |
| Mifepristone | 82 | Moduretic | 50 | Naltrexone AOP | |
| Milpharm | 131 | Molaxole | 24 | Naltrexone hydrochloride | 149 |
| Minerals | | Molnupiravir | 110 | Naltrexone Max Health | |
| Mini-Wright AFS Low Range | 263 | Moments | 77 | Naphazoline hydrochloride | 26 |
| Mini-Wright Standard | 263 | Mometasone furoate | 69 | Naprosyn SR 1000 | |
| Minidiab | 12 | Monogen | <mark>277</mark> | Naprosyn SR 750 | 110 |
| MiniMed 3.0 Reservoir | | Montelukast | 261 | Naproxen | 110 |
| MMT-332A | 21 | Montelukast Viatris | 261 | Narcaricin mite | |
| MiniMed Mio MMT-921A | 19 | Moroctocog alfa [Recombinan | t factor | Nasal Preparations | 26 |
| MiniMed Mio MMT-923A | 19 | VIII] | 38 | Natalizumab | 140 |
| MiniMed Mio MMT-925A | 19 | Morphine hydrochloride | | Natulan | |
| MiniMed Mio MMT-941A | 19 | Morphine sulphate | 127 | Nausafix | |
| MiniMed Mio MMT-943A | 19 | Motetis | 123 | Nausicalm | |
| MiniMed Mio MMT-945A | 19 | Mouth and Throat | 30 | Navelbine S29 | |
| MiniMed Mio MMT-965A | 19 | Movapo | 122 | Nefopam hydrochloride | 124 |
| MiniMed Mio MMT-975A | 19 | Moxifloxacin | 99 | Neo-Cytamen S29 | |
| MiniMed Quick-Set MMT-396A | | MSUD Anamix Infant | 288 | Neo-Mercazole | |
| MiniMed Quick-Set MMT-397A | | MSUD Anamix Junior | | Neocate Gold | |
| MiniMed Quick-Set MMT-398A | 19 | MSUD Anamix Junior LQ | 288 | Neocate Junior Unflavoured | 29 |
| MiniMed Quick-Set MMT-399A | 19 | MSUD Explore 5 | 288 | Neocate Junior Vanilla | |
| MiniMed Silhouette MMT-377A | 19 | MSUD Express 15 | | Neocate SYNEO | |
| MiniMed Silhouette MMT-378A | 19 | MSUD Lophlex LQ 20 | 288 | Neoral | |
| MiniMed Silhouette MMT-381A | 19 | MSUD Maxamum | | Neostigmine metilsulfate | |
| MiniMed Sure-T MMT-864A | 18 | Mucolytics | | Nepafenac | |
| MiniMed Sure-T MMT-866A | | Mucosoothe | | Nepro HP (strawberry) | |
| MiniMed Sure-T MMT-874A | 18 | Multiple Sclerosis Treatments | 139 | Nepro HP (vanilla) | |
| MiniMed Sure-T MMT-876A | 18 | Multivitamin renal | 32 | Neulactil | |

| NeuroTabs | 33 | NovoSeven RT | 37 | Olbetam | 52 |
|-----------------------------|-----|--------------------------------|-----|------------------------------|-----|
| Nevirapine | 111 | Nozinan | 135 | Olopatadine | 269 |
| Nevirapine Viatris | 111 | Nozinan (Swiss) | 135 | Olopatadine Teva | 269 |
| Nicorandil | 54 | Nozinan S29 | 135 | Olsalazine | 8 |
| Nicotine | | Nucala | | Omalizumab | |
| Nifedipine | 49 | Nuelin | 261 | Omeprazole | |
| Nifuran | 115 | Nuelin-SR | 261 | Omeprazole actavis 10 | |
| Nilotinib | 168 | Nupentin | | Omeprazole actavis 20 | |
| Nilstat | | Nusinersen | | Omeprazole actavis 40 | |
| Alimentary | 30 | Nutilis | 286 | Omeprazole Teva | |
| Genito-Urinary | 80 | Nutren Diabetes | 277 | Omnitrope | 8 |
| Infection | 102 | Nutrient Modules | 274 | Onbrez Breezhaler | |
| Nimenrix | 307 | Nutrini Energy Multi Fibre | 279 | Oncaspar LYO | 162 |
| Nintedanib | 260 | Nutrini Energy RTH | | OncoTICE | |
| Nipent | 162 | Nutrini Low Energy Multi Fibre | | Ondansetron | |
| Niraparib | | Nutrini Peptisorb | | One-Alpha | 3 |
| Nirmatrelvir with ritonavir | | Nutrini Peptisorb Energy | | One-Alpha S29 | |
| Nitrates | | Nutrini RTH | | Opdivo | 243 |
| Nitroderm TTS | | Nutrison 800 Complete Multi | | Ora-Blend | 273 |
| Nitrofurantoin | | Fibre | 284 | Ora-Blend SF | |
| Nitrolingual Pump Spray | | Nutrison Advanced Peptisorb | | Ora-Plus | |
| Nivestim | | Nutrison Concentrated | 286 | Ora-Sweet | |
| Nivolumab | | Nutrison Energy | | Ora-Sweet SF | |
| Nodia | | Nutrison Energy Multi Fibre | | Orabase | 30 |
| Noflam 250 | | Nutrison Multi Fibre | | Oral and Enteral Feeds | |
| Noflam 500 | | Nutrison RTH | | Oralcon 30 ED | |
| Non-Steroidal Anti-Inflamma | | Nyefax Retard | | Oramorph | |
| Drugs | | Nystatin | | Oramorph CDC S29 | 12 |
| Nonacog gamma, [Recombin | | Alimentary | 30 | Oratane | |
| Factor IX] | | Genito-Urinary | | Orgran | |
| Norethinderone - CDC | | Infection | | Ornidazole | |
| Norethisterone | | NZB Low Gluten Bread Mix | | Orphenadrine citrate | 12 |
| Genito-Urinary | 80 | -0- | | Ortho-tolidine | |
| Hormone | | Obinutuzumab | 213 | Oruvail SR | |
| Norflex | | Obstetric Preparations | | Osimertinib | |
| Norfloxacin | | Ocicure | | Osmolite RTH | |
| Noriday | | Ocrelizumab | | Other Endocrine Agents | |
| Noriday 28 | 80 | Ocrevus | | Other Oestrogen Preparations | |
| Norimin | | Octocog alfa [Recombinant fact | | Other Progestogen | |
| Normison | | VIII] (Advate) | | Preparations | 8 |
| Norpress | | Octocog alfa [Recombinant fact | | Other Skin Preparations | |
| Nortriptyline hydrochloride | | VIII] (Kogenate FS) | | Otodex | |
| Norvir | | Octreotide | | Ovestin | |
| Noumed Dexamfetamine | | Octreotide GH | | Genito-Urinary | 80 |
| Noumed Isoniazid | | Octreotide long-acting | | Hormone | |
| Noumed Paracetamol | | Oestradiol | | Oxaliplatin | |
| Noumed Pethidine | | Oestradiol valerate | | Oxaliplatin Accord | |
| Noumed Phenobarbitone | | Oestradiol with norethisterone | | Oxaliplatin Actavis 100 | 154 |
| Novadoz | | Oestriol | | Oxaliplatin Ebewe | |
| NovaSource Renal | | Genito-Urinary | 80 | Oxis Turbuhaler | |
| Novatretin | | Hormone | 87 | Oxpentifylline | |
| Novitium Sugar Free | | Oestrogens | | Oxybutynin | |
| NovoMix 30 FlexPen | 10 | Ofev | | Oxycodone Amneal | |
| NovoRapid | | Oil in water emulsion | | Oxycodone hydrochloride | |
| NovoRapid FlexPen | | Olanzapine | | Oxycodone Lucis | |
| NovoRapid Penfill | | Olaparib | | Oxycodone Sandoz | |
| 140401 tapia i Gillii | 1 1 | O14p4110 | | Oxyouddid dailadz | 121 |

| Oversedene Condez COO | 107 | Daniaillamina | 117 | PKU Anamix Junior Chocolate | 200 |
|-------------------------------|-----------------|----------------------------------|------------------|--------------------------------|-----|
| Oxycodone Sandoz S29 | | Penicillamine | | PKU Anamix Junior LQ | |
| OxyContin | | PenMix 30 | | | |
| OxyNorm | 127 | | | PKU Anamix Junior Orange | |
| Oxytocin | | PenMix 50 | | PKU Anamix Junior Vanilla | |
| Oxytocin BNM | | Pentasa | | PKU Build 10 | |
| Oxytocin Panpharma | 81 | Pentostatin [Deoxycoformycin | | PKU Build 20 Chocolate | 290 |
| Oxytocin with ergometrine | - | Pentoxifylline [Oxpentifylline]. | | PKU Build 20 Raspberry | |
| maleate | | Peptamen Junior | | Lemonade | |
| Ozurdex | 266 | Pepti-Junior | | PKU Build 20 Smooth | |
| -P- | | Perhexiline maleate | | PKU Build 20 Vanilla | |
| Pacifen | | Pericyazine | | PKU Explore 10 | |
| Pacimol | | Perindopril | | PKU Explore 5 | 289 |
| Paclitaxel | | Periset | | PKU Express 20 | |
| Paclitaxel Actavis | 162 | Periset ODT | | PKU First Spoon | |
| Paclitaxel Ebewe | 162 | Perjeta | | PKU Glytactin RTD 15 | |
| Padagis | | Permethrin | | PKU Glytactin RTD 15 Lite | |
| Paediatric Seravit | | Perrigo | | PKU GMPro LQ | 290 |
| Palbociclib | 170 | Pertuzumab | 216 | PKU GMPro Mix-In | 290 |
| Paliperidone | 138 | Peteha | 106 | PKU GMPro Ultra Lemonade | |
| Paliperidone palmitate | 138 | Pethidine hydrochloride | 128 | PKU GMPro Ultra Vanilla | 290 |
| Palivizumab | 215 | Pevaryl | 67 | PKU Lophlex LQ 10 | 289 |
| Pamidronate disodium | | Pexsig | 49 | PKU Lophlex LQ 20 | 289 |
| Pamisol | 118 | Pfizer Exemestane | 177 | PKU Lophlex Powder | 289 |
| Pamol | 125 | Pfizer S29 | 156 | PKU Lophlex Sensation 20 | 289 |
| Pancreatic enzyme | <mark>22</mark> | Pharmacy Services | <mark>270</mark> | PKU Restore Powder | 290 |
| Pantoprazole | | Pharmascience | | PKU sphere20 Banana | 290 |
| Panzop Relief | | Pheburane | 28 | PKU sphere20 Chocolate | |
| Papaverine hydrochloride | | Phenasen | 157 | PKU sphere20 Lemon | |
| Para-amino salicylic acid | | Phenergan Elixir | 256 | PKU sphere20 Red Berry | |
| Paracetamol | | Phenobarbitone | | PKU sphere20 Vanilla | |
| Paracetamol (Ethics) | | Phenobarbitone sodium | | PKU Start | |
| Paracetamol + Codeine | | Extemporaneous | 273 | Plaquenil | |
| (Relieve) | 127 | Nervous | | Plendil ER | |
| Paracetamol with codeine | | Phenoxybenzamine | | Pneumococcal (PCV13) conjugate | |
| Paraffin | | hydrochloride | 44 | vaccine | |
| Paraffin liquid with wool fat | | Phenoxymethylpenicillin (Pen | | Pneumococcal (PPV23) | |
| Parasiticidal Preparations | | V) | | polysaccharide vaccine | 310 |
| Parnate | | Phenylalanine50 | | Pneumovax 23 | |
| Paromomycin | | Phenytoin sodium | | Podophyllotoxin | |
| Paroxetine | 129 | Phillips Milk of Magnesia | | Polaramine | |
| Paser | | Phlexy 10 | | Poliomyelitis vaccine | |
| Paxam | | Phosphate Phebra | | Poloxamer | |
| Paxlovid | | Phosphorus | | Poly-Gel | |
| Pazopanib | | Phytomenadione | | Poly-Tears | |
| Pazopanib Teva | | Pilocarpine hydrochloride | | Poly-Visc | |
| Peak flow meter | | Pilocarpine nitrate | 260 | Polycal | |
| Pediasure | | Pimafucort | | Polyethylene glycol 400 and | 214 |
| | | Pimecrolimus | | propulane glycol 400 and | 000 |
| Pediasure Plus | | | | propylene glycol | 208 |
| Pediasure RTH | | Pine tar with trolamine laurilsu | | Pomalidomide | |
| Pegaspargase | | and fluorescein | | Pomolide | |
| Pegasys | 113 | Pinetarsol | | Ponstan | |
| Pegfilgrastim | 42 | Pioglitazone | | Posaconazole | |
| Pegylated interferon alfa-2a | | Pirfenidone | | Posaconazole Juno | |
| Pembrolizumab | | Pizotifen | | Potassium chloride | |
| Pemetrexed | | PKU Anamix Infant | | Potassium citrate | |
| Pemetrexed-AFT | 157 | PKU Anamix Junior | 289 | Potassium iodate | 33 |

INDEX: Generic Chemicals and Brands

| Povidone iodine | 71 | Puri-nethol | 156 | RINVOQ | 253 |
|-----------------------------|-----|--------------------------|-----|----------------------------------|------|
| Pradaxa | 41 | Puritan's Pride Vitamin | | Riodine | 71 |
| Pramipexole hydrochloride | 122 | B-2 100 mg | 27 | Risdiplam | 144 |
| Pravastatin | 52 | Pyrazinamide | 106 | Risedronate Sandoz | 119 |
| Praziquantel | 94 | Pyridostigmine bromide | | Risedronate sodium | |
| Prazosin | 44 | Pyridoxine hydrochloride | | Risperdal | 136 |
| Prazosin Mylan | 44 | Pyridoxine multichem | | Risperdal Consta | |
| Pred Forte | | Pyrimethamine | | Risperidone1 | |
| Prednisolone | | Pytazen SR | | Risperidone (Teva) | |
| Prednisolone acetate | | - Q - | | Risperidone Sandoz | |
| Prednisolone sodium | | Quantalan sugar free | 52 | Risperon | |
| phosphate | 267 | Quetapel | | Ritalin | |
| Prednisolone-AFT | | Quetiapine | | Ritalin LA | |
| Prednisone | | Quinapril | | Ritonavir | |
| Prednisone Clinect | | Qvar | | Rituximab (Mabthera) | |
| Pregabalin | | -R- | | Rituximab (Riximyo) | |
| Pregabalin Pfizer | | RA-Morph | 126 | Rivaroxaban | |
| Pregnancy Tests - hCG Urine | 01 | Ralicrom | | Rivastigmine | |
| | | | | | |
| Premarin | | Raloxifene hydrochloride | | Rivastigmine Patch BNM 10 | |
| Prevenar 13 | | Raltegravir potassium | | Rivastigmine Patch BNM 5 | |
| Priadel | | Ramipex | | Rivotril | |
| Primaquine | 104 | Ramipril | | Riximyo | |
| Primidone | | Ranbaxy-Cefaclor | | RIXUBIS | |
| Primidone Clinect | | Rapamune | | Rizamelt | |
| Primolut N | | Rasagiline | | Rizatriptan | |
| Priorix | | Reandron 1000 | | Robinul | |
| Probenecid | | Recombinant factor IX | , | Ronapreve | |
| Probenecid-AFT | | Recombinant factor VIIa | | Ropin | |
| Procarbazine hydrochloride | | Recombinant factor VIII | | Ropinirole hydrochloride | 122 |
| Prochlorperazine | 134 | Rectogesic | 8 | Rosuvastatin | 52 |
| Prochlorperazine Brown & | | Redipred | 85 | Rosuvastatin Viatris | 52 |
| Burk | 134 | Relieve | 116 | Rotarix | 311 |
| Proctofoam | 7 | Relistor | 23 | Rotavirus oral vaccine | 311 |
| Proctosedyl | 8 | Remicade | 203 | Roxane-Propranolol | 48 |
| Procyclidine hydrochloride | 123 | Renilon 7.5 | 280 | Roxithromycin | 96 |
| Progesterone | 87 | Resonium-A | 43 | Rubifen | 146 |
| Proglicem | | Resource Beneprotein | 276 | Rubifen SR | 146 |
| Progynova | | Respiratory Devices | | Rugby Capsaicin Topical Cream | |
| Prolia | | Respiratory Stimulants | | Musculoskeletal | 117 |
| Promethazine hydrochloride | 256 | Retinol palmitate | | Nervous | 124 |
| Propafenone hydrochloride | | ReTrieve | | Rurioctocog alfa pegol [Recombin | nant |
| Propranolol | | Retrovir | | factor VIII] | |
| Propylene glycol | | Revia | | Ruxolitinib | |
| Propylthiouracil | | Revolade | | Rydapt | |
| Prostacur | | Ribociclib | | Rythmodan | |
| Protaphane | | Riboflavin | | Rythmodan - Cheplafarm | |
| Protaphane Penfill | | Ribomustin | | Rytmonorm | |
| | | Ricit | | - S - | 47 |
| Protifar | | | | | 120 |
| Protionamide | | Rifabutin | | Sabril Sacubitril with valsartan | |
| Provera LID | | Rifadin | | | |
| Provera HD | 8/ | Rifadin Sanofi | | SalAir | |
| Psoriasis and Eczema | 70 | Rifampicin | | Salazopyrin | |
| Preparations | | Rifaximin | | Salazopyrin EN | |
| PTU | | Rifinah | | Salbutamol | 257 |
| Pulmicort Turbuhaler | | Rilutek | | Salbutamol with ipratropium | |
| Pulmozyme | 261 | Riluzole | 123 | bromide | 258 |

| Salicylic acid | 74 | Sodium citro-tartrate | 82 | Sulprix | 13 |
|--------------------------------|-------|--------------------------------|-----------|--------------------------------|------|
| Salmeterol | | Sodium cromoglicate | | Sumagran | 13 |
| Sandomigran | 133 | Alimentary | 8 | Sumatriptan | 13 |
| Sandostatin LAR | 175 | Sensory | 267 | Sunitinib | 17 |
| Sanofi Primaquine | 104 | Sodium Fusidate [fusidic acid] | | Sunitinib Pfizer | |
| Sapropterin dihydrochloride | 28 | Dermatological | 67 | Sunscreens | |
| Scalp Preparations | 74 | Infection | 100 | Sunscreens, proprietary | 7 |
| Scopolamine - Mylan | 134 | Sensory | 265 | Sustagen Hospital Formula | 28 |
| Sebizole | 74 | Sodium hyaluronate [Hyaluronic | 0 | Sustagen Hospital Formula | |
| Secukinumab | 229 | acid] | 269 | Active | 28 |
| Sedatives and Hypnotics | 142 | Sodium phenylbutyrate | 28 | Sustanon Ampoules | |
| Seebri Breezhaler | 258 | Sodium picosulfate | 24 | Sylvant | 23 |
| Senna | 24 | Sodium polystyrene sulphonate | 43 | Symbicort Turbuhaler 100/6 | 25 |
| Senokot | 24 | Sodium tetradecyl sulphate | 38 | Symbicort Turbuhaler 200/6 | 25 |
| SensoCard | 15 | Sodium valproate | | Symbicort Turbuhaler 400/12 | 25 |
| Serc | 133 | Sofradex | | Symmetrel | |
| Serenace | 135 | Soframycin | 265 | Sympathomimetics | |
| Seretide | 257 | Solgar2 | 25-27, 29 | Synacthen | 8 |
| Seretide Accuhaler | | Solifenacin succinate | | Synacthen Depot | |
| Serevent | 256 | Solifenacin succinate Max | | Synacthene Retard | 8 |
| Serevent Accuhaler | 256 | Health | 82 | Synagis | |
| Sertraline | 129 | Solifenacin Viatris | 82 | Synthroid | 8 |
| Setrona | 129 | Solu-Cortef | 84 | Syntometrine | |
| Sevredol | 127 | Solu-Medrol | 84 | Syrup (pharmaceutical grade) | |
| Sex Hormones Non | | Solu-Medrol-Act-O-Vial | 84 | Systane Unit Dose | |
| Contraceptive | 85 | Somatropin (Omnitrope) | 88 | · т- | |
| Shingles vaccine | | Sotalol | | Tacrolimus | |
| Shingrix | | Spacer device | 264 | Dermatological | 7 |
| SII-Onco-BCG | | Span-K | | Oncology | |
| Sildenafil | 59 | Spazmol | | Tacrolimus Sandoz | 25 |
| Siltuximab | 232 | Spinal Muscular Atrophy | | Tagrisso | |
| Simvastatin | | Spinraza | | Taliglucerase alfa | 2 |
| Simvastatin Mylan | 53 | Spiolto Respimat | 259 | Tambocor | 4 |
| Simvastatin Viatris | | Spiractin | | Tambocor German | |
| Sinemet | 122 | Spiriva | | Tamoxifen citrate | 17 |
| Sinemet CR | 122 | Spiriva Respimat | | Tamoxifen Sandoz | 170 |
| Sintetica Baclofen Intrathecal | | Spironolactone | | Tamsulosin hydrochloride | |
| Sirolimus | | Sporanox | | Tamsulosin-Rex | |
| Sirturo | | Sprycel | | Tandem Cartridge | |
| Siterone | | Stalevo | | Tandem t:slim X2 with Basal-IQ | |
| Slow-Lopresor | | Stelara | | Tandem t:slim X2 with | |
| Smith BioMed Rapid Pregnand | | Stemetil | | Control-IQ | 10 |
| Test | | Steril-Gene | | Tap water | |
| Sodibic | | SteroClear | | Taro | |
| Sodium acid phosphate | 24 | Stesolid | | Tasigna | |
| Sodium alginate | | Stimulants/ADHD Treatments | | Tasmar | |
| Sodium benzoate | | Stiripentol | | Taurine | |
| Sodium bicarbonate | | Stomahesive | | TCu 380 Plus Normal | |
| Blood | 42-43 | Strides Shasun | | Tecentriq | |
| Extemporaneous | | Stromectol | | Tecfidera | |
| Sodium calcium edetate | | Sucralfate | | Tegretol | |
| Sodium chloride | | Sulfadiazin-Heyl | | Tegretol AU | |
| Blood | 43 | Sulfadiazine Silver | | Tegretol CR | |
| Respiratory | | Sulfadiazine sodium | | Telfast | |
| Sodium citrate with sodium lau | | Sulfasalazine | | Temaccord | |
| sulphoacetate | | Sulphur | | Temazepam | |
| - 2.6 | 1 | | | a=opa | 1 10 |

INDEX: Generic Chemicals and Brands

| Temozolomide | 163 | Topical Products for Joint and | | TYR Explore 5 | 29 |
|------------------------------------|-----|----------------------------------|--------|------------------------------------|----------------|
| Temozolomide-Taro | 163 | Muscular Pain | 117 | TYR Lophlex LQ 20 | 29 |
| Tenofovir disoproxil | 107 | Topiramate | 132 | TYR Sphere 20 | 29 |
| Tenofovir Disoproxil Emtricitabine | е | Topiramate Actavis | 132 | Tyrosine1000 | |
| Viatr | 109 | Total parenteral nutrition (TPN) | 43 | Tysabri | 14 |
| Tenofovir Disoproxil Viatris | 107 | TPN | | . U - | |
| Tenoxicam | 116 | Tramadol hydrochloride | 128 | UK Synacthen | 8 |
| Tensipine MR10 | | Tramal SR 100 | 128 | Ultibro Breezhaler | 25 |
| Tepadina | 154 | Tramal SR 150 | 128 | Ultraproct | |
| Terbinafine | | Tramal SR 200 | 128 | Umeclidinium | 25 |
| Terbutaline sulphate | 257 | Trandate | 48 | Umeclidinium with vilanterol | |
| Teriflunomide | 141 | Tranexamic acid | 38 | Univent2 | 58, 26 |
| Teriflunomide Sandoz | | Tranylcypromine sulphate | 129 | Upadacitinib | 25 |
| Teriparatide | 119 | Trastuzumab (Herzuma) | 235 | Ural | |
| Teriparatide - Teva | 119 | Trastuzumab deruxtecan | | Urea | <mark>7</mark> |
| Testogel | | Trastuzumab emtansine | 238 | Urex Forte | 5 |
| Testosterone | | Travatan | 268 | Urinary Agents | 8 |
| Testosterone cipionate | | Travoprost | | Urinary Tract Infections | |
| Testosterone esters | | Treatments for Dementia | | UroFos | |
| Testosterone undecanoate | | Treatments for Substance | | Uromitexan | |
| Tetrabenazine | | Dependence | 148 | Ursodeoxycholic acid | |
| Tetrabromophenol | | Trelegy Ellipta | | Ursosan | |
| Tetracosactrin | | Trental 400 | | Ustekinumab | |
| Tetracycline | | Tretinoin | | Utrogestan | |
| Teva Lisinopril | | Dermatological | 66 | - V - | |
| Teva-Ketoconazole | | Oncology | | Vaccinations | 29 |
| Thalidomide | | Trexate | | Vaclovir | |
| Thalomid | | Triamcinolone acetonide | | Valaciclovir | |
| Theophylline | | Alimentary | 30 | Valganciclovir | |
| Thiamine hydrochloride | | Dermatological | | Valganciclovir Viatris | |
| Thiamine multichem | | Hormone | | Valine50 | |
| THIO-TEPA | | Triamcinolone acetonide with | | Vancomycin | |
| Thioguanine | | gramicidin, neomycin and ny | etatin | Vannair | |
| Thiotepa | | Dermatological | | Varenicline Pfizer | |
| Thyroid and Antithyroid Agents | | Sensory | | Varenicline tartrate | |
| Ticagrelor | | Trientine | | Varicella vaccine [Chickenpox | |
| Ticagrelor Sandoz | | Trientine Waymade | | vaccine] | 31 |
| Tilcotil | | Trikafta | | Varicella zoster vaccine [Shingles | |
| Timolol | | Trimethoprim | | vaccine] | |
| Tiotropium bromide | | Trimethoprim with | 101 | Varilrix | |
| Tiotropium bromide with | 200 | sulphamethoxazole | | Various | |
| olodaterol | 250 | [Co-trimoxazole] | 101 | VariSoft | |
| Tivicay | | Triovir | | Vasodilators | |
| Tixagevimab with cilgavimab | | Trisequens | | Vasopressin Agonists | |
| TMP | | Trisul | | Vasorex | |
| Tobramycin | 101 | Trophic Hormones | | Vebulis | |
| Infection | 100 | | | Vedafil | |
| • | | Tropicamide | | Vedolizumab | |
| Sensory | | I rulicity | | | |
| Tobramycin (Viatris) | | TruSteel | | Veletri | |
| Tobramycin BNM | | Tryzan | | Venclexta | |
| Tobrex | | Tuberculin PPD [Mantoux] test | | Venetoclax | |
| Tocilizumab | | Tubersol | | Venlafaxine | |
| Tofranil | | Two Cal HN | | Venomil | |
| Tolcapone | | TYR Anamix Infant | | VENOX | |
| Tolvaptan | | TYR Anamix Junior | 291 | Ventolin | |
| Topamax | 132 | TYR Anamix Junior LQ | 291 | Vepesid | 15 |

| Verapamil hydrochloride49 | Х |
|---|--------|
| Vermox94 | Χ |
| Versacloz135 | Х |
| Vesanoid164 | Χ |
| Vexazone12 | Х |
| Vfend103 | |
| Viaderm KC69 | Ζ |
| Victoza12 | Z |
| Vigabatrin132 | Z |
| Vigisom142 | Z |
| Vildagliptin12 | Z |
| Vildagliptin with metformin | Z |
| hydrochloride12 | Z |
| Vimpat130 | Z |
| Vinblastine sulphate165 | Z |
| Vincristine sulphate | Z |
| Vinorelbine 165 | Z |
| Vinorelbine Ebewe | _ |
| Vinorelbine Ebewe 103 Vinorelbine Te Arai 165 | Z |
| Viramune Suspension | Z |
| Viraniune Suspension | Z |
| ViruPOS | Z |
| Vit.D3 | |
| Vita-B1231 | Z Z |
| VitA-POS | _ |
| Vitabdeck32 | Z |
| Vital | Z |
| Vitamin B complex31 | Z |
| Vitamin B6 2531 | Z |
| Vitamins31–32 | Z |
| Vitarubin Depot Injection31 | Z |
| Vivonex TEN280 | Z |
| Voltaren | |
| Voltaren D116 | _ |
| Voltaren SR116 | Z |
| Volumatic264 | |
| Voriconazole103 | |
| Votrient | Z |
| Vttack | Z |
| Vyvanse145 | Z |
| - W - | Z |
| Warfarin sodium41 | Z |
| Wart Preparations75 | Z |
| Wasp venom allergy treatment255 | Z |
| Water | Z |
| Blood43 | Z |
| Extemporaneous273 | Z |
| White Soft Liquid Paraffin AFT70 | Z |
| Wool fat with mineral oil71 | Z |
| - X - | Z |
| Xaluprine156 | |
| Xarelto41 | |
| Xgeva117 | |
| Xifaxan10 | |
| XMET Maxamum288 | |
| Xolair214 | |
| Xolair AU214 | |

| XP Maxamum2 | 289 |
|-------------------------------|-----|
| Xylocaine 1 | 24 |
| Xylocaine 2% Jelly1 | 23 |
| Xylocard 5001 | 24 |
| Xvntha | 38 |
| - Z - | |
| Zapril | 44 |
| Zarontin1 | 30 |
| Zaroxolyn | |
| Zavedos1 | 59 |
| Zeffix1 | 07 |
| Zejula1 | |
| Zematop | 74 |
| Zetlam1 | 07 |
| Ziagen1 | 12 |
| Zidovudine [AZT]1 | 12 |
| Zidovudine [AZT] with | |
| lamivudine1 | 12 |
| Ziextenzo | |
| Ziextenzo AU | 42 |
| Zimybe | |
| Zinc and castor oil | 70 |
| Zinc sulphate | |
| Zincaps | 34 |
| Ziprasidone1 | |
| Zista2 | 255 |
| Zithromax | 95 |
| Zo-Rub HP1 | 24 |
| Zo-Rub Osteo1 | |
| Zoladex | |
| Zoledronic acid | |
| Hormone | 84 |
| Musculoskeletal1 | 20 |
| Zoledronic acid Viatris | |
| Hormone | 84 |
| Musculoskeletal1 | |
| Zopiclone1 | 43 |
| Zopiclone Actavis1 | 43 |
| Zostrix1 | 17 |
| Zostrix HP1 | 24 |
| Zuclopenthixol decanoate1 | 39 |
| Zuclopenthixol hydrochloride1 | 36 |
| Zusdone1 | 36 |
| Zyban1 | 49 |
| Zypine1 | 36 |
| Zypine ODT1 | 36 |
| Zyprexa Relprevv1 | 37 |
| Zytiga1 | 73 |
| Zyvox1 | 06 |
| | |