Alimentary Tract & Metabolism

Blood & Blood Forming Organs

Cardiovascular System

Genito Urinary System

Dermatologicals

General Rules

5

6

35

44

66

75

March 2025	
Volume 32	

Section A

Section B

Editors:

Kaye Wilson & Doris Chong email: enquiry@pharmac.govt.nz Telephone +64 4 460 4990 Level 9, 40 Mercer Street PO Box 10 254 Wellington

Freephone Information Line 0800 66 00 50 (9am – 5pm weekdays)

Circulation

You can register to have an electronic version of the Pharmaceutical Schedule (link to PDF copy) emailed to your nominated email address each month by subscribing at pharmac.govt.nz/subscribe.

Production

Typeset automatically from XML and T_EX. XML version of the Schedule available from schedule.pharmac.govt.nz/pub/schedule

Programmers

Anrik Drenth

CC

 $email: \ texschedule @pharmac.govt.nz \\$

©Pharmaceutical Management Agency

(i) ISSN 1179-3686

This work is licensed under the Creative Commons Attribution 4.0 International licence. In essence, you are free to copy, distribute and adapt it, as long as you attribute the work to Pharmac and abide by the other licence terms. To view a copy of this licence, visit: creativecommons.org/licenses/by/4.0/. Attribution to Pharmac should be in written form and not by reproduction of the Pharmac logo. While care has been taken in compiling this Schedule, Pharmac takes no responsibility for any errors or omissions, and shall not be liable for any consequences arising there from.

82
93
115
121
151
255
266
271
273
275
275

Introducing Pharmac

The Pharmaceutical Management Agency (Pharmac) makes decisions that help control Government spending on pharmaceuticals. This includes community pharmaceuticals, hospital pharmaceuticals, vaccines and increasingly, hospital medical devices. Pharmac negotiates prices, sets subsidy levels and conditions, and makes decisions on changes to the subsidised list.

Pharmac's role:

"to secure for eligible people in need of pharmaceuticals, the best health outcomes that are reasonably achievable from pharmaceutical treatment and from within the amount of funding provided."

Pae Ora (Healthy Futures) Act 2022

To ensure our decisions are as fair and robust as possible we use a decision-making process that incorporates clinical, economic and commercial issues. We also seek the views of users and the wider community through consultation. The processes we generally use are outlined in our Operating Policies and Procedures.

Further information about Pharmac and the way we make funding decisions can be found on the Pharmac website at https://pharmac.govt.nz/about.

Purpose of the Pharmaceutical Schedule

The purpose of the Schedule is to list:

- the Community Pharmaceuticals that are subsidised by the Government and to show the amount of the subsidy paid to contractors, as well as the manufacturer's price and any access conditions that may apply;
- the Hospital Pharmaceuticals that may be used in Health NZ Hospitals, as well as any access conditions that may apply; and
- the Pharmaceuticals, including Medical Devices, used in Health NZ Hospitals for which national prices have been negotiated by Pharmac.

The Schedule does not show the final cost to Government of subsidising each Community Pharmaceutical or to Health NZ Hospitals in purchasing each Pharmaceutical, since that will depend on any rebate and other arrangements Pharmac has with the supplier and, for Pharmaceuticals used in Health NZ Hospitals, on any logistics arrangements put in place.

This book contains sections A to D and Section I of the Pharmaceutical Schedule and lists the Pharmaceuticals funded for use in the community, including vaccines, as well as haemophilia and cancer treatments given in Health NZ Hospitals. Section H lists the Pharmaceuticals that that can be used in Health NZ Hospitals and is a separate publication.


The Pharmaceuticals in this book are listed by therapeutic group, which is based on the WHO Anatomical Therapeutic Chemical (ATC) system. The listings are displayed alphabetically under each heading.

The index lists both chemical entities and product brand names.

Explaining pharmaceutical entries

The Pharmaceutical Schedule lists pharmaceuticals subsidised by the Government, the subsidy, the supplier's price and the access conditions that may apply.

Example



Glossary

Units of Measure

gram g	
kilogram kg	
international unit iu	

Abbreviations

Capsule Cream Device Dispersible Effervescent Emulsion	Amp Cap Crm Dev Disp Eff Emul EC
Enteric Coated	EC

microgram me	cg
milligramn	ng
millilitreı	ml

millimole	mmol
unit	u

Gelatinous	Gel	SolutionSoln
Granules	Gran	SuppositorySupp
Infusion	Inf	TabletTab
Injection	Inj	Tincture Tinc
Liquid	Liq	Trans Dermal Delivery
Long Acting	LA	SystemTDDS
Ointment	Oint	-
Sachet	Sach	

Read the General Rules : https://pharmac.govt.nz/section-a.

SECTION B: ALIMENTARY TRACT AND METABOLISM

	Subsidy		Fully	Brand or Generic
	(Manufacturer's Price) \$	Per	Subsidised	Manufacturer
Antacids and Antiflatulents				
Antacids and Reflux Barrier Agents				
LGINIC ACID Sodium alginate 225 mg and magnesium alginate 87.5 mg pe sachet		30	1	Gaviscon Infant
ODIUM ALGINATE • Tab 500 mg with sodium bicarbonate 267 mg and calcium	1.00	CO		
carbonate 160 mg - peppermint flavour	1.80 (14.39)	60		Gaviscon Extra Strength
Oral liq 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg per 10 ml	1.50 (7.50)	500 m		Acidex
Phosphate Binding Agents				
LUMINIUM HYDROXIDE	12.56	100	1	Alu-Tab
ALCIUM CARBONATE Oral liq 1,250 mg per 5 ml (500 mg elemental per 5 ml) – Subsidy by endorsement		500 m 473 m		Roxane Calcium carbonate PAI \$29
Only when prescribed for patients unable to swallow calci inappropriate and the prescription is endorsed according!		ts or w	here calci	
Antidiarrhoeals				
Agents Which Reduce Motility				
OPERAMIDE HYDROCHLORIDE – Up to 30 cap available on a Tab 2 mg Cap 2 mg 	10.75	400 400		Nodia Diamide Relief
Rectal and Colonic Anti-inflammatories				
UDESONIDE Cap modified-release 3 mg – Special Authority see SA1886 below – Retail pharmacy		90	1	Budesonide Te Arai
SA1886 Special Authority for Subsidy itial application — (Crohn's disease) from any relevant practi e following criteria: oth:		alid foi	6 months	for applications meeting
 Mild to moderate ileal, ileocaecal or proximal Crohn's disea Any of the following: 	ise; and			
2.1 Diabetes; or				

continued...

6

Subsidy	Full	y Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per 🖌	Manufacturer	

continued...

- 2.2 Cushingoid habitus; or
- 2.3 Osteoporosis where there is significant risk of fracture; or
- 2.4 Severe acne following treatment with conventional corticosteroid therapy; or
- 2.5 History of severe psychiatric problems associated with corticosteroid treatment; or
- 2.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
- 2.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated).

Initial application — (collagenous and lymphocytic colitis (microscopic colitis)) from any relevant practitioner. Approvals valid for 6 months where patient has a diagnosis of microscopic colitis (collagenous or lymphocytic colitis) by colonoscopy with biopsies.

Initial application — (gut Graft versus Host disease) from any relevant practitioner. Approvals valid for 6 months where patient has a gut Graft versus Host disease following allogenic bone marrow transplantation*.

Note: Indication marked with * is an unapproved indication.

Initial application — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has autoimmune hepatitis*; and
- 2 Patient does not have cirrhosis; and
- 3 Any of the following:

- 3.1 Diabetes; or
- 3.2 Cushingoid habitus; or
- 3.3 Osteoporosis where there is significant risk of fracture; or
- 3.4 Severe acne following treatment with conventional corticosteroid therapy; or
- 3.5 History of severe psychiatric problems associated with corticosteroid treatment; or
- 3.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
- 3.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated); or
- 3.8 Adolescents with poor linear growth (where conventional corticosteroid use may limit further growth).
- Note: Indication marked with * is an unapproved indication.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

HIDROCORTISONE ACETATE		
Rectal foam 10%, CFC-Free (14 applications)57.09	15 g OP	 Colifoam
HYDROCORTISONE ACETATE WITH PRAMOXINE HYDROCHLORIDE		
Topical aerosol foam, 1% with pramoxine hydrochloride 1%	10 g OP	 Proctofoam S29
MESALAZINE		
Tab 400 mg	100	Asacol
Tab long-acting 500 mg56.10	100	Pentasa
Tab 800 mg	90	Asacol
-		Asacol S29 S29
Modified release granules, 1 g118.10	100 OP	Pentasa
Enema 1 g per 100 ml	7	Pentasa
Suppos 500 mg	20	Asacol
Suppos 1 g		Pentasa
(Asacol S29 S29 Tab 800 mg to be delisted 1 July 2025)		

(Asacol S29 S29 Tab 800 mg to be delisted 1 July 2025)

	Subsidy		Fully	
	(Manufacturer's Price) \$	Per	Subsidised	Generic Manufacturer
DLSALAZINE				
Tab 500 mg		60	1	Atnahs
,				Olsalazine S29
	93.37	100	1	Dipentum
Cap 250 mg	53.00	100	✓	Dipentum
ODIUM CROMOGLICATE				
Cap 100 mg	113.35	100	✓	Ralicrom
ULFASALAZINE				
🗧 Tab 500 mg		100		Salazopyrin
F Tab EC 500 mg	20.54	100	1	Salazopyrin EN
Local preparations for Anal and Rectal Disorders	\$			
Antihaemorrhoidal Preparations				
UOCORTOLONE CAPROATE WITH FLUOCORTOLONE PIVA			NE	
Oint 950 mcg, with fluocortolone pivalate 920 mcg, and				
cinchocaine hydrochloride 5 mg per g	13.05	30 g C	P 🗸	Ultraproct
Suppos 630 mcg, with fluocortolone pivalate 610 mcg, and				
cinchocaine hydrochloride 1 mg		12	1	Ultraproct
YDROCORTISONE WITH CINCHOCAINE				
Oint 5 mg with cinchocaine hydrochloride 5 mg per g		30 g C	P 🗸	Proctosedyl
Suppos 5 mg with cinchocaine hydrochloride 5 mg per g	9.90	12	~	Proctosedyl
Management of Anal Fissures				
LYCERYL TRINITRATE - Special Authority see SA1329 below	- Retail pharmacy			
Oint 0.2%		30 g C	P 🗸	Rectogesic
SA1329 Special Authority for Subsidy				
itial application from any relevant practitioner. Approvals valid		ewal u	nless notif	ied where the patient has
nronic anal fissure that has persisted for longer than three weeks				
Antispasmodics and Other Agents Altering Gut I	Motility			
	,			
Inj 200 mcg per ml, 1 ml ampoule – Up to 10 inj available on a	10.00	-		Dahimul
PSO		5	~	Robinul
YOSCINE BUTYLBROMIDE	0.05	00		
5 Tab 10 mg	2.25	20	~	Hyoscine Butylbromide
				(Adiramedica)
		400		

			(Aurameu
6.0	35	100	 Buscopan
Hyoscine Butylbromide (Adiramedica) to be Principal Supply on 1 A	April 2025		
* Inj 20 mg, 1 ml – Up to 5 inj available on a PSO	91	5	 Spazmol
(Buscopan Tab 10 mg to be delisted 1 April 2025)			
MEBEVERINE HYDROCHLORIDE			
* Tab 135 mg8.	50	90	 Colofac

	Subsidy	Fu	lly Brand or
	(Manufacturer's Price)	Subsidise	ed Generic
	\$	Per	Manufacturer
Antiulcerants			
Antisecretory and Cytoprotective			
MISOPROSTOL – Wastage claimable * Tab 200 mcg – Up to 120 tab available on a PSO	47.73	120	Cytotec
Helicobacter Pylori Eradication			
CLARITHROMYCIN Tab 500 mg – Subsidy by endorsement a) Maximum of 28 tab per prescription b) Subsidised only if prescribed for helicobacter pylori Note: the prescription is considered endorsed if cla inhibitor and either amoxicillin or metronidazole.	eradication and prescr	iption is endo	
H2 Antagonists			
FAMOTIDINE – Only on a prescription * Tab 20 mg	4.91	100 •	Famotidine Hovid 629
* Tab 40 mg	10.32	100 •	Famotidine Hovid S29
Inj 10 mg per ml, 4 ml – Subsidy by endorsement Subsidy by endorsement – Subsidised for patients rece			Mylan S29 care.
Proton Pump Inhibitors			
LANSOPRAZOLE			
* Cap 15 mg * Cap 30 mg			 Lanzol Relief Lanzol Relief
OMEPRAZOLE For omeprazole suspension refer Standard Formulae, page	273		
* Cap 10 mg	2.06		Omeprazole Teva <u>Omeprazole actavis</u> 10
* Cap 20 mg	2.02		Omeprazole Teva <u>Omeprazole actavis</u> 20
* Cap 40 mg	3.18		Omeprazole Teva Omeprazole actavis 40
 Powder – Only in combination Only in extemporaneously compounded omeprazole st 		5g •	Midwest
Inj 40 mg ampoule with diluent			<u>Dr Reddy's</u> <u>Omeprazole</u> Ocicure \$29
PANTOPRAZOLE * Tab EC 20 mg	1 99	90 •	Panzop Relief
* Tab EC 20 mg			Panzop Relief

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price \$	e) Per	Fully Subsidised	
Site Protective Agents				
COLLOIDAL BISMUTH SUBCITRATE				
Tab 120 mg	14.51	50	1	Gastrodenol S29
SUCRALFATE Tab 1 g	35.50	120		
·	(48.28)	120		Carafate
Bile and Liver Therapy				
RIFAXIMIN – Special Authority see SA1461 below – Retail pharr Tab 550 mg		56	1	<u>Xifaxan</u>
SA1461 Special Authority for Subsidy Initial application only from a gastroenterologist, hepatologist or hepatologist. Approvals valid for 6 months where the patient has tolerated doses of lactulose. Renewal only from a gastroenterologist, hepatologist or Practitior hepatologist. Approvals valid without further renewal unless notifi benefiting from treatment.	hepatic encephalop ner on the recomme	oathy de	espite an of a gast	adequate trial of maximum roenterologist or
Diabetes				
Hyperglycaemic Agents				
DIAZOXIDE – Special Authority see SA1320 below – Retail phar	macy			
Cap 25 mg	110.00	100		Proglicem S29
Cap 100 mg		100		Proglicem S29
Oral liq 50 mg per ml		30 ml O	P 🗸	e5 Pharma S29
SA1320 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valic hypoglycaemia caused by hyperinsulinism.	l for 12 months whe	ere use	d for the tr	reatment of confirmed
Renewal from any relevant practitioner. Approvals valid without f appropriate and the patient is benefiting from treatment. GLUCAGON HYDROCHLORIDE	further renewal unle	ess noti	ied where	e the treatment remains
Inj 1 mg syringe kit – Up to 5 kit available on a PSO	32.00	1	~	Glucagen Hypokit
Insulin - Short-acting Preparations				
INSULIN NEUTRAL				
▲ Inj human 100 u per ml		10 ml O		Actrapid
▲ Inj human 100 u per ml, 3 ml	42.66	5	1	Humulin R Actrapid Penfill Humulin R
Insulin - Intermediate-acting Preparations				
INSULIN ASPART WITH INSULIN ASPART PROTAMINE				
Inj 100 iu per ml, 3 ml prefilled pen	52.15	5	1	NovoMix 30 FlexPen

	Subsidy		Fully Brand or
	(Manufacturer's F	Price) Subs	idised Generic
	\$	Per	 Manufacturer
INSULIN ISOPHANE			* · · · · · · · · · · · · · · · · · · ·
Inj human 100 u per ml	17.68	10 ml OP	 Humulin NPH
			 Protaphane
▲ Inj human 100 u per ml, 3 ml	29.86	5	 Humulin NPH
		0	_
			 Protaphane Penfill
INSULIN ISOPHANE WITH INSULIN NEUTRAL			
Inj human with neutral insulin 100 u per ml	25.26	10 ml OP	🖌 Humulin 30/70
			✓ Mixtard 30
	40.00	-	_
Inj human with neutral insulin 100 u per ml, 3 ml		5	 Humulin 30/70
			PenMix 30
			 PenMix 50
(Mixtard 30 Inj human with neutral insulin 100 u per ml to be de	liated 1 June 2021	-)	
(PenMix 50 Inj human with neutral insulin 100 u per ml, 3 ml to	be delisted 1 June	e 2025)	
NSULIN LISPRO WITH INSULIN LISPRO PROTAMINE			
	al		
Inj lispro 25% with insulin lispro protamine 75% 100 u per n			.
3 ml		5	 Humalog Mix 25
Inj lispro 50% with insulin lispro protamine 50% 100 u per n	nl.		
3 ml		5	Humalog Mix 50
01111		5	
Insulin - Long-acting Preparations			
INSULIN GLARGINE			
Inj 100 u per ml, 10 ml	63.00	1	 Lantus
Inj 100 u per ml, 3 ml	94.50	5	✓ Lantus
 Inj 100 u per ml, 3 ml disposable pen 		5	✓ Lantus SoloStar
		5	
In sufficient Description Descriptions			
Insulin - Rapid Acting Preparations			
NOUNDAODADT			
INSULIN ASPART			_
Inj 100 u per ml, 10 ml		1	NovoRapid
Inj 100 u per ml, 3 ml	51.19	5	NovoRapid Penfill
▲ Inj 100 u per ml, 3 ml syringe		5	 NovoRapid FlexPen
		5	
NSULIN GLULISINE			
Inj 100 u per ml, 10 ml		1	 Apidra
▲ Inj 100 u per ml, 3 ml		5	 Apidra
Inj 100 u per ml, 3 ml disposable pen	46.07	5	 Apidra SoloStar
NSULIN LISPRO			
▲ Inj 100 u per ml, 10 ml	34 92	10 ml OP	 Humalog
Inj 100 u per ml, 3 ml		5	 Humalog
Alpha Glucosidase Inhibitors			
ACARROOF			
ACARBOSE			_
* Tab 50 mg	11.20	90	✓ <u>Accarb</u>
* Tab 100 mg		90	✓ Accarb
		-	
Oral Hypoglycaemic Agents			
GLIBENCLAMIDE			
W Tab C man	7 50		
* Tab 5 mg	7.50	100	 Daonil
-	7.50	100	V Daonii
* Tab 5 mg GLICLAZIDE * Tab 80 mg		100 500	 Daonii <u>Glizide</u>

A Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic Manufacturer
	\$	Per	/	Manufacturer
GLIPIZIDE * Tab 5 mg	6.96	100		Minidiab
•	0.00	100	• 1	WIIIIulab
METFORMIN HYDROCHLORIDE				
* Tab immediate-release 500 mg		1,000	✓ I	Metformin Viatris
* Tab immediate-release 850 mg	11.28	500	✓ I	Metformin Viatris
PIOGLITAZONE				
* Tab 15 mg	6.15	90	1	/exazone
* Tab 30 mg	7.25	90	 Image: A start of the start of	/exazone
* Tab 45 mg		90	 Image: A start of the start of	/exazone
VILDAGLIPTIN			-	
Tab 50 mg	25.00	60	1	Galvus
0		00	• (Jaivus
VILDAGLIPTIN WITH METFORMIN HYDROCHLORIDE				
Tab 50 mg with 1,000 mg metformin hydrochloride		60		Galvumet
Tab 50 mg with 850 mg metformin hydrochloride		60	✓ (Galvumet

GLP-1 Agonists

DULAGLUTIDE - Special Authority see SA2338 below - Retail pharmacy

Note: Not to be given in combination with another funded GI	LP-1 agonist or em	npagliflozin /	empagliflozin with metformin
hydrochloride unless receiving empagliflozin / empagliflozin v	with metformin hyd	Irochloride f	or the treatment of heart failure.
Inj 1.5mg per 0.5 ml prefilled pen		4	 Trulicity

➡SA2338 Special Authority for Subsidy

Note: Subsidy for patients with existing approvals prior to 1 May 2024. Approvals valid without further renewal unless notified. No new patients will be granted from 1 May 2024 until further notice.

LIRAGLUTIDE - Special Authority see SA2440 below - Retail pharmacy

a) Maximum of 9 inj per prescription

b)

- a) Note: Not to be given in combination with another funded GLP-1 agonist or empagliflozin / empagliflozin with metformin hydrochloride unless receiving empagliflozin / empagliflozin with metformin hydrochloride for the treatment of heart failure.
- b) Maximum of 1 pack of 3 (6 mg per ml, 3 ml) prefilled pens will be funded per month.

Inj 6 mg per ml, 3 ml prefilled pen		3	 Victoza
-------------------------------------	--	---	-----------------------------

⇒SA2440 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has type 2 diabetes; and
- 2 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of all of the following funded blood glucose lowering agents for a period of least 6 months, where clinically appropriate: empagliflozin, metformin, and vildagliptin; and
- 3 Any of the following:
 - 3.1 Patient is Māori or any Pacific ethnicity*; or
 - 3.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)*; or
 - 3.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator*; or
 - 3.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult*; or

continued...

Subsidy	F	ully	Brand or	
(Manufacturer's Price)	Subsidi	sed	Generic	
\$	Per	✓	Manufacturer	

continued...

3.5 Patient has diabetic kidney disease (see note b)*.

Notes: * Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

- a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack, ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.
- b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m2 in the presence of diabetes, without alternative cause identified.
- c) Funded GLP-1a treatment is not to be given in combination with (empagliflozin /empagliflozin with metformin hydrochloride) unless receiving (empagliflozin or empagliflozin in combination with metformin hydrochloride) for the treatment of heart failure.

SGLT2 Inhibitors

⇒SA2408 Special Authority for Subsidy

Initial application — (heart failure reduced ejection fraction) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has heart failure; and
- 2 Patient is in NYHA functional class II or III or IV; and
- 3 Either:
 - 3.1 Patient has a documented left ventricular ejection fraction (LVEF) of less than or equal to 40%; or
 - 3.2 An ECHO is not reasonably practicable, and in the opinion of the treating practitioner the patient would benefit from treatment; and
- 4 Patient is receiving concomitant optimal standard funded chronic heart failure treatment.

Initial application — (Type 2 Diabetes) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 Patient has previously received an initial approval for a GLP-1 agonist; or
- 2 All of the following:
 - 2.1 Patient has type 2 diabetes; and
 - 2.2 Any of the following:
 - 2.2.1 Patient is Maori or any Pacific ethnicity*; or
 - 2.2.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)*; or
 - 2.2.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator*; or
 - 2.2.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult*; or
 - 2.2.5 Patient has diabetic kidney disease (see note b)*; and
 - 2.3 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of at least one blood-glucose lowering agent (e.g. metformin, vildagliptin, or insulin) for at least 3 months.

Notes: * Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

- a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack, ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.
- b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in

continued...

AThree months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m2 in the presence of diabetes, without alternative cause.

c) Funded [empagliflozin / empagliflozin with metformin hydrochloride] treatment is not to be given in combination with a funded GLP-1 unless receiving (empagliflozin / empagliflozin with metformin hydrochloride] for the treatment of heart failure.

EMPAGLIFLOZIN - Special Authority see SA2408 on the prev	ious page – Retail p	harmacy	
* Tab 10 mg		30	 Jardiance
* Tab 25 mg		30	 Jardiance
EMPAGLIFLOZIN WITH METFORMIN HYDROCHLORIDE -	Special Authority se	e SA2408 o	n the previous page – Retail
pharmacy			
* Tab 5 mg with 1,000 mg metformin hydrochloride	58.56	60	 Jardiamet
* Tab 5 mg with 500 mg metformin hydrochloride	58.56	60	 Jardiamet
* Tab 12.5 mg with 1,000 mg metformin hydrochloride		60	 Jardiamet
* Tab 12.5 mg with 500 mg metformin hydrochloride		60	 Jardiamet

Diabetes Management

Ketone Testing

BLOOD KETONE DIAGNOSTIC TEST STRIP - Subsidy by endorsement

- a) Not on a BSO
- b) Maximum of 20 strip per prescription
- c) Up to 10 strip available on a PSO
- d) Patient has any of the following:
 - 1) type 1 diabetes; or
 - 2) permanent neonatal diabetes; or
 - 3) undergone a pancreatectomy; or
 - 4) cystic fibrosis-related diabetes; or

5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly.

Dual Blood Glucose and Blood Ketone Testing

DUAL BLOOD GLUCOSE AND BLOOD KETONE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A dual blood glucose and blood ketone diagnostic test meter is subsidised for a patient who has:
 - type 1 diabetes; or
 - 2) permanent neonatal diabetes; or
 - 3) undergone a pancreatectomy; or
 - 4) cystic fibrosis-related diabetes; or
 - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly. Only 1 meter per patient will be subsidised (no repeat prescriptions). For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Meter with 50 lancets	a lancing device and	10 blood glucose
-----------------------	----------------------	------------------

diagnostic test strips	 1 OP	CareSens Dua

14

sidy irer's Price) S	Fully Subsidised	Brand or Generic	
 B Per	✓	Manufacturer	
			-

Blood Glucose Testing

BLOOD GLUCOSE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A diagnostic blood glucose test meter is subsidised for a patient who:
 - 1) is receiving insulin or sulphonylurea therapy; or
 - 2) is pregnant with diabetes: or
 - 3) is on home TPN at risk of hypoglycaemia or hyperglycaemia; or
 - 4) has a genetic or an acquired disorder of glucose homeostasis, excluding type 1 or type 2 diabetes and metabolic syndrome.

The prescription must be endorsed accordingly. Only one CareSens meter per patient will be subsidised (no repeat prescriptions). Patients already using the CareSens N POP meter and CareSens N meter are not eligible for a new meter, unless they have:

- 1) type 1 diabetes; or
- 2) permanent neonatal diabetes: or
- 3) undergone a pancreatectomy; or
- 4) cvstic fibrosis-related diabetes.

For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Meter with 50 lancets, a lancing device and 10 diagnostic test

strips				 10.00	1 OP	 ✓ CareSens N ✓ CareSens N POP
				20.00		✓ CareSens N Premier
Note: Only 1	meter avai	lable per F	SO			

BLOOD GLUCOSE DIAGNOSTIC TEST STRIP - Up to 50 test available on a PSO

The number of test strips available on a prescription is restricted to 50 unless:

- 1) Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- 2) Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed: or
- Prescribed for a pregnant woman with diabetes and endorsed accordingly; or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

BLOOD GLUCOSE TEST STRIPS (VISUALLY IMPAIRED)

The number of test strips available on a prescription is restricted to 50 unless:

- 1) Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- 2) Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed: or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly: or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

50 test OP SensoCard

50 test OP

CareSens N

CareSens PRO

	Subsidy (Manufacturer's Price) \$	Su Per	Fully bsidised	Brand or Generic Manufacturer
Insulin Syringes and Needles				
Subsidy is available for disposable insulin syringes, need the supply of insulin or liraglutide or when prescribed for annotate the prescription as endorsed where there exists	a patient and the prescription	is endo	rsed acco	ordingly. Pharmacists may
NSULIN PEN NEEDLES - Maximum of 200 dev per pre	escription			
¥ 29 g × 12.7 mm		100	🗸 E	B-D Micro-Fine
米 31 g × 5 mm		100	🖌 E	B-D Micro-Fine
★ 31 g × 6 mm	9.50	100	🗸 E	Berpu
¥ 31 g × 8 mm		100	🗸 E	B-D Micro-Fine
米 32 g × 4 mm		100	🗸 E	B-D Micro-Fine
INSULIN SYRINGES, DISPOSABLE WITH ATTACHED	NEEDLE - Maximum of 200	dev per	prescript	ion
* Syringe 0.3 ml with 29 g × 12.7 mm needle		100		B-D Ultra Fine
	1.36	10		
	(1.99)		E	B-D Ultra Fine
* Syringe 0.3 ml with 31 g × 8 mm needle		100	🖌 E	B-D Ultra Fine II
	1.30	10		
	(1.99)		E	B-D Ultra Fine II
Syringe 0.5 ml with 29 g × 12.7 mm needle		100	🗸 E	B-D Ultra Fine
	1.36	10		
	(1.99)		-	3-D Ultra Fine
Syringe 0.5 ml with 31 g × 8 mm needle		100	🗸 E	B-D Ultra Fine II
	1.36	10		
	(1.99)		-	B-D Ultra Fine II
Syringe 1 ml with 29 g × 12.7 mm needle		100	✓ E	B-D Ultra Fine
	1.36	10	_	
	(1.99)		-	B-D Ultra Fine
Syringe 1 ml with 31 g × 8 mm needle		100	✓ E	B-D Ultra Fine II
	1.36	10	_	
	(1.99)		E	B-D Ultra Fine II

Insulin Pumps

INSULIN PUMP WITH ALGORITHM – Special Authority see SA2367 below – Retail pharmacy

- a) Maximum of 1 dev per prescription
- b) Only on a prescription

c) Maximum of 1 insulin pump per patient each four year pe	eriod.		
Min basal rate 0.02 U/h	8,970.00	1	 mylife YpsoPump with CamAPS FX
Min basal rate 0.1 U/h	7,653.00	1	 ✓ Tandem t:slim X2 with Basal-IQ ✓ Tandem t:slim X2 with Control-IQ

SA2367 Special Authority for Subsidy

Initial application — (type 1 diabetes) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

16

1 Any of the following:

continued...

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1.1 The patient has type 1 diabetes; or
- 1.2 The patient has permanent neonatal diabetes or specific monogenic diabetes subtypes with insulin deficiency, considered by the treating endocrinologist as likely to benefit; or
- 1.3 The patient has Type 3c diabetes considered by the treating endocrinologist as likely to benefit (Type 3c diabetes includes insulin deficiency due to pancreatectomy, insulin deficiency secondary to cystic fibrosis or pancreatitis); or
- 1.4 The patient has atypical inherited forms of diabetes; and
- 2 Patient has been evaluated by a diabetes multidisciplinary team for their suitability for insulin pump therapy; and
- 3 In the opinion of the treating relevant practitioner the patient would benefit from an Automated Insulin Delivery (AID) system.

Renewal — (type 1 diabetes) from any relevant practitioner. Approvals valid for 6 months where the patient is continuing to derive benefit according to the treatment plan agreed at induction.

Insulin Pump Consumables

⇒SA2380 Special Authority for Subsidy

Initial application — (type 1 diabetes) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has type 1 diabetes; or
 - 1.2 The patient has permanent neonatal diabetes or specific monogenic diabetes subtypes with insulin deficiency, considered by the treating endocrinologist as likely to benefit; or
 - 1.3 The patient has Type 3c diabetes considered by the treating endocrinologist as likely to benefit (Type 3c diabetes includes insulin deficiency due to pancreatectomy, insulin deficiency secondary to cystic fibrosis or pancreatitis); or
 - 1.4 The patient has atypical inherited forms of diabetes; and
- 2 Patient has been evaluated by a diabetes multidisciplinary team for their suitability for insulin pump therapy; and
- 3 In the opinion of the treating relevant practitioner the patient would benefit from an Automated Insulin Delivery (AID) system.

Renewal — (type 1 diabetes) from any relevant practitioner. Approvals valid for 2 years where the patient is continuing to derive benefit according to the treatment plan agreed at induction.

INSULIN PUMP CARTRIDGE - Special Authority see SA2380 above - Retail pharmacy

- a) Maximum of 5 sets per prescription
- b) Only on a prescription
- c) Maximum of 19 packs of cartridge sets will be funded per year.

 Tandem Cartridge

	Subsidy (Manufacturer's P \$	rice) Sut Per	Fully Brand or osidised Generic Manufacturer
NSULIN PUMP INFUSION SET (STEEL CANNULA) – Specia a) Maximum of 5 set per prescription	I Authority see SA	2380 on the p	previous page – Retail pharmacy
b) Only on a prescription			
c) Maximum of 19 infusion sets will be funded per year.			
* 6 mm steel needle; 60 cm tubing × 10	130.00	1 OP	 MiniMed Sure-T MMT-864A
* 6 mm steel needle; 80 cm tubing × 10	130.00	1 OP	 MiniMed Sure-T MMT-866A
* 8 mm steel needle; 60 cm tubing × 10	130.00	1 OP	 MiniMed Sure-T MMT-874A
* 8 mm steel needle; 80 cm tubing × 10	130.00	1 OP	 MiniMed Sure-T MMT-876A
(MiniMed Sure-T MMT-876A 8 mm steel needle; 80 cm tubing > INSULIN PUMP INFUSION SET (STEEL CANNULA, STRAIGH page – Retail pharmacy a) Maximum of 5 sets per prescription			,
 a) Maximum of 5 sets per prescription b) Only on a prescription c) Maximum of 19 infusion sets will be funded per year. 			
5.5 mm steel cannula; straight insertion; 45 cm line × 10 wir 10 needles		1 OP	✓ mylife Orbit micro
5.5 mm steel needle; straight insertion; 60 cm line × 10 with			
10 needles		1 OP	 mylife Orbit micro
5.5 mm steel needle; straight insertion; 80 cm line × 10 with 10 needles			. mulife Orhit miere
10 needles ★ 8.5 mm steel needle; straight insertion; 60 cm line × 10 with		1 OP	 mylife Orbit micro
10 needles		1 OP	mylife Orbit micro
 8.5 mm steel needle; straight insertion; 80 cm line × 10 with 			
10 needles		1 OP	 mylife Orbit micro
* 6 mm steel cannula; straight insertion; 80 cm line × 10 with			,
10 needles		1 OP	✓ TruSteel
₭ 8 mm steel cannula; straight insertion; 80 cm line × 10 with			
10 needles		1 OP	 TruSteel
* 6 mm steel cannula; straight insertion; 60 cm line × 10 with		4.00	
10 needles		1 OP	 TruSteel
✤ 8 mm steel cannula; straight insertion; 60 cm line × 10 with			

		Subsidy (Manufacturer's Pric	e) Sub:	Fully Bran sidised Gene	
_		\$	Per		lfacturer
INS	SULIN PUMP INFUSION SET (TEFLON CANNULA) - Spec	cial Authority see SA2	380 on pag	e 17 – Retail p	harmacy
	 a) Maximum of 5 set per prescription b) Only on a prescription 				
	c) Maximum of 19 infusion sets will be funded per year.				
*	13 mm teflon needle, 60 cm tubing × 10		1 OP	 MiniMe MMT· 	d Silhouette 381A
*	17 mm teflon needle, 110 cm tubing × 10		1 OP	✓ MiniMe MMT·	d Silhouette 377A
*	17 mm teflon needle, 60 cm tubing × 10		1 OP	✓ MiniMe MMT·	d Silhouette 378A
*	6 mm teflon needle, 110 cm tubing × 10		1 OP		d Quick-Set
*	6 mm teflon needle, 45 cm blue tubing × 10	130.00	1 OP	✓ MiniMe MMT·	
*	6 mm teflon needle, 45 cm pink tubing × 10		1 OP	✓ MiniMe MMT·	
*	6 mm teflon needle, 60 cm blue tubing \times 10		1 OP	✓ MiniMe MMT·	
*	6 mm teflon needle, 60 cm pink tubing × 10		1 OP	✓ MiniMe MMT	d Mio
*	6 mm teflon needle, 60 cm tubing × 10		1 OP		d Quick-Set
*	6 mm teflon needle, 80 cm blue tubing		1 OP	✓ MiniMe MMT·	
*	6 mm teflon needle, 80 cm clear tubing × 10		1 OP	✓ MiniMe MMT	d Mio
*	6 mm teflon needle, 80 cm pink tubing × 10		1 OP	✓ MiniMe MMT	d Mio
*	9 mm teflon needle, 110 cm tubing × 10		1 OP		d Quick-Set
*	9 mm teflon needle, 60 cm tubing × 10		1 OP		d Quick-Set
*	9 mm teflon needle, 80 cm clear tubing × 10		1 OP	✓ MiniMe MMT·	d Mio

(MiniMed Silhouette MMT-381A 13 mm teflon needle, 60 cm tubing × 10 to be delisted 1 October 2026) (MiniMed Silhouette MMT-377A 17 mm teflon needle, 110 cm tubing × 10 to be delisted 1 October 2026) (MiniMed Silhouette MMT-378A 17 mm teflon needle, 60 cm tubing × 10 to be delisted 1 October 2026) (MiniMed Quick-Set MMT-398A 6 mm teflon needle, 10 cm tubing × 10 to be delisted 1 October 2026) (MiniMed Mio MMT-941A 6 mm teflon needle, 45 cm blue tubing × 10 to be delisted 1 October 2026) (MiniMed Mio MMT-921A 6 mm teflon needle, 45 cm pink tubing × 10 to be delisted 1 October 2026) (MiniMed Mio MMT-921A 6 mm teflon needle, 60 cm blue tubing × 10 to be delisted 1 October 2026) (MiniMed Mio MMT-923A 6 mm teflon needle, 60 cm blue tubing × 10 to be delisted 1 October 2026) (MiniMed Mio MMT-933A 6 mm teflon needle, 60 cm blue tubing × 10 to be delisted 1 October 2026) (MiniMed Mio MMT-943A 6 mm teflon needle, 60 cm tubing × 10 to be delisted 1 October 2026) (MiniMed Mio MMT-945A 6 mm teflon needle, 80 cm clear tubing × 10 to be delisted 1 October 2026) (MiniMed Mio MMT-945A 6 mm teflon needle, 80 cm clear tubing × 10 to be delisted 1 October 2026) (MiniMed Mio MMT-945A 6 mm teflon needle, 80 cm pink tubing × 10 to be delisted 1 October 2026) (MiniMed Mio MMT-945A 6 mm teflon needle, 80 cm pink tubing × 10 to be delisted 1 October 2026) (MiniMed Mio MMT-945A 6 mm teflon needle, 80 cm pink tubing × 10 to be delisted 1 October 2026) (MiniMed Quick-Set MMT-396A 9 mm teflon needle, 60 cm tubing × 10 to be delisted 1 October 2026) (MiniMed Quick-Set MMT-397A 9 mm teflon needle, 60 cm tubing × 10 to be delisted 1 October 2026) (MiniMed Quick-Set MMT-397A 9 mm teflon needle, 60 cm tubing × 10 to be delisted 1 October 2026) (MiniMed Mio MMT-975A 9 mm teflon needle, 80 cm clear tubing × 10 to be delisted 1 October 2026)

	Subsidy (Manufacturer's Pri	iaa) Cub		rand or eneric
	(Manulactuler's Fil	Per Sub		anufacturer
INSULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLI	E INSERTION WITH	INSERTION	DEVICE)	- Special Authority see
SA2380 on page 17 – Retail pharmacy				
 a) Maximum of 5 sets per prescription b) Only on a prescription 				
c) Maximum of 19 infusion sets will be funded per year.				
* 13 mm teflon cannula; angle insertion; insertion device; 11	0 cm			
line × 10 with 10 needles		1 OP	🗸 Auto	Soft 30
* 13 mm teflon cannula; angle insertion; insertion device; 60				
line × 10 with 10 needles		1 OP	 Auto 	
INSULIN PUMP INFUSION SET (TEFLON CANNULA, FLEXI	BLE INSERTION WI	TH INSERTI	ON DEVICE	 Special Authority
see SA2380 on page 17 – Retail pharmacy a) Maximum of 5 set per prescription				
b) Only on a prescription				
c) Maximum of 19 infusion sets will be funded per year.				
* 6 mm teflon cannula; flexible insertion; insertion device; 4				
line × 10 with 10 needles		1 OP	🗸 myli	fe Inset soft
* 6 mm teflon cannula; flexible insertion; insertion device; 60 line with integrated inserter × 10 with 10 needles		1 OP	🖌 myli	fe Inset soft
 6 mm teflon cannula; flexible insertion; insertion device; 80 		101	• myn	
line × 10 with 10 needles		1 OP	🖌 myli	fe Inset soft
* 9 mm teflon cannula; flexible insertion; insertion device; 60) cm			
line × 10 with 10 needles		1 OP	🖌 myli	fe Inset soft
9 mm teflon cannula; flexible insertion; insertion device; 80		1 OP	. muli	fo lucat caft
line × 10 with 10 needles		-	•	fe Inset soft
INSULIN PUMP INFUSION SET (TEFLON CANNULA, STRAI see SA2380 on page 17 – Retail pharmacy	GHT INSERTION W	II H INSERI	ION DEVIC	 E) – Special Authority
a) Maximum of 5 sets per prescription				
b) Only on a prescription				
c) Maximum of 19 infusion sets will be funded per year.				
* 6 mm teflon cannula; straight insertion; insertion device;	100.00	1.00		0.400
110 cm line × 10 with 10 needles		1 OP	✓ Auto	Soft 90
line × 10 with 10 needles		1 OP	🗸 Auto	Soft 90
* 9 mm teflon cannula; straight insertion; insertion device;		-		
110 cm line × 10 with 10 needles		1 OP	🗸 Auto	Soft 90
* 9 mm teflon cannula; straight insertion; insertion device; 6		4.00		o <i>"</i> oo
line × 10 with 10 needles		1 OP	 Auto 	
INSULIN PUMP INFUSION SET (TEFLON CANNULA, VARIA	BLE INSERTION) -	- Special Aut	hority see S	A2380 on page 17 –
Retail pharmacy a) Maximum of 5 set per prescription				
b) Only on a prescription				
c) Maximum of 19 infusion sets will be funded per year.				
* 13 mm teflon cannula; variable insertion; 60 cm line × 10 v		4.05	• • • •	- <i>"</i>
10 needles		1 OP	 Varis 	δοπ

20

()	Subsidy /anufacturer's Price) \$	S Per	Fully ubsidised	Brand or Generic Manufacturer
INSULIN PUMP RESERVOIR – Special Authority see SA2380 on	age 17 – Retail ph	armac	у	
 a) Maximum of 9 sets per prescription b) Only on a prescription c) Maximum of 36 packs of resevoir sets will be funded per year 	ar			
 * 10 × 1.6 ml glass reservoir for YpsoPump 		1 OP	√ r	nylife YpsoPump Reservoir
* 10 × luer lock conversion cartridges 1.8 ml for Paradigm pumps	50.00	1 OP	✓	ADR Cartridge 1.8
Cartridge for 7 series pump; 3.0 ml × 10	50.00	1 OP	✓ 1	/iniMed 3.0 Reservoir MMT-332A
(ADR Cartridge 1.8 10 × luer lock conversion cartridges 1.8 ml for F (MiniMed 3.0 Reservoir MMT-332A Cartridge for 7 series pump; 3.0	0 / /			,
Continuous Glucose Monitor				

CO	NTINUOUS GLUCOSE MONITOR (INTEROPERABLE) – Special Authority	see SA2371 be	olow – Retail pharmacy
	Only on a prescription		
*	Sensor (9) and transmitter (Dexcom G6) – Maximum of 1 dev		
	per prescription	1 OP	Dexcom G6
	Maximum of 5 dev will be funded per year.		
*	Sensor (Dexcom G7) – Maximum of 9 dev per prescription	1	 Dexcom G7
	Maximum of 40 dev will be funded per year.		
*	Sensor (Freestyle Libre 3 Plus) – Maximum of 6 dev per		
	prescription	1	 Freestyle Libre
			3 Plus
*	, , , , , , , , , , , , , , , , , , , ,	1	

Maximum of 28 dev will be funded per year.

⇒SA2371 Special Authority for Subsidy

Initial application — (type 1 diabetes) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has type 1 diabetes; or
 - 1.2 The patient has permanent neonatal diabetes or specific monogenic diabetes subtypes with insulin deficiency, considered by the treating endocrinologist or relevant secondary health care professional as practicable, as likely to benefit; or
 - 1.3 The patient has Type 3c diabetes considered by the treating endocrinologist or relevant secondary health care professional as practicable, as likely to benefit (Type 3c diabetes includes insulin deficiency due to pancreatectomy, insulin deficiency secondary to cystic fibrosis or pancreatitis); or
 - 1.4 The patient has atypical inherited forms of diabetes; and
- 2 In the opinion of the treating relevant practitioner the patient would benefit from an Automated Insulin Delivery (AID) system.

Renewal — (type 1 diabetes) from any relevant practitioner. Approvals valid for 2 years where the patient is continuing to derive benefit according to the treatment plan agreed at induction.

CONTINUOUS GLUCOSE MONITOR (STANDALONE) - Special Authority see SA2370 on the next page - Retail pharma	су
Only on a prescription	

*	Sensor (Dexcom ONE+) – Maximum of 9 dev per prescription81.00	1	Dexcom ONE+
	Maximum of 40 dev will be funded per year.		
*	Sensor (Freestyle Libre 2) – Maximum of 7 dev per prescription92.83	1	Freestyle Libre 2
	Maximum of 29 dev will be funded per year.		

*Three months or six months, as applicable, dispensed all-at-once

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy (Manufacturer's Price \$	e) S Per	Fully Subsidised	Brand or Generic Manufacturer	
--	-------------	---------------------	-------------------------------------	--

⇒SA2370 Special Authority for Subsidy

Initial application — (type 1 diabetes) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 The patient has type 1 diabetes; or
- 2 The patient has permanent neonatal diabetes or specific monogenic diabetes subtypes with insulin deficiency, considered by the treating endocrinologist or relevant secondary health care professional as practicable, as likely to benefit; or
- 3 The patient has Type 3c diabetes considered by the treating endocrinologist or relevant secondary health care professional as practicable, as likely to benefit (Type 3c diabetes includes insulin deficiency due to pancreatectomy, insulin deficiency secondary to cystic fibrosis or pancreatitis); or
- 4 The patient has atypical inherited forms of diabetes.

Renewal — (type 1 diabetes) from any relevant practitioner. Approvals valid for 2 years where the patient is continuing to derive benefit according to the treatment plan agreed at induction.

Digestives Including Enzymes

PANCREATIC ENZYME

Cap pancreatin 150 mg (amylase 8,000 Ph Eur U, lipase 10,000 Ph Eur U, total protease 600 Ph Eur U)	34.93	100	✓ Creon 10000
Cap pancreatin 300 mg (amylase 18,000 Ph Eur U, lipase 25,000 Ph Eur U, total protease 1,000 Ph Eur U)	94.38	100	 Creon 25000
Modified release granules pancreatin 60.12 mg (amylase 3,600 Ph Eur U, lipase 5,000 Ph Eur U, protease 200 Ph Eur U)	34 93	20 g OP	✓ Creon Micro
URSODEOXYCHOLIC ACID – Special Authority see SA1739 below Cap 250 mg	– Retail pha	U	✓ Ursosan

➡SA1739 Special Authority for Subsidy

Initial application — (Alagille syndrome or progressive familial intrahepatic cholestasis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Either:

- 1 Patient has been diagnosed with Alagille syndrome; or
- 2 Patient has progressive familial intrahepatic cholestasis.

Initial application — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic severe drug induced cholestatic liver injury; and
- 2 Cholestatic liver injury not due to Total Parenteral Nutrition (TPN) use in adults; and
- 3 Treatment with ursodeoxycholic acid may prevent hospital admission or reduce duration of stay.

Initial application — (Primary biliary cholangitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Primary biliary cholangitis confirmed by antimitochondrial antibody titre (AMA) > 1:80, and raised cholestatic liver enzymes with or without raised serum IgM or, if AMA is negative, by liver biopsy; and
- 2 Patient not requiring a liver transplant (bilirubin > 100 umol/l; decompensated cirrhosis).

Initial application — (Pregnancy) from any relevant practitioner. Approvals valid for 6 months where the patient diagnosed with cholestasis of pregnancy.

Initial application — (Haematological Transplant) from any relevant practitioner. Approvals valid for 6 months for applications

continued...

	Subsidy		ully	Brand or
(Ma	(Manufacturer's Price)		sed	Generic
	\$	Per	1	Manufacturer

continued...

meeting the following criteria:

Both:

- 1 Patient at risk of veno-occlusive disease or has hepatic impairment and is undergoing conditioning treatment prior to allogenic stem cell or bone marrow transplantation; and
- 2 Treatment for up to 13 weeks.

Initial application — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Paediatric patient has developed abnormal liver function as indicated on testing which is likely to be induced by Total Parenteral Nutrition (TPN); and
- 2 Liver function has not improved with modifying the TPN composition.

Renewal — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 6 months where the patient continues to benefit from treatment.

Renewal — (Pregnancy/Primary biliary cholangitis) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months where the paediatric patient continues to require TPN and who is benefiting from treatment, defined as a sustained improvement in bilirubin levels.

Laxatives

Bulk-forming Agents

ISPAGHULA (PSYLLIUM) HUSK – Only on a prescription * Powder for oral soln	500 g OP	✓ Konsyl-D
Faecal Softeners		
DOCUSATE SODIUM – Only on a prescription		
* Tab 50 mg	100	 Coloxyl
* Tab 120 mg4.98	100	✓ Coloxyl
DOCUSATE SODIUM WITH SENNOSIDES		
Tab 50 mg with sennosides 8 mg	200	 Laxsol
POLOXAMER – Only on a prescription Not funded for use in the ear.		
* Oral drops 10%4.17	30 ml OP	✓ <u>Coloxyl</u>
Opioid Receptor Antagonists - Peripheral		
opioia neocotor Antagonioto - i cripiteral		

➡SA1691 Special Authority for Subsidy

Initial application — (Opioid induced constipation) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient is receiving palliative care; and
- 2 Either:
 - 2.1 Oral and rectal treatments for opioid induced constipation are ineffective; or
 - 2.2 Oral and rectal treatments for opioid induced constipation are unable to be tolerated.

	Subsidy		Fully	Brand or
	(Manufacturer's Price \$	e) Per	Subsidised	Generic Manufacturer
	\$	rei	•	Manulaciulei
Osmotic Laxatives				
GLYCEROL				
Suppos 2.8/4.0 g – Only on a prescription	10.39	20	~	Lax-suppositories Glycerol
ACTULOSE – Only on a prescription				
Oral liq 10 g per 15 ml	3.61	500 m	nl 🖌	Laevolac
ACROGOL 3350 WITH POTASSIUM CHLORIDE, SODIUM B	CARBONATE AND	SODIL	JM CHLOF	RIDE
Powder for oral soln 13.125 g with potassium chloride 46.6 r	ng,			
sodium bicarbonate 178.5 mg and sodium chloride 350.	7 mg8.50	30	1	Molaxole
SODIUM ACID PHOSPHATE – Only on a prescription				
Enema 16% with sodium phosphate 8%	2.50	1	1	Fleet Phosphate
				Enema
SODIUM CITRATE WITH SODIUM LAURYL SULPHOACETATE	E – Only on a presc	ription		
Enema 90 mg with sodium lauryl sulphoacetate 9 mg per ml	,			
5 ml	35.89	50	✓	Micolette
Stimulant Laxatives				
ISACODYL – Only on a prescription				
🗧 Tab 5 mg	5.80	200	1	Bisacodyl Viatris
 Suppos 10 mg 	4.14	10	1	Lax-Suppositories
ENNA – Only on a prescription				
 Tab, standardised 	2.17	100		
	(8.21)			Senokot
	0.43	20		
	(2.06)			Senokot
ODIUM PICOSULFATE – Special Authority see SA2053 below	v – Retail pharmacy			
Oral soln 7.5 mg per ml		30 ml C		Dulcolax SP Drop

Both:

1 The patient is a child with problematic constipation despite an adequate trial of other oral pharmacotherapies including macrogol where practicable; and

2 The patient would otherwise require a high-volume bowel cleansing preparation or hospital admission.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Metabolic Disorder Agents			
ALGLUCOSIDASE ALFA – Special Authority see SA1986 below – Retai Inj 50 mg vial		1 •	Myozyme
SA1986 Special Authority for Subsidy Initial application only from a metabolic physician. Approvals valid for 1 All of the following:	2 months for ap	plications r	meeting the following criteria:
1 The patient is aged up to 24 months at the time of initial application	n and has been	diagnosed	I with infantile Pompe disease;

continued...

and

24

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

- 2 Any of the following:
 - 2.1 Diagnosis confirmed by documented deficiency of acid alpha-glucosidase by prenatal diagnosis using chorionic villus biopsies and/or cultured amniotic cells; or
 - 2.2 Documented deficiency of acid alpha-glucosidase, and urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides; or
 - 2.3 Documented deficiency of acid alpha-glucosidase, and documented molecular genetic testing indicating a disease-causing mutation in the acid alpha-glucosidase gene (GAA gene); or
 - 2.4 Documented urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides, and molecular genetic testing indicating a disease-causing mutation in the GAA gene; and
- 3 Patient has not required long-term invasive ventilation for respiratory failure prior to starting enzyme replacement therapy (ERT); and
- 4 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by ERT or might be reasonably expected to compromise a response to ERT; and
- 5 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks.

Renewal only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks; and
- 3 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 4 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by ERT; and
- 5 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT; and
- 6 There is no evidence of life threatening progression of respiratory disease as evidenced by the needed for > 14 days of invasive ventilation; and
- 7 There is no evidence of new or progressive cardiomyopathy.

ARGININE - Special Authority see SA2042 below - Retail pharmacy

Tab 1,000 mg	 90	 Clinicians
Cap 500 mg	50	 Solgar
Powder	400 g	 Biomed

⇒SA2042 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to arginine supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to arginine supplementation; and

- 2 The treatment remains appropriate and the patient is benefiting from treatment.
- BETAINE Special Authority see SA1987 below Retail pharmacy

......575.00 180 g OP

Cystadane

⇒SA1987 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has a confirmed diagnosis of homocystinuria; and
- 2 Any of the following:
 - 2.1 A cystathionine beta-synthase (CBS) deficiency; or

continued...

Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Full Subsidise Per •	d Generic
continued			
2.2 A 5,10-methylene-tetrahydrofolate reductase (MT	HFR) deficiency; or		
2.3 A disorder of intracellular cobalamin metabolism;			
3 An appropriate homocysteine level has not been achieve	ed despite a sufficient to	ial of appropri	ate vitamin supplementatior
Renewal only from a metabolic physician. Approvals valid for 1	2 months where the tre	eatment remai	ns appropriate and the
patient is benefiting from treatment.			
COENZYME Q10 - Special Authority see SA2039 below - Ret	ail pharmacy		
Cap 120 mg	CBS		' Solgar
Cap 160 mg	CBS	60 🖌	Go Healthy
SA2039 Special Authority for Subsidy			
nitial application only from a metabolic physician. Approvals	valid for 6 months when	e patient has	a suspected inborn error of
netabolism that may respond to coenzyme Q10 supplementation			
Renewal only from a metabolic physician. Approvals valid for 2	4 months for applicatio	ns meeting th	e following criteria:
Both:			
 The patient has a confirmed diagnosis of an inborn error 	of metabolism that res	ponds to coen	zyme Q10 supplementation
and	<i>.</i>		
2 The treatment remains appropriate and the patient is be	nefiting from treatment.		
GALSULFASE – Special Authority see SA1988 below – Retail	pharmacy		
Inj 1 mg per ml, 5 ml vial	2,234.00	1 🗸	Naglazyme
SA1988 Special Authority for Subsidy			
nitial application only from a metabolic physician. Approvals	valid for 12 months for	applications m	neeting the following criteria:
Both:			
1 The patient has been diagnosed with mucopolysaccharic	losis VI; and		
2 Either:			
2.1 Diagnosis confirmed by demonstration of N-acety		atase (arylsulf	atase B) deficiency by eithe
enzyme activity assay in leukocytes or skin fibrob			
2.2 Detection of two disease causing mutations and	patient has a sibling wh	o is known to	have mucopolysaccharidosi
VI.			
Renewal only from a metabolic physician. Approvals valid for 1	2 months for applicatio	ns meeting th	e following criteria:
All of the following:			
1 The treatment remains appropriate for the patient and th			
2 Patient has not had severe infusion-related adverse read	tions which were not p	reventable by	appropriate pre-medication
and/or adjustment of infusion rates; and	ara diagona whara tha	long torm are	anaaia ia unlikalu ta ha
 Patient has not developed another life threatening or sev influenced by Enzyme Replacement Therapy (ERT); and 		long term pro	gnosis is unlikely to be
4 Patient has not developed another medical condition that		expected to co	moromise a response to
ERT.	t might reasonably be t		
DURSULFASE – Special Authority see SA1623 below – Retai	nharmacy		
Inj 2 mg per ml, 3 ml vial		1 🗸	Elaprase
SA1623 Special Authority for Subsidy		•	
nitial application only from a metabolic physician. Approvals	valid for 24 weeks for a	nnlications me	eting the following criteria:
All of the following:			soung the following ontella.
5	mu aanalu aaaha iidaai	a II), and	
1 The patient has been diagnosed with Hunter Syndrome	mucopolysaccharidosi	s ii); and	
0 Eithor			
2 Either:			
 2 Either: 2.1 Diagnosis confirmed by demonstration of idurona assay in cultured skin fibroblasts; or 	te 2-sulfatase deficienc	y in white blo	od cells by either enzyme

26

(Note a late o		Eul	ь. г	Dueural eu
	Subsidy acturer's Price)	S	Ful Subsidise		Brand or Generic
``````````````````````````````````````	\$	Per		1	Manufacturer
continued 2.2 Detection of a disease causing mutation in the iduronate 2-	-cultataco dor	no: and	4		
<ol> <li>Patient is going to proceed with a haematopoietic stem cell transp idursulfase would be bridging treatment to transplant; and</li> </ol>				3 mo	nths and treatment wit
<ul> <li>Patient has not required long-term invasive ventilation for respirate (ERT); and</li> </ul>	ory failure prio	or to st	arting E	nzym	e Replacement Thera
<ul> <li>5 Idursulfase to be administered for a total of 24 weeks (equivalent is greater than 0.5 mg/kg every week.</li> </ul>	to 12 weeks p	re- an	d 12 we	eks p	ost-HSCT) at doses n
LARONIDASE – Special Authority see SA1695 below – Retail pharmacy Inj 100 U per ml, 5 ml vial		1		Ald	urazyme
⇒SA1695 Special Authority for Subsidy	0.110	•			
<b>Initial application</b> only from a metabolic physician. Approvals valid for 2 All of the following:	4 weeks for a	pplica	tions m	eeting	the following criteria:
1 The patient has been diagnosed with Hurler Syndrome (mucopoly 2 Either:	sacchardosis	I-H); a	and		
<ol> <li>Diagnosis confirmed by demonstration of alpha-L-iduronida assay in cultured skin fibroblasts; or</li> </ol>	ase deficiency	' in wh	ite bloo	d cells	s by either enzyme
2.2 Detection of two disease causing mutations in the alpha-L- to have Hurler syndrome; and	iduronidase g	ene a	nd patie	ent has	s a sibling who is know
3 Patient is going to proceed with a haematopoietic stem cell transp laronidase would be bridging treatment to transplant; and	lant (HSCT) w	vithin t	he next	3 mo	nths and treatment wit
<ul> <li>Patient has not required long-term invasive ventilation for respirate (ERT); and</li> </ul>	ory failure prio	or to st	arting E	inzym	e Replacement Thera
5 Laronidase to be administered for a total of 24 weeks (equivalent than 100 units/kg every week.	to 12 weeks p	ore- an	d 12 po	st-HS	CT) at doses no great
EVOCARNITINE - Special Authority see SA2040 below - Retail pharm	acy				
Tab 500 mgC		30		Sol	•
Cap 250 mgC	BS	30		Sol	•
Cap 500 mgC		60		Bal	
		300			abolics
Oral lig 1 g per 10 ml	BS 1	18 ml	✓		nitor S29
era ind i giper re in this sector (					
			•	Nov	/itium Sugar
			•		vitium Sugar ree S29
Oral liq 500 mg per 10 mlC	BS 3	00 ml			ree S29
Oral liq 500 mg per 10 mlC	XBS 3	00 ml		F	ree S29
Oral liq 500 mg per 10 mlC SA2040 Special Authority for Subsidy nitial application only from a metabolic physician. Approvals valid for 6			v	Fi Bala	ree S29 ance
Oral liq 500 mg per 10 mlC → SA2040 Special Authority for Subsidy nitial application only from a metabolic physician. Approvals valid for 6 netabolism that may respond to carnitine supplementation. Renewal only from a metabolic physician. Approvals valid for 24 months	6 months wher	re patie	✔ ent has	Fi Bali a sus	ree S23 ance pected inborn error of
Oral liq 500 mg per 10 mlC SA2040 Special Authority for Subsidy nitial application only from a metabolic physician. Approvals valid for 6 netabolism that may respond to carnitine supplementation. Renewal only from a metabolic physician. Approvals valid for 24 months	o months wher for applicatio olism that resp	re patie Ins me ponds	ent has eting th	Fi Bali a sus ne follo	ree \$29 ance pected inborn error of owing criteria:
Oral liq 500 mg per 10 ml	on the test of	re patie Ins me ponds	ent has eting th	Fi Bali a sus ne follo	ree \$29 ance pected inborn error of owing criteria:
Oral liq 500 mg per 10 ml	or months wher for applicatio olism that respon treatment. pharmacy	re patie Ins me ponds	ent has eting th to carn	Fi Bala a sus ie follo itine s Cou Pur Vi	ree S29 ance pected inborn error of owing criteria: supplementation; and untry Life itan's Pride itamin
Oral liq 500 mg per 10 ml	6 months wher 1 for applicatio 1 olism that resp 1 om treatment. 1 pharmacy 2 BS	re patie Ins me ponds	ent has eting th to carn	Fi Bala a sus ie follo itine s Cou Pur Vi	ree S29 ance pected inborn error of owing criteria: supplementation; and untry Life itan's Pride itamin -2 100 mg S29

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per ✓	Manufacturer
Ŷ	101	

#### ⇒SA2041 Special Authority for Subsidy

Initial application only from a metabolic physician or neurologist. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to riboflavin supplementation.

Renewal only from a metabolic physician or neurologist. Approvals valid for 24 months for applications meeting the following criteria:

Both:

1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to riboflavin supplementation; and

2 The treatment remains appropriate and the patient is benefiting from treatment.

#### ➡SA1989 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 1 month for applications meeting the following criteria: All of the following:

- 1 Patient has phenylketonuria (PKU) and is pregnant or actively planning to become pregnant; and
- 2 Treatment with sapropterin is required to support management of PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

Renewal only from a metabolic physician or any relevant practitioner on the recommendation of a metabolic physician.

Approvals valid for 12 months for applications meeting the following criteria:

- All of the following:
  - 1 Either:
    - 1.1 Following the initial one-month approval, the patient has demonstrated an adequate response to a 2 to 4 week trial of sapropterin with a clinically appropriate reduction in phenylalanine levels to support management of PKU during pregnancy; or
    - 1.2 On subsequent renewal applications, the patient has previously demonstrated response to treatment with sapropterin and maintained adequate phenylalanine levels to support management of PKU during pregnancy; and
  - 2 Any of the following:
    - 2.1 Patient continues to be pregnant and treatment with sapropterin will not continue after delivery; or
    - 2.2 Patient is actively planning a pregnancy and this is the first renewal for treatment with sapropterin; or
    - 2.3 Treatment with sapropterin is required for a second or subsequent pregnancy to support management of their PKU during pregnancy; and
  - 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
  - 4 Sapropterin to be used alone or in combination with PKU dietary management; and
  - 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

SODIUM BENZOATE - Special Authority see SA1599 below - Retail pharmacy

Soln 100 mg per ml ......CBS 100 ml 🖌 Amzoate 😒

#### ⇒SA1599 Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

SODIUM PHENYLBUTYRATE - Special Authority see SA1990 on the next page -	- Retail pharmacy	/
Grans 483 mg per g2,016.00	174 g OP 🔹	Pheburane

28

	Subsidy (Manufacturer's Price) \$	Subsi Per	Fully dised	Brand or Generic Manufacturer
SA1990 Special Authority for Subsidy				
nitial application only from a metabolic physician. Approvals				•
ycle disorder involving a deficiency of carbamylphosphate syn vnthetase.	thetase, ornithine trans	scarbamyla	se or a	argininosuccinate
<b>Renewal</b> only from a metabolic physician. Approvals valid for atient is benefiting from treatment.	2 months where the tr	reatment re	mains	appropriate and the
AURINE - Special Authority see SA2043 below - Retail phan	macy			
Cap 500 mg	CBS	50	✓ S	Solgar
Cap 1,000 mg		90	✓ L	ife Extension
Powder	CBS	300 g	✓ L	ife Extension
SA2043 Special Authority for Subsidy		•		
<b>itial application</b> only from a metabolic physician. Approvals	valid for 6 months whe	ere patient l	nas a s	suspected specific
hitochondrial disorder that may respond taurine supplementation				
<b>Renewal</b> only from a metabolic physician Approvals valid for 2		ons meetin	a the f	ollowing criteria:

**Renewal** only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has confirmed diagnosis of a specific mitochondrial disorder which responds to taurine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

TRIENTINE – Special Authority see SA2324 below – Retail pharmacy

Cap 250 mg2,022.0	0 100	<ul> <li>Trientine Waymade</li> </ul>
-------------------	-------	---------------------------------------

#### ➡SA2324 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Wilson disease; and
- 2 Treatment with D-penicillamine has been trialled and discontinued because the person has experienced intolerable side effects or has not received sufficient benefit; and
- 3 Treatment with zinc has been trailled and discontinued because the person has experienced intolerable side effects or has not received sufficient benefit, or zinc is considered clinically inappropriate as the person has symptomatic liver disease and requires copper chelation.

### **Gaucher's Disease**

TALIGLUCERASE ALFA – Special Authority see SA2137 below – Retail pharmacy

Inj 200 unit vial...... 1,072.00 1 🖌 Elelyso

#### ⇒SA2137 Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has a diagnosis of symptomatic type 1 or type 3* Gaucher disease confirmed by the demonstration of specific deficiency of glucocerebrosidase in leukocytes or cultured skin fibroblasts, and genotypic analysis; and
- 2 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by
- enzyme replacement therapy (ERT) or the disease might be reasonably expected to compromise a response to ERT; and 3 Any of the following:
  - 3.1 Patient has haematological complications of Gaucher disease; or
  - 3.2 Patient has skeletal complications of Gaucher disease; or
  - 3.3 Patient has significant liver dysfunction or hepatomegaly attributable to Gaucher disease; or
  - 3.4 Patient has reduced vital capacity from clinically significant or progressive pulmonary disease due to Gaucher

continued...

AThree months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

#### continued...

- disease; or
- 3.5 Patient is a child and has experienced growth failure with significant decrease in percentile linear growth over a 6-12 month period; and
- 4 Taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units).

Note: Indication marked with * is an unapproved indication

**Renewal** only from a metabolic physician or any relevant practitioner on the recommendation of a metabolic physician. Approvals valid for 3 years for applications meeting the following criteria:

- All of the following:
  - 1 Patient has demonstrated a symptomatic improvement and has maintained improvements in the main symptom or symptoms for which therapy was started; and
  - 2 Patient has demonstrated a clinically objective improvement or no deterioration in haemoglobin levels, platelet counts and liver and spleen size; and
  - 3 Radiological (MRI) signs of bone activity performed at two years since initiation of treatment, and five yearly thereafter, demonstrate no deterioration shown by the MRI, compared with MRI taken immediately prior to commencement of therapy or adjusted dose; and
  - 4 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT; and
  - 5 Patient is adherent with regular treatment and taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units).

### Mouth and Throat

#### Agents Used in Mouth Ulceration

Agento obca in mouth officiation		
BENZYDAMINE HYDROCHLORIDE Soln 0.15% – Higher subsidy of \$22.60 per 500 ml with Endorsement	500 ml	Difflam
Additional subsidy by endorsement for a patient who has oral mucositis a prescription is endorsed accordingly.	as a result of tr	eatment for cancer, and the
CARMELLOSE SODIUM WITH GELATIN AND PECTIN		
Paste	56 g OP 15 g OP	<ul> <li>Stomahesive</li> </ul>
(7.90)		Orabase
1.52	5 g OP	Orabase
(3.60) Powder	28 g OP	Stomahesive
TRIAMCINOLONE ACETONIDE		
Paste 0.1%	5 g OP	✓ Kenalog in Orabase
Oropharyngeal Anti-infectives		
AMPHOTERICIN B		
Lozenges 10 mg5.86	20	<ul> <li>Fungilin</li> </ul>
MICONAZOLE		
Oral gel 20 mg per g5.19	40 g OP	<ul> <li>Decozol</li> </ul>
NYSTATIN		
Oral liq 100,000 u per ml2.22	24 ml OP	✓ Nilstat

30 fully subsidised Principal Supply S29 Unapproved medicine supplied under Section 29 Sole Subsidised Supply

	Subsidy (Manufacturer's Price	Per	Fully Subsidised	
	\$	Fei		Manulaciurei
Vitamins				
Vitamin B				
HYDROXOCOBALAMIN				
Inj 1 mg per ml, 1 ml ampoule – Up to 6 inj available on	a PSO2.46	3		Cobal-B12 S29 Vita-B12
	3.95			Hydroxocobalamin Panpharma
	4.10	5		Cobalin-H S29 Neo-Cytamen S29 S29
	8.20	10	1	Vitarubin Depot Injection S29
(Cobal-B12 529 Inj 1 mg per ml, 1 ml ampoule to be deliste (Vita-B12 Inj 1 mg per ml, 1 ml ampoule to be delisted 1 July (Cobalin-H 529 Inj 1 mg per ml, 1 ml ampoule to be delisted	2025) 1 1 July 2025)			
(Neo-Cytamen S29 s29 Inj 1 mg per ml, 1 ml ampoule to be (Vitarubin Depot Injection s29 Inj 1 mg per ml, 1 ml ampoul		105)		
PYRIDOXINE HYDROCHLORIDE	e lo be delisied i July 20	125)		
a) No more than 100 mg per dose				
<ul> <li>b) Only on a prescription</li> <li>* Tab 25 mg - No patient co-payment payable</li> <li>* Tab 50 mg</li> </ul>		90 500		<u>Vitamin B6 25</u> Pyridoxine
THIAMINE HYDROCHLORIDE – Only on a prescription				multichem
* Tab 50 mg	4.65	100	✓	Thiamine multichem
VITAMIN B COMPLEX * Tab, strong, BPC	11.25	500	1	Bplex
Vitamin C				
ASCORBIC ACID a) No more than 100 mg per dose				
<ul> <li>b) Only on a prescription</li> <li>★ Tab 100 mg</li> </ul>		500	1	<u>Cvite</u>
Vitamin D				
ALFACALCIDOL * Cap 0.25 mcg		100		One-Alpha
Cap 1 mcg     Oral drops 2 mcg per ml (One-Alpha S29 s29 Cap 0.25 mcg to be delisted 1 July 20.		100 0 ml C	<ul> <li>Image: A start of the start of</li></ul>	One-Alpha S29 S23 One-Alpha One-Alpha

	Subsidy		Fully Brand or
	(Manufacturer's P \$	rice) Subs Per	sidised Generic Manufacturer
CALCITRIOL	Ŷ	101	- manufacturor
* Cap 0.25 mcg	7.89	100	<ul> <li><u>Calcitriol-AFT</u></li> <li>Calcitriol-AFT</li> <li>S29 \$23</li> </ul>
* Cap 0.5 mcg	13.68	100	<ul> <li><u>Calcitriol-AFT</u></li> <li>Calcitriol-AFT</li> <li>S29 \$29</li> </ul>
COLECALCIFEROL * Cap 1.25 mg (50,000 iu) – Maximum of 12 cap per prescri * Oral liq 188 mcg per ml (7,500 iu per ml)		12 5 ml OP	✓ <u>Vit.D3</u> ✓ Clinicians
Multivitamin Preparations			
MULTIVITAMIN RENAL – Special Authority see SA1546 below * Cap		y 30	<ul> <li>Clinicians Renal Vit</li> </ul>
<ul> <li>SA1546 Special Authority for Subsidy         Initial application from any relevant practitioner. Approvals vathe following criteria:         Either:         <ol> <li>The patient has chronic kidney disease and is receiving</li> <li>The patient has chronic kidney disease grade 5, defined 15 ml/min/1.73 m² body surface area (BSA).</li> </ol> </li> </ul>	either peritoneal d as patient with an	ialysis or haem	nodialysis; or
MULTIVITAMINS – Special Authority see SA1036 below – Ret * Powder		200 g OP	✓ Paediatric Seravit
<ul> <li>SA1036 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals va inborn errors of metabolism.</li> <li>Renewal from any relevant practitioner. Approvals valid withou approval for multivitamins.</li> </ul>			
VITAMINS			
* Tab (BPC cap strength)		1,000	✓ <u>Mvite</u>
* Cap (fat soluble vitamins A, D, E, K) – Special Authority se SA1720 below – Retail pharmacy		60	<ul> <li>Vitabdeck</li> </ul>
SA1720 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals va the following criteria:	lid without further	renewal unless	s notified for applications meeting

Any of the following:

- 1 Patient has cystic fibrosis with pancreatic insufficiency; or
- 2 Patient is an infant or child with liver disease or short gut syndrome; or
- 3 Patient has severe malabsorption syndrome.

	Subsidy (Manufacturer's Price	e)	Fully Subsidised	Brand or Generic
	\$	Per	1	Manufacturer
Minerals				
Calcium				
CALCIUM CARBONATE				
* Tab 1.25 g (500 mg elemental)		250	-	<u>Calci-Tab 500</u>
* Tab eff 1.25 g (500 mg elemental) – Subsidy by endorseme	nt260.00	100	v	Calcium 500 mg Hexal S29
Subsidy by endorsement – Only when prescribed for pa considered unsuitable.	ediatric patients (< 5	years	) where cal	
CALCIUM GLUCONATE				
* Inj 10%, 10 ml ampoule		10	<b>√</b>	Max Health - Hamein S29
lodine				
POTASSIUM IODATE				
* Tab 253 mcg (150 mcg elemental iodine)	5.99	90	✓ ]	NeuroTabs
Iron				
FERROUS FUMARATE				
* Tab 200 mg (65 mg elemental)	3.49	100	✓ [	Ferro-tab
FERROUS FUMARATE WITH FOLIC ACID				
* Tab 310 mg (100 mg elemental) with folic acid 350 mcg	5.98	100	✓	Ferro-F-Tabs
FERROUS SULFATE * Tab long-acting 325 mg (105 mg elemental)	2 55	30	<b>~</b>	Ferrograd
<ul> <li>* Oral liq 30 mg (6 mg elemental) per 1 ml</li> </ul>		250 m		Ferro-Liquid
	13.10	500 m	nl 🖌 🗸	Ferodan
IRON (AS FERRIC CARBOXYMALTOSE) – Special Authority s Inj 50 mg per ml, 10 ml vial		Retail 1		Ferinject
SA2394 Special Authority for Subsidy				
Initial application — (Anaemia) from any relevant practitioner.	Approvals valid for	3 mon	ths for appl	ications meeting the
following criteria: All of the following:				
<ol> <li>Patient has been diagnosed with anaemia; and</li> <li>Any of the following:</li> </ol>				
2.1 Serum ferritin level is 20 mcg/L or less; or				
2.2 Both:				
2.2.1 Serum ferritin is between 20 and 50 mcg/L 2.2.2 C-Reactive Protein (CRP) is at least 5 mg/				
2.3 Patient has chronic inflammatory disease with syn	nptoms of anaemia c	lespite	normal iror	n levels; and
3 Any of the following:				
<ul> <li>3.1 Oral iron treatment has proven ineffective; or</li> <li>3.2 Oral iron treatment has resulted in dose-limiting in</li> <li>3.2 Papid correction of anomia is required</li> </ul>	tolerance; or			
<ol> <li>3.3 Rapid correction of anaemia is required.</li> <li>Renewal — (Anaemia) from any relevant practitioner. Approva</li> </ol>	le valid for 3 months	for an	nlications n	peeting the following
nenewai – (Anacinia) nom any relevant practitioner. Approva	is valid for 3 months	οιοιαρ	ρισαιοτίς Π	

continued...

Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

Subsidy	Fully		Brand or	
(Manufacturer's Price)	Subsidised		Generic	
\$	Per	1	Manufacturer	

#### continued...

criteria:

Both:

- 1 Patient continues to have anaemia with a serum ferritin level of 20 mcg/L, or less or between 20 and 50 mcg/L with CRP of at least 5 mg/L, or has chronic inflammatory disease with symptoms of anaemia despite normal iron levels; and
- 2 A trial (or re-trial) with oral iron is clinically inappropriate.

Initial application — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 Patient has been diagnosed with iron-deficiency anaemia; and
- 2 Any of the following:
  - 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
  - 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
  - 2.3 Patient has symptomatic heart failure, chronic kidney disease stage 3 or more or active inflammatory bowel disease and a trial of oral iron is unlikely to be effective; or
  - 2.4 Rapid correction of anaemia is required.

Renewal — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 Patient continues to have iron-deficiency anaemia; and
- 2 A re-trial with oral iron is clinically inappropriate.

**IRON POLYMALTOSE** 

* Inj 50 mg per ml, 2 ml ampoule	5	✓ Ferrosig
Magnesium		
MAGNESIUM HYDROXIDE Suspension 8%	355 ml	✓ Phillips Milk of Magnesia S29
MAGNESIUM SULPHATE           * Inj 2 mmol per ml, 5 ml ampoule           * Inj 2 mmol per ml, 10 ml ampoule	10 10	<ul> <li>✓ <u>Martindale</u></li> <li>✓ Inresa S29</li> </ul>
Zinc		
ZINC SULPHATE * Cap 137.4 mg (50 mg elemental)11.00	100	<ul> <li>Zincaps</li> </ul>

# **BLOOD AND BLOOD FORMING ORGANS**

Subsidised

Per

Fully

Subsidy (Manufacturer's Price) \$ Brand or Generic

Manufacturer

# Antianaemics

### Hypoplastic and Haemolytic

#### ⇒SA2266 Special Authority for Subsidy

**Initial application** — (chronic renal failure) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient in chronic renal failure; and
- 2 Haemoglobin is less than or equal to 100g/L; and
- 3 Any of the following:
  - 3.1 Both:
    - 3.1.1 Patient does not have diabetes mellitus; and
    - 3.1.2 Glomerular filtration rate is less than or equal to 30ml/min; or
  - 3.2 Both:
    - 3.2.1 Patient has diabetes mellitus; and
    - 3.2.2 Glomerular filtration rate is less than or equal to 45ml/min; or
  - 3.3 Patient is on haemodialysis or peritoneal dialysis.

Initial application — (myelodysplasia) from any specialist. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of myelodysplasia (MDS)*; and
- 2 Has had symptomatic anaemia with haemoglobin < 100g/L and is red cell transfusion-dependent; and
- 3 Patient has very low, low or intermediate risk MDS based on the WHO classification based prognostic scoring system for myelodysplastic syndrome (WPSS); and
- 4 Other causes of anaemia such as B12 and folate deficiency have been excluded; and
- 5 Patient has a serum epoetin level of < 500 IU/L; and
- 6 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.
- Note: Indication marked with * is an unapproved indication

**Renewal** — (chronic renal failure) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (myelodysplasia) from any specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient's transfusion requirement continues to be reduced with erythropoietin treatment; and
- 2 Transformation to acute myeloid leukaemia has not occurred; and
- 3 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Note: Indication marked with * is an unapproved indication

EPOETIN ALFA - Special Authority see SA2266 above - Retail pharmacy

Wastage claimable

Inj 1,000 iu in 0.5 ml, syringe	 6	<ul> <li>Binocrit</li> </ul>
Inj 2,000 iu in 1 ml, syringe	 6	<ul> <li>Binocrit</li> </ul>
Inj 3,000 iu in 0.3 ml, syringe	6	<ul> <li>Binocrit</li> </ul>
Inj 4,000 iu in 0.4 ml, syringe	6	<ul> <li>Binocrit</li> </ul>
Inj 5,000 iu in 0.5 ml, syringe	6	<ul> <li>Binocrit</li> </ul>
Inj 6,000 iu in 0.6 ml, syringe	6	<ul> <li>Binocrit</li> </ul>
Inj 8,000 iu in 0.8 ml, syringe	6	<ul> <li>Binocrit</li> </ul>
Inj 10,000 iu in 1 ml, syringe	6	<ul> <li>Binocrit</li> </ul>
Inj 40,000 iu in 1 ml, syringe	1	<ul> <li>Binocrit</li> </ul>

# **BLOOD AND BLOOD FORMING ORGANS**

	Subsidy (Manufacturer's Price) \$	Subsi Per	Fully idised	Brand or Generic Manufacturer
Megaloblastic				
FOLIC ACID * Tab 0.8 mg		1,000	✔ F	olic Acid multichem
* Tab 5 mg Oral liq 50 mcg per ml		100 5 ml OP		olic Acid Viatris iomed
Antifibrinolytics, Haemostatics and Local Scl	erosants			

#### EFTRENONACOG ALFA [RECOMBINANT FACTOR IX] - [Xpharm]

For patients with haemophilia B receiving prophylaxis treatment. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management group.

Inj 250 iu vial		<u> </u>	<ul> <li>Alprolix</li> </ul>
Inj 500 iu vial	1,225.00	1	Alprolix
Inj 1,000 iu vial	2,450.00	1	Alprolix
Inj 2,000 iu vial		1	<ul> <li>Alprolix</li> </ul>
Inj 3,000 iu vial		1	<ul> <li>Alprolix</li> </ul>
Inj 4,000 iu vial		1	<ul> <li>Alprolix</li> </ul>
ELTROMBOPAG – Special Authority see SA1743 be Wastage claimable	low – Retail pharmacy		
Tab 25 mg		28	Revolade
Tab 50 mg	-	28	<ul> <li>Revolade</li> </ul>

#### ► SA1743 Special Authority for Subsidy

Initial application — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 6 weeks for applications meeting the following criteria:

- All of the following:
  - 1 Patient has had a splenectomy; and
  - 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
  - 3 Any of the following:
    - 3.1 Patient has a platelet count of 20,000 to 30,000 platelets per microlitre and has evidence of significant mucocutaneous bleeding; or
    - 3.2 Patient has a platelet count of less than or equal to 20,000 platelets per microlitre and has evidence of active bleeding; or
    - 3.3 Patient has a platelet count of less than or equal to 10,000 platelets per microlitre.

**Initial application** — (idiopathic thrombocytopenic purpura - preparation for splenectomy) only from a haematologist. Approvals valid for 6 weeks where the patient requires eltrombopag treatment as preparation for splenectomy.

**Initial application** — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a significant and well-documented contraindication to splenectomy for clinical reasons; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Either:
  - 3.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per microliter; or
  - 3.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre

continued...

Subsidy	F	ully	Brand or	
(Manufacturer's Price)	Subsidi	sed	Generic	
\$	Per	✓	Manufacturer	

continued...

and significant mucocutaneous bleeding.

Initial application — (severe aplastic anaemia) only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

1 Two immunosuppressive therapies have been trialled and failed after therapy of at least 3 months duration; and

- 2 Either:
  - 2.1 Patient has severe aplastic anaemia with a platelet count of less than or equal to 20,000 platelets per microliter; or
  - 2.2 Patient has severe aplastic anaemia with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

Renewal — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 12 months where the patient has obtained a response (see Note) from treatment during the initial approval or subsequent renewal periods and further treatment is required.

Note: Response to treatment is defined as a platelet count of > 30,000 platelets per microlitre.

Renewal — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient's significant contraindication to splenectomy remains; and
- 2 The patient has obtained a response from treatment during the initial approval period: and
- 3 Patient has maintained a platelet count of at least 50,000 platelets per microlitre on treatment; and
- 4 Further treatment with eltrombopag is required to maintain response.

Renewal — (severe aplastic anaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has obtained a response from treatment of at least 20.000 platelets per microlitre above baseline during the initial approval period; and
- 2 Platelet transfusion independence for a minimum of 8 weeks during the initial approval period.

EMICIZUMAB - [Xpharm] - Special Authority see SA2272 below

Inj 30 mg in 1 ml vial		1	<ul> <li>Hemlibra</li> </ul>
Inj 60 mg in 0.4 ml vial	7,138.00	1	<ul> <li>Hemlibra</li> </ul>
Inj 105 mg in 0.7 ml vial		1	<ul> <li>Hemlibra</li> </ul>
Inj 150 mg in 1 ml vial	17,846.00	1	<ul> <li>Hemlibra</li> </ul>

#### ■SA2272 Special Authority for Subsidy

Initial application — (Severe Haemophilia A with or without FVIII inhibitors) only from a haematologist. Approvals valid without further renewal unless notified for applications meeting the following criteria: Both:

- - 1 Patient has severe congenital haemophilia A with a severe bleeding phenotype (endogenous factor VIII activity less than or equal to 2%); and
  - 2 Emicizumab is to be administered at a dose of no greater than 3 mg/kg weekly for 4 weeks followed by the equivalent of 1.5 mg/kg weekly.

#### EPTACOG ALFA [RECOMBINANT FACTOR VIIA] - [Xpharm]

For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group. Rare Clinical Circumstances Brand of bypassing agent for > 14 days predicted use. Access to funded treatment for > 14 days predicted use is by named patient application to the Haemophilia Treaters Group, subject to access criteria.

Inj 1 mg syringe1,178.30	1	NovoSeven RT
Inj 2 mg syringe2,356.60	1	NovoSeven RT
Inj 5 mg syringe	1	NovoSeven RT
Inj 8 mg syringe9,426.40	1	NovoSeven RT

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy		Fully	Brand or
	(Manufacturer's Pric		bsidised	Generic
	\$	Per		Manufacturer
ACTOR EIGHT INHIBITOR BYPASSING FRACTION - [Xp				
For patients with haemophilia. Preferred Brand of bypas				
is managed by the Haemophilia Treaters Group in conju		l Haemoph		
Inj 500 U		1	-	EIBA NF
Inj 1,000 U		1	-	EIBA NF
Inj 2,500 U	6,575.00	1	🗸 F	EIBA NF
IOROCTOCOG ALFA [RECOMBINANT FACTOR VIII] – [X	(pharm]			
For patients with haemophilia. Rare Clinical Circumstan	ces Brand of short half	-life recomb	oinant fa	ctor VIII. Access to fund
treatment is managed by the Haemophilia Treaters Grou	p in conjunction with th	ne National	Haemop	hilia Management Grou
subject to criteria.				
Inj 250 iu prefilled syringe		1	>	(yntha
Inj 500 iu prefilled syringe		1	>	(yntha
Inj 1,000 iu prefilled syringe		1		(yntha
Inj 2,000 iu prefilled syringe		1		(yntha
Inj 3,000 iu prefilled syringe	'	1		(yntha
ONACOG GAMMA, [RECOMBINANT FACTOR IX] - [Xph	-			•
For patients with haemophilia. Access to funded treatme		Haamonhili	a Troato	rs Group in conjunction
with the National Haemophilia Management Group.	sin is managed by the	пасттортни	a meate	
Inj 1,000 iu vial	970.00	1		RIXUBIS
Inj 2,000 iu vial		1		RIXUBIS
	,	1		RIXUBIS
Inj 3,000 iu vial		I	• •	IIXUDIS
CTOCOG ALFA [RECOMBINANT FACTOR VIII] (ADVATE				
For patients with haemophilia. Preferred Brand of short				
managed by the Haemophilia Treaters Group in conjunct				
Inj 500 iu vial		1	✓ ↓	Advate
Inj 1,000 iu vial		1	✓ ↓	Advate
Inj 2,000 iu vial		1	✓ ↓	Advate
Inj 3,000 iu vial	2,520.00	1	✓ ↓	Advate
CTOCOG ALFA [RECOMBINANT FACTOR VIII] (KOGENA	ATE FS) – [Xpharm]			
For patients with haemophilia. Rare Clinical Circumstan		-life recom	oinant fa	ctor VIII. Access to fund
treatment is managed by the Haemophilia Treaters Grou				
subject to criteria.				
Inj 250 iu vial	237 50	1	<b>V</b> H	Kogenate FS
Inj 500 iu vial		1		Kogenate FS
Inj 1,000 iu vial		1		Kogenate FS
Ini 2.000 iu vial		1		Kogenate FS
, ,	,	1		Kogenate FS
		1	• 1	togenale i 5
Inj 3,000 iu vial				
JRIOCTOCOG ALFA PEGOL [RECOMBINANT FACTOR	VIII] - [Xpharm]			
JRIOCTOCOG ALFA PEGOL [RECOMBINANT FACTOR For patients with haemophilia A receiving prophylaxis tre	VIII] – [Xpharm] eatment. Access to fun		ent is ma	anaged by the Haemopl
JRIOCTOCOG ALFA PEGOL [RECOMBINANT FACTOR ] For patients with haemophilia A receiving prophylaxis tree Treaters Group in conjunction with the National Haemop	VIII] – [Xpharm] eatment. Access to fun hilia Management grou	ıp.		
JRIOCTOCOG ALFA PEGOL [RECOMBINANT FACTOR For patients with haemophilia A receiving prophylaxis tre Treaters Group in conjunction with the National Haemop Inj 1,000 iu vial	VIII] – [Xpharm] eatment. Access to fun hilia Management grou 	ир. 1	<b>√</b> µ	Adynovate
JRIOCTOCOG ALFA PEGOL [RECOMBINANT FACTOR For patients with haemophilia A receiving prophylaxis tre Treaters Group in conjunction with the National Haemop	VIII] – [Xpharm] eatment. Access to fun hilia Management grou 	ıp.	<b>√</b> µ	
URIOCTOCOG ALFA PEGOL [RECOMBINANT FACTOR ] For patients with haemophilia A receiving prophylaxis tre Treaters Group in conjunction with the National Haemop Inj 1,000 iu vial Inj 2,000 iu vial.	VIII] – [Xpharm] eatment. Access to fun hilia Management grou 	ир. 1	<b>√</b> µ	Adynovate
URIOCTOCOG ALFA PEGOL [RECOMBINANT FACTOR ' For patients with haemophilia A receiving prophylaxis tre Treaters Group in conjunction with the National Haemop Inj 1,000 iu vial Inj 2,000 iu vial	VIII] – [Xpharm] eatment. Access to fun hilia Management grou 	ир. 1 1	<b>√</b> µ	Adynovate
URIOCTOCOG ALFA PEGOL [RECOMBINANT FACTOR ' For patients with haemophilia A receiving prophylaxis tre Treaters Group in conjunction with the National Haemop Inj 1,000 iu vial Inj 2,000 iu vial	VIII] – [Xpharm] eatment. Access to fun hilia Management grou 	ир. 1	✓    ✓	Adynovate Adynovate
URIOCTOCOG ALFA PEGOL [RECOMBINANT FACTOR ' For patients with haemophilia A receiving prophylaxis tre Treaters Group in conjunction with the National Haemop Inj 1,000 iu vial Inj 2,000 iu vial ODIUM TETRADECYL SULPHATE Inj 3% 2 ml	VIII] – [Xpharm] eatment. Access to fun hilia Management grou 	ир. 1 1	✓    ✓	Adynovate
URIOCTOCOG ALFA PEGOL [RECOMBINANT FACTOR ' For patients with haemophilia A receiving prophylaxis tre Treaters Group in conjunction with the National Haemop Inj 1,000 iu vial Inj 2,000 iu vial ODIUM TETRADECYL SULPHATE Inj 3% 2 ml RANEXAMIC ACID	VIII] – [Xpharm] eatment. Access to fun hilia Management grou 	ιρ. 1 1	✓ # ✓ # F	Adynovate Adynovate
URIOCTOCOG ALFA PEGOL [RECOMBINANT FACTOR ' For patients with haemophilia A receiving prophylaxis tre Treaters Group in conjunction with the National Haemop Inj 1,000 iu vial	VIII] – [Xpharm] eatment. Access to fun hilia Management grou 	ир. 1 1	✓ / ✓ / F	Adynovate Adynovate

	Subsidy	•	Fully	Brand or
(λ	Anufacturer's Price)	Su Per	bsidised	Generic
	\$	геі		Manufacturer
Vitamin K				
HYTOMENADIONE				
Inj 2 mg per 0.2 ml – Up to 5 inj available on a PSO	8.00	5	🗸 К	Conakion MM
Inj 10 mg per ml, 1 ml - Up to 5 inj available on a PSO	9.21	5	🗸 К	Conakion MM
Antithrombotic Agents				
Antiplatelet Agents				
SPIRIN				
← Tab 100 mg	12.65	990	✓ <u>E</u>	thics Aspirin EC
LOPIDOGREL				
<ul> <li>Tab 75 mg</li> </ul>	5.07	84	🗸 A	rrow - Clopid
IPYRIDAMOLE				
K Tab long-acting 150 mg		60	🗸 Р	ytazen SR
ICAGRELOR – Special Authority see SA1955 below – Retail phar				
Tab 90 mg		56	🗸 т	icagrelor Sandoz
SA1055 Special Authority for Subaidy			-	

#### ➡SA1955 Special Authority for Subsidy

**Initial application** — (acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

#### Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Initial application — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

### Both:

- 1 Either:
  - 1.1 Patient has had a neurological stenting procedure* in the last 60 days; or
  - 1.2 Patient is about to have a neurological stenting procedure performed*; and
- 2 Either:
  - 2.1 Patient has demonstrated clopidogrel resistance using the P2Y12 (VerifyNow) assay or another appropriate platelet function assay and requires antiplatelet treatment with ticagrelor; or
  - 2.2 Either:
    - 2.2.1 Clopidogrel resistance has been demonstrated by the occurrence of a new cerebral ischemic event; or
    - 2.2.2 Clopidogrel resistance has been demonstrated by the occurrence of transient ischemic attack symptoms referable to the stent.

Initial application — (Percutaneous coronary intervention with stent deployment) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

### All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic**.

**Initial application** — (Stent thrombosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has experienced cardiac stent thrombosis whilst on clopidogrel.

Renewal — (subsequent acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

	Subsidy	F	ully	Brand or
(Manu	ufacturer's Price)	Subsidi	sed	Generic
	\$	Per	✓	Manufacturer

applications meeting the following criteria:

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Renewal — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient is continuing to benefit from treatment; and
- 2 Treatment continues to be clinically appropriate.

Renewal — (Percutaneous coronary intervention with stent deployment) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic**.

Notes: indications marked with * are unapproved indications.

Note: ** Clopidogrel allergy is defined as a history of anaphylaxis, urticaria, generalised rash or asthma (in non-asthmatic patients) developing soon after clopidogrel is started and is considered unlikely to be caused by any other treatment.

## Heparin and Antagonist Preparations

ENOXAPARIN SODIUM - Special Authority see SA2152 below - Retail pharmacy

Inj 20 mg in 0.2 ml syringe		10	<ul> <li><u>Clexane</u></li> </ul>
Inj 40 mg in 0.4 ml syringe		10	<ul> <li>Clexane</li> </ul>
Inj 60 mg in 0.6 ml syringe		10	<ul> <li>Clexane</li> </ul>
Inj 80 mg in 0.8 ml syringe		10	<ul> <li>Clexane</li> </ul>
Inj 100 mg in 1 ml syringe		10	✓ Clexane
Inj 120 mg in 0.8 ml syringe		10	✓ Clexane Forte
Inj 150 mg in 1 ml syringe	100.70	10	<ul> <li>Clexane Forte</li> </ul>

### ⇒SA2152 Special Authority for Subsidy

Initial application — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patients pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Initial application — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 For the short-term treatment of venous thromboembolism prior to establishing a therapeutic level of oral anti-coagulant treatment; or
- 2 For the prophylaxis and treatment of venous thromboembolism in high risk surgery; or
- 3 To enable cessation/re-establishment of existing oral anticoagulant treatment pre/post surgery; or
- 4 For the prophylaxis and treatment of venous thromboembolism in Acute Coronary Syndrome surgical intervention; or
- 5 To be used in association with cardioversion of atrial fibrillation.

Initial application - (Short-term use during treatment of COVID-19 with nirmatrelvir with ritonavir) from any relevant

continued...

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

practitioner. Approvals valid for 2 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is receiving an anticoagulation treatment that has drug/drug interactions with ritonavir that increases risk of bleeding; and
- 2 Patient meets the Access Criteria for COVID-19 antivirals published on the Pharmac website*; and
- 3 Other antiviral treatments for COVID-19 have been considered and are not clinically suitable options.

Renewal — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patient's pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Renewal — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month where low molecular weight heparin treatment or prophylaxis is required for a second or subsequent event (surgery, ACS, cardioversion, or prior to oral anti-coagulation).

**HEPARIN SODIUM** 

Inj 1,000 iu per ml, 10 ml vial127	.44	25	<ul> <li>Pfizer S29</li> </ul>
Inj 1,000 iu per ml, 5 ml ampoule25	5.49	10	<ul> <li>Wockhardt S29</li> </ul>
103	3.70		Wockhardt PSF S29
127	<b>'</b> .44	50	<ul> <li>Pfizer</li> </ul>
Inj 5,000 iu per ml, 5 ml vial83	3.00	10	<ul> <li><u>Heparin Sodium</u></li> <li><u>Panpharma</u></li> </ul>
Inj 5,000 iu per ml, 1 ml70	).33	5	<ul> <li>Hospira</li> </ul>
Inj 25,000 iu per ml, 0.2 ml25		5	✓ Hospira
42	2.40		<ul> <li>Heparin DBL \$29</li> </ul>
482	2.20	50	<ul> <li>Heparin DBL \$29</li> </ul>
HEPARINISED SALINE			
Inj 10 iu per ml, 5 ml	6.91	50	<ul> <li>Pfizer</li> </ul>
Oral Anticoagulants			
DABIGATRAN			
Cap 75 mg – No more than 2 cap per day27	'.99	60	Pradaxa
Cap 110 mg27	<b>'</b> .99	60	<ul> <li>Pradaxa</li> </ul>
Cap 150 mg27	7.99	60	Pradaxa
RIVAROXABAN			
Tab 10 mg – No more than 1 tab per day 15	60	30	🗸 Xarelto

	Tab 10 mg – No more than 1 tab per day		30	Xarelto
	Tab 15 mg - Up to 14 tab available on a PSO	14.56	28	✓ Xarelto
	Tab 20 mg	14.56	28	✓ Xarelto
WA	ARFARIN SODIUM			
	Note: Marevan and Coumadin are not interchangeable.			
*	Tab 1 mg	3.46	50	<ul> <li>Coumadin</li> </ul>
	-	7.50	100	🗸 Marevan
*	Tab 2 mg	4.31	50	<ul> <li>Coumadin</li> </ul>
*	Tab 3 mg		100	<ul> <li>Marevan</li> </ul>
*	Tab 5 mg	5.93	50	<ul> <li>Coumadin</li> </ul>
	-	13.50	100	<ul> <li>Marevan</li> </ul>

	Subsidy		Fully	Brand or		
	(Manufacturer's Price) \$	Per	Subsidised	Generic Manufacturer		
Blood Colony-stimulating Factors						
FILGRASTIM – Special Authority see SA1259 below – Retail ph Inj 300 mcg per 0.5 ml prefilled syringe Inj 480 mcg per 0.5 ml prefilled syringe		10 10		livestim livestim		
■ SA1259 Special Authority for Subsidy Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following:						
<ol> <li>Prevention of neutropenia in patients undergoing high risk or equal to 20%*); or</li> <li>Desire and the end stars and machine the stars and end of the stars an</li></ol>						
<ol> <li>Peripheral blood stem cell mobilisation in patients underg</li> <li>Peripheral blood stem cell mobilisation or bone marrow de</li> <li>Treatment of severe chronic neutropenia (ANC &lt; 0.5 ×10</li> <li>Treatment of drug-induced prolonged neutropenia (ANC </li> </ol>	onation from healthy o		,			
Note: *Febrile neutropenia risk greater than or equal to 20% afte European Organisation for Research and Treatment of Cancer (I	•	other	risk factors	as defined by the		
PEGFILGRASTIM - Special Authority see SA1912 below - Retain	ail pharmacy					

- ✓ <u>Ziextenzo</u>
   ✓ Ziextenzo AU

1

### ⇒SA1912 Special Authority for Subsidy

**Initial application** only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where used for prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 5% *). Note: *Febrile neutropenia risk greater than or equal to 5% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

# Fluids and Electrolytes

### Intravenous Administration

GLUCOSE [DEXTROSE] * Inj 50%, 10 ml ampoule – Up to 5 inj available on a PSO * Inj 50%, 90 ml bottle – Up to 5 inj available on a PSO POTASSIUM CHLORIDE		5 1	✓ <u>Biomed</u> ✓ <u>Biomed</u>
✤ Inj 75 mg per ml, 10 ml	65.00	50	<ul> <li>✓ Juno</li> <li>✓ LumaCina</li> <li>✓ Pfizer ^{\$29}</li> </ul>
SODIUM BICARBONATE Inj 8.4%, 50 ml a) Up to 5 inj available on a PSO	24.70	1	<ul> <li>Biomed</li> </ul>
<ul><li>b) Not in combination</li><li>Inj 8.4%, 100 mla) Up to 5 inj available on a PSO</li></ul>	25.31	1	<ul> <li>Biomed</li> </ul>

b) Not in combination

	Subsidy		Fully Brand or
	(Manufacturer's P		idised Generic
	\$	Per	<ul> <li>Manufacturer</li> </ul>
SODIUM CHLORIDE	rupp overst whe	n upod in poniu	notion with an antibiatic intended
Not funded for use as a nasal drop. Not funded for nebulise for nebuliser use.	r use except whe	n used in conju	inction with an antibiotic intended
Inj 0.9%, bag – Up to 2000 ml available on a PSO		500 ml	<ul> <li>Baxter</li> </ul>
,,	1.58	1,000 ml	✓ Baxter
Only if prescribed on a prescription for renal dialysis, ma for emergency use. (500 ml and 1,000 ml packs)	aternity or post-na	atal care in the l	home of the patient, or on a PSC
Inj 23.4% (4 mmol/ml), 20 ml ampoule		5	<ul> <li>Biomed</li> </ul>
For Sodium chloride oral liquid formulation refer Standar			<b>4</b>
Inj 0.9%, 5 ml ampoule – Up to 5 inj available on a PSO		20	<ul> <li>Fresenius Kabi</li> </ul>
Inj 0.9%, 10 ml ampoule – Up to 5 inj available on a PSO		50	<ul> <li>Fresenius Kabi</li> </ul>
Inj 0.9%, 20 ml ampoule	5.00	20	<ul> <li>Fresenius Kabi</li> </ul>
TOTAL PARENTERAL NUTRITION (TPN)	000	4.00	
Infusion		1 OP	✓ TPN
WATER			
<ol> <li>On a prescription or Practitioner's Supply Order only w Schedule requiring a solvent or diluent; or</li> </ol>	hen on the same	form as an inje	ction listed in the Pharmaceutica
2) On a bulk supply order; or			
3) When used in the extemporaneous compounding of ey			
4) When used for the dilution of sodium chloride soln 7%	for cystic fibrosis	patients only.	
			<b>A a a a b b b</b>
Inj 10 ml ampoule – Up to 5 inj available on a PSO		50	✓ <u>Multichem</u>
Inj 20 ml ampoule – Up to 5 inj available on a PSO		50 20	<ul> <li>✓ <u>Multichem</u></li> <li>✓ <u>Fresenius Kabi</u></li> </ul>
Inj 20 ml ampoule – Up to 5 inj available on a PSO			
Inj 20 ml ampoule – Up to 5 inj available on a PSO	5.00		
Inj 20 ml ampoule – Up to 5 inj available on a PSO Oral Administration CALCIUM POLYSTYRENE SULPHONATE Powder	5.00	20	✓ Fresenius Kabi
Inj 20 ml ampoule – Up to 5 inj available on a PSO Oral Administration CALCIUM POLYSTYRENE SULPHONATE Powder		20	✓ Fresenius Kabi
Inj 20 ml ampoule – Up to 5 inj available on a PSO Oral Administration CALCIUM POLYSTYRENE SULPHONATE Powder COMPOUND ELECTROLYTES Powder for oral soln – Up to 5 sach available on a PSO		20 300 g OP	<ul> <li>✓ <u>Fresenius Kabi</u></li> <li>✓ Calcium Resonium</li> </ul>
Inj 20 ml ampoule – Up to 5 inj available on a PSO Oral Administration CALCIUM POLYSTYRENE SULPHONATE Powder COMPOUND ELECTROLYTES Powder for oral soln – Up to 5 sach available on a PSO		20 300 g OP	<ul> <li>✓ <u>Fresenius Kabi</u></li> <li>✓ Calcium Resonium</li> </ul>
Inj 20 ml ampoule – Up to 5 inj available on a PSO Oral Administration CALCIUM POLYSTYRENE SULPHONATE Powder COMPOUND ELECTROLYTES Powder for oral soln – Up to 5 sach available on a PSO COMPOUND ELECTROLYTES WITH GLUCOSE [DEXTROSE]		20 300 g OP 50	<ul> <li>✓ Fresenius Kabi</li> <li>✓ Calcium Resonium</li> <li>✓ Electral</li> </ul>
Inj 20 ml ampoule – Up to 5 inj available on a PSO Oral Administration CALCIUM POLYSTYRENE SULPHONATE Powder COMPOUND ELECTROLYTES Powder for oral soln – Up to 5 sach available on a PSO COMPOUND ELECTROLYTES WITH GLUCOSE [DEXTROSE] Soln with electrolytes		20 300 g OP 50	<ul> <li>Fresenius Kabi</li> <li>Calcium Resonium</li> <li><u>Electral</u></li> <li><u>Hydralyte -</u></li> </ul>
Inj 20 ml ampoule – Up to 5 inj available on a PSO Oral Administration CALCIUM POLYSTYRENE SULPHONATE Powder COMPOUND ELECTROLYTES Powder for oral soln – Up to 5 sach available on a PSO COMPOUND ELECTROLYTES WITH GLUCOSE [DEXTROSE] Soln with electrolytes		20 300 g OP 50	<ul> <li>Fresenius Kabi</li> <li>Calcium Resonium</li> <li><u>Electral</u></li> <li><u>Hydralyte -</u></li> </ul>
Inj 20 ml ampoule – Up to 5 inj available on a PSO Oral Administration CALCIUM POLYSTYRENE SULPHONATE Powder COMPOUND ELECTROLYTES Powder for oral soln – Up to 5 sach available on a PSO COMPOUND ELECTROLYTES WITH GLUCOSE [DEXTROSE] Soln with electrolytes PHOSPHORUS Tab eff 500 mg (16 mmol)		20 300 g OP 50 1,000 ml OP	<ul> <li>Fresenius Kabi</li> <li>Calcium Resonium</li> <li>Electral</li> <li>Hydralyte - Lemonade</li> </ul>
Inj 20 ml ampoule – Up to 5 inj available on a PSO Oral Administration CALCIUM POLYSTYRENE SULPHONATE Powder COMPOUND ELECTROLYTES Powder for oral soln – Up to 5 sach available on a PSO COMPOUND ELECTROLYTES WITH GLUCOSE [DEXTROSE] Soln with electrolytes PHOSPHORUS Tab eff 500 mg (16 mmol) POTASSIUM CHLORIDE		20 300 g OP 50 1,000 ml OP	<ul> <li>Fresenius Kabi</li> <li>Calcium Resonium</li> <li>Electral</li> <li>Hydralyte - Lemonade</li> </ul>
Inj 20 ml ampoule – Up to 5 inj available on a PSO Oral Administration CALCIUM POLYSTYRENE SULPHONATE Powder COMPOUND ELECTROLYTES Powder for oral soln – Up to 5 sach available on a PSO COMPOUND ELECTROLYTES WITH GLUCOSE [DEXTROSE] Soln with electrolytes PHOSPHORUS Tab eff 500 mg (16 mmol) POTASSIUM CHLORIDE * Tab eff 548 mg (14 m eq) with chloride 285 mg (8 m eq)		20 300 g OP 50 1,000 ml OP 100 60	<ul> <li>Fresenius Kabi</li> <li>Calcium Resonium</li> <li>Electral</li> <li><u>Hydralyte -</u> <u>Lemonade</u></li> <li>Phosphate Phebra</li> <li>Chlorvescent</li> </ul>
Inj 20 ml ampoule – Up to 5 inj available on a PSO Oral Administration CALCIUM POLYSTYRENE SULPHONATE Powder COMPOUND ELECTROLYTES Powder for oral soln – Up to 5 sach available on a PSO COMPOUND ELECTROLYTES WITH GLUCOSE [DEXTROSE] Soln with electrolytes PHOSPHORUS Tab eff 500 mg (16 mmol) POTASSIUM CHLORIDE * Tab eff 548 mg (14 m eq) with chloride 285 mg (8 m eq)		20 300 g OP 50 1,000 ml OP 100	<ul> <li>Fresenius Kabi</li> <li>Calcium Resonium</li> <li>Electral</li> <li><u>Hydralyte -</u> <u>Lemonade</u></li> <li>Phosphate Phebra</li> </ul>
Inj 20 ml ampoule – Up to 5 inj available on a PSO Oral Administration CALCIUM POLYSTYRENE SULPHONATE Powder COMPOUND ELECTROLYTES Powder for oral soln – Up to 5 sach available on a PSO COMPOUND ELECTROLYTES WITH GLUCOSE [DEXTROSE] Soln with electrolytes PHOSPHORUS Tab eff 500 mg (16 mmol) POTASSIUM CHLORIDE * Tab eff 548 mg (14 m eq) with chloride 285 mg (8 m eq) * Tab long-acting 600 mg (8 mmol) SODIUM BICARBONATE		20 300 g OP 50 1,000 ml OP 100 60 200	<ul> <li>Fresenius Kabi</li> <li>Calcium Resonium</li> <li>Electral</li> <li><u>Hydralyte - Lemonade</u></li> <li>Phosphate Phebra</li> <li>Chlorvescent</li> <li>Span-K</li> </ul>
Inj 20 ml ampoule – Up to 5 inj available on a PSO Oral Administration CALCIUM POLYSTYRENE SULPHONATE Powder COMPOUND ELECTROLYTES Powder for oral soln – Up to 5 sach available on a PSO COMPOUND ELECTROLYTES WITH GLUCOSE [DEXTROSE] Soln with electrolytes PHOSPHORUS Tab eff 500 mg (16 mmol) POTASSIUM CHLORIDE * Tab eff 548 mg (14 m eq) with chloride 285 mg (8 m eq) * Tab long-acting 600 mg (8 mmol)		20 300 g OP 50 1,000 ml OP 100 60	<ul> <li>Fresenius Kabi</li> <li>Calcium Resonium</li> <li>Electral</li> <li><u>Hydralyte - Lemonade</u></li> <li>Phosphate Phebra</li> <li>Chlorvescent</li> <li>Span-K</li> <li>Sodibic</li> </ul>
Inj 20 ml ampoule – Up to 5 inj available on a PSO Oral Administration CALCIUM POLYSTYRENE SULPHONATE Powder COMPOUND ELECTROLYTES Powder for oral soln – Up to 5 sach available on a PSO COMPOUND ELECTROLYTES WITH GLUCOSE [DEXTROSE] Soln with electrolytes PHOSPHORUS Tab eff 500 mg (16 mmol) POTASSIUM CHLORIDE * Tab long-acting 600 mg (8 mmol) SODIUM BICARBONATE Cap 840 mg		20 300 g OP 50 1,000 ml OP 100 60 200	<ul> <li>Fresenius Kabi</li> <li>Calcium Resonium</li> <li>Electral</li> <li><u>Hydralyte - Lemonade</u></li> <li>Phosphate Phebra</li> <li>Chlorvescent</li> <li>Span-K</li> </ul>
Inj 20 ml ampoule – Up to 5 inj available on a PSO Oral Administration CALCIUM POLYSTYRENE SULPHONATE Powder COMPOUND ELECTROLYTES Powder for oral soln – Up to 5 sach available on a PSO COMPOUND ELECTROLYTES WITH GLUCOSE [DEXTROSE] Soln with electrolytes PHOSPHORUS Tab eff 500 mg (16 mmol) POTASSIUM CHLORIDE * Tab eff 548 mg (14 m eq) with chloride 285 mg (8 m eq) * Tab long-acting 600 mg (8 mmol) SODIUM BICARBONATE		20 300 g OP 50 1,000 ml OP 100 60 200	<ul> <li>Fresenius Kabi</li> <li>Calcium Resonium</li> <li>Electral</li> <li><u>Hydralyte - Lemonade</u></li> <li>Phosphate Phebra</li> <li>Chlorvescent</li> <li>Span-K</li> <li>Sodibic</li> </ul>

	Subsidy (Manufacturer's Pric \$	e) S Per	Fully Subsidised	
Alpha-Adrenoceptor Blockers				
Alpha Adrenoceptor Blockers				
DOXAZOSIN				
★ Tab 2 mg		500		Doxazosin Clinect
₭ Tab 4 mg	20.94	500	~	Doxazosin Clinect
PHENOXYBENZAMINE HYDROCHLORIDE			_	_
₭ Сар 10 mg		30		BNM S29
	216.67	100	~	Dibenzyline S29
(Dibenzyline S29) Cap 10 mg to be delisted 1 July 2025)				
PRAZOSIN				
* Tab 1 mg	5.53	100	~	Arrotex-Prazosin S29 S29
	9.98		~	Minipress S29
* Tab 2 mg	7.00	100		Arrotex-Prazosin S29 S29
	13.29		-	Minipress S29
* Tab 5 mg	11.70	100	1	Arrotex-Prazosin S29 S29
	22.00		-	Minipress S29
* Cap 1 mg	15.40	100	-	Prazosin Mylan S29
* Cap 2 mg	15.58	100	✓	Prazosin Mylan S29
* Cap 5 mg	23.32	100	1	Prazosin Mylan S29
Agents Affecting the Renin-Angiotensin System	า			
ACE Inhibitors				
CAPTOPRIL				
Oral liq 5 mg per ml Oral liquid restricted to children under 12 years of age.		100 ml Ol	P 🗸	DP-Captopril
CILAZAPRIL – Subsidy by endorsement				
Subsidy by endorsement – Subsidised for patients who were endorsed accordingly. Pharmacists may annotate the presc dispensing of cilazapril.				
* Tab 0.5 mg	2 69	90	1	Zapril
* Tab 0.5 mg		90		Zapril
Tab C ma	40.05	00		

Tab 5 mg	
(Zapril Tab 0.5 mg to be delisted 1 April 2025)	
Zapril Tab 2.5 mg to be delisted 1 April 2025)	
(Zapril Tab 5 mg to be delisted 1 April 2025)	

ENALAPRIL MALEATE * *

*	Tab 5 mg1.75	90	<ul> <li>Acetec</li> </ul>
*	Tab 10 mg	90	✓ Acetec
	Tab 20 mg2.35	90	✓ Acetec

✓ Zapril

	Subsidy		Fully Brand or
	(Manufacturer's Price) \$	: Per	Subsidised Generic Manufacturer
	φ	rei	
_ISINOPRIL ₩ Tab 5 mg	11.07	90	<ul> <li>Ethics Lisinopril</li> </ul>
* Tab 5 Hig		90	✓ Teva Lisinopril
* Tab 10 mg		90	✓ Ethics Lisinopril
· · · · · · · · · · · · · · · · · · ·			<ul> <li>Teva Lisinopril</li> </ul>
* Tab 20 mg	14.69	90	<ul> <li>Ethics Lisinopril</li> </ul>
-			<ul> <li>Teva Lisinopril</li> </ul>
PERINDOPRIL			
* Tab 2 mg	1.79	30	✓ Coversyl
₭ Tab 4 mg	2.44	30	<ul> <li><u>Coversyl</u></li> </ul>
* Tab 8 mg	3.94	30	<ul> <li>Coversyl</li> </ul>
QUINAPRIL			
₭ Tab 5 mg		90	Arrow-Quinapril 5
* Tab 10 mg		90	<ul> <li>Arrow-Quinapril 10</li> </ul>
* Tab 20 mg	14.83	90	Arrow-Quinapril 20
RAMIPRIL			<b>4</b> -
* Cap 1.25 mg		90	Tryzan
₭ Cap 2.5 mg		90	✓ <u>Tryzan</u>
* Cap 5 mg		90 90	✓ <u>Tryzan</u>
* Cap 10 mg	17.03	90	<ul> <li><u>Tryzan</u></li> </ul>
Angiotensin II Antagonists			
CANDESARTAN CILEXETIL			
🖌 Tab 4 mg	2.68	90	<ul> <li>Candestar</li> </ul>
🖌 Tab 8 mg	2.67	90	✓ Candestar
₭ Tab 16 mg	4.22	90	Candestar
₭ Tab 32 mg	5.24	90	<ul> <li>Candestar</li> </ul>
OSARTAN POTASSIUM			
₭ Tab 12.5 mg		84	Losartan Actavis
₭ Tab 25 mg		84	<ul> <li>Losartan Actavis</li> </ul>
₭ Tab 50 mg		84	<ul> <li>Losartan Actavis</li> </ul>
K Tab 100 mg	4.57	84	<ul> <li>Losartan Actavis</li> </ul>
Angiotensin II Antagonists with Diuretics			
ANDESARTAN CILEXETIL WITH HYDROCHLOROTHIAZII	DE		
₭ Tab 16 mg with hydrochlorothiazide 12.5 mg		30	APO-Candesartan
			HCTZ 16/12.5
Fab 32 mg with hydrochlorothiazide 12.5 mg	5.25	30	<ul> <li>APO-Candesartan</li> </ul>
			HCTZ 32/12.5
OSARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE			
Fab 50 mg with hydrochlorothiazide 12.5 mg	4.00	30	✓ <u>Arrow-Losartan &amp;</u> <u>Hydrochlorothiazide</u>
Angiotensin II Antagonists with Neprilysin In	hibitors		
		Dete	il nhormooy
ACUBITRIL WITH VALSARTAN – Special Authority see SA	2302 on the next page -	Hetai	ii pharmacy

 SACUBITRIL WITH VALSARTAN – Special Authority see SA2302 on the next page – Retail pharmacy

 Tab 24.3 mg with valsartan 25.7 mg

 Tab 48.6 mg with valsartan 51.4 mg

 Tab 97.2 mg with valsartan 102.8 mg

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

Subsidy	S	Fully	Brand or
(Manufacturer's Price)		ubsidised	Generic
\$	Per	✓	Manufacturer

#### ⇒SA2302 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has heart failure; and
- 2 Any of the following:
  - 2.1 Patient is in NYHA/WHO functional class II; or
  - 2.2 Patient is in NYHA/WHO functional class III; or
  - 2.3 Patient is in NYHA/WHO functional class IV; and

3 Either:

- 3.1 Patient has a documented left ventricular ejection fraction (LVEF) of less than or equal to 35%; or
- 3.2 An ECHO is not reasonably practical, and in the opinion of the treating practitioner the patient would benefit from treatment; and
- 4 Patient is receiving concomitant optimal standard chronic heart failure treatments.

### Antiarrhythmics

For lignocaine hydrochloride refer to NERVOUS SYSTEM, Anaesthetics, Local, page 121

▲ Tab 100 mg		30	<ul> <li>Aratac</li> </ul>
▲ Tab 200 mg		30	✓ Aratac
Inj 50 mg per ml, 3 ml ampoule – Up to 10 inj available on		6	✓ Cordarone-X
	15.22	10	Max Health
ATROPINE SULPHATE			
Inj 600 mcg per ml, 1 ml ampoule – Up to 5 inj available on	а		
PSO		10	<ul> <li>Hikma S29</li> </ul>
			Juno \$29
			✓ Martindale
DIGOXIN			
* Tab 62.5 mcg – Up to 30 tab available on a PSO	7.80	240	Lanoxin PG
* Tab 250 mcg – Up to 30 tab available on a PSO		240	<ul> <li>Lanoxin</li> </ul>
* Oral liq 50 mcg per ml		60 ml	<ul> <li>Lanoxin</li> </ul>
			<ul> <li>Lanoxin Paediatric Elixir</li> </ul>
			<ul> <li>Lanoxin S29 S29</li> </ul>
DISOPYRAMIDE PHOSPHATE			
▲ Cap 100 mg		100	<ul> <li>Rythmodan</li> </ul>
	55.90	84	✓ Rythmodan -
	20100		Cheplafarm S29

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
LECAINIDE ACETATE				
Tab 50 mg		60		Flecainide BNM
Cap long-acting 100 mg	35.78	90	~	Flecainide Controlled Release Teva
Cap long-acting 200 mg	54.28	90	~	<u>Flecainide</u> <u>Controlled</u> Release Teva
Inj 10 mg per ml, 15 ml ampoule		5	✓	Almarytm S29
	108.16		1	Tambocor
			1	Tambocor German S29
IEXILETINE HYDROCHLORIDE				
Cap 150 mg		100	1	Teva S29
Cap 250 mg		100	1	Teva S29
Tab 150 mg	40.90	50	✓	Rytmonorm
Antihypotensives				
IIDODRINE – Special Authority see SA1474 below – Retail p				
Tab 2.5 mg		100		MAR-Midodrine S29 Midodrine Medsurge
Tab 5 mg	58.88	100		MAR-Midodrine S239 Midodrine Medsurge

#### ■SA1474 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years where patient has disabling orthostatic hypotension not due to drugs.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

# **Beta-Adrenoceptor Blockers**

## **Beta Adrenoceptor Blockers**

ATENOLOL		
* Tab 50 mg 11.0	0 500	<ul> <li>Viatris</li> </ul>
* Tab 100 mg	0 500	<ul> <li>Atenolol Viatris</li> </ul>
* Oral lig 25 mg per 5 ml		✓ Atenolol AFT
Restricted to children under 12 years of age.		
BISOPROLOL FUMARATE		
* Tab 2.5 mg1.3	6 90	Ipca-Bisoprolol
* Tab 5 mg	1 90	<ul> <li>Ipca-Bisoprolol</li> </ul>
* Tab 10 mg2.7	1 90	<ul> <li>Ipca-Bisoprolol</li> </ul>
CARVEDILOL		
* Tab 6.25 mg	4 60	Carvedilol Sandoz
* Tab 12.5 mg2.3	0 60	<ul> <li>Carvedilol Sandoz</li> </ul>
* Tab 25 mg		<ul> <li>Carvedilol Sandoz</li> </ul>

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Generic
LABETALOL				
* Tab 100 mg		100	1	Trandate
* Tab 200 mg		100	1	Trandate
* Inj 5 mg per ml, 20 ml ampoule		5		
	(88.60)			Trandate
METOPROLOL SUCCINATE				
* Tab long-acting 23.75 mg.	4.20	90	1	Myloc CR
* Tab long-acting 47.5 mg.		90		Myloc CR
* Tab long-acting 95 mg.		90	1	Myloc CR
* Tab long-acting 190 mg		90	1	Myloc CR
METOPROLOL TARTRATE				
* Tab 50 mg		100	1	IPCA-Metoprolol
* Tab 100 mg		60		IPCA-Metoprolol
* Tab long-acting 200 mg		28		Slow-Lopresor
* Inj 1 mg per ml, 5 ml vial		5		Metoprolol IV Mylan
<b>3 3 1 3 1 1</b>				Metoprolol IV Viatris
NADOLOL				
* Tab 40 mg		100	1	Nadolol BNM
* Tab 80 mg		100	1	Nadolol BNM
PROPRANOLOL				
* Tab 10 mg	7.04	100	1	Drofate
* Tab 40 mg		100	1	IPCA-Propranolol
* Cap long-acting 160 mg		100		Cardinol LA
* Oral liq 4 mg per ml – Special Authority see SA1327 below –				
Retail pharmacy		500 n	nl 🗸	Roxane-
· · ·				Propranolol S29

#### ⇒SA1327 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or

2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

#### SOTALOL

*	Tab 80 mg	500	🗸 Mylan
*	Tab 160 mg	100	<ul> <li>Mylan</li> </ul>

# **Calcium Channel Blockers**

### **Dihydropyridine Calcium Channel Blockers**

#### AMLODIPINE

*	Tab 2.5 mg 1	.45	90	<ul> <li>Vasorex</li> </ul>
	Tab 5 mg 1		90	✓ Vasorex
	Tab 10 mg 1		90	✓ Vasorex

48	🗸 fully subsidised
40	Principal Supply

	Subsidy		Fully	Brand or
	(Manufacturer's Price) \$	Per	Subsidised	Generic Manufacturer
ELODIPINE	*	-		
Tab long-acting 2.5 mg	2 18	30	1	Plendil ER
Tab long-acting 5 mg		90		Felo 5 ER
Tab long-acting 10 mg		90		Felo 10 ER
	0.95	30	•	
IFEDIPINE Tab long-acting 10 mg – Subsidy by endorsement		56	1	Tensipine MR10 \$29
				•
Subsidised for patients who were taking nifedipine	0 0 01			
endorsed accordingly. Pharmacists may annotate	the prescription as endors	ea wr	here there e	exists a record of prior
dispensing of nifedipine tab long-acting 10 mg.	17 70	100		Nuefey Deteral
Tab long-acting 20 mg Tab long-acting 30 mg		100	-	Nyefax Retard
Tab long-acting 30 mg	4./8	14	~	Mylan Italy (24 hr
				release) S29
	34.10	100	1	Mylan (24 hr
				release) S29
Tab long-acting 60 mg		100	1	Mylan (24 hr
5 5 5 ·····				release) S29
				, =
Other Calcium Channel Blockers				
TIAZEM HYDROCHLORIDE				
Cap long-acting 120 mg	65.35	500	✓	Diltiazem CD Clinect
Cap long-acting 180 mg	7.00	30	✓	Cardizem CD
Cap long-acting 240 mg	9.30	30	✓	Cardizem CD
RHEXILINE MALEATE				
Tab 100 mg		100	1	Pexsig
	7.01	100		Isoptin
5		100		
Tab 80 mg				Isoptin
Tab long-acting 120 mg		100		Isoptin Retard S29
	15 10	00		Isoptin SR
Tab long-acting 240 mg		30	~	Isoptin SR
Inj 2.5 mg per ml, 2 ml ampoule – Up to 5 inj available PSO		5		loontin
F9U	25.00	5	•	Isoptin
Centrally-Acting Agents				
ONIDINE			-	
Patch 2.5 mg, 100 mcg per day – Only on a prescription		4		Mylan
Patch 5 mg, 200 mcg per day – Only on a prescription		4		Mylan
Patch 7.5 mg, 300 mcg per day – Only on a prescription	on17.90	4	1	Mylan
ONIDINE HYDROCHLORIDE				
Tab 25 mcg		112	1	Clonidine Teva
Tab 150 mcg		100		Catapres
Inj 150 mcg per ml, 1 ml ampoule		5		Catapres
ETHYLDOPA				<b>.</b>
Tab 250 mg	15 10	100	1	Methyldopa Viatris
		100	•	mentyluopa viauts

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Pri \$	ice) Sub Per	Fully Brand or sidised Generic Manufacturer
Diuretics			
Loop Diuretics			
BUMETANIDE * Tab 1 mg * Inj 500 mcg per ml, 4 ml vial FUROSEMIDE [FRUSEMIDE]	7.95	100 5	<ul><li>✓ Burinex</li><li>✓ Burinex</li></ul>
Tab 40 mg – Up to 30 tab available on a PSO * Tab 500 mg * Oral liq 10 mg per ml * Inj 10 mg per ml, 25 ml ampoule * Inj 10 mg per ml, 2 ml ampoule – Up to 5 inj available on a PS	25.00 11.20 60.65	1,000 50 30 ml OP 6 5	<ul> <li>IPCA-Frusemide</li> <li>Urex Forte</li> <li>Lasix</li> <li>Lasix</li> <li>Furosemide-Baxter</li> </ul>
Potassium Sparing Diuretics			
AMILORIDE HYDROCHLORIDE Tab 5 mg	81.07 171.41	100 28	<ul> <li>✓ Padagis ^{\$29}</li> <li>✓ Wockhardt ^{\$29}</li> </ul>
Oral liq 1 mg per ml		25 ml OP	<ul> <li>Biomed</li> </ul>
EPLERENONE – Special Authority see SA1728 below – Retail ph Tab 25 mg Tab 50 mg		30 30	✓ <u>Inspra</u> ✓ <u>Inspra</u>
<ul> <li>SA1728 Special Authority for Subsidy         nitial application from any relevant practitioner. Approvals valid         he following criteria:             3oth:             1 Patient has heart failure with ejection fraction less than 40%             2 Either:         </li> </ul>		enewal unles	s notified for applications meeting
<ul><li>2.1 Patient is intolerant to optimal dosing of spironolactor</li><li>2.2 Patient has experienced a clinically significant adverted</li></ul>		on optimal do	sing of spironolactone.
SPIRONOLACTONE ★ Tab 25 mg ★ Tab 100 mg Oral liq 5 mg per ml	10.65	100 100 25 ml OP	✓ <u>Spiractin</u> ✓ <u>Spiractin</u> ✓ Biomed
Potassium Sparing Combination Diuretics			
AMILORIDE HYDROCHLORIDE WITH FUROSEMIDE * Tab 5 mg with furosemide 40 mg AMILORIDE HYDROCHLORIDE WITH HYDROCHLOROTHIAZID		28	✔ Frumil
	L		• · · · ·

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
Thiazide and Related Diuretics				
BENDROFLUMETHIAZIDE [BENDROFLUAZIDE] * Tab 2.5 mg – Up to 150 tab available on a PSO	51.50	500	<b>√</b> ,	Arrow- Bendrofluazide
May be supplied on a PSO for reasons other than emerge * Tab 5 mg		500	✓ <u>,</u>	<u>Arrow-</u> Bendrofluazide
CHLOROTHIAZIDE Oral liq 50 mg per ml CHLORTALIDONE [CHLORTHALIDONE]		5 ml C	)P 🖌	Biomed
Tab 25 mg     Minimum     INDAPAMIDE	6.95	50	•	Hygroton
Tab 2.5 mg METOLAZONE		90	•	Dapa-Tabs
Tab 5 mg	CBS	1 50		Metolazone s29 Zaroxolyn s29
(Metolazone see Tab 5 mg to be delisted 1 July 2025) Vasopressin receptor antagonists				
TOLVAPTAN – Special Authority see SA2166 below – Retail phar Tab 15 mg Tab 30 mg Tab 45 mg + 15 mg Tab 60 mg + 30 mg Tab 90 mg + 30 mg Tab 90 mg + 30 mg	873.50 873.50 1,747.00 1,747.00	28 OF 28 OF 56 OF 56 OF 56 OF		Jinarc Jinarc Jinarc Jinarc Jinarc

#### ⇒SA2166 Special Authority for Subsidy

**Initial application** — (autosomal dominant polycystic kidney disease) only from a renal physician or any relevant practitioner on the recommendation of a renal physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Patient has a confirmed diagnosis of autosomal dominant polycystic kidney disease; and
- 2 Patient has an estimated glomerular filtration rate (eGFR) of greater than or equal to 25 ml/min/1.73 m² at treatment initiation; and
- 3 Either:
  - 3.1 Patient's disease is rapidly progressing, with a decline in eGFR of greater than or equal to 5 mL/min/1.73 m² within one-year; or
  - 3.2 Patient's disease is rapidly progressing, with an average decline in eGFR of greater than or equal to 2.5 mL/min/1.73 m² per year over a five-year period.

Renewal — (autosomal dominant polycystic kidney disease) only from a renal physician or any relevant practitioner on the recommendation of a renal physician. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 Patient has not developed end-stage renal disease, defined as an eGFR of less than 15 mL/min/1.73 m²; and
- 2 Patient has not undergone a kidney transplant.

	Subsidy (Manufacturer's Price) \$	Per	Fully Brand or Subsidised Generic ✓ Manufacturer	
Lipid-Modifying Agents				
Fibrates				
BEZAFIBRATE ★ Tab 200 mg ★ Tab long-acting 400 mg		90 30	<ul> <li>✓ <u>Bezalip</u></li> <li>✓ <u>Bezalip</u> Retard</li> </ul>	<u>1</u>
Other Lipid-Modifying Agents				
ACIPIMOX ₭ Cap 250 mg		30	<ul> <li>Olbetam</li> </ul>	
Resins				
COLESTYRAMINE Powder for oral suspension 4 g sachet	61.50	50	<ul> <li>Colestyramine Mylan S29</li> <li>Quantalan sug free S29</li> </ul>	
HMG CoA Reductase Inhibitors (Statins)				
ATORVASTATIN Tab 10 mg	0.31 5.16	30 500	✓ <u>Lorstat</u> ✓ Lorstat	
Tab 20 mg	0.45 8.12	28 500	<ul> <li>✓ Lipitor</li> <li>✓ Lorstat</li> </ul>	
Tab 40 mg Tab 80 mg		500 30 500	✓ Lorstat ✓ Lorstat ✓ Lorstat	
PRAVASTATIN ₭ Tab 20 mg	7 16	100	✓ Clinect	
🖌 Tab 40 mg		100	✓ <u>Clinect</u>	
ROSUVASTATIN – Special Authority see SA2093 below – Re ₭ Tab 5 mg ₭ Tab 10 mg ₭ Tab 20 mg ₭ Tab 40 mg	1.29 1.69 2.71	30 30 30 30	<ul> <li>✓ <u>Rosuvastatin</u></li> <li>✓ <u>Rosuvastatin</u></li> <li>✓ <u>Rosuvastatin</u></li> <li>✓ <u>Rosuvastatin</u></li> </ul>	Viatris Viatris

### ► SA2093 Special Authority for Subsidy

Initial application — (cardiovascular disease risk) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Either:

1 Both:

- 1.1 Patient is considered to be at risk of cardiovascular disease; and
- 1.2 Patient is Māori or any Pacific ethnicity; or

2 Both:

2.1 Patient has a calculated risk of cardiovascular disease of at least 15% over 5 years; and

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🖌	Manufacturer	

continued...

2.2 LDL cholesterol has not reduced to less than 1.8 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initial application — (familial hypercholesterolemia) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

- Both:
  - 1 Patient has familial hypercholesterolemia (defined as a Dutch Lipid Criteria score greater than or equal to 6); and
  - 2 LDL cholesterol has not reduced to less than 1.8 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initial application — (established cardiovascular disease) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Both:

- 1 Any of the following:
  - 1.1 Patient has proven coronary artery disease (CAD); or
  - 1.2 Patient has proven peripheral artery disease (PAD); or
  - 1.3 Patient has experienced an ischaemic stroke; and
- 2 LDL cholesterol has not reduced to less than 1.4 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initial application — (recurrent major cardiovascular events) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Both:

- 1 Patient has experienced a recurrent major cardiovascular event (defined as myocardial infarction, ischaemic stroke, coronary revascularisation, hospitalisation for unstable angina) in the last 2 years; and
- 2 LDL cholesterol has not reduced to less than 1.0 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

#### SIMVASTATIN

*	Tab 10 mg1.68	90	<ul> <li>Simvastatin Mylan</li> </ul>
			<ul> <li>Simvastatin Viatris</li> </ul>
*	Tab 20 mg2.54	90	<ul> <li>Simvastatin Viatris</li> </ul>
*	Tab 40 mg4.11	90	<ul> <li>Simvastatin Viatris</li> </ul>
	Tab 80 mg8.81	90	<ul> <li>Simvastatin Viatris</li> </ul>

### **Selective Cholesterol Absorption Inhibitors**

EZETIMIBE * Tab 10 mg1.76	30	<ul> <li>Ezemibe Viatris</li> <li>Ezetimibe Sandoz</li> </ul>
(Ezemibe Viatris Tab 10 mg to be delisted 1 July 2025)		
EZETIMIBE WITH SIMVASTATIN		
Tab 10 mg with simvastatin 10 mg5.15	30	<ul> <li>Zimybe</li> </ul>
Tab 10 mg with simvastatin 20 mg6.15	30	<ul> <li>Zimybe</li> </ul>
Tab 10 mg with simvastatin 40 mg7.15	30	<ul> <li>Zimybe</li> </ul>
Tab 10 mg with simvastatin 80 mg8.15	30	<ul> <li>Zimybe</li> </ul>

	Subsidy		Fully Brand or
	(Manufacturer's		
	\$	Per	<ul> <li>Manufacturer</li> </ul>
Nitrates			
Nil ales			
GLYCERYL TRINITRATE			
✤ Oral pump spray, 400 mcg per dose – Up to 250 dose			
available on a PSO	7.48	250 dose OP	<ul> <li>Nitrolingual Pump</li> </ul>
			Spray
* Patch 25 mg, 5 mg per day		30	✓ Nitroderm TTS
* Patch 50 mg, 10 mg per day		30	<ul> <li>Nitroderm TTS</li> </ul>
ISOSORBIDE MONONITRATE	00.40	100	( Isono 00
<ul> <li>* Tab 20 mg</li> <li>* Tab long-acting 40 mg</li> </ul>		100 30	<ul> <li>✓ Ismo 20</li> <li>✓ Ismo 40 Retard</li> </ul>
* Tab long-acting 40 mg		30 90	✓ <u>Isilio 40 Retard</u> ✓ Duride
		50	• <u>Builde</u>
Sympathomimetics			
ADRENALINE			
Inj 1 in 1,000, 1 ml ampoule – Up to 5 inj available on a PS	SO 4.98	5	Aspen Adrenaline
	13.27	0	✓ DBL Adrenaline
	25.30	10	✓ Hamein S29
Inj 1 in 10,000, 10 ml ampoule - Up to 5 inj available on a	PSO27.00	5	<ul> <li>Hospira</li> </ul>
	49.00	10	<ul> <li>Aspen Adrenaline</li> </ul>
Vasodilators			
vasoullators			
HYDRALAZINE HYDROCHLORIDE			
* Tab 25 mg - Special Authority see SA1321 below - Retail			
pharmacy	CBS	1	<ul> <li>Hydralazine</li> </ul>
		56	<ul> <li>Onelink S29</li> </ul>
		84	AMDIPHARM \$29
		100	<ul> <li>Camber S29</li> </ul>
Inj 20 mg ampoule	25.90	5	<ul> <li>Apresoline</li> </ul>
SA1321 Special Authority for Subsidy			
Initial application from any relevant practitioner. Approvals va	alid without furthe	r renewal unless	notified for applications meeting
the following criteria:			
Eithe ave			
1 For the treatment of refractory hypertension; or	itrata in nationts	who are intolera	nt or have not responded to AC
<ol> <li>For the treatment of refractory hypertension; or</li> <li>For the treatment of heart failure in combination with a n</li> </ol>	itrate, in patients	who are intolera	nt or have not responded to AC
<ol> <li>For the treatment of refractory hypertension; or</li> <li>For the treatment of heart failure in combination with a n inhibitors and/or angiotensin receptor blockers.</li> </ol>	itrate, in patients	who are intolera	nt or have not responded to AC
<ol> <li>For the treatment of refractory hypertension; or</li> <li>For the treatment of heart failure in combination with a n inhibitors and/or angiotensin receptor blockers.</li> <li>MINOXIDIL</li> </ol>	·		
2 For the treatment of heart failure in combination with a n		60	✓ Minoxidil Roma S29
<ol> <li>For the treatment of refractory hypertension; or</li> <li>For the treatment of heart failure in combination with a n inhibitors and/or angiotensin receptor blockers.</li> <li>MINOXIDIL</li> <li>Tab 10 mg</li> </ol>	·		
<ol> <li>For the treatment of refractory hypertension; or</li> <li>For the treatment of heart failure in combination with a n inhibitors and/or angiotensin receptor blockers.</li> <li>MINOXIDIL</li> <li>Tab 10 mg</li> <li>NICORANDIL</li> </ol>	47.04 78.40	60 100	<ul> <li>✓ Minoxidil Roma ₅29</li> <li>✓ Loniten</li> </ul>
<ol> <li>For the treatment of refractory hypertension; or</li> <li>For the treatment of heart failure in combination with a n inhibitors and/or angiotensin receptor blockers.</li> <li>MINOXIDIL</li> <li>Tab 10 mg</li> <li>NICORANDIL</li> <li>Tab 10 mg</li> </ol>	47.04 78.40 21.73	60 100 60	<ul> <li>✓ Minoxidil Roma \$29</li> <li>✓ Loniten</li> <li>✓ Max Health</li> </ul>
<ol> <li>For the treatment of refractory hypertension; or</li> <li>For the treatment of heart failure in combination with a n inhibitors and/or angiotensin receptor blockers.</li> <li>MINOXIDIL</li> <li>Tab 10 mg</li> <li>MICORANDIL</li> <li>Tab 10 mg</li> <li>Tab 20 mg</li> </ol>	47.04 78.40 21.73	60 100	<ul> <li>✓ Minoxidil Roma ₅29</li> <li>✓ Loniten</li> </ul>
<ol> <li>For the treatment of refractory hypertension; or</li> <li>For the treatment of heart failure in combination with a n inhibitors and/or angiotensin receptor blockers.</li> <li>MINOXIDIL</li> <li>Tab 10 mg</li> <li>NICORANDIL</li> <li>Tab 10 mg</li> <li>Tab 20 mg</li> <li>PAPAVERINE HYDROCHLORIDE</li> </ol>	47.04 78.40 21.73 27.44	60 100 60 60	<ul> <li>Minoxidil Roma \$29</li> <li>Loniten</li> <li><u>Max Health</u></li> <li><u>Max Health</u></li> </ul>
<ol> <li>For the treatment of refractory hypertension; or</li> <li>For the treatment of heart failure in combination with a n inhibitors and/or angiotensin receptor blockers.</li> <li>MINOXIDIL</li> <li>Tab 10 mg</li> <li>NICORANDIL</li> <li>Tab 10 mg</li> <li>Tab 20 mg</li> <li>PAPAVERINE HYDROCHLORIDE</li> <li>* Inj 12 mg per ml, 10 ml ampoule</li> </ol>	47.04 78.40 21.73 27.44	60 100 60	<ul> <li>✓ Minoxidil Roma \$29</li> <li>✓ Loniten</li> <li>✓ Max Health</li> </ul>
<ol> <li>For the treatment of refractory hypertension; or</li> <li>For the treatment of heart failure in combination with a n inhibitors and/or angiotensin receptor blockers.</li> <li>MINOXIDIL</li> <li>Tab 10 mg</li> <li>NICORANDIL</li> <li>Tab 10 mg</li> <li>Mab 10 mg</li> <li>Tab 20 mg</li> <li>PAPAVERINE HYDROCHLORIDE</li> </ol>		60 100 60 60	<ul> <li>Minoxidil Roma \$29</li> <li>Loniten</li> <li><u>Max Health</u></li> <li><u>Max Health</u></li> </ul>

	Subsidy (Manufacturer's Price) \$	Sub Per	Fully osidised	Brand or Generic Manufacturer
Endothelin Receptor Antagonists				
AMBRISENTAN – Special Authority see SA2253 below – Reta Tab 5 mg Tab 10 mg SA2253 Special Authority for Subsidy nitial application — (PAH monotherapy) only from a respirator practitioner on the recommendation of a respiratory specialist, or applications meeting the following criteria: All of the following: 1 Patient has pulmonary arterial hypertension (PAH); and 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clir 3 PAH is in New York Heart Association/World Health Org		ologist. A	✓ A umatolog Approvals	s valid for 6 months for
<ul> <li>4 Any of the following:</li> <li>4.1 All of the following:</li> <li>4.1.1 PAH has been confirmed by right heart or</li> </ul>	atheterication: and			
<ul> <li>4.1.1 PAH has been confirmed by right heart ca</li> <li>4.1.2 A mean pulmonary artery pressure (PAPr</li> <li>4.1.3 A pulmonary capillary wedge pressure (P</li> <li>4.1.4 Pulmonary vascular resistance greater th cm⁻⁵); and</li> <li>4.1.5 Any of the following:</li> </ul>	n) greater than 20 mm CWP) less than or equa	al to 15 n	nmHg; ar	nd
4.1.5.1 PAH has been demonstrated to be	non-responsive in vaso			

- 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH 2022 (see note below for link to these guidelines) †; or
- 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
- 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
- 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
  - 5.1 Ambrisentan is to be used as PAH monotherapy; and
  - 5.2 Any of the following:
    - 5.2.1 Patient has experienced intolerable side effects with both sildenafil and bosentan; or
    - 5.2.2 Patient has an absolute contraindication to sildenafil and an absolute or relative contraindication to bosentan (e.g. due to current use of a combined oral contraceptive or liver disease); or
    - 5.2.3 Patient is a child with idiopathic PAH or PAH secondary to congenital heart disease.

**Initial application** — (PAH dual therapy) only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH has been confirmed by right heart catheterisation; and

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

Subsidy	Fully	Brand or
(Manufacturer's	Price) Subsidised	Generic
\$	Per 🗸	Manufacturer

- 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
- 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
- 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁵); and
- 4.1.5 Any of the following:
  - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH 2022 (see note below for link to these guidelines) †; or
  - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
  - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
- 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and

5 All of the following:

- 5.1 Ambrisentan is to be used as PAH dual therapy; and
- 5.2 Either:
  - 5.2.1 Patient has tried a PAH monotherapy (sildenafil or bosentan) for at least three months and has not experienced an acceptable response to treatment according to a validated risk stratification tool**; or
  - 5.2.2 Patient has tried PAH dual therapy including bosentan and has experienced intolerable side effects on bosentan; and
- 5.3 Both:
  - 5.3.1 Patient is presenting in NYHA/WHO functional class III or IV, and in the opinion of the treating clinician would benefit from initial dual therapy; and
  - 5.3.2 Patient has an absolute or relative contraindication to bosentan (e.g. due to current use of a combined oral contraceptive or liver disease).

**Initial application** — (PAH triple therapy) only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH has been confirmed by right heart catheterisation; and
    - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
    - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
    - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁵); and
    - 4.1.5 Any of the following:
      - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH 2022 (see note below for link to these guidelines) †; or
      - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or

continued...

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	I Generic	
\$	Per 🗸	Manufacturer	

- 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
- 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and

#### 5 Both:

- 5.1 Ambrisentan is to be used as PAH triple therapy; and
- 5.2 Any of the following:
  - 5.2.1 Patient is on the lung transplant list; or
  - 5.2.2 Both:
    - 5.2.2.1 Patient is presenting in NYHA/WHO functional class IV; and
    - 5.2.2.2 Patient has an absolute or relative contraindication to bosentan (e.g. due to current use of a combined oral contraceptive or liver disease); or
  - 5.2.3 Both:
    - 5.2.3.1 Patient has tried PAH dual therapy for at least three months and remains in an unacceptable risk category according to a validated risk stratification tool**; and
    - 5.2.3.2 Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario.

**Renewal** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 2 years where the patient is continuing to derive benefit from ambrisentan treatment according to a validated PAH risk stratification tool**.

Notes: † The European Respiratory Journal Guidelines can be found here: <u>2022 ECS/ERS Guidelines for the diagnosis and treatment of pulmonary hypertension PAH</u>

** the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

BOSENTAN - Special Authority see SA2254 below - Retail pharmacy

Tab 62.5 mg	100.00	60	<ul> <li><u>Bosentan Dr</u> Reddy's</li> </ul>
Tab 125 mg	100.00	60	✓ <u>Bosentan Dr</u> <u>Reddy's</u>

### ➡SA2254 Special Authority for Subsidy

**Initial application** — (PAH monotherapy) only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH has been confirmed by right heart catheterisation; and
    - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
    - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
    - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁵); and

Subsidy	Ful	y Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per 🔹	Manufacturer	

- 4.1.5 Any of the following:
  - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) †; or
  - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
  - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
- 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or
- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and

5 Both:

- 5.1 Bosentan is to be used as PAH monotherapy; and
- 5.2 Any of the following:
  - 5.2.1 Patient has experienced intolerable side effects on sildenafil; or
  - 5.2.2 Patient has an absolute contraindication to sildenafil; or
  - 5.2.3 Patient is a child with idiopathic PAH or PAH secondary to congenital heart disease.

**Initial application** — (PAH dual therapy) only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH has been confirmed by right heart catheterisation; and
    - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
    - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
    - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁵); and
    - 4.1.5 Any of the following:
      - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) †; or
      - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
      - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
  - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or
  - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Bosentan is to be used as part of PAH dual therapy; and
- 6 Either:
  - 6.1 Patient has tried a PAH monotherapy (sildenafil) for at least three months and has experienced an inadequate therapeutic response to treatment according to a validated risk stratification tool**; or
  - 6.2 Patient is presenting in NYHA/WHO functional class III or IV, and in the opinion of the treating clinician would likely

Subsidy	Fu	Illy Brand or	
(Manufacturer's	Price) Subsidis	ed Generic	
\$	Per	<ul> <li>Manufactu</li> </ul>	irer

benefit from initial dual therapy.

**Initial application** — (PAH triple therapy) only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH has been confirmed by right heart catheterisation; and
    - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
    - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
    - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁻⁵); and
    - 4.1.5 Any of the following:
      - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) †; or
      - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
      - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
  - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or
  - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
  - 5.1 Bosentan is to be used as part of PAH triple therapy; and
  - 5.2 Any of the following:
    - 5.2.1 Patient is on the lung transplant list; or
    - 5.2.2 Patient is presenting in NYHA/WHO functional class IV; or
    - 5.2.3 Both:
      - 5.2.3.1 Patient has tried PAH dual therapy for at least three months and has not experienced an acceptable response to treatment according to a validated risk stratification tool**; and
      - 5.2.3.2 Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario.

**Renewal** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 2 years where patient is continuing to derive benefit from bosentan treatment according to a validated PAH risk stratification tool**.

Notes: † The European Respiratory Journal Guidelines can be found here: <u>2022 ECS/ERS Guidelines for the diagnosis and</u> treatment of pulmonary hypertension PAH

** the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

## Phosphodiesterase Type 5 Inhibitors

SILDENAFIL - Special Authority see SA2255 on the next page - R	etail pharmacy		
Tab 25 mg	0.72	4	<ul> <li>Vedafil</li> </ul>
Tab 50 mg	1.45	4	✓ Vedafil
Tab 100 mg	11.22	12	✓ Vedafil

AThree months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

Subsidy		Fully	Brand or
(Manufacturer's Price)	5	Subsidised	Generic
\$	Per	~	Manufacturer

### ⇒SA2255 Special Authority for Subsidy

**Initial application** — (Raynaud's Phenomenon*) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has Raynaud's Phenomenon*; and
- 2 Patient has severe digital ischaemia (defined as severe pain requiring hospital admission or with a high likelihood of digital ulceration; digital ulcers; or gangrene); and
- 3 Patient is following lifestyle management (avoidance of cold exposure, sufficient protection, smoking cessation support, avoidance of sympathomimetic drugs); and
- 4 Patient is being treated with calcium channel blockers and nitrates (or these are contraindicated/not tolerated).

**Initial application — (Pulmonary arterial hypertension*)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH is confirmed by right heart catheterisation; and
    - 4.1.2 A mean pulmonary artery pressure (PAPm) of greater than 20 mmHg; and
    - 4.1.3 A pulmonary capillary wedge pressure (PCWP) that is less than or equal to 15 mmHg; and
    - 4.1.4 Pulmonary vascular resistance (PVR) of at least 2 Wood Units or greater than 160 International Units (dyn s cm⁵); and
    - 4.1.5 Any of the following:
      - 4.1.5.1 PAH is non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) †; or
      - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
      - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
  - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or
  - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures.

Initial application — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

60

- 1 Patient has a documented history of traumatic or non-traumatic spinal cord injury; and
- 2 Patient has erectile dysfunction secondary to spinal cord injury requiring pharmacological treatment.

Renewal — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Notes: Note: Indications marked with * are Unapproved Indications.

† The European Respiratory Journal Guidelines can be found here: <u>2022 ECS/ERS Guidelines for the diagnosis and treatment of</u> <u>pulmonary hypertension PAH</u>

** the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

	Subsidy (Manufacturer's Price) \$	Sı Per	Fully ubsidised	Brand or Generic Manufacturer
Prostacyclin Analogues				
EPOPROSTENOL – Special Authority see SA2256 below – Re Inj 500 mcg vial Inj 1.5 mg vial ■SA2256 Special Authority for Subsidy Initial application — (PAH dual therapy) only from a respirate practitioner on the recommendation of a respiratory specialist, of applications meeting the following criteria: All of the following:			✓ V umatologi	
<ol> <li>Patient has pulmonary arterial hypertension (PAH); and</li> <li>PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clir</li> <li>PAH is in New York Heart Association/World Health Org</li> <li>Any of the following:</li> </ol>	,		onal class	III or IV; and

- 4.1 All of the following:
  - 4.1.1 PAH has been confirmed by right heart catheterisation; and
  - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
  - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
  - 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁵); and
  - 4.1.5 Any of the following:
    - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) †; or
    - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
    - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
- 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 All of the following:
  - 5.1 Epoprostenol is to be used as part of PAH dual therapy with either sildenafil or an endothelin receptor antagonist; and
  - 5.2 Patient is presenting in NYHA/WHO functional class IV; and
  - 5.3 Patient has tried a PAH monotherapy for at least three months and remains in an unacceptable risk category according to a validated risk stratification tool.

**Initial application** — (PAH triple therapy) only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class III or IV; and
- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH has been confirmed by right heart catheterisation; and
    - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

- 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
- 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁵); and
- 4.1.5 Any of the following:
  - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) †; or
  - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
  - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
- 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and

5 Both:

- 5.1 Epoprostenol is to be used as PAH triple therapy; and
- 5.2 Any of the following:
  - 5.2.1 Patient is on the lung transplant list; or
  - 5.2.2 Patient is presenting in NYHA/WHO functional class IV; or
  - 5.2.3 Both:
    - 5.2.3.1 Patient has tried PAH dual therapy for at least three months and has not experienced an acceptable response to treatment according to a validated risk stratification tool; and
    - 5.2.3.2 Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario.

**Renewal** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 2 years where patient is continuing to derive benefit from epoprostenol treatment according to a validated PAH risk stratification tool**.

Notes: † The European Respiratory Journal Guidelines can be found here: <u>2022 ECS/ERS Guidelines for the diagnosis and</u> treatment of pulmonary hypertension PAH

** the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

ILOPROST - Special Authority see SA2257 below - Retail pharmacy

### ➡SA2257 Special Authority for Subsidy

**Initial application** — (PAH monotherapy) only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH has been confirmed by right heart catheterisation; and
    - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
    - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and

Subs	idy F	ully	Brand or
(Manufactur	er's Price) Subsid	ised	Generic
\$	Per	✓	Manufacturer

continued...

- 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁵); and
- 4.1.5 Any of the following:
  - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) †; or
  - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
  - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
- 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and

#### 5 Both:

- 5.1 Iloprost is to be used as PAH monotherapy; and
- 5.2 Either:
  - 5.2.1 Patient has experienced intolerable side effects on sildenafil and both the funded endothelin receptor antagonists (i.e. both bosentan and ambrisentan); or
  - 5.2.2 Patient has an absolute contraindication to sildenafil and an absolute or relative contraindication to endothelin receptor antagonists.

Initial application — (PAH dual therapy) only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH has been confirmed by right heart catheterisation; and
    - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
    - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
    - 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁵); and
    - 4.1.5 Any of the following:
      - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) †; or
      - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
      - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
  - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
  - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures ; and
- 5 All of the following:
  - 5.1 Iloprost is to be used as PAH dual therapy with either sildenafil or an endothelin receptor antagonist; and

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

- 5.2 Either:
  - 5.2.1 Patient has an absolute contraindication to or has experienced intolerable side effects on sildenafil; or
  - 5.2.2 Patient has an absolute or relative contraindication to or experienced intolerable side effects with a funded endothelin receptor antagonist; and
- 5.3 Either:
  - 5.3.1 Patient has tried a PAH monotherapy for at least three months and remains in an unacceptable risk category according to a validated risk stratification tool**; or
  - 5.3.2 Patient is presenting in NYHA/WHO functional class III or IV, and in the opinion of the treating clinician would benefit from initial dual therapy.

**Initial application — (PAH triple therapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH has been confirmed by right heart catheterisation; and
    - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
    - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
    - 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁵); and
    - 4.1.5 Any of the following:
      - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) †; or
      - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
      - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
  - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
  - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
  - 5.1 Iloprost is to be used as PAH triple therapy; and
  - 5.2 Any of the following:
    - 5.2.1 Patient is on the lung transplant list; or
    - 5.2.2 Patient is presenting in NYHA/WHO functional class IV; or
    - 5.2.3 Both:
      - 5.2.3.1 Patient has tried PAH dual therapy for at least three months and has not experienced an acceptable response to treatment according to a validated risk stratification tool**; and
      - 5.2.3.2 Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario.

**Renewal** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 2 years where patient is continuing to derive benefit from iloprost treatment according to a validated PAH risk stratification tool**.

continued...

 Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidi	sed	Generic
 <b>`</b> \$	Per	1	

continued...

Notes: † The European Respiratory Journal Guidelines can be found here: <u>2022 ECS/ERS Guidelines for the diagnosis and treatment of pulmonary hypertension PAH</u>

** the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

	Subsidy (Manufacturer's Price) \$	Sub Per	Fully osidised	Brand or Generic Manufacturer
Antiacne Preparations				
For systemic antibacterials, refer to INFECTIONS, Antibacterials,	page 93			
ADAPALENE				
a) Maximum of 30 g per prescription				
b) Only on a prescription				
Gel 0.1%		0 g OP	🗸 D	Differin
ISOTRETINOIN - Special Authority see SA2023 below - Retail pl	harmacy			
Cap 5 mg		60	✓ 0	Dratane
Cap 10 mg		120	✓ 0	Dratane
Cap 20 mg		120	<ul> <li>C</li> </ul>	Dratane

#### ⇒SA2023 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around isotretinoin and is competent to prescribe isotretinoin; and
- 3 Either:
  - 3.1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and that they must not become pregnant during treatment and for a period of one month after the completion of treatment; or
  - 3.2 Patient is not of child bearing potential.

**Renewal** from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and that they must not become pregnant during treatment and for a period of one month after the completion of treatment; or
- 2 Patient is not of child bearing potential.

#### TRETINOIN

Crm 0.5 mg per g $-$ Maximum of 50 g per prescrip	tion16.82	50 g OP	✓ <u>ReTrieve</u>	
Antibacterials Topical				
For systemic antibacterials, refer to INFECTIONS, Antit	pacterials, page 93			
HYDROGEN PEROXIDE * Crm 1%	8 56	10 g OP	✓ Crystaderm	
MUPIROCIN		10 9 01	• Crystaderin	
Oint 2%	6.60	15 g OP		
	(13.00)	-	Bactroban	
a) Only on a prescription				

b) Not in combination

				—
	Subsidy		Fully Brand or	
	(Manufacturer's F \$	Price) Subs Per	sidised Generic Manufacturer	
	φ	Fei	<ul> <li>Manufacturer</li> </ul>	
SODIUM FUSIDATE [FUSIDIC ACID]				
Crm 2%	1.69	5 g OP	Foban	
a) Maximum of 5 g per prescription				
b) Only on a prescription				
c) Not in combination			<b>4</b>	
Oint 2%	1.69	5 g OP	<ul> <li>Foban</li> </ul>	
<ul> <li>a) Maximum of 5 g per prescription</li> </ul>				
b) Only on a prescription				
c) Not in combination				
SULFADIAZINE SILVER				
Crm 1%	10.80	50 g OP	<ul> <li>Flamazine</li> </ul>	
	15.44		Ascend S29	
<ul> <li>a) Up to 250 g available on a PSO</li> </ul>				
<ul> <li>b) Not in combination</li> </ul>				
(Ascend S29 Crm 1% to be delisted 1 July 2025)				
· · · · ·				
Antifungals Topical				
For systemic antifungals, refer to INFECTIONS, Antifungals, pa	age 100			
AMOROLFINE				
a) Only on a prescription				
b) Not in combination				
Nail soln 5%	21.87	5 ml OP	<ul> <li>MycoNail</li> </ul>	
CLOTRIMAZOLE				
* Crm 1%	1.10	20 g OP	<ul> <li>Clomazol</li> </ul>	
a) Only on a prescription		- 5 -		
b) Not in combination				
* Soln 1%	4.36	20 ml OP		
	(7.55)		Canesten	
a) Only on a prescription	· · ·			
b) Not in combination				
ECONAZOLE NITRATE				
Crm 1%	8.04	20 g OP	Pevaryl	
a) Only on a prescription		g		
b) Not in combination				
c) Pevaryl to be Principal Supply on 1 June 2025				
Foaming soln 1%, 10 ml sachets		3		
······································	(18.64)	-	Pevaryl	
a) Only on a prescription	( )		,	
b) Not in combination				
· / · · · · · · · · · · · · · · · · · ·				

	Subsidy		Fully Brand or
	(Manufacturer's F \$	Price) Subs Per	sidised Generic Manufacturer
MICONAZOLE NITRATE	ψ	1 61	
* Crm 2%	0.90	15 g OP	<ul> <li>Multichem</li> </ul>
a) Only on a prescription			
b) Not in combination			
* Lotn 2%	4.36	30 ml OP	
	(10.03)		Daktarin
a) Only on a prescription			
<ul> <li>b) Not in combination</li> <li>Tinct 2%</li> </ul>	1.26	30 ml OP	
* TITCL 2 /8	(12.10)	30 IIII OF	Daktarin
a) Only on a prescription	(-=;)		
b) Not in combination			
Antipruritic Preparations			
CALAMINE			
a) Only on a prescription			
b) Not in combination			
Crm, aqueous, BP	3.45	100 g	healthE Calamine
haskh - Calenning Assessed to be Dringing! Complete			Aqueous
healthE Calamine Aqueous to be Principal Supply of	on T April 2025		
<ul><li>a) Only on a prescription</li><li>b) Not in combination</li></ul>			
Crm 10%	3.49	20 g OP	✓ Itch-Soothe
MENTHOL – Only in combination		_0 g 0.	<u></u>
<ol> <li>Only in combination with a dermatological base or</li> <li>With or without other dermatological galenicals.</li> </ol>	proprietary Topical C	Corticosteriod –	Plain
Crystals	6.02	25 a	✓ MidWest
Crystais	29.60	25 g 100 g	✓ MidWest
Corticosteroids Topical			
For systemic corticosteroids, refer to CORTICOSTEROIDS		NTS page 83	
Corticosteroids - Plain		NTO, page oo	
SETAMETHASONE DIPROPIONATE Crm 0.05%	2.06	15 g OP	✓ Diprosone
0111 0.00 %	36.00	50 g OP	✓ Diprosone
Oint 0.05%		15 g OP	✓ Diprosone
	36.00	50 g OP	<ul> <li>Diprosone</li> </ul>
Oint 0.05% in propylene glycol base	4.33	30 g OP	<ul> <li>Diprosone OV</li> </ul>
BETAMETHASONE VALERATE			
* Crm 0.1%		50 g OP	Beta Cream
* Oint 0.1%		50 g OP	<ul> <li>Beta Ointment</li> <li>Beta ouste</li> </ul>
Lotn 0.1% Betnovate to be Principal Supply on 1 May 2025		50 ml OP	<ul> <li>Betnovate</li> </ul>
CLOBETASOL PROPIONATE			
* Crm 0.05%	2.40	30 g OP	✓ Dermol
* Oint 0.05%		30 g OP	✓ Dermol

✓ fully subsidised Principal Supply

68

(\$29) Unapproved medicine supplied under Section 29 Sole Subsidised Supply

	<u> </u>			
	Subsidy (Manufacturer's P	rico) Sub	Fully	Brand or Generic
	(International Contents F	Per Suc		Manufacturer
CLOBETASONE BUTYRATE				
Crm 0.05%	5 29	30 g OP		
0111 0.05 /6	(10.00)	30 y OF		Eumovate
	(10.00)			Lunovale
HYDROCORTISONE	4 70			
* Crm 1% – Only on a prescription		30 g OP		Ethics
N. Develop Ochsin combination	20.40	500 g		Noumed
<ul> <li>Powder – Only in combination</li> <li>Up to 5% in a dermatological base (not proprietary Topic galenicals</li> </ul>	49.95 cal Corticosteriod	25 g – Plain) with		ABM out other dermatological
HYDROCORTISONE AND PARAFFIN LIQUID AND LANOLIN				
Lotn 1% with paraffin liquid 15.9% and lanolin 0.6% – Only	on			
a prescription		250 ml	1	DP Lotn HC
	12.00	200 111	•	
HYDROCORTISONE BUTYRATE		400 05		
Lipocream 0.1%		100 g OP		Locoid Lipocream
Oint 0.1%		100 g OP		Locoid
Milky emul 0.1%		100 ml OP	~	Locoid Crelo
METHYLPREDNISOLONE ACEPONATE				
Crm 0.1%	4.95	15 g OP	✓	Advantan
Oint 0.1%	4.95	15 g OP	✓	Advantan
MOMETASONE FUROATE				
Crm 0.1%		15 g OP	✓	Elocon Alcohol Free
	3.50	50 g OP		Elocon Alcohol Free
Oint 0.1%		15 g OP		Elocon
-	3.50	50 g OP		Elocon
Lotn 0.1%		30 ml OP		Elocon
TRIAMCINOLONE ACETONIDE				
Crm 0.02%	6 49	100 g OP	1	Aristocort
Oint 0.02%		100 g OP		Aristocort
On t 0.02 /0		100 9 01	•	Anatocon
Corticosteroids - Combination				
BETAMETHASONE VALERATE WITH SODIUM FUSIDATE [FU	ISIDIC ACID]			
Crm 0.1% with sodium fusidate (fusidic acid) 2%	3.49 (10.45)	15 g OP		Fucicort
<ul><li>a) Maximum of 15 g per prescription</li><li>b) Only on a prescription</li></ul>				
HYDROCORTISONE WITH MICONAZOLE - Only on a prescrip	otion			
* Crm 1% with miconazole nitrate 2%	2.85	15 g OP	✓	Micreme H
HYDROCORTISONE WITH NATAMYCIN AND NEOMYCIN – C Oint 1% with natamycin 1% and neomycin sulphate 0.5%		tion 15 g OP	1	Pimafucort
TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYC	IN AND NYSTAT	ΊN		
Crm 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 m				
and gramicidin 250 mcg per g - Only on a prescription .		15 g OP		
	(9.28)			Viaderm KC
	()			

	Subsidy (Manufacturer's P \$	rice) Subsi Per	Fully Brand or dised Generic ✓ Manufacturer
Barrier Creams and Emollients			
Barrier Creams			
DIMETHICONE			<b>4</b> • • • •
* Crm 5% pump bottle	4.30	500 ml OP	<ul> <li><u>healthE</u></li> <li>Dimethicone 5%</li> </ul>
* Crm 10% pump bottle	4.52	500 ml OP	<ul> <li>healthE</li> <li>Dimethicone 10%</li> </ul>
ZINC AND CASTOR OIL * Oint	4.05	500 a	✓ Evara
	4.25	500 g	
Emollients			
AQUEOUS CREAM * Crm	1 65	500 g	✓ <u>Evara</u>
CETOMACROGOL		500 g	
* Crm BP	2.29	500 g	<ul> <li><u>Cetomacrogol-AFT</u></li> </ul>
CETOMACROGOL WITH GLYCEROL Crm 90% with glycerol 10%	2 13	500 ml OP	✓ Evara
	3.50	1,000 ml OP	✓ Evara
EMULSIFYING OINTMENT * Oint BP	2 12	500 g	<ul> <li>Emulsifying</li> </ul>
		500 g	Ointment ADE
OIL IN WATER EMULSION	0.04	500 -	
* Crm	2.04 2.10	500 g	<ul> <li>Fatty Cream AFT</li> <li>Fatty Emulsion</li> </ul>
Fatty Emulsion Occom (Evers) to be Drivered Supply on a	1 April 0005		Cream (Evara)
Fatty Emulsion Cream (Evara) to be Principal Supply on (Fatty Cream AFT Crm to be delisted 1 April 2025)	1 April 2025		
PARAFFIN		500 00	
Oint liquid paraffin 50% with white soft paraffin 50%	4.94	500 g OP	<ul> <li>White Soft Liquid Paraffin AFT</li> </ul>
UREA			
* Crm 10% WOOL FAT WITH MINERAL OIL – Only on a prescription	1.37	100 g OP	healthE Urea Cream
* Lotn hydrous 3% with mineral oil	5.60	1,000 ml	
	(14.96) (20.53)		DP Lotion Alpha-Keri Lotion
	<b>1.40</b>	250 ml OP	
	(5.87) 5.60	1,000 ml	DP Lotion
	(23.91)		BK Lotion
	1.40 (7.73)	250 ml OP	BK Lotion
	· · · /		

	Subsidy		Fully Brand or
	(Manufacturer's F		idised Generic
	\$	Per	<ul> <li>Manufacturer</li> </ul>
Other Dermatological Bases			
PARAFFIN			
White soft – Only in combination	4.74	450 g	<ul> <li><u>EVARA White Soft</u></li> <li>Paraffin</li> </ul>
	19.00	2,500 g	✓ EVARA White Soft
Only in combination with a dermatological galenical or a	as a diluent for a p	proprietary Top	<u>Paraffin</u> ical Corticosteroid – Plain.
Minor Skin Infections			
OVIDONE IODINE			
Ovidone iodine Oint 10%	7 40	65 g OP	✓ Betadine
a) Maximum of 130 g per prescription		05 g 01	• Detaume
b) Only on a prescription			
Antiseptic Solution 10%		100 ml	✓ Riodine
Antiseptic soln 10%		15 ml	✓ Riodine
	6.99	500 ml	<ul> <li>Riodine</li> </ul>
Skin preparation, povidone iodine 10% with 30% alcohol	1.63	100 ml	
	(3.48)		Betadine Skin Prep
Parasiticidal Preparations			
IMETHICONE			
E Lotn 4%	4.25	200 ml OP	✓ <u>healthE</u> <u>Dimethicone 4%</u> <u>Lotion</u>
/ERMECTIN – Special Authority see SA2294 below – Retail p	harmacy		
Tab 3 mg - Up to 100 tab available on a PSO		4	<ul> <li>Stromectol</li> </ul>
1) PSO for institutional use only. Must be endorsed	with the name of	the institution f	or which the PSO is required ar
<ul><li>a valid Special Authority for patient of that instituti</li><li>2) Ivermectin available on BSO provided the BSO in</li><li>3) For the purposes of subsidy of ivermectin, instituti</li><li>facilities or prisons.</li></ul>	cludes a valid Sp		
SA2294 Special Authority for Subsidy nitial application — (Scabies) from any relevant practitioner. riteria: Either:	Approvals valid f	or 1 month for a	applications meeting the followi
1 The person has a severe scabies hyperinfestation (Crust 2 Both:	ted/ Norwegian sc	abies); or	
<ul><li>2.1 The person has a confirmed diagnosis of scabies</li><li>2.2 Either:</li></ul>	or is a close cont	act of a scabies	s case; and
2.2.1 The person is unable to complete topical the 2.2.2 Previous treatment with topical therapy ha	1.4.1	not cleared the	infestation.
nitial application — (Other parasitic infections) from any rel			

meeting the following criteria: Any of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Price)		sidised	Generic	
\$	Per	1	Manufacturer	

- 1 filariasis; or
- 2 cutaneous larva migrans (creeping eruption); or
- 3 strongyloidiasis.

**Renewal** — (Scabies) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria: Either:

1 The person has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or

2 Both:

- 2.1 The person has a confirmed diagnosis of scabies or is a close contact of a scabies case; and
- 2.2 Either:
  - 2.2.1 The person is unable to complete topical therapy; or
  - 2.2.2 Previous treatment with topical therapy has been tried and not cleared the infestation.

Renewal — (Other parasitic infections) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 filariasis; or
- 2 cutaneous larva migrans (creeping eruption); or
- 3 strongyloidiasis.

#### PERMETHRIN

Lotn 5%	30 ml OP	✓ <u>A-Scabies</u>	
---------	----------	--------------------	--

# **Psoriasis and Eczema Preparations**

ACITRETIN – Special Authority see SA2024 below – Retail pharmacy
------------------------------------------------------------------

Cap 10 mg	 60	✓ Novatretin
Cap 25 mg	60	✓ Novatretin

#### ⇒SA2024 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around acitretin and is competent to prescribe acitretin; and
- 3 Either:
  - 3.1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and that they must not become pregnant during treatment and for a period of three years after the completion of treatment; or
  - 3.2 Patient is not of child bearing potential.

**Renewal** from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and that they must not become pregnant during treatment and for a period of three years after the completion of treatment; or
- 2 Patient is not of child bearing potential.

#### BETAMETHASONE DIPROPIONATE WITH CALCIPOTRIOL

Foam spray 500 mcg with calcipotriol 50 mcg per g59.95	60 g OP	<ul> <li>Enstilar</li> </ul>
Gel 500 mcg with calcipotriol 50 mcg per g40.92	60 g OP	<ul> <li>Daivobet</li> </ul>
Oint 500 mcg with calcipotriol 50 mcg per g14.31	30 g OP	<ul> <li>Daivobet</li> </ul>

		_		
	Subsidy (Manufacturer's Pr \$	rice) Subs Per	Fully Brand or sidised Generic ✓ Manufacturer	
CALCIPOTRIOL				
Oint 50 mcg per g	40.00	120 g OP	<ul> <li>Daivonex</li> </ul>	
COAL TAR				
Soln BP – Only in combination		200 ml	<ul> <li>Midwest</li> </ul>	
<ol> <li>Up to 10% only in combination with a dermatological</li> <li>With or without other dermatological galenicals.</li> </ol>	al base or proprie	etary Topical (	Corticosteriod – Plain	
COAL TAR WITH ALLANTOIN, MENTHOL, PHENOL AND SULP	HUR			
Soln 5% with sulphur 0.5%, menthol 0.75%, phenol 0.5% and	l			
allantoin crm 2.5%		75 g OP		
	(8.00)		Egopsoryl TA	
	3.43	30 g OP		
	(4.35)		Egopsoryl TA	
COAL TAR WITH SALICYLIC ACID AND SULPHUR	4.07	05		
Soln 12% with salicylic acid 2% and sulphur 4% oint	4.97 7.95	25 g OP	<ul> <li>✓ Coco-Scalp</li> <li>✓ Coco-Scalp</li> </ul>	
		40 g OP	<ul> <li>Coco-Scalp</li> </ul>	
PIMECROLIMUS – Special Authority see SA1970 below – Retail	pnarmacy			
<ul> <li>a) Maximum of 15 g per prescription</li> <li>b) Note: a maximum of 15 g per prescription and no more the</li> </ul>	on one preserint	ion nor 12 wa	oko	
Cream 1%		15 a OP	Elidel	
SA1970 Special Authority for Subsidy		.e g e.		
of a dermatologist, paediatrician or ophthalmologist. Approvals va meeting the following criteria: Both: 1 Patient has atopic dermatitis on the eyelid; and 2 Patient has at least one of the following contraindications t documented epidermal atrophy, documented allergy to top	o topical corticos	steroids: perio	prificial dermatitis, rosacea	۱,
pressure.				
PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORES				
* Soln 2.3% with trolamine laurilsulfate and fluorescein sodium	5.41	500 ml	<ul> <li>Pinetarsol</li> </ul>	
SALICYLIC ACID	(		<b>.</b>	
Powder – Only in combination		250 g	<ul> <li>Midwest</li> </ul>	
<ol> <li>Only in combination with a dermatological base or p</li> <li>With or without other dermatological galenicals.</li> </ol>	proprietary Topic	al Corticostero	oid – Plain or collodion fle>	xible
SULPHUR				
Precipitated – Only in combination	6.35	100 g	<ul> <li>Midwest</li> </ul>	
<ol> <li>Only in combination with a dermatological base or p</li> <li>With or without other dermatological galenicals.</li> </ol>	proprietary Topic	al Corticostero	oid – Plain	
TACROLIMUS				
Oint 0.1% - Special Authority see SA2074 on the next page	_			
Retail pharmacy		30 g OP	✓ Zematop	
a) Maximum of 30 g per prescription		J	<b>+</b>	
b) Note: a maximum of 30 g per prescription and no mo	re than one pres	cription per 12	2 weeks.	

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

DERMATOLOGICALS

(Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer
--------------------------------------------------------------------

#### SA2074 Special Authority for Subsidy

**Initial application** only from a dermatologist, paediatrician or any relevant practitioner on the recommendation of a dermatologist, paediatrician, . Approvals valid without further renewal unless notified for applications meeting the following criteria: Both:

- 1 Patient has atopic dermatitis on the face; and
- 2 Patient has at least one of the following contraindications to topical corticosteroids: periorificial dermatitis, rosacea, documented epidermal atrophy or documented allergy to topical corticosteroids.

Scalp Preparations			
BETAMETHASONE VALERATE			
* Scalp app 0.1%		100 ml OP	<ul> <li>Beta Scalp</li> </ul>
CLOBETASOL PROPIONATE * Scalp app 0.05%	6.26	30 ml OP	✓ Dermol
HYDROCORTISONE BUTYRATE	0.20	50 mi Oi	• Dermor
Scalp lotn 0.1%	6.57	100 ml OP	✓ Locoid
KETOCONAZOLE			
Shampoo 2%		100 ml OP	Sebizole
a) Maximum of 100 ml per prescription	4.09		<ul> <li><u>Sebizole</u></li> </ul>
<ul> <li>b) Only on a prescription</li> </ul>			
Sunscreens			
SUNSCREENS, PROPRIETARY – Subsidy by endorsement Only if prescribed for a patient with severe photosensitivity s	econdary to a de	fined clinical co	ndition and the prescription is
endorsed accordingly.	6 50	200 g OP	<ul> <li>Marine Blue Lotion</li> </ul>
2011,		200 g 01	SPF 50+
Mout Dress and an			
Wart Preparations			
For salicylic acid preparations refer to PSORIASIS AND ECZEM	A PREPARATIO	NS, page 72	
PODOPHYLLOTOXIN			
Soln 0.5%		3.5 ml OP	<ul> <li>Condyline</li> </ul>
<ul> <li>a) Maximum of 3.5 ml per prescription</li> <li>b) Only on a prescription</li> </ul>			
<i>, ,</i> , ,			
Other Skin Preparations			
Antineoplastics			
FLUOROURACIL SODIUM			
Crm 5%	5.56	20 g OP	✓ Efudix
IMIQUIMOD			
Crm 5%, 250 mg sachet	21.72	24	<ul> <li>Perrigo</li> </ul>

Per ✓ Manufacturer
--------------------

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
Contraceptives - Non-hormonal				
Condoms				
ONDOMS				
€ 49 mm – Up to 144 dev available on a PSO		144	✓	Moments
53 mm	1.15	10	✓	Moments
	14.25	144	✓	Moments
<ul> <li>a) Maximum of 60 dev per prescription</li> </ul>				
<li>b) Up to 60 dev available on a PSO</li>				
53 mm, 0.05 mm thickness	1.15	10	✓	Moments
	14.25	144	✓	Moments
<ul> <li>a) Up to 60 dev available on a PSO</li> </ul>				
b) Maximum of 60 dev per prescription				
53 mm, chocolate, brown	1.15	10	1	Moments
	14.25	144	✓	Moments
a) Up to 60 dev available on a PSO				
b) Maximum of 60 dev per prescription				
53 mm, strawberry, red	1.15	10	1	Moments
	14.25	144	✓	Moments
a) Up to 60 dev available on a PSO				
b) Maximum of 60 dev per prescription				
€ 56 mm		10	1	Moments
	14.50	144		Moments
a) Maximum of 60 dev per prescription				
b) Up to 60 dev available on a PSO				
56 mm, 0.05 mm thickness	2 00	12	1	Gold Knight
	24.10	144		Gold Knight
a) Up to 60 dev available on a PSO	21.10		-	dola langit
b) Maximum of 60 dev per prescription				
56 mm, 0.05mm thickness (bulk pack)	20.17	144	1	Gold Knight
a) Maximum of 60 dev per prescription		1 7 7	•	dola kinght
b) Up to 60 dev available on a PSO				
56 mm, 0.08 mm thickness	1 15	10	1	Moments
	14.25	144		Moments
a) Lin to 60 day available and BCO	14.20	144	•	MONICIILS
a) Up to 60 dev available on a PSO				
b) Maximum of 60 dev per prescription	4.45	10		Moments
56 mm, 0.08 mm thickness, red		10 144		Moments
a) the te contain englishing on a DOO	14.25	144	•	woments
a) Up to 60 dev available on a PSO				
b) Maximum of 60 dev per prescription	1 70	10		Cold Knight
56 mm, chocolate		12	-	Gold Knight
	21.45	144	~	Gold Knight
a) Up to 60 dev available on a PSO				
b) Maximum of 60 dev per prescription	. ==			<b>6</b> 1 1 K 1 1 1
56 mm, strawberry		12		Gold Knight
	21.45	144	<i>✓</i>	Gold Knight
<ul> <li>a) Up to 60 dev available on a PSO</li> </ul>				
<ul> <li>b) Maximum of 60 dev per prescription</li> </ul>				
60 mm		12		Gold Knight XL
	21.89	144	1	Gold Knight XL
a) Maximum of 60 dev per prescription				
b) Uptin 6 upside allable on a PSO	S29 Unapproved	l med	icine supplie	ed under Section 29
6 60 mm Kalliparskippiy	Sdi7.78beidisod	c144	lv 🖌	Gold Knight XL

		Subsidy		Fully	Brand or
		(Manufacturer's Price)		sidised	Generic
		\$	Per	1	Manufacturer
a'	Maximum of 60 dev per prescription				
	Up to 60 dev available on a PSO				
Contra	ceptive Devices				
Contra	ceptive Devices				
INTRA-UT	ERINE DEVICE				
a) I Ir	to 40 dev available on a PSO				
, ,	ly on a PSO				
,	9.1 mm length × 23.2 mm width	20.00	4		hoice 380 7med
<b>木</b> 10D2	9. I IIIII leligui x 23.2 IIIII widui	29.00	I	• •	
					Nsha Silver/
					copper Short
* IUD 3	3.6 mm length × 29.9 mm width		1	✓ <u>T</u>	Cu 380 Plus
					Normal
* IUD 3	5.5 mm length × 19.6 mm width		1	✓ C	u 375 Standard
				_	

#### **Contraceptives - Hormonal**

#### Combined Oral Contraceptives

#### ⇒SA0500 Special Authority for Alternate Subsidy

**Initial application** from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

1 Either:

- 1.1 Patient is on a Social Welfare benefit; or
- 1.2 Patient has an income no greater than the benefit; and
- 2 Has tried at least one of the fully funded options and has been unable to tolerate it.

**Renewal** from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient is on a Social Welfare benefit; or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- on a Social Welfare benefit; or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

#### ETHINYLOESTRADIOL WITH DESOGESTREL

*	Tab 20 mcg with desogestrel 150 mcg and 7 inert tab - Up	to		
	84 tab available on a PSO		84	<ul> <li>Mercilon 28</li> </ul>

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
ETHINYLOESTRADIOL WITH LEVONORGESTREL				
* Tab 20 mcg with levonorgestrel 100 mcg and 7 inert tablets				
Up to 84 tab available on a PSO	1.50	84	✓ ]	Lo-Oralcon 20 ED
* Tab 30 mcg with levonorgestrel 150 mcg	6.62	63		
	(16.50)		I	Vicrogynon 30
<ul> <li>a) Higher subsidy of \$15.00 per 63 tab with Special Aut</li> <li>b) Up to 63 tab available on a PSO</li> <li>Tab 30 mcg with levonorgestrel 150 mcg and 7 inert tablets Up to 84 tab available on a PSO</li> </ul>	-	the p		ge Dralcon 30 ED
ETHINYLOESTRADIOL WITH NORETHISTERONE Tab 35 mcg with norethisterone 1 mg and 7 inert tab – Up to 84 tab available on a PSO		84		Alyacen Brevinor 1/28
Tab 35 mcg with norethisterone 500 mcg and 7 inert tab – U to 84 tab available on a PSO		84	✓	Norimin
Progestogen-only Contraceptives				

#### ⇒SA0500 Special Authority for Alternate Subsidy

**Initial application** from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

1 Either:

- 1.1 Patient is on a Social Welfare benefit; or
- 1.2 Patient has an income no greater than the benefit; and
- 2 Has tried at least one of the fully funded options and has been unable to tolerate it.

**Renewal** from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient is on a Social Welfare benefit; or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- on a Social Welfare benefit; or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

#### LEVONORGESTREL

* Tab 30 mcg – Up to 112 tab available on a PSO	22.00	112	<ul> <li>Microlut</li> </ul>
* Subdermal implant (2 × 75 mg rods) – Up to 3 pack available			
on a PSO	106.92	1	✓ Jadelle
MEDROXYPROGESTERONE ACETATE			
Inj 150 mg per ml, 1 ml syringe – Up to 5 inj available on a PSC	D 10.56	1	<ul> <li>Depo-Provera</li> </ul>

	Subsidy (Manufacturer's Price) \$	Per	Fully Brand or Subsidised Generic Manufacturer
NORETHISTERONE Tab 350 mcg – Up to 84 tab available on a PSO	12.25	84	<ul> <li>Norethinderone - CDC</li> <li>Noriday</li> <li>Noriday 28</li> </ul>
Emergency Contraceptives			
LEVONORGESTREL * Tab 1.5 mg a) Maximum of 2 tab per prescription	1.75	1	✓ <u>Levonorgestrel</u> <u>BNM</u>

b) Up to 5 tab available on a PSO

c) Note: Direct Provision by a pharmacist permitted under the provisions in Part I of Section A.

## Antiandrogen Oral Contraceptives

Prescribers may code prescriptions "contraceptive" (code "O") when used as indicated for contraception. The period of supply and prescription charge will be as per other contraceptives, as follows:

- A maximum \$5.00 prescription charge (patient co-payment) may apply.
- prescription may be written for up to six months supply.

Prescriptions coded in any other way are subject to any non contraceptive prescription charges that apply, and the non-contraceptive period of supply. ie. Prescriptions may be written for up to three months supply.

# CYPROTERONE ACETATE WITH ETHINYLOESTRADIOL

<ul> <li>Tab 2 mg with ethinyloestradiol 35 mcg and 7 inert tabs – Up to 168 tab available on a PSO5.08</li> </ul>	168	✓ <u>Ginet</u>
Gynaecological Anti-infectives		
ACETIC ACID WITH HYDROXYQUINOLINE AND RICINOLEIC ACID		
Jelly with glacial acetic acid 0.94%, hydroxyquinoline sulphate 0.025%, glycerol 5% and ricinoleic acid 0.75% with applicator8.43	100 g OP	
(24.87)	5 5 5	Aci-Jel
CLOTRIMAZOLE		
* Vaginal crm 1% with applicators	35 g OP	<ul> <li><u>Clomazol</u></li> </ul>
* Vaginal crm 2% with applicators	20 g OP	<ul> <li><u>Clomazol</u></li> </ul>
MICONAZOLE NITRATE		
* Vaginal crm 2% with applicator6.89	40 g OP	<ul> <li>Micreme</li> </ul>
NYSTATIN		
Vaginal crm 100,000 u per 5 g with applicator(s)5.70	75 g OP	✓ <u>Nilstat</u>
Myometrial and Vaginal Hormone Preparations		
ERGOMETRINE MALEATE		
Inj 500 mcg per ml, 1 ml ampoule – Up to 5 inj available on a		
PSO160.00	5	<ul> <li>DBL Ergometrine</li> </ul>
OESTRIOL		
* Crm 1 mg per g with applicator	15 g OP	✓ Ovestin
* Pessaries 500 mcg7.55	15	✓ Ovestin

AThree months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Pri		Fully Brand or dised Generic
	\$	Per	<ul> <li>Manufacturer</li> </ul>
OXVTOCIN Lin to E ini quailable on a BCO			
OXYTOCIN – Up to 5 inj available on a PSO	4.00	F	
Inj 5 iu per ml, 1 ml ampoule Inj 10 iu per ml, 1 ml ampoule		5 5	<ul> <li>✓ Oxytocin BNM</li> <li>✓ Oxytocin BNM</li> </ul>
Inj To iu per mi, T mi ampoule		-	✓ Oxytocin BNM ✓ Oxytocin
	11.96	10	,
			Panpharma
OXYTOCIN WITH ERGOMETRINE MALEATE - Up to 5 inj ava			
Inj 5 iu with ergometrine maleate 500 mcg per ml, 1 ml ampo	oule32.40	5	<ul> <li>Syntometrine</li> </ul>
Pregnancy Tests - hCG Urine			
BETA-HCG LOW SENSITIVITY URINE TEST KIT – Up to 15 te	st available on a P	SO	
Note: For use in abortion services only.		00	
Midstream	16.28	1 test OP	<ul> <li>CheckTop</li> </ul>
PREGNANCY TESTS - HCG URINE		1 1001 01	- onourop
a) Up to 200 test available on a PSO			
b) Only on a PSO	40.00		
Cassette		40 test OP	✓ <u>David One Step</u>
			Cassette
			Pregnancy Test
Urinary Agents			
offinally Agents			
For urinary tract Infections refer to INFECTIONS, Antibacterials,	page 112		
5-Alpha Reductase Inhibitors			
FINASTERIDE – Special Authority see SA0928 below – Retail p	hormooy		
* Tab 5 mg		100	✓ <u>Ricit</u>
	4.79	100	• <u>Hicit</u>
► SA0928 Special Authority for Subsidy			
Initial application from any relevant practitioner. Approvals vali	d without further re	enewal unless	notified for applications meeting
the following criteria: Both:			
1 Patient has symptomatic benign prostatic hyperplasia; an	a		
2 Either:			
2.1 The patient is intolerant of non-selective alpha blo			d; or
2.2 Symptoms are not adequately controlled with non-	selective alpha blo	ockers.	
Alpha-1A Adrenoreceptor Blockers			
TAMSULOSIN HYDROCHLORIDE - Special Authority see SA1	032 below – Betai	Inharmacy	
* Cap 400 mcg		100	<ul> <li>Tamsulosin-Rex</li> </ul>
		100	
► SA1032 Special Authority for Subsidy	al	امریک	a stiffe of few seconding times are stiffered
Initial application from any relevant practitioner. Approvals vali	a without further re	enewai uniéss	notified for applications meeting
the following criteria:			
Both:			

- Patient has symptomatic benign prostatic hyperplasia; and
   The patient is intolerant of non-selective alpha blockers or these are contraindicated.

	0.1.11		<b>-</b> "	<b>D</b> 1
	Subsidy (Manufacturer's Pric \$	ce) Sub Per	Fully sidised	Brand or Generic Manufacturer
Other Urinary Agents				
DXYBUTYNIN				
* Tab 5 mg	5.42	100	✓ A	llchemy Oxybutynin
POTASSIUM CITRATE				
Oral liq 3 mmol per ml – Special Authority see SA1083 b Retail pharmacy		200 ml OP	✔ В	liomed
nitial application from any relevant practitioner. Approvals				
Both: 1 The patient has recurrent calcium oxalate urolithiasis; 2 The patient has had more than two renal calculi in the Renewal from any relevant practitioner. Approvals valid for 2 penefitting from the treatment.	and two years prior to the	application.		
<ol> <li>The patient has recurrent calcium oxalate urolithiasis;</li> <li>The patient has had more than two renal calculi in the Renewal from any relevant practitioner. Approvals valid for 2 benefitting from the treatment.</li> <li>SODIUM CITRO-TARTRATE</li> </ol>	and two years prior to the a 2 years where the treat	application. Iment remain	ns appro	opriate and the patient
<ol> <li>The patient has recurrent calcium oxalate urolithiasis;</li> <li>The patient has had more than two renal calculi in the Renewal from any relevant practitioner. Approvals valid for 2 benefitting from the treatment.</li> <li>SODIUM CITRO-TARTRATE</li> <li>Grans eff 4 g sachets</li> </ol>	and two years prior to the a 2 years where the treat	application.		opriate and the patient
<ol> <li>The patient has recurrent calcium oxalate urolithiasis;</li> <li>The patient has had more than two renal calculi in the Renewal from any relevant practitioner. Approvals valid for 2 benefitting from the treatment.</li> <li>SODIUM CITRO-TARTRATE</li> </ol>	and two years prior to the a 2 years where the treat	application. Iment remain	ns appro ✓ <u>U</u>	opriate and the patient
<ol> <li>The patient has recurrent calcium oxalate urolithiasis;</li> <li>The patient has had more than two renal calculi in the Renewal from any relevant practitioner. Approvals valid for 2 benefitting from the treatment.</li> <li>SODIUM CITRO-TARTRATE</li> <li>Grans eff 4 g sachets</li> <li>SOLIFENACIN SUCCINATE</li> </ol>	and two years prior to the a 2 years where the treat	application. Iment remain 28	ns appro ✓ <u>U</u> ✓ S	ppriate and the patient Iral solifenacin succinate Max
<ol> <li>The patient has recurrent calcium oxalate urolithiasis;</li> <li>The patient has had more than two renal calculi in the Renewal from any relevant practitioner. Approvals valid for 2 benefitting from the treatment.</li> <li>SODIUM CITRO-TARTRATE</li> <li>Grans eff 4 g sachets</li> <li>SOLIFENACIN SUCCINATE</li> </ol>	and two years prior to the a e years where the treat 	application. Iment remain 28	ns appro ✓ <u>U</u> ✓ S	opriate and the patient Iral Solifenacin Succinate Max Health
<ol> <li>The patient has recurrent calcium oxalate urolithiasis;</li> <li>The patient has had more than two renal calculi in the Renewal from any relevant practitioner. Approvals valid for 2 benefitting from the treatment.</li> <li>SODIUM CITRO-TARTRATE</li> <li>Grans eff 4 g sachets</li> <li>SOLIFENACIN SUCCINATE Tab 5 mg</li> </ol>	and two years prior to the a e years where the treat 	application. Iment remain 28	ns appro	opriate and the patient Iral Solifenacin Succinate Max Health

Solifenacin succinate Max Health to be Principal Supply on 1 June 2025 (Solifenacin Viatris Tab 5 mg to be delisted 1 June 2025) (Solifenacin Viatris Tab 10 mg to be delisted 1 June 2025)

, j			
Detection of Substances in Urine			
ORTHO-TOLIDINE			
* Compound diagnostic sticks		50 test OP	
(	8.25)		Hemastix
TETRABROMOPHENOL			<b>*</b> • • • • •
* Blue diagnostic strips1	3.92	100 test OP	<ul> <li>Albustix</li> </ul>
Obstetric Preparations			
Antiprogesterones			
MIFEPRISTONE			
Tab 200 mg – Up to 15 tab available on a PSO	3.90	1	<ul> <li>Mifegyne</li> </ul>
	0.00	3	✓ Mifegyne

	Subsidy (Manufacturer's Price)	Su	Fully Brand or Subsidised Generic	
	\$	Per	1	Manufacturer
Calcium Homeostasis				
CALCITONIN				
* Inj 100 iu per ml, 1 ml ampoule	121.00	5	🗸 M	iacalcic
CINACALCET – Special Authority see SA2170 below – Retail ph	armacy			
Tab 30 mg – Wastage claimable		28	✓ C	inacalet Devatis
Tab 60 mg – Wastage claimable		28	✓ C	inacalet Devatis
SA2170 Special Authority for Subsidy			_	
nitial application — (parathyroid carcinoma or calciphylaxis)	only from a nephrol	ogist or	endocrinc	logist. Approvals valid f
months for applications mosting the following evitaria:		-		

6 months for applications meeting the following criteria:

Fither:

- 1 All of the following:
  - 1.1 The patient has been diagnosed with a parathyroid carcinoma (see Note); and
  - 1.2 The patient has persistent hypercalcaemia (serum calcium greater than or equal to 3 mmol/L) despite previous first-line treatments including sodium thiosulfate (where appropriate) and bisphosphonates; and
  - 1.3 The patient is symptomatic; or
- 2 All of the following:
  - 2.1 The patient has been diagnosed with calciphylaxis (calcific uraemic arteriolopathy); and
  - 2.2 The patient has symptomatic (e.g. painful skin ulcers) hypercalcaemia (serum calcium greater than or equal to 3 mmol/L): and
  - 2.3 The patient's condition has not responded to previous first-line treatments including bisphosphonates and sodium thiosulfate.

Renewal --- (parathyroid carcinoma or calciphylaxis) only from a nephrologist or endocrinologist. Approvals valid without further renewal unless notified for applications meeting the following criteria: Both:

- 1 The patient's serum calcium level has fallen to < 3mmol/L; and
- 2 The patient has experienced clinically significant symptom improvement.
- Note: This does not include parathyroid adenomas unless these have become malignant.

Initial application - (primary hyperparathyroidism) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has primary hyperparathyroidism; and
- 2 Either:
  - 2.1 Patient has hypercalcaemia of more than 3 mmol/L with or without symptoms: or
  - 2.2 Patient has hypercalcaemia of more than 2.85 mmol/L with symptoms; and
- 3 Surgery is not feasible or has failed; and
- 4 Patient has other comorbidities, severe bone pain, or calciphylaxis.

Initial application — (secondary or tertiary hyperparathyroidism) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

- All of the following:
  - 1 Either:

82

- 1.1 Patient has tertiary hyperparathyroidism and markedly elevated parathyroid hormone (PTH) with hypercalcaemia:
- 1.2 Patient has symptomatic secondary hyperparathyroidism and elevated PTH: and
- 2 Patient is on renal replacement therapy: and
- 3 Any of the following:
  - 3.1 Residual parathyroid tissue has not been localised despite repeat unsuccessful parathyroid explorations; or

continued...

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
continued				
<ul><li>3.2 Parathyroid tissue is surgically inaccessible; or</li><li>3.3 Parathyroid surgery is not feasible.</li></ul>				
Renewal — (secondary or tertiary hyperparathyroidism) from applications meeting the following criteria: Either:	any relevant practi	ioner.	Approvals	valid for 12 months for
<ol> <li>The patient has had a kidney transplant, and following a tre parathyroid hormone (PTH) level to support ongoing cessa</li> <li>The patient has not received a kidney transplant and trial o</li> </ol>	tion of treatment ha	s not l	been reach	ed; or
ZOLEDRONIC ACID				
lnj 4 mg per 5 ml, vial	15.65	1	✓ <u>;</u>	<u>Zoledronic acid</u> <u>Viatris</u>
Corticosteroids and Related Agents for Systemic	c Use			
BETAMETHASONE SODIUM PHOSPHATE WITH BETAMETHAS	SONE ACETATE			
Inj 3.9 mg with betamethasone acetate 3 mg per ml, 1 ml		5		
	(36.96)		(	Celestone Chronodose
EXAMETHASONE ₭ Tab 0.5 mg – Up to 60 tab available on a PSO	1.80	30	~ 1	Dexmethsone
<ul> <li>Tab 4 mg – Up to 30 tab available on a PSO</li> </ul>		30		Dexmethsone
Oral liq 1 mg per ml		5 ml C		Biomed
EXAMETHASONE PHOSPHATE Dexamethasone phosphate injection will not be funded for ora	Il use.			
k Inj 4 mg per ml, 1 ml ampoule – Up to 5 inj available on a PS		10	-	Hameln
k Inj 4 mg per ml, 2 ml ampoule − Up to 5 inj available on a PS	0 13.10	10	✓ 1	Hameln
EUDROCORTISONE ACETATE ₭ Tab 100 mcg	11.46	100	~	Florinef
YDROCORTISONE		100	• !	
★ Tab 5 mg	8.10	100	<b>√</b>	Douglas
k Tab 20 mg		100		Douglas
Inj 100 mg vial		1		Solu-Cortef
<ul><li>a) Not on a BSO</li><li>b) Up to 5 inj available on a PSO</li></ul>				
METHYLPREDNISOLONE	110.00	400		Mar al 1
₭ Tab 4 mg ₭ Tab 100 mg		100 20	-	Medrol Medrol
IETHYLPREDNISOLONE (AS SODIUM SUCCINATE)		20		
Inj 40 mg vial	22.30	1	<b>√</b> 9	Solu-Medrol-Act- O-Vial
Inj 125 mg vial	34.10	1	<b>√</b> 9	Solu-Medrol-Act- O-Vial
Inj 500 mg vial	43.01	1	<b>√</b> 9	Solu-Medrol-Act- O-Vial

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

	Subsidy	<u>)</u>	Fully	Brand or
	(Manufacturer's Pri \$	Ce) Subs	sidised ✓	Generic Manufacturer
ETHYLPREDNISOLONE ACETATE				
Inj 40 mg per ml, 1 ml vial	47.06	5	1	Depo-Medrol
REDNISOLONE		· ·		
<ul> <li>Oral liq 5 mg per ml – Up to 30 ml available on a PSO Restricted to children under 12 years of age.</li> </ul>	6.00	30 ml OP	•	Redipred
REDNISONE				
🗧 Tab 1 mg		500	1	Prednisone Clinect
🗧 Tab 2.5 mg	21.04	500	✓	Prednisone Clinect
<ul> <li>Tab 5 mg – Up to 30 tab available on a PSO</li> </ul>		500	✓	Prednisone Clinect
<ul> <li>Tab 20 mg – Up to 30 tab available on a PSO</li> </ul>	50.51	500	✓	Prednisone Clinect
ETRACOSACTRIN				
Inj 250 mcg per ml, 1 ml ampoule		1	✓	Synacthen
				UK Synacthen
Inj 1 mg per ml, 1 ml ampoule	690.00	1	1	Synacthen Depot
			✓	Synacthene
				Retard S29
RIAMCINOLONE ACETONIDE				
Inj 10 mg per ml, 1 ml ampoule	21 42	5	1	Kenacort-A 10
Inj 40 mg per ml, 1 ml ampoule		5		Kenacort-A 40
		Ū		Itenucont A 40
Sex Hormones Non Contraceptive				
Androgen Agonists and Antagonists				
YPROTERONE ACETATE				
Tab 50 mg	17.05	50	1	Siterone
Tab 100 mg		50		Siterone
ESTOSTERONE				••
Gel (transdermal) 16.2 mg per g	52.00	88 g OP	1	Testogel
		00 y OF	•	Testoger
ESTOSTERONE CIPIONATE				
Inj 100 mg per ml, 10 ml vial		1	~	Depo-Testosterone
ESTOSTERONE ESTERS				
Inj 250 mg per ml, 1 ml		1	✓	Sustanon Ampoules
ESTOSTERONE UNDECANOATE				
Cap 40 mg – Subsidy by endorsement	36.00	100	1	Steril-Gene S29
Subsidy by endorsement – subsidised for patients who				
1 November 2021 and the prescription is endorsed ac				
where there exists a record of prior dispensing of testo				
more more exists a record of prior disperiolity of toolo		1 1		Reandron 1000

		Subsidy		Fully	Brand or
		(Manufacturer's F \$	Price) Sub Per	sidised	Generic Manufacturer
		Ŷ	1.01	-	Manufacturor
Hormon	e Replacement Therapy - Systemic				
Oestrog	ens				
OESTRADIO					
✤ Tab 1 m	ng		28 OP	-	'atuatana
✤ Tab 2 m	ng	(11.10)	28 OP	E	strofem
* 1au 2 11	ıy	(11.10)	20 UF	F	strofem
* Gel (tra	nsdermal) 0.06% (750 mcg/actuation)	( -)	80 g OP		strogel
	5 mcg per day		8		stradiol TDP Mylan
		13.50			straderm MX S29
		14.50		_	stradot
		21.35		🗸 L	vllana
a)	No more than 2 patch per week				
	Only on a prescription				
Patch 5	0 mcg per day		8	🖌 E	stradiol TDP Mylan
				🗸 E	stradiol Viatris
		14.50		🖌 E	straderm MX S29
					stradiol Sandoz
					stradot
		21.55		✓ L	yllana
,	No more than 2 patch per week				
	Only on a prescription				
Patch 7	5 mcg per day		8		stradiol TDP Mylan
		14.50			stradiol Viatris stradiol Sandoz
		14.50			stradot
		22.37			yllana
a)	No more than 2 patch per week	22.07			y nunu
,	Only on a prescription				
	00 mcg per day		8	✓ E	stradiol TDP Mylan
					stradiol Viatris
		14.50		🖌 E	stradiol Sandoz
				🗸 E	stradot
		15.50		🖌 E	straderm MX S29
		22.77		🗸 L	yllana
a)	No more than 2 patch per week				
b)	Only on a prescription				
OESTRADIO	OL VALERATE				
	ng		84		rogynova
* Tab 2 m	ng		84	🗸 b	rogynova
OESTROGE					
* Conjuga	ated, equine tab 300 mcg	3.01	28		
		(19.25)		Р	remarin
* Conjuga	ated, equine tab 625 mcg		28		
		(19.25)		Р	remarin

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price \$	) Sub Per	Fully Brand or sidised Generic ✓ Manufacturer
Progestogens			
MEDROXYPROGESTERONE ACETATE			<b>4</b> -
* Tab 2.5 mg	6.56 8.75	30 56	<ul> <li>Provera</li> <li>Provera</li> </ul>
* Tab 5 mg		56	✓ Provera
No. Tale 40 years	20.13	100	<ul> <li>Provera</li> </ul>
* Tab 10 mg		30	Provera
Progestogen and Oestrogen Combined Prepa	rations		
OESTRADIOL WITH NORETHISTERONE			
* Tab 1 mg with 0.5 mg norethisterone acetate		28 OP	
Y Tab 0 may with 1 manager thistory as a state	(18.10)		Kliovance
* Tab 2 mg with 1 mg norethisterone acetate	5.40 (18.10)	28 OP	Kliogest
* Tab 2 mg with 1 mg norethisterone acetate (10), and 2 mg	( )		Nilogost
oestradiol tab (12) and 1 mg oestradiol tab (6)		28 OP	
	(18.10)		Trisequens
Other Oestrogen Preparations			
OESTRIOL			
* Tab 2 mg	7 70	30	✓ Ovestin
Other Progestogen Preparations			<u></u>
other Progestogen Preparations			
LEVONORGESTREL			<b>6</b>
<ul> <li>Intra-uterine device 52 mg.</li> <li>Intra-uterine device 52 mg.</li> </ul>		1	✓ Mirena
* Intra-uterine device 13.5 mg	215.60	1	Jaydess
MEDROXYPROGESTERONE ACETATE	199 57	100	Provera HD
Tab 100 mg		100	
NORETHISTERONE * Tab 5 mg – Up to 30 tab available on a PSO	5 /0	30	Primolut N
PROGESTERONE		50	
* Cap 100 mg	14.85	30	✓ Utrogestan
Thyroid and Antithyroid Agents			
CARBIMAZOLE			
* Tab 5 mg		100	✓ Neo-Mercazole
LEVOTHYROXINE			
* Tab 25 mcg	5.55	90	<ul> <li>Synthroid</li> </ul>
* Tab 50 mcg		28	<ul> <li>Mercury Pharma</li> </ul>
	5.79	90	<ul> <li>Synthroid</li> </ul>
* Tablet 50 mag	64.28	1,000	<ul> <li>Eltroxin</li> </ul>
* Tablet 50 mcg * Tab 100 mcg		200 28	<ul> <li>Eltroxin</li> <li>Mercury Pharma</li> </ul>
* Tab 100 mog	6.01	20 90	<ul> <li>✓ Synthroid</li> </ul>
	66.78	1,000	✓ Eltroxin
* Tablet 100 mcg	13.36	200	<ul> <li>Eltroxin</li> </ul>

	Subsidy (Manufacturer's Price) \$	Su Per	Fully Ibsidised	Brand or Generic Manufacturer
PROPYLTHIOURACIL – Special Authority see SA1199 below –	Retail pharmacy			
Tab 50 mg		100	🗸 b.	TU \$29

#### ⇒SA1199 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 The patient has hyperthyroidism; and
- 2 The patient is intolerant of carbimazole or carbimazole is contraindicated.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

## **Trophic Hormones**

#### Growth Hormones

SOMATROPIN (OMNITROPE) – Special Authority see SA2032 below –	Retail pharmad	су
* Inj 5 mg cartridge	30.21	1 <b>V</b> Omnitrope
* Inj 10 mg cartridge		1      Omnitrope
* Inj 15 mg cartridge		1 <b>V</b> Omnitrope

#### ⇒SA2032 Special Authority for Subsidy

Initial application — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria: Either:

- 1 Growth hormone deficiency causing symptomatic hypoglycaemia, or with other significant growth hormone deficient sequelae (e.g. cardiomyopathy, hepatic dysfunction) and diagnosed with GH < 5 mcg/l on at least two random blood samples in the first 2 weeks of life, or from samples during established hypoglycaemia (whole blood glucose < 2 mmol/l using a laboratory device); or
- 2 All of the following:
  - Height velocity < 25th percentile for age adjusted for bone age/pubertal status if appropriate over 6 or 12 months using the standards of Tanner and Davies (1985); and
  - 2.2 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
  - 2.3 Peak growth hormone value of < 5.0 mcg per litre in response to two different growth hormone stimulation tests. In children who are 5 years or older, GH testing with sex steroid priming is required; and</p>
  - 2.4 If the patient has been treated for a malignancy, they should be disease free for at least one year based upon follow-up laboratory and radiological imaging appropriate for the malignancy, unless there are strong medical reasons why this is either not necessary or appropriate; and
  - 2.5 Appropriate imaging of the pituitary gland has been obtained.

Renewal — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

- All of the following:
  - 1 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
  - 2 Height velocity is greater than or equal to 25th percentile for age (adjusted for bone age/pubertal status if appropriate) while on growth hormone treatment, as calculated over six months using the standards of Tanner and Davis (1985); and
  - 3 Height velocity is greater than or equal to 2.0 cm per year, as calculated over 6 months; and
  - 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone treatment has occurred; and
  - 5 No malignancy has developed since starting growth hormone.

Initial application — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

continued...

AThree months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

Subsidy (Manufacturer's Pr	ice)	Fully Subsidised	Brand or Generic	
\$	Per	1	Manufacturer	

#### continued...

All of the following:

- 1 The patient has a post-natal genotype confirming Turner Syndrome; and
- 2 Height velocity is < 25th percentile over 6-12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is < 14 years.

Renewal — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile for age (while on growth hormone calculated over 6 to 12 months using the Ranke's Turner Syndrome growth velocity charts); and
- 2 Height velocity is greater than or equal to 2 cm per year, calculated over six months; and
- 3 A current bone age is 14 years or under ; and
- 4 No serious adverse effect that the specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria: All of the following:

- 1 The patient's height is more than 3 standard deviations below the mean for age or for bone age if there is marked growth acceleration or delay; and
- 2 Height velocity is < 25th percentile for age (adjusted for bone age/pubertal status if appropriate), as calculated over 6 to 12 months using the standards of Tanner and Davies(1985); and
- 3 A current bone age is < 14 years or under (female patients) or < 16 years (male patients); and
- 4 The patient does not have severe chronic disease (including malignancy or recognized severe skeletal dysplasia) and is not receiving medications known to impair height velocity.

**Renewal** — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred.

Initial application — (short stature due to chronic renal insufficiency) only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 2 standard deviations below the mean; and
- 2 Height velocity is < 25th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is to 14 years or under (female patients) or to 16 years or under (male patients); and
- 4 The patient is metabolically stable, has no evidence of metabolic bone disease and absence of any other severe chronic disease; and
- 5 The patient is under the supervision of a specialist with expertise in renal medicine; and
- 6 Either:

88

6.1 The patient has a GFR less than or equal to 30 ml/min/1.73m² as measured by the Schwartz method (Height(cm)/plasma creatinine (umol/l)) × 40 = corrected GFR (ml/min/1.73m²) in a child who may or may not be receiving dialysis; or

continued...

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	

continued...

6.2 The patient has received a renal transplant and has received < 5mg/ m²/day of prednisone or equivalent for at least 6 months..

Renewal — (short stature due to chronic renal insufficiency) only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not experienced significant biochemical or metabolic deterioration confirmed by diagnostic results; and
- 7 The patient has not received renal transplantation since starting growth hormone treatment; and
- 8 If the patient requires transplantation, growth hormone prescription should cease before transplantation and a new application should be made after transplantation based on the above criteria.

**Initial application** — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of Prader-Willi syndrome that has been confirmed by genetic testing or clinical scoring criteria; and
- 2 The patient is aged six months or older; and
- 3 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
- 4 Sleep studies or overnight eximetry have been performed and there is no obstructive sleep disorder requiring treatment, or if an obstructive sleep disorder is found, it has been adequately treated under the care of a paediatric respiratory physician and/or ENT surgeon; and
- 5 Either:
  - 5.1 Both:
    - 5.1.1 The patient is aged two years or older; and
    - 5.1.2 There is no evidence of type II diabetes or uncontrolled obesity defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months; or
  - 5.2 The patient is aged between six months and two years and a thorough upper airway assessment is planned to be undertaken prior to treatment commencement and at six to 12 weeks following treatment initiation.

Renewal — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not developed type II diabetes or uncontrolled obesity as defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months.

Initial application - (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9

continued...

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

months for applications meeting the following criteria:

All of the following:

- 1 The patient has a medical condition that is known to cause growth hormone deficiency (e.g. surgical removal of the pituitary for treatment of a pituitary tumour); and
- 2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
- 3 The patient has severe growth hormone deficiency (see notes); and
- 4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
- 5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

Renewal — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 All of the following:
  - 1.1 The patient has been treated with somatropin for < 12 months; and
  - 1.2 There has been an improvement in Quality of Life defined as a reduction of at least 8 points on the Quality of Life Assessment of Growth Hormone Deficiency in Adults (QoL-AGHDA®) score from baseline; and
  - 1.3 Serum IGF-I levels have been increased within ±1SD of the mean of the normal range for age and sex; and
  - 1.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients, or 1 mg per day for female patients; or
- 2 All of the following:
  - 2.1 The patient has been treated with somatropin for more than 12 months; and
  - 2.2 The patient has not had a deterioration in Quality of Life defined as a 6 point or greater increase from their lowest QoL-AGHDA® score on treatment (other than due to obvious external factors such as external stressors); and
  - 2.3 Serum IGF-I levels have continued to be maintained within ±1SD of the mean of the normal range for age and sex (other than for obvious external factors); and
  - 2.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients or 1 mg per day for female patients; or
- 3 All of the following:
  - 3.1 The patient has had a Special Authority approval for somatropin for childhood deficiency in children and no longer meets the renewal criteria under this indication; and
  - 3.2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
  - 3.3 The patient has severe growth hormone deficiency (see notes); and
  - 3.4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
  - 3.5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

continued...

Subsidy	l	-ully	Brand or
(Manufacturer's Price)	Subsid	ised	Generic
 \$	Per	1	

continued...

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

#### **GnRH Analogues**

GOSERELIN			
Implant 3.6 mg, syringe	66.48	1	<ul> <li>Zoladex</li> </ul>
Implant 10.8 mg, syringe	138.23	1	✓ Zoladex
EUPRORELIN			
Additional subsidy by endorsement where the patient is a chi goserelin and the prescription is endorsed accordingly. Inj 3.75 mg prefilled dual chamber syringe – Higher subsidy		and is unable	to tolerate administration of
\$221.60 per 1 inj with Endorsement		1	
	(221.60)		Lucrin Depot 1-month
Inj 11.25 mg prefilled dual chamber syringe – Higher subside	( )		Edoin Doport Month
of \$591.68 per 1 inj with Endorsement		1	
	(591.68)	·	Lucrin Depot 3-month
	()		
Vasopressin Agonists			
DESMOPRESSIN			
Wafer 120 mcg	47.00	30	<ul> <li>Minirin Melt</li> </ul>
DESMOPRESSIN ACETATE			
Tab 100 mcg		30	<ul> <li>Minirin</li> </ul>
Tab 200 mcg		30	✓ Minirin
Nasal spray 10 mcg per dose		6 ml OP	<ul> <li>Desmopressin-</li> </ul>
			PH&T
Inj 4 mcg per ml, 1 ml	67.18	10	✓ Minirin
Other Endocrine Agents			
ABERGOLINE			
		0	✓ Dostinex
Tab 0.5 mg – Maximum of 2 tab per prescription; can be waived by Special Authority see SA2070 below	4 43	2	V Dostinex

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

1 Hyperprolactinemia; or

continued...

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Su Per	Fully Ibsidised	Brand or Generic Manufacturer
continued				
2 Acromegaly*; or				
3 Inhibition of lactation.				
Renewal — (for patients who have previously been funded practitioner. Approvals valid without further renewal unless noti which has expired and the treatment remains appropriate and the Note: Indication marked with * is an unapproved indication.	fied where the patient I	has prev	iously he	
CLOMIFENE CITRATE				
Tab 50 mg	29.84	10	🗸 N	lylan
				Clomiphen S29
METYRAPONE				
Cap 250 mg	558.00	50	🗸 N	letopirone

	Subsidy		Fully	Brand or
	(Manufacturer's Price)	Sul	osidised	Generic
		Per		Manufacturer
	\$	Per		Manufacturer
Anthelmintics				
ALBENDAZOLE - Special Authority see SA1318 below - Retai	Inharmacy			
Tab 400 mg		60	V F	skazole S29
	400.20	00	• -	
SA1318 Special Authority for Subsidy				
Initial application only from an infectious disease specialist or of	clinical microbiologist.	Approva	ıls valid f	or 6 months where the
patient has hydatids.				
Renewal only from an infectious disease specialist or clinical mi	crohiologist Approvs	ale valid fo	r 6 mont	he where the treatment
		lis valiu it		
remains appropriate and the patient is benefitting from the treatr	nent.			
MEBENDAZOLE – Only on a prescription				
Tab 100 mg	5 19	6	<b>1</b> V	ermox
0			• •	erniox
Oral liq 100 mg per 5 ml	2.18	15 ml		
	(7.83)		V	ermox
	· · · ·			
PRAZIQUANTEL			-	
Tab 600 mg		8	🗸 В	litricide
-				
Antibacterials				
Antibuotonuis				
a) For topical antibacterials, refer to DERMATOLOGICALS, page	20 66			
b) For anti-infective eye preparations, refer to SENSORY ORG.	ANS, page 266			
Or a had a second of a second of a second of a				
Cephalosporins and Cephamycins				
CEFACLOR MONOHYDRATE				
Cap 250 mg	25 85	100	V B	anbaxy-Cefaclor
Grans for oral lig 125 mg per 5 ml – Wastage claimable		100 ml		anbaxy-Cefaclor
Grans for oral lig 125 mg per 5 mi – Wastage claimable		100 111	• <u>n</u>	andaxy-celacion
CEFALEXIN				
Cap 250 mg	3.85	20	10	ephalexin ABM
		20		
Cap 500 mg				ephalexin ABM
Grans for oral liq 25 mg per ml – Wastage claimable	7.88	100 ml	🖌 F	lynn
Grans for oral lig 50 mg per ml – Wastage claimable		100 ml	🖌 F	lynn
· · · · · · · · · · · · · · · · · · ·	11.75			efalexin Sandoz
	11.75			cialexiii Gandoz
CEFAZOLIN – Subsidy by endorsement				
Only if prescribed for dialysis or cellulitis in accordance with	a Health NZ Hospital	approved	notoco	l and the prescription is
endorsed accordingly.		арр. ст с	. p. 0.000	
	0.00	-		
Inj 500 mg vial		5		efazolin-AFT
Inj 1 g vial	3.59	5	✓ 0	efazolin-AFT
Inj 2 g vial		5	✓ C	efazolin-AFT
		-	-	
CEFTRIAXONE – Subsidy by endorsement				
<ul> <li>a) Up to 10 inj available on a PSO</li> </ul>				
<ul> <li>b) Subsidised only if prescribed for a dialysis or cystic fibro</li> </ul>	cic nationt or the trea	tmont of	aonorrho	as or the treatment of
pelvic inflammatory disease, or the treatment of suspect	ed meningococcal dis	ease, and	a the pre	scription or PSO is
endorsed accordingly.				
Inj 500 mg vial	0.79	1	✓ C	eftriaxone-AFT
lnj 1 g vial		5	_	eftriaxone-AFT
		5	• •	
CEFUROXIME AXETIL – Subsidy by endorsement				
Only if prescribed for prophylaxis of endocarditis and the pre-	escription is endorsed	accordin	alv	
				acoud
Tab 250 mg		20	✓ A	scend-
				Cefuroxime S29

A Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

 $\ensuremath{\textbf{\#}}$  Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Subs Per	Fully idised	Brand or Generic Manufacturer
Macrolides				
AZITHROMYCIN – Maximum of 5 days treatment per prescriptio A maximum of 24 months of azithromycin treatment for non-or Authority. Tab 250 mg	cystic fibrosis bronch		l be su	
Tab 500 mg – Up to 8 tab available on a PSO		2		Zithromax
Grans for oral liq 200 mg per 5 ml (40 mg per ml) – Wastage claimable		15 ml	✓ Z	lithromax
SA1683 Special Authority for Waiver of Rule Initial application — (bronchiolitis obliterans syndrome, cyst	ic fibrosis and atyp	ical Mycol	bacteri	ium infections) only fro

a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following:

- 1 Patient has received a lung transplant, stem cell transplant, or bone marrow transplant and requires treatment for bronchiolitis obliterans syndrome*; or
- 2 Patient has received a lung transplant and requires prophylaxis for bronchiolitis obliterans syndrome*; or
- 3 Patient has cystic fibrosis and has chronic infection with Pseudomonas aeruginosa or Pseudomonas-related gram negative organisms*; or
- 4 Patient has an atypical Mycobacterium infection.
- Note: Indications marked with * are unapproved indications.

**Initial application — (non-cystic fibrosis bronchiectasis*)** only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 For prophylaxis of exacerbations of non-cystic fibrosis bronchiectasis*; and
- 2 Patient is aged 18 and under; and
- 3 Either:
  - 3.1 Patient has had 3 or more exacerbations of their bronchiectasis, within a 12 month period; or
  - 3.2 Patient has had 3 acute admissions to hospital for treatment of infective respiratory exacerbations within a 12 month period.

Note: Indications marked with * are unapproved indications.

Renewal — (non-cystic fibrosis bronchiectasis*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has completed 12 months of azithromycin treatment for non-cystic fibrosis bronchiectasis; and
- 2 Following initial 12 months of treatment, the patient has not received any further azithromycin treatment for non-cystic fibrosis bronchiectasis for a further 12 months, unless considered clinically inappropriate to stop treatment; and
- 3 The patient will not receive more than a total of 24 months' azithromycin cumulative treatment (see note).

The patient must not have had more than 1 prior approval.

Note: No further renewals will be subsidised. A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised. Indications marked with * are unapproved indications

CLARITHROMYCIN - Maximum of 500 mg per prescription; can be waived by Special Authority see SA1857 below

Tab 250 mg7.31	12	<ul> <li>Klaricid S29</li> </ul>
8.53	14	<ul> <li>Klacid</li> </ul>
Grans for oral liq 250 mg per 5 ml – Wastage claimable192.00	50 ml	<ul> <li>Klacid</li> </ul>

#### ⇒SA1857 Special Authority for Waiver of Rule

Initial application — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years for applications meeting the following criteria:

continued...

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🖌	Manufacturer

continued...

Either:

1 Atypical mycobacterial infection; or

2 Mycobacterium tuberculosis infection where there is drug-resistance or intolerance to standard pharmaceutical agents. **Initial application — (Helicobacter pylori eradication)** from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

1 For the eradication of helicobacter pylori in a patient unable to swallow tablets; and

2 For use only in combination with omeprazole and amoxicillin as part of a triple therapy regimen.

Initial application — (Prophylaxis of infective endocarditis) from any relevant practitioner. Approvals valid for 3 months where prophylaxis of infective endocarditis associated with surgical or dental procedures if amoxicillin is contra-indicated. Renewal — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment. ENTHROMYCIN (AS LACTORIONATE)

ERTTINOMICIN (AS LACTOBIONATE)			
Inj 1 g vial		1	Erythrocin IV
ERYTHROMYCIN ETHYL SUCCINATE			-
	25.00	100	<ul> <li>E-Mycin</li> </ul>
Tab 400 mg		100	
<ul> <li>a) Up to 20 tab available on a PSO</li> </ul>			
<li>b) Up to 2 x the maximum PSO quantity for RFPP</li>			
Grans for oral lig 200 mg per 5 ml	6.53	100 ml	E-Mycin
a) Up to 300 ml available on a PSO			
, ,			
b) Up to 2 x the maximum PSO quantity for RFPP			
c) Wastage claimable			
Grans for oral liq 400 mg per 5 ml	9.41	100 ml	E-Mycin
<ul> <li>a) Up to 200 ml available on a PSO</li> </ul>			
b) Wastage claimable			
, 6			
ROXITHROMYCIN			
Tab 150 mg	13.19	50	Arrow-
-			Roxithromycin
Tab 300 mg	25.00	50	Arrow-
· ~ · · · · · · · · · · · · · · · · · ·		50	Roxithromycin
			noxiunomycin

	Subsidy (Manufacturer's F		Fully Brand or idised Generic
Penicillins	\$	Per	Manufacturer
AMOXICILLIN Cap 250 mg	27 50	500	<ul> <li>Miro-Amoxicillin</li> </ul>
a) Up to 30 cap available on a PSO	27.50	500	
b) Up to 10 x the maximum PSO quantity for RFPP			
Cap 500 mg		500	<ul> <li>Miro-Amoxicillin</li> </ul>
a) Up to 30 cap available on a PSO			
b) Up to 10 x the maximum PSO quantity for RFPP			
Grans for oral liq 125 mg per 5 ml	2.22	100 ml	<ul> <li>Alphamox 125</li> </ul>
a) Up to 200 ml available on a PSO			
b) Wastage claimable			
Grans for oral liq 250 mg per 5 ml	2.81	100 ml	<ul> <li>Alphamox 250</li> </ul>
<ul> <li>a) Up to 300 ml available on a PSO</li> </ul>			
<li>b) Up to 10 x the maximum PSO quantity for RFPP</li>			
c) Wastage claimable			
Inj 250 mg vial		10	<ul> <li>Ibiamox</li> </ul>
Inj 500 mg vial		10	<ul> <li>Ibiamox</li> </ul>
Inj 1 g vial – Up to 5 inj available on a PSO	21.64	10	<ul> <li>Ibiamox</li> </ul>
AMOXICILLIN WITH CLAVULANIC ACID			
Tab 500 mg with clavulanic acid 125 mg – Up to 30 tab			
available on a PSO		10	<ul> <li>Curam Duo 500/125</li> </ul>
Grans for oral liq amoxicillin 25 mg with clavulanic acid 6.25 m			
per ml	8.50	100 ml	<ul> <li>Augmentin</li> </ul>
<ul> <li>a) Up to 200 ml available on a PSO</li> </ul>			
b) Wastage claimable			
c) Augmentin to be Principal Supply on 1 May 2025			
Grans for oral liq amoxicillin 50 mg with clavulanic acid 12.5 m	0	400 100	<i>(</i> <b>)</b>
per ml – Up to 200 ml available on a PSO		100 ml OP	<ul> <li>✓ Curam</li> <li>✓ Amoxiclay Devatis</li> </ul>
	5.61		Forte
Amoxiclav Devatis Forte to be Principal Supply on 1 June	2025		
Curam Grans for oral lig amoxicillin 50 mg with clavulanic acid 12		be delisted 1 J	lune 2025)
BENZATHINE BENZYLPENICILLIN	01-		-/
Inj 900 mg (1.2 million units) in 2.3 ml syringe – Up to 5 inj			
available on a PSO	432 37	10	<ul> <li>Bicillin LA</li> </ul>
		10	
BENZYLPENICILLIN SODIUM [PENICILLIN G]	0 16 50	10	- Condoz
Inj 600 mg (1 million units) vial – Up to 5 inj available on a PS	0	10	<ul> <li>Sandoz</li> </ul>

	Subsidy (Manufacturer's Price)		Fully Brand or Subsidised Generic
	\$	Per	<ul> <li>Manufacturer</li> </ul>
			<b>/</b>
Cap 250 mg – Up to 30 cap available on a PSO		250	<ul> <li>Flucloxacillin-AFT</li> </ul>
	22.58		✓ Staphlex
Cap 500 mg – Up to 30 cap available on a PSO		500	✓ Flucloxacillin-AFT
	72.71		✓ Staphlex
Grans for oral liq 25 mg per ml	4.89	100 ml	I ✓ <u>AFT</u>
a) Up to 200 ml available on a PSO			
b) Wastage claimable			
Grans for oral liq 50 mg per ml	5.89	100 ml	I ✓ <u>AFT</u>
<ul> <li>a) Up to 200 ml available on a PSO</li> </ul>			
b) Wastage claimable			
Inj 250 mg vial		10	<ul> <li>Flucloxin</li> </ul>
Inj 500 mg vial		10	✓ Flucloxin
Inj 1 g vial – Up to 5 inj available on a PSO	6.00	5	✓ Flucil
Flucloxacillin-AFT Cap 250 mg to be delisted 1 August 2025)			
Flucloxacillin-AFT Cap 500 mg to be delisted 1 August 2025)			
PHENOXYMETHYLPENICILLIN (PENICILLIN V)			
Cap 250 mg – Up to 30 cap available on a PSO	7.68	50	<ul> <li>Cilicaine VK</li> </ul>
Cap 500 mg		50	✓ Cilicaine VK
a) Up to 20 cap available on a PSO			<u></u>
b) Up to 2 x the maximum PSO guantity for RFPP			
Grans for oral liq 125 mg per 5 ml	3 40	100 ml	AFT
a) Up to 200 ml available on a PSO		100 111	<u> </u>
b) Wastage claimable			
Grans for oral liq 250 mg per 5 ml	4 24	100 ml	AFT
a) Up to 300 ml available on a PSO		100 111	
, ,			
<ul><li>b) Up to 2 x the maximum PSO quantity for RFPP</li><li>c) Wastage claimable</li></ul>			
c) wastage claimable			
Tetracyclines			
OXYCYCLINE			
Fab 100 mg – Up to 30 tab available on a PSO	64.43	500	<ul> <li>Doxine</li> </ul>
/INOCYCLINE HYDROCHLORIDE			
Tab 50 mg – Additional subsidy by Special Authority see			
SA1355 below – Retail pharmacy	5 79	60	
	(12.05)	00	Mino-tabs
₭ Cap 100 mg		100	
······································	(52.04)	100	Minomycin
	(02.01)		in in iterry on i

28

✓ Accord S29

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

2 For use only in combination with bismuth as part of a quadruple therapy regimen.

Initial application from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

1 For the eradication of helicobacter pylori following unsuccessful treatment with appropriate first-line therapy; and

■ SA1332 Special Authority for Subsidy

Both:

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
Other Antibiotics				
For topical antibiotics, refer to DERMATOLOGICALS, page 66 CIPROFLOXACIN Recommended for patients with any of the following: i) microbiologically confirmed and clinically significant ps ii) prostatitis; or iii) pyelonephritis; or iv) gonorrhoea.	eudomonas infection;	or		
Tab 250 mg – Up to 5 tab available on a PSO Tab 500 mg – Up to 5 tab available on a PSO Tab 750 mg CLINDAMYCIN	3.10	28 28 28	✓	Ipca-Ciprofloxacin Ipca-Ciprofloxacin Ipca-Ciprofloxacin
Cap hydrochloride 150 mg Inj 150 mg per ml, 4 ml ampoule		24 10		<u>Dalacin C</u> <u>Hameln</u>
COLISTIN SULPHOMETHATE – Retail pharmacy-Specialist – S Only if prescribed for dialysis or cystic fibrosis patient and th Inj 150 mg Inj 2 million iu, 10 ml vial	e prescription is endo		<b>v</b>	/. Colistin-Link Colomycin (529)
GENTAMICIN SULPHATE Inj 40 mg per ml, 2 ml vial – Subsidy by endorsement		5		Cidomycin P/Free \$29
Only if prescribed for a dialysis or cystic fibrosis patient endorsed accordingly.	or complicated urinary	/ trac	t infection a	
Inj 10 mg per ml, 1 ml ampoule – Subsidy by endorsement. Only if prescribed for a dialysis or cystic fibrosis patient endorsed accordingly.		5 / trac		<b>DBL Gentamicin</b> and the prescription is
Inj 10 mg per ml, 2 ml ampoule – Subsidy by endorsement. Only if prescribed for a dialysis or cystic fibrosis patient endorsed accordingly.		5 / trac		Wockhardt S29 and the prescription is
Inj 40 mg per ml, 2 ml ampoule – Subsidy by endorsement.	18.38	10		Gentamicin Amdipharm S29 Pfizer
	91.90	50	1	Gentamicin Noridem S29
Only if prescribed for a dialysis or cystic fibrosis patient endorsed accordingly.	or complicated urinary	/ trac	t infection a	and the prescription is
MOXIFLOXACIN – Special Authority see SA1740 below – Reta No patient co-payment payable Tab 400 mg		5	1	Avelox
■ SA1740 Special Authority for Subsidy Initial application — (Tuberculosis) only from a respiratory sp for applications meeting the following criteria: Any of the following:		iseas	e specialis	t. Approvals valid for 1 year
1 Both:				

continued...

Subsidy	Fully		Brand or	
(Manufacturer's Price)	Subsidised		Generic	
\$	Per	1	Manufacturer	

continued...

- 1.1 Active tuberculosis*; and
- 1.2 Any of the following:
  - 1.2.1 Documented resistance to one or more first-line medications; or
  - 1.2.2 Suspected resistance to one or more first-line medications (tuberculosis assumed to be contracted in an area with known resistance), as part of regimen containing other second-line agents; or
  - 1.2.3 Impaired visual acuity (considered to preclude ethambutol use); or
  - 1.2.4 Significant pre-existing liver disease or hepatotoxicity from tuberculosis medications; or
  - 1.2.5 Significant documented intolerance and/or side effects following a reasonable trial of first-line medications; or
- 2 Mycobacterium avium-intracellulare complex not responding to other therapy or where such therapy is contraindicated.*; or
- 3 Patient is under five years of age and has had close contact with a confirmed multi-drug resistant tuberculosis case.
- Note: Indications marked with * are unapproved indications.

**Renewal** only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Mycoplasma genitalium) only from a sexual health specialist or Practitioner on the recommendation of a sexual health specialist. Approvals valid for 1 month for applications meeting the following criteria: All of the following:

- 1 Has nucleic acid amplification test (NAAT) confirmed Mycoplasma genitalium* and is symptomatic; and
- 2 Either:
  - 2.1 Has tried and failed to clear infection using azithromycin; or
  - 2.2 Has laboratory confirmed azithromycin resistance; and
- 3 Treatment is only for 7 days.

**Initial application** — (Penetrating eye injury) only from an ophthalmologist. Approvals valid for 1 month where the patient requires prophylaxis following a penetrating eye injury and treatment is for 5 days only.

Note: Indications marked with * are unapproved indications.

PAROMOMYCIN – Special Authority see SA1689 below – Retail pharmacy

Cap 250 mg...... 126.00 16

#### ➡SA1689 Special Authority for Subsidy

Initial application only from an infectious disease specialist, clinical microbiologist or gastroenterologist. Approvals valid for 1 month for applications meeting the following criteria: Fither:

- 1 Patient has confirmed cryptosporidium infection; or
- 2 For the eradication of Entamoeba histolyica carriage.

Renewal only from an infectious disease specialist, clinical microbiologist or gastroenterologist. Approvals valid for 1 month for applications meeting the following criteria:

Either:

- 1 Patient has confirmed cryptosporidium infection; or
- 2 For the eradication of Entamoeba histolyica carriage.

PYRIMETHAMINE - Special Authority see SA1328 below - Retail pharmacy

30

✓ Daraprim S29

Humatin S29

#### ► SA1328 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 For the treatment of toxoplasmosis in patients with HIV for a period of 3 months; or
- 2 For pregnant patients for the term of the pregnancy; or
- 3 For infants with congenital toxoplasmosis until 12 months of age.

	Subsidy (Manufacturer's Price		Fully Subsidised	Generic
	\$	Per		Manufacturer
SODIUM FUSIDATE [FUSIDIC ACID]				
Tab 250 mg		36	~	Fucidin
SULFADIAZINE SODIUM – Special Authority see SA1331 below				
Tab 500 mg	150.70	100	<i>,</i>	Sulfadiazin-Heyl S29
	543.20	56	1	Wockhardt S29
<ul> <li>SA1331 Special Authority for Subsidy</li> <li>Initial application from any relevant practitioner. Approvals valid the following criteria:</li> <li>Any of the following:         <ol> <li>For the treatment of toxoplasmosis in patients with HIV for 2 For pregnant patients for the term of the pregnancy; or</li> </ol> </li> </ul>	without further rer	newal u		
3 For infants with congenital toxoplasmosis until 12 months of	of age.			
TOBRAMYCIN	0			
Inj 40 mg per ml, 2 ml vial – Subsidy by endorsement Only if prescribed for dialysis or cystic fibrosis patient and Solution for inhalation 60 mg per ml, 5 ml – Subsidy by		5 endor		Tobramycin (Viatris) dingly.
endorsement	395.00	56 dos	se 🗸	Tobramycin BNM
<ul><li>a) Wastage claimable</li><li>b) Only if prescribed for a cystic fibrosis patient and the patient and t</li></ul>				
TRIMETHOPRIM		1300 0	coordingry	
* Tab 300 mg – Up to 30 tab available on a PSO	27.83	50	1	ТМР
TRIMETHOPRIM WITH SULPHAMETHOXAZOLE [CO-TRIMOXA				
<ul> <li>* Tab trimethoprim 80 mg and sulphamethoxazole 400 mg – U to 30 tab available on a PSO</li> </ul>	p	500	~	Trisul
<ul> <li>Oral liq 8 mg sulphamethoxazole 40 mg per ml – Up to 200 n available on a PSO.</li> </ul>	nl	100 n		Deprim
VANCOMYCIN – Subsidy by endorsement				
Only if prescribed for a dialysis or cystic fibrosis patient or for difficile following metronidazole failure and the prescription is			is or for tre	eatment of Clostridium
Inj 500 mg vial		1	1	<u>Mylan</u>
Antifungals				
a) For topical antifungals refer to DERMATOLOGICALS, page 67	,			
b) For topical antifungals refer to GENITO URINARY, page 79				
FLUCONAZOLE	4.40	00		Madau
Cap 50 mg		28 1		<u>Mylan</u> Mylan
Cap 150 mg Cap 200 mg		1 28	-	<u>Mylan</u> Mylan
Powder for oral suspension 10 mg per ml – Special Authority		20		
see SA1359 below – Retail pharmacy		35 m	✓	Diflucan
■ SA1359 Special Authority for Subsidy Initial application — (Systemic candidiasis) from any relevant meeting the following criteria:	practitioner. Appre	ovals v	alid for 6 w	veeks for applications

continued...

trinued th: 1 Patient requires prophylaxis for, or treatment of systemic can 2 Patient is unable to swallow capsules. tial application — (Immunocompromised) from any relevant p eting the following criteria: of the following: 1 Patient is immunocompromised; and 2 Patient is at moderate to high risk of invasive fungal infection 3 Patient is unable to swallow capsules. newal — (Systemic candidiasis) from any relevant practitione	practitioner. App			
<ol> <li>Patient requires prophylaxis for, or treatment of systemic can 2 Patient is unable to swallow capsules.</li> <li>tial application — (Immunocompromised) from any relevant peting the following criteria: of the following:         <ol> <li>Patient is immunocompromised; and</li> <li>Patient is at moderate to high risk of invasive fungal infection</li> <li>Patient is unable to swallow capsules.</li> </ol> </li> <li>newal — (Systemic candidiasis) from any relevant practitione</li> </ol>	practitioner. App			
<ol> <li>Patient is unable to swallow capsules.</li> <li>tial application — (Immunocompromised) from any relevant peting the following criteria: of the following:         <ol> <li>Patient is immunocompromised; and</li> <li>Patient is at moderate to high risk of invasive fungal infection</li> <li>Patient is unable to swallow capsules.</li> </ol> </li> <li>newal — (Systemic candidiasis) from any relevant practitione</li> </ol>	practitioner. App			
eting the following criteria: of the following: 1 Patient is immunocompromised; and 2 Patient is at moderate to high risk of invasive fungal infection 3 Patient is unable to swallow capsules. newal — (Systemic candidiasis) from any relevant practitione	n; and			
<ol> <li>Patient is at moderate to high risk of invasive fungal infection</li> <li>Patient is unable to swallow capsules.</li> <li>newal — (Systemic candidiasis) from any relevant practitione</li> </ol>		d for 6 weeks	s for ap	oplications meeting the
	r. Approvals vali	d for 6 weeks	s for ap	oplications meeting the
owing criteria: th:				
<ol> <li>Patient requires prophylaxis for, or treatment of systemic can</li> <li>Patient is unable to swallow capsules.</li> </ol>	ndidiasis; and			
newal — (Immunocompromised) from any relevant practitione owing criteria: of the following:	er. Approvals va	lid for 6 mont	hs for a	applications meeting the
<ol> <li>Patient remains immunocompromised; and</li> <li>Patient remains at moderate to high risk of invasive fungal in</li> <li>Patient is unable to swallow capsules.</li> </ol>	nfection; and			
RACONAZOLE				
Cap 100 mg		15		trazole
	27.32	60	🗸 lt	racap S29
Oral liq 10 mg per ml – Special Authority see SA1322 below – Retail pharmacy		150 ml OP	🗸 it	traconazole Kent S29
			✓ s	Sporanox
poranox Oral liq 10 mg per ml to be delisted 1 April 2025)			Ū	F
SA1322 Special Authority for Subsidy				

practitioner on the recommendation of a infectious disease physician, clinical microbiologist or clinical immunologist. Approvals valid for 6 months where the patient has a congenital immune deficiency.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefitting from the treatment.

KETOCONAZOLE			
Tab 200 mg – PCT	CBS	30	<ul> <li>Burel S29</li> </ul>
		100	Strides Shasun S29
			<ul> <li>Taro S29</li> </ul>
			🗸 Teva-
			Ketoconazole S29
NYSTATIN			
Tab 500,000 u	14.16	50	
	(17.09)		Nilstat
Сар 500,000 и		50	
	(15.47)		Nilstat

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
POSACONAZOLE - Special Authority see SA2383 below - Retai	l pharmacy			
Tab modified-release 100 mg	206.00	24	✓ P	osaconazole Juno
Oral liq 40 mg per ml	342.51 10	5 ml C	)P 🖌 <u>D</u>	evatis

#### ► SA2383 Special Authority for Subsidy

**Initial application** only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

Either:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation chemotherapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppressive therapy*.

Renewal only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

Either:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation therapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppression* and requires on going posaconazole treatment.

Note: * Graft versus host disease (GVHD) on significant immunosuppression is defined as acute GVHD, grade II to IV, or extensive chronic GVHD, or if they were being treated with intensive immunosuppressive therapy consisting of either high-dose corticosteroids (1 mg or greater per kilogram of body weight per day for patients with acute GVHD or 0.8 mg or greater per kilogram every other day for patients with chronic GVHD), antithymocyte globulin, or a combination of two or more immunosuppressive agents or types of treatment.

**Initial application** — (Invasive fungal infection prophylaxis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient is at risk of invasive fungal infection; and
- 2 Either:
  - 2.1 Posaconazole is prescribed by, or recommended by a haematologist, transplant physician, infectious disease specialist, paediatric haematologist or paediatric oncologist; or
  - 2.2 Prescribing posaconazole is in accordance with a protocol or guideline that has been endorsed by the Health New Zealand Te Whatu Ora Hospital in the specific settings where there is a greater than 10% risk of invasive fungal infection (IFI).

Renewal — (Invasive fungal infection prophylaxis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient is at risk of invasive fungal infection; and
- 2 Either:
  - 2.1 Posaconazole is prescribed by, or recommended by a haematologist, transplant physician, infectious disease specialist, paediatric haematologist or paediatric oncologist; or
  - 2.2 Prescribing posaconazole is in accordance with a protocol or guideline that has been endorsed by the Health New Zealand Te Whatu Ora Hospital in the specific settings where there is a greater than 10% risk of invasive fungal infection (IFI).

TEF	RBI	NA	FI	NE	

* Tab 250 mg	8.97	84	✓ Deolate
VORICONAZOLE - Special Authority see SA2384 on the next page -	Retail pharma	су	
Tab 50 mg	71.00	56	<ul> <li>Vttack</li> </ul>
Tab 200 mg	263.00	56	<ul> <li>Vttack</li> </ul>
Powder for oral suspension 40 mg per ml – Wastage			
claimable1,	523.22	70 ml	<ul> <li>Vfend</li> </ul>

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

#### ⇒SA2384 Special Authority for Subsidy

Initial application - (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
  - 3.1 Patient has proven or probable invasive aspergillus infection; or
  - 3.2 Patient has possible invasive aspergillus infection: or
  - 3.3 Patient has fluconazole resistant candidiasis; or
  - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Renewal -- (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria: All of the followina:

- 1 Patient is immunocompromised: and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
  - 3.1 Patient continues to require treatment for proven or probable invasive aspergillus infection; or
  - 3.2 Patient continues to require treatment for possible invasive aspergillus infection; or
  - 3.3 Patient has fluconazole resistant candidiasis: or
  - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Initial application - (Invasive fungal infection prophylaxis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient is at risk of invasive fungal infection; and
- 2 Fither:
  - 2.1 Voriconazole is prescribed by, or recommended by a haematologist, transplant physician, infectious disease specialist, paediatric haematologist or paediatric oncologist; or
  - 2.2 Prescribing voriconazole is in accordance with a protocol or guideline that has been endorsed by the Health New Zealand - Te Whatu Ora Hospital in the specific settings where there is a greater than 10% risk of invasive fungal infection (IFI).

Renewal — (Invasive fungal infection prophylaxis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient is at risk of invasive fungal infection; and
- 2 Either:
  - 2.1 Voriconazole is prescribed by, or recommended by a haematologist, transplant physician, infectious disease specialist, paediatric haematologist or paediatric oncologist; or
  - 2.2 Prescribing voriconazole is in accordance with a protocol or guideline that has been endorsed by the Health New Zealand - Te Whatu Ora Hospital in the specific settings where there is a greater than 10% risk of invasive fungal infection (IFI).

## Antimalarials

PRIMAQUINE - Special Authority see SA1684 on the next page - Retail pharmacy 

100

✓ Sanofi Primaguine S29

	Subsidy	Fully	Brand or
(Man	ufacturer's Price)	Subsidised	Generic
	\$ F	Per 🗸	Manufacturer

#### ➡SA1684 Special Authority for Subsidy

Initial application only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 The patient has vivax or ovale malaria; and
- 2 Primaquine is to be given for a maximum of 21 days.

Renewal only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 The patient has relapsed vivax or ovale malaria; and

2 Primaquine is to be given for a maximum of 21 days.			
Antitrichomonal Agents			
METRONIDAZOLE Tab 200 mg – Up to 30 tab available on a PSO Tab 400 mg – Up to 15 tab available on a PSO Oral liq benzoate 200 mg per 5 ml Suppos 500 mg	4.29 25.00	250 21 100 ml 10	<ul> <li>✓ <u>Metronidamed</u></li> <li>✓ <u>Metronidamed</u></li> <li>✓ Flagyl-S</li> <li>✓ Flagyl</li> </ul>
ORNIDAZOLE Tab 500 mg		10	✓ Arrow-Ornidazole
Antituberculotics and Antileprotics			
Note: There is no co-payment charge for all pharmaceutical immigration status.	s listed in the Antituber	rculotics and	Antileprotics group regardless of
BEDAQUILINE – Special Authority see SA2244 below – Re No patient co-payment payable Tab 100mg		24 OP	✓ Sirturo
<ul> <li>SA2244 Special Authority for Subsidy</li> <li>Initial application — (multi-drug resistant tuberculosis) applications meeting the following criteria: Both:</li> </ul>			
<ol> <li>The person has multi-drug resistant tuberculosis (MD</li> <li>Ministry of Health's Tuberculosis Clinical Network has of the treatment regimen.</li> </ol>		al case and	recommends bedaquiline as part
<ul> <li>CLOFAZIMINE – Retail pharmacy-Specialist</li> <li>a) No patient co-payment payable</li> <li>b) Prescriptions must be written by, or on the recomme dermatologist.</li> <li>* Cap 50 mg</li> </ul>		s disease pł 100	nysician, clinical microbiologist or
CYCLOSERINE – Retail pharmacy-Specialist			
<ul> <li>a) No patient co-payment payable</li> <li>b) Prescriptions must be written by, or on the recomme respiratory physician.</li> </ul>			
Cap 250 mg	344.00	60	<ul> <li>Cyclorin S29</li> </ul>

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	
	\$	Per	1	Manufacturer
DAPSONE – Retail pharmacy-Specialist				
a) No patient co-payment payable				
<li>b) Prescriptions must be written by, or on the recommendation of the second state of the second state</li>	on of, an infectious d	isease	physiciar	n, clinical microbiologist or
dermatologist Tab 25 mg	268 50	100	1	Dapsone
Tab 100 mg		100		Dapsone
ETHAMBUTOL HYDROCHLORIDE – Retail pharmacy-Specialis				
a) No patient co-payment payable				
b) Prescriptions must be written by, or on the recommendati	on of, an infectious d	isease	physiciar	n, clinical microbiologist or
respiratory physician				, o
Tab 100 mg		100	✓	EMB Fatol S29
Tab 400 mg		56	1	Myambutol S29
ISONIAZID – Retail pharmacy-Specialist				
a) No patient co-payment payable				
<ul> <li>b) Prescriptions must be written by, or on the recommendation</li> </ul>	on of, an internal me	dicine	physician	, paediatrician, clinical
microbiologist, dermatologist or public health physician * Tab 100 mg	23.00	100	1	PSM
* Tab 100 mg	94.50	100		Isoniazid Teva S29
	327.41			Noumed Isoniazid
Noumed Isoniazid to be Principal Supply on 1 May 2025				
(PSM Tab 100 mg to be delisted 1 May 2025)				
ISONIAZID WITH RIFAMPICIN – Retail pharmacy-Specialist				
a) No patient co-payment payable				
b) Prescriptions must be written by, or on the recommendation	on of, an internal me	dicine	physician	, paediatrician, clinical
microbiologist, dermatologist or public health physician * Tab 100 mg with rifampicin 150 mg	00.00	100		Rifinah
<ul> <li>* Tab 100 mg with rifampicin 150 mg</li> <li>* Tab 150 mg with rifampicin 300 mg</li> </ul>		100		Rifinah
LINEZOLID – Special Authority see SA2234 below – Retail phar				
No patient co-payment payable	macy			
Tab 600 mg		10	1	<u>Zyvox</u>
Oral liq 20 mg per ml	1,879.00 1	150 ml		Zyvox
➡SA2234 Special Authority for Subsidy				
Initial application — (multi-drug resistant tuberculosis) from	any relevant practitio	ner. A	Approvals	valid for 18 months for
applications meeting the following criteria: Both:				
	R): and			
<ol> <li>The person has multi-drug resistant tuberculosis (MDR-TE</li> <li>Ministry of Health's Tuberculosis Clinical Network has rev</li> </ol>	<i>/</i> ··	ase ar	nd recomm	mends linezolid as part of
the treatment regimen.		uoc ui		
PARA-AMINO SALICYLIC ACID – Retail pharmacy-Specialist				
a) No patient co-payment payable				
b) Prescriptions must be written by, or on the recommendati	on of, an infectious d	isease	specialis	t, clinical microbiologist or
respiratory physician				-
Grans for oral liq 4 g sachet		30	1	Paser S29
PROTIONAMIDE – Retail pharmacy-Specialist				
a) No patient co-payment payable				
<li>b) Prescriptions must be written by, or on the recommendation reconstruction by a second second</li>	on of, an infectious d	isease	specialis	t, clinical microbiologist or
respiratory physician Tab 250 mg	305.00	100	1	Peteha S29
1 ab 200 mg		100	•	

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

		Subsidy		Fully	Brand or
		(Manufacturer's Pric		Subsidised	Generic
		\$	Per		Manufacturer
PYRAZINAMI	DE – Retail pharmacy-Specialist				
a) No pa	tient co-payment payable				
	riptions must be written by, or on the recommendati	on of, an infectious	s disease	physician	, clinical microbiologist o
	ng	64.95	100	✓ /	AFT-Pyrazinamide
RIFABUTIN -	- Retail pharmacy-Specialist				•
	tient co-payment payable				
b) Prescr	riptions must be written by, or on the recommendation interesting to the recommendation interesting to the recommendation of the recommend	on of, an infectious	s disease	physician	, respiratory physician or
	mg		30	✓ 1	Aycobutin
	<ul> <li>Subsidy by endorsement</li> </ul>				
	tient co-payment payable				
b) For co antimi Retail	nfirmed recurrent Staphylococcus aureus infection crobial based on susceptibilities and the prescription pharmacy - Specialist. Specialist must be an interr atrician, or public health physician.	n is endorsed acco	ordingly;	can be wai	ved by endorsement -
* Cap 150 r	mg		100	✓ ]	Rifadin
·	mg		100	-	<u>Rifadin</u> Rifadin Sanofi
* Oral liq 10	00 mg per 5 ml	12.60	60 ml	✓Į	Rifadin
Antivirals					
For eye prepa	rations refer to Eye Preparations, Anti-Infective Pre	parations, page 26	6		
Hepatitis	B Treatment				
ENTECAVIR * Tab 0.5 m	ıg		30	<b>√</b> [	Entecavir (Rex)
LAMIVUDINE	- Special Authority see SA1685 below - Retail pha	armacy			
Tab 100 r	ng		28	<ul> <li>Image: A second s</li></ul>	Zetlam
Oral liq 5	mg per ml		240 ml C	)P 🖌	Zeffix
⇒SA1685 S	pecial Authority for Subsidy				
nitial applica	tion only from a relevant specialist or medical practice of the treatment or prevention of the treatment o		mmenda	ation of a re	elevant specialist.
Renewal from	any relevant practitioner. Approvals valid for 2 yea		the treat	ment or pr	evention of hepatitis B.
Renewal from TENOFOVIR Tenofovir	any relevant practitioner. Approvals valid for 2 yea DISOPROXIL disoproxil prescribed under endorsement for the tre	ars where used for eatment of HIV is ir			
Renewal from TENOFOVIR Tenofovir antiretrovi	n any relevant practitioner. Approvals valid for 2 yea DISOPROXIL disoproxil prescribed under endorsement for the tre irals for the purposes of Special Authority SA2139.,	ars where used for eatment of HIV is ir page 109	ncluded i	n the coun	t of up to 4 subsidised
Renewal from TENOFOVIR Tenofovir antiretrovi	any relevant practitioner. Approvals valid for 2 yea DISOPROXIL disoproxil prescribed under endorsement for the tre	ars where used for eatment of HIV is ir page 109		n the coun	
Renewal from TENOFOVIR Tenofovir antiretrovi * Tab 245 r	n any relevant practitioner. Approvals valid for 2 yea DISOPROXIL disoproxil prescribed under endorsement for the tre irals for the purposes of Special Authority SA2139.,	ars where used for eatment of HIV is ir page 109	ncluded i	n the coun	t of up to 4 subsidised Tenofovir Disoproxil
Renewal from TENOFOVIR Tenofovir antiretrovi * Tab 245 r	n any relevant practitioner. Approvals valid for 2 yea DISOPROXIL disoproxil prescribed under endorsement for the tre irals for the purposes of Special Authority SA2139., ng (300 mg as a maleate)	ars where used for eatment of HIV is ir page 109	ncluded i	n the coun	t of up to 4 subsidised Tenofovir Disoproxil
Renewal from TENOFOVIR Tenofovir antiretrovi * Tab 245 r Herpesvir ACICLOVIR	n any relevant practitioner. Approvals valid for 2 yea DISOPROXIL disoproxil prescribed under endorsement for the tre irals for the purposes of Special Authority SA2139., ng (300 mg as a maleate)	ars where used for eatment of HIV is ir page 109 	ncluded i	n the coun	t of up to 4 subsidised Tenofovir Disoproxil
Renewal from TENOFOVIR Tenofovir antiretrovi * Tab 245 r Herpesvir ACICLOVIR * Tab dispe	n any relevant practitioner. Approvals valid for 2 yea DISOPROXIL disoproxil prescribed under endorsement for the tre irals for the purposes of Special Authority SA2139., ng (300 mg as a maleate)	ars where used for eatment of HIV is ir page 109 	ncluded i 30	n the coun	t of up to 4 subsidised Tenofovir Disoproxil <u>Viatris</u>
Renewal from TENOFOVIR Tenofovir antiretrovi * Tab 245 r Herpesvir ACICLOVIR * Tab dispe * Tab dispe	n any relevant practitioner. Approvals valid for 2 yea DISOPROXIL disoproxil prescribed under endorsement for the tre irals for the purposes of Special Authority SA2139., ng (300 mg as a maleate)	ars where used for eatment of HIV is ir page 109 	ncluded i 30 25	n the coun	t of up to 4 subsidised <u>Tenofovir Disoproxil</u> <u>Viatris</u> <u>Lovir</u>
Renewal from TENOFOVIR Tenofovir antiretrovi * Tab 245 r Herpesvir ACICLOVIR * Tab dispe * Tab dispe * Tab dispe	n any relevant practitioner. Approvals valid for 2 yea DISOPROXIL disoproxil prescribed under endorsement for the tre irals for the purposes of Special Authority SA2139., ng (300 mg as a maleate) <b>rus Treatments</b> ersible 200 mg ersible 400 mg	ars where used for eatment of HIV is ir page 109 	25 56	n the coun	t of up to 4 subsidised Tenofovir Disoproxil Viatris Lovir Lovir
Renewal from TENOFOVIR Tenofovir antiretrovi * Tab 245 r Herpesvir ACICLOVIR * Tab dispe * Tab dispe * Tab dispe VALACICLOV	n any relevant practitioner. Approvals valid for 2 yea DISOPROXIL disoproxil prescribed under endorsement for the tre irals for the purposes of Special Authority SA2139., ng (300 mg as a maleate) <b>rus Treatments</b> ersible 200 mg ersible 400 mg	ars where used for eatment of HIV is ir page 109 	25 56	n the coun	t of up to 4 subsidised Tenofovir Disoproxil Viatris Lovir Lovir

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
VALGANCICLOVIR – Special Authority see SA1993 below – Ret Tab 450 mg		60	✓ <u>V</u>	<u>alganciclovir</u> Viatris

#### ⇒SA1993 Special Authority for Subsidy

**Initial application** — (transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 3 months where the patient has undergone a solid organ transplant and requires valganciclovir for CMV prophylaxis.

Renewal — (transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

1 Both:

- 1.1 Patient has undergone a solid organ transplant and received anti-thymocyte globulin and requires valganciclovir therapy for CMV prophylaxis; and
- 1.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following anti-thymocyte globulin; or 2 Both:
  - 2.1 Patient has received pulse methylprednisolone for acute rejection and requires further valganciclovir therapy for CMV prophylaxis; and
  - 2.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following pulse methylprednisolone.

Initial application — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

- Both:
  - 1 Patient has undergone a solid organ transplant and received valganciclovir under Special Authority more than 2 years ago (27 months); and
  - 2 Patient has received anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Renewal — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months where the patient has received a further course of anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

**Initial application — (Lung transplant cytomegalovirus prophylaxis)** only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone a lung transplant; and
- 2 Either:
  - 2.1 The donor was cytomegalovirus positive and the patient is cytomegalovirus negative; or
  - 2.2 The recipient is cytomegalovirus positive; and
- 3 Patient has a high risk of CMV disease.

Initial application — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
  - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
  - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
  - 2.3 Patient has cytomegalovirus retinitis.

**Renewal — (Cytomegalovirus in immunocompromised patients)** only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
  - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or

continued...

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

Subsidy	Fully Subsidised		Brand or
(Manufacturer's Price)			Generic
\$	Per	✓	Manufacturer

continued...

- 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
- 2.3 Patient has cytomegalovirus retinitis.

Note: for the purpose of this Special Authority "immunocompromised" includes transplant recipients, patients with immunosuppressive diseases (e.g. HIV) or those receiving immunosuppressive treatment for other conditions.

## Hepatitis C Treatment

GLECAPREVIR WITH PIBRENTASVIR – [Xpharm] Note the supply of treatment is via Pharmac's approved of	liroot distribution sum	oly Eurthoric	lataile can be found on Dharmac's			
website https://pharmac.govt.nz/maviret	inect distribution sup	Jiy. Fuithert	ietalis can de lounu on Fhannac's			
Tab 100 mg with pibrentasvir 40 mg	24,750.00	84 OP	<ul> <li>Maviret</li> </ul>			
LEDIPASVIR WITH SOFOSBUVIR - [Xpharm] - Special Aut No patient co-payment payable	thority see SA1605 be	elow				
Tab 90 mg with sofosbuvir 400 mg	24,363.46	28	<ul> <li>Harvoni</li> </ul>			
■ SA1605 Special Authority for Subsidy						
Special Authority approved by the Hepatitis C Treatment Pan	el (HepCTP)					
Notes: By application to the Hepatitis C Treatment Panel (HepCTP).						
Applications will be considered by HepCTP and approved subject to confirmation of eligibility.						
Application details may be obtained from Pharmac's website http://www.pharmac.govt.nz/maviret or:						
The Coordinator, Hepatitis C Treatment Panel						
Pharmac, PO Box 10-254, WELLINGTON Tel: (04) 460 4990	),					
Email: <u>hepcpanel@pharmac.govt.nz</u>						

#### HIV Prophylaxis and Treatment

EMTRICITABINE WITH TENOFOVIR DISOPROXIL - Subsidy by endorsement; can be waived by Special Authority see SA2138 below

- a) Funding for emtricitabine with tenofovir disoproxil for use as PrEP, should be applied using Special Authority SA2138.
- b) Endorsement for treatment of conditions approved via Special Authority SA2139 (antiretrovirals for confirmed HIV. prevention of maternal transmission, post-exposure prophylaxis following exposure to HIV and percutaneous exposure): Prescription is deemed to be endorsed if emtricitabine with tenofovir disoproxil is co-prescribed with another antiretroviral subsidised under Special Authority SA2139 and the prescription is annotated accordingly by the Pharmacist or endorsed by the prescriber.

Note: Emtricitabine with tenofovir disoproxil prescribed under endorsement, for treatment of conditions approved via Special Authority SA2139 (antiretrovirals for confirmed HIV, prevention of maternal transmission, post-exposure prophylaxis following exposure to HIV and percutaneous exposure), is included in the count of up to 4 subsidised antiretrovirals, and counts as two antiretroviral medications, for the purposes of Special Authority SA2139, page 109 There is an approval process to become a named specialist to prescribe antiretroviral therapy in New Zealand. Further information is available on the Pharmac website.

*	Tab 200 mg with tenofovir disoproxil 245 mg (300 mg as a maleate)	15.45	30	✓ <u>Tenofovir Disoproxil</u> Emtricitabine Viatr			
*	Tab 200 mg with tenofovir disoproxil 245 mg (300.6 mg as a succinate)	15 45	30				
(Teva Tab 200 mg with tenofovir disoproxil 245 mg (300.6 mg as a succinate) to be delisted 1 August 2025)							
<u>`</u>	A2138 Special Authority for Subsidy	,		<b>3</b>			

Initial application from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria: Both:

continued...

Subsidy	Fu	lly Brand or	
(Manufacturer's Price)	Subsidis	ed Generic	
\$	Per	<ul> <li>Manufacturer</li> </ul>	

continued...

- 1 Patient has tested HIV negative, does not have signs or symptoms of acute HIV infection and has been assessed for HIV seroconversion; and
- 2 The Practitioner considers the patient is at elevated risk of HIV exposure and use of PrEP is clinically appropriate.

Notes: Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines:

https://ashm.org.au/HIV/PrEP/

Renewal from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 Patient has tested HIV negative, does not have signs or symptoms of acute HIV infection and has been assessed for HIV seroconversion; and
- 2 The Practitioner considers the patient is at elevated risk of HIV exposure and use of PrEP is clinically appropriate.

Notes: Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines:

https://ashm.org.au/HIV/PrEP/

### **COVID-19 Treatments**

NIRMATRELVIR WITH RITONAVIR - [Xpharm] - Subsidy by endorsement

- a) No patient co-payment payable
- b) Treatment is funded only if patient meets access criteria for oral antiviral COVID-19 treatments (as on <u>Pharmac's website</u>) and has been endorsed accordingly by the prescriber. The supply of treatment is via Pharmac's approved distribution process. Refer to the Pharmac website for more information about this and stock availability.
   Tab 150 mg with ritonavir 100 mg ......0.0 30 ✓ Paxlovid

### Antiretrovirals

### ⇒SA2139 Special Authority for Subsidy

**Initial application** — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the patient has confirmed HIV infection.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient is benefiting from treatment.

**Initial application** — (Prevention of maternal transmission) only from a named specialist. Approvals valid for 1 year for applications meeting the following criteria:

Either:

- 1 Prevention of maternal foetal transmission; or
- 2 Treatment of the newborn for up to eight weeks.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Some antiretrovirals are unapproved or contraindicated for this indication. Practitioners prescribing these medications should exercise their own skill, judgement, expertise and discretion, and make their own prescribing decisions with respect to the use of a Pharmaceutical for an indication for which it is not approved or contraindicated.

Subsidy		Fully	Brand or
(Manufacturer's Price)	Su	ubsidised	Generic
\$	Per	✓	Manufacturer

**Initial application** — (post-exposure prophylaxis following exposure to HIV) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
  - 2.1 Patient has had condomless anal intercourse or receptive vaginal intercourse with a known HIV positive person with an unknown or detectable viral load greater than 200 copies per ml; or
  - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
  - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required; or
  - 2.4 Patient has had condomless anal intercourse with a person from a high HIV prevalence country or risk group whose HIV status is unknown.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines for PEP (https://www.ashm.org.au/hiv/hiv-management/pep/).

Renewal — (second or subsequent post-exposure prophylaxis) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
  - 2.1 Patient has had condomless anal intercourse or receptive vaginal intercourse with a known HIV positive person with an unknown or detectable viral load greater than 200 copies per ml; or
  - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
  - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required; or
  - 2.4 Patient has had condomless anal intercourse with a person from a high HIV prevalence country or risk group whose HIV status is unknown.

**Initial application — (Percutaneous exposure)** only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Second or subsequent percutaneous exposure) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

### Non-nucleosides Reverse Transcriptase Inhibitors

EFAVIRENZ - Special Authority see SA2139 on the previous pag	e – Retail pha	rmacy	
Tab 600 mg	65.38	30	<ul> <li>Efavirenz</li> </ul>
			Milpharm S29
ETRAVIRINE - Special Authority see SA2139 on the previous page	ge – Retail ph	armacy	
Tab 200 mg	770.00	60	<ul> <li>Intelence</li> </ul>
NEVIRAPINE - Special Authority see SA2139 on the previous pa	<mark>ge</mark> – Retail ph	armacy	
Tab 200 mg	198.25	60	<ul> <li>Nevirapine Viatris</li> </ul>
Oral suspension 10 mg per ml	203.55	240 ml OP	✓ Viramune
			Suspension

	Subsidy (Manufacturer's Price) \$		Fully Brand or idised Generic ✓ Manufacturer
Nucleosides Reverse Transcriptase Inhibitors			
ABACAVIR SULPHATE - Special Authority see SA2139 on page	e 109 – Retail pharm	acy	
Tab 300 mg		60	<ul> <li>Ziagen</li> </ul>
ABACAVIR SULPHATE WITH LAMIVUDINE – Special Authority Note: abacavir with lamivudine (combination tablets) counts anti-retroviral Special Authority.			
Tab 600 mg with lamivudine 300 mg	29.50	30	✓ <u>Abacavir/</u> Lamivudine Viatris
EFAVIRENZ WITH EMTRICITABINE AND TENOFOVIR DISOPP	ROXIL - Special Aut	thority see S	
pharmacy Note: Efavirenz with emtricitabine and tenofovir disoproxil co anti-retroviral Special Authority	ounts as three anti-re	etroviral med	dications for the purposes of the
Tab 600 mg with emtricitabine 200 mg and tenofovir disoprox	cil		
245 mg (300 mg as a fumarate)		30	<ul> <li>Triovir S29</li> </ul>
Tab 600 mg with emtricitabine 200 mg and tenofovir disoproz 245 mg (300 mg as a maleate)		30	✓ Viatris
EMTRICITABINE – Special Authority see SA2139 on page 109 -			
Cap 200 mg		30	<ul> <li>Emtriva</li> </ul>
LAMIVUDINE - Special Authority see SA2139 on page 109 - Re	etail pharmacy		
Tab 150 mg		60	Lamivudine Viatris
Oral liq 10 mg per ml		10 ml OP	✓ 3TC
ZIDOVUDINE [AZT] – Special Authority see SA2139 on page 10 Cap 100 mg		100	Retrovir
Oral liq 10 mg per ml		00 ml OP	✓ Retrovir
ZIDOVUDINE [AZT] WITH LAMIVUDINE - Special Authority see Note: zidovudine [AZT] with lamivudine (combination tablets the anti-retroviral Special Authority.			
Tab 300 mg with lamivudine 150 mg	92.40	60	<ul> <li>Lamivudine/ Zidovudine Viatris</li> </ul>
Protease Inhibitors			
ATAZANAVIR SULPHATE - Special Authority see SA2139 on p	age 109 – Retail pha	armacy	
Cap 150 mg		60	<ul> <li>✓ <u>Atazanavir Mylan</u></li> <li>✓ Atazanavir Viatris</li> </ul>
Cap 200 mg		60	<ul> <li>Atazanavir Viatris</li> <li>Atazanavir Viatris</li> </ul>
DARUNAVIR – Special Authority see SA2139 on page 109 – Re			
Tab 400 mg		60	✓ Darunavir Viatris
Tab 600 mg		60	<ul> <li>Darunavir Viatris</li> </ul>
LOPINAVIR WITH RITONAVIR – Special Authority see SA2139 Tab 100 mg with ritonavir 25 mg		il pharmacy 60	<ul> <li>Lopinavir/Ritonavir</li> <li>Mylan</li> </ul>
Tab 200 mg with ritonavir 50 mg		120	<ul> <li><u>Lopinavir/Ritonavir</u></li> <li><u>Mylan</u></li> </ul>
RITONAVIR – Special Authority see SA2139 on page 109 – Ret Tab 100 mg		30	✓ Norvir

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Subsid Per	Fully Brand or lised Generic ✓ Manufacturer
Strand Transfer Inhibitors			
DOLUTEGRAVIR – Special Authority see SA2139 on page 105 Tab 50 mg		30	✓ Tivicay
DOLUTEGRAVIR WITH LAMIVUDINE – Special Authority see Tab 50 mg with lamivudine 300 mg		- Retail pha 30	vrmacy ✓ Dovato
RALTEGRAVIR POTASSIUM – Special Authority see SA2139 Tab 400 mg Tab 600 mg	1,090.00	harmacy 60 60	<ul><li>✓ Isentress</li><li>✓ Isentress HD</li></ul>
Immune Modulators			
<ul> <li>PEGYLATED INTERFERON ALFA-2A – Special Authority see Note: Pharmac will consider funding ribavirin for the small Special Authority criteria. Please contact the Hepatitis C Co Inj 180 mcg prefilled syringe</li></ul>	group of patients who l pordinator at Pharmac	nave a clinic	
Initial application — (chronic hepatitis C - genotype 1, 4, 5 c liver transplant) from any specialist. Approvals valid for 18 m			
Both: 1 Any of the following: 1.1 Patient has chronic hepatitis C, genotype 1, 4, 5 of			
<ul><li>1.2 Patient has chronic hepatitis C and is co-infected</li><li>1.3 Patient has chronic hepatitis C genotype 2 or 3 ar</li></ul>		transplant;	and
2 Maximum of 48 weeks therapy.	from a gootrooptorok	aiot infacti	aua diagona anggialist ar ganaral
Renewal — (Chronic hepatitis C - genotype 1 infection) only physician. Approvals valid for 18 months for applications meetin All of the following:			ous disease specialist of general
<ol> <li>Patient has chronic hepatitis C, genotype 1; and</li> <li>Patient has had previous treatment with pegylated interfer</li> <li>Either:</li> </ol>	eron and ribavirin; and		
<ul><li>3.1 Patient has responder relapsed; or</li><li>3.2 Patient was a partial responder; and</li></ul>			
<ol> <li>Patient is to be treated in combination with boceprevir; a</li> <li>Maximum of 48 weeks therapy.</li> </ol>	nd		
Initial application — (Chronic Hepatitis C - genotype 1 infec gastroenterologist, infectious disease specialist or general physi following criteria: All of the following:			
<ol> <li>Patient has chronic hepatitis C, genotype 1; and</li> <li>Patient has had previous treatment with pegylated interfer</li> <li>Any of the following:</li> </ol>	eron and ribavirin; and		
<ul> <li>3.1 Patient has responder relapsed; or</li> <li>3.2 Patient was a partial responder; or</li> <li>3.3 Patient received interferon treatment prior to 2004</li> </ul>	4: and		
<ul><li>4 Patient is to be treated in combination with boceprevir; at</li><li>5 Maximum of 48 weeks therapy.</li></ul>			

Subsidy	Full	/ Brand or	
(Manufacturer's Price	e) Subsidise	d Generic	
\$	Per 🖌	Manufacturer	

continued...

Initial application — (chronic hepatitis C - genotype 2 or 3 infection without co-infection with HIV) from any specialist. Approvals valid for 12 months for applications meeting the following criteria: Both:

- Patient has chronic hepatitis C, genotype 2 or 3 infection; and
  - 2 Maximum of 6 months therapy.

Initial application — (Hepatitis B) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

### All of the following:

- 1 Patient has confirmed Hepatitis B infection (HBsAg positive for more than 6 months); and
- 2 Patient is Hepatitis B treatment-naive; and
- 3 ALT > 2 times Upper Limit of Normal; and
- 4 HBV DNA < 10 log10 IU/ml; and
- 5 Either:
  - 5.1 HBeAg positive; or
  - 5.2 serum HBV DNA greater than or equal to 2,000 units/ml and significant fibrosis (Metavir Stage F2 or greater or moderate fibrosis); and
- 6 Compensated liver disease; and
- 7 No continuing alcohol abuse or intravenous drug use; and
- 8 Not co-infected with HCV, HIV or HDV; and
- 9 Neither ALT nor AST > 10 times upper limit of normal; and
- 10 No history of hypersensitivity or contraindications to pegylated interferon; and
- 11 Maximum of 48 weeks therapy.

**Initial application** — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 Patient has a cutaneous T cell lymphoma*; or
- 2 All of the following:
  - 2.1 Patient has a myeloproliferative disorder*; and
  - 2.2 Patient is intolerant of hydroxyurea; and
  - 2.3 Treatment with anagrelide and busulfan is not clinically appropriate; or
- 3 Both:
  - 3.1 Patient has a myeloproliferative disorder; and
  - 3.2 Patient is pregnant, planning pregnancy or lactating.

Renewal — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment; and
- 3 Either:
  - 3.1 Patient has a cutaneous T cell lymphoma*; or
  - 3.2 Both:
    - 3.2.1 Patient has a myeloproliferative disorder*; and
    - 3.2.2 Either:
      - 3.2.2.1 Remains intolerant of hydroxyurea and treatment with anagrelide and busulfan remains clinically inappropriate; or
      - 3.2.2.2 Patient is pregnant, planning pregnancy or lactating.

Note: Indications marked with * are unapproved indications.

continued...

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Su	lbsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Initial application — (post-allogenic bone marrow transplant) from any relevant practitioner. Approvals valid for 3 months where patient has received an allogeneic bone marrow transplant* and has evidence of disease relapse.

Renewal — (post-allogenic bone marrow transplant) from any relevant practitioner. Approvals valid for 3 months where patient is responding and ongoing treatment remains appropriate.

Note: Indications marked with * are unapproved indications.

#### 

#### ■ SA2406 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 months for applications meeting the following criteria: Both:

- 1 Patient has an acute, symptomatic, bacteriologically-proven uncomplicated urinary tract infection (UTI)/cystitis with Escherichia Coli; and
- 2 Either:
  - 2.1 Microbiological testing confirms the pathogen is resistant to all of: trimethoprim, nitrofurantoin, amoxicillin, cefaclor, cefalexin, amoxicillin with clavulanic acid, and norfloxacin; or
  - 2.2 The patient has a contraindication or intolerance to all of: trimethoprim, nitrofurantoin, amoxicillin, cefaclor, cefalexin, amoxicillin with clavulanic acid, and norfloxacin that the pathogen is susceptible to.

Renewal from any relevant practitioner. Approvals valid for 2 months for applications meeting the following criteria: Both:

- 1 Patient has an acute, symptomatic, bacteriologically-proven uncomplicated urinary tract infection (UTI)/cystitis with Escherichia Coli; and
- 2 Either:
  - 2.1 Microbiological testing confirms the pathogen is resistant to all of: trimethoprim, nitrofurantoin, amoxicillin, cefaclor, cefalexin, amoxicillin with clavulanic acid, and norfloxacin; or
  - 2.2 The patient has a contraindication or intolerance to all of: trimethoprim, nitrofurantoin, amoxicillin, cefaclor, cefalexin, amoxicillin with clavulanic acid, and norfloxacin that the pathogen is susceptible to.

## METHENAMINE (HEXAMINE) HIPPURATE

* Tab 1 g		100	<ul> <li>Hiprex</li> </ul>
NITROFURANTOIN			
* Tab 50 mg – Up to 30 tab available on a PSO		100	Nifuran
* Tab 100 mg		100	<ul> <li>Nifuran</li> </ul>
* Cap modified-release 100 mg - Up to 15 cap available	e on a		
PSO		100	<ul> <li>Macrobid</li> </ul>
NORFLOXACIN			
Tab 400 mg – Subsidy by endorsement		100	<ul> <li>Arrow-Norfloxacin</li> </ul>

Only if prescribed for a patient with an uncomplicated urinary tract infection that is unresponsive to a first line agent or with proven resistance to first line agents and the prescription is endorsed accordingly.

## MUSCULOSKELETAL SYSTEM

	Subsidy		Fully Brand or
	(Manufacturer's Price		idised Generic
	\$	Per	<ul> <li>Manufacturer</li> </ul>
Anticholinesterases			
Antichonnesterases			
NEOSTIGMINE METILSULFATE			
Inj 2.5 mg per ml, 1 ml ampoule		10	Max Health
PYRIDOSTIGMINE BROMIDE			
▲ Tab 60 mg		100	<ul> <li>Mestinon</li> </ul>
Non Storoidal Anti Inflommatory Druga			
Non-Steroidal Anti-Inflammatory Drugs			
DICLOFENAC SODIUM			
* Tab EC 25 mg		50	<ul> <li>Diclofenac Sandoz</li> </ul>
* Tab 50 mg dispersible		20	<ul> <li>Voltaren D</li> </ul>
* Tab EC 50 mg		50	<ul> <li>Diclofenac Sandoz</li> </ul>
* Tab long-acting 75 mg		100	Voltaren SR
Inj 25 mg per ml, 3 ml ampoule – Up to 5 inj available on a		5	✓ Voltaren
* Suppos 12.5 mg		10 10	<ul> <li>✓ Voltaren</li> <li>✓ Voltaren</li> </ul>
<ul> <li>Suppos 25 mg</li> <li>Suppos 50 mg – Up to 10 supp available on a PSO</li> </ul>		10	✓ Voltaren
* Suppos 50 mg - Op to 10 supp available of a 1 SO		10	✓ Voltaren
IBUPROFEN		10	Voluien
BOPROFEN * Tab 200 mg	21.40	1.000	✓ Relieve
★ Tab long-acting 800 mg		30	✓ <u>Inelieve</u> ✓ Brufen SR
	3.65	00	✓ Ibuprofen SR BNM
Ibuprofen SR BNM to be Principal Supply on 1 April 20			
* Oral lig 20 mg per ml		200 ml	<ul> <li>Ethics</li> </ul>
Ethics to be Principal Supply on 1 April 2025			
(Brufen SR Tab long-acting 800 mg to be delisted 1 April 2025)			
KETOPROFEN			
* Cap long-acting 200 mg	12.07	28	<ul> <li>Oruvail SR</li> </ul>
MEFENAMIC ACID			
* Cap 250 mg	1.25	50	
	(10.82)		Ponstan
	0.50	20	
	(7.50)		Ponstan
NAPROXEN			
* Tab 250 mg		500	Noflam 250
* Tab 500 mg		250	<ul> <li>Noflam 500</li> </ul>
* Tab long-acting 750 mg		28	Naprosyn SR 750
* Tab long-acting 1 g	11.50	28	Naprosyn SR 1000
TENOXICAM			
* Tab 20 mg		100	✓ <u>Tilcotil</u>
* Inj 20 mg vial	9.95	1	✓ AFT
NSAIDs Other			
CELECOXIB	o :-		
Cap 100 mg	3.45	60	✓ Celebrex
Cap 200 mg	2 00	20	<ul> <li>✓ <u>Celecoxib Pfizer</u></li> <li>✓ Celebrex</li> </ul>
Cap 200 mg		30	<ul> <li>Celebrex</li> <li>Celecoxib Pfizer</li> </ul>

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

	Subsidy		Fully Brand or
	(Manufacturer's Pric		sidised Generic
	\$	Per	<ul> <li>Manufacturer</li> </ul>
Topical Products for Joint and Muscular Pain			
APSAICIN			
Crm 0.025% – Special Authority see SA1289 below – Retail			
pharmacy		45 g OP	Zo-Rub Osteo S29
			✓ Zostrix
	13.00	60 g OP	<ul> <li>Rugby Capsaicin Topical</li> </ul>
			Cream S29
Rugby Capsaicin Topical Cream 👓 Crm 0.025% to be deliste	d 1 July 2025)		
SA1289 Special Authority for Subsidy			
nitial application from any relevant practitioner. Approvals valid			
osteoarthritis that is not responsive to paracetamol and oral non-	steroidal anti-inflarr	nmatories are	e contraindicated.
Antirheumatoid Agents			
HYDROXYCHLOROQUINE SULPHATE ★ Tab 200 mg	7 80	100	✓ Ipca-
* Tab 200 mg		100	Hydroxychloroquine
Ipca-Hydroxychloroquine to be Principal Supply on 1 Ma	8.78 v 2025		<ul> <li>Plaquenil</li> </ul>
(Plaquenil Tab 200 mg to be delisted 1 May 2025)	y 2023		
* Tab 10 mg	6.00	30	✓ <u>Arava</u>
* Tab 20 mg	6.00	30	✓ Arava
PENICILLAMINE			
Tab 125 mg		100	✓ D-Penamine
Tab 250 mg	110.12	100	<ul> <li>D-Penamine</li> </ul>
Drugs Affecting Bone Metabolism			
Alendronate for Osteoporosis			
ALENDRONATE SODIUM			
* Tab 70 mg	3.10	4	✓ Fosamax
ALENDRONATE SODIUM WITH COLECALCIFEROL			
* Tab 70 mg with colecalciferol 5,600 iu	1.99	4	✓ Fosamax Plus
Other Treatments			
DENOSUMAB - Special Authority see SA2441 on the next page			
Note: Denosumab inj 60 mg per 1 ml pre-filled syringe is Me		r use in oste	oporosis. Denosumab inj 120 m
per 1.7 ml vial is Medsafe approved for use in hypercalcaemi Inj 120 mg per 1.7 ml vial	• •	1	✓ Xgeva
Inj 60 mg per 1 ml prefilled syringe		1	✓ Ageva ✓ Prolia
		I	- 110114

## MUSCULOSKELETAL SYSTEM

(Manu	Subsidy	Fully	Brand or
	facturer's Price)	Subsidised	Generic
	\$ Pe	er 🖌	Manufacturer

### ➡SA2441 Special Authority for Subsidy

**Initial application** — (Osteoporosis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 The patient has established osteoporosis; and
- 2 Any of the following:
  - 2.1 History of one significant osteoporotic fracture demonstrated radiologically, with a documented T-Score less than or equal to -2.5, that incorporates BMD measured using dual-energy x-ray absorptiometry (DEXA); or
  - 2.2 History of one significant osteoporotic fracture, demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of logistical, technical or pathophysiological reasons; or
  - 2.3 History of two significant osteoporotic fractures demonstrated radiologically; or
  - 2.4 Documented T-Score less than or equal to -3.0; or
  - 2.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm that incorporates BMD measured using DEXA; and
- 3 Any of the following:
  - 3.1 Bisphosphonates are contraindicated because the patient's creatinine clearance or eGFR is less than 35 mL/min; or
  - 3.2 The patient has experienced at least two symptomatic new fractures or a BMD loss greater than 2% per year, after at least 12 months' continuous therapy with a funded antiresorptive agent; or
  - 3.3 Bisphosphonates result in intolerable side effects; or
  - 3.4 Intravenous bisphosphonates cannot be administered due to logistical or technical reasons.

Initial application — (Hypercalcaemia) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has hypercalcaemia of malignancy; and
- 2 Patient has severe renal impairment.

#### PAMIDRONATE DISODIUM

Inj 3 mg per ml, 10 ml vial	32.49	1	<ul> <li>Pamisol</li> </ul>
Inj 6 mg per ml, 10 ml vial		1	Pamisol
Inj 9 mg per ml, 10 ml vial		1	<ul> <li>Pamisol</li> </ul>
RALOXIFENE HYDROCHLORIDE - Special Authority see SA1779	<mark>below</mark> – Retail ph	armacy	
* Tab 60 mg	53.76	28	<ul> <li>Evista</li> </ul>

### ⇒SA1779 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Notes); or
- 2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
- 3 History of two significant osteoporotic fractures demonstrated radiologically; or
- 4 Documented T-Score less than or equal to -3.0 (see Notes); or
- 5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Notes); or

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsi	dised	Generic
\$	Per	1	Manufacturer

- 6 Patient has had a Special Authority approval for zoledronic acid (Underlying cause Osteoporosis) or has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) prior to 1 February 2019.
- Notes:
  - a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
  - b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for raloxifene funding.
  - c) Osteoporotic fractures are the incident events for severe (established) osteoporosis, and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
  - d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

#### RISEDRONATE SODIUM

Tab 35 mg2.50	4	<ul> <li>Risedronate Sandoz</li> </ul>
TERIPARATIDE – Special Authority see SA1139 below – Retail pharmacy		
Inj 250 mcg per ml, 2.4 ml 195.00	1	<ul> <li>Teriparatide - Teva</li> </ul>

### ➡SA1139 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 18 months for applications meeting the following criteria: All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 The patient has a documented T-score less than or equal to -3.0 (see Notes); and
- 3 The patient has had two or more fractures due to minimal trauma; and
- 4 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes).

#### Notes:

- a) The bone mineral density (BMD) measurement used to derive the T-score must be made using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: alendronate sodium tab 70 mg or tab 70 mg with colecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily; zoledronic acid 5 mg per year. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy.
- c) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.
- d) A maximum of 18 months of treatment (18 cartridges) will be subsidised.

### ZOLEDRONIC ACID

Inj 0.05 mg per ml, 100 ml, bag	22.53	100 ml OP	✓ Zoledronic Acid
			Viatris

## Hyperuricaemia and Antigout

ALI	OPURINOL			
*	Tab 100 mg	17.99	1,000	Ipca-Allopurinol
*	Tab 300 mg	22.50	500	<ul> <li>Ipca-Allopurinol</li> </ul>

# MUSCULOSKELETAL SYSTEM

	Subsidy		Fully	Brand or
	(Manufacturer's Price) \$	Per	Subsidised	Generic Manufacturer
	*	Fei		Wallulaciulei
BENZBROMARONE – Special Authority see SA1963 below –				
Tab 50 mg		100	1	Narcaricin mite S29
➡SA1963 Special Authority for Subsidy				
<b>Renewal</b> from any relevant practitioner. Approvals valid for 2 Both:	years for applications m	neetin	ig the follow	ving criteria:
<ol> <li>The treatment remains appropriate and the patient is b</li> <li>There is no evidence of liver toxicity and patient is cont tests.</li> </ol>				three months) liver function
COLCHICINE				
* Tab 500 mcg	6.00	100	✓	Colgout
FEBUXOSTAT - Special Authority see SA2054 below - Reta	il pharmacy			
Tab 80 mg		28	1	Febuxostat (Teva)
Tab 120 mg		28	✓	Febuxostat (Teva)
SA2054 Special Authority for Subsidy				
<b>Initial application — (Gout)</b> from any relevant practitioner. A	Approvals valid for 6 mo	nths	for applicati	ons meeting the following
criteria:				0 0
Both:				
1 Patient has been diagnosed with gout; and				
2 Any of the following:				
2.1 The patient has a serum urate level greater than	n 0.36 mmol/l despite tre	eatme	ent with allo	purinol at doses of at least
600 mg/day and addition of probenecid at doses	s of up to 2 g per day or	maxi	imum tolera	ited dose; or
2.2 The patient has experienced intolerable side eff	ects from allopurinol suc	ch tha	at treatment	discontinuation is required
and serum urate remains greater than 0.36 mm	ol/I despite use of probe	necio	d at doses o	of up to 2 g per day or
maximum tolerated dose; or				
2.3 The patient has renal impairment such that prob	enecid is contraindicate	ed or	likely to be	ineffective and serum urate
remains greater than 0.36 mmol/l despite optima				
2.4 The patient has previously had an initial Special				U U
Initial application — (Tumour lysis syndrome) only from a	haematologist or oncolo	ogist.	Approvals	valid for 6 weeks for
applications meeting the following criteria:				
Both:				
1 Patient is scheduled to receive cancer therapy carrying	•	ı risk	of tumour ly	sis syndrome; and

2 Patient has a documented history of allopurinol intolerance.

Renewal — (Gout) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from treatment.

Renewal — (Tumour lysis syndrome) only from a haematologist or oncologist. Approvals valid for 6 weeks where the treatment remains appropriate and the patient is benefitting from treatment.

PROBENECID

*	Tab 500 mg	100	Probenecid-AFT
N	luscle Relaxants		
BA	CLOFEN		
*	Tab 10 mg	100	<ul> <li>Pacifen</li> </ul>
	Inj 0.05 mg per ml, 1 ml ampoule - Subsidy by endorsement11.55	1	<ul> <li>Lioresal Intrathecal</li> </ul>
	Subsidised only for use in a programmable pump in patients where oral caused intolerable side effects and the prescription is endorsed according according to the prescription of the pr		ents have been ineffective or have
	Inj 2 mg per ml, 5 ml ampoule – Subsidy by endorsement	10	<ul> <li><u>Sintetica Baclofen</u> Intrathecal</li> </ul>
	Cubaidiand only for use in a programmable nump in patients where are	antionactic ca	anto hours has in offective or hours

Subsidised only for use in a programmable pump in patients where oral antispastic agents have been ineffective or have caused intolerable side effects and the prescription is endorsed accordingly.

## MUSCULOSKELETAL SYSTEM

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
DANTROLENE				
Cap 25 mg	112.13	100	✓	Dantrium
	145.77		✓	Dantrium S29 S29
Cap 50 mg	77.00	100	✓	Dantrium
(Dantrium Cap 25 mg to be delisted 1 April 2025)				
ORPHENADRINE CITRATE				
Tab 100 mg	23.25	100	1	Norflex

	Subsidy (Manufacturer's Price) \$	) Per	Fully Brand or Subsidised Generic Manufacturer
Agents for Parkinsonism and Related Disorders			
Dopamine Agonists and Related Agents			
AMANTADINE HYDROCHLORIDE			
▲ Cap 100 mg		60 100	<ul><li>✓ Symmetrel</li><li>✓ Symmetrel</li></ul>
APOMORPHINE HYDROCHLORIDE			
▲ Inj 10 mg per ml, 2 ml ampoule		5	<ul> <li>Movapo</li> </ul>
Inj 10 mg per ml, 5 ml ampoule	121.84	5	<ul> <li>Movapo</li> </ul>
ENTACAPONE			
▲ Tab 200 mg		100	<ul> <li>Entacapone Viatris</li> </ul>
	18.04		<ul> <li>Comtan</li> </ul>
(Comtan Tab 200 mg to be delisted 1 July 2025)			
EVODOPA WITH BENSERAZIDE			
* Tab dispersible 50 mg with benserazide 12.5 mg		100	🗸 Madopar Rapid
* Cap 50 mg with benserazide 12.5 mg		100	
<ul> <li>K Cap 100 mg with benserazide 25 mg</li> </ul>		100	<ul> <li>Madopar 125</li> </ul>
Cap long-acting 100 mg with benserazide 25 mg		100	
Cap 200 mg with benserazide 50 mg		100	<ul> <li>Madopar 250</li> </ul>
EVODOPA WITH CARBIDOPA			
₭ Tab 100 mg with carbidopa 25 mg		100	<ul> <li>Sinemet</li> </ul>
₭ Tab long-acting 200 mg with carbidopa 50 mg		100	✓ Sinemet CR
₭ Tab 250 mg with carbidopa 25 mg		100	✓ Sinemet
EVODOPA WITH CARBIDOPA AND ENTACAPONE			
₭ Tab 50 mg with carbidopa 12.5 mg and entacapone 200 mg		100	<ul> <li>Stalevo</li> </ul>
K Tab 100 mg with carbidopa 25 mg and entacapone 200 mg		100	<ul> <li>Stalevo</li> </ul>
₭ Tab 150 mg with carbidopa 37.5 mg and entacapone 200 mg		100	<ul> <li>Stalevo</li> </ul>
₭ Tab 200 mg with carbidopa 50 mg and entacapone 200 mg		100	<ul> <li>Stalevo</li> </ul>
PRAMIPEXOLE HYDROCHLORIDE			
Tab 0.25 mg	5.51	100	Ramipex
Tab 1 mg		100	
RASAGILINE			
₭ Tab 1 mg	53 50	30	✓ Azilect S29
ROPINIROLE HYDROCHLORIDE		00	ALICOL
Tab 0.25 mg	4.05	84	Ropin
Tab 0.25 mg		84	✓ Ropin
Tab 2 mg		84	✓ Ropin
Tab 5 mg		84	✓ Ropin
		•••	
Tab 100 mg	152 38	100	<ul> <li>Tasmar</li> </ul>
		100	• Tasmai
Anticholinergics			
BENZATROPINE MESYLATE			
	9 59	60	<ul> <li>Benztrop</li> </ul>
Tab 2 mg			
Tab 2 mg Inj 1 mg per ml, 2 ml		5	Phebra
5			

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

NERVOUS SYSTEM

	Subsidy (Manufacturer's Price) \$	Su Per	Fully bsidised	Brand or Generic Manufacturer
PROCYCLIDINE HYDROCHLORIDE Tab 5 mg	7.40	100	✓ к	emadrin
Agents for Essential Tremor, Chorea and Relate	d Disorders			
RILUZOLE – Special Authority see SA1403 below – Retail pharm Wastage claimable Tab 50 mg SA1403 Special Authority for Subsidy	117.00	56	_	l <mark>ilutek</mark>
Initial application only from a neurologist or respiratory specialist following criteria: All of the following: 1 The patient has amyotrophic lateral sclerosis with disease 2 The patient has at least 60 percent of predicted forced vital 3 The patient has not undergone a tracheostomy; and 4 The patient has not experienced respiratory failure; and 5 Any of the following: 5.1 The patient is ambulatory; or 5.2 The patient is able to use upper limbs; or 5.3 The patient is able to swallow.	duration of 5 years o I capacity within 2 m	r less; a onths pr	nd ior to the	initial application; and
<ul> <li>Renewal from any relevant practitioner. Approvals valid for 18 models and 18</li></ul>	onths for applications	s meetin	g the follo	owing criteria:
TETRABENAZINE Tab 25 mg	106.59	112	✓ <u>N</u>	lotetis
Anaesthetics Local				
LIDOCAINE [LIGNOCAINE] Gel 2%, tube – Subsidy by endorsement a) Up to 150 ml available on a PSO b) Subsidised only if prescribed for urethral or cervical ad Gel 2%, 11 ml urethral syringe – Subsidy by endorsement a) Up to 5 each available on a PSO b) Subsidised only if prescribed for urethral, cervical or re	dministration and the	10	otion is er ✓ <u>Ir</u>	nstillagel Lido

accordingly.

	Subsidy		Fully	Brand or
	(Manufacturer's Price	) Subsid	lised	Generic
	\$	Per	1	Manufacturer
LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE				
Oral (gel) soln 2%		200 ml	✓	Mucosoothe
Inj 1%, 5 ml ampoule - Up to 25 inj available on a PSO		25	✓	Lidocaine-Baxter
	17.50	50		
	(35.00)			Xylocaine
Inj 2%, 5 ml ampoule – Up to 5 inj available on a PSO		25	✓	Lidocaine-Baxter
Inj 1%, 20 ml ampoule - Up to 5 inj available on a PSO		5		
	(20.00)			Xylocaine
Inj 1%, 20 ml vial – Up to 5 inj available on a PSO		5	✓	Lidocaine-Baxter
Inj 2%, 20 ml vial – Up to 5 inj available on a PSO	14.00	5	✓	Lidocaine-Baxter
Inj 10%, 5 ml ampoule - Subsidy by endorsement		10		Xylocard 500 S29
Subsidised only for people receiving palliative care servi	ces where other and	algesic agent	s ha	ven't been effective.

### **Topical Local Anaesthetics**

### ⇒SA0906 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years where the patient is a child with a chronic medical condition requiring frequent injections or venepuncture.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

LIDOCAINE [LIGNOCAINE] – Special Authority see SA09	06 above – Retail pharm	nacy	
Crm 4%	5.40	5 g OP	🖌 LMX4
	27.00	30 g OP	🖌 LMX4
LIDOCAINE [LIGNOCAINE] WITH PRILOCAINE - Specia	I Authority see SA0906	<mark>above</mark> – Reta	il pharmacy
Crm 2.5% with prilocaine 2.5%	45.00	30 g OP	🖌 EMLA
Crm 2.5% with prilocaine 2.5% (5 g tubes)	45.00	5	🖌 EMLA

## Analgesics

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 115

### **Non-opioid Analgesics**

ASPIRIN * Tab dispersible 300 mg – Up to 30 tab available on a PSO	5.65	100	✓ Ethics Aspirin
CAPSAICIN – Subsidy by endorsement			
Subsidised only if prescribed for post-herpetic neuralgia or diab accordingly.	etic periphera	l neuropathy a	nd the prescription is endorsed
Crm 0.075%	11.95	45 g OP	<ul> <li>✓ Zo-Rub HP S29</li> <li>✓ Zostrix HP</li> </ul>
	15.14	57 g OP	<ul> <li>Rugby Capsaicin Topical Cream \$29</li> </ul>
(Rugby Capsaicin Topical Cream S29) Crm 0.075% to be delisted	1 July 2025)		
NEFOPAM HYDROCHLORIDE Tab 30 mg	23.40	90	<ul> <li>Acupan</li> </ul>

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

Subsidy (Manufacturer's Price) \$       Fully Subsidised Per       Fully Subsidised Fer       Brand or Generic Manufacturer         PARACETAMOL Tab 500 mg - blister pack
\$       Per       ✓ Manufacturer         PARACETAMOL       Tab 500 mg - blister pack
<ul> <li>PARACETAMOL Tab 500 mg - blister pack</li></ul>
<ul> <li>Tab 500 mg - blister pack</li></ul>
<ul> <li>a) Maximum of 300 tab per prescription; can be waived by endorsement</li> <li>b) Up to 30 tab available on a PSO</li> <li>c) <ol> <li>Subsidy by endorsement for higher quantities is available for patients with long term conditions who require regular daily dosing for one month or greater, and the prescription is annotated accordingly. Pharmacists ma annotate the prescription as endorsed where dispensing history supports a long-term condition.</li> <li>Maximum of 100 tab per dispensing for non-endorsed patients. If quantities prescribed for more than 100 tab (for non-endorsed patients), then dispense in repeat dispensings not exceeding 100 tab per dispensing.</li> </ol> </li> <li>Tab 500 mg - bottle pack – Maximum of 300 tab per prescription; can be waived by endorsement</li></ul>
<ul> <li>b) Up to 30 tab available on a PSO</li> <li>c) <ol> <li>Subsidy by endorsement for higher quantities is available for patients with long term conditions who require regular daily dosing for one month or greater, and the prescription is annotated accordingly. Pharmacists ma annotate the prescription as endorsed where dispensing history supports a long-term condition.</li> <li>Maximum of 100 tab per dispensing for non-endorsed patients. If quantities prescribed for more than 100 tab (for non-endorsed patients), then dispense in repeat dispensings not exceeding 100 tab per dispensing.</li> </ol> </li> <li>Tab 500 mg - bottle pack – Maximum of 300 tab per prescription; can be waived by endorsement</li></ul>
<ul> <li>c)         <ul> <li>Subsidy by endorsement for higher quantities is available for patients with long term conditions who require regular daily dosing for one month or greater, and the prescription is annotated accordingly. Pharmacists ma annotate the prescription as endorsed where dispensing history supports a long-term condition.</li> <li>Maximum of 100 tab per dispensing for non-endorsed patients. If quantities prescribed for more than 100 tab (for non-endorsed patients), then dispense in repeat dispensings not exceeding 100 tab per dispensing.</li> </ul> </li> <li>Tab 500 mg - bottle pack – Maximum of 300 tab per prescription; can be waived by endorsement</li></ul>
<ol> <li>Subsidy by endorsement for higher quantities is available for patients with long term conditions who require regular daily dosing for one month or greater, and the prescription is annotated accordingly. Pharmacists ma annotate the prescription as endorsed where dispensing history supports a long-term condition.</li> <li>Maximum of 100 tab per dispensing for non-endorsed patients. If quantities prescribed for more than 100 tab (for non-endorsed patients), then dispense in repeat dispensings not exceeding 100 tab per dispensing.</li> <li>Tab 500 mg - bottle pack – Maximum of 300 tab per prescription; can be waived by endorsement</li></ol>
<ul> <li>regular daily dosing for one month or greater, and the prescription is annotated accordingly. Pharmacists ma annotate the prescription as endorsed where dispensing history supports a long-term condition.</li> <li>2) Maximum of 100 tab per dispensing for non-endorsed patients. If quantities prescribed for more than 100 tab (for non-endorsed patients), then dispense in repeat dispensings not exceeding 100 tab per dispensing.</li> <li>Tab 500 mg - bottle pack – Maximum of 300 tab per prescription; can be waived by endorsement</li></ul>
<ul> <li>annotate the prescription as endorsed where dispensing history supports a long-term condition.</li> <li>2) Maximum of 100 tab per dispensing for non-endorsed patients. If quantities prescribed for more than 100 tab (for non-endorsed patients), then dispense in repeat dispensings not exceeding 100 tab per dispensing.</li> <li>Tab 500 mg - bottle pack – Maximum of 300 tab per prescription; can be waived by endorsement</li></ul>
<ul> <li>2) Maximum of 100 tab per dispensing for non-endorsed patients. If quantities prescribed for more than 100 tab (for non-endorsed patients), then dispense in repeat dispensings not exceeding 100 tab per dispensing.</li> <li>Tab 500 mg - bottle pack – Maximum of 300 tab per prescription; can be waived by endorsement</li></ul>
<ul> <li>Tab 500 mg - bottle pack – Maximum of 300 tab per prescription; can be waived by endorsement</li></ul>
<ul> <li>prescription; can be waived by endorsement</li></ul>
Paracetamol           1) Subsidy by endorsement for higher quantities is available for patients with long term conditions who require regularity dosing for one month or greater, and the prescription is annotated accordingly. Pharmacists may annotate prescription as endorsed where dispensing history supports a long-term condition.           2) Maximum of 100 tab per dispensing for non-endorsed patients. If quantities prescribed for more than 100 tabs (for non-endorsed patients), then dispense in repeat dispensings not exceeding 100 tab per dispensing.
<ol> <li>Subsidy by endorsement for higher quantities is available for patients with long term conditions who require regulative daily dosing for one month or greater, and the prescription is annotated accordingly. Pharmacists may annotate prescription as endorsed where dispensing history supports a long-term condition.</li> <li>Maximum of 100 tab per dispensing for non-endorsed patients. If quantities prescribed for more than 100 tabs (for non-endorsed patients), then dispense in repeat dispensings not exceeding 100 tab per dispensing.</li> </ol>
<ul> <li>daily dosing for one month or greater, and the prescription is annotated accordingly. Pharmacists may annotate prescription as endorsed where dispensing history supports a long-term condition.</li> <li>2) Maximum of 100 tab per dispensing for non-endorsed patients. If quantities prescribed for more than 100 tabs (f non-endorsed patients), then dispense in repeat dispensings not exceeding 100 tab per dispensing.</li> </ul>
<ul> <li>prescription as endorsed where dispensing history supports a long-term condition.</li> <li>2) Maximum of 100 tab per dispensing for non-endorsed patients. If quantities prescribed for more than 100 tabs (f non-endorsed patients), then dispense in repeat dispensings not exceeding 100 tab per dispensing.</li> </ul>
<ol> <li>Maximum of 100 tab per dispensing for non-endorsed patients. If quantities prescribed for more than 100 tabs (f non-endorsed patients), then dispense in repeat dispensings not exceeding 100 tab per dispensing.</li> </ol>
non-endorsed patients), then dispense in repeat dispensings not exceeding 100 tab per dispensing.
Oral lig 120 mg per 5 ml
(Ethics)
<ul> <li>a) Maximum of 600 ml per prescription; can be waived by endorsement</li> </ul>
b) Up to 200 ml available on a PSO
c) Not in combination
d)
1) Maximum of 200 ml per dispensing for non-endorsed patients. If quantities prescribed exceed 200 ml (for
non-endorsed patients), then dispense in repeat dispensing not exceeding 200 ml per dispensing.
<ol> <li>Subsidy by endorsement for higher quantities is available for patients with long term conditions who require require doily doily doing for one month or greater and the prescription is and aread a constituted accordingly.</li> </ol>
regular daily dosing for one month or greater and the prescription is endorsed or annotated accordingly. Pharmacists may annotate the prescription as endorsed where dispensing history supports a long-term
condition.
<ol> <li>Note: 200 ml presentations of paracetamol oral liquid may be supplied on BSO to a Vaccinator (other than a</li> </ol>
Pharmacist) under the provisions in Part I of Section A
4) Note: Direct Provision by a pharmacist of up to 200 ml permitted under the provisions in Part I of Section A ir
conjunction with immunisation of a child under 2 years of age with meningococcal B multicomponent vaccine.
Oral liq 250 mg per 5 ml Pamol
a) Maximum of 600 ml per prescription; can be waived by endorsement
b) Up to 200 ml available on a PSO
c) Not in combination
<ul> <li>d)</li> <li>1) Maximum of 200 ml per dispensing for non-endorsed patients. If quantities prescribed exceed 200 ml (for</li> </ul>
non-endorsed patients), then dispense in repeat dispensing not exceeding 200 ml per dispensing.
2) Subsidy by endorsement for higher quantities is available for patients with long term conditions who require
regular daily dosing for one month or greater and the prescription is endorsed or annotated accordingly.
Pharmacists may annotate the prescription as endorsed where dispensing history supports a long-term
condition.
3) Note: 200 ml presentations of paracetamol oral liquid may be supplied on BSO to a Vaccinator (other than a
Pharmacist) under the provisions in Part I of Section A
<ol> <li>Note: Direct Provision by a pharmacist of up to 200 ml permitted under the provisions in Part I of Section A in account of a shift under 2 upons of a split under 2 upons of a split maximum section.</li> </ol>
conjunction with immunisation of a child under 2 years of age with meningococcal B multicomponent vaccine. * Suppos 125 mg
[™] 000000 120 mg

—		Qubaidu		Euller	Brand or
		Subsidy (Manufacturer's Price)		Fully Subsidised	Generic
		(Manulacialer 5 Thee) \$	Per		Manufacturer
*	Suppos 250 mg	5,39	10	1	Gacet
*	Suppos 500 mg		50		Gacet
0	pioid Analgesics				
CO	DEINE PHOSPHATE - Safety medicine; prescriber may dete	ermine dispensing fre	quen	су	
	Tab 15 mg		100		Noumed
	Tab 30 mg		100		Noumed
	Tab 60 mg	13.89	100	✓	Noumed
D⊩	YDROCODEINE TARTRATE				
	Tab long-acting 60 mg	8.60	60	✓	DHC Continus
FE	NTANYL				
	a) Only on a controlled drug form				
	b) No patient co-payment payable				
	c) Safety medicine; prescriber may determine dispensing fre	equency			
	Inj 50 mcg per ml, 2 ml ampoule	4.25	10	✓	Boucher and Muir
	Boucher and Muir to be Principal Supply on 1 May 2025				
	Inj 50 mcg per ml, 10 ml ampoule	9.41	10	✓	Boucher and Muir
	Boucher and Muir to be Principal Supply on 1 May 2025				
	Patch 12.5 mcg per hour		5		Fentanyl Sandoz
	Patch 25 mcg per hour		5		Fentanyl Sandoz
	Patch 50 mcg per hour		5		Fentanyl Sandoz
	Patch 75 mcg per hour		5		Fentanyl Sandoz
	Patch 100 mcg per hour	16.37	5	<b>v</b>	Fentanyl Sandoz
ME	THADONE HYDROCHLORIDE				
	a) Only on a controlled drug form				
	b) No patient co-payment payable				
	c) Safety medicine; prescriber may determine dispensing fre				
	Tab 5 mg		10		Methadone BNM
	Oral liq 2 mg per ml		200 m		Biodone
	Oral liq 5 mg per ml		200 m		Biodone Forte
	Oral liq 10 mg per ml		200 m		Biodone Extra Forte
	Inj 10 mg per ml, 1 ml		10	<b>v</b>	AFT
MC	RPHINE HYDROCHLORIDE				
	a) Only on a controlled drug form				
	b) No patient co-payment payable				
	c) Safety medicine; prescriber may determine dispensing fre				
	Oral liq 1 mg per ml		200 m		RA-Morph
	Oral liq 2 mg per ml		200 m		RA-Morph
	Oral liq 5 mg per ml		200 m		RA-Morph
	Oral liq 10 mg per ml		200 m	nl 🗸	RA-Morph

(Mar DRPHINE SULPHATE a) Only on a controlled drug form b) No patient co-payment payable c) Safety medicine; prescriber may determine dispensing frequer	hufacturer's Price) \$	Per	Subsidised	Generic Manufacturer
<ul><li>a) Only on a controlled drug form</li><li>b) No patient co-payment payable</li></ul>				
b) No patient co-payment payable				
b) No patient co-payment payable				
	ncv			
Tab immediate-release 10 mg	2.80	10	✓	Sevredol
Tab immediate-release 20 mg	5.52	10	✓	Sevredol
Cap long-acting 10 mg	3.00	10	✓	m-Eslon
Cap long-acting 30 mg	4.30	10	✓	m-Eslon
Cap long-acting 60 mg	9.00	10	✓	m-Eslon
Cap long-acting 100 mg	10.50	10	✓	m-Eslon
Oral lig 2 mg per ml	16.31	100 m	l 🗸	Wockhardt S29
	29.80		✓	Oramorph
			✓	Oramorph CDC
				S29 S29
Inj 5 mg per ml, 1 ml ampoule – Up to 5 inj available on a PSO	5 38	5	1	Medsurge
Inj 10 mg per ml, 1 ml ampoule – Up to 5 inj available on a PSO.		5		Medsurge
Inj 15 mg per ml, 1 ml ampoule – Up to 5 inj available on a PSO.		5		Medsurge
Inj 30 mg per ml, 1 ml ampoule – Up to 5 inj available on a PSO.		5		Medsurge
YCODONE HYDROCHLORIDE		Ũ		<u>incucui go</u>
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing frequer		~~		
Tab controlled-release 5 mg		20		Oxycodone Sandoz
	3.77	28	•	Oxycodone Sandoz
				S29 S29
	4.04	30		OxyContin S29
Tab immediate-release 5 mg		100		Oxycodone Amneal
Tab controlled-release 10 mg		20		Oxycodone Sandoz
	3.77	28	✓	Oxycodone Sandoz
				S29 S29
Tab immediate-release 10 mg	18.77	100	✓	<b>Oxycodone Amneal</b>
Tab controlled-release 20 mg	3.41	20	✓	Oxycodone Sandoz
Tab immediate-release 20 mg	26.77	100	✓	Oxycodone Amneal
Tab controlled-release 40 mg	6.67	20	✓	Oxycodone Sandoz
Tab controlled-release 80 mg	12.99	20	✓	Oxycodone Sandoz
Oral liq 1 mg per ml	37.08	250 m	✓	Oxycodone Lucis
Inj 10 mg per ml, 1 ml ampoule	4.37	5	✓	Hameln
Inj 10 mg per ml, 2 ml ampoule	8.62	5	✓	Hameln
Inj 50 mg per ml, 1 ml ampoule	14.90	5	✓	Hameln
xycodone Sandoz S29 329 Tab controlled-release 5 mg to be deli	sted 1 July 2025	5)		
xyContin 529 Tab controlled-release 5 mg to be delisted 1 July 20				
xycodone Sandoz S29 s29 Tab controlled-release 10 mg to be de		25)		
	,	,		
RACETAMOL WITH CODEINE – Safety medicine; prescriber may		-		•
Tab paracetamol 500 mg with codeine phosphate 8 mg	27.50	1,000	~	Paracetamol + Codeine (Relieve)

(Manufacturer's Price)       Subsidised       Generic Manufacturer         PETHIDINE HYDROCHLORIDE       a) Only on a controlled drug form       b) No patient co-payment payable       c)         c) Sately medicine; prescriber may determine dispensing frequency       8.68       10       ✓ Noumed Pethidine         Inj 50 mg per ml, 1 ml ampoule – Up to 5 inj available on a PSO29.88       5       ✓ DBL Pethidine       Hydrochloride         TRAMADOL HYDROCHLORIDE       1.95       20       ✓ Tramal ST 100       Tab sustained-release 150 mg       2.95       20       ✓ Tramal ST 100         Tab sustained-release 150 mg       2.95       20       ✓ Tramal ST 100       Tab sustained-release 150 mg       3.80       20       ✓ Tramal ST 100         Tab sustained-release 100 mg       2.95       20       ✓ Tramal ST 100       Cartow-Antritiptylin         Tab sustained-release 200 mg       3.80       20       ✓ Tramal ST 100       Arrow-Antritiptylin         Tab 50 mg		Subsidy		Fully	Brand or
PETHIDINE HYDROCHLORIDE       a) Only on a controlled drug form         b) No patient co-payment payable       c) Safety medicine; prescriber may determine dispensing frequency         Tab 50 mg       main appoule – Up to 5 inj available on a PSO29.88       5       ✓ DBL Pethidine         Inj 50 mg per ml, 1 ml ampoule – Up to 5 inj available on a PSO30.72       5       ✓ DBL Pethidine         Hydrochloride       Tramal SR 100       ✓ Tramal SR 100       Tramal SR 100         Tab sustained-release 100 mg      95       20       ✓ Tramal SR 100         Tab sustained-release 100 mg      95       20       ✓ Tramal SR 100         Tab sustained-release 100 mg      95       20       ✓ Tramal SR 200         Cap 50 mg		(Manufacturer's Price)		Subsidised	Generic
a) Only on a controlled drug form b) No patient co-payment payable c) Safety medicine; prescriber may determine dispensing frequency Tab 50 mg		\$	Per	<i>✓</i>	Manufacturer
b) No patient co-payment payable c) Safety medicine; prescriber may determine dispensing frequency Tab 50 mg per ml, 1 ml ampoule – Up to 5 inj available on a PSO29.88 5 // DBL Pethidine Hydrochloride Inj 50 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO30.72 5 // DBL Pethidine Hydrochloride TRAMADOL HYDROCHLORIDE Tab sustained-release 100 mg					
c) Safety medicine; prescriber may determine dispensing frequency Tab 50 mg	, ,				
Tab 50 mg       Automation         Inj 50 mg per ml, 1 ml ampoule – Up to 5 inj available on a PSO29.88       0       ✓ Noumed Pethidine         Inj 50 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO30.72       5       ✓ DBL Pethidine         Hydrochloride       Tramal SR 100       ✓ Tramal SR 100       ✓ Tramal SR 100         Tab sustained-release 100 mg       1.95       20       ✓ Tramal SR 100         Tab sustained-release 100 mg       3.80       20       ✓ Tramal SR 200         Cap 50 mg       3.80       20       ✓ Tramal SR 120         Antidepressants       ✓ Arrow-Amitriptylin       Yarow-Amitriptylin         Tab 25 mg       1.99       100       ✓ Arrow-Amitriptylin         Tab 25 mg       1.99       00       ✓ Arrow-Amitriptylin         Tab 25 mg       1.99       00       ✓ Arrow-Amitriptylin         CLOMIPRAMINE HYDROCHLORIDE – Safety medicine; prescriber may determine dispensing frequency       ✓ Clomipramine Tew         Tab 25 mg       1.99       00       ✓ Arrow-Amitriptylin         CLOMIPRAMINE HYDROCHLORIDE – Safety medicine; prescriber may determine dispensing frequency       ✓ Clomipramine Tew         Tab 25 mg       10.17       30       ✓ Clomipramine Tew         Cap 10 mg       10.17       30       ✓ Antrow-Amitriptylin </td <td></td> <td></td> <td></td> <td></td> <td></td>					
Inj 50 mg per ml, 1 ml ampoule – Up to 5 inj available on a PSO29.88 5 / DBL Pethidine Hydrochloride Inj 50 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO30.72 5 / DBL Pethidine Hydrochloride TRAMADOL HYDROCHLORIDE Tab sustained-release 100 mg			10	1	Noumed Pothidine
Hydrochloride       Hydrochloride         Inj 50 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO30.72       5       ✓ DBL Pethidine         Hydrochloride       Hydrochloride       Hydrochloride         Tab sustained-release 150 mg       2.95       20       ✓ Tramal SR 150         Tab sustained-release 150 mg       2.95       20       ✓ Tramal SR 150         Tab sustained-release 200 mg       3.80       20       ✓ Tramal SR 150         Cap 50 mg       3.80       20       ✓ Tramal SR 200         Cap 50 mg       3.80       20       ✓ Tramal SR 200         Antidepressants       ✓       Arrow-Amitriptylin         Tab 25 mg       1.99       100       ✓ Arrow-Amitriptylin         Tab 25 mg					
Inj 50 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO30.72 5 ✓ DBL Pethidine Hydrochloride TRAMADOL HYDROCHLORIDE Tab sustained-release 100 mg		1 3029.00	5	•	
TRAMADOL HYDROCHLORIDE       1.95       20       ✓ Tramal SR 100         Tab sustained-release 100 mg	Ini 50 mg per ml. 2 ml ampoule. – Up to 5 ini available on a	PSO 30.72	5	1	•
TRAMADOL HYDROCHLORIDE       1.95       20			Ũ		
Tab sustained-release 100 mg       1.95       20       ✓ Tramal SR 100         Tab sustained-release 150 mg       2.95       20       ✓ Tramal SR 100         Tab sustained-release 200 mg       3.80       20       ✓ Tramal SR 200         Cap 50 mg       3.33       100       ✓ Arrow-Tramadol         Antidepressants         Arrow-Amitriptylin         Tab 25 mg       1.99       100       ✓ Arrow-Amitriptylin         Tab 25 mg       1.99       100       ✓ Arrow-Amitriptylin         Tab 50 mg       1.99       100       ✓ Arrow-Amitriptylin         Tab 50 mg       1.99       00       ✓ Arrow-Amitriptylin         CLOMIPRAMINE HYDROCHLORIDE – Safety medicine; prescriber may determine dispensing frequency       Tab 10 mg       0       ✓ Clomipramine Teva         Tab 25 mg       10.17       30       ✓ Clomipramine Teva       ✓ Clomipramine Teva         Cap 10 mg       55.0       28       ✓ Clomipramine Teva       ✓ Clomipramine Teva         Cap 25 mg       55.0       28 <td></td> <td></td> <td></td> <td></td> <td></td>					
Tab sustained-release 150 mg       2.95       20       ✓ Tramal SR 150         Tab sustained-release 200 mg       3.80       20       ✓ Tramal SR 200         Cap 50 mg       3.33       100       ✓ Arrow-Tramadol         Antidepressants        Arrow-Amitriptylin         Tab 50 mg       1.99       100       ✓ Arrow-Amitriptylin         Tab 50 mg       1.99       100       ✓ Arrow-Amitriptylin         Tab 50 mg       3.14       100       ✓ Arrow-Amitriptylin         Tab 50 mg       1.99       00       ✓ Arrow-Amitriptylin         Tab 50 mg       1.99       00       ✓ Arrow-Amitriptylin         Tab 50 mg       1.99       00       ✓ Arrow-Amitriptylin         CLOMIPRAMINE HYDROCHLORIDE – Safety medicine; prescriber may determine dispensing frequency       ✓ Clomipramine Teva         Tab 25 mg       0       ✓ Clomipramine Teva       ✓ Clomipramine Teva         Qa 25 mg       0       ✓ APO Clomipramine Teva       ✓ Clomipramine Teva         Qa 25 mg       10 mg to be delist		1.95	20	1	Tramal SR 100
Tab sustained-release 200 mg       3.30       20       ✓ Tramal SR 200         Cap 50 mg       3.33       100       ✓ Arrow-Tramadol         Antidepressants         Arrow-Tramadol         Cyclic and Related Agents            AMITRIPTYLINE – Safety medicine; prescriber may determine dispensing frequency Tab 10 mg         Arrow-Amitriptylin         Tab 50 mg       1.99       100         Arrow-Amitriptylin         CLOMIPRAMINE HYDROCHLORIDE – Safety medicine; prescriber may determine dispensing frequency         Arrow-Amitriptylin         CLOMIPRAMINE HYDROCHLORIDE – Safety medicine; prescriber may determine dispensing frequency         Arrow-Amitriptylin         CLOMIPRAMINE HYDROCHLORIDE – Safety medicine; prescriber may determine dispensing frequency         Arrow-Amitriptylin         CLOMIPRAMINE HYDROCHLORIDE – Safety medicine; prescriber may determine dispensing frequency            Tab 25 mg       .00        Application           Clomipramine Teva       .0.17       .00            Cap 10 mg       .00        Anfarani            Clomipramine Teva Tab 10 mg to	5				
Antidepressants         Cyclic and Related Agents         AMITRIPTYLINE – Safety medicine; prescriber may determine dispensing frequency Tab 10 mg       2.99       100       ✓ Arrow-Amitriptylin Tab 25 mg         Tab 25 mg       1.99       100       ✓ Arrow-Amitriptylin Tab 50 mg       ✓ Arrow-Amitriptylin CLOMIPRAMINE HYDROCHLORIDE – Safety medicine; prescriber may determine dispensing frequency         Tab 10 mg       10.17       30       ✓ Clomipramine Teva         Tab 25 mg       11.99       30       ✓ Clomipramine Teva         Tab 25 mg       10.17       30       ✓ Clomipramine Teva         Tab 25 mg       10.99       50       ✓ APO Clomipramine Teva         Tab 25 mg       35.50       28       ✓ Clomipramine Teva         Cap 10 mg       28 mg       35.50       28       ✓ Clomipramine Teva         Cap 25 mg       35.50       28       ✓ Clomipramine Teva         Clomipramine Teva Tab 25 mg to be delisted 1 July 2025)       (Clomipramine Teva Cap 10 mg to be delisted 1 July 2025)       (Clomipramine Teva Cap 25 mg to be delisted 1 July 2025)         Clomipramine Teva Cap 25 mg to be delisted 1 July 2025)       (Clomipramine Teva Cap 25 mg to be delisted 1 July 2025)       (Clomipramine Teva Cap 25 mg to be delisted 1 July 2025)         DOSULEPIN [DOTHIEPIN] HYDROCHLORIDE – Subsidy by endorsement       a) Safety medicine; prescriber may determine	•		20	1	Tramal SR 200
Cyclic and Related Agents         AMITRIPTYLINE – Safety medicine; prescriber may determine dispensing frequency         Tab 10 mg       2.99       100       ✓ Arrow-Amitriptylin         Tab 25 mg       1.99       100       ✓ Arrow-Amitriptylin         Tab 50 mg       3.14       100       ✓ Arrow-Amitriptylin         Tab 50 mg       3.14       100       ✓ Arrow-Amitriptylin         CLOMIPRAMINE HYDROCHLORIDE – Safety medicine; prescriber may determine dispensing frequency       Tab 10 mg       ✓ Clomipramine Teva         Tab 25 mg       10.17       30       ✓ Clomipramine Teva         Tab 25 mg       11.99       30       ✓ Clomipramine Teva         Tab 25 mg       39.97       100       ✓ Anafranil S20         Cap 10 mg       35.50       28       ✓ Clomipramine Teva         Cap 25 mg       35.50       28       ✓ Clomipramine Teva         (Clomipramine Teva Tab 10 mg to be delisted 1 July 2025)       (Clomipramine Teva Tab 25 mg to be delisted 1 July 2025)       (Clomipramine Teva Cap 10 mg to be delisted 1 July 2025)         (Clomipramine Teva Cap 25 mg to be delisted 1 July 2025)       (Clomipramine Teva Cap 25 mg to be delisted 1 July 2025)       (Clomipramine Teva Cap 25 mg to be delisted 1 July 2025)         (Clomipramine Teva Cap 25 mg to be delisted 1 July 2025)       (Clomipramine Teva Cap 25 m	Cap 50 mg	3.33	100	1	Arrow-Tramadol
Cyclic and Related Agents         AMITRIPTYLINE – Safety medicine; prescriber may determine dispensing frequency         Tab 10 mg       2.99       100       ✓ Arrow-Amitriptylin         Tab 25 mg       1.99       100       ✓ Arrow-Amitriptylin         Tab 50 mg       3.14       100       ✓ Arrow-Amitriptylin         Tab 50 mg       3.14       100       ✓ Arrow-Amitriptylin         CLOMIPRAMINE HYDROCHLORIDE – Safety medicine; prescriber may determine dispensing frequency       Tab 10 mg       ✓ Clomipramine Teva         Tab 25 mg       10.17       30       ✓ Clomipramine Teva         Tab 25 mg       11.99       30       ✓ Clomipramine Teva         Tab 25 mg       39.97       100       ✓ Anafranil S20         Cap 10 mg       35.50       28       ✓ Clomipramine Teva         Cap 25 mg       35.50       28       ✓ Clomipramine Teva         (Clomipramine Teva Tab 10 mg to be delisted 1 July 2025)       (Clomipramine Teva Tab 25 mg to be delisted 1 July 2025)       (Clomipramine Teva Cap 10 mg to be delisted 1 July 2025)         (Clomipramine Teva Cap 25 mg to be delisted 1 July 2025)       (Clomipramine Teva Cap 25 mg to be delisted 1 July 2025)       (Clomipramine Teva Cap 25 mg to be delisted 1 July 2025)         (Clomipramine Teva Cap 25 mg to be delisted 1 July 2025)       (Clomipramine Teva Cap 25 m					
AMITRIPTYLINE – Safety medicine; prescriber may determine dispensing frequency Tab 10 mg	Antidepressants				
Tab 10 mg       2.99       100       ✓ Arrow-Amitriptylin         Tab 25 mg       1.99       100       ✓ Arrow-Amitriptylin         Tab 50 mg       3.14       100       ✓ Arrow-Amitriptylin         CLOMIPRAMINE HYDROCHLORIDE – Safety medicine; prescriber may determine dispensing frequency       10.17       30       ✓ Clomipramine Teva         Tab 25 mg       11.99       30       ✓ Clomipramine Teva       16.99       50       ✓ APO Clomipramine Teva         Tab 25 mg       11.99       30       ✓ Clomipramine Teva       16.99       50       ✓ Anafranil see         Cap 10 mg       39.97       100       ✓ Anafranil see       ✓ Clomipramine Teva       28       ✓ Clomipramine Teva         Cap 10 mg       35.50       28       ✓ Clomipramine Teva       ✓ Clomipramine Teva       Clomipramine Teva         Clomipramine Teva Tab 10 mg to be delisted 1 July 2025)       (Clomipramine Teva Cap 10 mg to be delisted 1 July 2025)       ✓ Clomipramine Teva         Clomipramine Teva Cap 10 mg to be delisted 1 July 2025)       ✓ Clomipramine Teva       ✓ Description         Clomipramine Teva Cap 25 mg to be delisted 1 July 2025)       ✓ Clomipramine Teva       ✓ Description         Clomipramine Teva Cap 25 mg to be delisted 1 July 2025)       ✓ Clomipramine Teva       ✓ Description         DOSULEPIN [DOTHIEPIN] HYDR	Cyclic and Related Agents				
Tab 25 mg       1.99       100       ✓ Arrow-Amitriptylin         Tab 50 mg       3.14       100       ✓ Arrow-Amitriptylin         CLOMIPRAMINE HYDROCHLORIDE       – Safety medicine; prescriber may determine dispensing frequency       10.17       30       ✓ Clomipramine Teva         Tab 25 mg       11.99       30       ✓ Clomipramine Teva       16.99       50       ✓ APO Clomipramine Teva         Cap 10 mg       35.50       28       ✓ Clomipramine Teva       Clomipramine Teva       28       ✓ Clomipramine Teva         C(Comipramine Teva Tab 10 mg to be delisted 1 July 2025)       (Clomipramine Teva Tab 25 mg to be delisted 1 July 2025)       ✓ Clomipramine Teva       Clomipramine Teva         (Clomipramine Teva Cap 10 mg to be delisted 1 July 2025)       (Clomipramine Teva Cap 10 mg to be delisted 1 July 2025)       ✓ Clomipramine Teva       ✓ Clomipramine Teva         (Clomipramine Teva Cap 10 mg to be delisted 1 July 2025)       (Clomipramine Teva Cap 25 mg to be delisted 1 July 2025)       ✓ Clomipramine Teva       ✓ Clomipramine Teva         (Soutepeln [DOTHIEPIN] HYDROCHLORIDE – Subsidy by endorsement       a)       Safety medicine; prescriber may determine dispensing frequency       b)       Subsidy by endorsement – Subsidised for patients who were taking dosulepin [dothiepin] hydrochloride prior to 1 J       2019 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where       ✓ Dosulepin Viatris	MITRIPTYLINE - Safety medicine; prescriber may determine	dispensing frequency			
Tab 50 mg       3.14       100       ✓ Arrow-Amitriptylin         CLOMIPRAMINE HYDROCHLORIDE – Safety medicine; prescriber may determine dispensing frequency         Tab 10 mg       10.17       30       ✓ Clomipramine Teva         Tab 25 mg       11.99       30       ✓ Clomipramine Teva         16.99       50       ✓ APO Clomipramine       400         Cap 10 mg       35.50       28       ✓ Clomipramine Teva         Cap 25 mg       35.50       28       ✓ Clomipramine Teva         Clomipramine Teva Tab 10 mg to be delisted 1 July 2025)       (Clomipramine Teva Tab 25 mg to be delisted 1 July 2025)       (Clomipramine Teva Tab 25 mg to be delisted 1 July 2025)         (Clomipramine Teva Cap 10 mg to be delisted 1 July 2025)       (Clomipramine Teva Cap 25 mg to be delisted 1 July 2025)       (Clomipramine Teva Cap 25 mg to be delisted 1 July 2025)         (Clomipramine Teva Cap 10 mg to be delisted 1 July 2025)       (Clomipramine Teva Cap 25 mg to be delisted 1 July 2025)       (Clomipramine Teva Cap 25 mg to be delisted 1 July 2025)         DOSULEPIN [DOTHIEPIN] HYDROCHLORIDE – Subsidy by endorsement       a) Safety medicine; prescriber may determine dispensing frequency       b) Subsidy by endorsement – Subsidised for patients who were taking dosulepin [dothiepin] hydrochloride prior to 1 J 2019 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where exists a record of prior dispensing of dosulepin [dothiepin] hydrochloride. <tr< td=""><td></td><td></td><td></td><td></td><td></td></tr<>					
CLOMIPRAMINE HYDROCHLORIDE       – Safety medicine; prescriber may determine dispensing frequency         Tab 10 mg       10.17       30       ✓ Clomipramine Teva         Tab 25 mg       11.99       30       ✓ Clomipramine Teva         16.99       50       ✓ APO Clomipramine Teva         39.97       100       ✓ Anafranil S29         Cap 10 mg       35.50       28       ✓ Clomipramine Teva         Cap 25 mg       35.50       28       ✓ Clomipramine Teva         Clomipramine Teva Tab 10 mg to be delisted 1 July 2025)       (Clomipramine Teva Tab 25 mg to be delisted 1 July 2025)       ✓ Clomipramine Teva         (Clomipramine Teva Cap 10 mg to be delisted 1 July 2025)       (Clomipramine Teva Cap 25 mg to be delisted 1 July 2025)       ✓ Clomipramine Teva Cap 25 mg to be delisted 1 July 2025)         (Clomipramine Teva Cap 10 mg to be delisted 1 July 2025)       ✓ Clomipramine Teva Cap 25 mg to be delisted 1 July 2025)       ✓ Clomipramine Teva Cap 25 mg to be delisted 1 July 2025)         (COSULEPIN [DOTHIEPIN] HYDROCHLORIDE – Subsidy by endorsement       a) Safety medicine; prescriber may determine dispensing frequency       b) Subsidy by endorsement – Subsidised for patients who were taking dosulepin [dothiepin] hydrochloride prior to 1 J 2019 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where exists a record of prior dispensing of dosulepin [dothiepin] hydrochloride.         Tab 75 mg       3.85       30 </td <td></td> <td></td> <td></td> <td></td> <td></td>					
Tab 10 mg       10.17       30       ✓ Clomipramine Teva         Tab 25 mg       11.99       30       ✓ Clomipramine Teva         16.99       50       ✓ APO Clomipramine       39.97         Cap 10 mg       35.50       28       ✓ Clomipramine Teva         Cap 25 mg       35.50       28       ✓ Clomipramine Teva         Clomipramine Teva Tab 10 mg to be delisted 1 July 2025)       28       ✓ Clomipramine Teva         (Clomipramine Teva Tab 25 mg to be delisted 1 July 2025)       (Clomipramine Teva Cap 10 mg to be delisted 1 July 2025)       ✓ Clomipramine Teva         (Clomipramine Teva Cap 10 mg to be delisted 1 July 2025)       (Clomipramine Teva Cap 25 mg to be delisted 1 July 2025)       ✓ Clomipramine Teva         (Clomipramine Teva Cap 10 mg to be delisted 1 July 2025)       (Clomipramine Teva Cap 25 mg to be delisted 1 July 2025)       ✓ Clomipramine Teva         (OSULEPIN [DOTHIEPIN] HYDROCHLORIDE – Subsidy by endorsement       a) Safety medicine; prescriber may determine dispensing frequency       b) Subsidy by endorsement – Subsidised for patients who were taking dosulepin [dothiepin] hydrochloride prior to 1 J 2019 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where exists a record of prior dispensing of dosulepin [dothiepin] hydrochloride.         Tab 75 mg       3.85       30       ✓ Dosulepin Viatris Cap 25 mg.         Cap 25 mg       7.83       50       ✓ Dosul	5				
Tab 25 mg       11.99       30       ✓ Clomipramine Teva         16.99       50       ✓ APO Clomipramine         39.97       100       ✓ Anafranil \$29         Cap 10 mg       35.50       28       ✓ Clomipramine Teva         Cap 25 mg       35.50       28       ✓ Clomipramine Teva         (Clomipramine Teva Tab 10 mg to be delisted 1 July 2025)       35.50       28       ✓ Clomipramine Teva         (Clomipramine Teva Tab 25 mg to be delisted 1 July 2025)       (Clomipramine Teva Cap 10 mg to be delisted 1 July 2025)       ✓ Clomipramine Teva         (Clomipramine Teva Cap 10 mg to be delisted 1 July 2025)       (Clomipramine Teva Cap 25 mg to be delisted 1 July 2025)       ✓ Clomipramine Teva Cap 25 mg to be delisted 1 July 2025)         (OSULEPIN [DOTHIEPIN] HYDROCHLORIDE – Subsidy by endorsement       a) Safety medicine; prescriber may determine dispensing frequency       b) Subsidy by endorsement – Subsidised for patients who were taking dosulepin [dothiepin] hydrochloride prior to 1 J 2019 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where exists a record of prior dispensing of dosulepin [dothiepin] hydrochloride.         Tab 75 mg       3.85       30       ✓ Dosulepin Viatris Cap 25 mg         Cap 25 mg       7.83       50       ✓ Dosulepin			•		
16.99       50       ✓ APO Clomipramine         39.97       100       ✓ Anafranil \$29         Cap 10 mg       35.50       28       ✓ Clomipramine Teva         Cap 25 mg       35.50       28       ✓ Clomipramine Teva         Clomipramine Teva Tab 10 mg to be delisted 1 July 2025)       35.50       28       ✓ Clomipramine Teva         Clomipramine Teva Tab 25 mg to be delisted 1 July 2025)       Yestigation       Yestigation       Yestigation         Clomipramine Teva Cap 10 mg to be delisted 1 July 2025)       Yestigation       Yestigation       Yestigation         Clomipramine Teva Cap 10 mg to be delisted 1 July 2025)       Yestigation       Yestigation       Yestigation         Clomipramine Teva Cap 25 mg to be delisted 1 July 2025)       Yestigation       Yestigation       Yestigation         OSULEPIN [DOTHIEPIN] HYDROCHLORIDE – Subsidy by endorsement       a)       Safety medicine; prescriber may determine dispensing frequency       Yestigation       Yestigation       Yestigation         b)       Subsidy by endorsement – Subsidised for patients who were taking dosulepin [dothiepin] hydrochloride prior to 1 J       2019 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where       exists a record of prior dispensing of dosulepin [dothiepin] hydrochloride.         Tab 75 mg       3.85       30       ✓ Dosulepin Viatris <td></td> <td></td> <td></td> <td></td> <td></td>					
39.97       100 <ul> <li>Anafranil</li> <li>Sister and the prescription is endorsed accordingly.</li> <li>Pharmacists may annotate the prescription as endorsed where exists a record of prior dispensing of dosulepin [dothiepin] hydrochloride.</li> <li>Tab 75 mg</li> <li>Tab 75 mg</li></ul>	Tab 25 mg				
Cap 10 mg					•
Cap 25 mg	Cap 10 mg				
<ul> <li>(Clomipramine Teva Tab 10 mg to be delisted 1 July 2025)</li> <li>(Clomipramine Teva Tab 25 mg to be delisted 1 July 2025)</li> <li>(Anafranil ⁶²⁹ Tab 25 mg to be delisted 1 July 2025)</li> <li>(Clomipramine Teva Cap 10 mg to be delisted 1 July 2025)</li> <li>(Clomipramine Teva Cap 25 mg to be delisted 1 July 2025)</li> <li>(Clomipramine Teva Cap 25 mg to be delisted 1 July 2025)</li> <li>(DOSULEPIN [DOTHIEPIN] HYDROCHLORIDE – Subsidy by endorsement <ul> <li>a) Safety medicine; prescriber may determine dispensing frequency</li> <li>b) Subsidy by endorsement – Subsidised for patients who were taking dosulepin [dothiepin] hydrochloride prior to 1 J 2019 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where exists a record of prior dispensing of dosulepin [dothiepin] hydrochloride.</li> <li>Tab 75 mg</li></ul></li></ul>	1 5				
<ul> <li>(Clomipramine Teva Tab 25 mg to be delisted 1 July 2025)</li> <li>(Anafranil 329 Tab 25 mg to be delisted 1 July 2025)</li> <li>(Clomipramine Teva Cap 10 mg to be delisted 1 July 2025)</li> <li>(Clomipramine Teva Cap 25 mg to be delisted 1 July 2025)</li> <li>DOSULEPIN [DOTHIEPIN] HYDROCHLORIDE – Subsidy by endorsement <ul> <li>a) Safety medicine; prescriber may determine dispensing frequency</li> <li>b) Subsidy by endorsement – Subsidised for patients who were taking dosulepin [dothiepin] hydrochloride prior to 1 J 2019 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where exists a record of prior dispensing of dosulepin [dothiepin] hydrochloride.</li> <li>Tab 75 mg</li></ul></li></ul>			20	-	
<ul> <li>(Anafranil S29) Tab 25 mg to be delisted 1 July 2025)</li> <li>(Clomipramine Teva Cap 10 mg to be delisted 1 July 2025)</li> <li>(Clomipramine Teva Cap 25 mg to be delisted 1 July 2025)</li> <li>(DOSULEPIN [DOTHIEPIN] HYDROCHLORIDE – Subsidy by endorsement <ul> <li>a) Safety medicine; prescriber may determine dispensing frequency</li> <li>b) Subsidy by endorsement – Subsidised for patients who were taking dosulepin [dothiepin] hydrochloride prior to 1 J 2019 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where exists a record of prior dispensing of dosulepin [dothiepin] hydrochloride.</li> <li>Tab 75 mg</li> <li>Cap 25 mg</li> <li>Tab 75 mg</li></ul></li></ul>					
<ul> <li>Clomipramine Teva Cap 10 mg to be delisted 1 July 2025)</li> <li>Clomipramine Teva Cap 25 mg to be delisted 1 July 2025)</li> <li>DOSULEPIN [DOTHIEPIN] HYDROCHLORIDE – Subsidy by endorsement <ul> <li>a) Safety medicine; prescriber may determine dispensing frequency</li> <li>b) Subsidy by endorsement – Subsidised for patients who were taking dosulepin [dothiepin] hydrochloride prior to 1 J 2019 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where exists a record of prior dispensing of dosulepin [dothiepin] hydrochloride.</li> <li>Tab 75 mg</li></ul></li></ul>					
<ul> <li>DOSULEPIN [DOTHIEPIN] HYDROCHLORIDE – Subsidy by endorsement <ul> <li>a) Safety medicine; prescriber may determine dispensing frequency</li> <li>b) Subsidy by endorsement – Subsidised for patients who were taking dosulepin [dothiepin] hydrochloride prior to 1 J 2019 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where exists a record of prior dispensing of dosulepin [dothiepin] hydrochloride.</li> <li>Tab 75 mg</li></ul></li></ul>					
<ul> <li>a) Safety medicine; prescriber may determine dispensing frequency</li> <li>b) Subsidy by endorsement – Subsidised for patients who were taking dosulepin [dothiepin] hydrochloride prior to 1 J 2019 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where exists a record of prior dispensing of dosulepin [dothiepin] hydrochloride.</li> <li>Tab 75 mg</li></ul>	Clomipramine Teva Cap 25 mg to be delisted 1 July 2025)				
<ul> <li>b) Subsidy by endorsement – Subsidised for patients who were taking dosulepin [dothiepin] hydrochloride prior to 1 J 2019 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where exists a record of prior dispensing of dosulepin [dothiepin] hydrochloride.</li> <li>Tab 75 mg</li></ul>	OSULEPIN [DOTHIEPIN] HYDROCHLORIDE – Subsidy by e	ndorsement			
2019 and the prescription is endorsed accordingly.       Pharmacists may annotate the prescription as endorsed where exists a record of prior dispensing of dosulepin [dothiepin] hydrochloride.         Tab 75 mg       30       ✓ Dosulepin Viatris Cap 25 mg         Cap 25 mg       7.83       50       ✓ Dosulepin	a) Safety medicine; prescriber may determine dispensing fr	requency			
exists a record of prior dispensing of dosulepin [dothiepin] hydrochloride. Tab 75 mg	b) Subsidy by endorsement - Subsidised for patients who	were taking dosulepin	[doth	iepin] hydi	ochloride prior to 1 June
Tab 75 mg       30       ✓ Dosulepin Viatris         Cap 25 mg       7.83       50       ✓ Dosulepin	1 1 0,		e the	prescriptio	n as endorsed where there
Cap 25 mg					<b>_</b>
	0				•
Viatris \$29	Cap 25 mg		50	~	•
MIPRAMINE HYDROCHLORIDE – Safety medicine; prescriber may determine dispensing frequency					
Tab 10 mg5.48 50 ✓ Tofranil	Tab 10 mg				
10.96 100 ✓ Tofranil	Tob 05 mg				
Tab 25 mg4.93 28 ✓ Imipramine	iau ∠o mg	4.93	28	•	•
Crescent \$29		0.00	<b>F</b> 0		
8.80 50 <b>✓ Tofranil</b>		8.80	50	<b>v</b>	Iotranii

A Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

	Subsidy		Fully	Brand or
	(Manufacturer's Price) \$	Sub Per	sidised	Generic Manufacturer
	÷			
ORTRIPTYLINE HYDROCHLORIDE – Safety medicine; pres				
Tab 10 mg		100		orpress
Tab 25 mg	6.29	180	✓ N	orpress
Monoamine-Oxidase Inhibitors (MAOIs) - Non	Selective			
ANYLCYPROMINE SULPHATE				
Tab 10 mg		50	🗸 Р	arnate
Monoamine-Oxidase Type A Inhibitors				
IOCLOBEMIDE				
F Tab 150 mg	23.60	60	✓ <u>A</u>	urorix
F Tab 300 mg		60	✓ <u>A</u>	urorix
Selective Serotonin Reuptake Inhibitors				
ITALOPRAM HYDROBROMIDE				
← Tab 20 mg	2.86	84	<b>√</b> C	elapram
SCITALOPRAM			_	
€ Tab 10 mg	0 79	28	🖌 ir	oca-Escitalopram
• 140 10 mg	1.07	20		scitalopram
	1.07		• -	(Ethics)
- Tab 20 mg	1 /0	28	🖌 Ir	ca-Escitalopram
•		20	• 1	
	0.50			
<ul> <li>Tab dispersible 20 mg, scored – Subsidy by endorsement.</li> </ul>	2.50	28	✓ F	luox
Subsidised by endorsement				and a the set of a second second set
<ol> <li>When prescribed for a patient who cannot swallor accordingly are</li> </ol>	w whole tablets of caps	sules and	the pres	scription is endorsed
accordingly; or 2) When prescribed in a daily dose that is not a mult	tiple of 20 mg in which	aaaa tha	orocorin	tion is deemed to be
endorsed. Note: Tablets should be combined wi				
endorsed. Note. Tablets should be combined wi			ilai 101	ny uoses.
← Cap 20 mg	3 13	90	۸ 🗸	rrow-Fluoxetine
		00	• •	
AROXETINE		00		!
• Tab 20 mg	4.11	90	ΨL	oxamine
ERTRALINE				
<ul> <li>Tab 50 mg</li> </ul>		30	_	etrona
Fab 100 mg	1.74	30	✓ <u>s</u>	etrona
Other Antidepressants				
IRTAZAPINE				
Tab 30 mg	2.60	30	🗸 N	oumed
Tab 45 mg		30		oumed
ENLAFAXINE				
Cap 37.5 mg	8 29	84	✓ F	nlafax XR
Cap 75 mg		28		nlafax XR
	10.32	20 84		nlafax XR
Cap 150 mg		28		nlafax XR
	13.95	20 84		nlafax XR
	10.90	04	- C	

	Subsidy		Fully Brand or
	(Manufacturer's Price	) Sub	sidised Generic
	\$	Per	<ul> <li>Manufacturer</li> </ul>
Antiepilepsy Drugs			
Agents for Control of Status Epilepticus			
DIAZEPAM - Safety medicine; prescriber may determine dis	nensing frequency		
Inj 5 mg per ml, 2 ml ampoule – Subsidy by endorseme		5	✓ Hospira
a) Up to 5 inj available on a PSO	1E7.0E	U	· Hospita
b) Only on a PSO			
<ul> <li>c) PSO must be endorsed "not for anaesthetic proc</li> </ul>	oduros"		
Rectal tubes 5 mg – Up to 5 tube available on a PSO		5	✓ Stesolid
		5	• <u>otesonu</u>
HENYTOIN SODIUM			
Inj 50 mg per ml, 2 ml ampoule – Up to 5 inj available or			_
PSO		5	<ul> <li>Hospira</li> </ul>
Inj 50 mg per ml, 5 ml ampoule – Up to 5 inj available or			
PSO	154.01	5	<ul> <li>Hospira</li> </ul>
Control of Epilepsy			
ARBAMAZEPINE			
🖌 Tab 200 mg		100	<ul> <li>Tegretol</li> </ul>
			<ul> <li>Tegretol AU</li> </ul>
Fab long-acting 200 mg		100	<ul> <li>Tegretol CR</li> </ul>
5 5 5	33.96	200	<ul> <li>Tegretol CR</li> </ul>
₭ Tab 400 mg		100	✓ Tegretol
K Tab long-acting 400 mg		100	<ul> <li>Tegretol CR</li> </ul>
🖌 Oral liq 20 mg per ml		250 ml	<ul> <li>Tegretol</li> </ul>
CLOBAZAM – Safety medicine; prescriber may determine di			·
Tab 10 mg		50	✓ Frisium
0		50	• Thistum
CLONAZEPAM – Safety medicine; prescriber may determine			
Oral drops 2.5 mg per ml		0 ml OP	<ul> <li>Rivotril</li> </ul>
THOSUXIMIDE			
Cap 250 mg		56	<ul> <li>Essential</li> </ul>
			Ethosuximide S29
	140.88	100	<ul> <li>Zarontin</li> </ul>
Oral lig 250 mg per 5 ml		200 ml	<ul> <li>Zarontin</li> </ul>
ABAPENTIN			
Note: Not subsidised in combination with subsidised pre	aahalin		
<ul> <li>Cap 100 mg</li> </ul>	0	100	<ul> <li>Nupentin</li> </ul>
Cap 100 mg		100	✓ Nupentin
k Cap 400 mg		100	✓ Nupentin
		100	
ACOSAMIDE – Special Authority see SA2267 on the next	<b>v</b> 1 7	4.4	/ Winnet
Tab 50 mg		14	✓ Vimpat
Tab 100 mg		14	✓ Vimpat
Tab 150 mg	200.24	56	✓ Vimpat
Tab 150 mg		14	✓ Vimpat
Tab 000 mm	300.40	56	✓ Vimpat
Tab 200 mg		56	<ul> <li>Vimpat</li> </ul>

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	~	Manufacturer

### ➡SA2267 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Patient has focal epilepsy; and
- 2 Seizures are not adequately controlled by, or patient has experienced unacceptable side effects from, optimal treatment with all of the following: sodium valproate, topiramate, levetiracetam and any two of carbamazepine, lamotrigine and phenytoin sodium (see Note).

Note: Those of childbearing potential are not required to trial phenytoin sodium, sodium valproate, or topiramate. Those who can father children are not required to trial sodium valproate.

**Renewal** from any relevant practitioner. Approvals valid for 24 months where the patient has demonstrated a significant and sustained improvement in seizure rate or severity and/or quality of life compared with that prior to starting lacosamide treatment.

LA	MOTRIGINE				
	Tab dispersible 2 mg		30	1	Lamictal
	Tab dispersible 5 mg		30	1	Lamictal
*	Tab dispersible 25 mg	4.20	56	1	Logem
*	Tab dispersible 50 mg	5.11	56	1	Logem
*	Tab dispersible 100 mg	6.75	56	1	Logem
١F	VETIRACETAM				
	Tab 250 mg		60	1	Everet
	Tab 500 mg		60		Everet
	Tab 750 mg		60		Everet
	Tab 1,000 mg		60		Everet
	Oral lig 100 mg per ml		300 ml OP	1	Levetiracetam-AFT
	Inj 100 mg per ml, 5 ml vial		10		Levetiracetam-AFT
PH	IENOBARBITONE				
	For phenobarbitone oral liquid refer Standard Formulae,	age 273			
	Tab 15 mg	•	500	1	Noumed
		2.0.00			Phenobarbitone
	Tab 30 mg	398 50	500	1	Noumed
			000	-	Phenobarbitone
БЦ	IENYTOIN SODIUM				<u></u>
*	Tab 50 mg	75.00	200	1	Dilantin Infatab
ጥ	Cap 30 mg		200		Dilantin
	Cap 100 mg		200		Dilantin
*	Oral liq 30 mg per 5 ml		500 ml		Dilantin Paediatric
			500 111	•	Ditantin'i aculatic
PH	EGABALIN	onontin			
*	Note: Not subsidised in combination with subsidised gab		56		Dreachalin Dfiner
*	Cap 25 mg		00		Pregabalin Pfizer
	0	7.80	50		Milpharm S29
*	Cap 75 mg		56		Pregabalin Pfizer
		8.10			Milpharm S29
*	Cap 150 mg	4.01	56		Lyrica
	o				Pregabalin Pfizer
*	Cap 300 mg	7.38	56	~	Pregabalin Pfizer
PF	IMIDONE				
*	Tab 250 mg		100	1	Primidone Clinect

	Subsidy (Manufacturer's Price)		Fully	
			Subsidised	
	\$	Per		Manufacturer
SODIUM VALPROATE				
Tab 100 mg		100	✓	Epilim Crushable
Tab 200 mg EC	27.44	100	✓	Epilim
Tab 500 mg EC		100	✓	Epilim
* Oral liq 200 mg per 5 ml		300 m	nl 🗸	Epilim S/F Liquid
			✓	Epilim Syrup
* Inj 100 mg per ml, 4 ml	41.50	1	✓	Epilim IV
STIRIPENTOL - Special Authority see SA2268 below - Retail ph	armacy			
Cap 250 mg	509.29	60	1	Diacomit
Powder for oral lig 250 mg sachet		60	1	Diacomit

### ➡SA2268 Special Authority for Subsidy

TOPIRAMATE

Initial application only from a paediatric neurologist or Practitioner on the recommendation of a paediatric neurologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient has confirmed diagnosis of Dravet syndrome; and
- 2 Seizures have been inadequately controlled by appropriate courses of sodium valproate, clobazam and at least two of the following: topiramate, levetiracetam, ketogenic diet.

Note: Those of childbearing potential are not required to trial sodium valproate or topiramate. Those who can father children are not required to trial sodium valproate.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where the patient continues to benefit from treatment as measured by reduced seizure frequency from baseline.

▲ Tab 25 mg	11.07	60	Arrow-Topiramate
C C			<ul> <li>Topiramate Actavis</li> </ul>
	26.04		<ul> <li>Topamax</li> </ul>
▲ Tab 50 mg	18.81	60	Arrow-Topiramate
Ĵ			<ul> <li>Topiramate Actavis</li> </ul>
	44.26		<ul> <li>Topamax</li> </ul>
▲ Tab 100 mg	31.99	60	<ul> <li>Arrow-Topiramate</li> </ul>
			<ul> <li>Topiramate Actavis</li> </ul>
	75.25		<ul> <li>Topamax</li> </ul>
▲ Tab 200 mg	55.19	60	<ul> <li>Arrow-Topiramate</li> </ul>
·			<ul> <li>Topiramate Actavis</li> </ul>
	129.85		<ul> <li>Topamax</li> </ul>
Sprinkle cap 15 mg	20.84	60	<ul> <li>Topamax</li> </ul>
▲ Sprinkle cap 25 mg		60	<ul> <li>Topamax</li> </ul>
VIGABATRIN - Special Authority see SA2088 below - Retail pharr			
Tab 500 mg	,	100	<ul> <li>Sabril</li> </ul>
<ul> <li>Powder for oral soln 500 mg per sachet</li> </ul>		60	✓ Sabril

### ► SA2088 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Any of the following:
  - 1.1 Patient has infantile spasms; or
  - 1.2 Both:
    - 1.2.1 Patient has epilepsy; and
    - 1.2.2 Either:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	sidised	Generic	
\$	Per	1	Manufacturer	

- 1.2.2.1 Seizures are not adequately controlled with optimal treatment with other antiepilepsy agents; or
- 1.2.2.2 Seizures are controlled adequately but the patient has experienced unacceptable side effects from optimal treatment with other antiepilepsy agents; or
- 1.3 Patient has tuberous sclerosis complex; and

2 Either:

- 2.1 Patient is, or will be, receiving regular automated visual field testing (ideally before starting therapy and on a 6-monthly basis thereafter); or
- 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields...

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient has demonstrated a significant and sustained improvement in seizure rate or severity and or quality of life; and 2 Either:
  - 2.1 Patient is receiving regular automated visual field testing (ideally every 6 months) on an ongoing basis for duration of treatment with vigabatrin; or
  - 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields...

## **Antimigraine Preparations**

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 115

### Acute Migraine Treatment

Principal Supply

RIZATRIPT Tab or	ΓAN odispersible 10 mg	4.84	30	✓ Rizamelt		
SUMATRIF						
	) mg		90	✓ <u>Sumagran</u>		
	00 mg ng per ml, 0.5 ml prefilled pen  – Maximum of 1		90	<ul> <li>Sumagran</li> </ul>		
	escription	••	2 OP	✓ <u>Clustran</u>		
Prophy	laxis of Migraine					
	drenoceptor Blockers refer to CARDIOVASCUI	_AR SYSTEM, page 47				
		00.01	100	. Condominuon		
	)0 mcg	23.21	100	<ul> <li>Sandomigran</li> </ul>		
Antinau	usea and Vertigo Agents					
or Antispa	asmodics refer to ALIMENTARY TRACT, page	8				
PREPITA	NT – Special Authority see SA0987 below – R	etail pharmacy				
Cap 2	$\times$ 80 mg and 1 $\times$ 125 mg	21.90	3 OP	Emend Tri-Pack		
	Special Authority for Subsidy					
	lication from any relevant practitioner. Approv					
•	c chemotherapy and/or anthracycline-based ch rom any relevant practitioner. Approvals valid f					
	apy and/or anthracycline-based chemotherapy			dergoing nighty entetogenic		
		for the deathorit of mang	lanoy.			
	3 mg		100	✓ Serc		
	5			<u></u>		
	✓ fully subsidised	S29 Unappro	ved medicine	e supplied under Section 29		
132	Principal Supply	Sole Subsidised Supply				

Sole Subsidised Supply

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
CYCLIZINE HYDROCHLORIDE Tab 50 mg	0.66	10	✓ <u>N</u>	lausicalm
CYCLIZINE LACTATE				
Inj 50 mg per ml, 1 ml ampoule – Up to 10 inj available on a PSO	16.36	10	✓ <u>⊦</u>	lameln
DOMPERIDONE				
* Tab 10 mg	4.00	100	✓ [	<u>Domperidone</u> <u>Viatris</u>
HYOSCINE HYDROBROMIDE				
Inj 400 mcg per ml, 1 ml ampoule Patch 1 mg per 72 hours – Special Authority see SA1998	93.00	10	🗸 N	Martindale S29
below – Retail pharmacy		10	<b>√</b> S	Scopolamine - Mylan

### ► SA1998 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Control of intractable nausea, vomiting, or inability to swallow saliva in the treatment of malignancy or chronic disease where the patient cannot tolerate or does not adequately respond to oral anti-nausea agents; or
- 2 Control of clozapine-induced hypersalivation where trials of at least two other alternative treatments have proven ineffective.

Renewal from any relevant practitioner. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

METOCLOPRAMIDE HYDROCHLORIDE		
* Tab 10 mg – Up to 30 tab available on a PSO1.	57 100	<ul> <li><u>Metoclopramide</u></li> <li><u>Actavis 10</u></li> </ul>
* Inj 5 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO7.	00 10	✓ Baxter
ONDANSETRON		
* Tab 4 mg2.	27 50	<ul> <li>Periset</li> </ul>
Tab disp 4 mg – Up to 10 tab available on a PSO0.		<ul> <li>Periset ODT</li> </ul>
* Tab 8 mg	10 50	✓ Periset
Tab disp 8 mg – Up to 10 tab available on a PSO0.	90 10	<ul> <li>Periset ODT</li> </ul>
PROCHLORPERAZINE		
* Tab 3 mg buccal5.	97 50	
(30.		Buccastem
(30.	00)	Max Health \$29
(30.	00)	Prochlorperazine
		Brown & Burk S29
(30.	00)	Prochlorperazine Max Health
* Tab 5 mg – Up to 30 tab available on a PSO25.	00 250	Nausafix
<ul> <li>Inj 12.5 mg per ml, 1 ml – Up to 5 inj available on a PSO25.</li> <li>(Buccastem Tab 3 mg buccal to be delisted 1 July 2025)</li> <li>(Max Health \$20 Tab 3 mg buccal to be delisted 1 July 2025)</li> </ul>		✓ Stemetil

(Prochlorperazine Brown & Burk S29 Tab 3 mg buccal to be delisted 1 July 2025)

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy	<b>`</b>	Fully	
	(Manufacturer's Price		Subsidised	
	\$	Per		Manufacturer
ntipsychotics				
ieneral				
IISULPRIDE – Safety medicine; prescriber may determine di				
Tab 100 mg		30		Sulprix
Tab 200 mg		60		Sulprix
Tab 400 mg		60	•	<u>Sulprix</u>
IPIPRAZOLE – Safety medicine; prescriber may determine c	dispensing frequency			
Tab 5 mg		30	✓	Aripiprazole Sandoz
			✓	Ascend
				Aripiprazole S29
Tab 10 mg		30	1	Aripiprazole Sandoz
Tab 15 mg		30		Aripiprazole Sandoz
Tab 20 mg		30		Aripiprazole Sandoz
Tab 30 mg		30		Aripiprazole Sandoz
scend Aripiprazole S29 Tab 5 mg to be delisted 1 July 2025)		2.		1-1
LORPROMAZINE HYDROCHLORIDE – Safety medicine; pi				
Tab 25 mg – Up to 30 tab available on a PSO		100		Largactil
Tab 100 mg – Up to 30 tab available on a PSO		100		Largactil
Inj 25 mg per ml, 2 ml – Up to 5 inj available on a PSO		10	~	Largactil
OZAPINE – Hospital pharmacy [HP4]				
Safety medicine; prescriber may determine dispensing frequ	Jency			
Tab 25 mg	6.69	50	✓	Clopine
			✓	Clozaril
	13.37	100	✓	Clopine
			✓	Clozaril
Tab 50 mg	8.67	50	✓	Clopine
Ũ	17.33	100	✓	Clopine
Tab 100 mg		50	✓	Clopine
5			✓	Clozaril
	34.65	100	✓	Clopine
				Clozaril
Tab 200 mg		50	✓	Clopine
č	69.30	100		Clopine
Suspension 50 mg per ml	147.30	100 m	l 🗸	Versacloz
LOPERIDOL – Safety medicine; prescriber may determine d				
Tab 500 mcg – Up to 30 tab available on a PSO		100	1	Serenace
Tab 1.5 mg – Up to 30 tab available on a PSO		100		Serenace
Tab 5 mg – Up to 30 tab available on a PSO		50		Serenace
1 ab 5 mg - 0p to 50 tab available off a F 50		100		Serenace
Oral lig 2 mg per ml – Up to 200 ml available on a PSO		100 m		Serenace
Inj 5 mg per ml, 1 ml ampoule – Up to 5 inj available on a PSO		100 m		Serenace
			•	Jerenaue
VOMEPROMAZINE – Safety medicine; prescriber may deter			_	
Tab 25 mg (33.8 mg as a maleate)		100		Nozinan (Swiss)
Tab 25 mg as a maleate		100		Nozinan
Tab 100 mg (135 mg as a maleate)	41.75	100		Nozinan (Swiss)
		100		Nozinan

			Fully	Brand or
	(Manufacturer's Pric \$	e) Sub Per	sidised	Generic Manufacturer
	•	-		
EVOMEPROMAZINE HYDROCHLORIDE – Safety medicine		10	-	
Inj 25 mg per ml, 1 ml ampoule	24.48	10	-	Nozinan S29 S29 Wockhardt
Nazinan \$20 520 Ini 25 ma par ml. 1 ml ampaula to bo dalici	ad 1 July 2025)		•	WOCKHAIUL
Nozinan S29 S29 Inj 25 mg per ml, 1 ml ampoule to be delist	• •			
ITHIUM CARBONATE – Safety medicine; prescriber may det				Duiodal
Tab long-acting 400 mg		100 100	-	Priadel Douglas
Cap 250 mg		100	•	Douglas
DLANZAPINE – Safety medicine; prescriber may determine di				<b>_</b> .
Tab 2.5 mg		30		Zypine
Tab 5 mg		30		Zypine
Tab orodispersible 5 mg		28		Zypine ODT
Tab 10 mg		30	-	Zypine Zypine ODT
Tab orodispersible 10 mg		28	v	Zypine ODT
PERICYAZINE – Safety medicine; prescriber may determine of				
Tab 2.5 mg	13.61	100		Neulactil
Tab 10 mg		100	~	Neulactil
QUETIAPINE - Safety medicine; prescriber may determine dis	spensing frequency			
Tab 25 mg	2.36	90	✓	Quetapel
Tab 100 mg	6.40	90	✓	Quetapel
Tab 200 mg		90	✓	Quetapel
Tab 300 mg	15.83	90	✓	Quetapel
RISPERIDONE - Safety medicine; prescriber may determine (	dispensing frequency			
Tab 0.5 mg		20	1	Risperdal
	2.17	60		Risperidone (Teva)
	4.01			Risperidone
				Sandoz S29
Tab 1 mg	2 44	60	1	Risperdal
100 T Hig	£	00		Risperidone (Teva)
	3.68		-	Risperidone
	0.00		-	Sandoz S29
Tab 2 mg	0.70	60	1	Risperdal
Tab 2 mg		00		Risperidone (Teva)
	5.38			Risperidone
	5.50		•	Sandoz S29
Tab 0 mg	4.50	60	./	
Tab 3 mg	4.50	60		Risperdal
	8.57			Risperidone (Teva)
	8.57		v	Risperidone
				Sandoz S29
Tab 4 mg	6.25	60		Risperdal
	40.00	00 I	-	Risperidone (Teva)
Oral liq 1 mg per ml		30 ml	•	Risperon
IPRASIDONE – Safety medicine; prescriber may determine of	dispensing frequency			
Cap 20 mg	17.90	60		Zusdone
Cap 40 mg		60		Zusdone
Cap 60 mg		60		Zusdone
Cap 80 mg		60	~	Zusdone

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Subsic Per	Fully lised	
Depot Injections				
ARIPIPRAZOLE – Special Authority see SA2395 below – Retail p Safety medicine; prescriber may determine dispensing frequen Inj 300 mg vial	ncy	1		Abilify Maintena Abilify Maintena S29 529
Inj 400 mg vial	341.96	1		Abilify Maintena Abilify Maintena S29 S29

## SA2395 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Fither:

- 1 Either:
  - 1.1 The patient has had an initial Special Authority approval for risperidone depot injection, paliperidone depot injection or olanzapine depot injection: or
  - 1.2 All of the following:
    - 1.2.1 The patient has schizophrenia or other psychotic disorder; and
    - 1.2.2 The patient has received treatment with oral atvoical antipsychotic agents but has been unable to adhere: and
    - 1.2.3 The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months; or
- 2 Patient has been unable to access olanzapine depot injection due to supply issues with olanzapine depot injection, or otherwise would have been started on olanzapine depot injection but has been unable to due to supply issues with olanzapine depot injection.

Notes: The Olanzapine depot injection Special Authority criteria that apply to criterion 2 in this Aripiprazole Special Authority application are as follows:

- The patient has had an initial Special Authority approval for paliperidone depot injection or risperidone depot injection; or
- All of the following:
  - The patient has schizophrenia; and
  - The patient has not been able to adhere with treatment using oral atvpical antipsychotic agents: and
  - The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months.

FLUPENTHIXOL DECANOATE - Safety medicine; prescriber may determine dispensing frequency

Inj 20 mg per ml, 1 ml – Up to 5 inj available on a PSO	5	<ul> <li>Fluanxol</li> </ul>
Inj 20 mg per ml, 2 ml – Up to 5 inj available on a PSO20.90	5	<ul> <li>Fluanxol</li> </ul>
Inj 100 mg per ml, 1 ml – Up to 5 inj available on a PSO40.87	5	<ul> <li>Fluanxol</li> </ul>

HALOPERIDOL DECANOATE - Safety medicine; prescriber may determine dispensing frequency Haldol

- Inj 50 mg per ml, 1 ml Up to 5 inj available on a PSO......28.39

OLANZAPINE - Special Authority see SA2313 on the next page - Retail pharmacy

a) Safety medicine; prescriber may determine dispensing frequency

<li>b) Note – no new patients to be initiated on olanzap</li>	pine.		
Inj 210 mg vial		1	<ul> <li>Zyprexa Relprevv</li> </ul>
Inj 300 mg vial		1	<ul> <li>Zyprexa Relprevv</li> </ul>
Inj 405 mg vial		1	<ul> <li>Zyprexa Relprevv</li> </ul>
, ,			••

5

5

✓ Haldol Concentrate

Decanoas S29

Haldol

	Subsidy		Fully	Brand or
(Ma	anufacturer's Price)	Subs	sidised	Generic
	\$	Per	✓	Manufacturer

### ⇒SA2313 Special Authority for Subsidy

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of olanzapine depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

PALIPERIDONE - Special Authority see SA2396 below - Retail pharmacy

Safety medicine; prescriber may determine dispensing f	requency		
Inj 25 mg syringe		1	Invega Sustenna
Inj 50 mg syringe		1	Invega Sustenna
Inj 75 mg syringe		1	Invega Sustenna
Inj 100 mg syringe		1	Invega Sustenna
Inj 150 mg syringe	435.12	1	<ul> <li>Invega Sustenna</li> </ul>

### ⇒SA2396 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for risperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
  - 2.1 The patient has schizophrenia or other psychotic disorder; and
  - 2.2 Has been unable to adhere to treatment using oral atypical antipsychotic agents; and
  - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

**Renewal** from any relevant practitioner. Approvals valid for 12 months where the initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

PALIPERIDONE PALMITATE - Special Aut	thority see <mark>SA2167 below</mark> – Retail pharm	acy	
Inj 175 mg syringe		1	🗸 Invega Trinza
Inj 263 mg syringe	1,072.26	1	Invega Trinza
Inj 350 mg syringe		1	🗸 Invega Trinza
Inj 525 mg syringe		1	🗸 Invega Trinza

#### ⇒SA2167 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has schizophrenia; and
- 2 The patient has had an initial Special Authority approval for paliperidone once-monthly depot injection.

**Renewal** from any relevant practitioner. Approvals valid for 12 months where the initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

RISPERIDONE – Special Authority see SA2397 below – Retail pharmacy

Safety medicine; prescriber may determine dispensing frequency

Inj 25 mg vial	. 135.98	1	Risperdal Consta
Inj 37.5 mg vial	178.71	1	Risperdal Consta
Inj 50 mg vial		1	<ul> <li>Risperdal Consta</li> </ul>

#### ➡SA2397 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

1 The patient has had an initial Special Authority approval for paliperidone depot injection or olanzapine depot injection; or

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sub	sidised	Generic
\$	Per	1	Manufacturer

- 2 All of the following:
  - 2.1 The patient has schizophrenia or other psychotic disorder; and
  - 2.2 Has not been able to adhere with treatment using oral atypical antipsychotic agents; and
  - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

**Renewal** from any relevant practitioner. Approvals valid for 12 months where the initiation of risperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

ZUCLOPENTHIXOL DECANOATE - Safety medicine; prescriber may determine dispensing frequency

Inj 200 mg per ml, 1 ml – Up to 5 inj available on a PSO..... 19.80 5 Clopixol

Anxiolytics		
BUSPIRONE HYDROCHLORIDE * Tab 5 mg * Tab 10 mg	100 100	<ul> <li>✓ <u>Buspirone Viatris</u></li> <li>✓ <u>Buspirone Viatris</u></li> </ul>
CLONAZEPAM – Safety medicine; prescriber may determine Tab 500 mcg Tab 2 mg	 100 100	<ul><li>✓ Paxam</li><li>✓ Paxam</li></ul>
DIAZEPAM – Safety medicine; prescriber may determine dis Tab 2 mg Tab 5 mg	 500 500	<ul> <li>✓ <u>Arrow-Diazepam</u></li> <li>✓ <u>Arrow-Diazepam</u></li> </ul>
LORAZEPAM – Safety medicine; prescriber may determine of Tab 1 mg Tab 2.5 mg	 250 100	✓ <u>Ativan</u> ✓ <u>Ativan</u>

## **Multiple Sclerosis Treatments**

### ⇒SA2274 Special Authority for Subsidy

Initial application — (Multiple Sclerosis - dimethyl fumarate, fingolimod, glatiramer acetate, interferon beta-1-alpha, interferon beta-1-beta, natalizumab and teriflunomide) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 All of the following:
  - 1.1 Diagnosis of multiple sclerosis (MS) meets the McDonald 2017 diagnostic criteria for MS and has been confirmed by a neurologist; and
  - 1.2 Patient has an EDSS score between 0 6.0; and
  - 1.3 Patient has had at least one significant attack of MS in the previous 12 months or two significant attacks in the past 24 months; and
  - 1.4 All of the following:
    - 1.4.1 Each significant attack must be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the attack, but the neurologist/physician must be satisfied that the clinical features were characteristic); and
    - 1.4.2 Each significant attack is associated with characteristic new symptom(s)/sign(s) or substantially worsening of previously experienced symptoms(s)/sign(s); and
    - 1.4.3 Each significant attack has lasted at least one week and has started at least one month after the onset of a

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

previous attack (where relevant); and

- 1.4.4 Each significant attack can be distinguished from the effects of general fatigue; and is not associated with a fever (T> 37.5°C); and
- 1.4.5 Either:
  - 1.4.5.1 Each significant attack is severe enough to change either the EDSS or at least one of the Kurtze Functional System scores by at least 1 point; or
  - 1.4.5.2 Each significant attack is a recurrent paroxysmal symptom of multiple sclerosis (tonic seizures/spasms, trigeminal neuralgia, Lhermitte's symptom); and
- 1.5 Evidence of new inflammatory activity on an MRI scan within the past 24 months; and
- 1.6 Any of the following:
  - 1.6.1 A sign of that new inflammatory activity on MRI scanning (in criterion 5 immediately above) is a gadolinium enhancing lesion; or
  - 1.6.2 A sign of that new inflammatory activity is a lesion showing diffusion restriction; or
  - 1.6.3 A sign of that new inflammatory is a T2 lesion with associated local swelling; or
  - 1.6.4 A sign of that new inflammatory activity is a prominent T2 lesion that clearly is responsible for the clinical features of a recent attack that occurred within the last 2 years; or
  - 1.6.5 A sign of that new inflammatory activity is new T2 lesions compared with a previous MRI scan; or
- 2 Patient has an active approval for ocrelizumab and does not have primary progressive MS.
- Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

Renewal — (Multiple Sclerosis - dimethyl fumarate, fingolimod, glatiramer acetate, interferon beta-1-alpha, interferon beta-1-beta, natalizumab and teriflunomide) from any relevant practitioner. Approvals valid for 12 months where patient has had an EDSS score of 0 to 6.0 (inclusive) with or without the use of unilateral or bilateral aids at any time in the last six months (ie the patient has walked 100 metres or more with or without aids in the last six months).

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

DIMETHYL FUMARATE – Special Authority see SA2274 on the previous page – Retail pharmacy

a) Wastage claimable

<ul> <li>b) Note: Treatment on two or more funded multiple scler Cap 120 mg</li></ul>	520.00 2,000.00	14 56	t permitted. ✓ Tecfidera ✓ Tecfidera
FINGOLIMOD – Special Authority see SA2274 on the previou a) Wastage claimable			
<li>b) Note: Treatment on two or more funded multiple scler Cap 0.5 mg</li>			t permitted.  Gilenya
GLATIRAMER ACETATE – Special Authority see SA2274 on Note: Treatment on two or more funded multiple sclerosis Inj 40 mg prefilled syringe	s treatments simultaneou	usly is not pe	,
INTERFERON BETA-1-ALPHA – Special Authority see SA22 Note: Treatment on two or more funded multiple sclerosis Inj 6 million iu prefilled syringe	s treatments simultaneou	usly is not pe	
Injection 6 million iu per 0.5 ml pen injector	1,170.00		<ul> <li>Avonex Pen</li> </ul>
INTERFERON BETA-1-BETA – Special Authority see SA227 Note: Treatment on two or more funded multiple sclerosis Inj 8 million iu per 1 ml	s treatments simultaneou	usly is not pe	
NATALIZUMAB – Special Authority see SA2274 on the previo Note: Treatment on two or more funded multiple sclerosis Inj 20 mg per ml, 15 ml vial	s treatments simultaneou		rmitted. ✔ Tysabri

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

		Subsidy (Manufacturer's Price) \$	Full Subsidise Per •	5
TERIFLUNOMIDE	- Special Authority see SA2274 on page	138 – Retail pharmacy		
a) Wastage cla				
, 0	ment on two or more funded multiple scl	erosis treatments simultan	eouslv is not r	permitted.
,				Teriflunomide
0				Sandoz
		659.90	✓	' Aubagio
Teriflunomi	de Sandoz to be Principal Supply on 1 A	pril 2025		-
(Aubagio Tab 14 mg	g to be delisted 1 April 2025)			
Multiple Scler	osis Treatments - Other			
OCRELIZUMAB -	Special Authority see SA2273 below – R	etail pharmacy		
	nt on two or more funded multiple scleros		slv is not pern	nitted.
	l, 10 ml vial			Ocrevus
	Authority for Subsidy			
nitial application -	– (Multiple Sclerosis - ocrelizumab) fi	rom any relevant practition	er. Approvals	valid for 12 months for
	the following criteria:		on reproteit	
Either:	<b>3</b> • • • •			
1 All of the foll	owing:			
1.1 Diagr	osis of multiple sclerosis (MS) meets the	e McDonald 2017 diagnost	ic criteria for N	IS and has been confirmed
by a l	neurologist; and			
1.2 Patie	nt has an EDSS score between 0 - 6.0; a	and		
1.3 Patie	nt has had at least one significant attack	of MS in the previous 12 n	nonths or two	significant attacks in the pas
	onths; and			
	the following:			
1.4.1	Each significant attack must be confirm			
	not necessarily have been seen by the	•	e neurologist/	physician must be satisfied
1.4.0	that the clinical features were characte	<i>,,</i>	atam(a)/aian/a	
1.4.2	<ul> <li>Each significant attack is associated w of previously experienced symptoms(s)</li> </ul>	, ,	prom(s)/sign(s	) or substantially worsening
149	Each significant attack has lasted at le		tod at loast on	a month after the onset of a
1.4.0	previous attack (where relevant); and	asi one week and has siar		
144	Each significant attack can be distingu	ished from the effects of a	eneral fatique:	and is not associated with a
	fever (T> 37.5°C); and		onora rangao,	
1.4.5	Either:			
	1.4.5.1 Each significant attack is severe	enough to change either t	he EDSS or a	t least one of the Kurtze
	Functional System scores by at	least 1 point; or		
	1.4.5.2 Each significant attack is a recu			lerosis (tonic
	seizures/spasms, trigeminal neu	iralgia, Lhermitte's symptoi	m); and	
1.5 Evide	nce of new inflammatory activity on an N	IRI scan within the past 24	months; and	
1.6 Any o	f the following:			
1.6.1	A sign of that new inflammatory activity	on MRI scanning (in crite	rion 5 immedia	ately above) is a gadolinium
	enhancing lesion; or			
	A sign of that new inflammatory activity			
	A sign of that new inflammatory is a T2		0,	
1.6.4	A sign of that new inflammatory activity			esponsible for the clinical
1.6.6	features of a recent attack that occurre			viewe MPL coopy or
1.6.5	A sign of that new inflammatory activity	is new 12 lesions compai	ieu with a prev	nous with scan; or
0 Detternt	in active Special Authority approval for e	the end alternation of ferror ends in the	الأراب الممطامين	and a second at the second and

Sut	ıbsidy	Fully	Brand or	
(Manufacti	turer's Price) Sub	sidised	Generic	
	\$ Per	1	Manufacturer	

beta-1-alpha, interferon beta-1-beta, natalizumab or teriflunomide.

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

Renewal — (Multiple Sclerosis - ocrelizumab) from any relevant practitioner. Approvals valid for 12 months where patient has had an EDSS score of 0 to 6.0 (inclusive) with or without the use of unilateral or bilateral aids at any time in the last six months (ie the patient has walked 100 metres or more with or without aids in the last six months).

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

Initial application — (Primary Progressive Multiple Sclerosis) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Diagnosis of primary progressive multiple sclerosis (PPMS) meets the 2017 McDonald criteria and has been confirmed by a neurologist; and
- 2 Patient has an EDSS 2.0 (score equal to or greater than 2 on pyramidal functions) to EDSS 6.5; and
- 3 Patient has no history of relapsing remitting multiple sclerosis.

Renewal — (Primary Progressive Multiple Sclerosis) from any relevant practitioner. Approvals valid for 12 months where patient has had an EDSS score of less than or equal to 6.5 at any time in the last six months (ie patient has walked 20 metres with bilateral assistance/aids, without rest in the last six months).

## Sedatives and Hypnotics

MELATONIN – Special Authority see SA1666 below – Retail pharmacy

✓ Viaisom

30

### ⇒SA1666 Special Authority for Subsidy

**Initial application** only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with persistent and distressing insomnia secondary to a neurodevelopmental disorder (including, but not limited to, autism spectrum disorder or attention deficit hyperactivity disorder)*; and
- 2 Behavioural and environmental approaches have been tried and were unsuccessful, or are inappropriate; and
- 3 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day; and
- 4 Patient is aged 18 years or under*.

**Renewal** only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is aged 18 years or under*; and
- 2 Patient has demonstrated clinically meaningful benefit from funded modified-release melatonin (clinician determined); and
- 3 Patient has had a trial of funded modified-release melatonin discontinuation within the past 12 months and has had a recurrence of persistent and distressing insomnia; and
- 4 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day.

Note: Indications marked with * are unapproved indications.

	Subsidy		Fully	
(	Manufacturer's Price) \$	Per	Subsidised	
	*	1.61	•	Manulacturei
AIDAZOLAM – Safety medicine; prescriber may determine disper laid may negative for the manual set.		10		Midanalam Davidan
Inj 1 mg per ml, 5 ml ampoule		10		Midazolam-Baxter
	16.75		•	Midazolam Viatris
Inj 1 mg per ml, 5 ml plastic ampoule – Up to 10 inj available				
on a PSO		10		Pfizer
On a PSO for status epilepticus use only. PSO must be e	endorsed for status e	epilep	oticus use	only.
Inj 5 mg per ml, 1 ml plastic ampoule – Up to 10 inj available				
on a PSO		10		Midazolam-Pfizer
On a PSO for status epilepticus use only. PSO must be e		epilep		,
Inj 5 mg per ml, 3 ml ampoule	4.75	5		Midazolam-Baxter
	5.50		✓	Midazolam Viatris
Inj 5 mg per ml, 3 ml plastic ampoule – Up to 5 inj available o	n			
a PSO		5	✓	Pfizer
On a PSO for status epilepticus use only. PSO must be e	ndorsed for status e	epilep	ticus use	only.
Midazolam Viatris Inj 1 mg per ml, 5 ml ampoule to be delisted 1 l	Mav 2025)			
Midazolam Viatris Inj 5 mg per ml, 3 ml ampoule to be delisted 1	• •			
HENOBARBITONE SODIUM - Special Authority see SA1386 be	• •	2011		
· · · · ·		•		Marcella alle and
Inj 200 mg per ml, 1 ml ampoule SA1386 Special Authority for Subsidy	113.37	10	•	Max Health S29
he following criteria: Both: 1 For the treatment of terminal agitation that is unresponsive 2 The applicant is part of a multidisciplinary team working in p	•	ł		
EMAZEPAM – Safety medicine; prescriber may determine dispe		05		Manualaan
Tab 10 mg		25	•	Normison
OPICLONE – Safety medicine; prescriber may determine dispen Tab 7.5 mg	• • •	500	1	Zopiclone Actavis
Spinal Muscular Atrophy				
USINERSEN – PCT only – Special Authority see SA2174 below				
Inj 12 mg per 5 ml vial	120,000.00	1	1	Spinraza
SA2174 Special Authority for Subsidy				
nitial application — (spinal muscular atrophy (SMA)) from any	relevant practition	er. A	oprovals v	alid for 12 months for
pplications meeting the following criteria:	· · · · · · · · · · · · · · · · · · ·			
Il of the following:				
1 Patient has genetic documentation of homozygous SMN1 g	ene deletion, homo	zvqo	us SMN1	point mutation, or compour
heterozygous mutation; and	· ·			· · ·
2 Patient is 18 years of age or under; and				
3 Either:				
3.1 Patient has experienced the defined signs and symp	ntoms of SMA type I	llor	Illa prior	to three years of age: or
3.2 Both:		, 0	ina prior	to anot yours of ago, of
3.2.1 Patient is pre-symptomatic: and				

- 3.2.1 Patient is pre-symptomatic; and
- 3.2.2 Patient has three or less copies of SMN2.

Renewal — (spinal muscular atrophy (SMA)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Subsidy	I	Fully	Brand or	
(Manufacturer's Price)	Subsid	lised	Generic	
\$	Per	✓	Manufacturer	

All of the following:

- 1 There has been demonstrated maintenance of motor milestone function since treatment initiation; and
- 2 Patient does not require invasive permanent ventilation (at least 16 hours per day) in the absence of a potentially reversible cause while being treated with nusinersen; and
- 3 Nusinersen not to be administered in combination other SMA disease modifying treatments or gene therapy.

#### RISDIPLAM - [Xpharm] - Special Authority see SA2203 below

Note: the supply of risdiplam is via Pharmac's approved direct distribution supply. Further details can be found on Pharmac's website https://pharmac.govt.nz/risdiplam

Powder for oral soln 750 mcg per ml, 60 mg per bottle......14,100.00 80 ml OP 🖌 Evrysdi

### ⇒SA2203 Special Authority for Subsidy

**Initial application — (spinal muscular atrophy (SMA))** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has genetic documentation of homozygous SMN1 gene deletion, homozygous SMN1 point mutation, or compound heterozygous mutation; and
- 2 Patient is 18 years of age or under; and
- 3 Either:
  - 3.1 Patient has experienced the defined signs and symptoms of SMA type I, II or IIIa prior to three years of age; or
  - 3.2 Both:
    - 3.2.1 Patient is pre-symptomatic; and
    - 3.2.2 Patient has three or less copies of SMN2.

Renewal — (spinal muscular atrophy (SMA)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 There has been demonstrated maintenance of motor milestone function since treatment initiation; and
- 2 Patient does not require invasive permanent ventilation (at least 16 hours per day) in the absence of a potentially reversible cause while being treated with risdiplam; and
- 3 Risdiplam not to be administered in combination other SMA disease modifying treatments or gene therapy.

## Stimulants/ADHD Treatments

#### ATOMOXETINE

Cap 10 mg		28	APO-Atomoxetine
Cap 18 mg		28	✓ APO-Atomoxetine
Cap 25 mg		28	✓ APO-Atomoxetine
Cap 40 mg		28	✓ APO-Atomoxetine
Cap 60 mg		28	<ul> <li>APO-Atomoxetine</li> </ul>
Cap 80 mg		28	✓ APO-Atomoxetine
Cap 100 mg	65.71	28	✓ APO-Atomoxetine
DEXAMEETAMINE SI II EATE - Special Authority	see SA2410 on the payt page -	Rotail nhar	macy

DEXAMFETAMINE SULFATE - Special Authority see SA2410 on the next page - Retail pharmacy

a) Only on a controlled drug form

<ul> <li>Safety medicine; prescriber may determine dispensing free</li> </ul>	quency		
Tab 5 mg	29.80	100	Noumed

Dexamfetamine

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

### ⇒SA2410 Special Authority for Subsidy

Initial application — (ADHD in patients aged 5 years or over) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid without further renewal unless notified for applications meeting the following criteria: All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) in patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
  - 3.1 Applicant is a paediatrician or psychiatrist; or
  - 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients aged under 5 years) only from a paediatrician or psychiatrist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) in patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

**Initial application** — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid without further renewal unless notified where the patient suffers from narcolepsy.

LISDEXAMFETAMINE DIMESILATE - Special Authority see SA2415 below - Retail pharmacy

a) Only on a controlled drug form

b) Safety medicine; prescriber may determine dispensing free	quency		
Cap 30 mg - No more than 1 cap per day		30	<ul> <li>Vyvanse</li> </ul>
Cap 50 mg		30	✓ Vyvanse
Cap 70 mg	60.00	30	<ul> <li>Vyvanse</li> </ul>

### ⇒SA2415 Special Authority for Subsidy

Initial application only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 Patient is currently on treatment with lisdexamfetamine dimesilate and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
  - 2.1 ADHD (Attention Deficit and Hyperactivity Disorder); and
  - 2.2 Diagnosed according to DSM-V or ICD 11 criteria; and
  - 2.3 Either:
    - 2.3.1 Applicant is a paediatrician or psychiatrist; or
    - 2.3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing; and
  - 2.4 Any of the following:
    - 2.4.1 Patient is taking a currently subsidised formulation of atomoxetine or methylphenidate hydrochloride (extended-release) and has not received sufficient benefit or has experienced intolerable side effects; or
    - 2.4.2 Patient is taking a currently subsidised formulation of dexamfetamine sulfate (immediate-release) which has not been effective due to significant administration and/or treatment adherence difficulties; or
    - 2.4.3 There is significant concern regarding the risk of diversion or abuse of immediate release dexamfetamine sulfate; or
    - 2.4.4 Patient is taking a currently subsidised formulation of methylphenidate hydrochloride (immediate-release or sustained release) which has not been effective due to significant administration and/or treatment

Subsidy (Manufacturer's Pri \$	ice) Subs Per	Fully sidised	Brand or Generic Manufacturer
continued			
adherence difficulties; or			
2.4.5 There is significant concern regarding the risk of diversion or	abuse of imm	ediate i	release methylphenidate
hydrochloride; or			
2.4.6 Both:			
2.4.6.1 Patient would have been prescribed a subsidised form (extended-release) but has been unable to access due hydrochloride (extended-release); and	e to supply iss	ues with	h methylphenidate
2.4.6.2 Other alternative stimulant presentations (methylpheni and	date or dexan	ıfetamir	ne) are not appropriate;
2.5 Lisdexamfetamine dimesilate is not to be used in combination with a	nother funded	I methy	Iphenidate presentation.
METHYLPHENIDATE HYDROCHLORIDE - Special Authority see SA2411 below	– Retail phar	macv	
a) Only on a controlled drug form	i totali pitali	indeg	
b) Safety medicine; prescriber may determine dispensing frequency			
Tab immediate-release 5 mg	30	✓ R	lubifen
Tab immediate-release 10 mg	30	🗸 R	ubifen
4.00		🗸 R	litalin
Tab extended-release 18 mg7.75	30	✓ M	lethylphenidate ER - Teva
Tab immediate-release 20 mg7.85	30	🗸 R	lubifen
Tab sustained-release 20 mg10.95	30	🗸 R	ubifen SR
Tab extended-release 27 mg11.45	30	✓ M	lethylphenidate ER - Teva
Tab extended-release 36 mg15.50	30	✓ M	lethylphenidate ER - Teva
Tab extended-release 54 mg22.25	30	✓ M	lethylphenidate ER - Teva

### ⇒SA2411 Special Authority for Subsidy

**Initial application** — (ADHD in patients aged 5 years or over) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid without further renewal unless notified for applications meeting the following criteria: All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) in patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
  - 3.1 Applicant is a paediatrician or psychiatrist; or
  - 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients aged under 5 years) only from a paediatrician or psychiatrist. Approvals valid without further renewal unless notified for applications meeting the following criteria: Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) in patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

**Initial application** — (Narcolepsy*) only from a neurologist or respiratory specialist. Approvals valid without further renewal unless notified where the patient suffers from narcolepsy.

Note: *narcolepsy is not a registered indication for Methylphenidate ER - Teva.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
METHYLPHENIDATE HYDROCHLORIDE EXTENDED-RELEAS	SE - Special Authority	/ see	SA2446 b	elow – Retail pharmacy
a) Only on a controlled drug form				
b) Safety medicine; prescriber may determine dispensing fro	equency			
Tab extended-release 18 mg		30	✓	Concerta
Tab extended-release 27 mg	65.44	30	✓	Concerta
Tab extended-release 36 mg	71.93	30	✓	Concerta
Tab extended-release 54 mg		30	✓	Concerta
Cap modified-release 10 mg		30	✓	Ritalin LA
Cap modified-release 20 mg	27.72	30	✓	Ritalin LA
Cap modified-release 30 mg		30	✓	Ritalin LA
Cap modified-release 40 mg		30	✓	Ritalin LA

### ► SA2446 Special Authority for Subsidy

**Initial application** — (ADHD) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid without further renewal unless notified for applications meeting the following criteria: Either:

- 1 All of the following:
  - 1.1 ADHD (Attention Deficit and Hyperactivity Disorder); and
  - 1.2 Diagnosed according to DSM-IV or ICD 10 criteria; and
  - 1.3 Either:
    - 1.3.1 Applicant is a paediatrician or psychiatrist; or
    - 1.3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing; and
  - 1.4 Either:
    - 1.4.1 Patient is taking a currently subsidised formulation of methylphenidate hydrochloride (immediate-release or sustained-release) which has not been effective due to significant administration and/or difficulties with adherence; or
    - 1.4.2 There is significant concern regarding the risk of diversion or abuse of immediate-release methylphenidate hydrochloride; or
- 2 Both:
  - 2.1 Patient meets the Special Authority criteria for SA2411 methylphenidate hydrochloride; and
  - 2.2 Patient is unable to access other methylphenidate hydrochloride presentations under Special Authority criteria SA2411 due to an out of stock (see note).

Note: Criterion 2 is to permit short-term funding to cover an out-of-stock on tab extended-release Methylphenidate ER – Teva and tab sustained-release 20 mg Rubifen SR subsidised under

SA2411 (https://schedule.pharmac.govt.nz/2025/02/01/SA2411.pdf).

#### MODAFINIL - Special Authority see SA2413 below - Retail pharmacy

Tab 100 mg		30	<ul> <li>Modafinil Max Health</li> </ul>
Modafinil Max Health to be Principal Supply on 1 Ma	29.13 y 2025	60	<ul> <li>Modavigil</li> </ul>

(Modavigil Tab 100 mg to be delisted 1 May 2025)

### ► SA2413 Special Authority for Subsidy

Initial application only from a neurologist or respiratory specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

1 The patient has a diagnosis of narcolepsy and has excessive daytime sleepiness associated with narcolepsy occurring

Subsi	idy Fully	Brand or
(Manufacture	er's Price) Subsidised	I Generic
\$	Per 🗸	Manufacturer

continued...

almost daily for three months or more; and

- 2 Either:
  - 2.1 The patient has a multiple sleep latency test with a mean sleep latency of less than or equal to 10 minutes and 2 or more sleep onset rapid eye movement periods; or
  - 2.2 The patient has at least one of: cataplexy, sleep paralysis or hypnagogic hallucinations; and
- 3 Either:
  - 3.1 An effective dose of a subsidised formulation of methylphenidate or dexamfetamine has been trialled and discontinued because of intolerable side effects; or
  - 3.2 Methylphenidate and dexamfetamine are contraindicated.

### **Treatments for Dementia**

DONEPEZIL HYDROCHLORIDE			
* Tab 5 mg		84	Ipca-Donepezil
* Tab 10 mg	5.50	84	Ipca-Donepezil
RIVASTIGMINE - Special Authority see SA1488 below - Re	etail pharmacy		
Patch 4.6 mg per 24 hour		30	Rivastigmine Patch
			BNM 5
	90.00		Exelon Patch 5
Patch 9.5 mg per 24 hour		30	Rivastigmine Patch
			BNM 10
	90.00		Exelon Patch 10
(Exelon Patch 5 Patch 4.6 mg per 24 hour to be delisted 1 Ju	ıne 2025)		

(Exelon Patch 10 Patch 9.5 mg per 24 hour to be delisted 1 June 2025)

#### ⇒SA1488 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The patient has been diagnosed with dementia; and
- 2 The patient has experienced intolerable nausea and/or vomiting from donepezil tablets.

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

1 The treatment remains appropriate; and

2 The patient has demonstrated a significant and sustained benefit from treatment.

### **Treatments for Substance Dependence**

BUPRENORPHINE WITH NALOXONE – Special Authority see SA1203 below – Retail pharmacy

<ul> <li>a) No patient co-payment payable</li> <li>b) Safety medicine; prescriber may determine dispensing frequency</li> </ul>		
Tab sublingual 2 mg with naloxone 0.5 mg	28	✓ <u>Buprenorphine</u> Naloxone BNM
Tab sublingual 8 mg with naloxone 2 mg	28	✓ <u>Buprenorphine</u> Naloxone BNM

### ⇒SA1203 Special Authority for Subsidy

**Initial application** (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Subsidy	Ful		
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per 🖌	Manufacturer	

#### continued...

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health..

Initial application — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient will not be receiving methadone; and
- 3 Patient is currently enrolled in an opioid substitution treatment program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient has previously trialled but failed detoxification with buprenorphine with naloxone with relapse back to opioid use and another attempt is planned; and
- 3 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is or has been receiving maintenance therapy with buprenorphine with naloxone (and is not receiving methadone); and
- 2 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health or is a medical practitioner authorised by the service to manage treatment in this patient.

Renewal — (Maintenance treatment where the patient has previously had an initial application for detoxification) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Patient received but failed detoxification with buprenorphine with naloxone; and
- 2 Maintenance therapy with buprenorphine with naloxone is planned (and patient will not be receiving methadone); and
- 3 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

### BUPROPION HYDROCHLORIDE

Tab modified-release 150 mg		30	✓ Zyban
DISULFIRAM Tab 200 mg	236 40	100	✓ Antabuse
NALTREXONE HYDROCHLORIDE – Special Authority see S			pharmacy
Tab 50 mg	77.77	28	<ul> <li>Naltrexone AOP \$29</li> </ul>
	83.33	30	✓ Naltraccord
	102.60		<ul> <li>Naltrexone Max Health \$29</li> </ul>
	138.88	50	Revia S29

(Revia \$29) Tab 50 mg to be delisted 1 July 2025)

Subsidy	- Fully	Brand or
(Manufacturer's	s Price) Subsidised	Generic
\$	Per 🗸	Manufacturer

#### ⇒SA1408 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient is currently enrolled in a recognised comprehensive treatment programme for alcohol dependence; and
- 2 Applicant works in or with a community Alcohol and Drug Service contracted to Health NZ or accredited against the New Zealand Alcohol and Other Drug Sector Standard or the National Mental Health Sector Standard.

**Renewal** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Compliance with the medication (prescriber determined); and
- 2 Any of the following:
  - 2.1 Patient is still unstable and requires further treatment; or
  - 2.2 Patient achieved significant improvement but requires further treatment; or
  - 2.3 Patient is well controlled but requires maintenance therapy.

#### NICOTINE

a) Nicotine will not be funded in amounts less than 4 weeks of treatment.

b) Note: Direct Provision by a pharmacist permitted under the provisions	in Part I of Sect	ion A.
Patch 7 mg - Up to 28 patch available on a PSO 19.62	28	<ul> <li>Habitrol</li> </ul>
Patch 14 mg – Up to 28 patch available on a PSO21.57	28	<ul> <li>Habitrol</li> </ul>
Patch 14 mg for direct distribution only - [Xpharm]12.49	7	<ul> <li>Habitrol</li> </ul>
Patch 21 mg – Up to 28 patch available on a PSO24.72	28	<ul> <li>Habitrol</li> </ul>
Patch 21 mg for direct distribution only - [Xpharm]13.19	7	<ul> <li>Habitrol</li> </ul>
Lozenge 1 mg – Up to 216 loz available on a PSO22.53	216	<ul> <li>Habitrol</li> </ul>
Lozenge 1 mg for direct distribution only - [Xpharm] 12.89	36	<ul> <li>Habitrol</li> </ul>
Lozenge 2 mg – Up to 216 loz available on a PSO24.68	216	<ul> <li>Habitrol</li> </ul>
Lozenge 2 mg for direct distribution only - [Xpharm] 13.25	36	<ul> <li>Habitrol</li> </ul>
Gum 2 mg (Fruit) – Up to 204 piece available on a PSO23.02	204	<ul> <li>Habitrol</li> </ul>
Gum 2 mg (Fruit) for direct distribution only - [Xpharm]17.57	96	<ul> <li>Habitrol</li> </ul>
Gum 2 mg (Mint) – Up to 204 piece available on a PSO23.02	204	<ul> <li>Habitrol</li> </ul>
Gum 2 mg (Mint) for direct distribution only - [Xpharm] 17.57	96	<ul> <li>Habitrol</li> </ul>
Gum 4 mg (Fruit) – Up to 204 piece available on a PSO25.98	204	<ul> <li>Habitrol</li> </ul>
Gum 4 mg (Fruit) for direct distribution only - [Xpharm]23.87		<ul> <li>Habitrol</li> </ul>
Gum 4 mg (Mint) – Up to 204 piece available on a PSO25.98	204	<ul> <li>Habitrol</li> </ul>
Gum 4 mg (Mint) for direct distribution only - [Xpharm]23.87	96	<ul> <li>Habitrol</li> </ul>

VARENICLINE TARTRATE - Special Authority see SA1845 below - Retail pharmacy

- a) A maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval, including the starter pack
- b) Varenicline will not be funded in amounts less than 4 weeks of treatment.

c) The 6-month time period in which a patient can receive a funded 12-week course of varenicline tartrate starts from the date the Special Authority is approved.

Tab 0.5 mg × 11 and 1 mg × 42	16.67	53 OP	Varenicline Pfizer
Tab 1 mg		56	<ul> <li>Varenicline Pfizer</li> </ul>

#### ⇒SA1845 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme,

	Subsidy	Fully	Brand or
(Man	ufacturer's Price)	Subsidised	Generic
	\$ I	Per 🖌	Manufacturer

continued...

which includes prescriber or nurse monitoring; and

- 3 Either:
  - 3.1 The patient has tried but failed to quit smoking after at least two separate trials of nicotine replacement therapy, at least one of which included the patient receiving comprehensive advice on the optimal use of nicotine replacement therapy; or
  - 3.2 The patient has tried but failed to quit smoking using bupropion or nortriptyline; and
- 4 The patient has not had a Special Authority for varenicline approved in the last 6 months; and
- 5 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 6 The patient is not pregnant; and
- 7 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

**Renewal** from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 It has been 6 months since the patient's previous Special Authority was approved; and
- 4 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 5 The patient is not pregnant; and
- 6 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

The patient must not have had an approval in the past 6 months.

Notes: a maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval.

This includes the 4-week 'starter' pack.

	Subsidy	Cub	Fully	Brand or
	(Manufacturer's Price) \$	Per	sidised ✓	Generic Manufacturer
Chemotherapeutic Agents				
Alkylating Agents				
BENDAMUSTINE HYDROCHLORIDE - PCT only - Specialist -	- Special Authority se	e SA2398	below	
Inj 25 mg vial		1	✔ В	endamustine Sandoz
	77.00		🗸 R	ibomustin
Inj 100 mg vial	200.20	1	-	endamustine Sandoz
	308.00		🗸 R	ibomustin
Inj 1 mg for ECP		1 mg	🗸 В	axter

#### ⇒SA2398 Special Authority for Subsidy

**Initial application** — (CLL*) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has chronic lymphocytic leukaemia requiring treatment; and
- 2 Patient has ECOG performance status of 0-2; and
- 3 Bendamustine is to be administered at a maximum dose of 100 mg/m² on days 1 and 2 every 4 weeks for a maximum of 6 cycles.

Note: Indication marked with a * includes indications that are unapproved. 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL).

Initial application — (Indolent, Low-grade lymphomas) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: All of the following:

- 1 The patient has indolent low grade NHL requiring treatment; and
- 2 The patient has ECOG performance status of 0-2; and
- 3 Any of the following:
  - 3.1 Both:
    - 3.1.1 Patient is treatment naive; and
    - 3.1.2 Bendamustine is to be administered for a maximum of 6 cycles (in combination with rituximab when CD20+); or
  - 3.2 Both:
    - 3.2.1 Patient is refractory to or has relapsed within 12 months of a rituximab containing combined chemo-immunotherapy regimen; and
    - 3.2.2 Bendamustine is to be administered in combination with obinutuzumab for a maximum of 6 cycles; or
  - 3.3 All of the following:
    - 3.3.1 The patient has not received prior bendamustine therapy; and
    - 3.3.2 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
    - 3.3.3 Patient has had a rituximab treatment-free interval of 12 months or more; or
  - 3.4 Bendamustine is to be administered as monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Renewal — (Indolent, Low-grade lymphomas) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: Either:

1 Both:

1.1 Patient is refractory to or has relapsed within 12 months of rituximab in combination with bendamustine; and

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

S	Subsidy	Fully	Brand or
(Manufa	acturer's Price) Sub	sidised	Generic
	\$ Per	1	Manufacturer

continued...

1.2 Bendamustine is to be administered in combination with obinutuzumab for a maximum of 6 cycles; or

2 Both:

2.1 Patients have not received a bendamustine regimen within the last 12 months; and

2.2 Either:

DUOLUEAN DOT DIVIL

2.2.1 Both:

- 2.2.1.1 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
- 2.2.1.2 Patient has had a rituximab treatment-free interval of 12 months or more; or
- 2.2.2 Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Note: 'indolent, low-grade lymphomas' includes follicular, mantle cell, marginal zone and lymphoplasmacytic/ Waldenstrom's macroglobulinaemia.

**Initial application — (Hodgkin's lymphoma*)** only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has Hodgkin's lymphoma requiring treatment; and
- 2 Patient has a ECOG performance status of 0-2; and
- 3 Patient has received one prior line of chemotherapy; and
- 4 Patient's disease relapsed or was refractory following prior chemotherapy; and

~

5 Bendamustine is to be administered in combination with gemcitabine and vinorelbine (BeGeV) at a maximum dose of no greater than 90 mg/m2 twice per cycle, for a maximum of four cycles.

Note: Indications marked with * are unapproved indications.

BUSULFAN – PCT – Retail pharmacy-Specialist			
Tab 2 mg		100	<ul> <li>Myleran</li> </ul>
CARBOPLATIN – PCT only – Specialist			
Inj 10 mg per ml, 45 ml vial	25.73	1	<ul> <li>Carboplatin Accord</li> </ul>
			<ul> <li>DBL Carboplatin</li> </ul>
			S29 S29
	32.59		<ul> <li>DBL Carboplatin</li> </ul>
	48.50		<ul> <li>Carbaccord</li> </ul>
Inj 1 mg for ECP	0.06	1 mg	<ul> <li>Baxter</li> </ul>
CARMUSTINE – PCT only – Specialist			
Inj 100 mg vial	710.00	1	BiCNU
			BiCNU S29 S29
			Novadoz S29
Inj 100 mg for ECP	710.00	100 mg OP	<ul> <li>Baxter</li> </ul>
(BiCNU S29 S29 Inj 100 mg vial to be delisted 1 July 2025)		-	
(Novadoz sego Inj 100 mg vial to be delisted 1 July 2025)			
CHLORAMBUCIL – PCT – Retail pharmacy-Specialist			
Tab 2 mg	29.06	25	Leukeran FC
	20100	_0	
CISPLATIN – PCT only – Specialist	0.45	1	<ul> <li>Cisplatin Accord</li> </ul>
Inj 1 mg per ml, 50 ml vial	15.00	I	<ul> <li>Cisplatin Accord</li> <li>Cisplatin Ebewe</li> </ul>
Inj 1 mg per ml, 100 ml vial		1	<ul> <li>Cisplatin Ebewe</li> <li>Cisplatin Accord</li> </ul>
	21.00	'	<ul> <li>Cisplatin Accord</li> <li>Cisplatin Ebewe</li> </ul>
	29.66		✓ DBL Cisplatin
Inj 1 mg for ECP		1 mg	✓ Baxter
		. mg	Bunton

(N	Subsidy (Manufacturer's Price)		Fully Subsidised	
<i>t</i>	\$	Per	1	Manufacturer
CYCLOPHOSPHAMIDE				
Tab 50 mg – PCT – Retail pharmacy-Specialist	145.00	50	1	Cyclonex
Inj 1 g vial – PCT – Retail pharmacy-Specialist	47.46	1	1	Endoxan
	127.80	6	1	Cytoxan
Inj 2 g vial – PCT only – Specialist		1		Endoxan
Inj 1 mg for ECP – PCT only – Specialist	0.05	1 mg	~	Baxter
FOSFAMIDE – PCT only – Specialist				
Inj 1 g	96.00	1	1	Holoxan
lnj 2 g	180.00	1	✓	Holoxan
Inj 1 mg for ECP	0.10	1 mg	~	Baxter
OMUSTINE – PCT – Retail pharmacy-Specialist				
Cap 40 mg	880.00	20	1	Medac S29
MELPHALAN				
Tab 2 mg – PCT – Retail pharmacy-Specialist	40.70	25	1	Alkeran
Inj 50 mg – PCT only – Specialist		1	1	Megval S29
				Melpha
	67.80			Alkeran
Megval 🖘 Inj 50 mg to be delisted 1 July 2025)				
DXALIPLATIN – PCT only – Specialist				
Inj 100 mg vial	25.01	1	1	Oxaliplatin Actavis
		•	-	100
	110.00		1	Oxaliplatin Ebewe
Inj 5 mg per ml, 20 ml vial		1		Alchemy Oxaliplatin
	46.32			Oxaliplatin Accord
Inj 1 mg for ECP	0.35	1 mg	1	Baxter
THIOTEPA – PCT only – Specialist				
Inj 15 mg vial	CBS	1	1	Bedford S29
		•		Max Health S29
				THIO-TEPA S29
	398.00			Tepadina
Inj 100 mg vial		1		Max Health S29
ing 100 mg viai	1,800.00	I		Tepadina
	1,000.00		•	repaulla
Antimetabolites				
AZACITIDINE – PCT only – Specialist – Special Authority see SA2	141 below			
Inj 100 mg vial		1	1	Azacitidine Dr
· -				Reddy's
Inj 1 mg for ECP	0.54	1 mg	1	Baxter
SA2141 Special Authority for Subsidy				

#### ial Author Subsidy

Initial application only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

1 Any of the following:

- 1.1 The patient has International Prognostic Scoring System (IPSS) intermediate-2 or high risk myelodysplastic syndrome; or
- 1.2 The patient has chronic myelomonocytic leukaemia (10%-29% marrow blasts without myeloproliferative disorder);

	ubsidy		Fully	
(Manufac	cturer's Price) \$	Per	Subsidised	Generic Manufacturer
ontinued				
or				
<ol> <li>The patient has acute myeloid leukaemia with 20-30% blast Health Organisation Classification (WHO); and</li> </ol>	is and multi-l	ineag	e dysplasi	ia, according to World
<ul><li>2 The patient has performance status (WHO/ECOG) grade 0-2; and</li><li>3 The patient has an estimated life expectancy of at least 3 months.</li></ul>				
tenewal only from a haematologist or medical practitioner on the recomments for applications meeting the following criteria: Both:	nendation of	a hae	matologis	t. Approvals valid for 1
<ol> <li>No evidence of disease progression; and</li> <li>The treatment remains appropriate and patient is benefitting from t</li> </ol>	reatment.			
CALCIUM FOLINATE				
Tab 15 mg – PCT – Retail pharmacy-Specialist	5.33	10	1	DBL Leucovorin Calcium
Inj 3 mg per ml, 1 ml – PCT – Retail pharmacy-Specialist17	7 10	5	1	Hospira
Inj 10 mg per ml, 5 ml vial – PCT – Retail pharmacy-Specialist		1		Calcium Folinate Sandoz
			~	Calcium Folinate Sandoz S29 S29
		-		Eurofolic S29
Inj 50 mg – PCT – Retail pharmacy-Specialist72	5.48 0.90	5 10		Leucovorin
inj 50 mg – PCT – Retail pharmacy-Specialist	2.80	10	v	
lei 10 ma annul 10 mInial - DOT anha Canaialist	10	4		Pharmacia S29
Inj 10 mg per ml, 10 ml vial – PCT only – Specialist		1		Calcium Folinate Sandoz
	7.45	5		Eurofolic S29
Inj 100 mg - PCT only - Specialist7	7.33	1	1	Calcium Folinate Ebewe
94	1.90	10	~	Leucovorin Pharmacia S29
Inj 300 mg – PCT only – Specialist21	.55	1	1	Leucovorin DBL S29
22	2.51		~	Calcium Folinate Ebewe
Inj 10 mg per ml, 35 ml vial – PCT only – Specialist25	5.14	1	1	Calcium Folinate Sandoz
			1	Calcium Folinate Sandoz S29 S29
Inj 1 g – PCT only – Specialist67	7.51	1	~	Calcium Folinate Ebewe
Inj 10 mg per ml, 100 ml vial – PCT only – Specialist72	2.00	1	1	Calcium Folinate Sandoz
			1	Eurofolic S29
Inj 1 mg for ECP – PCT only – Specialist	).06	1 mg	~	Baxter
APECITABINE – Retail pharmacy-Specialist				
Tab 150 mg	9.80	60	✓	Capecitabine Viatris
Tab 500 mg	6.50	120		Capecitabine Viatris
CLADRIBINE – PCT only – Specialist				
Inj 1 mg per ml, 10 ml749	9.96	1	✓	Leustatin
Inj 10 mg for ECP749		mg C		Baxter

	Subsidy (Manufacturer's	Price) Subs	Fully Brand or idised Generic
	(inditidated of 0 \$	Per	<ul> <li>Manufacturer</li> </ul>
CYTARABINE			
Inj 20 mg per ml, 5 ml vial – PCT – Retail pharmacy-Speci	ialist472.00	5	<ul> <li>Pfizer</li> </ul>
Inj 100 mg per ml, 20 ml vial – PCT – Retail			
pharmacy-Specialist		1	<ul> <li>Cytarabine DBL</li> </ul>
			<ul> <li>Pfizer</li> </ul>
			Pfizer S29 S29
Inj 1 mg for ECP – PCT only – Specialist	0.29	10 mg	<ul> <li>Baxter</li> </ul>
Inj 100 mg intrathecal syringe for ECP - PCT only - Speci	ialist94.40	100 mg OP	<ul> <li>Baxter</li> </ul>
FLUDARABINE PHOSPHATE		-	
Tab 10 mg - PCT - Retail pharmacy-Specialist		20	<ul> <li>Fludara Oral</li> </ul>
Inj 50 mg vial – PCT only – Specialist		1	✓ Fludarabine
,,,		-	Sagent S29
	634.00	5	✓ Fludarabine Ebewe
Inj 50 mg for ECP – PCT only – Specialist		50 mg OP	✓ Baxter
FLUOROUBACIL		oo nig or	Build
Ini 50 mg per ml, 20 ml vial – PCT only – Specialist	10 51	1	<ul> <li>Fluorouracil Accord</li> </ul>
Inj 50 mg per ml, 50 ml vial – PCT only – Specialist		1	<ul> <li>Fluorouracii Accor</li> <li>Fluorouracii Accor</li> </ul>
Inj 50 mg per ml, 100 ml vial – PCT only – Specialist		1	<ul> <li>Fluorouracil Accor</li> <li>Fluorouracil Accor</li> </ul>
Inj 1 mg for ECP – PCT only – Specialist		100 mg	<ul> <li>✓ Pluorouracii Accor</li> <li>✓ Baxter</li> </ul>
		Too mg	
GEMCITABINE HYDROCHLORIDE – PCT only – Specialist			
Inj 43.3 mg per ml (equivalent to 38 mg per ml gemcitabine	,		
26.3 ml vial		1	✓ DBL Gemcitabine
Inj 1 g		1	<ul> <li>Gemcitabine Ebew</li> <li>Bowter</li> </ul>
Inj 1 mg for ECP	0.02	1 mg	<ul> <li>Baxter</li> </ul>
RINOTECAN HYDROCHLORIDE – PCT only – Specialist			
Inj 20 mg per ml, 5 ml vial		1	<ul> <li>Accord</li> </ul>
	71.44		<ul> <li>Irinotecan Actavis</li> </ul>
			100
	100.00		<ul> <li>Irinotecan-Rex</li> </ul>
Inj 1 mg for ECP	0.54	1 mg	<ul> <li>Baxter</li> </ul>
MERCAPTOPURINE			
Tab 50 mg – PCT – Retail pharmacy-Specialist	25.90	25	<ul> <li>Puri-nethol</li> </ul>
Oral suspension 20 mg per ml - Retail pharmacy-Speciali	st –		
Special Authority see SA1725 below		100 ml OP	<ul> <li>Allmercap</li> </ul>
· · · ·			•

### ➡SA1725 Special Authority for Subsidy

Initial application only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where the patient requires a total dose of less than one full 50 mg tablet per day.

Renewal only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where patient still requires a total dose of less than one full 50 mg tablet per day.

	Subsidy		Fully Brand or
	(Manufacturer's Pr		idised Generic
	\$	Per	<ul> <li>Manufacturer</li> </ul>
ETHOTREXATE			
Tab 2.5 mg – PCT – Retail pharmacy-Specialist	7.80	90	<ul> <li>Trexate</li> </ul>
<ul> <li>Tab 10 mg – PCT – Retail pharmacy-Specialist</li> </ul>		90	✓ Trexate
<ul> <li>Inj 2.5 mg per ml, 2 ml – PCT – Retail pharmacy-Specialist</li> </ul>		5	<ul> <li>Methotrexate DBL</li> </ul>
<ul> <li>Inj 7.5 mg prefilled syringe</li> </ul>		1	<ul> <li>Methotrexate</li> </ul>
			Sandoz
<ul> <li>Inj 10 mg prefilled syringe</li> </ul>	19.09	1	<ul> <li>Methotrexate</li> </ul>
			Sandoz
Inj 15 mg prefilled syringe	24 53	1	✓ Methotrexate
	24.00		Sandoz
	10.04		
<ul> <li>Inj 20 mg prefilled syringe</li> </ul>		1	✓ <u>Methotrexate</u>
			Sandoz
<ul> <li>Inj 25 mg prefilled syringe</li> </ul>	20.72	1	<ul> <li>Methotrexate</li> </ul>
			<u>Sandoz</u>
<ul> <li>Inj 30 mg prefilled syringe</li> </ul>	55.00	1	<ul> <li>Methotrexate</li> </ul>
			Sandoz
<ul> <li>Inj 25 mg per ml, 2 ml vial – PCT – Retail pharmacy-Specia</li> </ul>	alist30.00	5	<ul> <li>Methotrexate DBL</li> </ul>
, ., ,,,,,,,,,,		-	Onco-Vial
Inj 25 mg per ml, 20 ml vial – PCT – Retail pharmacy-Spec	cialist 45.00	1	✓ DBL Methotrexate
	Janst		Onco-Vial
( Ini 100 mg nor ml 10 ml DCT Datail sharmony Crossial	int 05.00	4	
Inj 100 mg per ml, 10 ml – PCT – Retail pharmacy-Special	IST	1	<ul> <li>Methotrexate Ebewe</li> </ul>
<ul> <li>Inj 100 mg per ml, 50 ml vial – PCT – Retail</li> </ul>			
pharmacy-Specialist		1	<ul> <li>Methotrexate Ebewe</li> </ul>
<ul> <li>Inj 1 mg for ECP – PCT only – Specialist</li> </ul>		1 mg	<ul> <li>Baxter</li> </ul>
<ul> <li>Inj 5 mg intrathecal syringe for ECP – PCT only – Specialis</li> </ul>	st4.73	5 mg OP	<ul> <li>Baxter</li> </ul>
EMETREXED – PCT only – Specialist			
Inj 100 mg vial		1	Pemetrexed-AFT
	60.89	-	<ul> <li>Juno Pemetrexed</li> </ul>
Inj 500 mg vial		1	✓ Pemetrexed-AFT
	217.77		✓ Juno Pemetrexed
Inj 1 mg for ECP		1 mg	✓ Baxter
		i ing	
HIOGUANINE – PCT – Retail pharmacy-Specialist			
Tab 40 mg		25	<ul> <li>Lanvis</li> </ul>
Other Cytotoxic Agents			
MSACRINE - PCT only - Specialist			
MSACRINE – PCT only – Specialist	1 500 00	0	Ameridia -
Inj 50 mg per ml, 1.5 ml ampoule	-	6	Amsidine S29
	4,736.00		<ul> <li>Amsidine S29</li> </ul>
Inj 75 mg	1,250.00	5	<ul> <li>AmsaLyo S29</li> </ul>
-	6,218.00		AmsaLyo S29
NAGRELIDE HYDROCHLORIDE – PCT – Retail pharmacy-S		100	Acaulio
Cap 0.5 mg	1,1/5.8/	100	<ul> <li>Agrylin</li> </ul>
RSENIC TRIOXIDE – PCT only – Specialist			
Inj 1 mg per ml, 10 ml vial	4,817.00	10	<ul> <li>Phenasen</li> </ul>
Inj 10 mg for ECP		10 mg OP	<ul> <li>Baxter</li> </ul>
LEOMYCIN SULPHATE – PCT only – Specialist			
Inj 15,000 iu, vial	185 16	1	<ul> <li>DBL Bleomycin</li> </ul>
inj 10,000 lu, vlai		I	Sulfate
		1,000 iu	✓ Baxter
Inj 1,000 iu for ECP			

156 ✓ fully subsidised Principal Supply S29 Unapproved medicine supplied under Section 29 Sole Subsidised Supply

	Subsidy		Fully Brand or
	(Manufacturer's F		idised Generic
	\$	Per	<ul> <li>Manufacturer</li> </ul>
BORTEZOMIB - PCT only - Specialist - Special Authority see S	SA2355 below		
Inj 3.5 mg vial	74.93	1	<ul> <li>DBL Bortezomib</li> </ul>
Inj 1 mg for ECP	22.26	1 mg	<ul> <li>Baxter</li> </ul>
■ SA2355 Special Authority for Subsidy			
Initial application - (plasma cell dyscrasia) from any relevant	t practitioner. A	pprovals valid w	vithout further renewal unless
notified where the patient has plasma cell dyscrasia, not including	g Waldenström r	nacroglobulinae	emia, requiring treatment.
DACARBAZINE – PCT only – Specialist			
Inj 200 mg vial	72.11	1	<ul> <li>DBL Dacarbazine</li> </ul>
Inj 200 mg for ECP	72.11	200 mg OP	<ul> <li>Baxter</li> </ul>
DACTINOMYCIN [ACTINOMYCIN D] - PCT only - Specialist			
Inj 0.5 mg vial		1	<ul> <li>Cosmegen</li> </ul>
Inj 0.5 mg for ECP		0.5 mg OP	✓ Baxter
DAUNORUBICIN – PCT only – Specialist		-	
Inj 2 mg per ml, 10 ml		1	✓ Pfizer
Inj 20 mg for ECP		20 mg OP	✓ Baxter
DOCETAXEL – PCT only – Specialist		- 5 -	
Inj 20 mg	48 75	1	<ul> <li>Docetaxel Sandoz</li> </ul>
Inj 10 mg per ml, 8 ml vial		1	✓ DBL Docetaxel
Inj 20 mg per ml, 4 ml vial		1	✓ Docetaxel
			Accord S29
Inj 80 mg	195.00	1	✓ Docetaxel Sandoz
Inj 1 mg for ECP		1 mg	✓ Baxter
DOXORUBICIN HYDROCHLORIDE – PCT only – Specialist		i iig	Bunton
Inj 2 mg per ml, 5 ml vial	10.00	1	Doxorubicin Ebewe
Inj 2 mg per ml, 25 ml vial		1	<ul> <li>Doxorubicin Ebewe</li> <li>Doxorubicin Ebewe</li> </ul>
11) z 119 per 111, zo 111 viai	11.50	I	<ul> <li>Arrow-Doxorubicin</li> </ul>
Inj 2 mg per ml, 50 ml vial		1	<ul> <li>Anow-Doxorubicin</li> <li>Doxorubicin Ebewe</li> </ul>
Inj 2 mg per ml, 100 ml vial		1	✓ Arrow-Doxorubicin
	69.99		✓ Doxorubicin Ebewe
Inj 1 mg for ECP		1 mg	✓ Baxter
EPIRUBICIN HYDROCHLORIDE - PCT only - Specialist		5	
Inj 2 mg per ml, 5 ml vial	25.00	1	<ul> <li>Epirubicin Ebewe</li> </ul>
Inj 2 mg per ml, 25 ml vial		1	<ul> <li>Epirubicin Ebewe</li> </ul>
Inj 2 mg per ml, 100 ml vial		1	<ul> <li>Epirubicin Ebewe</li> </ul>
Inj 1 mg for ECP		1 mg	✓ Baxter
ETOPOSIDE		0	
Cap 50 mg – PCT – Retail pharmacy-Specialist	340.73	20	✓ Vepesid
Cap 100 mg – PCT – Retail pharmacy-Specialist		10	✓ Vepesid
Inj 20 mg per ml, 5 ml vial – PCT – Retail pharmacy-Special		1	✓ Rex Medical
Inj 1 mg for ECP – PCT only – Specialist		1 mg	✓ Baxter
ETOPOSIDE PHOSPHATE – PCT only – Specialist		Ū	
Inj 100 mg (of etoposide base)		1	<ul> <li>Etopophos</li> </ul>
Inj 1 mg (of etoposide base) for ECP		1 mg	✓ Baxter
HYDROXYUREA [HYDROXYCARBAMIDE] – PCT – Retail pha		0	
Cap 500 mg		100	<ul> <li>Devatis</li> </ul>
		100	- Devans
IBRUTINIB – Special Authority see SA2168 on the next page –		00	. Imbuudaa
Tab 140 mg	,	30	<ul> <li>Imbruvica</li> <li>Imbruvica</li> </ul>
Tab 420 mg	9,052.00	30	<ul> <li>Imbruvica</li> </ul>

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

Subsidy		Fully	Brand or
(Manufacturer's Price)	_	Subsidised	Generic
\$	Per	/	Manufacturer

### ⇒SA2168 Special Authority for Subsidy

**Initial application** — (chronic lymphocytic leukaemia (CLL)) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic lymphocytic leukaemia (CLL) requiring therapy; and
- 2 Patient has not previously received funded ibrutinib; and
- 3 Ibrutinib is to be used as monotherapy; and
- 4 Any of the following:
  - 4.1 Both:
    - 4.1.1 There is documentation confirming that patient has 17p deletion or TP53 mutation; and
    - 4.1.2 Patient has experienced intolerable side effects with venetoclax monotherapy; or
  - 4.2 All of the following:
    - 4.2.1 Patient has received at least one prior immunochemotherapy for CLL; and
    - 4.2.2 Patient's CLL has relapsed within 36 months of previous treatment; and
    - 4.2.3 Patient has experienced intolerable side effects with venetoclax in combination with rituximab regimen; or
  - 4.3 Patient's CLL is refractory to or has relapsed within 36 months of a venetoclax regimen.

Renewal — (chronic lymphocytic leukaemia (CLL)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

1 No evidence of clinical disease progression; and

2 The treatment remains appropriate and the patient is benefitting from treatment.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL) and B-cell prolymphocytic leukaemia (B-PLL)*. Indications marked with * are Unapproved indications.

IDARUBICIN HYDROCHLORIDE

Inj 5 mg vial – PCT only – Specialist Inj 10 mg vial – PCT only – Specialist Inj 1 mg for ECP – PCT only – Specialist	233.64	1 1 1 mg	<ul><li>✓ Zavedos</li><li>✓ Zavedos</li><li>✓ Baxter</li></ul>
LENALIDOMIDE (VIATRIS) - Special Authority see SA23	53 below – Retail pharm	acy	
Cap 5 mg		21	✓ <u>Lenalidomide</u> <u>Viatris</u>
Cap 10 mg		21	✓ <u>Lenalidomide</u> <u>Viatris</u>
Cap 15 mg	62.13	21	✓ <u>Lenalidomide</u> <u>Viatris</u>
Cap 25 mg	65.09	21	✓ <u>Lenalidomide</u> Viatris

#### ⇒SA2353 Special Authority for Subsidy

Initial application — (Plasma cell dyscrasia) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has plasma cell dyscrasia, not including Waldenström macroglobulinaemia, requiring treatment; and
- 2 Patient is not refractory to prior lenalidomide use.

**Initial application — (Myelodysplastic syndrome)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 Patient has low or intermediate-1 risk myelodysplastic syndrome (based on IPSS or an IPSS-R score of less than 3.5)

	Subsidy (Manufacturer's Price	e) (	Fully Subsidised	Brand or Generic
	\$	Per	<ul> <li>✓</li> </ul>	Manufacturer
continued				
associated with a deletion 5q cytogenetic abnormality; and	l			
2 Patient has transfusion-dependent anaemia.				
Renewal — (Myelodysplastic syndrome) from any relevant pra the following criteria: Both:	ctitioner. Approva	ls valid f	for 12 mon	ths for applications meeting
<ol> <li>Patient has not needed a transfusion in the last 4 months;</li> <li>No evidence of disease progression.</li> </ol>	and			
MESNA				
Tab 400 mg – PCT – Retail pharmacy-Specialist	314.00	50	<ul> <li>Image: A second s</li></ul>	Uromitexan
Tab 600 mg - PCT - Retail pharmacy-Specialist	448.50	50	✓	Uromitexan
Inj 100 mg per ml, 4 ml ampoule - PCT only - Specialist	177.45	15	✓	Uromitexan
Inj 100 mg per ml, 10 ml ampoule – PCT only – Specialist	407.40	15	✓	Uromitexan
Inj 1 mg for ECP – PCT only – Specialist	2.96	100 mg	, 🖌	Baxter
MITOMYCIN C – PCT only – Specialist				
Inj 5 mg vial	517.65	1	✓	Mitomycin (Fresenius Kabi) ^{S29}
	526.00		✓	Mitomycin (Sagent) 829
	641.70		✓ ,	Accord S29
Inj 20 mg vial	1.250.00	1	1	Omegapharm S29
		•		Teva
Inj 1 mg for ECP		1 mg	✓	Baxter
MITOZANTRONE – PCT only – Specialist		0		
Inj 2 mg per ml, 10 ml vial	97 50	1	<ul> <li>Image: A second s</li></ul>	Mitozantrone Ebewe
Inj 1 mg for ECP		1 mg		Baxter
NIRAPARIB – Special Authority see SA2325 below – Retail phar Wastage claimable				
Tab 100 mg	13,393.50	84	✓ :	Zejula
Cap 100 mg		56		Zejula
	13,393.50	84	✓ :	Zejula
SA2325 Special Authority for Subsidy				

SA2325 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has advanced high-grade serous* epithelial ovarian, fallopian tube, or primary peritoneal cancer; and
- 2 Patient has received at least one line** of treatment with platinum-based chemotherapy; and
- 3 Patient has experienced a partial or complete response to the preceding treatment with platinum-based chemotherapy; and
- 4 Patient has not previously received funded treatment with a PARP inhibitor; and
- 5 Either:
  - 5.1 Treatment will be commenced within 12 weeks of the patient's last dose of the preceding platinum-based regimen; or
  - 5.2 Patient commenced treatment with niraparib prior to 1 May 2024; and
- 6 Treatment to be administered as maintenance treatment; and
- 7 Treatment not to be administered in combination with other chemotherapy.

Renewal from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

continued...

AThree months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

Subsidy		Fully	Brand or
(Manufacturer's Price)	5	Subsidised	Generic
\$	Per	~	Manufacturer

continued...

All of the following:

- 1 No evidence of progressive disease; and
- 2 Treatment to be administered as maintenance treatment; and
- 3 Treatment not to be administered in combination with other chemotherapy; and
- 4 Either:
  - 4.1 Treatment with niraparib to cease after a total duration of 36 months from commencement; or
  - 4.2 Treatment with niraparib is being used in the second-line or later maintenance setting.

Notes: * "high-grade serous" includes tumours with high-grade serous features or a high-grade serous component. **A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments

OLAPARIB - Retai	pharmacy-Specialist - Specia	I Authority see SA2163 below
------------------	------------------------------	------------------------------

🗸 Lynparza	56	 Tab 100 mg
<ul> <li>Lynparza</li> </ul>	56	 Tab 150 mg

#### ⇒SA2163 Special Authority for Subsidy

**Initial application — (Ovarian cancer)** only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Patient has a high-grade serous* epithelial ovarian, fallopian tube, or primary peritoneal cancer; and
- 2 There is documentation confirming pathogenic germline BRCA1 or BRCA2 gene mutation; and
- 3 Either:
  - 3.1 All of the following:
    - 3.1.1 Patient has newly diagnosed, advanced disease; and
    - 3.1.2 Patient has received one line** of previous treatment with platinum-based chemotherapy; and
    - 3.1.3 Patient's disease must have experienced a partial or complete response to the first-line platinum-based regimen; or
  - 3.2 All of the following:
    - 3.2.1 Patient has received at least two lines** of previous treatment with platinum-based chemotherapy; and
    - 3.2.2 Patient has platinum sensitive disease defined as disease progression occurring at least 6 months after the last dose of the penultimate line** of platinum-based chemotherapy; and
    - 3.2.3 Patient's disease must have experienced a partial or complete response to treatment with the immediately preceding platinum-based regimen; and
    - 3.2.4 Patient has not previously received funded olaparib treatment; and
- 4 Treatment will be commenced within 12 weeks of the patient's last dose of the immediately preceding platinum-based regimen; and
- 5 Treatment to be administered as maintenance treatment; and
- 6 Treatment not to be administered in combination with other chemotherapy.

Renewal — (Ovarian cancer) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from treatment; and
- 2 Either:
  - 2.1 No evidence of progressive disease; or
  - 2.2 Evidence of residual (not progressive) disease and the patient would continue to benefit from treatment in the clinician's opinion; and
- 3 Treatment to be administered as maintenance treatment; and
- 4 Treatment not to be administered in combination with other chemotherapy; and

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subs	idised	Generic
 \$	Per	1	Manufacturer

continued...

5 Either:

- 5.1 Both:
  - 5.1.1 Patient has received one line** of previous treatment with platinum-based chemotherapy; and
  - 5.1.2 Documentation confirming that the patient has been informed and acknowledges that the funded treatment period of olaparib will not be continued beyond 2 years if the patient experiences a complete response to treatment and there is no radiological evidence of disease at 2 years: or
- 5.2 Patient has received at least two lines** of previous treatment with platinum-based chemotherapy.

Notes: *Note "high-grade serous" includes tumours with high-grade serous features or a high-grade serous component. **A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

PACLITAXEL – PCT only – Specialist			
Inj 30 mg		5	Paclitaxel Ebewe
Inj 6 mg per ml, 16.7 ml vial		1	<ul> <li>Anzatax</li> </ul>
	24.00		Paclitaxel Ebewe
	91.67		Paclitaxel Actavis
lnj 150 mg		1	Paclitaxel Ebewe
, 0	137.50		<ul> <li>Anzatax</li> </ul>
			Paclitaxel Actavis
Inj 6 mg per ml, 50 ml vial		1	<ul> <li>Anzatax</li> </ul>
	44.00		Paclitaxel Ebewe
	275.00		Paclitaxel Actavis
Inj 1 mg for ECP	0.17	1 mg	<ul> <li>Baxter</li> </ul>
PEGASPARGASE - PCT only - Special Authority see	SA1979 below		
Inj 750 iu per ml, 5 ml vial		1	<ul> <li>Oncaspar LYO</li> </ul>

#### ► SA1979 Special Authority for Subsidy

Initial application — (Acute lymphoblastic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has newly diagnosed acute lymphoblastic leukaemia: and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

Initial application - (Lymphoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months where the patient has lymphoma requiring L-asparaginase containing protocols (e.g. SMILE).

Renewal — (Acute lymphoblastic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: Both:

1 The patient has relapsed acute lymphoblastic leukaemia: and

2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

#### PENTOSTATIN [DEOXYCOFORMYCIN] - PCT only - Specialist

Inj 10 mg	CBS	1	<ul> <li>Nipent S29</li> </ul>
POMALIDOMIDE - Special Authority see SA2354 on th	e next page – Retail pharm	nacy	
Cap 1 mg		14	Pomolide
	71.18	21	Pomolide
Cap 2 mg		14	Pomolide
	142.35	21	Pomolide
Cap 3 mg		14	Pomolide
	213.53	21	Pomolide
Cap 4 mg		14	Pomolide
	284.71	21	Pomolide

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Subs Per	Fully idised	Brand or Generic Manufacturer
SA2354 Special Authority for Subsidy				
Initial application — (Relapsed/refractory plasma cell dyscra	sia) from any relevar	nt practitio	ner. Ap	provals valid for 6 months
for applications meeting the following criteria:				
Both:				
<ol> <li>Patient has relapsed or refractory plasma cell dyscrasia, r treatment; and</li> </ol>	not including Waldens	tröm macr	oglobul	inaemia, requiring
2 Patient has not received prior funded pomalidomide.				
Renewal — (Relapsed/refractory plasma cell dyscrasia) from there is no evidence of disease progression.	any relevant practitic	oner. App	rovals v	alid for 12 months where
PROCARBAZINE HYDROCHLORIDE - PCT - Retail pharmacy	-Specialist			
Cap 50 mg		50	🗸 N	atulan S29
TEMOZOLOMIDE – Special Authority see SA2275 below – Reta				
Cap 5 mg		5	🖌 Т	emaccord
oup o mg		0	-	emozolomide-
			•	Taro S29
Cap 20 mg		5	🗸 T	emaccord
	18.30		🗸 A	po-Temozolomide
Cap 100 mg		5	🖌 T	emaccord
	40.20		🗸 A	po-Temozolomide
Cap 140 mg		5	🗸 T	emaccord
Cap 250 mg		5	🗸 T	emaccord

### ➡SA2275 Special Authority for Subsidy

**Initial application** — (gliomas) only from a relevant specialist. Approvals valid for 12 months where the patient has a glioma. **Renewal** — (gliomas) only from a relevant specialist. Approvals valid for 12 months where treatment remains appropriate and patient is benefitting from treatment.

**Initial application** — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with metastatic or unresectable well-differentiated neuroendocrine tumour*; and
- 2 Temozolomide is to be given in combination with capecitabine; and
- 3 Temozolomide is to be used in 28 day treatment cycles for a maximum of 5 days treatment per cycle at a maximum dose of 200 mg/m² per day; and
- 4 Temozolomide to be discontinued at disease progression.

Renewal — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

**Initial application** — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 9 months where the patient has relapsed/refractory Ewing's sarcoma.

Renewal — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indication marked with a * is an unapproved indication. Temozolomide is not subsidised for the treatment of relapsed high grade glioma.

	Subsidy (Manufacturer's Price) \$	Su Per	Fully bsidised	Brand or Generic Manufacturer
THALIDOMIDE - Retail pharmacy-Specialist - Special Authority	see SA2356 below			
Cap 50 mg		28	🗸 T	halomid
Cap 100 mg	756.00	28	🗸 T	halomid

#### ⇒SA2356 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months where the patient has plasma cell dyscrasia, not including Waldenström macroglobulinaemia, requiring treatment.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has obtained a response from treatment during the initial approval period.

Notes: Prescription must be written by a registered prescriber in the thalidomide risk management programme operated by the supplier.

Maximum dose of 400 mg daily as monotherapy or in a combination therapy regimen.

#### TRETINOIN

Cap 10 mg – PCT – Retail pharmacy-Specialist	479.50	100	<ul> <li>Vesanoid</li> </ul>
VENETOCLAX - Retail pharmacy-Specialist - Special Authority see	SA1868 below		
Tab 14 × 10 mg, 7 × 50 mg, 21 × 100 mg	1,771.86	42 OP	<ul> <li>Venclexta</li> </ul>
Tab 10 mg		2 OP	<ul> <li>Venclexta</li> </ul>
Tab 50 mg	239.44	7 OP	<ul> <li>Venclexta</li> </ul>
Tab 100 mg - Wastage claimable	8,209.41	120	Venclexta

#### ➡SA1868 Special Authority for Subsidy

**Initial application** — (relapsed/refractory chronic lymphocytic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic lymphocytic leukaemia requiring treatment; and
- 2 Patient has received at least one prior therapy for chronic lymphocytic leukaemia; and
- 3 Patient has not previously received funded venetoclax; and
- 4 The patient's disease has relapsed within 36 months of previous treatment; and
- 5 Venetoclax to be used in combination with six 28-day cycles of rituximab commencing after the 5-week dose titration schedule with venetoclax; and
- 6 Patient has an ECOG performance status of 0-2.

Renewal — (relapsed/refractory chronic lymphocytic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment; and
- 2 Venetoclax is to be discontinued after a maximum of 24 months of treatment following the titration schedule unless earlier discontinuation is required due to disease progression or unacceptable toxicity.

Initial application — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has previously untreated chronic lymphocytic leukaemia; and
- 2 There is documentation confirming that patient has 17p deletion by FISH testing or TP53 mutation by sequencing; and
- 3 Patient has an ECOG performance status of 0-2.

Renewal — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where the treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL)* and B-cell prolymphocytic leukaemia (B-PLL)*. Indications marked with * are Unapproved indications.

	Subsidy		Fully	Brand or
	(Manufacturer's Price)	Su	bsidised	Generic
	\$	Per	1	Manufacturer
INBLASTINE SULPHATE				
Inj 1 mg per ml, 10 ml vial - PCT - Retail pharmacy-Spec	ialist270.37	5	<ul> <li>Image: A second s</li></ul>	Hospira
Inj 1 mg for ECP – PCT only – Specialist	6.00	1 mg	<ul> <li>I</li> </ul>	Baxter
INCRISTINE SULPHATE				
Inj 1 mg per ml, 1 ml vial - PCT - Retail pharmacy-Specia	alist74.52	5	<ul> <li>Image: A second s</li></ul>	DBL Vincristine
				Sulfate
Inj 1 mg per ml, 2 ml vial – PCT – Retail pharmacy-Specia	alist102.73	5	<ul> <li>Image: A second s</li></ul>	DBL Vincristine
				Sulfate
Inj 1 mg for ECP – PCT only – Specialist	12.60	1 mg	<ul> <li>I</li> </ul>	Baxter
INORELBINE				
Cap 20 mg		1	1	/inorelbine Te Arai
Cap 30 mg	40.00	1	<ul> <li>Image: A second s</li></ul>	/inorelbine Te Arai
Cap 80 mg		1	-	/inorelbine Te Arai
Inj 10 mg per ml, 1 ml vial – PCT only – Specialist		1		Vinorelbine Ebewe
Inj 10 mg per ml, 5 ml vial – PCT only – Specialist	168.00	1		Navelbine S29 S29
	210.00			Vinorelbine Ebewe
Inj 1 mg for ECP – PCT only – Specialist	3.80	1 mg	✓	Baxter
Protein-tyrosine Kinase Inhibitors				
FOTINIP _ Details to serve the site of the Oracical Anthonism	044070 h d			
LECTINIB – Retail pharmacy-Specialist – Special Authority s	see SA1870 below			
Wastage claimable Cap 150 mg	7 025 00	224	1	Alecensa
		224	• 1	AIECEIISa
»SA1870 Special Authority for Subsidy	reatitioner on the record	amandati	an af a r	ale cont en esiglist
itial application only from a medical oncologist or medical p pprovals valid for 6 months for applications meeting the follow		Imenuali	011 01 a 1	elevant specialist.
Il of the following:	wing ontona.			
<ol> <li>Patient has locally advanced, or metastatic, unresectable</li> </ol>	le non-small cell lung	rancer: a	nd	
2 There is documentation confirming that the patient has				nent using an appropriat
ALK test; and		<u>j</u>		
3 Patient has an ECOG performance score of 0-2.				
enewal only from a medical oncologist or medical practitione	r on the recommendation	on of a re	levant s	pecialist. Approvals vali
r 6 months for applications meeting the following criteria:				•••
oth:				
1 No evidence of progressive disease according to RECI	ST criteria; and			
2 The patient is benefitting from and tolerating treatment.				

DASATINIB - Special Authority see SA2385 below - Retail pharmacy

### ⇒SA2385 Special Authority for Subsidy

Initial application only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

1 The patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis or accelerated phase; or

Subsidy		Fully	Brand or	
(Manufacturer's Price	e) 5	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

2 The patient has a diagnosis of Philadelphia chromosome-positive acute lymphoid leukaemia (Ph+ ALL); or

3 Both:

3.1 The patient has a diagnosis of CML in chronic phase; and

- 3.2 Any of the following:
  - 3.2.1 Patient has documented treatment failure* with imatinib; or
  - 3.2.2 Patient has experienced treatment-limiting toxicity with imatinib precluding further treatment with imatinib; or
  - 3.2.3 Patient has high-risk chronic-phase CML defined by the Sokal or EURO scoring system.

**Renewal** only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Lack of treatment failure while on dasatinib*; and
- 2 Dasatinib treatment remains appropriate and the patient is benefiting from treatment.

Note: *treatment failure for CML as defined by Leukaemia Net Guidelines.

ERLOTINIB - Retail pharmacy-Specialist - Special Author	ity see SA2422 below		
Tab 100 mg		30	<ul> <li>Alchemy</li> </ul>
Tab 150 mg		30	<ul> <li>Alchemy</li> </ul>

#### ➡SA2422 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 There is documentation confirming that the disease expresses activating mutations of EGFR; and
- 3 Any of the following:
  - 3.1 Patient is treatment naive; or
  - 3.2 Patient has received prior treatment in the adjuvant setting and/or while awaiting EGFR results; or

3.3 Both:

- 3.3.1 The patient has discontinued osimertinib or gefitinib due to intolerance; and
- 3.3.2 The cancer did not progress while on osimertinib or gefitinib.

Renewal from any relevant practitioner. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

GEFITINIB - Retail pharmacy-Specialist - Special Authority see SA2423 below

Tab 250 mg	0 🖌 Ire	essa
------------	---------	------

#### ► SA2423 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 Any of the following:
  - 2.1 Patient is treatment naive; or
  - 2.2 Patient has received prior treatment in the adjuvant setting and/or while awaiting EGFR results; or
  - 2.3 Both:

IMATINIB MESILATE

- 2.3.1 The patient has discontinued osimertinib or erlotinib due to intolerance; and
- 2.3.2 The cancer did not progress whilst on osimertinib or erlotinib; and
- 3 There is documentation confirming that disease expresses activating mutations of EGFR.

Renewal from any relevant practitioner. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

*	Cap 100 mg	44.93	60	Imatinib-Rex
*	Cap 400 mg	69.76	30	<ul> <li>Imatinib-Rex</li> </ul>

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
LENVATINIB – Special Authority see SA2442 below – Retail pha Wastage claimable	armacy			
Cap 4 mg	3.407.40	30	🗸 L	envima
Cap 10 mg	3,407.40	30	🗸 L	envima
➡ SA2442 Special Authority for Subsidy				

**Initial application** — (thyroid cancer) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Patient is currently on treatment with lenvatinib and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
  - 2.1 The patient has locally advanced or metastatic differentiated thyroid cancer; and
  - 2.2 Either:
    - 2.2.1 Patient must have symptomatic progressive disease prior to treatment; or
    - 2.2.2 Patient must progressive disease at critical anatomical sites with a high risk of morbidity or mortality where local control cannot be achieved by other measures; and
  - 2.3 Any of the following:
    - 2.3.1 A lesion without iodine uptake in a RAI scan; or
    - 2.3.2 Receiving cumulative RAI greater than or equal to 600 mCi; or
    - 2.3.3 Experiencing disease progression after a RAI treatment within 12 months; or
    - 2.3.4 Experiencing disease progression after two RAI treatments administered within 12 months of each other; and
  - 2.4 Patient has thyroid stimulating hormone (TSH) adequately supressed; and
  - 2.5 Patient is not a candidate for radiotherapy with curative intent; and
  - 2.6 Surgery is clinically inappropriate; and
  - 2.7 Patient has an ECOG performance status of 0-2.

Renewal — (thyroid cancer) from any relevant practitioner. Approvals valid for 6 months where there is no evidence of disease progression.

**Initial application** — (unresectable hepatocellular carcinoma) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has unresectable hepatocellular carcinoma; and
- 2 Patient has preserved liver function (Childs-Pugh A); and
- 3 Transarterial chemoembolisation (TACE) is unsuitable; and
- 4 Patient has an ECOG performance status of 0-2; and
- 5 Either:
  - 5.1 Patient has not received prior systemic therapy for their disease in the palliative setting; or
  - 5.2 Both:
    - 5.2.1 Patient has experienced treatment-limiting toxicity from treatment with atezolizumab with bevacizumab; and
    - 5.2.2 No disease progression since initiation of atezolizumab with bevacizumab.

Renewal — (unresectable hepatocellular carcinoma) from any relevant practitioner. Approvals valid for 6 months where there is no evidence of disease progression.

Initial application — (renal cell carcinoma) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 All of the following:
  - 1.1 The patient has metastatic renal cell carcinoma; and
  - 1.2 The disease is of predominant clear-cell histology; and
  - 1.3 The patient has documented disease progression following one previous line of treatment; and

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Generic
<ul> <li>continued         <ol> <li>The patient has an ECOG performance status of 0</li> <li>Lenvatinib is to be used in combination with everol</li> </ol> </li> <li>All of the following:         <ol> <li>Patient has received funded treatment with nivolun carcinoma; and</li> <li>Patient has experienced treatment limiting toxicity</li> <li>Lenvatinib is to be used in combination with everol</li> <li>Patient has experienced treatment limiting toxicity</li> <li>Lenvatinib is to be used in combination with everol</li> <li>There is no evidence of disease progression.</li> </ol> </li> <li>Renewal — (renal cell carcinoma) from any relevant practitione disease progression.</li> <li>MIDOSTAURIN – PCT only – Special Authority see SA2342 bekeed Cap 25 mg</li></ul>	imus; or nab for the second lir from treatment with r imus; and er. Approvals valid for er. 10,981.00 d for 9 months for app n positive; and emotherapy for acute	ivolu or 4 n 56 plicat	mab; and nonths whe ions meeti loid leukae	ere there is no evidence of <b>Rydapt</b> ng the following criteria: emia; and
NILOTINIB – Special Authority see SA2301 below – Retail pharr Wastage claimable Cap 150 mg Cap 200 mg	4,680.00 6,532.00 r 6 months for applica		s meeting t	-
<ol> <li>Patient has a diagnosis of chronic myeloid leukaemia (CM and</li> <li>Either:         <ol> <li>Patient has documented CML treatment failure* wi</li> <li>Patient has experienced treatment limiting toxicity and</li> <li>Maximum nilotinib dose of 800 mg/day; and</li> <li>Subsidised for use as monotherapy only.</li> </ol> </li> <li>Note: *treatment failure as defined by Leukaemia Net Guidelines         Renewal only from a haematologist. Approvals valid for 6 month             All of the following:             <ol> <li>Lack of treatment failure while on nilotinib as defined by L</li> <li>Nilotinib treatment remains appropriate and the patient is             <ol> <li>Maximum nilotinib dose of 800 mg/day; and</li> <li>Subsidised for use as monotherapy only.</li> </ol> </li> </ol></li></ol>	th a tyrosine kinase i with a tyrosine kinase s. Is for applications me eukaemia Net Guidel	nhibit e inhil eeting lines;	or (TKI); o bitor (TKI) the follow and	r precluding further treatment;

OSIMERTINIB – Special Authority see SA2418 on the	next page – Retail pharmacy		
Tab 40 mg	9,310.00	30	🗸 Tagrisso
Tab 80 mg	9,310.00	30	<ul> <li>Tagrisso</li> </ul>

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy	Fi	ully	Brand or
(	Manufacturer's Price)	Subsidis	ed	Generic
	\$	Per	×	Manufacturer

### ⇒SA2418 Special Authority for Subsidy

**Initial application** — (NSCLC – first line) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Patient is currently on treatment with osimertinib and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
  - 2.1 Patient has locally advanced or metastatic, incurable, non-squamous non-small cell lung cancer (NSCLC); and
  - 2.2 Any of the following:
    - 2.2.1 Patient is treatment naïve; or
    - 2.2.2 Patient has received prior chemotherapy in the adjuvant setting and/or while awaiting EGFR results; or 2.2.3 Both:
      - 2.2.3.1 The patient has discontinued gefitinib or erlotinib due to intolerance; and
      - 2.2.3.2 The cancer did not progress while on gefitinib or erlotinib; and
  - 2.3 There is documentation confirming that the cancer expresses activating mutations of EGFR; and
  - 2.4 Patient has an ECOG performance status 0-3; and
  - 2.5 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Renewal — (NSCLC – first line) from any relevant practitioner. Approvals valid for 6 months where response to or stable disease with treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period.

Initial application — (NSCLC – second line) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

#### Either:

- 1 Patient is currently on treatment with osimertinib and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
  - 2.1 Patient has locally advanced or metastatic, incurable, non-squamous non-small cell lung cancer (NSCLC); and
  - 2.2 Patient has an ECOG performance status 0-3; and
  - 2.3 The patient must have received previous treatment with erlotinib or gefitinib; and
  - 2.4 There is documentation confirming that the cancer expresses T790M mutation of EGFR following progression on or after erlotinib or gefitinib; and
  - 2.5 The treatment must be given as monotherapy; and
  - 2.6 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Renewal — (NSCLC – second line) from any relevant practitioner. Approvals valid for 6 months where response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period.

PALBOCICLIB - Special Authority see SA2345 below - Retail pharmacy

Wastage claimable			
Tab 75 mg		21	<ul> <li>Ibrance</li> </ul>
Tab 100 mg		21	<ul> <li>Ibrance</li> </ul>
5	-	21	<ul> <li>Ibrance</li> </ul>
Tab 125 mg	4,000.00	21	<ul> <li>Ibranc</li> </ul>

#### ⇒SA2345 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Either:

- 1 All of the following:
  - 1.1 Patient has unresectable locally advanced or metastatic breast cancer; and
  - 1.2 There is documentation confirming disease is hormone-receptor positive and HER2-negative; and
  - 1.3 Patient has an ECOG performance score of 0-2; and
  - 1.4 Either:

(Manufacturer's Price) \$	Subsi Per	dised ✓	Generic Manufacturer	
	Per	1	Manufacturer	
prior endocrine thera	py; or			
	ndocrine le	evels c	onsistent with a	
1 /				
	static disea	ise; an	ıd	
n a CDK4/6 inhibitor;	or			
,				
ction to ribociclib tha	t cannot be	e mana	aged by dose reductions	5
	monting t	ha falle	wing oritorio:	
onuns ior applications	meeting t		Jwing chiena.	
narther: and				
·				
	30	<b>у</b> р	azonanih Teva	
	50			
1,004.70		•	othent	
	30	🗸 Р	azopanib Teva	
2,669.40				
	Illy or induced, with e -potential state; and ic treatment for metas docrine partner; and h a CDK4/6 inhibitor; ribociclib; and ction to ribociclib tha docrine partner; and nitiation of ribociclib. onths for applications partner; and h of palbociclib. rmacy 172.88 1,334.70 	Illy or induced, with endocrine le- potential state; and ic treatment for metastatic disea docrine partner; and h a CDK4/6 inhibitor; or ribociclib; and ction to ribociclib that cannot be docrine partner; and nitiation of ribociclib. onths for applications meeting the partner; and h of palbociclib. macy 	Illy or induced, with endocrine levels c         -potential state; and         ic treatment for metastatic disease; ar         docrine partner; and         h a CDK4/6 inhibitor; or         ribociclib; and         ctorine partner; and         nitiation of ribociclib that cannot be mana         docrine partner; and         nitiation of ribociclib.         onths for applications meeting the follow         partner; and         n of palbociclib.         macy         1,334.70	Illy or induced, with endocrine levels consistent with a -potential state; and ic treatment for metastatic disease; and docrine partner; and n a CDK4/6 inhibitor; or         ribociclib; and corrine partner; and nitiation of ribociclib. onths for applications meeting the following criteria:         partner; and not partner; and nitiation of ribociclib. onths for applications meeting the following criteria:         partner; and not partner; and nitiation of ribociclib. onths for applications meeting the following criteria:         partner; and n of palbociclib.         macy

(Votrient Tab 400 mg to be delisted 1 May 2025)

### ⇒SA2429 Special Authority for Subsidy

**Initial application** only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria: Either:

- 1 All of the following:
  - 1.1 The patient has metastatic renal cell carcinoma of predominantly clear cell histology; and
  - 1.2 Either:
    - 1.2.1 The patient is treatment naive; or
    - 1.2.2 The patient has only received prior cytokine treatment; and
  - 1.3 The patient has an ECOG performance score of 0-2; and
    - The patient has intermediate or poor prognosis defined as:
  - 1.4 Any of the following:
    - 1.4.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
    - 1.4.2 Haemoglobin level < lower limit of normal; or
    - 1.4.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
    - 1.4.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
    - 1.4.5 Karnofsky performance score of less than or equal to 70; or
    - 1.4.6 2 or more sites of organ metastasis; and
  - 1.5 Pazopanib to be used for a maximum of 3 months; or
- 2 All of the following:

continued...

AThree months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

Subsidy	Ful	y Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per •	Manufacturer	

continued...

- 2.1 The patient has metastatic renal cell carcinoma; and
- 2.2 The patient has discontinued sunitinib within 3 months of starting treatment due to intolerance; and
- 2.3 The cancer did not progress whilst on sunitinib; and
- 2.4 Pazopanib to be used for a maximum of 3 months.

Renewal only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months where there is no evidence of disease progression.

RIBOCICLIB - Special Authority see SA2343 below - Retail pharmacy

Wastage claimable

Tab 200 mg		21	🗸 Kisqali
C C	3,767.00	42	🗸 Kisqali
	5,650.00	63	🗸 Kisqali

#### ⇒SA2343 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Either:

- 1 All of the following:
  - 1.1 Patient has unresectable locally advanced or metastatic breast cancer; and
  - 1.2 There is documentation confirming disease is hormone-receptor positive and HER2-negative; and
  - 1.3 Patient has an ECOG performance score of 0-2; and
  - 1.4 Any of the following:
    - 1.4.1 Disease has relapsed or progressed during prior endocrine therapy; or
    - 1.4.2 Both:
      - 1.4.2.1 Patient is amenorrhoeic, either naturally or induced, with endocrine levels consistent with a postmenopausal or without menstrual-potential state; and
      - 1.4.2.2 Patient has not received prior systemic endocrine treatment for metastatic disease; or

1.4.3 Both:

- 1.4.3.1 Patient commenced treatment with ribociclib in combination with an endocrine partner prior to 1 July 2024; and
- 1.4.3.2 There is no evidence of progressive disease; and
- 1.5 Treatment to be used in combination with an endocrine partner; and
- 1.6 Patient has not received prior funded treatment with a CDK4/6 inhibitor; or
- 2 All of the following:
  - 2.1 Patient has an active Special Authority approval for palbociclib; and
  - 2.2 Patient has experienced a grade 3 or 4 adverse reaction to palbociclib that cannot be managed by dose reductions and requires treatment discontinuation; and
  - 2.3 Treatment must be used in combination with an endocrine partner; and
  - 2.4 There is no evidence of progressive disease since initiation of palbociclib.

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 Treatment must be used in combination with an endocrine partner; and
- 2 There is no evidence of progressive disease since initiation of ribociclib.

RUXOLITINIB - Special Authority see SA1890 on the next page - Retail pharmacy

Wastage claimable

Tab 5 mg2,500.00	56	🖌 Jakavi
Tab 10mg5,000.00	56	🖌 Jakavi
Tab 15 mg5,000.00	56	🖌 Jakavi
Tab 20 mg5,000.00	56	🗸 Jakavi

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	1	Manufacturer	

#### ⇒SA1890 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has primary myelofibrosis or post-polycythemia vera myelofibrosis or post-essential thrombocythemia myelofibrosis; and
- 2 Either:
  - 2.1 A classification of risk of intermediate-2 or high-risk myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS; or
  - 2.2 Both:
    - 2.2.1 A classification of risk of intermediate-1 myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS; and
    - 2.2.2 Patient has severe disease-related symptoms that are resistant, refractory or intolerant to available therapy; and
- 3 A maximum dose of 20 mg twice daily is to be given.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 A maximum dose of 20 mg twice daily is to be given.

SUNITINIB - Special Authority see SA2430 below - Retail pharmacy

Cap 12.5 mg	 28	<ul> <li>Sunitinib Pfizer</li> </ul>
Cap 25 mg	 28	<ul> <li>Sunitinib Pfizer</li> </ul>
Cap 50 mg	 28	<ul> <li>Sunitinib Pfizer</li> </ul>

### ➡SA2430 Special Authority for Subsidy

**Initial application** — (RCC) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 The patient has metastatic renal cell carcinoma of predominantly clear cell histology; and
- 2 Any of the following:
  - 2.1 The patient is treatment naive; or
  - 2.2 The patient has only received prior cytokine treatment; or
  - 2.3 The patient has only received prior treatment with an investigational agent within the confines of a bona fide clinical trial which has Ethics Committee approval; or
  - 2.4 Both:
    - 2.4.1 The patient has discontinued pazopanib within 3 months of starting treatment due to intolerance; and
    - 2.4.2 The cancer did not progress whilst on pazopanib; and
- 3 The patient has an ECOG performance score of 0-2; and
- 4 Sunitinib to be used for a maximum of 2 cycles.

**Initial application** — **(GIST)** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal tumour (GIST); and
- 2 Either:
  - 2.1 The patient's disease has progressed following treatment with imatinib; or
  - 2.2 The patient has documented treatment-limiting intolerance, or toxicity to, imatinib.

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

Subsidy		Fully	Brand or	
(Manufacturer's F	Price) Sub	osidised	Generic	
\$	Per	~	Manufacturer	

continued...

**Renewal** — (RCC) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months where there is no evidence of disease progression.

**Renewal** — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

The patient has responded to treatment or has stable disease as determined by Choi's modified CT response evaluation criteria as follows:

- 1 Any of the following:
  - 1.1 The patient has had a complete response (disappearance of all lesions and no new lesions); or
  - 1.2 The patient has had a partial response (a decrease in size of 10% or more or decrease in tumour density in Hounsfield Units (HU) of 15% or more on CT and no new lesions and no obvious progression of non measurable disease); or
  - 1.3 The patient has stable disease (does not meet criteria the two above) and does not have progressive disease and no symptomatic deterioration attributed to tumour progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: It is recommended that response to treatment be assessed using Choi's modified CT response evaluation criteria (J Clin Oncol, 2007, 25:1753-1759). Progressive disease is defined as either: an increase in tumour size of 10% or more and not meeting criteria of partial response (PR) by tumour density (HU) on CT; or: new lesions, or new intratumoral nodules, or increase in the size of the existing intratumoral nodules.

Renewal — (GIST pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal (GIST); and
- 2 The patient is clinically benifiting from treatment and continued treatment remains appropriate; and
- 3 Sunitinib is to be discontinued at progression; and
- 4 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

### **Endocrine Therapy**

For GnRH ANALOGUES - refer to HORMONE PREPARATIONS, Trophic Hormones, page 87

ABIRATERONE ACETATE - Retail pharmacy-Specialist - Special Authority see SA2118 below

Wastage claimable

#### ⇒SA2118 Special Authority for Subsidy

**Initial application** only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has prostate cancer; and
- 2 Patient has metastases; and
- 3 Patient's disease is castration resistant; and
- 4 Either:

#### 4.1 All of the following:

- 4.1.1 Patient is symptomatic; and
- 4.1.2 Patient has disease progression (rising serum PSA) after second line anti-androgen therapy; and
- 4.1.3 Patient has ECOG performance score of 0-1; and
- 4.1.4 Patient has not had prior treatment with taxane chemotherapy; or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 4.2 All of the following:
  - 4.2.1 Patient's disease has progressed following prior chemotherapy containing a taxane; and
  - 4.2.2 Patient has ECOG performance score of 0-2; and
  - 4.2.3 Patient has not had prior treatment with abiraterone.

Renewal — (abiraterone acetate) only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Significant decrease in serum PSA from baseline; and
- 2 No evidence of clinical disease progression; and
- 3 No initiation of taxane chemotherapy with abiraterone; and
- 4 The treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 2 Abiraterone acetate to be discontinued at progression; and
- 3 No initiation of taxane chemotherapy with abiraterone; and
- 4 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

#### BICALUTAMIDE

Tab 50 mg	4.18	28	<ul> <li>Binarex</li> </ul>
FLUTAMIDE			
Tab 250 mg		90	Prostacur S29
C C	119.50	100	<ul> <li>Flutamin</li> </ul>
FULVESTRANT – Retail pharmacy-Specialist – Special	Authority see SA1895 bel	ow	
Inj 50 mg per ml, 5 ml prefilled syringe		2	Faslodex

#### ➡SA1895 Special Authority for Subsidy

**Initial application** only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has oestrogen-receptor positive locally advanced or metastatic breast cancer; and
- 2 Patient has disease progression following prior treatment with an aromatase inhibitor or tamoxifen for their locally advanced or metastatic disease; and
- 3 Treatment to be given at a dose of 500 mg monthly following loading doses; and
- 4 Treatment to be discontinued at disease progression.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains appropriate and patient is benefitting from treatment; and
- 2 Treatment to be given at a dose of 500 mg monthly; and
- 3 There is no evidence of disease progression.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
OCTREOTIDE				
Inj 50 mcg per ml, 1 ml vial	27.58	5	1	Omega S29
Inj 100 mcg per ml, 1 ml vial		5	1	Omega S29
Inj 500 mcg per ml, 1 ml vial		5	1	Omega S29
Inj 50 mcg per ml, 1 ml ampoule		5	~	Max Health
Inj 100 mcg per ml, 1 ml ampoule		5	1	Octreotide GH S29 Max Health Octreotide GH S29
Inj 500 mcg per ml, 1 ml ampoule	113.10	5	1 1	Sun Pharma 529 Max Health Octreotide GH 529 Sun Pharma 529
TAMOXIFEN CITRATE * Tab 10 mg * Tab 20 mg		60 60	1 1	

### Long-acting Somatostatin Analogues

### ➡SA2445 Special Authority for Subsidy

**Initial application — (Malignant Bowel Obstruction)** from any relevant practitioner. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 The patient has nausea* and vomiting* due to malignant bowel obstruction*; and
- 2 Treatment with antiemetics, rehydration, antimuscarinic agents, corticosteroids and analgesics for at least 48 hours has not been successful; and
- 3 Treatment to be given for up to 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 3 months where the treatment remains appropriate and the patient is benefiting from treatment.

**Initial application** — (Acromegaly) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has acromegaly; and
- 2 Either:
  - 2.1 Treatment with surgery and radiotherapy is not suitable or was unsuccessful; or
  - 2.2 Treatment is for an interim period while awaiting the beneficial effects of radiotherapy; and
- 3 Treatment with a dopamine agonist has been unsuccessful.

Renewal — (Acromegaly) from any relevant practitioner. Approvals valid for 2 years where iGF1 levels have decreased since starting treatment.

Note: In patients with acromegaly, treatment should be discontinued if IGF1 levels have not decreased 3 months after treatment. In patients treated with radiotherapy treatment should be withdrawn every 2 years, for 1 month, for assessment of remission. Treatment should be stopped where there is biochemical evidence of remission (normal IGF1 levels) following treatment withdrawal for at least 4 weeks

Initial application — (pre-operative acromegaly) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

Subsidy (Manufacturer's Price) \$	Fu Subsidis Per	ully sed	Brand or Generic Manufacturer
 Ψ	1.01	-	Manalaotaroi

continued...

- 1 Patient has acromegaly; and
- 2 Patient has a large pituitary tumour, greater than 10 mm at its widest; and
- 3 Patient is scheduled to undergo pituitary surgery in the next six months.

Initial application — (Other Indications) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 VIPomas and Glucagonomas for patients who are seriously ill in order to improve their clinical state prior to definitive surgery; or
- 2 Both:
  - 2.1 Gastrinoma; and
  - 2.2 Either:
    - 2.2.1 Surgery has been unsuccessful; or
    - 2.2.2 Patient has metastatic disease after treatment with H2 antagonist or proton pump inhibitors has been unsuccessful; or
- 3 Both:
  - 3.1 Insulinomas; and
  - 3.2 Surgery is contraindicated or has not been successful; or
- 4 For pre-operative control of hypoglycaemia and for maintenance therapy; or
- 5 Both:
  - 5.1 Carcinoid syndrome (diagnosed by tissue pathology and/or urinary 5HIAA analysis); and
  - 5.2 Disabling symptoms not controlled by maximal medical therapy.

Note: The use of a long-acting somatostatin analogue in patients with fistulae, oesophageal varices, miscellaneous diarrhoea and hypotension will not be funded under Special Authority

Renewal — (Other Indications) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

us page – Retail pharr	nacy	
	1	<ul> <li>Mytolac</li> </ul>
646.70	1	<ul> <li>Mytolac</li> </ul>
145 on the previous pa	<mark>ge</mark> – Retail	pharmacy
	1	<ul> <li>Sandostatin LAR</li> </ul>
	1	Sandostatin LAR
670.80	1	<ul> <li>Sandostatin LAR</li> </ul>

### Aromatase Inhibitors

ANASTROZOLE <b>*</b> Tab 1 mg	.4.39	30	✓ Anatrole
EXEMESTANE * Tab 25 mg	.9.86	30	✓ Pfizer Exemestane
LETROZOLE			
* Tab 2.5 mg	.4.36	28	<ul> <li>Accord S29</li> </ul>
	4.67	30	✓ Letrole

### Immunosuppressants

#### Cytotoxic Immunosuppressants

AZ	ATHIOPRINE			
*	Tab 25 mg	7.36	60	<ul> <li><u>Azamun</u></li> </ul>
*	Tab 50 mg	8.10	100	✓ <u>Azamun</u>

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

	Subsidy		Fully	Brand or
	(Manufacturer's Price	,	ubsidised	Generic
	\$	Per		Manufacturer
MYCOPHENOLATE MOFETIL				
Tab 500 mg		50	✓ (	Cellcept
Cap 250 mg		100	✓ (	Cellcept
Powder for oral liq 1 g per 5 ml - Subsidy by endorsement		165 ml OF	· <b>√</b> (	Cellcept
the prescription is endorsed accordingly. Fusion Proteins				• •
ETANERCEPT – Special Authority see SA2399 below – Retail p	harmacy			
Inj 25 mg		4	✓ E	Inbrel
Inj 25 mg autoinjector		4	🖌 E	nbrel
Inj 50 mg autoinjector		4	🖌 E	nbrel
Inj 50 mg prefilled syringe		4	🖌 E	nbrel
SA2399 Special Authority for Subsidy				

**Initial application** — (adult-onset Still's disease) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 Either:
    - 1.1.1 The patient has had an initial Special Authority approval for adalimumab for adult-onset Still's disease (AOSD); or
    - 1.1.2 The patient has been started on tocilizumab for AOSD in a Health NZ Hospital; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or tocilizumab; or
    - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or tocilizumab such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
  - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
  - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal anti-inflammatory drugs (NSAIDs) and methotrexate; and
  - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

**Renewal — (adult-onset Still's disease)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 Applicant is a rheumatologist; or
  - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 The patient has a sustained improvement in inflammatory markers and functional status.

Initial application — (ankylosing spondylitis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for ankylosing spondylitis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
 \$	Per 🗸	

continued...

- 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for ankylosing spondylitis; or
- 2 All of the following:
  - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis present for more than six months; and
  - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
  - 2.3 Patient has bilateral sacroiliitis demonstrated by plain radiographs, CT or MRI scan; and
  - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis: and
  - 2.5 Either:
    - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
    - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender (see Notes); and
  - 2.6 A Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of initial application.

Average normal chest expansion corrected for age and gender:

- 18-24 years Male: 7.0 cm; Female: 5.5 cm
- 25-34 years Male: 7.5 cm; Female: 5.5 cm
- 35-44 years Male: 6.5 cm; Female: 4.5 cm
- 45-54 years Male: 6.0 cm; Female: 5.0 cm
- 55-64 years Male: 5.5 cm; Female: 4.0 cm
- 65-74 years Male: 4.0 cm; Female: 4.0 cm

75+ years - Male: 3.0 cm; Female: 2.5 cm

**Renewal** — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

- All of the following:
  - 1 Either:
    - 1.1 Applicant is a rheumatologist; or
    - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
  - 2 Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
  - 3 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
  - 4 Etanercept to be administered at doses no greater than 50 mg every 7 days.

**Initial application — (polyarticular course juvenile idiopathic arthritis)** only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
- 1.2 Either:
  - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsid	ised	Generic
\$	Per	1	Manufacturer

continued...

- 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for polyarticular course JIA; or
- 2 All of the following:
  - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
  - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
  - 2.3 Any of the following:
    - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
    - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
    - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Renewal — (polyarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
  - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Fither:

1 Both:

- The patient has had an initial Special Authority approval for adalimumab for oligoarticular course juvenile idiopathic arthritis (JIA); and
- 1.2 Either:
  - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
  - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for oligoarticular course JIA; or
- 2 All of the following:
  - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
  - 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
  - 2.3 Any of the following:
    - 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
    - 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose); or
    - 2.3.3 High disease activity (cJADAS10 score greater than 4) after a 6-month trial of methotrexate.

Renewal — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

Subsidy		Fully	Brand or
(Manufacturer's Price)	9	Subsidised	Generic
\$	Per	1	Manufacturer

continued...

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
  - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

### Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab or secukinumab for psoriatic arthritis; and 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab or secukinumab; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab or secukinumab to meet the renewal criteria for adalimumab or secukinumab for psoriatic arthritis; or
- 2 All of the following:
  - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
  - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
  - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
  - 2.4 Either:
    - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
    - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
  - 2.5 Any of the following:
    - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
    - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
    - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

**Renewal** — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

1 Either:

- 1.1 Applicant is a rheumatologist; or
- 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidised		Generic
\$	Per	✓	Manufacturer

#### continued...

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Indications marked with * are unapproved indications.

**Renewal** — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

Initial application — (Arthritis - rheumatoid) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for rheumatoid arthritis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects; or
    - 1.2.2 The patient has received insufficient benefit to meet the renewal criteria for rheumatoid arthritis; or
- 2 All of the following:
  - 2.1 Patient has had rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
  - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
  - 2.3 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
  - 2.4 Patient has tried and not responded to at least three months of methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses unless contraindicated); and
  - 2.5 Either:
    - 2.5.1 Patient has tried and not responded to at least three months of methotrexate in combination with the maximum tolerated dose of ciclosporin; or
    - 2.5.2 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with methotrexate; and
  - 2.6 Either:
    - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
    - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip.

**Renewal — (Arthritis - rheumatoid)** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:

Subsidy	Fully	Brand or	_
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

- 2.1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

**Initial application — (severe chronic plaque psoriasis)** only from a dermatologist or any relevant practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria: Either:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for adalimumab for severe chronic plaque psoriasis; and
- 1.2 Either:
  - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
  - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe chronic plaque psoriasis; or
- 2 All of the following:
  - 2.1 Any of the following:
    - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
    - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; or
    - 2.1.3 Patient has severe chronic localised genital or flexural plaque psoriasis where the plaques or lesions have been present for at least 6 months from the time of initial diagnosis, and with a Dermatology Life Quality Index (DLQI) score greater than 10; and
  - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
  - 2.3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
  - 2.4 The most recent PASI or DLQI assessment is no more than 1 month old at the time of application.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand, foot, genital or flexural areas at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and for the face, palm of a hand or sole of a foot the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 Any of the following:

1.1 Both:

- 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
- 1.1.2 Either:
  - 1.1.2.1 Following each prior etanercept treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-treatment baseline value; or

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Su	lbsidised	Generic	
\$	Per	1	Manufacturer	

continued...

- 1.1.2.2 Following each prior etanercept treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline value; or
- 1.2 Both:
  - 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
  - 1.2.2 Either:
    - 1.2.2.1 Following each prior etanercept treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
    - 1.2.2.2 Following each prior etanercept treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; or

#### 1.3 Both:

- 1.3.1 Patient had severe chronic localised genital or flexural plaque psoriasis at the start of treatment; and
- 1.3.2 Either:
  - 1.3.2.1 The patient has experienced a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; or
  - 1.3.2.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing etanercept; and
- 2 Etanercept to be administered at doses no greater than 50 mg every 7 days.
- Note: A treatment course is defined as a minimum of 12 weeks of etanercept treatment

Initial application — (undifferentiated spondyloarthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undifferentiated peripheral spondyloarthritis* with active peripheral joint arthritis in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day (or maximum tolerated dose); and
- 4 Patient has tried and not responded to at least three months of leflunomide at a dose of up to 20 mg daily (or maximum tolerated dose); and
- 5 Any of the following:
  - 5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
  - 5.2 Patient has an elevated enythrocyte sedimentation rate (ESR) greater than 25 mm per hour measured no more than one month prior to the date of this application; or
  - 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Note: Indications marked with * are unapproved indications.

**Renewal** — (undifferentiated spondyloarthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 Applicant is a rheumatologist; or
  - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:

Subsidy (Manufacturer's Price)	Subs	Fully sidised	Brand or Generic
 \$	Per	~	Manufacturer

continued...

- 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg dose every 7 days.

### **Immune Modulators**

ANTITHYMOCYTE GLOBULIN (EQUINE) - PCT only - S		-	( 170.14
Inj 50 mg per ml, 5 ml	4,439.17	5	🗸 ATGAM
BACILLUS CALMETTE-GUERIN (BCG) VACCINE - PCT	only – Specialist		
Subsidised only for bladder cancer.	<i>y</i> 1		
Inj 2-8 × 100 million CFU		1	<ul> <li>OncoTICE</li> </ul>
Inj 40 mg per ml, vial		3	SII-Onco-BCG S29

### **Monoclonal Antibodies**

ADALIMUMAB (AMGEVITA) - Special Authority see SA240	0 below – Retail pharma	су	
Inj 20 mg per 0.4 ml prefilled syringe		1	<ul> <li>Amgevita</li> </ul>
Inj 40 mg per 0.8 ml prefilled pen		2	<ul> <li>Amgevita</li> </ul>
Inj 40 mg per 0.8 ml prefilled syringe		2	<ul> <li>Amgevita</li> </ul>

#### ⇒SA2400 Special Authority for Subsidy

**Initial application** — (Behcet's disease - severe) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Both:

- 1 The patient has severe Behcet's disease* that is significantly impacting the patient's quality of life; and
- 2 Either:
  - 2.1 The patient has severe ocular, neurological, and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s); or
  - 2.2 The patient has severe gastrointestinal, rheumatological, and/or mucocutaneous symptoms and has not responded adequately to two or more treatments appropriate for the particular symptom(s).

Note: Indications marked with * are unapproved indications.

Initial application — (Hidradenitis suppurativa) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has hidradenitis suppurativa Hurley Stage II or Hurley Stage III lesions in distinct anatomic areas; and
- 2 Patient has tried, but had an inadequate response to at least a 90 day trial of systemic antibiotics or has demonstrated intolerance to or has contraindications for systemic antibiotics; and
- 3 Patient has 3 or more active lesions; and
- 4 The patient has a DLQI of 10 or more and the assessment is no more than 1 month old at time of application.

Renewal — (Hidradenitis suppurativa) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline; and
- 2 The patient has a DLQI improvement of 4 or more from baseline.

Initial application - (Plaque psoriasis - severe chronic) only from a dermatologist or any relevant practitioner on the

Subsid	dy Fu	ly Brand or
(Manufacture)	r's Price) Subsidis	ed Generic
\$	Per	Manufacturer

### continued...

recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria: Either:

- 1 Both:
  - 1.1 Patient has had an initial Special Authority approval for etanercept for severe chronic plaque psoriasis; and
  - 1.2 Either:
    - 1.2.1 Patient has experienced intolerable side effects; or
    - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for etanercept for severe chronic plaque psoriasis; or
- 2 All of the following:
  - 2.1 Any of the following:
    - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a PASI score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
    - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; or
    - 2.1.3 Patient has severe chronic localised genital or flexural plaque psoriasis where the plaques or lesions have been present for at least 6 months from the time of initial diagnosis, and with a Dermatology Life Quality Index (DLQI) score greater than 10; and
  - 2.2 Patient has tried, but had an inadequate response to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
  - 2.3 A PASI assessment or DLQI assessment has been completed for at least the most recent prior treatment course but no longer than 1 month following cessation of each prior treatment course and is no more than 1 month old at the time of application.

# Renewal — (Plaque psoriasis - severe chronic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 Both:
  - 1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced a 75% or more reduction in PASI score, or is sustained at this level, when compared with the pre-treatment baseline value; or
    - 1.2.2 The patient has a DLQI improvement of 5 or more, when compared with the pre-treatment baseline value; or
- 2 Both:
  - 2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
  - 2.2 Either:
    - 2.2.1 The patient has experienced reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
    - 2.2.2 The patient has experienced reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre treatment baseline value; or
- 3 Both:
  - 3.1 Patient had severe chronic localised genital or flexural plaque psoriasis at the start of treatment; and
  - 3.2 Either:
    - 3.2.1 The patient has experienced a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; or

Subsidy		Fully	Brand or	
(Manufacturer's Pric	ce)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

3.2.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing adalimumab.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

- Both:
  - 1 Patient has pyoderma gangrenosum*; and
  - 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporin, azathioprine, or methotrexate) and has not received an adequate response.
- Note: Indications marked with * are unapproved indications.

Initial application — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active Crohn's disease; and
- 2 Any of the following:
  - 2.1 Patient has a CDAI score of greater than or equal to 300, or HBI score of greater than or equal to 10; or
  - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
  - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
  - 2.4 Patient has an ileostomy or colostomy and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Renewal — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 CDAI score has reduced by 100 points from the CDAI score, or HBI score has reduced by 3 points, from when the patient was initiated on adalimumab; or
- 2 CDAI score is 150 or less, or HBI is 4 or less; or
- 3 The patient has demonstrated an adequate response to treatment, but CDAI score and/or HBI score cannot be assessed.

Initial application — (Crohn's disease - children) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

1 Paediatric patient has active Crohn's disease; and

2 Either:

- 2.1 Patient has a PCDAI score of greater than or equal to 30; or
- 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Renewal — (Crohn's disease - children) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab; or
- 2 PCDAI score is 15 or less; or
- 3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed.

**Initial application** — (Crohn's disease - fistulising) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

1 Patient has confirmed Crohn's disease; and

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	~	Manufacturer

continued...

- 2 Any of the following:
  - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
  - 2.2 Patient has one or more rectovaginal fistula(e); or
  - 2.3 Patient has complex peri-anal fistula; and
- 3 A Baseline Fistula Assessment has been completed and is no more than 1 month old at the time of application.

Renewal — (Crohn's disease - fistulising) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 The number of open draining fistulae have decreased from baseline by at least 50%; or
- 2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain.

**Initial application — (Ocular inflammation - chronic)** from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

#### Either:

- 1 The patient has had an initial Special Authority approval for infliximab for chronic ocular inflammation; or
- 2 Both:
  - 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
  - 2.2 Any of the following:
    - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective; or
    - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
    - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Renewal — (Ocular inflammation - chronic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

- Any of the following:
  - 1 The patient has had a good clinical response following 12 weeks' initial treatment; or
  - 2 Following each 2 year treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
  - 3 Following each 2 year treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

**Initial application — (Ocular inflammation - severe)** from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

1 Patient has had an initial Special Authority approval for infliximab for severe ocular inflammation; or

2 Both:

- 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
- 2.2 Any of the following:
  - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
  - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
  - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidi	sed	Generic
 \$	Per	✓	

#### continued...

Renewal — (Ocular inflammation - severe) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 2 year treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
- 3 Following each 2 year treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.</p>

Initial application — (ankylosing spondylitis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 Patient has had an initial Special Authority approval for etanercept for ankylosing spondylitis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects; or
    - 1.2.2 The patient has received insufficient benefit to meet the renewal criteria for ankylosing spondylitis; or
- 2 All of the following:
  - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis for more than six months; and
  - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
  - 2.3 Patient has bilateral sacroiliitis demonstrated by radiology imaging; and
  - 2.4 Patient has not responded adequately to treatment with two or more NSAIDs, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
  - 2.5 Either:
    - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following BASMI measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
    - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender; and
  - 2.6 A BASDAI of at least 6 on a 0-10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment and is no more than 1 month old at the time of application.

Renewal — (ankylosing spondylitis) from any relevant practitioner. Approvals valid for 2 years where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less.

Initial application — (Arthritis - oligoarticular course juvenile idiopathic) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Fither:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for etanercept for oligoarticular course juvenile idiopathic arthritis (JIA); and
- 1.2 Either:
  - 1.2.1 Patient has experienced intolerable side effects; or
  - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for oligoarticular course JIA; or
- 2 All of the following:
  - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
  - 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

Subsidy (Manufacturer's Price)	:	Fully Subsidised	Brand or Generic
\$	Per	1	Manufacturer

#### continued...

- 2.3 Either:
  - 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
  - 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose).

Renewal — (Arthritis - oligoarticular course juvenile idiopathic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

- Either:
  - 1 Following initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
  - 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - polyarticular course juvenile idiopathic) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 Patient has had an initial Special Authority approval for etanercept for polyarticular course juvenile idiopathic arthritis (JIA); and
  - 1.2 Either:
    - 1.2.1 Patient has experienced intolerable side effects; or
    - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for polyarticular course JIA; or
- 2 All of the following:
  - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
  - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
  - 2.3 Any of the following:
    - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
    - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
    - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Renewal — (Arthritis - polyarticular course juvenile idiopathic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - psoriatic) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

#### Either:

1 Both:

- 1.1 Patient has had an initial Special Authority approval for etanercept or secukinumab for psoriatic arthritis; and 1.2 Either:
  - 1.2.1 The patient has experienced intolerable side effects; or
  - 1.2.2 The patient has received insufficient benefit from to meet the renewal criteria for psoriatic arthritis; or
- 2 All of the following:

	Subsidy	F	ully	Brand or
(Man	ufacturer's Price)	Subsidis	sed	Generic
	\$	Per	✓	Manufacturer

continued...

- 2.1 Patient has had active psoriatic arthritis for six months duration or longer; and
- 2.2 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
- 2.3 Patient has tried and not responded to at least three months of sulfasalazine or leflunomide at maximum tolerated doses (unless contraindicated); and
- 2.4 Either:
  - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
  - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2.5 Any of the following:
  - 2.5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
  - 2.5.2 Patient has an ESR greater than 25 mm per hour; or
  - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (Arthritis - psoriatic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in swollen joint count from baseline and a clinically significant response in the opinion of the physician; or
- 2 Patient demonstrates at least a continuing 30% improvement in swollen joint count from baseline and a clinically significant response in the opinion of the treating physician.

Initial application — (Arthritis - rheumatoid) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for etanercept for rheumatoid arthritis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects; or
    - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for rheumatoid arthritis; or
- 2 All of the following:
  - 2.1 Patient has had rheumatoid arthritis (either confirmed by radiology imaging, or the patient is CCP antibody positive) for six months duration or longer; and
  - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
  - 2.3 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
  - 2.4 Patient has tried and not responded to at least three months of methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate at maximum tolerated doses (unless contraindicated); and
  - 2.5 Either:
    - 2.5.1 Patient has tried and not responded to at least three months of methotrexate in combination with the maximum tolerated dose of ciclosporin; or
    - 2.5.2 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with methotrexate; and
  - 2.6 Either:
    - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or

(	Subsidy	Fully	Brand or
(Manufa	acturer's Price)	Subsidised	Generic
	\$ Per	· · ·	Manufacturer

continued...

2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip.

**Renewal** — (Arthritis - rheumatoid) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Initial application — (Still's disease - adult-onset (AOSD)) only from a rheumatologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for etanercept and/or tocilizumab for AOSD; and
  - 1.2 Either:
    - 1.2.1 Patient has experienced intolerable side effects from etanercept and/or tocilizumab; or
    - 1.2.2 Patient has received insufficient benefit from at least a three-month trial of etanercept and/or tocilizumab; or
- 2 All of the following:
  - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria; and
  - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, NSAIDs and methotrexate; and
  - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Initial application — (ulcerative colitis) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active ulcerative colitis; and
- 2 Either:
  - 2.1 Patient's SCCAI score is greater than or equal to 4; or
  - 2.2 Patient's PUCAI score is greater than or equal to 20; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from prior therapy with immunomodulators and systemic corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 The SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on biologic therapy; or
- 2 The PUCAI score has reduced by 10 points or more from the PUCAI score when the patient was initiation on biologic therapy.

**Initial application — (undifferentiated spondyloarthritis)** only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undifferentiated peripheral spondyloarthritis* with active peripheral joint arthritis in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2 Patient has tried and not responded to at least three months of each of methotrexate, sulfasalazine and leflunomide, at maximum tolerated doses (unless contraindicated); and
- 3 Any of the following:

Subsidy (Manufacturer's Price)	Su	Fully bsidised	Brand or Generic
 \$	Per	1	Manufacturer

continued...

- 3.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
- 3.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application; or
- 3.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Note: Indications marked with * are unapproved indications

Renewal — (undifferentiated spondyloarthritis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response in the opinion of the treating physician.

**Initial application** — (inflammatory bowel arthritis – axial) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has axial inflammatory pain for six months or more; and
- 3 Patient is unable to take NSAIDs; and
- 4 Patient has unequivocal sacroiliitis demonstrated by radiological imaging or MRI; and
- 5 Patient has not responded adequately to prior treatment consisting of at least 3 months of an exercise regime supervised by a physiotherapist; and
- 6 A BASDAI of at least 6 on a 0-10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment.

**Renewal — (inflammatory bowel arthritis – axial)** from any relevant practitioner. Approvals valid for 2 years where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less.

Initial application — (inflammatory bowel arthritis – peripheral) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has active arthritis in at least four joints from the following: hip, knee, ankle, subtalar, tarsus, forefoot, wrist, elbow, shoulder, sternoclavicular; and
- 3 Patient has tried and not experienced a response to at least three months of methotrexate, or azathioprine at a maximum tolerated dose (unless contraindicated); and
- 4 Patient has tried and not experienced a response to at least three months of sulfasalazine at a maximum tolerated dose (unless contraindicated); and
- 5 Any of the following:
  - 5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
  - 5.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application; or
  - 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

**Renewal — (inflammatory bowel arthritis – peripheral)** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

Subsidy	<u></u>	Fully	Brand or
(Manufacturer's	,	idised	Generic
<u>م</u>	Per	•	Manufacturer

continued...

- 1 Following initial treatment, patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 Patient has experienced at least a continuing 30% improvement in active joint count from baseline in the opinion of the treating physician.

ADALIMUMAB (HUMIRA - ALTERNATIVE BRAND) - Special Authority see SA2157 below - Retail pharmacy

Inj 20 mg per 0.2 ml prefilled syringe1,599.96	2	🗸 Humira
Inj 40 mg per 0.4 ml prefilled pen1,599.96	2	<ul> <li>HumiraPen</li> </ul>
Inj 40 mg per 0.4 ml prefilled syringe1,599.96	2	<ul> <li>Humira</li> </ul>

#### ⇒SA2157 Special Authority for Subsidy

**Initial application** — (Behcet's disease – severe) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Behcet's disease – severe) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient has had a good clinical response to treatment with measurably improved quality of life; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Hidradenitis suppurativa) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Either:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 7 days. Fortnightly dosing has been considered.

**Renewal — (Hidradenitis suppurativa)** only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline; and
- 2 The patient has a Dermatology Quality of Life Index improvement of 4 or more from baseline; and
- 3 Adalimumab is to be administered at doses no greater than 40mg every 7 days. Fortnightly dosing has been considered.

Initial application — (Psoriasis - severe chronic plaque) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🖌	Manufacturer

continued...

- 1 Either:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Psoriasis - severe chronic plaque) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both: 1 Fither:

- - 1.1 Both:
    - 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
    - 1.1.2 Either:
      - 1.1.2.1 Following each prior adalimumab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-adalimumab treatment baseline value; or
      - 1.1.2.2 Following each prior adalimumab treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline value; or
  - 1.2 Both:
    - 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
    - 1.2.2 Either:
      - 1.2.2.1 Following each prior adalimumab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
      - 1.2.2.2 Following each prior adalimumab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-adalimumab treatment baseline value; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

**Initial application — (Pyoderma gangrenosum)** only from a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 A maximum of 8 doses.

**Renewal** — (Pyoderma gangrenosum) only from a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient has demonstrated clinical improvement and continues to require treatment; and
- 2 A maximum of 8 doses.

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	1	Manufacturer

continued...

Initial application — (Crohn's disease - adult) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Any of the following:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevitat; or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
  - 1.3 Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment; and
  - 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
  - 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Crohn's disease - adult) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Any of the following:
  - 1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on adalimumab; or
  - 1.2 CDAI score is 150 or less; or
  - 1.3 The patient has demonstrated an adequate response to treatment, but CDAI score cannot be assessed; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Crohn's disease - children) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Any of the following:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
  - 1.3 Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Crohn's disease - children) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Any of the following:
  - 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab; or
  - 1.2 PCDAI score is 15 or less; or
  - 1.3 The patient has demonstrated an adequate response to treatment, but PCDAI score cannot be assessed; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Crohn's disease - fistulising) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

1 Any of the following:

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

- 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
- 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
- 1.3 Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Crohn's disease - fistulising) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

- 1 Either:
  - 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
  - 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

**Initial application — (Ocular inflammation – chronic)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with Amgevita, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
  - 1.3 Patient has uveitis and is considered to be at risk of vision loss if they were to change treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Ocular inflammation – chronic) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 The patient has had a good clinical response following 12 weeks' initial treatment; or
  - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
  - 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

**Initial application** — (Ocular inflammation – severe) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or

Subsidy	Fu	y Brand or
(Manufacturer's	Price) Subsidise	d Generic
\$	Per	Manufacturer

continued...

- 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with Amgevita, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
- 1.3 Patient has uveitis and is considered to be at risk of vision loss if they were to change treatment; and
- Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
   Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Ocular inflammation – severe) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 The patient has had a good clinical response following 3 initial doses; or
  - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
  - 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Either:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita); and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

**Initial application** — (Arthritis – oligoarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meting the following criteria:

All of the following:

- 1 Either:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

Renewal — (Arthritis – oligoarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months where the patient demonstrates at

Subsidy		Fully	Brand or	
(Manufacturer's Price)	5	Subsidised	Generic	
\$	Per	1	Manufacturer	

continued...

least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

**Initial application** — (Arthritis - polyarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

Renewal — (Arthritis - polyarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months where the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - psoriatic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Either:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Arthritis - psoriatic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Arthritis – rheumatoid) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

1 Either:

- 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
- 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and 4 Either:
  - 4.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

4.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

Renewal — (Arthritis – rheumatoid) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 2 Either:
  - 2.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or
  - 2.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

Initial application — (Still's disease – adult-onset (AOSD)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Either:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

Renewal — (Still's disease – adult-onset (AOSD)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months where the patient has demonstrated a sustained improvement in inflammatory markers and functional status.

AFLIBERCEPT – Special Authority see SA1772 below – Retail pharmacy

Inj 40 mg per ml, 0.1 ml vial...... 1,250.00 1 🗸 Eylea

#### ⇒SA1772 Special Authority for Subsidy

**Initial application** — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 All of the following:
  - 1.1 Any of the following:
    - 1.1.1 Wet age-related macular degeneration (wet AMD); or
    - 1.1.2 Polypoidal choroidal vasculopathy; or
    - 1.1.3 Choroidal neovascular membrane from causes other than wet AMD; and
  - 1.2 Either:
    - 1.2.1 The patient has developed severe endophthalmitis or severe posterior uveitis following treatment with bevacizumab; or
    - 1.2.2 There is worsening of vision or failure of retina to dry despite three intraocular injections of bevacizumab four weeks apart; and
  - 1.3 There is no structural damage to the central fovea of the treated eye; and
  - 1.4 Patient has not previously been treated with ranibizumab for longer than 3 months; or
- 2 Either:
  - 2.1 Patient has current approval to use ranibizumab for treatment of wAMD and was found to be intolerant to ranibizumab within 3 months; or
  - 2.2 Patient has previously* (*before June 2018) received treatment with ranibizumab for wAMD and disease was stable

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

while on treatment.

**Initial application** — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has centre involving diabetic macular oedema (DMO); and
- 2 Patient's disease is non responsive to 4 doses of intravitreal bevacizumab when administered 4-6 weekly; and
- 3 Patient has reduced visual acuity between 6/9 6/36 with functional awareness of reduction in vision; and
- 4 Patient has DMO within central OCT (ocular coherence tomography) subfield > 350 micrometers; and
- 5 There is no centre-involving sub-retinal fibrosis or foveal atrophy.

**Renewal — (wet age related macular degeneration)** only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Documented benefit must be demonstrated to continue; and
- 2 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 3 There is no structural damage to the central fovea of the treated eye.

Renewal — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 There is stability or two lines of Snellen visual acuity gain; and
- 2 There is structural improvement on OCT scan (with reduction in intra-retinal cysts, central retinal thickness, and sub-retinal fluid); and
- 3 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 4 There is no centre-involving sub-retinal fibrosis or foveal atrophy; and
- 5 After each consecutive 12 months treatment with (2nd line anti-VEGF agent), patient has retrialled with at least one injection of bevacizumab and had no response.

BENRALIZUMAB – Special Authority see SA2151 below – Retail pharmacy

#### ⇒SA2151 Special Authority for Subsidy

**Initial application** — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Patient must have a diagnosis of severe eosinophilic asthma documented by a respiratory physician or clinical immunologist; and
- 3 Conditions that mimic asthma eg. vocal cord dysfunction, central airway obstruction, bronchiolitis etc. have been excluded; and
- 4 Patient has a blood eosinophil count of greater than 0.5 × 10°9 cells/L in the last 12 months; and
- 5 Patient must be adherent to optimised asthma therapy including inhaled corticosteroids (equivalent to at least 1000 mcg per day of fluticasone propionate) plus long-acting beta-2 agonist, or budesonide/formoterol as part of the anti-inflammatory reliever therapy plus maintenance regimen, unless contraindicated or not tolerated; and
- 6 Either:
  - 6.1 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral corticosteroids; or
  - 6.2 Patient has received continuous oral corticosteroids of at least the equivalent of 10 mg per day over the previous 3 months; and
- 7 Treatment is not to be used in combination with subsidised mepolizumab; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 8 Patient has an Asthma Control Test (ACT) score of 10 or less. Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 52 weeks after the first dose to assess response to treatment; and
- 9 Either:
  - 9.1 Patient has not previously received an anti-IL5 biological therapy for their severe eosinophilic asthma; or
  - 9.2 Both:
    - 9.2.1 Patient was refractory or intolerant to previous anti-IL5 biological therapy; and
    - 9.2.2 Patient was not eligible to continue treatment with previous anti-IL5 biological therapy and discontinued within 12 months of commencing treatment.

Renewal — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 Either:
  - 2.1 Exacerbations have been reduced from baseline by 50% as a result of treatment with benralizumab; or
  - 2.2 Reduction in continuous oral corticosteroid use by 50% or by 10 mg/day while maintaining or improving asthma control.

BEVACIZUMAB - PCT only - Special Authority see SA2444 below

Inj 25 mg per ml, 4 ml vial		1	Vegzelma
Inj 25 mg per ml, 16 ml vial		1	<ul> <li>Vegzelma</li> </ul>
Inj 1 mg for ECP	0.71	1 mg	<ul> <li>Baxter</li> </ul>

#### ► SA2444 Special Authority for Subsidy

Initial application — (unresectable hepatocellular carcinoma) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Patient is currently on treatment with bevacizumab, and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
  - 2.1 Patient has locally advanced or metastatic, unresectable hepatocellular carcinoma; and
  - 2.2 Patient has preserved liver function (Child-Pugh A); and
  - 2.3 Transarterial chemoembolisation (TACE) is unsuitable; and
  - 2.4 Any of the following:
    - 2.4.1 Patient has not received prior systemic therapy for the treatment of hepatocellular carcinoma; or
    - 2.4.2 Patient received funded lenvatinib before 1 March 2025; or
    - 2.4.3 Both:
      - 2.4.3.1 Patient has experienced treatment-limiting toxicity from treatment with lenvatinib; and
      - 2.4.3.2 No disease progression since initiation of lenvatinib; and
  - 2.5 Patient has an ECOG performance status of 0-2; and
  - 2.6 To be given in combination with atezolizumab.

**Renewal — (unresectable hepatocellular carcinoma)** from any relevant practitioner. Approvals valid for 6 months where there is no evidence of disease progression.

Initial application — (advanced or metastatic ovarian cancer) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

1 Either:

- 1.1 The patient has FIGO Stage IV epithelial ovarian, fallopian tube, or primary peritoneal cancer; or
- 1.2 Both:

Subsidy (Manufacturer's Price)	Sub	Fully sidised	Brand or Generic
 \$	Per	~	Manufacturer

continued...

1.2.1 The patient has previously untreated advanced (FIGO Stage IIIB or IIIC) epithelial ovarian, fallopian tube, or primary peritoneal cancer; and

1.2.2 Either:

1.2.2.1 Debulking surgery is inappropriate; or

- 1.2.2.2 The cancer is sub-optimally debulked (maximum diameter of any gross residual disease greater than 1cm); and
- 2 Bevacizumab to be administered at a maximum dose of 7.5 mg/kg every three weeks; and
- 3 18 weeks concurrent treatment with chemotherapy is planned.

**Renewal — (advanced or metastatic ovarian cancer)** from any relevant practitioner. Approvals valid for 4 months where there is no evidence of disease progression.

**Initial application** — (Recurrent Respiratory Papillomatosis) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Maximum of 6 doses; and
- 2 The patient has recurrent respiratory papillomatosis; and
- 3 The treatment is for intra-lesional administration.

Renewal — (Recurrent Respiratory Papillomatosis) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Maximum of 6 doses; and
- 2 The treatment is for intra-lesional administration; and
- 3 There has been a reduction in surgical treatments or disease regrowth as a result of treatment.

Initial application — (Ocular Conditions) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 Ocular neovascularisation; or
- 2 Exudative ocular angiopathy.

#### BRENTUXIMAB VEDOTIN - PCT only - Special Authority see SA2289 below

#### ⇒SA2289 Special Authority for Subsidy

**Initial application** — (relapsed/refractory Hodgkin lymphoma) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

1 Either:

- 1.1 Both:
  - 1.1.1 Patient has relapsed/refractory CD30-positive Hodgkin lymphoma after two or more lines of chemotherapy; and
  - 1.1.2 Patient is ineligible for autologous stem cell transplant; or
- 1.2 Both:
  - 1.2.1 Patient has relapsed/refractory CD30-positive Hodgkin lymphoma; and
  - 1.2.2 Patient has previously undergone autologous stem cell transplant; and
- 2 Patient has not previously received funded brentuximab vedotin; and
- 3 Response to brentuximab vedotin treatment is to be reviewed after a maximum of 6 treatment cycles; and
- 4 Brentuximab vedotin to be administered at doses no greater than 1.8 mg/kg every 3 weeks.

# **Renewal — (relapsed/refractory Hodgkin lymphoma)** from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Si	ubsidised	Generic	
\$	Per	✓	Manufacturer	

#### continued...

All of the following:

- 1 Patient has achieved a partial or complete response to brentuximab vedotin after 6 treatment cycles; and
- 2 Treatment remains clinically appropriate and the patient is benefitting from treatment and treatment is being tolerated; and
- 3 Patient is to receive a maximum of 16 total cycles of brentuximab vedotin treatment.

# **Initial application** — (anaplastic large cell lymphoma) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has relapsed/refractory CD30-positive systemic anaplastic large cell lymphoma; and
- 2 Patient has an ECOG performance status of 0-1; and
- 3 Patient has not previously received brentuximab vedotin; and
- 4 Response to brentuximab vedotin treatment is to be reviewed after a maximum of 6 treatment cycles; and
- 5 Brentuximab vedotin to be administered at doses no greater than 1.8 mg/kg every 3 weeks.

# Renewal — (anaplastic large cell lymphoma) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has achieved a partial or complete response to brentuximab vedotin after 6 treatment cycles; and
- 2 Treatment remains clinically appropriate and the patient is benefitting from treatment and treatment is being tolerated; and
- 3 Patient is to receive a maximum of 16 total cycles of brentuximab vedotin treatment.

CASIRIVIMAB AND IMDEVIMAB - [Xpharm] - Special Authority see SA2096 below

Inj 120 mg per ml casirivimab, 11.1 ml vial (1) and inj 120 mg

per ml imdevimab, 11.1 ml vial (1)	0.00	1 OP	<ul> <li>Ronapreve</li> </ul>
------------------------------------	------	------	-------------------------------

#### ► SA2096 Special Authority for Subsidy

Initial application — (Treatment of profoundly immunocompromised patients) from any relevant practitioner. Approvals valid for 2 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed (or probable) COVID-19; and
- 2 The patient is in the community with mild to moderate disease severity*; and
- 3 Patient is profoundly immunocompromised** and is at risk of not having mounted an adequate response to vaccination against COVID-19 or is unvaccinated; and
- 4 Patient's symptoms started within the last 10 days; and
- 5 Patient is not receiving high flow oxygen or assisted/mechanical ventilation; and
- 6 Casirivimab and imdevimab is to be administered at a maximum dose of no greater than 2,400 mg.

Notes: * Mild to moderate disease severity as described on the Ministry of Health Website

** Examples include B-cell depletive illnesses or patients receiving treatment that is B-Cell depleting.

CETUXIMAB – PCT only – Specialist – Special Authority see SA2401 below		
Inj 5 mg per ml, 20 ml vial	1	🗸 Erbitux
Inj 5 mg per ml, 100 ml vial	1	🗸 Erbitux
Inj 1 mg for ECP	1 mg	<ul> <li>Baxter</li> </ul>

#### ⇒SA2401 Special Authority for Subsidy

**Initial application** — (head and neck cancer, locally advanced) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has locally advanced, non-metastatic, squamous cell cancer of the head and neck; and
- 2 Cisplatin is contraindicated or has resulted in intolerable side effects; and
- 3 Patient has an ECOG performance score of 0-2; and
- 4 To be administered in combination with radiation therapy.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

**Initial application** — (colorectal cancer, metastatic) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has metastatic colorectal cancer located on the left side of the colon (see Note); and
- 2 There is documentation confirming disease is RAS and BRAF wild-type; and
- 3 Patient has an ECOG performance score of 0-2; and
- 4 Patient has not received prior funded treatment with cetuximab; and
- 5 Either:
  - 5.1 Cetuximab is to be used in combination with chemotherapy; or
  - 5.2 Chemotherapy is determined to not be in the best interest of the patient based on clinician assessment.

Renewal — (colorectal cancer, metastatic) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where there is no evidence of disease progression.

Note: Left-sided colorectal cancer comprises of the distal one-third of the transverse colon, the splenic flexure, the descending colon, the sigmoid colon, or the rectum.

GEMTUZUMAB OZOGAMICIN – PCT only – Specialist – Special Authority see SA2269 below

Inj 5 mg vial ...... 12,973.00 1 🗸 Mylotarg

#### ► SA2269 Special Authority for Subsidy

Initial application only from a haematologist, paediatric haematologist or paediatric oncologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has not received prior chemotherapy for this condition; and
- 2 Patient has de novo CD33-positive acute myeloid leukaemia; and
- 3 Patient does not have acute promyelocytic leukaemia; and
- 4 Gemtuzumab ozogamicin will be used in combination with standard anthracycline and cytarabine (AraC); and
- 5 Patient is being treated with curative intent; and
- 6 Patient's disease risk has been assessed by cytogenetic testing to be good or intermediate; and
- 7 Patient must be considered eligible for standard intensive remission induction chemotherapy with standard anthracycline and cytarabine (AraC); and
- 8 Gemtuzumab ozogamicin to be funded for one course only (one dose at 3 mg per m² body surface area or up to 2 vials of 5 mg as separate doses).

Note: Acute myeloid leukaemia excludes acute promyelocytic leukaemia and acute myeloid leukaemia that is secondary to another haematological disorder (eg myelodysplasia or myeloproliferative disorder).

INFLIXIMAB – PCT only – Special Authority see SA2402 below		
Inj 100 mg	1	<ul> <li>Remicade</li> </ul>
Inj 1 mg for ECP4.40	1 mg	<ul> <li>Baxter</li> </ul>

#### ⇒SA2402 Special Authority for Subsidy

**Initial application** — (Crohn's disease (adults)) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active Crohn's disease; and
- 2 Any of the following:
  - 2.1 Patient has a CDAI score of greater than or equal to 300 or HBI score of greater than or equal to 10; or
  - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
  - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
  - 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sub	sidised	Generic
\$	Per	1	Manufacturer

continued...

3 Patient has tried but has experienced an inadequate response to, or intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Renewal — (Crohn's disease (adults)) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 CDAI score has reduced by 100 points from the CDAI score, or HBI score has reduced by 3 points, from when the patient was initiated on infliximab; or
  - 1.2 CDAI score is 150 or less, or HBI is 4 or less; or
  - 1.3 The patient has demonstrated an adequate response to treatment but CDAI score and/or HBI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (Crohn's disease (children)) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has active Crohn's disease; and
- 2 Either:
  - 2.1 Patient has a PCDAI score of greater than or equal to 30; or
  - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but experienced an inadequate response to, or intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Renewal — (Crohn's disease (children)) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on infliximab; or
  - 1.2 PCDAI score is 15 or less; or
  - 1.3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (Graft vs host disease) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has steroid-refractory acute graft vs. host disease of the gut.

**Initial application** — (Pulmonary sarcoidosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has life-threatening pulmonary sarcoidosis diagnosed by a multidisciplinary team that is refractory to other treatments.

Initial application — (acute fulminant ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 weeks for applications meeting the following criteria: Both:

- 1 Patient has acute, fulminant ulcerative colitis; and
- 2 Treatment with intravenous or high dose oral corticosteroids has not been successful.

**Initial application — (ankylosing spondylitis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria:

Subsidy		Fully	Brand or
(Manufacturer's Price	e) 5	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and 2 Either:
  - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
  - 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

**Renewal** — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Following 12 weeks of infliximab treatment, BASDAI has improved by 4 or more points from pre-infliximab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 3 Infliximab to be administered at doses no greater than 5 mg/kg every 6-8 weeks.

Initial application — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria: Either:

1 Boti

1 Both:

- 1.1 The patient has had an initial Special Authority approval for adalimumab for chronic ocular inflammation; and
- 1.2 Either:
  - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
  - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for chronic ocular inflammation; or

2 Both:

- 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
- 2.2 Any of the following:
  - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective; or
  - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
  - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Renewal — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (fistulising Crohn's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Su	Ibsidised	Generic	
\$	Per	1	Manufacturer	

continued...

- 1 Patient has confirmed Crohn's disease; and
- 2 Any of the following:
  - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
  - 2.2 Patient has one or more rectovaginal fistula(e); or
  - 2.3 Patent has complex peri-anal fistula.

Renewal — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Either:
  - 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
  - 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline (in the case of adult patients, as demonstrated by a reduction in the Fistula Assessment score), together with less induration and patient reported pain; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

#### **Initial application** — (neurosarcoidosis) only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with neurosarcoiosis by a multidisciplinary team; and
- 2 Patient has CNS involvement; and
- 3 Patient has steroid-refractory disease; and
- 4 Either:
  - 4.1 IV cyclophosphamide has been tried; or
  - 4.2 Treatment with IV cyclophosphamide is clinically inappropriate.

Renewal — (neurosarcoidosis) only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria:

Either:

- 1 A withdrawal period has been tried and the patient has relapsed; or
- 2 All of the following:
  - 2.1 A withdrawal period has been considered but would not be clinically appropriate; and
  - 2.2 There has been a marked reduction in prednisone dose; and
  - 2.3 Either:
    - 2.3.1 There has been an improvement in MRI appearances; or
    - 2.3.2 Marked improvement in other symptomology.

Initial application — (plaque psoriasis) only from a dermatologist or any relevant practitioner on the recommendation of a dermatologist. Approvals valid for 3 months for applications meeting the following criteria: Either:

1 Both:

1.1 The patient has had an initial Special Authority approval for adalimumab, etanercept or secukinumab for severe chronic plaque psoriasis; and

1.2 Either:

- 1.2.1 Patient has experienced intolerable side effects from adalimumab, etanercept or secukinumab; or
- 1.2.2 Patient has received insufficient benefit from adalimumab, etanercept or secukinumab to meet the renewal criteria for adalimumab, etanercept or secukinumab for severe chronic plaque psoriasis; or
- 2 All of the following:

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidi	sed	Generic
 \$	Per	✓	

#### continued...

- 2.1 Any of the following:
  - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
  - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; or
  - 2.1.3 Patient has severe chronic localised genital or flexural plaque psoriasis where the plaques or lesions have been present for at least 6 months from the time of initial diagnosis, and with a Dermatology Life Quality Index (DLQI) score greater than 10; and
- 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
- 2.3 A PASI assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 2.4 The most recent PASI assessment is no more than 1 month old at the time of initiation.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand, foot, genital or flexural areas at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and for the face, palm of a hand or sole of a foot the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (plaque psoriasis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 Both:
    - 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
    - 1.1.2 Following each prior infliximab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-infliximab treatment baseline value; or
  - 1.2 Both:
    - 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
    - 1.2.2 Either:
      - 1.2.2.1 Following each prior infliximab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
      - 1.2.2.2 Following each prior infliximab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-infliximab treatment baseline value; or
  - 1.3 Both:
    - 1.3.1 Patient had severe chronic localised genital or flexural plaque psoriasis at the start of treatment; and
    - 1.3.2 Either:
      - 1.3.2.1 The patient has experienced a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; or
      - 1.3.2.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing infliximab; and

 Price) S	Fully ubsidised	Brand or Generic	
\$ Per	1	Manufacturer	

continued...

2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

**Initial application — (previous use)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient was being treated with infliximab prior to 1 February 2019; and
- 2 Any of the following:
  - 2.1 Rheumatoid arthritis; or
  - 2.2 Ankylosing spondylitis; or
  - 2.3 Psoriatic arthritis; or
  - 2.4 Severe ocular inflammation; or
  - 2.5 Chronic ocular inflammation; or
  - 2.6 Crohn's disease (adults); or
  - 2.7 Crohn's disease (children); or
  - 2.8 Fistulising Crohn's disease; or
  - 2.9 Severe fulminant ulcerative colitis; or
  - 2.10 Severe ulcerative colitis; or
  - 2.11 Plaque psoriasis; or
  - 2.12 Neurosarcoidosis; or
  - 2.13 Severe Behcet's disease.

**Initial application — (psoriatic arthritis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

#### Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis; and
- 2 Either:
  - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept and/or secukinumab; or
  - 2.2 Following 3-4 months' initial treatment with adalimumab and/or etanercept and/or secukinumab, the patient did not meet the renewal criteria for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis.

**Renewal — (psoriatic arthritis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

1 Either:

- 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior infliximab treatment in the opinion of the treating physician; and

2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

**Initial application — (rheumatoid arthritis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and 2 Either:
  - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
  - 2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept; and
- 3 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

**Renewal** — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Infliximab to be administered at doses no greater than 3 mg/kg every 8 weeks.

Initial application — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe Behcet's disease which is significantly impacting the patient's quality of life (see Notes); and
- 2 Either:
  - 2.1 The patient has severe ocular, neurological and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s) (see Notes); or
  - 2.2 The patient has severe gastrointestinal, rheumatologic and/or mucocutaneous symptoms and has not responded adequately to two or more treatment appropriate for the particular symptom(s) (see Notes); and
- 3 The patient is experiencing significant loss of quality of life.

Notes: Behcet's disease diagnosed according to the International Study Group for Behcet's Disease. Lancet 1990;335(8697):1078-80. Quality of life measured using an appropriate quality of life scale such as that published in Gilworth et al J Rheumatol. 2004;31:931-7.

Treatments appropriate for the particular symptoms are those that are considered standard conventional treatments for these symptoms, for example intravenous/oral steroids and other immunosuppressants for ocular symptoms; azathioprine, steroids, thalidomide, interferon alpha and ciclosporin for mucocutaneous symptoms; and colchicine, steroids and methotrexate for rheumatological symptoms.

Renewal — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has had a good clinical response to initial treatment with measurably improved quality of life; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Renewal — (fulminant ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Where maintenance treatment is considered appropriate, infliximab should be used in combination with immunomodulators and reassessed every 6 months; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria: Either:

1 Both:

1.1 The patient has had an initial Special Authority approval for adalimumab for severe ocular inflammation; and

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sul	osidised	Generic
\$	Per	✓	Manufacturer

#### continued...

- 1.2 Either:
  - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
  - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe ocular inflammation; or
- 2 Both:
  - 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
  - 2.2 Any of the following:
    - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
    - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
    - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Renewal — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (ulcerative colitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active ulcerative colitis; and
- 2 Either:
  - 2.1 Patients SCCAI is greater than or equal to 4; or
  - 2.2 Patients PUCAI score is greater than or equal to 20; and
- 3 Patient has tried but has experienced an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and systemic corticosteroids.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 The SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on infliximab; or
  - 1.2 The PUCAI score has reduced by 10 points or more from the PUCAI score when the patient was initiated on infliximab; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

**Initial application — (pyoderma gangrenosum)** only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g.
- prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Note: Indications marked with * are unapproved indications.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

Initial application — (inflammatory bowel arthritis – axial) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has had axial inflammatory pain for six months or more; and
- 3 Patient is unable to take NSAIDs; and
- 4 Patient has unequivocal sacroiliitis demonstrated by radiological imaging or MRI; and
- 5 Patient's disease has not responded adequately to prior treatment consisting of at least 3 months of an exercise regime supervised by a physiotherapist; and
- 6 Patient has a BASDAI of at least 6 on a 0 10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment .

Renewal — (inflammatory bowel arthritis – axial) from any relevant practitioner. Approvals valid for 2 years where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10-point scale, or an improvement in BASDAI of 50%, whichever is less.

**Initial application** — (inflammatory bowel arthritis – peripheral) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has active arthritis in at least four joints from the following: hip, knee, ankle, subtalar, tarsus, forefoot, wrist, elbow, shoulder, sternoclavicular; and
- 3 Patient has tried and not experienced a response to at least three months of methotrexate or azathioprine at a maximum tolerated dose (unless contraindicated); and
- 4 Patient has tried and not experienced a response to at least three months of sulfasalazine at a maximum tolerated dose (unless contraindicated); and
- 5 Any of the following:
  - 5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
  - 5.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application; or
  - 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (inflammatory bowel arthritis – peripheral) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

- Either:
  - 1 Following initial treatment, patient has experienced at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or

	Subsidy (Manufacturer's Price) \$	F Subsidi Per	ully Brand or sed Generic Manufacturer
continued			
2 Patient has experienced at least a continuing 30% impr treating physician.	rovement in active joint of	count from ba	aseline in the opinion of the
MEPOLIZUMAB – Special Authority see SA2331 below – Ret	ail pharmacy		
Inj 100 mg prefilled pen		1	<ul> <li>Nucala</li> </ul>
SA2331 Special Authority for Subsidy Initial application — (Severe eosinophilic asthma) only fro for 12 months for applications meeting the following criteria: All of the following:	m a respiratory physicia	n or clinical i	mmunologist. Approvals valid
1 Patient must be aged 12 years or older; and			
<ol> <li>Patient must have a diagnosis of severe eosinophilic as immunologist; and</li> </ol>	sthma documented by a	respiratory p	hysician or clinical
3 Conditions that mimic asthma eg. vocal cord dysfunction excluded; and	on, central airway obstru	uction, broncl	niolitis etc. have been
<ul> <li>4 Patient has a blood eosinophil count of greater than 0.5</li> <li>5 Patient must be adherent to optimised asthma therapy per day of fluticasone propionate) plus long acting beta maintenance and reliever therapy regimen, unless cont</li> <li>6 Either:</li> </ul>	including inhaled cortico -2 agonist, or budesonic	osteroids (equ de/formoterol	uivalent to at least 1000 mcg
<ul> <li>6.1 Patient has had at least 4 exacerbations needin exacerbation is defined as either documented us corticosteroids; or</li> <li>6.2 Patient has received continuous oral corticoster 3 months; and</li> </ul>	se of oral corticosteroids	s for at least	3 days or parenteral
<ul> <li>7 Treatment is not to be used in combination with subsidi</li> <li>8 Patient has an Asthma Control Test (ACT) score of 10 using the ACT and oral corticosteroid dose must be mathe first dose to assess response to treatment; and</li> <li>9 Either:</li> </ul>	or less. Baseline measu		
<ul><li>9.1 Patient has not previously received an anti-IL5 t</li><li>9.2 Both:</li></ul>	piological therapy for the	ir severe eos	sinophilic asthma; or
<ul><li>9.2.1 Patient was refractory or intolerant to pre</li><li>9.2.2 Patient was not eligible to continue treatr</li><li>within 12 months of commencing treatment</li></ul>	ment with previous anti-l		
Renewal — (Severe eosinophilic asthma) only from a respi years for applications meeting the following criteria: Both:	ratory physician or clinic	al immunolo	jist. Approvals valid for 2
1 An increase in the Asthma Control Test (ACT) score of 2 Either:	at least 5 from baseline	; and	
<ul><li>2.1 Exacerbations have been reduced from baseline</li><li>2.2 Reduction in continuous oral corticosteroid use control.</li></ul>			•
Initial application — (eosinophilic granulomatosis with pol practitioner on the recommendation of a relevant specialist. A criteria: All of the following:			

All of the following:

1 The patient has eosinophilic granulomatosis with polyangiitis; and

 Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
 \$	Per 🗸	

continued...

- 2 The patient has trialled and not received adequate benefit from at least one of the following for at least three months (unless contraindicated to all): azathioprine, cyclophosphamide, leflunomide, methotrexate, mycophenolate, or rituximab; and
- 3 Fither:

- 3.1 The patient has trialled prednisone for a minimum of three months and is unable to maintain disease control at doses below 7.5 mg per day; or
- 3.2 Corticosteroids are contraindicated.

Renewal - (eosinophilic granulomatosis with polyangiitis) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months where patient has no evidence of clinical disease progression. 

OBINUTUZUMAB – PCT only – Specialist – Special Author	ority see SA2155 on the i	next page	
Inj 25 mg per ml, 40 ml vial		1	🗸 Gazyva
Inj 1 mg for ECP	6.21	1 mg	<ul> <li>Baxter</li> </ul>

Subsidy		Fully	Brand or
(Manufacturer's Price)	S	Subsidised	Generic
\$	Per	1	Manufacturer

### ⇒SA2155 Special Authority for Subsidy

**Initial application** — (chronic lymphocytic leukaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C CD20+ chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is obinutuzumab treatment naive; and
- 3 The patient is not eligible for full dose FCR due to comorbidities with a score > 6 on the Cumulative Illness Rating Scale (CIRS) or reduced renal function (creatinine clearance < 70mL/min); and</p>
- 4 Patient has adequate neutrophil and platelet counts* unless the cytopenias are a consequence of marrow infiltration by CLL; and
- 5 Patient has good performance status; and
- 6 Obinutuzumab to be administered at a maximum cumulative dose of 8,000 mg and in combination with chlorambucil for a maximum of 6 cycles.

Notes: Chronic lymphocytic leukaemia includes small lymphocytic lymphoma. Comorbidity refers only to illness/impairment other than CLL induced illness/impairment in the patient. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with obinutuzumab is expected to improve symptoms and improve ECOG score to < 2.

* Neutrophil greater than or equal to  $1.5 \times 10^{9}$ /L and platelets greater than or equal to  $75 \times 10^{9}$ /L.

Initial application — (follicular / marginal zone lymphoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: All of the following:

- 1 Either:
  - 1.1 Patient has follicular lymphoma; or
  - 1.2 Patient has marginal zone lymphoma; and
- 2 Patient is refractory to or has relapsed within 12 months of a rituximab containing combined chemo-immunotherapy regimen*; and
- 3 Patient has an ECOG performance status of 0-2; and
- 4 Patient has been previously treated with no more than four chemotherapy regimens; and
- 5 Obinutuzumab to be administered at a maximum dose of 1000 mg for a maximum of 6 cycles in combination with chemotherapy*.
- Note: * includes unapproved indications

**Renewal — (follicular / marginal zone lymphoma)** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 24 months for applications meeting the following criteria: All of the following:

- 1 Patient has no evidence of disease progression following obinutuzumab induction therapy; and
- 2 Obinutuzumab to be administered at a maximum of 1000 mg every 2 months for a maximum of 2 years; and
- 3 Obinutuzumab to be discontinued at disease progression.

OMALIZUMAB - Special Authority see SA1744 below - Retail pharmacy

Inj 150 mg prefilled syringe		1	<ul> <li>✓ Xolair</li> <li>✓ Xolair AU</li> </ul>
Inj 150 mg vial	450.00	1	<ul> <li>Xolair</li> </ul>

### ⇒SA1744 Special Authority for Subsidy

**Initial application** — (severe asthma) only from a respiratory specialist or clinical immunologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 6 years or older; and
- 2 Patient has a diagnosis of severe asthma; and

Subsidy		Fully	Brand or	
(Manufacturer's Pr	rice) S	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 3 Past or current evidence of atopy, documented by skin prick testing or RAST; and
- 4 Total serum human immunoglobulin E (IgE) between 76 IU/mL and 1300 IU/ml at baseline; and
- 5 Proven adherence with optimal inhaled therapy including high dose inhaled corticosteroid (budesonide 1,600 mcg per day or fluticasone propionate 1,000 mcg per day or equivalent), plus long-acting beta-2 agonist therapy (at least salmeterol 50 mcg bd or eformoterol 12 mcg bd) for at least 12 months, unless contraindicated or not tolerated; and
- 6 Either:
  - 6.1 Patient has received courses of systemic corticosteroids equivalent to at least 28 days treatment in the past 12 months, unless contraindicated or not tolerated; or
  - 6.2 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral steroids; and
- 7 Patient has an Asthma Control Test (ACT) score of 10 or less; and
- 8 Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 26 weeks after the first dose to assess response to treatment.

**Initial application — (severe chronic spontaneous urticaria)** only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Either:
  - 2.1 Both:
    - 2.1.1 Patient is symptomatic with Urticaria Activity Score 7 (UAS7) of 20 or above; and
    - 2.1.2 Patient has a Dermatology life quality index (DLQI) of 10 or greater; or
  - 2.2 Patient has a Urticaria Control Test (UCT) of 8 or less; and
- 3 Any of the following:
  - 3.1 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and ciclosporin (> 3 mg/kg day) for at least 6 weeks; or
  - 3.2 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and at least 3 courses of systemic corticosteroids (> 20 mg prednisone per day for at least 5 days) in the previous 6 months; or
  - 3.3 Patient has developed significant adverse effects whilst on corticosteroids or ciclosporin; and

4 Either:

- 4.1 Treatment to be stopped if inadequate response* following 4 doses; or
- 4.2 Complete response* to 6 doses of omalizumab.

Renewal — (severe asthma) only from a clinical immunologist or respiratory specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 A reduction in the maintenance oral corticosteroid dose or number of exacerbations of at least 50% from baseline.

Renewal — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria: Either:

1 Patient has previously adequately responded* to 6 doses of omalizumab; or

2 Both:

- 2.1 Patient has previously had a complete response* to 6 doses of omalizumab; and
- 2.2 Patient has relapsed after cessation of omalizumab therapy.

Note: *Inadequate response defined as less than 50% reduction in baseline UAS7 and DLQI score, or an increase in Urticaria Control Test (UCT) score of less than 4 from baseline. Patient is to be reassessed for response after 4 doses of omalizumab.

	Subsidy	Fully	Brand or
	(Manufacturer's Price) \$	Subsidised Per 🖌	Generic Manufacturer
continued	d DL OL loop than ar	aqual ta Er ar LIC	T of 16 Polonce of
Complete response is defined as UAS7 less than or equal to 6 ar chronic urticaria on stopping prednisone/ciclosporin does not just			T of 16. Relapse of
PALIVIZUMAB – PCT only – Special Authority see SA2419 below Inj 100 mg per ml, 1 ml vial		1 🗸 5	Synagis
SA2419 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid Both:	d for 6 months for app	plications meeting	the following criteria:
<ol> <li>Palivizumab to be administered during the annual RSV se</li> <li>Either:</li> </ol>	ason; and		
2.1 Both:			
2.1.1 Infant was born in the last 12 months; and	dovo' gostation: or		
2.1.2 Infant was born at less than 32 weeks zero 2.2 Both:	days gestation; or		
2.2.1 Child was born in the last 24 months; and			
2.2.2 Any of the following:			
2.2.2.1 Child has severe lung, airway, neurol ventilatory/respiratory support (see N	•		requires ongoing
2.2.2.2 Both:	ote A) in the commu	nity, or	
2.2.2.2.1 Child has haemodynamically s 2.2.2.2.2 Any of the following:	ignificant heart disea	se; and	
2.2.2.2.1 Child has unoperated si (see Note B); or	mple congenital hear	t disease with sig	nificant left to right shunt
2.2.2.2.2 Child has unoperated or 2.2.2.2.2.3 Child has severe pulmor 2.2.2.2.2.4 Child has moderate or s	nary hypertension (se	ee Note C); or	
2.2.2.3 Child has severe combined immune stem cell transplant; or		. , .	<i>/</i> ·
2.2.2.4 Child has inborn errors of immunity ( respiratory infections, confirmed by a	,	ease susceptibility	y to life-threatening viral
Renewal from any relevant practitioner. Approvals valid for 6 mc All of the following:	onths for applications	meeting the follo	wing criteria:
1 Palivizumab to be administered during the annual RSV se	ason; and		
<ul><li>2 Child was born in the last 24 months; and</li><li>3 Any of the following:</li></ul>			
3.1 Child has severe lung, airway, neurological or neur	omuscular disease th	hat requires ongo	ing ventilatory/respiratory
support (see Note A) in the community, or 3.2 Both:			
3.2.1 Child has haemodynamically significant hea 3.2.2 Any of the following:	art disease; and		
3.2.2.1 Child has unoperated simple congen or	ital heart disease with	h significant left to	o right shunt (see Note B);
3.2.2.2 Child has unoperated or surgically pa 3.2.2.3 Child has severe pulmonary hyperter			ise; or
3.2.2.4 Child has moderate or severe left ver			
3.3 Child has severe combined immune deficiency, con transplant; or	nfirmed by an immun	ologist, but has n	ot received a stem cell

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	1	Manufacturer	

continued...

3.4 Child has inborn errors of immunity (see Note E) that increase susceptibility to life-threatening viral respiratory infections, confirmed by an immunologist.

Notes:

- a) Ventilatory/respiratory support includes those on home oxygen, CPAP/VPAP and those with tracheostomies in situ managed at home
- b) Child requires/will require heart failure medication, and/or child has significant pulmonary hypertension, and/or infant will require surgical palliation/definitive repair within the next 3 months
- c) Mean pulmonary artery pressure more than 25 mmHg
- d) LV Ejection Fraction less than 40%
- e) Inborn errors of immunity include, but are not limited to, IFNAR deficiencies

PERTUZUMAB - PCT only - Specialist - Special Authority see SA2276 below

Inj 30 mg per ml, 14 ml vial		1	<ul> <li>Perjeta</li> </ul>
Inj 420 mg for ECP	3,927.00	420 mg OP	<ul> <li>Baxter</li> </ul>

### ⇒SA2276 Special Authority for Subsidy

**Initial application** — (metastatic breast cancer) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Either:
  - 2.1 Patient is chemotherapy treatment naïve; or
  - 2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
- 3 The patient has good performance status (ECOG grade 0-1); and
- 4 Pertuzumab to be administered in combination with trastuzumab; and
- 5 Pertuzumab maximum first dose of 840 mg, followed by maximum of 420 mg every 3 weeks; and
- 6 Pertuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

1 Both:

- 1.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 1.2 The cancer has not progressed at any time point during the previous 12 months whilst on pertuzumab and trastuzumab; or
- 2 All of the following:
  - 2.1 Patient has previously discontinued treatment with pertuzumab and trastuzumab for reasons other than severe toxicity or disease progression; and
  - 2.2 Patient has signs of disease progression; and
  - 2.3 Disease has not progressed during previous treatment with pertuzumab and trastuzumab.

### RITUXIMAB (MABTHERA) - PCT only - Specialist - Special Authority see SA1976 on the next page

Inj 100 mg per 10 ml vial1,075.50	2	Mabthera
Inj 500 mg per 50 ml vial2,688.30	1	<ul> <li>Mabthera</li> </ul>
Inj 1 mg for ECP5.64	1 mg	<ul> <li>Baxter (Mabthera)</li> </ul>

Subsidy		Fully	Brand or
(Manufacturer's Price)	Su	lbsidised	Generic
\$	Per	✓	Manufacturer

### ➡SA1976 Special Authority for Subsidy

Initial application — (rheumatoid arthritis - TNF inhibitors contraindicated) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Treatment with a Tumour Necrosis Factor alpha inhibitor is contraindicated; and
- 2 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
- 5 Any of the following:
  - 5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or
  - 5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
  - 5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
- 6 Either:
  - 6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
  - 6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 7 Either:
  - 7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
  - 7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months; and
- 8 Either:
  - 8.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
  - 8.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 9 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Initial application — (rheumatoid arthritis - prior TNF inhibitor use) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Both:
  - 1.1 The patient has had an initial community Special Authority approval for at least one of etanercept and/or adalimumab for rheumatoid arthritis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
    - 1.2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
  - 2.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
  - 2.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 3 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Subsi	idy Fu	ly Brand or
(Manufacture	er's Price) Subsidise	d Generic
\$	Per	Manufacturer

continued...

Renewal — (rheumatoid arthritis - re-treatment in 'partial responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 At 4 months following the initial course of rituximab infusions the patient had between a 30% and 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 1.2 At 4 months following the second course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 1.3 At 4 months following the third and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and

3 Either:

- 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
- 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

**Renewal — (rheumatoid arthritis - re-treatment in 'responders' to rituximab)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

1 Either:

- 1.1 At 4 months following the initial course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 1.2 At 4 months following the second and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Either:
  - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
  - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

RITUXIMAB (RIXIMYO) - PCT only - Specialist - Special Authority see SA2233 below

Inj 100 mg per 10 ml vial		2	Riximyo
Inj 500 mg per 50 ml vial		1	Riximyo
Inj 1 mg for ECP	1.38	1 mg	<ul> <li>Baxter (Riximyo)</li> </ul>

### ➡SA2233 Special Authority for Subsidy

**Initial application** — (ABO-incompatible organ transplant) from any relevant practitioner. Approvals valid without further renewal unless notified where patient is to undergo an ABO-incompatible solid organ transplant*.

Note: Indications marked with * are unapproved indications.

Initial application — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 The total rituximab dose would not exceed the equivalent of 375  $mg/m^2$  of body-surface area per week for a total of 4 weeks; and
- 3 Any of the following:

Subsidy	Full	Brand or
(Manufacturer's	Price) Subsidise	Generic
\$	Per 🖌	Manufacturer

continued...

- 3.1 Induction therapy with daily oral or pulse intravenous cyclophosphamide has failed to achieve significant improvement of disease after at least 3 months; or
- 3.2 Patient has previously had a cumulative dose of cyclophosphamide > 15 g or a further repeat 3 month induction course of cyclophosphamide would result in a cumulative dose > 15 g; or
- 3.3 Cyclophosphamide and methotrexate are contraindicated; or
- 3.4 Patient is a female of child-bearing potential; or
- 3.5 Patient has a previous history of haemorrhagic cystitis, urological malignancy or haematological malignancy.
- Note: Indications marked with * are unapproved indications.

Renewal — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 Patient has previously responded to treatment with rituximab but is now experiencing an acute flare of vasculitis; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

**Initial application** — (Antibody-mediated organ transplant rejection) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has been diagnosed with antibody-mediated organ transplant rejection*. Note: Indications marked with * are unapproved indications.

**Initial application — (Chronic lymphocytic leukaemia)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C chronic lymphocytic leukaemia (CLL) requiring treatment; and
- 2 Any of the following:
  - 2.1 The patient is rituximab treatment naive; or
  - 2.2 Either:
    - 2.2.1 The patient is chemotherapy treatment naive; or
    - 2.2.2 Both:
      - 2.2.2.1 The patient's disease has relapsed following no more than three prior lines of chemotherapy treatment; and
      - 2.2.2.2 The patient has had a treatment-free interval of 12 months or more if previously treated with fludarabine and cyclophosphamide chemotherapy; or
  - 2.3 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax; and
- 3 The patient has good performance status; and
- 4 Either:
  - 4.1 The patient does not have chromosome 17p deletion CLL; or
  - 4.2 Rituximab treatment is to be used in combination with funded venetoclax for relapsed/refractory chronic lymphocytic leukaemia; and
- 5 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles; and
- 6 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration), bendamustine or venetoclax.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with rituximab is expected to improve symptoms and improve ECOG score to < 2.

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

Renewal — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

1 Either:

- 1.1 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax; or
- 1.2 All of the following:
  - 1.2.1 The patient's disease has relapsed following no more than one prior line of treatment with rituximab for CLL; and
  - 1.2.2 The patient has had an interval of 36 months or more since commencement of initial rituximab treatment; and
  - 1.2.3 The patient does not have chromosome 17p deletion CLL; and
  - 1.2.4 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration) or bendamustine; and
- 2 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initial application — (Neuromyelitis Optica Spectrum Disorder(NMOSD)) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks; and
- 2 Either:
  - 2.1 The patient has experienced a severe episode or attack of NMOSD (rapidly progressing symptoms and clinical investigations supportive of a severe attack of NMOSD); or
  - 2.2 All of the following:
    - 2.2.1 The patient has experienced a breakthrough attack of NMOSD; and
    - 2.2.2 The patient is receiving treatment with mycophenolate; and
    - 2.2.3 The patients is receiving treatment with corticosteroids.

Renewal — (Neuromyelitis Optica Spectrum Disorder) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks; and
- 2 The patients has responded to the most recent course of rituximab; and
- 3 The patient has not received rituximab in the previous 6 months.

**Initial application — (Post-transplant)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 2 To be used for a maximum of 8 treatment cycles.

Note: Indications marked with * are unapproved indications.

Renewal — (Post-transplant) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

continued...

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

5	Subsidy	Fully	Brand or
(Manufa	acturer's Price) Sub	sidised	Generic
	\$ Per	✓	Manufacturer

continued...

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 3 To be used for no more than 6 treatment cycles.
- Note: Indications marked with * are unapproved indications.

Initial application — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
- 2 Either:
  - 2.1 Treatment with corticosteroids and at least one other immunosuppressant for at least a period of 12 months has been ineffective; or
  - 2.2 Both:
    - 2.2.1 Treatment with at least one other immunosuppressant for a period of at least 12 months; and
    - 2.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

Renewal — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Either:
  - 3.1 The patient has relapsed despite treatment with corticosteroids and at least one other immunosuppressant for a period of at least 12 months; or
  - 3.2 Both:
    - 3.2.1 The patient's myasthenia gravis has relapsed despite treatment with at least one immunosuppressant for a period of at least 12 months; and
    - 3.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

## Initial application — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS))

only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is a child with SDNS* or FRNS*; and
- 2 Treatment with steroids for at least a period of 3 months has been ineffective or associated with evidence of steroid toxicity; and
- 3 Treatment with ciclosporin for at least a period of 3 months has been ineffective and/or discontinued due to unacceptable side effects; and
- 4 Treatment with mycophenolate for at least a period of 3 months with no reduction in disease relapses; and
- 5 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	~	Manufacturer	

continued...

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Patient is a child with SRNS* where treatment with steroids and ciclosporin for at least 3 months have been ineffective; and
- 2 Treatment with tacrolimus for at least 3 months has been ineffective; and
- 3 Genetic causes of nephrotic syndrome have been excluded; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

**Initial application** — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 All of the following:
  - 1.1 The patient has treatment naive aggressive CD20 positive NHL; and
  - 1.2 To be used with a multi-agent chemotherapy regimen given with curative intent; and
  - 1.3 To be used for a maximum of 8 treatment cycles; or
- 2 Both:
  - 2.1 The patient has aggressive CD20 positive NHL with relapsed disease following prior chemotherapy; and
  - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Renewal — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has relapsed refractory/aggressive CD20 positive NHL; and
- 3 To be used with a multi-agent chemotherapy regimen given with curative intent; and
- 4 To be used for a maximum of 4 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

**Initial application** — (haemophilia with inhibitors) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria: Any of the following:

Subsidy (Manufacturer's Price)	ę	Fully Subsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

- 1 Patient has mild congenital haemophilia complicated by inhibitors; or
- 2 Patient has severe congenital haemophilia complicated by inhibitors and has failed immune tolerance therapy; or
- 3 Patient has acquired haemophilia.

**Renewal** — (haemophilia with inhibitors) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for haemophilia with inhibitors; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment.

**Initial application — (immune thrombocytopenic purpura (ITP))** only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Either:
  - 1.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per microlitre; or
  - 1.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding; and
- 2 Any of the following:
  - 2.1 Treatment with steroids and splenectomy have been ineffective; or
  - 2.2 Treatment with steroids has been ineffective and splenectomy is an absolute contraindication; or
  - Other treatments including steroids have been ineffective and patient is being prepared for elective surgery (e.g. splenectomy); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.
- Note: Indications marked with * are unapproved indications.

Renewal — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

- Either:
  - 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
  - 2 All of the following:
    - 2.1 Patient was previously treated with rituximab for immune thrombocytopenic purpura*; and
    - 2.2 An initial response lasting at least 12 months was demonstrated; and
    - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (indolent, low-grade lymphomas or hairy cell leukaemia*) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria: Either:

1 Both:

- The patient has indolent low grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
- 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
  - 2.1 The patient has indolent, low grade lymphoma or hairy cell leukaemia* requiring first-line systemic chemotherapy; and
  - 2.2 To be used for a maximum of 6 treatment cycles.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsi	idised	Generic	
\$	Per	1	Manufacturer	

continued...

Note: 'Indolent, Iow-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant. Renewal — (indolent, Iow-grade lymphomas or hairy cell leukaemia*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has indolent, low-grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
- 3 To be used for no more than 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Initial application — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient has autoimmune pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder.

Note: Indications marked with * are unapproved indications.

Renewal — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient was previously treated with rituximab for pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder and demonstrated an initial response lasting at least 12 months.

Note: Indications marked with * are unapproved indications.

**Initial application** — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Patient has cold haemagglutinin disease*; and
- 2 Patient has severe disease which is characterized by symptomatic anaemia, transfusion dependence or disabling circulatory symptoms; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
  - 2.1 Patient was previously treated with rituximab for severe cold haemagglutinin disease*; and
  - 2.2 An initial response lasting at least 12 months was demonstrated; and
  - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Both:

1 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks; and

2 Either:

2.1 Patient has thrombotic thrombocytopenic purpura* and has experienced progression of clinical symptoms or persistent thrombocytopenia despite plasma exchange; or

2.2 Patient has acute idiopathic thrombotic thrombocytopenic purpura* with neurological or cardiovascular pathology. Note: Indications marked with * are unapproved indications.

Subsidy	Fu	lly Brand or
(Manufacturer's	Price) Subsidise	ed Generic
\$	Per	<ul> <li>Manufacturer</li> </ul>

### continued...

Renewal — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for thrombotic thrombocytopenic purpura*; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe, immediately life- or organ-threatening SLE*; and
- 2 The disease has proved refractory to treatment with steroids at a dose of at least 1 mg/kg; and
- 3 The disease has relapsed following prior treatment for at least 6 months with maximal tolerated doses of azathioprine, mycophenolate mofetil and high dose cyclophosphamide, or cyclophosphamide is contraindicated; and
- 4 Maximum of four 1000 mg infusions of rituximab.
- Note: Indications marked with * are unapproved indications.

Renewal — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient's SLE* achieved at least a partial response to the previous round of prior rituximab treatment; and
- 2 The disease has subsequently relapsed; and
- 3 Maximum of two 1000 mg infusions of rituximab.

Note: Indications marked with * are unapproved indications.

Initial application — (warm autoimmune haemolytic anaemia (warm AIHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Patient has warm autoimmune haemolytic anaemia*; and
- 2 One of the following treatments has been ineffective: steroids (including if patient requires ongoing steroids at doses equivalent to > 5 mg prednisone daily), cytotoxic agents (e.g. cyclophosphamide monotherapy or in combination), intravenous immunoglobulin; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

**Renewal — (warm autoimmune haemolytic anaemia (warm AIHA))** only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Either:

1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or

2 All of the following:

- 2.1 Patient was previously treated with rituximab for warm autoimmune haemolytic anaemia*; and
- 2.2 An initial response lasting at least 12 months was demonstrated; and
- 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

**Initial application** — (severe antisynthetase syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed antisynthetase syndrome; and
- 2 Patient has severe, immediately life or organ threatening disease, including interstitial lung disease; and
- 3 Either:
  - 3.1 Treatment with at least 3 immunosuppressants (oral steroids, cyclophosphamide, methotrexate, mycophenolate, ciclosporin, azathioprine) has not be effective at controlling active disease; or
  - 3.2 Rapid treatment is required due to life threatening complications; and
- 4 Maximum of four 1,000mg infusions of rituximab.

Renewal — (severe antisynthetase syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in inflammatory markers, muscle strength and pulmonary function; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 Maximum of two cycles of  $2 \times 1,000$  mg infusions of rituximab given two weeks apart.

Initial application — (graft versus host disease) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has refractory graft versus host disease following transplant; and
- 2 Treatment with at least 3 immunosuppressants (oral steroids, ciclosporin, tacrolimus, mycophenolate, sirolimus) has not be effective at controlling active disease; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

**Initial application** — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has severe chronic inflammatory demyelinating polyneuropathy (CIPD); and
- 2 Either:
  - 2.1 Both:
    - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
    - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
  - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function compared to baseline; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

#### continued...

Initial application — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has severe anti-NMDA receptor autoimmune encephalitis; and
- 2 Either:
  - 2.1 Both:
    - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
    - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
  - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 The patient has experienced a relapse and now requires further treatment; and
- 4 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Initial application — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has CD20+ low grade or follicular B-cell NHL with relapsed disease following prior chemotherapy; and
  - 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
  - 2.1 The patient has CD20+ low grade or follicular B-cell NHL requiring first-line systemic chemotherapy; and
  - 2.2 To be used for a maximum of 6 treatment cycles.

Renewal — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 Rituximab is to be used for maintenance in CD20+ low grade or follicular B-cell NHL following induction with first-line systemic chemotherapy; and
- 2 Patient is intended to receive rituximab maintenance therapy for 2 years at a dose of 375 mg/m2 every 8 weeks (maximum of 12 cycles).

**Initial application** — (Membranous nephropathy) only from a nephrologist or any relevant practitioner on the recommendation of a nephrologist. Approvals valid for 6 weeks for applications meeting the following criteria: All of the following:

1 Either:

- 1.1 Patient has biopsy-proven primary/idiopathic membranous nephropathy*; or
- 1.2 Patient has PLA2 antibodies with no evidence of secondary cause, and an eGFR of > 60ml/min/1.73m2; and
- 2 Patient remains at high risk of progression to end-stage kidney disease despite more than 3 months of treatment with conservative measures (see Note); and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	Subsidised	Generic	
\$	Per	1	Manufacturer	

continued...

3 The total rituximab dose would not exceed the equivalent of 375mg/m2 of body surface area per week for a total of 4 weeks.

**Renewal** — (Membranous nephropathy) only from a nephrologist or any relevant practitioner on the recommendation of a nephrologist. Approvals valid for 6 weeks for applications meeting the following criteria: All of the following:

1 Patient was previously treated with rituximab for membranous nephropathy*; and

2 Either:

- 2.1 Treatment with rituximab was previously successful, but the condition has relapsed, and the patient now requires repeat treatment; or
- 2.2 Patient achieved partial response to treatment and requires repeat treatment (see Note); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Notes:

- a) Indications marked with * are unapproved indications.
- b) High risk of progression to end-stage kidney disease defined as > 5g/day proteinuria.
- c) Conservative measures include renin-angiotensin system blockade, blood-pressure management, dietary sodium and protein restriction, treatment of dyslipidaemia, and anticoagulation agents unless contraindicated or the patient has experienced intolerable side effects.
- d) Partial response defined as a reduction of proteinuria of at least 50% from baseline, and between 0.3 grams and 3.5 grams per 24 hours.

Initial application — (B-cell acute lymphoblastic leukaemia/lymphoma*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient has newly diagnosed B-cell acute lymphoblastic leukaemia/lymphoma*; and
- 2 Treatment must be in combination with an intensive chemotherapy protocol with curative intent; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m² per dose for a maximum of 18 doses.

Note: Indications marked with * are unapproved indications.

**Initial application — (desensisation prior to transplant)** from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires desensitisation prior to mismatched allogenic stem cell transplant*; and
- 2 Patient would receive no more than two doses at 375 mg/m2 of body-surface area.

Note: Indications marked with * are unapproved indications.

**Initial application — (pemiphigus*)** only from a dermatologist or relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 All of the following:
  - 1.1 Patient has severe rapidly progressive pemphigus; and
  - 1.2 Is used in combination with systemic corticosteroids (20 mg/day); and
  - 1.3 Any of the following:
    - 1.3.1 Skin involvement is at least 5% body surface area; or
    - 1.3.2 Significant mucosal involvement (10 or more mucosal erosions) or diffuse gingivitis or confluent large erosions; or
    - 1.3.3 Involvement of two or more mucosal sites; or

2 Both:

2.1 Patient has pemphigus; and

continued...

AThree months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsid	lised	Generic
\$	Per	1	Manufacturer

continued...

2.2 Patient has not experienced adequate clinical benefit from systemic corticosteroids (20 mg/day) in combination with a steroid sparing agent, unless contraindicated.

Note: Indications marked with * are unapproved indications.

Renewal — (pemiphigus*) only from a dermatologist or relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has experienced adequate clinical benefit from rituximab treatment, with improvement in symptoms and healing of skin ulceration and reduction in corticosteroid requirement; and
- 2 Patient has not received rituximab in the previous 6 months.
- Note: Indications marked with * are unapproved indications.

Initial application — (immunoglobulin G4-related disease (IgG4-RD*)) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed diagnosis of IgG4-RD*; and
- 2 Either:
  - 2.1 Treatment with corticosteroids and/or disease modifying anti-rheumatic drugs for at least 3 months has been ineffective in lowering corticosteroid dose below 5 mg per day (prednisone equivalent) without relapse; or
  - 2.2 Treatment with corticosteroids and/or disease modifying anti-rheumatic drugs is contraindicated or associated with evidence of toxicity or intolerance; and
- 3 Total rituximab dose used should not exceed a maximum of two 1000 mg infusions of rituximab given two weeks apart.

Note: Indications marked with * are unapproved indications.

Renewal — (immunoglobulin G4-related disease (IgG4-RD*)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 Either:

- 1.1 Treatment with rituximab for IgG4-RD* was previously successful and patient's disease has demonstrated sustained response, but the condition has relapsed; or
- 1.2 Patient is receiving maintenance treatment for IgG4-RD*; and
- 2 Rituximab re-treatment not to be given within 6 months of previous course of treatment; and
- 3 Maximum of two 1000 mg infusions of rituximab given two weeks apart.

Note: Indications marked with * are unapproved indications.

SECUKINUMAB - Special Authority see SA2403 below - Retail p	oharmacy		
Inj 150 mg per ml, 1 ml prefilled syringe		1	<ul> <li>Cosentyx</li> </ul>
	1,599.00	2	<ul> <li>Cosentyx</li> </ul>

### ⇒SA2403 Special Authority for Subsidy

Initial application — (severe chronic plaque psoriasis – second-line biologic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

1 The patient has had an initial Special Authority approval for adalimumab or etanercept, or has trialled infliximab in a Health NZ Hospital, for severe chronic plaque psoriasis; and

2 Either:

- 2.1 The patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
- 2.2 The patient has received insufficient benefit from adalimumab, etanercept or infliximab; and
- 3 A Psoriasis Area and Severity Index (PASI) assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
 (Manulacturer's Frice)	Per 🗸	

continued...

4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

**Initial application — (severe chronic plaque psoriasis – first-line biologic)** only from a dermatologist or any relevant practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
  - 1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; or
  - 1.3 Patient has severe chronic localised genital or flexural plaque psoriasis where the plaques or lesions have been present for at least 6 months from the time of initial diagnosis, and with a Dermatology Life Quality Index (DLQI) score greater than 10; and
- 2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
- 3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Note: A treatment course is defined as a minimum of 12 weeks of treatment. "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand. foot, genital or flexural areas, at least 2 of the 3 PASI symptom sub scores for erythema, thickness and scaling are rated as severe or very severe, and for the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis – first and second-line biologic) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 Either:
    - 1.1.1 Patient's PASI score has reduced by 75% or more (PASI 75) as compared to baseline PASI prior to commencing secukinumab; or
    - 1.1.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing secukinumab; or

1.2 Both:

- 1.2.1 Patient had severe chronic localised genital or flexural plaque psoriasis at the start of treatment; and
- 1.2.2 Either:
  - 1.2.2.1 The patient has experienced a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; or
  - 1.2.2.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing secukinumab; and
- 2 Secukinumab to be administered at a maximum dose of 300 mg monthly.

**Initial application** — (ankylosing spondylitis – second-line biologic) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria: Both:

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and 2 Either:
  - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
  - 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

Renewal — (ankylosing spondylitis – second-line biologic) only from a rheumatologist or medical practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Following 12 weeks initial treatment of secukinumab treatment, BASDAI has improved by 4 or more points from pre-secukinumab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefitted from treatment and that continued treatment is appropriate; and
- 3 Secukinumab to be administered at doses no greater than 150 mg monthly.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 Patient has had an initial Special Authority approval for adalimumab, etanercept or infliximab for psoriatic arthritis; and
  - 1.2 Either:
    - 1.2.1 Patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
    - 1.2.2 Patient has received insufficient benefit from adalimumab, etanercept or infliximab to meet the renewal criteria for adalimumab, etanercept or infliximab for psoriatic arthritis; or
- 2 All of the following:
  - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
  - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
  - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
  - 2.4 Either:
    - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
    - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
  - 2.5 Any of the following:
    - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
    - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
    - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

**Renewal** — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Fither
  - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior secukinumab treatment in the opinion of the treating physician; and

1 mg

Baxter

	Subsidy (Manufacturer's Price) \$	Fu Subsidis Per	ully Brand or sed Generic Manufacturer
continued			
2 Secukinumab to be administered at doses no greater than	n 300 mg monthly.		
SILTUXIMAB – Special Authority see SA1596 below – Retail ph. Note: Siltuximab is to be administered at doses no greater the Inj 100 mg vial	han 11 mg/kg every 3		✓ Sylvant
Inj 400 mg vial			✓ Sylvant
<ul> <li>SA1596 Special Authority for Subsidy</li> <li>Initial application only from a haematologist or rheumatologist.</li> <li>following criteria:</li> <li>All of the following:         <ol> <li>Patient has severe HHV-8 negative idiopathic multicentric</li> <li>Treatment with an adequate trial of corticosteroids has pro- 3 Siltuximab is to be administered at doses no greater than</li> </ol> </li> <li>Renewal only from a haematologist or rheumatologist. Approval</li> </ul>	Castleman's Disease oven ineffective; and 11 mg/kg every 3 wee	e; and eks.	
and the patient has sustained improvement in inflammatory mark			atment remains appropriate
TOCILIZUMAB - PCT only - Special Authority see SA2404 belo			
Inj 20 mg per ml, 4 ml vial		-	<ul> <li>Actemra</li> </ul>
Inj 20 mg per ml, 10 ml vial		-	<ul> <li>Actemra</li> </ul>
Inj 20 mg per ml, 20 ml vial	1,100.00	1	<ul> <li>Actemra</li> </ul>

#### Either: 1 Both:

■ SA2404 Special Authority for Subsidy

unless notified for applications meeting the following criteria:

1.1 The patient has developed grade 3 or 4 cytokine release syndrome associated with the administration of blinatumomab for the treatment of acute lymphoblastic leukaemia; and

Initial application — (cytokine release syndrome) from any relevant practitioner. Approvals valid without further renewal

- 1.2 Tocilizumab is to be administered at doses no greater than 8 mg/kg IV for a maximum of 3 doses (if less than 30kg, maximum of 12 mg/kg); or
- 2 All of the following:
  - 2.1 The patient is enrolled in the Malaghan Institute of Medical Research ENABLE trial programme; and
  - 2.2 The patient has developed CRS or Immune Effector Cell-Associated Neurotoxicity Syndrome (ICANS) following CAR T-cell therapy for the treatment of relapsed or refractory B-cell non-Hodgkin lymphoma; and
  - 2.3 Tocilizumab is to be administered according to the consensus guidelines for CRS or ICANS for CAR T-cell therapy at doses no greater than 8 mg/kg IV for a maximum of 3 doses.

**Initial application — (previous use)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 Patient was being treated with tocilizumab prior to 1 February 2019; and

- 2 Any of the following:
  - 2.1 rheumatoid arthritis; or
  - 2.2 systemic juvenile idiopathic arthritis; or
  - 2.3 adult-onset Still's disease; or
  - 2.4 polyarticular juvenile idiopathic arthritis; or
  - 2.5 idiopathic multicentric Castleman's disease.

Initial application — (Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	1	Manufacturer

continued...

the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
  - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
  - 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
- 3 Either:
  - 3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or 3.2 Both:
    - 3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a Health NZ Hospital; and 3.2.2 Either:
      - 3.2.2.1 The patient has experienced intolerable side effects from rituximab; or
      - 3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

**Initial application — (Rheumatoid Arthritis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 2 Tocilizumab is to be used as monotherapy; and
- 3 Either:
  - 3.1 Treatment with methotrexate is contraindicated; or
  - 3.2 Patient has tried and did not tolerate oral and/or parenteral methotrexate; and
- 4 Either:
  - 4.1 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of ciclosporin alone or in combination with another agent; or
  - 4.2 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of leflunomide alone or in combination with another agent; and
- 5 Either:
  - 5.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 active, swollen, tender joints; or
  - 5.2 Patient has persistent symptoms of poorly controlled and active disease in at least four active joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 6 Either:
  - 6.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
  - 6.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Initial application — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient diagnosed with systemic juvenile idiopathic arthritis; and
- 2 Patient has tried and not responded to a reasonable trial of all of the following, either alone or in combination: oral or parenteral methotrexate; non-steroidal anti-inflammatory drugs (NSAIDs); and systemic corticosteroids.

**Initial application — (adult-onset Still's disease)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

### continued...

Either:

1 Both:

- 1.1 Either:
  - 1.1.1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for adult-onset Still's disease (AOSD); or
  - 1.1.2 The patient has been started on tocilizumab for AOSD in a Health NZ Hospital; and
- 1.2 Either:
  - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
  - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
  - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
  - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal antiinflammatory drugs (NSAIDs) and methotrexate; and
  - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

**Initial application — (polyarticular juvenile idiopathic arthritis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: Either:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for both etanercept and adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
- 1.2 The patient has experienced intolerable side effects, or has received insufficient benefit from, both etanercept and adalimumab; or
- 2 All of the following:
  - 2.1 Treatment with a tumour necrosis factor alpha inhibitor is contraindicated; and
  - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
  - 2.3 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
  - 2.4 Any of the following:
    - 2.4.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
    - 2.4.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
    - 2.4.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Initial application — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Tocilizumab to be administered at doses no greater than 8 mg/kg IV every 3-4 weeks.

**Initial application — (moderate to severe COVID-19)** from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed (or probable) COVID-19; and
- 2 Oxygen saturation of < 92% on room air, or requiring supplemental oxygen; and
- 3 Patient is receiving adjunct systemic corticosteroids, or systemic corticosteroids are contraindicated; and

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

Subsidy		Fully	Brand or
(Manufacturer's Price)	5	Subsidised	Generic
\$	Per	1	Manufacturer

continued...

- 4 Tocilizumab is to be administered at doses no greater than 8mg/kg IV for a maximum of one dose; and
- 5 Tocilizumab is not to be administered in combination with barcitinib.

**Renewal — (Rheumatoid Arthritis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

- Either:
  - 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Renewal — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Either:

- 1 Following up to 6 months' initial treatment, the patient has achieved at least an American College of Rheumatology paediatric 30% improvement criteria (ACR Pedi 30) response from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing ACR Pedi 30 response from baseline.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months where the patient has a sustained improvement in inflammatory markers and functional status. Renewal — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
  - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Renewal — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has a sustained improvement in inflammatory markers and functional status.

TRASTUZUMAB (HERZUMA) - PCT only - Special Authority see SA2293 below

Inj 150 mg vial		1	<ul> <li>Herzuma</li> </ul>
Inj 440 mg vial		1	<ul> <li>Herzuma</li> </ul>
Inj 1 mg for ECP	0.70	1 mg	✓ Baxter

### ► SA2293 Special Authority for Subsidy

**Initial application** — (early breast cancer) from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria:

Both:

- 1 The patient has early breast cancer expressing HER-2 IHC 3+ or ISH + (including FISH or other current technology); and
- 2 Maximum cumulative dose of 106 mg/kg (12 months' treatment).

Renewal — (early breast cancer*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

#### Either:

- 1 All of the following:
  - 1.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and

Sub	bsidy F	ully E	Brand or
(Manufactu	urer's Price) Subsidi	sed C	Generic
	\$ Per	<ul> <li>N</li> </ul>	Manufacturer

#### continued...

- 1.2 The patient received prior adjuvant trastuzumab treatment for early breast cancer; and
- 1.3 Any of the following:
  - 1.3.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
  - 1.3.2 The patient discontinued lapatinib within 3 months due to intolerable side effects and the cancer did not progress whilst on lapatinib; or
- 1.3.3 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and 1.4 Fither:
- 1.4 Either:
  - 1.4.1 Trastuzumab will not be given in combination with pertuzumab; or
  - 1.4.2 All of the following:
    - 1.4.2.1 Trastuzumab to be administered in combination with pertuzumab; and
    - 1.4.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
    - 1.4.2.3 The patient has good performance status (ECOG grade 0-1); and
- 1.5 Trastuzumab to be discontinued at disease progression; or
- 2 All of the following:
  - 2.1 Patient has previously discontinued treatment with trastuzumab in the metastatic setting for reasons other than severe toxicity or disease progression; and
  - 2.2 Patient has signs of disease progression; and
  - 2.3 Disease has not progressed during previous treatment with trastuzumab.

Note: * For patients with relapsed HER-2 positive disease who have previously received adjuvant trastuzumab for early breast cancer

Initial application — (metastatic breast cancer) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Either:
  - 2.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
  - 2.2 The patient discontinued lapatinib within 3 months due to intolerable side effects and the cancer did not progress whilst on lapatinib; and
- 3 Either:
  - 3.1 Trastuzumab will not be given in combination with pertuzumab; or
  - 3.2 All of the following:
    - 3.2.1 Trastuzumab to be administered in combination with pertuzumab; and
    - 3.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
    - 3.2.3 The patient has good performance status (ECOG grade 0-1); and
- 4 Trastuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 All of the following:
  - 1.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
  - 1.2 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidi	sed	Generic
\$	Per	✓	

#### continued...

- 1.3 Trastuzumab to be discontinued at disease progression; or
- 2 All of the following:
  - 2.1 Patient has previously discontinued treatment with trastuzumab for reasons other than severe toxicity or disease progression; and
  - 2.2 Patient has signs of disease progression; and
  - 2.3 Disease has not progressed during previous treatment with trastuzumab.

Initial application — (gastric, gastro-oesophageal junction and oesophageal cancer) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has locally advanced or metastatic gastric, gastro-oesophageal junction or oesophageal cancer expressing HER-2 IHC 2+ FISH+ or IHC3+ (or other current technology); and
- 2 Patient has an ECOG score of 0-2.

Renewal — (gastric, gastro-oesophageal junction and oesophageal cancer) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 2 Trastuzumab to be discontinued at disease progression.

TRASTUZUMAB DERUXTECAN – PCT only – Special Authority see SA2420 below

Inj 100 mg per ml, 1 ml vial	2,550.00	1	🗸 Enhertu
Inj 1 mg for ECP	27.05	1 mg	<ul> <li>Baxter</li> </ul>

### ⇒SA2420 Special Authority for Subsidy

Initial application only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: Fither:

- 1 Patient is currently on treatment with trastuzumab deruxtecan and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
  - Patient has metastatic breast cancer expressing HER-2 IHC3+ or ISH+ (including FISH or other current technology); and
  - 2.2 Patient has previously received trastuzumab and chemotherapy, separately or in combination; and
  - 2.3 Either:
    - 2.3.1 The patient has received prior therapy for metastatic disease; or
    - 2.3.2 The patient developed disease recurrence during, or within six months of completing adjuvant therapy; and
  - 2.4 Patient has a good performance status (ECOG 0-1); and
  - 2.5 Patient has not received prior funded trastuzumab deruxtecan treatment; and
  - 2.6 Treatment to be discontinued at disease progression.

**Renewal** only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The cancer has not progressed at any time point during the previous approval period whilst on trastuzumab deruxtecan; and
- 2 Treatment to be discontinued at disease progression.

Note: Prior or adjuvant therapy includes anthracycline, other chemotherapy, biological drugs, or endocrine therapy.

TRASTUZUMAB EMTANSINE - PCT only - Specialis	st – Special Authority see SA2	424 on the	next page
Inj 100 mg vial	2,320.00	1	Kadcyla
Inj 160 mg vial		1	Kadcyla
Inj 1 mg for ECP	24.52	1 mg	<ul> <li>Baxter</li> </ul>

Subsidy	Fu	ılly	Brand or
(Manufacturer's Price)	Subsidis	ed	Generic
 \$	Per	1	Manufacturer

### ⇒SA2424 Special Authority for Subsidy

**Initial application** — (early breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

1 Patient has early breast cancer expressing HER2 IHC3+ or ISH+; and

- 2 Documentation of pathological invasive residual disease in the breast and/or axiliary lymph nodes following completion of surgery; and
- 3 Patient has completed systemic neoadjuvant therapy with trastuzumab and chemotherapy prior to surgery; and
- 4 Disease has not progressed during neoadjuvant therapy; and
- 5 Patient has left ventricular ejection fraction of 45% or greater; and
- 6 Adjuvant treatment with trastuzumab emtansine to be commenced within 12 weeks of surgery; and
- 7 Trastuzumab emtansine to be discontinued at disease progression; and
- 8 Total adjuvant treatment duration must not exceed 42 weeks (14 cycles).

**Initial application** — (metastatic breast cancer) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Patient has previously received trastuzumab and chemotherapy, separately or in combination; and
- 3 Either:
  - 3.1 The patient has received prior therapy for metastatic disease*; or
- 3.2 The patient developed disease recurrence during, or within six months of completing adjuvant therapy*; and
- 4 Patient has a good performance status (ECOG 0-1); and
- 5 Either:
  - 5.1 Patient does not have symptomatic brain metastases; or
  - 5.2 Patient has brain metastases and has received prior local CNS therapy; and
- 6 Either:
  - 6.1 Patient has not received prior funded trastuzumab emtansine or trastuzumab deruxtecan treatment; or
  - 6.2 Both:
    - 6.2.1 Patient has discontinued trastuzumab deruxtecan due to intolerance; and
    - 6.2.2 The cancer did not progress while on trastuzumab deruxtecan; and
- 7 Treatment to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- Both:
  - 1 The cancer has not progressed at any time point during the previous approval period whilst on trastuzumab emtansine; and
  - 2 Treatment to be discontinued at disease progression.

Note: Prior or adjuvant therapy includes anthracycline, other chemotherapy, biological drugs, or endocrine therapy.

USTEKINUMAB - Special Authority see SA2182 below - Retail pharmacy

### ➡SA2182 Special Authority for Subsidy

**Initial application** — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

1 Patient is currently on treatment with ustekinumab commenced prior to 1 February 2023 and met all remaining criteria (criterion 2) below at the time of commencing treatment; or

Subsidy		Fully	Brand or	
(Manufacturer's Price)	5	Subsidised	Generic	
\$	Per	✓	Manufacturer	

#### continued...

2 Both:

- 2.1 Patient has active Crohn's disease; and
- 2.2 Either:

2.2.1 Patient has had an initial approval for prior biologic therapy for Crohn's disease and has experienced intolerable side effects or insufficient benefit to meet renewal criteria; or

2.2.2 Both:

- 2.2.2.1 Patient meets the initiation criteria for prior biologic therapies for Crohn's disease; and
- 2.2.2.2 Other biologics for Crohn's disease are contraindicated.

Renewal — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 CDAI score has reduced by 100 points, or HBI score has reduced by 3 points, from when the patient was initiated on biologic therapy; or
  - 1.2 CDAI score is 150 or less, or HBI is 4 or less; or
  - 1.3 The patient has experienced an adequate response to treatment, but CDAI score and/or HBI score cannot be assessed; and
- 2 Ustekinumab to be administered at a dose no greater than 90 mg every 8 weeks.

Initial application — (Crohn's disease - children*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Patient is currently on treatment with ustekinumab commenced prior to 1 February 2023 and met all remaining criteria (criterion 2) below at the time of commencing treatment; or
- 2 Both:
  - 2.1 Patient has active Crohn's disease; and
  - 2.2 Either:
    - 2.2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria; or
    - 2.2.2 Both:
      - 2.2.2.1 Patient meets the initiation criteria for prior biologic therapies for Crohn's disease; and
      - 2.2.2.2 Other biologics for Crohn's disease are contraindicated.

Note: Indication marked with * is an unapproved indication.

Renewal — (Crohn's disease - children*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 PCDAI score has reduced by 10 points from when the patient was initiated on biologic therapy; or
  - 1.2 PCDAI score is 15 or less; or
  - 1.3 The patient has experienced an adequate response to treatment, but CDAI score cannot be assessed; and
- 2 Ustekinumab to administered at a dose no greater than 90 mg every 8 weeks.

Note: Indication marked with * is an unapproved indication.

**Initial application — (ulcerative colitis)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Patient is currently on treatment with ustekinumab commenced prior to 1 February 2023 and met all remaining criteria (criterion 2) below at the time of commencing treatment; or
- 2 Both:

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sub	sidised	Generic
\$	Per	1	Manufacturer

continued...

2.1 Patient has active ulcerative colitis; and

2.2 Either:

2.2.1 Patient has had an initial approval for prior biologic therapy for ulcerative colitis and has experienced intolerable side effects or insufficient benefit to meet renewal criteria; or

2.2.2 Both:

2.2.2.1 Patient meets the initiation criteria for prior biologic therapies for ulcerative colitis; and 2.2.2.2 Other biologics for ulcerative colitis are contraindicated.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

1 Either:

1.1 The SCCAI score has reduced by 2 points or more from the SCCAI score since initiation on biologic therapy; or

PUCAI score has reduced by 10 points or more from the PUCAI score since initiation on biologic therapy*; and
 Ustekinumab will be used at a dose no greater than 90 mg intravenously every 8 weeks.

Note: Criterion marked with * is for an unapproved indication.

VEDOLIZUMAB - PCT only - Special Authority see SA2183 below

### ➡SA2183 Special Authority for Subsidy

**Initial application** — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active Crohn's disease; and
- 2 Any of the following:
  - 2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria (unless contraindicated); or
  - 2.2 Patient has a CDAI score of greater than or equal to 300, or HBI score of greater than or equal to 10; or
  - 2.3 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
  - 2.4 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
  - 2.5 Patient has an ileostomy or colostomy, and has intestinal inflammation; and

3 Any of the following:

- 3.1 Patient has tried but experienced an inadequate response to (including lack of initial response and/or loss of initial response) from prior therapy with immunomodulators and corticosteroids; or
- 3.2 Patient has experienced intolerable side effects from immunomodulators and corticosteroids; or
- 3.3 Immunomodulators and corticosteroids are contraindicated.

Renewal — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 CDAI score has reduced by 100 points, or HBI score has reduced by 3 points, from when the patient was initiated on biologic therapy; or
  - 1.2 CDAI score is 150 or less, or HBI is 4 or less; or
  - 1.3 The patient has experienced an adequate response to treatment, but CDAI score and/or HBI score cannot be assessed; and
- 2 Vedolizumab to administered at a dose no greater than 300 mg every 8 weeks.

Initial application - (Crohn's disease - children*) from any relevant practitioner. Approvals valid for 6 months for applications

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Su	bsidised	Generic	
\$	Per	1	Manufacturer	

continued...

meeting the following criteria:

All of the following:

- 1 Paediatric patient has active Crohn's disease; and
- 2 Any of the following:
  - 2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria (unless contraindicated); or
  - 2.2 Patient has a Paediatric Crohn's Disease Activity Index (PCDAI) score of greater than or equal to 30; or
  - 2.3 Patient has extensive small intestine disease; and
- 3 Any of the following:
  - 3.1 Patient has tried but experienced an inadequate response to (including lack of initial response and/or loss of initial response) from prior therapy with immunomodulators and corticosteroids; or
  - 3.2 Patient has experienced intolerable side effects from immunomodulators and corticosteroids; or
  - 3.3 Immunomodulators and corticosteroids are contraindicated.

Note: Indication marked with * is an unapproved indication.

Renewal — (Crohn's disease - children*) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 PCDAI score has reduced by 10 points from when the patient was initiated on biologic therapy; or
  - 1.2 PCDAI score is 15 or less; or
  - 1.3 The patient has experienced an adequate response to treatment, but CDAI score cannot be assessed; and
- 2 Vedolizumab to administered at a dose no greater than 300mg every 8 weeks.

Note: Indication marked with * is an unapproved indication.

**Initial application — (ulcerative colitis)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active ulcerative colitis; and
- 2 Any of the following:
  - 2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria (unless contraindicated); or
  - 2.2 Patient has a SCCAI score is greater than or equal to 4; or
  - 2.3 Patient's PUCAI score is greater than or equal to 20*; and
- 3 Any of the following:
  - 3.1 Patient has tried but experienced an inadequate response to (including lack of initial response and/or loss of initial response) from prior therapy with immunomodulators and corticosteroids; or
  - 3.2 Patient has experienced intolerable side effects from immunomodulators and corticosteroids; or
  - 3.3 Immunomodulators and corticosteroids are contraindicated.

Note: Indication marked with * is an unapproved indication.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

#### Both:

1 Either:

- 1.1 The SCCAI score has reduced by 2 points or more from the SCCAI score since initiation on biologic therapy; or
- 1.2 The PUCAI score has reduced by 10 points or more from the PUCAI score since initiation on biologic therapy *; and
- 2 Vedolizumab will be used at a dose no greater than 300 mg intravenously every 8 weeks.

Note: Indication marked with * is an unapproved indication.

	Subsidy (Manufacturer's Price) \$	Subs Per	Fully idised	Brand or Generic Manufacturer
Programmed Cell Death-1 (PD-1) Inhibitors	φ		•	
ATEZOLIZUMAB – PCT only – Specialist – Special Authority see	SA2443 below			
Inj 60 mg per ml, 20 ml vial	9,503.00	1		ecentriq
Inj 1 mg for ECP	8.08	1 mg	✓ B	axter
■ SA2443 Special Authority for Subsidy	nenetherenu) only	from a ma	diaalar	a alagiat ar any relevant
Initial application — (non-small cell lung cancer second line r practitioner on the recommendation of a medical oncologist. Appl				
criteria:		1013 101 44	piloatio	no meeting the following
All of the following:				
1 Patient has locally advanced or metastatic non-small cell lu	ung cancer; and			
2 Patient has not received prior funded treatment with an im				
3 For patients with non-squamous histology there is docume			ease do	es not express activating
mutations of EGFR or ALK tyrosine kinase unless not pose	sible to ascertain; and	b		
<ul> <li>4 Patient has an ECOG 0-2; and</li> <li>5 Patient has documented disease progression following treater</li> </ul>	atmont with at loast t		of plati	num basad abamatharany
and	alment with at least t	wo cycles	oi piati	num-based chemotherapy
6 Atezolizumab is to be used as monotherapy at a dose of 1	200 ma every three v	veeks (or	equival	ent) for a maximum of
16 weeks; and		(		
7 Baseline measurement of overall tumour burden is docume	ented clinically and ra	adiologica	lly.	
Renewal — (non-small cell lung cancer second line monother				
on the recommendation of a medical oncologist. Approvals valid	for 4 months for appl	ications m	eeting	the following criteria:
All of the following:				
1 Any of the following:				
1.1 Patient's disease has had a complete response to t				
<ul><li>1.2 Patient's disease has had a partial response to trea</li><li>1.3 Patient has stable disease; and</li></ul>	itment; or			
2 Response to treatment in target lesions has been determin	ed by comparable ra	diologic a	eedeem	ent following the most
recent treatment period; and	ice by comparable re	laiologic a	3303311	ient following the most
3 No evidence of disease progression; and				
4 The treatment remains clinically appropriate and patient is	benefitting from treat	tment; and	ł	
5 Atezolizumab to be used at a maximum dose of 1200 mg e				
6 Treatment with atezolizumab to cease after a total duration	of 24 months from o	commence	ement (	or equivalent of 35 cycles
dosed every 3 weeks).	(			and a solid for 0 month. If
Initial application — (unresectable hepatocellular carcinoma)	from any relevant p	ractitioner	. Appro	ovais valid for 6 months for
applications meeting the following criteria: Either:				
1 Patient is currently on treatment with atezolizumab and me	t all remaining criteri	a prior to	comme	ncing treatment: or
2 All of the following:				nong touthon, of
2.1 Patient has locally advanced or metastatic, unresed	table hepatocellular	carcinoma	a; and	
2.2 Patient has preserved liver function (Child-Pugh A)				

- 2.3 Transarterial chemoembolisation (TACE) is unsuitable; and
- 2.4 Any of the following:
  - 2.4.1 Patient has not received prior systemic therapy for the treatment of hepatocellular carcinoma; or
  - 2.4.2 Patient received funded lenvatinib before 1 March 2025; or
  - 2.4.3 Both:
    - 2.4.3.1 Patient has experienced treatment-limiting toxicity from treatment with lenvatinib; and
    - 2.4.3.2 No disease progression since initiation of lenvatinib; and

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

(Man	Subsidy ufacturer's Price)	Subsi	Fully dised	Brand or Generic
· · · · · · · · · · · · · · · · · · ·	\$	Per	1	Manufacturer

continued...

- 2.5 Patient has an ECOG performance status of 0-2; and
- 2.6 To be given in combination with bevacizumab.

Renewal — (unresectable hepatocellular carcinoma) from any relevant practitioner. Approvals valid for 6 months where there is no evidence of disease progression.

Inj 50 mg per ml, 10 ml vial		1	🗸 Imfinzi
Inj 50 mg per ml, 2.4 ml vial		1	🗸 Imfinzi
Inj 1 mg for ECP	9.59	1 mg	<ul> <li>Baxter</li> </ul>

### ⇒SA2425 Special Authority for Subsidy

**Initial application** — (Non-small cell lung cancer) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Either:
  - 1.1 Patient has histologically or cytologically documented stage III, locally advanced, unresectable non-small cell lung cancer (NSCLC); or
  - 1.2 Patient has histologically or cytologically documented stage IIb (T1N2a only), locally advanced, unresectable non-small cell lung cancer (NSCLC); and
- 2 Patient has received two or more cycles of platinum-based chemotherapy concurrently with definitive radiation therapy; and
- 3 Patient has no disease progression following the second or subsequent cycle of platinum-based chemotherapy with definitive radiation therapy treatment; and
- 4 Patient has a ECOG performance status of 0 or 1; and
- 5 Patient has completed last radiation dose within 8 weeks of starting treatment with durvalumab; and
- 6 Patient must not have received prior PD-1 or PD-L1 inhibitor therapy for this condition; and

7 Either:

- 7.1 Durvalumab is to be used at a maximum dose of no greater than 10 mg/kg every 2 weeks; or
- 7.2 Durvalumab is to be used at a flat dose of 1500 mg every 4 weeks; and
- 8 Treatment with durvalumab to cease upon signs of disease progression.

Renewal — (Non-small cell lung cancer) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

1 The treatment remains clinically appropriate and the patient is benefitting from treatment; and

- 2 Either:
  - 2.1 Durvalumab is to be used at a maximum dose of no greater than 10 mg/kg every 2 weeks; or
  - 2.2 Durvalumab is to be used at a flat dose of 1500 mg every 4 weeks; and
- 3 Treatment with durvalumab to cease upon signs of disease progression; and
- 4 Total continuous treatment duration must not exceed 12 months.

NIVOLUMAB - PCT only - Specialist - Special Authority see SA2405 below

Inj 10 mg per ml, 4 ml vial1,051.98	1	<ul> <li>Opdivo</li> </ul>
Inj 10 mg per ml, 10 ml vial2,629.96	1	<ul> <li>Opdivo</li> </ul>
Inj 1 mg for ECP	1 mg	<ul> <li>Baxter</li> </ul>

### ➡SA2405 Special Authority for Subsidy

Initial application only from a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 2 Baseline measurement of overall tumour burden is documented clinically and radiologically; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Either:
  - 4.1 Patient has not received funded pembrolizumab; or
  - 4.2 Both:
    - 4.2.1 Patient has received an initial Special Authority approval for pembrolizumab and has discontinued pembrolizumab within 12 weeks of starting treatment due to intolerance; and
    - 4.2.2 The cancer did not progress while the patient was on pembrolizumab; and
- 5 Documentation confirming that the patient has been informed and acknowledges that funded treatment with nivolumab will not be continued if their disease progresses.

Renewal — (less than 24 months on treatment) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: Either

1 All of the following:

- 1.1 Any of the following:
  - 1.1.1 Patient's disease has had a complete response to treatment; or
  - 1.1.2 Patient's disease has had a partial response to treatment; or
  - 1.1.3 Patient has stable disease; and
- 1.2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period; and
- 1.3 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
- 2 All of the following:
  - 2.1 Patient has previously discontinued treatment with nivolumab for reasons other than severe toxicity or disease progression; and
  - 2.2 Patient has signs of disease progression; and
  - 2.3 Disease has not progressed during previous treatment with nivolumab.

**Renewal — (more than 24 months on treatment)** only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: Both:

- 1 Patient has been on treatment for more than 24 months; and
- 2 Either:
  - 2.1 All of the following:
    - 2.1.1 Any of the following:
      - 2.1.1.1 Patient's disease has had a complete response to treatment; or
      - 2.1.1.2 Patient's disease has had a partial response to treatment; or
      - 2.1.1.3 Patient has stable disease; and
    - 2.1.2 Response to treatment in target lesions has been determined by comparable radiologic or clinical assessment following the most recent treatment period; and
    - 2.1.3 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
  - 2.2 All of the following:
    - 2.2.1 Patient has previously discontinued treatment with nivolumab for reasons other than severe toxicity or disease progression; and
    - 2.2.2 Patient has signs of disease progression; and
    - 2.2.3 Disease has not progressed during previous treatment with nivolumab.

**Initial application** — (Renal cell carcinoma) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria: Either:

Subsidy		Fully	Brand or	
(Manufacturer's F	Price)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

#### continued...

- 1 Patient is currently on treatment with nivolumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
  - 2.1 Patient has metastatic renal-cell carcinoma; and
  - 2.2 The disease is of predominant clear-cell histology; and
  - 2.3 Patient has an ECOG performance score of 0-2; and
  - 2.4 Patient has documented disease progression following one or two previous regimens of antiangiogenic therapy; and
  - 2.5 Nivolumab is to be used as monotherapy at a maximum dose of 240 mg every 2 weeks (or equivalent) and discontinued at disease progression.

Renewal — (Renal cell carcinoma) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Any of the following:
  - 1.1 Patient's disease has had a complete response to treatment; or
  - 1.2 Patient's disease has had a partial response to treatment; or
  - 1.3 Patient has stable disease; and
- 2 No evidence of disease progression; and
- 3 Nivolumab is to be used as monotherapy at a maximum dose of 240 mg every 2 weeks (or equivalent) and discontinued at disease progression.

PEMBROLIZUMAB - PCT only - Specialist - Special Authority see SA2386 below

Inj 25 mg per ml, 4 ml vial	4,680.00	1	<ul> <li>Keytruda</li> </ul>
Inj 1 mg for ECP	47.74	1 mg	<ul> <li>Baxter</li> </ul>

### ⇒SA2386 Special Authority for Subsidy

**Initial application** — (unresectable or metastatic melanoma) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Baseline measurement of overall tumour burden is documented clinically and radiologically; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Either:
  - 4.1 Patient has not received funded nivolumab; or
  - 4.2 Both:
    - 4.2.1 Patient has received an initial Special Authority approval for nivolumab and has discontinued nivolumab within 12 weeks of starting treatment due to intolerance; and
    - 4.2.2 The cancer did not progress while the patient was on nivolumab; and
- 5 Documentation confirming that the patient has been informed and acknowledges that funded treatment with pembrolizumab will not be continued if their disease progresses.

Renewal — (unresectable or metastatic melanoma, less than 24 months on treatment) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 All of the following:
  - 1.1 Any of the following:
    - 1.1.1 Patient's disease has had a complete response to treatment; or
    - 1.1.2 Patient's disease has had a partial response to treatment; or
    - 1.1.3 Patient has stable disease; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🖌	Manufacturer

continued...

- 1.2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period; and
- 1.3 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
- 2 All of the following:
  - 2.1 Patient has previously discontinued treatment with pembrolizumab for reasons other than severe toxicity or disease progression; and
  - 2.2 Patient has signs of disease progression; and
  - 2.3 Disease has not progressed during previous treatment with pembrolizumab.

Renewal — (unresectable or metastatic melanoma, more than 24 months on treatment) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 Patient has been on treatment for more than 24 months; and
- 2 Either:
  - 2.1 All of the following:
    - 2.1.1 Any of the following:
      - 2.1.1.1 Patient's disease has had a complete response to treatment; or
      - 2.1.1.2 Patient's disease has had a partial response to treatment; or
      - 2.1.1.3 Patient has stable disease; and
    - 2.1.2 Response to treatment in target lesions has been determined by comparable radiologic or clinical assessment following the most recent treatment period; and
    - 2.1.3 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
  - 2.2 All of the following:
    - 2.2.1 Patient has previously discontinued treatment with pembrolizumab for reasons other than severe toxicity or disease progression; and
    - 2.2.2 Patient has signs of disease progression; and
    - 2.2.3 Disease has not progressed during previous treatment with pembrolizumab.

**Initial application — (non-small cell lung cancer first-line monotherapy)** only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-small cell lung cancer; and
- 2 Patient has not had chemotherapy for their disease in the palliative setting; and
- 3 Patient has not received prior funded treatment with an immune checkpoint inhibitor for NSCLC; and
- 4 For patients with non-squamous histology there is documentation confirming that the disease does not express activating mutations of EGFR or ALK tyrosine kinase unless not possible to ascertain; and
- 5 Pembrolizumab to be used as monotherapy; and
- 6 Either:
  - 6.1 There is documentation confirming the disease expresses PD-L1 at a level greater than or equal to 50% as determined by a validated test unless not possible to ascertain; or
  - 6.2 Both:
    - 6.2.1 There is documentation confirming the disease expresses PD-L1 at a level greater than or equal to 1% as determined by a validated test unless not possible to ascertain; and
    - 6.2.2 Chemotherapy is determined to be not in the best interest of the patient based on clinician assessment; and
- 7 Patient has an ECOG 0-2; and
- 8 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks; and

Subsidy		Fully	Brand or	
(Manufacturer's Price	)	Subsidised	Generic	
\$	Per	1	Manufacturer	

continued...

9 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Renewal — (non-small cell lung cancer first line monotherapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Any of the following:
  - 1.1 Patient's disease has had a complete response to treatment; or
  - 1.2 Patient's disease has had a partial response to treatment; or
  - 1.3 Patient has stable disease; and
- 2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period; and
- 3 No evidence of disease progression; and
- 4 The treatment remains clinically appropriate and patient is benefitting from treatment; and
- 5 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 6 Treatment with pembrolizumab to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

**Initial application — (non-small cell lung cancer first-line combination therapy)** only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-small cell lung cancer; and
- 2 The patient has not had chemotherapy for their disease in the palliative setting; and
- 3 Patient has not received prior funded treatment with an immune checkpoint inhibitor for NSCLC; and
- 4 For patients with non-squamous histology there is documentation confirming that the disease does not express activating mutations of EGFR or ALK tyrosine kinase unless not possible to ascertain; and
- 5 Pembrolizumab to be used in combination with platinum-based chemotherapy; and
- 6 Patient has an ECOG 0-2; and
- 7 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks; and
- 8 Baseline measurement of overall tumour burden is documented clinically and radiologically.

**Renewal — (non-small cell lung cancer first line combination therapy)** only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 Patient's disease has had a complete response to treatment; or
  - 1.2 Patient's disease has had a partial response to treatment; or
  - 1.3 Patient has stable disease; and
- 2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period; and
- 3 No evidence of disease progression; and
- 4 The treatment remains clinically appropriate and patient is benefitting from treatment; and
- 5 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 6 Treatment with pembrolizumab to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

**Initial application** — (breast cancer, advanced) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: Either:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🖌	Manufacturer

continued...

- 1 Patient is currently on treatment with pembrolizumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
  - 2.1 Either:
    - 2.1.1 Patient has recurrent or de novo unresectable, inoperable locally advanced triple-negative breast cancer (that does not express ER, PR or HER2 IHC3+ or ISH+ [including FISH or other technology]); or
    - 2.1.2 Patient has recurrent or de novo metastatic triple-negative breast cancer (that does not express ER, PR or HER2 IHC3+ or ISH+ [including FISH or other technology]); and
  - 2.2 Patient is treated with palliative intent; and
  - 2.3 Patient's cancer has confirmed PD-L1 Combined Positive Score (CPS) is greater than or equal to 10; and
  - 2.4 Patient has received no prior systemic therapy in the palliative setting; and
  - 2.5 Patient has an ECOG score of 0-2; and
  - 2.6 Pembrolizumab is to be used in combination with chemotherapy; and
  - 2.7 Baseline measurement of overall tumour burden is documented clinically and radiologically; and
  - 2.8 Pembrolizumab is to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks.

Renewal — (breast cancer, advanced) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 Patient's disease has had a complete response to treatment; or
  - 1.2 Patient's disease has had a partial response to treatment; or
  - 1.3 Patient has stable disease; and
- 2 No evidence of disease progression; and
- 3 Response to treatment in target lesions has been determined by a comparable radiologic assessment following the most recent treatment period; and
- 4 Pembrolizumab is to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 5 Treatment with pembrolizumab is to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

**Initial application** — (head and neck squamous cell carcinoma) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria: Either:

- 1 Patient is currently on treatment with pembrolizumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
  - 2.1 Patient has recurrent or metastatic head and neck squamous cell carcinoma of mucosal origin (excluding nasopharyngeal carcinoma) that is incurable by local therapies; and
  - 2.2 Patient has not received prior systemic therapy in the recurrent or metastatic setting; and
  - 2.3 Patient has a positive PD-L1 combined positive score (CPS) of greater than or equal to 1; and
  - 2.4 Patient has an ECOG performance score of 0-2; and
  - 2.5 Either:
    - 2.5.1 Pembrolizumab to be used in combination with platinum-based chemotherapy; or
    - 2.5.2 Pembrolizumab to be used as monotherapy; and
  - 2.6 Pembrolizumab is to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks.

Renewal — (head and neck squamous cell carcinoma) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

1 Any of the following:

Subsidy (Manufacturer's Price)	Fu Subsidis		
 \$	Per	<ul> <li>Manufacturer</li> </ul>	

continued...

- 1.1 Patient's disease has had a complete response to treatment; or
- 1.2 Patient's disease has had a partial response to treatment; or
- 1.3 Patient has stable disease; and
- 2 No evidence of disease progression; and
- 3 Pembrolizumab is to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 4 Treatment with pembrolizumab is to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initial application — (MSI-H/dMMR advanced colorectal cancer) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria: Either:

- 1 Patient is currently on treatment with pembrolizumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
  - 2.1 Either:
    - 2.1.1 Patient has deficient mismatch repair (dMMR) or microsatellite instability-high (MSI-H) metastatic colorectal cancer; or
    - 2.1.2 Patient has deficient mismatch repair (dMMR) or microsatellite instability-high (MSI-H) unresectable colorectal cancer; and
  - 2.2 Patient is treated with palliative intent; and
  - 2.3 Patient has not previously received funded treatment with pembrolizumab; and
  - 2.4 Patient has an ECOG performance score of 0-2; and
  - 2.5 Baseline measurement of overall tumour burden is documented clinically and radiologically; and
  - 2.6 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks.

Renewal — (MSI-H/dMMR advanced colorectal cancer) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 3 Treatment with pembrolizumab is to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

**Initial application** — (Urothelial carcinoma) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria: Either:

- 1 Patient is currently on treatment with pembrolizumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
  - 2.1 Patient has inoperable locally advanced (T4) or metastatic urothelial carcinoma; and
  - 2.2 Patient has an ECOG performance score of 0-2; and
  - 2.3 Patient has documented disease progression following treatment with chemotherapy; and
  - 2.4 Pembrolizumab to be used as monotherapy at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks.

**Renewal** — (Urothelial carcinoma) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Any of the following:
  - 1.1 Patient's disease has had a complete response to treatment; or
  - 1.2 Patient's disease has had a partial response to treatment; or
  - 1.3 Patient has stable disease; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🖌	Manufacturer

continued...

- 2 No evidence of disease progression; and
- 3 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 4 Treatment with pembrolizumab is to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

**Initial application** — (relapsed/refractory Hodgkin lymphoma) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria: Either:

- 1 Patient is currently on treatment with pembrolizumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
  - 2.1 Either:

2.1.1 Both:

- 2.1.1.1 Patient has relapsed/refractory Hodgkin lymphoma after two or more lines of chemotherapy; and
- 2.1.1.2 Patient is ineligible for autologous stem cell transplant; or
- 2.1.2 Patient has relapsed/refractory Hodgkin lymphoma and has previously undergone an autologous stem cell transplant; and
- 2.2 Patient has not previously received funded pembrolizumab; and
- 2.3 Pembrolizumab to be administered at doses no greater than 200 mg once every 3 weeks.

Renewal — (relapsed/refractory Hodgkin lymphoma) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient has received a partial or complete response to pembrolizumab; and
- 2 Treatment with pembrolizumab is to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

### Other Immunosuppressants

CICLOSPORIN			
Cap 25 mg		50	Neoral
Cap 50 mg		50	Neoral
Cap 100 mg		50	Neoral
Oral liq 100 mg per ml		50 ml OP	<ul> <li>Neoral</li> </ul>
EVEROLIMUS - Special Authority see SA2414 below - Re	tail pharmacy		
Wastage claimable			
Tab 10 mg	6,512.29	30	<ul> <li>Afinitor</li> </ul>
Tab 5 mg	4,555.76	30	<ul> <li>Afinitor</li> </ul>

### SA2414 Special Authority for Subsidy

Initial application only from a neurologist or oncologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

1 Patient has tuberous sclerosis; and

2 Patient has progressively enlarging sub-ependymal giant cell astrocytomas (SEGAs) that require treatment.

**Renewal** only from a neurologist or oncologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Documented evidence of SEGA reduction or stabilisation by MRI within the last 3 months; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Everolimus to be discontinued at progression of SEGAs.

Subsidy		Fully	Brand or	
(Manufacturer's Price	e)	Subsidised	Generic	
\$	Per	1	Manufacturer	

#### continued...

**Initial application** — (renal cell carcinoma) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 All of the following:
  - 1.1 The patient has metastatic renal cell carcinoma; and
  - 1.2 The disease is of predominant clear-cell histology; and
  - 1.3 The patient has documented disease progression following one previous line of treatment; and
  - 1.4 The patient has an ECOG performance status of 0-2; and
  - 1.5 Everolimus is to be used in combination with lenvatinib; or
- 2 All of the following:
  - 2.1 Patient has received funded treatment with nivolumab for the second line treatment of metastatic renal cell carcinoma; and
  - 2.2 Patient has experienced treatment limiting toxicity from treatment with nivolumab; and
  - 2.3 Everolimus is to be used in combination with lenvatinib; and
  - 2.4 There is no evidence of disease progression.

Renewal — (renal cell carcinoma) from any relevant practitioner. Approvals valid for 4 months where there is no evidence of disease progression.

SIROLIMUS - Special Authority see SA2270 below - Retail pharmacy

Tab 1 mg		100	<ul> <li>Rapamune</li> </ul>
Tab 2 mg		100	<ul> <li>Rapamune</li> </ul>
Oral liq 1 mg per ml	-	60 ml OP	<ul> <li>Rapamune</li> </ul>

### ► SA2270 Special Authority for Subsidy

Initial application from any medical practitioner. Approvals valid without further renewal unless notified where the drug is to be used for rescue therapy for an organ transplant recipient.

Notes: Rescue therapy defined as unresponsive to calcineurin inhibitor treatment as defined by refractory rejection; or intolerant to calcineurin inhibitor treatment due to any of the following:

- GFR< 30 ml/min; or
- · Rapidly progressive transplant vasculopathy; or
- Rapidly progressive obstructive bronchiolitis; or
- HUS or TTP; or
- Leukoencepthalopathy; or
- Significant malignant disease

Initial application — (severe non-malignant lymphovascular malformations*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe non-malignant lymphovascular malformation*; and
- 2 Any of the following:
  - 2.1 Malformations are not adequately controlled by sclerotherapy and surgery; or
  - 2.2 Malformations are widespread/extensive and sclerotherapy and surgery are not considered clinically appropriate; or
  - 2.3 Sirolimus is to be used to reduce malformation prior to consideration of surgery; and
- 3 Patient is being treated by a specialist lymphovascular malformation multi-disciplinary team; and
- 4 Patient has measurable disease as defined by RECIST version 1.1 (see Note).

Renewal — (severe non-malignant lymphovascular malformations*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 Either:

## ONCOLOGY AGENTS AND IMMUNOSUPPRESSANTS

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

- 1.1 Patient's disease has had either a complete response or a partial response to treatment, or patient has stable disease according to RECIST version 1.1 (see Note); or
- 1.2 Patient's disease has stabilised or responded clinically and disease response to treatment has been clearly documents in patient notes; and
- 2 No evidence of progressive disease; and
- 3 The treatment remains clinically appropriate and the patient is benefitting from the treatment.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer et al. Eur J Cancer 2009;45:228-47)

Indications marked with * are unapproved indications

Initial application — (renal angiomyolipoma(s) associated with tuberous sclerosis complex*) only from a nephrologist or urologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient has tuberous sclerosis complex*; and
- 2 Evidence of renal angiomyolipoma(s) measuring 3 cm or greater and that have shown interval growth.
- Renewal --- (renal angiomyolipoma(s) associated with tuberous sclerosis complex*) from any relevant practitioner.

Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Documented evidence of renal angiomyolipoma reduction or stability by magnetic resonance imaging (MRI) or ultrasound; and
- 2 Demonstrated stabilisation or improvement in renal function; and
- 3 The patient has not experienced angiomyolipoma haemorrhage or significant adverse effects to sirolimus treatment; and
- 4 The treatment remains appropriate and the patient is benefitting from treatment.
- Note: Indications marked with * are unapproved indications

Initial application — (refractory seizures associated with tuberous sclerosis complex*) only from a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has epilepsy with a background of documented tuberous sclerosis complex; and
- 2 Either:
  - 2.1 Both:
    - 2.1.1 Vigabatrin has been trialled and has not adequately controlled seizures; and
    - 2.1.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least two of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); or
  - 2.2 Both:
    - 2.2.1 Vigabatrin is contraindicated; and
    - 2.2.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least three of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); and
- 3 Seizures have a significant impact on quality of life; and
- 4 Patient has been assessed and surgery is considered inappropriate for this patient, or the patient has been assessed and would benefit from mTOR inhibitor treatment prior to surgery.

Note: Those of childbearing age potential are not required to trial phenytoin sodium, sodium valproate, or topiramate. Those who can father children are not required to trial sodium valproate.

Renewal — (refractory seizures associated with tuberous sclerosis complex*) only from a neurologist. Approvals valid for 12 months where demonstrated significant and sustained improvement in seizure rate (e.g. 50% reduction in seizure frequency) or severity and/or patient quality of life compared with baseline prior to starting sirolimus treatment. Note: Indications marked with * are unapproved indications

# ONCOLOGY AGENTS AND IMMUNOSUPPRESSANTS

	Subsidy (Manufacturer's Price) \$		Fully Subsidised	Generic
TACROLIMUS - Special Authority see SA2271 below - Retail ph	narmacy			
Cap 0.5 mg		100	✓	Tacrolimus Sandoz
Cap 0.75 mg		100	✓	Tacrolimus Sandoz
Cap 1 mg		100	✓	Tacrolimus Sandoz
Cap 5 mg	248.20	50	1	Tacrolimus Sandoz

#### SA2271 Special Authority for Subsidy

**Initial application** — (organ transplant) only from a relevant specialist. Approvals valid without further renewal unless notified where the patient is an organ transplant recipient.

Note: Subsidy applies for either primary or rescue therapy.

**Initial application** — (non-transplant indications*) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

- Both: 1 Patient requires long-term systemic immunosuppression; and
  - 2 Fither:
    - ither: 2.1 Ciclosporin has been trialled and discontinued treatment because of unacceptable side effects or inadequate
    - clinical response; or
    - 2.2 Patient is a child with nephrotic syndrome*.

Note: Indications marked with * are unapproved indications

### **JAK** inhibitors

RINVOQ

28

#### ⇒SA2079 Special Authority for Subsidy

Initial application — (Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

#### All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
  - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
  - 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
- 3 Either:
  - 3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or 3.2 Both:
    - 3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a Health NZ Hospital; and 3.2.2 Fither:
      - 3.2.2.1 The patient has experienced intolerable side effects from rituximab; or
      - 3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

**Renewal** — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Either:

- 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer \$ Antiallergy Preparations Allergic Emergencies ADRENALINE - Special Authority see SA2185 below - Retail pharmacy a) Maximum of 2 ini per prescription b) Additional prescriptions limited to replacement of up to two devices prior to expiry, or replacement of used device for treatment of anaphylaxis. 1 OP Epipen Jr 1 OP Epipen ⇒SA2185 Special Authority for Subsidy Initial application — (anaphylaxis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Both: 1 Either: 1.1 Patient has experienced an anaphylactic reaction which has resulted in presentation to a hospital or emergency department: or 1.2 Patient has been assessed to be at significant risk of anaphylaxis by a relevant practitioner; and 2 Patient is not to be prescribed more than two devices in initial prescription. ICATIBANT - Special Authority see SA1558 below - Retail pharmacy Ini 10 ma per ml. 3 ml prefilled svringe......2.668.00 1 Firazvr ⇒SA1558 Special Authority for Subsidy Initial application only from a clinical immunologist or relevant specialist. Approvals valid for 12 months for applications meeting

the following criteria:

Both:

- 1 Supply for anticipated emergency treatment of larvngeal/oro-pharyngeal or severe abdominal attacks of acute hereditary angioedema (HAE) for patients with confirmed diagnosis of C1-esterase inhibitor deficiency; and
- 2 The patient has undergone product training and has agreed upon an action plan for self-administration.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

# Allergy Desensitisation

### ■ SA1367 Special Authority for Subsidy

Initial application only from a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 RAST or skin test positive; and
- 2 Patient has had severe generalised reaction to the sensitising agent.

Renewal only from a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

	Subsidy		Fully	Brand or
	(Manufacturer's Price		Subsidised	
	(Manalacialer 51 field	Per		Manufacturer
	· ·		-	
BEE VENOM ALLERGY TREATMENT – Special Authority see S	SA1367 on the previ	ous pa	<mark>ge</mark> – Reta	il pharmacy
Initiation kit - 1 vial freeze dried venom with diluent		1 OP	✓	VENOX S29
Initiation kit - 5 vials freeze dried venom with diluent		1 OP	1	VENOX S29
Maintenance kit - 1 vial freeze dried venom with diluent	305.00	1 OP	1	VENOX S29
Maintenance kit - 6 vials 120 mcg freeze dried venom, with		1 01	-	
diluent	005 00	1 OP		Venomil S29
		TOP	•	venomii 529
Treatment kit - 1 vial 550 mcg freeze dried venom, 1 diluent				
9 ml, 3 diluent 1.8 ml		1 OP	1	Albey
Treatment kit - 1 vial 550 mcg freeze dried venom, with dilue	nt305.00	1 OP	✓	Hymenoptera S29
(VENOX S29 Initiation kit - 5 vials freeze dried venom with dilue	ent to be delisted 1 N	/av 202	25)	
			,	te the best set
WASP VENOM ALLERGY TREATMENT – Special Authority see	e SA1367 on the pre	evious p	bage – Re	tall pharmacy
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze				
dried polistes venom, 1 diluent 9 ml, 3 diluent 1.8 ml		1 OP	✓	Albey
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze				
dried venom, with diluent		1 OP	1	Hymenoptera S29
Treatment kit (Paper wasp venom) - 6 vials 120 mcg freeze				, ,
dried venom, with diluent	305.00	1 OP	1	Venomil S29
		101	•	Venonini
Treatment kit (Yellow Jacket venom) - 1 vial 550 mcg freeze		4.00		
dried venom, with diluent		1 OP	<b>v</b>	Hymenoptera S29
Treatment kit (Yellow jacket venom) - 1 vial 550 mcg freeze				
dried vespula venom, 1 diluent 9 ml, 3 diluent 1.8 ml		1 OP	1	Albey
Treatment kit (Yellow jacket venom) - 6 vials 120 mcg freeze	•			
dried venom, with diluent		1 OP	1	Venomil S29
Antihistamines				
Antimistanimies				
CETIRIZINE HYDROCHLORIDE				
* Tab 10 mg	1 71	100	1	Zista
* Oral lig 1 mg per ml		200 m		Histaclear
	0.00	200 11		Instacleal
DEXTROCHLORPHENIRAMINE MALEATE				
* Tab 2 mg	2.02	40		
	(8.40)			Polaramine
	1.01	20		
	(5.99)			Polaramine
* Oral lig 2 mg per 5 ml		100 m	I	
	(10.29)			Polaramine
	( )			
FEXOFENADINE HYDROCHLORIDE	4.04	00		
* Tab 60 mg		20		
	(8.23)			Telfast
* Tab 120 mg		30	✓	Fexaclear
	14.22			
	(26.44)			Telfast
* Tab 180 mg	4.10	30	1	Fexaclear
(Telfast Tab 120 mg to be delisted 1 July 2025)				
LORATADINE				
-	4 70	100		Lavafin
* Tab 10 mg		100		Lorafix
* Oral liq 1 mg per ml	1.43	100 m	I 🗸	Haylor syrup

_

	Subsidy (Manufacturer's Prio \$	Fully ce) Subsidised Per <b>⁄</b>	Generic
PROMETHAZINE HYDROCHLORIDE			
* Tab 10 mg	1.39	50 🗸	Allersoothe
* Tab 25 mg	1.58	50 🗸	Allersoothe
* Oral liq 1 mg per 1 ml	3.39		Allersoothe
	10.47	<ul> <li>Image: A start of the start of</li></ul>	Phenergan Elixir
* Inj 25 mg per ml, 2 ml ampoule – Up to 5 inj available o	n a PSO21.09	5 🗸	Hospira
Inhaled Corticosteroids			
BECLOMETHASONE DIPROPIONATE			
Aerosol inhaler, 50 mcg per dose		00 dose OP 🗸	Qvar
Aerosol inhaler, 50 mcg per dose CFC-free		00 dose OP 🗸	Beclazone 50
Aerosol inhaler, 100 mcg per dose		00 dose OP 🗸	Qvar
Aerosol inhaler, 100 mcg per dose CFC-free		00 dose OP 🗸 🗸	Beclazone 100
Aerosol inhaler, 250 mcg per dose CFC-free		00 dose OP 🗸 🗸	Beclazone 250
BUDESONIDE			
Powder for inhalation, 100 mcg per dose		00 dose OP 🗸	Pulmicort
,, ,			Turbuhaler
Powder for inhalation, 200 mcg per dose		00 dose OP	Pulmicort
· · · · · · · · · · · · · · · · · · ·			Turbuhaler
Powder for inhalation, 400 mcg per dose	32.00 2	00 dose OP	Pulmicort
r ender for initialition, ree meg per dece	E		Turbuhaler
FLUTICASONE			
Aerosol inhaler, 50 mcg per dose	7 10 1	20 dose OP 🗸	Flixotide
Powder for inhalation, 50 mcg per dose	······		Flixotide Accuhaler
Powder for inhalation, 100 mcg per dose			Flixotide Accuhaler
Aerosol inhaler, 125 mcg per dose			Flixotide
Aerosol inhaler, 250 mcg per dose			Flixotide
Powder for inhalation, 250 mcg per dose			Flixotide Accuhaler
. ender ter innalation, zoo mog per dooo minimisti			

# Inhaled Long-acting Beta-adrenoceptor Agonists

# EFORMOTEROL FUMARATE DIHYDRATE

ezhaler
ezhaler
aler

	Subsidy (Manufacturer's \$	Price) Subsi Per	Fully Brand or dised Generic ✔ Manufactu	rer
Inhaled Corticosteroids with Long-Acting Beta-	Adrenocep	tor Agonists		
BUDESONIDE WITH EFORMOTEROL				
Powder for inhalation 160 mcg with 4.5 mcg eformoterol	with			
fumarate per dose (equivalent to 200 mcg budesonide w 6 mcg eformoterol fumarate metered dose)		120 dose OP	🗸 DuoResp Sp	hiromax
Powder for inhalation 320 mcg with 9 mcg eformoterol fumar		120 0000 01	· Buoncop of	in onlax
per dose (equivalent to 400 mcg budesonide with 12 mc				
eformoterol fumarate metered dose) - No more than 2	-			
dose per day		120 dose OP	<ul> <li>DuoResp Sp</li> </ul>	piromax
Aerosol inhaler 100 mcg with eformoterol fumarate 6 mcg		120 dose OP 120 dose OP	<ul> <li>Vannair</li> <li>Symbicort</li> </ul>	
Powder for inhalation 100 mcg with eformoterol fumarate 6 n	ncg 33.74	120 00se OP	Turbuhale	er 100/6
Aerosol inhaler 200 mcg with eformoterol fumarate 6 mcg		120 dose OP	✓ Vannair	
Powder for inhalation 200 mcg with eformoterol fumarate 6 n		120 dose OP	✓ Symbicort	
	-		Turbuhale	er 200/6
Powder for inhalation 400 mcg with eformoterol fumarate				
12 mcg – No more than 2 dose per day		60 dose OP	<ul> <li>Symbicort</li> </ul>	
			Turbuhale	er 400/12
FLUTICASONE FUROATE WITH VILANTEROL	44.00			
Powder for inhalation 100 mcg with vilanterol 25 mcg		30 dose OP	<ul> <li>Breo Ellipta</li> </ul>	
FLUTICASONE WITH SALMETEROL Aerosol inhaler 50 mcg with salmeterol 25 mcg	25 70	120 dose OP	<ul> <li>Seretide</li> </ul>	
Aerosol inhaler 125 mcg with salmeterol 25 mcg		120 dose OP 120 dose OP	<ul> <li>Seretide</li> <li>Seretide</li> </ul>	
Powder for inhalation 100 mcg with salmeterol 50 mcg – No		120 0000 01	· Ocicilae	
more than 2 dose per day		60 dose OP	<ul> <li>Seretide Ac</li> </ul>	cuhaler
Powder for inhalation 250 mcg with salmeterol 50 mcg - No				
more than 2 dose per day		60 dose OP	<ul> <li>Seretide Ac</li> </ul>	cuhaler
Beta-Adrenoceptor Agonists				
SALBUTAMOL				
Oral liq 400 mcg per ml	50.00	150 ml	<ul> <li>Ventolin</li> </ul>	
Ventolin to be Principal Supply on 1 May 2025	400.00	10	Mandalla	
Infusion 1 mg per ml, 5 ml Inj 500 mcg per ml, 1 ml – Up to 5 inj available on a PSO		10 5	<ul> <li>✓ Ventolin</li> <li>✓ Ventolin</li> </ul>	
		J	• ventonn	
Inhaled Beta-Adrenoceptor Agonists				
SALBUTAMOL				
Aerosol inhaler, 100 mcg per dose CFC free – Up to 1000		000 I 07	( o ) · · ·	
dose available on a PSO	4.18 (6.80)	200 dose OP	<ul> <li>SalAir</li> <li>Ventolin</li> </ul>	
Nebuliser soln, 1 mg per ml, 2.5 ml ampoule – Up to 30 neb	( )		VENIOIIII	
available on a PSO		20	<ul> <li>Asthalin</li> </ul>	
Nebuliser soln, 2 mg per ml, 2.5 ml ampoule – Up to 30 neb		-		
available on a PSO		20	<ul> <li>Asthalin</li> </ul>	
TERBUTALINE SULPHATE				
Powder for inhalation, 200 mcg per dose (equivalent to				
250 mcg metered dose), breath activated	22.20	120 dose OP	🗸 Bricanyl Tu	rbuhaler

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per <b>v</b>	
Anticholinergic Agents	Ψ		
IPRATROPIUM BROMIDE Aerosol inhaler, 20 mcg per dose CFC-free – Up to 400 dose available on a PSO Nebuliser soln, 250 mcg per ml, 2 ml ampoule – Up to 40 nel available on a PSO		10 20	Atrovent Pharmascience S29 Univent
(Pharmascience ⁶²⁹ Nebuliser soln, 250 mcg per ml, 2 ml ampo		· /	
Inhaled Beta-Adrenoceptor Agonists with Antich	noiinergic Agei	nts	
SALBUTAMOL WITH IPRATROPIUM BROMIDE Aerosol inhaler, 100 mcg with ipratropium bromide, 20 mcg p dose CFC-free Nebuliser soln, 2.5 mg with ipratropium bromide 0.5 mg per vial, 2.5 ml ampoule – Up to 20 neb available on a PSO			Duolin HFA
(Duolin Cipla S29 Nebuliser soln, 2.5 mg with ipratropium bromic		1	Duolin Cipla S29
Long-Acting Muscarinic Antagonists			
<ul> <li>GLYCOPYRRONIUM – Subsidy by endorsement <ul> <li>a) Inhaled glycopyrronium treatment will not be subsidised if umeclidinium.</li> <li>b) Glycopyrronium powder for inhalation 50 mcg per dose is having COPD using spirometry if spirometry is possible, a Powder for inhalation 50 mcg per dose</li></ul></li></ul>	subsidised only for nd the prescription i 61.00 30 o receiving treatmer ve been diagnosed cordingly. Patients endorsed. 	patients who ha is endorsed acco dose OP	ve been diagnosed as ordingly. Seebri Breezhaler ed inhaled glycopyrronium or D using spirometry if bium dispensed before Spiriva Spiriva Respimat ed glycopyrronium or ve been diagnosed as having
<ul> <li>b) Glycopyrronium powder for inhalation 50 mcg per dose is having COPD using spirometry if spirometry is possible, a Powder for inhalation 50 mcg per dose</li></ul>	nd the prescription i 	is endorsed acco dose OP nt with subsidise as having COPI who had tiotrop 30 dose dose OP subsidised inhal patients who have rsed accordingly	ordingly. Seebri Breezhaler ed inhaled glycopyrronium D using spirometry if bium dispensed before Spiriva Spiriva Respimat ed glycopyrronium or ve been diagnosed as ha

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	✓	Manufacturer	

### Long-Acting Muscarinic Antagonists with Long-Acting Beta-Adrenoceptor Agonists

Combination long acting muscarinic antagonist and long acting beta-2 agonist will not be subsidised if patient is also receiving treatment with a combination inhaled corticosteroid and long acting beta-2 agonist.

#### ⇒SA1584 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

1 Patient has been stabilised on a long acting muscarinic antagonist; and

2 The prescriber considers that the patient would receive additional benefit from switching to a combination product. **Renewal** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both: 1 Patient is compliant with the medication: and

Patient is compliant with the medication, and
 Patient has experienced improved COPD symptom control (prescriber determined).

GLYCOPYRRONIUM WITH INDACATEROL - Special Authority se	e SA1584 al	bove – Retail pha	rmacy
Powder for Inhalation 50 mcg with indacaterol 110 mcg	81.00	30 dose OP	<ul> <li>Ultibro Breezhaler</li> </ul>
TIOTROPIUM BROMIDE WITH OLODATEROL - Special Authority	see SA1584	4 <mark>above –</mark> Retail p	harmacy
Soln for inhalation 2.5 mcg with olodaterol 2.5 mcg	81.00	60 dose OP	<ul> <li>Spiolto Respimat</li> </ul>
UMECLIDINIUM WITH VILANTEROL - Special Authority see SA15	i84 above –	Retail pharmacy	
Powder for inhalation 62.5 mcg with vilanterol 25 mcg	77.00	30 dose OP	<ul> <li>Anoro Ellipta</li> </ul>

### Inhaled Corticosteroid with Long-Acting Muscarinic Antagonist and Beta Agonist

BUDESONIDE WITH GLYCOPYRRONIUM AND EFORMOTEROL - Special Authority see SA2421 below - Retail pharmacy

Aerosol inhaler budesonide 160 mcg with glycopyrronium

#### ⇒SA2421 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

1 Patient has a diagnosis of COPD confirmed by spirometry or spirometry has been attempted and technically acceptable results are not possible; and

2 Either:

- 2.1 Both:
  - 2.1.1 Patient is currently receiving an inhaled corticosteroid with long acting beta-2 agonist (ICS/LABA) or a long acting muscarinic antagonist with long acting beta-2 agonist (LAMA/LABA); and
  - 2.1.2 Any of the following:

Clinical criteria:

- 2.1.2.1 Patient has a COPD Assessment Test (CAT) score greater than 10; or
- 2.1.2.2 Patient has had 2 or more exacerbations in the previous 12 months; or
- 2.1.2.3 Patient has had one exacerbation requiring hospitalisation in the previous 12 months; or
- 2.1.2.4 Patient has had an eosinophil count greater than or equal to 0.3 × 10°9 cells/L in the previous 12 months; or
- 2.2 Patient is currently receiving multiple inhaler triple therapy (inhaled corticosteroid with long-acting muscarinic antagonist and long-acting beta-2 agonist ICS/LAMA/LABA) and met at least one of the clinical criteria above prior to commencing multiple inhaler therapy.

### FLUTICASONE FUROATE WITH UMECLIDINIUM AND VILANTEROL – Special Authority see SA2326 below – Retail pharmacy Powder for inhalation fluticasone furoate 100 mcg with

umeclidinium 62.5 mcg and vilanterol 25 mcg...... 104.24 30 dose OP 🖌 Trelegy Ellipta

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

#### ⇒SA2326 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has a diagnosis of COPD confirmed by spirometry or spirometry has been attempted and technically acceptable results are not possible; and
- 2 Either:
  - 2.1 Both:
    - 2.1.1 Patient is currently receiving an inhaled corticosteroid with long acting beta-2 agonist (ICS/LABA) or a long acting muscarinic antagonist with long acting beta-2 agonist (LAMA/LABA); and
    - 2.1.2 Any of the following:

Clinical criteria:

- 2.1.2.1 Patient has a COPD Assessment Test (CAT) score greater than 10; or
- 2.1.2.2 Patient has had 2 or more exacerbations in the previous 12 months; or
- 2.1.2.3 Patient has had one exacerbation requiring hospitalisation in the previous 12 months; or
- 2.1.2.4 Patient has had an eosinophil count greater than or equal to 0.3 × 10⁹ cells/L in the previous 12 months; or
- 2.2 Patient is currently receiving multiple inhaler triple therapy (inhaled corticosteroid with long acting muscarinic antagonist and long acting beta-2 agonist ICS/LAMA/LABA) and met at least one of the clinical criteria above prior to commencing multiple inhaler triple therapy.

### Antifibrotics

NINTEDANIB - Special Authority see SA2012 below - Retail pharmacy

Note: Nintedanib not subsidised in combination with subsidised pirfenidone.

Cap 100 mg	2,554.00	60 OP	<ul> <li>Ofev</li> </ul>
Cap 150 mg	3,870.00	60 OP	<ul> <li>Ofev</li> </ul>

#### ⇒SA2012 Special Authority for Subsidy

**Initial application** — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Nintedanib is to be discontinued at disease progression (See Note); and
- 4 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 5 Any of the following:
  - 5.1 The patient has not previously received treatment with pirfenidone; or
  - 5.2 Patient has previously received pirfenidone, but discontinued pirfenidone within 12 weeks due to intolerance; or
  - 5.3 Patient has previously received pirfenidone, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with pirfenidone).

Renewal — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 3 Nintedanib is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	Generic	
DIDEENIDONIE Datail sharmany Crasiclist Crasicl Authority	ۍ موه 240012 holow	Per	•	Manufacturer	
PIRFENIDONE – Retail pharmacy-Specialist – Special Authority Note: Pirfenidone is not subsidised in combination with subs					
Tab 801 mg	3,645.00	90 OF	P ✓	Esbriet	
Tab 267 mg	1,215.00	90	1	Esbriet	
■SA2013 Special Authority for Subsidy					

#### 013 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note); and
- 4 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 5 Any of the following:
  - 5.1 The patient has not previously received treatment with nintedanib: or
  - 5.2 Patient has previously received nintedanib, but discontinued nintedanib within 12 weeks due to intolerance; or
  - 5.3 Patient has previously received nintedanib, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with nintedanib).

Renewal — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Pirfenidone is not to be used in combination with subsidised nintedanib: and
- 3 Pirfenidone is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

### Leukotriene Receptor Antagonists

#### MONTELUKAST

*	Tab 4 mg	28	<ul> <li>Montelukast Viatris</li> </ul>
	Tab 5 mg	28	<ul> <li>Montelukast Viatris</li> </ul>
*	Tab 10 mg2.90	28	✓ Montelukast Viatris

### Methylxanthines

#### AMINOPHYLLINE

Inj 25 mg per ml, 10 ml ampoule – Up to 5 inj available on a PSO	180.00	5	<ul> <li>DBL Aminophylline</li> </ul>
THEOPHYLLINE			
* Tab long-acting 250 mg		100	<ul> <li>Nuelin-SR</li> </ul>
* Oral liq 80 mg per 15 ml		500 ml	<ul> <li>Nuelin</li> </ul>

### **Mucolytics**

DORNASE ALFA - Special Authority see SA1978 on the next pa	<mark>ige</mark> – Retail pharma	асу
Nebuliser soln. 2.5 mg per 2.5 ml ampoule		6

Pulmozvme

#### SA1978 Special Authority for Subsidy

Initial application - (cystic fibrosis) only from a respiratory physician or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🖌	Manufacturer	

continued

All of the followina:

- 1 Patient has a confirmed diagnosis of cystic fibrosis; and
- 2 Patient has previously undergone a trial with, or is currently being treated with, hypertonic saline; and
- 3 Any of the following:
  - 3.1 Patient has required one or more hospital inpatient respiratory admissions in the previous 12 month period; or
  - 3.2 Patient has had 3 exacerbations due to CF, requiring oral or intravenous (IV) antibiotics in the previous 12 month period: or
  - 3.3 Patient has had 1 exacerbation due to CF, requiring oral or IV antibiotics in the previous 12 month period and a Brasfield score of < 22/25: or
  - 3.4 Patient has a diagnosis of allergic bronchopulmonary aspergillosis (ABPA).

Renewal — (cystic fibrosis) only from a respiratory physician or paediatrician. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient continues to benefit from treatment.

ELEXACAFTOR WITH TEZACAFTOR, IVACAFTOR AND IVACAFTOR - PCT only - Special Authority see SA2196 below

Tab elexacaftor 50 mg with tezacaftor 25 mg, ivacaftor 37.5 mg					
(56) and ivacaftor 75 mg (28)	27,647.39	84 OP	🗸 Trikafta		
Tab elexacaftor 100 mg with tezacaftor 50 mg, ivacaftor 75 mg					
(56) and ivacaftor 150 mg (28)	27,647.39	84 OP	🗸 Trikafta		

#### ➡SA2196 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the followina:

- 1 Patient has been diagnosed with cystic fibrosis; and
- 2 Patient is 6 years of age or older; and
- 3 Fither:
  - 3.1 Patient has two cystic fibrosis-causing mutations in the cystic fibrosis transmembrane regulator (CFTR) gene (one from each parental allele); or
  - 3.2 Patient has a sweat chloride value of at least 60 mmol/L by quantitative pilocarpine iontophoresis or by Macroduct sweat collection system; and

4 Fither

- 4.1 Patient has a heterozygous or homozygous F508del mutation; or
- 4.2 Patient has a G551D mutation or other mutation responsive in vitro to elexacaftor/tezacaftor/ivacaftor (see note a); and
- 5 The treatment must be the sole funded CFTR modulator therapy for this condition: and

6 Treatment with elexacaftor/tezacaftor/ivacaftor must be given concomitantly with standard therapy for this condition.

Notes:

a) Eligible mutations are listed in the Food and Drug Administration (FDA) Trikafta prescribing information https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/212273s004lbl.pdf

#### IVACAFTOR - PCT only - Specialist - Special Authority see SA2017 below

Tab 150 mg		56	Kalydeco
Oral granules 50 mg, sachet		56	Kalydeco
Oral granules 75 mg, sachet	-	56	<ul> <li>Kalydeco</li> </ul>

#### ■ SA2017 Special Authority for Subsidy

Initial application only from a respiratory specialist or paediatrician. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

	Subsidy	Fu	Ily Brand o	r
(N	lanufacturer's Price)	Subsidis	ed Generic	:
	\$	Per	<ul> <li>Manufa</li> </ul>	cturer

continued...

- 1 Patient has been diagnosed with cystic fibrosis; and
- 2 Either:
  - 2.1 Patient must have G551D mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene on at least 1 allele; or
  - 2.2 Patient must have other gating (class III) mutation (G1244E, G1349D, G178R, G551S, S1251N, S1255P, S549N and S549R) in the CFTR gene on at least 1 allele; and
- 3 Patients must have a sweat chloride value of at least 60 mmol/L by quantitative pilocarpine iontophoresis or by Macroduct sweat collection system; and
- 4 Treatment with ivacaftor must be given concomitantly with standard therapy for this condition; and
- 5 Patient must not have an acute upper or lower respiratory infection, pulmonary exacerbation, or changes in therapy (including antibiotics) for pulmonary disease in the last 4 weeks prior to commencing treatment with ivacaftor; and
- 6 The dose of ivacaftor will not exceed one tablet or one sachet twice daily; and
- 7 Applicant has experience and expertise in the management of cystic fibrosis.

#### SODIUM CHLORIDE

Not funded for use as a nasal drop. Soln 7%25.73	90 ml OP	✓ Biomed
Nasal Preparations		
Allergy Prophylactics		
BUDESONIDE		
Metered aqueous nasal spray, 50 mcg per dose2.59	200 dose OP	✓ <u>SteroClear</u>
Metered aqueous nasal spray, 100 mcg per dose2.89	200 dose OP	<ul> <li><u>SteroClear</u></li> </ul>
FLUTICASONE PROPIONATE		
Metered aqueous nasal spray, 50 mcg per dose1.98	120 dose OP	<ul> <li>Flixonase Hayfever &amp; Allergy</li> </ul>
IPRATROPIUM BROMIDE		
Aqueous nasal spray, 0.03%5.23	15 ml OP	<ul> <li>Univent</li> </ul>
Respiratory Devices		
MASK FOR SPACER DEVICE		
a) Up to 50 dev available on a PSO		
b) Only on a PSO		
c) Only for children aged six years and under		
Small	1	<ul> <li>e-chamber Mask</li> </ul>
PEAK FLOW METER		
a) Up to 25 dev available on a PSO		
b) Only on a PSO Low range9.54	1	Mini-Wright AFS
Low range	·	Low Range
Normal range9.54	1	<ul> <li>Mini-Wright Standard</li> </ul>

	Subsidy (Manufacturer's Price) \$	Sul Per	Fully osidised	Brand or Generic Manufacturer
SPACER DEVICE				
a) Up to 50 dev available on a PSO				
b) Only on a PSO				
220 ml (single patient)	3.65	1	🗸 e	-chamber Turbo
510 ml (single patient)	5.95	1	✔ e	-chamber La Grande
800 ml	6.50	1	🗸 V	olumatic
Respiratory Stimulants				
CAFFEINE CITRATE Oral liq 20 mg per ml (10 mg base per ml)		5 ml OP	✓ В	liomed

	Subsidy		Fully Brand or
	(Manufacturer's Pi	rice) Subs	idised Generic
	\$	Per	✓ Manufacturer
Ear Preparations			
FLUMETASONE PIVALATE			
Ear drops 0.02% with clioquinol 1%	4.46	7.5 ml OP	<ul> <li>Locacorten-Viaform ED's</li> </ul>
			<ul> <li>Locorten-Vioform</li> </ul>
TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYC	IN AND NYSTAT	IN	
Ear drops 1 mg with nystatin 100,000 u, neomycin sulphate			
2.5 mg and gramicidin 250 mcg per g	5.16	7.5 ml OP	<ul> <li>Kenacomb</li> </ul>
Ear/Eye Preparations			
DEXAMETHASONE WITH FRAMYCETIN AND GRAMICIDIN			
Ear/Eye drops 500 mcg with framycetin sulphate 5 mg and			
gramicidin 50 mcg per ml	4.50	8 ml OP	
	(9.27)		Otodex S29
	(9.27)		Sofradex
FRAMYCETIN SULPHATE	4.40	0	
Ear/Eye drops 0.5%	4.13 (8.65)	8 ml OP	Soframycin
	(8.05)		Sonaniyoni
Eye Preparations			
Eye preparations are only funded for use in the eye, unless expli-	citly stated otherv	vise.	
Anti-Infective Preparations			
ACICLOVIR			_
* Eye oint 3%	15.89	4.5 g OP	✓ <u>ViruPOS</u>
CHLORAMPHENICOL			
Eye oint 1%		5 g OP	Devatis     Oblansia
Eye drops 0.5% Funded for use in the ear*. Indications marked with * ar		10 ml OP	✓ Chlorsig
CIPROFLOXACIN	e unapproved me	iloations.	
Eye drops 0.3% – Subsidy by endorsement	10.85	5 ml OP	<ul> <li>Ciprofloxacin Teva</li> </ul>
When prescribed for the treatment of bacterial keratitis of			
for the second line treatment of chronic suppurative otitis			
Note: Indication marked with a * is an unapproved indic	ation.		
SODIUM FUSIDATE [FUSIDIC ACID]			
Eye drops 1%	5.29	5 g OP	<ul> <li>Fucithalmic</li> </ul>
			<ul> <li>Fucithalmic S29 S29</li> </ul>
TOBRAMYCIN	10.45	25 a OP	
Eye oint 0.3% Eye drops 0.3%		3.5 g OP 5 ml OP	<ul> <li>✓ Tobrex</li> <li>✓ Tobrex</li> </ul>
			100107

	Subsidy (Manufacturer's Pr \$	ice) Subs Per	Fully Brand or idised Generic ✓ Manufacturer
Corticosteroids and Other Anti-Inflammatory F	Preparations		
DEXAMETHASONE			
* Eye oint 0.1%	5.86	3.5 g OP	<ul> <li>Maxidex</li> </ul>
* Eye drops 0.1%		5 ml OP	<ul> <li>Maxidex</li> </ul>
Ocular implant 700 mcg - Special Authority see SA1680 b	elow		
- Retail pharmacy		1	<ul> <li>Ozurdex</li> </ul>
➡SA1680 Special Authority for Subsidy			
Initial application — (Diabetic macular oedema) only from a meeting the following criteria: All of the following: 1 Patient has diabetic macular oedema with pseudophakid 2 Patient has reduced visual acuity of between 6/9 - 6/48 v 3 Either: 3.1 Patient's disease has progressed despite 3 inject 3.2 Patient is unsuitable or contraindicated to treatme 4 Dexamethasone implants are to be administered not mo	c lens; and with functional awar tions with bevacizur ent with anti-VEGF	reness of redu nab; or agents; and	iction in vision; and
maximum of 3 implants per eye per year.	. ,	,	
Renewal — (Diabetic macular oedema) only from an ophthal the following criteria: Both:	mologist. Approval	ls valid for 12	months for applications meeting
1 Patient's vision is stable or has improved (prescriber det	ermined): and		
<ol> <li>Praticity vision is stable of has improved (prescriber det 2 Dexamethasone implants are to be administered not mo maximum of 3 implants per eye per year.</li> </ol>		once every 4 r	nonths into each eye, and up to
Initial application — (Women of child bearing age with diab valid for 12 months for applications meeting the following criterial and the following criteria		ma) only fror	n an ophthalmologist. Approvals

All of the following:

- 1 Patient has diabetic macular oedema; and
- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- 3 Patient is of child bearing potential and has not yet completed a family; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Patient is of child bearing potential and has not yet completed a family; and
- 3 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

#### DEXAMETHASONE WITH NEOMYCIN SULPHATE AND POLYMYXIN B SULPHATE

¥ Eye oint 0.1% with neomycin sulphate 0.35% and polymyxin b sulphate 6,000 u per g	3.5 g OP	✓ Maxitrol
¥ Eye drops 0.1% with neomycin sulphate 0.35% and polymyxin b sulphate 6,000 u per ml	5 ml OP	✓ Maxitrol
DICLOFENAC SODIUM		
Eye drops 0.1%, single dose1.85 5.54	10 dose 30 dose	<ul> <li>Diclofenac Devatis</li> <li>Diclofenac Devatis</li> </ul>
FLUOROMETHOLONE		
* Eye drops 0.1%	5 ml OP	<ul><li>✓ FML</li><li>✓ Flucon</li></ul>

Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

SENSORY ORGANS

# SENSORY ORGANS

	Subsidy (Manufacturer's P	rice) Subs	Fully Brand or sidised Generic
	(Manulacturer 3 1 \$	Per	Manufacturer
EVOCABASTINE			
Eye drops 0.5 mg per ml		4 ml OP	
	(10.34)		Livostin
ODOXAMIDE	0.74		
Eye drops 0.1%	8.71	10 ml OP	<ul> <li>Lomide</li> </ul>
	0.00	0	. Hauna
Eye drops 0.3% llevro Eye drops 0.3% to be delisted 1 July 2025)	8.80	3 ml OP	✓ Ilevro
PREDNISOLONE ACETATE			
Eye drops 1%	6 92	10 ml OP	Prednisolone-AFT
	7.00	5 ml OP	Pred Forte
REDNISOLONE SODIUM PHOSPHATE – Special Authority s	ee SA1715 below	– Retail pharr	nacy
Eye drops 0.5%, single dose (preservative free)		20 dose	✓ Minims
			Prednisolone
SA1715 Special Authority for Subsidy			
nitial application only from an ophthalmologist or optometrist.	Approvals valid for	or 6 months for	r applications meeting the
ollowing criteria:			
Both:			
<ol> <li>Patient has severe inflammation; and</li> <li>Patient has a confirmed allergic reaction to preservative i</li> </ol>	in eve drone		
Renewal from any relevant practitioner. Approvals valid for 6 m		reatment rema	ains appropriate and the patien
enefiting from treatment.			
ODIUM CROMOGLICATE			
Eye drops 2%	2.62	10 ml OP	✓ Allerfix
Glaucoma Preparations - Beta Blockers			
BETAXOLOL			
₭ Eye drops 0.25%		5 ml OP	<ul> <li>Betoptic S</li> </ul>
₭ Eye drops 0.5%		5 ml OP	<ul> <li>Betoptic</li> </ul>
Betoptic S Eye drops 0.25% to be delisted 1 December 2025)			
Betoptic Eye drops 0.5% to be delisted 1 December 2025)			
IMOLOL			
₭ Eye drops 0.25%		5 ml OP	Arrow-Timolol
€ Eye drops 0.5%	2.50	5 ml OP	Arrow-Timolol
	Inhibitors		
Glaucoma Preparations - Carbonic Anhydrase	minipitors		
Glaucoma Preparations - Carbonic Anhydrase			
•		100	Diamox
CETAZOLAMIDE		100	<ul> <li>Diamox</li> </ul>
CETAZOLAMIDE K Tab 250 mg		100 5 ml OP	<ul> <li>✓ Diamox</li> <li>✓ <u>Azopt</u></li> </ul>
CETAZOLAMIDE ≰ Tab 250 mg RINZOLAMIDE ≰ Eye drops 1% DORZOLAMIDE WITH TIMOLOL	17.03		
CETAZOLAMIDE ≰ Tab 250 mg RINZOLAMIDE ≰ Eye drops 1%	17.03		
CETAZOLAMIDE ≰ Tab 250 mg RINZOLAMIDE ≰ Eye drops 1% DORZOLAMIDE WITH TIMOLOL		5 ml OP	✓ <u>Azopt</u>
CETAZOLAMIDE Tab 250 mg RINZOLAMIDE Eye drops 1% ORZOLAMIDE WITH TIMOLOL Eve drops 2% with timolol 0.5% Glaucoma Preparations - Prostaglandin Analog		5 ml OP	✓ <u>Azopt</u>
CETAZOLAMIDE ≰ Tab 250 mg RINZOLAMIDE ≰ Eye drops 1% DORZOLAMIDE WITH TIMOLOL ≰ Eye drops 2% with timolol 0.5%		5 ml OP	✓ <u>Azopt</u>

### SENSORY ORGANS

	Subsidy (Manufacturer's F \$	Price) Subs Per	Fully Brand or sidised Generic ✓ Manufacturer
LATANOPROST			
¥ Eye drops 0.005%	2.08	2.5 ml OP	✓ <u>Teva</u>
TRAVOPROST			
* Eye drops 0.004%	6.80	2.5 ml OP	✓ <u>Travatan</u>
Glaucoma Preparations - Other			
BRIMONIDINE TARTRATE			
* Eye drops 0.2%	5.16	5 ml OP	Arrow-Brimonidine
BRIMONIDINE TARTRATE WITH TIMOLOL MALEATE			
* Eye drops 0.2% with timolol maleate 0.5%	7.13	5 ml OP	<ul> <li>Combigan</li> </ul>
LATANOPROST WITH TIMOLOL			-
* Eye drops 0.005% with timolol 0.5%	4.95	2.5 ml OP	Arrow - Lattim
PILOCARPINE HYDROCHLORIDE			
* Eye drops 1%	4.26	15 ml OP	Isopto Carpine
* Eye drops 2%		15 ml OP	Isopto Carpine
* Eye drops 4%		15 ml OP	<ul> <li>Isopto Carpine</li> </ul>
Subsidised for oral use pursuant to the Standard Form	ulae.		
PILOCARPINE NITRATE			
* Eye drops 2% single dose – Special Authority see SA0895			
below – Retail pharmacy		20 dose	<ul> <li>Minims Pilocarpine</li> </ul>
SA0895 Special Authority for Subsidy			

**Initial application** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

1 Patient has to use an unpreserved solution due to an allergy to the preservative; or

2 Patient wears soft contact lenses.

Note: Minims for a general practice are considered to be "tools of trade" and are not approved as special authority items. **Renewal** from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

### **Mydriatics and Cycloplegics**

ATROPINE SULPHATE * Eye drops 1%	 15 ml OP	✓ Atropt
CYCLOPENTOLATE HYDROCHLORIDE * Eye drops 1%	 15 ml OP	<ul> <li>Cyclogyl</li> </ul>
TROPICAMIDE           *         Eye drops 0.5%           *         Eye drops 1%	15 ml OP 15 ml OP	<ul><li>✓ Mydriacyl</li><li>✓ Mydriacyl</li></ul>
Preparations for Tear Deficiency		

For acetylcysteine eye drops refer Standard Formulae, page 273		
HYPROMELLOSE		
* Eye drops 0.5%	15 ml OP	<ul> <li>Methopt</li> </ul>
HYPROMELLOSE WITH DEXTRAN		
* Eye drops 0.3% with dextran 0.1%2.30	15 ml OP	<ul> <li>Poly-Tears</li> </ul>

	Subsidy (Manufacturer's Pric \$	e) Per	Fully Subsidised	Brand or Generic Manufacturer
Preservative Free Ocular Lubricants				
► SA2431 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals vali the following criteria: Both:	d without further re	newal ur	nless notifie	d for applications meeting
<ol> <li>Confirmed diagnosis by slit lamp or Schirmer test of sever</li> <li>Either:</li> <li>2.1 Patient is using eye drops more than four times da</li> <li>2.2 Patient has had a confirmed allergic reaction to pro-</li> </ol>	ily on a regular bas	sis; or		
CARBOMER – Special Authority see SA2431 above – Retail ph Ophthalmic gel 0.3%, 0.5 g (Poly-Gel Ophthalmic gel 0.3%, 0.5 g to be delisted 1 July 2025)	8.25	30	✔ P	oly-Gel
POLYETHYLENE GLYCOL 400 AND PROPYLENE GLYCOL – Eye drops 0.4% and propylene glycol 0.3%, 0.8 ml		ee <mark>SA2</mark> 4 30		- Retail pharmacy systane Unit Dose
SODIUM HYALURONATE [HYALURONIC ACID] – Special Auth Eye drops 1 mg per ml Hylo-Fresh has a 6 month expiry after opening. The Ph month is not relevant and therefore only the prescribed of	13.58 armacy Procedures	10 ml O Manua	P ✓ <u>H</u> I restriction	l <b>ylo-Fresh</b> allowing one bottle per
Other Eye Preparations				
NAPHAZOLINE HYDROCHLORIDE * Eye drops 0.1%	5.65	15 ml O	P <b>√ <u>A</u></b>	Ibalon
OLOPATADINE Eye drops 0.1%	2.17	5 ml Ol	∽ ✓ ₫	Nopatadine Teva
PARAFFIN LIQUID WITH WOOL FAT * Eye oint 3% with wool fat 3%	3.63	3.5 g O	P <b>√ P</b>	oly-Visc
RETINOL PALMITATE				

5 g OP

VitA-POS

VARIOUS

	(Manufacturer's P	rice) Subs Per	sidised Generic Manufacturer
<i>Vov:</i>	\$	Fei	• Manulacturer
Various			
HARMACY SERVICES Brand switch feea) May only be claimed once per patient.	4.50	1 fee	<ul> <li>BSF Dasatinib-Teva</li> </ul>
b) The Pharmacode for BSF Dasatinib-Teva is 2700441	- see also page	164	
Immunisation administration fee - flu		1 fee	<ul> <li>Immunisation - Flu</li> </ul>
Immunisation administration fee - other	0.00	1 fee	<ul> <li>Immunisation Other</li> </ul>
Immunisation co-administration fee - flu and shingles	0.00	1 fee	<ul> <li>Immunisation Flu and Shingles</li> </ul>
3SF Dasatinib-Teva Brand switch fee to be delisted 1 June 2025	;)		
Agents Used in the Treatment of Poisonings			
Antidotes			
CETYLCYSTEINE			
Inj 200 mg per ml, 10 ml ampoule		10	<ul> <li>DBL Acetylcysteine</li> </ul>
	52.88		✓ Martindale Pharma
DBL Acetylcysteine to be Principal Supply on 1 April 2029 Martindale Pharma Inj 200 mg per ml, 10 ml ampoule to be delist		2025)	
ALOXONE HYDROCHLORIDE			
a) Up to 10 inj available on a PSO			
b) Only on a PSO			
Inj 400 mcg per ml, 1 ml ampoule	13.29	5	<ul> <li>DBL Naloxone Hydrochloride</li> </ul>
	35.26	10	✓ Hameln
DBL Naloxone Hydrochloride to be Principal Supply on 1 Hameln Inj 400 mcg per ml, 1 ml ampoule to be delisted 1 April 2	April 2025		
Removal and Elimination			
HARCOAL			
• Oral liq 50 g per 250 ml	43.50	250 ml OP	<ul> <li>Carbosorb-X</li> </ul>
<ul> <li>a) Up to 250 ml available on a PSO</li> <li>b) Only on a PSO</li> </ul>			
EFERASIROX – Special Authority see SA1492 below – Retail p	oharmacv		
Wastage claimable	,		
Tab 125 mg dispersible	276.00	28	<ul> <li>Exjade</li> </ul>
Tab 250 mg dispersible		28	<ul> <li>Exjade</li> </ul>
Tab 500 mg dispersible		28	<ul> <li>Exjade</li> </ul>
SA1492 Special Authority for Subsidy			
itial application only from a haematologist. Approvals valid for	2 vears for appl	ications meeti	ng the following criteria:
Il of the following:	- )		

 $2\;$  Deferasirox is to be given at a daily dose not exceeding 40 mg/kg/day; and

3 Any of the following:

3.1 Treatment with maximum tolerated doses of deferiprone monotherapy or deferiprone and desferrioxamine

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

Subsidy Fu		Fully	Brand or
(Manufacturer's Price) Subsidis		sidised	Generic
\$	Per	1	

- combination therapy have proven ineffective as measured by serum ferritin levels, liver or cardiac MRI T2*; or
- 3.2 Treatment with deferiprone has resulted in severe persistent vomiting or diarrhoea; or
- 3.3 Treatment with deferiprone has resulted in arthritis; or
- 3.4 Treatment with deferiprone is contraindicated due to a history of agranulocytosis (defined as an absolute neutrophil count (ANC) of < 0.5 cells per μL) or recurrent episodes (greater than 2 episodes) of moderate neutropenia (ANC 0.5 1.0 cells per μL).</p>

Renewal only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria:

- Either:
  - 1 For the first renewal following 2 years of therapy, the treatment has been tolerated and has resulted in clinical improvement in all three parameters namely serum ferritin, cardiac MRI T2* and liver MRI T2* levels; or
  - 2 For subsequent renewals, the treatment has been tolerated and has resulted in clinical stability or continued improvement in all three parameters namely serum ferritin, cardiac MRI T2* and liver MRI T2* levels.

DEFERIPRONE – Special Authority see SA1480 below – Retail pharmacy

Tab 500 mg	100	<ul> <li>Ferriprox</li> </ul>
Oral liq 100 mg per 1 ml	 250 ml OP	<ul> <li>Ferriprox</li> </ul>

#### ► SA1480 Special Authority for Subsidy

**Initial application** only from a haematologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; or
- 2 The patient has been diagnosed with chronic iron overload due to acquired red cell aplasia.

DESFERRIOXAMINE MESILATE  * Inj 500 mg vial	151.31	10	<ul> <li>Deferoxamine Pfizer</li> <li>S29 S29</li> </ul>
	332.88		✓ DBL Desferrioxamine Mesylate for Inj BP
SODIUM CALCIUM EDETATE			
* Inj 200 mg per ml, 5 ml	53.31	6	
	(156.71)		Calcium Disodium Versenate

# **Standard Formulae**

ACETYLCYSTEINE EYE DROPS Acetylcysteine inj 200 mg per ml, 10 ml	qs	PHENOBARBITONE SODIUM PAEDIATRIC ORAL mg per ml)	LIQUID (10
Suitable eye drop base	qs	Phenobarbitone Sodium	400 mg
CODEINE LINCTUS (3 mg per 5 ml) Codeine phosphate	60 mg	Glycerol BP Water	4 ml to 40 ml
Glycerol	40 ml	PILOCARPINE ORAL LIQUID	
Preservative	qs	Pilocarpine 4% eye drops	qs
Water	to 100 ml	Preservative	qs
CODEINE LINCTUS (15 mg per 5 ml)		Water (Preservative should be used if quantity supplied is	to 500 ml
Codeine phosphate Glycerol	300 mg 40 ml	than 5 days.)	
Preservative	qs	SALIVA SUBSTITUTE FORMULA	
Water	to 100 ml	Methylcellulose	5 g
FOLINIC MOUTHWASH		Preservative Water	qs to 500 ml
Calcium folinate 15 mg tab	1 tab	(Preservative should be used if quantity supplied is	
Preservative	qs	than 5 days. Maximum 500 ml per prescription.)	
Water	to 500 ml		
(Preservative should be used if quantity supplied is than 5 days. Maximum 500 ml per prescription.)	for more	SODIUM CHLORIDE ORAL LIQUID Sodium chloride inj 23.4%, 20 ml Water	qs qs
METHYL HYDROXYBENZOATE 10% SOLUTION		(Only funded if prescribed for treatment of hyponatr	
Methyl hydroxybenzoate	10 g		,
Propylene glycol	to 100 ml	VANCOMYCIN ORAL SOLUTION (25 mg per ml) Vancomycin 500 mg injection	5 vials
(Use 1 ml of the 10% solution per 100 ml of oral liqu	lid mixture)	Glycerin with sucrose suspension	37.5 ml
OMEPRAZOLE SUSPENSION		Water	to 100 ml
Omeprazole capsules or powder	qs	(Only funded if prescribed for treatment of Clostridiu	um difficile
Sodium bicarbonate powder BP	8.4 g	following metronidazole failure)	
Water	to 100 ml		
PHENOBARBITONE ORAL LIQUID Phenobarbitone Sodium	1 g		
Glycerol BP	70 ml		
Water	to 100 ml		

# EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

	Subsidy		Fully	Brand or
	(Manufacturer's Pric	e) Subs	idised	Generic
	\$	Per	✓	Manufacturer
Extemporaneously Compounded Preparations	and Galenicals	\$		
CODEINE PHOSPHATE - Safety medicine; prescriber may det	ermine dispensing f	requency		
Powder – Only in combination		25 q		
	(90.09)	20 9	г	Douglas
Only in extemporaneously compounded codeine linctus	( /			Douglas
	•			
COLLODION FLEXIBLE				
Note: This product is no longer being manufactured by the	supplier and will be	delisted from	n the S	chedule at a date to be
determined.				
Collodion flexible		100 ml	✓ I	PSM
COMPOUND HYDROXYBENZOATE - Only in combination				
Only in extemporaneously compounded oral mixtures.				
	20.00	100 ml		Midwest
Soln		100 mi	• 1	wiuwest
GLYCERIN WITH SODIUM SACCHARIN – Only in combination				
Only in combination with Ora-Plus or when used in the vanc	omycin oral Iquuid S	Standard For	mulae	
Suspension		473 ml	✓ (	Ora-Sweet SF
GLYCERIN WITH SUCROSE – Only in combination				
Only in combination with Ora-Plus or when used in the vanc	omvoin oral lauuid 9	Standard For	mulaa	
Suspension		473 ml		Ora-Sweet
•		4/3 111	• (	Old-Sweel
GLYCEROL				
* Liquid – Only in combination	3.23	500 ml	✓ I	healthE Glycerol BP
Only in extemporaneously compounded oral liquid prep	arations.			
METHYL HYDROXYBENZOATE				
Powder	8 98	25 g	<b>1</b>	Midwest
	0.00	20 g	• •	Mawest
METHYLCELLULOSE				
Powder		100 g	-	MidWest
Suspension – Only in combination		473 ml	✓ (	Ora-Plus
METHYLCELLULOSE WITH GLYCERIN AND SODIUM SACCH	ARIN – Only in co	mbination		
Suspension		473 ml	✓ (	Ora-Blend SF
METHYLCELLULOSE WITH GLYCERIN AND SUCROSE - On		470		
Suspension		473 ml	•	Ora-Blend
PHENOBARBITONE SODIUM				
Powder – Only in combination		10 g	✓ I	MidWest
-	325.00	100 g	✓ 1	MidWest
Only in children up to 12 years		Ū		
PROPYLENE GLYCOL				
Only in extemporaneously compounded methyl hydroxyben:	zoato 10% colution			
Liq		500 ml		Midwest
		500 mi	• 1	wiuwest
SODIUM BICARBONATE				
Powder BP – Only in combination		500 g	✓ I	Midwest
Only in extemporaneously compounded omeprazole an	d lansoprazole susp	pension.		
SYRUP (PHARMACEUTICAL GRADE) - Only in combination				
Only in extemporaneously compounded oral liquid preparati	one			
Lig		500 ml		Midwest
	14.90	300 111	• 1	MIGWESI
WATER				
Tap – Only in combination	0.00	1 ml	✓ 1	Tap water

### SECTION D: SPECIAL FOODS

Fully

Subsidy (Manufacturer's Price)

\$

Subsidised Per ✓ Brand or Generic Manufacturer

# **Nutrient Modules**

#### Carbohydrate

#### ⇒SA1930 Special Authority for Subsidy

**Initial application** — (Cystic fibrosis or kidney disease) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria: Fither:

- 1 cystic fibrosis; or
- 2 chronic kidney disease.

**Initial application** — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 cancer in children; or
- 2 cancers affecting alimentary tract where there are malabsorption problems in patients over the age of 20 years; or
- 3 faltering growth in an infant/child; or
- 4 bronchopulmonary dysplasia; or
- 5 premature and post premature infant; or
- 6 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Initial application** — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified where the patient has inborn errors of metabolism. **Renewal** — (Cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general

practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

. Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

CARBOHYDRATE SUPPLEMENT – Special Authority see SA1930 above – Hospital pharmacy [HP3]

Powder	 	6.72	400 g OP	<ul> <li>Polycal</li> </ul>

### **Carbohydrate And Fat**

#### ⇒SA1376 Special Authority for Subsidy

**Initial application** — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria: Both:

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic	
\$	Per		Manufacturer	

- 1 Infant or child aged four years or under; and
- 2 cystic fibrosis.

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 infant or child aged four years or under; and
- 2 Any of the following:
  - 2.1 cancer in children; or
  - 2.2 faltering growth; or
  - 2.3 bronchopulmonary dysplasia; or
  - 2.4 premature and post premature infants.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

CARBOHYDRATE AND FAT S	SUPPLEMENT - Special Author	ity see SA1376 on t	the previous page	ge -	Hospital pharmacy [HP3]
Powder (neutral)			400 g OP	1	Duocal Super
			-		Soluble Powder

### Fat

#### ⇒SA2204 Special Authority for Subsidy

Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified where the patient has an inborn error of metabolism. Initial application — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 faltering growth in an infant/child; or
- 2 bronchopulmonary dysplasia; or
- 3 fat malabsorption; or
- 4 lymphangiectasia; or
- 5 short bowel syndrome; or
- 6 infants with necrotising enterocolitis; or
- 7 biliary atresia; or
- 8 for use in a ketogenic diet; or
- 9 chyle leak; or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

- 10 ascites; or
- 11 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal — (Indications other than inborn errors of metabolism)** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT SUPPLEMENT - Special Authority see SA2204 on the	he previous page – Hos	spital pharmacy	[HP3]
Emulsion (neutral)		200 ml OP	<ul> <li>Calogen</li> </ul>
	38.44	500 ml OP	<ul> <li>Calogen</li> </ul>
Emulsion (strawberry)	15.38		<ul> <li>Calogen</li> </ul>
Oil		500 ml OP	<ul> <li>MCT oil (Nutricia)</li> </ul>
MCT Emulsion, 250 ml		4 OP	<ul> <li>Liquigen</li> </ul>

### Protein

#### ⇒SA1524 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 protein losing enteropathy; or
- 2 high protein needs; or
- 3 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

PROTEIN SUPPLEMENT - Sp	pecial Authority see SA1524 above - Hospital ph	narmacy [HP3]	
Powder		227 g OP	<ul> <li>Resource</li> </ul>
			Beneprotein
	13.82	225 g OP	<ul> <li>Protifar</li> </ul>

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

# **Oral and Enteral Feeds**

### **Diabetic Products**

#### ⇒SA1095 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is a type I or and II diabetic who is suffering weight loss and malnutrition that requires nutritional support. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

DIABETIC ENTERAL FEED 1KCAL/ML - Special Authority see	SA1095 above -	- Hospital pharn	nacy [HP3]
Liquid	4.65	500 ml OP	<ul> <li>Glucerna Select</li> </ul>
DIABETIC ORAL FEED 1KCAL/ML - Special Authority see SA1	095 above – Ho	spital pharmacy	[HP3]
Liquid (strawberry)	2.25	200 ml OP	✓ Diasip
Liquid (vanilla)	2.10	200 ml OP	<ul> <li>Nutren Diabetes</li> </ul>
	2.25		🗸 Diasip

### **Fat Modified Products**

#### ⇒SA2205 Special Authority for Subsidy

Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified where the patient has an inborn error of metabolism. Initial application — (Indications other than errors of inborn metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Patient has a chyle leak; or
- 2 Modified as a modular feed, made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule, for adults.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT MODIFIED FEED	- Special Authority see	e SA2205 above – Hospital pha	armacy [HP3]	
Powder			0 400 g OP	🗸 Monogen

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsic	lised	Generic
\$	Per	✓	Manufacturer

### Paediatric Products For Children Awaiting Liver Transplant

#### ⇒SA1098 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) who requires a liver transplant.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

### Paediatric Products For Children With Chronic Renal Failure

#### ⇒SA1099 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) with acute or chronic kidney disease.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL FEED 1KCAL/ML - Spec	cial Authority see SA1099 above - H	Hospital pharmacy	/ [HP3]
Powder		400 g OP	<ul> <li>Kindergen</li> </ul>

### **Paediatric Products**

#### ⇒SA1379 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Child is aged one to ten years; and
- 2 Any of the following:
  - 2.1 the child is being fed via a tube or a tube is to be inserted for the purposes of feeding; or
  - 2.2 any condition causing malabsorption; or
  - 2.3 faltering growth in an infant/child; or
  - 2.4 increased nutritional requirements; or
  - 2.5 the child is being transitioned from TPN or tube feeding to oral feeding.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for

	Subsidy (Manufacturer's Price \$	Fully e) Subsidised Per ✓	Brand or Generic Manufacturer
continued applications meeting the following criteria:			
Both:			
<ol> <li>The treatment remains appropriate and the patient is ben</li> <li>General Practitioners must include the name of the dietitia practitioner and date contacted.</li> </ol>			egistered general
PAEDIATRIC ENTERAL FEED 1.5KCAL/ML – Special Authority Liquid			Hospital pharmacy [HP3] <b>Nutrini Energy RTH</b>
PAEDIATRIC ENTERAL FEED 1KCAL/ML – Special Authority s Liquid		500 ml OP 🛛 🖌 🖡	ospital pharmacy [HP3] Pediasure RTH Nutrini RTH
PAEDIATRIC ENTERAL FEED WITH FIBRE 1.5KCAL/ML - Sp pharmacy [HP3]	ecial Authority see S	SA1379 on the prev	vious page – Hospital
Liquid	7.14 5	500 ml OP 🖌 🖌	lutrini Energy Multi Fibre
PAEDIATRIC ORAL FEED 1.5KCAL/ML - Special Authority see	e SA1379 on the pre-	vious page – Hosp	ital pharmacy [HP3]
Liquid (strawberry)	1.90 2	200 ml OP 🖌 🖌 🖌	ortini
Liquid (vanilla)			ortini
	8.67 5	500 ml OP 🛛 🖌 🖌	Pediasure Plus
PAEDIATRIC ORAL FEED 1KCAL/ML - Special Authority see S	SA1379 on the previo	ous page – Hospita	al pharmacy [HP3]
Liquid (chocolate)			Pediasure
Liquid (strawberry)			Pediasure
Liquid (vanilla)			Pediasure
	1.66 2	250 ml OP 🖌 🖌 F	Pediasure
PAEDIATRIC ORAL FEED WITH FIBRE 1.5KCAL/ML - Special	I Authority see SA13	379 on the previous	s <mark>page</mark> – Hospital
pharmacy [HP3]			
Liquid (unflavoured)			Fortini Multi Fibre
Liquid (chocolate)			Fortini Multi Fibre Fortini Multi Fibre
Liquid (strawberry) Liquid (vanilla)			Fortini Multi Fibre
PEPTIDE-BASED ORAL FEED – Special Authority see SA1379 Powder			nacy [HP3] <b>Peptamen Junior</b>

### **Renal Products**

### ⇒SA1101 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has acute or chronic kidney disease.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

RENAL ORAL FEED 1.8 KCAL/ML	- Special Authority see SA1101 above - I	Hospital pharmacy	[HP3]
Liquid		220 ml OP	<ul> <li>Nepro HP (strawberry)</li> </ul>
			✓ Nepro HP (vanilla)

	Subsidy (Manufacturer's Price) \$	Sul Per	Fully bsidised	Brand or Generic Manufacturer
RENAL ORAL FEED 2 KCAL/ML - Special Authority see SA110				
Liquid, 200 ml bottle		4 OP	🗸 N	ovaSource Renal
Liquid (apricot) 125 ml		4 OP	🗸 R	enilon 7.5
Liquid (caramel) 125 ml	13.72	4 OP	✓ R	enilon 7.5

### **Specialised And Elemental Products**

#### ⇒SA1377 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 malabsorption; or
- 2 short bowel syndrome; or
- 3 enterocutaneous fistulas; or
- 4 eosinophilic oesophagitis; or
- 5 inflammatory bowel disease; or
- 6 patients with multiple food allergies requiring enteral feeding.

Notes: Each of these products is highly specialised and would be prescribed only by an expert for a specific disorder. The alternative is hospitalisation.

Elemental 028 Extra is more expensive than other products listed in this section and should only be used where the alternatives have been tried first and/or are unsuitable.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL SEMI-ELEMENTAL FEED 1.5KCAL/ML – Spe Liquid		e SA1377 abov 1,000 ml OP	
ORAL ELEMENTAL FEED 0.8KCAL/ML – Special Authority see Liquid (grapefruit), 250 ml carton Liquid (pineapple & orange), 250 ml carton Liquid (summer fruits), 250 ml carton	179.46 179.46	– Hospital phan 18 OP 18 OP 18 OP 18 OP	macy [HP3] ✓ Elemental 028 Extra ✓ Elemental 028 Extra ✓ Elemental 028 Extra
ORAL ELEMENTAL FEED 1KCAL/ML – Special Authority see S Powder (unflavoured)		Hospital pharma 80 g OP	acy [HP3] ✓ Vivonex TEN
SEMI-ELEMENTAL ENTERAL FEED 1KCAL/ML – Special Auth Liquid		7 above – Hosp 500 ml OP	ital pharmacy [HP3] Vitrison Advanced Peptisorb

### Paediatric Products For Children With Low Energy Requirements

#### ⇒SA1196 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

Subsidy	ç	Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	1	Manufacturer

- 1 Child aged one to eight years; and
- 2 The child has a low energy requirement but normal protein and micronutrient requirements.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Liquid	6.27	500 ml OP	🗸 Nutri	ni Low En
			Mu	lti Fibre

### **Standard Supplements**

#### ⇒SA1859 Special Authority for Subsidy

Initial application — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age; and
- 2 Any of the following:
  - 2.1 The patient has a condition causing malabsorption; or
  - 2.2 The patient has failure to thrive; or
  - 2.3 The patient has increased nutritional requirements; and
- 3 Nutrition goal has been set (eg reach a specific weight or BMI).

Renewal — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 A nutrition goal has been set (eg reach a specific weight or BMI).

Initial application — (Children - exclusive enteral nutrition for Crohn's disease) only from a gastroenterologist or dietitian on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 Dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

**Renewal** — (Children - exclusive enteral nutrition for Crohn's disease) from any relevant practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 General Practitioners and dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

Initial application — (Adults) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- Any of the following:
  - Patient is Malnourished
  - 1.1 Patient has a body mass index (BMI) of less than 18.5 kg/m²; or
  - 1.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
  - Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months; and
- 2 Any of the following:
  - Patient has not responded to first-line dietary measures over a 4 week period by:
  - 2.1 Increasing their food intake frequency (eg snacks between meals); or
  - 2.2 Using high-energy foods (e.g. milkshakes, full fat milk, butter, cream, cheese, sugar etc); or
  - 2.3 Using over the counter supplements (e.g. Complan); and
- 3 A nutrition goal has been set (e.g. to reach a specific weight or BMI).

**Renewal** — (Adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 A nutrition goal has been set (eg reach a specific weight or BMI); and
- 2 Any of the following:
  - Patient is Malnourished
  - 2.1 Patient has a body mass index (BMI) of less than 18.5 kg/m²; or
  - 2.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
  - 2.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months.

Initial application — (Short-term medical condition) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube or a nasogastric tube is to be inserted for feeding; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Is undergoing a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
  - 5.1 Pregnant; and
  - 5.2 Any of the following:
    - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or</p>
    - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
    - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

Renewal — (Short-term medical condition) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Has undergone a bone marrow transplant; or

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	✓	Manufacturer	

- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
  - 5.1 Pregnant; and
  - 5.2 Any of the following:
    - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or</p>
    - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
    - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

**Initial application** — (Long-term medical condition) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis; or
- 3 Liver disease; or
- 4 Chronic Renal failure; or
- 5 Inflammatory bowel disease; or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome; or
- 8 Bowel fistula; or
- 9 Severe chronic neurological conditions; or
- 10 Epidermolysis bullosa; or
- 11 AIDS (CD4 count < 200 cells/mm³); or
- 12 Chronic pancreatitis.

#### Renewal — (Chronic disease OR tube feeding for patients who have previously been funded under Special Authority

forms SA0702 or SA0583) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis; or
- 3 Liver disease; or
- 4 Chronic Renal failure: or
- 5 Inflammatory bowel disease; or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome; or
- 8 Bowel fistula; or
- 9 Severe chronic neurological conditions.

ENTERAL FEED 1.5KCAL/ML - Special Authority see SA1859 on page 282	- Hospital pharmac	y [HP3]
Liquid2.17	7 250 ml OP	<ul> <li>Ensure Plus HN</li> </ul>
8.68	3 1,000 ml OP	<ul> <li>Ensure Plus HN RTH</li> </ul>
9.00	)	<ul> <li>Nutrison Energy</li> </ul>

### SPECIAL FOODS

	Subsidy (Manufacturer's   \$		Illy Brand or ed Generic Manufacturer
ENTERAL FEED 1KCAL/ML - Special Authority see SA1859 on	page 282 Ho	cnital pharmaov [4	100
Liquid			Isosource Standard
	6.56	,	Osmolite RTH
	6.90		Nutrison RTH
ENTERAL FEED WITH FIBRE 0.83 KCAL/ML - Special Authorit	ty see <mark>SA1859</mark> o	on page 282 - Hos	pital pharmacy [HP3]
Liquid	9.05	1,000 ml OP	<ul> <li>Nutrison</li> </ul>
			800 Complete
			Multi Fibre
ENTERAL FEED WITH FIBRE 1 KCAL/ML - Special Authority s	00 CA1950 on r	Loopito	I phormooy [UD2]
Liquid		,	Jevity RTH
	7.21		Nutrison Multi Fibre
ENTERAL FEED WITH FIBRE 1.2KCAL/ML - Special Authority	see SA1859 on	page 282 - Hospit	al pharmacy [HP3]
Liquid	7.87	1,000 ml OP	Jevity Plus RTH
ENTERAL FEED WITH FIBRE 1.5KCAL/ML - Special Authority			•
Liquid	8.68		Jevity HiCal RTH
			<ul> <li>Nutrison Energy</li> </ul>
			Multi Fibre
ORAL FEED (POWDER) - Special Authority see SA1859 on page	ge 282 – Hospit	al pharmacy [HP3]	
Powder (chocolate).		840 g OP	Sustagen Hospital
		5	Formula
	26.00	850 g OP	Ensure
Powder (vanilla)			Sustagen Hospital
	14.00	040 y Oi	Formula Active
	00.00	050 - 00	
	26.00	850 g OP	Ensure
ORAL FEED 1.5KCAL/ML – Special Authority see SA1859 on pa	age 282 – Hospi	ital pharmacy [HP3	]
Additional subsidy by endorsement is available for patients b	eing bolus fed t	hrough a feeding tu	ibe, who have severe
epidermolysis bullosa, or as exclusive enteral nutrition in chil	dren under the a	age of 18 years for	the treatment of Crohn's
disease, or for patients with COPD and hypercapnia, defined	l as CO2 value e	exceeding 55mmH	a. The prescription must be
endorsed accordingly.		<b>J</b>	
Liquid (banana) – Higher subsidy of up to \$1.76 per 200 ml			
with Endorsement	0.72	200 ml OP	
		200 IIII OF	Ensure Plus
	(1.56)		
	(1.76)		Fortisip
Liquid (chocolate) – Higher subsidy of up to \$1.76 per 200 n			
with Endorsement		200 ml OP	
	(1.56)		Ensure Plus
	(1.76)		Fortisip
Liquid (fruit of the forest) – Higher subsidy of \$1.56 per 200	ml		
with Endorsement		200 ml OP	
	(1.56)		Ensure Plus
Liquid (strawberry) – Higher subsidy of \$1.76 per 200 ml wit	( )		
Endorsement	0.72	200 ml OP	
		200 ml OP	Eartiain
	(1.76)		Fortisip
Liquid (vanilla) – Higher subsidy of up to \$1.76 per 200 ml w			
Endorsement		237 ml OP	
	(1.65)		Ensure Plus
	0.72	200 ml OP	
	(1.56)		Ensure Plus
	(1.76)		Fortisip
	(		

	Subsidy (Manufacturer's F \$	Price) Su Per	Fully bsidised	Brand or Generic Manufacturer
ORAL FEED WITH FIBRE 1.5 KCAL/ML – Special Authority see Additional subsidy by endorsement is available for patients b epidermolysis bullosa. The prescription must be endorsed av Liquid (chocolate) – Higher subsidy of \$1.76 per 200 ml with	eing bolus fed th ccordingly.			
Endorsement		200 ml OP		ortisip Multi Fibre
Liquid (strawberry) – Higher subsidy of \$1.76 per 200 ml with Endorsement		200 ml OP		ortisip Multi Fibre
Liquid (vanilla) – Higher subsidy of \$1.76 per 200 ml with Endorsement	0.72 (1.76)	200 ml OP		ortisip Multi Fibre

# **High Calorie Products**

### ⇒SA1195 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria: All of the following:

- 1 Cystic fibrosis; and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements.

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Any of the following:
  - 1.1 any condition causing malabsorption; or
  - 1.2 faltering growth in an infant/child; or
  - 1.3 increased nutritional requirements; or
  - 1.4 fluid restricted; and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements or is fluid restricted.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer \$ ENTERAL FEED 2 KCAL/ML - Special Authority see SA1195 on the previous page - Hospital pharmacy [HP3] 500 ml OP Nutrison Concentrated 13.64 1.000 ml OP Ensure Two Cal HN RTH ORAL FEED 2 KCAL/ML - Special Authority see SA1195 on the previous page - Hospital pharmacy [HP3] Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, or who have severe epidermolysis bullosa. The prescription must be endorsed accordingly. Liquid (vanilla) - Higher subsidy of \$2.34 per 200 ml with 200 ml OP (2.34)Two Cal HN

SPECIAL FOODS

Healtheries Simple **Baking Mix** 

# Food Thickeners

#### SA1106 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 vear where the patient has motor neurone disease with swallowing disorder.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FOOD THICKENER	- Special Authority see SA1106 above - Hospital pharmad	y [HP3]	
Powder		300 g OP	<ul> <li>Nutilis</li> </ul>
	24.00	380 g OP	<ul> <li>Aptamil Feed</li> </ul>
		-	Thickener

### Gluten Free Foods

The funding of gluten free foods is no longer being actively managed by Pharmac from 1 April 2011. This means that we are no longer considering the listing of new products, or making subsidy, or other changes to the existing listings. As a result we anticipate that the range of funded items will reduce over time. Management of Coeliac disease with a gluten free diet is necessary for good outcomes. A range of gluten free options are available through retail outlets.

#### ■ SA1729 Special Authority for Subsidy

Initial application — (all patients) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Either:

- 1 Gluten enteropathy has been diagnosed by biopsy; or
- 2 Patient suffers from dermatitis herpetiformis.

Initial application — (paediatric patients diagnosed by ESPGHAN criteria) only from a paediatric gastroenterologist. Approvals valid without further renewal unless notified where the paediatric patient fulfils ESPGHAN criteria for biopsy free diagnosis of coeliac disease.

GLUTEN FREE BAKING MIX - Special Authority see SA1729 above - Hospital ph	narmacy [HP3]
Powder	1,000 g OP
(5.15)	•

	\$		<ul> <li>Manufacturer</li> </ul>
		Per	
GLUTEN FREE BREAD MIX - Special Authority see SA1729 or			armacy [HP3]
Powder		1,000 g OP	
	(7.32)		NZB Low Gluten
			Bread Mix
	3.51		
	(10.87)		Horleys Bread Mix
GLUTEN FREE FLOUR – Special Authority see SA1729 on the	previous page - H	-lospital pharma	icy [HP3]
Powder		2,000 g OP	
	(18.10)	· · ·	Horleys Flour
GLUTEN FREE PASTA - Special Authority see SA1729 on the	provious page - H	locnital nharma	
Buckwheat Spirals		250 g OP	cy [III 5]
	(3.11)	200 y 01	Orgran
Corn and Vegetable Shells	(- )	250 g OP	Orgian
Contraitu vegetable Shelis	(2.92)	250 y OF	Orgran
Corn and Vegetable Spirals	( )	250 g OP	Olgiali
Contrand vegetable Spirals		250 y OF	Orgran
Rice and Corn Lasagne Sheets	(2.92)	200 g OP	Olgiali
Rice and Com Lasayne Sheets		200 y OF	Oraron
Rice and Corn Macaroni	(3.82)	050 ~ OD	Orgran
		250 g OP	0
Dies and Care Denne	(2.92)	050 - 00	Orgran
Rice and Corn Penne		250 g OP	0
Disa and Maine Deste Cairola	(2.92)	050 - 00	Orgran
Rice and Maize Pasta Spirals		250 g OP	0
Disc and Millet Onivola	(2.92)	050 - 00	Orgran
Rice and Millet Spirals		250 g OP	0
D'ar and any analysis's college	(3.11)	075	Orgran
Rice and corn spaghetti noodles		375 g OP	0
	(2.92)		Orgran
Vegetable and Rice Spirals		250 g OP	0
the Para law work down when the	(2.92)	000 × 00	Orgran
Italian long style spaghetti		220 g OP	0
	(3.11)		Orgran
(Orgran Buckwheat Spirals to be delisted 1 July 2025) (Orgran Corn and Vegetable Shells to be delisted 1 July 2025)			

(Orgran Corn and Vegetable Shells to be delisted 1 July 2025) (Orgran Corn and Vegetable Shells to be delisted 1 July 2025) (Orgran Rice and Corn Lasagne Sheets to be delisted 1 July 2025) (Orgran Rice and Corn Macaroni to be delisted 1 July 2025) (Orgran Rice and Corn Penne to be delisted 1 July 2025) (Orgran Rice and Maize Pasta Spirals to be delisted 1 July 2025) (Orgran Rice and Millet Spirals to be delisted 1 July 2025) (Orgran Rice and Corn spaghetti noodles to be delisted 1 July 2025) (Orgran Vegetable and Rice Spirals to be delisted 1 July 2025) (Orgran Italian long style spaghetti to be delisted 1 July 2025)

### Foods And Supplements For Inherited Metabolic Disease

### ⇒SA2357 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid without further renewal unless notified where patient requires dietary management of inherited metabolic disorders.

## **Supplements For Homocystinuria**

AMINOACID FORMULA WITHOUT METHIONINE – Special Authority see SA2357 on the previous page – Hospital pharmacy [HP3]

Powder (neutral), 36 g sachets	750.30	30	HCU Anamix Junior
Powder, 12.5 g sachets		30	HCU Explore 5
Powder, 25 g sachets	1,048.95	30	<ul> <li>HCU Express 15</li> </ul>
Powder (neutral), can		500 g OP	<ul> <li>XMET Maxamum</li> </ul>
Powder (unflavoured), can		400 g OP	<ul> <li>HCU Anamix Infant</li> </ul>
Liquid (juicy berries), 125 ml bottle		30	HCU Lophlex LQ
Liquid (orange), 125 ml bottle		36	<ul> <li>HCU Anamix Junior</li> </ul>
			LQ

## Supplements For MSUD and short chain enoyl coA hydratase deficiency

AMINOACID FORMULA WITHOUT VALINE, LEUCINE AND ISOLEUCINE – Special Authority see SA2357 on the previous page – Hospital pharmacy [HP3]

Powder (neutral) 36 g sachets	750.00 30	) 🗸	MSUD Anamix Junior
Powder, 12.5 g sachets	349.65 30	) 🗸	MSUD Explore 5
Powder, 25 g sachets		) 🗸	MSUD Express 15
Powder (neutral), can		OP 🗸	MSUD Maxamum
Powder (orange), can		OP 🗸	MSUD Maxamum
Powder (unflavoured), can	260.00 400 g	OP 🗸	MSUD Anamix Infant
Liquid (orange) 125 ml bottles	941.40 36	6 🗸	MSUD Anamix Junior LQ
Liquid (juicy berries) 125 ml pouches	1,684.80 30	0 🗸	MSUD Lophlex LQ 20

	Subsidy		Fully	
	(Manufacturer's Pric \$	e) Per	Subsidised	Generic Manufacturer
Supplements For PKU				
MINOACID FORMULA WITHOUT PHENYLALANINE - Spe	cial Authority see SA	2357 on	page 288	- Hospital pharmacy [HP3
Tabs		75 OF	<ul> <li>✓</li> </ul>	Phlexy 10
Powder (Lemon), 34 g sachets		30		PKU Express 20
Powder (Neutral), 12.5 g sachets		30	✓	PKU Explore 5
Powder (Neutral), 34 g sachets		30	✓	PKU Express 20
Powder (Orange), 25 g sachets		30	✓	PKU Explore 10
Powder (Orange), 34 g sachets		30	✓	PKU Express 20
Powder (Raspberry), 25 g sachets		30	1	PKU Explore 10
Powder (Tropical), 34 g sachets		30	1	PKU Express 20
Powder (berry) 28 g sachets	936.00	30	~	PKU Lophlex Powder
Powder (chocolate) 36 g sachet		30	~	PKU Anamix Junior Chocolate
Powder (neutral) 28 g sachets	936.00	30	~	PKU Lophlex Powder
Powder (neutral) 36 g sachets	303.00	30	1	PKU Anamix Junior
Powder (orange) 28 g sachets		30		PKU Lophlex Powder
Powder (orange) 36 g sachet		30	1	PKU Anamix Junior Orange
Powder (unflavoured) 12.5 g sachets	234.00	30	1	PKU First Spoon
Powder (vanilla) 36 g sachet		30		PKU Anamix Junior Vanilla
Infant formula	174.72	400 g C	)P 🖌	PKU Anamix Infant
Powder (orange)		500 g C		XP Maxamum
Powder (unflavoured)		500 g C		XP Maxamum
Liquid (berry)		125 ml (		PKU Anamix Junior
Liquid (orange)		125 ml (	OP 🗸	PKU Anamix Junior
Liquid (forest berries), 250 ml carton	540.00	18 OF	) 🖌	Easiphen Liquid
Liquid (juicy tropical) 125 ml		30 OF		PKU Lophlex LQ 20
Oral semi-solid (berries) 109 g		36 OF		PKU Lophlex Sensation 20
Powder (neutral), 400 g can	715 16	4 OP	1	PKU Start
Liquid (juicy berries) 62.5 ml.		4 OF 60 OF		PKU Lophlex LQ 10
Liquid (juicy berries) 62.5 ml.		30 OF		PKU Lophlex LQ 20
Liquid (juicy bernes) 125 ml		30 OF		PKU Lophlex LQ 20
Liquiu (juicy orange) 125 mi		30 OF	•	PRO LOphiex LQ 20

## SPECIAL FOODS

	Subsidy		Fully	
	(Manufacturer's Price \$	) Per	Subsidised	
	,		-	
YCOMACROPEPTIDE AND AMINO ACID CONTAINS SC ge 288 – Hospital pharmacy [HP3]	OME PHENYLALANINE	– Spe	cial Autho	ority see SA2357 on
Powder (Banana) 35 g sachets	930.00	30	1	PKU
		00	•	sphere20 Banana
Powder (Berry), 20 g sachets	110.28	60	1	PKU Restore
		00	•	Powder
Powder (Chocolate) 32 g sachets	808 56	30	1	PKU Build
		00	•	20 Chocolate
Powder (Chocolate) 35 g sachets	930.00	30	1	PKU
		00	•	sphere20 Chocolate
				Sprici C20 Onocolate
Powder (Lemon) 35 g sachets		30	1	PKU
· · · ·				sphere20 Lemon
Powder (Lemonade) 33.4 g sachets		30	1	PKU GMPro Ultra
				Lemonade
Powder (Neutral), 15 g sachets		30	1	PKU Build 10
Powder (Orange), 20 g sachets		60	1	PKU Restore
				Powder
Powder (Raspberry Lemonade) 31 g sachets		30	1	PKU Build
				20 Raspberry
				Lemonade
Powder (Smooth) 31 g sachets		30	1	PKU Build
				20 Smooth
Powder (Vanilla) 33 g sachets		30	✓	PKU Build 20 Vanilla
Powder (neutral), 40 g sachets	673.92	30	1	Glytactin Bettermilk
Powder (unflavoured) 12.5 g sachets		30	~	PKU GMPro Mix-In
Powder (vanilla) 33.4 g sachets		30	✓	PKU GMPro Ultra
				Vanilla
Powder (Red Berry) 35 g sachets	930.00	30	✓	PKU sphere20 Red
				Berry
Powder (Vanilla) 35 g sachets	930.00	30	✓	PKU
				sphere20 Vanilla
Liquid (neutral), 250 ml carton		18		PKU GMPro LQ
Liquid (original), 250 ml carton		30 OF	· ·	PKU Glytactin RTD
				15
Liquid (Coffee Mocha), 250 ml carton		30 OF	· 🗸	PKU Glytactin RTD
				15 Lite
Liquid (chocolate), 250 ml carton		30 OF	> 🗸	PKU Glytactin RTD
				15
Liquid (vanilla), 250 ml carton		30 OF	> 🗸	PKU Glytactin RTD
				15 Lite

# Foods

LOW PROTEIN BAKING MIX - Special Authority see SA2357 on p	age 288 – Ho	ospital pharmac	y [HP3]
Powder	8.55	500 g OP	🖌 Loprofin Mix

	Suboidy		Fully Propd or
	Subsidy (Manufacturer's Pri		Fully Brand or dised Generic
	(Manulacialei 3 i ii	Per	<ul> <li>Manufacturer</li> </ul>
	000 Lloopitol m		1
LOW PROTEIN PASTA – Special Authority see SA2357 on page			
Animal shapes		500 g OP	✓ Loprofin
Lasagne		250 g OP	<ul> <li>Loprofin</li> </ul>
Low protein rice pasta		500 g OP	<ul> <li>Loprofin</li> </ul>
Macaroni	6.19	250 g OP	<ul> <li>Loprofin</li> </ul>
Penne		500 g OP	<ul> <li>Loprofin</li> </ul>
Spaghetti		500 g OP	<ul> <li>Loprofin</li> </ul>
Spirals	12.39	500 g OP	<ul> <li>Loprofin</li> </ul>
Supplements for Tyrosinaemia			
AMINOACID FORMULA WITHOUT PHENYLALANINE AND TYF	ROSINE – Specia	I Authority see	SA2357 on page 288 – Hospita
pharmacy [HP3]	040 GE	20	TVD Evalere E
Powder (Neutral), 12.5 g sachets		30	✓ TYR Explore 5
Powder (neutral) 36 g sachets		30	<ul> <li>TYR Anamix Junior</li> </ul>
Powder, can		400 g OP	<ul> <li>TYR Anamix Infant</li> </ul>
Liquid (juicy berries) 125 ml pouches	1,684.80	30	<ul> <li>TYR Lophlex LQ 20</li> </ul>
Liquid (orange) 125 ml bottle	941.40	36	<ul> <li>TYR Anamix Junior</li> </ul>
			LQ
GLYCOMACROPEPTIDE AND AMINO ACID CONTAINS SOME	TYROSINE AND	PHENYLALA	NINE – Special Authority see
SA2357 on page 288 – Hospital pharmacy [HP3]			
Powder (Red Berry), 35 g sachets	1,398.60	30	<ul> <li>TYR Sphere 20</li> </ul>
Powder (Vanilla), 35 g sachets	1,398.60	30	<ul> <li>TYR Sphere 20</li> </ul>
Supplements for Organic Acidaemias AMINOACID FORMULA WITHOUT ISOLEUCINE, METHIONINE	, THREONINE A	ND VALINE -	Special Authority see SA2357
on page 288 – Hospital pharmacy [HP3]			
Powder, can		400 g OP	<ul> <li>MMA/PA Anamix Infant</li> </ul>
AMINOACID FORMULA WITHOUT METHIONINE, THREONINE Hospital pharmacy [HP3]	AND VALINE -	Special Author	ity see SA2357 on page 288 -
Powder (neutral), 18 g sachets	750.30	30	<ul> <li>MMA/PA Anamix Junior</li> </ul>
Powder, 12.5 g sachets	349.65	30	MMA/PA Explore 5
Powder, 25 g sachets		30	✓ MMA/PA Express 15
Supplements for Glutaric Aciduria type 1			
			nital pharmany [UD2]
AMINOACID FORMULA WITHOUT LYSINE - Special Authority			
Powder (neutral), 18 g sachets		30	✓ GA1 Anamix Junior
Powder, 12.5 g sachets		30	✓ GA Explore 5
Powder, can		400 g OP	<ul> <li>GA1 Anamix Infant</li> </ul>
Supplements for Glycogen Storage Disease			
HIGH AMYLOPECTIN CORN-STARCH – Special Authority see		<mark>288</mark> – Hospital 30	pharmacy [HP3] ✓ Glycosade
Single dose amino acids			
ARGININE – Special Authority see SA2357 on page 288 – Hosp Powder, 4 g sachets		?3] 30	✓ Arginine2000

### SPECIAL FOODS

	Subsidy (Manufacturer's Price) \$		Fully Brand or ised Generic Manufacturer
CITRULLINE – Special Authority see SA2357 on page 288 – Hos Powder, 4 g sachets		3] 30	✓ Citrulline1000
ISOLEUCINE – Special Authority see SA2357 on page 288 – Hos Powder, 4 g sachets		'3] 30	✓ Isoleucine50
LEUCINE – Special Authority see SA2357 on page 288 – Hospita Powder, 4 g sachets		30	✓ Leucine100
PHENYLALANINE – Special Authority see SA2357 on page 288 · Powder, 4 g sachets		y [HP3] 30	<ul> <li>Phenylalanine50</li> </ul>
TYROSINE – Special Authority see SA2357 on page 288 – Hospi Powder, 4 g sachets		30	✓ Tyrosine1000
VALINE – Special Authority see SA2357 on page 288 – Hospital Powder, 4 g sachets		30	✓ Valine50
Other Fat Modified Products			
ELEMENTAL FEED WITH HIGH MEDIUM CHAIN TRIGLYCERID pharmacy [HP3]	·	ority see SA2	2357 on page 288 – Hospital
Powder (neutral), 100 g sachets	47.01	10	<ul> <li>Emsogen</li> </ul>
Carbohydrate and Fat with added vitamins and r	ninerals		
PROTEIN FREE SUPPLEMENT CONTAINING CARBOHYDRATI Authority see SA2357 on page 288 – Hospital pharmacy [HP3]	E, FAT WITH ADDE	D VITAMINS	SAND MINERALS - Special
Powder (neutral), can		00 g OP	<ul> <li>Energivit</li> </ul>
Essential Amino Acids			
ESSENTIAL AMINOACID FORMULA – Special Authority see SA Powder (neutral), can		- Hospital ph 00 g OP	armacy [HP3] <b>✓ Essential Amino</b> Acid Mix
Infant Formulae			

## For Williams Syndrome

### ➡SA1110 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is an infant suffering from Williams Syndrome and associated hypercalcaemia.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

LOW CALCIUM INFANT FORMULA	- Special Authority see SA1110 abov	e – Hospital	pharmacy [H	P3]

Powder	46.18	400 g OP	<ul> <li>Locasol</li> </ul>

	Subsidy (Manufacturer's Pric	e) Subs	Fully	Brand or Generic
	(Manulaciale) 31 no \$	Per	siuiseu ✓	Manufacturer
Gastrointestinal and Other Malabsorptive Probl	ems			
AMINO ACID FORMULA – Special Authority see SA2092 below Powder		cy [HP3] 400 g OP		Alfamino
Powder (unflavoured)	55.61	400 g OP	✓ N	Alfamino Junior Neocate Gold Neocate Junior Unflavoured
	65.72		✓ E	Neocate SYNEO Elecare Elecare LCP
Powder (vanilla)	55.61 65.72	400 g OP	-	Veocate Junior Vanilla Elecare

### ► SA2092 Special Authority for Subsidy

**Initial application** — (Infants under 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 History of anaphylaxis to cow's milk protein formula or dairy products; or
- 2 Eosinophilic oesophagitis; or
- 3 Ultra-short gut; or
- 4 Severe Immune deficiency; or
- 5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
- 6 Both:
  - 6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
  - 6.2 Either:
    - 6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or
    - 6.2.2 Patient has IgE mediated allergy.

Initial application — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

1 Either:

- 1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or
- 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
  - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products; or
  - 2.2 Eosinophilic oesophagitis; or
  - 2.3 Ultra-short gut; or
  - 2.4 Severe Immune deficiency; or
  - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
  - 2.6 Both:
    - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
    - 2.6.2 Either:
      - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

number; or

2.6.2.2 Patient has IgE mediated allergy.

Renewal — (Infants up to 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

1 Both:

- 1.1 Patient has IgE mediated allergy; and
- 1.2 All of the following:
  - 1.2.1 Patient remains allergic to cow's milk; and
  - 1.2.2 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy or extensively hydrolysed infant formula has been undertaken; and
  - 1.2.3 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
  - 1.2.4 Amino acid formula is required for a nutritional deficit; and
  - 1.2.5 It has been more than three months from the previous approval; or

2 Both:

- Patient has non IgE mediated severe gastrointestinal intolerance (including eosinophilic oesophagitis, ultra-short gut and severe immune deficiency); and
- 2.2 All of the following:
  - 2.2.1 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy, or extensively hydrolysed infant formula has been undertaken; and
  - 2.2.2 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
  - 2.2.3 Amino acid formula is required for a nutritional deficit; and
  - 2.2.4 It has been more than three months from the previous approval.

Renewal — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

1 Either:

- 1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or
- 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
  - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products; or
  - 2.2 Eosinophilic oesophagitis; or
  - 2.3 Ultra-short gut; or
  - 2.4 Severe Immune deficiency; or
  - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or

2.6 Both:

- 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
- 2.6.2 Either:
  - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or
  - 2.6.2.2 Patient has IgE mediated allergy.

Initial application — (for patients who have a current funding under Special Authority form SA1557) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 months for applications meeting the following criteria:

continued...

Subsidy (Manufacturer's Price)	s	Fully Subsidised	Brand or Generic
\$	Per	1	Manufacturer

continued...

All of the following:

- 1 Patient has a valid Special Authority approval for extensively hydrolysed formula (SA1557); and
- 2 Extensively hydrolysed formula (Aptamil Gold+ Pepti Junior, AllerPro SYNEO 1 and 2) is unable to be supplied at this time; and
- 3 The approval only applies to funded dispensings of Neocate Gold and Neocate Syneo.

Note: This criteria is short term funding to cover an out-of-stock situation on some extensively hydrolysed formula powder funded under Special Authority form SA1557. There is no renewal criteria under this restriction.

ENTERAL LIQUID PEPTIDE FORMULA	- Special Authority see SA1953 below	- Hospital phari	macy [HP3]
Liquid 1 kcal/ml		500 ml OP	<ul> <li>Nutrini Peptisorb</li> </ul>
Liquid 1.5 kcal/ml		500 ml OP	<ul> <li>Nutrini Peptisorb</li> </ul>
			Energy

### ⇒SA1953 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has impaired gastrointestinal function and either cannot tolerate polymeric feeds, or polymeric feeds are unsuitable; and
- 2 Any of the following:
  - 2.1 Severe malabsorption; or
  - 2.2 Short bowel syndrome; or
  - 2.3 Intractable diarrhoea; or
  - 2.4 Biliary atresia; or
  - 2.5 Cholestatic liver diseases causing malabsorption; or
  - 2.6 Cystic fibrosis; or
  - 2.7 Proven fat malabsorption; or
  - 2.8 Severe intestinal motility disorders causing significant malabsorption; or
  - 2.9 Intestinal failure; or
  - 2.10 Both:
    - 2.10.1 The patient is currently receiving funded amino acid formula; and
    - 2.10.2 The patient is to be trialled on, or transitioned to, an enteral liquid peptide formula; and
- 3 Either:
  - 3.1 A semi-elemental or partially hydrolysed powdered feed has been reasonably trialled and considered unsuitable; or
  - 3.2 For step down from intravenous nutrition.

Note: A reasonable trial is defined as a 2-4 week trial.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 An assessment as to whether the patient can be transitioned to a cows milk protein or soy infant formula or extensively hydrolysed formula has been undertaken; and
- 2 The outcome of the assessment is that the patient continues to require an enteral liquid peptide formula; and
- 3 General practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

EXTENSIVELY HYDROLYSED FORMULA	- Special Authority see SA1557 on t	he next page -	Hospital pharmacy [HP3]
Powder		450 g OP	<ul> <li>Pepti-Junior</li> </ul>
	36.20	900 g OP	Allerpro Syneo 1
		-	Allerpro Syneo 2

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsid	dised	Generic	
\$	Per	1	Manufacturer	

### ➡SA1557 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 Both:
  - 1.1 Cows milk formula is inappropriate due to severe intolerance or allergy to its protein content; and
  - 1.2 Either:
    - 1.2.1 Soy milk formula has been reasonably trialled without resolution of symptoms; or
    - 1.2.2 Soy milk formula is considered clinically inappropriate or contraindicated; or
- 2 Severe malabsorption; or
- 3 Short bowel syndrome; or
- 4 Intractable diarrhoea; or
- 5 Biliary atresia; or
- 6 Cholestatic liver diseases causing malsorption; or
- 7 Cystic fibrosis; or
- 8 Proven fat malabsorption; or
- 9 Severe intestinal motility disorders causing significant malabsorption; or
- 10 Intestinal failure; or
- 11 All of the following:
  - 11.1 For step down from Amino Acid Formula; and
  - 11.2 The infant is currently receiving funded amino acid formula; and
  - 11.3 The infant is to be trialled on, or transitioned to, an extensively hydrolysed formula; and
  - 11.4 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.
- Note: A reasonable trial is defined as a 2-4 week trial, or signs of an immediate IgE mediated allergic reaction.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 An assessment as to whether the infant can be transitioned to a cows milk protein or soy infant formula has been undertaken; and
- 2 The outcome of the assessment is that the infant continues to require an extensively hydrolysed infant formula; and
- 3 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

## Fluid Restricted

PAEDIATRIC ORAL/ENTERAL FEED 1 KCAL/ML	- Special Authority see SA1698	below -	- Hospital pharmacy [HP3]
Liquid			OP 🖌 Infatrini

#### ⇒SA1698 Special Authority for Subsidy

Initial application only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Patient is fluid restricted or volume intolerant and has been diagnosed with faltering growth; and
- 2 Patient is under the care of a paediatrician or dietitian who has recommended treatment with a high energy infant formula; and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

continued...

Subsidy (Manufacturer's Price)	Qui	Fully bsidised	Brand or Generic	
(Manulacturers Flice)	Per		Manufacturer	

continued...

Note: "Volume intolerant' patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

**Renewal** only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient continues to be fluid restricted or volume intolerant and has faltering growth; and
- 2 Patient is under the care of a hospital paediatrician or dietitian who has recommended treatment with a high energy infant formula; and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: 'Volume intolerant' patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

## **Ketogenic Diet**

#### ⇒SA1197 Special Authority for Subsidy

**Initial application** only from a metabolic physician or paediatric neurologist. Approvals valid for 3 months where the patient has intractable epilepsy, pyruvate dehydrogenase deficiency or glucose transported type-1 deficiency and other conditions requiring a ketogenic diet.

Renewal only from a metabolic physician or paediatric neurologist. Approvals valid for 2 years where the patient is on a ketogenic diet and the patient is benefiting from the diet.

HIGH FAT LOW CARBOHYDRATE FORMULA – Special Authority see SA119	7 above – Hospita	l pharmacy [HP3]
Powder (unflavoured)	300 g OP	<ul> <li>KetoCal 4:1</li> </ul>
	-	<ul> <li>Ketocal 3:1</li> </ul>
Powder (vanilla)	300 g OP	<ul> <li>KetoCal 4:1</li> </ul>

## SECTION I: NATIONAL IMMUNISATION SCHEDULE

	Subsidy (Manufacturer's Price) \$	Su Per	Fully ubsidised	Brand or Generic Manufacturer
Vaccinations				
<ul> <li>BACILLUS CALMETTE-GUERIN VACCINE – [Xpharm]</li> <li>For infants at increased risk of tuberculosis. Increased risk is</li> <li>1) living in a house or family with a person with current or p</li> <li>2) having one or more household members or carers who equal to 40 per 100,000 for 6 months or longer; or</li> <li>3) during their first 5 years will be living 3 months or longer</li> <li>Note a list of countries with high rates of TB are available at v www.bcgatlas.org/index.php.</li> <li>Inj Mycobacterium bovis BCG (Bacillus Calmette-Guerin),</li> </ul>	sast history of TB; or within the last 5 years in a country with a ra ww.health.govt.nz/tu	ate of T berculo	B > or equ osis (searc	ual to 40 per 100,000 th for downloads) or
Danish strain 1331, live attenuated, vial with diluent	0.00	10	✓ <u>B</u>	CG Vaccine AJV
COVID-19 VACCINE – [Xpharm]				
Inj 3 mcg bretovameran per 0.3 ml, 0.48 ml vial; infant vaccin yellow cap		10	<b>√</b> c	omirnaty Omicron
Up to three doses for previously unvaccinated children a	ed 6 months - 4 vea	rs at hio	ah risk of s	(JN.1) severe illness.
	, ,		5	
Inj 10 mcg bretovameran per 0.3 ml, 0.48 ml vial; paediatric vaccine, light blue cap	0.00	10	✓ C	omirnaty Omicron (JN.1)
Either:				
<ol> <li>One dose for previously unvaccinated children age</li> <li>Up to three doses for immunocompromised children</li> </ol>		l.		
Inj 30 mcg bretovameran per 0.3 ml, 0.48 ml vial; adult vaccir	ie,			
light grey cap	0.00	10	✓ C	omirnaty Omicron (JN.1)
Any of the following:				, ,
<ol> <li>One dose for previously unvaccinated people aged</li> <li>Up to three doses for immunocompromised people</li> <li>Up to two doses for previously unvaccinated people</li> <li>Up to four doses for people aged 16-29 at high risk</li> <li>One dose for previously unvaccinated people aged</li> </ol>	aged 12-15 years old 16-29 years old; or of severe illness; or	l; or		
6) One additional dose every 6 months for previously	vaccinated people ag	ed 30 y	ears and	over – additional dose is

6) One additional dose every 6 months for previously vaccinated people aged 30 years and over – additional dose is given at least 6 months after last dose.

Subsid	, . ,
(Manufacturer	r's Price) Subsidised Generic
\$	Per   Manufacturer

#### DIPHTHERIA, TETANUS AND PERTUSSIS VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- C)
- A) Funded for any of the following criteria:
  - 1) A single dose for pregnant women in the second or third trimester of each pregnancy; or
  - 2) A single dose for parents or primary caregivers of infants admitted to a Neonatal Intensive Care Unit or Specialist Care Baby Unit for more than 3 days, who had not been exposed to maternal vaccination at least 14 days prior to birth; or
  - A course of up to four doses is funded for children from age 7 up to the age of 18 years inclusive to complete full primary immunisation; or
  - 4) An additional four doses (as appropriate) are funded for (re-)immunisation for patients post haematopoietic stem cell transplantation or chemotherapy; pre or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or
  - 5) A single dose for vaccination of patients aged from 65 years old; or
  - 6) A single dose for vaccination of patients aged from 45 years old who have not had 4 previous tetanus doses; or
  - 7) For vaccination of previously unimmunised or partially immunised patients; or
  - 8) For revaccination following immunosuppression; or
  - 9) For boosting of patients with tetanus-prone wounds.

Notes: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.

- B) Contractors will be entitled to claim payment from the Funder for the supply of diphtheria, tetanus and pertussis vaccine to patients eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the diphtheria, tetanus and pertussis vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraphs 1 9 above.

Inj 2 IU diphtheria toxoid with 20 IU tetanus toxoid, 8 mcg pertussis toxoid, 8 mcg pertussis filamentous

	, .	31				
haemagglu	utinin and	2.5 mca	pertactin	in	0.5	ml prefilled

syringe	0.00	10	<ul> <li>Boostrix</li> </ul>

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🖌	Manufacturer

10

#### DIPHTHERIA, TETANUS, PERTUSSIS AND POLIO VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- c)
- A) Funded for any of the following:
  - 1) A single dose for children up to the age of 7 who have completed primary immunisation; or
  - A course of four vaccines is funded for catch up programmes for children (to the age of 10 years) to complete full primary immunisation; or
  - 3) An additional four doses (as appropriate) are funded for (re-)immunisation for people post HSCT, or chemotherapy; pre- or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or
  - 4) Five doses will be funded for children requiring solid organ transplantation.
- B) Contractors will be entitled to claim payment from the Funder for the supply of Diphtheria, tetanus, pertussis and polio vaccine to people eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Diphtheria, tetanus, pertussis and polio vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.

Inj 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg

pertussis toxoid, 25 mcg pertussis filamentous

haemagglutinin, 8 mcg pertactin and 80 D-antigen units

poliomyelitis virus in 0.5ml syringe ......0.00

DIPHTHERIA, TETANUS, PERTUSSIS, POLIO, HEPATITIS B AND HAEMOPHILUS INFLUENZAE TYPE B VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- C)
- A) Funded for children meeting any of the following criteria
  - 1) Up to four doses for children under the age of 10 years for primary immunisation; or
  - An additional four doses (as appropriate) for (re-)immunisation of children under the age of 18 years post haematopoietic stem cell transplantation; or
  - 3) An additional four doses (as appropriate) for (re-)immunisation of children under the age of 10 years who are post chemotherapy; pre or post splenectomy; undergoing renal dialysis and other severely immunosuppressive regimens; or
  - 4) Up to five doses for children under the age of 10 years receiving solid organ transplantation.
- B) Contractors will be entitled to claim payment from the Funder for the supply of Diphtheria, tetanus, pertussis, polio, hepatitis B and haemophilus influenzae type b vaccine to people eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Diphtheria, tetanus, pertussis, polio, hepatitis B and haemophilus influenzae type b vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Note: A course of up-to four vaccines is funded for catch up programmes for children (up to and under the age of 10 years) to complete full primary immunisation. Please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes.

Inj 30IU diphtheria with 40IU tetanus and 25mcg pertussis toxoids, 25mcg pertussis filamentous haemagglutinin, 8mcg pertactin, 80D-AgU polio virus, 10mcg hepatitis B antigen, 10mcg H. influenzae type b with tetanus toxoid 20-40mcg in 0.5ml syringe......0.00

10



✓ Infanrix IPV

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy	Fully	Brand or
	(Manufacturer's Price)	Subsidised	Generic
	\$	Per 🗸	Manufacturer
HAEMOPHILUS INFLUENZAE TYPE B VACCINE			
a) Only on a prescription			
b) No patient co-payment payable			
c)			
A) One dose for people meeting any of the following:			
<ol> <li>For primary vaccination in children; or</li> </ol>			
<ol><li>An additional dose (as appropriate) is funded in</li></ol>	for (re-)immunisation f	for people post h	aematopoietic stem cell
transplantation, or chemotherapy; functional a			
transplant, pre or post cochlear implants, rena			
<ol><li>For use in testing for primary immunodeficience</li></ol>	cy diseases, on the re-	commendation o	f an internal medicine
physician or paediatrician.			
<ul> <li>B) Contractors will be entitled to claim payment from the</li> </ul>			
vaccine to people eligible under the above criteria p			
for subsidised immunisation, and they may only do	so in respect of the Ha	aemophilus influe	enzae type b vaccine listed
in the Pharmaceutical Schedule.			
<ul> <li>Contractors may only claim for populations within the second secon</li></ul>		ered by their con	itract, which may be a
sub-set of the population described in paragraph A			
Inj 10 mcg vial with diluent syringe	0.00	· • •	Act-HIB
HEPATITIS A VACCINE – [Xpharm]			
Funded for patients meeting any of the following criteria:			
<ol> <li>Two vaccinations for use in transplant patients; or</li> </ol>			
<ol><li>Two vaccinations for use in children with chronic liver d</li></ol>			
<ol><li>One dose of vaccine for close contacts of known hepat</li></ol>	itis A cases.		
Inj 1440 ELISA units in 1 ml syringe	0.00	1	lavrix 1440
ing 1440 LEIGA units in 1 nii synnye	0.00	· • •	IAVIIA 1440

Inj 1440 ELISA units in 1 ml syringe	0.00	1	Havrix 1440
Inj 720 ELISA units in 0.5 ml syringe	0.00	1	<ul> <li>Havrix Junior</li> </ul>

		Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised			
HEPATITIS E	RECOMBINANT VACCINE – [Xpharm]						
	g per 0.5 ml prefilled syringe	0.00	1	1	Engerix-B		
Funded for patients meeting any of the following criteria:							
1) for household or sexual contacts of known acute hepatitis B patients or hepatitis B carriers; or							
	for children born to mothers who are hepatitis B su						
3)	for children up to and under the age of 18 years ind	clusive who are consid	dered	not to hav	ve achieved a positive		
,	serology and require additional vaccination or requ	ire a primary course c	of vac	cination; o	r		
4)	for HIV positive patients; or						
	for hepatitis C positive patients; or						
6)	for patients following non-consensual sexual interc	ourse; or					
7)	for patients prior to planned immunosuppression for	or greater than 28 days	s; or				
8)	8) for patients following immunosuppression; or						
,	for solid organ transplant patients; or						
	for post-haematopoietic stem cell transplant (HSC)	<ul><li>patients; or</li></ul>					
11)	following needle stick injury.						
Ini 20 ma	g per 1 ml prefilled syringe	0.00	1	1	Engerix-B		
	ded for patients meeting any of the following criteria:						
1)	for household or sexual contacts of known acute he	epatitis B patients or h	nepati	tis B carrie	ers: or		
	for children born to mothers who are hepatitis B su						
	for children up to and under the age of 18 years inc				e achieved a positive		
,	serology and require additional vaccination or requ	ire a primary course o	of vac	cination; o	r		
4)	for HIV positive patients; or						
5)	5) for hepatitis C positive patients; or						
6)	6) for patients following non-consensual sexual intercourse; or						
7)	<ol><li>for patients prior to planned immunosuppression for greater than 28 days; or</li></ol>						
8)	<ol><li>for patients following immunosuppression; or</li></ol>						
9)	for solid organ transplant patients; or						
	for post-haematopoietic stem cell transplant (HSC)	<ul><li>F) patients; or</li></ul>					
	following needle stick injury; or						
,	for dialysis patients; or						
13)	for liver or kidney transplant patients						

13) for liver or kidney transplant patients.

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✔	Brand or Generic Manufacturer
HUMAN PAPILLOMAVIRUS (6, 11, 16, 18, 31, 33, 45, 52 AND	58) VACCINE [HPV]		
a) Maximum of 1 inj per prescription			
b) Only on a prescription			
<ul> <li>c) No patient co-payment payable</li> <li>d)</li> </ul>			
<ul> <li>a) A) Any of the following:</li> <li>1) Maximum of two doses for children age</li> <li>2) Maximum of three doses for people me</li> <li>1) People aged 15 to 26 years inclu</li> </ul>	eting any of the followi		
<ol> <li>2) Either: People aged 9 to 26 years inclusi</li> <li>1) Confirmed HIV infection; or</li> <li>2) Received a transplant (inclu</li> </ol>	ding stem cell): or		
<ol><li>Maximum of four doses for people age</li></ol>			
<ul> <li>B) Contractors will be entitled to claim payment</li> </ul>	from the Funder for the	e supply of Huma	n papillomavirus vaccine

- B) Contractors will be entitled to claim payment from the Funder for the supply of Human papillomavirus vaccine to people eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Human papillomavirus vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraphs A above.

Inj 270 mcg in 0.5 ml syringe	0.00	10	<ul> <li>Gardasil 9</li> </ul>
-------------------------------	------	----	--------------------------------

	Subsidy (Manufacturer's Price) \$	S Per	Fully ubsidised	Brand or Generic Manufacturer
INFLUENZA VACCINE Inj 60 mcg in 0.5 ml syringe (quadrivalent vaccine)	120.00	10		nfluvac Tetra (2025 formulation)

	Subsidy (Manufacturer's Price) \$	Subs Per	Fully sidised	Brand or Generic Manufacturer
<ul> <li>a) Maximum of 1 inj per prescription</li> </ul>				
<ul> <li>b) Only on a prescription</li> </ul>				
<ul> <li>No patient co-payment payable</li> </ul>				
d)				
A) INFLUENZA VACCINE				
is available each year for patients who meet th	e following criteria, as	set by Ph	armac:	
a) all people 65 years of age and over; or				
b) people under 65 years of age who:				
<ul> <li>i) have any of the following cardiovase a) ischaemic heart disease, or</li> </ul>	cular diseases.			
b) congestive heart failure, or				
c) rheumatic heart disease, or				
d) congenital heart disease, or				
e) cerebo-vascular disease; or				
ii) have either of the following chronic	respiratory diseases:			
a) asthma, if on a regular preven	itative therapy, or			
<li>b) other chronic respiratory disea</li>	ase with impaired lung	function;	or	
iii) have diabetes; or				
iv) have chronic renal disease; or				
v) have any cancer, excluding basal a		ncers if no	t invasi	ve; or
vi) have any of the following other cond	ditions:			
<ul><li>a) autoimmune disease, or</li><li>b) immune suppression or immu</li></ul>	na dafiaianay, ar			
c) HIV, or	ne deliciency, or			
d) transplant recipients, or				
e) neuromuscular and CNS dise	ases/disorders or			
f) haemoglobinopathies, or				
g) are children on long term aspi	rin, or			
h) have a cochlear implant, or				
i) errors of metabolism at risk of	major metabolic deco	mpensatio	on, or	

- j) pre and post splenectomy, or
- k) Down syndrome, or
- vii) are pregnant; or
- children 4 years of age and under who have been hospitalised for respiratory illness or have a history of significant respiratory illness; or
- d) people under 65 years of age who:
  - i) have any of the following serious mental health conditions:
    - a) schizophrenia, or
    - b) major depressive disorder, or
    - c) bipolar disorder, or
    - d) schizoaffective disorder, or
- ii) are currently accessing secondary or tertiary mental health and addiction services; or

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy,
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Contractors will be entitled to claim payment for the supply of influenza vaccine to patients eligible under the above criteria pursuant to their contract with Health NZ for subsidised immunisation, and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

#### MEASLES, MUMPS AND RUBELLA VACCINE

- a) Only on a prescription
- b) No patient co-payment payable

C)

#### A) Measles, mumps and rubella vaccine

A maximum of two doses for any patient meeting the following criteria:

- 1) For primary vaccination in children; or
- 2) For revaccination following immunosuppression; or
- 3) For any individual susceptible to measles, mumps or rubella; or
- 4) A maximum of three doses for children who have had their first dose prior to 12 months.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes. Although a price is listed for the vaccine, doctors can still order measles mumps and rubella vaccine free of charge, as with other Schedule vaccines.

- B) Contractors will be entitled to claim payment for the supply of measles, mumps and rubella vaccine to patients eligible under the above criteria pursuant to their contract with Health NZ for subsidised immunisation, and they may only do so in respect of the measles, mumps and rubella vaccine listed in the Pharmaceutical Schedule.
- C) Contractors can only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Inj, measles virus 1,000 CCID50, mumps virus 5,012 CCID50, Rubella virus 1,000 CCID50; prefilled syringe/ampoule of

diluent 0.5 ml ......0.00 10

✓ Priorix

	Subsidy		Fully	Brand or
(	Manufacturer's Price) \$	Sub: Per	sidised	Generic Manufacturer
MENINGOCOCCAL (GROUPS A, C, Y AND W-135) CONJUGATE	•			Manufacturer
Inj 10 mcg of each meningococcal polysaccharide conjugated	WIGOINE			
to a total of approximately 55 mcg of tetanus toxoid carrier				
per 0.5 ml vial		1	🗸 M	enQuadfi
a) Only on a prescription				
b) No patient co-payment payable				
c)				
A) Any of the following:				
<ol> <li>Up to three doses and a booster every five with functional or anatomic asplenia, HIV, c solid organ transplant; or</li> <li>One dose for close contacts of meningococ</li> <li>One dose for person who has previously has</li> </ol>	omplement deficien cal cases of any gr d meningococcal d	ncy (acquin oup; or lisease of	red or inf	nerited), or pre or post
<ol><li>A maximum of two doses for bone marrow to</li></ol>				
<ol> <li>A maximum of two doses for person pre- ar</li> </ol>	nd post-immunosup	pression*;	or	
B) Both:	and a straight of the straight			
<ol> <li>Person is aged between 13 and 25 years, in</li> <li>Either:</li> </ol>	nciusive; and			
<ol> <li>One dose for individuals who are ente in boarding school hostels, tertiary ed residences, or prisons; or</li> <li>One dose for individuals who turn 13 y</li> <li>C) Contractors will be entitled to claim payment from</li> </ol>	ucation halls of resivery vears of age while lit	dence, mi iving in bo	litary bar arding s	racks, Youth Justice chool hostels.
W-135 vaccine to patients eligible under the above (Health NZ) for subsidised immunisation, and the W-135 vaccine listed in the Pharmaceutical Sche	y may only do so in			
<ul> <li>D) Contractors may only claim for patient population may be a sub-set of the population described in p Note: children under seven years of age require two d</li> </ul>	s within the criteria paragraphs A-B abo	ove.		
primary series and then five yearly. *Immunosuppression due to steroid or other immunosu				
28 days.				
Inj 5 mcg of each meningococcal polysaccharide conjugated to	)			
a total of approximately 44 mcg of tetanus toxoid carrier				
per 0.5 ml vial – [Xpharm]	0.00	1	🗸 Ni	imenrix
A) Both:				
<ol> <li>The child is under 12 months of age; and</li> <li>Any of the following:</li> </ol>				
<ol> <li>A maximum of three doses (dependant or</li> </ol>	a ago at first doso)	for nationt	e nro- ar	nd nost- splanactomy and
for patients with functional or anatomic as pre- or post- solid organ transplant; or				
<ol> <li>A maximum of three doses (dependant or of any group; or</li> </ol>	-			
<ol> <li>A maximum of three doses (dependant or meningococcal disease of any group; or</li> </ol>	о ,			
<ol> <li>A maximum of three doses (dependant or 5) A maximum of three doses (dependant or post-immunosuppression*.</li> </ol>	<b>o</b> ,			ansplant patients; or
Note: infants from 6 weeks to less than 6 months of age r	equire a 2+1 sched	lule. infant	s from 6	months to less than

Note: infants from 6 weeks to less than 6 months of age require a 2+1 schedule, infants from 6 months to less than 12 months of age require a 1+1 schedule. Refer to the Immunisation Handbook for recommended booster schedules with meningococcal ACWY vaccine.

*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	5	Subsidised	Generic	
\$	Per	1	Manufacturer	

### MENINGOCOCCAL B MULTICOMPONENT VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- c) Any of the following:
  - A) Three doses for children up to 12 months of age (inclusive) for primary immunisation; or
  - B) Up to three doses (dependent on age at first dose) for a catch-up programme for children from 13 months to
    - 59 months of age (inclusive) for primary immunisation, from 1 March 2023 to 31 August 2025; or
  - C) Both:
    - 1) Person is one year of age or over; and
    - 2) Any of the following:
      - i) up to two doses and a booster every five years for patients pre- and post-splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre- or post-solid organ transplant; or
      - ii) up to two doses for close contacts of meningococcal cases of any group; or
      - iii) up to two doses for person who has previously had meningococcal disease of any group; or
      - iv) up to two doses for bone marrow transplant patients; or
      - v) up to two doses for person pre- and post-immunosuppression*; or
  - D) Both:
    - 1) Person is aged between 13 and 25 years (inclusive); and
    - 2) Either:
      - Two doses for individuals who are entering within the next three months, or in their first year of living in boarding school hostels, tertiary education halls of residence, military barracks, Youth Justice residences or prisons; or
      - ii) Two doses for individuals who turn 13 years of age while living in boarding school hostels.
  - E) Contractors will be entitled to claim payment from the Funder for the supply of Meningococcal B multicomponent vaccine to patients eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Meningococcal B multicomponent vaccine listed in the Pharmaceutical Schedule.
  - F) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraphs A-D above.

*Immunosuppression due to corticosteroid or other immunosuppressive therapy must be for a period of greater than 28 days.

Inj 175 mcg per 0.5 ml prefilled syringe	0.00	1	<ul> <li>Bexsero</li> </ul>
		10	Bexsero

Subsidy		Fully	Brand or
(Manufacturer's Price)	S	ubsidised	Generic
\$	Per	1	Manufacturer

### PNEUMOCOCCAL (PCV13) CONJUGATE VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- C)

A) Any of the following:

- 1) A course of three doses for previously unvaccinated children up to the age of 59 months inclusive; or
- 2) Two doses are funded for high risk individuals (over the age of 12 months and under 18 years) who have previously received two doses of the primary course of PCV10; or
- 3) Up to an additional four doses (as appropriate) are funded for the (re)immunisation of high risk children aged under 5 years with any of the following:
  - a) on immunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient immune response; or
  - b) primary immune deficiencies; or
  - c) HIV infection; or
  - d) renal failure, or nephrotic syndrome; or
  - e) who are immune-suppressed following organ transplantation (including haematopoietic stem cell transplant); or
  - f) cochlear implants or intracranial shunts; or
  - g) cerebrospinal fluid leaks; or
  - receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage of 20 mg or greater; or
  - i) chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or
  - j) pre term infants, born before 28 weeks gestation; or
  - k) cardiac disease, with cyanosis or failure; or
  - I) diabetes; or
  - m) Down syndrome; or
  - n) who are pre-or post-splenectomy, or with functional asplenia; or
- 4) Up to an additional four doses (as appropriate) are funded for the (re-)immunisation of individuals 5 years and over with HIV, pre or post haematopoietic stem cell transplantation, or chemotherapy; pre- or post splenectomy; functional asplenia, pre- or post- solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, intracranial shunts, cerebrospinal fluid leaks or primary immunodeficiency; or
- 5) For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.
- B) Contractors will be entitled to claim payment from the Funder for the supply of Pneumococcal (PCV13) conjugate vaccine to people eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Pneumococcal (PCV13) conjugate vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Note: please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes Inj 30.8 mcg of pneumococcal polysaccharide serotypes 1, 3, 4,

5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and 23F in 0.5ml	
syringe0.00	

 10
 ✓
 Prevenar 13

 1
 ✓
 Prevenar 13

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$ Pe	er ✓	Manufacturer

#### PNEUMOCOCCAL (PPV23) POLYSACCHARIDE VACCINE – [Xpharm] Either:

 Up to three doses (as appropriate) for patients with HIV, for patients post haematopoietic stem cell transplant, or chemotherapy; pre- or post-splenectomy or with functional asplenia, pre- or post-solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, or primary immunodeficiency; or

#### 2) All of the following:

- a) Patient is a child under 18 years for (re-)immunisation; and
- b) Treatment is for a maximum of two doses; and
- c) Any of the following:
  - i) on immunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient immune response; or
  - ii) with primary immune deficiencies; or
  - iii) with HIV infection; or
  - iv) with renal failure, or nephrotic syndrome; or
  - v) who are immune-suppressed following organ transplantation (including haematopoietic stem cell transplant); or
  - vi) with cochlear implants or intracranial shunts; or
  - vii) with cerebrospinal fluid leaks; or
  - viii) receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage of 20 mg or greater; or
  - ix) with chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or
  - x) pre term infants, born before 28 weeks gestation; or
  - xi) with cardiac disease, with cyanosis or failure; or
  - xii) with diabetes; or
  - xiii) with Down syndrome; or
  - xiv) who are pre-or post-splenectomy, or with functional asplenia.

#### Inj 575 mcg in 0.5 ml prefilled syringe (25 mcg of each

23 pneumococcal serotype)0.00	1	Pneumovax 23
POLIOMYELITIS VACCINE – [Xpharm]		
Up to three doses for patients meeting either of the following:		
1) For partially vaccinated or previously unvaccinated individuals; or		
<ol><li>For revaccination following immunosuppression.</li></ol>		
Note: Please refer to the Immunisation Handbook for appropriate schedule for	or catch-up p	rogrammes.
Inj 80D antigen units in 0.5 ml syringe0.00	1	✓ IPOL

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	

### ROTAVIRUS ORAL VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- c)
- A) Maximum of two doses for people meeting the following:
  - 1) first dose to be administered in infants aged under 14 weeks of age; and
  - 2) no vaccination being administered to children aged 24 weeks or over.
- B) Contractors will be entitled to claim payment from the Funder for the supply of Rotavirus oral vaccine to people eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Rotavirus oral vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.
- Oral susp live attenuated human rotavirus

1,000,000 CCID50 per dose, squeezable tube0	.00	10	<ul> <li>Rotarix</li> </ul>
Oral susp live attenuated human rotavirus			
1,000,000 CCID50 per dose, prefilled oral applicator0	.00	10	✓ <u>Rotarix</u>

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	~	Manufacturer

#### VARICELLA VACCINE [CHICKENPOX VACCINE]

- a) Only on a prescription
- b) No patient co-payment payable

A) Either:

c)

- 1) Maximum of one dose for primary vaccination for either:
  - a) Any infant born on or after 1 April 2016; or
  - b) For previously unvaccinated children turning 11 years old on or after 1 July 2017, who have not previously had a varicella infection (chickenpox), or
- 2) Maximum of two doses for any of the following:
  - a) Any of the following for non-immune individuals:
    - i) with chronic liver disease who may in future be candidates for transplantation; or
    - ii) with deteriorating renal function before transplantation; or
    - iii) prior to solid organ transplant; or
    - iv) prior to any elective immunosuppression*; or
    - v) for post exposure prophylaxis who are immune competent inpatients; or
  - b) For individuals at least 2 years after bone marrow transplantation, on advice of their specialist; or
  - c) For individuals at least 6 months after completion of chemotherapy, on advice of their specialist; or
  - For HIV positive non immune to varicella with mild or moderate immunosuppression on advice of HIV specialist; or
  - For individuals with inborn errors of metabolism at risk of major metabolic decompensation, with no clinical history of varicella; or
  - f) For household contacts of paediatric patients who are immunocompromised, or undergoing a procedure leading to immune compromise where the household contact has no clinical history of varicella; or
  - g) For household contacts of adult patients who have no clinical history of varicella and who are severely immunocompromised, or undergoing a procedure leading to immune compromise where the household contact has no clinical history of varicella.
- B) Contractors will be entitled to claim payment from the Funder for the supply of Varicella vaccine [Chickenpox vaccine] vaccine to people eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Varicella vaccine [Chickenpox vaccine] listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraphs A above.

* immunosuppression due to steroid or other immunosuppressive therapy must be for a treatment period of greater than 28 days

Inj 2000 PFU prefilled syringe plus vial ......0.00 10 🗸 Varilrix

			Subsidy (Manufacturer's Price) \$	S Per	Fully ubsidised	Brand or Generic Manufacturer
ARICELLA ZO	STER VACC	INE [SHINGLES VACCIN	E]			
a) Only on a	a prescriptior	1				
b) No patier	nt co-paymer	nt payable				
c)						
,		ents meeting the following	criteria:			
1	<ol> <li>Either:</li> </ol>					
		doses for all people aged				
	,		s of age or older with any of the		0	
	,		pietic stem cell transplant or cel	lular the	erapy; or	
	,	pre- or post-solid organ				
		haematological malignar	controlled HIV infection; or			
			ease modifying anti-rheumatic d	lruae (F	MARDe -	- targeted synthetic
	0)	· ·	synthetic) for polymyalgia rheu	• •		• •
		rheumatoid arthritis; or	oynaload) for polynlyaigia mea	matioa,	oyotonno	lapus crythematosus of
	f)	end stage kidney diseas	e (CKD 4 or 5): or			
		primary immunodeficien				
B) Co			ent from the Funder for the sup	ply of V	aricella zo	oster vaccine (Shingles
vac	ccine) to patie	ents eligible under the abo	ve criteria pursuant to their con	tract w	ith Health	New Zealand (Health NZ)
for	subsidised in	nmunisation, and they ma	y only do so in respect of the V	aricella	zoster va	ccine [Shingles vaccine]
liste	ed in the Pha	armaceutical Schedule.				
,			pulations within the criteria that	are cov	/ered by t	heir contract, which may b
2.0	ub-cot of the	population described in p	araaranh A ahaya			

Inj 50 mcg per 0.5 ml vial plus vial0.00	1	<ul> <li>Shingrix</li> </ul>
	10	<ul> <li>Shingrix</li> </ul>

# Diagnostic Agents

TUBERCULIN PPD [MANTOUX] TEST – [Xpharm]			
Inj 5 TU per 0.1 ml, 1 ml vial	0.00	1	✓ <u>Tubersol</u>

#### - Symbols -

3TC111
- A -
A-Scabies
Abacavir sulphate 111
Abacavir sulphate with
lamivudine 111
Abacavir/Lamivudine Viatris111
Abilify Maintena
Abilify Maintena S29136
Abiraterone acetate
Acarbose11
Accarb11
Acetazolamide268
Acetec
Acetic acid with hydroxyquinoline and
ricinoleic acid
Acetylcysteine271
Aci-Jel
Aciclovir
Infection106
Sensory266
Acidex
Acipimox52
Acitretin72
Act-HIB
Actemra233
Actinomycin D157
Actrapid10
Actrapid Penfill10
Acupan
Adalimumab (Amgevita) 183
Adalimumab (Humira - Alternative
brand) 192
Adapalene66
Adcetris201
ADR Cartridge 1.821
Adrenaline
Cardiovascular 54
Respiratory255
Advantan 69
Advate
Adynovate38
Afinitor
Aflibercept198
AFT-Pyrazinamide106
Agents Affecting the
Renin-Angiotensin System 44
Agents for Parkinsonism and Related
Disorders 121
Agents Used in the Treatment of
Poisonings 271
Agrylin
Albalon

Albendazole9	
Albey250	
Albustix	
Alchemy Oxaliplatin 15	3
Alchemy Oxybutynin8	
Aldurazyme2	
Alecensa164	
Alectinib164	
Alendronate sodium110	6
Alendronate sodium with colecalciferol	_
Alfacalcidol	
Alfamino	
Alfamino Junior	
Alginic acid	6
Alglucosidase alfa24	
Alkeran	
Allerfix	
Allerpro Syneo 1	
Allerpro Syneo 2	6
Allersoothe25	
Allmercap15	
Allopurinol118	
Almarytm4	7
Alpha-Adrenoceptor Blockers44	4
Alpha-Keri Lotion	0
Alphamox 1259	6
Alphamox 2509	6
Alprolix	
Alu-Tab	6
Aluminium hydroxide	
Alyacen	B
Amantadine hydrochloride12	1
Ambrisentan	5
Ambrisentan Viatris5	5
Amgevita183	3
Amiloride hydrochloride	0
Amiloride hydrochloride with	
furosemide50	0
Amiloride hydrochloride with	
hydrochlorothiazide 50	0
Aminophylline	2
Amiodarone hydrochloride40	6
Amisulpride134	4
Amitriptyline 12	
Amlodipine4	
Amorolfine6	
Amoxicillin	6
Amoxicillin with clavulanic acid9	6
Amoxiclav Devatis Forte9	6
Amphotericin B 30	
Amsacrine150	6
AmsaLyo150	
Amsidine150	6

Amzoate	28
Anaesthetics	.122
Anafranil	.127
Anagrelide hydrochloride	.156
Analgesics	.123
Anastrozole	.175
Anatrole	.175
Anoro Ellipta	
Antabuse	
Antacids and Antiflatulents	6
Anthelmintics	
Antiacne Preparations	66
Antiallergy Preparations	255
Antianaemics	
Antiandrogen Oral	
Contraceptives	79
Antiarrhythmics	46
Antibacterials	07
Antibacterials Topical	66
Anticholinergic Agents	250
Anticholinesterases	115
Antidepressants	127
Antidiarrhoeals	. 127
Antiepilepsy Drugs	120
Antifibrinolytics, Haemostatics and	. 129
Local Sclerosants	26
Antifibrotics	
Antifungals	
Antifungals Topical	
Antihistamines Antihypotensives	.250
Antimalarials	. 103
Antimigraine Preparations	.132
Antinausea and Vertigo Agents	.132
Antipruritic Preparations	
Antipsychotics	.134
Antiretrovirals	.109
Antirheumatoid Agents	.116
Antispasmodics and Other Agents	_
Altering Gut Motility	8
Antithrombotic Agents	39
Antithymocyte globulin	
(equine)	. 183
Antitrichomonal Agents	. 104
Antituberculotics and	
Antileprotics	
Antiulcerants	
Antivirals	. 106
Anxiolytics	. 138
Anzatax	. 161
Apidra	11
Apidra SoloStar	11
APO Clomipramine	
APO-Atomoxetine	.143

_

Apo-Azithromycin94
APO-Candesartan HCTZ
16/12.5
APO-Candesartan HCTZ
32/12.5
Apo-Temozolomide 162
Apomorphine hydrochloride121
Aprepitant 132
Apresoline54
Aptamil Feed Thickener
Aqueous cream70
Aratac
Arava
Arginine25
Arginine2000
Aripiprazole
Aripiprazole Sandoz 134
Aristocort69
Arrotex-Prazosin S2944
Arrow - Clopid
Arrow - Lattim
Arrow-Amitriptyline127
Arrow-Bendrofluazide
Arrow-Brimonidine
Arrow-Diazepam
Arrow-Doxorubicin
Arrow-Fluoxetine
Arrow-Losartan &
Hydrochlorothiazide
Arrow-Norfloxacin
Arrow-Ornidazole104
Arrow-Quinapril 1045
Arrow-Quinapril 20
Arrow-Quinapril 5
Arrow-Roxithromycin
Arrow-Timolol
Arrow-Topiramate
Arrow-Tramadol
Arsenic trioxide
Asacol
Asacol S29
Ascend
Ascend Aripiprazole
Ascend-Cefuroxime
Ascorbic acid
Aspen Adrenaline
Aspirin
Blood
Nervous
Asthalin
Atazanavir Mylan111
Atazanavir sulphate
Atazanavir Viatris
Atenolol
Atenolol AFT47
Atenolol Viatris

Atezolizumab243
ATGAM183
Ativan138
Atnahs Olsalazine8
Atomoxetine143
Atorvastatin52
Atropine sulphate
Cardiovascular
Sensory269
Atropt
Atrovent
Aubagio140
Augmentin96
Aurorix
AutoSoft 3020
AutoSoft 9020
Avelox
Avonex139
Avonex Pen139
Azacitidine 153
Azacitidine Dr Reddy's 153
Azamun 175
Azathioprine175
Azilect
Azithromycin94
Azopt
AZT111
- B -
B-D Micro-Fine16
B-D Ultra Fine16
B-D Ultra Fine II16
Bacillus Calmette-Guerin (BCG)
vaccine
Bacillus Calmette-Guerin
vaccine
Baclofen119
Bactroban
Balance
Barrier Creams and Emollients70
BCG Vaccine AJV
Beclazone 100
Beclazone 250
Beclazone 50257
Beclomethasone dipropionate
Bedaquiline104
Bee venom allergy treatment
Bendamustine hydrochloride
Bendamustine Sandoz
Bendrofluazide
Bendroflumethiazide
[Bendrofluazide]
Benralizumab
Benzathine benzylpenicillin
Benzatropine mesylate
Benzbromarone
Benztrop
201121 0P

Benzydamine hydrochloride	30
Benzylpenicillin sodium [Penicillin	
G]	96
Beta Cream	68
Beta Ointment	
Beta Scalp	
Beta-Adrenoceptor Agonists2	
Beta-Adrenoceptor Blockers	47
Beta-hCG low sensitivity urine test	
kit	៱
Betadine	
Betadine Skin Prep	71
Betaferon	30
Betahistine dihydrochloride1	
Betaine	25
Betamethasone dipropionate	62
Betamethasone dipropionate with	00
calcipotriol	70
Betamethasone sodium phosphate	12
with betamethasone acetate	ەم
Betamethasone valerate	
Betamethasone valerate with sodium	/4
fusidate [fusidic acid]	<u></u>
Betaxolol	
Betnovate	
Betoptic	80
Betoptic S2	
Bevacizumab2	
Bexsero	
Bezafibrate	
Bezalip Bezalip Retard	52 50
Bicalutamide	
Bicillin LA	
BiCNU1	
BiCNU S291 Bile and Liver Therapy	52 4 0
Dile and Liver Therapy	10
Biltricide	93 60
Binarex1 Binocrit	/ 3 05
Biodone	
Biodone Extra Forte	
Bisacodyl	
Bisacodyl Viatris	
Bisoprolol fumarate	
BK Lotion	
Bleomycin sulphate	90
Blood Colony-stimulating	
Factors	42
Blood glucose diagnostic test	4.5
meter	15
Blood glucose diagnostic test	4.5
strip	15
Blood glucose test strips (visually	

impaired) 15
Blood Ketone Diagnostic Test
Strip 14
Boostrix
Bortezomib
Bosentan
Bosentan Dr Reddy's
Bplex
Brentuximab Vedotin
Breo Ellipta
Brevinor 1/28
Breztri Aerosphere
Driegrad Turbubeler
Bricanyl Turbuhaler
Brimonidine tartrate
Brimonidine tartrate with timolol
maleate
Brinzolamide
Brufen SR115
BSF Dasatinib-Teva271
Buccastem133
Budesonide
Alimentary6
Respiratory257, 264
Budesonide Te Arai6
Budesonide with eformoterol 258
Budesonide with glycopyrronium and
eformoterol 260
Bumetanide50
Buprenorphine Naloxone BNM 147
Buprenorphine with naloxone
Bupropion hydrochloride148
Burel 101
Burinex
Buscopan8
Buspirone hydrochloride138
Buspirone Viatris
Busulfan
- C -
Cabergoline
Caffeine citrate
Calamine
Calci-Tab 500
Calcipotriol73
Calcitonin
Calcitriol
Calcitriol-AFT
Calcitriol-AFT S29
Calcium 500 mg Hexal
Calcium carbonate
Calcium carbonate PAI
Calcium Channel Blockers
Calcium Disodium Versenate
Calcium folinate
Calcium Folinate Ebewe
Calcium Folinate Sandoz
Calcium Folinate Sandoz S29 154

Calcium gluconate	33
Calcium Homeostasis	00
Calcium polystyrene sulphonate	
Calcium polystyrene sulphonate	40
Calcium Resonium	
Calogen2	
Camber	54
Candesartan cilexetil	45
Candesartan cilexetil with	
hydrochlorothiazide	45
Candestar	45
Canesten	67
Capecitabine1	54
Capecitabine Viatris1	54
Capsaicin	
Musculoskeletal1	16
Nervous1	
Captopril	
Carafate	
Carbaccord1	
Carbamazepine	
Carbimazole	
Carbomer	
Carboplatin 1	
Carboplatin Accord	52
Carbosorb-X	
Cardinol LA	
Cardizem CD	
CareSens Dual	14
CareSens N	15
CareSens N POP	15
CareSens N Premier	15
CareSens PRO	15
Carmellose sodium with gelatin and	
pectin	30
Carmustine 1	52
Carnitor	27
Carvedilol	
Carvedilol Sandoz	47
Casirivimab and imdevimab2	
Catapres	
Cefaclor monohydrate	93
Cefalexin	
Cefalexin Sandoz	93
Cefazolin	
Cefazolin-AFT	
Ceftriaxone	00
Ceftriaxone-AFT	02
Cefuroxime axetil	00
Celapram1	
Celebrex 1	
Celecoxib1	
Celecoxib Pfizer1	
Celestone Chronodose	
Cellcept1	
Centrally-Acting Agents	
Cephalexin ABM	93

Cetirizine hydrochloride	256
Cetomacrogol	
Cetomacrogol with glycerol	70
Cetomacrogol-AFT	70
Cetuximab	202
Charcoal	
CheckTop	
Chemotherapeutic Agents	151
Chickenpox vaccine	313
Chlorambucil	152
Chloramphenicol	266
Chlorothiazide	
Chlorpromazine hydrochloride	134
Chlorsig	
Chlortalidone [Chlorthalidone]	200
Chlorthalidone	
Chlorvescent	43
Choice 380 7med Nsha Silver/copp	10
Short	
Ciclosporin	251
Cidomycin P/Free	08 08
Cilazapril	
Cilicaine VK	44
Cinacalcet	
Cinacalet Devatis	20
Ciprofloxacin	02
Infection	08
Sensory	
Ciprofloxacin Teva	266
Cisplatin	
Cisplatin Accord	152
Cisplatin Ebewe	152
Citalopram hydrobromide	100
Citrulline1000	202
Cladribine	290
Clarithromycin	104
Alimentary	0
Infection	
Clexane	
Clexane Forte	
Clindamycin	
Clinicians	
Clinicians Renal Vit	20, 02 00
Clobazam	
Clobetasol propionate	129
Clobetasone butyrate	0, 74 60
Clofazimine	104
Clomazol	104
Dermatological	67
Genito-Urinary	
Clomifene citrate	
Clomipramine hydrochloride	
Clonipramine Teva	
Clonazepam 129	
Clonidine Clonidine hydrochloride	
	49

Clonidine Teva49
Clopidogrel
Clopine
Clopixol
Clotrimazole
Dermatological67
Genito-Urinary79
Clozapine134
Clozaril
Clustran
Co-trimoxazole 100
Coal tar
Coal tar with allantoin, menthol,
phenol and sulphur 73
Coal tar with salicylic acid and
sulphur
Cobal-B12
Cobalin-H
Coco-Scalp
Codeine phosphate
Extemporaneous
Nervous
Coenzyme Q10
Colchicine
Colecalciferol
Colestyramine
Colestyramine - Mylan
Colgout
Colifoam
Colistin sulphomethate
Colistin-Link
Collodion flexible
Colloidal bismuth subcitrate10
Colofac
Colomycin
Coloxyl
Combigan
Comirnaty Omicron (JN.1)
Compound electrolytes
Compound electrolytes with glucose
[Dextrose]
Compound hydroxybenzoate
Comtan
Concerta
Condoms
Condyline
Continuous glucose monitor
(interoperable) 21
Continuous glucose monitor
(standalone)
Contraceptives - Hormonal
Contraceptives - Non-hormonal
Copaxone
Cordarone-X 46
Cordarone-X

Corticosteroids Topical	68
Cosentyx	.230
Cosmegen	. 157
Coumadin	41
Country Life	27
Coversyl	45
COVID-19 vaccine	.299
Creon 10000	
Creon 25000	
Creon Micro	
Crotamiton	
Crystaderm	
Cu 375 Standard	
Curam	
Curam Duo 500/125	90
Cvite	
Cyclizine hydrochloride	
Cyclizine lactate	100
Cyclogyl	. 100
Cyclonex	. 153
Cyclopentolate hydrochloride	
Cyclophosphamide	. 153
Cyclorin	. 104
Cycloserine	
Cyklokapron	
Cyproterone acetate	84
Cyproterone acetate with	
ethinyloestradiol	79
Cystadane	25
Cytarabine	
Cytotec	
Cytoxan	. 153
- D -	
D-Penamine	
Dabigatran	41
Dacarbazine	. 157
Dactinomycin [Actinomycin D]	
Daivobet	72
Daivonex	
Daktarin	
Dalacin C	98
Dantrium	. 120
Dantrium S29	. 120
Dantrolene	. 120
Daonil	11
Dapa-Tabs	
Dapsone	. 105
Daraprim	99
Darunavir	
Darunavir Viatris	
Dasatinib	
Dasatinib-Teva	
Daunorubicin	
David One Step Cassette Pregnanc	
Test	
DBL Acetylcysteine	271

DBL Adrenaline54
DBL Aminophylline
DBL Bleomycin Sulfate156
DBL Bortezomib157
DBL Carboplatin152
DBL Carboplatin S29152
DBL Cisplatin
DBL Dacarbazine
DBL Desferrioxamine Mesylate for Inj
BP
DBL Docetaxel
DBL Ergometrine
DBL Gemcitabine
DBL Gentamicin
DBL Leucovorin Calcium
DBL Methotrexate Onco-Vial
DBL Naloxone Hydrochloride
DBL Pethidine Hydrochloride
DBL Vincristine Sulfate
Decozol
Deferasirox
Deferiprone
Deferoxamine Pfizer S29272
Denosumab
Deolate
Deoxycoformycin
Depo-Medrol
Depo-Provera
Depo-Testosterone
Deprim
Dermol
Desferrioxamine mesilate
Desmopressin
Desmopressin acetate
Desmopressin-PH&T
Detection of Substances in
Urine
Dexamethasone
Hormone
Sensory
Dexamethasone phosphate
Dexamethasone with framycetin and
gramicidin
Dexamethasone with neomycin
sulphate and polymyxin B
sulphate
Dexamfetamine sulfate
Dexcom G6
Dexcom G7
Dexcom ONE+
Dexmethsone
Dextrochlorpheniramine
maleate
Dextrose
DHC Continus
Diabetes 10

Diabetes Management	14
Diacomit	. 131
Diagnostic Agents	.314
Diamide Relief	6
Diamox	.268
Diasip	.278
Diazepam129,	
Diazoxide	
Dibenzyline	44
Diclofenac Devatis	
Diclofenac Sandoz	
Diclofenac sodium	
Musculoskeletal	.115
Sensory	
Differin	
Difflam	
Diflucan	
Digestives Including Enzymes	
Digoxin	
Dihydrocodeine tartrate	125
Dilantin	
Dilantin Infatab	130
Dilantin Paediatric	130
Diltiazem CD Clinect	
Diltiazem hydrochloride	49
Dimethicone	
Dimethyl fumarate	139
Dipentum	
Dipolitani	
Diphtheria tetanus and pertussis	•••••
Diphtheria, tetanus and pertussis	
Diphtheria, tetanus and pertussis vaccine	
Diphtheria, tetanus and pertussis vaccine Diphtheria, tetanus, pertussis and	300
Diphtheria, tetanus and pertussis vaccine Diphtheria, tetanus, pertussis and polio vaccine	300 301
Diphtheria, tetanus and pertussis vaccine Diphtheria, tetanus, pertussis and polio vaccine Diphtheria, tetanus, pertussis, polio,	300 301
Diphtheria, tetanus and pertussis vaccine Diphtheria, tetanus, pertussis and polio vaccine Diphtheria, tetanus, pertussis, polio, hepatitis B and haemophilus	. 300 . 301
Diphtheria, tetanus and pertussis vaccine Diphtheria, tetanus, pertussis and polio vaccine Diphtheria, tetanus, pertussis, polio, hepatitis B and haemophilus influenzae type B vaccine	. 300 . 301 . 301
Diphtheria, tetanus and pertussis vaccine Diphtheria, tetanus, pertussis and polio vaccine Diphtheria, tetanus, pertussis, polio, hepatitis B and haemophilus influenzae type B vaccine Diprosone	. 300 . 301 
Diphtheria, tetanus and pertussis vaccine Diphtheria, tetanus, pertussis and polio vaccine Diphtheria, tetanus, pertussis, polio, hepatitis B and haemophilus influenzae type B vaccine Diprosone Diprosone OV	. 300 . 301 68 68
Diphtheria, tetanus and pertussis vaccine Diphtheria, tetanus, pertussis and polio vaccine Diphtheria, tetanus, pertussis, polio, hepatitis B and haemophilus influenzae type B vaccine Diprosone Diprosone OV Dipyridamole	. 300 . 301 68 68 39
Diphtheria, tetanus and pertussis vaccine Diphtheria, tetanus, pertussis and polio vaccine Diphtheria, tetanus, pertussis, polio, hepatitis B and haemophilus influenzae type B vaccine Diprosone Diprosone OV Dipyridamole Disopyramide phosphate	. 300 . 301 68 68 39 46
Diphtheria, tetanus and pertussis vaccine Diphtheria, tetanus, pertussis and polio vaccine Diphtheria, tetanus, pertussis, polio, hepatitis B and haemophilus influenzae type B vaccine Diprosone Diprosone OV Dipyridamole Disopyramide phosphate Disulfiram	. 300 . 301 68 68 39 46 148
Diphtheria, tetanus and pertussis vaccine Diphtheria, tetanus, pertussis and polio vaccine Diphtheria, tetanus, pertussis, polio, hepatitis B and haemophilus influenzae type B vaccine Diprosone Diprosone OV Dipyridamole Disopyramide phosphate Disulfiram	. 300 . 301 68 68 39 46 . 148 50
Diphtheria, tetanus and pertussis vaccine Diphtheria, tetanus, pertussis and polio vaccine Diphtheria, tetanus, pertussis, polio, hepatitis B and haemophilus influenzae type B vaccine Diprosone Diprosone OV Dipyridamole Disopyramide phosphate Disulfiram Diuretics Docetaxel	. 300 . 301 68 68 39 46 . 148 50 . 157
Diphtheria, tetanus and pertussis vaccine Diphtheria, tetanus, pertussis and polio vaccine Diphtheria, tetanus, pertussis, polio, hepatitis B and haemophilus influenzae type B vaccine Diprosone OV Diprosone OV Disopyramide phosphate Disulfiram Diuretics Docetaxel Docetaxel Accord	. 300 . 301 68 68 39 46 . 148 50 . 157 . 157
Diphtheria, tetanus and pertussis vaccine Diphtheria, tetanus, pertussis and polio vaccine Diphtheria, tetanus, pertussis, polio, hepatitis B and haemophilus influenzae type B vaccine Diprosone OV Diprosone OV Dipyridamole Disulfiram Diuretics Docetaxel Docetaxel Accord Docetaxel Sandoz	. 300 . 301 68 39 46 . 148 50 . 157 . 157 . 157
Diphtheria, tetanus and pertussis vaccine Diphtheria, tetanus, pertussis and polio vaccine Diphtheria, tetanus, pertussis, polio, hepatitis B and haemophilus influenzae type B vaccine Diprosone OV Diprosone OV Dipyridamole Disopyramide phosphate Disulfiram Diuretics Docetaxel Docetaxel Accord Docetaxel Sandoz Docusate sodium	. 300 . 301 68 39 46 . 148 50 . 157 . 157 . 157
Diphtheria, tetanus and pertussis vaccine Diphtheria, tetanus, pertussis and polio vaccine Diphtheria, tetanus, pertussis, polio, hepatitis B and haemophilus influenzae type B vaccine Diprosone OV Diprosone OV Dipyridamole Disopyramide phosphate Disulfiram Disulfiram Docetaxel Accord Docetaxel Sandoz Docusate sodium Docusate sodium with	. 300 . 301 68 68 68 39 46 . 148 50 . 157 . 157 23
Diphtheria, tetanus and pertussis vaccine Diphtheria, tetanus, pertussis and polio vaccine Diphtheria, tetanus, pertussis, polio, hepatitis B and haemophilus influenzae type B vaccine Diprosone OV Diprosone OV Dipyridamole Disopyramide phosphate Disulfiram Diuretics Docetaxel Docetaxel Accord Docetaxel Sandoz Docusate sodium with sennosides	. 300 . 301 68 68 68 68 68 46 . 148 50 . 157 . 157 . 157 23 23
Diphtheria, tetanus and pertussis vaccine Diphtheria, tetanus, pertussis and polio vaccine Diphtheria, tetanus, pertussis, polio, hepatitis B and haemophilus influenzae type B vaccine Diprosone OV Dipyridamole Disopyramide phosphate Disopyramide phosphate Disulfiram Disulfiram Diuretics Docetaxel Docetaxel Accord Docetaxel Sandoz Docusate sodium Docusate sodium with sennosides Dolutegravir	. 300 . 301 68 68 68 46 . 148 50 . 157 . 157 . 157 23 23 . 112
Diphtheria, tetanus and pertussis vaccine Diphtheria, tetanus, pertussis and polio vaccine Diphtheria, tetanus, pertussis, polio, hepatitis B and haemophilus influenzae type B vaccine Diprosone OV Dipyridamole Disopyramide phosphate Disopyramide phosphate Disulfiram Disulfiram Diuretics Docetaxel Docetaxel Accord Docetaxel Sandoz Docusate sodium Docusate sodium with sennosides Dolutegravir Dolutegravir with lamivudine	. 300 . 301 68 68 39 46 148 50 157 23 23 23 23 23
Diphtheria, tetanus and pertussis vaccine Diphtheria, tetanus, pertussis and polio vaccine Diphtheria, tetanus, pertussis, polio, hepatitis B and haemophilus influenzae type B vaccine Diprosone OV Diporsone OV Dipyridamole Disopyramide phosphate Disopyramide phosphate Disulfiram Disulfiram Disulfiram Docetaxel Docetaxel Docetaxel Accord Docetaxel Sandoz Docusate sodium with sennosides Dolutegravir Dolutegravir with lamivudine Domperidone	. 300 . 301 68 68 39 46 148 50 157 23 23 23 23 23
Diphtheria, tetanus and pertussis vaccine Diphtheria, tetanus, pertussis and polio vaccine Diphtheria, tetanus, pertussis, polio, hepatitis B and haemophilus influenzae type B vaccine Diprosone OV Diporosone OV Diporidamole Disopyramide phosphate Disopyramide phosphate Disulfiram Disulfiram Disulfiram Docetaxel Docetaxel Accord Docetaxel Sandoz Docetaxel Sandoz Docusate sodium with sennosides Dolutegravir Dolutegravir with lamivudine Domperidone Viatris	. 300 . 301 68 39 46 .148 50 .157 .157 .157 23 23 .112 .112 .113 .133
Diphtheria, tetanus and pertussis vaccine Diphtheria, tetanus, pertussis and polio vaccine Diphtheria, tetanus, pertussis, polio, hepatitis B and haemophilus influenzae type B vaccine Diprosone OV Dipordamole Disopyramide phosphate Disulfiram Disulfiram Disulfiram Disulfiram Docetaxel Docetaxel Docetaxel Accord Docetaxel Sandoz Docusate sodium with sennosides Dolutegravir Dolutegravir with lamivudine Domperidone Domperidone Viatris Donepezil hydrochloride	. 300 . 301 68 39 46 .148 50 .157 .157 .157 23 23 .112 .112 .113 .133 .147
Diphtheria, tetanus and pertussis vaccine Diphtheria, tetanus, pertussis and polio vaccine Diphtheria, tetanus, pertussis, polio, hepatitis B and haemophilus influenzae type B vaccine Diprosone OV Dipordamole Disopyramide phosphate Disopyramide phosphate Disulfiram Disulfiram Divertics Docetaxel Docetaxel Docetaxel Accord Docetaxel Sandoz Docetaxel Sandoz Docusate sodium with sennosides Dolutegravir with lamivudine Donperidone Domperidone Viatris Donepezil hydrochloride Dormase alfa	. 300 . 301 68 68 39 46 148 50 157 23 127 23 127 23 23 23 23 23 23 23 23 23
Diphtheria, tetanus and pertussis vaccine Diphtheria, tetanus, pertussis and polio vaccine Diphtheria, tetanus, pertussis, polio, hepatitis B and haemophilus influenzae type B vaccine Diprosone OV Diprosone OV Dipyridamole Disopyramide phosphate Disopyramide phosphate Disulfiram Disulfiram Diuretics Docetaxel Docetaxel Accord Docetaxel Sandoz Docusate sodium Docusate sodium Docusate sodium with sennosides Dolutegravir Dolutegravir with lamivudine Domperidone Domperidone Viatris Donege alfa Dortimopt	. 300 . 301 68 68 39 46 148 50 23 23 23 23 23 23 23 23 23 23 23 23 23
Diphtheria, tetanus and pertussis vaccine Diphtheria, tetanus, pertussis and polio vaccine Diphtheria, tetanus, pertussis, polio, hepatitis B and haemophilus influenzae type B vaccine Diprosone OV Dipordamole Disopyramide phosphate Disopyramide phosphate Disulfiram Disulfiram Divertics Docetaxel Docetaxel Docetaxel Accord Docetaxel Sandoz Docetaxel Sandoz Docusate sodium with sennosides Dolutegravir with lamivudine Donperidone Domperidone Viatris Donepezil hydrochloride Dormase alfa	. 300 . 301 68 68 39 46 50 .157 23 23 23 23 23 23 23 23 23 23 23 23 23 23 23 23 23 23 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 24 

Dosulepin [Dothiepin]	
hydrochloride 12	
Dosulepin Viatris12	
Dothiepin12	7
Dovato	
Doxazosin	
Doxazosin Clinect	4
Doxine	
Doxorubicin Ebewe	
Doxorubicin Lbeweining 15	' 7
Doxycycline	1
DP Lotion	
DP Lotn HC	
DP-Captopril 44	4
Dr Reddy's Omeprazole	9
Drofate4	8
Drugs Affecting Bone	
Metabolism 110	6
Dual blood glucose and blood ketone	
diagnostic test meter 14	4
Dulaglutide12	
Dulcolax SP Drop	4
Duocal Super Soluble Powder	6
Duolin	
Duolin Cipla	
Duolin HFA	
DuoResp Spiromax250	
Duride	
Durvalumab244	4
- E -	
- E - e-chamber La Grande	5
- E - e-chamber La Grande26 e-chamber Mask	5 4
- E - e-chamber La Grande	5 4
- E - e-chamber La Grande	5 4 5
- E - e-chamber La Grande26 e-chamber Mask	5 4 5 5
- E - e-chamber La Grande	54 55 0
- E - e-chamber La Grande	54 550 6
- E - e-chamber La Grande	5455066
- E - e-chamber La Grande	54550660
- E - e-chamber La Grande	545506607
- E - e-chamber La Grande	5455066070
- E - e-chamber La Grande	5455066070
- E - e-chamber La Grande	54550660700
- E - e-chamber La Grande	54550660700 1
- E - e-chamber La Grande	54550660700 1
- E - e-chamber La Grande	54550660700 17
- E - e-chamber La Grande	54550660700 176
- E - e-chamber La Grande	54550660700 1764
- E - e-chamber La Grande	54550660700 1764
- E - e-chamber La Grande	54550660700 17 643
- E - e-chamber La Grande	54550660700 17 6436
- E - e-chamber La Grande	54550660700 17 64364
- E - e-chamber La Grande	54550660700 17 643644
- E - e-chamber La Grande	54550660700 17 6436443
- E - e-chamber La Grande	54550660700 17 64364439
- E - e-chamber La Grande	54550660700 17 64364439
- E - e-chamber La Grande	54550660700 17 643644391

Elidel73
Elocon69
Elocon Alcohol Free69
Eltrombopag36
Eltroxin
EMB Fatol105
Emend Tri-Pack132
Emicizumab37
EMLA123
Empagliflozin 14
Empagliflozin with metformin
hydrochloride 14
Emsogen
Emtricitabine111
Emtricitabine with tenofovir
disoproxil 108
Emtriva 111
Emulsifying ointment70
Emulsifying Ointment ADE70
Enalapril maleate44
Enbrel 176
Endocrine Therapy 172
Endoxan
Energivit
Engerix-B
Enhertu
Enlafax XR128
Enoxaparin sodium40
Enstilar72
Ensure
Ensure Plus 285
Ensure Plus HN284
Ensure Plus HN RTH284
Ensure Two Cal HN RTH 287
Entacapone 121
Entacapone Viatris121
Entecavir106
Entecavir (Rex)106
Entresto 24/26 45
Entresto 49/51 45
Entresto 97/103 45
Entyvio241
Epilim131
Epilim Crushable131
Epilim IV131
Epilim S/F Liquid131
Epilim Syrup 131
Epipen255
Epipen Jr255
Epirubicin Ebewe157
Epirubicin hydrochloride 157
Eplerenone50
Epoetin alfa35
Epoprostenol61
Eptacog alfa [Recombinant factor
VIIa]37

Erbitux
Ergometrine maleate79
Erlotinib165
Erythrocin IV95
Erythromycin (as lactobionate)95
Erythromycin ethyl succinate95
Esbriet
Escitalopram128
Escitalopram (Ethics)128
Eskazole
Essential Amino Acid Mix
Essential Ethosuximide
Estraderm MX85
Estradiol Sandoz
Estradiol TDP Mylan85
Estradiol Viatris
Estradot
Estrofem
Estrogel
Etanercept
Ethambutol hydrochloride
Ethics Aspirin
Ethics Aspirin EC
Ethics Lisinopril
Ethipulo estradial with
Ethinyloestradiol with desogestrel
Ethinyloestradiol with
levenorgestrel 78
levonorgestrel
Ethinyloostradiol with
Ethinyloestradiol with norethisterone
Ethinyloestradiol with         78           norethisterone         78           Ethosuximide         129           Etopophos         157           Etoposide         157           Etoposide phosphate         157           Etravirine         110           Eurooside phosphate         69           Eurofolic         154           Evara         70           EVARA White Soft Paraffin         71           Everet         130           Everolimus         251           Evista         117           Evrysdi         143           Exelon Patch 10         147           Exelon Patch 5         147
Ethinyloestradiol with norethisterone
Ethinyloestradiol with       78         norethisterone
Ethinyloestradiol with norethisterone
Ethinyloestradiol with norethisterone
Ethinyloestradiol with norethisterone

-F-	
Factor eight inhibitor bypassing	
fraction	38
Famotidine	9
Famotidine Hovid	9
Fasenra	. 199
Faslodex	.173
Fatty Cream AFT	70
Fatty Emulsion Cream (Evara)	70
Febuxostat	. 119
Febuxostat (Teva)	.119
FEIBA NF	38
Felo 10 ER	49
Felo 5 ER	49
Felodipine	49
Fentanyl	. 125
Fentanyl Sandoz	. 125
Ferinject	33
Ferodan	33
Ferriprox	
Ferro-F-Tabs	
Ferro-Liquid	33
Ferro-tab	33
Ferrograd	33
Ferrosig	34
Ferrous fumarate	33
Ferrous fumarate with folic acid	
Ferrous sulfate	
Fexaclear	. 256
Fexofenadine hydrochloride	
Fibro-vein	38
Filgrastim	
Finasteride	80
Fingolimod	. 139
Firazyr	.255
Flagyl	. 104
FlagyI-S	
Flamazine	
Flecainide acetate	
Flecainide BNM	47
Flecainide Controlled Release	
Teva	47
Fleet Phosphate Enema	24
Flixonase Hayfever & Allergy	
Flixotide	.257
Flixotide Accuhaler	
Florinef	83
Fluanxol	
Flucil	
Flucloxacillin	
Flucloxacillin-AFT	
Flucloxin	
Flucon	
Fluconazole	
Fludara Oral	. 155
Fludarabine Ebewe	. 155

Fludarabine phosphate	155
Fludarabine Sagent	155
Fludrocortisone acetate	100
Fluids and Electrolytes	40
Fluids and Electrolytes	42
Flumetasone pivalate	266
Fluocortolone caproate with	
fluocortolone pivalate and	
cinchocaine	
Fluorometholone	
Fluorouracil	155
Fluorouracil Accord	
Fluorouracil sodium	74
Fluox	128
Fluoxetine hydrochloride	128
Flupenthixol decanoate	136
Flutamide	173
Flutamin	173
Fluticasone	257
Fluticasone furoate with	
umeclidinium and vilanterol	260
Fluticasone furoate with	200
vilanterol	050
Fluticasone propionate	264
Fluticasone with salmeterol	258
Flynn	
FML	267
Foban	
Folic acid	36
Folic Acid multichem	
Folic Acid Viatris	
Food Thickeners	287
Foods And Supplements For	
Inherited Metabolic Disease	288
Fortini	
Fortini Multi Fibre	280
Fortisip	285
Fortisip Multi Fibre	286
Fosamax	
Fosamax Plus	
Fosfomycin	
Framycetin sulphate	114
Freestyle Libre 2	200
Freestyle Libre 3 Plus	21
Frisium	
Frumil	
Frusemide	
Fucicort	
Fucidin	
Fucithalmic	
Fucithalmic S29	266
Fulvestrant	
Fungilin	
Furosemide [Frusemide]	50
Furosemide-Baxter	
fusidic acid	
Dermatological	67 60
- onnatorogroui	., 00

Infection 100
Sensory266
- G -
GA Explore 5
GA1 Anamix Infant
GA1 Anamix Junior
Gabapentin
Gacet
Galsulfase
Galvumet
Galvus
Gardasil 9
Gastrodenol10
Gaviscon Extra Strength
Gaviscon Extra Strength
Gaviscon Infant
Gazyva
Gefitinib
Gemcitabine Ebewe 155
Gemcitabine hydrochloride155
Gemtuzumab ozogamicin203
Gentamicin Amdipharm
Gentamicin Noridem98
Gentamicin sulphate98
Gilenya 139
Ginet
Glatiramer acetate 139
Glecaprevir with pibrentasvir 108
Glibenclamide11
Gliclazide11
Glipizide12
Glizide11
Glucagen Hypokit 10
Glucagon hydrochloride10
Glucerna Select
Glucose [Dextrose]42
Gluten Free Foods
Glycerin with sodium saccharin 274
Glycerin with sucrose
Glycerol
Alimentary24
Extemporaneous
Glyceryl trinitrate
Alimentary
Cardiovascular
Glycopyrronium
Glycopyrronium bromide
Glycopyrronium with
indacaterol
Glycosade
Glytactin Bettermilk
Gold Knight
Gold Knight XL
Goserelin
Gynaecological Anti-infectives79
-н-
Habitrol149

Haemophilus influenzae type B
vaccine 302
Haldol 136
Haldol Concentrate 136
Haldol Decanoas 136
Haloperidol 134
Haloperidol decanoate136
Harvoni 108
Havrix 1440 302
Havrix Junior
Haylor syrup256
HCU Anamix Infant
HCU Anamix Junior 289
HCU Anamix Junior LQ 289
HCU Explore 5289
HCU Express 15289
HCU Lophlex LQ289
healthE Calamine Aqueous
healthE Dimethicone 10%70
healthE Dimethicone 4% Lotion71
healthE Dimethicone 5%70
healthE Glycerol BP274
healthE Urea Cream70
Healtheries Simple Baking Mix 287
Hemastix81
Hemlibra
Heparin sodium41
Heparin Sodium Panpharma41
Heparinised saline
Heparon Junior
Hepatitis A vaccine
Hepatitis B recombinant
vaccine 303
Herzuma236
Hikma
Hiprex
Histaclear256
Holoxan153
Horleys Bread Mix
Horleys Flour
Hormone Replacement Therapy -
Systemic 85
HPV
Humalog11
Humalog Mix 2511
Humalog Mix 5011
Human papillomavirus (6, 11, 16, 18,
31, 33, 45, 52 and 58) vaccine
[HPV]
Humatin
Humira192
HumiraPen192
Humulin 30/70 11
Humulin NPH11
Humulin R10
Hyaluronic acid270

Hydralazine54
Hydralazine hydrochloride54
Hydralyte - Lemonade43
Hydrocortisone
Dermatological
Hormone
Hydrocortisone acetate7
Hydrocortisone acetate with
pramoxine hydrochloride7
Hydrocortisone and paraffin liquid
and lanolin 69
Hydrocortisone butyrate69, 74
Hydrocortisone with cinchocaine
Hydrocortisone with miconazole69
Hydrocortisone with natamycin and
neomycin
Hydrogen peroxide66
Hydroxocobalamin
Hydroxocobalamin Panpharma
hydroxycarbamide
Hydroxychloroquine sulphate
Hydroxyurea
[hydroxycarbamide] 157
Hygroton
Hylo-Fresh270
Hymenoptera256
Hyoscine butylbromide8
Hyoscine Butylbromide
(Adiramedica)
Hyoscine hydrobromide
Hyperuricaemia and Antigout
Hypromellose
Hypromellose with dextran
-l-
Ibiamox96
Ibrance
Ibrutinib157
Ibuprofen 115
Ibuprofen SR BNM 115
Icatibant255
Idarubicin hydrochloride158
Idursulfase
Ifosfamide153
llevro
lloprost62
Imatinib mesilate165
Imatinib-Rex 165
Imbruvica 157
Imfinzi
Imipramine Crescent
Imipramine hydrochloride
Imiquimod74
Immune Modulators112
Immunisation - Flu271
Immunisation Flu and Shingles 271
Immunisation Other271

Immunosuppressants 175
Incruse Ellipta259
Indacaterol257
Indapamide51
Infanrix IPV
Infanrix-hexa301
Infant Formulae
Infatrini
Infliximab
Influenza vaccine
Influvac Tetra
(2025 formulation) 305
Inhaled Corticosteroids257
Inhaled Long-acting
Beta-adrenoceptor Agonists 257
Inresa
Inspra
Instillagel Lido122
Insulin aspart11
Insulin aspart with insulin aspart
protamine
Insulin glargine11
Insulin glulisine
Insulin isophane11
Insulin isophane with insulin
neutral11
Insulin lispro
Insulin lispro with insulin lispro
protamine
Insulin neutral
Insulin pen needles
Insulin pump cartridge17
Insulin pump infusion set (steel
cannula)18
Insulin pump infusion set (steel
cannula, straight insertion) 18
Insulin pump infusion set (teflon
cannula)19
Insulin pump infusion set (teflon
cannula, angle insertion with
insertion device) 20
Insulin pump infusion set (teflon
cannula, flexible insertion with
insertion device) 20
Insulin pump infusion set (teflon
cannula, straight insertion with
insertion device) 20
Insulin pump infusion set (teflon
cannula, variable insertion)
Insulin pump reservoir21
Insulin pump with algorithm
Insulin syringes, disposable with
attached needle 16
Intelence 110
Interferon beta-1-alpha 139
Interferon beta-1-beta139

Intra-uterine device	.77
Invega Sustenna	137
Invega Trinza	137
Ipca-Allopurinol	
Ipca-Bisoprolol	110
Ipca-Ciprofloxacin	.47
Ipca-Donepezil	147
lpca-Escitalopram	128
IPCA-Frusemide	. 50
Ipca-Hydroxychloroquine	116
IPCA-Metoprolol	. 48
IPCA-Propranolol	
IPOL	311
Ipratropium bromide259,	264
Iressa	165
Irinotecan Actavis 100	155
Irinotecan hydrochloride	155
Irinotecan-Rex	155
Iron (as ferric carboxymaltose)	100
Iron polymaltose	. 00 04
Isentress	112
Isentress HD	
Ismo 20	
Ismo 40 Retard	
Isoleucine50	
Isoniazid	105
Isoniazid Teva	105
Isoniazid with rifampicin	105
Isoptin	
Isoptin Retard	.49
Isoptin SR	.49
Isopto Carpine	
Isosorbide mononitrate	
Isosource Standard	
Isotretinoin	
Ispaghula (psyllium) husk	23
Itch-Soothe	68
Itracap	.00 101
Itraconazole	
Itraconazole Kent	101
Itrazole	101
lvacaftor	
Ivermectin	.71
- J -	
Jadelle	
Jakavi	
Jardiamet	.14
Jardiance	
Jaydess	.86
Jevity HiCal RTH	
Jevity Plus RTH	
Jevity RTH	285
Jinarc	.51
Juno Pemetrexed	
- K -	.00
Kadcyla	228
าเล้นบ้าน	200

Kalydeco263
Kemadrin 122
Kenacomb
Kenacort-A 1084
Kenacort-A 4084
Kenalog in Orabase
Ketocal 3:1
KetoCal 4:1
Ketoconazole
Dermatological
Infection
Ketogenic Diet
Ketoprofen 115
KetoSens14
Keutrude 046
Keytruda
Kindergen
Kisqali
Klacid
Alimentary
Infection
Klaricid94
Kliogest86
Kliovance86
Kogenate FS
Konakion MM
Konsyl-D23
Kuvan
-L-
Labetalol48
Lacosamide 129
Lactulose
Laevolac
Lamictal
Lamivudine
Lamivudine Viatris 111
Lamivudine/Zidovudine Viatris111
Lamotrigine
Lamprene 104
Lanoxin
Lanoxin Paediatric Elixir
Lanoxin PG
Lanoxin S29
Lanoxin 329
Lansoprazole
Lantus
Lantus SoloStar11
Lanvis
Lanzol Relief
Largactil
Laronidase27
Lasix
Latanoprost269
Latanoprost with timolol269
Lax-Suppositories24
Lax-suppositories Glycerol24

	_
Laxsol	.23
Ledipasvir with sofosbuvir	
Leflunomide	116
Lenalidomide (Viatris)	158
Lenalidomide Viatris	158
Lenvatinib	166
Lenvima	
Letrole	175
Letrozole	
Leucine100	
Leukeran FC	152
Leukotriene Receptor	
Antagonists	262
Leuprorelin	.91
Leustatin	
Levetiracetam	130
Levetiracetam-AFT	
Levocabastine	268
Levocarnitine	
Levodopa with benserazide	
Levodopa with carbidopa	121
Levodopa with carbidopa and	
entacapone	121
Levomepromazine	134
Levomepromazine	
hydrochloride	135
Levonorgestrel	
Genito-Urinary78	
Hormone	
Levonorgestrel BNM	
Levothyroxine	. 86
Lidocaine [Lignocaine] 122-	123
Lidocaine [Lignocaine] hydrochloride	
hydrochloride	123
Lidocaine [Lignocaine] with	
prilocaine	
Lidocaine-Baxter	
Life Extension	.29
Lignocaine 122-	123
Linezolid	105
Lioresal Intrathecal	119
Lipid-Modifying Agents	
Lipitor	. 52
Liquigen	
Liraglutide	.12
Lisdexamfetamine dimesilate	
Lisinopril	.45
Lithium carbonate	
Livostin	
LMX4	123
Lo-Oralcon 20 ED	.78
Locacorten-Viaform ED's	266
Local preparations for Anal and	~
Rectal Disorders	
Locasol	
Locoid69	, 74

Loopid Crolo	
Locoid Crelo	
Locoid Lipocream	
Locorten-Vioform	266
Lodoxamide	268
Logem	
Lomide	268
Lomustine	153
Loniten	
Loperamide hydrochloride	
Lopinavir with ritonavir	
Lopinavir/Ritonavir Mylan	
Loprofin	292
Loprofin Mix	291
Lorafix	
Loratadine	
Lorazepam	
Lorstat	
Losartan Actavis	
Losartan potassium	.45
Losartan potassium with	
hydrochlorothiazide	. 45
Lovir	
Loxamine	
Lucrin Depot 1-month	
Lucrin Depot 3-month	
LumaCina	
Lumigan	
Lyllana	. 85
Lynparza	160
Lyrica	130
- M -	
m-Eslon	126
Mabthera	217
Macrobid	11/
	114
Macrogol 3350 with potassium	114
Macrogol 3350 with potassium chloride, sodium bicarbonate and	
Macrogol 3350 with potassium chloride, sodium bicarbonate and sodium chloride	. 24
Macrogol 3350 with potassium chloride, sodium bicarbonate and sodium chloride Madopar 125	. 24 121
Macrogol 3350 with potassium chloride, sodium bicarbonate and sodium chloride Madopar 125 Madopar 250	. 24 121 121
Macrogol 3350 with potassium chloride, sodium bicarbonate and sodium chloride Madopar 125 Madopar 250 Madopar 62.5	. 24 121 121 121
Macrogol 3350 with potassium chloride, sodium bicarbonate and sodium chloride Madopar 125 Madopar 250 Madopar 62.5 Madopar HBS	. 24 121 121 121 121
Macrogol 3350 with potassium chloride, sodium bicarbonate and sodium chloride Madopar 125 Madopar 250 Madopar 62.5 Madopar HBS Madopar Rapid	. 24 121 121 121 121 121
Macrogol 3350 with potassium chloride, sodium bicarbonate and sodium chloride Madopar 125 Madopar 250 Madopar 62.5 Madopar HBS Madopar Rapid Magnesium hydroxide	. 24 121 121 121 121 121 . 34
Macrogol 3350 with potassium chloride, sodium bicarbonate and sodium chloride	. 24 121 121 121 121 121 . 34 . 34
Macrogol 3350 with potassium chloride, sodium bicarbonate and sodium chloride	. 24 121 121 121 121 . 34 . 34 314
Macrogol 3350 with potassium chloride, sodium bicarbonate and sodium chloride	. 24 121 121 121 121 . 34 . 34 314 . 47
Macrogol 3350 with potassium chloride, sodium bicarbonate and sodium chloride	. 24 121 121 121 121 .34 .34 314 .47 .41
Macrogol 3350 with potassium chloride, sodium bicarbonate and sodium chloride	. 24 121 121 121 121 .34 .34 .47 .41 .74
Macrogol 3350 with potassium chloride, sodium bicarbonate and sodium chloride	. 24 121 121 121 121 . 34 . 34 . 34 . 47 . 41 . 74 271
Macrogol 3350 with potassium chloride, sodium bicarbonate and sodium chloride	. 24 121 121 121 121 .34 .34 314 .47 .41 .74 271 264
Macrogol 3350 with potassium chloride, sodium bicarbonate and sodium chloride	. 24 121 121 121 121 .34 .34 .41 .74 271 264 108
Macrogol 3350 with potassium chloride, sodium bicarbonate and sodium chloride	. 24 121 121 121 121 .34 .34 .47 .41 .74 271 264 108 267
Macrogol 3350 with potassium chloride, sodium bicarbonate and sodium chloride	. 24 121 121 121 121 . 34 . 41 . 74 271 264 108 267 267
Macrogol 3350 with potassium chloride, sodium bicarbonate and sodium chloride	. 24 121 121 121 121 . 34 . 41 . 74 271 264 108 267 267
Macrogol 3350 with potassium chloride, sodium bicarbonate and sodium chloride	. 24 121 121 121 121 .34 .34 .47 .47 .47 .47 .264 108 267 267 267

Mebendazole	93
Mebeverine hydrochloride	<mark>8</mark>
Medac	153
Medrol	83
Medroxyprogesterone acetate	
Genito-Urinary	78
Hormone	86
Mefenamic acid	
Megval	
Melatonin	
Melpha	
Melphalan	
Meningococcal (groups A, C, Y and	100
W-135) conjugate vaccine	200
Meningococcal B multicomponent	300
	200
vaccine	
MenQuadfi	
Menthol	
Mepolizumab	212
Mercaptopurine	
Mercilon 28	
Mesalazine	
Mesna	
Mestinon	115
Metabolic Disorder Agents	
Metabolics	
Metformin hydrochloride	12
Metformin Viatris	12
Methadone BNM	125
Methadone hydrochloride	125
Methenamine (hexamine)	
hippurate	114
Methopt	269
Methotrexate	
Methotrexate DBL Onco-Vial	156
Methotrexate Ebewe	
Methotrexate Sandoz	
Methyl hydroxybenzoate	
Methylcellulose	274
Methylcellulose with glycerin and	214
sodium saccharin	27/
Methylcellulose with glycerin and	214
SUCROSE	074
Methyldopa	
Methyldopa Viatris	
Methylnaltrexone bromide	23
Methylphenidate ER - Teva	
Methylphenidate hydrochloride	145
Methylphenidate hydrochloride	
extended-release	
Methylprednisolone	83
Methylprednisolone (as sodium	
succinate)	
Methylprednisolone aceponate	
Methylprednisolone acetate	
Methylxanthines	262

Metoclopramide Actavis 10 133
Metoclopramide hydrochloride 133
Metolazone51
Metopirone92
Metoprolol IV Mylan48
Metoprolol IV Viatris
Metoprolol succinate48
Metoprolol tartrate
Metronidamed104
Metronidazole 104
Metyrapone
Mexiletine hydrochloride
Miacalcic
Micolette
Miconazole
Miconazole nitrate
Dermatological
Genito-Urinary
Micreme
Micreme H
Microgynon 30
Midazolam
Midazolam Viatris
Midazolam-Pfizer
Midodrine
Midostaurin
Mifepristone
Minerals
Mini-Wright AFS Low Range
Mini-Wright Standard
Minidiab
MiniMad 2.0 Pessenvoir
MiniMed 3.0 Reservoir MMT-332A21
MiniMed Mio MMT-921A
MiniMed Mio MMT-921A
MiniMed Mio MMT-925A
MiniMed Mio MMT-925A
MiniMed Mio MMT-943A
MiniMed Mio MMT-945A
MiniMed Mio MMT-945A
MiniMed Mio MMT-975A
MiniMed Quick-Set MMT-396A
MiniMed Quick-Set MMT-397A
MiniMed Quick-Set MMT-398A
MiniMed Quick-Set MMT-398A
MiniMed Silhouette MMT-377A
MiniMed Silhouette MMT-378A
MiniMed Silhouette MMT-376A
MiniMed Sure-T MMT-864A
MiniMed Sure-T MMT-866A
MiniMed Sure-T MMT-874A
MiniMed Sure-T MMT-874A

Minims Pilocarpine	260
Minims Prednisolone	.209
Minipress	
Minirin	
Minirin Melt	91
Mino-tabs	97
Minocycline hydrochloride	97
Minomycin	
Minor Skin Infections	71
Minoxidil	54
Minoxidil Roma	
Mirena	
Miro-Amoxicillin	96
Mirtazapine	
Misoprostol	9
Mitomycin (Fresenius Kabi)	. 159
Mitomycin (Sagent)	.159
Mitomycin C	
Mitozantrone	
Mitozantrone Ebewe	
Mixtard 30	
MMA/PA Anamix Infant	1 1
MMA/PA Anamix Infant	.292
MMA/PA Anamix Junior	.292
MMA/PA Explore 5	.292
MMA/PA Express 15	. 292
Moclobemide	.128
Modafinil	.146
Modafinil Max Health	
Modavigil	
Moduretic	50
Molaxole	
Moments	
Mometasone furoate	
Monogen	
Montelukast	
Montelukast Viatris	.262
Moroctocog alfa [Recombinant facto	or
VIII]	38
Morphine hydrochloride	.125
Morphine sulphate	126
Motetis	
Mouth and Throat	
Movapo	
Moxifloxacin	
MSUD Anamix Infant	
MSUD Anamix Junior	.289
MSUD Anamix Junior LQ	.289
MSUD Explore 5	
MSUD Express 15	
MSUD Lophlex LQ 20	
MSUD Maxamum	
Mucolytics	
Mucosoothe	
Multiple Sclerosis Treatments	.138
Multivitamin renal	
Multivitamins	32

	=
Mupirocin6	6
Muscle Relaxants 11	9
Mvite	2
Myambutol 10	5
Mycobutin	6
MycoNail	
Mycophenolate mofetil	6
Mydriacyl	
Mylan (24 hr release)	0
Mylan Clomiphen	
Mylan Italy (24 hr release)	2 م
Myleran	2
mylife Inset soft	0
mylife Orbit micro1	
mylife YpsoPump Reservoir	1
mylife YpsoPump with CamAPS	_
FX	
Myloc CR4	
Mylotarg20	3
Myometrial and Vaginal Hormone	
Preparations7	9
Myozyme24	4
Mytolac 17	5
- N -	
Nadolol4	8
Nadolol BNM 4	8
Naglazyme	
Naloxone hydrochloride	1
Naltraccord	י פ
Naltrexone AOP14	0
Naltrexone hydrochloride14	0
Naltrexone Max Health	0
Naphazoline hydrochloride	0
Naprosyn SR 1000 11	5
Naprosyn SR 750 11	5
Naproxen 11	5
Narcaricin mite11	9
Nasal Preparations	4
Natalizumab13	9
Natulan162	2
Nausafix13	3
Nausicalm13	3
Navelbine S29 164	4
Nefopam hydrochloride 12	3
Neo-Cytamen S293	1
Neo-Mercazole8	
Neocate Gold	
Neocate Junior Unflavoured	
Neocate Junior Vanilla	
Neocate SYNEO	
Neoral	
Neostigmine metilsulfate	
Nepafenac	đ
Nepro HP (strawberry)	
Nepro HP (vanilla)	
Neulactil13	5

NeuroTabs
Nevirapine110
Nevirapine Viatris
Nicorandil
Nicotine
Nifedipine
Nifuran
Nilotinib
Nilstat
Alimentary
Genito-Urinary
Infection 101
Nimenrix
Nintedanib
Nipent
Niraparib159
Nirmatrelvir with ritonavir 109
Nitrates54
Nitroderm TTS 54
Nitrofurantoin 114
Nitrolingual Pump Spray54
Nivestim42
Nivolumab244
Nodia6
Noflam 250 115
Noflam 500 115
Non-Steroidal Anti-Inflammatory
Drugs 115
Nonacog gamma, [Recombinant
Nonacog gamma, [Recombinant
Nonacog gamma, [Recombinant Factor IX]
Nonacog gamma, [Recombinant
Nonacog gamma, [Recombinant Factor IX]

NovoSeven RT37
Nozinan134
Nozinan (Swiss)134
Nozinan S29
Nucala
Nuelin
Nuelin-SR
Nupentin 129
Nusinersen142
Nutilis
Nutren Diabetes278
Nutrient Modules275
Nutrini Energy Multi Fibre
Nutrini Energy RTH 280
Nutrini Low Energy Multi Fibre
Nutrini Peptisorb
Nutrini Peptisorb Energy
Nutrini RTH
Nutrison 800 Complete Multi
Fibre
Nutrison Advanced Peptisorb
Nutrison Concentrated
Nutrison Energy
Nutrison Energy Multi Fibre
Nutrison Multi Fibre
Nutrison RTH
Nyefax Retard
Nystatin
Alimentary
Genito-Urinary
Infection 101
NZB Low Gluten Bread Mix
- O -
Objecture and 010
Obinutuzumab
Obstetric Preparations81
Obstetric Preparations
Obstetric Preparations
Obstetric Preparations
Obstetric Preparations
Obstetric Preparations       81         Ocicure       9         Ocrelizumab       140         Ocrevus       140         Octocog alfa [Recombinant factor       141         VIII] (Advate)       38
Obstetric Preparations       81         Ocicure       9         Ocrelizumab       140         Ocrevus       140         Octocog alfa [Recombinant factor       180         VIII] (Advate)       38         Octocog alfa [Recombinant factor       38
Obstetric Preparations       81         Ocicure       9         Ocrelizumab       140         Ocrevus       140         Octocog alfa [Recombinant factor         VIII] (Advate)       38         Octocog alfa [Recombinant factor         VIII] (Advate)       38         Octocog alfa [Recombinant factor         VIII] (Kogenate FS)       38
Obstetric Preparations       81         Ocicure       9         Ocrelizumab       140         Octocog alfa [Recombinant factor       140         VIII] (Advate)       38         Octocog alfa [Recombinant factor       140         VIII] (Advate)       38         Octocog alfa [Recombinant factor       174
Obstetric Preparations       81         Ocicure       9         Ocrelizumab       140         Ocrovus       140         Octocog alfa [Recombinant factor         VIII] (Advate)       38         Octocog alfa [Recombinant factor         VIII] (Kogenate FS)       38         Octreotide       174         Octreotide GH       174
Obstetric Preparations       81         Ocicure       9         Ocrelizumab       140         Ocrovus       140         Octocog alfa [Recombinant factor       141         VIII] (Advate)       38         Octocog alfa [Recombinant factor       141         VIII] (Advate)       38         Octocog alfa [Recombinant factor       111         VIII] (Kogenate FS)       38         Octreotide       174         Octreotide GH       174         Octreotide long-acting       175
Obstetric Preparations       81         Ocicure       9         Ocrelizumab       140         Ocrovus       140         Octocog alfa [Recombinant factor       140         VIII] (Advate)       38         Octocog alfa [Recombinant factor       111]         VIII] (Advate)       38         Octocog alfa [Recombinant factor       111]         VIII] (Kogenate FS)       38         Octreotide       174         Octreotide GH       174         Octreotide long-acting       175         Oestradiol       85
Obstetric Preparations       81         Ocicure       9         Ocrelizumab       140         Ocrevus       140         Octocog alfa [Recombinant factor       111] (Advate)         VIII] (Advate)       38         Octocog alfa [Recombinant factor       111] (Kogenate FS)         VIII] (Kogenate FS)       38         Octreotide       174         Octreotide GH       174         Octreotide Iong-acting       175         Oestradiol       85         Oestradiol valerate       85
Obstetric Preparations       81         Ocicure       9         Ocrelizumab       140         Ocrevus       140         Octocog alfa [Recombinant factor       111] (Advate)         VIII] (Advate)       38         Octocog alfa [Recombinant factor       111] (Kogenate FS)         VIII] (Kogenate FS)       38         Octreotide       174         Octreotide GH       174         Octreotide long-acting       175         Oestradiol       85         Oestradiol valerate       85         Oestradiol with norethisterone       86
Obstetric Preparations       81         Ocicure       9         Ocrelizumab       140         Ocrevus       140         Octocog alfa [Recombinant factor       140         VIII] (Advate)       38         Octocog alfa [Recombinant factor       111]         VIII] (Kogenate FS)       38         Octreotide       174         Octreotide GH       174         Octrotide long-acting       175         Oestradiol       85         Oestradiol with norethisterone       86         Oestradiol with norethisterone       86         Oestradiol       85
Obstetric Preparations       81         Ocicure       9         Ocrelizumab       140         Ocrevus       140         Octocog alfa [Recombinant factor       141         VIII] (Advate)       38         Octocog alfa [Recombinant factor       111]         VIII] (Kogenate FS)       38         Octrootide       174         Octrootide GH       174         Octrootide long-acting       175         Oestradiol       85         Oestradiol valerate       85         Oestradiol with norethisterone       86         Oestroil       87         Oestroil       79
Obstetric Preparations       81         Ocicure       9         Ocrelizumab       140         Ocrevus       140         Octocog alfa [Recombinant factor       141         VIII] (Advate)       38         Octocog alfa [Recombinant factor       111         VIII] (Kogenate FS)       38         Octrootide       174         Octrootide GH       174         Octrootide long-acting       175         Oestradiol       85         Oestradiol valerate       85         Oestradiol with norethisterone       86         Oestriol       79         Hormone       86
Obstetric Preparations       81         Ocicure       9         Ocrelizumab       140         Ocrevus       140         Octocog alfa [Recombinant factor       140         VIII] (Advate)       38         Octocog alfa [Recombinant factor       111         VIII] (Kogenate FS)       38         Octreotide       174         Octreotide GH       174         Octrotide long-acting       175         Oestradiol       85         Oestradiol with norethisterone       86         Oestrol       79         Hormone       86         Oestrogens       85
Obstetric Preparations       81         Ocicure       9         Ocrelizumab       140         Ocrevus       140         Octocog alfa [Recombinant factor       140         VIII] (Advate)       38         Octocog alfa [Recombinant factor       140         VIII] (Kogenate FS)       38         Octreotide       174         Octreotide GH       174         Octreotide long-acting       175         Oestradiol valerate       85         Oestradiol with norethisterone       86         Oestriol       79         Hormone       86         Oestrogens       85
Obstetric Preparations       81         Ocicure       9         Ocrelizumab       140         Ocrevus       140         Octocog alfa [Recombinant factor       110         VIII] (Advate)       38         Octocog alfa [Recombinant factor       111         VIII] (Kogenate FS)       38         Octreotide       174         Octreotide GH       174         Octreotide long-acting       175         Oestradiol       85         Oestradiol with norethisterone       86         Oestrogens       85         Ofev       261         Oil in water emulsion       70
Obstetric Preparations       81         Ocicure       9         Ocrelizumab       140         Ocrevus       140         Octocog alfa [Recombinant factor       140         VIII] (Advate)       38         Octocog alfa [Recombinant factor       140         VIII] (Kogenate FS)       38         Octreotide       174         Octreotide GH       174         Octreotide long-acting       175         Oestradiol valerate       85         Oestradiol with norethisterone       86         Oestriol       79         Hormone       86         Oestrogens       85

Olbetam
Olopatadine
Olopatadine Teva
Olsalazine
Omalizumab214
Omeprazole
Omeprazole actavis 10
Omeprazole actavis 20
Omeprazole actavis 20
Omeprazole Teva
Omnitrope
Onbrez Breezhaler25
Oncaspar LYO16
OncoTICE
Ondansetron13
One-Alpha
One-Alpha S293
Opdivo24
Ora-Blend
Ora-Blend SF274
Ora-Plus274
Ora-Sweet274
Ora-Sweet SF274
Orabase
Oral and Enteral Feeds27
Oralcon 30 ED7
Oramorph 120
Oramorph CDC S29 120
Oratane60
Orgran
Ornidazole 104
Orphenadrine citrate120
Ortho-tolidine
Oruvail SR11
Osimertinib
Osmolite RTH
Other Endocrine Agents
Other Destroyen Preparations
Other Progestogen Preparations
Preparations
Other Skin Preparations
Ovestin
Genito-Urinary7
Hormone8
Oxaliplatin15
Oxaliplatin Accord15
Oxaliplatin Actavis 100 15
Oxaliplatin Ebewe15
Oxis Turbuhaler25
Oxpentifylline
Oxybutynin8
Oxycodone Amneal
Oxycodone hydrochloride
Oxycodone Lucis
Oxycodone Sandoz 120

Oxycodone Sandoz S29126
OxyContin126
Oxytocin80
Oxytocin BNM80
Oxytocin Panpharma80
Oxytocin with ergometrine
maleate
Ozurdex
- P -
Pacifen119
Pacimol124
Paclitaxel
Paclitaxel Actavis
Paclitaxel Ebewe
Padagis
Paediatric Seravit
Palbociclib
Paliperidone
Paliperidone palmitate
Palivizumab
Pamidronate disodium
Pamisol117
Pamol
Pancreatic enzyme
Pantoprazole
Panzop Relief
Papaverine hydrochloride54
Para-amino salicylic acid 105
Paracetamol 124
Paracetamol (Ethics) 124
Paracetamol + Codeine
(Relieve) 126
Paracetamol with codeine 126
Paraffin
Paraffin liquid with wool fat
Parasiticidal Preparations71
Parnate 128
Paromomycin99
Paroxetine
Paser 105
Paxam
Paxlovid109
Pazopanib169
Pazopanib Teva169
Peak flow meter
Pediasure
Pediasure Plus 280
Pediasure RTH280
Pegaspargase161
Pegasys112
Pegfilgrastim42
Pegylated interferon alfa-2a 112
Pembrolizumab246
Pemetrexed 156
Pemetrexed-AFT 156
Penicillamine116

Penicillin G	96
PenMix 30	11
PenMix 50	
Pentasa	
Pentostatin [Deoxycoformycin]	. 161
Pentoxifylline [Oxpentifylline]	54
Peptamen Junior	.280
Pepti-Junior	296
Perhexiline maleate	
Pericyazine	135
Perindopril	45
Periset	
Periset ODT	133
Perjeta	217
Permethrin	72
Perrigo	74
Pertuzumab	
Peteha	105
Pethidine hydrochloride	127
Pevaryl	. 127
Pexsig	07
Pfizer Exemestane	175
Pfizer S29	155
Pharmacy Services	. 100
Pharmascience	.2/1
Pheburane	.209
Phenasen	
Phenergan Elixir	. 150
Phenobarbitone	.20/
Phenobarbitone sodium	. 130
Extemporaneous	074
Nervous	.2/4
	. 142
Phenoxybenzamine hydrochloride	
Phenoxymethylpenicillin (Penicillin	44
	07
V)	97
Phenylalanine50	
Phenytoin sodium 129	-130
Phillips Milk of Magnesia	34
Phlexy 10	.290
Phosphate Phebra	43
Phosphorus	43
Phytomenadione	39
Pilocarpine hydrochloride	.269
Pilocarpine nitrate	.269
Pimafucort	
Pimecrolimus	73
Pine tar with trolamine laurilsulfate	
and fluorescein	
Pinetarsol	
Pioglitazone	
Pirfenidone	
Pizotifen	
PKU Anamix Infant	
PKU Anamix Junior	
PKU Anamix Junior Chocolate	.290

PKU Anamix Junior LQ	.290
PKU Anamix Junior Orange	.290
PKU Anamix Junior Vanilla	.290
PKU Build 10	.291
PKU Build 20 Chocolate	
PKU Build 20 Raspberry	
Lemonade	291
PKU Build 20 Smooth	291
PKU Build 20 Vanilla	291
PKU Explore 10	
PKU Explore 5	200
PKU Express 20	200
PKU First Spoon	200
PKU Glytactin RTD 15	201
PKU Glytactin RTD 15 Lite	.291
	.291
PKU GMPro LQ	.291
PKU GMPro Mix-In	
PKU GMPro Ultra Lemonade	.291
PKU GMPro Ultra Vanilla	.291
PKU Lophlex LQ 10	.290
PKU Lophlex LQ 20	. 290
PKU Lophlex Powder	. 290
PKU Lophlex Sensation 20	. 290
PKU Restore Powder	
PKU sphere20 Banana	.291
PKU sphere20 Chocolate	.291
PKU sphere20 Lemon	
PKU sphere20 Red Berry	.291
PKU sphere20 Vanilla	.291
PKU Start	.290
Plaquenil	. 116
Plendil ER	49
Pneumococcal (PCV13) conjugate	
vaccine	. 310
Pneumococcal (PPV23)	
polysaccharide vaccine	. 311
Pneumovax 23	.311
Podophyllotoxin	74
Polaramine	.256
Poliomyelitis vaccine	
Poloxamer	23
Polv-Gel	.270
Poly-Gel Poly-Tears	.269
Poly-Visc	270
Polycal	275
Polyethylene glycol 400 and	
propylene glycol	270
Pomalidomide	
Pomolide	
Ponstan	
Posaconazole	
Posaconazole Juno	
Potassium chloride4	
Potassium citrate	
Potassium iodate	
Povidone iodine	
	/ 1

Pradaxa41
Pramipexole hydrochloride 121
Pravastatin
Praziquantel
Prazosin44
Prazosin Mylan44
Pred Forte268
Prednisolone84
Prednisolone acetate268
Prednisolone sodium
phosphate
Prednisolone-AFT
Prednisone84
Prednisone Clinect84
Pregabalin130
Pregabalin Pfizer 130
Pregnancy Tests - hCG Urine
Premarin
Prevenar 13
Priadel
Primaquine 103
Primidone130
Primidone Clinect 130
Primolut N86
Priorix
Probenecid119
Probenecid-AFT 119
Procarbazine hydrochloride
Prochlorperazine
Prochlamorazina Drown 9
Prochlorperazine Brown & Burk
Burk 133
Prochlorperazine Max Health 133
Proctofoam7
Proctosedyl8
Procyclidine hydrochloride122
Progesterone
Proglicem
Progynova
Prolia
Promethazine hydrochloride257
Propafenone hydrochloride47
Propranolol48
Propylene glycol
Propylthiouracil87
Prostacur
Protaphane11
Protaphane Penfill
Protifar
Protionamide
Provera86
Provera HD86
Psoriasis and Eczema
Preparations72
PTU
Pulmicort Turbuhaler
Pulmozyme
1 anno2,110

Puri-nethol155
Puritan's Pride Vitamin
B-2 100 mg 27
Pyrazinamide106
Pyridostigmine bromide115
Pyridoxine hydrochloride
Pyridoxine multichem
Pyrimethamine
Pytazen SR
- Q -
Quantalan sugar free
Quetapel
Quetiapine 135
Quinapril
Qvar
RA-Morph 125
Ralicrom8
Raloxifene hydrochloride 117
Raltegravir potassium112
Ramipex
Ramipril
Ranbaxy-Cefaclor93
Rapamune252
Rasagiline121
Reandron 100084
Recombinant factor IX 36, 38
Recombinant factor VIIa37
Recombinant factor VIII
Rectogesic8
Redipred84
Relieve115
Relistor
Remicade
Renilon 7.5
Resonium-A43
Resource Beneprotein277
Respiratory Devices
Respiratory Stimulants
Retinol palmitate270
ReTrieve66
Retrovir
Revia
Revolade
Ribociclib
Riboflavin
Ribomustin
Bicit
Rifabutin
Rifadin
Rifadin Sanofi 106
Rifampicin106 Rifaximin
Rifinah
Rilinan
Riluzole122

RINVOQ	254
Riodine	
Risdiplam	143
Risedronate Sandoz	118
Risedronate sodium	
Risperdal	135
Risperdal Consta	137
Risperidone 135, 1	137
Risperidone (Teva)	125
Risperidone Sandoz	100
Risperon	100
Ritalin	100
Ritalin LA	
Ritonavir	
Rituximab (Mabthera)	217
Rituximab (Riximyo)	
Rivaroxaban	
Rivastigmine	
Rivastigmine Patch BNM 10	
Rivastigmine Patch BNM 5	
Rivotril	129
Riximyo	219
RIXUBIS	.38
Rizamelt	132
Rizatriptan	
Robinul	
Ronapreve	202
Ropin	
Ropinirole hydrochloride	121
Rosuvastatin	
Rosuvastatin Viatris	52
Rotarix	
Rotavirus oral vaccine	
Roxane-Propranolol	
Roxithromycin	.40
Rubifen	. 90
Rubifen SR	145
Rugby Capsaicin Topical Cream	
Musculoskeletal	
Nervous	
Rurioctocog alfa pegol [Recombinant	~~
factor VIII]	
Ruxolitinib	
Rydapt	
Rythmodan	.46
Rythmodan - Cheplafarm	
Rytmonorm	.47
- S -	
Sabril	
Sacubitril with valsartan	.45
SalAir	
Salazopyrin	8
Salazopyrin EN	8
Salbutamol	258
Salbutamol with ipratropium	
bromide	250

Salicylic acid73
Salmeterol
Sandomigran 132
Sandostatin LAR 175
Sanofi Primaquine 103
Sapropterin dihydrochloride28
Scalp Preparations74
Scopolamine - Mylan 133
Sebizole74
Secukinumab230
Sedatives and Hypnotics 141
Seebri Breezhaler259
Senna24
Senokot
SensoCard15
Serc
Serenace
Seretide
Seretide Accuhaler
Serevent
Serevent Accuhaler
Servent Accurate
Setrona
Sevredol 126
Sex Hormones Non
Contraceptive
Shingles vaccine
Shingrix
SII-Onco-BCG
Sildenafil59
Siltuximab233
Simvastatin53
Simvastatin Mylan53
Simvastatin Viatris53
Sinemet 121
Sinemet CR 121
Sintetica Baclofen Intrathecal 119
Sirolimus252
Sirturo 104
Siterone84
Slow-Lopresor
Sodibic
Sodium acid phosphate24
Sodium alginate6
Sodium benzoate28
Sodium bicarbonate
Blood
Extemporaneous274
Sodium calcium edetate
Sodium chloride
Blood
Beeniratory 264
Respiratory264 Sodium citrate with sodium lauryl
sulphoacetate
Sodium citro-tartrate
JUUIUIII CILIU-LAI LI ALE

Sodium cromoglicate	_
Alimentary	8
Sensory26	8
Sodium Fusidate [fusidic acid]	
Dermatological	7
Infection	
Sensory	
Sodium hyaluronate [Hyaluronic	Č
acid]27	0
Sodium phenylbutyrate2	Ř
Sodium picosulfate	
Sodium polystyrene sulphonate 4	3
Sodium tetradecyl sulphate	R
Sodium valproate	1
Sofradex	
Soframycin	6
Solgar	a
Solifenacin succinate	
Solifenacin succinate Max	1
Health	4
Solifenacin Viatris	÷
Solu-Cortef	
Solu-Medrol8	
Solu-Medrol-Act-O-Vial	ა ი
Somatropin (Omnitrope)	3 7
Sotalol	
Spacer device	о 0
Spazmol14 Spinal Muscular Atrophy14	
Spinraza14	
Spiolto Respimat	2
Spiractin	0
Spiriva	9
Spiriva Respimat	
Spironolactone	0
Sporanox	1
Stalevo	
Staphlex	
Stelara	
Stemetil	
Steril-Gene	
SteroClear	
Stesolid	9
Stimulants/ADHD Treatments	3
Stiripentol13	
Stomahesive	0
Strides Shasun10	
Stromectol	
Sucralfate1	
Sulfadiazin-Heyl10	0
Sulfadiazine Silver	7
Sulfadiazine sodium10	0
Sulfasalazine	
Sulphur7	3

Sulprix	
Ouiprix	134
Sumagran	132
Sumatriptan	
Sunitinib	
Sunitinib Pfizer	171
Sunscreens	74
Sunscreens, proprietary	74
Sustagen Hospital Formula	285
Sustagen Hospital Formula	
Active	285
Sustanon Ampoules	
Sylvant	
Symbicort Turbuhaler 100/6	258
Symbicort Turbuhaler 200/6	258
Symbicort Turbuhaler 400/12	258
Symmetrel	121
Sympathomimetics	. 54
Synacthen	. 84
Synacthen Depot	. 84
Synacthene Retard	. 84
Synagis	216
Synthroid	. 86
Syntometrine	80
Syrup (pharmaceutical grade)	274
Systane Unit Dose	270
́-т-	
Tacrolimus	
Dermatological	. 73
Oncology	254
Tacrolimus Sandoz	254
Tagrisso	107
	167
Taliglucerase alfa	. 29
Taliglucerase alfa Tambocor	29 47
Taliglucerase alfa Tambocor Tambocor German	29 47 47
Taliglucerase alfa Tambocor Tambocor German Tamoxifen citrate	29 47 47 174
Taliglucerase alfa Tambocor Tambocor German Tamoxifen citrate Tamoxifen Sandoz	29 47 47 174 174
Taliglucerase alfa Tambocor Tambocor German Tamoxifen citrate Tamoxifen Sandoz Tamsulosin hydrochloride	29 47 47 174 174 80
Taliglucerase alfa Tambocor Tambocor German Tamoxifen citrate Tamoxifen Sandoz Tamsulosin hydrochloride Tamsulosin-Rex	29 47 47 174 174 80 80
Taliglucerase alfa Tambocor Tambocor German Tamoxifen citrate Tamoxifen Sandoz Tamsulosin hydrochloride Tamsulosin-Rex Tandem Cartridge	29 47 174 174 80 80 17
Taliglucerase alfa Tambocor Tambocor German Tamoxifen citrate Tamoxifen Sandoz Tamsulosin hydrochloride Tamsulosin-Rex Tandem Cartridge Tandem t:slim X2 with Basal-IQ Tandem t:slim X2 with	29 47 174 174 80 80 17 16
Taliglucerase alfa Tambocor Tambocor German Tamoxifen citrate Tamoxifen Sandoz Tamsulosin hydrochloride Tamsulosin-Rex Tandem Cartridge Tandem t:slim X2 with Basal-IQ Tandem t:slim X2 with	29 47 174 174 80 80 17 16
Taliglucerase alfa Tambocor Tambocor German Tamoxifen citrate Tamoxifen Sandoz Tamsulosin hydrochloride Tamsulosin hydrochloride Tamsulosin-Rex Tandem Cartridge Tandem t:slim X2 with Basal-IQ Tandem t:slim X2 with Control-IQ	29 47 174 174 80 80 17 16
Taliglucerase alfa Tambocor Tambocor German Tamoxifen citrate Tamoxifen Sandoz Tamsulosin hydrochloride Tamsulosin hydrochloride Tamsulosin-Rex Tandem Cartridge Tandem t:slim X2 with Basal-IQ Tandem t:slim X2 with Control-IQ Tap water	29 47 47 174 174 80 80 17 16 274
Taliglucerase alfa Tambocor Tambocor German Tamoxifen citrate Tamoxifen Sandoz Tamsulosin hydrochloride Tamsulosin-Rex Tandem Cartridge Tandem t:slim X2 with Basal-IQ Tandem t:slim X2 with Control-IQ Tap water Taro	29 47 174 174 80 17 16 274 101
Taliglucerase alfa Tambocor Tambocor German Tamoxifen citrate Tamoxifen Sandoz. Tamsulosin hydrochloride Tamsulosin-Rex Tandem Cartridge Tandem t:slim X2 with Basal-IQ Tandem t:slim X2 with Control-IQ. Tap water Taro Tasigna	29 47 174 174 80 17 16 274 101 167
Taliglucerase alfa Tambocor Tambocor German Tamoxifen citrate Tamoxifen Sandoz Tamsulosin hydrochloride Tamsulosin-Rex Tandem Cartridge Tandem t:slim X2 with Basal-IQ Tandem t:slim X2 with Basal-IQ Tandem t:slim X2 with Control-IQ Tap water Taro Tasigna Tasigna	29 47 174 174 80 80 80 17 16 274 101 167 121
Taliglucerase alfa Tambocor Tambocor German Tamoxifen citrate Tamoxifen Sandoz Tamsulosin hydrochloride Tamsulosin-Rex Tandem Cartridge Tandem Cartridge Tandem t:slim X2 with Basal-IQ Tandem t:slim X2 with Control-IQ Tap water Taro Tasigna Tasigna Tasigna Taurine	29 47 47 174 174 80 17 16 274 101 167 121 29
Taliglucerase alfa Tambocor Tambocor German Tamoxifen citrate Tamoxifen Sandoz Tamsulosin hydrochloride Tamsulosin-Rex Tandem Cartridge Tandem Cartridge Tandem t:slim X2 with Basal-IQ Tandem t:slim X2 with Control-IQ Tap water Taro Tasigna Tasigna Tasigna Tasigna Taurine TCu 380 Plus Normal	29 47 47 174 174 80 80 17 16 274 101 167 121 29 77
Taliglucerase alfa Tambocor Tambocor German Tamoxifen citrate Tamoxifen Sandoz Tamsulosin hydrochloride Tamsulosin-Rex Tandem Cartridge Tandem Cartridge Tandem t:slim X2 with Basal-IQ Tandem t:slim X2 with Basal-IQ Tandem t:slim X2 with Control-IQ Tandem t:slim X2 with Control-IQ Taro Tasigna Tasigna Tasigna Tasigna Tasigna Taurine TCu 380 Plus Normal Tecentriq	29 47 47 174 174 80 80 80 17 16 274 101 167 121 29 77 243
Taliglucerase alfa Tambocor Tambocor German Tamoxifen citrate Tamoxifen Sandoz Tamsulosin hydrochloride Tamsulosin-Rex Tandem Cartridge Tandem t.slim X2 with Basal-IQ Tandem t.slim X2 with Basal-IQ Tandem t.slim X2 with Control-IQ Tandem t.slim X2 with Control-IQ Tag water Taro Taro Tasigna Tasmar Taurine TCu 380 Plus Normal Tecfidera	29 47 47 174 174 174 80 80 17 16 274 101 167 121 29 77 243 139
Taliglucerase alfa Tambocor Tambocor German Tamoxifen citrate Tamoxifen Sandoz Tamsulosin hydrochloride Tamsulosin hydrochloride Tamsulosin-Rex Tamdem Cartridge Tandem t:slim X2 with Basal-IQ Tandem t:slim X2 with Basal-IQ Tandem t:slim X2 with Control-IQ Tandem t:slim X2 with Control-IQ Tag water Taro Taro Tasigna Tasigna Tasigna Taurine TCu 380 Plus Normal Tecefidera Tegretol	29 47 47 174 174 174 80 80 17 16 274 101 167 121 29 77 243 139 129
Taliglucerase alfa Tambocor Tambocor German Tamoxifen citrate Tamoxifen Sandoz Tamsulosin hydrochloride Tamsulosin hydrochloride Tamsulosin Rex Tamdem Cartridge Tandem tslim X2 with Basal-IQ Tandem tslim X2 with Basal-IQ Tandem tslim X2 with Control-IQ Tandem tslim X2 with Control-IQ Tag water Taro Taro Tasigna Tasigna Tasigna Taurine TCu 380 Plus Normal Tecentriq Tecefidera Tegretol Tegretol AU	29 47 174 174 174 80 17 16 274 101 167 121 29 77 243 139 129 129
Taliglucerase alfa	29 47 174 174 174 80 80 17 16 274 101 167 121 29 77 243 139 129 129
Taliglucerase alfa Tambocor Tambocor German Tamoxifen citrate Tamoxifen Sandoz Tamsulosin hydrochloride Tamsulosin hydrochloride Tamsulosin Rex Tamdem Cartridge Tandem tslim X2 with Basal-IQ Tandem tslim X2 with Basal-IQ Tandem tslim X2 with Control-IQ Tandem tslim X2 with Control-IQ Tag water Taro Taro Tasigna Tasigna Tasigna Taurine TCu 380 Plus Normal Tecentriq Tecefidera Tegretol Tegretol AU	29 47 174 174 174 80 80 17 16 274 101 167 121 29 77 243 139 129 129 256

Temozolomide 162
Temozolomide-Taro162
Tenofovir disoproxil 106
Tenofovir Disoproxil Emtricitabine
Viatr
Tenofovir Disoproxil Viatris
Tenoxicam
Tensipine MR10
Tepadina
Terbinafine
Terbutaline sulphate
Teriflunomide
Teriflunomide Sandoz
Teriparatide
Teriparalide
Teriparatide - Teva
Testogel
Testosterone
Testosterone cipionate
Testosterone esters
Testosterone undecanoate
Tetrabenazine122
Tetrabromophenol81
Tetracosactrin84
Tetracycline97
Teva Lisinopril45
Teva-Ketoconazole101
Thalidomide163
Thalomid163
Theophylline
Thiamine hydrochloride31
Thiamine multichem31
THIO-TEPA 153
Thioguanine156
Thiotepa153
Thyroid and Antithyroid Agents
Ticagrelor
Ticagrelor Sandoz
Tilcotil
Timolol
Tiotropium bromide259
Tiotropium bromide with
olodaterol
Tivicay
TMP
Tobramycin
Infection 100
Sensory266
Tobramycin (Viatris) 100
Tobramycin BNM
Tobrex
Tocilizumab
Tofranil
Tolcapone
Tolvaptan
Topamax
Topical Products for Joint and

Muscular Pain	
Topiramate	131
Topiramate Actavis	131
Total parenteral nutrition (TPN)	43
TPN	43
Tramadol hydrochloride	127
Tramal SR 100	127
Tramal SR 150	127
Tramal SR 200	107
Trandate	127
	40
Tranexamic acid	
Tranylcypromine sulphate	128
Trastuzumab (Herzuma)	236
Trastuzumab deruxtecan	
Trastuzumab emtansine	238
Travatan	269
Travoprost	269
Treatments for Dementia	147
Treatments for Substance	
Dependence	147
Trelegy Ellipta	260
Trental 400	200
Tretinoin	
Dermatological	66
Oncology	160
Trexate	100
	100
Triamcinolone acetonide	~~
Alimentary	30
Dermatological	09
Hormone	84
Hormone	84
Hormone Triamcinolone acetonide with gramicidin, neomycin and nysta	<mark>84</mark> tin
Hormone Triamcinolone acetonide with gramicidin, neomycin and nysta Dermatological	84 tin 69
Hormone Triamcinolone acetonide with gramicidin, neomycin and nysta Dermatological Sensory	84 tin 69 266
Hormone Triamcinolone acetonide with gramicidin, neomycin and nysta Dermatological Sensory Trientine	84 tin 69 266 29
Hormone Triamcinolone acetonide with gramicidin, neomycin and nysta Dermatological Sensory Trientine Trientine Waymade	84 tin 69 266 29 29
Hormone Triamcinolone acetonide with gramicidin, neomycin and nysta Dermatological Sensory Trientine Trientine Waymade Trikafta	84 tin 69 266 29 29 263
Hormone Triamcinolone acetonide with gramicidin, neomycin and nysta Dermatological Sensory Trientine Trientine Waymade Trikafta	84 tin 69 266 29 29 263
Hormone Triamcinolone acetonide with gramicidin, neomycin and nysta Dermatological Sensory Trientine Trientine Waymade	84 tin 69 266 29 29 263
Hormone Triamcinolone acetonide with gramicidin, neomycin and nysta Dermatological Sensory Trientine Trientine Waymade Trikafta Trimethoprim Trimethoprim with	84 tin 69 266 29 29 263
Hormone Triamcinolone acetonide with gramicidin, neomycin and nysta Dermatological Sensory Trientine Waymade Trikafta Trimethoprim Trimethoprim with sulphamethoxazole	84 tin 69 266 29 29 263 100
Hormone Triamcinolone acetonide with gramicidin, neomycin and nysta Dermatological Sensory. Trientine Waymade Trinetha Waymade Trikafta Trimethoprim Trimethoprim with sulphamethoxazole [Co-trimoxazole]	84 tin 69 266 29 29 263 100
Hormone Triamcinolone acetonide with gramicidin, neomycin and nysta Dermatological Sensory. Trientine. Trientine Waymade Trikafta Trimethoprim Trimethoprim with sulphamethoxazole [Co-trimoxazole] Triovir.	84 tin 69 266 29 29 29 20 100 100 111
Hormone Triamcinolone acetonide with gramicidin, neomycin and nysta Dermatological Sensory. Trientine Waymade Trinethoprim Waymade Trimethoprim Trimethoprim with sulphamethoxazole [Co-trimoxazole] Triovir Trisequens	84 tin 69 266 29 263 263 100 100 111
Hormone Triamcinolone acetonide with gramicidin, neomycin and nysta Dermatological Sensory Trientine Waymade Trikafta Trimethoprim with sulphamethoxazole [Co-trimoxazole] Triovir Trisequens Trisul	84 tin 69 266 29 263 263 100 110 111 86 100
Hormone Triamcinolone acetonide with gramicidin, neomycin and nysta Dermatological Sensory Trientine Waymade Trihentine Waymade Trikafta Trimethoprim Trimethoprim with sulphamethoxazole [Co-trimoxazole] Triovir Trisequens Trisul Trophic Hormones	84 tin 69 266 29 29 263 100 110 111 86 100 87
Hormone Triamcinolone acetonide with gramicidin, neomycin and nysta Dermatological Sensory Trientine Waymade Trihafta Trimethoprim Trimethoprim with sulphamethoxazole [Co-trimoxazole] Trisequens Trisel Trophic Hormones Tropicamide	
Hormone Triamcinolone acetonide with gramicidin, neomycin and nysta Dermatological Sensory Trientine Waymade Trinethoprim Waymade Trimethoprim with sulphamethoxazole [Co-trimoxazole] Triovir Trisequens Trisul Trophic Hormones Tropicamide Truilicity	
Hormone Triamcinolone acetonide with gramicidin, neomycin and nysta Dermatological Sensory Trientine Waymade Trikafta Trimethoprim Trimethoprim with sulphamethoxazole [Co-trimoxazole] Triovi Trisequens Trisul Trophic Hormones Tropicamide Trulicity TruSteel	
Hormone Triamcinolone acetonide with gramicidin, neomycin and nysta Dermatological Sensory Trientine Waymade Trikafta Trikafta Trimethoprim with sulphamethoxazole [Co-trimoxazole] Trisequens Trisul Tropic Hormones Tropicamide Trulicity TruSteel Tryzan	84 tin 69 266 29 29 263 100 110 
Hormone Triamcinolone acetonide with gramicidin, neomycin and nysta Dermatological Sensory Trientine Waymade Trinethoprim Waymade Trikafta Trimethoprim with sulphamethoxazole [Co-trimoxazole] Triovir Trisequens Trisul Trophic Hormones Trophic Hormones Trophic Hormones Trophic Hormones Trulicity TruSteel Tryzan Tuberculin PPD [Mantoux] test	84 tin 69 226 29 263 100 110 100 110 
Hormone Triamcinolone acetonide with gramicidin, neomycin and nysta Dermatological Sensory Trientine Waymade Trikafta Trimethoprim with sulphamethoxazole [Co-trimoxazole] Trisequens Trisul Tropic Hormones Tropicamide Trulicity TruSteel Tryzan Tuberculin PPD [Mantoux] test	
Hormone Triamcinolone acetonide with gramicidin, neomycin and nysta Dermatological Sensory Trientine Waymade Trinethoprim With sulphamethoxazole [Co-trimoxazole] Trisequens Trisul Tropic Hormones Tropicamide Trulicity Trusteel Tryzan Tuberculin PPD [Mantoux] test Tubersol Two Cal HN	84 tin 
Hormone Triamcinolone acetonide with gramicidin, neomycin and nysta Dermatological Sensory Trientine Waymade Trikafta Trimethoprim with sulphamethoxazole [Co-trimoxazole] Triovir Trisequens Trisul Trophic Hormones Trophic Hormones Trophic Hormones Trusul Trubersol Two Cal HN TYR Anamix Infant	84 tin 
Hormone Triamcinolone acetonide with gramicidin, neomycin and nysta Dermatological Sensory Trientine Trinethoprim Waymade Trikafta Trimethoprim with sulphamethoxazole [Co-trimoxazole] Triovir Trisequens Trisul Trophic Hormones Tropicamide Trulicity Trulicity Tryzan Tubercolin PPD [Mantoux] test Tubersol Two Cal HN TYR Anamix Infant TYR Anamix Junior	
Hormone Triamcinolone acetonide with gramicidin, neomycin and nysta Dermatological Sensory Trientine Waymade Trinethoprim With sulphamethoxazole [Co-trimoxazole] Trisequens Trisul Trophic Hormones Trophic Hormones Trophic Hormones Trusteel Trusteel Tryzan Tubersol Two Cal HN TYR Anamix Infant	

TYR Lophlex LQ 20	. 292
TYR Sphere 20	.292
Tyrosine1000	.293
Tysabri	139
- U -	
UK Synacthen	84
Ultibro Breezhaler	260
Ultraproct	002.
Umeclidinium Umeclidinium with vilanterol	.209
Umecilainium with vilanterol	.260
Univent	264
Upadacitinib	
Ural	
Urea	70
Urex Forte	50
Urinary Agents	80
Urinary Tract Infections	.114
UroFos	.114
Uromitexan	159
Ursodeoxycholic acid	22
Ursosan	. 22
Ustekinumab	239
Utrogestan	86
- V -	
Vaccinations	200
Vaclovir	106
Valaciclovir	100
Valaciciovii	100
Valganciclovir	107
Valganciclovir Viatris	. 107
Valine50	.293
Vancomycin	.100
Vannair	.258
Varenicline Pfizer	. 149
Varenicline tartrate	. 149
Varicella vaccine [Chickenpox	
vaccine]	313
Varicella zoster vaccine [Shingles	
vaccine]	314
Varilrix	313
Various	.271
VariSoft	20
Vasodilators	54
Vasopressin Agonists	
Vasorex	48
Vebulis	62
Vedafil	59
Vedolizumab	241
Vegzelma	200
Vegzeinia	200 . 1 م
Veneleyte	100
Venclexta	103
Venetoclax	163
Venlafaxine	
Venomil	.256
VENOX	.256
Ventolin	
Vepesid	.157

Verapamil hydrochloride49
Vermox
Versacloz134
Vesanoid163
Vexazone12
Vfend 102
Viaderm KC
Victoza12
Vigabatrin 131
Vigisom
Vildagliptin
Vildagliptin with metformin
hydrochloride
Vimpat 129
Vinblastine sulphate
Vincristine sulphate
Vinorelbine
Vinorelbine Ebewe164
Vinorelbine Te Arai164
Viramune Suspension 110
ViruPOS
Vit.D3
Vita-B12
VitA-POS
Vitabdeck
Vital
Vitamin B complex
Vitamin B6 25
Vitamins
Vitarubin Depot Injection
Vivonex TEN
Voltaren
Voltaren D
Voltaren SR 115
Volumatic
Voriconazole
Votrient
Vttack
Vyvanse144 - W -
- w - Warfarin sodium41
Wart Preparations
Wasp venom allergy treatment
Wasp venom allergy treatment256 Water
Blood
Extemporaneous
White Soft Liquid Paraffin AFT70
Wool fat with mineral oil
- X -
Xaluprine
Xarelto
Xgeva
Xifaxan
XMET Maxamum
Xolair

KP Maxamum	290
Kylocaine	123
Kylocaine 2% Jelly	122
Kylocard 500	
Kyntha	
-Z-	
Zapril	44
Zarontin	129
Zaroxolyn	51
Zavedos	158
Zeffix	106
Zejula	159
Zematop	73
Zetlam	106
Ziagen	.111
Zidovudine [AZT]	.111
Zidovudine [AZT] with	
lamivudine	111
Ziextenzo	42
Ziextenzo AU	
Zimybe	
Zinc and castor oil	70
Zinc sulphate	
Zincaps	
Ziprasidone	135
Zista	
Zithromax	94
Zo-Rub HP	
Zo-Rub Osteo	.116
Zoladex	
Zoledronic acid	
Hormone	83
Musculoskeletal	118
Zoledronic acid Viatris	
Hormone	83
Musculoskeletal	118
Zopiclone	.142
Zopiclone Actavis	.142
Zostrix	.116
Zostrix HP	123
Zuclopenthixol decanoate	138
Zuclopenthixol hydrochloride	.135
Zusdone	135
Zyban	.148
Żypine	.135
Zypine ODT	135
Zyprexa Relprevv	136
Zytiga	
Źyvox	

Xolair AU ......214





