

APPLICANT (stamp or sticker acceptable) **PATIENT NHI:** **REFERRER** Reg No:

Reg No: First Names: First Names:

Name: Surname: Surname:

Address: DOB: Address:

..... Address:

.....

Fax Number: Fax Number:

Iloprost

Initial application — PAH monotherapy

Applications only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months.

Prerequisites(tick boxes where appropriate)

- ☐ Patient has pulmonary arterial hypertension (PAH)
- and ☐ PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications
- and ☐ PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV
- and
- ☐ PAH has been confirmed by right heart catheterisation

and ☐ A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair)

and ☐ A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg

and ☐ A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁻⁵)

and

☐ PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) †

or ☐ Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**

or ☐ Patient has PAH other than idiopathic / heritable or drug-associated type
- or ☐ Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease
- or ☐ Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures
- and
- ☐ Iloprost is to be used as PAH monotherapy

and

☐ Patient has experienced intolerable side effects on sildenafil and both the funded endothelin receptor antagonists (i.e. both bosentan and ambrisentan)

or ☐ Patient has an absolute contraindication to sildenafil and an absolute or relative contraindication to endothelin receptor antagonists

I confirm the above details are correct and that in signing this form I understand I may be audited.

Signed: Date:

Post application to Ministry of Health, Private Bag 3015, Wanganui – email: customerservice@health.govt.nz

APPLICANT (stamp or sticker acceptable) **PATIENT** NHI: **REFERRER** Reg No:
Reg No: First Names: First Names:
Name: Surname: Surname:
Address: DOB: Address:
..... Address:
.....
Fax Number: Fax Number:

Iloprost - continued

Initial application — PAH dual therapy

Applications only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months.

Prerequisites(tick boxes where appropriate)

- ☐ Patient has pulmonary arterial hypertension (PAH)
and ☐ PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications
and ☐ PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV
and
- ☐ PAH has been confirmed by right heart catheterisation
and ☐ A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair)
and ☐ A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg
and ☐ A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁻⁵)
and
- ☐ PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) †
or ☐ Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**
or ☐ Patient has PAH other than idiopathic / heritable or drug-associated type
- or ☐ Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease
or ☐ Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures
- and ☐ Iloprost is to be used as PAH dual therapy with either sildenafil or an endothelin receptor antagonist
and
- ☐ Patient has an absolute contraindication to or has experienced intolerable side effects on sildenafil
or ☐ Patient has an absolute or relative contraindication to or experienced intolerable side effects with a funded endothelin receptor antagonist
- and
- ☐ Patient has tried a PAH monotherapy for at least three months and remains in an unacceptable risk category according to a validated risk stratification tool**
or ☐ Patient is presenting in NYHA/WHO functional class III or IV, and in the opinion of the treating clinician would benefit from initial dual therapy

I confirm the above details are correct and that in signing this form I understand I may be audited.

Signed: Date:

Post application to Ministry of Health, Private Bag 3015, Wanganui – email: customerservice@health.govt.nz

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
.....	Address:	
.....		
Fax Number:		Fax Number:

Iloprost - continued

Initial application — PAH triple therapy

Applications only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months.

Prerequisites(tick boxes where appropriate)

- ☐ Patient has pulmonary arterial hypertension (PAH)
and
☐ PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications
and
☐ PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV
and
- ☐ PAH has been confirmed by right heart catheterisation
and
☐ A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair)
and
☐ A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg
and
☐ A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁻⁵)
and

☐ PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) †
or
☐ Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**
or
☐ Patient has PAH other than idiopathic / heritable or drug-associated type
- or
☐ Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease
or
☐ Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures
- and
☐ Iloprost is to be used as PAH triple therapy
and
- ☐ Patient is on the lung transplant list
or
☐ Patient is presenting in NYHA/WHO functional class IV
or

☐ Patient has tried PAH dual therapy for at least three months and has not experienced an acceptable response to treatment according to a validated risk stratification tool**
and
☐ Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario

I confirm the above details are correct and that in signing this form I understand I may be audited.

Signed: Date:

Post application to Ministry of Health, Private Bag 3015, Wanganui – email: customerservice@health.govt.nz

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
.....	Address:	
.....		
Fax Number:		Fax Number:

Iloprost - continued

Renewal

Current approval Number (if known):.....

Applications only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 2 years.

Prerequisites(tick box where appropriate)

☐ Patient is continuing to derive benefit from iloprost treatment according to a validated PAH risk stratification tool**

Note: † The European Respiratory Journal Guidelines can be found here: [2022 ECS/ERS Guidelines for the diagnosis and treatment of pulmonary hypertension PAH](#)

** the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

I confirm the above details are correct and that in signing this form I understand I may be audited.

Signed: Date:

Post application to Ministry of Health, Private Bag 3015, Wanganui – email: customerservice@health.govt.nz