December 2025 Volume 32

Editors:

Kaye Wilson, Doris Chong email: enquiry@pharmac.govt.nz Telephone +64 4 460 4990 Level 9, 40 Mercer Street PO Box 10 254 Wellington

Freephone Information Line 0800 66 00 50 (9am – 5pm weekdays)

Circulation

You can register to have an electronic version of the Pharmaceutical Schedule (link to PDF copy) emailed to your nominated email address each month by subscribing at pharmac.govt.nz/subscribe.

Production

Typeset automatically from XML and T_EX. XML version of the Schedule available from schedule.pharmac.govt.nz/pub/schedule

Programmers

Anrik Drenth

email: texschedule@pharmac.govt.nz
@Pharmaceutical Management Agency



ISSN 1179-3686

This work is licensed under the Creative Commons Attribution 4.0 International licence. In essence, you are free to copy, distribute and adapt it, as long as you attribute the work to Pharmac and abide by the other licence terms. To view a copy of this licence, visit: creativecommons.org/licenses/by/4.0/. Attribution to Pharmac should be in written form and not by reproduction of the Pharmac logo. While care has been taken in compiling this Schedule, Pharmac takes no responsibility for any errors or omissions, and shall not be liable for any consequences arising there from.

	Introducing Pharmac	2
Section A	General Rules	5
Section B	Alimentary Tract & Metabolism	6
	Blood & Blood Forming Organs	36
	Cardiovascular System	46
	Dermatologicals	67
	Genito Urinary System	76
	Hormone Preparations – Systemic	82
	Infections – Agents For Systemic Use	92
	Musculoskeletal System	115
	Nervous System	121
	Oncology Agents & Immunosuppressants	150
	Respiratory System & Allergies	264
	Sensory Organs	274
	Various	279
Section C	Extemporaneous Compounds (ECPs)	281
Section D	Special Foods	283
Section I	National Immunisation Schedule	306
	Index	323

Introducing Pharmac

Introducing Pharmac

The Pharmaceutical Management Agency (Pharmac) makes decisions that help control Government spending on pharmaceuticals. This includes community pharmaceuticals, hospital pharmaceuticals, vaccines and increasingly, hospital medical devices. Pharmac negotiates prices, sets subsidy levels and conditions, and makes decisions on changes to the subsidised list.

Pharmac's role:

"to secure for eligible people in need of pharmaceuticals, the best health outcomes that are reasonably achievable from pharmaceutical treatment and from within the amount of funding provided."

Pae Ora (Healthy Futures) Act 2022

To ensure our decisions are as fair and robust as possible we use a decision-making process that incorporates clinical, economic and commercial issues. We also seek the views of users and the wider community through consultation. The processes we generally use are outlined in our Operating Policies and Procedures.

Further information about Pharmac and the way we make funding decisions can be found on the Pharmac website at https://pharmac.govt.nz/about.

Purpose of the Pharmaceutical Schedule

The purpose of the Schedule is to list:

- the Community Pharmaceuticals that are subsidised by the Government and to show the amount of the subsidy paid to contractors, as well as the manufacturer's price and any access conditions that may apply;
- the Hospital Pharmaceuticals that may be used in Health NZ Hospitals, as well as any access conditions that may apply; and
- the Pharmaceuticals, including Medical Devices, used in Health NZ Hospitals for which national prices have been negotiated by Pharmac.

The Schedule does not show the final cost to Government of subsidising each Community Pharmaceutical or to Health NZ Hospitals in purchasing each Pharmaceutical, since that will depend on any rebate and other arrangements Pharmac has with the supplier and, for Pharmaceuticals used in Health NZ Hospitals, on any logistics arrangements put in place.

This book contains sections A to D and Section I of the Pharmaceutical Schedule and lists the Pharmaceuticals funded for use in the community, including vaccines, as well as haemophilia and cancer treatments given in Health NZ Hospitals. Section H lists the Pharmaceuticals that that can be used in Health NZ Hospitals and is a separate publication.


The Pharmaceuticals in this book are listed by therapeutic group, which is based on the WHO Anatomical Therapeutic Chemical (ATC) system. The listings are displayed alphabetically under each heading.

The index lists both chemical entities and product brand names.

Explaining pharmaceutical entries

The Pharmaceutical Schedule lists pharmaceuticals subsidised by the Government, the subsidy, the supplier's price and the access conditions that may apply.

Example



Glossary

Units of Measure

gramg kilogramkg international unitiu	mi mi mi
Abbreviations	
AmpouleAmp	Ge
CapsuleCap	Gr
Cream	Inf
DeviceDev	Ini
DispersibleDisp	Lic
EffervescentEff	Lo
EmulsionEmul	Oi
Enteric Coated EC	Sa

microgrammilligrammillilitre	mg
Gelatinous	
Granules	
Infusion	Inf
Injection	Inj
Liquid	Liq
Long Acting	LA
Ointment	Oint
Sachet	Sach

millimoleunit	
Solution	Supp Tab
Trans Dermal Delivery System	TDDS

Read the General Rules: https://pharmac.govt.nz/section-a.

SECTION B: ALIMENTARY TRACT AND METABOLISM

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	Brand or Generic Manufacturer
Antacids and Antiflatulents				
Antacids and Reflux Barrier Agents				
ALGINIC ACID Sodium alginate 225 mg and magnesium alginate 87.5 mg persachet		30	✓ (Gaviscon Infant
SODIUM ALGINATE Tab 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg - peppermint flavour	1.80 (17.99)	60	(Gaviscon Extra Strength
* Oral liq 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg per 10 ml		500 m		Acidex
Phosphate Binding Agents				
ALUMINIUM HYDROXIDE * Tab 600 mg CALCIUM CARBONATE	12.56	100	✓ /	Alu-Tab
Oral liq 1,250 mg per 5 ml (500 mg elemental per 5 ml) – Subsidy by endorsement		500 m 173 m		Roxane Calcium carbonate PAI 829
Only when prescribed for patients unable to swallow calc inappropriate and the prescription is endorsed according		s or v	where calciu	
Antidiarrhoeals				
Agents Which Reduce Motility				
LOPERAMIDE HYDROCHLORIDE − Up to 30 cap available on a * Tab 2 mg * Cap 2 mg Diamide Relief to be Principal Supply on 1 February 202	10.75 12.00	400 400	•	Nodia Diamide Relief
Rectal and Colonic Anti-inflammatories				
BUDESONIDE Cap modified-release 3 mg - Special Authority see SA2535	22.20	00		Pudaganida Ta Avai

⇒SA2535 Special Authority for Subsidy

Initial application — (Crohn's disease) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

1 Mild to moderate ileal, ileocaecal or proximal Crohn's disease; and

2 Any of the following:

continued...

✓ Budesonide Te Arai

90

Subsidy	Fully		Brand or	
(Manufacturer's Price)	Sub	sidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 2.1 Diabetes; or
- 2.2 Cushingoid habitus; or
- 2.3 Osteoporosis where there is significant risk of fracture; or
- 2.4 Severe acne following treatment with conventional corticosteroid therapy; or
- 2.5 History of severe psychiatric problems associated with corticosteroid treatment; or
- 2.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
- 2.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated).

Initial application — (collagenous and lymphocytic colitis (microscopic colitis)) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has a diagnosis of microscopic colitis (collagenous or lymphocytic colitis) by colonoscopy with biopsies.

Initial application — (gut Graft versus Host disease) from any relevant practitioner. Approvals valid for 6 months where patient has a gut Graft versus Host disease following allogenic bone marrow transplantation*.

Note: Indication marked with * is an unapproved indication.

Initial application — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has autoimmune hepatitis*; and
- 2 Patient does not have cirrhosis: and
- 3 Any of the following:
 - 3.1 Diabetes; or
 - 3.2 Cushingoid habitus: or
 - 3.3 Osteoporosis where there is significant risk of fracture; or
 - 3.4 Severe acne following treatment with conventional corticosteroid therapy; or
 - 3.5 History of severe psychiatric problems associated with corticosteroid treatment; or
 - 3.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
 - 3.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated); or
- 3.8 Adolescents with poor linear growth (where conventional corticosteroid use may limit further growth).

Note: Indication marked with * is an unapproved indication.

Renewal — (gut Graft versus Host disease) from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

HYDROCORTISONE ACETATE			
Rectal foam 10%, CFC-Free (14 applications)57.09	15 g OP	✓ Colifoam	
HYDROCORTISONE ACETATE WITH PRAMOXINE HYDROCHLORIDE			
Topical aerosol foam, 1% with pramoxine hydrochloride 1%26.55	10 g OP	✓ Proctofoam S29	
MESALAZINE			
Tab 400 mg49.50	100	✓ Asacol	
71.00	90	✓ Octasa S29	
Tab long-acting 500 mg56.10	100	✓ Pentasa	
Tab 800 mg85.50	90	✓ Asacol	
		✓ Asacol S29 S29	
Tab 1,600 mg85.50	60	✓ Asacol S29	
Modified release granules, 1 g118.10	100 OP	✓ Pentasa	
Enema 1 g per 100 ml41.30	7	✓ Pentasa	
Suppos 500 mg22.80	20	✓ Asacol	
Suppos 1 g50.96	28	✓ Pentasa	

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Generic
OLSALAZINE				
Tab 500 mg	56.02	60	•	Atnahs
				Olsalazine S29
	93.37	100	1	Dipentum
Cap 250 mg	53.00	100	1	Dipentum
SODIUM CROMOGLICATE Cap 100 mg	113.35	100	✓	Ralicrom
SULFASALAZINE				
* Tab 500 mg		100		Salazopyrin
* Tab EC 500 mg	20.54	100	•	Salazopyrin EN

Local preparations for Anal and Rectal Disorders

Antihaemorrhoidal Preparations

FLUOCORTOLONE CAPROATE WITH FLUOCORTOLONE PIVALATE AND CINCHOCAINE				
Oint 950 mcg, with fluocortolone pivalate 920 mcg, and				
cinchocaine hydrochloride 5 mg per g13.05	30 g OP	Ultraproct		
Suppos 630 mcg, with fluocortolone pivalate 610 mcg, and				
cinchocaine hydrochloride 1 mg8.61	12	Ultraproct		
HYDROCORTISONE WITH CINCHOCAINE				
Oint 5 mg with cinchocaine hydrochloride 5 mg per g15.00	30 g OP	✓ Proctosedyl		
Suppos 5 mg with cinchocaine hydrochloride 5 mg per g9.90	12	✓ Proctosedyl		

Management of Anal Fissures

⇒SA1329 Special Authority for Subsidy

CL VCODVDDONII IM DDOMIDE

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has a chronic anal fissure that has persisted for longer than three weeks.

Antispasmodics and Other Agents Altering Gut Motility

40.00	-	/ Dateback
19.00	5	✓ Robinul
2.25	20	 Hyoscine
		Butylbromide (Adiramedica)
1.91	5	✓ Spazmol
8.50	90	✓ Colofac
		2.25 20

Antiulcerants

Antisecretory and Cytoprotective

Subsidy		Fully	Brand or	
(Manufacturer's Price	e)	Subsidised	Generic	
\$	Per	1	Manufacturer	

Helicobacter Pylori Eradication

CLARITHROMYCIN

Tab 500 mg − Subsidy by endorsement......14.58 14 ✓ Klacid

- a) Maximum of 28 tab per prescription
- Subsidised only if prescribed for helicobacter pylori eradication and prescription is endorsed accordingly.
 Note: the prescription is considered endorsed if clarithromycin is prescribed in conjunction with a proton pump inhibitor and either amoxicillin or metronidazole.

H2 Antagonists

FA	MOTIDINE - Only on a prescription			
*	Tab 20 mg	4.86	100	✓ Famotidine
				Hovid S29
*	Tab 40 mg	10.27	100	✓ Famotidine Hovid
	•			MY (\$29)
*	Inj 10 mg per ml, 4 ml - Subsidy by endorsement	CBS	10	✓ Mylan S29
	Subsidy by endorsement – Subsidised for patients receiving			e care.

Proton Pump Inhibitors

LAN	NSOPRAZOLE		
*	Cap 15 mg4.04	100	✓ Lanzol Relief
*	Cap 30 mg5.43	100	✓ Lanzol Relief
ОМ	EPRAZOLE		
0	For omeprazole suspension refer Standard Formulae, page 281		
*	Cap 10 mg	90	✓ Omeprazole Teva
•			✓ Omeprazole actavis
			10
*	Cap 20 mg2.02	90	✓ Omeprazole Teva
-			✓ Omeprazole actavis
			20
*	Cap 40 mg3.18	90	Omeprazole Teva
	, ,		 Omeprazole actavis
			40
*	Powder - Only in combination52.00	5 g	✓ Midwest
	Only in extemporaneously compounded omeprazole suspension.	· ·	
*	Inj 40 mg ampoule with diluent	5	✓ Dr Reddy's
	, ,		Omeprazole
			✓ Ocicure S29
DAI	NTOPRAZOLE		
*		90	✓ Panzop Relief
*	Tab EC 20 mg	90	•
不	Tab EC 40 mg2.70	90	✓ Panzop Relief
C	ite Protective Agents		
3	ito i Tottotiivo Agenta		

COLLOIDAL BISMUTH SUBCITRATE			
Tab 120 mg	14.51	50	✓ Gastrodenol S29
SUCRALFATE			
Tab 1 g	35.50	120	
-	(48.28)		Carafate

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer

Bile and Liver Therapy

RIFAXIMIN - Special Authority see SA1461 below - Retail pharmacy 56

Xifaxan

⇒SA1461 Special Authority for Subsidy

Initial application only from a gastroenterologist, hepatologist or Practitioner on the recommendation of a gastroenterologist or hepatologist. Approvals valid for 6 months where the patient has hepatic encephalopathy despite an adequate trial of maximum tolerated doses of lactulose.

Renewal only from a gastroenterologist, hepatologist or Practitioner on the recommendation of a gastroenterologist or hepatologist. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient is benefiting from treatment.

Diabetes

Hyperglycaemic Agents

DIAZOXIDE - Special Authority see SA1320 below - Retail pharmacy

Cap 25 mg	110.00	100	✓ Proglicem S29
Cap 100 mg	280.00	100	✓ Proglicem S29
Oral liq 50 mg per ml	620.00	30 ml OP	✓ e5 Pharma S29

⇒SA1320 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months where used for the treatment of confirmed hypoglycaemia caused by hyperinsulinism.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient is benefiting from treatment.

GLUCAGON HYDROCHLORIDE

Inj 1 mg syringe kit - Up to 5 kit available on a PSO......32.00 ✓ Glucagen Hypokit

Insulin - Short-acting Preparations

INSL	11 11	N M	IFI	ITE	ΔΙ

\blacktriangle	Inj human 100 u per ml, 3 ml	42.66	5	 Actrapid Penfill
				✓ Humulin R
•	Ini human 100 u nor ml. 10 ml vial	25.26	1 OP	✓ Actranid

Humulin R

Insulin - Intermediate-acting Preparations

▲ Inj 100 iu per ml, 3 ml prefilled pen52.15	5	✓ NovoMix 30 FlexPen
INSULIN DEGLUDEC WITH INSULIN ASPART		
▲ Inj degludec 70 u with insulin aspart 30 u, 100 u per ml, 3 ml80.00	5	✓ Ryzodeg

Inj degludec 70 u with insulin aspart 30 u, 100 u per ml, 3 ml.........80.00

IIV	BULIN ISOPHANE		
\blacktriangle	Inj human 100 u per ml, 3 ml29.86	5	✓ Humulin NPH
			Protaphane Penfill
\blacktriangle	Inj human 100 u per ml, 10 ml vial17.68	1 OP	✓ Humulin NPH

Protaphane

70/30 Penfill

	Subsidy		Fully Brand or
	(Manufacturer's Price)		,
	\$	Per	✓ Manufacturer
NSULIN ISOPHANE WITH INSULIN NEUTRAL			
Inj human with neutral insulin 100 u per ml, 3 ml	42.66	5	✓ Humulin 30/70 ✓ PenMix 30
Inj human with neutral insulin 100 u per ml, 10 ml vial	25.26	1 OP	✓ Humulin 30/70
PenMix 30 Inj human with neutral insulin 100 u per ml, 3 ml to be	delisted 1 May 202		- Humani ooro
NSULIN LISPRO WITH INSULIN LISPRO PROTAMINE			
Inj lispro 25% with insulin lispro protamine 75% 100 u per ml,			
3 ml	42.66	5	Humalog Mix 25
Inj lispro 50% with insulin lispro protamine 50% 100 u per ml,		_	
3 ml	42.66	5	✓ Humalog Mix 50
Insulin - Long-acting Preparations			
SULIN GLARGINE			
Inj 100 u per ml, 10 ml	63.00	1	✓ Lantus
Inj 100 u per ml, 3 ml		5	✓ Lantus
Inj 100 u per ml, 3 ml disposable pen	94.50	5	✓ Lantus SoloStar
Insulin - Rapid Acting Preparations			
•			
NSULIN ASPART	20.02	4	✓ NavaDanid
、 Inj 100 u per ml, 10 ml		1 5	✓ NovoRapid✓ NovoRapid Penfill
Inj 100 u per ml, 3 ml syringe		5	✓ NovoRapid FlexPen
NSULIN GLULISINE			
Inj 100 u per ml, 10 ml	27.03	1	✓ Apidra
Inj 100 u per ml, 3 ml		5	✓ Apidra
Inj 100 u per ml, 3 ml disposable pen		5	✓ Apidra SoloStar
NSULIN LISPRO			
Inj 100 u per ml, 3 ml	59.52	5	✓ Humalog
Inj 100 u per ml, 10 ml vial	34.92	1 OP	✓ Humalog
Alpha Glucosidase Inhibitors			
CARBOSE			
€ Tab 50 mg	11.20	90	✓ Accarb
€ Tab 100 mg	17.38	90	✓ Accarb
Oral Hypoglycaemic Agents			
iLIBENCLAMIDE			
₹ Tab 5 mg	7.50	100	✓ Daonil
GLICLAZIDE			
€ Tab 80 mg	20.10	500	✓ Glizide
SLIPIZIDE			
₹ Tab 5 mg	6.86	100	✓ Minidiab
METFORMIN HYDROCHLORIDE			
₹ Tab immediate-release 500 mg	14.74	1,000	✓ Metformin Viatris
* Tab immediate-release 850 mg	11.28	500	✓ Metformin Viatris

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

5 90 25 90		exazone
5 90 25 90		
25 90	✓ V	
	• <u>v</u>	exazone
00 90	✓ <u>V</u> e	exazone
00 60	✓ Galleria	alvus
00 60	✓ G	alvumet
	✓ Galleria	alvumet
	00 60	00 60 ✓ G

GLP-1 Agonists

DULAGLUTIDE - Special Authority see SA2509 below - Retail pharmacy

Note: Not to be given in combination with another funded GLP-1 agonist or empagliflozin / empagliflozin with metformin hydrochloride unless receiving empagliflozin / empagliflozin with metformin hydrochloride for the treatment of heart failure.

Inj 1.5mg per 0.5 ml prefilled pen115.23 4 ✓ Trulicity

⇒SA2509 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has type 2 diabetes; and
- 2 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of all of the following funded blood glucose lowering agents for a period of least 6 months, where clinically appropriate: empagliflozin, metformin, and vildagliptin; and
- 3 Any of the following:
 - 3.1 Patient is Māori or any Pacific ethnicity*; or
 - 3.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)*; or
 - 3.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator*: or
 - 3.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a voung adult*: or
 - 3.5 Patient has diabetic kidney disease (see note b)*.

Notes: * Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

- a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack, ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.
- b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m2 in the presence of diabetes, without alternative cause identified.
- c) Funded GLP-1a treatment is not to be given in combination with funded (empagliflozin /empagliflozin with metformin hydrochloride) unless receiving funded (empagliflozin or empagliflozin in combination with metformin hydrochloride) for the treatment of heart failure.

LIRAGLUTIDE - Special Authority see SA2510 on the next page - Retail pharmacy

- a) Maximum of 9 inj per prescription
- b)
- a) Note: Not to be given in combination with another funded GLP-1 agonist or empagliflozin / empagliflozin with metformin hydrochloride unless receiving empagliflozin / empagliflozin with metformin hydrochloride for the treatment of heart failure.
- b) Maximum of 1 pack of 3 (6 mg per ml. 3 ml) prefilled pens will be funded per month.

Subs	,	Fully	Brand or
(Manufactur		lised	Generic
\$	Per	1	Manufacturer

⇒SA2510 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has type 2 diabetes; and
- 2 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of all of the following funded blood glucose lowering agents for a period of least 6 months, where clinically appropriate: empagliflozin, metformin, and vildaoliptin; and
- 3 Any of the following:
 - 3.1 Patient is Māori or any Pacific ethnicity*; or
 - 3.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)*; or
 - 3.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator*; or
 - 3.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult*; or
 - 3.5 Patient has diabetic kidney disease (see note b)*.

Notes: * Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

- a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack, ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.
- b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m2 in the presence of diabetes, without alternative cause identified.
- c) Funded GLP-1a treatment is not to be given in combination with funded (empagliflozin /empagliflozin with metformin hydrochloride) unless receiving funded (empagliflozin or empagliflozin in combination with metformin hydrochloride) for the treatment of heart failure.

SGI T2 Inhibitors

⇒SA2408 Special Authority for Subsidy

Initial application — **(heart failure reduced ejection fraction)** from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has heart failure; and
- 2 Patient is in NYHA functional class II or III or IV; and
- 3 Either:
 - 3.1 Patient has a documented left ventricular ejection fraction (LVEF) of less than or equal to 40%; or
 - 3.2 An ECHO is not reasonably practicable, and in the opinion of the treating practitioner the patient would benefit from treatment; and
- 4 Patient is receiving concomitant optimal standard funded chronic heart failure treatment.

Initial application — (Type 2 Diabetes) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 Patient has previously received an initial approval for a GLP-1 agonist; or
- 2 All of the following:
 - 2.1 Patient has type 2 diabetes; and

continued...

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 2.2 Any of the following:
 - 2.2.1 Patient is Māori or any Pacific ethnicity*; or
 - 2.2.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)*: or
 - 2.2.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator*; or
 - 2.2.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult*; or
 - 2.2.5 Patient has diabetic kidney disease (see note b)*; and
- 2.3 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of at least one blood-glucose lowering agent (e.g. metformin, vildagliptin, or insulin) for at least 3 months.

Notes: * Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

- a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack, ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.
- b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m2 in the presence of diabetes, without alternative cause.
- c) Funded [empagliflozin / empagliflozin with metformin hydrochloride] treatment is not to be given in combination with a funded GLP-1 unless receiving (empagliflozin / empagliflozin with metformin hydrochloride] for the treatment of heart failure.

EMPAGLIFLOZIN - Special Authority see SA2408 on the previous page - Retail pharmacy

*	Tab 10 mg58.56	30	Jardiance
*	Tab 25 mg58.56	30	Jardiance

EMPAGLIFLOZIN WITH METFORMIN HYDROCHLORIDE - Special Authority see SA2408 on the previous page - Retail pharmacy

*	Tab 5 mg with 1,000 mg metformin hydrochloride	58.56	60	Jardiamet
*	Tab 5 mg with 500 mg metformin hydrochloride	58.56	60	✓ Jardiamet
*	Tab 12.5 mg with 1,000 mg metformin hydrochloride	58.56	60	✓ Jardiamet
*	Tab 12.5 mg with 500 mg metformin hydrochloride	58 56	60	✓ .lardiamet

Diabetes Management

Ketone Testing

BLOOD KETONE DIAGNOSTIC TEST STRIP - Subsidy by endorsement

- a) Not on a BSO
- b) Maximum of 20 strip per prescription
- c) Up to 10 strip available on a PSO
- d) Patient has any of the following:
 - 1) type 1 diabetes; or
 - 2) permanent neonatal diabetes; or
 - 3) undergone a pancreatectomy; or
 - cvstic fibrosis-related diabetes: or
 - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

Dual Blood Glucose and Blood Ketone Testing

DUAL BLOOD GLUCOSE AND BLOOD KETONE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A dual blood glucose and blood ketone diagnostic test meter is subsidised for a patient who has:
 - 1) type 1 diabetes; or
 - 2) permanent neonatal diabetes: or
 - 3) undergone a pancreatectomy; or
 - 4) cystic fibrosis-related diabetes; or
 - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly. Only 1 meter per patient will be subsidised (no repeat prescriptions). For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Meter with 50 lancets, a lancing device and 10 blood glucose

Blood Glucose Testing

BLOOD GLUCOSE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A diagnostic blood glucose test meter is subsidised for a patient who:
 - 1) is receiving insulin or sulphonylurea therapy; or
 - 2) is pregnant with diabetes; or
 - 3) is on home TPN at risk of hypoglycaemia or hyperglycaemia; or
 - has a genetic or an acquired disorder of glucose homeostasis, excluding type 1 or type 2 diabetes and metabolic syndrome.

The prescription must be endorsed accordingly. Only one CareSens meter per patient will be subsidised (no repeat prescriptions). Patients already using the CareSens N POP meter and CareSens N meter are not eligible for a new meter, unless they have:

- 1) type 1 diabetes; or
- 2) permanent neonatal diabetes; or
- 3) undergone a pancreatectomy; or
- 4) cystic fibrosis-related diabetes.

For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Meter with 50 lancets, a lancing device and 10 diagnostic test

rips.......10.00 1 OP ✓ CareSens N

✓ CareSens N POP

20.00 ✓ CareSens N Premier

Note: Only 1 meter available per PSO

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	•	Manufacturer

BLOOD GLUCOSE DIAGNOSTIC TEST STRIP - Up to 50 test available on a PSO

The number of test strips available on a prescription is restricted to 50 unless:

- Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the
 prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed; or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly; or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

Test strips10.56	50 test OP	CareSens N
		✓ CareSens PRO

BLOOD GLUCOSE TEST STRIPS (VISUALLY IMPAIRED)

The number of test strips available on a prescription is restricted to 50 unless:

- Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the
 prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed; or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly; or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

Insulin Syringes and Needles

Subsidy is available for disposable insulin syringes, needles, and pen needles if prescribed on the same form as the one used for the supply of insulin or liraglutide or when prescribed for a patient and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of insulin or liraglutide.

INSULIN PEN NEEDLES - Maximum of 200 dev per prescription

*	29 g × 12.7 mm	10.95	100	✓ B-D Micro-Fine
	31 g × 5 mm		100	✓ B-D Micro-Fine
	31 g × 6 mm		100	✓ Berpu
	31 g × 8 mm		100	✓ B-D Micro-Fine
	32 a × 4 mm	10.95	100	✓ B-D Micro-Fine

		Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
INS	SULIN SYRINGES, DISPOSABLE WITH ATTACHED NEEDLE	E – Maximum of 200	dev p	per prescrip	tion
*	Syringe 0.3 ml with 29 g × 12.7 mm needle	13.56	100	√ E	B-D Ultra Fine
		1.36	10		
		(1.99)		E	3-D Ultra Fine
*	Syringe 0.3 ml with 31 g × 8 mm needle	13.56	100	✓ E	B-D Ultra Fine II
		1.30	10		
		(1.99)		E	3-D Ultra Fine II
*	Syringe 0.5 ml with 29 g x 12.7 mm needle	13.56	100	✓ [B-D Ultra Fine
		1.36	10		
		(1.99)		E	3-D Ultra Fine
*	Syringe 0.5 ml with 31 g × 8 mm needle	13.56	100	✓ E	B-D Ultra Fine II
		1.36	10		
		(1.99)		E	3-D Ultra Fine II
*	Syringe 1 ml with 29 g x 12.7 mm needle	13.56	100	✓ E	B-D Ultra Fine
		1.36	10		
		(1.99)		E	3-D Ultra Fine
*	Syringe 1 ml with 31 g × 8 mm needle	13.56	100	✓ [B-D Ultra Fine II
		1.36	10		
		(1.99)		E	3-D Ultra Fine II

Insulin Pumps

INSULIN PUMP WITH ALGORITHM - Special Authority see SA2367 below - Retail pharmacy

- a) Maximum of 1 dev per prescription
- b) Only on a prescription
- ✓ mylife YpsoPump
 with CamAPS FX
- ✓ Tandem t:slim
 X2 with Basal-IQ
- ✓ Tandem t:slim X2 with Control-IQ+

⇒SA2367 Special Authority for Subsidy

Initial application — (type 1 diabetes) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has type 1 diabetes; or
 - 1.2 The patient has permanent neonatal diabetes or specific monogenic diabetes subtypes with insulin deficiency, considered by the treating endocrinologist as likely to benefit; or
 - 1.3 The patient has Type 3c diabetes considered by the treating endocrinologist as likely to benefit (Type 3c diabetes includes insulin deficiency due to pancreatectomy, insulin deficiency secondary to cystic fibrosis or pancreatitis); or
 - 1.4 The patient has atypical inherited forms of diabetes; and
- 2 Patient has been evaluated by a diabetes multidisciplinary team for their suitability for insulin pump therapy; and
- 3 In the opinion of the treating relevant practitioner the patient would benefit from an Automated Insulin Delivery (AID) system.

Renewal — (type 1 diabetes) from any relevant practitioner. Approvals valid for 6 months where the patient is continuing to derive benefit according to the treatment plan agreed at induction.

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	1	Manufacturer	

Insulin Pump Consumables

⇒SA2536 Special Authority for Subsidy

Initial application — (type 1 diabetes) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has type 1 diabetes: or
 - 1.2 The patient has permanent neonatal diabetes or specific monogenic diabetes subtypes with insulin deficiency, considered by the treating endocrinologist as likely to benefit; or
 - 1.3 The patient has Type 3c diabetes considered by the treating endocrinologist as likely to benefit (Type 3c diabetes includes insulin deficiency due to pancreatectomy, insulin deficiency secondary to cystic fibrosis or pancreatitis); or
 - 1.4 The patient has atypical inherited forms of diabetes; and
- 2 Patient has been evaluated by a diabetes multidisciplinary team for their suitability for insulin pump therapy; and
- 3 In the opinion of the treating relevant practitioner the patient would benefit from an Automated Insulin Delivery (AID) system.

INSULIN PUMP CARTRIDGE - Special Authority see SA2536 above - Retail pharmacy

- a) Maximum of 50 cart per prescription
- b) Only on a prescription
- c) Maximum of 190 cartridges will be funded per year.

INSULIN PUMP INFUSION SET (STEEL CANNULA) - Special Authority see \$A2536 above - Retail pharmacy

- a) Maximum of 5 set per prescription
- b) Only on a prescription
- c) Maximum of 19 infusion sets will be funded per year.

*	6 mm steel needle; 60 cm tubing × 10	1 OP	✓ MiniMed Sure-T MMT-864A
*	6 mm steel needle; 80 cm tubing × 10130.00	1 OP	MiniMed Sure-T MMT-866A
*	8 mm steel needle; 60 cm tubing × 10130.00	1 OP	✓ MiniMed Sure-T MMT-874A
*	8 mm steel needle; 80 cm tubing × 10130.00	1 OP	✓ MiniMed Sure-T MMT-876A

(MiniMed Sure-T MMT-864A 6 mm steel needle; 60 cm tubing \times 10 to be delisted 1 October 2026) (MiniMed Sure-T MMT-866A 6 mm steel needle; 80 cm tubing \times 10 to be delisted 1 October 2026) (MiniMed Sure-T MMT-874A 8 mm steel needle; 60 cm tubing \times 10 to be delisted 1 October 2026) (MiniMed Sure-T MMT-876A 8 mm steel needle; 80 cm tubing \times 10 to be delisted 1 October 2026)

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	Subsidised	Generic	
\$	Per 🗸		Manufacturer	
	_			

INSULIN PUMP INFUSION SET (STEEL CANNULA, STRAIGHT INSERTION) – Special Authority see SA2536 on the previous page – Retail pharmacy

- a) Maximum of 5 sets per prescription
- b) Only on a prescription

	c) Maximum of 19 infusion sets will be funded per year.			
*	5.5 mm steel cannula; straight insertion; 45 cm line x 10 with 10 needles	136.00	1 OP	✓ mylife Orbit micro
*	5.5 mm steel needle; straight insertion; 60 cm line × 10 with 10 needles	136.00	1 OP	✓ mylife Orbit micro
*	5.5 mm steel needle; straight insertion; 80 cm line × 10 with 10 needles	136.00	1 OP	✓ mylife Orbit micro
*	8.5 mm steel needle; straight insertion; 60 cm line × 10 with 10 needles	136.00	1 OP	✓ mylife Orbit micro
*	8.5 mm steel needle; straight insertion; 80 cm line × 10 with 10 needles	136.00	1 OP	✓ mylife Orbit micro
*	6 mm steel cannula; straight insertion; 80 cm line × 10 with 10 needles	182.00	1 OP	✓ TruSteel
*	8 mm steel cannula; straight insertion; 80 cm line × 10 with 10 needles	182.00	1 OP	✓ TruSteel
*	6 mm steel cannula; straight insertion; 60 cm line × 10 with 10 needles		1 OP	✓ TruSteel
*	8 mm steel cannula; straight insertion; 60 cm line × 10 with 10 needles		1 OP	✓ TruSteel
	1011000100	102.00	. 01	- 11401001

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

INSULIN PUMP INFUSION SET (TEFLON CANNULA) - Special Authority see SA2536 on page 18 - Retail pharmacy

- a) Maximum of 5 set per prescription
- b) Only on a prescription

	c) Maximum of 19 infusion sets will be funded per year.		
*	13 mm teflon needle, 60 cm tubing × 10	1 OP	MiniMed Silhouette MMT-381A
*	17 mm teflon needle, 110 cm tubing × 10130.00	1 OP	MiniMed Silhouette MMT-377A
*	17 mm teflon needle, 60 cm tubing × 10130.00	1 OP	MiniMed Silhouette MMT-378A
*	6 mm teflon needle, 110 cm tubing × 10130.00	1 OP	MiniMed Quick-Set MMT-398A
*	6 mm teflon needle, 45 cm blue tubing × 10130.00	1 OP	✓ MiniMed Mio MMT-941A
*	6 mm teflon needle, 45 cm pink tubing × 10130.00	1 OP	✓ MiniMed Mio MMT-921A
*	6 mm teflon needle, 60 cm blue tubing × 10130.00	1 OP	✓ MiniMed Mio MMT-943A
*	6 mm teflon needle, 60 cm pink tubing × 10130.00	1 OP	✓ MiniMed Mio MMT-923A
*	6 mm teflon needle, 60 cm tubing × 10	1 OP	MiniMed Quick-Set MMT-399A
*	6 mm teflon needle, 80 cm blue tubing130.00	1 OP	✓ MiniMed Mio MMT-945A
*	6 mm teflon needle, 80 cm clear tubing × 10130.00	1 OP	✓ MiniMed Mio MMT-965A
*	6 mm teflon needle, 80 cm pink tubing × 10130.00	1 OP	✓ MiniMed Mio MMT-925A
*	9 mm teflon needle, 110 cm tubing × 10130.00	1 OP	MiniMed Quick-Set MMT-396A
*	9 mm teflon needle, 60 cm tubing × 10	1 OP	MiniMed Quick-Set MMT-397A
*	9 mm teflon needle, 80 cm clear tubing × 10130.00	1 OP	✓ MiniMed Mio MMT-975A

(MiniMed Silhouette MMT-381A 13 mm teflon needle, 60 cm tubing × 10 to be delisted 1 October 2026) (MiniMed Silhouette MMT-377A 17 mm teflon needle, 110 cm tubing × 10 to be delisted 1 October 2026) (MiniMed Silhouette MMT-378A 17 mm teflon needle, 60 cm tubing × 10 to be delisted 1 October 2026) (MiniMed Quick-Set MMT-398A 6 mm teflon needle, 110 cm tubing × 10 to be delisted 1 October 2026) (MiniMed Mio MMT-941A 6 mm teflon needle, 45 cm pink tubing × 10 to be delisted 1 February 2026) (MiniMed Mio MMT-921A 6 mm teflon needle, 45 cm pink tubing × 10 to be delisted 1 February 2026) (MiniMed Mio MMT-943A 6 mm teflon needle, 60 cm blue tubing × 10 to be delisted 1 February 2026) (MiniMed Mio MMT-943A 6 mm teflon needle, 60 cm pink tubing × 10 to be delisted 1 February 2026) (MiniMed Quick-Set MMT-399A 6 mm teflon needle, 60 cm blue tubing to be delisted 1 October 2026) (MiniMed Mio MMT-945A 6 mm teflon needle, 80 cm blue tubing to be delisted 1 February 2026) (MiniMed Mio MMT-95A 6 mm teflon needle, 80 cm clear tubing × 10 to be delisted 1 February 2026) (MiniMed Mio MMT-95A 9 mm teflon needle, 110 cm tubing × 10 to be delisted 1 October 2026) (MiniMed Quick-Set MMT-396A 9 mm teflon needle, 60 cm tubing × 10 to be delisted 1 October 2026) (MiniMed Quick-Set MMT-397A 9 mm teflon needle, 60 cm tubing × 10 to be delisted 1 October 2026) (MiniMed Quick-Set MMT-397A 9 mm teflon needle, 60 cm tubing × 10 to be delisted 1 February 2026) (MiniMed Mio MMT-975A 9 mm teflon needle, 60 cm tubing × 10 to be delisted 1 February 2026)

		Subsidy (Manufacturer's Price)	Per	Fully Subsidised	Brand or Generic Manufacturer
SA	SULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE IN 2536 on page 18 – Retail pharmacy a) Maximum of 5 sets per prescription b) Only on a prescription c) Maximum of 19 infusion sets will be funded per year.		SERTI	ON DEVICE) - Special Authority see
	13 mm teflon cannula; angle insertion; insertion device; 110 c line x 10 with 10 needles	182.00	1 OP	✓ Au	itoSoft 30
*	13 mm teflon cannula; angle insertion; insertion device; 60 cm line \times 10 with 10 needles		1 OP	✓ Au	itoSoft 30
see	BULIN PUMP INFUSION SET (TEFLON CANNULA, FLEXIBLE SA2536 on page 18 – Retail pharmacy a) Maximum of 5 set per prescription b) Only on a prescription c) Maximum of 19 infusion sets will be funded per year.		INSEF	RTION DEVI	CE) - Special Authority
	6 mm teflon cannula; flexible insertion; insertion device; 46 cr line × 10 with 10 needles	157.00	1 OP	✓ my	life Inset soft
*	6 mm teflon cannula; flexible insertion; insertion device; 60 cr line with integrated inserter \times 10 with 10 needles		1 OP	✓ my	life Inset soft
*	line x 10 with 10 needles	157.00	1 OP	✓ my	life Inset soft
*	line x 10 with 10 needles	157.00	1 OP	✓ my	life Inset soft
*	line × 10 with 10 needles		1 OP	✓ my	life Inset soft
see	SULIN PUMP INFUSION SET (TEFLON CANNULA, STRAIGH SA2536 on page 18 – Retail pharmacy a) Maximum of 5 sets per prescription b) Only on a prescription c) Maximum of 19 infusion sets will be funded per year. 6 mm teflon cannula; straight insertion; insertion device;	IT INSERTION WITH	1 INSE	RTION DEV	ICE) - Special Authority
	110 cm line × 10 with 10 needles		1 OP	✓ Au	toSoft 90
	line × 10 with 10 needles		1 OP	✓ Au	itoSoft 90
*	110 cm line × 10 with 10 needles		1 OP	✓ Au	itoSoft 90
*	9 mm teflon cannula; straight insertion; insertion device; 60 cl line × 10 with 10 needles		1 OP	✓ Au	itoSoft 90
Re	SULIN PUMP INFUSION SET (TEFLON CANNULA, VARIABLIA pharmacy a) Maximum of 5 set per prescription b) Only on a prescription c) Maximum of 19 infusion sets will be funded per year. 13 mm teflon cannula; variable insertion; 60 cm line × 10 with 10 needles	, ,	pecial A	·	SA2536 on page 18 –
	-		-		

Subsidy		Fully	Brand or
(Manufacturer's Price)	S	ubsidised	Generic
 \$	Per	✓	Manufacturer

INSULIN PUMP RESERVOIR - Special Authority see SA2536 on page 18 - Retail pharmacy

- a) Maximum of 90 cart per prescription
- b) Only on a prescription

*	10 × 1.6 ml glass reservoir for YpsoPump50.00	10 OP	✓ mylife YpsoPump Reservoir
*	10 × luer lock conversion cartridges 1.8 ml for paradigm pumps50.00	10 OP	✓ ADR Cartridge 1.8
*	Cartridge for 7 series pump; 3.0 ml × 1050.00	10 OP	✓ MiniMed 3.0 Reservoir

(ADR Cartridge 1.8 10 x luer lock conversion cartridges 1.8 ml for paradigm pumps to be delisted 1 October 2026) (MiniMed 3.0 Reservoir MMT-332A Cartridge for 7 series pump: 3.0 ml x 10 to be delisted 1 October 2026)

Continuous Glucose Monitor

CONTINUOUS GLUCOSE MONITOR (INTEROPERABLE) - Special Authority see \$A2537 below - Retail pharmacy Only on a prescription

-,-	ochool (b) and transmitter (bextoom do) - waximam or r dev		
	per prescription990.00	1 OP	✓ Dexcom G6
	Maximum of 5 dev will be funded per year.		
*	Sensor (Dexcom G7) - Maximum of 9 dev per prescription110.00	1	✓ Dexcom G7
	Maximum of 40 dev will be funded per year.		
*	Sensor (Freestyle Libre 3 Plus) - Maximum of 6 dev per		
	prescription99.46	1	✓ Freestyle Libre
			3 Plus

Maximum of 28 dev will be funded per year.

* Sensor (9) and transmitter (Dexcom G6) - Maximum of 1 dev

⇒SA2537 Special Authority for Subsidy

Initial application — (type 1 diabetes) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has type 1 diabetes: or
 - 1.2 The patient has permanent neonatal diabetes or specific monogenic diabetes subtypes with insulin deficiency, considered by the treating endocrinologist or relevant secondary health care professional as practicable, as likely to benefit: or
 - 1.3 The patient has Type 3c diabetes considered by the treating endocrinologist or relevant secondary health care professional as practicable, as likely to benefit (Type 3c diabetes includes insulin deficiency due to pancreatectomy, insulin deficiency secondary to cystic fibrosis or pancreatitis); or
 - 1.4 The patient has atypical inherited forms of diabetes; and
- 2 In the opinion of the treating relevant practitioner the patient would benefit from an Automated Insulin Delivery (AID) system.

MMT-332A

	Subsidy (Manufacturer's Price) \$	Subs	Fully sidised	Brand or Generic Manufacturer
CONTINUOUS GLUCOSE MONITOR (STANDALONE) - Speci	T			
Only on a prescription				
* Sensor (Dexcom ONE+) – Maximum of 9 dev per prescription Maximum of 40 dev will be funded per year.	on81.00	1	✓ D	excom ONE+
* Sensor (Freestyle Libre 2 Plus) - Maximum of 6 dev per				
prescription	99.46	1	√ F	reestyle Libre 2 Plus
Maximum of 28 dev will be funded per year.				
* Sensor (Freestyle Libre 2) – Maximum of 7 dev per prescrip Maximum of 29 dev will be funded per year. (Freestyle Libre 2 Sensor (Freestyle Libre 2) to be delisted 1 Ma		1	√ F	reestyle Libre 2

⇒SA2538 Special Authority for Subsidy

Initial application — **(type 1 diabetes)** from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 The patient has type 1 diabetes; or
- 2 The patient has permanent neonatal diabetes or specific monogenic diabetes subtypes with insulin deficiency, considered by the treating endocrinologist or relevant secondary health care professional as practicable, as likely to benefit; or
- 3 The patient has Type 3c diabetes considered by the treating endocrinologist or relevant secondary health care professional as practicable, as likely to benefit (Type 3c diabetes includes insulin deficiency due to pancreatectomy, insulin deficiency secondary to cystic fibrosis or pancreatitis); or
- 4 The patient has atypical inherited forms of diabetes.

Digestives Including Enzymes

PANCREATIC ENZYME

Cap pancreatin 150 mg (amylase 8,000 Ph Eur U, lipase			
10,000 Ph Eur U, total protease 600 Ph Eur U)	34.93	100	✓ Creon 10000
Cap pancreatin 300 mg (amylase 18,000 Ph Eur U, lipase			
25,000 Ph Eur U, total protease 1,000 Ph Eur U)	94.38	100	✓ Creon 25000
Modified release granules pancreatin 60.12 mg (amylase			
3,600 Ph Eur U, lipase 5,000 Ph Eur U, protease 200 F	Ph		
Eur U)	34.93	20 g OP	Creon Micro
URSODEOXYCHOLIC ACID - Special Authority see SA2448 b	oelow – Retail pha	rmacy	
Cap 250 mg	33.95	100	Ursosan

⇒SA2448 Special Authority for Subsidy

Initial application — (Alagille syndrome or progressive familial intrahepatic cholestasis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Either:

- 1 Patient has been diagnosed with Alagille syndrome; or
- 2 Patient has progressive familial intrahepatic cholestasis.

Initial application — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic severe drug induced cholestatic liver injury; and
- 2 Cholestatic liver injury not due to Total Parenteral Nutrition (TPN) use in adults; and
- 3 Treatment with ursodeoxycholic acid may prevent hospital admission or reduce duration of stay.

Initial application — (Primary biliary cholangitis) from any relevant practitioner. Approvals valid for 6 months for applications

continued...

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

meeting the following criteria:

Both:

- 1 Primary biliary cholangitis confirmed by antimitochondrial antibody titre (AMA) > 1:80, and raised cholestatic liver enzymes with or without raised serum IgM or, if AMA is negative, by liver biopsy; and
- 2 Patient not requiring a liver transplant (bilirubin > 100 umol/l; decompensated cirrhosis).

Initial application — (**Pregnancy**) from any relevant practitioner. Approvals valid for 6 months where the patient diagnosed with cholestasis of pregnancy.

Initial application — (Haematological Transplant) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient at risk of veno-occlusive disease or has hepatic impairment and is undergoing conditioning treatment prior to allogenic stem cell or bone marrow transplantation; and
- 2 Treatment for up to 13 weeks.

Initial application — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Paediatric patient has developed abnormal liver function as indicated on testing which is likely to be induced by Total Parenteral Nutrition (TPN); and
- 2 Liver function has not improved with modifying the TPN composition.

Renewal — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 6 months where the patient continues to benefit from treatment.

Renewal — (**Pregnancy/Primary biliary cholangitis**) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months where the paediatric patient continues to require TPN and who is benefiting from treatment, defined as a sustained improvement in bilirubin levels.

Initial application — (prevention of sinusoidal obstruction syndrome) from any relevant practitioner. Approvals valid without further renewal unless notified where the individual has leukaemia/lymphoma and requires prophylaxis for medications/therapies with a high risk of sinusoidal obstruction syndrome.

Laxatives

Bulk-forming Agents		
ISPAGHULA (PSYLLIUM) HUSK – Only on a prescription * Powder for oral soln22.10	500 g OP	✓ Konsyl-D
Faecal Softeners		
DOCUSATE SODIUM — Only on a prescription # Tab 50 mg	100 100	✓ <u>Coloxyl</u> ✓ <u>Coloxyl</u> ✓ Solax
3.50 (Laxsol Tab 50 mg with sennosides 8 mg to be delisted 1 May 2026)	200	✓ Laxsol
POLOXAMER – Only on a prescription Not funded for use in the ear. * Oral drops 10%	30 ml OP	✓ Coloxyl

Subsidy		Fully	Brand or	
(Manufacturer's Price) 5	Subsidised	Generic	
\$	Per	✓	Manufacturer	

Opioid Receptor Antagonists - Peripheral

⇒SA1691 Special Authority for Subsidy

Initial application — (Opioid induced constipation) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient is receiving palliative care; and
- 2 Either:
 - 2.1 Oral and rectal treatments for opioid induced constipation are ineffective; or
 - 2.2 Oral and rectal treatments for opioid induced constipation are unable to be tolerated.

Osmotic Laxatives

GLYCEROL			
* Suppos 2.8/4.0 g - Only on a prescription	12.39	20	✓ Lax-suppositories Glycerol
Lax-suppositories Glycerol to be Principal Supply on 1 February	uary 2026		
LACTULOSE - Only on a prescription			
* Oral liq 10 g per 15 ml Laevolac to be Principal Supply on 1 February 2026	6.16	500 ml	✓ Laevolac
MACROGOL 3350 WITH POTASSIUM CHLORIDE, SODIUM BICAF Powder for oral soln 13.125 g with potassium chloride 46.6 mg,	RBONATE AND	SODIUM C	HLORIDE
sodium bicarbonate 178.5 mg and sodium chloride 350.7 mg	g 8.50	30	✓ APO Health Macrogol S29
	10.15		✓ Molaxole
	12.19		✓ Movicol
(APO Health Macrogol S29) Powder for oral soln 13.125 g with pota sodium chloride 350.7 mg to be delisted 1 January 2026)	ssium chloride 4	16.6 mg, soc	lium bicarbonate 178.5 mg and
SODIUM ACID PHOSPHATE - Only on a prescription			
Enema 16% with sodium phosphate 8%	3.70	1	✓ Fleet Phosphate Enema
SODIUM CITRATE WITH SODIUM LAURYL SULPHOACETATE – Enema 90 mg with sodium lauryl sulphoacetate 9 mg per ml,	Only on a presc	ription	
5 mlMicolette to be Principal Supply on 1 February 2026	36.89	50	✓ Micolette

Stimulant Laxatives

BISACODYL - Only on a prescription * Tab 5 mg * Suppos 10 mg		200 10	✓ Bisacodyl Viatris✓ Lax-Suppositories
SENNA – Only on a prescription * Tab, standardised		100	
rab, standardised	(9.38) 0.43	20	Senokot
	(2.06)		Senokot

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price)		Fully dised	Brand or Generic
	` \$	Per	1	Manufacturer
SODIUM PICOSULFATE - Special Authority see SA2053 below	 Retail pharmacy 			
Oral soln 7.5 mg per ml	7.40 30) ml OP	✓ D	ulcolax SP Drop

⇒SA2053 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The patient is a child with problematic constipation despite an adequate trial of other oral pharmacotherapies including macrogol where practicable; and
- 2 The patient would otherwise require a high-volume bowel cleansing preparation or hospital admission.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Metabolic Disorder Agents

ALGLUCOSIDASE ALFA - Special Authority see SA1986 below - Retail pharmacy ✓ Myozyme

⇒SA1986 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient is aged up to 24 months at the time of initial application and has been diagnosed with infantile Pompe disease; and
- 2 Any of the following:
 - 2.1 Diagnosis confirmed by documented deficiency of acid alpha-glucosidase by prenatal diagnosis using chorionic villus biopsies and/or cultured amniotic cells; or
 - 2.2 Documented deficiency of acid alpha-glucosidase, and urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides; or
 - 2.3 Documented deficiency of acid alpha-glucosidase, and documented molecular genetic testing indicating a disease-causing mutation in the acid alpha-glucosidase gene (GAA gene); or
 - 2.4 Documented urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides, and molecular genetic testing indicating a disease-causing mutation in the GAA gene; and
- 3 Patient has not required long-term invasive ventilation for respiratory failure prior to starting enzyme replacement therapy (ERT): and
- 4 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by ERT or might be reasonably expected to compromise a response to ERT; and
- 5 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks.

Renewal only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks; and
- 3 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 4 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by ERT; and
- 5 Patient has not developed another medical condition that might reasonably be expected to compromise a response to
- 6 There is no evidence of life threatening progression of respiratory disease as evidenced by the needed for > 14 days of invasive ventilation: and
- 7 There is no evidence of new or progressive cardiomyopathy.

ARGININE - Special Authority see SA2042 on the next p	page – Retail pharmacy		
Tab 1,000 mg	CBS	90	Clinicians
Cap 500 mg	CBS	50	✓ Solgar
Powder	CBS	400 g	✓ Biomed

Subsidy (Manufactured Price)		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

⇒SA2042 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to arginine supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to arginine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

⇒SA1987 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has a confirmed diagnosis of homocystinuria; and
- 2 Any of the following:
 - 2.1 A cystathionine beta-synthase (CBS) deficiency; or
 - 2.2 A 5,10-methylene-tetrahydrofolate reductase (MTHFR) deficiency; or
 - 2.3 A disorder of intracellular cobalamin metabolism; and
- 3 An appropriate homocysteine level has not been achieved despite a sufficient trial of appropriate vitamin supplementation. **Renewal** only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

COENTYME 010 - Special Authority see \$42039 below - Retail pharmacy.

		pecial Authority see SA2039 below – Retail pharmacy	COENZYME Q10 - Sp
✓ Solgar	30	CBS	Cap 120 mg
✓ Go Healthy	60	CBS	Cap 160 mg

⇒SA2039 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to coenzyme Q10 supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to coenzyme Q10 supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

GALSULFASE – Special Authority see SA1988 below – Retail pharmacy
Inj 1 mg per ml, 5 ml vial.......2,234.00 1 ✓ Naglazyme

⇒SA1988 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has been diagnosed with mucopolysaccharidosis VI; and
- 2 Either:
 - 2.1 Diagnosis confirmed by demonstration of N-acetyl-galactosamine-4-sulfatase (arylsulfatase B) deficiency by either enzyme activity assay in leukocytes or skin fibroblasts; or
 - 2.2 Detection of two disease causing mutations and patient has a sibling who is known to have mucopolysaccharidosis VI

Renewal only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication

continued...

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
<u> </u>	Per	✓	Manufacturer

continued...

and/or adjustment of infusion rates; and

- 3 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by Enzyme Replacement Therapy (ERT); and
- 4 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT.

IDURSULFASE - Special Authority see SA1623 below - Retail pharmacy

⇒SA1623 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hunter Syndrome (mucopolysaccharidosis II); and
- 2 Fither:
 - 2.1 Diagnosis confirmed by demonstration of iduronate 2-sulfatase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts; or
 - 2.2 Detection of a disease causing mutation in the iduronate 2-sulfatase gene; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with idursulfase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT); and
- 5 Idursulfase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 weeks post-HSCT) at doses no greater than 0.5 mg/kg every week.

(Novitium Sugar Free S29 Oral lig 1 g per 10 ml to be delisted 1 February 2026)

1 ✓ Aldurazyme

⇒SA1695 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hurler Syndrome (mucopolysacchardosis I-H); and
- 2 Fither:
 - 2.1 Diagnosis confirmed by demonstration of alpha-L-iduronidase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts; or
 - 2.2 Detection of two disease causing mutations in the alpha-L-iduronidase gene and patient has a sibling who is known to have Hurler syndrome; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with laronidase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT); and
- 5 Laronidase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 post-HSCT) at doses no greater than 100 units/kg every week.

LEVO	DCARNITINE - Special Authority see SA2040 on the n	ext page – Retail pha	rmacy	
1	Гаb 500 mg	CBS	30	✓ Solgar
(Cap 250 mg	CBS	30	✓ Solgar
(Cap 500 mg	CBS	60	✓ Balance
			300	Metabolics
(Oral liq 1 g per 10 ml	CBS	118 ml	✓ Lacuna S29
				✓ Novitium Sugar
				Free S29
(Oral liq 500 mg per 10 ml	CBS	300 ml	✓ Balance

Subsidy (Manufacturer's Price)	Sub	Fully sidised	Brand or Generic
\$	Per	•	Manufacturer

⇒SA2040 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to carnitine supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to carnitine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

RIBOFLAVIN - Special Authority see SA2041 below - R	etail pharmacy		
Tab 100 mg	CBS	100	Country Life
y			✓ Puritan's Pride Vitamin B-2 100 mg \$29
Cap 100 mg	CBS	100	✓ Solgar

⇒SA2041 Special Authority for Subsidy

Initial application only from a metabolic physician or neurologist. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to riboflavin supplementation.

Renewal only from a metabolic physician or neurologist. Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to riboflavin supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

⇒SA1989 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 1 month for applications meeting the following criteria: All of the following:

- 1 Patient has phenylketonuria (PKU) and is pregnant or actively planning to become pregnant; and
- 2 Treatment with sapropterin is required to support management of PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

Renewal only from a metabolic physician or any relevant practitioner on the recommendation of a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Following the initial one-month approval, the patient has demonstrated an adequate response to a 2 to 4 week trial of sapropterin with a clinically appropriate reduction in phenylalanine levels to support management of PKU during pregnancy; or
 - 1.2 On subsequent renewal applications, the patient has previously demonstrated response to treatment with sapropterin and maintained adequate phenylalanine levels to support management of PKU during pregnancy; and
- 2 Any of the following:
 - 2.1 Patient continues to be pregnant and treatment with sapropterin will not continue after delivery; or
 - 2.2 Patient is actively planning a pregnancy and this is the first renewal for treatment with sapropterin; or
 - 2.3 Treatment with sapropterin is required for a second or subsequent pregnancy to support management of their PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and

continued...

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic	
 \$	Per		Manufacturer	

continued...

- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

SODIUM BENZOATE - Special Authority see SA1599 below - Retail pharmacy

Soln 100 mg per mlCBS 100 ml ✓ Amzoate ©29

⇒SA1599 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

SODIUM PHENYLBUTYRATE - Special Authority see SA1990 below - Retail pharmacy
Grans 483 mg per g......2,016.00 174 g OP

✓ Pheburane

⇒SA1990 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder involving a deficiency of carbamylphosphate synthetase, ornithine transcarbamylase or argininosuccinate synthetase.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

TAURINE - Special Authority see SA2043 below - Retail pharmacy

Cap 500 mg	CBS	50	✓ Solgar
Cap 1,000 mg	CBS	90	✓ Life Extension
Powder	CBS	300 g	✓ Life Extension

⇒SA2043 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected specific mitochondrial disorder that may respond taurine supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has confirmed diagnosis of a specific mitochondrial disorder which responds to taurine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

TRIENTINE – Special Authority see SA2324 below – Retail pharmacy
Cap 250 mg.......2,022.00 100 ✓ Trientine Waymade

⇒SA2324 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Wilson disease; and
- 2 Treatment with D-penicillamine has been trialled and discontinued because the person has experienced intolerable side effects or has not received sufficient benefit; and
- 3 Treatment with zinc has been trailled and discontinued because the person has experienced intolerable side effects or has not received sufficient benefit, or zinc is considered clinically inappropriate as the person has symptomatic liver disease and requires copper chelation.

Gaucher's Disease

(Ma	Subsidy nufacturer's Price)	F Subsid	ully	Brand or Generic
	\$	Per	1	Manufacturer

⇒SA2137 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has a diagnosis of symptomatic type 1 or type 3* Gaucher disease confirmed by the demonstration of specific deficiency of glucocerebrosidase in leukocytes or cultured skin fibroblasts, and genotypic analysis; and
- 2 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by enzyme replacement therapy (ERT) or the disease might be reasonably expected to compromise a response to ERT; and
- 3 Any of the following:
 - 3.1 Patient has haematological complications of Gaucher disease; or
 - 3.2 Patient has skeletal complications of Gaucher disease; or
 - 3.3 Patient has significant liver dysfunction or hepatomegaly attributable to Gaucher disease; or
 - 3.4 Patient has reduced vital capacity from clinically significant or progressive pulmonary disease due to Gaucher disease: or
 - 3.5 Patient is a child and has experienced growth failure with significant decrease in percentile linear growth over a 6-12 month period; and
- 4 Taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units).

Note: Indication marked with * is an unapproved indication

Renewal only from a metabolic physician or any relevant practitioner on the recommendation of a metabolic physician. Approvals valid for 3 years for applications meeting the following criteria:

All of the following:

- 1 Patient has demonstrated a symptomatic improvement and has maintained improvements in the main symptom or symptoms for which therapy was started; and
- 2 Patient has demonstrated a clinically objective improvement or no deterioration in haemoglobin levels, platelet counts and liver and spleen size; and
- 3 Radiological (MRI) signs of bone activity performed at two years since initiation of treatment, and five yearly thereafter, demonstrate no deterioration shown by the MRI, compared with MRI taken immediately prior to commencement of therapy or adjusted dose; and
- 4 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT; and
- 5 Patient is adherent with regular treatment and taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units).

Mouth and Throat

Agents Used in Mouth Ulceration

BENZYDAMINE HYDROCHLORIDE

Soln 0.15% - Higher subsidy of \$22.60 per 500 ml with

Additional subsidy by endorsement for a patient who has oral mucositis as a result of treatment for cancer, and the prescription is endorsed accordingly.

CARMELLOSE SODIUM WITH GELATIN AND PECTIN

Paste	17.20	56.7 g OP	✓ Stomahesive
	4.55	15 g OP	
	(7.90)	ŭ	Orabase
	1.52	5 g OP	
	(3.60)	•	Orabase
Powder	8.48 [′]	28 g OP	
	(10.95)	ŭ	Stomahesive

Difflam

	Subsidy (Manufacturer's P \$	rice) Subs Per	Fully Brand or didised Generic Manufacturer
RIAMCINOLONE ACETONIDE Paste 0.1%	5.49	5 g OP	✓ Kenalog in Orabase
Oropharyngeal Anti-infectives			
MPHOTERICIN B Lozenges 10 mg	5.86	20	✓ Fungilin
AICONAZOLE Oral gel 20 mg per g		40 g OP	✓ Decozol
IYSTATIN Oral liq 100,000 u per ml		24 ml OP	✓ Nilstat
Vitamins		241111 01	- Motat
Vitamin B			
HYDROXOCOBALAMIN ★ Inj 1 mg per ml, 1 ml ampoule – Up to 6 inj available on a	PSO3.95	3	✓ <u>Hydroxocobalamin</u> Panpharma
PYRIDOXINE HYDROCHLORIDE a) No more than 100 mg per dose b) Only on a prescription ★ Tab 25 mg — No patient co-payment payable	3.43	90	✓ Vitamin B6 25
₹ Tab 50 mg		500	✓ Pyridoxine multichem
HIAMINE HYDROCHLORIDE - Only on a prescription Tab 50 mg	4.65	100	✓ Thiamine multichem
/ITAMIN B COMPLEX ★ Tab, strong, BPC	11.25	500	✓ Bplex
Vitamin C			
ASCORBIC ACID a) No more than 100 mg per dose b) Only on a prescription			
★ Tab 100 mg Cvite to be Principal Supply on 1 March 2026	16.00	500	✓ Cvite
Vitamin D			
LFACALCIDOL K Cap 0.25 mcg K Cap 1 mcg K Oral drops 2 mcg per ml	87.98	100 100 20 ml OP	✓ One-Alpha ✓ One-Alpha ✓ One-Alpha
ALCITRIOL Cap 0.25 mcg		100	✓ Calcitriol XL S29
₭ Cap 0.5 mcg	13.68	100	✓ Calcitriol-AFT ✓ Calcitriol XL S29 ✓ Calcitriol-AFT

	Subsidy (Manufacturer's Pric	e) Subs Per	Fully sidised	Brand or Generic Manufacturer
COLECALCIFEROL * Cap 1.25 mg (50,000 iu) — Maximum of 12 cap per prescrip * Oral liq 188 mcg per ml (7,500 iu per ml)		12 5 ml OP		/it.D3 Clinicians
Multivitamin Preparations				
MULTIVITAMIN RENAL – Special Authority see SA1546 below * Cap		30	√ (Clinicians Renal Vit
▶ \$A1546 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valithe following criteria: Either:	id without further re	newal unless	notifie	d for applications meeting
The patient has chronic kidney disease and is receiving e The patient has chronic kidney disease grade 5, defined a 15 ml/min/1.73 m² body surface area (BSA).				
MULTIVITAMINS – Special Authority see SA1036 below – Reta * Powder		200 g OP	✓ F	Paediatric Seravit
■ SA1036 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals vali inborn errors of metabolism. Renewal from any relevant practitioner. Approvals valid without approval for multivitamins.				·
VITAMINS * Tab (BPC cap strength)		1,000	✓ N	/lvite
* Cap (fat soluble vitamins A, D, E, K) – Special Authority see SA1720 below – Retail pharmacy		60	✓ ∨	/itabdeck
■ SA1720 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valithe following criteria: Any of the following: 1 Patient has cystic fibrosis with pancreatic insufficiency; or 2 Patient is an infant or child with liver disease or short gut 3 Patient has severe malabsorption syndrome.	r	newal unless	notifie	d for applications meeting
Minerals				
Calcium				
CALCIUM CARBONATE * Tab 1.25 g (500 mg elemental) * Tab eff 1.25 g (500 mg elemental) – Subsidy by endorseme	7.28 ent260.00	250 100		Calci-Tab 500 Calcium 500 mg Hexal §29
Subsidy by endorsement – Only when prescribed for pa considered unsuitable. CALCIUM GLUCONATE	ediatric patients (<	5 years) whe	ere calc	

10

✓ Max Health - HameIn \$29

* Inj 10%, 10 ml ampoule......32.00

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price)	Su Per	Fully bsidised	Brand or Generic Manufacturer
lodine				
POTASSIUM IODATE * Tab 253 mcg (150 mcg elemental iodine)	5.99	90	✓ <u>N</u>	<u>euroTabs</u>
Iron				
FERROUS FUMARATE * Tab 200 mg (65.7 mg elemental)	3.49	100	✓ <u>F</u>	erro-tab
FERROUS FUMARATE WITH FOLIC ACID * Tab 310 mg (100 mg elemental) with folic acid 350 mcg	5.98	100	√ <u>F</u>	erro-F-Tabs
FERROUS SULFATE Tab long-acting 325 mg (105 mg elemental) Note: No new patients to be initiated on ferrous sulfate t		30	√ F	errograd
* Oral liq 30 mg (6 mg elemental) per 1 ml	10.25	250 ml 500 ml		erro-Liquid erodan
Ferro-Liquid to be Principal Supply on 1 February 2026 (Ferrograd Tab long-acting 325 mg (105 mg elemental) to be deli (Ferodan Oral liq 30 mg (6 mg elemental) per 1 ml to be delisted	,			
IRON (AS FERRIC CARBOXYMALTOSE) – Special Authority se Inj 50 mg per ml, 10 ml vial		Retail pha	•-	erinject

⇒SA2394 Special Authority for Subsidy

Initial application — (Anaemia) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with anaemia; and
- 2 Any of the following:
 - 2.1 Serum ferritin level is 20 mcg/L or less; or
 - 2.2 Both:
 - 2.2.1 Serum ferritin is between 20 and 50 mcg/L; and
 - 2.2.2 C-Reactive Protein (CRP) is at least 5 mg/L; or
 - 2.3 Patient has chronic inflammatory disease with symptoms of anaemia despite normal iron levels; and
- 3 Any of the following:
 - 3.1 Oral iron treatment has proven ineffective; or
 - 3.2 Oral iron treatment has resulted in dose-limiting intolerance; or
 - 3.3 Rapid correction of anaemia is required.

Renewal — (Anaemia) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient continues to have anaemia with a serum ferritin level of 20 mcg/L, or less or between 20 and 50 mcg/L with CRP of at least 5 mg/L, or has chronic inflammatory disease with symptoms of anaemia despite normal iron levels; and
- 2 A trial (or re-trial) with oral iron is clinically inappropriate.

Initial application — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

1 Patient has been diagnosed with iron-deficiency anaemia; and

continued...

(1)	Subsidy		Fully	Brand or
	(Manufacturer's Price)		ubsidised	Generic
	\$	Per	•	Manufacturer

continued...

2 Any of the following:

IRON POLYMALTOSE

- 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
- 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
- 2.3 Patient has symptomatic heart failure, chronic kidney disease stage 3 or more or active inflammatory bowel disease and a trial of oral iron is unlikely to be effective; or
- 2.4 Rapid correction of anaemia is required.

Renewal — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient continues to have iron-deficiency anaemia; and
- 2 A re-trial with oral iron is clinically inappropriate.

* Inj 50 mg per ml, 2 ml ampoule	5	✓ Ferrosig
Magnesium		
MAGNESIUM HYDROXIDE Suspension 8%33.60	355 ml	✓ Phillips Milk of Magnesia \$29
MAGNESIUM SULPHATE ★ Inj 2 mmol per ml, 5 ml ampoule 37.53 ★ Inj 2 mmol per ml, 10 ml ampoule 75.06	10 10	✓ <u>Martindale</u> ✓ Inresa \$29
Zinc		

ZINC SULPHATE		
* Cap 137.4 mg (50 mg elemental)11.00	100	✓ Zincaps
		✓ Zincaps S29 S29

BLOOD AND BLOOD FORMING ORGANS

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

Antianaemics

Hypoplastic and Haemolytic

⇒SA2539 Special Authority for Subsidy

Initial application — (chronic renal failure) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient in chronic renal failure: and
- 2 Haemoglobin is less than or equal to 100g/L; and
- 3 Any of the following:
 - 3.1 Both:
 - 3.1.1 Patient does not have diabetes mellitus; and
 - 3.1.2 Glomerular filtration rate is less than or equal to 30ml/min; or
 - 3.2 Both:
 - 3.2.1 Patient has diabetes mellitus; and
 - 3.2.2 Glomerular filtration rate is less than or equal to 45ml/min; or
 - 3.3 Patient is on haemodialysis or peritoneal dialysis.

Initial application — (myelodysplasia) from any specialist. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of myelodysplasia (MDS)*; and
- 2 Has had symptomatic anaemia with haemoglobin < 100g/L and is red cell transfusion-dependent; and
- 3 Patient has very low, low or intermediate risk MDS based on the WHO classification based prognostic scoring system for myelodysplastic syndrome (WPSS); and
- 4 Other causes of anaemia such as B12 and folate deficiency have been excluded; and
- 5 Patient has a serum epoetin level of < 500 IU/L; and
- 6 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Note: Indication marked with * is an unapproved indication

Renewal — (myelodysplasia) from any specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient's transfusion requirement continues to be reduced with erythropoietin treatment; and
- 2 Transformation to acute myeloid leukaemia has not occurred; and
- 3 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Note: Indication marked with * is an unapproved indication

EPOETIN ALFA - Special Authority see SA2539 above - Retail pharmacy

Wastage claimable			
Inj 1,000 iu in 0.5 ml, syringe	250.00	6	Binocrit
Inj 2,000 iu in 1 ml, syringe		6	✓ Binocrit
Inj 3,000 iu in 0.3 ml, syringe		6	✓ Binocrit
Inj 4,000 iu in 0.4 ml, syringe		6	✓ Binocrit
Inj 5,000 iu in 0.5 ml, syringe		6	✓ Binocrit
Inj 6,000 iu in 0.6 ml, syringe		6	✓ Binocrit
Inj 8,000 iu in 0.8 ml, syringe		6	✓ Binocrit
Inj 10,000 iu in 1 ml, syringe		6	✓ Binocrit
Inj 40,000 iu in 1 ml, syringe		1	✓ Binocrit

	Subsidy (Manufacturer's Price) \$	Sub Per	Fully osidised	Brand or Generic Manufacturer
Megaloblastic				
FOLIC ACID * Tab 0.8 mg	26.60	1,000	-	olic Acid multichem
* Tab 5 mg Oral liq 50 mcg per ml		100 5 ml OP	_	olic Acid Viatris iomed

Antifibrinolytics, Haemostatics and Local Sclerosants

EFTRENONACOG ALFA [RECOMBINANT FACTOR IX] - [Xpharm]

For patients with haemophilia B receiving prophylaxis treatment. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management group.

Treaters Group in conjunction with the National Haen	nophilia Management gro	up.	
Inj 250 iu vial	612.50	1	Alprolix
Inj 500 iu vial	1,225.00	1	✓ Alprolix
Inj 1,000 iu vial		1	✓ Alprolix
Inj 2,000 iu vial	4,900.00	1	✓ Alprolix
Inj 3,000 iu vial	7,350.00	1	✓ Alprolix
Inj 4,000 iu vial	·	1	✓ Alprolix
ELTROMBOPAG - Special Authority see SA1743 below	- Retail pharmacy		
Wastage claimable			
Tab 25 mg	1,550.00	28	✓ Revolade
Tab 50 mg	3.100.00	28	✓ Revolade

⇒SA1743 Special Authority for Subsidy

Initial application — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has had a splenectomy; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Any of the following:
 - 3.1 Patient has a platelet count of 20,000 to 30,000 platelets per microlitre and has evidence of significant mucocutaneous bleeding; or
 - 3.2 Patient has a platelet count of less than or equal to 20,000 platelets per microlitre and has evidence of active bleeding; or
 - 3.3 Patient has a platelet count of less than or equal to 10,000 platelets per microlitre.

Initial application — (idiopathic thrombocytopenic purpura - preparation for splenectomy) only from a haematologist. Approvals valid for 6 weeks where the patient requires eltrombopag treatment as preparation for splenectomy. Initial application — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist.

Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a significant and well-documented contraindication to splenectomy for clinical reasons; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Either:
 - 3.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per microliter: or
 - 3.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre

Subsidy	Fully Subsidised		Brand or
(Manufacturer's Price)			Generic
<u> </u>	Per		Manufacturer

continued...

and significant mucocutaneous bleeding.

Initial application — (severe aplastic anaemia) only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Two immunosuppressive therapies have been trialled and failed after therapy of at least 3 months duration; and
- 2 Either:
 - 2.1 Patient has severe aplastic anaemia with a platelet count of less than or equal to 20,000 platelets per microliter; or
 - 2.2 Patient has severe aplastic anaemia with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

Renewal — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 12 months where the patient has obtained a response (see Note) from treatment during the initial approval or subsequent renewal periods and further treatment is required.

Note: Response to treatment is defined as a platelet count of > 30,000 platelets per microlitre.

Renewal — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient's significant contraindication to splenectomy remains; and
- 2 The patient has obtained a response from treatment during the initial approval period; and
- 3 Patient has maintained a platelet count of at least 50,000 platelets per microlitre on treatment; and
- 4 Further treatment with eltrombopag is required to maintain response.

Renewal — (severe aplastic anaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has obtained a response from treatment of at least 20,000 platelets per microlitre above baseline during the initial approval period; and
- 2 Platelet transfusion independence for a minimum of 8 weeks during the initial approval period.

EMICIZUMAB - [Xpharm] - Special Authority see \$A2272 below

✓ Hemlibra	1	3,570.00	Inj 30 mg in 1 ml vial
✓ Hemlibra	1	7,138.00	Inj 60 mg in 0.4 ml vial
✓ Hemlibra	1	•	Inj 105 mg in 0.7 ml vial
✓ Hemlibra	1	17,846.00	Inj 150 mg in 1 ml vial

⇒SA2272 Special Authority for Subsidy

Initial application — (Severe Haemophilia A with or without FVIII inhibitors) only from a haematologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has severe congenital haemophilia A with a severe bleeding phenotype (endogenous factor VIII activity less than or equal to 2%); and
- 2 Emicizumab is to be administered at a dose of no greater than 3 mg/kg weekly for 4 weeks followed by the equivalent of 1.5 mg/kg weekly.

EPTACOG ALFA [RECOMBINANT FACTOR VIIA] - [Xpharm]

For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group. Rare Clinical Circumstances Brand of bypassing agent for > 14 days predicted use. Access to funded treatment for > 14 days predicted use is by named patient application to the Haemophilia Treaters Group, subject to access criteria.

Inj 1 mg syringe	1,178.30	1	✓ NovoSeven RT
Inj 2 mg syringe	2,356.60	1	✓ NovoSeven RT
Inj 5 mg syringe	·	1	✓ NovoSeven RT
Inj 8 mg syringe	·	1	✓ NovoSeven RT

			D 1 O11	
	Subsidy		Fully	Brand or
	(Manufacturer's Price)	Sul Per	bsidised •	Generic Manufacturer
FACTOR EIGHT INHIBITOR BYPASSING FRACTION - [Xphar	*			manufacturo.
For patients with haemophilia. Preferred Brand of bypassing		predicted	duse A	ccess to funded treatment
is managed by the Haemophilia Treaters Group in conjunction				
Inj 500 Ü		1		EIBA NF
Inj 1,000 U	2,630.00	1	√ F	EIBA NF
Inj 2,500 U	6,575.00	1	√ F	EIBA NF
MOROCTOCOG ALFA [RECOMBINANT FACTOR VIII] - [Xpha	rm]			
For patients with haemophilia. Rare Clinical Circumstances	Brand of short half-life			
treatment is managed by the Haemophilia Treaters Group in	conjunction with the	National	Haemop	hilia Management Group,
subject to criteria.				
Inj 250 iu prefilled syringe		1		(yntha
Inj 500 iu prefilled syringe		1		(yntha
Inj 1,000 iu prefilled syringe		1		(yntha
Inj 2,000 iu prefilled syringe		1		(yntha (yntha
Inj 3,000 iu prefilled syringe	·	ı	• ^	хупипа
NONACOG GAMMA, [RECOMBINANT FACTOR IX] - [Xpharm			- .	
For patients with haemophilia. Access to funded treatment i	s managed by the Ha	emophili	a Treate	rs Group in conjunction
with the National Haemophilia Management Group.	070.00	1		NVIIDIO
Inj 1,000 iu vial Inj 2,000 iu vial		1		RIXUBIS RIXUBIS
Inj 3,000 iu vial		1		RIXUBIS
• •	·	'	• 1	IIXODIO
OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (ADVATE) —			anna ta	fundad traatmant is
For patients with haemophilia. Preferred Brand of short half managed by the Haemophilia Treaters Group in conjunction				
Inj 500 iu vial		1		Advate
Inj 1,000 iu vial		1	_	Advate
Inj 2,000 iu vial		1		Advate
Inj 3,000 iu vial	,	1		Advate
OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (KOGENATE				
For patients with haemophilia. Rare Clinical Circumstances	Brand of short half-life	e recomb	oinant fac	ctor VIII. Access to funded
treatment is managed by the Haemophilia Treaters Group in	conjunction with the	National	Haemon	hilia Management Group
subject to criteria.	,			,
Inj 250 iu vial	237.50	1	✓ K	Cogenate FS
Inj 500 iu vial	475.00	1	✓ K	(ogenate FS
Inj 1,000 iu vial		1		(ogenate FS
Inj 2,000 iu vial	,	1		Cogenate FS
Inj 3,000 iu vial		1	✓ K	Cogenate FS
RURIOCTOCOG ALFA PEGOL [RECOMBINANT FACTOR VIII]				
For patients with haemophilia A receiving prophylaxis treatm		d treatme	ent is ma	anaged by the Haemophilia
Treaters Group in conjunction with the National Haemophilia	0 0 1			
Inj 1,000 iu vial		1		Adynovate
Inj 2,000 iu vial	2,400.00	1	✓ A	Adynovate
SODIUM TETRADECYL SULPHATE				
* Inj 3% 2 ml		5		
	(140.00)		F	Fibro-vein
TRANEXAMIC ACID				
Tab 500 mg	9.93	60	✓ N	Mercury Pharma

	Subsidy		Fully	Brand or
	(Manufacturer's Price)	S	Subsidised	Generic
	\$	Per	✓	Manufacturer
Vitamin K				
PHYTOMENADIONE				
Inj 2 mg per 0.2 ml - Up to 5 inj available on a PSO	8.00	5	✓ K	onakion MM Paediatric
Inj 10 mg per ml, 1 ml - Up to 5 inj available on a PSO	9.21	5	✓ K	Conakion MM
Antithrombotic Agents				
Antiplatelet Agents				
ASPIRIN				
* Tab 100 mg	12.65	990	✓ E	thics Aspirin EC
CLOPIDOGREL				
* Tab 75 mg	5.07	84	✓ A	rrow - Clopid
DIPYRIDAMOI F			_	<u> </u>
Note: No new patients to be initiated on dipyridamole.				
Tab long-acting 150 mg	13 93	60	✓ P	vtazen SR
Cap modified-release 200 mg		60		ipyridamole -
				Strides S29
(Pytazen SR Tab long-acting 150 mg to be delisted 1 January 2	1026)			Oli Idoo
(1 ytazen on rab long-acting 100 mg to be delisted 1 January 2	020)			

⇒SA2530 Special Authority for Subsidy

Initial application — (acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

56

✓ Ticagrelor Sandoz

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

TICAGRELOR - Special Authority see SA2530 below - Retail pharmacy

Renewal — (subsequent acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Initial application — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 Either:
 - 1.1 Patient has had a neurological stenting procedure* in the last 60 days; or
 - 1.2 Patient is about to have a neurological stenting procedure performed*; and
- 2 Either:
 - 2.1 Patient has demonstrated clopidogrel resistance using the P2Y12 (VerifyNow) assay or another appropriate platelet function assay and requires antiplatelet treatment with ticagrelor; or
 - - 2.2.1 Clopidogrel resistance has been demonstrated by the occurrence of a new cerebral ischemic event; or

	Subsidy	Fully	Brand or
(Manuf	acturer's Price)	Subsidised	Generic
	\$ Pe	er 🗸	Manufacturer

continued...

2.2.2 Clopidogrel resistance has been demonstrated by the occurrence of transient ischemic attack symptoms referable to the stent.

Renewal — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient is continuing to benefit from treatment; and
- 2 Treatment continues to be clinically appropriate.

Initial application — (Percutaneous coronary intervention with stent deployment) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic**.

Renewal — (Percutaneous coronary intervention with stent deployment) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic**.

Initial application — (Stent thrombosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has experienced cardiac stent thrombosis whilst on clopidogrel.

Initial application — (acute minor stroke or high-risk transient ischemic attack (TIA)*) from any relevant practitioner.

Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with a minor stroke (NIHSS† score 3 or less), high-risk TIA (ABCD2 score 4 or more) or Crescendo TIA: and
- 2 Either:
 - 2.1 Patient is expected to be a poor metaboliser of clopidogrel, with documented clinical rationale; or
 - 2.2 Patient is allergic to clopidogrel**; and
- 3 Ticagrelor to be prescribed for a maximum of 21 days following minor stroke or TIA.

Renewal — (subsequent minor stroke or TIA, or Crescendo TIA) from any relevant practitioner. Approvals valid for 1 month where patient has been diagnosed with a minor stroke (NIHSS score 3 or less) or high-risk transient ischemic attack (ABCD2 score 4 or more) or Crescendo TIA.

Notes: indications marked with * are unapproved indications.

Note:** Clopidogrel allergy is defined as a history of anaphylaxis, urticaria, generalised rash or asthma (in non-asthmatic patients) developing soon after clopidogrel is started and is considered unlikely to be caused by any other treatment.

Note: NIHSS† National Institutes of Health Stroke Scale.

Heparin and Antagonist Preparations

ENOXAPARIN SODIUM - Special Authority see SA2152	on the next page - Retail	pharmacy	
Inj 20 mg in 0.2 ml syringe	21.90	10	✓ Clexane
Inj 40 mg in 0.4 ml syringe	29.74	10	✓ Clexane
Inj 60 mg in 0.6 ml syringe		10	✓ Clexane
Inj 80 mg in 0.8 ml syringe		10	✓ Clexane
Inj 100 mg in 1 ml syringe		10	✓ Clexane
Inj 120 mg in 0.8 ml syringe		10	✓ Clexane Forte
Inj 150 mg in 1 ml syringe	100.70	10	✓ Clexane Forte

	,	ully Brand or	
(Manufact	turer's Price) Subsidi	sed Generic	
	\$ Per	 Manufac 	turer

⇒SA2152 Special Authority for Subsidy

Initial application — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patients pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Initial application — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 For the short-term treatment of venous thromboembolism prior to establishing a therapeutic level of oral anti-coagulant treatment; or
- 2 For the prophylaxis and treatment of venous thromboembolism in high risk surgery; or
- 3 To enable cessation/re-establishment of existing oral anticoagulant treatment pre/post surgery; or
- 4 For the prophylaxis and treatment of venous thromboembolism in Acute Coronary Syndrome surgical intervention; or
- 5 To be used in association with cardioversion of atrial fibrillation.

Initial application — (Short-term use during treatment of COVID-19 with nirmatrelvir with ritonavir) from any relevant practitioner. Approvals valid for 2 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is receiving an anticoagulation treatment that has drug/drug interactions with ritonavir that increases risk of bleeding; and
- 2 Patient meets the Access Criteria for COVID-19 antivirals published on the Pharmac website*; and
- 3 Other antiviral treatments for COVID-19 have been considered and are not clinically suitable options.

Renewal — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patient's pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Renewal — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month where low molecular weight heparin treatment or prophylaxis is required for a second or subsequent event (surgery, ACS, cardioversion, or prior to oral anti-coagulation).

127 44

25

✓ Pfizor S29

HEPARIN SODIUM

Ini 1 000 ju per ml 10 ml vial

111 1,000 ta per 1111, 10 1111 viai		20	· I IIZCI CES
Inj 1,000 iu per ml, 5 ml ampoule	25.49	10	✓ Wockhardt S29
	103.70		✓ Wockhardt PSF S29
	127.44	50	✓ Pfizer
Inj 10 iu per ml, 5 ml ampoule (flushing solution)	19.38	10	✓ Wockhardt S29
Inj 5,000 iu per ml, 5 ml vial	83.00	10	Heparin SodiumPanpharma
Inj 5,000 iu per ml, 1 ml	70.33	5	✓ Hospira
Inj 5,000 iu per ml, 5 ml ampoule		50	✓ Pfizer
Inj 25,000 iu per ml, 0.2 ml	25.78	5	✓ Hospira
(Pfizer S29 Inj 1,000 iu per ml, 10 ml vial to be delisted 1 January	uary 2026)		
(Wockhardt \$29 Inj 1,000 iu per ml, 5 ml ampoule to be delist	ed 1 January 2026)		
(Wockhardt PSF \$23) Inj 1,000 iu per ml, 5 ml ampoule to be (Heparin Sodium Panpharma Inj 5,000 iu per ml, 5 ml vial to be	,	,	
HEPARINISED SALINE			
Inj 10 iu per ml, 5 ml	96.91	50	✓ Pfizer

	Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
	\$	Per		Manufacturer
Oral Anticoagulants				
DABIGATRAN				
Cap 75 mg - No more than 2 cap per day	27.99	60	✓ F	Pradaxa
Cap 110 mg		60	✓ [Pradaxa
Cap 150 mg		60	✓ [Pradaxa
RIVAROXABAN				
Tab 10 mg - No more than 1 tab per day	15.60	30	✓)	Karelto
Tab 15 mg - Up to 14 tab available on a PSO		28	√)	Karelto
Tab 20 mg		28	√)	Karelto
WARFARIN SODIUM				
Note: Marevan and Coumadin are not interchangeable.				
* Tab 1 mg	3.46	50	✓ (Coumadin
ŭ	7.50	100	✓ I	Marevan
* Tab 2 mg	4.31	50	✓ (Coumadin
* Tab 3 mg	12.00	100	✓ [Marevan
* Tab 5 mg	5.93	50	✓ (Coumadin
	13.50	100	✓ I	Marevan
Blood Colony-stimulating Factors				
FILGRASTIM – Special Authority see SA1259 below – Retail phar	rmany			
Inj 300 mcg per 0.5 ml prefilled syringe	•	10	1	Nivestim
Inj 480 mcg per 0.5 ml prefilled syringe		10	-	Nivestim
mij 700 mog por 0.0 mi promed symige	100.72	10	• <u>i</u>	11VCSUIII

⇒SA1259 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 20%*); or
- 2 Peripheral blood stem cell mobilisation in patients undergoing haematological transplantation; or
- 3 Peripheral blood stem cell mobilisation or bone marrow donation from healthy donors for transplantation; or
- 4 Treatment of severe chronic neutropenia (ANC < 0.5 ×10⁹/L); or
- 5 Treatment of drug-induced prolonged neutropenia (ANC < 0.5 ×10⁹/L).

Note: *Febrile neutropenia risk greater than or equal to 20% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

PEGFILGRASTIM - Special Authority see SA1912 below - Retail pharmacy

✓ 7ievtenzo Inj 6 mg per 0.6 ml syringe69.50 Ziextenzo to be Principal Supply on 1 February 2026

⇒SA1912 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where used for prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 5%*). Note: *Febrile neutropenia risk greater than or equal to 5% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	Brand or Generic Manufacturer
Fluids and Electrolytes				
Intravenous Administration				
GLUCOSE [DEXTROSE] * Inj 50%, 10 ml ampoule – Up to 5 inj available on a PSO * Inj 50%, 90 ml bottle – Up to 5 inj available on a PSO		5 1	_	Biomed Biomed
POTASSIUM CHLORIDE * Inj 75 mg per ml, 10 ml	65.00	50	✓ L	uno umaCina fizer ^{S29}
SODIUM BICARBONATE Inj 8.4%, 50 ml	24.70	1	√ E	Biomed
b) Not in combination Inj 8.4%, 100 ml a) Up to 5 inj available on a PSO b) Not in combination	25.31	1	√ E	Biomed
SODIUM CHLORIDE Not funded for use as a nasal drop. Not funded for nebuliser for nebuliser use.	r use except when us	ed in (conjunction	with an antibiotic intended
Inj 23.4% (4 mmol/ml), 20 ml ampoule For Sodium chloride oral liquid formulation refer Standar		5 1	√ E	Biomed
Inj 0.9%, 5 ml ampoule – Up to 5 inj available on a PSO Fresenius Kabi to be Principal Supply on 1 February 202		20	√ F	resenius Kabi
Inj 0.9%, 10 ml ampoule – Up to 5 inj available on a PSO Fresenius Kabi to be Principal Supply on 1 February 202		50	√ F	resenius Kabi
Inj 0.9%, 20 ml ampouleFresenius Kabi to be Principal Supply on 1 February 202		20	√ F	resenius Kabi
Inj 0.9%, 1,000 ml bag — Up to 2 bag available on a PSO Only if prescribed on a prescription for renal dialysis, ma		1 are in	_	Saxter of the patient, or on a PSO
for emergency use. (500 ml and 1,000 ml packs) Inj 0.9%, 500 ml bag — Up to 4 bag available on a PSO Only if prescribed on a prescription for renal dialysis, ma for emergency use. (500 ml and 1,000 ml packs)		1 are in	_	Baxter of the patient, or on a PSO
TOTAL PARENTERAL NUTRITION (TPN)	CRS	1 OP	√ T	'DN
WATER 1) On a prescription or Practitioner's Supply Order only will Schedule requiring a solvent or diluent; or 2) On a bulk supply order; or	hen on the same form		-	
3) When used in the extemporaneous compounding of ey4) When used for the dilution of sodium chloride soln 7%		ents o	nly.	
Inj 10 ml ampoule - Up to 5 inj available on a PSO		50		resenius Kabi Iultichem
Inj 20 ml ampoule - Up to 5 inj available on a PSO	5.00	20	√ F	resenius Kabi

	Subsidy (Manufacturer's Price \$	e) Subs Per	Fully sidised	Brand or Generic Manufacturer
Oral Administration				
CALCIUM POLYSTYRENE SULPHONATE Powder	169.85	300 g OP	✓ 0	alcium Resonium
Powder for oral soln — Up to 5 sach available on a PSO	9.50	50	√ <u>E</u>	lectral
COMPOUND ELECTROLYTES WITH GLUCOSE [DEXTROSE] Soln with electrolytes		1 OP	✓ H	lydralyte - Lemonade
PHOSPHORUS Tab eff 500 mg (16 mmol)	82.50	100	√ P	hosphate Phebra
POTASSIUM CHLORIDE * Tab eff 548 mg (14 m eq) with chloride 285 mg (8 m eq)	5.26 (17.10)	60	C	chlorvescent
* Tab long-acting 600 mg (8 mmol)	, ,	200	-	pan-K
SODIUM BICARBONATE Cap 840 mg	8.52	100	-	odibic
SODIUM POLYSTYRENE SULPHONATE Powder	84.65	454 g OP	✓ R	lesonium-A

	Subsidy		Fully	Brand or
	(Manufacturer's Price	١ ،	Subsidised	
	\$	Per	Jubsidised •	Manufacturer
	Ψ	1 61		Manuacturei
Aluba Advanasantas Disalassa				
Alpha-Adrenoceptor Blockers				
Alpha Adrenoceptor Blockers				
DOVAZOCINI				
DOXAZOSIN ************************************	47.05	500	,	Daniel Ollowsk
* Tab 2 mg		500		Doxazosin Clinect
* Tab 4 mg	20.94	500	/	Doxazosin Clinect
PHENOXYBENZAMINE HYDROCHLORIDE				
* Cap 10 mg	65.00	30	1	BNM S29
,	05.00	30	•	DINIVI 323
PRAZOSIN				
* Tab 1 mg	5.53	100	✓	Arrotex-Prazosin
•				S29 S29
	0.00			
ate. T. L. O.	9.98	465		Minipress S29
* Tab 2 mg	7.00	100	•	Arrotex-Prazosin
				S29 S29
	13.29		1	Minipress S29
* Tab 5 mg		100		Arrotex-Prazosin
* Tab 5 Hig	11.70	100	•	
				S29 S29
	22.00		✓	Minipress S29
* Cap 1 mg	15 40	100	1	Prazosin Mylan S29
, ,				•
* Cap 2 mg		100		Prazosin Mylan S29
* Cap 5 mg	23.32	100	/	Prazosin Mylan S29
Agents Affecting the Renin-Angiotensin System				
Agents Affecting the nemin-Anglotensin System				
ACE labibitare				
ACE Inhibitors				
CAPTOPRIL				
* Oral liq 5 mg per ml	96.00 1	00 ml O	D 1	DP-Captopril
. •,	00.00	00 1111 0	Γ Ψ	DF-Captoprii
Oral liquid restricted to children under 12 years of age.				
ENALAPRIL MALEATE				
* Tab 5 mg	1.75	90	/	Acetec
* Tab 10 mg		90		Acetec
* Tab 20 mg		90		Acetec
•	2.33	90	•	ACEIEC
LISINOPRIL				
* Tab 5 mg	12.00	90	✓	Teva Lisinopril
Teva Lisinopril to be Principal Supply on 1 March 2026				•
* Tab 10 mg	12.00	90	1	Teva Lisinopril
	12.00	30	•	reva Lisinopini
Teva Lisinopril to be Principal Supply on 1 March 2026	40.00		,	
* Tab 20 mg	16.00	90	•	Teva Lisinopril
Teva Lisinopril to be Principal Supply on 1 March 2026				
PERINDOPRIL				
* Tab 2 mg	1 70	30	J	Coversyl
•				Coveravi
* Tab 4 mg		30		Coversyl
* Tab 8 mg	3.94	30	•	Coversyl
QUINAPRIL				
* Tab 5 mg	10 24	90	1	Arrow-Quinapril 5
* Tab 10 mg		90	_	Arrow-Quinapril 10
本 Tab 20 mg	14.00	90		Arrow-Quinapril 10

* Tab 20 mg14.83

✓ Arrow-Quinapril 20

90

•	Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
	\$	Per	1	Manufacturer
AMIPRIL				
Cap 1.25 mg	17.25	90	✓ -	Γryzan
Cap 2.5 mg	16.50	90	✓.	Γryzan
- Cap 5 mg	16.88	90	✓.	Γryzan
Cap 10 mg	17.63	90	/	Гryzan
Angiotensin II Antagonists				
ANDESARTAN CILEXETIL				
Tab 4 mg	2.68	90	✓ (Candestar
: Tab 8 mg	2.67	90	✓ (Candestar
: Tab 16 mg	4.22	90	✓ (Candestar
Tab 32 mg	5.24	90	1	Candestar
OSARTAN POTASSIUM				
: Tab 12.5 mg	2.00	84	✓ I	osartan Actavis
Tab 25 mg		84	√ i	Losartan Actavis
- Tab 50 mg	2.86	84	√ i	osartan Actavis
Tab 100 mg		84	✓ [Losartan Actavis
Angiotensin II Antagonists with Diuretics				
ANDESARTAN CILEXETIL WITH HYDROCHLOROTHIA	AZIDE			
Tab 16 mg with hydrochlorothiazide 12.5 mg	4.10	30	✓	APO-Candesartan HCTZ 16/12.5
Tab 32 mg with hydrochlorothiazide 12.5 mg	5.25	30	•	APO-Candesartan HCTZ 32/12.5
OSARTAN POTASSIUM WITH HYDROCHLOROTHIAZII	DE			
Tab 50 mg with hydrochlorothiazide 12.5 mg	4.00	30	✓ /	Arrow-Losartan & Hydrochlorothiazide

Angiotensin II Antagonists with Neprilysin Inhibitors

SACUBITRIL WITH VALSARTAN - Special Authority see SA			
Tab 24.3 mg with valsartan 25.7 mg	190.00	56	✓ Entresto 24/26
Tab 48.6 mg with valsartan 51.4 mg	190.00	56	✓ Entresto 49/51
Tab 97.2 mg with valsartan 102.8 mg	190.00	56	✓ Entresto 97/103

⇒SA2302 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has heart failure: and
- 2 Any of the following:
 - 2.1 Patient is in NYHA/WHO functional class II; or
 - 2.2 Patient is in NYHA/WHO functional class III; or
 - 2.3 Patient is in NYHA/WHO functional class IV; and
- 3 Either:
 - 3.1 Patient has a documented left ventricular ejection fraction (LVEF) of less than or equal to 35%; or
 - 3.2 An ECHO is not reasonably practical, and in the opinion of the treating practitioner the patient would benefit from treatment; and
- 4 Patient is receiving concomitant optimal standard chronic heart failure treatments.

	Subsidy		Fully Brand or
	(Manufacturer's Price)	Per	osidised Generic Manufacturer
	Ψ	1 01	Wandiactarci
Antiarrhythmics			
For lignocaine hydrochloride refer to NERVOUS SYSTEM, Anae	esthetics, Local, page	121	
AMIODARONE HYDROCHLORIDE			
▲ Tab 100 mg	4.95	30	✓ Aratac
Aratac to be Principal Supply on 1 February 2026			
▲ Tab 200 mg	5.86	30	✓ Aratac
Aratac to be Principal Supply on 1 February 2026			
Inj 50 mg per ml, 3 ml ampoule – Up to 10 inj available on a PSO		10	✓ Max Health
Max Health to be Principal Supply on 1 February 2026	17.90	10	▼ Iviax ricalui
ATROPINE SULPHATE			
* Inj 600 mcg per ml, 1 ml ampoule – Up to 5 inj available on	а		
PSO		10	✓ Hikma S29
			✓ <u>Martindale</u>
DIGOXIN			
* Tab 62.5 mcg - Up to 30 tab available on a PSO	8.58	240	Lanoxin PG
Lanoxin PG to be Principal Supply on 1 February 2026	40.75	0.40	4 Lancardo
* Tab 250 mcg - Up to 30 tab available on a PSO Lanoxin to be Principal Supply on 1 February 2026	18./5	240	✓ Lanoxin
* Oral lig 50 mcg per ml	16.60	60 ml	✓ Lanoxin
			✓ Lanoxin S29 S29
DISOPYRAMIDE PHOSPHATE			
▲ Cap 100 mg	55.90	84	Rythmodan -
			Cheplafarm S29
			Rythmodan
			Neon S29
FLECAINIDE ACETATE			
Tab 50 mg		60 90	 ✓ <u>Flecainide BNM</u> ✓ Flecainide
▲ Cap long-acting 100 mg	33.70	90	Controlled
			Release Teva
▲ Cap long-acting 200 mg	54.28	90	✓ Flecainide
			Controlled
1:40	400 70	_	Release Teva
Inj 10 mg per ml, 15 ml ampoule	102.79 108.16	5	✓ Almarytm S29✓ Tambocor
	100.10		✓ Tambocor
			German S29
MEXILETINE HYDROCHLORIDE			
▲ Cap 150 mg	162.00	100	✓ Teva S29
▲ Cap 250 mg		100	✓ Teva S29
PROPAFENONE HYDROCHLORIDE			
▲ Tab 150 mg	40.90	50	✓ Rytmonorm

Subsidy

Fully

Brand or

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	/	Manufacturer

Antihypotensives

		see SA1474 below – Retail pharmacy	MIDODRINE - Special Authority see S
✓ <u>Midodrine</u>	100	36.68	Tab 2.5 mg
Medsurge ✓ Midodrine	100	58.88	Tah 5 mg
Medsurge	100		Tab 5 mg

⇒SA1474 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years where patient has disabling orthostatic hypotension not due to drugs.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Beta-Adrenoceptor Blockers

Beta Adrenoceptor Blockers

ATENOLOL			
* Tab 50 mg	11.00	500	✓ Viatris
* Tab 100 mg		500	✓ Atenolol Viatris
* Oral liq 25 mg per 5 ml		300 ml OP	✓ Atenolol AFT
Restricted to children under 12 years of age.	49.00	300 1111 01	A ALCHOIDI AI I
, ,			
BISOPROLOL FUMARATE	1.00	00	/ Inco Dicensolal
* Tab 2.5 mg		90	✓ <u>Ipca-Bisoprolol</u>
* Tab 5 mg		90	✓ <u>Ipca-Bisoprolol</u>
* Tab 10 mg	2./1	90	✓ <u>Ipca-Bisoprolol</u>
CARVEDILOL			
* Tab 6.25 mg		60	Carvedilol Sandoz
* Tab 12.5 mg	2.30	60	Carvedilol Sandoz
* Tab 25 mg	2.95	60	 Carvedilol Sandoz
LABETALOL			
* Tab 100 mg	14.50	100	✓ Trandate
	49.54		✓ Biocon S29
* Tab 200 mg	27.00	100	✓ Trandate
·	42.07		✓ Presolol S29
* Inj 5 mg per ml, 20 ml ampoule	59.06	5	
, 31	(88.60)		Trandate
METOPROLOL SUCCINATE			
* Tab long-acting 23.75 mg	4.20	90	✓ Myloc CR
* Tab long-acting 47.5 mg		90	✓ Myloc CR
* Tab long-acting 95 mg		90	✓ Myloc CR
* Tab long-acting 190 mg		90	✓ Myloc CR
METOPROLOL TARTRATE			
* Tab 50 mg	5.66	100	✓ IPCA-Metoprolol
* Tab 100 mg		60	✓ IPCA-Metoprolol
* Tab long-acting 200 mg		28	✓ Slow-Lopresor
* Inj 1 mg per ml, 5 ml vial		5	✓ Metoprolol IV Mylan
., ,		J	✓ Metoprolol IV Viatris

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price \$	e) Per	Fully Subsidised	Brand or Generic Manufacturer
NADOLOL				
* Tab 40 mg	19.19	100	✓	Nadolol BNM
* Tab 80 mg	30.39	100	✓	Nadolol BNM
PROPRANOLOL				
* Tab 10 mg	7.04	100	✓	Drofate
* Tab 40 mg		100	✓]	IPCA-Propranolol
* Cap long-acting 160 mg		100	1	Cardinol LA
* Oral liq 4 mg per ml - Special Authority see SA1327 below -	-			
Retail pharmacy	CBS	500 n	nl 🗸 l	Hikma-
				Propranolol S29
			✓	Roxane- Propranolol \$29

⇒SA1327 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

SOTAL OL

*	Tab 80 mg	22.50	300	✓ Sotalol Viatris S29
	v	40.00	500	✓ Mylan
	Mylan to be Principal Supply on 1 February 2026			•
*	Tab 160 mg	20.00	100	✓ Mylan
	Mylan to be Principal Supply on 1 February 2026			•

(Sotalol Viatris §29 Tab 80 mg to be delisted 1 March 2026)

Calcium Channel Blockers

Dihydropyridine Calcium Channel Blockers

AMI	LODIPINE			
*	Tab 2.5 mg	1.45	90	✓ Vasorex
*	Tab 5 mg	1.21	90	✓ Vasorex
*	Tab 10 mg	1.31	90	✓ Vasorex
FEL	ODIPINE			
*	Tab long-acting 2.5 mg	2.18	30	✓ Plendil ER
*	Tab long-acting 5 mg	6.57	90	✓ Felo 5 ER
	Tab long-acting 10 mg		90	✓ Felo 10 ER

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
FEDIPINE				
Tab long-acting 10 mg - Subsidy by endorsement	19.42	56	✓	Tensipine MR10 S29
Subsidised for patients who were taking nifedipine tab endorsed accordingly. Pharmacists may annotate the dispensing of nifedipine tab long-acting 10 mg.	prescription as endors	ed wh	nere there	exists a record of prior
Tab long-acting 20 mg		100		Nyefax Retard
Tab long-acting 30 mg	4.78	14	•	Mylan Italy (24 hr release) \$29
	34.10	100	/	Mylan (24 hr release) \$29
Tab long-acting 60 mg	52.81	100	•	Mylan (24 hr release) S29
Other Calcium Channel Blockers				<u> </u>
LTIAZEM HYDROCHLORIDE				
Cap long-acting 120 mg	65.35	500	1	Diltiazem CD Clinect
Cap long-acting 180 mg	7.00	30	1	Cardizem CD
Cap long-acting 240 mg	9.30	30	•	Cardizem CD
ERHEXILINE MALEATE				
Tab 100 mg	62.90	100	1	Pexsig
RAPAMIL HYDROCHLORIDE				
Tab 40 mg		100	_	Isoptin
Tab 80 mg		100	_	Isoptin
Tab long-acting 120 mg	36.02	100	_	Isoptin Retard S29
Tab long-acting 240 mg	15 10	30		Isoptin SR Isoptin SR
		50	•	isopiiii Sii
Inj 2.5 mg per ml, 2 ml ampoule – Up to 5 inj available on PSO		5	✓	Isoptin
Centrally-Acting Agents				
ONIDINE				
Patch 2.5 mg, 100 mcg per day – Only on a prescription		4		Mylan
Patch 5 mg, 200 mcg per day — Only on a prescription		4 4	_	Mylan Mylan
Patch 7.5 mg, 300 mcg per day — Only on a prescription	17.90	4	•	<u>Mylan</u>
ONIDINE HYDROCHLORIDE Tab 25 mcg	29.74	112	/	Clonidine Teva
Clonidine Teva to be Principal Supply on 1 February 2			_	Olollianio 101a
Tab 150 mcg		100	1	Catapres
Inj 150 mcg per ml, 1 ml ampoule	14.10	5	•	Catapres
ETHYLDOPA				
Tab 250 mg	15.10	100	✓	Methyldopa Viatris
Diuretics				
oop Diuretics				
JMETANIDE	46.55			
Tab 1 mg Inj 500 mcg per ml, 4 ml vial		100 5		Burinex Burinex

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

(Subsidy Manufacturer's Pri	ce) Subsi	Fully dised	Brand or Generic
,	\$	Per	1	Manufacturer
UROSEMIDE [FRUSEMIDE]				
Tab 40 mg - Up to 30 tab available on a PSO		1,000		IPCA-Frusemide
€ Tab 500 mg		50		Urex Forte
Gral liq 10 mg per ml		30 ml OP		Lasix
Inj 10 mg per ml, 25 ml ampoule		6		Lasix
Inj 10 mg per ml, 2 ml ampoule - Up to 5 inj available on a PS	SO2.40	5	/	Furosemide-Baxter
Potassium Sparing Diuretics				
MILORIDE HYDROCHLORIDE				
Tab 5 mg	81.07	100	1	Padagis S29
-	171.41	28		Wockhardt \$29
Oral liq 1 mg per ml		25 ml OP		Biomed
PLERENONE – Special Authority see SA1728 below – Retail ph		-		
Tab 25 mg		30	/	Inspra
Tab 50 mg		30		Inspra
SA1728 Special Authority for Subsidy	20.00	00	•	<u>шоріа</u>
e following criteria: th: 1 Patient has heart failure with ejection fraction less than 40%	s; and			
Either: 2.1 Patient is intolerant to optimal dosing of spironolacto 2.2 Patient has experienced a clinically significant adver		n optimal dosi	ng of	f spironolactone.
PIRONOLACTONE	4.00	400	,	Only atte
Tab 25 mgSpiractin to be Principal Supply on 1 March 2026	4.20	100	•	Spiractin
Spiractiff to be Principal Supply on 1 March 2026 ← Tab 100 mg	11.40	100	./	Cniroctin
Spiractin to be Principal Supply on 1 March 2026	11.40	100	•	Spiractin
Oral liq 5 mg per ml	35.70	25 ml OP	/	Biomed
Potassium Sparing Combination Diuretics				
MILORIDE HYDROCHLORIDE WITH FUROSEMIDE				
	0.00	28	1	Frumil
Tab 5 mg with furosemide 40 mg	8.63			
· ·		20		
MILORIDE HYDROCHLORIDE WITH HYDROCHLOROTHIAZID	E		,	Moduratio
MILORIDE HYDROCHLORIDE WITH HYDROCHLOROTHIAZID Tab 5 mg with hydrochlorothiazide 50 mg	E	50	1	Moduretic
MILORIDE HYDROCHLORIDE WITH HYDROCHLOROTHIAZID Tab 5 mg with hydrochlorothiazide 50 mg	E		/	Moduretic
MILORIDE HYDROCHLORIDE WITH HYDROCHLOROTHIAZID Tab 5 mg with hydrochlorothiazide 50 mg Thiazide and Related Diuretics ENDROFLUMETHIAZIDE [BENDROFLUAZIDE]	E 5.00	50		
MILORIDE HYDROCHLORIDE WITH HYDROCHLOROTHIAZID Tab 5 mg with hydrochlorothiazide 50 mg Thiazide and Related Diuretics ENDROFLUMETHIAZIDE [BENDROFLUAZIDE]	E 5.00			Moduretic Arrow- Bendrofluazide
MILORIDE HYDROCHLORIDE WITH HYDROCHLOROTHIAZID Tab 5 mg with hydrochlorothiazide 50 mg Thiazide and Related Diuretics ENDROFLUMETHIAZIDE [BENDROFLUAZIDE] Tab 2.5 mg - Up to 150 tab available on a PSO	E5.00	50		Arrow-
MILORIDE HYDROCHLORIDE WITH HYDROCHLOROTHIAZID Tab 5 mg with hydrochlorothiazide 50 mg Thiazide and Related Diuretics ENDROFLUMETHIAZIDE [BENDROFLUAZIDE] Tab 2.5 mg – Up to 150 tab available on a PSO	E5.00	50	✓	Arrow- Bendrofluazide
MILORIDE HYDROCHLORIDE WITH HYDROCHLOROTHIAZID Tab 5 mg with hydrochlorothiazide 50 mg Thiazide and Related Diuretics ENDROFLUMETHIAZIDE [BENDROFLUAZIDE] Tab 2.5 mg – Up to 150 tab available on a PSO	E5.00	50	✓	Arrow- Bendrofluazide Arrow-
MILORIDE HYDROCHLORIDE WITH HYDROCHLOROTHIAZID Tab 5 mg with hydrochlorothiazide 50 mg Thiazide and Related Diuretics ENDROFLUMETHIAZIDE [BENDROFLUAZIDE] Tab 2.5 mg – Up to 150 tab available on a PSO	E5.00	50	✓	Arrow- Bendrofluazide
• • • • • • • • • • • • • • • • • • • •	E5.00	50	✓	Arrow- Bendrofluazide Arrow-

Oral liq 50 mg per ml30.67

25 ml OP

✓ Biomed

	Subsidy (Manufacturer's Price) \$	Sub Per	Fully sidised	Brand or Generic Manufacturer
CHLORTALIDONE [CHLORTHALIDONE] * Tab 25 mg Hygroton to be Principal Supply on 1 February 2026	6.95	50	✓ H	ygroton
INDAPAMIDE * Tab 2.5 mg METOLAZONE	16.00	90	✓ <u>D</u>	apa-Tabs
Tab 5 mg	CBS	50	✓ Z	aroxolyn §29
Vasopressin receptor antagonists				
TOLVAPTAN – Special Authority see SA2166 below – Retail ph. Tab 15 mg Tab 30 mg Tab 45 mg + 15 mg Tab 60 mg + 30 mg Tab 90 mg + 30 mg	873.50 873.50 1,747.00 1,747.00	28 OP 28 OP 56 OP 56 OP 56 OP	✓ Ji ✓ Ji ✓ Ji	inarc inarc inarc inarc inarc

⇒SA2166 Special Authority for Subsidy

Initial application — (autosomal dominant polycystic kidney disease) only from a renal physician or any relevant practitioner on the recommendation of a renal physician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of autosomal dominant polycystic kidney disease; and
- 2 Patient has an estimated glomerular filtration rate (eGFR) of greater than or equal to 25 ml/min/1.73 m² at treatment initiation; and
- 3 Either:
 - 3.1 Patient's disease is rapidly progressing, with a decline in eGFR of greater than or equal to 5 mL/min/1.73 m² within one-year; or
 - 3.2 Patient's disease is rapidly progressing, with an average decline in eGFR of greater than or equal to 2.5 mL/min/1.73 m² per year over a five-year period.

Renewal — (autosomal dominant polycystic kidney disease) only from a renal physician or any relevant practitioner on the recommendation of a renal physician. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has not developed end-stage renal disease, defined as an eGFR of less than 15 mL/min/1.73 m²; and
- 2 Patient has not undergone a kidney transplant.

Lipid-Modifying Agents		
Fibrates		
BEZAFIBRATE * Tab 200 mg	90 30	 ✓ Bezalip ✓ Bezalip Retard
Other Lipid-Modifying Agents		
ACIPIMOX	30	✓ Olbetam

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
Resins				
COLESTYRAMINE Powder for oral suspension 4 g sachet	61.50	50	√ 0	Colestyramine -

ATORVASTATIN		
* Tab 10 mg	30	✓ Lorstat
5.16	500	✓ Lorstat
* Tab 20 mg8.12	500	✓ Lorstat
* Tab 40 mg	500	✓ Lorstat
* Tab 80 mg25.39	500	✓ Lorstat
PRAVASTATIN		
* Tab 20 mg7.16	100	✓ Clinect
* Tab 40 mg	100	✓ Clinect
ROSUVASTATIN – Special Authority see SA2093 below – Retail pharmacy		
* Tab 5 mg	30	Rosuvastatin Viatris
* Tab 10 mg	30	✓ Rosuvastatin Viatris
* Tab 20 mg2.71	30	✓ Rosuvastatin Viatris
4.21		✓ Rosuvastatin-
		Sandoz
* Tab 40 mg4.55	30	✓ Rosuvastatin Viatris

⇒SA2093 Special Authority for Subsidy

Initial application — (cardiovascular disease risk) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient is considered to be at risk of cardiovascular disease; and
 - 1.2 Patient is Māori or any Pacific ethnicity; or
- 2 Both:
 - 2.1 Patient has a calculated risk of cardiovascular disease of at least 15% over 5 years; and
 - 2.2 LDL cholesterol has not reduced to less than 1.8 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initial application — (familial hypercholesterolemia) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has familial hypercholesterolemia (defined as a Dutch Lipid Criteria score greater than or equal to 6); and
- 2 LDL cholesterol has not reduced to less than 1.8 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initial application — (established cardiovascular disease) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

1 Any of the following:

✓ fully subsidised

Principal Supply

continued...

Mylan ©29

✓ Quantalan sugar
free ©29

		CARDIOV	'ASC	ULAR SYSTEM
	Subsidy (Manufacturer's Pri \$	ice) Subsi	Fully dised	Brand or Generic Manufacturer
continued				
1.1 Patient has proven coronary artery disease (CAL1.2 Patient has proven peripheral artery disease (PA				
1.3 Patient has experienced an ischaemic stroke; an				
2 LDL cholesterol has not reduced to less than 1.4 mmol/l and/or simvastatin.	itre with treatment v	vith the maxim	um tole	erated dose of atorvastat
nitial application — (recurrent major cardiovascular events) from any relevan	t practitioner.	Appro	vals valid without further
enewal unless notified for applications meeting the following cr	iteria:			
Both:				
1 Patient has experienced a recurrent major cardiovascula			arction	n, ischaemic stroke,
coronary revascularisation, hospitalisation for unstable a 2 LDL cholesterol has not reduced to less than 1.0 mmol/l			um tol	arated does of atomisetat
and/or simvastatin.	ille willi liealillelil v	viui uie iiiaxiiii	uiii toit	eraleu uose or alorvasia
SIMVASTATIN				
₹ Tab 10 mg	1.68	90	✓ S	imvastatin Mylan
				imvastatin Viatris
★ Tab 20 mg		90	_	imvastatin Viatris
* Tab 40 mg		90		imvastatin Viatris
* Tab 80 mg	8.81	90	✓ <u>S</u>	imvastatin Viatris
Selective Cholesterol Absorption Inhibitors				
ZETIMIBE				
米 Tab 10 mg	1.76	30	✓ E	zetimibe Sandoz
ZETIMIBE WITH SIMVASTATIN				
Tab 10 mg with simvastatin 10 mg		30		imybe
Tab 10 mg with simvastatin 20 mg		30		imybe
Tab 10 mg with simvastatin 40 mg Tab 10 mg with simvastatin 80 mg		30 30		imybe imybe
Tab To Tilg Will Sillivastatill 60 Hig	0.15	30	• 2	illiybe
Nitrates				
GLYCERYL TRINITRATE				
★ Oral pump spray, 400 mcg per dose – Up to 250 dose				
available on a PSO	7.48	250 dose OP	✓ N	litrolingual Pump Spray
* Patch 25 mg, 5 mg per day	15.73	30	✓ N	litroderm TTS
⊁ Patch 50 mg, 10 mg per day		30		litroderm TTS
J, - J,,				-

100

30

90

5

5

10

49.00

✓ Ismo 20 ✓ Ismo 40 Retard

✓ Duride

✓ Hospira

✓ Aspen Adrenaline✓ DBL Adrenaline

✓ Aspen Adrenaline

Tab long-acting 60 mg......13.50

Inj 1 in 1,000, 1 ml ampoule - Up to 5 inj available on a PSO4.98

Inj 1 in 10,000, 10 ml ampoule - Up to 5 inj available on a PSO27.00

ISOSORBIDE MONONITRATE

Sympathomimetics

ADRENALINE

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidised		Generic
 \$	Per	•	Manufacturer

Vasodilators

HYDRALAZINE HYDROCHLORIDE

*	Tab 25 mg - Special Authority see SA1321 below - Retail			
	pharmacy	CBS	1	 Hydralazine
			56	✓ Onelink S29
			84	✓ AMDIPHARM \$29
			100	✓ Camber S29
*	Ini 20 ma ampoule	25.00	5	✓ Anresoline

⇒SA1321 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 For the treatment of refractory hypertension; or
- 2 For the treatment of heart failure in combination with a nitrate, in patients who are intolerant or have not responded to ACE inhibitors and/or angiotensin receptor blockers.

MINOXIDIL

▲ Tab 10 mg	47.04	60	Minoxidil Roma S29
-	78.40	100	✓ Loniten
NICORANDIL			
▲ Tab 10 mg	27.81	60	✓ Max Health
Max Health to be Principal Supply on 1 February 2026			
▲ Tab 20 mg	35.12	60	Max Health
Max Health to be Principal Supply on 1 February 2026			
PAPAVERINE HYDROCHLORIDE			
* Inj 12 mg per ml, 10 ml ampoule	257.12	5	✓ Hospira
PENTOXIFYLLINE [OXPENTIFYLLINE]			
Tab 400 mg	44.37	50	✓ Trental 400

Endothelin Receptor Antagonists

AMBRISENTAN - Special Authority see SA2486 below - Retai	l pharmacy			
Tab 5 mg	200.00	30	1	Ambrisentan Viatris
Tab 10 mg	200.00	30	1	Ambrisentan Viatris

⇒SA2486 Special Authority for Subsidy

Initial application — (PAH monotherapy) only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and

Subsidy (Manufacturer's Price)	Fully Subsidised	
 \$	Per 🗸	Manufacturer

continued...

- 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
- 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁵); and
- 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH 2022 (see note below for link to these guidelines) †; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
- 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
 - 5.1 Ambrisentan is to be used as PAH monotherapy; and
 - 5.2 Any of the following:
 - 5.2.1 Patient has experienced intolerable side effects with both sildenafil and bosentan; or
 - 5.2.2 Patient has an absolute contraindication to sildenafil and an absolute or relative contraindication to bosentan (e.g. due to current use of a combined oral contraceptive or liver disease); or
 - 5.2.3 Patient is a child with idiopathic PAH or PAH secondary to congenital heart disease.

Initial application — **(PAH dual therapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁻⁵); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH 2022 (see note below for link to these guidelines) †; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**: or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
 - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
 - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
` \$	Por 🗸	Manufacturer

continued...

- 5.1 Ambrisentan is to be used as PAH dual therapy; and
- 5.2 Any of the following:
 - 5.2.1 Patient has tried bosentan (either as PAH monotherapy, or PAH dual therapy with sildenafil) for at least three months and has not experienced an acceptable response to treatment according to a validated risk stratification tool**; or
 - 5.2.2 Patient has experienced intolerable side effects on bosentan; or
 - 5.2.3 Patient has an absolute or relative contraindication to bosentan (e.g. due to current use of a combined oral contraceptive or liver disease); or
 - 5.2.4 Patient is presenting in NYHA/WHO functional class III or IV, and would benefit from initial dual therapy in the opinion of the treating clinician and has an absolute or relative contraindication to bosentan (eg. due to current liver disease or use of a combined oral contraceptive).

Initial application — **(PAH triple therapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁵); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH 2022 (see note below for link to these guidelines) †; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
 - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
 - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
 - 5.1 Ambrisentan is to be used as PAH triple therapy; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is on the lung transplant list; or
 - 5.2.2 Both:
 - 5.2.2.1 Patient is presenting in NYHA/WHO functional class IV; and
 - 5.2.2.2 Patient has an absolute or relative contraindication to bosentan (e.g. due to current use of a combined oral contraceptive or liver disease); or
 - 5.2.3 Both:
 - 5.2.3.1 Patient has tried PAH dual therapy for at least three months and remains in an unacceptable risk category according to a validated risk stratification tool**; and

Subsidy		Fully	Brand or	
(Manufacturer's Pri	ce)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

5.2.3.2 Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario.

Renewal only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 2 years where the patient is continuing to derive benefit from ambrisentan treatment according to a validated PAH risk stratification tool**.

Notes: † The European Respiratory Journal Guidelines can be found here: 2022 ECS/ERS Guidelines for the diagnosis and treatment of pulmonary hypertension PAH

** the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

 BOSENTAN - Special Authority see SA2254 below - Retail pharmacy
 100.00
 60
 ✓ Bosentan Dr Reddy's

 Tab 125 mg
 100.00
 60
 ✓ Bosentan Dr Reddy's

⇒SA2254 Special Authority for Subsidy

Initial application — **(PAH monotherapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁻⁵); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these quidelines) †: or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**: or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
 - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or
 - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
 - 5.1 Bosentan is to be used as PAH monotherapy; and
 - 5.2 Any of the following:
 - 5.2.1 Patient has experienced intolerable side effects on sildenafil; or
 - 5.2.2 Patient has an absolute contraindication to sildenafil; or
 - 5.2.3 Patient is a child with idiopathic PAH or PAH secondary to congenital heart disease.

-			
	Subsidy	Fully	Brand or
	(Manufacturer's Price)	Subsidised	Generic
	` ¢ ′	Por 🗸	Manufacturor

continued...

Initial application — **(PAH dual therapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*: and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁵); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) †; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
 - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or
 - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Bosentan is to be used as part of PAH dual therapy; and
- 6 Either:
 - 6.1 Patient has tried a PAH monotherapy (sildenafil) for at least three months and has experienced an inadequate therapeutic response to treatment according to a validated risk stratification tool**; or
 - 6.2 Patient is presenting in NYHA/WHO functional class III or IV, and in the opinion of the treating clinician would likely benefit from initial dual therapy.

Initial application — **(PAH triple therapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁵); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or

Subsidy		Fully	Brand or
(Manufacturer's Price) \$	Per	Subsidised 🗸	Generic Manufacturer

continued...

- nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) †; or
- 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
- 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
- 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease: or
- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
 - 5.1 Bosentan is to be used as part of PAH triple therapy; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is on the lung transplant list; or
 - 5.2.2 Patient is presenting in NYHA/WHO functional class IV; or
 - 5.2.3 Both:
 - 5.2.3.1 Patient has tried PAH dual therapy for at least three months and has not experienced an acceptable response to treatment according to a validated risk stratification tool**: and
 - 5.2.3.2 Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario.

Renewal only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 2 years where patient is continuing to derive benefit from bosentan treatment according to a validated PAH risk stratification tool**.

Notes: † The European Respiratory Journal Guidelines can be found here: 2022 ECS/ERS Guidelines for the diagnosis and treatment of pulmonary hypertension PAH

** the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

Phosphodiesterase Type 5 Inhibitors

SILDENAFIL – Special Authority see SA2255 below – Retail pl	harmacy		
Tab 25 mg	0.72	4	✓ Vedafil
Tab 50 mg		4	✓ Vedafil
Tab 100 mg		12	✓ Vedafil

⇒SA2255 Special Authority for Subsidy

Initial application — (Raynaud's Phenomenon*) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has Raynaud's Phenomenon*; and
- 2 Patient has severe digital ischaemia (defined as severe pain requiring hospital admission or with a high likelihood of digital ulceration; digital ulcers; or gangrene); and
- 3 Patient is following lifestyle management (avoidance of cold exposure, sufficient protection, smoking cessation support, avoidance of sympathomimetic drugs); and
- 4 Patient is being treated with calcium channel blockers and nitrates (or these are contraindicated/not tolerated).

Initial application — (Pulmonary arterial hypertension*) only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH is confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) of greater than 20 mmHg; and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) that is less than or equal to 15 mmHg; and
 - 4.1.4 Pulmonary vascular resistance (PVR) of at least 2 Wood Units or greater than 160 International Units (dyn s cm⁵); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH is non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) †; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**: or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
 - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease: or
 - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures.

Initial application — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has a documented history of traumatic or non-traumatic spinal cord injury; and
- 2 Patient has erectile dysfunction secondary to spinal cord injury requiring pharmacological treatment.

Renewal — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Notes: Note: Indications marked with * are Unapproved Indications.

† The European Respiratory Journal Guidelines can be found here: 2022 ECS/ERS Guidelines for the diagnosis and treatment of pulmonary hypertension PAH

** the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

Prostacyclin Analogues

EPOPROSTENOL – Special Authority see SA2256 below –	Retail pharmacy		
Inj 500 mcg vial	36.61	1	✓ Veletri
Inj 1.5 mg vial	73.21	1	✓ Veletri

⇒SA2256 Special Authority for Subsidy

Initial application — **(PAH dual therapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class III or IV; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ıbsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁵); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) †; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
 - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
 - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
 - 5 All of the following:
 - 5.1 Epoprostenol is to be used as part of PAH dual therapy with either sildenafil or an endothelin receptor antagonist; and
 - 5.2 Patient is presenting in NYHA/WHO functional class IV; and
 - 5.3 Patient has tried a PAH monotherapy for at least three months and remains in an unacceptable risk category according to a validated risk stratification tool.

Initial application — **(PAH triple therapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class III or IV: and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁻⁵); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these quidelines) †; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
 - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsic	lised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
 - 5.1 Epoprostenol is to be used as PAH triple therapy; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is on the lung transplant list; or
 - 5.2.2 Patient is presenting in NYHA/WHO functional class IV; or
 - 5.2.3 Both:
 - 5.2.3.1 Patient has tried PAH dual therapy for at least three months and has not experienced an acceptable response to treatment according to a validated risk stratification tool; and
 - 5.2.3.2 Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario.

Renewal only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 2 years where patient is continuing to derive benefit from epoprostenol treatment according to a validated PAH risk stratification tool**.

Notes: † The European Respiratory Journal Guidelines can be found here: 2022 ECS/ERS Guidelines for the diagnosis and treatment of pulmonary hypertension PAH

** the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

ILOPROST - Special Authority see SA2257 below - Retail pharmacy

30 ✓ Vebulis

⇒SA2257 Special Authority for Subsidy

Initial application — **(PAH monotherapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁵); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) †; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
 - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
 - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major

Subsidy		Fully	Brand or	
(Manufacturer's Pr	ice)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and

- 5 Both:
 - 5.1 Iloprost is to be used as PAH monotherapy; and
 - 5.2 Fither:
 - 5.2.1 Patient has experienced intolerable side effects on sildenafil and both the funded endothelin receptor antagonists (i.e. both bosentan and ambrisentan); or
 - 5.2.2 Patient has an absolute contraindication to sildenafil and an absolute or relative contraindication to endothelin receptor antagonists.

Initial application — **(PAH dual therapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁵); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these quidelines) †: or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**: or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
 - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
 - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 All of the following:
 - 5.1 Iloprost is to be used as PAH dual therapy with either sildenafil or an endothelin receptor antagonist; and
 - 5.2 Either:
 - 5.2.1 Patient has an absolute contraindication to or has experienced intolerable side effects on sildenafil: or
 - 5.2.2 Patient has an absolute or relative contraindication to or experienced intolerable side effects with a funded endothelin receptor antagonist; and
 - 5.3 Fither:
 - 5.3.1 Patient has tried a PAH monotherapy for at least three months and remains in an unacceptable risk category according to a validated risk stratification tool**; or
 - 5.3.2 Patient is presenting in NYHA/WHO functional class III or IV, and in the opinion of the treating clinician would benefit from initial dual therapy.

Initial application — **(PAH triple therapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

|--|

continued...

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁻⁵); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these quidelines) †: or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
 - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
 - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
 - 5.1 Iloprost is to be used as PAH triple therapy; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is on the lung transplant list; or
 - 5.2.2 Patient is presenting in NYHA/WHO functional class IV; or
 - 5.2.3 Both:
 - 5.2.3.1 Patient has tried PAH dual therapy for at least three months and has not experienced an acceptable response to treatment according to a validated risk stratification tool**; and
 - 5.2.3.2 Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario.

Renewal only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 2 years where patient is continuing to derive benefit from iloprost treatment according to a validated PAH risk stratification tool**.

Notes: † The European Respiratory Journal Guidelines can be found here: 2022 ECS/ERS Guidelines for the diagnosis and treatment of pulmonary hypertension PAH

** the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

Subsidy Fully Brand or
(Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

Antiacne Preparations

For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 92

ADAPALENE

- a) Maximum of 30 g per prescription
- b) Only on a prescription

22.89	30 g OP	Differin
pharmacy		
11.26	60	Oratane
18.75	120	✓ Oratane
26.73	120	✓ Oratane
	11.26 18.75	pharmacy 6011.26 6018.75 120

⇒SA2449 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, paediatrician, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around isotretinoin and is competent to prescribe isotretinoin; and
- 3 Any of the following:
 - 3.1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and that they must not become pregnant during treatment and for a period of one month after the completion of treatment; or
 - 3.2 Patient is not of child bearing potential: or
 - 3.3 Patient is a child and it is considered not appropriate to exclude pregnancy or start contraceptives or undertake pregnancy-related isotretinoin counselling.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and that they must not become pregnant during treatment and for a period of one month after the completion of treatment: or
- 2 Patient is not of child bearing potential; or
- 3 Patient is a child and it is considered not appropriate to exclude pregnancy or start contraceptives or undertake pregnancy-related isotretinoin counselling.

TRETINOIN

Antibacterials Topical

For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 92

HYDROGEN PEROXIDE	
* Crm 1%	
Crystadorm to be Principal Supply on 1 January 2006	

MUPIROCIN

15 g OP

Bactroban

Crystaderm

- a) Only on a prescription
- b) Not in combination

DERMATOLOGICALS

	Subsidy		Fully B	rand or	_
	(Manufacturer's F		sidised G	ieneric	
CODILIM FLICIDATE IFLICIDIO ACIDA	\$	Per	✓ N	lanufacturer	_
SODIUM FUSIDATE [FUSIDIC ACID] Crm 2%	1.69	5 g OP	✓ Fob	an	
a) Maximum of 5 g per prescription		- 	. 55		
b) Only on a prescription					
c) Not in combination Oint 2%	1 60	5 g OP	✓ Fob	an .	
a) Maximum of 5 g per prescription	1.09	3 y Oi	• <u>1 0 0 0</u>	<u>all</u>	
b) Only on a prescription					
c) Not in combination					
SULFADIAZINE SILVER	40.00	50 . 05	<i>(</i> -:		
Crm 1%	10.80	50 g OP	✓ Flan	nazine	
a) Up to 250 g available on a PSOb) Not in combination					
,					
Antifungals Topical					
For systemic antifungals, refer to INFECTIONS, Antifungals	s, page 99				
AMOROLFINE	•				
a) Only on a prescription					
b) Not in combination					
Nail soln 5%	21.87	5 ml OP	✓ Myc	<u>oNail</u>	
CLOTRIMAZOLE * Crm 1%	1 10	20 g OP	✓ Clor	nazol	
a) Only on a prescription		20 g Oi	• 0101	iiuzoi	
b) Not in combination					
* Soln 1%		20 ml OP	0		
a) Only on a prescription	(11.58)		Can	esten	
b) Not in combination					
ECONAZOLE NITRATE					
Crm 1%	8.04	20 g OP	✓ Peva	aryl	
a) Only on a prescription					
b) Not in combination Foaming soln 1%, 10 ml sachets	9.89	3			
. Saming Sont 176, 10 th Sachets	(18.64)	3	Peva	aryl	
a) Only on a prescription	, ,				
b) Not in combination					
MICONAZOLE NITRATE	0.00	15 ~ OD		iaham	
* Crm 2%	0.90	15 g OP	✓ <u>Mult</u>	ichem	
b) Not in combination					
* Lotn 2%		30 ml OP	_		
a) Only on a managing time	(10.03)		Dakt	arin	
a) Only on a prescription b) Not in combination					
* Tinct 2%	4.36	30 ml OP			
	(12.10)		Dakt	arin	
a) Only on a prescription					
b) Not in combination					

✓ <u>healthE Calamine</u>

Aqueous

✓ Itch-Soothe

✓ MidWest

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per ✓	Manufacturer

100 g

20 g OP

100 g

36.80

Antipruritic Preparations

CALAN	ЛIN	١E
-------	-----	----

- a) Only on a prescription
- b) Not in combination

CROTAMITON

- a) Only on a prescription
- b) Not in combination

MENTHOL – Only in combination

1) Only in combination with a dermatological base or proprietary Topical Corticosteriod – Plain

2) With or without other dermatological galenicals.

Crystals......12.60

25 g ✓ MidWest

Corticosteroids Topical

For systemic corticosteroids, refer to CORTICOSTEROIDS AND RELATED AGENTS, page 83

Corticosteroids - Plain

BETAMETHASONE DIPROPIONATE			
Crm 0.05%	2.96	15 g OP	✓ Diprosone
	36.00	50 g OP	✓ Diprosone
Oint 0.05%	2.96	15 g OP	✓ Diprosone
	36.00	50 g OP	✓ Diprosone
Oint 0.05% in propylene glycol base	4.33	30 g OP	✓ Diprosone OV
BETAMETHASONE VALERATE			
* Crm 0.1%	5.85	50 g OP	✓ Beta Cream
* Oint 0.1%		50 g OP	✓ Beta Ointment
* Lotn 0.1%		50 ml OP	✓ Betnovate
CLOBETASOL PROPIONATE			
* Crm 0.05%	2.75	30 g OP	✓ Dermol
Dermol to be Principal Supply on 1 February 2026		30 g OF	Definion
* Oint 0.05%	2.60	30 g OP	✓ Dermol
Dermol to be Principal Supply on 1 February 2026	3.00	30 g OF	Definion
CLOBETASONE BUTYRATE			
Crm 0.05%		30 g OP	
	(10.00)		Eumovate
HYDROCORTISONE			
* Crm 1% - Only on a prescription	1.78	30 g OP	✓ Ethics
, , ,	20.40	500 g	✓ Noumed
Noumed to be Principal Supply on 1 February 2026		•	
* Powder – Only in combination	49.95	25 g	✓ ABM
Up to 5% in a dermatological base (not proprietary Togalenicals	oical Corticosterio	d – Plain) with o	or without other dermatological

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's I	Price) Subs	Fully sidised	
IYDROCORTISONE AND PARAFFIN LIQUID AND LANOLIN				
Lotn 1% with paraffin liquid 15.9% and lanolin 0.6% - Only o	n			
a prescription	12.83	250 ml	1	DP Lotn HC
IYDROCORTISONE BUTYRATE				
Lipocream 0.1%		100 g OP		Locoid Lipocream
Oint 0.1%		100 g OP		Locoid
Milky emul 0.1%	12.33	100 ml OP		Locoid Crelo
METHYLPREDNISOLONE ACEPONATE				
Crm 0.1%	4.95	15 g OP	1	<u>Advantan</u>
Oint 0.1%	4.95	15 g OP	1	<u>Advantan</u>
MOMETASONE FUROATE				
Crm 0.1%	2.25	15 g OP	1	Elocon Alcohol Free
	3.50	50 g OP	1	Elocon Alcohol Free
Oint 0.1%	2.25	15 g OP		Elocon
	3.50	50 g OP	1	Elocon
Lotn 0.1%	4.99	30 ml OP	1	Elocon
RIAMCINOLONE ACETONIDE				
Crm 0.02%	6.49	100 g OP	1	Aristocort
Oint 0.02%	6.54	100 g OP	1	Aristocort
Crm 0.1% with sodium fusidate (fusidic acid) 2%	SIDIC ACID] 3.49 (10.45)	15 g OP		Fucicort
Crm 0.1% with sodium fusidate (fusidic acid) 2% a) Maximum of 15 g per prescription b) Only on a prescription	3.49 (10.45)	15 g OP		Fucicort
Crm 0.1% with sodium fusidate (fusidic acid) 2%	3.49 (10.45)	ŭ	_	
Crm 0.1% with sodium fusidate (fusidic acid) 2%	3.49 (10.45) tion 2.85	15 g OP	,	Fucicort Micreme H
Crm 0.1% with sodium fusidate (fusidic acid) 2%	3.49 (10.45) tion 2.85 nly on a prescri	15 g OP		Micreme H
Crm 0.1% with sodium fusidate (fusidic acid) 2%	3.49 (10.45) tion 2.85 nly on a prescri 4.34	15 g OP ption 15 g OP		
Crm 0.1% with sodium fusidate (fusidic acid) 2%	3.49 (10.45) tion 2.85 nly on a prescri 4.34 N AND NYSTA	15 g OP ption 15 g OP		Micreme H
A) Maximum of 15 g per prescription b) Only on a prescription IYDROCORTISONE WITH MICONAZOLE – Only on a prescription CORTISONE WITH MICONAZOLE – Only on a prescription IYDROCORTISONE WITH NATAMYCIN AND NEOMYCIN – On Oint 1% with natamycin 1% and neomycin sulphate 0.5% RIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYCII Crm 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 mg	3.49 (10.45) tion 2.85 nly on a prescri 4.34 N AND NYSTA	15 g OP ption 15 g OP TIN		Micreme H
Crm 0.1% with sodium fusidate (fusidic acid) 2%	3.49 (10.45) tion 2.85 nly on a prescri 4.34 N AND NYSTA	15 g OP ption 15 g OP		Micreme H Pimafucort
A) Maximum of 15 g per prescription b) Only on a prescription IYDROCORTISONE WITH MICONAZOLE – Only on a prescription CORTISONE WITH MICONAZOLE – Only on a prescription IYDROCORTISONE WITH NATAMYCIN AND NEOMYCIN – On Oint 1% with natamycin 1% and neomycin sulphate 0.5% RIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYCII Crm 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 mg	3.49 (10.45) tion 2.85 nly on a prescri 4.34 N AND NYSTA	15 g OP ption 15 g OP TIN		Micreme H
A) Maximum of 15 g per prescription b) Only on a prescription IYDROCORTISONE WITH MICONAZOLE – Only on a prescription CORTISONE WITH MICONAZOLE – Only on a prescription IYDROCORTISONE WITH NATAMYCIN AND NEOMYCIN – On Oint 1% with natamycin 1% and neomycin sulphate 0.5% RIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYCII Crm 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 mg	3.49 (10.45) tion 2.85 nly on a prescri 4.34 N AND NYSTA	15 g OP ption 15 g OP TIN		Micreme H Pimafucort
a) Maximum of 15 g per prescription b) Only on a prescription lYDROCORTISONE WITH MICONAZOLE − Only on a prescription c Crm 1% with miconazole nitrate 2%	3.49 (10.45) tion 2.85 nly on a prescri 4.34 N AND NYSTA	15 g OP ption 15 g OP TIN		Micreme H Pimafucort
a) Maximum of 15 g per prescription b) Only on a prescription lYDROCORTISONE WITH MICONAZOLE — Only on a prescription c Crm 1% with miconazole nitrate 2%	3.49 (10.45) tion 2.85 nly on a prescri 4.34 N AND NYSTA	15 g OP ption 15 g OP TIN		Micreme H Pimafucort
a) Maximum of 15 g per prescription b) Only on a prescription IYDROCORTISONE WITH MICONAZOLE – Only on a prescription Corm 1% with miconazole nitrate 2%	3.49 (10.45) tion 2.85 nly on a prescri 4.34 N AND NYSTA' (9.28)	15 g OP ption 15 g OP TIN	•	Micreme H Pimafucort Viaderm KC
a) Maximum of 15 g per prescription b) Only on a prescription lYDROCORTISONE WITH MICONAZOLE — Only on a prescription c Crm 1% with miconazole nitrate 2%	3.49 (10.45) tion2.85 nly on a prescri4.34 N AND NYSTA	15 g OP ption 15 g OP TIN 15 g OP	•	Micreme H Pimafucort Viaderm KC
A) Maximum of 15 g per prescription b) Only on a prescription lYDROCORTISONE WITH MICONAZOLE — Only on a prescription c Crm 1% with miconazole nitrate 2%	3.49 (10.45) tion2.85 nly on a prescri4.34 N AND NYSTA	15 g OP ption 15 g OP TIN 15 g OP	•	Micreme H Pimafucort Viaderm KC healthE Dimethicone 5%
A) Maximum of 15 g per prescription b) Only on a prescription lYDROCORTISONE WITH MICONAZOLE — Only on a prescription c Crm 1% with miconazole nitrate 2%	3.49 (10.45) tion2.85 nly on a prescri4.34 N AND NYSTA	15 g OP ption 15 g OP TIN 15 g OP	•	Micreme H Pimafucort Viaderm KC healthE Dimethicone 5% healthE
a) Maximum of 15 g per prescription b) Only on a prescription lYDROCORTISONE WITH MICONAZOLE — Only on a prescription c Crm 1% with miconazole nitrate 2%		15 g OP ption 15 g OP TIN 15 g OP	<i>y y y</i>	Micreme H Pimafucort Viaderm KC healthE Dimethicone 5% healthE

	Subsidy		Fully	Brand or
	(Manufacturer's	Price) Sub	sidised	Generic
	\$	Per	√	Manufacturer
	Ψ	1 01		Manadataro
Emollients				
Linoments				
AQUEOUS CREAM				
* Crm	1 65	500 g	1	Evara
	1.00	300 g	•	Lvara
CETOMACROGOL				
* Crm BP	2.29	500 g	/	Cetomacrogol-AFT
CETOMACROGOL WITH GLYCEROL				
Crm 90% with glycerol 10%	1 02	460 g OP	1	Evara
Citil 50 % with gryceror 10 %		•	_	
	3.25	920 g OP	•	<u>Evara</u>
EMULSIFYING OINTMENT				
* Oint BP	3.13	500 g	1	Evara Emulsifying
T. OIR DI		000 g	-	Ointment
				Omunent
OIL IN WATER EMULSION				
* Crm	2.10	500 g	1	Fatty Emulsion
		ŭ		Cream (Evara)
DADACCINI				
PARAFFIN			_	
Oint liquid paraffin 50% with white soft paraffin 50%	4.94	500 g OP		White Soft Liquid
				Paraffin AFT
UREA				
* Crm 10%	1 27	100 a OB	./	healthE Urea Cream
	1.37	100 g OP	•	nealthe orea Cream
WOOL FAT WITH MINERAL OIL - Only on a prescription				
* Lotn hydrous 3% with mineral oil	5.60	1,000 ml		
, ,	(14.96)	,		DP Lotion
	(20.53)			Alpha-Keri Lotion
	, ,	050 ml OD		Alpha-Reff Lotion
	1.40	250 ml OP		DD 1 .:
	(5.87)			DP Lotion
	5.60	1,000 ml		
	(23.91)			BK Lotion
	1.40	250 ml OP		
	(7.73)			BK Lotion
	(1.13)			DIX LOUIDII
Other Dermetalogical Pages				
Other Dermatological Bases				
PARAFFIN				
	4.74	450 ~		EVADA White Cott
White soft - Only in combination	4./4	450 g	•	EVARA White Soft
				<u>Paraffin</u>
	19.00	2,500 g	1	EVARA White Soft
		, 0		Paraffin
Only in combination with a dermatological galenical or	an a diluant for a	nronrioton, To	niaal C	
Only in combination with a defination gical galefical of	as a unuent for a	proprietary ro	picai C	orticosteroid – Flairi.
Miner Olde Infections				
Minor Skin Infections				
DOMBONE IODINE				
POVIDONE IODINE			_	
Oint 10%	7.40	65 g OP		Betadine
a) Maximum of 130 g per prescription				
b) Only on a prescription				
Antiseptic Solution 10%	4.00	100 ml	1	Riodine
Antiseptic soln 10%		15 ml		Riodine
	6.99	500 ml	/	Riodine
Skin preparation, povidone iodine 10% with 30% alcohol	1.63	100 ml		
	(3.48)			Betadine Skin Prep
	(00)			

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once



Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

Parasiticidal Preparations

DIMETHICONE

IVERMECTIN - Special Authority see SA2511 below - Retail pharmacy

Tab 3 mg − Up to 100 tab available on a PSO......17.20 4 Stromectol

- PSO for institutional use only. Must be endorsed with the name of the institution for which the PSO is required and a valid Special Authority for patient of that institution.
- 2) Ivermectin available on BSO provided the BSO includes a valid Special Authority for a patient of the institution.
- For the purposes of subsidy of ivermectin, institution means age related residential care facilities, disability care facilities or prisons.

⇒SA2511 Special Authority for Subsidy

Initial application — (Scables) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Either:

- 1 The person has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
- 2 Both:
 - 2.1 The person has a confirmed diagnosis of scabies or is a close contact of a scabies case; and
 - 2.2 Fither:
 - 2.2.1 The person is unable to complete topical therapy; or
 - 2.2.2 Previous treatment with topical therapy has been tried and not cleared the infestation.

Initial application — (Other parasitic infections) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 filariasis; or
- 2 cutaneous larva migrans (creeping eruption); or
- 3 strongyloidiasis; or
- 4 The individual has a travel or residence history that requires presumptive parasite treatment.

Renewal — (Scables) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria: Either:

- 1 The person has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
- 2 Both:
 - 2.1 The person has a confirmed diagnosis of scabies or is a close contact of a scabies case; and
 - 2.2 Either:
 - 2.2.1 The person is unable to complete topical therapy; or
 - 2.2.2 Previous treatment with topical therapy has been tried and not cleared the infestation.

Renewal — (Other parasitic infections) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 filariasis: or
- 2 cutaneous larva migrans (creeping eruption); or
- 3 strongyloidiasis.

PERMETHRIN

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

Psoriasis and Eczema Preparations

ACITRETIN - Special Authority see SA2024 below - Retail ph	narmacy		
Cap 10 mg	26.20	60	Novatretin
Cap 25 mg	57.37	60	✓ Novatretin

⇒SA2024 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around acitretin and is competent to prescribe acitretin; and
- 3 Either:
 - 3.1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and that they must not become pregnant during treatment and for a period of three years after the completion of treatment; or
 - 3.2 Patient is not of child bearing potential.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and that they must not become pregnant during treatment and for a period of three years after the completion of treatment; or
- 2 Patient is not of child bearing potential.

BETAMETHASONE DIPROPIONATE WITH CALCIPOTRIOL
Foam spray 500 mcg with calcinotriol 50 mcg per g

Foam spray 500 mcg with calcipotriol 50 mcg per g	59.95	60 g OP	Enstilar
Gel 500 mcg with calcipotriol 50 mcg per g	40.92	60 g OP	Daivobet
Oint 500 mcg with calcipotriol 50 mcg per g		30 g OP	✓ Daivobet
CALCIPOTRIOL			
Oint 50 mcg per g	40.00	120 g OP	Daivonex
COAL TAR			
Soln BP - Only in combination	46.00	200 ml	✓ Midwest
1) I In to 10% only in combination with a dermatologic	ical hace or propri	atany Tonical C	orticostariod - Pla

- 1) Up to 10% only in combination with a dermatological base or proprietary Topical Corticosteriod Plain
- 2) With or without other dermatological galenicals.

COAL TAR WITH ALLANTOIN, MENTHOL, PHENOL AND SULPHUR

3011 3 % With Sulphur 0.3 %, mention 0.73 %, priend 0.3 % and			
allantoin crm 2.5%	6.59	75 g OP	
	(8.00)	-	Egopsoryl TA
	3.43	30 g OP	•
	(4.35)	-	Egopsoryl TA
COAL TAR WITH SALICYLIC ACID AND SULPHUR			
Soln 12% with salicylic acid 2% and sulphur 4% oint	7.95	40 g OP	✓ Coco-Scalp

PIMECROLIMUS – Special Authority see SA1970 on the next page – Retail pharmacy

- a) Maximum of 15 g per prescription
- b) Note: a maximum of 15 g per prescription and no more than one prescription per 12 weeks

b) Trotor a maximum or rog por procemption and no more than a	,,,o b. 000br	po=oc	,
Cream 1%	33.00	15 g OP	Elidel



Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer

⇒SA1970 Special Authority for Subsidy

Initial application only from a dermatologist, paediatrician, ophthalmologist or any relevant practitioner on the recommendation of a dermatologist, paediatrician or ophthalmologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has atopic dermatitis on the eyelid; and
- 2 Patient has at least one of the following contraindications to topical corticosteroids: periorificial dermatitis, rosacea, documented epidermal atrophy, documented allergy to topical corticosteroids, cataracts, glaucoma, or raised intraocular pressure.

PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORESCEIN - Only on a prescription

* Soln 2.3% with trolamine laurilsulfate and fluorescein sodium...........5.41 500 ml Pinetarsol SALICYLIC ACID

250 g ✓ Midwest Powder – Only in combination......29.00

1) Only in combination with a dermatological base or proprietary Topical Corticosteroid - Plain or collodion flexible

2) With or without other dermatological galenicals.

SULPHUR

100 a ✓ Midwest

- 1) Only in combination with a dermatological base or proprietary Topical Corticosteroid Plain
- 2) With or without other dermatological galenicals.

(Midwest Precipitated to be delisted 1 October 2028)

TACROLIMUS

Oint 0.1% - Special Authority see SA2074 below - Retail 30 g OP ✓ Zematop

- a) Maximum of 30 g per prescription
- b) Note: a maximum of 30 g per prescription and no more than one prescription per 12 weeks.

⇒SA2074 Special Authority for Subsidy

Initial application only from a dermatologist, paediatrician or any relevant practitioner on the recommendation of a dermatologist, paediatrician, . Approvals valid without further renewal unless notified for applications meeting the following criteria: Both:

- 1 Patient has atopic dermatitis on the face: and
- 2 Patient has at least one of the following contraindications to topical corticosteroids: periorificial dermatitis, rosacea, documented epidermal atrophy or documented allergy to topical corticosteroids.

Scalp Preparations

BETAMETHASONE VALERATE			
* Scalp app 0.1%	12.95	100 ml OP	✓ Beta Scalp
CLOBETASOL PROPIONATE			
* Scalp app 0.05%	6.90	30 ml OP	Dermol
Dermol to be Principal Supply on 1 February 2026			
HYDROCORTISONE BUTYRATE			
Scalp lotn 0.1%	6.57	100 ml OP	✓ Locoid
KETOCONAZOLE			
Shampoo 2%	4.09	100 ml OP	✓ Sebizole
a) Maximum of 100 ml per prescription			

DERMATOLOGICALS

Subsidy (Manufacturer's Price) \$

Subsidised Per 🗸

Fully

Brand or Generic Manufacturer

Sunscreens

SUNSCREENS, PROPRIETARY - Subsidy by endorsement

Only if prescribed for a patient with severe photosensitivity secondary to a defined clinical condition and the prescription is endorsed accordingly.

200 g OP

✓ Marine Blue Lotion SPF 50+

Wart Preparations

For salicylic acid preparations refer to PSORIASIS AND ECZEMA PREPARATIONS, page 73

PODOPHYLLOTOXIN

✓ Condyline S29 S29

- a) Maximum of 3.5 ml per prescription
- b) Only on a prescription

Other Skin Preparations

Antineoplastics

FLUOROURACIL SODIUM

IMIQUIMOD

GENITO-URINARY SYSTEM

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

Contraceptives - Non-hormonal

Condoms

COND		14.05	144	✓ Moments
	mm – Up to 144 dev available on a PSOmm		10	✓ Moments
• 55		14.25	144	✓ Moments
	a) Maximum of 60 dev per prescription	14.20	177	- momento
	b) Up to 60 dev available on a PSO			
÷ 53	mm, 0.05 mm thickness	1.15	10	✓ Moments
	, , , , , , , , , , , , , , , , , , , ,	14.25	144	✓ Moments
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription			
53	mm, chocolate, brown	1.15	10	✓ Moments
		14.25	144	✓ Moments
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription			
53	mm, strawberry, red	1.15	10	✓ Moments
		14.25	144	✓ Moments
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription			
56	mm		10	✓ Moments
		14.50	144	✓ Moments
	a) Maximum of 60 dev per prescription			
	b) Up to 60 dev available on a PSO			
56	mm, 0.05 mm thickness		12	✓ Gold Knight
		24.10	144	Gold Knight
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription			
56	mm, 0.05mm thickness (bulk pack)	20.17	144	Gold Knight
	a) Maximum of 60 dev per prescription			
	b) Up to 60 dev available on a PSO			
56	mm, 0.08 mm thickness		10	✓ Moments
		14.25	144	✓ Moments
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription			
56	mm, 0.08 mm thickness, red		10	✓ Moments
		14.25	144	✓ Moments
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription	4 70	40	A Cald Madella
56	mm, chocolate		12	✓ Gold Knight
	a) Ha ta 00 day aya'labla ay a BOO	21.45	144	✓ Gold Knight
	a) Up to 60 dev available on a PSO			
EC	b) Maximum of 60 dev per prescription	1 70	10	✓ Cald Value
56	mm, strawberry	1.79 21.45	12 144	✓ Gold Knight
	a) Ha ta CO day available an a DCO	∠1.45	144	✓ Gold Knight
	a) Up to 60 dev available on a PSO			
60	b) Maximum of 60 dev per prescription mm	1 00	12	✓ Gold Knight XL
60		21.89	12 144	✓ Gold Knight XL ✓ Gold Knight XL
	a) Maximum of 60 day now proporting	21.09	144	- Goid Killyill AL
	a) Maximum of 60 dev per prescription			

=					
		Subsidy (Manufacturer's Price)	Su	Fully bsidised	Brand or Generic
		\$	Per	√	Manufacturer
	a) Maximum of 60 dev per prescription				
	b) Up to 60 dev available on a PSO				
_					
C	ontraceptive Devices				
INT	RA-UTERINE DEVICE				
	a) Up to 40 dev available on a PSO				
	b) Only on a PSO				
*	IUD 29.1 mm length × 23.2 mm width	29.80	1	₽ C	hoice 380 7med Nsha Silver/
					copper Short
*	IUD 33.6 mm length × 29.9 mm width	26.80	1	✓ T	Cu 380 Plus
					Normal
*	IUD 35.5 mm length × 19.6 mm width	33.00	1	✓ C	u 375 Standard
^	ontraceptives - Hormonal				
_	ontraceptives - normonal				
C	ombined Oral Contraceptives				
FT	HINYLOESTRADIOL WITH DESOGESTREL				
	Tab 20 mcg with desogestrel 150 mcg and 7 inert tab - Up to	n			
•	84 tab available on a PSO		84	✓ M	lercilon 28
ET	HINYLOESTRADIOL WITH LEVONORGESTREL				
*	Tab 20 mcg with levonorgestrel 100 mcg and 7 inert tablets -				
	Up to 84 tab available on a PSO		84	✓ L	o-Oralcon 20 ED
*	Tab 30 mcg with levonorgestrel 150 mcg and 7 inert tablets - Up to 84 tab available on a PSO		84	./ 0	ralcon 30 ED
гт	HINYLOESTRADIOL WITH NORETHISTERONE	2.30	04	• 0	TAICOII 30 ED
LI	Tab 35 mcg with norethisterone 1 mg and 7 inert tab – Up to	1			
	84 tab available on a PSO		84	✓ A	lyacen
					revinor 1/28
	Tab 35 mcg with norethisterone 500 mcg and 7 inert tab - U				
	to 112 tab available on a PSO		84		orimin
(ΔΙ	yacen Tab 35 mcg with norethisterone 1 mg and 7 inert tab to	29.32 he delisted 1 January	112	♥ N	orimin
(7 1)	vacen rab oo meg war noreanoterone r mg and r men tab to	be denoted I dandary	2020)		
P	rogestogen-only Contraceptives				
DE	SOGESTREL				
	Tab 75 mcg - Up to 84 tab available on a PSO	24.50	84	√ C	erazette
LE'	VONORGESTREL				
	Tab 30 mcg - Up to 112 tab available on a PSO		112		licrolut
	Intra-uterine device 52 mg — Up to 25 dev available on a PS0	O269.50	1	✓ M	lirena
*	Intra-uterine device 13.5 mg – Up to 10 dev available on a PSO	215.60	1	./ 1	aydess
*	Subdermal implant (2 × 75 mg rods) – Up to 40 impl availabl		'	• 0	ayuess
•	on a PSO		2 OP	√ Ja	adelle
ME	DROXYPROGESTERONE ACETATE			_	_
	Inj 150 mg per ml, 1 ml syringe - Up to 5 inj available on a P	SO 10.56	1	✓ D	epo-Provera

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	Generic
	\$	Per	✓	Manufacturer
NORETHISTERONE				
Tab 350 mcg - Up to 84 tab available on a PSO	12.25	84	•	Norethinderone -
			,	CDC
				Noriday
			•	Noriday 28
(Norethinderone - CDC Tab 350 mcg to be delisted 1 January 20.	26)			
Emergency Contraceptives				

LEVONORGESTREL

- Levonorgestrel **BNM**
 - a) Maximum of 2 tab per prescription
 - b) Up to 5 tab available on a PSO
 - c) Note: Direct Provision by a pharmacist permitted under the provisions in Part I of Section A.

Antiandrogen Oral Contraceptives

Prescribers may code prescriptions "contraceptive" (code "O") when used as indicated for contraception. The period of supply and prescription charge will be as per other contraceptives, as follows:

- A maximum \$5.00 prescription charge (patient co-payment) may apply.
- prescription may be written for up to six months supply.

Prescriptions coded in any other way are subject to any non contraceptive prescription charges that apply, and the non-contraceptive period of supply. ie. Prescriptions may be written for up to three months supply.

CYPROTERONE ACETATE WITH ETHINYLOESTRADIOL

Tab 2 mg with ethinyloestradiol 35 mcg and 7 inert tabs - Up

✓ Ginet 168

Gynaecological Anti-infectives

ACETIC ACID WITH HYDROXYQUINOLINE AND RICINOLEIC ACID
Jelly with glacial acetic acid 0.94%, hydroxyguinoline sulphate

0.025%, glycerol 5% and ricinoleic acid 0.75% with applicator.... 8.43 100 a OP (24.87)Aci-Jel

CLOTRIMAZOLE

35 a OP ✓ Clomazol 20 g OP Clomazol

MICONAZOLE NITRATE

40 g OP Micreme

NYSTATIN

75 g OP Nilstat

Myometrial and Vaginal Hormone Preparations

ERGOMETRINE MALEATE

inj 500 mcg per mi, i mi ampoule – op to 5 mj avallable	UII a		
PSO	160.00	5	 DBL Ergometrine
OESTRIOI			

Crm 1 mg per g with applicator......6.95 15 q OP Ovestin Ovestin * Pessaries 500 mcg......7.55 15

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

GENITO-URINARY SYSTEM

	Subsidy (Manufacturer's Price) \$	Sul Per	Fully bsidised	Brand or Generic Manufacturer
OXYTOCIN – Up to 5 inj available on a PSO	5.00	_		Annata dia DAMA
Inj 5 iu per ml, 1 ml ampoule Oxytocin BNM to be Principal Supply on 1 March 2026	5.98	5	• 0	Oxytocin BNM
Inj 10 iu per ml, 1 ml ampoule Oxytocin BNM to be Principal Supply on 1 March 2026	7.18	5	√ 0	Oxytocin BNM
OXYTOCIN WITH ERGOMETRINE MALEATE – Up to 5 inj avai Inj 5 iu with ergometrine maleate 500 mcg per ml, 1 ml ampo Syntometrine to be Principal Supply on 1 February 2026	ule41.47	5	√ \$	syntometrine

Pregnancy Tests - hCG Urine

BETA-HCG LOW SENSITIVITY URINE TEST KIT - Up to 15 test available on a PSO

Note: For use in abortion services only.

PREGNANCY TESTS - HCG URINE

- a) Up to 200 test available on a PSO
- b) Only on a PSO

Cassette Pregnancy Test

Urinary Agents

For urinary tract Infections refer to INFECTIONS, Antibacterials, page 112

5-Alpha Reductase Inhibitors

FINASTERIDE – Special Authority see SA0928 below – Retail pharmacy

⇒SA0928 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 Eitner:
 - 2.1 The patient is intolerant of non-selective alpha blockers or these are contraindicated; or
 - 2.2 Symptoms are not adequately controlled with non-selective alpha blockers.

Alpha-1A Adrenoreceptor Blockers

TAMSULOSIN HYDROCHLORIDE - Special Authority see SA1032 below - Retail pharmacy

Tamsulosin-Rex to be Principal Supply on 1 February 2026

⇒SA1032 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 The patient is intolerant of non-selective alpha blockers or these are contraindicated.

	Subsidy (Manufacturer's P \$	rice) Subsi Per		
Other Urinary Agents				
OXYBUTYNIN				
* Tab 5 mg	5.42	100	✓ Alchem Oxvb	y utynin
POTASSIUM CITRATE			,	•
Oral liq 3 mmol per ml — Special Authority see SA1083 belov Retail pharmacy		200 ml OP	✓ Biomed	
⇒SA1083 Special Authority for Subsidy				
Initial application from any relevant practitioner. Approvals valid Both:	I for 12 months f	or applications	meeting the f	ollowing criteria:
1 The patient has recurrent calcium oxalate urolithiasis; and				
2 The patient has had more than two renal calculi in the two			. annvanviata	and the notiont is
Renewal from any relevant practitioner. Approvals valid for 2 yea benefitting from the treatment.	us where the tre	aunent remains	appropriate	and the patient is
SODIUM CITRO-TARTRATE				
* Grans eff 4 g sachets	3.50	28	✓ Ural	
SOLIFENACIN SUCCINATE				
Tab 5 mg	1.95	30	✓ Solifena	<u>acin</u>
				nate Max
T 40	0.50	••	Healtl	_
Tab 10 mg	3.53	30	✓ Solifena	
			Healtl	nate Max n
			Hour	<u>-</u>
Detection of Substances in Urine				
ORTHO-TOLIDINE				
* Compound diagnostic sticks	7.50	50 test OP		
. •	(8.25)		Hemast	X
TETRABROMOPHENOL				
* Blue diagnostic strips	13.92	100 test OP	✓ Albusti:	(
Obstetric Preparations				
Antiprogesterance				
Antiprogesterones				

MIFEPRISTONE ✓ Mifegyne✓ Mifegyne Tab 200 mg - Up to 15 tab available on a PSO......83.90 1 180.00

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy		Fully	Brand or	
(Manufacturer's Pric	e)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

Calcium Homeostasis

CALCITONIN * Inj 100 iu per ml, 1 ml ampoule	121.00	5	✓ Miacalcic ✓ Miacalcic S29 S29
CINACALCET - Special Authority see SA2170 below - Retail p	harmacy		
Tab 30 mg - Wastage claimable	25.24	28	✓ Cinacalet Devatis
Tab 60 mg - Wastage claimable	50.47	28	✓ Cinacalet Devatis

⇒SA2170 Special Authority for Subsidy

Initial application — (parathyroid carcinoma or calciphylaxis) only from a nephrologist or endocrinologist. Approvals valid for 6 months for applications meeting the following criteria: Fither:

- 1 All of the following:
 - 1.1 The patient has been diagnosed with a parathyroid carcinoma (see Note); and
 - 1.2 The patient has persistent hypercalcaemia (serum calcium greater than or equal to 3 mmol/L) despite previous first-line treatments including sodium thiosulfate (where appropriate) and bisphosphonates: and
 - 1.3 The patient is symptomatic; or
- 2 All of the following:
 - 2.1 The patient has been diagnosed with calciphylaxis (calcific uraemic arteriolopathy); and
 - 2.2 The patient has symptomatic (e.g. painful skin ulcers) hypercalcaemia (serum calcium greater than or equal to 3 mmol/L); and
 - 2.3 The patient's condition has not responded to previous first-line treatments including bisphosphonates and sodium thiosulfate.

Renewal — (parathyroid carcinoma or calciphylaxis) only from a nephrologist or endocrinologist. Approvals valid without further renewal unless notified for applications meeting the following criteria: Both:

- 1 The patient's serum calcium level has fallen to < 3mmol/L: and
- 2 The patient has experienced clinically significant symptom improvement.

Note: This does not include parathyroid adenomas unless these have become malignant.

Initial application — (primary hyperparathyroidism) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has primary hyperparathyroidism; and
- 2 Fither:
 - 2.1 Patient has hypercalcaemia of more than 3 mmol/L with or without symptoms; or
 - 2.2 Patient has hypercalcaemia of more than 2.85 mmol/L with symptoms; and
- 3 Surgery is not feasible or has failed; and
- 4 Patient has other comorbidities, severe bone pain, or calciphylaxis.

Initial application — (secondary or tertiary hyperparathyroidism) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Patient has tertiary hyperparathyroidism and markedly elevated parathyroid hormone (PTH) with hypercalcaemia:
 - 1.2 Patient has symptomatic secondary hyperparathyroidism and elevated PTH; and
- 2 Patient is on renal replacement therapy; and
- 3 Any of the following:

Subsidy (Manufacturer's Price)	Subs	Fully	Brand or Generic
\$	Per	/	Manufacturer

continued...

- 3.1 Residual parathyroid tissue has not been localised despite repeat unsuccessful parathyroid explorations; or
- 3.2 Parathyroid tissue is surgically inaccessible; or
- 3.3 Parathyroid surgery is not feasible.

Renewal — (secondary or tertiary hyperparathyroidism) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 The patient has had a kidney transplant, and following a treatment free interval of at least 12 weeks a clinically acceptable parathyroid hormone (PTH) level to support ongoing cessation of treatment has not been reached; or
- 2 The patient has not received a kidney transplant and trial of withdrawal of cinacalcet is clinically inappropriate.

ZOLEDRONIC ACID

EEDITOTTO			
Inj 4 mg per 5 ml, vial	15.65	1	✓ Zoledronic acid Injection
			Mylan S29
			✓ Zoledronic acid
			Viatris

Corticosteroids and Related Agents for Systemic Use

BETAMETHASONE SODIUM PHOSPHATE WITH BETAMETHASONE ACETA* * Inj 3.9 mg with betamethasone acetate 3 mg per ml, 1 ml	TE 5	Celestone Chronodose
DEXAMETHASONE		
* Tab 0.5 mg – Up to 60 tab available on a PSO1.80	30	✓ Dexmethsone
* Tab 4 mg - Up to 30 tab available on a PSO	30	✓ <u>Dexmethsone</u>
Oral liq 1 mg per ml53.86	25 ml OP	✓ Biomed
DEXAMETHASONE PHOSPHATE		
Dexamethasone phosphate injection will not be funded for oral use.		
* Inj 4 mg per ml, 1 ml ampoule – Up to 5 inj available on a PSO6.88	10	Dexamethasone Medsurge
7.86		✓ Hameln
Dexamethasone Medsurge to be Principal Supply on 1 March 2026		
* Inj 4 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO10.98	10	Dexamethasone Medsurge
13.10		✓ Hameln
Dexamethasone Medsurge to be Principal Supply on 1 March 2026		
(Hameln Inj 4 mg per ml, 1 ml ampoule to be delisted 1 March 2026)		
(Hameln Inj 4 mg per ml, 2 ml ampoule to be delisted 1 March 2026)		
FLUDROCORTISONE ACETATE		
* Tab 100 mcg8.05	100	✓ Florinef
HYDROCORTISONE		
* Tab 5 mg	100	✓ Douglas
* Tab 20 mg	100	✓ Douglas
* Inj 100 mg vial	1	✓ Solu-Cortef
a) Not on a BSO		
b) Up to 5 inj available on a PSO		
METHYLPREDNISOLONE		
* Tab 4 mg	100	✓ Medrol
* Tab 100 mg223.10	20	✓ Medrol

	Subsidy (Manufacturer's Price \$) Per	Fully Brand or Subsidised Generic Manufacturer
METHYLPREDNISOLONE (AS SODIUM SUCCINATE)			
Inj 40 mg vial	22.30	1	✓ Solu-Medrol-Act- O-Vial
Inj 125 mg vial	34.10	1	✓ Solu-Medrol-Act- O-Vial
Inj 500 mg vial	43.01	1	✓ Solu-Medrol-Act- O-Vial
Inj 1 g vial	52.54	1	✓ Solu-Medrol
METHYLPREDNISOLONE ACETATE Inj 40 mg per ml, 1 ml vial	47.06	5	✓ Depo-Medrol
	47.00	J	• Depo-Medioi
REDNISOLONE ★ Oral liq 5 mg per ml — Up to 30 ml available on a PSO Restricted to children under 12 years of age.	6.00	30 ml C	OP ✓ <u>Redipred</u>
REDNISONE			
€ Tab 1 mg		500	
F Tab 2.5 mg		500	
Tab 5 mg - Up to 30 tab available on a PSO		500	
Tab 20 mg − Up to 30 tab available on a PSO ETRACOSACTRIN	50.51	500	✓ Prednisone Clinect
f Inj 250 mcg per ml, 1 ml ampoule	86.25	1	✓ Synacthen
Finj 1 mg per ml, 1 ml ampoule	690.00	1	✓ UK Synacthen ✓ Synacthen Depot ✓ Synacthene Retard \$29
RIAMCINOLONE ACETONIDE			
Inj 10 mg per ml, 1 ml ampoule	21.42	5	✓ Kenacort-A 10
Inj 40 mg per ml, 1 ml ampoule		5	✓ Kenacort-A 40
Sex Hormones Non Contraceptive			
Androgen Agonists and Antagonists			
YPROTERONE ACETATE			
Tab 50 mg		50	✓ <u>Siterone</u>
Tab 100 mg	31.00	50	✓ <u>Siterone</u>
ESTOSTERONE Gel (transdermal) 16.2 mg per g, 88 g	52.00	60 OF	P ✓ Testogel
ESTOSTERONE CIPIONATE		00 01	<u> 100togor</u>
Inj 100 mg per ml, 10 ml vial	85.00	1	✓ Depo-Testosterone
ESTOSTERONE ESTERS Inj 250 mg per ml, 1 ml	12.98	1	✓ Sustanon Ampoules
ESTOSTERONE UNDECANOATE		•	
Cap 40 mg - Subsidy by endorsement	36.00	100	✓ Steril-Gene S29
Subsidy by endorsement – subsidised for patients who	were taking testoster	one ur	ndecanoate cap 40mg prior to
1 November 2021 and the prescription is endorsed acc	cordingly. Pharmacist	s may	annotate the prescription as endor
where there exists a record of prior dispensing of testo			
Inj 250 mg per ml, 4 ml vial	86.00	1	✓ Reandron 1000

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

	\$ Per	 Manutac
Harmana Bankaamant Tharany, Cystamia		
Hormone Replacement Therapy - Systemic		

Destrogens			
ESTRADIOL			
Tab 1 mg	4.12	28 OP	
	(11.10)		Estrofem
Tab 2 mg	4.12	28 OP	
	(11.10)		Estrofem
Gel (transdermal) 0.06% (750 mcg/actuation)		80 g OP	✓ Estrogel
Patch 25 mcg per day		8	Estradiol TDP Mylar
	16.23		✓ Estradot
a) Brand switch fee payable (Pharmacode 27b) No more than 2 patch per week	717573) - see page 279 for	details	
c) Only on a prescription			
Patch 50 mcg per day	9.26	8	 Estradiol TDP Mylar
	15.79		✓ Estradot
 a) Brand switch fee payable (Pharmacode 27 b) No more than 2 patch per week 	717573) - see page 279 for	details	
c) Only on a prescription Patch 75 mcg per day	10.00	8	✓ Estradiol TDP Myla
Falcii 75 mcg per day	16.53	0	✓ Estradioi TDP Mylai ✓ Estradot
a) Brand switch fee payable (Pharmacode 27b) No more than 2 patch per weekc) Only on a prescription	, , ,	details	
Patch 100 mcg per day	10.59	8	 Estradiol TDP Mylar
	16.18		Estradot
a) Brand switch fee payable (Pharmacode 27b) No more than 2 patch per weekc) Only on a prescription	717573) - see page 279 for	details	
ESTRADIOL VALERATE			
Tab 1 mg	12.36	84	✓ Progynova
Tab 2 mg	12.36	84	✓ Progynova
ESTROGENS			
Conjugated, equine tab 300 mcg	3.01	28	
	(19.25)		Premarin
Conjugated, equine tab 625 mcg	` ,	28	
•	(19.25)		Premarin

ME	DROXYPROGESTERONE ACETATE		
*	Tab 2.5 mg6.56	30	✓ Provera
	8.75	56	✓ Provera
*	Tab 5 mg9.80	56	✓ Provera
	20.13	100	✓ Provera
*	Tab 10 mg	30	✓ Provera

	Subsidy (Manufacturer's Price \$	e) Subs Per	Fully Brand or sidised Generic Manufacturer
Progestogen and Oestrogen Combined Prepara	ations		
OESTRADIOL WITH NORETHISTERONE			
* Tab 1 mg with 0.5 mg norethisterone acetate		28 OP	
	(18.10)		Kliovance
* Tab 2 mg with 1 mg norethisterone acetate		28 OP	Kliagast
* Tab 2 mg with 1 mg norethisterone acetate (10), and 2 mg	(18.10)		Kliogest
oestradiol tab (12) and 1 mg oestradiol tab (6)	5.40	28 OP	
destraction tab (12) and 1 mg destraction tab (0)	(18.10)	20 01	Trisequens
	(10.10)		mocqueno
Other Oestrogen Preparations			
OESTRIOL			
* Tab 2 mg	7.70	30	✓ Ovestin
Other Progestogen Preparations			
MEDROXYPROGESTERONE ACETATE Tab 100 mg	100 57	100	✓ Provera HD
· ·	133.37	100	▼ Provera nD
NORETHISTERONE ** Tab 5 mg	E 40	30	✓ Primolut N
* Tab 5 mg - Up to 30 tab available on a PSO	5.49	30	♥ Primolut N
PROGESTERONE	14.05	20	./ Ilitra mantan
* Cap 100 mg	14.65	30	✓ Utrogestan
Thyroid and Antithyroid Agents			
CARBIMAZOLE			
* Tab 5 mg	7.56	100	✓ Neo-Mercazole
LEVOTHYROXINE			
* Tab 25 mcg	5.55	90	✓ Synthroid
* Tab 50 mcg	1.71	28	Mercury Pharma
	5.79	90	✓ Synthroid
	64.28	1,000	✓ Eltroxin
* Tablet 50 mcg		200	✓ Eltroxin
* Tab 100 mcg		28	✓ Mercury Pharma
	6.01 66.78	90 1.000	✓ Synthroid✓ Eltroxin
* Tablet 100 mcg		200	✓ Eltroxin
<u> </u>		200	LIUVAIII
PROPYLTHIOURACIL - Special Authority see SA1199 below -		100	./ DTIL coo
Tab 50 mg	35.00	100	✓ PTU S29

⇒SA1199 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 The patient has hyperthyroidism; and
- 2 The patient is intolerant of carbimazole or carbimazole is contraindicated.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
•	Por		Manufacturer

Trophic Hormones

Growth Hormones

SO	MATROPIN (OMNITROPE) - Special Auth	nority see SA2032 below - Retail pharr	nacy	
*	Inj 5 mg cartridge	80.21	1	✓ Omnitrope
				✓ Omnitrope AU S29
*	Inj 10 mg cartridge	80.21	1	✓ Omnitrope
*	Inj 15 mg cartridge	139.50	1	✓ Omnitrope

⇒SA2032 Special Authority for Subsidy

Initial application — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

Either:

- 1 Growth hormone deficiency causing symptomatic hypoglycaemia, or with other significant growth hormone deficient sequelae (e.g. cardiomyopathy, hepatic dysfunction) and diagnosed with GH < 5 mcg/l on at least two random blood samples in the first 2 weeks of life, or from samples during established hypoglycaemia (whole blood glucose < 2 mmol/l using a laboratory device); or</p>
- 2 All of the following:
 - 2.1 Height velocity < 25th percentile for age adjusted for bone age/pubertal status if appropriate over 6 or 12 months using the standards of Tanner and Davies (1985); and
 - 2.2 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
 - 2.3 Peak growth hormone value of < 5.0 mcg per litre in response to two different growth hormone stimulation tests. In children who are 5 years or older, GH testing with sex steroid priming is required; and
 - 2.4 If the patient has been treated for a malignancy, they should be disease free for at least one year based upon follow-up laboratory and radiological imaging appropriate for the malignancy, unless there are strong medical reasons why this is either not necessary or appropriate; and
 - 2.5 Appropriate imaging of the pituitary gland has been obtained.

Renewal — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 2 Height velocity is greater than or equal to 25th percentile for age (adjusted for bone age/pubertal status if appropriate) while on growth hormone treatment, as calculated over six months using the standards of Tanner and Davis (1985); and
- 3 Height velocity is greater than or equal to 2.0 cm per year, as calculated over 6 months; and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a post-natal genotype confirming Turner Syndrome; and
- 2 Height velocity is < 25th percentile over 6-12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is < 14 years.

Renewal — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 Height velocity is greater than or equal to 50th percentile for age (while on growth hormone calculated over 6 to 12 months using the Ranke's Turner Syndrome growth velocity charts); and

Subsidy (Manufacturer's Price)	Sub	Fully sidised	Brand or Generic
 \$	Per	1	Manufacturer

continued...

- 2 Height velocity is greater than or equal to 2 cm per year, calculated over six months; and
- 3 A current bone age is 14 years or under; and
- 4 No serious adverse effect that the specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 3 standard deviations below the mean for age or for bone age if there is marked growth acceleration or delay; and
- 2 Height velocity is < 25th percentile for age (adjusted for bone age/pubertal status if appropriate), as calculated over 6 to 12 months using the standards of Tanner and Davies(1985); and
- 3 A current bone age is < 14 years or under (female patients) or < 16 years (male patients); and
- 4 The patient does not have severe chronic disease (including malignancy or recognized severe skeletal dysplasia) and is not receiving medications known to impair height velocity.

Renewal — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

- All of the following:
 - 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
 - 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
 - 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
 - 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred.

Initial application — **(short stature due to chronic renal insufficiency)** only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 2 standard deviations below the mean; and
- 2 Height velocity is < 25th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and</p>
- 3 A current bone age is to 14 years or under (female patients) or to 16 years or under (male patients); and
- 4 The patient is metabolically stable, has no evidence of metabolic bone disease and absence of any other severe chronic disease; and
- 5 The patient is under the supervision of a specialist with expertise in renal medicine; and
- 6 Either:
 - 6.1 The patient has a GFR less than or equal to 30 ml/min/1.73m² as measured by the Schwartz method (Height(cm)/plasma creatinine (umol/l)) × 40 = corrected GFR (ml/min/1.73m²) in a child who may or may not be receiving dialysis; or
 - 6.2 The patient has received a renal transplant and has received < 5mg/ m²/day of prednisone or equivalent for at least 6 months...

Renewal — (short stature due to chronic renal insufficiency) only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and

Subsidy	F	ully	Brand or	
(Manufacturer's Price)	Subsidi	sed	Generic	
\$	Per	1	Manufacturer	

continued...

- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not experienced significant biochemical or metabolic deterioration confirmed by diagnostic results; and
- 7 The patient has not received renal transplantation since starting growth hormone treatment; and
- 8 If the patient requires transplantation, growth hormone prescription should cease before transplantation and a new application should be made after transplantation based on the above criteria.

Initial application — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of Prader-Willi syndrome that has been confirmed by genetic testing or clinical scoring criteria; and
- 2 The patient is aged six months or older; and
- 3 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
- 4 Sleep studies or overnight oximetry have been performed and there is no obstructive sleep disorder requiring treatment, or if an obstructive sleep disorder is found, it has been adequately treated under the care of a paediatric respiratory physician and/or ENT surgeon; and
- 5 Fither:
 - 5.1 Both:
 - 5.1.1 The patient is aged two years or older; and
 - 5.1.2 There is no evidence of type II diabetes or uncontrolled obesity defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months; or
 - 5.2 The patient is aged between six months and two years and a thorough upper airway assessment is planned to be undertaken prior to treatment commencement and at six to 12 weeks following treatment initiation.

Renewal — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not developed type II diabetes or uncontrolled obesity as defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months.

Initial application — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a medical condition that is known to cause growth hormone deficiency (e.g. surgical removal of the pituitary for treatment of a pituitary tumour); and
- 2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
- 3 The patient has severe growth hormone deficiency (see notes); and
- 4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
- 5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life

	Subsidy		Fully	Brand or
(N	fanufacturer's Price)	5	Subsidised	Generic
	\$	Per	•	Manufacturer

continued...

questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

Renewal — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria: Any of the following:

- 1 All of the following:
 - 1.1 The patient has been treated with somatropin for < 12 months; and
 - 1.2 There has been an improvement in Quality of Life defined as a reduction of at least 8 points on the Quality of Life Assessment of Growth Hormone Deficiency in Adults (QoL-AGHDA®) score from baseline; and
 - 1.3 Serum IGF-I levels have been increased within ±1SD of the mean of the normal range for age and sex; and
 - 1.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients, or 1 mg per day for female patients; or
- 2 All of the following:
 - 2.1 The patient has been treated with somatropin for more than 12 months; and
 - 2.2 The patient has not had a deterioration in Quality of Life defined as a 6 point or greater increase from their lowest QoL-AGHDA® score on treatment (other than due to obvious external factors such as external stressors); and
 - 2.3 Serum IGF-I levels have continued to be maintained within ±1SD of the mean of the normal range for age and sex (other than for obvious external factors); and
 - 2.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients or 1 mg per day for female patients; or
- 3 All of the following:
 - 3.1 The patient has had a Special Authority approval for somatropin for childhood deficiency in children and no longer meets the renewal criteria under this indication; and
 - 3.2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
 - 3.3 The patient has severe growth hormone deficiency (see notes); and
 - 3.4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
 - 3.5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		osidised	Generic
	\$	Per		Manufacturer
continued doses of corticosteroid and levothyroxine.				
GnRH Analogues				
GOSERELIN				
Implant 3.6 mg, syringe Implant 10.8 mg, syringe		1 1	-	<u>Zoladex</u> Zoladex
LEUPRORELIN			_	
Additional subsidy by endorsement where the patient is a ch goserelin and the prescription is endorsed accordingly.	nild or adolescent and	l is unable	to toler	rate administration of
Inj 3.75 mg prefilled dual chamber syringe - Higher subsidy				
\$221.60 per 1 inj with Endorsement		1		
	(221.60)		l	Lucrin Depot 1-month
Inj 11.25 mg prefilled dual chamber syringe – Higher subsid				
of \$591.68 per 1 inj with Endorsement		1		ucrin Donot 2 month
	(591.68)		Į.	Lucrin Depot 3-month
Vasopressin Agonists				
DESMOPRESSIN				
Wafer 120 mcg	47 00	30	√ I	Minirin Melt
DESMOPRESSIN ACETATE		00		
Tab 100 mcg	25.00	30	√ I	Minirin
Tab 200 mcg		30		Minirin
Inj 4 mcg per ml, 1 ml		10		Minirin
▲ Nasal spray 10 mcg per dose, 6 ml	34.95	60 OP	√ <u>I</u>	<u>Desmopressin-</u> <u>PH&T</u>
Other Endessine Avents				
Other Endocrine Agents				
CABERGOLINE				
Tab 0.5 mg – Maximum of 2 tab per prescription; can be		_		
waived by Special Authority see SA2070 below		2		Dostinex
	17.94	8	•	Dostinex
Special Authority for Waiver of Rule	id without fruthou room	موامين اميين	a natifi	ad for applications mosting
Initial application from any relevant practitioner. Approvals valithe following criteria:	ia without further rene	ewai unies	ss noune	ed for applications meeting
Any of the following:				
1 Hyperprolactinemia; or				
2 Acromegaly*; or				
3 Inhibition of lactation.				
Renewal — (for patients who have previously been funded u	under Special Autho	rity form	SA103	1) from any relevant
practitioner. Approvals valid without further renewal unless notif				
which has expired and the treatment remains appropriate and th	e patient is benefiting	from trea	atment.	
Note: Indication marked with * is an unapproved indication.				
CLOMIFENE CITRATE				
Tab 50 mg	29.84	10	√ I	Mylan Clomiphen S29
METYRAPONE				
Cap 250 mg	558.00	50	✓ I	Metopirone
-				

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.
★Three months or six months, as applicable, dispensed all-at-once

Subsidy	Full	y Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per 🗸	Manufacturer	

Anthelmintics

ALBENDAZOLE - Special Authority see SA2512 below - Retail pharmacy

60 ✓ Eskazole S29

⇒SA2512 Special Authority for Subsidy

Initial application only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: Either:

- 1 The individual has hydatids; or
- 2 The individual has a travel or residence history that requires presumptive parasite treatment.

Renewal only from an infectious disease specialist or clinical microbiologist. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefitting from the treatment.

MEBENDAZOLE - Only on a prescription

Tab 100 mg	5.18	6	✓ Vermox
Oral liq 100 mg per 5 ml		15 ml	
	(7.83)		Vermox
PRAZIQUANTEL			
Tab 600 mg	68.00	8	✓ Biltricide
•	87.68		✓ Distoside S29
(Biltricide Tab 600 mg to be delisted 1 April 2026)			

Antibacterials

- a) For topical antibacterials, refer to DERMATOLOGICALS, page 67
- b) For anti-infective eye preparations, refer to SENSORY ORGANS, page 274

Cephalosporins and Cephamycins

CEFACLOR MONOHYDRATE			
Cap 250 mg	29.73	100	Ranbaxy-Cefactor
Ranbaxy-Cefaclor to be Principal Supply on 1 February 2	2026		•
Grans for oral liq 125 mg per 5 ml - Wastage claimable	5.83	100 ml	✓ Ranbaxy-Cefaclor
Ranbaxy-Cefaclor to be Principal Supply on 1 February 2	2026		
CEFALEXIN			
Cap 250 mg	3.85	20	 Cephalexin ABM
, ,	3.90		✓ Cefalexin Lupin
Cap 500 mg	3.33	20	 Cefalexin Sandoz
	5.85		 Cephalexin ABM
Grans for oral liq 25 mg per ml - Wastage claimable	7.88	100 ml	✓ Flynn
Grans for oral lig 50 mg per ml - Wastage claimable	10.38	100 ml	✓ Flynn
	11.75		✓ Cefalexin Sandoz
(Cephalexin ABM Cap 250 mg to be delisted 1 July 2026)			

(Cephalexin ABM Cap 500 mg to be delisted 1 July 2026)

CEFAZOLIN - Subsidy by endorsement

Only if prescribed for dialysis or cellulitis in accordance with a Health NZ Hospital approved protocol and the prescription is endorsed accordingly.

Inj 500 mg vial	5	✓ Cefazolin-AFT
Inj 1 g vial	5	✓ Cefazolin-AFT
Inj 2 g vial7.09	5	✓ Cefazolin-AFT

Cefuroxime \$29

	Subsidy (Manufacturer's Price)		Fully lised	Brand or Generic Manufacturer
CEFTRIAXONE – Subsidy by endorsement				
a) Up to 10 inj available on a PSO				
 Subsidised only if prescribed for a dialysis or cystic fibrosi pelvic inflammatory disease, or the treatment of suspected endorsed accordingly. 				
Inj 500 mg vial		1	✓ (Ceftriaxone-AFT
Inj 1 g vial Ceftriaxone-AFT to be Principal Supply on 1 February 20	3.49	5	✓ (Ceftriaxone-AFT
CEFUROXIME AXETIL - Subsidy by endorsement Only if prescribed for prophylaxis of endocarditis and the pres	scription is endorsed	accordingly		
Tab 250 mg	CBS	20	√	Ascend-

Macrolides

AZITHROMYCIN – Maximum of 5 days treatment per prescription; can be waived by Special Authority see SA1683 below
A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised on Special
Authority.

Tab 250 mg

Tab 250 mg	8.19	30	✓ Apo-Azithromyci
Tab 500 mg - Up to 8 tab available on a PSO		2	✓ Zithromax
Zithromax to be Principal Supply on 1 January 2026			
Grans for oral liq 200 mg per 5 ml (40 mg per ml) - Wastage			
claimable	16.97	15 ml	✓ Zithromax

⇒SA1683 Special Authority for Waiver of Rule

Initial application — (bronchiolitis obliterans syndrome, cystic fibrosis and atypical Mycobacterium infections) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following:

- 1 Patient has received a lung transplant, stem cell transplant, or bone marrow transplant and requires treatment for bronchiolitis obliterans syndrome*; or
- 2 Patient has received a lung transplant and requires prophylaxis for bronchiolitis obliterans syndrome*; or
- 3 Patient has cystic fibrosis and has chronic infection with Pseudomonas aeruginosa or Pseudomonas-related gram negative organisms*; or
- 4 Patient has an atypical Mycobacterium infection.

Note: Indications marked with * are unapproved indications.

Initial application — (non-cystic fibrosis bronchiectasis*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 For prophylaxis of exacerbations of non-cystic fibrosis bronchiectasis*; and
- 2 Patient is aged 18 and under; and
- 3 Either:
 - 3.1 Patient has had 3 or more exacerbations of their bronchiectasis, within a 12 month period; or
 - 3.2 Patient has had 3 acute admissions to hospital for treatment of infective respiratory exacerbations within a 12 month period.

Note: Indications marked with * are unapproved indications.

Renewal — (non-cystic fibrosis bronchiectasis*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

Subsidy		Fully	Brand or
(Manufacturer's Price	e)	Subsidised	Generic
\$	Per	/	Manufacturer

continued...

- 1 The patient has completed 12 months of azithromycin treatment for non-cystic fibrosis bronchiectasis; and
- 2 Following initial 12 months of treatment, the patient has not received any further azithromycin treatment for non-cystic fibrosis bronchiectasis for a further 12 months, unless considered clinically inappropriate to stop treatment; and
- 3 The patient will not receive more than a total of 24 months' azithromycin cumulative treatment (see note).

The patient must not have had more than 1 prior approval.

Note: No further renewals will be subsidised. A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised. Indications marked with * are unapproved indications

CLARITHROMYCIN - Maximum of 500 mg per prescription; can be waived by Special Authority see SA1857 below

Tab 250 mg7.31	12	✓ Klaricid \$29
8.53	14	✓ Klacid
Grans for oral liq 250 mg per 5 ml - Wastage claimable192.00	50 ml	✓ Klacid

⇒SA1857 Special Authority for Waiver of Rule

Initial application — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years for applications meeting the following criteria:

Fither:

ilci.

- 1 Atypical mycobacterial infection; or
- 2 Mycobacterium tuberculosis infection where there is drug-resistance or intolerance to standard pharmaceutical agents.

Initial application — (Helicobacter pylori eradication) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 For the eradication of helicobacter pylori in a patient unable to swallow tablets; and
- 2 For use only in combination with omeprazole and amoxicillin as part of a triple therapy regimen.

Initial application — (Prophylaxis of infective endocarditis) from any relevant practitioner. Approvals valid for 3 months where prophylaxis of infective endocarditis associated with surgical or dental procedures if amoxicillin is contra-indicated. Renewal — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

ERYTHROMYCIN (AS LACTOBIONATE) Inj 1 g vial	10.00	1	✓ Erythrocin IV
ERYTHROMYCIN ETHYL SUCCINATE	10.00	1	Elyunochilv
Tab 400 mg	35.82	100	✓ E-Mycin
a) Up to 20 tab available on a PSOb) Up to 2 x the maximum PSO quantity for RFPP			·
Grans for oral liq 200 mg per 5 ml	6.53	100 ml	✓ E-Mycin
a) Up to 300 ml available on a PSOb) Up to 2 x the maximum PSO quantity for RFPPc) Wastage claimable			
Grans for oral liq 400 mg per 5 ml	9.41	100 ml	E-Mycin
a) Up to 200 ml available on a PSOb) Wastage claimable			
ROXITHROMYCIN			
Tab 150 mg	13.19	50	Arrow- Roxithromycin
Tab 300 mg	25.00	50	✓ <u>Arrow-</u> <u>Roxithromycin</u>

	Subsidy (Manufacturer's Price \$	e) S Per	Fully ubsidised	Brand or Generic Manufacturer
Penicillins				
AMOXICILLIN				
Cap 250 mg	54.00	500	1	Miro-Amoxicillin
a) Up to 30 cap available on a PSO				
b) Up to 10 x the maximum PSO quantity for RFPP				
c) Miro-Amoxicillin to be Principal Supply on 1 Februar	v 2026			
Cap 500 mg		500	1	Miro-Amoxicillin
a) Up to 30 cap available on a PSO				
b) Up to 10 x the maximum PSO quantity for RFPP				
Grans for oral liq 125 mg per 5 ml	2.22	100 ml	/	Alphamox 125
a) Up to 200 ml available on a PSO				
b) Wastage claimable				
Grans for oral liq 250 mg per 5 ml	2.81	100 ml	1	Alphamox 250
a) Up to 300 ml available on a PSO				
b) Up to 10 x the maximum PSO quantity for RFPP				
c) Wastage claimable				
Inj 250 mg vial	15.97	10	✓	Ibiamox
Inj 500 mg vial		10	1	Ibiamox
Inj 1 g vial - Up to 5 inj available on a PSO	21.64	10	1	lbiamox
AMOXICILLIN WITH CLAVULANIC ACID				
Tab 500 mg with clavulanic acid 125 mg - Up to 30 tab				
available on a PSO	1.59	10	1	Curam Duo 500/125
Grans for oral lig amoxicillin 25 mg with clavulanic acid 6.25				
per ml	•	100 ml	/	Augmentin
a) Up to 200 ml available on a PSO				
b) Wastage claimable				
Grans for oral lig amoxicillin 50 mg with clavulanic acid 12.5	ma			
per ml – Up to 200 ml available on a PSO		00 ml OF	•	Amoxiclav Devatis
,				Forte
BENZATHINE BENZYLPENICILLIN				
Inj 900 mg (1.2 million units) vial	13 21	1	1	Benzetacil S29
Inj 900 mg (1.2 million units) in 2.3 ml syringe – Up to 5 inj	40.24	'	•	Delizetacii 020
available on a PSO	132 37	10	1	Bicillin LA
	402.07	10	•	DICIIIII LA
BENZYLPENICILLIN SODIUM [PENICILLIN G]	10.50	40		Condon
Inj 600 mg (1 million units) vial – Up to 5 inj available on a F	'SU 16.50	10	•	<u>Sandoz</u>
FLUCLOXACILLIN			_	
Cap 250 mg – Up to 30 cap available on a PSO		250		<u>Staphlex</u>
Cap 500 mg – Up to 30 cap available on a PSO		500		Staphlex
Grans for oral liq 25 mg per ml	4.89	100 ml	•	<u>AFT</u>
a) Up to 200 ml available on a PSO				
b) Wastage claimable	F 00	100!		AFT
Grans for oral liq 50 mg per ml	5.89	100 ml	•	<u>AFT</u>
a) Up to 200 ml available on a PSO				
b) Wastage claimable	40.60	10	.1	Eluciovin
Inj 250 mg vial		10 10		Flucloxin Flucloxin
Inj 500 mg vialInj 1 g vial – Up to 5 inj available on a PSO		5		Flucioxin Flucil
ing i g viai – up to 5 ing available on a F30	0.00	J	•	<u>ı ıucıı</u>

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	
PHENOXYMETHYLPENICILLIN (PENICILLIN V) Cap 250 mg – Up to 30 cap available on a PSO Cap 500 mg		50 50		Cilicaine VK Cilicaine VK
a) Up to 20 cap available on a PSO b) Up to 2 x the maximum PSO quantity for RFPP Grans for oral liq 125 mg per 5 ml	5.75	100 ml	✓	AFT
b) Wastage claimable c) AFT to be Principal Supply on 1 February 2026 Grans for oral liq 250 mg per 5 ml	5.89	100 ml	•	AFT
c) Wastage claimable d) AFT to be Principal Supply on 1 February 2026				

Tetracyclines

DO	XYCYCLINE			
*	Tab 100 mg - Up to 30 tab available on a PSO	64.43	500	Doxine
MIN	NOCYCLINE HYDROCHLORIDE			
*	Tab 50 mg - Additional subsidy by Special Authority see			
	SA1355 below – Retail pharmacy	5.79	60	
		(12.05)		Mino-tabs
*	Cap 100 mg	19.32	100	
		(52.04)		Minomycin

⇒SA1355 Special Authority for Manufacturers Price

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has rosacea.

⇒SA2513 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 For the eradication of helicobacter pylori following unsuccessful treatment with appropriate first-line therapy; and
- 2 For use only in combination with bismuth as part of a quadruple therapy regimen.

Renewal from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 For the eradication of helicobacter pylori following unsuccessful treatment with, or noncompletion of second line therapy; and
- 2 For use only in combination with bismuth as part of a quadruple therapy regimen.

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	Brand or Generic Manufacturer
Other Antibiotics	<u>.</u>			
For topical antibiotics, refer to DERMATOLOGICALS, page 67				
CIPROFLOXACIN Recommended for patients with any of the following: i) microbiologically confirmed and clinically significant ps ii) prostatitis: or	eudomonas infection;	or		
iii) pyelonephritis; or iv) gonorrhoea.				
Tab 250 mg - Up to 5 tab available on a PSO	1.95	28	✓ <u>I</u> p	oca-Ciprofloxacin
Tab 500 mg - Up to 5 tab available on a PSO Tab 750 mg		28 28	-	oca-Ciprofloxacin
CLINDAMYCIN	4.00	20	• 17	oca-cipronoxaciii
Cap hydrochloride 150 mg	4.94	24	✓ D	alacin C
Inj 150 mg per ml, 4 ml ampoule		10		lameln Palacin C
Dalacin C to be Principal Supply on 1 March 2026 (Hameln Inj 150 mg per ml, 4 ml ampoule to be delisted 1 March	2026)			
COLISTIN SULPHOMETHATE – Retail pharmacy-Specialist – Only if prescribed for dialysis or cystic fibrosis patient and th			accordingly.	
Inj 2 million iu, 10 ml vial	216.67	10	✓ C	colomycin S29
GENTAMICIN SULPHATE				
Inj 40 mg per ml, 2 ml vial - Subsidy by endorsement	36.70	5	✓ C	idomycin P/Free S29
Only if prescribed for a dialysis or cystic fibrosis patient endorsed accordingly.	or complicated urinary	trac	t infection an	nd the prescription is
Inj 10 mg per ml, 1 ml ampoule – Subsidy by endorsement. Only if prescribed for a dialysis or cystic fibrosis patient		5 trac	_	BL Gentamicin and the prescription is
endorsed accordingly.				
Inj 40 mg per ml, 2 ml ampoule - Subsidy by endorsement.	18.38	10	✓ P	fizer

Inj 40 mg per ml, 2 ml ampoule – Subsidy by endorsement......18.38

Only if prescribed for a dialysis or cystic fibrosis patient or complicated urinary tract infection and the prescription is endorsed accordingly.

MOXIFLOXACIN - Special Authority see SA1740 below - Retail pharmacy

No patient co-payment payable

Tab 400 mg42.00 5 ✓ Avelox

⇒SA1740 Special Authority for Subsidy

Initial application — (Tuberculosis) only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 Active tuberculosis*; and
 - 1.2 Any of the following:
 - 1.2.1 Documented resistance to one or more first-line medications; or
 - 1.2.2 Suspected resistance to one or more first-line medications (tuberculosis assumed to be contracted in an area with known resistance), as part of regimen containing other second-line agents; or

(manufacturer's Price) Subsidised Generic \$ Per ✓ Manufacturer		Subsidy (Manufacturer's Price) \$		Fully bsidised	Brand or Generic Manufacturer	
--	--	---	--	----------------	-------------------------------------	--

continued...

- 1.2.3 Impaired visual acuity (considered to preclude ethambutol use); or
- 1.2.4 Significant pre-existing liver disease or hepatotoxicity from tuberculosis medications; or
- 1.2.5 Significant documented intolerance and/or side effects following a reasonable trial of first-line medications:
- 2 Mycobacterium avium-intracellulare complex not responding to other therapy or where such therapy is contraindicated.*; or
- 3 Patient is under five years of age and has had close contact with a confirmed multi-drug resistant tuberculosis case.

Note: Indications marked with * are unapproved indications.

Renewal only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Mycoplasma genitalium) only from a sexual health specialist or Practitioner on the recommendation of a sexual health specialist. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Has nucleic acid amplification test (NAAT) confirmed Mycoplasma genitalium* and is symptomatic; and
- 2 Either:
 - 2.1 Has tried and failed to clear infection using azithromycin; or
 - 2.2 Has laboratory confirmed azithromycin resistance; and
- 3 Treatment is only for 7 days.

Initial application — (Penetrating eye injury) only from an ophthalmologist. Approvals valid for 1 month where the patient requires prophylaxis following a penetrating eye injury and treatment is for 5 days only.

Note: Indications marked with * are unapproved indications.

PAROMOMYCIN - Special Authority see SA1689 below - Retail pharmacy Cap 250 mg......126.00

16

✓ Humatin S29

⇒SA1689 Special Authority for Subsidy

Initial application only from an infectious disease specialist, clinical microbiologist or gastroenterologist. Approvals valid for 1 month for applications meeting the following criteria:

Fither:

- 1 Patient has confirmed cryptosporidium infection; or
- 2 For the eradication of Entamoeba histolytica carriage.

Renewal only from an infectious disease specialist, clinical microbiologist or gastroenterologist. Approvals valid for 1 month for applications meeting the following criteria:

Either:

- 1 Patient has confirmed cryptosporidium infection; or
- 2 For the eradication of Entamoeba histolytica carriage.

SODIUM FUSIDATE (FUSIDIC ACID)

Tab 250 mg	135.70	36	Fucidin
SULFADIAZINE SODIUM - Special Authority see SA1331 belo	ow – Retail pharmacy		
Tab 500 mg	150.70	100	✓ Sulfadiazin-Heyl S29

⇒SA1331 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 For the treatment of toxoplasmosis in patients with HIV for a period of 3 months; or
- 2 For pregnant patients for the term of the pregnancy; or
- 3 For infants with congenital toxoplasmosis until 12 months of age.

	Subsidy (Manufacturer's Pric \$	e) Sub	sidised	Brand or Generic Manufacturer
TOBRAMYCIN				
Inj 40 mg per ml, 2 ml vial - Subsidy by endorsement	15.50	5	✓ <u>Tob</u>	oramycin (Viatris)
Only if prescribed for dialysis or cystic fibrosis patient ar	nd the prescription is	s endorsed a	accordingl	у.
Solution for inhalation 60 mg per ml, 5 ml - Subsidy by				
endorsement	395.00	56 dose	✓ <u>Tok</u>	ramycin BNM
a) Wastage claimable				
b) Only if prescribed for a cystic fibrosis patient and the	prescription is ende	orsed accor	dingly.	
TRIMETHOPRIM				
* Tab 300 mg - Up to 30 tab available on a PSO	27.83	50	✓ TM	<u>P</u>
TRIMETHOPRIM WITH SULPHAMETHOXAZOLE [CO-TRIMO)	(AZOLFI			
* Tab trimethoprim 80 mg and sulphamethoxazole 400 mg -	•			
to 30 tab available on a PSO		500	✓ Tris	sul
* Oral liq 8 mg sulphamethoxazole 40 mg per ml - Up to 200				
available on a PSO		100 ml	✓ Der	orim
VANCOMYCIN - Subsidy by endorsement				
Only if prescribed for a dialysis or cystic fibrosis patient or fo	or prophylavic of and	locarditic or	for treatm	ant of Clastridium
difficile following metronidazole failure and the prescription is			ioi iicaiiii	Cit of Olostifatain
Inj 500 mg vial		יעיפי. 1	✓ Mvl	an
11, 000 11g 10a		•	. —	ncomycin Viatris
(Mylan Inj 500 mg vial to be delisted 1 March 2026)			<u>, u</u>	,

Antifungals

- a) For topical antifungals refer to DERMATOLOGICALS, page 68
- b) For topical antifungals refer to GENITO URINARY, page 79

FLUCONAZOLE

Cap 50 mg4.10	28	✓ Mylan
Cap 150 mg	1	✓ Mylan
Cap 200 mg8.90	28	✓ Mylan
Powder for oral suspension 10 mg per ml - Special Authority		
see SA1359 below – Retail pharmacy129.02	35 ml	Diflucan
Wastage claimable		

⇒SA1359 Special Authority for Subsidy

Initial application — **(Systemic candidiasis)** from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

Initial application — (Immunocompromised) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Patient is at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

Renewal — (Systemic candidiasis) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

Renewal — (Immunocompromised) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient remains immunocompromised: and
- 2 Patient remains at moderate to high risk of invasive fungal infection; and

Tab 200 mg - PCT......CBS

3 Patient is unable to swallow capsules.

ITRACONAZOLE

Cap 100 mg	6.83	15	Itraconazole
, •			Cresent S29
			✓ Itrazole
Oral liq 10 mg per ml - Special Authority see SA1322 below -			
Retail pharmacy14	1.80	150 ml OP	Itraconazole
			Kent S29

⇒SA1322 Special Authority for Subsidy

Initial application only from an infectious disease specialist, clinical microbiologist, clinical immunologist or any relevant practitioner on the recommendation of a infectious disease physician, clinical microbiologist or clinical immunologist. Approvals valid for 6 months where the patient has a congenital immune deficiency.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefitting from the treatment.

KETOCONAZOLE

		100	✓ Strides Shasun \$29 ✓ Taro \$29 ✓ Teva- Ketoconazole \$29
NYSTATIN			
Tab 500,000 u	14.16	50	
	(17.09)		Nilstat
Cap 500,000 u	12.81	50	
	(15.47)		Nilstat
POSACONAZOLE - Special Authority see SA2383 below - Re	etail pharmacy		
Tab modified-release 100 mg	123.60	24	✓ Posaconazole Juno
Oral lig 40 mg per ml	308.26	105 ml OP	✓ Devatis

⇒SA2383 Special Authority for Subsidy

Initial application only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

Either:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation chemotherapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppressive therapy*.

Renewal only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the

continued...

✓ Burel S29

continued...

following criteria:

Either:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation therapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppression* and requires on going posaconazole treatment.

Note: * Graft versus host disease (GVHD) on significant immunosuppression is defined as acute GVHD, grade II to IV, or extensive chronic GVHD, or if they were being treated with intensive immunosuppressive therapy consisting of either high-dose corticosteroids (1 mg or greater per kilogram of body weight per day for patients with acute GVHD or 0.8 mg or greater per kilogram every other day for patients with chronic GVHD), antithymocyte globulin, or a combination of two or more immunosuppressive agents or types of treatment.

Initial application — (Invasive fungal infection prophylaxis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient is at risk of invasive fungal infection; and
- 2 Either:
 - 2.1 Posaconazole is prescribed by, or recommended by a haematologist, transplant physician, infectious disease specialist, paediatric haematologist or paediatric oncologist; or
 - 2.2 Prescribing posaconazole is in accordance with a protocol or guideline that has been endorsed by the Health New Zealand - Te Whatu Ora Hospital in the specific settings where there is a greater than 10% risk of invasive fungal infection (IFI).

Renewal — (Invasive fungal infection prophylaxis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

_ _ _

- 1 The patient is at risk of invasive fungal infection; and
- 2 Either:
 - 2.1 Posaconazole is prescribed by, or recommended by a haematologist, transplant physician, infectious disease specialist, paediatric haematologist or paediatric oncologist; or
 - 2.2 Prescribing posaconazole is in accordance with a protocol or guideline that has been endorsed by the Health New Zealand - Te Whatu Ora Hospital in the specific settings where there is a greater than 10% risk of invasive fungal infection (IFI).

TERBINAFINE

* Tab 250 mg	8.97	84	✓ Deolate
VORICONAZOLE - Special Authority see SA2384 below - F	Retail pharmacy		
Tab 50 mg	71.00	56	✓ Vttack
Tab 200 mg	263.00	56	✓ Vttack
Powder for oral suspension 40 mg per ml – Wastage			
claimable	1,523.22	70 ml	✓ Vfend

⇒SA2384 Special Authority for Subsidy

Initial application — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
 - 3.1 Patient has proven or probable invasive aspergillus infection; or

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 3.2 Patient has possible invasive aspergillus infection; or
- 3.3 Patient has fluconazole resistant candidiasis; or
- 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Renewal — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
 - 3.1 Patient continues to require treatment for proven or probable invasive aspergillus infection; or
 - 3.2 Patient continues to require treatment for possible invasive aspergillus infection; or
 - 3.3 Patient has fluconazole resistant candidiasis; or
 - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Initial application — (Invasive fungal infection prophylaxis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient is at risk of invasive fungal infection; and
- 2 Either:
 - 2.1 Voriconazole is prescribed by, or recommended by a haematologist, transplant physician, infectious disease specialist, paediatric haematologist or paediatric oncologist; or
 - 2.2 Prescribing voriconazole is in accordance with a protocol or guideline that has been endorsed by the Health New Zealand - Te Whatu Ora Hospital in the specific settings where there is a greater than 10% risk of invasive fungal infection (IFI)

Renewal — (Invasive fungal infection prophylaxis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

.

- 1 The patient is at risk of invasive fungal infection; and
- 2 Either:
 - 2.1 Voriconazole is prescribed by, or recommended by a haematologist, transplant physician, infectious disease specialist, paediatric haematologist or paediatric oncologist; or
 - 2.2 Prescribing voriconazole is in accordance with a protocol or guideline that has been endorsed by the Health New Zealand - Te Whatu Ora Hospital in the specific settings where there is a greater than 10% risk of invasive fungal infection (IFI).

Antimalarials

⇒SA1684 Special Authority for Subsidy

Initial application only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 The patient has vivax or ovale malaria; and
- 2 Primaquine is to be given for a maximum of 21 days.

Renewal only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting

Subsidy (Manufacturer's Price)	Subsic	Fully lised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

the following criteria:

Both:

- 1 The patient has relapsed vivax or ovale malaria; and
- 2 Primaquine is to be given for a maximum of 21 days.

Antitrichomonal Agents

METRONIDAZOLE			
Tab 200 mg - Up to 30 tab available on a PSO	25.86	250	✓ <u>Metronidamed</u>
Tab 400 mg - Up to 15 tab available on a PSO	4.29	21	✓ <u>Metronidamed</u>
Oral liq benzoate 200 mg per 5 ml	25.00	100 ml	✓ Flagyl-S
Suppos 500 mg	24.48	10	✓ Flagyl
ORNIDAZOLE			
Tab 500 mg	36.52	10	✓ <u>Arrow-Ornidazole</u>

Antituberculotics and Antileprotics

Note: There is no co-payment charge for all pharmaceuticals listed in the Antituberculotics and Antileprotics group regardless of immigration status.

BEDAQUILINE - Special Authority see SA2244 below - Retail pharmacy

No patient co-payment payable

Tab 100mg3,084.51 24 OP ✓ Sirturo

⇒SA2244 Special Authority for Subsidy

Initial application — (multi-drug resistant tuberculosis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The person has multi-drug resistant tuberculosis (MDR-TB); and
- 2 Ministry of Health's Tuberculosis Clinical Network has reviewed the individual case and recommends bedaquiline as part of the treatment regimen.

CLOFAZIMINE - Retail pharmacy-Specialist

- a) No patient co-payment payable
- Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or dermatologist.

CYCLOSERINE - Retail pharmacy-Specialist

- a) No patient co-payment payable
- b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or respiratory physician.

Cap 250 mg.......344.00 60 **✓ Cyclorin** 529

DAPSONE - Retail pharmacy-Specialist

- a) No patient co-payment payable
- Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or dermatologist

Tab 25 mg	100	Dapsone
Tab 100 mg329.50	100	✓ Dapsone

		Subsidy (Manufacture de Dries)		Fully	
		(Manufacturer's Price) \$	Per	Subsidised	Generic Manufacturer
TI	HAMBUTOL HYDROCHLORIDE - Retail pharmacy-Speci	alist			
	a) No patient co-payment payable				
	b) Prescriptions must be written by, or on the recommend	lation of, an infectious d	iseas	e physicia	n, clinical microbiologist
	respiratory physician	05.70	100		FMD Fatal 000
	Tab 100 mg		100	_	EMB Fatol \$29
	Tab 400 mg	49.34	56	•	Myambutol S29
3O	NIAZID – Retail pharmacy-Specialist				
	a) No patient co-payment payableb) Prescriptions must be written by, or on the recommend	lation of an internal ma	dicino	nhycioiar	nandiatrinian clinical
	microbiologist, dermatologist or public health physiciar		JICITIE	priysiciai	i, paediatriciari, ciirlicai
k	Tab 100 mg		100	/	Isoniazid Teva S29
	3	327.41		✓	Noumed Isoniazid
30	NIAZID WITH RIFAMPICIN - Retail pharmacy-Specialist				
	a) No patient co-payment payable				
	b) Prescriptions must be written by, or on the recommend	dation of, an internal me	dicine	physiciar	, paediatrician, clinical
	microbiologist, dermatologist or public health physiciar			_	
	Tab 100 mg with rifampicin 150 mg		100		Rifinah
	Tab 150 mg with rifampicin 300 mg		100		Rifinah
	Cap 100 mg with rifampicin 150 mg		100	•	Rifamazid S29
IN	EZOLID - Special Authority see SA2234 below - Retail pl	narmacy			
	No patient co-payment payable	104.60	10	./	7.0.44
	Tab 600 mg Oral lig 20 mg per ml		10 I50 m		Zyvox Zyvox
_			100 11		Lyvox
	SA2234 Special Authority for Subsidy				
٠i+	al application (multi-drug recistant tuberculesis) fro	om any rolovant practitio	nor		valid for 10 months for
	al application — (multi-drug resistant tuberculosis) fro	om any relevant practition	ner.	Approvais	valid for 18 months for
pp	lications meeting the following criteria:	om any relevant practitio	ner.	Approvals	valid for 18 months for
pp	lications meeting the following criteria:	, ,	ner.	Approvals	valid for 18 months for
	lications meeting the following criteria: h:	-TB); and			
op	lications meeting the following criteria: h: 1 The person has multi-drug resistant tuberculosis (MDR	-TB); and			
ot	lications meeting the following criteria: h: The person has multi-drug resistant tuberculosis (MDR Ministry of Health's Tuberculosis Clinical Network has I	-TB); and eviewed the individual c			
ot	lications meeting the following criteria: h: 1 The person has multi-drug resistant tuberculosis (MDR 2 Ministry of Health's Tuberculosis Clinical Network has a the treatment regimen.	-TB); and eviewed the individual c			
ot	lications meeting the following criteria: h: 1 The person has multi-drug resistant tuberculosis (MDR 2 Ministry of Health's Tuberculosis Clinical Network has a the treatment regimen. RA-AMINO SALICYLIC ACID — Retail pharmacy-Specialis a) No patient co-payment payable b) Prescriptions must be written by, or on the recommend	-TB); and eviewed the individual o	ase a	and recom	mends linezolid as part
op ot	lications meeting the following criteria: h: 1 The person has multi-drug resistant tuberculosis (MDR 2 Ministry of Health's Tuberculosis Clinical Network has a the treatment regimen. RA-AMINO SALICYLIC ACID — Retail pharmacy-Specialis a) No patient co-payment payable b) Prescriptions must be written by, or on the recommend respiratory physician	-TB); and reviewed the individual of t dation of, an infectious d	ase a	and recom	mends linezolid as part
op ot	lications meeting the following criteria: h: 1 The person has multi-drug resistant tuberculosis (MDR 2 Ministry of Health's Tuberculosis Clinical Network has a the treatment regimen. RA-AMINO SALICYLIC ACID — Retail pharmacy-Specialis a) No patient co-payment payable b) Prescriptions must be written by, or on the recommend respiratory physician Grans for oral liq 4 g sachet	-TB); and reviewed the individual of t dation of, an infectious d	ase a	and recom	mends linezolid as part
pp ot	lications meeting the following criteria: h: 1 The person has multi-drug resistant tuberculosis (MDR 2 Ministry of Health's Tuberculosis Clinical Network has a the treatment regimen. RA-AMINO SALICYLIC ACID — Retail pharmacy-Specialis a) No patient co-payment payable b) Prescriptions must be written by, or on the recomment respiratory physician Grans for oral liq 4 g sachet DTIONAMIDE — Retail pharmacy-Specialist	-TB); and reviewed the individual of t dation of, an infectious d	ase a	and recom	mends linezolid as part
pp ot	lications meeting the following criteria: h: 1 The person has multi-drug resistant tuberculosis (MDR 2 Ministry of Health's Tuberculosis Clinical Network has a the treatment regimen. RA-AMINO SALICYLIC ACID — Retail pharmacy-Specialis a) No patient co-payment payable b) Prescriptions must be written by, or on the recommend respiratory physician Grans for oral liq 4 g sachet	-TB); and reviewed the individual of t dation of, an infectious d280.00	ase asiseas	e specialis	mends linezolid as part st, clinical microbiologist Paser \$29
op ot	lications meeting the following criteria: h: 1 The person has multi-drug resistant tuberculosis (MDR 2 Ministry of Health's Tuberculosis Clinical Network has a the treatment regimen. RA-AMINO SALICYLIC ACID — Retail pharmacy-Specialis a) No patient co-payment payable b) Prescriptions must be written by, or on the recommend respiratory physician Grans for oral liq 4 g sachet	-TB); and reviewed the individual of t dation of, an infectious d280.00	ase asiseas	e specialis	mends linezolid as part st, clinical microbiologist Paser \$29
ot AF	lications meeting the following criteria: h: 1 The person has multi-drug resistant tuberculosis (MDR 2 Ministry of Health's Tuberculosis Clinical Network has a the treatment regimen. RA-AMINO SALICYLIC ACID — Retail pharmacy-Specialis a) No patient co-payment payable b) Prescriptions must be written by, or on the recommend respiratory physician Grans for oral liq 4 g sachet	-TB); and reviewed the individual of the detection of, an infectious duttion of the detection of	ase asiseas 30	e specialis	mends linezolid as part st, clinical microbiologist Paser \$29 st, clinical microbiologist
op ot AF	lications meeting the following criteria: h: 1 The person has multi-drug resistant tuberculosis (MDR 2 Ministry of Health's Tuberculosis Clinical Network has a the treatment regimen. RA-AMINO SALICYLIC ACID — Retail pharmacy-Specialis a) No patient co-payment payable b) Prescriptions must be written by, or on the recommend respiratory physician Grans for oral liq 4 g sachet	-TB); and reviewed the individual of the detection of, an infectious duttion of the detection of	ase asiseas	e specialis	mends linezolid as part st, clinical microbiologist Paser \$29
AF	lications meeting the following criteria: h: 1 The person has multi-drug resistant tuberculosis (MDR 2 Ministry of Health's Tuberculosis Clinical Network has a the treatment regimen. RA-AMINO SALICYLIC ACID — Retail pharmacy-Specialis a) No patient co-payment payable b) Prescriptions must be written by, or on the recommend respiratory physician Grans for oral liq 4 g sachet	-TB); and reviewed the individual of the detection of, an infectious duttion of the detection of	ase asiseas 30	e specialis	mends linezolid as part st, clinical microbiologist Paser \$29 st, clinical microbiologist
pp ot Al	lications meeting the following criteria: h: 1 The person has multi-drug resistant tuberculosis (MDR 2 Ministry of Health's Tuberculosis Clinical Network has a the treatment regimen. RA-AMINO SALICYLIC ACID — Retail pharmacy-Specialis a) No patient co-payment payable b) Prescriptions must be written by, or on the recommend respiratory physician Grans for oral liq 4 g sachet	-TB); and reviewed the individual of t dation of, an infectious d280.00 dation of, an infectious d305.00	iseas 30 iseas	e specialis e specialis	mends linezolid as part st, clinical microbiologist Paser \$29 st, clinical microbiologist Peteha \$29
pp ot Al	lications meeting the following criteria: h: 1 The person has multi-drug resistant tuberculosis (MDR 2 Ministry of Health's Tuberculosis Clinical Network has a the treatment regimen. RA-AMINO SALICYLIC ACID — Retail pharmacy-Specialis a) No patient co-payment payable b) Prescriptions must be written by, or on the recommend respiratory physician Grans for oral liq 4 g sachet	-TB); and reviewed the individual of t dation of, an infectious d280.00 dation of, an infectious d305.00	iseas 30 iseas	e specialis e specialis	mends linezolid as part st, clinical microbiologist Paser \$29 st, clinical microbiologist Peteha \$29
pp ot Al Yi	lications meeting the following criteria: h: 1 The person has multi-drug resistant tuberculosis (MDR 2 Ministry of Health's Tuberculosis Clinical Network has a the treatment regimen. RA-AMINO SALICYLIC ACID — Retail pharmacy-Specialis a) No patient co-payment payable b) Prescriptions must be written by, or on the recommend respiratory physician Grans for oral liq 4 g sachet	-TB); and reviewed the individual of the lation of, an infectious dution	iseas 30 iseas	e specialis e specialis e physicia	mends linezolid as part st, clinical microbiologist Paser \$29 st, clinical microbiologist Peteha \$29

	INFECTIONS - A	ACENTS.	EOR 9	SVSTEMIC LISE
	Subsidy (Manufacturer's Price		Fully sidised	Brand or Generic Manufacturer
RIFABUTIN - Retail pharmacy-Specialist				
a) No patient co-payment payable b) Prescriptions must be written by, or on the recomme gastroenterologist	ndation of, an infectious	disease ph	ysician,	respiratory physician or
# Cap 150 mg	353.71	30	✓ N	lycobutin
RIFAMPICIN – Subsidy by endorsement				
 a) No patient co-payment payable b) For confirmed recurrent Staphylococcus aureus inferantimicrobial based on susceptibilities and the prescine Retail pharmacy - Specialist. Specialist must be an paediatrician, or public health physician. 	ription is endorsed accor	dingly; can	be waiv	ved by endorsement -
★ Cap 150 mg	58.54	100	_	<u>lifadin</u>
≮ Cap 300 mg	122.06	100	_	<u>lifadin</u>
♦ Oral liq 100 mg per 5 ml	12.60	60 ml		ifadin Sanofi <u>ifadin</u>
Antivirals				
or eye preparations refer to Eye Preparations, Anti-Infective	e Preparations, page 27	4		
Hepatitis B Treatment				
NTECAVIR * Tab 0.5 mg	10.04	30	./ =	ntecavir (Rex)
· ·		30	▼ ⊑	illecavii (nex)
AMIVUDINE - Special Authority see SA1685 below - Reta Tab 100 mg		28	17	etlam
Oral lig 5 mg per ml		40 ml OP		effix
⇒SA1685 Special Authority for Subsidy				
nitial application only from a relevant specialist or medical pprovals valid for 1 year where used for the treatment or pr		nmendatior	of a re	levant specialist.
lenewal from any relevant practitioner. Approvals valid for ENOFOVIR DISOPROXIL	2 years where used for	the treatme	nt or pre	evention of hepatitis B.
Tenofovir disoproxil prescribed under endorsement for the antiretrovirals for the purposes of Special Authority SA2		cluded in th	e count	of up to 4 subsidised
* Tab 245 mg (300 mg as a maleate)		30	√ <u>T</u>	enofovir Disoproxil Viatris

	and the control of the purposes of openial realisting or	.= , page			
*	Tab 245 mg (300 mg as a maleate)	13.80	30	✓ Tenofovir Disopr	oxil
				<u>Viatris</u>	
*	Tab 245 mg (300 mg as a fumarate)	13.80	30	✓ Ricovir S29	

30

✓ Vaclovir

Herpesvirus Treatments		
ACICLOVIR		
* Tab dispersible 200 mg2.05	25	Lovir
Lovir to be Principal Supply on 1 February 2026		
* Tab dispersible 400 mg7.55	56	Lovir
Lovir to be Principal Supply on 1 February 2026		<i>a</i>
* Tab dispersible 800 mg7.43	35	✓ Lovir
Lovir to be Principal Supply on 1 February 2026		
VALACICLOVIR		_
Tab 500 mg	30	✓ Vaclovir

	Subsidy (Manufacturer's Price)	Fully Subsidised		Brand or Generic	
	. \$	Per	1	Manufacturer	
VALGANCICLOVIR – Special Authority see SA2514 below – Retail pharmacy					
Tab 450 mg	140.89	60	✓ <u>v</u>	<u>alganciclovir</u> Viatris	

⇒SA2514 Special Authority for Subsidy

Initial application — (transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 3 months where the patient has undergone a solid organ transplant and requires valganciclovir for CMV prophylaxis.

Renewal — (transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has undergone a solid organ transplant and received anti-thymocyte globulin and requires valganciclovir therapy for CMV prophylaxis: and
 - 1.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following anti-thymocyte globulin; or
- 2 Both:
 - 2.1 Patient has received pulse methylprednisolone for acute rejection and requires further valganciclovir therapy for CMV prophylaxis; and
 - 2.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following pulse methylprednisolone.

Initial application — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 Patient has undergone a solid organ transplant and received valganciclovir under Special Authority more than 2 years ago (27 months): and
- 2 Patient has received anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Renewal — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months where the patient has received a further course of anti-thymocyte globulin and requires valganciclovir for CMV

Initial application — (Lung transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone a lung transplant; and
- 2 Either:
 - 2.1 The donor was cytomegalovirus positive and the patient is cytomegalovirus negative; or
 - 2.2 The recipient is cytomegalovirus positive; and
- 3 Patient has a high risk of CMV disease.

Renewal — (Lung transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone a lung re-transplant; and
- 2 Either:
 - 2.1 The donor was cytomegalovirus positive and the patient is cytomegalovirus negative; or
 - 2.2 The recipient is cytomegalovirus positive: and
- 3 Patient has a high risk of CMV disease.

Initial application — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
 - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or

continued...

- 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
- 2.3 Patient has cytomegalovirus retinitis.

Renewal — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
 - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
 - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
 - 2.3 Patient has cytomegalovirus retinitis.

Note: for the purpose of this Special Authority "immunocompromised" includes transplant recipients, patients with immunosuppressive diseases (e.g. HIV) or those receiving immunosuppressive treatment for other conditions.

Hepatitis C Treatment

GLECAPREVIR WITH PIBRENTASVIR - [Xpharm]

Note the supply of treatment is via Pharmac's approved direct distribution supply. Further details can be found on Pharmac's website https://pharmac.govt.nz/maviret

Tab 100 mg with pibrentasvir 40 mg24,750.00 84 OP ✓ Maviret

LEDIPASVIR WITH SOFOSBUVIR - [Xpharm] - Special Authority see SA1605 below

No patient co-payment payable

Tab 90 mg with sofosbuvir 400 mg......24,363.46 28 **✓ Harvoni**

⇒SA1605 Special Authority for Subsidy

Special Authority approved by the Hepatitis C Treatment Panel (HepCTP)

Notes: By application to the Hepatitis C Treatment Panel (HepCTP).

Applications will be considered by HepCTP and approved subject to confirmation of eligibility.

Application details may be obtained from Pharmac's website http://www.pharmac.govt.nz/harvoni or:

The Coordinator, Hepatitis C Treatment Panel

Pharmac, PO Box 10-254, WELLINGTON Tel: (04) 460 4990,

Email: hepcpanel@pharmac.govt.nz

HIV Prophylaxis and Treatment

EMTRICITABINE WITH TENOFOVIR DISOPROXIL — Subsidy by endorsement; can be waived by Special Authority see SA2520 on the next page

- a) Funding for emtricitabine with tenofovir disoproxil for use as PrEP or PEP, should be applied using Special Authority SA2520.
- b) Endorsement for treatment of conditions approved via Special Authority SA2139 (antiretrovirals for confirmed HIV, prevention of maternal transmission, post-exposure prophylaxis following exposure to HIV (when co-prescribed with other antiretrovirals) and percutaneous exposure): Prescription is deemed to be endorsed if emtricitabine with tenofovir disoproxil is co-prescribed with another antiretroviral subsidised under Special Authority SA2139 and the prescription is annotated accordingly by the Pharmacist or endorsed by the prescriber.

Note: Emtricitabine with tenofovir disoproxil prescribed under endorsement, for treatment of conditions approved via Special Authority SA2139 (antiretrovirals for confirmed HIV, prevention of maternal transmission, post-exposure prophylaxis following exposure to HIV and percutaneous exposure), is included in the count of up to 4 subsidised antiretrovirals, and counts as two antiretroviral medications, for the purposes of Special Authority SA2139, page 109 There is an approval process to become a named specialist to prescribe antiretroviral therapy in New Zealand. Further information is available on the Pharmac website.

* Tab 200 mg with tenofovir disoproxil 245	mg (300 mg as a

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

⇒SA2520 Special Authority for Subsidy

Initial application — (Pre-exposure prophylaxis) from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 Patient has tested HIV negative, does not have signs or symptoms of acute HIV infection and has been assessed for HIV seroconversion: and
- 2 The Practitioner considers the patient is at elevated risk of HIV exposure and use of PrEP is clinically appropriate.

Notes: Refer to local health pathways or the Australasian Society for HIV. Viral Hepatitis and Sexual Health Medicine clinical auidelines:

https://ashm.org.au/HIV/PrEP/

Renewal — (Pre-exposure prophylaxis) from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 Patient has tested HIV negative, does not have signs or symptoms of acute HIV infection and has been assessed for HIV seroconversion: and
- 2 The Practitioner considers the patient is at elevated risk of HIV exposure and use of PrEP is clinically appropriate.

Notes: Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines:

https://ashm.org.au/HIV/PrEP/

Initial application — (post-exposure prophylaxis following exposure to HIV) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria: Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had condomless anal intercourse or receptive vaginal intercourse with a known HIV positive person with an unknown or detectable viral load greater than 200 copies per ml; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is appropriate; or
 - 2.4 Patient has had condomless anal intercourse with a person from a high HIV prevalence country or risk group whose HIV status is unknown.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Refer to local health pathways or the Australasian Society for HIV. Viral Hepatitis and Sexual Health Medicine clinical guidelines for PEP (https://www.ashm.org.au/hiv/hiv-management/pep/).

Renewal — (second or subsequent post-exposure prophylaxis) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria: Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had condomless anal intercourse or receptive vaginal intercourse with a known HIV positive person with an unknown or detectable viral load greater than 200 copies per ml; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is appropriate; or
 - 2.4 Patient has had condomless anal intercourse with a person from a high HIV prevalence country or risk group whose HIV status is unknown.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

COVID-19 Treatments

NIRMATRELVIR WITH RITONAVIR - PCT - Subsidy by endorsement

Subsidised for patients meeting access criteria for oral COVID-19 antiviral treatments (as on Pharmac's website) and where the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed when supplying by Direct Provision under the provisions in Part I of Section A of the Pharmaceutical Schedule.

REMDESIVIR - PCT only

Antiretrovirals

⇒SA2139 Special Authority for Subsidy

Initial application — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the patient has confirmed HIV infection.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Prevention of maternal transmission) only from a named specialist. Approvals valid for 1 year for applications meeting the following criteria:

Either:

- 1 Prevention of maternal foetal transmission: or
- 2 Treatment of the newborn for up to eight weeks.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Some antiretrovirals are unapproved or contraindicated for this indication. Practitioners prescribing these medications should exercise their own skill, judgement, expertise and discretion, and make their own prescribing decisions with respect to the use of a Pharmaceutical for an indication for which it is not approved or contraindicated.

Initial application — (post-exposure prophylaxis following exposure to HIV) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had condomless anal intercourse or receptive vaginal intercourse with a known HIV positive person with an unknown or detectable viral load greater than 200 copies per ml; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required; or
 - 2.4 Patient has had condomless anal intercourse with a person from a high HIV prevalence country or risk group whose HIV status is unknown.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals.

	,	ully Brand or	
(Manufactu	ırer's Price) Subsidise	sed Generic Manufacturer	
4	י ו ווי י	Wandadaca	

continued...

Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines for PEP (https://www.ashm.org.au/hiv/hiv-management/pep/).

Renewal — (second or subsequent post-exposure prophylaxis) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had condomless anal intercourse or receptive vaginal intercourse with a known HIV positive person with an unknown or detectable viral load greater than 200 copies per ml; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required; or
 - 2.4 Patient has had condomless anal intercourse with a person from a high HIV prevalence country or risk group whose HIV status is unknown.

Initial application — (**Percutaneous exposure**) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Second or subsequent percutaneous exposure) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Non-nucleosides Reverse Transcriptase Inhibitors

EFAVIRENZ – Special Authority see SA2139 on the previous p Note: No new patients to be initiated on efavirenz.	age – Retail pha	rmacy	
Tab 600 mg	65.38	30	✓ Efavirenz
(Efavirenz Milpharm \$29 Tab 600 mg to be delisted 1 Novemb	er 2026)		Milpharm S29
ETRAVIRINE – Special Authority see SA2139 on the previous	,	armacv	
Tab 200 mg		60	✓ Intelence
NEVIRAPINE - Special Authority see SA2139 on the previous	page – Retail ph	armacy	
Tab 200 mg	198.25	60	✓ Nevirapine Viatris
Oral suspension 10 mg per ml	203.55	240 ml OP	✓ Viramune Suspension

Nucleosides Reverse Transcriptase Inhibitors

ABACAVIR SULPHATE - Special Authority see SA2139 on the pre	evious page – F	Retail pharma	acy
Tab 300 mg	180.00	60	✓ Ziagen
ABACAVIR SULPHATE WITH LAMIVUDINE – Special Authority son Note: abacavir with lamivudine (combination tablets) counts as			
anti-retroviral Special Authority.			
Tab 600 mg with lamivudine 300 mg	35.00	30	Abacavir/ Lamivudine

Abacavir/Lamivudine Viatris to be Principal Supply on 1 February 2026

Viatris

		osidy urer's Price) \$	F Subsidi Per	ully ised	Brand or Generic Manufacturer
EFAVIRENZ WITH EMTRICITABINE AND TENOFOVIR DISOP pharmacy Note: Efavirenz with emtricitabine and tenofovir disoproxil of		•	•		
anti-retroviral Special Authority Tab 600 mg with emtricitabine 200 mg and tenofovir disopro					
245 mg (300 mg as a fumarate)		88	30		TEEVIR S29 Triovir S29
Tab 600 mg with emtricitabine 200 mg and tenofovir disopro	106.		30		Viatris
(Triovir \$29 Tab 600 mg with emtricitabine 200 mg and tenofor January 2026)	·		ou mg as a	TUM	arate) to de delisted i
EMTRICITABINE - Special Authority see SA2139 on page 109 Cap 200 mg LAMIVUDINE - Special Authority see SA2139 on page 109 - R	307.	20	30	•	Emtriva
Tab 150 mg Oral liq 10 mg per ml	98.	00	60 O ml OP	_ '	Lamivudine Viatris 3TC
ZIDOVUDINE [AZT] – Special Authority see SA2139 on page 1 Cap 100 mg	<mark>09</mark> – Retail	pharmacy	100		Retrovir
Oral liq 10 mg per ml	30.	45 200	ml OP		Retrovir nacv
Note: zidovudine [AZT] with lamivudine (combination tablet the anti-retroviral Special Authority.	s) counts a	s two anti-re		dica	tions for the purposes of
Tab 300 mg with lamivudine 150 mg	92.	40	60	•	Lamivudine/ Zidovudine Viatris
Protease Inhibitors					
ATAZANAVIR SULPHATE – Special Authority see SA2139 on Cap 150 mg	102.		macy 60	•	Atazanavir Viatris
Atazanavir Viatris to be Principal Supply on 1 February Cap 200 mg	152.	30	60	1	Atazanavir Viatris
DARUNAVIR – Special Authority see SA2139 on page 109 – R Tab 400 mg	etail pharm		60		Darunavir Viatris
Tab 600 mg LOPINAVIR WITH RITONAVIR – Special Authority see SA2138	on page 1	09 – Retail			Darunavir Viatris
Tab 200 mg with ritonavir 50 mg			120		Lopinavir/Ritonavir Mylan
RITONAVIR – Special Authority see SA2139 on page 109 – Re Tab 100 mg		•	30	•	Norvir
Strand Transfer Inhibitors					
DOLUTEGRAVIR - Special Authority see SA2139 on page 109 Tab 50 mg	1,090.	00	30		Tivicay
DOLUTEGRAVIR WITH LAMIVUDINE - Special Authority see Tab 50 mg with lamivudine 300 mg			- Retail phai 30	_	cy Dovato
RALTEGRAVIR POTASSIUM – Special Authority see SA2139 (Tab 400 mg Tab 600 mg	1,090.	00	harmacy 60 60		Isentress Isentress HD

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

Subsidy	Full	y Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per 🗸	Manufacturer	

Immune Modulators

PEGYLATED INTERFERON ALFA-2A - Special Authority see SA2034 below - Retail pharmacy

Note: Pharmac will consider funding ribavirin for the small group of patients who have a clinical need for ribavirin and meet Special Authority criteria. Please contact the Hepatitis C Coordinator at Pharmac on 0800-023-588 option 4.

1.355.71

✓ Pegasys S29 S29

Inj 135 mcg prefilled syringe......887.35 ✓ Pegasys (S29) S29 Inj 180 mcg prefilled syringe.......748.50 ✓ Pegasys

⇒SA2034 Special Authority for Subsidy

Initial application — (chronic hepatitis C - genotype 1, 4, 5 or 6 infection or co-infection with HIV or genotype 2 or 3 post liver transplant) from any specialist. Approvals valid for 18 months for applications meeting the following criteria: Both:

- 1 Any of the following:
 - 1.1 Patient has chronic hepatitis C, genotype 1, 4, 5 or 6 infection; or
 - 1.2 Patient has chronic hepatitis C and is co-infected with HIV; or
 - 1.3 Patient has chronic hepatitis C genotype 2 or 3 and has received a liver transplant; and
- 2 Maximum of 48 weeks therapy.

Renewal — (Chronic hepatitis C - genotype 1 infection) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria: All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Fither:
 - 3.1 Patient has responder relapsed; or
 - 3.2 Patient was a partial responder; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application — (Chronic Hepatitis C - genotype 1 infection treatment more than 4 years prior) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Any of the following:
 - 3.1 Patient has responder relapsed; or
 - 3.2 Patient was a partial responder; or
 - 3.3 Patient received interferon treatment prior to 2004; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application — (chronic hepatitis C - genotype 2 or 3 infection without co-infection with HIV) from any specialist. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 Patient has chronic hepatitis C, genotype 2 or 3 infection; and
- 2 Maximum of 6 months therapy.

Initial application — (Hepatitis B) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

Subsidy	Full	/ Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per 🗸	Manufacturer Manufacturer	

continued...

- 1 Patient has confirmed Hepatitis B infection (HBsAg positive for more than 6 months); and
- 2 Patient is Hepatitis B treatment-naive; and
- 3 ALT > 2 times Upper Limit of Normal; and
- 4 HBV DNA < 10 log10 IU/ml; and
- 5 Either:
 - 5.1 HBeAg positive; or
 - 5.2 serum HBV DNA greater than or equal to 2,000 units/ml and significant fibrosis (Metavir Stage F2 or greater or moderate fibrosis); and
- 6 Compensated liver disease; and
- 7 No continuing alcohol abuse or intravenous drug use; and
- 8 Not co-infected with HCV, HIV or HDV; and
- 9 Neither ALT nor AST > 10 times upper limit of normal; and
- 10 No history of hypersensitivity or contraindications to pegylated interferon; and
- 11 Maximum of 48 weeks therapy.

Initial application — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 Patient has a cutaneous T cell lymphoma*: or
- 2 All of the following:
 - 2.1 Patient has a myeloproliferative disorder*; and
 - 2.2 Patient is intolerant of hydroxyurea; and
 - 2.3 Treatment with anagrelide and busulfan is not clinically appropriate; or
- 3 Both:
 - 3.1 Patient has a myeloproliferative disorder; and
 - 3.2 Patient is pregnant, planning pregnancy or lactating.

Renewal — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment; and
- 3 Either:
 - 3.1 Patient has a cutaneous T cell lymphoma*: or
 - 3.2 Both:
 - 3.2.1 Patient has a myeloproliferative disorder*; and
 - 3.2.2 Fither:
 - 3.2.2.1 Remains intolerant of hydroxyurea and treatment with anagrelide and busulfan remains clinically inappropriate; or
 - 3.2.2.2 Patient is pregnant, planning pregnancy or lactating.

Note: Indications marked with * are unapproved indications.

Initial application — (post-allogenic bone marrow transplant) from any relevant practitioner. Approvals valid for 3 months where patient has received an allogeneic bone marrow transplant* and has evidence of disease relapse.

Renewal — (post-allogenic bone marrow transplant) from any relevant practitioner. Approvals valid for 3 months where patient is responding and ongoing treatment remains appropriate.

Note: Indications marked with * are unapproved indications.

Urinary Tract Infections

✓ UroFos

Subsidy Fully		Brand or	
(Manufacturer's Price)	Subsidised		Generic
\$	Per	•	Manufacturer

⇒SA2406 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 months for applications meeting the following criteria: Both:

- 1 Patient has an acute, symptomatic, bacteriologically-proven uncomplicated urinary tract infection (UTI)/cystitis with Escherichia Coli; and
- 2 Either:
 - 2.1 Microbiological testing confirms the pathogen is resistant to all of: trimethoprim, nitrofurantoin, amoxicillin, cefaclor, cefalexin, amoxicillin with clavulanic acid, and norfloxacin; or
 - 2.2 The patient has a contraindication or intolerance to all of: trimethoprim, nitrofurantoin, amoxicillin, cefaclor, cefalexin, amoxicillin with clavulanic acid, and norfloxacin that the pathogen is susceptible to.

Renewal from any relevant practitioner. Approvals valid for 2 months for applications meeting the following criteria: Both:

- 1 Patient has an acute, symptomatic, bacteriologically-proven uncomplicated urinary tract infection (UTI)/cystitis with Escherichia Coli; and
- 2 Either:
 - 2.1 Microbiological testing confirms the pathogen is resistant to all of: trimethoprim, nitrofurantoin, amoxicillin, cefaclor, cefalexin, amoxicillin with clavulanic acid, and norfloxacin; or
 - 2.2 The patient has a contraindication or intolerance to all of: trimethoprim, nitrofurantoin, amoxicillin, cefaclor, cefalexin, amoxicillin with clavulanic acid, and norfloxacin that the pathogen is susceptible to.

*	Tab 1 g	19.95	100	✓ Hiprex
NΙΊ	ROFURANTOIN			
*	Tab 50 mg - Up to 30 tab available on a PSO	22.20	100	✓ Nifuran
*	Tab 100 mg	37.50	100	✓ Nifuran
*	Cap modified-release 100 mg - Up to 15 cap available on a			
	PSO	81.20	100	✓ <u>Macrobid</u>
NO	RFLOXACIN			
	Tab 400 mg - Subsidy by endorsement	245.00	100	✓ Arrow-Norfloxacin

Only if prescribed for a patient with an uncomplicated urinary tract infection that is unresponsive to a first line agent or with proven resistance to first line agents and the prescription is endorsed accordingly.

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	
	\$	Per		Manufacturer
nticholinesterases				
mucholinesterases				
OSTIGMINE METILSULFATE				
Inj 2.5 mg per ml, 1 ml ampoule	48.25	10	1	Max Health
RIDOSTIGMINE BROMIDE				
Tab 60 mg	50.28	100	1	Mestinon
145 00 mg		100		mootmon
lon-Steroidal Anti-Inflammatory Drugs				
CLOFENAC SODIUM			_	
Tab EC 25 mg		50		Diclofenac Sandoz
Tab 50 mg dispersible		20		Voltaren D
Tab EC 50 mg		50		Diclofenac Sandoz
Tab long-acting 75 mg		100		Voltaren SR
Inj 25 mg per ml, 3 ml ampoule – Up to 5 inj available on a	PSO 13.20	5		Voltaren
Suppos 12.5 mg	2.04	10		Voltaren
Suppos 25 mg		10		Voltaren
Suppos 50 mg - Up to 10 supp available on a PSO	4.22	10		Voltaren
Suppos 100 mg	7.00	10	•	Voltaren
JPROFEN				
Tab 200 mg	21.40	1.000	1	Relieve
Tab long-acting 800 mg		30	1	Ibuprofen SR BNM
Oral lig 20 mg per ml		200 m		Ethics
TOPROFEN				
Cap long-acting 200 mg	12.07	28	1	Oruvail SR
	12.07	20	•	Oluvali Sh
FENAMIC ACID				
Cap 250 mg		50		_
	(10.82)			Ponstan
	0.50	20		_
	(7.50)			Ponstan
PROXEN				
Tab 250 mg	39.23	500	1	Noflam 250
Tab 500 mg	34.45	250	1	Noflam 500
Tab long-acting 750 mg	10.40	28	1	Naprosyn SR 750
Tab long-acting 1 g	11.50	28		Naprosyn SR 1000
NOXICAM				
Tab 20 mg	23 50	100	1	Tilcotil
Tilcotil to be Principal Supply on 1 February 2026	20.00	100	•	i iiootii
Inj 20 mg vial	9.95	1	1	AFT
11] 20 11g via				Al 1
SAIDs Other				
ELECOXIB				
Cap 100 mg		60		Celecoxib Pfizer
	3.60		•	Celebrex
Celebrex to be Principal Supply on 1 February 2026				
				A
Cap 200 mg	3.20	30		Celebrex Celecoxib Pfizer

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	sidised	Generic	
\$	Per	1	Manufacturer	

✓ Zostriy

Topical Products for Joint and Muscular Pain

CAPSAICIN

Crm 0.025% - Special Authority see SA1289 below - Retail 45 g OP ✓ Zo-Rub Osteo pharmacy......9.75

⇒SA1289 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has osteoarthritis that is not responsive to paracetamol and oral non-steroidal anti-inflammatories are contraindicated.

Antirheumatoid Agents

HADDOAACHI ODOOTIIVE GLII BRIVEE

7.80	100	✓ <u>Ipca-</u> <u>Hydroxychloroquine</u>
6.00	30	✓ Arava
	30	✓ Arava
67.23	100	✓ D-Penamine
110.12	100	✓ D-Penamine
		6.00 30 30 30 30 30 30 30 30 30 30 30 30 3

Drugs Affecting Bone Metabolism

Alendronate for Osteoporosis

ALENDRONATE SODI	UM
------------------	----

Toh 70 mg

~	7ab 70 mg	7	· I OSalliax
ALI	ENDRONATE SODIUM WITH COLECALCIFEROL		
*	Tab 70 mg with colecalciferol 5,600 iu	4	✓ Fosamax Plus

Other Treatments

DENOSUMAB - Special Authority see SA2441 below - Retail pharmacy

Note: Denosumab ini 60 mg per 1 ml pre-filled syringe is Medsafe approved for use in osteoporosis. Denosumab ini 120 mg per 1.7 ml vial is Medsafe approved for use in hypercalcaemia of malignancy.

3 10

Inj 120 mg per 1.7 ml vial	375.00	1	✓ Xgeva
Inj 60 mg per 1 ml prefilled syringe	e187.50	1	Prolia

⇒SA2441 Special Authority for Subsidy

Initial application — (Osteoporosis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 The patient has established osteoporosis; and
- 2 Any of the following:

✓ fully subsidised

Principal Supply

- 2.1 History of one significant osteoporotic fracture demonstrated radiologically, with a documented T-Score less than or egual to -2.5, that incorporates BMD measured using dual-energy x-ray absorptiometry (DEXA); or
- 2.2 History of one significant osteoporotic fracture, demonstrated radiologically, and either the patient is elderly, or

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic	
\$	Per	•	Manufacturer	

continued...

- densitometry scanning cannot be performed because of logistical, technical or pathophysiological reasons; or
- 2.3 History of two significant osteoporotic fractures demonstrated radiologically; or
- 2.4 Documented T-Score less than or equal to -3.0; or
- 2.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm that incorporates BMD measured using DEXA; and
- 3 Any of the following:
 - 3.1 Bisphosphonates are contraindicated because the patient's creatinine clearance or eGFR is less than 35 mL/min; or
 - 3.2 The patient has experienced at least two symptomatic new fractures or a BMD loss greater than 2% per year, after at least 12 months' continuous therapy with a funded antiresorptive agent; or
 - 3.3 Bisphosphonates result in intolerable side effects; or
 - 3.4 Intravenous bisphosphonates cannot be administered due to logistical or technical reasons.

Initial application — (Hypercalcaemia) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has hypercalcaemia of malignancy; and
- 2 Patient has severe renal impairment.

PAMIDRONATE DISODIUM

Inj 3 mg per ml, 10 ml vial	32.49	1	Pamisol
Inj 6 mg per ml, 10 ml vial	88.11	1	✓ Pamisol
Inj 9 mg per ml, 10 ml vial	94.34	1	✓ Pamisol
RALOXIFENE HYDROCHLORIDE - Special Authority see SA1779		pharmacy	
* Tab 60 mg	53.76	28	Evista

(Evista Tab 60 mg to be delisted 1 April 2026) SA1779 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Notes); or
- 2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
- 3 History of two significant osteoporotic fractures demonstrated radiologically; or
- 4 Documented T-Score less than or equal to -3.0 (see Notes); or
- 5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Notes); or
- 6 Patient has had a Special Authority approval for zoledronic acid (Underlying cause Osteoporosis) or has had a Special Authority approval for alendronate (Underlying cause - Osteoporosis) prior to 1 February 2019.

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
 Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for raloxifene funding.

	Subsidy (Manufacturer's Price) \$	Su Per	Fully bsidised	Brand or Generic Manufacturer
continued c) Osteoporotic fractures are the incident events for severe definitions of osteoporosis and fragility fracture. The WH -2.5 with one or more associated fragility fractures. Fragi forces that would not ordinarily cause fracture (minimal trifall from a standing height or less. d) A vertebral fracture is defined as a 20% or greater reduct relative to the posterior height of that body, or a 20% or g body above or below the affected vertebral body.	O defines severe (est ility fractures are fract auma). The WHO ha ion in height of the an	ablished ures that s quantif terior or) osteopo t occur as fied this a mid porti	orosis as a T-score below a result of mechanical as forces equivalent to a on of a vertebral body
RISEDRONATE SODIUM Tab 35 mgRisedronate Sandoz to be Principal Supply on 1 Februa	3.00 ary 2026	4	√ F	lisedronate Sandoz
TERIPARATIDE – Special Authority see SA1139 below – Retail Inj 250 mcg per ml, 2.4 ml		1	√ T	eriparatide - Teva
All of the following: 1 The patient has severe, established osteoporosis; and 2 The patient has a documented T-score less than or equal 3 The patient has had two or more fractures due to minimal 4 The patient has experienced at least one symptomatic ne funded antiresorptive agent at adequate doses (see Note	I trauma; and w fracture after at lea		onths' cor	ntinuous therapy with a
Notes: a) The bone mineral density (BMD) measurement used to d absorptiometry (DXA). Quantitative ultrasound and quant b) Antiresorptive agents and their adequate doses for the pusodium tab 70 mg or tab 70 mg with colecalciferol 5,600 i zoledronic acid 5 mg per year. If an intolerance of a seve during the use of one antiresorptive agent, an alternate at the minimum requirement of 12 months' continuous thera c) A vertebral fracture is defined as a 20% or greater reduct relative to the posterior height of that body, or a 20% or g body above or below the affected vertebral body. d) A maximum of 18 months of treatment (18 cartridges) will	erive the T-score musitative computed tomurposes of this Specia u once weekly; raloxiverity necessitating perntiresorptive agent mapy.	ography I Authori iene hyd manent ust be tri terior or	(QCT) are ity are de- rochloride treatment alled so to mid porti	e not acceptable fined as: alendronate e tab 60 mg once daily; withdrawal develops hat the patient achieves on of a vertebral body
ZOLEDRONIC ACID Inj 0.05 mg per ml, 100 ml, bag	19.45	1	✓ Z	oledronic Acid Viatris
Zoledronic Acid Viatris to be Principal Supply on 1 February P	uary 2026			
Hyperuricaemia and Antigout				
ALLOPURINOL * Tab 100 mg	17 00	1,000	√ lı	oca-Allopurinol

BENZBROMARONE - Special Authority see SA1963 on the next page - Retail pharmacy Tab 50 mg32.00

500

100

✓ Ipca-Allopurinol

✓ Narcaricin mite S29

Subsid	dy Fi	ılly Brand or	
(Manufacture	r's Price) Subsidis	ed Generic	
\$	Per	 Manufacturer 	

⇒SA1963 Special Authority for Subsidy

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

- 1 The treatment remains appropriate and the patient is benefitting from the treatment; and
- 2 There is no evidence of liver toxicity and patient is continuing to receive regular (at least every three months) liver function tests

COLCHICINE

* Tab 500 mcg	6.00	100	✓ Colgout
FEBUXOSTAT – Special Authority see SA2555 below –	- Retail pharmacy		•
Tab 80 mg	4.73	28	✓ Febuxostat (Teva)
Tab 120 mg	11.78	28	✓ Febuxostat (Teva)

⇒SA2555 Special Authority for Subsidy

Initial application — (Gout) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- Patient has been diagnosed with gout: and
 - 2 Any of the following:
 - 2.1 The patient has a serum urate level greater than 0.36 mmol/l despite treatment with allopurinol at doses of at least 600 mg/day and addition of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.2 The patient has experienced intolerable side effects from allopurinol such that treatment discontinuation is required and serum urate remains greater than 0.36 mmol/l despite use of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.3 The patient has renal impairment such that probenecid is contraindicated or likely to be ineffective and serum urate remains greater than 0.36 mmol/l despite optimal treatment with allopurinol; or
 - 2.4 The patient has previously had an initial Special Authority approval for benzbromarone for treatment of gout...

Initial application — (Tumour lysis syndrome) only from a haematologist or oncologist. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient is scheduled to receive cancer therapy carrying an intermediate or high risk of tumour lysis syndrome; and
- 2 Patient has a documented history of allopurinol intolerance.

Renewal — (Tumour lysis syndrome) only from a haematologist or oncologist. Approvals valid for 6 weeks where the treatment remains appropriate and the patient is benefitting from treatment.

PROBENECID

Muscle Relaxants

BACLOFEN * Tab 10 mg

-1-	1 ab 10 mg		100	· I dolloll
	Inj 0.05 mg per ml, 1 ml ampoule - Subsidy by endorsement	11.55	1	✓ Lioresal Intrathecal
	Subsidised only for use in a programmable pump in patients	where oral ar	ntispastic age	nts have been ineffective or have
	caused intolerable side effects and the prescription is endor	sed according	ly.	
	Ini 2 mg nor ml 5 ml amnoula - Subsidy by andorsament	/Q0 Q1	10	✓ Sintatica Raclofon

Inj 2 mg per ml, 5 ml ampoule − Subsidy by endorsement...........490.91 10 ✓ Sintetica Baclofen Intrathecal

Subsidised only for use in a programmable pump in patients where oral antispastic agents have been ineffective or have caused intolerable side effects and the prescription is endorsed accordingly.

3 70

✓ Dacifon

DANTROLENE

Cap 25 mg145.77	100	✓ Dantrium S29 S29
Cap 50 mg77.00	100	✓ Dantrium

	Subsidy (Manufacturer's Price)	Sub	Fully sidised	Brand or Generic
	\$	Per	•	Manufacturer
ORPHENADRINE CITRATE				
Tab 100 mg	23.25	100	✓ No	orflex

Subsidy (Manufacturer's Price)

Fully Subsidised Per

Brand or Generic Manufacturer

Agents for Parkinsonism and Related Disorders

AMANTADINE HYDROCHLORIDE		
▲ Cap 100 mg38.24	60	✓ Symmetrel
63.73	100	✓ Symmetrel
APOMORPHINE HYDROCHLORIDE		
▲ Inj 10 mg per ml, 2 ml ampoule	5	✓ Movapo
▲ Inj 10 mg per ml, 5 ml ampoule	5	✓ Movapo
ENTACAPONE		·
▲ Tab 200 mg13.73	100	✓ Entacapone Viatris
LEVODOPA WITH BENSERAZIDE		
* Tab dispersible 50 mg with benserazide 12.5 mg	100	✓ Madopar Rapid
* Cap 50 mg with benserazide 12.5 mg	100	✓ Madopar 62.5
* Cap 100 mg with benserazide 25 mg	100	✓ Madopar 125
* Cap long-acting 100 mg with benserazide 25 mg	100	✓ Madopar HBS
* Cap 200 mg with benserazide 50 mg26.25	100	✓ Madopar 250
LEVODOPA WITH CARBIDOPA		•
* Tab 100 mg with carbidopa 25 mg	100	✓ Sinemet
* Tab long-acting 200 mg with carbidopa 50 mg44.99	100	✓ Sinemet CR
* Tab 250 mg with carbidopa 25 mg	100	✓ Sinemet
LEVODOPA WITH CARBIDOPA AND ENTACAPONE		
* Tab 50 mg with carbidopa 12.5 mg and entacapone 200 mg27.01	100	✓ Stalevo
* Tab 100 mg with carbidopa 25 mg and entacapone 200 mg34.18	100	✓ Stalevo
* Tab 150 mg with carbidopa 37.5 mg and entacapone 200 mg44.96	100	✓ Stalevo
* Tab 200 mg with carbidopa 50 mg and entacapone 200 mg51.23	100	✓ Stalevo
PRAMIPEXOLE HYDROCHLORIDE		
▲ Tab 0.25 mg	100	✓ Ramipex
▲ Tab 1 mg17.73	100	✓ Ramipex
RASAGILINE		
* Tab 1 mg53.50	30	✓ Azilect S29
ROPINIROLE HYDROCHLORIDE		
▲ Tab 0.25 mg	84	✓ Ropin
▲ Tab 1 mg	84	✓ Ropin
▲ Tab 2 mg	84	✓ Ropin
▲ Tab 5 mg	84	✓ Ropin
TOLCAPONE		- 1
▲ Tab 100 mg152.38	100	✓ Tasmar
_ 132 133 mg		racinal

Anticholinergics

BENZATROPINE MESYLATE			
Tab 2 mg	10.99	60	✓ Benztrop
Inj 1 mg per ml, 2 ml	95.00	5	✓ Phebra
a) Up to 10 inj available on a PSO			
b) Only on a PSO			

PROCYCLIDINE HYDROCHLORIDE

Tab 5 mg7.40 100 **✓ Kemadrin**



Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

Agents for Essential Tremor, Chorea and Related Disorders

RILUZOLE - Special Authority see SA1403 below - Retail pharmacy

Wastage claimable

Tab 50 mg117.00 56 **✓ Rilutek**

⇒SA1403 Special Authority for Subsidy

Initial application only from a neurologist or respiratory specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has amyotrophic lateral sclerosis with disease duration of 5 years or less; and
- 2 The patient has at least 60 percent of predicted forced vital capacity within 2 months prior to the initial application; and
- 3 The patient has not undergone a tracheostomy; and
- 4 The patient has not experienced respiratory failure; and
- 5 Any of the following:
 - 5.1 The patient is ambulatory; or
 - 5.2 The patient is able to use upper limbs; or
 - 5.3 The patient is able to swallow.

Renewal from any relevant practitioner. Approvals valid for 18 months for applications meeting the following criteria: All of the following:

- 1 The patient has not undergone a tracheostomy; and
- 2 The patient has not experienced respiratory failure; and
- 3 Any of the following:
 - 3.1 The patient is ambulatory; or
 - 3.2 The patient is able to use upper limbs; or
 - 3.3 The patient is able to swallow.

TETRABENAZINE

Anaesthetics

Local

LIDOCAINE [LIGNOCAINE]

Gel 2%, tube − Subsidy by endorsement14.50 30 ml **✓ Xylocaine 2% Jelly**

- a) Up to 150 ml available on a PSO
- b) Subsidised only if prescribed for urethral or cervical administration and the prescription is endorsed accordingly.

Gel 2%, 11 ml urethral syringe − Subsidy by endorsement..............65.45

- a) Up to 5 each available on a PSO
- Subsidised only if prescribed for urethral, cervical or rectal administration and the prescription is endorsed accordingly.
- c) Instillagel Lido to be Principal Supply on 1 February 2026

	Subsidy		Fully	Brand or
	(Manufacturer's Price		bsidised	Generic
	\$	Per		Manufacturer
LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE				
Oral (gel) soln 2%	30.80	200 ml	✓)	Xylocaine Viscous
	44.00		✓ I	Mucosoothe
Inj 1%, 5 ml ampoule - Up to 25 inj available on a PSO	15.00	25	✓ [Lidocaine-Baxter
	17.50	50		
	(35.00))	Xylocaine
Inj 2%, 5 ml ampoule - Up to 5 inj available on a PSO	27.50	25	✓ [Lidocaine-Baxter
Inj 1%, 20 ml ampoule - Up to 5 inj available on a PSO	12.00	5		
	(20.00))	Xylocaine
Inj 1%, 20 ml vial - Up to 5 inj available on a PSO	19.50	5	✓ [Lidocaine-Baxter
Inj 2%, 20 ml vial - Up to 5 inj available on a PSO	14.00	5	✓ [Lidocaine-Baxter
Inj 10%, 5 ml ampoule - Subsidy by endorsement	CBS	10	✓)	Xylocard 500 S29
Subsidised only for people receiving palliative care ser (Mucosoothe Oral (gel) soln 2% to be delisted 1 April 2026)		algesic ag	ents hav	ven't been effective.

Topical Local Anaesthetics

⇒SA0906 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years where the patient is a child with a chronic medical condition requiring frequent injections or venepuncture.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

LIDOCAINE [LIGNOCAINE] - Special Authority see SA0906 above - Retail pharmacy

5 g OP ✓ LMX4 30.00 30 g OP ✓ LMX4

LIDOCAINE [LIGNOCAINE] WITH PRILOCAINE - Special Authority see \$A0906 above - Retail pharmacy Crm 2.5% with prilocaine 2.5%......45.00 30 g OP ✓ EMLA **✓** EMLA 5

Analgesics

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 115

Non-opioid Analgesics

ASF	PΙF	₹IN	
	_		ı

*	Tab dispersible 300 mg – Up to 30 tab available on a PSO5.65	100	Ethics Aspirin
CA	PSAICIN - Subsidy by endorsement		
	Subsidised only if prescribed for post-herpetic neuralgia or diabetic peripheral r	neuropathy	and the prescription is endo

orsed accordingly. ✓ Zo-Rub HP

45 g OP ✓ Zostrix HP

NEFOPAM HYDROCHLORIDE

Tab 30 mg23.40 90 ✓ Acupan

		Subsidy		Fully	Brand or
		(Manufacturer's Price		ubsidised	Generic
_		\$	Per		Manufacturer
PA	RACETAMOL Tab 500 mg hijatay agaly	10.75	1 000		la alma I
	Tab 500 mg - blister pack		1,000	▼ <u>F</u>	Pacimol Pacimol
	a) Maximum of 300 tab per prescription; can be waiveb) Up to 30 tab available on a PSO	a by endorsement			
	c)				
	1) Subsidy by endorsement for higher quantities regular daily dosing for one month or greater, annotate the prescription as endorsed where 2) Maximum of 100 tab per dispensing for none (for non-endorsed patients), then dispense in Tab 500 mg - bottle pack – Maximum of 300 tab per prescription; can be waived by endorsement	and the prescription dispensing history si ndorsed patients. If repeat dispensings	upports a quantities not exceed 1,000	ted accord long-term s prescribe ding 100 to	dingly. Pharmacists may condition. If for more than 100 tabs ab per dispensing. Counsel Cou
	daily dosing for one month or greater, and the pre				rmacists may annotate the
	prescription as endorsed where dispensing histor 2) Maximum of 100 tab per dispensing for non-endorsed patients), then dispense in repeat	y supports a long-te orsed patients. If qua	rm conditi antities pre	on. escribed fo	or more than 100 tabs (for
	Oral liq 120 mg per 5 ml	3.98	200 ml	√ P	aracetamol (Ethics)
	a) Maximum of 600 ml per prescription; can be waived b) Up to 200 ml available on a PSO c) Not in combination d)	d by endorsement			
	 Maximum of 200 ml per dispensing for non-er non-endorsed patients), then dispense in repe Subsidy by endorsement for higher quantities regular daily dosing for one month or greater and Pharmacists may annotate the prescription as condition. 	eat dispensing not ex- is available for pation and the prescription s endorsed where dis	ceeding 2 ents with lo is endorse spensing h	200 ml per ong term o ed or anno nistory sup	dispensing. onditions who require stated accordingly. sports a long-term
	3) Note: 200 ml presentations of paracetamol of		pplied on E	BSO to a V	accinator (other than a
	Pharmacist) under the provisions in Part I of S 4) Note: Direct Provision by a pharmacist of up conjunction with immunisation of a child unde Oral liq 250 mg per 5 ml	to 200 ml permitted r 2 years of age with3.35		coccal B n	
	 a) Maximum of 600 ml per prescription; can be waived b) Up to 200 ml available on a PSO c) Not in combination d) 	d by endorsement			
	Maximum of 200 ml per dispensing for non-er				
	non-endorsed patients), then dispense in repe 2) Subsidy by endorsement for higher quantities regular daily dosing for one month or greater a Pharmacists may annotate the prescription as condition.	is available for patie and the prescription	ents with lo	ong term c ed or anno	onditions who require stated accordingly.
	 Note: 200 ml presentations of paracetamol or Pharmacist) under the provisions in Part I of S 		plied on E	BSO to a V	accinator (other than a
	 Note: Direct Provision by a pharmacist of up conjunction with immunisation of a child unde 	to 200 ml permitted r 2 years of age with	meningo	coccal B n	nulticomponent vaccine.
*	Suppos 125 mg	4.29	10	✓ 9	<u>iacet</u>

				INE	1VUUS STSTEIN
		Subsidy (Manufacturer's Price)	Per	Fully Subsidised	Brand or Generic Manufacturer
*	Suppos 250 mg		10		Gacet
*	Suppos 500 mg	16.55	50	•	<u>Gacet</u>
C	pioid Analgesics				
CC	DEINE PHOSPHATE - Safety medicine; prescriber may dete	ermine dispensing fre	quen	су	
	Tab 15 mg	5.82	100	′ 🗸	Noumed
	Tab 30 mg	6.88	100		Noumed
	Tab 60 mg	13.89	100		Noumed
DII	HYDROCODEINE TARTRATE				
	Tab long-acting 60 mg	9.20	60	1	DHC Continus
	DHC Continus to be Principal Supply on 1 February 202				
FF	NTANYL				
-	a) Only on a controlled drug form				
	b) No patient co-payment payable				
	c) Safety medicine; prescriber may determine dispensing fre	equency			
	Inj 50 mcg per ml, 2 ml ampoule		10	1	Boucher and Muir
	Inj 50 mcg per ml, 10 ml ampoule		10		Boucher and Muir
	Patch 12 mcg per hour		5		Fentanyl Sandoz
	Patch 25 mcg per hour		5		Fentanyl Sandoz
	Patch 50 mcg per hour	9.28	5	1	Fentanyl Sandoz
	Patch 75 mcg per hour	15.50	5	✓	Fentanyl Sandoz
	Patch 100 mcg per hour	16.37	5	✓	Fentanyl Sandoz
ME	THADONE HYDROCHLORIDE				
	a) Only on a controlled drug form				
	b) No patient co-payment payable				
	c) Safety medicine; prescriber may determine dispensing fre	equency			
	Tab 5 mg		10	✓	Methadone BNM
	Oral liq 2 mg per ml	7.80	200 n	nl 🗸	<u>Biodone</u>
	Oral liq 5 mg per ml	7.80	200 n	nl 🗸	Biodone Forte
	Oral liq 10 mg per ml	9.65	200 n		Biodone Extra Forte
	Inj 10 mg per ml, 1 ml	72.99	10	✓	AFT
MC	DRPHINE HYDROCHLORIDE				
	a) Only on a controlled drug form				
	b) No patient co-payment payable				
	c) Safety medicine; prescriber may determine dispensing fre				
	Oral liq 1 mg per ml		200 m		RA-Morph
	Oral liq 2 mg per ml		200 m		RA-Morph
	Oral liq 5 mg per ml		200 m		RA-Morph
	Ovel lie 10 men men vel	40.05	200		DA Massala

✓ RA-Morph

200 ml

Oral liq 10 mg per ml40.25

(1)	Subsidy Manufacturer's Price)		Fully Subsidised	
,	\$	Per	✓	Manufacturer
ORPHINE SULPHATE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing frequency	uency			
Tab immediate-release 10 mg	2.80	10	✓	Sevredol
Tab immediate-release 20 mg		10		Sevredol
Cap long-acting 10 mg		10		m-Eslon
Cap long-acting 30 mg		10		m-Eslon
Cap long-acting 60 mg		10		m-Eslon
Cap long-acting 100 mg	10.50	10	•	m-Eslon
Oral liq 2 mg per ml	16.31	100 m	✓	Wockhardt S29
	29.80		✓	Oramorph
			✓	Oramorph CDC
				S29 S29
Inj 5 mg per ml, 1 ml ampoule – Up to 5 inj available on a PSC Medsurge to be Principal Supply on 1 February 2026	5.96	5	1	Medsurge
Inj 10 mg per ml, 1 ml ampoule – Up to 5 inj available on a PS Medsurge to be Principal Supply on 1 February 2026	O4.99	5	•	Medsurge
Inj 15 mg per ml, 1 ml ampoule - Up to 5 inj available on a PS	O 6.93	5	•	Medsurge
Medsurge to be Principal Supply on 1 February 2026 Inj 30 mg per ml, 1 ml ampoule – Up to 5 inj available on a PS Medsurge to be Principal Supply on 1 February 2026	O7.28	5	•	Medsurge
(YCODONE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing frequ	uency			
Tab controlled-release 5 mg		20	✓	Oxycodone Sando
Tab immediate-release 5 mg	13.77	100	✓	Oxycodone Amne
Tab controlled-release 10 mg	2.49	20	✓	Oxycodone Sando
Tab immediate-release 10 mg	18.77	100	✓	Oxycodone Amne
Tab controlled-release 20 mg	3.41	20	✓	Oxycodone Sando
Tab immediate-release 20 mg	26.77	100		Oxycodone Amne
Tab controlled-release 40 mg	6.67	20		Oxycodone Sando
Tab controlled-release 80 mg		20		Oxycodone Sando
Oral liq 1 mg per ml	37.08	250 m		Oxycodone Lucis
				Rosemont
Inj 10 mg per ml, 1 ml ampoule		5		<u>Hameln</u>
Inj 10 mg per ml, 2 ml ampoule		5		<u>Hameln</u>
Inj 50 mg per ml, 1 ml ampoule	14.90	5	•	<u>Hameln</u>
xycodone Lucis Oral liq 1 mg per ml to be delisted 1 June 2026)				
RACETAMOL WITH CODEINE - Safety medicine; prescriber m	ay determine disp	ensino	frequenc	V
Tab paracetamol 500 mg with codeine phosphate 8 mg		1,000		Paracetamol +

Paracetamol + Codeine (Relieve) to be Principal Supply on 1 February 2026

Codeine (Relieve)

			IVL	11VO03 3131EW
	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	I Generic
PETHIDINE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing free		40		Navona d Dathidina
Tab 50 mg Noumed Pethidine to be Principal Supply on 1 February 2		10	•	Noumed Pethidine
Inj 50 mg per ml, 1 ml ampoule – Up to 5 inj available on a P		5	1	DBL Pethidine
ing 50 mg per mi, 1 mi ampoule - Op to 5 mg available on a 1	0020.00	J	•	Hydrochloride
Inj 50 mg per ml, 2 ml ampoule - Up to 5 inj available on a Ps	SO30.72	5	/	DBL Pethidine
.,		-		Hydrochloride
TRAMADOL HYDROCHLORIDE				•
Tab sustained-release 100 mg	1.95	20	1	Tramal SR 100
Tab sustained-release 150 mg	2.95	20	✓	Tramal SR 150
Tab sustained-release 200 mg	3.80	20	✓	Tramal SR 200
Cap 50 mg	3.33	100	/	Arrow-Tramadol
Autidonycoconto				
Antidepressants				
Cyclic and Related Agents				
•				
AMITRIPTYLINE - Safety medicine; prescriber may determine di	spensing frequency			
Tab 10 mg		100		Arrow-Amitriptyline
Tab 25 mg		100		Arrow-Amitriptyline
Tab 50 mg		100		Arrow-Amitriptyline
CLOMIPRAMINE HYDROCHLORIDE - Safety medicine; prescrit				
Tab 25 mg		50		APO Clomipramine
Cap 10 mg	35.50	28	•	Clomipramine Teva
(Clomipramine Teva Cap 10 mg to be delisted 1 April 2026)				
DOSULEPIN [DOTHIEPIN] HYDROCHLORIDE – Subsidy by end				
a) Safety medicine; prescriber may determine dispensing free				
b) Subsidy by endorsement – Subsidised for patients who we				
2019 and the prescription is endorsed accordingly. Pharm exists a record of prior dispensing of dosulepin [dothiepin]		tne	prescriptio	n as endorsed where ther
Tab 75 mg		30	/	Dosulepin Viatris
Cap 25 mg		50		Dosulepin
34p = 3 g		•		Viatris S29
IMIPRAMINE HYDROCHLORIDE - Safety medicine; prescriber r	nav determine disne	nsinc	ı frequenc	
Tab 10 mg		50		Tofranil
1.22 . 0	10.96	100		Tofranil
Tab 25 mg	4.93	28	1	Imipramine
•				Crescent S29
	8.80	50	/	Tofranil
NORTRIPTYLINE HYDROCHLORIDE				
a) Brand switch fee payable (Pharmacode 2715740) - see pa	age 279 for details			
b) Safety medicine; prescriber may determine dispensing free				
Tab 10 mg		50	1	Allegron
•	2.46	100		Norpress
Tab 25 mg	2.95	50		Allegron
	6.29	180	1	Norpress
(Norpress Tab 10 mg to be delisted 1 March 2026)				
(Norpress Tab 25 mg to be delisted 1 March 2026)				

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
Monoamine-Oxidase Inhibitors (MAOIs) - Non	Selective			
TRANYLCYPROMINE SULPHATE * Tab 10 mg	22.94	50	/	Parnate
Monoamine-Oxidase Type A Inhibitors				
MOCLOBEMIDE * Tab 150 mg* * Tab 300 mg		60 60		Aurorix Aurorix
Selective Serotonin Reuptake Inhibitors				
CITALOPRAM HYDROBROMIDE * Tab 20 mg ESCITALOPRAM	3.55	84	•	Celapram
* Tab 10 mg	1.07	28	•	Ipca-Escitalopram Escitalopram (Ethics)
* Tab 20 mg FLUOXETINE HYDROCHLORIDE		28		<u>Ipca-Escitalopram</u>
a) Subsidised by endorsement 1) When prescribed for a patient who cannot swa		28 apsu		Fluox e prescription is endorsed
a) Subsidised by endorsement 1) When prescribed for a patient who cannot swa accordingly; or 2) When prescribed in a daily dose that is not a rendorsed. Note: Tablets should be combined b) Fluox to be Principal Supply on 1 March 2026 Cap 20 mg	allow whole tablets or of multiple of 20 mg in whole to facili with capsules to facili	apsu	ase the pre	e prescription is endorsed
a) Subsidised by endorsement 1) When prescribed for a patient who cannot swa accordingly; or 2) When prescribed in a daily dose that is not a rendorsed. Note: Tablets should be combined b) Fluox to be Principal Supply on 1 March 2026 Cap 20 mg	allow whole tablets or or multiple of 20 mg in wh d with capsules to facili3.50	apsuich cata	ase the preincrementa	e prescription is endorsed escription is deemed to be I 10 mg doses. Arrow-Fluoxetine
a) Subsidised by endorsement 1) When prescribed for a patient who cannot swa accordingly; or 2) When prescribed in a daily dose that is not a rendorsed. Note: Tablets should be combined b) Fluox to be Principal Supply on 1 March 2026 Cap 20 mg	allow whole tablets or of multiple of 20 mg in whole tables to facili	eapsuich catate i	ase the preincrementa	e prescription is endorsed escription is deemed to be I 10 mg doses. Arrow-Fluoxetine Paxtine
a) Subsidised by endorsement 1) When prescribed for a patient who cannot swa accordingly; or 2) When prescribed in a daily dose that is not a rendorsed. Note: Tablets should be combined b) Fluox to be Principal Supply on 1 March 2026 * Cap 20 mg	allow whole tablets or or multiple of 20 mg in wh d with capsules to facili3.50	apsuich catate i	ase the preincrementa	e prescription is endorsed escription is deemed to be I 10 mg doses. Arrow-Fluoxetine
a) Subsidised by endorsement 1) When prescribed for a patient who cannot swa accordingly; or 2) When prescribed in a daily dose that is not a rendorsed. Note: Tablets should be combined b) Fluox to be Principal Supply on 1 March 2026 ** Cap 20 mg	allow whole tablets or of multiple of 20 mg in whole tables to facili	eapsuich catate i	ase the preincrementa	e prescription is endorsed escription is deemed to be I 10 mg doses. Arrow-Fluoxetine Paxtine
a) Subsidised by endorsement 1) When prescribed for a patient who cannot swa accordingly; or 2) When prescribed in a daily dose that is not a rendorsed. Note: Tablets should be combined b) Fluox to be Principal Supply on 1 March 2026 ** Cap 20 mg	allow whole tablets or of multiple of 20 mg in whole tables to facili	eapsucich catate i 90 30 90 30	ase the preincrementa	e prescription is endorsed escription is deemed to be I 10 mg doses. Arrow-Fluoxetine Paxtine Loxamine Setrona
a) Subsidised by endorsement 1) When prescribed for a patient who cannot swa accordingly; or 2) When prescribed in a daily dose that is not a rendorsed. Note: Tablets should be combined b) Fluox to be Principal Supply on 1 March 2026 ** Cap 20 mg	allow whole tablets or of multiple of 20 mg in whole tables to facili	eapsucich catate i 90 30 90 30	ase the preincrementa	e prescription is endorsed escription is deemed to be I 10 mg doses. Arrow-Fluoxetine Paxtine Loxamine Setrona
a) Subsidised by endorsement 1) When prescribed for a patient who cannot swa accordingly; or 2) When prescribed in a daily dose that is not a rendorsed. Note: Tablets should be combined b) Fluox to be Principal Supply on 1 March 2026 * Cap 20 mg	allow whole tablets or of multiple of 20 mg in whole tables to facili	eapsucich catate i 90 30 90 30	ase the preincrementa	e prescription is endorsed escription is deemed to be I 10 mg doses. Arrow-Fluoxetine Paxtine Loxamine Setrona Setrona Noumed
a) Subsidised by endorsement 1) When prescribed for a patient who cannot swa accordingly; or 2) When prescribed in a daily dose that is not a rendorsed. Note: Tablets should be combined by Fluox to be Principal Supply on 1 March 2026 ** Cap 20 mg	allow whole tablets or comultiple of 20 mg in whole tablets to facility and the capsules are capsules caps	apsuich catate i 90 30 90 30 30 30	ase the preincrementa	e prescription is endorsed escription is deemed to be I 10 mg doses. Arrow-Fluoxetine Paxtine Loxamine Setrona Setrona
a) Subsidised by endorsement 1) When prescribed for a patient who cannot swa accordingly; or 2) When prescribed in a daily dose that is not a rendorsed. Note: Tablets should be combined by Fluox to be Principal Supply on 1 March 2026 ** Cap 20 mg	allow whole tablets or comultiple of 20 mg in whole tablets or comultiple of 20 mg in whole with capsules to facility	30 30 30 30 30	ales and the	e prescription is endorsed escription is deemed to be I 10 mg doses. Arrow-Fluoxetine Paxtine Loxamine Setrona Setrona Noumed Noumed
1) When prescribed for a patient who cannot swa accordingly; or 2) When prescribed in a daily dose that is not a rendorsed. Note: Tablets should be combined b) Fluox to be Principal Supply on 1 March 2026 ** Cap 20 mg	allow whole tablets or comultiple of 20 mg in whole tablets to facility and the capsules to facility an	30 30 30	ales and the ase the preincrementa	e prescription is endorsed escription is deemed to be I 10 mg doses. Arrow-Fluoxetine Paxtine Loxamine Setrona Setrona Noumed

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

Antiepilepsy Drugs

Agents for Control of Status Epilepticus

DIAZEPAM – Safety medicine; prescriber may determine dispensing frequency Inj 5 mg per ml, 2 ml ampoule – Subsidy by endorsement27.92	5	✓ Hospira
a) Up to 5 inj available on a PSO		
b) Only on a PSO		
 c) PSO must be endorsed "not for anaesthetic procedures". 		
Rectal tubes 5 mg - Up to 5 tube available on a PSO54.58	5	Stesolid
PHENYTOIN SODIUM		
* Inj 50 mg per ml, 2 ml ampoule - Up to 5 inj available on a		
PSO104.58	5	Hospira
* Inj 50 mg per ml, 5 ml ampoule - Up to 5 inj available on a		•
PSO154.01	5	✓ Hospira
(Hospira Inj 50 mg per ml, 2 ml ampoule to be delisted 1 February 2026)		

Control of Epilepsy

CARBAMAZEPINE			
* Tab 200 mg	14.53	100	Tegretol
* Tab long-acting 200 mg	16.98	100	Tegretol CR
	33.96	200	Tegretol CR
* Tab 400 mg		100	Tegretol
* Tab long-acting 400 mg		100	Tegretol CR
* Oral liq 20 mg per ml	26.37	250 ml	Tegretol
CLOBAZAM - Safety medicine; prescriber may determine	dispensing frequency		
Tab 10 mg	9.12	50	✓ Frisium
CLONAZEPAM - Safety medicine; prescriber may determine	ne dispensing frequen	icy	
Oral drops 2.5 mg per ml		10 ml OP	✓ Rivotril
ETHOSUXIMIDE			
Cap 250 mg	140.88	100	✓ Zarontin
Oral liq 250 mg per 5 ml		200 ml	✓ Zarontin
GABAPENTIN			
Note: Not subsidised in combination with subsidised pr	regabalin		
* Cap 100 mg	6.45	100	✓ Nupentin
* Cap 300 mg	8.45	100	✓ Nupentin
* Cap 400 mg	10.26	100	✓ Nupentin
LACOSAMIDE - Special Authority see SA2267 on the next	page – Retail pharma	асу	
▲ Tab 50 mg	25.04	14	✓ Vimpat
▲ Tab 100 mg	50.06	14	✓ Vimpat
-	200.24	56	✓ Vimpat
▲ Tab 150 mg	75.10	14	✓ Vimpat
	300.40	56	✓ Vimpat
▲ Tab 200 mg	400.55	56	✓ Vimpat

Subsidy Fully (Manufacturer's Price) Subsidised Per

Brand or Generic

Manufacturer

⇒SA2267 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Patient has focal epilepsy: and
- 2 Seizures are not adequately controlled by, or patient has experienced unacceptable side effects from, optimal treatment with all of the following: sodium valproate, topiramate, levetiracetam and any two of carbamazepine, lamotrigine and phenytoin sodium (see Note).

Note: Those of childbearing potential are not required to trial phenytoin sodium, sodium valproate, or topiramate. Those who can father children are not required to trial sodium valproate.

Renewal from any relevant practitioner. Approvals valid for 24 months where the patient has demonstrated a significant and sustained improvement in seizure rate or severity and/or quality of life compared with that prior to starting lacosamide treatment.

	MOTRIGINE	y or mo compared	with that phor to	Jolai	ang accountac accan
▲	Tab dispersible 2 mg	55.00	30	/	Lamictal
<u> </u>	Tab dispersible 5 mg		30	/	Lamictal
*	Tab dispersible 25 mg		56	1	Logem
*	Tab dispersible 50 mg		56		Logem
	Tab dispersible 100 mg		56	1	Logem
۱F۱	/ETIRACETAM				
	Tab 250 mg	5.84	60	/	Everet
	Tab 500 mg		60	/	Everet
	Tab 750 mg		60	1	Everet
	Tab 1,000 mg	21.82	60	/	Everet
	Oral liq 100 mg per ml	44.78	300 ml OP	1	Levetiracetam-AFT
	Inj 100 mg per ml, 5 ml vial	38.95	10	•	Levetiracetam-AFT
PHI	ENOBARBITONE				
	For phenobarbitone oral liquid refer Standard Formulae, p	age 281			
	Tab 15 mg	•	500	/	Noumed
					Phenobarbitone
	Tab 30 mg	398.50	500	1	Noumed
	ů				Phenobarbitone
PHI	ENYTOIN SODIUM				
	Tab 50 mg	75.00	200	/	Dilantin Infatab
	Cap 30 mg		200		Dilantin
	Cap 100 mg		200	1	Dilantin
*	Oral lig 30 mg per 5 ml		500 ml	•	Dilantin Paediatric
PRI	EGABALIN				
	Note: Not subsidised in combination with subsidised gaba	epentin			
*	Cap 25 mg	•	56	/	Lyrica
-					Pregabalin Pfizer
*	Cap 75 mg	2.65	56		Lyrica
				/	Pregabalin Pfizer
*	Cap 150 mg	4.01	56	1	Lyrica
				1	Pregabalin Pfizer
*	Cap 300 mg	7.38	56	1	Lyrica
				✓	Pregabalin Pfizer
PR	MIDONE				
*	Tab 250 mg	37.35	100	1	Primidone Clinect

	Subsidy		Fully	Brand or
	(Manufacturer's Pric	e) Si	ubsidised	Generic
	\$	Per	1	Manufacturer
ODIUM VALPROATE				
Tab 100 mg	13.65	100	✓ E	pilim Crushable
Tab 200 mg EC	27.44	100	✓ E	pilim
Tab 500 mg EC	52.24	100	✓ E	pilim
F Oral liq 200 mg per 5 ml	20.48	300 ml	✓ E	pilim S/F Liquid
,			✓ E	pilim Syrup
k Inj 100 mg per ml, 4 ml	41.50	1	✓ E	pilim IV
STIRIPENTOL - Special Authority see SA2268 belo	ow – Retail pharmacy			-
Cap 250 mg		60	✓ D	iacomit
Powder for oral liq 250 mg sachet		60	✓ D	iacomit

⇒SA2268 Special Authority for Subsidy

Initial application only from a paediatric neurologist or Practitioner on the recommendation of a paediatric neurologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has confirmed diagnosis of Dravet syndrome; and
- 2 Seizures have been inadequately controlled by appropriate courses of sodium valproate, clobazam and at least two of the following: topiramate, levetiracetam, ketogenic diet.

Note: Those of childbearing potential are not required to trial sodium valproate or topiramate. Those who can father children are not required to trial sodium valproate.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where the patient continues to benefit from treatment as measured by reduced seizure frequency from baseline.

TOPIRAMATE

▲ Tab 25 mg11.07	60	✓ Arrow-Topiramate
•		✓ Topiramate Actavis
26.04		✓ Topamax
▲ Tab 50 mg18.81	60	✓ Arrow-Topiramate
·		✓ Topiramate Actavis
44.26		✓ Topamax
▲ Tab 100 mg31.99	60	✓ Arrow-Topiramate
		✓ Topiramate Actavis
75.25		✓ Topamax
▲ Tab 200 mg55.19	60	Arrow-Topiramate
		Topiramate Actavis
129.85		✓ Topamax
▲ Sprinkle cap 15 mg20.84	60	✓ Topamax
▲ Sprinkle cap 25 mg26.04	60	✓ Topamax
VIGABATRIN - Special Authority see SA2088 below - Retail pharmacy		
▲ Tab 500 mg119.30	100	✓ Sabril
▲ Powder for oral soln 500 mg per sachet71.58	60	✓ Sabril S29

⇒SA2088 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Any of the following:
 - 1.1 Patient has infantile spasms; or
 - 1.2 Both:
 - 1.2.1 Patient has epilepsy; and
 - 1.2.2 Either:



|--|

continued...

- 1.2.2.1 Seizures are not adequately controlled with optimal treatment with other antiepilepsy agents; or
- 1.2.2.2 Seizures are controlled adequately but the patient has experienced unacceptable side effects from optimal treatment with other antiepilepsy agents; or
- 1.3 Patient has tuberous sclerosis complex; and
- 2 Either:
 - 2.1 Patient is, or will be, receiving regular automated visual field testing (ideally before starting therapy and on a 6-monthly basis thereafter): or
 - 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields...

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient has demonstrated a significant and sustained improvement in seizure rate or severity and or quality of life; and
- 2 Either:
 - 2.1 Patient is receiving regular automated visual field testing (ideally every 6 months) on an ongoing basis for duration of treatment with vigabatrin; or
 - 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields...

Antimigraine Preparations

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 115

Acute Migraine Treatment

Tab orodispersible 10 mg4	1.84	30	Rizamelt
SUMATRIPTAN			
Tab 50 mg14	.41	90	Sumagran
Tab 100 mg22	2.68	90	Sumagran
Inj 12 mg per ml, 0.5 ml prefilled pen - Maximum of 10 inj per			-
prescription29	9.80 2	OP ✓	Clustran

Prophylaxis of Migraine

For Beta Adrenoceptor Blockers refer to CARDIOVASCULAR SYSTEM, page 49

PIZOTIFFN

★ Tab 500 mcg......23.21 100 **✓ Sandomigran**

Antinausea and Vertigo Agents

For Antispasmodics refer to ALIMENTARY TRACT, page 8

APREPITANT – Special Authority see SA0987 below – Retail pharmacy

Cap 2 × 80 mg and 1 × 125 mg.......21.90 3 OP ✓ <u>Emend Tri-Pack</u>

⇒SA0987 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months where the patient is undergoing highly emetogenic chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.

Renewal from any relevant practitioner. Approvals valid for 12 months where the patient is undergoing highly emetogenic chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.

BETAHISTINE DIHYDROCHLORIDE

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Generic
CYCLIZINE HYDROCHLORIDE	0.00	- 10		
Tab 50 mg	0.66	10	•	<u>Nausicalm</u>
CYCLIZINE LACTATE				
Inj 50 mg per ml, 1 ml ampoule – Up to 10 inj available on a PSO	16.36	10	•	Hameln
DOMPERIDONE				
* Tab 10 mg	3.80	100	•	<u>Domperidone</u> <u>Viatris</u>
HYOSCINE HYDROBROMIDE				
* Inj 400 mcg per ml, 1 ml ampoule	93.00	10	1	Martindale \$29
Patch 1 mg per 72 hours – Special Authority see SA1998				
below – Retail pharmacy	88.50	10	•	Scopolamine Transdermal System Viatris

⇒SA1998 Special Authority for Subsidy

METOCI OPRAMIDE HYDROCHI ORIDE

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Control of intractable nausea, vomiting, or inability to swallow saliva in the treatment of malignancy or chronic disease where the patient cannot tolerate or does not adequately respond to oral anti-nausea agents; or
- 2 Control of clozapine-induced hypersalivation where trials of at least two other alternative treatments have proven ineffective.

Renewal from any relevant practitioner. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

* Tab 10 mg – Up to 30 tab available on a PSO1.57	100	✓ <u>Metoclopramide</u> Actavis 10
* Inj 5 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO5.48 7.00	10	✓ Medsurge✓ Baxter
(Baxter Inj 5 mg per ml, 2 ml ampoule to be delisted 1 April 2026)		
ONDANSETRON		
* Tab 4 mg1.95	50	✓ Periset
Tab disp 4 mg - Up to 10 tab available on a PSO0.56	10	✓ Periset ODT
* Tab 8 mg3.50	50	✓ Periset
Tab disp 8 mg - Up to 10 tab available on a PSO0.90	10	✓ Periset ODT
PROCHLORPERAZINE		
* Tab 3 mg buccal	50	
(30.00)		Prochlorperazine maleate (Brown & Burk)
* Tab 5 mg - Up to 30 tab available on a PSO25.00	250	✓ Nausafix
* Inj 12.5 mg per ml, 1 ml – Up to 5 inj available on a PSO25.81	10	✓ Stemetil

Antipsychotics

General

AMISULPRIDE - Safety medicine; prescriber may determine dis	spensing frequency	,	
Tab 100 mg	5.84	30	✓ Sulprix
Tab 200 mg	14.47	60	✓ Sulprix
Tab 400 mg	35.06	60	✓ Sulprix

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	I Generic
	\$	Per	•	Manufacturer
ARIPIPRAZOLE – Safety medicine; prescriber may determine of	dispensing frequency			
Tab 5 mg		30	1	Aripiprazole Sandoz
Tab 10 mg		30		Aripiprazole Sandoz
Tab 15 mg		30		Aripiprazole Sandoz
•		30		
Tab 20 mg				Aripiprazole Sandoz
Tab 30 mg		30		Aripiprazole Sandoz
CHLORPROMAZINE HYDROCHLORIDE - Safety medicine; p	rescriber may determi	ine dis	spensing fr	requency
Tab 25 mg - Up to 30 tab available on a PSO	15.62	100	✓	Largactil
Tab 100 mg - Up to 30 tab available on a PSO	36.73	100	✓	Largactil
Inj 25 mg per ml, 2 ml - Up to 5 inj available on a PSO	30.79	10	✓	Largactil
CLOZAPINE – Hospital pharmacy [HP4]				•
Safety medicine; prescriber may determine dispensing frequency	IODOV.			
, , , , , , , , , , , , , , , , , , , ,	•	EΟ	./	Clanina
Tab 25 mg	0.09	50		Clopine
				Clozaril
	13.37	100		Clopine
				Clozaril
Tab 50 mg	8.67	50		Clopine
	17.33	100		Clopine
Tab 100 mg	17.33	50	✓	Clopine
			✓	Clozaril
	34.65	100	✓	Clopine
			/	Clozaril
Tab 200 mg	34.65	50	1	Clopine
3	69.30	100		Clopine
Suspension 50 mg per ml	173.30	100 m		Versacloz
, , , , , , , , , , , , , , , , , , , ,				
HALOPERIDOL – Safety medicine; prescriber may determine of		400	,	0
Tab 500 mcg – Up to 30 tab available on a PSO		100		Serenace
Tab 1.5 mg — Up to 30 tab available on a PSO		100		Serenace
Tab 5 mg - Up to 30 tab available on a PSO		50		Serenace
	29.72	100		Serenace
Oral liq 2 mg per ml - Up to 200 ml available on a PSO		100 m		Serenace
Inj 5 mg per ml, 1 ml ampoule – Up to 5 inj available on a F	PSO21.55	10	•	Serenace
LEVOMEPROMAZINE - Safety medicine; prescriber may deter	rmine dispensina frea	uencv		
Tab 25 mg (33.8 mg as a maleate)		100		Nozinan (Swiss)
Tab 25 mg as a maleate		100		Nozinan
Tab 100 mg (135 mg as a maleate)		100	_	Nozinan (Swiss)
Tab 100 mg as a maleate		100		Nozinan (Owiss)
•				
LEVOMEPROMAZINE HYDROCHLORIDE – Safety medicine;				
Inj 25 mg per ml, 1 ml ampoule	23.26	10	•	<u>Wockhardt</u>
LITHIUM CARBONATE - Safety medicine; prescriber may dete	ermine dispensing fred	quenc	٧	
Tab long-acting 400 mg		100		Priadel
Cap 250 mg		100		Douglas
,				
OLANZAPINE – Safety medicine; prescriber may determine dis		20		7. mina
Tab 2.5 mg		30		Zypine
Tab 5 mg		30		Zypine
Tab orodispersible 5 mg		28		Zypine ODT
Tab 10 mg		28		<u>Zypine</u>
	1.93	30		Zypine
Tab orodispersible 10 mg	2.89	28	•	Zypine ODT

	Subsidy		Fully	
	(Manufacturer's Price)		Subsidised	
	\$	Per		Manufacturer
ERICYAZINE - Safety medicine; prescriber may determine	dispensing frequency			
Tab 2.5 mg	13.61	100	✓	Neulactil
Tab 10 mg	48.45	100	•	Neulactil
UETIAPINE - Safety medicine; prescriber may determine of	dispensing frequency			
Tab 25 mg	,	30	✓	Quetiapine
•				Viatris \$29
	2.36	90	1	Quetapel
	13.11	500		Quetiapine
				Viatris S29
Tab 100 mg	6.40	90	1	Quetapel
Tab 200 mg		90		Quetapel
Tab 300 mg		90	_	Quetapel
ISPERIDONE – Safety medicine; prescriber may determine				
Tab 0.5 mg		60	1	Risperidone (Teva)
Tab 1 mg		60		Risperidone (Teva)
Tab 2 mg		60		Risperidone (Teva)
Tab 3 mg		60		Risperidone (Teva)
Tab 4 mg		60		Risperdal
•			1	Risperidone (Teva)
Oral liq 1 mg per ml	10.29	30 m	/	Risperon
	34.30	100 m	nl 🗸	Risperon
PRASIDONE - Safety medicine; prescriber may determine	dispensing frequency			-
Cap 20 mg	,	60	1	Zusdone
Cap 40 mg		60		Zusdone
Cap 60 mg		60		Zusdone
Cap 80 mg		60	1	Zusdone
JCLOPENTHIXOL HYDROCHLORIDE - Safety medicine;		ne disa	nensing fre	equency
	procediber may determin	io uio	Johnship IIC	2quoi io y

Depot Injections

⇒SA2395 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 Either:
 - 1.1 The patient has had an initial Special Authority approval for risperidone depot injection, paliperidone depot injection or olanzapine depot injection; or
 - 1.2 All of the following:
 - 1.2.1 The patient has schizophrenia or other psychotic disorder; and
 - 1.2.2 The patient has received treatment with oral atypical antipsychotic agents but has been unable to adhere; and
 - 1.2.3 The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months; or



(Manu	Subsidy	Fully	Brand or
	facturer's Price)	Subsidised	Generic
	\$ P	Per 🗸	Manufacturer

continued...

2 Patient has been unable to access olanzapine depot injection due to supply issues with olanzapine depot injection, or otherwise would have been started on olanzapine depot injection but has been unable to due to supply issues with olanzapine depot injection.

Notes: The Olanzapine depot injection Special Authority criteria that apply to criterion 2 in this Aripiprazole Special Authority application are as follows:

- The patient has had an initial Special Authority approval for paliperidone depot injection or risperidone depot injection; or
- All of the following:
 - The patient has schizophrenia; and
 - The patient has not been able to adhere with treatment using oral atypical antipsychotic agents; and
 - The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months.

FLUPENTHIXOL DECANOATE - Safety medicine; prescriber may determine dispensing frequency

Inj 20 mg per ml, 1 ml - Up to 5 inj available on a PSO	5	✓ Fluanxol
Inj 20 mg per ml, 2 ml - Up to 5 inj available on a PSO20.90	5	✓ Fluanxol
Inj 100 mg per ml, 1 ml - Up to 5 inj available on a PSO40.87	5	✓ Fluanxol

HALOPERIDOL DECANOATE - Safety medicine; prescriber may determine dispensing frequency

	oogoqo	
Inj 50 mg per ml, 1 ml - Up to 5 inj available on a PSO28.39	5	✓ Haldol
Inj 100 mg per ml, 1 ml - Up to 5 inj available on a PSO55.90	5	✓ Haldol Concentrate
•		✓ Haldol

Decanoas \$29

OLANZAPINE - Special Authority see SA2313 below - Retail pharmacy

- a) Safety medicine; prescriber may determine dispensing frequency
- b) Note no new patients to be initiated on olanzapine.

Inj 210 mg vial	252.00	1	Zyprexa Relprevv
Inj 300 mg vial	414.00	1	✓ Zyprexa Relprevv
Inj 405 mg vial	504.00	1	✓ Zyprexa Relprevv

⇒SA2313 Special Authority for Subsidy

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of olanzapine depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

PALIPERIDONE - Special Authority see SA2396 below - Retail pharmacy

Safety medicine; prescriber may determine dispensing frequency

Salety medicine, prescriber may determine di	spensing nequency		
Inj 25 mg syringe	194.25	1	✓ Invega Sustenna
Inj 50 mg syringe	271.95	1	✓ Invega Sustenna
Inj 75 mg syringe	357.42	1	✓ Invega Sustenna
Inj 100 mg syringe	435.12	1	✓ Invega Sustenna
Inj 150 mg syringe	435.12	1	✓ Invega Sustenna

⇒SA2396 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for risperidone depot injection or olanzapine depot injection or aripiprazole depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 Has been unable to adhere to treatment using oral atypical antipsychotic agents; and

NERVOUS SYSTEM

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsid		Generic
\$	Per	1	Manufacturer

continued...

2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

PALIPERIDONE PALMITATE – Special Authority	/ see SA2167 below – Retail pharm	acy	
Inj 175 mg syringe	815.85	1	Invega Trinza
Inj 263 mg syringe	1,072.26	1	✓ Invega Trinza
Inj 350 mg syringe		1	✓ Invega Trinza
Inj 525 mg syringe	1,305.36	1	✓ Invega Trinza

⇒SA2167 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has schizophrenia; and
- 2 The patient has had an initial Special Authority approval for paliperidone once-monthly depot injection.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

⇒SA2397 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or olanzapine depot injection or aripiprazole depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 Has not been able to adhere with treatment using oral atypical antipsychotic agents; and
 - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of risperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

ZUCLOPENTHIXOL DECANOATE - Safety medicine; prescriber may determine dispensing frequency
Inj 200 mg per ml, 1 ml - Up to 5 inj available on a PSO......19.80 5

✓ Clopixol

Anxiolytics			
BUSPIRONE HYDROCHLORIDE			
* Tab 5 mg	13.95	100	Buspirone Viatris
* Tab 10 mg	12.50	100	 Buspirone Viatris
CLONAZEPAM - Safety medicine; prescriber may determine	dispensing frequency	/	
Tab 500 mcg	5.64	100	✓ Paxam
Tab 2 mg	10.78	100	✓ Paxam

NERVOUS SYSTEM

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
DIAZEPAM – Safety medicine; prescriber may determine dispension 2 mg	95.00	500 500	_	Arrow-Diazepam Arrow-Diazepam
LORAZEPAM – Safety medicine; prescriber may determine dispersal 1 mg	10.20	250 100	_	Ativan Ativan

Multiple Sclerosis Treatments

⇒SA2274 Special Authority for Subsidy

Initial application — (Multiple Sclerosis - dimethyl fumarate, fingolimod, glatiramer acetate, interferon beta-1-alpha, interferon beta-1-beta, natalizumab and teriflunomide) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 Diagnosis of multiple sclerosis (MS) meets the McDonald 2017 diagnostic criteria for MS and has been confirmed by a neurologist; and
 - 1.2 Patient has an EDSS score between 0 6.0; and
 - 1.3 Patient has had at least one significant attack of MS in the previous 12 months or two significant attacks in the past 24 months: and
 - 1.4 All of the following:
 - 1.4.1 Each significant attack must be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the attack, but the neurologist/physician must be satisfied that the clinical features were characteristic); and
 - 1.4.2 Each significant attack is associated with characteristic new symptom(s)/sign(s) or substantially worsening of previously experienced symptoms(s)/sign(s); and
 - 1.4.3 Each significant attack has lasted at least one week and has started at least one month after the onset of a previous attack (where relevant); and
 - 1.4.4 Each significant attack can be distinguished from the effects of general fatigue; and is not associated with a fever (T> 37.5°C); and
 - 1.4.5 Fither:
 - 1.4.5.1 Each significant attack is severe enough to change either the EDSS or at least one of the Kurtze Functional System scores by at least 1 point; or
 - 1.4.5.2 Each significant attack is a recurrent paroxysmal symptom of multiple sclerosis (tonic seizures/spasms, trigeminal neuralgia, Lhermitte's symptom); and
 - 1.5 Evidence of new inflammatory activity on an MRI scan within the past 24 months; and
 - 1.6 Any of the following:
 - 1.6.1 A sign of that new inflammatory activity on MRI scanning (in criterion 5 immediately above) is a gadolinium enhancing lesion; or
 - 1.6.2 A sign of that new inflammatory activity is a lesion showing diffusion restriction; or
 - 1.6.3 A sign of that new inflammatory is a T2 lesion with associated local swelling; or
 - 1.6.4 A sign of that new inflammatory activity is a prominent T2 lesion that clearly is responsible for the clinical features of a recent attack that occurred within the last 2 years; or
 - 1.6.5 A sign of that new inflammatory activity is new T2 lesions compared with a previous MRI scan; or
- 2 Patient has an active approval for ocrelizumab and does not have primary progressive MS.

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

Renewal — (Multiple Sclerosis - dimethyl fumarate, fingolimod, glatiramer acetate, interferon beta-1-alpha, interferon

			NER'	VOUS SYSTEM
	Subsidy (Manufacturer's Price) \$	Subsi Per	Fully dised	Brand or Generic Manufacturer
continued beta-1-beta, natalizumab and teriflunomide) from any relevant had an EDSS score of 0 to 6.0 (inclusive) with or without the use of the patient has walked 100 metres or more with or without aids in Note: Treatment on two or more funded multiple sclerosis treatment DIMETHYL FUMARATE – Special Authority see SA2274 on the page 3. Wastage claimable b) Note: Treatment on two or more funded multiple sclerosis	of unilateral or bilatera the last six months). ents simultaneously is previous page – Reta	al aids at a s not perm il pharmac	iny timo itted. Sy	e in the last six months (ie
Cap 120 mg	520.00	14 56	✓ T	ecfidera ecfidera
FINGOLIMOD – Special Authority see SA2274 on the previous pa a) Wastage claimable b) Note: Treatment on two or more funded multiple sclerosis Cap 0.5 mg	treatments simultane	,	•	nitted. i ilenya
GLATIRAMER ACETATE – Special Authority see SA2274 on the Note: Treatment on two or more funded multiple sclerosis treating 40 mg prefilled syringe	atments simultaneou		ermitte	ed. opaxone
INTERFERON BETA-1-ALPHA – Special Authority see SA2274 c Note: Treatment on two or more funded multiple sclerosis tree Inj 6 million iu prefilled syringe	atments simultaneou		ermitte	•
INTERFERON BETA-1-BETA – Special Authority see SA2274 on Note: Treatment on two or more funded multiple sclerosis tre- Inj 8 million iu per 1 ml	atments simultaneou		permitte	
NATALIZUMAB – Special Authority see SA2274 on the previous Note: Treatment on two or more funded multiple sclerosis treating 20 mg per ml, 15 ml vial	atments simultaneou	,		ed. ysabri

TERIFLUNOMIDE - Special Authority see SA2274 on the previous page - Retail pharmacy

a) Wastage claimable

b) Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

28 ✓ Teriflunomide Sandoz

Multiple Sclerosis Treatments - Other

OCRELIZUMAB - Special Authority see SA2273 below - Retail pharmacy

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

Inj 30 mg per ml, 10 ml vial.......8,450.00 1 ✓ Ocrevus

✓ Ocrevus SC 1

⇒SA2273 Special Authority for Subsidy

Initial application — (Multiple Sclerosis - ocrelizumab) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 Diagnosis of multiple sclerosis (MS) meets the McDonald 2017 diagnostic criteria for MS and has been confirmed by a neurologist; and
 - 1.2 Patient has an EDSS score between 0 6.0; and
 - 1.3 Patient has had at least one significant attack of MS in the previous 12 months or two significant attacks in the past



Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

24 months: and

- 1.4 All of the following:
 - 1.4.1 Each significant attack must be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the attack, but the neurologist/physician must be satisfied that the clinical features were characteristic); and
 - 1.4.2 Each significant attack is associated with characteristic new symptom(s)/sign(s) or substantially worsening of previously experienced symptoms(s)/sign(s); and
 - 1.4.3 Each significant attack has lasted at least one week and has started at least one month after the onset of a previous attack (where relevant); and
 - 1.4.4 Each significant attack can be distinguished from the effects of general fatigue; and is not associated with a fever (T> 37.5°C); and
 - 1.4.5 Either:
 - 1.4.5.1 Each significant attack is severe enough to change either the EDSS or at least one of the Kurtze Functional System scores by at least 1 point; or
 - 1.4.5.2 Each significant attack is a recurrent paroxysmal symptom of multiple sclerosis (tonic seizures/spasms, trigeminal neuralgia, Lhermitte's symptom); and
- 1.5 Evidence of new inflammatory activity on an MRI scan within the past 24 months; and
- 1.6 Any of the following:
 - 1.6.1 A sign of that new inflammatory activity on MRI scanning (in criterion 5 immediately above) is a gadolinium enhancing lesion; or
 - 1.6.2 A sign of that new inflammatory activity is a lesion showing diffusion restriction; or
 - 1.6.3 A sign of that new inflammatory is a T2 lesion with associated local swelling; or
 - 1.6.4 A sign of that new inflammatory activity is a prominent T2 lesion that clearly is responsible for the clinical features of a recent attack that occurred within the last 2 years; or
 - 1.6.5 A sign of that new inflammatory activity is new T2 lesions compared with a previous MRI scan; or
- 2 Patient has an active Special Authority approval for either dimethyl fumarate, fingolimod, glatiramer acetate, interferon beta-1-alpha. interferon beta-1-beta. natalizumab or teriflunomide.

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

Renewal — (Multiple Sclerosis - ocrelizumab) from any relevant practitioner. Approvals valid for 12 months where patient has had an EDSS score of 0 to 6.0 (inclusive) with or without the use of unilateral or bilateral aids at any time in the last six months (ie the patient has walked 100 metres or more with or without aids in the last six months).

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

Initial application — (Primary Progressive Multiple Sclerosis) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Diagnosis of primary progressive multiple sclerosis (PPMS) meets the 2017 McDonald criteria and has been confirmed by a neurologist; and
- 2 Patient has an EDSS 2.0 (score equal to or greater than 2 on pyramidal functions) to EDSS 6.5; and
- 3 Patient has no history of relapsing remitting multiple sclerosis.

Renewal — (Primary Progressive Multiple Sclerosis) from any relevant practitioner. Approvals valid for 12 months where patient has had an EDSS score of less than or equal to 6.5 at any time in the last six months (ie patient has walked 20 metres with bilateral assistance/aids, without rest in the last six months).

Sedatives and Hypnotics

MELATONIN – Special Authority see SA2523 on the next page – Retail pharmacy

Tab modified-release 2 mg - No more than 5 tab per day......5.80

Restricted to patients aged 18 years or under.

30

✓ <u>Vigisom</u>

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidi	sed	Generic
\$	Per	✓	Manufacturer

⇒SA2523 Special Authority for Subsidy

Initial application only from a psychiatrist, paediatrician, neurologist, respiratory specialist or any relevant practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with persistent and distressing insomnia secondary to a neurodevelopmental disorder (including, but not limited to, autism spectrum disorder or attention deficit hyperactivity disorder)*; and
- 2 Behavioural and environmental approaches have been tried and were unsuccessful, or are inappropriate; and
- 3 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day; and
- 4 Patient is aged 18 years or under*.

Renewal only from a psychiatrist, paediatrician, neurologist, respiratory specialist or any relevant practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is aged 18 years or under*; and
- 2 Patient has demonstrated clinically meaningful benefit from funded modified-release melatonin (clinician determined); and
- 3 Patient has had a trial of funded modified-release melatonin discontinuation within the past 12 months and has had a recurrence of persistent and distressing insomnia; and
- 4 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day.

Note: Indications marked with * are unapproved indications. MIDAZOLAM - Safety medicine: prescriber may determine dispensing frequency 10 ✓ Midazolam-Baxter Inj 1 mg per ml, 5 ml plastic ampoule - Up to 10 inj available ✓ Midazolam-Pfizer 10 On a PSO for status epilepticus use only. PSO must be endorsed for status epilepticus use only. Inj 5 mg per ml, 1 ml plastic ampoule - Up to 10 inj available 10 ✓ Midazolam-Pfizer On a PSO for status epilepticus use only. PSO must be endorsed for status epilepticus use only. Inj 5 mg per ml, 3 ml ampoule4.75 ✓ Midazolam-Baxter Ini 5 mg per ml. 3 ml plastic ampoule - Up to 5 ini available on ✓ Midazolam-Pfizer On a PSO for status epilepticus use only. PSO must be endorsed for status epilepticus use only. PHENOBARBITONE SODIUM - Special Authority see SA1386 below - Retail pharmacy Inj 200 mg per ml, 1 ml ampoule113.37 ✓ Max Health \$29 ⇒SA1386 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 For the treatment of terminal agitation that is unresponsive to other agents; and
- 2 The applicant is part of a multidisciplinary team working in palliative care.

TEMAZEPAM - Safety medicine: prescriber may determine dispensing frequency 25 Normison ZOPICLONE - Safety medicine: prescriber may determine dispensing frequency 500 Zopiclone Actavis

Spinal Muscular Atrophy

NUSINERSEN - PCT only - Special Authority see SA2174 on the next page ✓ Spinraza 1



Subsidy	F	ully	Brand or	
(Manufacturer's Price)	Subsid	sed	Generic	
\$	Per	✓	Manufacturer	

⇒SA2174 Special Authority for Subsidy

Initial application — (spinal muscular atrophy (SMA)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has genetic documentation of homozygous SMN1 gene deletion, homozygous SMN1 point mutation, or compound heterozygous mutation; and
- 2 Patient is 18 years of age or under; and
- 3 Either:
 - 3.1 Patient has experienced the defined signs and symptoms of SMA type I, II or IIIa prior to three years of age; or
 - 3.2 Both:
 - 3.2.1 Patient is pre-symptomatic; and
 - 3.2.2 Patient has three or less copies of SMN2.

Renewal — (spinal muscular atrophy (SMA)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 There has been demonstrated maintenance of motor milestone function since treatment initiation; and
- 2 Patient does not require invasive permanent ventilation (at least 16 hours per day) in the absence of a potentially reversible cause while being treated with nusinersen; and
- 3 Nusinersen not to be administered in combination other SMA disease modifying treatments or gene therapy.

RISDIPLAM - [Xpharm] - Special Authority see SA2203 below

Note: the supply of risdiplam is via Pharmac's approved direct distribution supply. Further details can be found on Pharmac's website https://pharmac.govt.nz/risdiplam

Powder for oral soln 750 mcg per ml, 60 mg per bottle......14,100.00 80 ml OP ✓ Evrysdi

⇒SA2203 Special Authority for Subsidy

Initial application — (spinal muscular atrophy (SMA)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has genetic documentation of homozygous SMN1 gene deletion, homozygous SMN1 point mutation, or compound heterozygous mutation; and
- 2 Patient is 18 years of age or under; and
- 3 Either:
 - 3.1 Patient has experienced the defined signs and symptoms of SMA type I, II or IIIa prior to three years of age; or
 - 3.2 Both:
 - 3.2.1 Patient is pre-symptomatic; and
 - 3.2.2 Patient has three or less copies of SMN2.

Renewal — (spinal muscular atrophy (SMA)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 There has been demonstrated maintenance of motor milestone function since treatment initiation; and
- 2 Patient does not require invasive permanent ventilation (at least 16 hours per day) in the absence of a potentially reversible cause while being treated with risdiplam; and
- 3 Risdiplam not to be administered in combination other SMA disease modifying treatments or gene therapy.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
Stimulants/ADHD Treatments				
ATOMOXETINE				
Cap 10 mg	43.02	28	✓	APO-Atomoxetine
Cap 18 mg	45.57	28	✓	APO-Atomoxetine
Cap 25 mg	44.30	28	✓	APO-Atomoxetine
Cap 40 mg	46.21	28	✓	APO-Atomoxetine
Cap 60 mg	51.31	28	✓	APO-Atomoxetine
Cap 80 mg		28	✓	APO-Atomoxetine
Cap 100 mg	65.71	28	1	APO-Atomoxetine
DEXAMFETAMINE SULFATE – Special Authority see SA2410 I a) Only on a controlled drug form b) Safety medicine; prescriber may determine dispensing from	·	су		

⇒SA2410 Special Authority for Subsidy

Initial application — (ADHD in patients aged 5 years or over) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid without further renewal unless notified for applications meeting the following criteria: All of the following:

100

✓ Noumed

Dexamfetamine

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) in patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and

Tab 5 mg29.80

- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients aged under 5 years) only from a paediatrician or psychiatrist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) in patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid without further renewal unless notified where the patient suffers from narcolepsy.

LISDEXAMFETAMINE DIMESILATE - Special Authority see SA2415 below - Retail pharmacy

- a) Only on a controlled drug form

⇒SA2415 Special Authority for Subsidy

Initial application only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 Patient is currently on treatment with lisdexamfetamine dimesilate and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer

continued...

- 2.1 ADHD (Attention Deficit and Hyperactivity Disorder); and
- 2.2 Diagnosed according to DSM-V or ICD 11 criteria; and
- 2.3 Either:
 - 2.3.1 Applicant is a paediatrician or psychiatrist; or
 - 2.3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing; and
- 2.4 Any of the following:
 - 2.4.1 Patient is taking a currently subsidised formulation of atomoxetine or methylphenidate hydrochloride (extended-release) and has not received sufficient benefit or has experienced intolerable side effects; or
 - 2.4.2 Patient is taking a currently subsidised formulation of dexamfetamine sulfate (immediate-release) which has not been effective due to significant administration and/or treatment adherence difficulties; or
 - 2.4.3 There is significant concern regarding the risk of diversion or abuse of immediate release dexamfetamine sulfate: or
 - 2.4.4 Patient is taking a currently subsidised formulation of methylphenidate hydrochloride (immediate-release or sustained release) which has not been effective due to significant administration and/or treatment adherence difficulties: or
 - 2.4.5 There is significant concern regarding the risk of diversion or abuse of immediate release methylphenidate hydrochloride; or
 - 2.4.6 Both:

a) Only on a controlled drug form

- 2.4.6.1 Patient would have been prescribed a subsidised formulation of methylphenidate hydrochloride (extended-release) but has been unable to access due to supply issues with methylphenidate hydrochloride (extended-release); and
- 2.4.6.2 Other alternative stimulant presentations (methylphenidate or dexamfetamine) are not appropriate:
- 2.5 Lisdexamfetamine dimesilate is not to be used in combination with another funded methylphenidate presentation.

METHYLPHENIDATE HYDROCHLORIDE - Special Authority see \$A2546 on the next page - Retail pharmacy

Tab modified-release 27 mg16.25	
	30
Tab modified-release 36 mg21.25	30
Tab modified-release 54 mg24.25	30
Tab immediate-release 5 mg	30 30 30

Tab sustained-release 20 mg.......10.95

Tab extended-release 36 mg......21.25

Tab extended-release 54 mg......24.25

b) Safety medicine; prescriber may determine dispensing frequency

1	Methylphenidate
	Sandoz XR

- ✓ Methylphenidate Sandoz XR
- ✓ Methylphenidate Sandoz XR
- ✓ Methylphenidate Sandoz XR
- ✓ Rubifen
- ✓ Rubifen
- ✓ Ritalin
- Methylphenidate ER - Teva
- ✓ Rubifen
- ✓ Rubifen SR
- ✓ Methylphenidate ER - Teva
- ✓ Methylphenidate ER
 - Teva
- ✓ Methylphenidate ER - Teva

30

30

30

30

30

NERVOUS SYSTEM

Subsidy		Fully	Brand or
(Manufacturer's Price)	S	Subsidised	Generic
\$	Per	✓	Manufacturer

⇒SA2546 Special Authority for Subsidy

Initial application — (ADHD in patients aged 5 years or over) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid without further renewal unless notified for applications meeting the following criteria: All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) in patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Fither
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients aged under 5 years) only from a paediatrician or psychiatrist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) in patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy*) only from a neurologist or respiratory specialist. Approvals valid without further renewal unless notified where the patient suffers from narcolepsy.

Note: *narcolepsy is not a registered indication for Methylphenidate ER - Teva or Methylphenidate Sandoz XR.

METHYLPHENIDATE HYDROCHLORIDE EXTENDED-RELEASE - Special Authority see SA2450 below - Retail pharmacy

- a) Only on a controlled drug form
- b) Safety medicine; prescriber may determine dispensing frequency

Tab extended-release 18 mg	58.96	30	Concerta
Tab extended-release 27 mg		30	✓ Concerta
Tab extended-release 36 mg	71.93	30	✓ Concerta
Tab extended-release 54 mg	86.24	30	✓ Concerta
Cap modified-release 10 mg	19.41	30	Ritalin LA
Cap modified-release 20 mg	27.72	30	Ritalin LA
Cap modified-release 30 mg		30	Ritalin LA
Cap modified-release 40 mg	38.67	30	Ritalin LA

⇒SA2450 Special Authority for Subsidy

Initial application — (ADHD) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid without further renewal unless notified for applications meeting the following criteria:

Fither:

- 1 All of the following:
 - 1.1 ADHD (Attention Deficit and Hyperactivity Disorder); and
 - 1.2 Diagnosed according to DSM-IV or ICD 10 criteria; and
 - 1.3 Either:
 - 1.3.1 Applicant is a paediatrician or psychiatrist; or
 - 1.3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing; and
 - 1.4 Fither:
 - 1.4.1 Patient is taking a currently subsidised formulation of methylphenidate hydrochloride (immediate-release or sustained-release) which has not been effective due to significant administration and/or difficulties with adherence; or
 - 1.4.2 There is significant concern regarding the risk of diversion or abuse of immediate-release methylphenidate hydrochloride; or

NERVOUS SYSTEM

Subsid	ly Fu	y Brand or
(Manufacture)	,	
\$	Per	Manufacturer

continued...

- 2 Both:
 - 2.1 Patient meets the Special Authority criteria for SA2411 methylphenidate hydrochloride; and
 - 2.2 Patient is unable to access other methylphenidate hydrochloride presentations under Special Authority criteria SA2411 due to an out of stock (see note).

Note: Criterion 2 is to permit short-term funding to cover an out-of-stock on tab extended-release Methylphenidate ER – Teva and tab sustained-release 20 mg Rubifen SR subsidised under

SA2411 (https://schedule.pharmac.govt.nz/2025/02/01/SA2411.pdf).

Initial application — (Narcolepsy*) only from a neurologist or respiratory specialist. Approvals valid without further renewal unless notified where the patient suffers from narcolepsy.

Note: *narcolepsy is not a registered indication for Concerta or Ritalin LA.

MODAFINIL - Special Authority see SA2451 below - Retail pharmacy

⇒SA2451 Special Authority for Subsidy

Initial application only from a neurologist or respiratory specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient has a diagnosis of narcolepsy and has excessive daytime sleepiness associated with narcolepsy occurring almost daily for three months or more; and
 - 1.2 Either:
 - 1.2.1 The patient has a multiple sleep latency test with a mean sleep latency of less than or equal to 10 minutes and 2 or more sleep onset rapid eye movement periods; or
 - 1.2.2 The patient has at least one of: cataplexy, sleep paralysis or hypnagogic hallucinations; and
 - 1.3 Fither:
 - 1.3.1 An effective dose of a subsidised formulation of methylphenidate or dexamfetamine has been trialled and discontinued because of intolerable side effects: or
 - 1.3.2 Methylphenidate and dexamfetamine are contraindicated; or
- 2 Both:
 - 2.1 Patient meets the Special Authority criteria for methylphenidate hydrochloride or methylphenidate hydrochloride extended-release for narcolepsy; and
 - 2.2 Patient is unable to access methylphenidate hydrochloride presentations due to an out of stock (see note).

Note: Criterion 2 is to permit short-term funding to cover an out-of-stock of methylphenidate hydrochloride or methylphenidate hydrochloride extended release.

Treatments for Dementia

DONEPEZIL HYDROCHLORIDE			
* Tab 5 mg	3.70	84	✓ Ipca-Donepezil
* Tab 10 mg	5.50	84	✓ Ipca-Donepezil
RIVASTIGMINE - Special Authority see SA2524 on the	next page – Retail pharmacy		
Patch 4.6 mg per 24 hour	49.40	30	✓ Rivastigmine Patch
			<u>BNM 5</u>
Patch 9.5 mg per 24 hour	49.40	30	✓ Rivastigmine Patch
			BNM 10

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	•	Manufacturer

⇒SA2524 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The patient has been diagnosed with dementia; and
- 2 The patient is contraindicated to or has experienced intolerable side effects from donepezil tablets.

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate; and
- 2 The patient has demonstrated a significant and sustained benefit from treatment.

Treatments for Substance Dependence

BUPRENORPHINE WITH NALOXONE - Special Authority see SA1203 below - Retail pharmacy

- a) No patient co-payment payable
- b) Safety medicine; prescriber may determine dispensing frequency

Tab sublingual 2 mg with naloxone 0.5 mg	,	28	✓ Buprenorphine Naloxone BNM
Tab sublingual 8 mg with naloxone 2 mg	26.86	28	✓ Buprenorphine Naloxone BNM

⇒SA1203 Special Authority for Subsidy

Initial application — (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health...

Initial application — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient will not be receiving methadone; and
- 3 Patient is currently enrolled in an opioid substitution treatment program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient has previously trialled but failed detoxification with buprenorphine with naloxone with relapse back to opioid use and another attempt is planned; and
- 3 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 Patient is or has been receiving maintenance therapy with buprenorphine with naloxone (and is not receiving methadone); and



Subsidy (Manufacturer's Price)	Suk	Fully	Brand or Generic
 \$	Per	√	Manufacturer

continued...

- 2 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health or is a medical practitioner authorised by the service to manage treatment in this patient.

Renewal — (Maintenance treatment where the patient has previously had an initial application for detoxification) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient received but failed detoxification with buprenorphine with naloxone; and
- 2 Maintenance therapy with buprenorphine with naloxone is planned (and patient will not be receiving methadone); and
- 3 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

BUPROPION HYDROCHLORIDE

Tab modified-release 150 mg	15.00	30	✓ Zyban
DISULFIRAM Tab 200 mg	236.40	100	✓ Antabuse
NALTREXONE HYDROCHLORIDE - Special Authority se	ee SA1408 below – Retail	pharmacy	
Tab 50 mg	83.33	30	✓ Naltraccord

⇒SA1408 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient is currently enrolled in a recognised comprehensive treatment programme for alcohol dependence; and
- 2 Applicant works in or with a community Alcohol and Drug Service contracted to Health NZ or accredited against the New Zealand Alcohol and Other Drug Sector Standard or the National Mental Health Sector Standard.

Renewal from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Compliance with the medication (prescriber determined); and
- 2 Any of the following:
 - 2.1 Patient is still unstable and requires further treatment: or
 - 2.2 Patient achieved significant improvement but requires further treatment; or
 - 2.3 Patient is well controlled but requires maintenance therapy.

NICOTINE

a) Nicotine will not be funded in amounts less than 4 weeks of treatment.

b) Note: Direct Provision by a pharmacist permitted under the provisi	ons in Part I of S	ection A.
Patch 7 mg - Up to 28 patch available on a PSO19	.62 28	Habitrol
Patch 14 mg - Up to 28 patch available on a PSO21	.57 28	Habitrol
Patch 14 mg for direct distribution only - [Xpharm]12	.49 7	Habitrol
Patch 21 mg - Up to 28 patch available on a PSO24	.72 28	Habitrol
Patch 21 mg for direct distribution only - [Xpharm]13	.19 7	Habitrol
Lozenge 1 mg - Up to 216 loz available on a PSO22	.53 216	Habitrol
Lozenge 1 mg for direct distribution only - [Xpharm]12	.89 36	Habitrol
Lozenge 2 mg - Up to 216 loz available on a PSO24	.68 216	Habitrol
Lozenge 2 mg for direct distribution only - [Xpharm]13	.25 36	Habitrol
Gum 2 mg (Fruit) - Up to 204 piece available on a PSO23	.02 204	Habitrol
Gum 2 mg (Fruit) for direct distribution only - [Xpharm]17	.57 96	Habitrol
Gum 2 mg (Mint) - Up to 204 piece available on a PSO23	.02 204	Habitrol
Gum 2 mg (Mint) for direct distribution only - [Xpharm]17.	.57 96	Habitrol
Gum 4 mg (Fruit) - Up to 204 piece available on a PSO25	.98 204	Habitrol
Gum 4 mg (Fruit) for direct distribution only - [Xpharm]23	.87 96	Habitrol
Gum 4 mg (Mint) - Up to 204 piece available on a PSO25	.98 204	Habitrol
Gum 4 mg (Mint) for direct distribution only - [Xpharm]23	.87 96	✓ Habitrol

NERVOUS SYSTEM

Subsidy	Full	y Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per 🗸	 Manufacturer 	

VARENICLINE TARTRATE - Special Authority see SA1845 below - Retail pharmacy

- a) A maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval, including the starter pack
- b) Varenicline will not be funded in amounts less than 4 weeks of treatment.
- c) The 6-month time period in which a patient can receive a funded 12-week course of varenicline tartrate starts from the date the Special Authority is approved.

Tab 0.5 mg × 11 and 1 mg × 42	16.67	53 OP	Champix
Tab 1 mg	17.62	56	Champix

⇒SA1845 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 Either:
 - 3.1 The patient has tried but failed to quit smoking after at least two separate trials of nicotine replacement therapy, at least one of which included the patient receiving comprehensive advice on the optimal use of nicotine replacement therapy; or
 - 3.2 The patient has tried but failed to quit smoking using bupropion or nortriptyline; and
- 4 The patient has not had a Special Authority for varenicline approved in the last 6 months; and
- 5 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 6 The patient is not pregnant; and
- 7 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

Renewal from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 It has been 6 months since the patient's previous Special Authority was approved; and
- 4 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 5 The patient is not pregnant; and
- 6 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

The patient must not have had an approval in the past 6 months.

Notes: a maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval.

This includes the 4-week 'starter' pack.

				_
Subsidy		Fully	Brand or	_
(Manufacturer's Price)	Subs	sidised	Generic	
\$	Per	1	Manufacturer	

Chemotherapeutic Agents

Alkylating Agents

BENDAMUSTINE HYDROCHLORIDE - PCT only - Specialist - Special Authorit	y see <mark>SA2398</mark>	3 below
Inj 25 mg vial50.05	1	Bendamustine Sandoz
77.00		✓ Ribomustin
Inj 100 mg vial200.20	1	✓ Bendamustine Sandoz
308.00		✓ Ribomustin
Inj 1 mg for ECP2.11	1 mg	✓ Baxter

⇒SA2398 Special Authority for Subsidy

Initial application — (CLL*) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has chronic lymphocytic leukaemia requiring treatment; and
- 2 Patient has ECOG performance status of 0-2; and
- 3 Bendamustine is to be administered at a maximum dose of 100 mg/m² on days 1 and 2 every 4 weeks for a maximum of 6 cycles.

Note: Indication marked with a * includes indications that are unapproved. 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL).

Initial application — (Indolent, Low-grade lymphomas) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: All of the following:

- 1 The patient has indolent low grade NHL requiring treatment; and
- 2 The patient has ECOG performance status of 0-2; and
- 3 Any of the following:
 - 3.1 Both:
 - 3.1.1 Patient is treatment naive; and
 - 3.1.2 Bendamustine is to be administered for a maximum of 6 cycles (in combination with rituximab when CD20+); or
 - 3.2 Both:
 - 3.2.1 Patient is refractory to or has relapsed within 12 months of a rituximab containing combined chemo-immunotherapy regimen; and
 - 3.2.2 Bendamustine is to be administered in combination with obinutuzumab for a maximum of 6 cycles; or
 - 3.3 All of the following:
 - 3.3.1 The patient has not received prior bendamustine therapy; and
 - 3.3.2 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
 - 3.3.3 Patient has had a rituximab treatment-free interval of 12 months or more; or
 - 3.4 Bendamustine is to be administered as monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Renewal — (Indolent, Low-grade lymphomas) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: Either:

- 1 Both:
 - 1.1 Patient is refractory to or has relapsed within 12 months of rituximab in combination with bendamustine; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	sidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1.2 Bendamustine is to be administered in combination with obinutuzumab for a maximum of 6 cycles; or
- 2 Both:
 - 2.1 Patients have not received a bendamustine regimen within the last 12 months; and
 - 2.2 Fither:
 - 2.2.1 Both:
 - 2.2.1.1 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
 - 2.2.1.2 Patient has had a rituximab treatment-free interval of 12 months or more; or
 - 2.2.2 Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Note: 'indolent, low-grade lymphomas' includes follicular, mantle cell, marginal zone and lymphoplasmacytic/ Waldenstrom's macroglobulinaemia.

Initial application — (**Hodgkin's lymphoma***) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has Hodgkin's lymphoma requiring treatment; and
- 2 Patient has a ECOG performance status of 0-2; and
- 3 Patient has received one prior line of chemotherapy; and
- 4 Patient's disease relapsed or was refractory following prior chemotherapy; and
- 5 Bendamustine is to be administered in combination with gemcitabine and vinorelbine (BeGeV) at a maximum dose of no greater than 90 mg/m2 twice per cycle, for a maximum of four cycles.

Note: Indications marked with * are unapproved indications.

BUSULFAN - PCT - Retail pharmacy-Specialist

CARBOPLATIN − PCT only − Specialist Inj 10 mg per ml, 45 ml vial	Tab 2 mg	89.25	100	✓ Myleran
Second	CARBOPLATIN - PCT only - Specialist			
Align Alig	Inj 10 mg per ml, 45 ml vial	25.73	1	✓ DBL Carboplatin
Inj 1 mg for ECP		32.59		✓ DBL Carboplatin
(DBL Carboplatin S29 № Inj 10 mg per ml, 45 ml vial to be delisted 1 January 2026) CARMUSTINE - PCT only - Specialist Inj 100 mg vial		48.50		Carbaccord
CARMUSTINE − PCT only − Specialist 710.00 1 ✓ BiCNU Inj 100 mg vial 710.00 100 mg OP ✓ Baxter CHLORAMBUCIL − PCT − Retail pharmacy-Specialist 29.06 25 ✓ Leukeran FC CISPLATIN − PCT only − Specialist 9.45 1 ✓ Cisplatin Accord Inj 1 mg per ml, 50 ml vial 9.45 1 ✓ Cisplatin Ebewe Inj 1 mg per ml, 100 ml vial 18.90 1 ✓ Cisplatin Accord 21.00 29.66 ✓ DBL Cisplatin	Inj 1 mg for ECP	0.06	1 mg	✓ Baxter
Inj 100 mg vial	(DBL Carboplatin S29 S29 Inj 10 mg per ml, 45 ml vial to be	delisted 1 January	2026)	
Inj 100 mg vial	CARMUSTINE - PCT only - Specialist			
Inj 100 mg for ECP		710.00	1	✓ BiCNU
Tab 2 mg 29.06 25 ✓ Leukeran FC CISPLATIN − PCT only − Specialist 9.45 1 ✓ Cisplatin Accord Inj 1 mg per ml, 50 ml vial 15.00 ✓ Cisplatin Ebewe Inj 1 mg per ml, 100 ml vial 18.90 1 ✓ Cisplatin Accord 21.00 ✓ Cisplatin Ebewe 29.66 ✓ DBL Cisplatin			100 mg OP	✓ Baxter
CISPLATIN − PCT only − Specialist Inj 1 mg per ml, 50 ml vial	CHLORAMBUCIL - PCT - Retail pharmacy-Specialist			
Inj 1 mg per ml, 50 ml vial 9.45 1 ✓ Cisplatin Accord 15.00 ✓ Cisplatin Ebewe Inj 1 mg per ml, 100 ml vial 18.90 1 ✓ Cisplatin Accord 21.00 ✓ Cisplatin Ebewe 29.66 ✓ DBL Cisplatin	Tab 2 mg	29.06	25	✓ Leukeran FC
Inj 1 mg per ml, 50 ml vial 9.45 1 ✓ Cisplatin Accord 15.00 ✓ Cisplatin Ebewe Inj 1 mg per ml, 100 ml vial 18.90 1 ✓ Cisplatin Accord 21.00 ✓ Cisplatin Ebewe 29.66 ✓ DBL Cisplatin	CISPLATIN - PCT only - Specialist			
15.00		9.45	1	 Cisplatin Accord
21.00 ✓ Cisplatin Ebewe 29.66 ✓ DBL Cisplatin	, ,			✓ Cisplatin Ebewe
29.66 ✓ DBL Cisplatin	Inj 1 mg per ml, 100 ml vial	18.90	1	✓ Cisplatin Accord
·	, •	21.00		✓ Cisplatin Ebewe
Inj 1 mg for ECP		29.66		✓ DBL Cisplatin
	Inj 1 mg for ECP	0.19	1 mg	✓ Baxter

A)	Subsidy Manufacturer's Price)		Fully Subsidised	Generic
	\$	Per		Manufacturer
CYCLOPHOSPHAMIDE			_	_
Tab 50 mg - PCT - Retail pharmacy-Specialist		50		Cyclonex
Inj 1 g vial - PCT - Retail pharmacy-Specialist		1		Endoxan
	127.80	6		Cytoxan
Inj 2 g vial – PCT only – Specialist		_ 1	_	<u>Endoxan</u>
Inj 1 mg for ECP - PCT only - Specialist	0.05	1 mg	•	Baxter
FOSFAMIDE - PCT only - Specialist				
Inj 1 g	96.00	1	✓	Holoxan
Inj 2 g		1		Holoxan
Inj 1 mg for ECP	0.10	1 mg	•	Baxter
OMUSTINE - PCT - Retail pharmacy-Specialist				
Cap 40 mg	880.00	20	1	Medac S29
/ELPHALAN				
Tab 2 mg - PCT - Retail pharmacy-Specialist	40.70	25	1	Alkeran
Inj 50 mg - PCT only - Specialist		1		Melpha
.,	67.80			Alkeran
DXALIPLATIN - PCT only - Specialist				
Inj 100 mg vial	25.01	1	1	Oxaliplatin Actavis
ing 100 mg viai	25.01	'	•	100
	110.00		✓	Oxaliplatin Ebewe
Inj 5 mg per ml, 20 ml vial	33.35	1	✓	Alchemy Oxaliplatin
	46.32		✓	Oxaliplatin Accord
Inj 1 mg for ECP	0.35	1 mg	/	Baxter
HIOTEPA - PCT only - Specialist				
Inj 15 mg vial	CBS	1	1	Bedford \$29
, - 3			1	Max Health \$29
				THIO-TEPA \$29
	398.00			Tepadina
Inj 100 mg vial		1		Max Health \$29
ing 100 mg viai	1,800.00	'		Tepadina
	1,000.00			Териина
Antimetabolites				
ZACITIDINE - PCT only - Specialist - Special Authority see SA2	479 below			
Ini 100 mg vial		1	1	Azacitidine Dr

		pecialist – Special Authority see SA2479 below	AZACITIDINE - PCT only - Spec
Azacitidine Dr	1	50.00	Inj 100 mg vial
Reddy's			
✓ Baxter	1 mg		Inj 1 mg for ECP

⇒SA2479 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The individual has intermediate or high risk MDS based on an internationally recognised scoring system; or
 - 1.2 The individual has chronic myelomonocytic leukaemia (based on an intermediate or high risk score from an internationally recognised scoring system or 10%-29% marrow blasts without myeloproliferative disorder); or
 - 1.3 The individual has acute myeloid leukaemia according to World Health Organisation Classification (WHO); and
- 2 The individual has an estimated life expectancy of at least 3 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where there is no evidence of disease progression.

	Subsidy		Fully	
(M	lanufacturer's Pri \$	ice) Subs Per	sidised •	
ALOUM FOLINATE	Ψ	1 61		Ivianulaciurei
ALCIUM FOLINATE	105.00	10	./	DBI Laviaguarin
Tab 15 mg - PCT - Retail pharmacy-Specialist	135.33	10	•	DBL Leucovorin
lai Carra annual 1 ant - DOT - Botail altra ann ann Carrainlist	17.10	-		Calcium
Inj 3 mg per ml, 1 ml – PCT – Retail pharmacy-Specialist		5		Hospira
Inj 10 mg per ml, 5 ml vial – PCT – Retail pharmacy-Specialist.		5		Eurofolic S29
Inj 50 mg - PCT - Retail pharmacy-Specialist	72.80	10	•	Leucovorin
				Pharmacia S29
Inj 10 mg per ml, 10 ml vial - PCT only - Specialist	163.35	5	1	Eurofolic S29
Inj 100 mg - PCT only - Specialist	94.90	10	1	Leucovorin
				Pharmacia S29
Inj 300 mg - PCT only - Specialist	21.55	1	•	Leucovorin DBL S29
Inj 10 mg per ml, 100 ml vial - PCT only - Specialist	139.48	1	/	Eurofolic §29
Inj 1 mg for ECP - PCT only - Specialist	0.14	1 mg	1	Baxter
PECITABINE - Retail pharmacy-Specialist				
Tab 150 mg	10.92	60	1	Capecitabine Viatris
Capecitabine Viatris to be Principal Supply on 1 February 2				
Tab 500 mg	50.96	120	1	Capecitabine Viatris
Capecitabine Viatris to be Principal Supply on 1 February 2				
ADRIBINE - PCT only - Specialist				
Inj 1 mg per ml, 10 ml	749 96	1	1	Leustatin
Inj 10 mg for ECP		10 mg OP		Baxter
	7 40.00	10 mg Oi	•	Duxtor
TARABINE	470.00	-	./	Diller
Inj 20 mg per ml, 5 ml vial – PCT – Retail pharmacy-Specialist.	472.00	5	•	Pfizer
Inj 100 mg per ml, 20 ml vial – PCT – Retail	40.00		,	
pharmacy-Specialist	48.80	1		Cytarabine DBL
Lei A man for EOD DOT and a Constallat	0.00	40		Pfizer
Inj 1 mg for ECP – PCT only – Specialist		10 mg	_	Baxter
Inj 100 mg intrathecal syringe for ECP - PCT only - Specialist.	94.40	100 mg OP	•	Baxter
JDARABINE PHOSPHATE			_	
Tab 10 mg - PCT - Retail pharmacy-Specialist		20		Fludara Oral
Inj 50 mg vial - PCT only - Specialist		5		Fludarabine Ebewe
Inj 50 mg for ECP - PCT only - Specialist	126.80	50 mg OP	•	Baxter
JOROURACIL				
Inj 50 mg per ml, 20 ml vial - PCT only - Specialist	10.51	1	1	Fluorouracil Accord
Inj 50 mg per ml, 50 ml vial - PCT only - Specialist		1	1	Fluorouracil Accord
Inj 50 mg per ml, 100 ml vial - PCT only - Specialist	19.36	1		Fluorouracil Accord
Inj 1 mg for ECP - PCT only - Specialist	0.41	100 mg	1	Baxter
MCITABINE HYDROCHLORIDE - PCT only - Specialist				
Inj 43.3 mg per ml (equivalent to 38 mg per ml gemcitabine),				
26.3 ml vial	18.94	1	1	DBL Gemcitabine
Inj 1 g		1		Gemcitabine Ebewe
Inj 1 mg for ECP		1 mg		Baxter
NOTECAN HYDROCHLORIDE - PCT only - Specialist		· ····ອ		
Inj 20 mg per ml, 5 ml vial	52 57	1	ſ	Accord
nij 20 nig per nii, 5 nii viai	52.57 71.44	ı		Irinotecan Actavis
	71.44		•	100
	100.00		./	Irinotecan-Rex
let 00 man annual 05 miletel	100.00			
Inj 20 mg per ml, 25 ml vial		1		Accord \$29
Inj 1 mg for ECP	0.54	1 mg	/	Baxter

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

		Subsidy (Manufacturer's \$	Price) Subs	Fully sidised	Generic
ME	RCAPTOPURINE				
	Tab 50 mg - PCT - Retail pharmacy-Specialist	19.50	25	1	Puri-nethol
	Oral suspension 20 mg per ml - Retail pharmacy-Specialist	_			
	Special Authority see SA1725 below		100 ml OP	1	Allmercap
				1	Xaluprine S29
>	SA1725 Special Authority for Subsidy				·
	tial application only from a paediatric haematologist or paedia	atric oncologist.	Approvals vali	d for 1	2 months where the patie
ec	uires a total dose of less than one full 50 mg tablet per day.				
	newal only from a paediatric haematologist or paediatric onco	logist. Approva	als valid for 12 n	nonths	where patient still requir
	otal dose of less than one full 50 mg tablet per day.	0 11			
ИF	THOTREXATE				
	Tab 2.5 mg - PCT - Retail pharmacy-Specialist	7.80	90	1	Trexate
*	Tab 10 mg - PCT - Retail pharmacy-Specialist		90		Trexate
*	Inj 2.5 mg per ml, 2 ml - PCT - Retail pharmacy-Specialist.		5		Methotrexate DBL
*	Inj 7.5 mg prefilled syringe		1	1	Methotrexate
	, 31 , 3				Sandoz
*	Inj 10 mg prefilled syringe	19.09	1	1	Methotrexate
	,		-		Sandoz
*	Inj 15 mg prefilled syringe	24 53	1	1	Methotrexate
•	ing to mg promod dyninge		•		Sandoz
*	Inj 20 mg prefilled syringe	16 64	1	1	Methotrexate
•	ing 20 mg promod syrings			-	Sandoz
*	Inj 25 mg prefilled syringe	20.72	1	1	Methotrexate
~	ing 25 mg premied synnige	20.72	'	•	Sandoz
*	Inj 30 mg prefilled syringe	55.00	1	1	Methotrexate
Τ.	ing 50 mg premied synnige	55.00	'	•	Sandoz
*	Inj 25 mg per ml, 2 ml vial - PCT - Retail pharmacy-Special	ict 20.00	5	1	Methotrexate DBL
•	inj 25 mg per mi, 2 mi viai – POT – netali pilamacy-special	151	5	•	Onco-Vial
Ł	Ini 05 mg nor ml 00 ml vial DCT Datail pharmagy Chaoir	aliat 45.00	1	./	DBL Methotrexate
*	Inj 25 mg per ml, 20 ml vial - PCT - Retail pharmacy-Specia	diist45.00	ı	•	Onco-Vial
v	Ini 100 ma nev ml 10 ml DCT Detail pharmacu Chaeialia	+ 05.00	1	./	
*	Inj 100 mg per ml, 10 ml – PCT – Retail pharmacy-Specialis	125.00	ı	V	Methotrexate Ebewe
*	Inj 100 mg per ml, 50 ml vial – PCT – Retail	07.00		,	Made during the Fire
	pharmacy-Specialist		1		Methotrexate Ebewe
*	Inj 1 mg for ECP – PCT only – Specialist		1 mg		Baxter Baxter
*	Inj 5 mg intrathecal syringe for ECP - PCT only - Specialist	19.06	5 mg OP	V	Daxier
PΕ	METREXED – PCT only – Specialist				
	Inj 100 mg vial		1		Pemetrexed-AFT
	Let FOO worded	60.89			Juno Pemetrexed
	Inj 500 mg vial		1		Pemetrexed-AFT
	Ini 1 mm for FCD	217.77	4		Juno Pemetrexed
	Inj 1 mg for ECP	0.11	1 mg	•	Baxter
Ή	IOGUANINE - PCT - Retail pharmacy-Specialist				
	Tab 40 mg	126.31	25	1	Lanvis

Other Cytotoxic Agents

AMSACRINE - PCT only - Specialist			
Inj 50 mg per ml, 1.5 ml ampoule	4,736.00	6	✓ Amsidine S29
Inj 75 mg	6,218.00	5	✓ AmsaLyo S29

	Subsidy		Fully	Brand or
	(Manufacturer's		sidised	
	\$	Per		Manufacturer
ANAGRELIDE HYDROCHLORIDE – PCT – Retail pharmacy				
Cap 0.5 mg	1,175.87	100	/	Agrylin
ARSENIC TRIOXIDE - PCT only - Specialist				
Inj 1 mg per ml, 10 ml vial		10		Phenasen
Inj 10 mg for ECP	481.70	10 mg OP	/	Baxter
BLEOMYCIN SULPHATE - PCT only - Specialist			_	
Inj 15,000 iu, vial	185.16	1	/	DBL Bleomycin
1:4000: (500	44.00	4 000 '	,	Sulfate
Inj 1,000 iu for ECP		1,000 iu	•	Baxter
BORTEZOMIB - PCT only - Specialist - Special Authority se			_	
Inj 3.5 mg vial		. 1	_	DBL Bortezomib
Inj 1 mg for ECP	22.26	1 mg	•	Baxter
⇒SA2355 Special Authority for Subsidy				
Initial application — (plasma cell dyscrasia) from any rele				
notified where the patient has plasma cell dyscrasia, not inclu	ding Waldenström	macroglobulinae	emia,	requiring treatment.
DACARBAZINE - PCT only - Specialist		_	_	
Inj 200 mg vial		1		DBL Dacarbazine
Inj 200 mg for ECP		200 mg OP	•	Baxter
DACTINOMYCIN [ACTINOMYCIN D] - PCT only - Specialis			_	
Inj 0.5 mg vial		1		Cosmegen
Inj 0.5 mg for ECP	255.00	0.5 mg OP	/	Baxter
DAUNORUBICIN - PCT only - Specialist			_	
Inj 2 mg per ml, 10 ml		1		Pfizer
Inj 18.7 mg for ECP		18.7 mg OP		Baxter
Inj 20 mg for ECP		20 mg OP		Baxter Pfizer
Inj 18.7 mg vial	171.93	1	•	Pilzer
(Pfizer Inj 2 mg per ml, 10 ml to be delisted 1 January 2026) (Baxter Inj 20 mg for ECP to be delisted 1 January 2026)				
DOCETAXEL – PCT only – Specialist	40.75	1	./	Deceteval Conden
Inj 20 mgInj 10 mg per ml, 8 ml vial		1		Docetaxel Sandoz DBL Docetaxel
Inj 20 mg per ml, 4 ml vial		1		Docetaxel
11) 20 11g por 111, 4 111 viai	20.00	•	•	Accord \$29
Inj 80 mg	195.00	1	/	Docetaxel Sandoz
Inj 1 mg for ECP		1 mg		Baxter
DOXORUBICIN HYDROCHLORIDE – PCT only – Specialist		9		
Inj 2 mg per ml, 5 ml vial		1	/	Doxorubicin Ebewe
Inj 2 mg per ml, 25 ml vial		i		Doxorubicin Ebewe
11) 2 119 por 111, 20 111 val	17.00	•		Arrow-Doxorubicin
Inj 2 mg per ml, 50 ml vial		1		Doxorubicin Ebewe
Inj 2 mg per ml, 100 ml vial		1	1	Arrow-Doxorubicin
	69.99		1	Doxorubicin Ebewe
			1	Baxter
Inj 1 mg for ECP		1 mg	•	Daxier
		1 mg	·	baxter
Inj 1 mg for ECP EPIRUBICIN HYDROCHLORIDE – PCT only – Specialist Inj 2 mg per ml, 5 ml vial	0.35	1 mg 1		Epirubicin Ebewe
EPIRUBICIN HYDROCHLORIDE - PCT only - Specialist	25.00	v	/	
EPIRUBICIN HYDROCHLORIDE - PCT only - Specialist Inj 2 mg per ml, 5 ml vial		1	\ \ \ \ \	Epirubicin Ebewe

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
ETOPOSIDE				
Cap 50 mg - PCT - Retail pharmacy-Specialist	340.73	20	/	Vepesid
Cap 100 mg - PCT - Retail pharmacy-Specialist	340.73	10	/	Vepesid
Inj 20 mg per ml, 5 ml vial - PCT - Retail pharmacy-Speciali	st7.90	1	✓	Rex Medical
Inj 1 mg for ECP - PCT only - Specialist	0.09	1 mg	•	Baxter
ETOPOSIDE PHOSPHATE - PCT only - Specialist				
Inj 100 mg (of etoposide base)	40.00	1	1	Etopophos
Inj 1 mg (of etoposide base) for ECP	0.47	1 mg	/	Baxter
HYDROXYUREA [HYDROXYCARBAMIDE] - PCT - Retail phart	macy-Specialist			
Cap 500 mg		100	1	Devatis
IBRUTINIB - Special Authority see SA2480 below - Retail pharm	nacy			
Tab 140 mg	3,217.00	30	1	Imbruvica
Tab 420 mg	9,652.00	30	1	Imbruvica

⇒SA2480 Special Authority for Subsidy

Initial application — (chronic lymphocytic leukaemia (CLL)) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Individual has chronic lymphocytic leukaemia (CLL) requiring therapy; and
- 2 Individual has not previously received funded ibrutinib; and
- 3 Ibrutinib is to be used as monotherapy; and
- 4 Any of the following:
 - 4.1 Both:
 - 4.1.1 There is documentation confirming that the individual has 17p deletion or TP53 mutation; and
 - 4.1.2 Individual has experienced intolerable side effects with venetoclax monotherapy; or
 - 4.2 All of the following:
 - 4.2.1 Individual has received at least one prior immunochemotherapy for CLL; and
 - 4.2.2 Individual's CLL has relapsed; and
 - 4.2.3 Individual has experienced intolerable side effects with venetoclax in combination with rituximab regimen; or
 - 4.3 Individual's CLL is refractory to or has relapsed following a venetoclax regimen.

Renewal — (chronic lymphocytic leukaemia (CLL)) from any relevant practitioner. Approvals valid for 12 months where there is no evidence of disease progression.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL) and B-cell prolymphocytic leukaemia (B-PLL)*. Indications marked with * are Unapproved indications.

IDADI	IDICINI	HYDROCHI	ODIDE
II IARI	IRIC IIV	HYDROCHI	()

Inj 5 mg vial – PCT only – Specialist	109.74	1	✓ Zavedos
Inj 10 mg vial - PCT only - Specialist	233.64	1	✓ Zavedos
Inj 1 mg for ECP - PCT only - Specialist	25.77	1 mg	✓ Baxter
LENALIDOMIDE (VIATRIS) - Special Authority see SA235	3 on the next page – Re	etail pharma	су
Cap 5 mg	76.92	21	✓ <u>Lenalidomide</u>
			<u>Viatris</u>
Cap 10 mg	50.30	21	✓ <u>Lenalidomide</u>
			<u>Viatris</u>
Cap 15 mg	62.13	21	✓ <u>Lenalidomide</u>
			<u>Viatris</u>
Cap 25 mg	65.09	21	✓ <u>Lenalidomide</u>
			Viatris

Subsidy	ı	ully	Brand or
(Manufacturer's Price)	Subsid	ised	Generic
\$	Per	1	Manufacturer

⇒SA2353 Special Authority for Subsidy

Initial application — (Plasma cell dyscrasia) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has plasma cell dyscrasia, not including Waldenström macroglobulinaemia, requiring treatment; and
- 2 Patient is not refractory to prior lenalidomide use.

Initial application — (Myelodysplastic syndrome) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has low or intermediate-1 risk myelodysplastic syndrome (based on IPSS or an IPSS-R score of less than 3.5) associated with a deletion 5g cytogenetic abnormality; and
- 2 Patient has transfusion-dependent anaemia.

Renewal — (Myelodysplastic syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has not needed a transfusion in the last 4 months; and
- 2 No evidence of disease progression.

М	ES	N	Δ
IVI	LO	I۷	М

Tab 400 mg - PCT - Retail pharmacy-Specialist31			Uromitexan
Tab 600 mg – PCT – Retail pharmacy-Specialist44			Uromitexan
Inj 100 mg per ml, 4 ml ampoule - PCT only - Specialist17			Uromitexan
Inj 100 mg per ml, 10 ml ampoule - PCT only - Specialist40	7.40	15	Uromitexan
Inj 1 mg for ECP - PCT only - Specialist	2.96 10	0 mg 🗸	Baxter
MITOMYCIN C - PCT only - Specialist			
Inj 5 mg vial51	7.65	1	Accord S29
. •		1	Mitomycin
			(Fresenius
			Kabi) S29
52	6.00	✓	Mitomycin
			(Sagent) S29
Inj 20 mg vial1,12	9.94	1	Teva
Inj 1 mg for ECP11		mg 🗸	Baxter
MITOZANTRONE - PCT only - Specialist			
Inj 2 mg per ml, 10 ml vial9	7.50	1	Mitozantrone Ebewe
Inj 1 mg for ECP		mg 🗸	Baxter
NIRAPARIB - Special Authority see SA2325 below - Retail pharmacy			
Wastage claimable			
Tab 100 mg	3.50	84	Zejula
Cap 100 mg			Zejula
ουρ 100 mg	0.07	•	Lojuiu

⇒SA2325 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has advanced high-grade serous* epithelial ovarian, fallopian tube, or primary peritoneal cancer; and
- 2 Patient has received at least one line** of treatment with platinum-based chemotherapy; and
- 3 Patient has experienced a partial or complete response to the preceding treatment with platinum-based chemotherapy; and

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsic	lised	Generic
\$	Per	1	Manufacturer

continued...

- 4 Patient has not previously received funded treatment with a PARP inhibitor; and
- 5 Either
 - 5.1 Treatment will be commenced within 12 weeks of the patient's last dose of the preceding platinum-based regimen;
 or
 - 5.2 Patient commenced treatment with niraparib prior to 1 May 2024; and
- 6 Treatment to be administered as maintenance treatment; and
- 7 Treatment not to be administered in combination with other chemotherapy.

Renewal from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 No evidence of progressive disease: and
- 2 Treatment to be administered as maintenance treatment; and
- 3 Treatment not to be administered in combination with other chemotherapy; and
- 4 Either:
 - 4.1 Treatment with niraparib to cease after a total duration of 36 months from commencement; or
 - 4.2 Treatment with niraparib is being used in the second-line or later maintenance setting.

Notes: * "high-grade serous" includes tumours with high-grade serous features or a high-grade serous component.

**A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments

OLAPARIB - Retail pharmacy-Specialist - Special Au	uthority see SA2163 below		
Tab 100 mg	3,701.00	56	✓ Lynparza
Tab 150 mg	3,701.00	56	✓ Lynparza

⇒SA2163 Special Authority for Subsidy

Initial application — (Ovarian cancer) only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a high-grade serous* epithelial ovarian, fallopian tube, or primary peritoneal cancer; and
- 2 There is documentation confirming pathogenic germline BRCA1 or BRCA2 gene mutation; and
- 3 Fither:
 - 3.1 All of the following:
 - 3.1.1 Patient has newly diagnosed, advanced disease; and
 - 3.1.2 Patient has received one line** of previous treatment with platinum-based chemotherapy; and
 - 3.1.3 Patient's disease must have experienced a partial or complete response to the first-line platinum-based regimen; or
 - 3.2 All of the following:
 - 3.2.1 Patient has received at least two lines** of previous treatment with platinum-based chemotherapy; and
 - 3.2.2 Patient has platinum sensitive disease defined as disease progression occurring at least 6 months after the last dose of the penultimate line** of platinum-based chemotherapy; and
 - 3.2.3 Patient's disease must have experienced a partial or complete response to treatment with the immediately preceding platinum-based regimen; and
 - 3.2.4 Patient has not previously received funded olaparib treatment; and
- 4 Treatment will be commenced within 12 weeks of the patient's last dose of the immediately preceding platinum-based regimen; and
- 5 Treatment to be administered as maintenance treatment; and
- 6 Treatment not to be administered in combination with other chemotherapy.

Renewal — (Ovarian cancer) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 12 months for applications meeting the following criteria:

Subsidy		Fully	Brand or
(Manufacturer's Price)	S	ubsidised	Generic
 \$	Per	✓	Manufacturer

continued...

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from treatment; and
- 2 Either:
 - 2.1 No evidence of progressive disease; or
 - 2.2 Evidence of residual (not progressive) disease and the patient would continue to benefit from treatment in the clinician's opinion; and
- 3 Treatment to be administered as maintenance treatment; and
- 4 Treatment not to be administered in combination with other chemotherapy; and
- 5 Either:
 - 5.1 Both:
 - 5.1.1 Patient has received one line** of previous treatment with platinum-based chemotherapy; and
 - 5.1.2 Documentation confirming that the patient has been informed and acknowledges that the funded treatment period of olaparib will not be continued beyond 2 years if the patient experiences a complete response to treatment and there is no radiological evidence of disease at 2 years; or
 - 5.2 Patient has received at least two lines** of previous treatment with platinum-based chemotherapy.

Notes: *Note "high-grade serous" includes tumours with high-grade serous features or a high-grade serous component.
**A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

ouppoint a dumonio.		
PACLITAXEL - PCT only - Specialist		
Inj 30 mg47.30	5	✓ Paclitaxel Ebewe
Inj 6 mg per ml, 16.7 ml vial19.59	1	✓ Anzatax
24.00		✓ Paclitaxel Ebewe
91.67		✓ Paclitaxel Actavis
Inj 150 mg26.69	1	✓ Paclitaxel Ebewe
137.50		✓ Anzatax
		✓ Paclitaxel Actavis
Inj 6 mg per ml, 50 ml vial37.89	1	✓ Anzatax
44.00		✓ Paclitaxel Ebewe
275.00		✓ Paclitaxel Actavis
Inj 1 mg for ECP0.17	1 mg	✓ Baxter
PEGASPARGASE – PCT only – Special Authority see SA1979 below		
Inj 750 iu per ml, 5 ml vial	1	Oncaspar LYO

⇒SA1979 Special Authority for Subsidy

Initial application — (Acute lymphoblastic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has newly diagnosed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

Initial application — (Lymphoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months where the patient has lymphoma requiring L-asparaginase containing protocols (e.g. SMILE).

Renewal — (Acute lymphoblastic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Roth:

- 1 The patient has relapsed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

PENTOSTATIN [DEOXYCOFORMYCIN] - PCT only - Specialist			
Ini 10 ma	CBS	1	✓ Ninent S29

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer	
POMALIDOMIDE - Special Authority see SA2354 below - Retai	l pharmacy				
Cap 1 mg	47.45	14	√ إ	Pomolide Pomolide	
	71.18	21	√ إ	Pomolide Pomolide	
Cap 2 mg	94.90	14	✓ [Pomolide	
	142.35	21	✓ [Pomolide	
Cap 3 mg	142.35	14	✓ [Pomolide Pomolide	
	213.53	21	√ إ	Pomolide Pomolide	
Cap 4 mg	189.81	14	√ إ	Pomolide Pomolide	
	284.71	21	✓]	Pomolide	

⇒SA2354 Special Authority for Subsidy

Initial application — (Relapsed/refractory plasma cell dyscrasia) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has relapsed or refractory plasma cell dyscrasia, not including Waldenström macroglobulinaemia, requiring treatment; and
- 2 Patient has not received prior funded pomalidomide.

Renewal — (Relapsed/refractory plasma cell dyscrasia) from any relevant practitioner. Approvals valid for 12 months where there is no evidence of disease progression.

50	✓ Natulan S29
5	✓ Temaccord
	✓ Temozolomide-
	Taro S29
5	✓ Temaccord
	✓ Apo-Temozolomide
5	✓ Temaccord
	✓ Apo-Temozolomide
5	✓ Temaccord
5	✓ Temaccord
	5 5 5

⇒SA2275 Special Authority for Subsidy

Initial application — (gliomas) only from a relevant specialist. Approvals valid for 12 months where the patient has a glioma. Renewal — (gliomas) only from a relevant specialist. Approvals valid for 12 months where treatment remains appropriate and patient is benefitting from treatment.

Initial application — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with metastatic or unresectable well-differentiated neuroendocrine tumour*; and
- 2 Temozolomide is to be given in combination with capecitabine; and
- 3 Temozolomide is to be used in 28 day treatment cycles for a maximum of 5 days treatment per cycle at a maximum dose of 200 mg/m² per day; and
- 4 Temozolomide to be discontinued at disease progression.

Renewal — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Roth:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued

Initial application — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 9 months where the patient has relapsed/refractory Ewing's sarcoma.

Renewal — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indication marked with a * is an unapproved indication. Temozolomide is not subsidised for the treatment of relapsed high grade glioma.

THALIDOMIDE - Retail pharmacy-Specialist - Special Autho	rity see SA2356 belov	V	
Cap 50 mg	378.00	28	Thalomid
Cap 100 mg	756.00	28	Thalomid

⇒SA2356 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months where the patient has plasma cell dyscrasia, not including Waldenström macroglobulinaemia, requiring treatment.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has obtained a response from treatment during the initial approval period.

Notes: Prescription must be written by a registered prescriber in the thalidomide risk management programme operated by the supplier.

Maximum dose of 400 mg daily as monotherapy or in a combination therapy regimen.

TRETINOIN

Cap 10 mg - PCT - Retail pharmacy-Specialist479.50	100	Vesanoid
Wastage claimable		
ENETOCLAX - Retail pharmacy-Specialist - Special Authority see SA2481 below		

VENETOCLAX - Retail pharmacy-Specialist - Special Authority see SA2481 below

Tab 14 × 10 mg, 7 × 50 mg, 21 × 100 mg1,771.86		✓ Venclexta
Tab 10 mg13.68		✓ Venclexta
Tab 50 mg	7 OP	✓ Venclexta
Tab 100 mg - Wastage claimable8,209.41	120	✓ Venclexta

⇒SA2481 Special Authority for Subsidy

Initial application — (relapsed/refractory chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 Individual has chronic lymphocytic leukaemia requiring treatment; and
- 2 Individual has received at least one prior therapy for chronic lymphocytic leukaemia; and
- 3 Individual has not previously received funded venetoclax; and
- 4 The individual's disease has relapsed; and
- 5 Venetoclax to be used in combination with six 28-day cycles of rituximab commencing after the 5-week dose titration schedule with venetoclax; and
- 6 Individual has an ECOG performance status of 0-2.

Renewal — (relapsed/refractory chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Treatment remains clinically appropriate and the individual is benefitting from and tolerating treatment; and
- 2 Venetoclax is to be discontinued after a maximum of 24 months of treatment following the titration schedule unless earlier discontinuation is required due to disease progression or unacceptable toxicity.

Initial application — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation*) from any

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
 \$	Per	√	

continued...

relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

- All of the following:
 - 1 Individual has previously untreated chronic lymphocytic leukaemia; and
 - 2 There is documentation confirming that individual has 17p deletion by FISH testing or TP53 mutation by sequencing; and
 - 3 Individual has an ECOG performance status of 0-2.

Renewal — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation*) from any relevant practitioner. Approvals valid for 6 months where the treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL)* and B-cell prolymphocytic leukaemia (B-PLL)*. Indications marked with * are Unapproved indications

Initial application — (previously untreated acute myeloid leukaemia) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Either:

- 1 The individual is currently on treatment with venetoclax and met all remaining special authority criteria prior to commencing treatment: or
- 2 All of the following:

VINBLASTINE SULPHATE

- 2.1 Individual has previously untreated acute myeloid leukaemia (see note a), according to World Health Organization (WHO) Classification: and
- 2.2 Venetoclax not to be used in combination with standard intensive remission induction chemotherapy; and
- 2.3 Venetoclax to be used in combination with azacitidine or low dose cytarabine.

Renewal — (previously untreated acute myeloid leukaemia) from any relevant practitioner. Approvals valid for 6 months where there is no evidence of disease progression. Notes:

- a) 'Acute myeloid leukaemia' includes myeloid sarcoma*
- b) Indications marked with * are Unapproved indications

Inj 1 mg per ml, 10 ml vial - PCT - Retail pharmacy-Specialist270	.37
Inj 1 mg for ECP - PCT only - Specialist6	.00
VINCRISTINE SULPHATE	
Ini 1 mg per ml. 1 ml vial – PCT – Retail pharmacy-Specialist74	.52

ing i mg for Eor i or only opecialist	i ilig	Daxiei
VINCRISTINE SULPHATE		
Inj 1 mg per ml, 1 ml vial - PCT - Retail pharmacy-Specialist74.52	5	DBL Vincristine Sulfate
Inj 1 mg per ml, 2 ml vial - PCT - Retail pharmacy-Specialist 102.73	5	DBL Vincristine Sulfate
Inj 1 mg for ECP - PCT only - Specialist12.60	1 mg	✓ Baxter
VINORELBINE		
Cap 20 mg32.10	1	✓ Vinorelbine Te Arai
Vinorelbine Te Arai to be Principal Supply on 1 February 2026		
Cap 30 mg42.80	1	✓ Vinorelbine Te Arai
Vinorelbine Te Arai to be Principal Supply on 1 February 2026		
Cap 80 mg80.00	1	✓ Vinorelbine Te Arai
Vinorelbine Te Arai to be Principal Supply on 1 February 2026		
Inj 10 mg per ml, 1 ml vial - PCT only - Specialist42.00	1	✓ Vinorelbine Ebewe
Inj 10 mg per ml, 5 ml vial - PCT only - Specialist155.00	1	✓ Vinorelbine Ebewe
168.00		✓ Navelbine S29 S29

1 mg

5

1 ma

✓ Hospira

✓ Baxter

✓ Baxter

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic

\$ Per ✓ Manufacturer

Protein-tyrosine Kinase Inhibitors

ALECTINIB - Retail pharmacy-Specialist - Special Authority see SA1870 below

Wastage claimable

⇒SA1870 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-small cell lung cancer; and
- 2 There is documentation confirming that the patient has an ALK tyrosine kinase gene rearrangement using an appropriate ALK test; and
- 3 Patient has an ECOG performance score of 0-2.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of progressive disease according to RECIST criteria: and
- 2 The patient is benefitting from and tolerating treatment.

AXITINIB - Special Authority see SA2458 below - Retail pharmacy

Wastage claimable

Tab 1 mg	536.40	28	Inlyta
Tab 5 mg	2,682.00	28	Inlyta

⇒SA2458 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 The disease is of predominant clear cell histology; and
- 3 The patient has documented disease progression following one previous line of treatment; and
- 4 The patient has ECOG performance status of 0-2.

Renewal from any relevant practitioner. Approvals valid for 4 months where there is no evidence of disease progression..

CRIZOTINIB - Special Authority see SA2547 below - Retail pharmacy

Cap 200 mg	7,250.00	60	Xalkori
Cap 250 mg	7,250.00	60	Xalkori

⇒SA2547 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Individual has locally advanced or metastatic, unresectable, non-squamous non-small cell lung cancer; and
- 2 Either:
 - 2.1 The individual has not received entrectinib: or
 - 2.2 Both:
 - 2.2.1 The individual has received an initial Special Authority approval for entrectinib and has discontinued entrectinib due to intolerance; and
 - 2.2.2 The cancer did not progress while the individual was on entrectinib; and
- 3 There is documentation confirming that the patient has a ROS1 rearrangement using an appropriate ROS1 test; and
- 4 Individual has ECOG performance score of 0-3; and
- 5 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Subsidy		Fully	Brand or
(Manufacturer's Pric	ce)	Subsidised	Generic
\$	Pe	r 🗸	Manufacturer

continued...

Renewal from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

- 1 Response to treatment has been determined by comparable radiological assessment following the most recent treatment period; and
- 2 No evidence of disease progression.

DABRAFENIB - Special Authority see SA2548 below - Retail pharmacy

Cap 50 mg	6,320.86	120	Tafinlar
Cap 75 mg	9,481.29	120	Tafinlar

⇒SA2548 Special Authority for Subsidy

Initial application — (stage III or IV resected melanoma - adjuvant) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 The individual has resected stage IIIB, IIIC, IIID or IV melanoma (excluding uveal) (see note a); or
 - 1.2 Both:
 - 1.2.1 The individual has received neoadjuvant treatment with a PD-1/PD-L1 inhibitor; and
 - 1.2.2 Adjuvant treatment with dabrafenib is required; and
- 2 The individual has not received prior funded systemic treatment in the adjuvant setting for stage IIIB, IIIC, IIID or IV melanoma; and
- 3 Treatment must be adjuvant to complete surgical resection; and
- 4 Treatment must be initiated within 13 weeks of surgical resection, unless delay is necessary due to post-surgery recovery (see note b); and
- 5 The individual has a confirmed BRAF mutation: and
- 6 Dabrafenib must be administered in combination with trametinib; and
- 7 The individual has ECOG performance score 0-2.

Notes:

- a) Stage IIIB, IIIC, IIID or IV melanoma defined as per American Joint Committee on Cancer (AJCC) 8th Edition
- b) Initiating treatment within 13 weeks of complete surgical resection means 13 weeks after resection (primary or lymphadenectomy)

Renewal — (stage III or IV resected melanoma - adjuvant) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Any of the following:

- 1 All of the following:
 - 1.1 No evidence of disease recurrence: and
 - 1.2 Dabrafenib must be administered in combination with trametinib; and
 - 1.3 Treatment to be discontinued at signs of disease recurrence or at completion of 12 months' total treatment course, including any systemic neoadjuvant treatment; or
- 2 All of the following:
 - 2.1 The individual has received adjuvant treatment with a BRAF/MEK inhibitor; and
 - 2.2 The individual has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
 - 2.3 The individual meets initial application criteria for dabrafenib for unresectable or metastatic melanoma; or
- 3 All of the following:
 - 3.1 The individual has received adjuvant treatment with a BRAF/MEK inhibitor; and
 - 3.2 The individual has received a BRAF/MFK inhibitor for unresectable or metastatic melanoma; and
 - 3.3 The individual meets renewal criteria for dabrafenib for unresectable or metastatic melanoma.

Initial application — (unresectable or metastatic melanoma) from any relevant practitioner. Approvals valid for 4 months for

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

applications meeting the following criteria:

All of the following:

- 1 The individual has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Baseline measurement of overall tumour burden is documented clinically and radiologically; and
- 3 The individual has ECOG performance score 0-2; and
- 4 The individual has confirmed BRAF mutation; and
- 5 Dabrafenib must be administered in combination with trametinib: and
- 6 Any of the following:
 - 6.1 The individual has been diagnosed in the metastatic or unresectable stage III or IV setting; or
 - 6.2 The individual did not receive treatment in the adjuvant setting with a BRAF/MEK inhibitor; or
 - 6.3 All of the following:
 - 6.3.1 The individual received treatment in the adjuvant setting with a BRAF/MEK inhibitor; and
 - 6.3.2 The individual did not experience disease recurrence while on treatment with that BRAF/MEK inhibitor; and
 - 6.3.3 The individual did not experience disease recurrence within six months of completing adjuvant treatment with a BRAF/MFK inhibitor.

Renewal — (unresectable or metastatic melanoma) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The individual's disease has had a complete response to treatment; or
 - 1.2 The individual's disease has had a partial response to treatment: or
 - 1.3 The individual has stable disease with treatment; and
- 2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period.

DASATINIB - Special Authority see SA2385 below - Retail pharmacy

Wastage claimable

Tab 20 mg	132.88	60	✓ Dasatinib-Teva
Tab 50 mg	304.13	60	✓ Dasatinib-Teva
Tab 70 mg	415.75	60	✓ Dasatinib-Teva

⇒SA2385 Special Authority for Subsidy

Initial application only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis or accelerated phase; or
- 2 The patient has a diagnosis of Philadelphia chromosome-positive acute lymphoid leukaemia (Ph+ ALL); or
- 3 Both:
 - 3.1 The patient has a diagnosis of CML in chronic phase; and
 - 3.2 Any of the following:
 - 3.2.1 Patient has documented treatment failure* with imatinib: or
 - 3.2.2 Patient has experienced treatment-limiting toxicity with imatinib precluding further treatment with imatinib; or
 - 3.2.3 Patient has high-risk chronic-phase CML defined by the Sokal or EURO scoring system.

Renewal only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Lack of treatment failure while on dasatinib*; and
- 2 Dasatinib treatment remains appropriate and the patient is benefiting from treatment.

Note: *treatment failure for CML as defined by Leukaemia Net Guidelines.

ENTRECTINIB – Special Authority see SA2532 on the next page – Retail pharmacy

Cap 200 mg9,610.00 90 ✔ Rozlytrek

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

⇒SA2532 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Individual has locally advanced or metastatic, unresectable, non-squamous non-small cell lung cancer; and
- 2 Fither
 - 2.1 The individual has not received crizotinib; or
 - 22 Roth
 - 2.2.1 The individual has received an initial Special Authority approval for crizotinib and has discontinued crizotinib due to intolerance: and
 - 2.2.2 The cancer did not progress while the individual was on crizotinib; and
- 3 There is documentation confirming that the patient has a ROS1 rearrangement using an appropriate ROS1 test; and
- 4 Individual has ECOG performance score of 0-3; and
- 5 Baseline measurement of overall tumour burden is documented clinically and radiologicallyy.

Renewal from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Response to treatment has been determined by comparable radiological assessment following the most recent treatment period; and
- 2 No evidence of disease progression.

ERLOTINIB - Retail pharmacy-Specialist - Special Authority	see SA2422 below		
Tab 100 mg	280.84	30	✓ Alchemy
Tab 150 mg	484.24	30	✓ Alchemy

⇒SA2422 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 There is documentation confirming that the disease expresses activating mutations of EGFR; and
- 3 Any of the following:
 - 3.1 Patient is treatment naive; or
 - 3.2 Patient has received prior treatment in the adjuvant setting and/or while awaiting EGFR results; or
 - 3.3 Both:
 - 3.3.1 The patient has discontinued osimertinib or gefitinib due to intolerance; and
 - 3.3.2 The cancer did not progress while on osimertinib or gefitinib.

Renewal from any relevant practitioner. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

GEFITINIB − Retail pharmacy-Specialist − Special Authority see SA2423 below
Tab 250 mg918.00 30 ✓ Iressa

⇒SA2423 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 Any of the following:
 - 2.1 Patient is treatment naive: or
 - 2.2 Patient has received prior treatment in the adjuvant setting and/or while awaiting EGFR results; or
 - 2.3 Both:
 - 2.3.1 The patient has discontinued osimertinib or erlotinib due to intolerance; and
 - 2.3.2 The cancer did not progress whilst on osimertinib or erlotinib; and
- 3 There is documentation confirming that disease expresses activating mutations of EGFR.

Renewal from any relevant practitioner. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

	Subsidy Manufacturer's Price) \$	S Per	Fully ubsidised	Brand or Generic Manufacturer
IMATINIB MESILATE				
* Cap 100 mg	44.93	60	✓ <u>In</u>	matinib-Rex
* Cap 400 mg	69.76	30	✓ In	natinib-Rex
LENVATINIB – Special Authority see SA2442 below – Retail pharr Wastage claimable	macy			
Cap 4 mg	2 407 40	30	1 1	envima
Cap 10 mg		30	_	envima
- CADA40 Created Avide suits for Cycleside				

SA2442 Special Authority for Subsidy

Initial application — **(thyroid cancer)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Patient is currently on treatment with lenvatinib and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 The patient has locally advanced or metastatic differentiated thyroid cancer; and
 - 2.2 Either:
 - 2.2.1 Patient must have symptomatic progressive disease prior to treatment; or
 - 2.2.2 Patient must progressive disease at critical anatomical sites with a high risk of morbidity or mortality where local control cannot be achieved by other measures; and
 - 2.3 Any of the following:
 - 2.3.1 A lesion without iodine uptake in a RAI scan: or
 - 2.3.2 Receiving cumulative RAI greater than or equal to 600 mCi; or
 - 2.3.3 Experiencing disease progression after a RAI treatment within 12 months; or
 - 2.3.4 Experiencing disease progression after two RAI treatments administered within 12 months of each other; and
 - 2.4 Patient has thyroid stimulating hormone (TSH) adequately supressed; and
 - 2.5 Patient is not a candidate for radiotherapy with curative intent; and
 - 2.6 Surgery is clinically inappropriate; and
 - 2.7 Patient has an ECOG performance status of 0-2.

Renewal — (thyroid cancer) from any relevant practitioner. Approvals valid for 6 months where there is no evidence of disease progression.

Initial application — (unresectable hepatocellular carcinoma) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has unresectable hepatocellular carcinoma; and
- 2 Patient has preserved liver function (Childs-Pugh A); and
- 3 Transarterial chemoembolisation (TACE) is unsuitable: and
- 4 Patient has an ECOG performance status of 0-2; and
- 5 Either:
 - 5.1 Patient has not received prior systemic therapy for their disease in the palliative setting; or
 - 5.2 Both:
 - 5.2.1 Patient has experienced treatment-limiting toxicity from treatment with atezolizumab with bevacizumab; and
 - 5.2.2 No disease progression since initiation of atezolizumab with bevacizumab.

Renewal — (unresectable hepatocellular carcinoma) from any relevant practitioner. Approvals valid for 6 months where there is no evidence of disease progression.

Initial application — (renal cell carcinoma) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

1 All of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsic	lised	Generic	
\$	Per	•	Manufacturer	

continued...

- 1.1 The patient has metastatic renal cell carcinoma; and
- 1.2 The disease is of predominant clear-cell histology; and
- 1.3 The patient has documented disease progression following one previous line of treatment; and
- 1.4 The patient has an ECOG performance status of 0-2; and
- 1.5 Lenvatinib is to be used in combination with everolimus; or
- 2 All of the following:
 - 2.1 Patient has received funded treatment with nivolumab for the second line treatment of metastatic renal cell carcinoma; and
 - 2.2 Patient has experienced treatment limiting toxicity from treatment with nivolumab; and
 - 2.3 Lenvatinib is to be used in combination with everolimus; and
 - 2.4 There is no evidence of disease progression.

Renewal — (renal cell carcinoma) from any relevant practitioner. Approvals valid for 4 months where there is no evidence of disease progression.

56 **✓ Rydapt**

⇒SA2342 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria: All of the following:

- 1 Patient has a diagnosis of acute myeloid leukaemia; and
- 2 Condition must be FMS tyrosine kinase 3 (FLT3) mutation positive; and
- 3 Patient must not have received a prior line of intensive chemotherapy for acute myeloid leukaemia; and
- 4 Patient is to receive standard intensive chemotherapy in combination with midostaurin only; and
- 5 Midostaurin to be funded for a maximum of 4 cycles.

NILOTINIB - Special Authority see SA2301 below - Retail pharmacy

⇒SA2301 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis, high risk chronic phase, or in chronic phase; and
- 2 Either:
 - 2.1 Patient has documented CML treatment failure* with a tyrosine kinase inhibitor (TKI); or
 - 2.2 Patient has experienced treatment limiting toxicity with a tyrosine kinase inhibitor (TKI) precluding further treatment; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

Note: *treatment failure as defined by Leukaemia Net Guidelines.

Renewal only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Lack of treatment failure while on nilotinib as defined by Leukaemia Net Guidelines; and
- 2 Nilotinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

OSIMERTINIB – Special Authority see SA2418 on the next page – Retail pharmacy		
Tab 40 mg9,310.00	30	

 Tab 40 mg
 9,310.00
 30
 ✓ Tagrisso

 Tab 80 mg
 9,310.00
 30
 ✓ Tagrisso

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA2418 Special Authority for Subsidy

Initial application — (NSCLC – first line) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Patient is currently on treatment with osimertinib and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 Patient has locally advanced or metastatic, incurable, non-squamous non-small cell lung cancer (NSCLC); and
 - 2.2 Any of the following:
 - 2.2.1 Patient is treatment naïve; or
 - 2.2.2 Patient has received prior chemotherapy in the adjuvant setting and/or while awaiting EGFR results; or
 - 222 Both
 - 2.2.3.1 The patient has discontinued gefitinib or erlotinib due to intolerance; and
 - 2.2.3.2 The cancer did not progress while on gefitinib or erlotinib; and
 - 2.3 There is documentation confirming that the cancer expresses activating mutations of EGFR; and
 - 2.4 Patient has an ECOG performance status 0-3; and
 - 2.5 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Renewal — (NSCLC – first line) from any relevant practitioner. Approvals valid for 6 months where response to or stable disease with treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period.

Initial application — (NSCLC – second line) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Patient is currently on treatment with osimertinib and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 Patient has locally advanced or metastatic, incurable, non-squamous non-small cell lung cancer (NSCLC); and
 - 2.2 Patient has an ECOG performance status 0-3; and
 - 2.3 The patient must have received previous treatment with erlotinib or gefitinib; and
 - 2.4 There is documentation confirming that the cancer expresses T790M mutation of EGFR following progression on or after erlotinib or gefitinib; and
 - 2.5 The treatment must be given as monotherapy; and
 - 2.6 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Renewal — (NSCLC – second line) from any relevant practitioner. Approvals valid for 6 months where response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period.

PALBOCICLIB - Special Authority see SA2345 below - Retail pharmacy

Wastage claimable			
Tab 75 mg	1,200.00	21	Palbociclib Pfizer
Tab 100 mg	1,200.00	21	Palbociclib Pfizer
Tab 125 mg	1,200.00	21	Palbociclib Pfizer

⇒SA2345 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Either:

- 1 All of the following:
 - 1.1 Patient has unresectable locally advanced or metastatic breast cancer; and
 - 1.2 There is documentation confirming disease is hormone-receptor positive and HER2-negative; and
 - 1.3 Patient has an ECOG performance score of 0-2; and
 - 1.4 Either:
 - 1.4.1 Disease has relapsed or progressed during prior endocrine therapy; or

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1.4.2 Both:
 - 1.4.2.1 Patient is amenorrhoeic, either naturally or induced, with endocrine levels consistent with a postmenopausal or without menstrual-potential state; and
 - 1.4.2.2 Patient has not received prior systemic treatment for metastatic disease; and
- 1.5 Treatment must be used in combination with an endocrine partner; and
- 1.6 Patient has not received prior funded treatment with a CDK4/6 inhibitor; or
- 2 All of the following:
 - 2.1 Patient has an active Special Authority approval for ribociclib; and
 - 2.2 Patient has experienced a grade 3 or 4 adverse reaction to ribociclib that cannot be managed by dose reductions and requires treatment discontinuation; and
 - 2.3 Treatment must be used in combination with an endocrine partner; and
 - 2.4 There is no evidence of progressive disease since initiation of ribociclib.

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 Treatment must be used in combination with an endocrine partner; and
- 2 There is no evidence of progressive disease since initiation of palbociclib.

PAZOPANIB – Special Authority see SA2429 below – F	Retail pharmacy		
Tab 200 mg	172.88	30	✓ Pazopanib Teva
Tab 400 mg	464.00	30	✓ Pazopanib Teva

⇒SA2429 Special Authority for Subsidy

Initial application only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient has metastatic renal cell carcinoma of predominantly clear cell histology; and
 - 1.2 Either:
 - 1.2.1 The patient is treatment naive; or
 - 1.2.2 The patient has only received prior cytokine treatment; and
 - 1.3 The patient has an ECOG performance score of 0-2; and The patient has intermediate or poor prognosis defined as:
 - 1.4 Any of the following:
 - 1.4.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
 - 1.4.2 Haemoglobin level < lower limit of normal; or
 - 1.4.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
 - 1.4.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
 - 1.4.5 Karnofsky performance score of less than or equal to 70; or
 - 1.4.6 2 or more sites of organ metastasis; and
 - 1.5 Pazopanib to be used for a maximum of 3 months; or
- 2 All of the following:
 - 2.1 The patient has metastatic renal cell carcinoma; and
 - 2.2 The patient has discontinued sunitinib within 3 months of starting treatment due to intolerance; and
 - 2.3 The cancer did not progress whilst on sunitinib; and
 - 2.4 Pazopanib to be used for a maximum of 3 months.

Renewal only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months where there is no evidence of disease progression.

	Subsidy (Manufacturer's Price) \$	S Per	Fully ubsidised	Brand or Generic Manufacturer	
RIBOCICLIB – Special Authority see SA2495 below – Retail Wastage claimable	pharmacy				
Tab 200 mg	1,883.00	21	✓ K	isqali	
•	3,767.00	42	✓ K	isqali	
	5,650.00	63	✓ K	isqali	

⇒SA2495 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Fither:

- 1 All of the following:
 - 1.1 Patient has unresectable locally advanced or metastatic breast cancer; and
 - 1.2 There is documentation confirming disease is hormone-receptor positive and HER2-negative; and
 - 1.3 Patient has an ECOG performance score of 0-2; and
 - 1.4 Either:
 - 1.4.1 Disease has relapsed or progressed during prior endocrine therapy; or
 - 1.4.2 Both:
 - 1.4.2.1 Patient is amenorrhoeic, either naturally or induced, with endocrine levels consistent with a postmenopausal or without menstrual-potential state; and
 - 1.4.2.2 Patient has not received prior systemic endocrine treatment for metastatic disease; and
 - 1.5 Treatment to be used in combination with an endocrine partner; and
 - 1.6 Patient has not received prior funded treatment with a CDK4/6 inhibitor; or
- 2 All of the following:
 - 2.1 Patient has an active Special Authority approval for palbociclib; and
 - 2.2 Patient has experienced a grade 3 or 4 adverse reaction to palbociclib that cannot be managed by dose reductions and requires treatment discontinuation; and
 - 2.3 Treatment must be used in combination with an endocrine partner; and
 - 2.4 There is no evidence of progressive disease since initiation of palbociclib.

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 Treatment must be used in combination with an endocrine partner; and
- 2 There is no evidence of progressive disease since initiation of ribociclib.

RUXOLITINIB - Special Authority see SA1890 below - Retail pharmacy

Wastage claimable		
Tab 5 mg2,500.00	56	✓ Jakavi
Tab 10mg5,000.00	56	Jakavi
Tab 15 mg	56	Jakavi
Tab 20 mg5,000.00	56	🗸 Jakavi

⇒SA1890 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has primary myelofibrosis or post-polycythemia vera myelofibrosis or post-essential thrombocythemia myelofibrosis; and
- 2 Fither:
 - 2.1 A classification of risk of intermediate-2 or high-risk myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS; or
 - 2.2 Both:
 - 2.2.1 A classification of risk of intermediate-1 myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted

(M	Subsidy anufacturer's Price)	Su	Fully bsidised	Brand or Generic
	\$	Per	1	Manufacturer

continued...

DIPSS: and

- 2.2.2 Patient has severe disease-related symptoms that are resistant, refractory or intolerant to available therapy; and
- 3 A maximum dose of 20 mg twice daily is to be given.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 A maximum dose of 20 mg twice daily is to be given.

SUNITINIB - Special Authority see SA2452 below - Retail ph	armacy		
Cap 12.5 mg	103.11	28	 Sunitinib Rex
	208.38		 Sunitinib Pfizer
Sunitinib Rex to be Principal Supply on 1 March 2026			
Cap 25 mg	203.15	28	 Sunitinib Rex
, ,	416.77		 Sunitinib Pfizer
Sunitinib Rex to be Principal Supply on 1 March 2026			
Cap 50 mg	343.19	28	 Sunitinib Rex
•	694 62		✓ Sunitinib Pfizer

(Sunitinib Pfizer Cap 12.5 mg to be delisted 1 March 2026) (Sunitinib Pfizer Cap 25 mg to be delisted 1 March 2026) (Sunitinib Pfizer Cap 50 mg to be delisted 1 May 2026)

⇒SA2452 Special Authority for Subsidy

Initial application — (RCC) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 The patient has not previously received funded sunitinib.

Initial application — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal tumour (GIST); and
- 2 Either:
 - 2.1 The patient's disease has progressed following treatment with imatinib; or
 - 2.2 The patient has documented treatment-limiting intolerance, or toxicity to, imatinib.

Renewal — (RCC) from any relevant practitioner. Approvals valid for 4 months where there is no evidence of disease progression.

Renewal — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

The patient has responded to treatment or has stable disease as determined by Choi's modified CT response evaluation criteria as follows:

- 1 Any of the following:
 - 1.1 The patient has had a complete response (disappearance of all lesions and no new lesions); or
 - 1.2 The patient has had a partial response (a decrease in size of 10% or more or decrease in tumour density in Hounsfield Units (HU) of 15% or more on CT and no new lesions and no obvious progression of non measurable disease); or

Subsidy		Fully	Brand or	
(Manufacturer's F	rice) Sub	sidised	Generic	
<u> </u>	Per	1	Manufacturer	

continued...

- 1.3 The patient has stable disease (does not meet criteria the two above) and does not have progressive disease and no symptomatic deterioration attributed to tumour progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: It is recommended that response to treatment be assessed using Choi's modified CT response evaluation criteria (J Clin Oncol, 2007, 25:1753-1759). Progressive disease is defined as either: an increase in tumour size of 10% or more and not meeting criteria of partial response (PR) by tumour density (HU) on CT; or: new lesions, or new intratumoral nodules, or increase in the size of the existing intratumoral nodules.

Renewal — (GIST pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal (GIST); and
- 2 The patient is clinically benifiting from treatment and continued treatment remains appropriate; and
- 3 Sunitinib is to be discontinued at progression; and
- 4 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

TRAMETINIB – Special Authority see SA2549 below – Retail pharmacy

Tab 0.5 mg	2,370.32	30	Mekinist
Tab 2 mg	9,481.29	30	✓ Mekinist

⇒SA2549 Special Authority for Subsidy

Initial application — (stage III or IV resected melanoma - adjuvant) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 The individual has resected stage IIIB, IIIC, IIID or IV melanoma (excluding uveal) (see note a); or
 - 1.2 Both:
 - 1.2.1 The individual has received neoadjuvant treatment with a PD-1/PD-L1 inhibitor; and
 - 1.2.2 Adjuvant treatment with trametinib is required; and
- 2 The individual has not received prior funded systemic treatment in the adjuvant setting for stage IIIB, IIIC, IIID or IV melanoma; and
- 3 Treatment must be adjuvant to complete surgical resection; and
- 4 Treatment must be initiated within 13 weeks of surgical resection, unless delay is necessary due to post-surgery recovery (see note b); and
- 5 The individual has a confirmed BRAF mutation; and
- 6 Trametinib must be administered in combination with dabrafenib; and
- 7 The individual has ECOG performance score 0-2.

Notes:

- a) Stage IIIB, IIIC, IIID or IV melanoma defined as per American Joint Committee on Cancer (AJCC) 8th Edition
- b) Initiating treatment within 13 weeks of complete surgical resection means 13 weeks after resection (primary or lymphadenectomy)

Renewal — (stage III or IV resected melanoma - adjuvant) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Any of the following:

- 1 All of the following:
 - 1.1 No evidence of disease recurrence: and
 - 1.2 Trametinib must be administered in combination with dabrafenib; and
 - 1.3 Treatment to be discontinued at signs of disease recurrence or at completion of 12 months' total treatment course, including any systemic neoadjuvant treatment; or

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer	
--	--

continued...

- 2 All of the following:
 - 2.1 The individual has received adjuvant treatment with a BRAF/MEK inhibitor; and
 - 2.2 The individual has metastatic or unresectable melanoma (excluding uveal) stage III or IV: and
 - 2.3 The individual meets initial application criteria for trametinib for unresectable or metastatic melanoma; or
 - 3 All of the following:
 - 3.1 The individual has received adjuvant treatment with a BRAF/MEK inhibitor; and
 - 3.2 The individual has received a BRAF/MEK inhibitor for unresectable or metastatic melanoma; and
 - 3.3 The individual meets renewal criteria for trametinib for unresectable or metastatic melanoma.

Initial application — (unresectable or metastatic melanoma) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The individual has metastatic or unresectable melanoma (excluding uveal melanoma) stage III or IV; and
- 2 Baseline measurement of overall turnour burden is documented clinically and radiologically; and
- 3 The individual has ECOG performance score 0-2; and
- 4 The individual has confirmed BRAF mutation; and
- 5 Trametinib must be administered in combination with dabrafenib; and
- 6 Any of the following:
 - 6.1 The individual has been diagnosed in the metastatic or unresectable stage III or IV setting; or
 - 6.2 The individual did not receive treatment in the adjuvant setting with a BRAF/MEK inhibitor; or
 - 6.3 All of the following:
 - 6.3.1 The individual received treatment in the adjuvant setting with a BRAF/MEK inhibitor; and
 - 6.3.2 The individual did not experience disease recurrence while on treatment with that BRAF/MEK inhibitor; and
 - 6.3.3 The individual did not experience disease recurrence within six months of completing adjuvant treatment with a BRAF/MEK inhibitor.

Renewal — (unresectable or metastatic melanoma) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The individual's disease has had a complete response to treatment; or
 - 1.2 The individual's disease has had a partial response to treatment; or
 - 1.3 The individual has stable disease with treatment; and
- 2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period.

Endocrine Therapy

For GnRH ANALOGUES - refer to HORMONE PREPARATIONS, Trophic Hormones, page 87

ABIRATERONE ACETATE - Retail pharmacy-Specialist - Special Authority see SA2118 below

Wastage claimable

Tab 250 mg4,276.19 120 ✓ Zvtiga

⇒SA2118 Special Authority for Subsidy

Initial application only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has prostate cancer; and
- 2 Patient has metastases; and

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per ✓	Manufacturer	

continued...

- 3 Patient's disease is castration resistant; and
- 4 Fither:
 - 4.1 All of the following:
 - 4.1.1 Patient is symptomatic: and
 - 4.1.2 Patient has disease progression (rising serum PSA) after second line anti-androgen therapy; and
 - 4.1.3 Patient has ECOG performance score of 0-1; and
 - 4.1.4 Patient has not had prior treatment with taxane chemotherapy; or
 - 4.2 All of the following:
 - 4.2.1 Patient's disease has progressed following prior chemotherapy containing a taxane; and
 - 4.2.2 Patient has ECOG performance score of 0-2; and
 - 4.2.3 Patient has not had prior treatment with abiraterone.

Renewal — (abiraterone acetate) only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Significant decrease in serum PSA from baseline; and
- 2 No evidence of clinical disease progression; and
- 3 No initiation of taxane chemotherapy with abiraterone; and
- 4 The treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 2 Abiraterone acetate to be discontinued at progression; and
- 3 No initiation of taxane chemotherapy with abiraterone; and
- 4 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

BICALUTAMIDE			4
Tab 50 mg	4.18	28	✓ Binarex
FLUTAMIDE			
Tab 250 mg	107.55	90	✓ Prostacur S29
•	119.50	100	✓ Flutamin
FULVESTRANT - Retail pharmacy-Specialist - Special Auth	ority see SA1895 bel	OW	
Inj 50 mg per ml, 5 ml prefilled syringe	181.00	2	✓ Fulvestrant EVER Pharma
	1,068.00		✓ Faslodex

(Faslodex Inj 50 mg per ml, 5 ml prefilled syringe to be delisted 1 May 2026)

⇒SA1895 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has oestrogen-receptor positive locally advanced or metastatic breast cancer; and
- 2 Patient has disease progression following prior treatment with an aromatase inhibitor or tamoxifen for their locally advanced or metastatic disease: and
- 3 Treatment to be given at a dose of 500 mg monthly following loading doses; and
- 4 Treatment to be discontinued at disease progression.

Sub: (Manufactu	,	,	
\$	Per	✓ Manufacturer	

continued...

Renewal only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains appropriate and patient is benefitting from treatment; and
- 2 Treatment to be given at a dose of 500 mg monthly; and
- 3 There is no evidence of disease progression.

OCTREOTIDE

OTTLOTIBL			
Inj 100 mcg per ml, 1 ml ampoule	32.71	5	✓ Max Health
			✓ Octreotide GH S29
			✓ Sun Pharma S29
Inj 100 mcg per ml, 1 ml vial	48.50	5	✓ Omega S29
Inj 50 mcg per ml, 1 ml vial		5	✓ Omega S29
Inj 500 mcg per ml, 1 ml vial		5	✓ Omega S29
Inj 50 mcg per ml, 1 ml ampoule	27.58	5	✓ Max Health
			✓ Octreotide GH S29
Inj 500 mcg per ml, 1 ml ampoule	113.10	5	✓ Max Health
			✓ Octreotide GH S29
			✓ Sun Pharma S29
AMOXIFEN CITRATE			
Tab 10 mg	15.00	60	✓ Tamoxifen Sandoz
Tab 20 mg	5.32	60	✓ Tamoxifen Sandoz

Long-acting Somatostatin Analogues

⇒SA2445 Special Authority for Subsidy

Initial application — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

TA * *

- 1 The patient has nausea* and vomiting* due to malignant bowel obstruction*; and
- 2 Treatment with antiemetics, rehydration, antimuscarinic agents, corticosteroids and analgesics for at least 48 hours has not been successful; and
- 3 Treatment to be given for up to 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 3 months where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Acromegaly) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has acromegaly; and
- 2 Either:
 - 2.1 Treatment with surgery and radiotherapy is not suitable or was unsuccessful; or
 - 2.2 Treatment is for an interim period while awaiting the beneficial effects of radiotherapy; and
- 3 Treatment with a dopamine agonist has been unsuccessful.

Renewal — (Acromegaly) from any relevant practitioner. Approvals valid for 2 years where iGF1 levels have decreased since starting treatment.

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Note: In patients with acromegaly, treatment should be discontinued if IGF1 levels have not decreased 3 months after treatment. In patients treated with radiotherapy treatment should be withdrawn every 2 years, for 1 month, for assessment of remission.

Treatment should be stopped where there is biochemical evidence of remission (normal IGF1 levels) following treatment withdrawal for at least 4 weeks

Initial application — (pre-operative acromegaly) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has acromegaly; and
- 2 Patient has a large pituitary tumour, greater than 10 mm at its widest; and
- 3 Patient is scheduled to undergo pituitary surgery in the next six months.

Initial application — (Other Indications) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 VIPomas and Glucagonomas for patients who are seriously ill in order to improve their clinical state prior to definitive surgery: or
- 2 Both:
 - 2.1 Gastrinoma: and
 - 2.2 Fither:
 - 2.2.1 Surgery has been unsuccessful; or
 - 2.2.2 Patient has metastatic disease after treatment with H2 antagonist or proton pump inhibitors has been unsuccessful; or
- 3 Both:
 - 3.1 Insulinomas: and
 - 3.2 Surgery is contraindicated or has not been successful; or
- 4 For pre-operative control of hypoglycaemia and for maintenance therapy; or
- 5 Both:
 - 5.1 Carcinoid syndrome (diagnosed by tissue pathology and/or urinary 5HIAA analysis); and
 - 5.2 Disabling symptoms not controlled by maximal medical therapy.

Note: The use of a long-acting somatostatin analogue in patients with fistulae, oesophageal varices, miscellaneous diarrhoea and hypotension will not be funded under Special Authority

Renewal — (Other Indications) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

LANREOTIDE - Special Authority see SA2445 on the prev	ious page – Retail phar	macy	
Inj 60 mg per 0.5 ml, 0.5 ml syringe	382.77	1	✓ Mytolac
			✓ Mytolac S29 S29
Inj 90 mg per 0.5 ml, 0.5 ml syringe	562.92	1	✓ Mytolac
Inj 120 mg per 0.5 ml, 0.5 ml syringe	646.70	1	✓ Mytolac
OCTREOTIDE LONG-ACTING - Special Authority see SA	2445 on the previous pa	age – Retail	pharmacy
Inj depot 10 mg prefilled syringe	438.40	1	✓ Sandostatin LAR
Inj depot 20 mg prefilled syringe	583.70	1	✓ Sandostatin LAR
Inj depot 30 mg prefilled syringe	670.80	1	✓ Sandostatin LAR
Aromatase Inhibitors			

ANASTROZOLE		

* Tab 1 mg4.39	30	✓ Anatrole
EXEMESTANE		
* Tab 25 mg	30	✓ Pfizer Exemestane

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	Brand or Generic Manufacturer
LETROZOLE * Tab 2.5 mg	4.36 4.67	28 30		Accord S29 Letrole

Immunosuppressants

Cytotoxic Immunosuppressants

AZATHIOPRINE		
* Tab 25 mg	60	✓ Azamun
Azamun to be Principal Supply on 1 February 2026		
* Tab 50 mg	100	Azamun
Azamun to be Principal Supply on 1 February 2026		
MYCOPHENOLATE MOFETIL		
Tab 500 mg35.90	50	✓ Cellcept
Cap 250 mg35.90	100	✓ Cellcept
Powder for oral liq 1 g per 5 ml - Subsidy by endorsement187.25	165 ml OP	✓ Cellcept
Microphonologic polynophy for and limited to substitute and anti-formational and	بلايينا المتنبية الملاحليات	ممانيم متما مما معامات

Mycophenolate powder for oral liquid is subsidised only for patients unable to swallow tablets and capsules, and when the prescription is endorsed accordingly.

Fusion Proteins

ETANERCEPT - Special Authority see SA2399 below - F	Retail pharmacy		
Inj 25 mg	690.00	4	Enbrel
Inj 25 mg autoinjector	690.00	4	Enbrel
Inj 50 mg autoinjector		4	Enbrel
Inj 50 mg prefilled syringe		4	Enbrel

⇒SA2399 Special Authority for Subsidy

Initial application — (adult-onset Still's disease) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Either:
 - 1.1.1 The patient has had an initial Special Authority approval for adalimumab for adult-onset Still's disease (AOSD): or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a Health NZ Hospital; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or tocilizumab; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or tocilizumab such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal anti-inflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

Subsidy		Fully	Brand or	_
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 The patient has a sustained improvement in inflammatory markers and functional status.

Initial application — (ankylosing spondylitis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for ankylosing spondylitis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for ankylosing spondylitis; or
- 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis present for more than six months; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.3 Patient has bilateral sacroiliitis demonstrated by plain radiographs, CT or MRI scan; and
 - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
 - 2.5 Either:
 - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
 - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender (see Notes); and
 - 2.6 A Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of initial application.

Average normal chest expansion corrected for age and gender:

18-24 years - Male: 7.0 cm: Female: 5.5 cm

25-34 years - Male: 7.5 cm; Female: 5.5 cm

35-44 years - Male: 6.5 cm: Female: 4.5 cm

45-54 years - Male: 6.0 cm; Female: 5.0 cm

55-64 years - Male: 5.5 cm; Female: 4.0 cm

65-74 years - Male: 4.0 cm; Female: 4.0 cm

75+ years - Male: 3.0 cm; Female: 2.5 cm

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	/	Manufacturer

continued...

- 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 3 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 4 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (polyarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for polyarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
 - 2.3 Any of the following:
 - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose): or
 - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Renewal — (polyarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance: and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for oligoarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for oligoarticular course JIA; or
- 2 All of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
<u> </u>	Per	✓	Manufacturer	

continued...

- 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
- 2.3 Any of the following:
 - 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose): or
 - 2.3.3 High disease activity (cJADAS10 score greater than 4) after a 6-month trial of methotrexate.

Renewal — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Fither:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab or secukinumab for psoriatic arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab or secukinumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab or secukinumab to meet the renewal criteria for adalimumab or secukinumab for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
 - 2.4 Either:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist.

	Subsidy	Fully	Brand or
(Manu	facturer's Price)	Subsidised	Generic
	\$ Pe	er 🗸	Manufacturer

continued...

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Fither:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Indications marked with * are unapproved indications.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

Initial application — (Arthritis - rheumatoid) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for rheumatoid arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects; or
 - 1.2.2 The patient has received insufficient benefit to meet the renewal criteria for rheumatoid arthritis; or
- 2 All of the following:
 - 2.1 Patient has had rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
 - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.3 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
 - 2.4 Patient has tried and not responded to at least three months of methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses unless contraindicated); and
 - 2.5 Either:
 - 2.5.1 Patient has tried and not responded to at least three months of methotrexate in combination with the maximum tolerated dose of ciclosporin; or

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 2.5.2 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with methotrexate; and
- 2.6 Fither:
 - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
 - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip.

Renewal — (Arthritis - rheumatoid) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Fither:
 - 2.1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (severe chronic plaque psoriasis) only from a dermatologist or any relevant practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria: Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe chronic plaque psoriasis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe chronic plaque psoriasis; or
- 2 All of the following:
 - 2.1 Any of the following:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.3 Patient has severe chronic localised genital or flexural plaque psoriasis where the plaques or lesions have been present for at least 6 months from the time of initial diagnosis, and with a Dermatology Life Quality Index (DLQI) score greater than 10; and
 - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
 - 2.3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
 - 2.4 The most recent PASI or DLQI assessment is no more than 1 month old at the time of application.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand, foot, genital or flexural areas at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and for the face, palm of a hand or sole of a foot the skin

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 Both:
 - 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 1.1.2 Either:
 - 1.1.2.1 Following each prior etanercept treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-treatment baseline value; or
 - 1.1.2.2 Following each prior etanercept treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline value; or
 - 1.2 Both:
 - 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 1.2.2 Either:
 - 1.2.2.1 Following each prior etanercept treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values: or
 - 1.2.2.2 Following each prior etanercept treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; or
 - 1.3 Both:
 - 1.3.1 Patient had severe chronic localised genital or flexural plague psoriasis at the start of treatment; and
 - 1.3.2 Either:
 - 1.3.2.1 The patient has experienced a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; or
 - 1.3.2.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing etanercept; and
- 2 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Note: A treatment course is defined as a minimum of 12 weeks of etanercept treatment

Initial application — (undifferentiated spondyloarthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undifferentiated peripheral spondyloarthritis* with active peripheral joint arthritis in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day (or maximum tolerated dose); and
- 4 Patient has tried and not responded to at least three months of leflunomide at a dose of up to 20 mg daily (or maximum tolerated dose); and
- 5 Any of the following:
 - 5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour measured no more than one month prior to the date of this application; or

Su	ubsidy	Fully	Brand or
(Manufac	cturer's Price) Subsic	lised	Generic
	\$ Per	✓	Manufacturer

continued...

5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Note: Indications marked with * are unapproved indications.

Renewal — (undifferentiated spondyloarthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or

ANTITHYMOCYTE GLOBULIN (FOUINE) - PCT only - Specialist

- 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg dose every 7 days.

Immune Modulators

/ (TITITITION OF THE GEODOLING (EGOINE) TOTOING	poolanot		
Inj 50 mg per ml, 5 ml	4,439.17	5	✓ ATGAM
BACILLUS CALMETTE-GUERIN (BCG) VACCINE - PCT	only - Specialist		
Subsidised only for bladder cancer.			
Inj 2-8 × 100 million CFU	149.37	1	✓ OncoTICE
Inj 40 mg per ml, vial	182.45	3	✓ SII-Onco-BCG S29

Monoclonal Antibodies

	/	525 below – Retail pharma	ADALIMUMAB (AMGEVITA) – Special Authority see SA25
✓ Amgevita	1	190.00	Inj 20 mg per 0.4 ml prefilled syringe
✓ Amgevita	2	375.00	Inj 40 mg per 0.8 ml prefilled pen
✓ Amgevita	2	375.00	Ini 40 mg per 0.8 ml prefilled syringe

⇒SA2525 Special Authority for Subsidy

Initial application — (Behcet's disease - severe) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

Jouri.

- 1 The patient has severe Behcet's disease* that is significantly impacting the patient's quality of life; and
- 2 Either:
 - 2.1 The patient has severe ocular, neurological, and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s); or
 - 2.2 The patient has severe gastrointestinal, rheumatological, and/or mucocutaneous symptoms and has not responded adequately to two or more treatments appropriate for the particular symptom(s).

Note: Indications marked with * are unapproved indications.

Initial application — (Hidradenitis suppurativa) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

1 Patient has hidradenitis suppurativa Hurley Stage II or Hurley Stage III lesions in distinct anatomic areas; and

	Subsidy	Fully	Brand or
(Manu	facturer's Price)	Subsidised	Generic
	\$ Pe	er 🗸	Manufacturer

continued...

- 2 Patient has tried, but had an inadequate response to at least a 90 day trial of systemic antibiotics or has demonstrated intolerance to or has contraindications for systemic antibiotics; and
- 3 Patient has 3 or more active lesions; and
- 4 The patient has a DLQI of 10 or more and the assessment is no more than 1 month old at time of application.

Renewal — (Hidradenitis suppurativa) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline; and
- 2 The patient has a DLQI improvement of 4 or more from baseline.

Initial application — (Plaque psoriasis - severe chronic) only from a dermatologist or any relevant practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria: Fither:

- 1 Both:
 - 1.1 Patient has had an initial Special Authority approval for etanercept for severe chronic plaque psoriasis; and
 - 1.2 Fither:
 - 1.2.1 Patient has experienced intolerable side effects; or
 - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for etanercept for severe chronic plaque psoriasis: or
- 2 All of the following:
 - 2.1 Any of the following:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a PASI score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.3 Patient has severe chronic localised genital or flexural plaque psoriasis where the plaques or lesions have been present for at least 6 months from the time of initial diagnosis, and with a Dermatology Life Quality Index (DLQI) score greater than 10; and
 - 2.2 Patient has tried, but had an inadequate response to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
 - 2.3 A PASI assessment or DLQI assessment has been completed for at least the most recent prior treatment course but no longer than 1 month following cessation of each prior treatment course and is no more than 1 month old at the time of application.

Renewal — (Plaque psoriasis - severe chronic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced a 75% or more reduction in PASI score, or is sustained at this level, when compared with the pre-treatment baseline value; or
 - 1.2.2 The patient has a DLQI improvement of 5 or more, when compared with the pre-treatment baseline value; or
- 2 Both:
 - 2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 2.2 Fither:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per ✓	Manufacturer

continued...

- 2.2.1 The patient has experienced reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
- 2.2.2 The patient has experienced reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre treatment baseline value; or
- 3 Both:
 - 3.1 Patient had severe chronic localised genital or flexural plague psoriasis at the start of treatment; and
 - 3.2 Either:
 - 3.2.1 The patient has experienced a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; or
 - 3.2.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing adalimumab.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporin, azathioprine, or methotrexate) and has not received an adequate response.

Note: Indications marked with * are unapproved indications.

Initial application — **(Crohn's disease - adults)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active Crohn's disease: and
- 2 Any of the following:
 - 2.1 Patient has a CDAI score of greater than or equal to 300, or HBI score of greater than or equal to 10; or
 - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
 - 2.4 Patient has an ileostomy or colostomy and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Renewal — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 CDAI score has reduced by 100 points from the CDAI score, or HBI score has reduced by 3 points, from when the patient was initiated on adalimumab; or
- 2 CDAI score is 150 or less, or HBI is 4 or less; or
- 3 The patient has demonstrated an adequate response to treatment, but CDAI score and/or HBI score cannot be assessed.

Initial application — (Crohn's disease - children) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has active Crohn's disease; and
- 2 Either:
 - 2.1 Patient has a PCDAI score of greater than or equal to 30; or
 - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub: Per	sidised •	Generic Manufacturer	

continued...

Renewal — (Crohn's disease - children) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab; or
- 2 PCDAI score is 15 or less; or
- 3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed.

Initial application — (Crohn's disease - fistulising) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Crohn's disease: and
- 2 Any of the following:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
 - 2.2 Patient has one or more rectovaginal fistula(e); or
 - 2.3 Patient has complex peri-anal fistula; and
- 3 A Baseline Fistula Assessment has been completed and is no more than 1 month old at the time of application.

Renewal — (Crohn's disease - fistulising) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 The number of open draining fistulae have decreased from baseline by at least 50%; or
- 2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain.

Initial application — (Ocular inflammation - chronic) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

_....

- 1 The patient has had an initial Special Authority approval for infliximab for chronic ocular inflammation; or
- 2 Both
 - 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
 - 2.2 Any of the following:
 - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective: or
 - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
 - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Renewal — (Ocular inflammation - chronic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 12 weeks' initial treatment; or
- 2 Following each 2 year treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 2 year treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.</p>

Initial application — (Ocular inflammation - severe) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	1	Manufacturer	

continued...

Either:

- 1 Patient has had an initial Special Authority approval for infliximab for severe ocular inflammation; or
- 2 Both:
 - 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
 - 2.2 Any of the following:
 - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
 - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
 - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Renewal — (Ocular inflammation - severe) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 2 year treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 2 year treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Initial application — (ankylosing spondylitis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 Patient has had an initial Special Authority approval for etanercept for ankylosing spondylitis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects; or
 - 1.2.2 The patient has received insufficient benefit to meet the renewal criteria for ankylosing spondylitis; or
- 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis for more than six months; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.3 Patient has bilateral sacroiliitis demonstrated by radiology imaging; and
 - 2.4 Patient has not responded adequately to treatment with two or more NSAIDs, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
 - 2.5 Either:
 - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following BASMI measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
 - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender; and
 - 2.6 A BASDAI of at least 6 on a 0-10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment and is no more than 1 month old at the time of application.

Renewal — (ankylosing spondylitis) from any relevant practitioner. Approvals valid for 2 years where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less.

Initial application — (Arthritis - oligoarticular course juvenile idiopathic) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

Subsidy (Manufacturer's Price)	Su	Fully bsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for oligoarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 Fither:
 - 1.2.1 Patient has experienced intolerable side effects: or
 - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for oligoarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
 - 2.3 Either:
 - 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose).

Renewal — (Arthritis - oligoarticular course juvenile idiopathic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - polyarticular course juvenile idiopathic) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has had an initial Special Authority approval for etanercept for polyarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 Fither:
 - 1.2.1 Patient has experienced intolerable side effects; or
 - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for polyarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
 - 2.3 Any of the following:
 - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Renewal — (Arthritis - polyarticular course juvenile idiopathic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

1 Following initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline: or

Subsidy Fully Brand or
(Manufacturer's Price) Subsidised Generic
\$ Per ✓ Manufacturer

continued...

2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - psoriatic) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 Patient has had an initial Special Authority approval for etanercept or secukinumab for psoriatic arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects; or
 - 1.2.2 The patient has received insufficient benefit from to meet the renewal criteria for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine or leflunomide at maximum tolerated doses (unless contraindicated); and
 - 2.4 Fither:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.5 Any of the following:
 - 2.5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an ESR greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (Arthritis - psoriatic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in swollen joint count from baseline and a clinically significant response in the opinion of the physician; or
- 2 Patient demonstrates at least a continuing 30% improvement in swollen joint count from baseline and a clinically significant response in the opinion of the treating physician.

Initial application — (Arthritis - rheumatoid) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for rheumatoid arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for rheumatoid arthritis; or
- 2 All of the following:
 - 2.1 Patient has had rheumatoid arthritis (either confirmed by radiology imaging, or the patient is CCP antibody positive) for six months duration or longer; and
 - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sub	sidised	Generic
\$	Per	•	Manufacturer

continued...

- 2.3 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
- 2.4 Patient has tried and not responded to at least three months of methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate at maximum tolerated doses (unless contraindicated); and
- 2.5 Either:
 - 2.5.1 Patient has tried and not responded to at least three months of methotrexate in combination with the maximum tolerated dose of ciclosporin (unless contraindicated); or
 - 2.5.2 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide (unless contraindicated) alone or in combination with methotrexate; and
- 2.6 Fither:
 - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
 - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip.

Renewal — (Arthritis - rheumatoid) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Initial application — (Still's disease - adult-onset (AOSD)) only from a rheumatologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept and/or tocilizumab for AOSD; and
 - 1.2 Fither:
 - 1.2.1 Patient has experienced intolerable side effects from etanercept and/or tocilizumab; or
 - 1.2.2 Patient has received insufficient benefit from at least a three-month trial of etanercept and/or tocilizumab; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria; and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, NSAIDs and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Initial application — (ulcerative colitis) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active ulcerative colitis; and
- 2 Fither:
 - 2.1 Patient's SCCAI score is greater than or equal to 4: or
 - 2.2 Patient's PUCAI score is greater than or equal to 20; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from prior therapy with immunomodulators and systemic corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Fither:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	osidised	Generic	
\$	Per	1	Manufacturer	

continued...

- 1 The SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on biologic therapy; or
- 2 The PUCAI score has reduced by 10 points or more from the PUCAI score when the patient was initiation on biologic therapy.

Initial application — (undifferentiated spondyloarthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undifferentiated peripheral spondyloarthritis* with active peripheral joint arthritis in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2 Patient has tried and not responded to at least three months of each of methotrexate, sulfasalazine and leflunomide, at maximum tolerated doses (unless contraindicated); and
- 3 Any of the following:
 - 3.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application: or
 - 3.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application; or
 - 3.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Note: Indications marked with * are unapproved indications

Renewal — (undifferentiated spondyloarthritis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response in the opinion of the treating physician.

Initial application — (inflammatory bowel arthritis – axial) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has axial inflammatory pain for six months or more; and
- 3 Patient is unable to take NSAIDs; and
- 4 Patient has unequivocal sacroiliitis demonstrated by radiological imaging or MRI; and
- 5 Patient has not responded adequately to prior treatment consisting of at least 3 months of an exercise regime supervised by a physiotherapist; and
- 6 A BASDAI of at least 6 on a 0-10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment.

Renewal — (inflammatory bowel arthritis – axial) from any relevant practitioner. Approvals valid for 2 years where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less.

Initial application — (inflammatory bowel arthritis – peripheral) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has active arthritis in at least four joints from the following: hip, knee, ankle, subtalar, tarsus, forefoot, wrist, elbow, shoulder, sternoclavicular; and
- 3 Patient has tried and not experienced a response to at least three months of methotrexate, or azathioprine at a maximum tolerated dose (unless contraindicated); and

bsidy urer's Price) Subs	Fully	Brand or Generic
 \$ Per	•	Manufacturer

continued...

- 4 Patient has tried and not experienced a response to at least three months of sulfasalazine at a maximum tolerated dose (unless contraindicated); and
- 5 Any of the following:
 - 5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application: or
 - 5.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application: or
 - 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (inflammatory bowel arthritis – peripheral) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 Patient has experienced at least a continuing 30% improvement in active joint count from baseline in the opinion of the treating physician.

ADALIMUMAB (HUMIRA - ALTERNATIVE BRAND) - Special Authority see SA2157 below - Retail pharmacy

Inj 20 mg per 0.2 ml prefilled syringe	595.50	2	✓ Humira
Inj 40 mg per 0.4 ml prefilled pen	595.50	2	✓ HumiraPen
Inj 40 mg per 0.4 ml prefilled syringe	595.50	2	Humira

⇒SA2157 Special Authority for Subsidy

Initial application — (Behcet's disease – severe) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Behcet's disease – severe) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient has had a good clinical response to treatment with measurably improved guality of life; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Hidradenitis suppurativa) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
 - 4 Adalimumab to be administered at doses no greater than 40 mg every 7 days. Fortnightly dosing has been considered.

Renewal — (Hidradenitis suppurativa) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline: and
- 2 The patient has a Dermatology Quality of Life Index improvement of 4 or more from baseline; and
- 3 Adalimumab is to be administered at doses no greater than 40mg every 7 days. Fortnightly dosing has been considered.

Initial application — (Psoriasis - severe chronic plaque) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Psoriasis - severe chronic plaque) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Fither:
 - 1.1 Both:
 - 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 1.1.2 Either:
 - 1.1.2.1 Following each prior adalimumab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-adalimumab treatment baseline value: or
 - 1.1.2.2 Following each prior adalimumab treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline value; or
 - 1.2 Both:
 - 1.2.1 Patient had severe chronic plague psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment: and
 - 1.2.2 Fither:
 - 1.2.2.1 Following each prior adalimumab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values: or
 - 1.2.2.2 Following each prior adalimumab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-adalimumab treatment baseline value: and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Pvoderma gangrenosum) only from a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

1 Fither:

Subsidy (Manufacturer's Price)	Fı Subsidis	ılly ed	Brand or Generic	_
\$	Per	✓	Manufacturer	

continued...

- 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks
- 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 A maximum of 8 doses.

Renewal — (Pyoderma gangrenosum) only from a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient has demonstrated clinical improvement and continues to require treatment; and
- 2 A maximum of 8 doses.

Initial application — (Crohn's disease - adult) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevitat; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
 - 1.3 Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment: and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Crohn's disease - adult) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on adalimumab; or
 - 1.2 CDAI score is 150 or less; or
- 1.3 The patient has demonstrated an adequate response to treatment, but CDAI score cannot be assessed; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Crohn's disease - children) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
 - 1.3 Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment: and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per ✓	Manufacturer

continued...

Renewal — (Crohn's disease - children) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab; or
 - 1.2 PCDAI score is 15 or less: or
 - 1.3 The patient has demonstrated an adequate response to treatment, but PCDAI score cannot be assessed; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Crohn's disease - fistulising) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
 - 1.3 Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Crohn's disease - fistulising) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
 - 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Ocular inflammation – chronic) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with Amgevita, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
 - 1.3 Patient has uveitis and is considered to be at risk of vision loss if they were to change treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Ocular inflammation – chronic) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has had a good clinical response following 12 weeks' initial treatment; or

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	/	Manufacturer

continued...

- 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
- 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and</p>
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Ocular inflammation – severe) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with Amgevita, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
 - 1.3 Patient has uveitis and is considered to be at risk of vision loss if they were to change treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Ocular inflammation – severe) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has had a good clinical response following 3 initial doses; or
 - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
 - 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and</p>
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita); and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

Initial application — (Arthritis – oligoarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

Renewal — (Arthritis – oligoarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months where the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline

Initial application — (Arthritis - polyarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

Renewal — (Arthritis - polyarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months where the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - psoriatic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Arthritis - psoriatic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Subsidy		Fully	Brand or	
(Manufacturer's Pr	ice) S	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Initial application — (Arthritis – rheumatoid) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Fither:
 - 4.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or
 - 4.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

Renewal — (Arthritis – rheumatoid) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 2 Fithe
 - 2.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or
 - 2.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

Initial application — (Still's disease – adult-onset (AOSD)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

Renewal — (Still's disease – adult-onset (AOSD)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months where the patient has demonstrated a sustained improvement in inflammatory markers and functional status.

AFLIBERCEPT – Special Authority see SA2550 below – Retail pharmacy
Inj 40 mg per ml, 0.1 ml vial......1,250.00

1 ✓ Eylea

⇒SA2550 Special Authority for Subsidy

Initial application — (diabetic macular oedema) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has centre involving diabetic macular oedema (DMO); and
- 2 Patient's disease is non responsive to 4 doses of intravitreal bevacizumab when administered 4-6 weekly; and
- 3 Patient has reduced visual acuity between 6/9 6/36 with functional awareness of reduction in vision; and
- 4 Patient has DMO within central OCT (ocular coherence tomography) subfield > 350 micrometers; and

Subsidy	F	ully	Brand or	_
(Manufacturer's Price)	Subsidi	sed	Generic	
\$	Per	1	Manufacturer	

continued...

- 5 There is no centre-involving sub-retinal fibrosis or foveal atrophy; and
- 6 Patient has not previously been treated with faricimab for longer than 3 months.

Renewal — (diabetic macular oedema) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 There is stability or two lines of Snellen visual acuity gain; and
- 2 There is structural improvement on OCT scan (with reduction in intra-retinal cysts, central retinal thickness, and sub-retinal fluid): and
- 3 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 4 There is no centre-involving sub-retinal fibrosis or foveal atrophy.

Initial application — (wet age related macular degeneration) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Wet age-related macular degeneration (wet AMD); or
 - 1.1.2 Polypoidal choroidal vasculopathy; or
 - 1.1.3 Choroidal neovascular membrane from causes other than wet AMD: and
 - 1.2 Either:
 - 1.2.1 The patient has developed severe endophthalmitis or severe posterior uveitis following treatment with bevacizumab: or
 - 1.2.2 There is worsening of vision or failure of retina to dry despite three intraocular injections of bevacizumab four weeks apart; and
 - 1.3 There is no structural damage to the central fovea of the treated eye; and
 - 1.4 Patient has not previously been treated with ranibizumab or faricimab for longer than 3 months; or
- 2 Either:
 - 2.1 Patient has current approval to use ranibizumab or faricimab for treatment of wAMD and was found to be intolerant within 3 months; or
 - 2.2 Patient has previously* (*before June 2018) received treatment with ranibizumab for wAMD and disease was stable while on treatment.

Renewal — (wet age related macular degeneration) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Documented benefit must be demonstrated to continue; and
- 2 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 3 There is no structural damage to the central fovea of the treated eye.

BENRALIZUMAB - Special Authority see SA2151 below - Retail pharmacy

⇒SA2151 Special Authority for Subsidy

Initial application — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Patient must have a diagnosis of severe eosinophilic asthma documented by a respiratory physician or clinical immunologist; and
- 3 Conditions that mimic asthma eg. vocal cord dysfunction, central airway obstruction, bronchiolitis etc. have been

Subsidy		Fully	Brand or	_
(Manufacturer's Price))	Subsidised	Generic	
\$	Per	1	Manufacturer	

continued...

excluded: and

- 4 Patient has a blood eosinophil count of greater than 0.5 x 10^9 cells/L in the last 12 months; and
- 5 Patient must be adherent to optimised asthma therapy including inhaled corticosteroids (equivalent to at least 1000 mcg per day of fluticasone propionate) plus long-acting beta-2 agonist, or budesonide/formoterol as part of the anti-inflammatory reliever therapy plus maintenance regimen, unless contraindicated or not tolerated; and
- 6 Either:
 - 6.1 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral corticosteroids; or
 - 6.2 Patient has received continuous oral corticosteroids of at least the equivalent of 10 mg per day over the previous 3 months; and
- 7 Treatment is not to be used in combination with subsidised mepolizumab; and
- 8 Patient has an Asthma Control Test (ACT) score of 10 or less. Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 52 weeks after the first dose to assess response to treatment; and
- 9 Fither:
 - 9.1 Patient has not previously received an anti-IL5 biological therapy for their severe eosinophilic asthma; or
 - 9.2 Both:
 - 9.2.1 Patient was refractory or intolerant to previous anti-IL5 biological therapy; and
 - 9.2.2 Patient was not eligible to continue treatment with previous anti-IL5 biological therapy and discontinued within 12 months of commencing treatment.

Renewal — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 Either:
 - 2.1 Exacerbations have been reduced from baseline by 50% as a result of treatment with benralizumab; or
 - 2.2 Reduction in continuous oral corticosteroid use by 50% or by 10 mg/day while maintaining or improving asthma control.

BEVACIZUMAB - PCT only - Special Authority see SA2453 below

✓ Vegzelma	1	Inj 25 mg per ml, 4 ml vial69.00
✓ Vegzelma	1	Inj 25 mg per ml, 16 ml vial276.00
✓ Baxter	1 mg	Inj 1 mg for ECP

⇒SA2453 Special Authority for Subsidy

Initial application — (unresectable hepatocellular carcinoma) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Patient is currently on treatment with bevacizumab, and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 Patient has locally advanced or metastatic, unresectable hepatocellular carcinoma; and
 - 2.2 Patient has preserved liver function (Child-Pugh A); and
 - 2.3 Transarterial chemoembolisation (TACE) is unsuitable; and
 - 2.4 Any of the following:
 - 2.4.1 Patient has not received prior systemic therapy for the treatment of hepatocellular carcinoma; or
 - 2.4.2 Patient received funded lenvatinib before 1 March 2025; or
 - 2.4.3 Both:
 - 2.4.3.1 Patient has experienced treatment-limiting toxicity from treatment with lenvatinib; and

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

- 2.4.3.2 No disease progression since initiation of lenvatinib; and
- 2.5 Patient has an ECOG performance status of 0-2; and
- 2.6 To be given in combination with atezolizumab.

Renewal — (unresectable hepatocellular carcinoma) from any relevant practitioner. Approvals valid for 6 months where there is no evidence of disease progression.

Initial application — (advanced or metastatic ovarian cancer) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 The patient has FIGO Stage IV epithelial ovarian, fallopian tube, or primary peritoneal cancer; or
 - 1.2 Both:
 - 1.2.1 The patient has previously untreated advanced (FIGO Stage IIIB or IIIC) epithelial ovarian, fallopian tube, or primary peritoneal cancer; and
 - 1.2.2 Either:
 - 1.2.2.1 Debulking surgery is inappropriate; or
 - 1.2.2.2 The cancer is sub-optimally debulked (maximum diameter of any gross residual disease greater than 1cm); and
- 2 Bevacizumab to be administered at a maximum dose of 15 mg/kg every three weeks; and
- 3 18 weeks concurrent treatment with chemotherapy is planned.

Renewal — (advanced or metastatic ovarian cancer) from any relevant practitioner. Approvals valid for 4 months where there is no evidence of disease progression.

Initial application — (Recurrent Respiratory Papillomatosis) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Maximum of 6 doses: and
- 2 The patient has recurrent respiratory papillomatosis; and
- 3 The treatment is for intra-lesional administration.

Renewal — (Recurrent Respiratory Papillomatosis) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Maximum of 6 doses; and
- 2 The treatment is for intra-lesional administration; and
- 3 There has been a reduction in surgical treatments or disease regrowth as a result of treatment.

Initial application — (Ocular Conditions) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 Ocular neovascularisation: or
- 2 Exudative ocular angiopathy.

BRENTUXIMAB VEDOTIN - PCT only - Special Authority see SA2289 below

⇒SA2289 Special Authority for Subsidy

Initial application — (relapsed/refractory Hodgkin lymphoma) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Both:

Sub: (Manufactu	,	,	
\$	Per	✓ Manufacturer	

continued...

- 1.1.1 Patient has relapsed/refractory CD30-positive Hodgkin lymphoma after two or more lines of chemotherapy; and
- 1.1.2 Patient is ineligible for autologous stem cell transplant; or
- 1.2 Both:
 - 1.2.1 Patient has relapsed/refractory CD30-positive Hodgkin lymphoma; and
 - 1.2.2 Patient has previously undergone autologous stem cell transplant; and
- 2 Patient has not previously received funded brentuximab vedotin; and
- 3 Response to brentuximab vedotin treatment is to be reviewed after a maximum of 6 treatment cycles; and
- 4 Brentuximab vedotin to be administered at doses no greater than 1.8 mg/kg every 3 weeks.

Renewal — (relapsed/refractory Hodgkin lymphoma) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has achieved a partial or complete response to brentuximab vedotin after 6 treatment cycles; and
- 2 Treatment remains clinically appropriate and the patient is benefitting from treatment and treatment is being tolerated; and
- 3 Patient is to receive a maximum of 16 total cycles of brentuximab vedotin treatment.

Initial application — (anaplastic large cell lymphoma) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has relapsed/refractory CD30-positive systemic anaplastic large cell lymphoma; and
- 2 Patient has an ECOG performance status of 0-1; and
- 3 Patient has not previously received brentuximab vedotin; and
- 4 Response to brentuximab vedotin treatment is to be reviewed after a maximum of 6 treatment cycles; and
- 5 Brentuximab vedotin to be administered at doses no greater than 1.8 mg/kg every 3 weeks.

Renewal — (anaplastic large cell lymphoma) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has achieved a partial or complete response to brentuximab vedotin after 6 treatment cycles; and
- 2 Treatment remains clinically appropriate and the patient is benefitting from treatment and treatment is being tolerated; and
- 3 Patient is to receive a maximum of 16 total cycles of brentuximab vedotin treatment.

⇒SA2401 Special Authority for Subsidy

Initial application — (head and neck cancer, locally advanced) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, non-metastatic, squamous cell cancer of the head and neck; and
- 2 Cisplatin is contraindicated or has resulted in intolerable side effects; and
- 3 Patient has an ECOG performance score of 0-2; and
- 4 To be administered in combination with radiation therapy.

Initial application — (colorectal cancer, metastatic) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has metastatic colorectal cancer located on the left side of the colon (see Note); and
- 2 There is documentation confirming disease is RAS and BRAF wild-type; and

Subsidy	Out	Fully	Brand or	
(Manufacturer's Price)	Sub	sidised	Generic	
\$	Per	•	Manufacturer	

continued...

- 3 Patient has an ECOG performance score of 0-2; and
- 4 Patient has not received prior funded treatment with cetuximab; and
- 5 Either:
 - 5.1 Cetuximab is to be used in combination with chemotherapy; or
 - 5.2 Chemotherapy is determined to not be in the best interest of the patient based on clinician assessment.

Renewal — **(colorectal cancer, metastatic)** only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where there is no evidence of disease progression.

Note: Left-sided colorectal cancer comprises of the distal one-third of the transverse colon, the splenic flexure, the descending colon, the sigmoid colon, or the rectum.

FARICIMAB - Special Authority see SA2533 below - Retail pharmacy

⇒SA2533 Special Authority for Subsidy

Initial application — (diabetic macular oedema) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has centre involving diabetic macular oedema (DMO); and
- 2 Patient's disease is nonresponsive to 4 doses of intravitreal bevacizumab when administered 4-6 weekly; and
- 3 Patient has reduced visual acuity between 6/9 6/36 with functional awareness of reduction in vision; and
- 4 Patient has DMO within central OCT (ocular coherence tomography) subfield > 350 micrometers; and
- 5 There is no centre-involving sub-retinal fibrosis or foveal atrophy; and
- 6 Patient has not previously been treated with aflibercept for longer than 3 months.

Renewal — (diabetic macular oedema) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 There is stability or two lines of Snellen visual acuity gain; and
- 2 There is structural improvement on OCT scan (with reduction in intra-retinal cysts, central retinal thickness, and sub-retinal fluid); and
- 3 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 4 There is no centre-involving sub-retinal fibrosis or foveal atrophy.

Initial application — (wet age related macular degeneration) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 Wet age-related macular degeneration (wet AMD); or
 - 1.2 Polypoidal choroidal vasculopathy; or
 - 1.3 Choroidal neovascular membrane from causes other than wet AMD; and
- 2 Either:
 - 2.1 The patient has developed severe endophthalmitis or severe posterior uveitis following treatment with bevacizumab; or
 - 2.2 There is worsening of vision or failure of retina to dry despite three intraocular injections of bevacizumab four weeks apart; and
- 3 There is no structural damage to the central fovea of the treated eve; and
- 4 Patient has not previously been treated with ranibizumab or aflibercept for longer than 3 months.

Renewal — (wet age related macular degeneration) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 2 There is no structural damage to the central fovea of the treated eye.

	Subsidy (Manufacturer's Price) Sub	Fully sidised	Brand or Generic	
	\$	Per	1	Manufacturer	
GEMTUZUMAB OZOGAMICIN - PCT only - Specialist - Special Authority see SA2269 below					
Inj 5 mg vial	12,973.00	1	✓ N	lylotarg	

⇒SA2269 Special Authority for Subsidy

Initial application only from a haematologist, paediatric haematologist or paediatric oncologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has not received prior chemotherapy for this condition; and
- 2 Patient has de novo CD33-positive acute myeloid leukaemia; and
- 3 Patient does not have acute promyelocytic leukaemia; and
- 4 Gemtuzumab ozogamicin will be used in combination with standard anthracycline and cytarabine (AraC); and
- 5 Patient is being treated with curative intent; and
- 6 Patient's disease risk has been assessed by cytogenetic testing to be good or intermediate; and
- 7 Patient must be considered eligible for standard intensive remission induction chemotherapy with standard anthracycline and cytarabine (AraC): and
- 8 Gemtuzumab ozogamicin to be funded for one course only (one dose at 3 mg per m² body surface area or up to 2 vials of 5 mg as separate doses).

Note: Acute myeloid leukaemia excludes acute promyelocytic leukaemia and acute myeloid leukaemia that is secondary to another haematological disorder (eg myelodysplasia or myeloproliferative disorder).

INFLIXIMAB - PCT only - Special Authority see SA2487 below

Inj 100 mg	428.00	1	Remicade
Inj 1 mg for ECP	4.40	1 mg	✓ Baxter

⇒SA2487 Special Authority for Subsidy

Initial application — (Crohn's disease (adults)) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has a CDAI score of greater than or equal to 300 or HBI score of greater than or equal to 10; or
 - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection;
 - 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but has experienced an inadequate response to, or intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Renewal — (Crohn's disease (adults)) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points from the CDAI score, or HBI score has reduced by 3 points, from when the patient was initiated on infliximab; or
 - 1.2 CDAI score is 150 or less, or HBI is 4 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment but CDAI score and/or HBI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (Crohn's disease (children)) from any relevant practitioner. Approvals valid for 6 months for applications

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

meeting the following criteria:

All of the following:

- 1 Paediatric patient has active Crohn's disease: and
- 2 Either:
 - 2.1 Patient has a PCDAI score of greater than or equal to 30; or
 - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but experienced an inadequate response to, or intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Renewal — (Crohn's disease (children)) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on infliximab; or
 - 1.2 PCDAI score is 15 or less: or
 - 1.3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — **(Graft vs host disease)** from any relevant practitioner. Approvals valid without further renewal unless notified where patient has steroid-refractory acute graft vs. host disease of the gut.

Initial application — **(Pulmonary sarcoidosis)** from any relevant practitioner. Approvals valid without further renewal unless notified where patient has life-threatening pulmonary sarcoidosis diagnosed by a multidisciplinary team that is refractory to other treatments.

Initial application — (acute fulminant ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 weeks for applications meeting the following criteria:
Both:

- 1 Patient has acute, fulminant ulcerative colitis; and
- 2 Treatment with intravenous or high dose oral corticosteroids has not been successful.

Initial application — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria:
Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Following 12 weeks of infliximab treatment, BASDAI has improved by 4 or more points from pre-infliximab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 3 Infliximab to be administered at doses no greater than 5 mg/kg every 6-8 weeks.

Initial application — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

Subsidy	rice)	Fully	Brand or
(Manufacturer's Pr		Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for chronic ocular inflammation; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for chronic ocular inflammation; or

2 Both:

- 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
- 2.2 Any of the following:
 - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective; or
 - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
 - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Renewal — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (fistulising Crohn's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has confirmed Crohn's disease: and
- 2 Any of the following:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
 - 2.2 Patient has one or more rectovaginal fistula(e); or
 - 2.3 Patent has complex peri-anal fistula.

Renewal — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
 - 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline (in the case of adult patients, as demonstrated by a reduction in the Fistula Assessment score), together with less induration and patient reported pain; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

be used for patients treated with this dose prior to 1 February 2019.

Initial application — (neurosarcoidosis) only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with neurosarcoiosis by a multidisciplinary team; and
- 2 Patient has CNS involvement; and
- 3 Patient has steroid-refractory disease; and
- 4 Either:
 - 4.1 IV cyclophosphamide has been tried; or
 - 4.2 Treatment with IV cyclophosphamide is clinically inappropriate.

Renewal — (neurosarcoidosis) only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria:

Either:

- 1 A withdrawal period has been tried and the patient has relapsed; or
- 2 All of the following:
 - 2.1 A withdrawal period has been considered but would not be clinically appropriate; and
 - 2.2 There has been a marked reduction in prednisone dose; and
 - 2.3 Fither:
 - 2.3.1 There has been an improvement in MRI appearances: or
 - 2.3.2 Marked improvement in other symptomology.

Initial application — (plaque psoriasis) only from a dermatologist or any relevant practitioner on the recommendation of a dermatologist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab, etanercept or secukinumab for severe chronic plaque psoriasis; and
 - 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects from adalimumab, etanercept or secukinumab; or
 - 1.2.2 Patient has received insufficient benefit from adalimumab, etanercept or secukinumab to meet the renewal criteria for adalimumab, etanercept or secukinumab for severe chronic plaque psoriasis: or
- 2 All of the following:
 - 2.1 Any of the following:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.3 Patient has severe chronic localised genital or flexural plaque psoriasis where the plaques or lesions have been present for at least 6 months from the time of initial diagnosis, and with a Dermatology Life Quality Index (DLQI) score greater than 10; and
 - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
 - 2.3 A PASI assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course: and
 - 2.4 The most recent PASI assessment is no more than 1 month old at the time of initiation.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

continued...

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand, foot, genital or flexural areas at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and for the face, palm of a hand or sole of a foot the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (plaque psoriasis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 Both:
 - 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 1.1.2 Following each prior infliximab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-infliximab treatment baseline value; or
 - 1.2 Both:
 - 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 1.2.2 Either:
 - 1.2.2.1 Following each prior infliximab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values: or
 - 1.2.2.2 Following each prior infliximab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-infliximab treatment baseline value: or
 - 1.3 Both:
 - 1.3.1 Patient had severe chronic localised genital or flexural plaque psoriasis at the start of treatment; and
 - 1.3.2 Either:
 - 1.3.2.1 The patient has experienced a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; or
 - 1.3.2.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing infliximab; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient was being treated with infliximab prior to 1 February 2019; and
- 2 Any of the following:
 - 2.1 Rheumatoid arthritis; or
 - 2.2 Ankylosing spondylitis; or
 - 2.3 Psoriatic arthritis: or
 - 2.4 Severe ocular inflammation: or
 - 2.5 Chronic ocular inflammation: or
 - 2.6 Crohn's disease (adults); or
 - 2.7 Crohn's disease (children); or
 - 2.8 Fistulising Crohn's disease; or
 - 2.9 Severe fulminant ulcerative colitis: or
 - 2.10 Severe ulcerative colitis; or

bsidy	Fully	Brand or
turer's Price) Subsi	dised	Generic
 \$ Per	✓	

continued...

- 2.11 Plague psoriasis; or
- 2.12 Neurosarcoidosis; or
- 2.13 Severe Behcet's disease.

Initial application — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis; and
- 2 Fither:
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept and/or secukinumab; or
 - 2.2 Following 3-4 months' initial treatment with adalimumab and/or etanercept and/or secukinumab, the patient did not meet the renewal criteria for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior infliximab treatment in the opinion of the treating physician; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Fither
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept; and
- 3 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance.

Renewal — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Infliximab to be administered at doses no greater than 3 mg/kg every 8 weeks.

Initial application — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe Behcet's disease which is significantly impacting the patient's quality of life (see Notes); and
- 2 Fither:

	Subsidy	Fully	Brand or
(Manu	facturer's Price)	Subsidised	Generic
	\$ Pe	er 🗸	Manufacturer

continued...

- 2.1 The patient has severe ocular, neurological and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s) (see Notes); or
- 2.2 The patient has severe gastrointestinal, rheumatologic and/or mucocutaneous symptoms and has not responded adequately to two or more treatment appropriate for the particular symptom(s) (see Notes); and
- 3 The patient is experiencing significant loss of quality of life.

Notes: Behcet's disease diagnosed according to the International Study Group for Behcet's Disease. Lancet 1990:335(8697):1078-80. Quality of life measured using an appropriate quality of life scale such as that published in Gilworth et al J Rheumatol. 2004:31:931-7.

Treatments appropriate for the particular symptoms are those that are considered standard conventional treatments for these symptoms, for example intravenous/oral steroids and other immunosuppressants for ocular symptoms; azathioprine, steroids, thalidomide, interferon alpha and ciclosporin for mucocutaneous symptoms; and colchicine, steroids and methotrexate for rheumatological symptoms.

Renewal — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has had a good clinical response to initial treatment with measurably improved quality of life; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Renewal — (fulminant ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Where maintenance treatment is considered appropriate, infliximab should be used in combination with immunomodulators and reassessed every 6 months; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe ocular inflammation; and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe ocular inflammation: or
- 2 Both:
 - 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
 - 2.2 Any of the following:
 - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
 - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
 - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Renewal — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

1 The patient has had a good clinical response following 3 initial doses: or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (ulcerative colitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active ulcerative colitis: and
- 2 Either:
 - 2.1 Patients SCCAI is greater than or equal to 4; or
 - 2.2 Patients PUCAI score is greater than or equal to 20; and
- 3 Patient has tried but has experienced an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and systemic corticosteroids.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 The SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on infliximab: or
 - 1.2 The PUCAI score has reduced by 10 points or more from the PUCAI score when the patient was initiated on infliximab; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Note: Indications marked with * are unapproved indications.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

Initial application — (inflammatory bowel arthritis – axial) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has had axial inflammatory pain for six months or more; and

 Subsidy Manufacturer's Price)	٤	Fully Subsidised	Brand or Generic
 \$	Per	1	Manufacturer

continued...

- 3 Patient is unable to take NSAIDs; and
- 4 Patient has unequivocal sacroiliitis demonstrated by radiological imaging or MRI; and
- 5 Patient's disease has not responded adequately to prior treatment consisting of at least 3 months of an exercise regime supervised by a physiotherapist; and
- 6 Patient has a BASDAI of at least 6 on a 0 10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment.

Renewal — (inflammatory bowel arthritis – axial) from any relevant practitioner. Approvals valid for 2 years where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10-point scale, or an improvement in BASDAI of 50%, whichever is less.

Initial application — (inflammatory bowel arthritis – peripheral) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has active arthritis in at least four joints from the following: hip, knee, ankle, subtalar, tarsus, forefoot, wrist, elbow, shoulder, sternoclavicular; and
- 3 Patient has tried and not experienced a response to at least three months of methotrexate or azathioprine at a maximum tolerated dose (unless contraindicated); and
- 4 Patient has tried and not experienced a response to at least three months of sulfasalazine at a maximum tolerated dose (unless contraindicated); and
- 5 Any of the following:
 - 5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application: or
 - 5.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application; or
 - 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (inflammatory bowel arthritis – peripheral) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, patient has experienced at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 Patient has experienced at least a continuing 30% improvement in active joint count from baseline in the opinion of the treating physician.

Initial application — (immune checkpoint inhibitor toxicity in malignancy*) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The individual requires treatment for moderate to severe autoimmune toxicity following immune checkpoint inhibitor treatment for malignancy; and
- 2 The individual has received insufficient benefit from use of corticosteroids; and
- 3 Infliximab is to be administered at up to 5mg/kg for up to four doses.

Renewal — (immune checkpoint inhibitor toxicity in malignancy*) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 The individual has shown clinical improvement and ongoing treatment is required; and
- 2 Infliximab is to be administered at up to 5mg/kg for up to a total of 8 doses.

Note: Indications marked with * are unapproved indications.

INOTUZUMAB OZOGAMICIN - PCT only - Specialist - Special Authority see SA2460 on the next page

Inj 1 mg vial14,4	457.00	1	Besponsa
Inj 1 mg for ECP14,4	457.00 1 i	mg 🗸	Baxter

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer

⇒SA2460 Special Authority for Subsidy

Initial application only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has relapsed or refractory CD22-positive B-cell acute lymphoblastic leukaemia/lymphoma, including minimal residual disease: and
- 2 Patient has ECOG performance status of 0-2; and
- 3 Either:
 - 3.1 Both:
 - 3.1.1 Patient has Philadelphia chromosome positive B-Cell ALL; and
 - 3.1.2 Patient has previously received a tyrosine kinase inhibitor; or
 - 3.2 Patient has received one prior line of treatment involving intensive chemotherapy; and
- 4 Treatment is to be administered for a maximum of 3 cycles.

Renewal only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient is not proceeding to a stem cell transplant; and
- 2 Either:
 - 2.1 Patient has experienced complete disease response; or
 - 2.2 Patient has experienced complete remission with incomplete haematological recovery; and
- 3 Treatment with inotuzumab ozogamicin is to cease after a total duration of 6 cycles.

MEPOLIZUMAB − Special Authority see SA2331 below − Retail pharmacy
Inj 100 mg prefilled pen1,638.00 1 ✓ Nucala

⇒SA2331 Special Authority for Subsidy

Initial application — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Patient must have a diagnosis of severe eosinophilic asthma documented by a respiratory physician or clinical immunologist; and
- 3 Conditions that mimic asthma eg. vocal cord dysfunction, central airway obstruction, bronchiolitis etc. have been excluded; and
- 4 Patient has a blood eosinophil count of greater than 0.5 x 10^9 cells/L in the last 12 months; and
- 5 Patient must be adherent to optimised asthma therapy including inhaled corticosteroids (equivalent to at least 1000 mcg per day of fluticasone propionate) plus long acting beta-2 agonist, or budesonide/formoterol as part of the single maintenance and reliever therapy regimen, unless contraindicated or not tolerated; and
- 6 Either:
 - 6.1 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral corticosteroids: or
 - 6.2 Patient has received continuous oral corticosteroids of at least the equivalent of 10 mg per day over the previous 3 months: and
- 7 Treatment is not to be used in combination with subsidised benralizumab; and
- 8 Patient has an Asthma Control Test (ACT) score of 10 or less. Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 52 weeks after the first dose to assess response to treatment; and

continued...

- 9 Either:
 - 9.1 Patient has not previously received an anti-IL5 biological therapy for their severe eosinophilic asthma; or
 - 9.2 Both:
 - 9.2.1 Patient was refractory or intolerant to previous anti-IL5 biological therapy; and
 - 9.2.2 Patient was not eligible to continue treatment with previous anti-IL5 biological therapy and discontinued within 12 months of commencing treatment.

Renewal — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 Either:
 - 2.1 Exacerbations have been reduced from baseline by 50% as a result of treatment with mepolizumab; or
 - 2.2 Reduction in continuous oral corticosteroid use by 50% or by 10 mg/day while maintaining or improving asthma control.

Initial application — (eosinophilic granulomatosis with polyangiitis) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has eosinophilic granulomatosis with polyangiitis; and
- 2 The patient has trialled and not received adequate benefit from at least one of the following for at least three months (unless contraindicated to all): azathioprine, cyclophosphamide, leflunomide, methotrexate, mycophenolate, or rituximab; and
- 3 Either:
 - 3.1 The patient has trialled prednisone for a minimum of three months and is unable to maintain disease control at doses below 7.5 mg per day; or
 - 3.2 Corticosteroids are contraindicated.

Renewal — (eosinophilic granulomatosis with polyangiitis) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months where patient has no evidence of clinical disease progression.

OBINUTUZUMAB - PCT only - Specialist - Special Aut	thority see SA2551 below		
Inj 25 mg per ml, 40 ml vial	5,910.00 1	✓ Gaz	yva
Inj 1 mg for ECP	6.21 1 m	ng 🗸 Baxi	ter

⇒SA2551 Special Authority for Subsidy

Initial application — (chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C CD20+ chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is obinutuzumab treatment naive; and
- 3 The patient is not eligible for full dose FCR due to comorbidities with a score > 6 on the Cumulative Illness Rating Scale (CIRS) or reduced renal function (creatinine clearance < 70mL/min); and</p>
- 4 Patient has adequate neutrophil and platelet counts* unless the cytopenias are a consequence of marrow infiltration by CLL; and
- 5 Patient has good performance status; and
- 6 Obinutuzumab to be administered at a maximum cumulative dose of 8,000 mg and in combination with chlorambucil for a maximum of 6 cycles.

Notes: Chronic lymphocytic leukaemia includes small lymphocytic lymphoma. Comorbidity refers only to illness/impairment other

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sı	ubsidised	Generic
\$	Per	✓	Manufacturer

continued...

than CLL induced illness/impairment in the patient. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with obinutuzumab is expected to improve symptoms and improve ECOG score to < 2.

* Neutrophil greater than or equal to 1.5×10^9 /L and platelets greater than or equal to 75×10^9 /L.

Initial application — (follicular / marginal zone lymphoma) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient has follicular lymphoma; or
 - 1.2 Patient has marginal zone lymphoma; and
- 2 Patient is refractory to or has relapsed within 12 months of a rituximab containing combined chemo-immunotherapy regimen*; and
- 3 Patient has an ECOG performance status of 0-2; and
- 4 Patient has been previously treated with no more than four chemotherapy regimens; and
- 5 Obinutuzumab to be administered at a maximum dose of 1000 mg for a maximum of 6 cycles in combination with chemotherapy*.

Note: * includes unapproved indications

Renewal — (follicular / marginal zone lymphoma) from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 Patient has no evidence of disease progression following obinutuzumab induction therapy; and
- 2 Obinutuzumab to be administered at a maximum of 1000 mg every 2 months for a maximum of 2 years; and
- 3 Obinutuzumab to be discontinued at disease progression.

OMALIZUMAB - Special Authority see SA1744 below -	Retail pharmacy		
Inj 150 mg prefilled syringe	450.00	1	✓ Xolair
Inj 150 mg vial	450.00	1	✓ Xolair AU ✓ Xolair

⇒SA1744 Special Authority for Subsidy

Initial application — (severe asthma) only from a respiratory specialist or clinical immunologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 6 years or older; and
- 2 Patient has a diagnosis of severe asthma; and
- 3 Past or current evidence of atopy, documented by skin prick testing or RAST; and
- 4 Total serum human immunoglobulin E (IgE) between 76 IU/mL and 1300 IU/ml at baseline; and
- 5 Proven adherence with optimal inhaled therapy including high dose inhaled corticosteroid (budesonide 1,600 mcg per day or fluticasone propionate 1,000 mcg per day or equivalent), plus long-acting beta-2 agonist therapy (at least salmeterol 50 mcg bd or eformoterol 12 mcg bd) for at least 12 months, unless contraindicated or not tolerated; and
- 6 Either:
 - 6.1 Patient has received courses of systemic corticosteroids equivalent to at least 28 days treatment in the past 12 months, unless contraindicated or not tolerated; or
 - 6.2 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral steroids; and
- 7 Patient has an Asthma Control Test (ACT) score of 10 or less; and
- 8 Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 26 weeks after the first dose to assess response to treatment.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

Initial application — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Fither:
 - 2.1 Both:
 - 2.1.1 Patient is symptomatic with Urticaria Activity Score 7 (UAS7) of 20 or above; and
 - 2.1.2 Patient has a Dermatology life quality index (DLQI) of 10 or greater; or
 - 2.2 Patient has a Urticaria Control Test (UCT) of 8 or less; and
- 3 Any of the following:
 - 3.1 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and ciclosporin (> 3 mg/kg day) for at least 6 weeks; or
 - 3.2 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and at least 3 courses of systemic corticosteroids (> 20 mg prednisone per day for at least 5 days) in the previous 6 months; or
 - 3.3 Patient has developed significant adverse effects whilst on corticosteroids or ciclosporin; and
- 4 Fither:
 - 4.1 Treatment to be stopped if inadequate response* following 4 doses: or
 - 4.2 Complete response* to 6 doses of omalizumab.

Renewal — (severe asthma) only from a clinical immunologist or respiratory specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 A reduction in the maintenance oral corticosteroid dose or number of exacerbations of at least 50% from baseline.

Renewal — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

- Fither:
 - 1 Patient has previously adequately responded* to 6 doses of omalizumab; or
 - 2 Both:
 - 2.1 Patient has previously had a complete response* to 6 doses of omalizumab; and
 - 2.2 Patient has relapsed after cessation of omalizumab therapy.

Note: *Inadequate response defined as less than 50% reduction in baseline UAS7 and DLQI score, or an increase in Urticaria Control Test (UCT) score of less than 4 from baseline. Patient is to be reassessed for response after 4 doses of omalizumab. Complete response is defined as UAS7 less than or equal to 6 and DLQI less than or equal to 5; or UCT of 16. Relapse of chronic urticaria on stopping prednisone/ciclosporin does not justify the funding of omalizumab.

PALIVIZUMAB - PCT only - Special Authority see SA2419 below

⇒SA2419 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Palivizumab to be administered during the annual RSV season; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Infant was born in the last 12 months; and
 - 2.1.2 Infant was born at less than 32 weeks zero days' gestation; or
 - 2.2 Both:
 - 2.2.1 Child was born in the last 24 months; and

|--|

continued...

- 2.2.2 Any of the following:
 - 2.2.2.1 Child has severe lung, airway, neurological or neuromuscular disease that requires ongoing ventilatory/respiratory support (see Note A) in the community: or
 - 2.2.2.2 Both:
 - 2.2.2.2.1 Child has haemodynamically significant heart disease; and
 - 2.2.2.2.2 Any of the following:
 - 2.2.2.2.2.1 Child has unoperated simple congenital heart disease with significant left to right shunt (see Note B); or
 - 2.2.2.2.2. Child has unoperated or surgically palliated complex congenital heart disease; or
 - 2.2.2.2.3 Child has severe pulmonary hypertension (see Note C); or
 - 2.2.2.2.2.4 Child has moderate or severe left ventricular (LV) failure (see Note D); or
 - 2.2.2.3 Child has severe combined immune deficiency, confirmed by an immunologist, but has not received a stem cell transplant; or
 - 2.2.2.4 Child has inborn errors of immunity (see Note E) that increase susceptibility to life-threatening viral respiratory infections, confirmed by an immunologist.

Renewal from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Palivizumab to be administered during the annual RSV season; and
- 2 Child was born in the last 24 months; and
- 3 Any of the following:
 - 3.1 Child has severe lung, airway, neurological or neuromuscular disease that requires ongoing ventilatory/respiratory support (see Note A) in the community; or
 - 3.2 Both:
 - 3.2.1 Child has haemodynamically significant heart disease; and
 - 3.2.2 Any of the following:
 - 3.2.2.1 Child has unoperated simple congenital heart disease with significant left to right shunt (see Note B);
 - 3.2.2.2 Child has unoperated or surgically palliated complex congenital heart disease; or
 - 3.2.2.3 Child has severe pulmonary hypertension (see Note C); or
 - 3.2.2.4 Child has moderate or severe left ventricular (LV) failure (see Note D); or
 - 3.3 Child has severe combined immune deficiency, confirmed by an immunologist, but has not received a stem cell transplant; or
 - 3.4 Child has inborn errors of immunity (see Note E) that increase susceptibility to life-threatening viral respiratory infections, confirmed by an immunologist.

Notes:

- a) Ventilatory/respiratory support includes those on home oxygen, CPAP/VPAP and those with tracheostomies in situ managed at home
- b) Child requires/will require heart failure medication, and/or child has significant pulmonary hypertension, and/or infant will require surgical palliation/definitive repair within the next 3 months
- c) Mean pulmonary artery pressure more than 25 mmHg
- d) LV Ejection Fraction less than 40%
- e) Inborn errors of immunity include, but are not limited to, IFNAR deficiencies

PERTUZUMAB - PCT only - Specialist - Special Authori	ity see SA2276 on the r	next page	
Inj 30 mg per ml, 14 ml vial	3,927.00	1	Perjeta
Inj 420 mg for ECP	3,927.00	420 mg OP	✓ Baxter

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA2276 Special Authority for Subsidy

Initial application — (metastatic breast cancer) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Either:
 - 2.1 Patient is chemotherapy treatment naïve; or
 - 2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
- 3 The patient has good performance status (ECOG grade 0-1); and
- 4 Pertuzumab to be administered in combination with trastuzumab; and
- 5 Pertuzumab maximum first dose of 840 mg, followed by maximum of 420 mg every 3 weeks; and
- 6 Pertuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
 - 1.2 The cancer has not progressed at any time point during the previous 12 months whilst on pertuzumab and trastuzumab; or
- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with pertuzumab and trastuzumab for reasons other than severe toxicity or disease progression; and
 - 2.2 Patient has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with pertuzumab and trastuzumab.

	2534 below	ly – Special Authority see SA2	PERTUZUMAB WITH TRASTUZUMAB - PCT only - S
✓ Phesgo	1	al7,707.00	Inj 600 mg with trastuzumab 600 mg, 10 ml vial
✓ Phesgo	1	vial12,894.00	Inj 1,200 mg with trastuzumab 600 mg, 15 ml vial

⇒SA2534 Special Authority for Subsidy

Initial application — (metastatic breast cancer) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The individual has received an initial Special Authority approval for intravenous pertuzumab and trastuzumab for metastatic breast cancer; and
 - 1.2 Pertuzumab with trastuzumab to be administered subcutaneously at a maximum dose of 600 mg pertuzumab with 600 mg trastuzumab every three weeks (or equivalent); or
- 2 All of the following:
 - 2.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
 - 2.2 Either:
 - 2.2.1 Patient is chemotherapy treatment naïve; or
 - 2.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast

continued...

cancer: and

- 2.3 The patient has good performance status (ECOG grade 0-1); and
- 2.4 Loading dose of pertuzumab with trastuzumab to be administered subcutaneously at a maximum dose of 1200 mg pertuzumab with 600 mg trastuzumab, respectively; and
- 2.5 Maintenance doses of pertuzumab with trastuzumab to be administered subcutaneously at a maximum dose of 600 mg pertuzumab with 600 mg trastuzumab every three weeks (or equivalent); and
- 2.6 Pertuzumab with trastuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

1 Both:

- 1.1 The individual has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 1.2 The cancer has not progressed at any time point during the previous 12 months whilst on pertuzumab and trastuzumab; or
- 2 All of the following:
 - 2.1 Individual has previously discontinued treatment with pertuzumab with trastuzumab for reasons other than severe toxicity or disease progression; and
 - 2.2 Individual has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with pertuzumab with trastuzumab.

RITUXIMAB (MABTHERA) - PCT only - Specialist - 3	Special Authority see SA2552 be	elow	
Inj 100 mg per 10 ml vial	1,075.50	2	Mabthera

Inj 500 mg per 50 ml vial	2,688.30	1	✓ Mabthera
Inj 1 mg for ECP	5.64	1 mg	✓ Baxter (Mabthera)

⇒SA2552 Special Authority for Subsidy

Initial application — (rheumatoid arthritis - TNF inhibitors contraindicated) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Treatment with a Tumour Necrosis Factor alpha inhibitor is contraindicated; and
- 2 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroguine sulphate (at maximum tolerated doses); and
- 5 Any of the following:
 - 5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
 - 5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
- 6 Fither:
 - 6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
 - 6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and

Subsidy	Fı	ılly	Brand or
(Manufacturer's Price)	Subsidis	ed	Generic
 \$	Per	✓	

continued...

- 7 Either:
 - 7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months; and
- 8 Either:
 - 8.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 8.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 9 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Initial application — (rheumatoid arthritis - prior TNF inhibitor use) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Both:
 - 1.1 The patient has had an initial community Special Authority approval for at least one of etanercept and/or adalimumab for rheumatoid arthritis: and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 1.2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept for rheumatoid arthritis: and
- 2 Either:
 - 2.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 2.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 3 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Renewal — (rheumatoid arthritis - re-treatment in 'partial responders' to rituximab) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 At 4 months following the initial course of rituximab infusions the patient had between a 30% and 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 At 4 months following the second course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.3 At 4 months following the third and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Renewal — (rheumatoid arthritis - re-treatment in 'responders' to rituximab) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 At 4 months following the initial course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or

Subsidy (Manufacturer's Pr	rice)	Fully Subsidised	Brand or Generic	
<u> </u>	Per	1	Manufacturer	

continued...

- 1.2 At 4 months following the second and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Fither:
 - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

RITUXIMAB (RIXIMYO) - PCT only - Specialist - Special Authority see SA2497 below

Inj 100 mg per 10 ml vial	275.33	2	✓ Riximyo
Inj 500 mg per 50 ml vial	688.20	1	✓ Riximyo
Inj 1 mg for ECP	1.38	1 mg	✓ Baxter (Riximyo)

⇒SA2497 Special Authority for Subsidy

Initial application — (ABO-incompatible organ transplant) from any relevant practitioner. Approvals valid without further renewal unless notified where patient is to undergo an ABO-incompatible solid organ transplant*.

Note: Indications marked with * are unapproved indications.

Initial application — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks; and
- 3 Any of the following:
 - 3.1 Induction therapy with daily oral or pulse intravenous cyclophosphamide has failed to achieve significant improvement of disease after at least 3 months; or
 - 3.2 Patient has previously had a cumulative dose of cyclophosphamide > 15 g or a further repeat 3 month induction course of cyclophosphamide would result in a cumulative dose > 15 g; or
 - 3.3 Cyclophosphamide and methotrexate are contraindicated; or
 - 3.4 Patient is a female of child-bearing potential; or
 - 3.5 Patient has a previous history of haemorrhagic cystitis, urological malignancy or haematological malignancy.

Note: Indications marked with * are unapproved indications.

Renewal — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 Patient has previously responded to treatment with rituximab but is now experiencing an acute flare of vasculitis; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks

Note: Indications marked with * are unapproved indications.

Initial application — **(Antibody-mediated organ transplant rejection)** from any relevant practitioner. Approvals valid without further renewal unless notified where patient has been diagnosed with antibody-mediated organ transplant rejection*.

Note: Indications marked with * are unapproved indications.

Initial application — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 The patient has progressive Binet stage A, B or C chronic lymphocytic leukaemia (CLL) requiring treatment; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 2 Any of the following:
 - 2.1 The patient is rituximab treatment naive; or
 - 2.2 Either:
 - 2.2.1 The patient is chemotherapy treatment naive: or
 - 2.2.2 Both:
 - 2.2.2.1 The patient's disease has relapsed following no more than three prior lines of chemotherapy treatment: and
 - 2.2.2.2 The patient has had a treatment-free interval of 12 months or more if previously treated with fludarabine and cyclophosphamide chemotherapy; or
 - 2.3 The patient's disease has relapsed and rituximab treatment is to be used in combination with funded venetoclax; and
 - 3 The patient has good performance status; and
 - 4 Either:
 - 4.1 The patient does not have chromosome 17p deletion CLL; or
 - 4.2 Rituximab treatment is to be used in combination with funded venetoclax for relapsed/refractory chronic lymphocytic leukaemia; and
 - 5 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles; and
 - 6 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration), bendamustine or venetoclax.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with rituximab is expected to improve symptoms and improve ECOG score to < 2.

Renewal — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 The patient's disease has relapsed and rituximab treatment is to be used in combination with funded venetoclax; or
 - 1.2 All of the following:
 - 1.2.1 The patient's disease has relapsed following no more than one prior line of treatment with rituximab for CLL; and
 - 1.2.2 The patient has had an interval of 36 months or more since commencement of initial rituximab treatment;
 - 1.2.3 The patient does not have chromosome 17p deletion CLL; and
 - 1.2.4 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration) or bendamustine; and
 - 2 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initial application — (Neuromyelitis Optica Spectrum Disorder(NMOSD)) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks; and

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidis	sed	Generic
\$	Per	✓	Manufacturer

continued...

- 2 Either:
 - 2.1 The patient has experienced a severe episode or attack of NMOSD (rapidly progressing symptoms and clinical investigations supportive of a severe attack of NMOSD); or
 - 2.2 All of the following:
 - 2.2.1 The patient has experienced a breakthrough attack of NMOSD; and
 - 2.2.2 The patient is receiving treatment with mycophenolate; and
 - 2.2.3 The patients is receiving treatment with corticosteroids.

Renewal — (Neuromyelitis Optica Spectrum Disorder) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks; and
- 2 The patients has responded to the most recent course of rituximab; and
- 3 The patient has not received rituximab in the previous 6 months.

Initial application — (Post-transplant) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 2 To be used for a maximum of 8 treatment cycles.

Note: Indications marked with * are unapproved indications.

Renewal — (Post-transplant) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 3 To be used for no more than 6 treatment cycles.

Note: Indications marked with * are unapproved indications.

Initial application — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1.000 mg doses given two weeks apart; and
- 2 Either:
 - 2.1 Treatment with corticosteroids and at least one other immunosuppressant for at least a period of 12 months has been ineffective; or
 - 2.2 Both:
 - 2.2.1 Treatment with at least one other immunosuppressant for a period of at least 12 months; and
 - 2.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

Renewal — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Fither:

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 3.1 The patient has relapsed despite treatment with corticosteroids and at least one other immunosuppressant for a period of at least 12 months; or
- 3.2 Both:
 - 3.2.1 The patient's myasthenia gravis has relapsed despite treatment with at least one immunosuppressant for a period of at least 12 months; and
 - 3.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

Initial application — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is a child with SDNS* or FRNS*: and
- 2 Treatment with steroids for at least a period of 3 months has been ineffective or associated with evidence of steroid toxicity; and
- 3 Treatment with ciclosporin for at least a period of 3 months has been ineffective and/or discontinued due to unacceptable side effects: and
- 4 Treatment with mycophenolate for at least a period of 3 months with no reduction in disease relapses; and
- 5 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Patient is a child with SRNS* where treatment with steroids and ciclosporin for at least 3 months have been ineffective;
- 2 Treatment with tacrolimus for at least 3 months has been ineffective; and
- 3 Genetic causes of nephrotic syndrome have been excluded; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Note: Indications marked with * are unapproved indications.

Initial application — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Fither:

- 1 All of the following:
 - 1.1 The patient has treatment naive aggressive CD20 positive NHL; and
 - 1.2 To be used with a multi-agent chemotherapy regimen given with curative intent; and
 - 1.3 To be used for a maximum of 8 treatment cycles; or
- 2 Both:
 - 2.1 The patient has aggressive CD20 positive NHL with relapsed disease following prior chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Renewal — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has relapsed refractory/aggressive CD20 positive NHL; and
- 3 To be used with a multi-agent chemotherapy regimen given with curative intent; and
- 4 To be used for a maximum of 4 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Initial application — (haemophilia with inhibitors) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria: Any of the following:

- 1 Patient has mild congenital haemophilia complicated by inhibitors; or
- 2 Patient has severe congenital haemophilia complicated by inhibitors and has failed immune tolerance therapy; or
- 3 Patient has acquired haemophilia.

Renewal — (haemophilia with inhibitors) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for haemophilia with inhibitors; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment.

Initial application — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Fither:
 - 1.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per microlitre: or
 - 1.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding; and
- 2 Any of the following:
 - 2.1 Treatment with steroids and splenectomy have been ineffective; or
 - 2.2 Treatment with steroids has been ineffective and splenectomy is an absolute contraindication; or
 - 2.3 Other treatments including steroids have been ineffective and patient is being prepared for elective surgery (e.g. splenectomy); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sı	ubsidised	Generic
\$	Per	1	Manufacturer

continued...

Note: Indications marked with * are unapproved indications.

Renewal — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for immune thrombocytopenic purpura*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (indolent, low-grade lymphomas or hairy cell leukaemia*) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has indolent low grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy: and
 - 1.2 To be used for a maximum of 6 treatment cycles: or
- 2 Both:
 - 2.1 The patient has indolent, low grade lymphoma or hairy cell leukaemia* requiring first-line systemic chemotherapy:
 - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Renewal — (indolent, low-grade lymphomas or hairy cell leukaemia*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has indolent, low-grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
- 3 To be used for no more than 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairv cell leukaemia' also includes hairv cell leukaemia variant.

Initial application — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient has autoimmune pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder.

Note: Indications marked with * are unapproved indications.

Renewal — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient was previously treated with rituximab for pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder and demonstrated an initial response lasting at least 12 months.

Note: Indications marked with * are unapproved indications.

Initial application — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Patient has cold haemagglutinin disease*; and
- 2 Patient has severe disease which is characterized by symptomatic anaemia, transfusion dependence or disabling circulatory symptoms: and

Subsidy (Manufacturer's Price)	Fully Subsidised		
(ivialitulacture) 5 Filoe)	Per 🗸	Manufacturer	
Φ	rei •	Manuacturei	

continued...

3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks

Note: Indications marked with * are unapproved indications.

Renewal — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Fither:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for severe cold haemagglutinin disease*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Both:

- 1 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks; and
- 2 Either:
 - 2.1 Patient has thrombotic thrombocytopenic purpura* and has experienced progression of clinical symptoms or persistent thrombocytopenia despite plasma exchange; or
 - 2.2 Patient has acute idiopathic thrombotic thrombocytopenic purpura* with neurological or cardiovascular pathology.

Note: Indications marked with * are unapproved indications.

Renewal — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for thrombotic thrombocytopenic purpura*: and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe, immediately life- or organ-threatening SLE*; and
- 2 The disease has proved refractory to treatment with steroids at a dose of at least 1 mg/kg; and
- 3 The disease has relapsed following prior treatment for at least 6 months with maximal tolerated doses of azathioprine, mycophenolate mofetil and high dose cyclophosphamide, or cyclophosphamide is contraindicated; and
- 4 Maximum of four 1000 mg infusions of rituximab.

Note: Indications marked with * are unapproved indications.

Renewal — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

1 Patient's SLE* achieved at least a partial response to the previous round of prior rituximab treatment; and

(Subsidy	F	ully	Brand or
	(Manufacturer's Price)	Subsidi	sed	Generic
·	\$	Per	•	Manufacturer

continued...

- 2 The disease has subsequently relapsed; and
- 3 Maximum of two 1000 mg infusions of rituximab.

Note: Indications marked with * are unapproved indications.

Initial application — (warm autoimmune haemolytic anaemia (warm AIHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has warm autoimmune haemolytic anaemia*; and
- 2 One of the following treatments has been ineffective: steroids (including if patient requires ongoing steroids at doses equivalent to > 5 mg prednisone daily), cytotoxic agents (e.g. cyclophosphamide monotherapy or in combination), intravenous immunoglobulin; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (warm autoimmune haemolytic anaemia (warm AIHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for warm autoimmune haemolytic anaemia*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (severe antisynthetase syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed antisynthetase syndrome; and
- 2 Patient has severe, immediately life or organ threatening disease, including interstitial lung disease; and
- 3 Either:
 - 3.1 Treatment with at least 3 immunosuppressants (oral steroids, cyclophosphamide, methotrexate, mycophenolate, ciclosporin, azathioprine) has not be effective at controlling active disease; or
 - 3.2 Rapid treatment is required due to life threatening complications; and
- 4 Maximum of four 1,000mg infusions of rituximab.

Renewal — (severe antisynthetase syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in inflammatory markers, muscle strength and pulmonary function; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 Maximum of two cycles of $2 \times 1,000$ mg infusions of rituximab given two weeks apart.

Initial application — (graft versus host disease) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has refractory graft versus host disease following transplant; and
- 2 Treatment with at least 3 immunosuppressants (oral steroids, ciclosporin, tacrolimus, mycophenolate, sirolimus) has not be effective at controlling active disease; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	sidised	Generic	
\$	Per	1	Manufacturer	

continued...

4 weeks.

Initial application — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has severe chronic inflammatory demyelinating polyneuropathy (CIPD); and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
 - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
 - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function compared to baseline; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Initial application — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has severe anti-NMDA receptor autoimmune encephalitis; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
 - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
 - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 The patient has experienced a relapse and now requires further treatment; and
- 4 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Initial application — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

Fither:

ubsidy	Fully	Brand or
turer's Price) Sub	sidised	Generic
 \$ Per	1	

continued...

- 1 Both:
 - 1.1 The patient has CD20+ low grade or follicular B-cell NHL with relapsed disease following prior chemotherapy; and
 - 1.2 To be used for a maximum of 6 treatment cycles; or
 - 2 Both:
 - 2.1 The patient has CD20+ low grade or follicular B-cell NHL requiring first-line systemic chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Renewal — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 Rituximab is to be used for maintenance in CD20+ low grade or follicular B-cell NHL following induction with first-line systemic chemotherapy; and
- 2 Patient is intended to receive rituximab maintenance therapy for 2 years at a dose of 375 mg/m2 every 8 weeks (maximum of 12 cycles).

Initial application — (Membranous nephropathy) only from a nephrologist or any relevant practitioner on the recommendation of a nephrologist. Approvals valid for 6 weeks for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 Patient has biopsy-proven primary/idiopathic membranous nephropathy*; or
 - 1.2 Patient has PLA2 antibodies with no evidence of secondary cause, and an eGFR of > 60ml/min/1.73m2; and
- 2 Patient remains at high risk of progression to end-stage kidney disease despite more than 3 months of treatment with conservative measures (see Note): and
- 3 The total rituximab dose would not exceed the equivalent of 375mg/m2 of body surface area per week for a total of

Renewal — (Membranous nephropathy) only from a nephrologist or any relevant practitioner on the recommendation of a nephrologist. Approvals valid for 6 weeks for applications meeting the following criteria: All of the following:

- 1 Patient was previously treated with rituximab for membranous nephropathy*: and
- 2 Either:
 - 2.1 Treatment with rituximab was previously successful, but the condition has relapsed, and the patient now requires repeat treatment; or
 - 2.2 Patient achieved partial response to treatment and requires repeat treatment (see Note); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Notes:

- a) Indications marked with * are unapproved indications.
- b) High risk of progression to end-stage kidney disease defined as > 5g/day proteinuria.
- c) Conservative measures include renin-angiotensin system blockade, blood-pressure management, dietary sodium and protein restriction, treatment of dyslipidaemia, and anticoagulation agents unless contraindicated or the patient has experienced intolerable side effects.
- d) Partial response defined as a reduction of proteinuria of at least 50% from baseline, and between 0.3 grams and 3.5 grams per 24 hours.

Initial application — (B-cell acute lymphoblastic leukaemia/lymphoma*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient has newly diagnosed B-cell acute lymphoblastic leukaemia/lymphoma*; and
- 2 Treatment must be in combination with an intensive chemotherapy protocol with curative intent; and

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

3 The total rituximab dose would not exceed the equivalent of 375 mg/m² per dose for a maximum of 18 doses.

Note: Indications marked with * are unapproved indications.

Initial application — (desensisation prior to transplant) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires desensitisation prior to mismatched allogenic stem cell transplant*; and
- 2 Patient would receive no more than two doses at 375 mg/m2 of body-surface area.

Note: Indications marked with * are unapproved indications.

Initial application — (pemiphigus*) only from a dermatologist or relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 Patient has severe rapidly progressive pemphigus; and
 - 1.2 Is used in combination with systemic corticosteroids (20 mg/day); and
 - 1.3 Any of the following:
 - 1.3.1 Skin involvement is at least 5% body surface area; or
 - 1.3.2 Significant mucosal involvement (10 or more mucosal erosions) or diffuse gingivitis or confluent large erosions; or
 - 1.3.3 Involvement of two or more mucosal sites; or
- 2 Both:
 - 2.1 Patient has pemphigus; and
 - 2.2 Patient has not experienced adequate clinical benefit from systemic corticosteroids (20 mg/day) in combination with a steroid sparing agent, unless contraindicated.

Note: Indications marked with * are unapproved indications.

Renewal — (pemiphigus*) only from a dermatologist or relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has experienced adequate clinical benefit from rituximab treatment, with improvement in symptoms and healing of skin ulceration and reduction in corticosteroid requirement; and
- 2 Patient has not received rituximab in the previous 6 months.

Note: Indications marked with * are unapproved indications.

Initial application — (immunoglobulin G4-related disease (IgG4-RD*)) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed diagnosis of IgG4-RD*; and
- 2 Either:
 - 2.1 Treatment with corticosteroids and/or disease modifying anti-rheumatic drugs for at least 3 months has been ineffective in lowering corticosteroid dose below 5 mg per day (prednisone equivalent) without relapse; or
 - 2.2 Treatment with corticosteroids and/or disease modifying anti-rheumatic drugs is contraindicated or associated with evidence of toxicity or intolerance; and
- 3 Total rituximab dose used should not exceed a maximum of two 1000 mg infusions of rituximab given two weeks apart.

Note: Indications marked with * are unapproved indications.

Renewal — (immunoglobulin G4-related disease (IgG4-RD*)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 Fither:

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sub	sidised	Generic
\$	Per	•	Manufacturer

continued...

- 1.1 Treatment with rituximab for IgG4-RD* was previously successful and patient's disease has demonstrated sustained response, but the condition has relapsed; or
- 1.2 Patient is receiving maintenance treatment for IgG4-RD*; and
- 2 Rituximab re-treatment not to be given within 6 months of previous course of treatment; and
- 3 Maximum of two 1000 mg infusions of rituximab given two weeks apart.

Note: Indications marked with * are unapproved indications.

SECUKINUMAB - Special Authority see SA2488 below - Retail pharmacy

Inj 150 mg per ml, 1 ml prefilled syringe.......799.50 ✓ Cosentyx 1.599.00 ✓ Cosentvx

⇒SA2488 Special Authority for Subsidy

Initial application — (severe chronic plaque psoriasis – second-line biologic) only from a dermatologist or any relevant practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab or etanercept, or has trialled infliximab in a Health NZ Hospital, for severe chronic plague psoriasis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
 - 2.2 The patient has received insufficient benefit from adalimumab, etanercept or infliximab; and
- 3 A Psoriasis Area and Severity Index (PASI) assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Initial application — (severe chronic plaque psoriasis – first-line biologic) only from a dermatologist or any relevant practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plagues have been present for at least 6 months from the time of initial diagnosis; or
 - 1.3 Patient has severe chronic localised genital or flexural plague psoriasis where the plagues or lesions have been present for at least 6 months from the time of initial diagnosis, and with a Dermatology Life Quality Index (DLQI) score greater than 10; and
- 2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin: and
- 3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course: and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Note: A treatment course is defined as a minimum of 12 weeks of treatment. "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plague psoriasis of the face, hand. foot, genital or flexural areas, at least 2 of the 3 PASI symptom sub scores for erythema, thickness and scaling are rated as severe or very severe, and for the face, palm of a hand or sole of a foot the skin area affected is 30% or more of the face, palm of

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis – first and second-line biologic) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 Fither:
 - 1.1.1 Patient's PASI score has reduced by 75% or more (PASI 75) as compared to baseline PASI prior to commencing secukinumab; or
 - 1.1.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing secukinumab; or
 - 1.2 Both:
 - 1.2.1 Patient had severe chronic localised genital or flexural plaque psoriasis at the start of treatment; and
 - 1.2.2 Either:
 - 1.2.2.1 The patient has experienced a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; or
 - 1.2.2.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing secukinumab; and
- 2 Secukinumab to be administered at a maximum dose of 300 mg monthly.

Initial application — **(ankylosing spondylitis – second-line biologic)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

Renewal — (ankylosing spondylitis – second-line biologic) only from a rheumatologist or medical practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Following 12 weeks initial treatment of secukinumab treatment, BASDAI has improved by 4 or more points from pre-secukinumab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefitted from treatment and that continued treatment is appropriate; and
- 3 Secukinumab to be administered at doses no greater than 300 mg monthly.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has had an initial Special Authority approval for adalimumab, etanercept or infliximab for psoriatic arthritis; and
 - 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
 - 1.2.2 Patient has received insufficient benefit from adalimumab, etanercept or infliximab to meet the renewal criteria for adalimumab, etanercept or infliximab for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	•	Manufacturer

continued...

- 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
- 2.4 Either:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior secukinumab treatment in the opinion of the treating physician; and
- 2 Secukinumab to be administered at doses no greater than 300 mg monthly.

SILTUXIMAB - Special Authority see SA1596 below - Retail pharmacy

⇒SA1596 Special Authority for Subsidy

Initial application only from a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's Disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Siltuximab is to be administered at doses no greater than 11 mg/kg every 3 weeks.

Renewal only from a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has sustained improvement in inflammatory markers and functional status.

TOCILIZUMAB - PCT only - Special Authority see SA2489 below

Inj 20 mg per ml, 4 ml vial	220.00	1	✓ Actemra
Inj 20 mg per ml, 10 ml vial		1	✓ Actemra
Inj 20 mg per ml, 20 ml vial		1	✓ Actemra
Inj 1 mg for ECP	2.85	1 mg	Baxter

⇒SA2489 Special Authority for Subsidy

Initial application — (cytokine release syndrome) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

Subsidy	Fu	lly Brand or	
(Manufacturer's Price)	Subsidis	ed Generic	
\$	Per	 Manufacturer 	

continued...

- 1 Both:
 - 1.1 The patient has developed grade 3 or 4 cytokine release syndrome associated with the administration of blinatumomab for the treatment of acute lymphoblastic leukaemia; and
 - 1.2 Tocilizumab is to be administered at doses no greater than 8 mg/kg IV for a maximum of 3 doses (if less than 30kg, maximum of 12 mg/kg); or
- 2 All of the following:
 - 2.1 The patient is enrolled in the Malaghan Institute of Medical Research ENABLE trial programme; and
 - 2.2 The patient has developed CRS or Immune Effector Cell-Associated Neurotoxicity Syndrome (ICANS) following CAR T-cell therapy for the treatment of relapsed or refractory B-cell non-Hodgkin lymphoma; and
 - 2.3 Tocilizumab is to be administered according to the consensus guidelines for CRS or ICANS for CAR T-cell therapy at doses no greater than 8 mg/kg IV for a maximum of 3 doses.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient was being treated with tocilizumab prior to 1 February 2019; and
- 2 Any of the following:
 - 2.1 rheumatoid arthritis; or
 - 2.2 systemic juvenile idiopathic arthritis; or
 - 2.3 adult-onset Still's disease: or
 - 2.4 polyarticular juvenile idiopathic arthritis; or
 - 2.5 idiopathic multicentric Castleman's disease.

Initial application — (Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
- 3 Either:
 - 3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or
 - 3.2 Both:
 - 3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a Health NZ Hospital; and
 - 3.2.2 Either:
 - 3.2.2.1 The patient has experienced intolerable side effects from rituximab; or
 - 3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Initial application — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 2 Tocilizumab is to be used as monotherapy; and
- 3 Either:
 - 3.1 Treatment with methotrexate is contraindicated; or
 - 3.2 Patient has tried and did not tolerate oral and/or parenteral methotrexate; and

Subsidy		ully	Brand or	
(Manufacturer's Price)	Subsid	ised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 4 Either:
 - 4.1 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of ciclosporin alone or in combination with another agent; or
 - 4.2 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of leflunomide alone or in combination with another agent; and
- 5 Either:
 - 5.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 active, swollen, tender joints;
 - 5.2 Patient has persistent symptoms of poorly controlled and active disease in at least four active joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 6 Fither:
 - 6.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 6.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Initial application — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient diagnosed with systemic juvenile idiopathic arthritis; and
- 2 Patient has tried and not responded to a reasonable trial of all of the following, either alone or in combination: oral or parenteral methotrexate; non-steroidal anti-inflammatory drugs (NSAIDs); and systemic corticosteroids.

Initial application — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Either:

- 1 Both:
 - 1.1 Either:
 - 1.1.1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a Health NZ Hospital; and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal antiinflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Initial application — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for both etanercept and adalimumab for polyarticular course iuvenile idiopathic arthritis (JIA); and
 - 1.2 The patient has experienced intolerable side effects, or has received insufficient benefit from, both etanercept and adalimumab: or
- 2 All of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	1	Manufacturer	

continued...

- 2.1 Treatment with a tumour necrosis factor alpha inhibitor is contraindicated; and
- 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
- 2.3 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2.4 Any of the following:
 - 2.4.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.4.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.4.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Initial application — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Tocilizumab to be administered at doses no greater than 8 mg/kg IV every 3-4 weeks.

Initial application — (moderate to severe COVID-19) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed (or probable) COVID-19; and
- 2 Oxygen saturation of < 92% on room air, or requiring supplemental oxygen; and
- 3 Patient is receiving adjunct systemic corticosteroids, or systemic corticosteroids are contraindicated; and
- 4 Tocilizumab is to be administered at doses no greater than 8mg/kg IV for a maximum of one dose; and
- 5 Tocilizumab is not to be administered in combination with barcitinib.

Renewal — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Either:

- 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Renewal — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Following up to 6 months' initial treatment, the patient has achieved at least an American College of Rheumatology paediatric 30% improvement criteria (ACR Pedi 30) response from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing ACR Pedi 30 response from baseline.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months where the patient has a sustained improvement in inflammatory markers and functional status. Renewal — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
 \$	Per	1	

continued...

improvement in physician's global assessment from baseline; or

2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Renewal — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has a sustained improvement in inflammatory markers and functional status.

Initial application — (immune checkpoint inhibitor toxicity in malignancy*) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The individual requires treatment for moderate to severe autoimmune toxicity following immune checkpoint inhibitor treatment for malignancy; and
- 2 The individual has received insufficient benefit from use of corticosteroids; and
- 3 Tocilizumab is to be administered at a maximum dose of 8 mg/kg fortnightly.

Renewal — (immune checkpoint inhibitor toxicity in malignancy*) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 The individual has shown clinical improvement and ongoing treatment is required; and
- 2 Tocilizumab is to be administered at a maximum dose of 8 mg/kg fortnightly.

Note: Indications marked with * are unapproved indications.

TRASTUZUMAB (HERZUMA) - PCT only - Special Authority see SA2293 below

Inj 150 mg vial	100.00	1	✓ <u>Herzuma</u>
Inj 440 mg vial	293.35	1	✓ <u>Herzuma</u>
Inj 1 mg for ECP	0.70	1 mg	Baxter

⇒SA2293 Special Authority for Subsidy

Initial application — (early breast cancer) from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria:

Both:

- 1 The patient has early breast cancer expressing HER-2 IHC 3+ or ISH + (including FISH or other current technology); and
- 2 Maximum cumulative dose of 106 mg/kg (12 months' treatment).

Renewal — (early breast cancer*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
 - 1.2 The patient received prior adjuvant trastuzumab treatment for early breast cancer; and
 - 1.3 Any of the following:
 - 1.3.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
 - 1.3.2 The patient discontinued lapatinib within 3 months due to intolerable side effects and the cancer did not progress whilst on lapatinib; or
 - 1.3.3 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
 - 1.4 Either:
 - 1.4.1 Trastuzumab will not be given in combination with pertuzumab; or
 - 1.4.2 All of the following:
 - 1.4.2.1 Trastuzumab to be administered in combination with pertuzumab; and
 - 1.4.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free

Subsidy	Full	y Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per 🗸	 Manufacturer 	

continued...

interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and

- 1.4.2.3 The patient has good performance status (ECOG grade 0-1); and
- 1.5 Trastuzumab to be discontinued at disease progression; or
- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with trastuzumab in the metastatic setting for reasons other than severe toxicity or disease progression; and
 - 2.2 Patient has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with trastuzumab.

Note: * For patients with relapsed HER-2 positive disease who have previously received adjuvant trastuzumab for early breast cancer

Initial application — (metastatic breast cancer) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Either:
 - 2.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
 - 2.2 The patient discontinued lapatinib within 3 months due to intolerable side effects and the cancer did not progress whilst on lapatinib; and
- 3 Either:
 - 3.1 Trastuzumab will not be given in combination with pertuzumab; or
 - 3.2 All of the following:
 - 3.2.1 Trastuzumab to be administered in combination with pertuzumab; and
 - 3.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer: and
 - 3.2.3 The patient has good performance status (ECOG grade 0-1); and
- 4 Trastuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Fither:

- 1 All of the following:
 - 1.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
 - 1.2 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
 - 1.3 Trastuzumab to be discontinued at disease progression; or
- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with trastuzumab for reasons other than severe toxicity or disease progression; and
 - 2.2 Patient has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with trastuzumab.

Initial application — (gastric, gastro-oesophageal junction and oesophageal cancer) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

1 The patient has locally advanced or metastatic gastric, gastro-oesophageal junction or oesophageal cancer expressing HER-2 IHC 2+ FISH+ or IHC3+ (or other current technology); and

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

2 Patient has an ECOG score of 0-2.

Renewal — (gastric, gastro-oesophageal junction and oesophageal cancer) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 2 Trastuzumab to be discontinued at disease progression.

TRASTUZUMAB DERUXTECAN - PCT only - Special Authority see SA2420 below

Inj 100 mg per ml, 1 ml vial	 	2,550	.00	1	✓ Enher	tu
Inj 1 mg for ECP	 	27	.05	1 mg	✓ Baxter	r

⇒SA2420 Special Authority for Subsidy

Initial application only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Patient is currently on treatment with trastuzumab deruxtecan and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 2.1 Patient has metastatic breast cancer expressing HER-2 IHC3+ or ISH+ (including FISH or other current
 - technology); and
 2.2 Patient has previously received trastuzumab and chemotherapy, separately or in combination; and
 - 2.3 Fither:
 - 2.3.1 The patient has received prior therapy for metastatic disease; or
 - 2.3.2 The patient developed disease recurrence during, or within six months of completing adjuvant therapy; and
 - 2.4 Patient has a good performance status (ECOG 0-1); and
 - 2.5 Patient has not received prior funded trastuzumab deruxtecan treatment; and
 - 2.6 Treatment to be discontinued at disease progression.

Renewal only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The cancer has not progressed at any time point during the previous approval period whilst on trastuzumab deruxtecan; and
- 2 Treatment to be discontinued at disease progression.

Note: Prior or adjuvant therapy includes anthracycline, other chemotherapy, biological drugs, or endocrine therapy.

TRASTUZUMAB EMTANSINE - PCT only - Specialist - Special Authority see SA2424 below

Inj 100 mg vial	2,320.00	1	Kadcyla
Inj 160 mg vial	3,712.00	1	✓ Kadcyla
Inj 1 mg for ECP	24.52	1 mg	✓ Baxter

⇒SA2424 Special Authority for Subsidy

Initial application — (early breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has early breast cancer expressing HER2 IHC3+ or ISH+; and
- 2 Documentation of pathological invasive residual disease in the breast and/or axiliary lymph nodes following completion of surgery; and
- 3 Patient has completed systemic neoadjuvant therapy with trastuzumab and chemotherapy prior to surgery; and
- 4 Disease has not progressed during neoadjuvant therapy; and

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

- 5 Patient has left ventricular ejection fraction of 45% or greater; and
- 6 Adjuvant treatment with trastuzumab emtansine to be commenced within 12 weeks of surgery; and
- 7 Trastuzumab emtansine to be discontinued at disease progression; and
- 8 Total adjuvant treatment duration must not exceed 42 weeks (14 cycles).

Initial application — (metastatic breast cancer) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Patient has previously received trastuzumab and chemotherapy, separately or in combination; and
- 3 Fither:
 - 3.1 The patient has received prior therapy for metastatic disease*; or
 - 3.2 The patient developed disease recurrence during, or within six months of completing adjuvant therapy*; and
- 4 Patient has a good performance status (ECOG 0-1); and
- 5 Either:
 - 5.1 Patient does not have symptomatic brain metastases; or
 - 5.2 Patient has brain metastases and has received prior local CNS therapy; and
- 6 Either:
 - 6.1 Patient has not received prior funded trastuzumab emtansine or trastuzumab deruxtecan treatment; or
 - 6.2 Both:
 - 6.2.1 Patient has discontinued trastuzumab deruxtecan due to intolerance; and
 - 6.2.2 The cancer did not progress while on trastuzumab deruxtecan; and
- 7 Treatment to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The cancer has not progressed at any time point during the previous approval period whilst on trastuzumab emtansine; and
- 2 Treatment to be discontinued at disease progression.

Note: Prior or adjuvant therapy includes anthracycline, other chemotherapy, biological drugs, or endocrine therapy.

USTEKINUMAB - Special Authority see SA2182 below - Retail pharmacy

Inj 90 mg per ml, 1 ml pre-filled syringe.......4,162.00 1 ✓ Stelara

⇒SA2182 Special Authority for Subsidy

Initial application — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Patient is currently on treatment with ustekinumab commenced prior to 1 February 2023 and met all remaining criteria (criterion 2) below at the time of commencing treatment; or
- 2 Both:
 - 2.1 Patient has active Crohn's disease; and
 - 2.2 Either:
 - 2.2.1 Patient has had an initial approval for prior biologic therapy for Crohn's disease and has experienced intolerable side effects or insufficient benefit to meet renewal criteria: or
 - 2.2.2 Both:
 - 2.2.2.1 Patient meets the initiation criteria for prior biologic therapies for Crohn's disease; and
 - 2.2.2.2 Other biologics for Crohn's disease are contraindicated.

Renewal — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 12 months for applications meeting

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

the following criteria:

Both:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points, or HBI score has reduced by 3 points, from when the patient was initiated on biologic therapy; or
 - 1.2 CDAI score is 150 or less, or HBI is 4 or less; or
 - 1.3 The patient has experienced an adequate response to treatment, but CDAI score and/or HBI score cannot be assessed; and
- 2 Ustekinumab to be administered at a dose no greater than 90 mg every 8 weeks.

Initial application — (Crohn's disease - children*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Patient is currently on treatment with ustekinumab commenced prior to 1 February 2023 and met all remaining criteria (criterion 2) below at the time of commencing treatment; or
- 2 Both:
 - 2.1 Patient has active Crohn's disease: and
 - 2.2 Either:
 - 2.2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria: or
 - 2.2.2 Both:
 - 2.2.2.1 Patient meets the initiation criteria for prior biologic therapies for Crohn's disease; and
 - 2.2.2.2 Other biologics for Crohn's disease are contraindicated.

Note: Indication marked with * is an unapproved indication.

Renewal — (Crohn's disease - children*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from when the patient was initiated on biologic therapy; or
 - 1.2 PCDAI score is 15 or less; or
 - 1.3 The patient has experienced an adequate response to treatment, but CDAI score cannot be assessed; and
- 2 Ustekinumab to administered at a dose no greater than 90 mg every 8 weeks.

Note: Indication marked with * is an unapproved indication.

Initial application — (ulcerative colitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Patient is currently on treatment with ustekinumab commenced prior to 1 February 2023 and met all remaining criteria (criterion 2) below at the time of commencing treatment; or
- 2 Both:
 - 2.1 Patient has active ulcerative colitis: and
 - 2.2 Either:
 - 2.2.1 Patient has had an initial approval for prior biologic therapy for ulcerative colitis and has experienced intolerable side effects or insufficient benefit to meet renewal criteria; or
 - 2.2.2 Both:
 - 2.2.2.1 Patient meets the initiation criteria for prior biologic therapies for ulcerative colitis; and
 - 2.2.2.2 Other biologics for ulcerative colitis are contraindicated.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

Both:

- 1 Either:
 - 1.1 The SCCAI score has reduced by 2 points or more from the SCCAI score since initiation on biologic therapy; or
 - 1.2 PUCAI score has reduced by 10 points or more from the PUCAI score since initiation on biologic therapy*; and
- 2 Ustekinumab will be used at a dose no greater than 90 mg intravenously every 8 weeks.

Note: Criterion marked with * is for an unapproved indication.

VEDOLIZUMAB - PCT only - Special Authority see SA2183 below

⇒SA2183 Special Authority for Subsidy

Initial application — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active Crohn's disease: and
- 2 Any of the following:
 - 2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria (unless contraindicated); or
 - 2.2 Patient has a CDAI score of greater than or equal to 300, or HBI score of greater than or equal to 10; or
 - 2.3 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.4 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
 - 2.5 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Any of the following:
 - 3.1 Patient has tried but experienced an inadequate response to (including lack of initial response and/or loss of initial response) from prior therapy with immunomodulators and corticosteroids; or
 - 3.2 Patient has experienced intolerable side effects from immunomodulators and corticosteroids; or
 - 3.3 Immunomodulators and corticosteroids are contraindicated.

Renewal — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points, or HBI score has reduced by 3 points, from when the patient was initiated on biologic therapy; or
 - 1.2 CDAI score is 150 or less, or HBI is 4 or less; or
 - 1.3 The patient has experienced an adequate response to treatment, but CDAI score and/or HBI score cannot be assessed: and
- 2 Vedolizumab to administered at a dose no greater than 300 mg every 8 weeks.

Initial application — **(Crohn's disease - children*)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria (unless contraindicated); or
 - 2.2 Patient has a Paediatric Crohn's Disease Activity Index (PCDAI) score of greater than or equal to 30; or
 - 2.3 Patient has extensive small intestine disease; and
- 3 Any of the following:

Subsidy	Fı	ılly	Brand or
(Manufacturer's Price)	Subsidis	ed	Generic
\$	Per	/	Manufacturer

continued...

- 3.1 Patient has tried but experienced an inadequate response to (including lack of initial response and/or loss of initial response) from prior therapy with immunomodulators and corticosteroids; or
- 3.2 Patient has experienced intolerable side effects from immunomodulators and corticosteroids; or
- 3.3 Immunomodulators and corticosteroids are contraindicated.

Note: Indication marked with * is an unapproved indication.

Renewal — (Crohn's disease - children*) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from when the patient was initiated on biologic therapy; or
 - 1.2 PCDAI score is 15 or less; or
 - 1.3 The patient has experienced an adequate response to treatment, but CDAI score cannot be assessed; and
- 2 Vedolizumab to administered at a dose no greater than 300mg every 8 weeks.

Note: Indication marked with * is an unapproved indication.

Initial application — (ulcerative colitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active ulcerative colitis; and
- 2 Any of the following:
 - 2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria (unless contraindicated); or
 - 2.2 Patient has a SCCAI score is greater than or equal to 4; or
 - 2.3 Patient's PUCAI score is greater than or equal to 20*; and
- 3 Any of the following:
 - 3.1 Patient has tried but experienced an inadequate response to (including lack of initial response and/or loss of initial response) from prior therapy with immunomodulators and corticosteroids; or
 - 3.2 Patient has experienced intolerable side effects from immunomodulators and corticosteroids; or
 - 3.3 Immunomodulators and corticosteroids are contraindicated.

Note: Indication marked with * is an unapproved indication.


Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 The SCCAI score has reduced by 2 points or more from the SCCAI score since initiation on biologic therapy; or
 - 1.2 The PUCAI score has reduced by 10 points or more from the PUCAI score since initiation on biologic therapy *; and
- 2 Vedolizumab will be used at a dose no greater than 300 mg intravenously every 8 weeks.

Note: Indication marked with * is an unapproved indication.

Programmed Cell Death-1 (PD-1) Inhibitors



⇒SA2443 Special Authority for Subsidy

Initial application — (non-small cell lung cancer second line monotherapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

continued...

All of the following:

- 1 Patient has locally advanced or metastatic non-small cell lung cancer; and
- 2 Patient has not received prior funded treatment with an immune checkpoint inhibitor for NSCLC; and
- 3 For patients with non-squamous histology there is documentation confirming that the disease does not express activating mutations of EGFR or ALK tyrosine kinase unless not possible to ascertain; and
- 4 Patient has an ECOG 0-2: and
- 5 Patient has documented disease progression following treatment with at least two cycles of platinum-based chemotherapy; and
- 6 Atezolizumab is to be used as monotherapy at a dose of 1200 mg every three weeks (or equivalent) for a maximum of 16 weeks; and
- 7 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Renewal — (non-small cell lung cancer second line monotherapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Any of the following:
 - 1.1 Patient's disease has had a complete response to treatment; or
 - 1.2 Patient's disease has had a partial response to treatment; or
 - 1.3 Patient has stable disease: and
- 2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period: and
- 3 No evidence of disease progression; and
- 4 The treatment remains clinically appropriate and patient is benefitting from treatment; and
- 5 Atezolizumab to be used at a maximum dose of 1200 mg every three weeks (or equivalent); and
- 6 Treatment with atezolizumab to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initial application — (unresectable hepatocellular carcinoma) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Patient is currently on treatment with atezolizumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 Patient has locally advanced or metastatic, unresectable hepatocellular carcinoma; and
 - 2.2 Patient has preserved liver function (Child-Pugh A); and
 - 2.3 Transarterial chemoembolisation (TACE) is unsuitable; and
 - 2.4 Any of the following:
 - 2.4.1 Patient has not received prior systemic therapy for the treatment of hepatocellular carcinoma; or
 - 2.4.2 Patient received funded lenvatinib before 1 March 2025; or
 - 2.4.3 Both:
 - 2.4.3.1 Patient has experienced treatment-limiting toxicity from treatment with lenvatinib; and
 - 2.4.3.2 No disease progression since initiation of lenvatinib; and
 - 2.5 Patient has an ECOG performance status of 0-2; and
 - 2.6 To be given in combination with bevacizumab.

Renewal — (unresectable hepatocellular carcinoma) from any relevant practitioner. Approvals valid for 6 months where there is no evidence of disease progression.

ALUMAB – PCT only – Specialist – Special Autho	rity see SA2425 on the ne	xt page	
50 mg per ml, 10 ml vial	4,700.00	1	Imfinzi
50 mg per ml, 2.4 ml vial		1	Imfinzi
1 mg for ECP	9.59	1 mg	✓ Baxter

Subside	/ Fu	ılly Brand or
(Manufacturer	,	sed Generic
	Per	✓ Manufacturer

⇒SA2425 Special Authority for Subsidy

Initial application — (Non-small cell lung cancer) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 Patient has histologically or cytologically documented stage III, locally advanced, unresectable non-small cell lung cancer (NSCLC): or
 - 1.2 Patient has histologically or cytologically documented stage IIb (T1N2a only), locally advanced, unresectable non-small cell lung cancer (NSCLC); and
- 2 Patient has received two or more cycles of platinum-based chemotherapy concurrently with definitive radiation therapy; and
- 3 Patient has no disease progression following the second or subsequent cycle of platinum-based chemotherapy with definitive radiation therapy treatment; and
- 4 Patient has a ECOG performance status of 0 or 1; and
- 5 Patient has completed last radiation dose within 8 weeks of starting treatment with durvalumab; and
- 6 Patient must not have received prior PD-1 or PD-L1 inhibitor therapy for this condition; and
- 7 Fither:
 - 7.1 Durvalumab is to be used at a maximum dose of no greater than 10 mg/kg every 2 weeks; or
 - 7.2 Durvalumab is to be used at a flat dose of 1500 mg every 4 weeks; and
- 8 Treatment with durvalumab to cease upon signs of disease progression.

Renewal — (Non-small cell lung cancer) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The treatment remains clinically appropriate and the patient is benefitting from treatment; and
- 2 Either:
 - 2.1 Durvalumab is to be used at a maximum dose of no greater than 10 mg/kg every 2 weeks; or
 - 2.2 Durvalumab is to be used at a flat dose of 1500 mg every 4 weeks; and
- 3 Treatment with durvalumab to cease upon signs of disease progression; and
- 4 Total continuous treatment duration must not exceed 12 months.

IPILIMUMAB - PCT only - Specialist - Special Authority see SA2461 below		
Inj 5 mg per ml, 10 ml vial	1	✓ Yervoy
Inj 5 mg per ml, 40 ml vial20,000.00	1	✓ Yervoy
Inj 1 mg for ECP106.00	1 mg	✓ Baxter

⇒SA2461 Special Authority for Subsidy

Initial application — **(renal cell carcinoma)** from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 The patient is currently on treatment with ipilimumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 The patient has metastatic renal cell carcinoma; and
 - 2.2 The patient is treatment naive; and
 - 2.3 The patient has ECOG performance status 0-2; and
 - 2.4 The disease is predominantly of clear cell histology; and
 - 2.5 Any of the following:
 - 2.5.1 The patient has sarcomatoid histology; or
 - 2.5.2 Haemoglobin levels less than the lower limit of normal; or

Subsidy (Manufacturer's Pric \$	e) Per	Fully Subsidised	Brand or Generic Manufacturer	
---------------------------------------	-----------	---------------------	-------------------------------------	--

continued...

- 2.5.3 Corrected serum calcium level greater than 10 mg/dL (2.5 mmol/L); or
- 2.5.4 Neutrophils greater than the upper limit of normal; or
- 2.5.5 Platelets greater than the upper limit of normal; or
- 2.5.6 Interval of less than 1 year from original diagnosis to the start of systemic therapy; or
- 2.5.7 Karnofsky performance score of less than or equal to 70; and
- 2.6 Ipilimumab is to be used at a maximum dose of 1 mg/kg for up to four cycles in combination with nivolumab...

NIVOLUMAB - PCT only - Specialist - Special Authority see SA2490 below

Inj 10 mg per ml, 4 ml vial	1,051.98	1	Opdivo
Inj 10 mg per ml, 10 ml vial	2,629.96	1	Opdivo
Inj 1 mg for ECP	27.22	1 mg	✓ Baxter

⇒SA2490 Special Authority for Subsidy

Initial application — (unresectable or metastatic melanoma) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The individual has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Baseline measurement of overall tumour burden is documented clinically and radiologically; and
- 3 The individual has ECOG performance 0-2; and
- 4 Either:
 - 4.1 The individual has not received funded pembrolizumab; or
 - 4.2 Both:
 - 4.2.1 The individual has received an initial Special Authority approval for pembrolizumab and has discontinued pembrolizumab within 12 weeks of starting treatment due to intolerance; and
 - 4.2.2 The cancer did not progress while the indvidual was on pembrolizumab; and
- 5 Any of the following:
 - 5.1 The individual has been diagnosed in the metastatic or unresectable stage III or IV setting; or
 - 5.2 The individual did not receive treatment in the perioperative setting with a PD-1/PD-L1 inhibitor; or
 - 5.3 All of the following:
 - 5.3.1 The individual received treatment in the perioperative setting with a PD-1/PD-L1 inhibitor; and
 - 5.3.2 The individual did not experience disease recurrence while on treatment with that PD-1/PD-L1 inhibitor; and
 - 5.3.3 The individual did not experience disease recurrence within six months of completing perioperative treatment with a PD-1/PD-L1 inhibitor.

Renewal — (unresectable or metastatic melanoma, less than 24 months on treatment) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Any of the following:
 - 1.1.1 The individual's disease has had a complete response to treatment; or
 - 1.1.2 The individual's disease has had a partial response to treatment; or
 - 1.1.3 The individual has stable disease; and
 - 1.2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period: or
- 2 All of the following:
 - 2.1 The individual has previously discontinued treatment with nivolumab for reasons other than severe toxicity or disease progression; and
 - 2.2 The individual has signs of disease progression; and

Subsidy (Manufacturer's Price)	Fully Subsidised		
\$	Per 🗸	Manufacturer	

continued...

2.3 Disease has not progressed during previous treatment with nivolumab.

Renewal — (unresectable or metastatic melanoma, more than 24 months on treatment) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 The individual has been on treatment for more than 24 months; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Any of the following:
 - 2.1.1.1 The individual's disease has had a complete response to treatment; or
 - 2.1.1.2 The individual's disease has had a partial response to treatment; or
 - 2.1.1.3 The individual has stable disease: and
 - 2.1.2 Response to treatment in target lesions has been determined by comparable radiologic or clinical assessment following the most recent treatment period; or
 - 2.2 All of the following:
 - 2.2.1 The individual has previously discontinued treatment with nivolumab for reasons other than severe toxicity or disease progression; and
 - 2.2.2 The individual has signs of disease progression; and
 - 2.2.3 Disease has not progressed during previous treatment with nivolumab.

Initial application — (renal cell carcinoma, first line) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Patient is currently on treatment with nivolumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 The patient has metastatic renal cell carcinoma; and
 - 2.2 The patient is treatment naive: and
 - 2.3 The patient has ECOG performance status 0-2; and
 - 2.4 The disease is predominantly of clear cell histology; and
 - 2.5 Any of the following:
 - 2.5.1 The patient has sarcomatoid histology; or
 - 2.5.2 Haemoglobin levels less than the lower limit of normal: or
 - 2.5.3 Corrected serum calcium level greater than 10 mg/dL (2.5 mmol/L); or
 - 2.5.4 Neutrophils greater than the upper limit of normal; or
 - 2.5.5 Platelets greater than the upper limit of normal; or
 - 2.5.6 Interval of less than 1 year from original diagnosis to the start of systemic therapy; or
 - 2.5.7 Karnofsky performance score of less than or equal to 70; and
 - 2.6 Nivolumab is to be used in combination with ipilimumab for the first four treatment cycles at a maximum dose of 3 mg/kg; and
 - 2.7 Nivolumab is to be used as monotherapy at a maximum maintenance dose of 240 mg every 2 weeks (or equivalent).

Initial application — (Renal cell carcinoma, second line) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has metastatic renal-cell carcinoma; and
- 2 The disease is of predominant clear-cell histology; and
- 3 Patient has ECOG performance status 0-2; and
- 4 Patient has documented disease progression following one or two previous regimens of antiangiogenic therapy; and

continued...

- 5 Patient has not previously received a funded immune checkpoint inhibitor; and
- 6 Nivolumab is to be used as monotherapy at a maximum dose of 240 mg every 2 weeks (or equivalent) and discontinued at disease progression.

Renewal — (Renal cell carcinoma) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 Patient's disease has had a complete response to treatment; or
 - 1.2 Patient's disease has had a partial response to treatment; or
 - 1.3 Patient has stable disease; and
- 2 No evidence of disease progression; and
- 3 Nivolumab is to be used as monotherapy at a maximum dose of 240 mg every 2 weeks (or equivalent) and discontinued at disease progression.

PEMBROLIZUMAB - PCT only - Specialist - Special A	uthority see SA2553 below		
Inj 25 mg per ml, 4 ml vial	4,680.00	1	✓ Keytruda
Inj 1 mg for ECP	47.74 1	mg	✓ Baxter

⇒SA2553 Special Authority for Subsidy

Initial application — (stage III or IV resectable melanoma - neoadjuvant) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The individual has resectable stage IIIB, IIIC, IIID or IV melanoma (excluding uveal) (see note); and
- 2 The individual has not received prior funded systemic treatment in the perioperative setting for their stage IIIB, IIIC, IIID or IV melanoma; and
- 3 Treatment must be prior to complete surgical resection; and
- 4 Pembrolizumab must be administered as monotherapy; and
- 5 The individual has ECOG performance score 0-2; and
- 6 Pembrolizumab to be administered at a fixed dose of 200 mg every 3 weeks (or equivalent).

Renewal — (stage III or IV resectable melanoma - neoadjuvant) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 The individual has received neoadjuvant treatment with an immune checkpoint inhibitor; and
 - 1.2 The individual meets initial application criteria for pembrolizumab for stage III or IV resected melanoma adjuvant; or
- 2 Both:
 - 2.1 The individual has received neoadjuvant and adjuvant treatment with an immune checkpoint inhibitor; and
 - 2.2 The individual meets renewal criteria for pembrolizumab for stage III or IV resected melanoma adjuvant; or
- 3 All of the following:
 - 3.1 The individual has received neoadiuvant and adjuvant treatment with an immune checkpoint inhibitor; and
 - 3.2 The individual has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
 - 3.3 The individual meets initial application criteria for pembrolizumab for unresectable or metastatic melanoma; or
- 4 All of the following:
 - 4.1 The individual has received neoadjuvant and adjuvant treatment with an immune checkpoint inhibitor; and
 - 4.2 The individual has received treatment with an immune checkpoint inhibitor for unresectable or metastatic melanoma; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	1	Manufacturer	

continued...

4.3 The individual meets renewal criteria for pembrolizumab for unresectable or metastatic melanoma.

Notes:

- a) Stage IIIB, IIIC, IIID or IV melanoma defined as per American Joint Committee on Cancer (AJCC) 8th Edition
- b) Initiating treatment within 13 weeks of complete surgical resection means either 13 weeks after resection (primary or lymphadenectomy) or 13 weeks prior to the scheduled date of the resection (primary or lymphadenectomy)

Initial application — (stage III or IV resected melanoma - adjuvant) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 The individual has resected stage IIIB, IIIC, IIID or IV melanoma (excluding uveal) (see note a); and
- 2 Adjuvant treatment with pembrolizumab is required; and
- 3 The individual has not received prior funded systemic treatment in the adjuvant setting for stage IIIB, IIIC, IIID or IV melanoma; and
- 4 Treatment must be in addition to complete surgical resection; and
- 5 Treatment must be initiated within 13 weeks of complete surgical resection, unless delay is necessary due to post-surgery recovery (see note b); and
- 6 Pembrolizumab must be administered as monotherapy; and
- 7 The individual has ECOG performance score 0-2; and
- 8 Pembrolizumab to be administered at a fixed dose of 200 mg every 3 weeks (or equivalent).

Notes:

- a) Stage IIIB, IIIC, IIID or IV melanoma defined as per American Joint Committee on Cancer (AJCC) 8th Edition
- b) Initiating treatment within 13 weeks of complete surgical resection means 13 weeks after resection (primary or lymphadenectomy)

Renewal — (stage III or IV resected melanoma - adjuvant) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria: Any of the following:

- 1 All of the following:
 - 1.1 No evidence of disease recurrence: and
 - 1.2 Pembrolizumab must be administered as monotherapy; and
 - 1.3 Pembrolizumab to be administered at a fixed dose of 200 mg every three weeks (or equivalent) for a maximum of 12 months total treatment course, including any systemic neoadjuvant treatment; and
 - 1.4 Treatment to be discontinued at signs of disease recurrence or at completion of 12 months total treatment course (equivalent to 18 cycles at a dose of 200 mg every 3 weeks), including any systemic neoadjuvant treatment; or
- 2 All of the following:
 - 2.1 The individual has received adjuvant treatment with an immune checkpoint inhibitor; and
 - 2.2 The individual has metastatic or unresectable melanoma (excluding uveal) stage III or IV: and
 - 2.3 The individual meets initial application criteria for pembrolizumab for unresectable or metastatic melanoma; or
- 3 All of the following:
 - 3.1 The individual has received adjuvant treatment with an immune checkpoint inhibitor; and
 - 3.2 The individual has received treatment with an immune checkpoint inhibitor for unresectable or metastatic melanoma; and
 - 3.3 The individual meets renewal criteria for pembrolizumab for unresectable or metastatic melanoma.

Initial application — (unresectable or metastatic melanoma) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The individual has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Baseline measurement of overall turnour burden is documented clinically and radiologically; and
- 3 The individual has ECOG performance score of 0-2; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 4 Either:
 - 4.1 The individual has not received funded nivolumab; or
 - 4.2 Both:
 - 4.2.1 The individual has received an initial Special Authority approval for nivolumab and has discontinued nivolumab within 12 weeks of starting treatment due to intolerance; and
 - 4.2.2 The cancer did not progress while the individual was on nivolumab; and
 - 5 Any of the following:
 - 5.1 The individual has been diagnosed in the metastatic or unresectable stage III or IV setting; or
 - 5.2 The individual did not receive treatment in the perioperative setting with a PD-1/PD-L1 inhibitor; or
 - 5.3 All of the following:
 - 5.3.1 The individual received treatment in the perioperative setting with a PD-1/PD-L1 inhibitor; and
 - 5.3.2 The individual did not experience disease recurrence while on treatment with that PD-1/PD-L1 inhibitor; and
 - 5.3.3 The individual did not experience disease recurrence within six months of completing perioperative treatment with a PD-1/PD-I 1 inhibitor.

Renewal — (unresectable or metastatic melanoma, less than 24 months on treatment) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Any of the following:
 - 1.1.1 The individual's disease has had a complete response to treatment; or
 - 1.1.2 The individual's disease has had a partial response to treatment; or
 - 1.1.3 The individual has stable disease; and
 - 1.2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period; or
- 2 All of the following:
 - 2.1 The individual has previously discontinued treatment with pembrolizumab for reasons other than severe toxicity or disease progression; and
 - 2.2 The individual has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with pembrolizumab.

Renewal — (unresectable or metastatic melanoma, more than 24 months on treatment) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 The individual has been on treatment for more than 24 months; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Any of the following:
 - 2.1.1.1 The individual's disease has had a complete response to treatment; or
 - 2.1.1.2 The individual's disease has had a partial response to treatment; or
 - 2.1.1.3 The individual has stable disease; and
 - 2.1.2 Response to treatment in target lesions has been determined by comparable radiologic or clinical assessment following the most recent treatment period; or
 - 2.2 All of the following:
 - 2.2.1 The individual has previously discontinued treatment with pembrolizumab for reasons other than severe toxicity or disease progression; and

(Manufacturer's Price) Subsidi		Fully Subsidised	Brand or Generic
\$	Per	1	Manufacturer

continued...

- 2.2.2 The individual has signs of disease progression; and
- 2.2.3 Disease has not progressed during previous treatment with pembrolizumab.

Initial application — (non-small cell lung cancer first-line monotherapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-small cell lung cancer; and
- 2 Patient has not had chemotherapy for their disease in the palliative setting; and
- 3 Patient has not received prior funded treatment with an immune checkpoint inhibitor for NSCLC; and
- 4 For patients with non-squamous histology there is documentation confirming that the disease does not express activating mutations of EGFR or ALK tyrosine kinase unless not possible to ascertain; and
- 5 Pembrolizumab to be used as monotherapy; and
- 6 Either:
 - 6.1 There is documentation confirming the disease expresses PD-L1 at a level greater than or equal to 50% as determined by a validated test unless not possible to ascertain; or
 - 6.2 Both:
 - 6.2.1 There is documentation confirming the disease expresses PD-L1 at a level greater than or equal to 1% as determined by a validated test unless not possible to ascertain; and
 - 6.2.2 Chemotherapy is determined to be not in the best interest of the patient based on clinician assessment; and
- 7 Patient has an ECOG 0-2: and
- 8 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks; and
- 9 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Renewal — (non-small cell lung cancer first line monotherapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 Patient's disease has had a complete response to treatment: or
 - 1.2 Patient's disease has had a partial response to treatment; or
 - 1.3 Patient has stable disease; and
- 2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period; and
- 3 No evidence of disease progression; and
- 4 The treatment remains clinically appropriate and patient is benefitting from treatment; and
- 5 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 6 Treatment with pembrolizumab to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initial application — (non-small cell lung cancer first-line combination therapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-small cell lung cancer; and
- 2 The patient has not had chemotherapy for their disease in the palliative setting; and
- 3 Patient has not received prior funded treatment with an immune checkpoint inhibitor for NSCLC; and
- 4 For patients with non-squamous histology there is documentation confirming that the disease does not express activating mutations of EGFR or ALK tyrosine kinase unless not possible to ascertain; and
- 5 Pembrolizumab to be used in combination with platinum-based chemotherapy; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 6 Patient has an ECOG 0-2; and
- 7 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks; and
- 8 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Renewal — (non-small cell lung cancer first line combination therapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 Patient's disease has had a complete response to treatment; or
 - 1.2 Patient's disease has had a partial response to treatment; or
 - 1.3 Patient has stable disease; and
- 2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period: and
- 3 No evidence of disease progression; and
- 4 The treatment remains clinically appropriate and patient is benefitting from treatment; and
- 5 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 6 Treatment with pembrolizumab to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initial application — (breast cancer, advanced) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: Fither:

- 1 Patient is currently on treatment with pembrolizumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has recurrent or de novo unresectable, inoperable locally advanced triple-negative breast cancer (that does not express ER, PR or HER2 IHC3+ or ISH+ [including FISH or other technology]); or
 - 2.1.2 Patient has recurrent or de novo metastatic triple-negative breast cancer (that does not express ER, PR or HER2 IHC3+ or ISH+ [including FISH or other technology]); and
 - 2.2 Patient is treated with palliative intent: and
 - 2.3 Patient's cancer has confirmed PD-L1 Combined Positive Score (CPS) is greater than or equal to 10; and
 - 2.4 Patient has received no prior systemic therapy in the palliative setting; and
 - 2.5 Patient has an ECOG score of 0-2; and
 - 2.6 Pembrolizumab is to be used in combination with chemotherapy; and
 - 2.7 Baseline measurement of overall tumour burden is documented clinically and radiologically; and
 - 2.8 Pembrolizumab is to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks.

Renewal — (breast cancer, advanced) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 Patient's disease has had a complete response to treatment; or
 - 1.2 Patient's disease has had a partial response to treatment; or
 - 1.3 Patient has stable disease: and
 - 2 No evidence of disease progression; and
 - 3 Response to treatment in target lesions has been determined by a comparable radiologic assessment following the most recent treatment period: and

Subsidy (Manufactured a Drice)		Fully	Brand or
 (Manufacturer's Price) \$	Subside Per	JISEO 🗸	Generic Manufacturer

continued...

- 4 Pembrolizumab is to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 5 Treatment with pembrolizumab is to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initial application — (head and neck squamous cell carcinoma) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria: Either:

- 1 Patient is currently on treatment with pembrolizumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 Patient has recurrent or metastatic head and neck squamous cell carcinoma of mucosal origin (excluding nasopharyngeal carcinoma) that is incurable by local therapies; and
 - 2.2 Patient has not received prior systemic therapy in the recurrent or metastatic setting; and
 - 2.3 Patient has a positive PD-L1 combined positive score (CPS) of greater than or equal to 1; and
 - 2.4 Patient has an ECOG performance score of 0-2; and
 - 2.5 Fither:
 - 2.5.1 Pembrolizumab to be used in combination with platinum-based chemotherapy; or
 - 2.5.2 Pembrolizumab to be used as monotherapy; and
- 2.6 Pembrolizumab is to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks.

Renewal — (head and neck squamous cell carcinoma) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 Patient's disease has had a complete response to treatment; or
 - 1.2 Patient's disease has had a partial response to treatment; or
 - 1.3 Patient has stable disease: and
- 2 No evidence of disease progression; and
- 3 Pembrolizumab is to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 4 Treatment with pembrolizumab is to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initial application — (MSI-H/dMMR advanced colorectal cancer) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria: Either:

- 1 Individual is currently on treatment with pembrolizumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 Fither:
 - 2.1.1 Individual has deficient mismatch repair (dMMR) or microsatellite instability-high (MSI-H) metastatic colorectal cancer; or
 - 2.1.2 Individual has deficient mismatch repair (dMMR) or microsatellite instability-high (MSI-H) unresectable colorectal cancer; and
 - 2.2 Individual is treated with palliative intent; and
 - 2.3 Individual has not previously received funded treatment with pembrolizumab for MSI-H/dMMR advanced colorectal cancer; and
 - 2.4 Individual has an ECOG performance score of 0-2; and
 - 2.5 Baseline measurement of overall tumour burden is documented clinically and radiologically; and
 - 2.6 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks.

Renewal — (MSI-H/dMMR advanced colorectal cancer) from any relevant practitioner. Approvals valid for 4 months for

Subsidy		Fully	Brand or	
(Manufacturer's Price)	9	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued

applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 3 Treatment with pembrolizumab is to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initial application — (Urothelial carcinoma) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 Patient is currently on treatment with pembrolizumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 Patient has inoperable locally advanced (T4) or metastatic urothelial carcinoma; and
 - 2.2 Patient has an ECOG performance score of 0-2; and
 - 2.3 Patient has documented disease progression following treatment with chemotherapy; and
 - 2.4 Pembrolizumab to be used as monotherapy at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks.

Renewal — (Urothelial carcinoma) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 Patient's disease has had a complete response to treatment; or
 - 1.2 Patient's disease has had a partial response to treatment; or
 - 1.3 Patient has stable disease; and
- 2 No evidence of disease progression; and
- 3 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 4 Treatment with pembrolizumab is to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initial application — (relapsed/refractory Hodgkin lymphoma) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria: Fither:

- 1 Individual is currently on treatment with pembrolizumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Both:
 - 2.1.1.1 Individual has relapsed/refractory Hodgkin lymphoma after two or more lines of chemotherapy; and
 - 2.1.1.2 Individual is ineligible for autologous stem cell transplant; or
 - 2.1.2 Individual has relapsed/refractory Hodgkin lymphoma and has previously undergone an autologous stem cell transplant; and
 - 2.2 Individual has not previously received funded pembrolizumab for relapsed/refractory Hodgkin lymphoma; and
 - 2.3 Pembrolizumab to be administered at doses no greater than 200 mg once every 3 weeks.

Renewal — (relapsed/refractory Hodgkin lymphoma) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient has received a partial or complete response to pembrolizumab; and
- 2 Treatment with pembrolizumab is to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

	(Manufacturer's Price) \$	Subs Per	idised	Generic Manufacturer
Other Immunecumpressents				

Cuboldy

E. ili.

Drand or

Other Immunosuppressants

44.63	50	Neoral
88.91	50	Neoral
177.81	50	Neoral
198.13	50 ml OP	Neoral
il pharmacy		
6,512.29	30	Afinitor
4,555.76	30	Afinitor

⇒SA2414 Special Authority for Subsidy

Initial application only from a neurologist or oncologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has tuberous sclerosis; and
- 2 Patient has progressively enlarging sub-ependymal giant cell astrocytomas (SEGAs) that require treatment.

Renewal only from a neurologist or oncologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Documented evidence of SEGA reduction or stabilisation by MRI within the last 3 months; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Everolimus to be discontinued at progression of SEGAs.

Initial application — **(renal cell carcinoma)** from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient has metastatic renal cell carcinoma; and
 - 1.2 The disease is of predominant clear-cell histology; and
 - 1.3 The patient has documented disease progression following one previous line of treatment; and
 - 1.4 The patient has an ECOG performance status of 0-2; and
 - 1.5 Everolimus is to be used in combination with lenvatinib; or
- 2 All of the following:
 - 2.1 Patient has received funded treatment with nivolumab for the second line treatment of metastatic renal cell carcinoma; and
 - 2.2 Patient has experienced treatment limiting toxicity from treatment with nivolumab; and
 - 2.3 Everolimus is to be used in combination with lenvatinib; and
 - 2.4 There is no evidence of disease progression.

Renewal — (renal cell carcinoma) from any relevant practitioner. Approvals valid for 4 months where there is no evidence of disease progression.

SIROLIMUS - Special Authority see SA2270 below - Re	etail pharmacy		
Tab 1 mg	749.99	100	Rapamune
Tab 2 mg	1,499.99	100	✓ Rapamune
Oral liq 1 mg per ml	449.99	60 ml OP	✓ Rapamune

⇒SA2270 Special Authority for Subsidy

Initial application from any medical practitioner. Approvals valid without further renewal unless notified where the drug is to be used for rescue therapy for an organ transplant recipient.

Notes: Rescue therapy defined as unresponsive to calcineurin inhibitor treatment as defined by refractory rejection; or intolerant

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

to calcineurin inhibitor treatment due to any of the following:

- GFR< 30 ml/min; or
- Rapidly progressive transplant vasculopathy; or
- · Rapidly progressive obstructive bronchiolitis; or
- . HUS or TTP: or
- · Leukoencepthalopathy; or
- · Significant malignant disease

Initial application — (severe non-malignant lymphovascular malformations*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe non-malignant lymphovascular malformation*; and
- 2 Any of the following:
 - 2.1 Malformations are not adequately controlled by sclerotherapy and surgery; or
 - 2.2 Malformations are widespread/extensive and sclerotherapy and surgery are not considered clinically appropriate; or
 - 2.3 Sirolimus is to be used to reduce malformation prior to consideration of surgery; and
- 3 Patient is being treated by a specialist lymphovascular malformation multi-disciplinary team; and
- 4 Patient has measurable disease as defined by RECIST version 1.1 (see Note).

Renewal — (severe non-malignant lymphovascular malformations*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient's disease has had either a complete response or a partial response to treatment, or patient has stable disease according to RECIST version 1.1 (see Note); or
 - 1.2 Patient's disease has stabilised or responded clinically and disease response to treatment has been clearly documents in patient notes; and
- 2 No evidence of progressive disease; and
- 3 The treatment remains clinically appropriate and the patient is benefitting from the treatment.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer et al. Eur J Cancer 2009;45:228-47)

Indications marked with * are unapproved indications

Initial application — (renal angiomyolipoma(s) associated with tuberous sclerosis complex*) only from a nephrologist or urologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has tuberous sclerosis complex*: and
- 2 Evidence of renal angiomyolipoma(s) measuring 3 cm or greater and that have shown interval growth.

Renewal — (renal angiomyolipoma(s) associated with tuberous sclerosis complex*) from any relevant practitioner.

Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Documented evidence of renal angiomyolipoma reduction or stability by magnetic resonance imaging (MRI) or ultrasound;
- 2 Demonstrated stabilisation or improvement in renal function; and
- 3 The patient has not experienced angiomyolipoma haemorrhage or significant adverse effects to sirolimus treatment; and
- 4 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indications marked with * are unapproved indications

Initial application — (refractory seizures associated with tuberous sclerosis complex*) only from a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

 Subsidy Fully (Manufacturer's Price) Subsidised		. ,	Brand or Generic
 \$	Per	1	Manufacturer

continued...

- 1 Patient has epilepsy with a background of documented tuberous sclerosis complex; and
- 2 Fither:
 - 2.1 Both:
 - 2.1.1 Vigabatrin has been trialled and has not adequately controlled seizures; and
 - 2.1.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least two of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); or
 - 2.2 Both:
 - 2.2.1 Vigabatrin is contraindicated; and
 - 2.2.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least three of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); and
- 3 Seizures have a significant impact on quality of life; and
- 4 Patient has been assessed and surgery is considered inappropriate for this patient, or the patient has been assessed and would benefit from mTOR inhibitor treatment prior to surgery.

Note: Those of childbearing age potential are not required to trial phenytoin sodium, sodium valproate, or topiramate. Those who can father children are not required to trial sodium valproate.

Renewal — (refractory seizures associated with tuberous sclerosis complex*) only from a neurologist. Approvals valid for 12 months where demonstrated significant and sustained improvement in seizure rate (e.g. 50% reduction in seizure frequency) or severity and/or patient quality of life compared with baseline prior to starting sirolimus treatment.

Note: Indications marked with * are unapproved indications

TACROLIMUS – Special Authority see SA2455 below – Retail pharmacy

Cap 0.5 mg	49.60	100	✓ Tacrolimus Sandoz
Cap 0.75 mg		100	✓ Tacrolimus Sandoz
Cap 1 mg	84.30	100	✓ Tacrolimus Sandoz
Cap 5 mg	.248.20	50	✓ Tacrolimus Sandoz

⇒SA2455 Special Authority for Subsidy

Initial application — (organ transplant) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Fither:

- 1 The individual is an organ transplant recipient; or
- 2 The individual is receiving induction therapy for an organ transplant.

Note: Subsidy applies for either primary or rescue therapy.

Initial application — (non-transplant indications*) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient requires long-term systemic immunosuppression; and
- 2 Either:
 - 2.1 Ciclosporin has been trialled and discontinued treatment because of unacceptable side effects or inadequate clinical response; or
 - 2.2 Patient is a child with nephrotic syndrome*.

Note: Indications marked with * are unapproved indications

JAK inhibitors

UPADACITINIB – Special Authority see SA2483 on the nex	<mark>kt page – Retail pharmacy</mark>	1	
Tab modified-release 15 mg	1,271.00	28	✓ Rinvoq
Tab modified-release 30 mg	2,033.00	28	✓ Rinvoq
Tab modified-release 45 mg		28	✓ Rinvoq

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA2483 Special Authority for Subsidy

Initial application — (Rheumatoid Arthritis (previously treated with adalimumab or etanercept)) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The individual has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Fither:
 - 2.1 The individual has experienced intolerable side effects with adalimumab and/or etanercept; or
 - 2.2 The individual has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis: and
- 3 Any of the following:
 - 3.1 Rituximab is not clinically appropriate; or
 - 3.2 The individual is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or
 - 3.3 Both:
 - 3.3.1 The individual has been started on rituximab for rheumatoid arthritis in a Health NZ Hospital: and
 - 3.3.2 Either:
 - 3.3.2.1 The individual has experienced intolerable side effects with rituximab; or
 - 3.3.2.2 At four months following the initial course of rituximab the individual has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Renewal — (Rheumatoid Arthritis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Following 6 months' initial treatment, the individual has experienced at least a 50% decrease in active joint count from baseline: or
- 2 On subsequent reapplications, the individual has experienced at least a continuing 30% improvement in active joint count from baseline.

Initial application — (atopic dermatitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Individual is currently on treatment with upadacitinib for atopic dermatitis and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 Individual has moderate to severe atopic dermatitis, severity as defined by an Eczema Area and Severity Index (EASI) score of greater than or equal to 16 or a Dermatology Life Quality Index (DLQI) score of greater than or equal to 10; and
 - 2.2 Individual has received insufficient benefit from topical therapy (including topical corticosteroids or topical calcineurin inhibitors) for a 28-day trial within the last 6 months, unless contraindicated to all; and
 - 2.3 Individual has trialled and received insufficient benefit from at least one systemic therapy for a minimum of three months (eq ciclosporin, azathioprine, methotrexate or mycophenolate mofetil), unless contraindicated to all; and
 - 2.4 An EASI assessment or DLQI assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
 - 2.5 The most recent EASI or DQLI assessment is no more than 1 month old at the time of application.

Renewal — (atopic dermatitis) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Fither:

- 1 Individual has received a 75% or greater reduction in EASI score (EASI 75) as compared to baseline EASI prior to commencing upadacitinib; or
- 2 Individual has received a DLQI improvement of 4 or more as compared to baseline DLQI prior to commencing upadacitinib.

Subsidy (Manufacturer's Price)	Full Subsidise		
\$	Per 🗸	Manufacturer	

continued...

Initial application — (Crohn's disease - adult) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Individual is currently on treatment with upadacitinib for Crohn's disease and met all remaining criteria prior to commencing treatment: or
- 2 Both:
 - 2.1 Individual has active Crohn's disease; and
 - 2.2 Fither:
 - 2.2.1 Individual has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria; or
 - 2.2.2 Both:
 - 2.2.2.1 Individual meets the initiation criteria for prior biologic therapies for Crohn's disease; and
 - 2.2.2.2 Other biologic therapies for Crohn's disease are contraindicated.

Renewal — (Crohn's disease - adult) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 CDAI score has reduced by 100 points from the CDAI score when the individual was initiated on biologic therapy; or
- 2 HBI score has reduced by 3 points from when individual was initiated on biologic therapy; or
- 3 CDAI score is 150 or less; or
- 4 HBI score is 4 or less: or
- 5 The individual has experienced an adequate response to treatment, but CDAI score cannot be assessed.

Initial application — (Crohn's disease - children*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Individual is currently on treatment with upadacitinib for Crohn's disease and met all remaining criteria prior to commencing
- 2 Both:
 - 2.1 Child has active Crohn's disease: and
 - 2.2 Either:
 - 2.2.1 Child has had an initial approval for prior biologic therapy for Crohn's disease and has experienced intolerable side effects or insufficient benefit to meet renewal criteria: or
 - 2.2.2 Both:
 - 2.2.2.1 Child meets the initiation criteria for prior biologic therapies for Crohn's disease; and
 - 2.2.2.2 Other biologic therapies for Crohn's disease are contraindicated.

Renewal — (Crohn's disease - children*) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 PCDAI score has reduced by 10 points from the child was initiated on treatment; or
- 2 PCDAI score is 15 or less; or
- 3 The child has experienced an adequate response to treatment, but PCDAI score cannot be assessed.

Note: Indications marked with * are unapproved indications.

Initial application — (ulcerative colitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Individual is currently on treatment with upadacitinib for ulcerative colitis and met all remaining criteria prior to commencing treatment: or
- 2 Roth:

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
\$	Per	1	Manufacturer

continued...

- 2.1 Individual has active ulcerative colitis; and
- 2.2 Either:
 - 2.2.1 Individual has had an initial approval for prior biologic therapy for ulcerative colitis and has experienced intolerable side effects or insufficient benefit to meet renewal criteria; or
 - 2.2.2 Both:
 - 2.2.2.1 Individual meets the initiation criteria for prior biologic therapies for ulcerative colitis; and 2.2.2.2 Other biologic therapies for ulcerative colitis are contraindicated.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 The SCCAI score has reduced by 2 points or more from the SCCAI score when the individual was initiated on treatment; or
- 2 PUCAI score has reduced by 10 points or more from the PUCAI score when the individual was initiated on treatment.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

Antiallergy Preparations

Allergic Emergencies

ADRENALINE - Special Authority see SA2185 below - Retail pharmacy

- a) Maximum of 2 ini per prescription
- Additional prescriptions limited to replacement of up to two devices prior to expiry, or replacement of used device for treatment of anaphylaxis.

Inj 0.15 mg per 0.3 ml auto-injector	85.50	1 OP	 Epipen Jr
Inj 0.3 mg per 0.3 ml auto-injector	85.50	1 OP	✓ Epipen

⇒SA2185 Special Authority for Subsidy

Initial application — (anaphylaxis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Patient has experienced an anaphylactic reaction which has resulted in presentation to a hospital or emergency department; or
 - 1.2 Patient has been assessed to be at significant risk of anaphylaxis by a relevant practitioner; and
- 2 Patient is not to be prescribed more than two devices in initial prescription.

⇒SA1558 Special Authority for Subsidy

Initial application only from a clinical immunologist or relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Supply for anticipated emergency treatment of laryngeal/oro-pharyngeal or severe abdominal attacks of acute hereditary angioedema (HAE) for patients with confirmed diagnosis of C1-esterase inhibitor deficiency; and
- 2 The patient has undergone product training and has agreed upon an action plan for self-administration.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Allergy Desensitisation

⇒SA1367 Special Authority for Subsidy

Initial application only from a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 RAST or skin test positive; and
- 2 Patient has had severe generalised reaction to the sensitising agent.

Renewal only from a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

BEE VENOM ALLERGY TREATMENT – Special Authority see SA1367 above – Retail ph	armacy
Initiation kit - 1 vial freeze dried venom with diluent305.00	✓ VENOX \$29
Maintenance kit - 1 vial freeze dried venom with diluent305.00	✓ VENOX \$29
Maintenance kit - 6 vials 120 mcg freeze dried venom, with	
diluent285.00 1 OF	✓ Venomil S29
Treatment kit - 1 vial 550 mcg freeze dried venom, 1 diluent	
9 ml, 3 diluent 1.8 ml	○ ✓ Albey
Treatment kit - 1 vial 550 mcg freeze dried venom, with diluent305.00	✓ Hymenoptera S29

	Subsidy		Fully	Brand or	
	(Manufacturer's Price		sidised	Generic	
	\$	Per		Manufacturer	
WASP VENOM ALLERGY TREATMENT - Special Authority se	ee SA1367 on the p	revious page	– Reta	ail pharmacy	
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze					
dried polistes venom, 1 diluent 9 ml, 3 diluent 1.8 ml	382.23	1 OP	✓ /	Albey	
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze					
dried venom, with diluent	305.00	1 OP	✓	Hymenoptera S29	
Treatment kit (Paper wasp venom) - 6 vials 120 mcg freeze)				
dried venom, with diluent	305.00	1 OP	/ \	Venomil S29	
Treatment kit (Yellow Jacket venom) - 1 vial 550 mcg freezo					
dried venom, with diluent	305.00	1 OP	√	Hymenoptera S29	
Treatment kit (Yellow jacket venom) - 1 vial 550 mcg freeze)				
dried vespula venom, 1 diluent 9 ml, 3 diluent 1.8 ml	431.24	1 OP	✓ /	Albey	
Treatment kit (Yellow jacket venom) - 6 vials 120 mcg freez					
dried venom, with diluent	305.00	1 OP	/ \	Venomil S29	
Audibistaninas					
Antihistamines					
CETIRIZINE HYDROCHLORIDE					
* Tab 10 mg	1.71	100	1	<u>Zista</u>	
* Oral liq 1 mg per ml		200 ml	✓	Histaclear	
DEXTROCHLORPHENIRAMINE MALEATE					
* Tab 2 mg	2.02	40			
•	(8.40)		F	Polaramine	
	1.01	20			
	(5.99)		F	Polaramine	
* Oral liq 2 mg per 5 ml		100 ml			
	(10.29)		ı	Polaramine	
FEXOFENADINE HYDROCHLORIDE					
* Tab 60 mg		20			
st. T 100	(8.23)	20		Telfast -	
* Tab 120 mg		30	-	Fexaclear	
* Tab 180 mg	4.10	30	V į	Fexaclear	
LORATADINE					
* Tab 10 mg		100	_	Lorafix	
* Oral liq 1 mg per ml	1.43	100 ml	✓ I	Haylor syrup	
PROMETHAZINE HYDROCHLORIDE			_		
* Tab 10 mg		100	_	Allersoothe	
* Tab 25 mg		100	_	Allersoothe	
* Oral liq 1 mg per 1 ml		100 ml		Allersoothe	
* Inj 25 mg per ml, 2 ml ampoule – Up to 5 inj available on a	P5021.09	5	•	Hospira	
Inhaled Corticosteroids					
BECLOMETHASONE DIPROPIONATE					
Aerosol inhaler, 50 mcg per dose	14.01 2	200 dose OP	1	Qvar	
Aerosol inhaler, 50 mcg per dose CFC-free		200 dose OP	✓	Beclazone 50	
Aerosol inhaler, 100 mcg per dose		200 dose OP	1	Qvar	
Aerosol inhaler, 100 mcg per dose CFC-free		200 dose OP	✓	Beclazone 100	
Aerosol inhaler, 250 mcg per dose CFC-free	22.67 2	200 dose OP	✓	Beclazone 250	

	Subsidy (Manufacturer's	Price) Subs	Fully	
	\$	Per	✓	Manufacturer
UDESONIDE				
Powder for inhalation, 100 mcg per dose	17.00	200 dose OP	•	Pulmicort Turbuhaler
Powder for inhalation, 200 mcg per dose	19.00	200 dose OP	•	Pulmicort Turbuhaler
Powder for inhalation, 400 mcg per dose	32.00	200 dose OP	•	Pulmicort Turbuhaler
LUTICASONE				
Aerosol inhaler, 50 mcg per dose	7.19	120 dose OP	1	Flixotide
Powder for inhalation, 50 mcg per dose		60 dose OP	1	Flixotide Accuhaler
Powder for inhalation, 100 mcg per dose		60 dose OP	1	Flixotide Accuhaler
Aerosol inhaler, 125 mcg per dose	13.60	120 dose OP		Flixotide
Aerosol inhaler, 250 mcg per dose		120 dose OP		Flixotide
Powder for inhalation, 250 mcg per dose		60 dose OP		Flixotide Accuhaler
nhaled Long-acting Beta-adrenoceptor Agonis	sts			
FORMOTEROL FUMARATE DIHYDRATE				
Powder for inhalation 4.5 mcg per dose, breath activated				
(equivalent to eformoterol fumarate 6 mcg metered dos	e) 10.32	60 dose OP		
(Squitalone to distinistion influence of mog motorod dos	(16.90)	00 0000 01		Oxis Turbuhaler
ID A CATEDOL	(10.30)			Oxio Turburiaici
IDACATEROL				
Powder for inhalation 150 mcg		30 dose OP		Onbrez Breezhaler
Powder for inhalation 300 mcg	61.00	30 dose OP	/	Onbrez Breezhaler
ALMETEROL				
Aerosol inhaler CFC-free, 25 mcg per dose	26.25	120 dose OP	1	Serevent
Powder for inhalation, 50 mcg per dose, breath activated		60 dose OP		Serevent Accuhaler
· • • · · ·				
nhaled Corticosteroids with Long-Acting Beta	-Aarenocepi	or Agonists		
UDESONIDE WITH EFORMOTEROL				
Powder for inhalation 160 mcg with 4.5 mcg eformoterol				
	with			
fumarate per dose (equivalent to 200 mcg budesonide				
fumarate per dose (equivalent to 200 mcg budesonide to 6 mcg eformoterol fumarate metered dose) – Up to 120	0	120 dose OP	•	DuoResp Spiromax
fumarate per dose (equivalent to 200 mcg budesonide v 6 mcg eformoterol fumarate metered dose) — Up to 120 dose available on a PSO	0 41.50	120 dose OP	•	DuoResp Spiromax
fumarate per dose (equivalent to 200 mcg budesonide of mcg eformoterol fumarate metered dose) – Up to 120 dose available on a PSO	0 41.50 arate	120 dose OP	/	DuoResp Spiromax
fumarate per dose (equivalent to 200 mcg budesonide of mcg eformoterol fumarate metered dose) – Up to 120 dose available on a PSO	0 41.50 arate cg	120 dose OP	•	DuoResp Spiromax
fumarate per dose (equivalent to 200 mcg budesonide of mcg eformoterol fumarate metered dose) – Up to 120 dose available on a PSO	0 41.50 arate cg			
fumarate per dose (equivalent to 200 mcg budesonide of mcg eformoterol fumarate metered dose) – Up to 120 dose available on a PSO	0 41.50 arate cg 82.50	120 dose OP		DuoResp Spiromax DuoResp Spiromax
fumarate per dose (equivalent to 200 mcg budesonide of mcg eformoterol fumarate metered dose) — Up to 120 dose available on a PSO	0 41.50 arate cg 82.50 - Up	120 dose OP	/	DuoResp Spiromax
fumarate per dose (equivalent to 200 mcg budesonide of mcg eformoterol fumarate metered dose) – Up to 120 dose available on a PSO	0 41.50 arate cg 82.50 - Up		/	
fumarate per dose (equivalent to 200 mcg budesonide of mcg eformoterol fumarate metered dose) — Up to 120 dose available on a PSO	041.50 arate cg82.50 - Up18.23	120 dose OP	/	DuoResp Spiromax
fumarate per dose (equivalent to 200 mcg budesonide of mcg eformoterol fumarate metered dose) — Up to 120 dose available on a PSO	041.50 arate cg82.50 - Up18.23 mcg	120 dose OP	<i>y</i>	DuoResp Spiromax
fumarate per dose (equivalent to 200 mcg budesonide of mcg eformoterol fumarate metered dose) — Up to 120 dose available on a PSO	041.50 arate cg82.50 - Up18.23 mcg	120 dose OP	<i>y</i>	DuoResp Spiromax Vannair Symbicort
fumarate per dose (equivalent to 200 mcg budesonide of mcg eformoterol fumarate metered dose) — Up to 120 dose available on a PSO	041.50 arate cg82.50 - Up18.23 mcg33.74	120 dose OP	<i>y</i>	DuoResp Spiromax
fumarate per dose (equivalent to 200 mcg budesonide of mcg eformoterol fumarate metered dose) — Up to 120 dose available on a PSO	041.50 arate cg82.50 - Up18.23 mcg33.74 Up	120 dose OP 120 dose OP 120 dose OP	/ /	DuoResp Spiromax Vannair Symbicort Turbuhaler 100/6
6 mcg eformoterol fumarate metered dose) – Up to 120 dose available on a PSO	041.50 arate cg82.50 - Up18.23 mcg33.74 - Up21.40	120 dose OP	/ /	DuoResp Spiromax Vannair Symbicort
fumarate per dose (equivalent to 200 mcg budesonide of mcg eformoterol fumarate metered dose) — Up to 120 dose available on a PSO	0	120 dose OP 120 dose OP 120 dose OP 120 dose OP	/ /	DuoResp Spiromax Vannair Symbicort Turbuhaler 100/6 Vannair
fumarate per dose (equivalent to 200 mcg budesonide of mcg eformoterol fumarate metered dose) — Up to 120 dose available on a PSO	0	120 dose OP 120 dose OP 120 dose OP	/ /	DuoResp Spiromax Vannair Symbicort Turbuhaler 100/6 Vannair Symbicort
fumarate per dose (equivalent to 200 mcg budesonide of mcg eformoterol fumarate metered dose) — Up to 120 dose available on a PSO	0	120 dose OP 120 dose OP 120 dose OP 120 dose OP	/ /	DuoResp Spiromax Vannair Symbicort Turbuhaler 100/6 Vannair
fumarate per dose (equivalent to 200 mcg budesonide of mcg eformoterol fumarate metered dose) — Up to 120 dose available on a PSO	0	120 dose OP 120 dose OP 120 dose OP 120 dose OP	/ /	DuoResp Spiromax Vannair Symbicort Turbuhaler 100/6 Vannair Symbicort
fumarate per dose (equivalent to 200 mcg budesonide of mcg eformoterol fumarate metered dose) — Up to 120 dose available on a PSO	0	120 dose OP 120 dose OP 120 dose OP 120 dose OP	\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	DuoResp Spiromax Vannair Symbicort Turbuhaler 100/6 Vannair Symbicort

	Subsidy		Fully Brand or
	(Manufacturer's I	Price) Subsi Per	idised Generic Manufacturer
	Φ	rei	Wanuacturer
FLUTICASONE FUROATE WITH VILANTEROL Rounday for inhalation 100 mag with vilanteral 25 mag	44.00	30 dose OP	✓ Breo Ellipta
Powder for inhalation 100 mcg with vilanterol 25 mcg	44.00	30 dose OF	▼ breo Empla
FLUTICASONE WITH SALMETEROL Aerosol inhaler 50 mcg with salmeterol 25 mcg	25.70	120 dose OP	✓ Seretide
Aerosol inhaler 35 mcg with salmeterol 25 mcg		120 dose OP	✓ Seretide
Powder for inhalation 100 mcg with salmeterol 50 mcg – No		.20 0000 0.	00.0
more than 2 dose per day	33.74	60 dose OP	Seretide Accuhaler
Powder for inhalation 250 mcg with salmeterol 50 mcg - No			
more than 2 dose per day	44.08	60 dose OP	Seretide Accuhaler
Beta-Adrenoceptor Agonists			
Deta-Autenoceptor Agonists			
SALBUTAMOL			
Oral liq 400 mcg per ml		150 ml	✓ <u>Ventolin</u>
Infusion 1 mg per ml, 5 ml Inj 500 mcg per ml, 1 ml – Up to 5 inj available on a PSO		10 5	✓ Ventolin✓ Ventolin
ing 500 mag per mi, 1 mil – op to 5 mg available on a 1 50	100.00	J	Ventonii
Inhaled Beta-Adrenoceptor Agonists			
SALBUTAMOL			
Aerosol inhaler, 100 mcg per dose CFC free – Up to 1000			
dose available on a PSO	4.18	200 dose OP	✓ SalAir
	(7.45)		Ventolin
Nebuliser soln, 1 mg per ml, 2.5 ml ampoule - Up to 30 neb			_
available on a PSO	8.96	20	✓ Asthalin
N. I			✓ UK Cipla S29
Nebuliser soln, 2 mg per ml, 2.5 ml ampoule – Up to 30 neb available on a PSO	0.42	20	✓ Asthalin
available on a F30	9.40	20	✓ UK Cipla S29
TERBUTALINE SULPHATE			• Olt Olpiu •
Powder for inhalation, 200 mcg per dose (equivalent to			
250 mcg metered dose), breath activated	22.20	120 dose OP	✓ Bricanyl Turbuhaler
			•
Anticholinergic Agents			
IPRATROPIUM BROMIDE			
Aerosol inhaler, 20 mcg per dose CFC-free - Up to 400 dose	е		
available on a PSO		200 dose OP	✓ Atrovent
Nebuliser soln, 250 mcg per ml, 2 ml ampoule - Up to 40 ne			
available on a PSO	11./3	20	✓ Accord S29 ✓ Univent
			- Offiverit
Inhaled Beta-Adrenoceptor Agonists with Antic	holinergic A	gents	
SALBUTAMOL WITH IPRATROPIUM BROMIDE			
Aerosol inhaler, 100 mcg with ipratropium bromide, 20 mcg p	er		
dose CFC-free		200 dose OP	✓ Duolin HFA
Nebuliser soln, 2.5 mg with ipratropium bromide 0.5 mg per			
vial, 2.5 ml ampoule - Up to 20 neb available on a PSO	11.04	20	✓ Duolin

RESPIRATORY SYSTEM AND ALLERGIES				
	Subsidy (Manufacturer's P \$	rice) Per	Fully Subsidised	Brand or Generic Manufacturer
Long-Acting Muscarinic Antagonists				
GLYCOPYRRONIUM – Subsidy by endorsement a) Inhaled glycopyrronium treatment will not be subsidised if umeclidinium. b) Glycopyrronium powder for inhalation 50 mcg per dose is a having COPD using spirometry if spirometry is possible, ar Powder for inhalation 50 mcg per dose	subsidised only	for patien	ts who have	been diagnosed as
 TIOTROPIUM BROMIDE – Subsidy by endorsement a) Tiotropium treatment will not be subsidised if patient is also umeclidinium. b) Tiotropium bromide is subsidised only for patients who have spirometry is possible, and the prescription is endorsed act 1 October 2018 with a valid Special Authority are deemed Powder for inhalation, 18 mcg per dose	ve been diagnos cordingly. Patie endorsed. 50.37	sed as hav	ving COPD unad tiotropiuse ✓ S	using spirometry if
UMECLIDINIUM — Subsidy by endorsement a) Umeclidinium will not be subsidised if patient is also receive tiotropium bromide. b) Umeclidinium powder for inhalation 62.5 mcg per dose is see COPD using spirometry if spirometry is possible, and the period powder for inhalation 62.5 mcg per dose	subsidised only prescription is er	for patient ndorsed a 30 dose	s who have ccordingly.	been diagnosed as having
Long-Acting Muscarinic Antagonists with Long-	Acting Beta	-Adreno	oceptor A	gonists
Combination long acting muscarinic antagonist and long acting be treatment with a combination inhaled corticosteroid and long acting SA2554 Special Authority for Subsidy	g beta-2 agonis	t.	·	v
Initial application from any relevant practitioner. Approvals valid the following criteria: Both: 1 Patient has been stabilised on a long acting muscarinic ant 2 The prescriber considers that the patient would receive add	tagonist; and			
GLYCOPYRRONIUM WITH INDACATEROL – Special Authority s Powder for Inhalation 50 mcg with indacaterol 110 mcg	see SA2554 abo		ail pharmacy	•
TIOTROPIUM BROMIDE WITH OLODATEROL – Special Author Soln for inhalation 2.5 mcg with olodaterol 2.5 mcg	ity see SA2554 81.00	60 dose	OP 🗸 🗸 S	acy piolto Respimat
UMECLIDINIUM WITH VILANTEROL – Special Authority see SA Powder for inhalation 62.5 mcg with vilanterol 25 mcg		etail phar 30 dose		noro Ellipta
Inhaled Corticosteroid with Long-Acting Muscar	inic Antago	nist and	d Beta Ag	onist

innaled Corticosteroid with Long-Acting Muscarinic Antagonist and Beta Agonist

BUDESONIDE WITH GLYCOPYRRONIUM AND EFORMOTEROL – Special Authority see SA2421 on the next page – Retail pharmacy

Aerosol inhaler budesonide 160 mcg with glycopyrronium

7.2 mcg and formoterol 5 mcg per dose.......79.15 120 dose OP ✓ Breztri Aerosphere

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

(Manufacturer's Pri

⇒SA2421 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has a diagnosis of COPD confirmed by spirometry or spirometry has been attempted and technically acceptable results are not possible; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient is currently receiving an inhaled corticosteroid with long acting beta-2 agonist (ICS/LABA) or a long acting muscarinic antagonist with long acting beta-2 agonist (LAMA/LABA); and
 - 2.1.2 Any of the following:

Clinical criteria:

- 2.1.2.1 Patient has a COPD Assessment Test (CAT) score greater than 10; or
- 2.1.2.2 Patient has had 2 or more exacerbations in the previous 12 months; or
- 2.1.2.3 Patient has had one exacerbation requiring hospitalisation in the previous 12 months; or
- 2.1.2.4 Patient has had an eosinophil count greater than or equal to 0.3×10^9 cells/L in the previous 12 months; or
- 2.2 Patient is currently receiving multiple inhaler triple therapy (inhaled corticosteroid with long-acting muscarinic antagonist and long-acting beta-2 agonist ICS/LAMA/LABA) and met at least one of the clinical criteria above prior to commencing multiple inhaler therapy.

FLUTICASONE FUROATE WITH UMECLIDINIUM AND VILANTEROL – Special Authority see SA2326 below – Retail pharmacy Powder for inhalation fluticasone furoate 100 mcg with

umeclidinium 62.5 mcg and vilanterol 25 mcg......104.24 30 dose OP ✓ Trelegy Ellipta

⇒SA2326 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has a diagnosis of COPD confirmed by spirometry or spirometry has been attempted and technically acceptable results are not possible; and
- 2 Fither:
 - 2.1 Both:
 - 2.1.1 Patient is currently receiving an inhaled corticosteroid with long acting beta-2 agonist (ICS/LABA) or a long acting muscarinic antagonist with long acting beta-2 agonist (LAMA/LABA); and
 - 2.1.2 Any of the following:

Clinical criteria:

- 2.1.2.1 Patient has a COPD Assessment Test (CAT) score greater than 10; or
- 2.1.2.2 Patient has had 2 or more exacerbations in the previous 12 months; or
- 2.1.2.3 Patient has had one exacerbation requiring hospitalisation in the previous 12 months; or
- 2.1.2.4 Patient has had an eosinophil count greater than or equal to 0.3×10^9 cells/L in the previous 12 months; or
- 2.2 Patient is currently receiving multiple inhaler triple therapy (inhaled corticosteroid with long acting muscarinic antagonist and long acting beta-2 agonist ICS/LAMA/LABA) and met at least one of the clinical criteria above prior to commencing multiple inhaler triple therapy.

Antifibrotics

NINTEDANIB – Special Authority see SA2012 on the next page – Retail pharmacy

Troto: Trintodariib fiot odbordiood iii oombiildaron Wari od	ooiaiooa piiroriiaorio.		
Cap 100 mg	2,554.00	60 OP	Ofev
Cap 150 mg	3,870.00	60 OP	Ofev

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
. •	Por 🗸	Manufacturer

⇒SA2012 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Nintedanib is to be discontinued at disease progression (See Note); and
- 4 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 5 Any of the following:
 - 5.1 The patient has not previously received treatment with pirfenidone; or
 - 5.2 Patient has previously received pirfenidone, but discontinued pirfenidone within 12 weeks due to intolerance; or
 - 5.3 Patient has previously received pirfenidone, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with pirfenidone).

Renewal — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 3 Nintedanib is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

PIRFENIDONE - Retail pharmacy-Specialist - Special Authority see SA2013 below

Note: Pirfenidone is not subsidised in combination with subsidised nintedanib.

Tab 801 mg3,6	45.00 9	0 OP 💆	Esbriet
Tab 267 mg	15.00	90	Esbriet

⇒SA2013 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note); and
- 4 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 5 Any of the following:
 - 5.1 The patient has not previously received treatment with nintedanib; or
 - 5.2 Patient has previously received nintedanib, but discontinued nintedanib within 12 weeks due to intolerance; or
 - 5.3 Patient has previously received nintedanib, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with nintedanib).

Renewal — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

	Subsidy (Manufacturer's Price) \$	Sul Per	Fully bsidised	Brand or Generic Manufacturer
Leukotriene Receptor Antagonists				
MONTELUKAST * Tab 4 mg * Tab 5 mg * Tab 10 mg	3.10	28 28 28	✓ <u>M</u>	ontelukast Viatris ontelukast Viatris ontelukast Viatris
Methylxanthines				
AMINOPHYLLINE * Inj 25 mg per ml, 10 ml ampoule – Up to 5 inj available on a PSO		5	✓ DI	BL Aminophylline
THEOPHYLLINE * Tab long-acting 250 mg * Oral liq 80 mg per 15 ml		100 500 ml	✓ Ni ✓ Ni	uelin-SR uelin
Mucolytics				
DORNASE ALFA – Special Authority see SA1978 below – Retain Nebuliser soln, 2.5 mg per 2.5 ml ampoule		6	✓ Pu	ulmozyme
⇒SA1978 Special Authority for Subsidy Initial application — (cystic fibrosis) only from a respiratory pl	hysician or paediatrici	ian. App	rovals val	lid for 12 months for

- 1 Patient has a confirmed diagnosis of cystic fibrosis; and
- 2 Patient has previously undergone a trial with, or is currently being treated with, hypertonic saline; and
- 3 Any of the following:

All of the following:

applications meeting the following criteria:

- 3.1 Patient has required one or more hospital inpatient respiratory admissions in the previous 12 month period; or
- 3.2 Patient has had 3 exacerbations due to CF, requiring oral or intravenous (IV) antibiotics in the previous 12 month period: or
- 3.3 Patient has had 1 exacerbation due to CF, requiring oral or IV antibiotics in the previous 12 month period and a Brasfield score of < 22/25; or</p>
- 3.4 Patient has a diagnosis of allergic bronchopulmonary aspergillosis (ABPA).

Renewal — (cystic fibrosis) only from a respiratory physician or paediatrician. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient continues to benefit from treatment.

ELEXACAFTOR WITH TEZACAFTOR, IVACAFTOR AND IVACAFTOR - PCT only - Special Authority see SA2456 below

Tab elexacaftor 50 mg with tezacaftor 25 mg, ivacaftor 37.5 mg

(56) and ivacaftor 75 mg (28)27,647.39 84 O

84 OP **✓ Trikafta**

Tab elexacaftor 100 mg with tezacaftor 50 mg, ivacaftor 75 mg

⇒SA2456 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with cystic fibrosis; and
- 2 Patient is 6 years of age or older; and
- 3 Either:

Subsidy		Fully	Brand or	
(Manufacturer's Pri	ce)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 3.1 Patient has two cystic fibrosis-causing mutations in the cystic fibrosis transmembrane regulator (CFTR) gene (one from each parental allele); or
- 3.2 Patient has a sweat chloride value of at least 60 mmol/L by quantitative pilocarpine iontophoresis or by Macroduct sweat collection system; and
- 4 Fither:
 - 4.1 Patient has a heterozygous or homozygous F508del mutation; or
 - 4.2 Patient has a G551D mutation or other mutation responsive in vitro to elexacaftor/tezacaftor/ivacaftor (see note a); and
- 5 The treatment must be the sole funded CFTR modulator therapy for this condition; and
- 6 Treatment with elexacaftor/tezacaftor/ivacaftor must be given concomitantly with standard therapy for this condition.

Notes:

 a) Eligible mutations are listed in the Food and Drug Administration (FDA) Trikafta prescribing information https://nctr-crs.fda.gov/fdalabel/services/spl/set-ids/f354423a-85c2-41c3-a9db-0f3aee135d8d/spl-doc

IVACAFTOR - PCT only - Specialist - Special Author	ority see SA2017 below		
Tab 150 mg	29,386.00	56	✓ Kalydeco
Oral granules 50 mg, sachet	29,386.00	56	✓ Kalydeco
Oral granules 75 mg, sachet	29,386.00	56	✓ Kalydeco

⇒SA2017 Special Authority for Subsidy

Initial application only from a respiratory specialist or paediatrician. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with cystic fibrosis; and
- 2 Either:
 - 2.1 Patient must have G551D mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene on at least 1 allele; or
 - 2.2 Patient must have other gating (class III) mutation (G1244E, G1349D, G178R, G551S, S1251N, S1255P, S549N and S549R) in the CFTR gene on at least 1 allele; and
- 3 Patients must have a sweat chloride value of at least 60 mmol/L by quantitative pilocarpine iontophoresis or by Macroduct sweat collection system; and
- 4 Treatment with ivacaftor must be given concomitantly with standard therapy for this condition; and
- 5 Patient must not have an acute upper or lower respiratory infection, pulmonary exacerbation, or changes in therapy (including antibiotics) for pulmonary disease in the last 4 weeks prior to commencing treatment with ivacaftor; and
- 6 The dose of ivacaftor will not exceed one tablet or one sachet twice daily; and
- 7 Applicant has experience and expertise in the management of cystic fibrosis.

SODIUM CHLORIDE

Not funded for use as a nasal drop.

Nasal Preparations

Allergy Prophylactics

FLUTICASONE PROPIONATE Metered aqueous nasal spray, 50 mcg per dose2.57	120 dose OP	✓ Flixonase Hayfever & Allergy
Metered aqueous nasal spray, 50 mcg per dose		✓ <u>SteroClear</u>
BUDESONIDE Material agreeing pagel apray, 50 mag par dage.	200 doos OB	✓ SteroClear

25 ml OP

✓ Biomed

	Subsidy	` .	Fully	Brand or
	(Manufacturer's Pric	ce) Subs Per	idised •	Generic Manufacturer
IPRATROPIUM BROMIDE	Ψ	1 01		Manufacturor
Aqueous nasal spray, 0.03%	5.22	15 ml OP	/ 11	Inivent
Aqueous nasar spray, 0.00 /6		13 1111 01	• 0	illivelit
Respiratory Devices				
MASK FOR SPACER DEVICE				
a) Up to 50 dev available on a PSO				
b) Only on a PSO				
c) Only for children aged six years and under	0.70		,	
Small	2.70	1	∨ e	-chamber Mask
PEAK FLOW METER				
a) Up to 25 dev available on a PSO				
b) Only on a PSO	0.54			U-1 W-1-1- AFO
Low range	9.54	1	• IV	lini-Wright AFS Low Range
Normal range	0.54	1	✓ N	lini-Wright
Normal range		'	- 14	Standard
SPACER DEVICE				
a) Up to 50 dev available on a PSO				
b) Only on a PSO				
220 ml (single patient)	3.65	1	√ e	-chamber Turbo
510 ml (single patient)		1	√ e	-chamber La
				Grande
800 ml	6.50	1	✓ V	olumatic
Description Office lands				
Respiratory Stimulants				
CAFFEINE CITRATE				

Oral liq 20 mg per ml (10 mg base per ml)......16.91

	Subsidy		Fully Brand or
	(Manufacturer's F \$	rice) Subs Per	idised Generic Manufacturer
			manadatat
Ear Preparations			
FLUMETASONE PIVALATE			
Ear drops 0.02% with clioquinol 1%	4.46	7.5 ml OP	✓ Locacorten-Viaform
			ED's
			✓ Locorten-Vioform
TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYC	IN AND NYSTAT	ΓIN	
Ear drops 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 mg and gramicidin 250 mcg per g	5 16	7.5 ml OP	✓ Kenacomb
2.5 mg and gramiolain 200 mag par g		7.0 1111 01	Rendomb
Ear/Eye Preparations			
DEXAMETHASONE WITH FRAMYCETIN AND GRAMICIDIN			
Ear/Eye drops 500 mcg with framycetin sulphate 5 mg and			
gramicidin 50 mcg per ml	4.50	8 ml OP	
	(9.27)		Sofradex
FRAMYCETIN SULPHATE			
Ear/Eye drops 0.5%		8 ml OP	Cofromusin
	(8.65)		Soframycin
Eye Preparations			
Eye preparations are only funded for use in the eye, unless expli	citly stated other	wise.	
Anti-Infective Preparations			
·			
ACICLOVIR * Eye oint 3%	15 90	4.5 g OP	✓ ViruPOS
CHLORAMPHENICOL	15.69	4.5 g OF	VIIUPO3
Eye oint 1%	1.55	5 g OP	✓ Devatis
Devatis to be Principal Supply on 1 February 2026		0 9 01	5074110
Eye drops 0.5%		10 ml OP	✓ Chlorsig
A Provided Course to the court to discrete a conduct of the	1.84	al to alternations	✓ Chlorafast
 a) Funded for use in the ear*. Indications marked with b) Chlorafast to be Principal Supply on 1 March 2026 	are unapproved	a indications.	
(Chlorsig Eye drops 0.5% to be delisted 1 March 2026)			
CIPROFLOXACIN			
Eye drops 0.3% - Subsidy by endorsement	10.85	5 ml OP	✓ Ciprofloxacin Teva
When prescribed for the treatment of bacterial keratitis			
for the second line treatment of chronic suppurative otiti Note: Indication marked with a * is an unapproved indic		*; and the preso	cription is endorsed accordingly.
• •	alion.		
SODIUM FUSIDATE [FUSIDIC ACID] Eye drops 1%	5 29	5 g OP	✓ Fucithalmic
_yo diopo 1 /o		0 9 01	✓ Fucithalmic Canada
			(ON) S29
			✓ Fucithalmic
			Spain S29
TOBRAMYCIN			.
Eye oint 0.3%		3.5 g OP 5 ml OP	✓ Tobrex✓ Tobrex
Eye drops 0.3%	11.48	5 IIII OP	▼ TODIEX

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	Subsidised	Generic	
\$	Per	1	Manufacturer	

Corticosteroids and Other Anti-Inflammatory Preparations

DE	XAMETHASONE			
*	Eye oint 0.1%	5.86	3.5 g OP	Maxidex
*	Eye drops 0.1%	4.50	5 ml OP	Maxidex
	Ocular implant 700 mcg - Special Authority see SA1680 below			
	- Retail pharmacy1,4	44.50	1	Ozurdex

⇒SA1680 Special Authority for Subsidy

Initial application — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has diabetic macular oedema with pseudophakic lens: and
- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- 3 Fither
 - 3.1 Patient's disease has progressed despite 3 injections with bevacizumab; or
 - 3.2 Patient is unsuitable or contraindicated to treatment with anti-VEGF agents; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Initial application — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has diabetic macular oedema; and
- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- 3 Patient is of child bearing potential and has not yet completed a family; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Patient is of child bearing potential and has not yet completed a family; and
- 3 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

DEXAMETHASONE WITH NEOMYCIN SULPHATE AND POLYMYXIN B SULPHATE

* Eve sint 0.19/ with reconvoir authors 0.259/ and polymorin b

sulphate 6,000 u per q	' ' '	3.5 g OP	✓ Maxitrol
* Eye drops 0.1% with neomycin sulphate 0.35%		0.0 g Oi	· maximor
b sulphate 6,000 u per ml	, , ,	5 ml OP	✓ Maxitrol
DICLOFENAC SODIUM			
Eye drops 0.1%, single dose	1.85	10 dose	✓ Diclofenac Devatis
	5.54	30 dose	✓ Diclofenac Devatis
FLUOROMETHOLONE			
* Eye drops 0.1%	3.09	5 ml OP	✓ FML
•	5.20		✓ Flucon

	Subsidy (Manufacturer's P	rice) Subs	Fully sidised	Brand or Generic Manufacturer
LEVOCABASTINE				
Eye drops 0.5 mg per ml	8.71 (10.34)	4 ml OP	L	ivostin
LODOXAMIDE				
Eye drops 0.1%	8.71	10 ml OP	✓ L	omide
PREDNISOLONE ACETATE	0.00	10 ml OD		huaduiaalama AFT
Eye drops 1%	7.00	10 ml OP 5 ml OP		rednisolone-AFT red Forte
PREDNISOLONE SODIUM PHOSPHATE - Special Authority		-		
Eye drops 0.5%, single dose (preservative free)		20 dose		linims Prednisolone
➤ SA1715 Special Authority for Subsidy nitial application only from an ophthalmologist or optometrist. ollowing criteria: Both:	Approvals valid fo	or 6 months for	r applica	ations meeting the
Patient has severe inflammation; and Patient has a confirmed allergic reaction to preservative Renewal from any relevant practitioner. Approvals valid for 6 n		reatment rema	ins app	propriate and the patient
penefiting from treatment.				
SODIUM CROMOGLICATE				
Eye drops 2%Allerfix to be Principal Supply on 1 March 2026	2.91	10 ml OP	✓ A	Allerfix
Glaucoma Preparations - Beta Blockers				
ΓΙΜΟLOL * Eye drops 0.25%	2 42	5 ml OP	√ ∆	arrow-Timolol
* Eye drops 0.5%		5 ml OP	_	rrow-Timolol
Glaucoma Preparations - Carbonic Anhydrase	Inhibitors			
ACETAZOLAMIDE				
* Tab 250 mg	13.96	100	✓ <u>N</u>	<u>ledsurge</u>
BRINZOLAMIDE	E 11	E ml OD	./ ^	
* Eye drops 1%		5 ml OP	V A	zopt
* Eye drops 2% with timolol 0.5%	3.58	5 ml OP	✓ <u>D</u>	<u>Oortimopt</u>
Glaucoma Preparations - Prostaglandin Analo	gues			
BIMATOPROST				
★ Eye drops 0.03%	5.15	3 ml OP	√ L	<u>umigan</u>
ATANOPROST				
★ Eye drops 0.005%	2.08	2.5 ml OP	√ <u>T</u>	eva
FRAVOPROST * Eye drops 0.004%	6.00	0 E ml OD	./ +	royeten
· ·	0.80	2.5 ml OP	<u> </u>	<u>ravatan</u>
Glaucoma Preparations - Other				
BRIMONIDINE TARTRATE ★ Eye drops 0.2%	5.16	5 ml OP	✓ A	arrow-Brimonidine
•			_	

	Subsidy (Manufacturer's F \$	Price) Subs Per	Fully Brand or idised Generic Manufacturer
BRIMONIDINE TARTRATE WITH TIMOLOL MALEATE	Ψ	1 61	• Manuacturer
* Eye drops 0.2% with timolol maleate 0.5%	7.13	5 ml OP	✓ Combigan
LATANOPROST WITH TIMOLOL			
* Eye drops 0.005% with timolol 0.5%	4.95	2.5 ml OP	✓ Arrow - Lattim
PILOCARPINE HYDROCHLORIDE			
* Eye drops 1%	4.26	15 ml OP	✓ Isopto Carpine
* Eye drops 2%	5.35	15 ml OP	✓ Isopto Carpine
* Eye drops 4%	7.99	15 ml OP	✓ Isopto Carpine
Subsidised for oral use pursuant to the Standard Formu	lae.		
PILOCARPINE NITRATE			
* Eye drops 2% single dose – Special Authority see SA0895 below – Retail pharmacy	35.90	20 dose	✓ Minims Pilocarpine
⇒SA0895 Special Authority for Subsidy			

| ⇒SA0895 | Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient has to use an unpreserved solution due to an allergy to the preservative; or
- 2 Patient wears soft contact lenses.

Note: Minims for a general practice are considered to be "tools of trade" and are not approved as special authority items. Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Mydriatics and Cycloplegics ATROPINE SUI PHATE 15 ml OP ✓ Atropt CYCLOPENTOLATE HYDROCHLORIDE 15 ml OP Cyclogyl **TROPICAMIDE** * Eve drops 0.5% 20.52 ✓ Mydriacyl 15 ml OP ✓ Mydriacyl 15 ml OP **Preparations for Tear Deficiency** For acetylcysteine eye drops refer Standard Formulae, page 281 **HYPROMELLOSE** 15 ml OP ✓ Methopt HYPROMELLOSE WITH DEXTRAN 15 ml OP ✓ Poly-Tears

Preservative Free Ocular Lubricants

⇒SA2431 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Confirmed diagnosis by slit lamp or Schirmer test of severe secretory dry eye; and
- - 2.1 Patient is using eye drops more than four times daily on a regular basis; or
 - 2.2 Patient has had a confirmed allergic reaction to preservative in eye drop.

SENSORY ORGANS

	Subsidy (Manufacturer's Price) \$) Subs Per	Fully idised	Brand or Generic Manufacturer
POLYETHYLENE GLYCOL 400 AND PROPYLENE GLYCOL – pharmacy				
Eye drops 0.4% and propylene glycol 0.3%, 0.8 ml SODIUM HYALURONATE [HYALURONIC ACID] – Special Auti		30 In the previo		e – Retail pharmacy
Eye drops 1 mg per ml	armacy Procedures		riction	
Other Eye Preparations				
NAPHAZOLINE HYDROCHLORIDE * Eye drops 0.1%	5.65 1	5 ml OP	✓ <u>A</u>	<u>llbalon</u>
OLOPATADINE Eye drops 0.1% Olopatadine Teva to be Principal Supply on 1 March 20.		5 ml OP	✓ 0	Diopatadine Teva
PARAFFIN LIQUID WITH WOOL FAT * Eye oint 3% with wool fat 3% RETINOL PALMITATE	3.63	3.5 g OP	√ P	Poly-Visc

Eye oint 138 mcg per g......3.80

5 g OP

✓ VitA-POS

			VA	RIOUS
	Subsidy (Manufacturer's Price) \$		Fully Brand or dised Generic Manufactur	or.
	D	Per	Manulaciur	er
Various				
PHARMACY SERVICES				
* Brand switch fee	4.50	1 fee	✓ BSF Allegro ✓ BSF Estradio Mylan	
 a) May only be claimed once per patient. 				
b) The Pharmacode for BSF Allegron is 2715740 - see al		0.5		
c) The Pharmacode for BSF Estradiol TDP Mylan is 2717			/	
* Immunisation administration fee - flu		1 fee	✓ Immunisatio	
* Immunisation administration fee - other		1 fee	✓ Immunisatio	
* Immunisation co-administration fee - flu and shingles	0.00	1 fee	Immunisation and Shing	
* Paxlovid fee (Pharmacist initiated)	0.00	1 fee	✓ Paxlovid fee	
(BSF Allegron Brand switch fee to be delisted 1 February 2026)				
(BSF Estradiol TDP Mylan Brand switch fee to be delisted 1 March	n 2026)			
Agents Used in the Treatment of Poisonings				
Agents osca in the Treatment of Folsonings				
Antidotes				
ACETYLCYSTEINE				
Inj 200 mg per ml, 10 ml ampoule	42.99	10	✓ DBL Acetylc	ysteine
Inj 200 mg per ml, 10 ml vial	42.99	10	Hikma	
			Acetylcyst	eine S29
NALOVONE HVDDOCHI ODIDE				
NALOXONE HYDROCHLORIDE				
a) Up to 10 inj available on a PSO				
b) Only on a PSO	10.00	_	✓ DDI Nalawa	
* Inj 400 mcg per ml, 1 ml ampoule	13.29	5	✓ <u>DBL Naloxo</u> Hydrochlo	
			Hydrocino	iiue
Removal and Elimination				
CHARCOAL				
* Oral liq 50 g per 250 ml	59.85 25	50 ml OP	✓ Carbosorb-)	(
a) Up to 250 ml available on a PSOb) Only on a PSO				
DEFERASIROX – Special Authority see SA1492 below – Retail pl	harmacy			
Wastage claimable	патнасу			
Tab 125 mg dispersible	276.00	28	✓ Exjade	
Tab 250 mg dispersible		28	✓ Exjade	
Tab 500 mg dispersible		28	✓ Exjade	
⇒SA1492 Special Authority for Subsidy	•		•	
OATTOE Opcolar Authority for Cubolay				

Initial application only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; and
- 2 Deferasirox is to be given at a daily dose not exceeding 40 mg/kg/day; and



Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
 (Manufacturer's Frice)	Per	Jubsiuiseu ✓	Manufacturer

continued...

- 3 Any of the following:
 - 3.1 Treatment with maximum tolerated doses of deferiprone monotherapy or deferiprone and desferrioxamine combination therapy have proven ineffective as measured by serum ferritin levels. liver or cardiac MRI T2*: or
 - 3.2 Treatment with deferiprone has resulted in severe persistent vomiting or diarrhoea; or
 - 3.3 Treatment with deferiprone has resulted in arthritis: or
 - 3.4 Treatment with deferiprone is contraindicated due to a history of agranulocytosis (defined as an absolute neutrophil count (ANC) of < 0.5 cells per μL) or recurrent episodes (greater than 2 episodes) of moderate neutropenia (ANC 0.5 1.0 cells per μL).</p>

Renewal only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria: Fither:

- 1 For the first renewal following 2 years of therapy, the treatment has been tolerated and has resulted in clinical improvement in all three parameters namely serum ferritin, cardiac MRI T2* and liver MRI T2* levels; or
- 2 For subsequent renewals, the treatment has been tolerated and has resulted in clinical stability or continued improvement in all three parameters namely serum ferritin, cardiac MRI T2* and liver MRI T2* levels.

DEFERIPRONE - Special Authority see SA1480 below - F	Retail pharmacy		
Tab 500 mg	533.17	100	✓ Ferriprox
Oral liq 100 mg per 1 ml	266.59	250 ml OP	✓ Ferriprox

⇒SA1480 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Fither:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; or
- 2 The patient has been diagnosed with chronic iron overload due to acquired red cell aplasia.

DESFERRIOXAMINE MESILATE

* Inj 500 mg vial	332.88	10	 ✓ DBL Desferrioxamine Mesylate for Inj BP
SODIUM CALCIUM EDETATE			
* Inj 200 mg per ml, 5 ml	53.31	6	
	(156.71)		Calcium Disodium Versenate

Standard Formulae

Water

ACETYLCYSTEINE EYE DROPS Acetylcysteine inj 200 mg per ml, 10 ml	qs	PHENOBARBITONE SODIUM PAEDIATRIC ORAL mg per ml)	LIQUID (10
Suitable eye drop base	qs	Phenobarbitone Sodium	400 mg
CODEINE LINCTUS (3 mg per 5 ml) Codeine phosphate Glycerol Preservative Water CODEINE LINCTUS (15 mg per 5 ml) Codeine phosphate	60 mg 40 ml qs to 100 ml	Glycerol BP Water PILOCARPINE ORAL LIQUID Pilocarpine 4% eye drops Preservative Water (Preservative should be used if quantity supplied is	4 ml to 40 ml qs qs to 500 ml for more
Glycerol Preservative Water FOLINIC MOUTHWASH	40 ml qs to 100 ml	than 5 days.) SALIVA SUBSTITUTE FORMULA Methylcellulose Preservative Water	5 g qs to 500 ml
Calcium folinate 15 mg tab Preservative Water	1 tab qs to 500 ml	(Preservative should be used if quantity supplied is than 5 days. Maximum 500 ml per prescription.)	
(Preservative should be used if quantity supplied is than 5 days. Maximum 500 ml per prescription.)	for more	SODIUM CHLORIDE ORAL LIQUID Sodium chloride inj 23.4%, 20 ml	qs
METHYL HYDROXYBENZOATE 10% SOLUTION Methyl hydroxybenzoate Propylene glycol (Use 1 ml of the 10% solution per 100 ml of oral liqu	10 g to 100 ml	Water (Only funded if prescribed for treatment of hyponatra VANCOMYCIN ORAL SOLUTION (25 mg per ml) Vancomycin 500 mg injection	qs aemia) 5 vials
OMEPRAZOLE SUSPENSION Omeprazole capsules or powder Sodium bicarbonate powder BP Water	qs 8.4 g to 100 ml	Glycerin with sucrose suspension Water (Only funded if prescribed for treatment of Clostridiu following metronidazole failure)	37.5 ml to 100 ml
PHENOBARBITONE ORAL LIQUID Phenobarbitone Sodium Glycerol BP	1 g 70 ml		

to 100 ml

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

Subsidy (Manufacturer's Price) \$

Fully Subsidised Per

Brand or Generic Manufacturer

Extemporaneously Compounded Preparations and Galenicals

COLLODION FLEXIBLE Note: This product is no longer being manufactured by the support of the sup	oplier and will b	e delisted fron	n the Schedule at a date to be
determined. Collodion flexible	19.30	100 ml	✓ PSM
COMPOUND HYDROXYBENZOATE — Only in combination Only in extemporaneously compounded oral mixtures. Soln	36.00	100 ml	✓ Midwest
GLYCERIN WITH SODIUM SACCHARIN – Only in combination Suspension	38.00	473 ml	✓ Ora-Sweet SF
GLYCERIN WITH SUCROSE – Only in combination Suspension	38.00	473 ml	✓ Ora-Sweet
GLYCEROL * Liquid – Only in combination Only in extemporaneously compounded oral liquid prepara		500 ml	✓ healthE Glycerol BP
METHYL HYDROXYBENZOATE Powder	11.00	25 g	✓ Midwest
METHYLCELLULOSE Powder Suspension – Only in combination		100 g 473 ml	✓ MidWest✓ Ora-Plus
(MidWest Powder to be delisted 1 February 2028) METHYLCELLULOSE WITH GLYCERIN AND SODIUM SACCHAF Suspension	,	combination 473 ml	✓ Ora-Blend SF
METHYLCELLULOSE WITH GLYCERIN AND SUCROSE - Only Suspension		473 ml	✓ Ora-Blend
PHENOBARBITONE SODIUM Powder – Only in combination Only in children up to 12 years	125.00	10 g	✓ MidWest
PROPYLENE GLYCOL Only in extemporaneously compounded methyl hydroxybenzoa Lig		n. 500 ml	✓ Midwest
SODIUM BICARBONATE Powder BP - Only in combination	13.50	500 g	✓ Midwest
Only in extemporaneously compounded omeprazole and la SYRUP (PHARMACEUTICAL GRADE) – Only in combination Only in extemporaneously compounded oral liquid preparations	·	ispension.	
Liq		500 ml	✓ Midwest
WATER Tap – Only in combination	0.00	1 ml	✓ Tap water

Subsidy (Manufacturer's Price) \$

Subsidised Per

Fully

Brand or Generic Manufacturer

Nutrient Modules

Carbohydrate

⇒SA1930 Special Authority for Subsidy

Initial application — (Cystic fibrosis or kidney disease) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Fither:

- 1 cystic fibrosis; or
- 2 chronic kidney disease.

Initial application — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

1 cancer in children: or

Both:

- 2 cancers affecting alimentary tract where there are malabsorption problems in patients over the age of 20 years; or
- 3 faltering growth in an infant/child; or
- 4 bronchopulmonary dysplasia; or
- 5 premature and post premature infant; or
- 6 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified where the patient has inborn errors of metabolism. Renewal — (Cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

CARBOHYDRATE SUPPLEMENT - Special Authority see SA1930 above - Hospital pharmacy [HP3]

Carbohydrate And Fat

⇒SA1376 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

continued...

✓ fully subsidised 283



Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	1	Manufacturer

continued...

- 1 Infant or child aged four years or under; and
- 2 cystic fibrosis.

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 infant or child aged four years or under; and
- 2 Any of the following:
 - 2.1 cancer in children; or
 - 2.2 faltering growth; or
 - 2.3 bronchopulmonary dysplasia; or
 - 2.4 premature and post premature infants.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Fat

⇒SA2204 Special Authority for Subsidy

Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified where the patient has an inborn error of metabolism. Initial application — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 faltering growth in an infant/child; or
- 2 bronchopulmonary dysplasia; or
- 3 fat malabsorption; or
- 4 lymphangiectasia; or
- 5 short bowel syndrome: or
- 6 infants with necrotising enterocolitis; or
- 7 biliary atresia; or
- 8 for use in a ketogenic diet; or
- 9 chyle leak; or

Subsidy		Fully	Brand or	
(Manufacturer's Pri	ce)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 10 ascites: or
 - 11 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal — (Indications other than inborn errors of metabolism)** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT SUPPLEMENT - Special Authority see SA2204 on the previous page - Hospital pharmacy [HP3]

Emulsion (neutral)	15.38	200 ml OP	Calogen
Emulsion (strawberry)	15.38	200 ml OP	✓ Calogen
Oil	37.50	500 ml OP	✓ MCT oil (Nutricia)
MCT Emulsion, 250 ml	143.65		✓ Liquigen

Protein

⇒SA1524 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 protein losing enteropathy; or
- 2 high protein needs; or
- 3 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

PROTEIN SUPPLEMENT -	- Special Authority see SA1524 above - Hospital ph	narmacy [HP3]	
Powder	8.95	227 g OP	✓ Resource
			Beneprotein
	13.82	225 g OP	✓ Protifar

✓ fully subsidised 285

Subsidy (Manufacturer's Price) Fully Subsidised Brand or Generic Manufacturer

Oral and Enteral Feeds

Diabetic Products

⇒SA1095 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is a type I or and II diabetic who is suffering weight loss and malnutrition that requires nutritional support. Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

DIABETIC ENTERAL FEED 1KCAL/ML - Special Authority see SA1095 above - Hospital pharmacy [HP3]					
Liquid, 500 ml bottle	4.65	1 OP	✓ Glucerna Select		
DIABETIC ORAL FEED 1KCAL/ML - Special Authority see SA109	5 above – Hos	pital pharmac	y [HP3]		
Liquid (strawberry), 200 ml bottle	2.25	1 OP	✓ Diasip		
Liquid (vanilla), 200 ml bottle	2.10	1 OP	✓ Nutren Diabetes		
	2.25		✓ Diasip		

Fat Modified Products

⇒SA2205 Special Authority for Subsidy

Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified where the patient has an inborn error of metabolism. Initial application — (Indications other than errors of inborn metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Patient has a chyle leak; or
- 2 Modified as a modular feed, made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule, for adults.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Subsidy (Manufacturer's Price) \$ Fully Subsidised Per

Brand or Generic Manufacturer

Paediatric Products For Children Awaiting Liver Transplant

⇒SA1098 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) who requires a liver transplant.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL FEED 1KCAL/ML - Special Authority see SA1098 above - Hospital pharmacy [HP3]

Paediatric Products For Children With Chronic Renal Failure

⇒SA1099 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) with acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL FEED 1KCAL/ML - Special Authority see SA1099 above - Hospital pharmacy [HP3]

Paediatric Products

⇒SA1379 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Child is aged one to ten years; and
- 2 Any of the following:
 - 2.1 the child is being fed via a tube or a tube is to be inserted for the purposes of feeding; or
 - 2.2 any condition causing malabsorption; or
 - 2.3 faltering growth in an infant/child; or
 - 2.4 increased nutritional requirements; or
 - 2.5 the child is being transitioned from TPN or tube feeding to oral feeding.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for

continued...

✓ fully subsidised 287

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

PAEDIATRIC ENTERAL FEED 1.5KCAL/ML - Special Authority see SA Liquid, 500 ml bottle	•		- Hospital pharmacy [HP3] Nutrini Energy RTH
PAEDIATRIC ENTERAL FEED 1KCAL/ML - Special Authority see SA13 Liquid, 500 ml bottle		OP 🗸	Hospital pharmacy [HP3] Pediasure RTH Nutrini RTH
PAEDIATRIC ENTERAL FEED WITH FIBRE 1.5KCAL/ML - Special Autl pharmacy [HP3]	nority see SA1	1379 on the pr	revious page - Hospital
Liquid, 500 ml bottle	7.14 1	OP 🗸	Nutrini Energy Multi Fibre
PAEDIATRIC ORAL FEED 1.5KCAL/ML - Special Authority see SA1379	on the previo	us page – Ho	spital pharmacy [HP3]
Liquid (strawberry), 200 ml bottle			Fortini
Liquid (vanilla), 200 ml bottle	1.90 1	OP 🗸	Fortini
Liquid (vanilla), 500 ml bottle		OP 🗸	Pediasure Plus
PAEDIATRIC ORAL FEED 1KCAL/ML - Special Authority see SA1379 o	n the previous	page – Hosp	pital pharmacy [HP3]
Liquid (chocolate), 200 ml bottle			Pediasure
Liquid (strawberry), 200 ml bottle		OP 🗸	Pediasure
Liquid (vanilla), 200 ml bottle		OP 🗸	Pediasure
PAEDIATRIC ORAL FEED WITH FIBRE 1.5KCAL/ML - Special Authorit pharmacy (HP3)	y see SA1379	on the previo	us page – Hospital
Liquid (chocolate), 200 ml bottle	1.90 1	OP 🗸	Fortini Multi Fibre
Liquid (strawberry), 200 ml bottle		OP 🗸	Fortini Multi Fibre
Liquid (unflavoured), 200 ml bottle		OP 🗸	Fortini Multi Fibre
Liquid (vanilla), 200 ml bottle	1.90 1	OP 🗸	Fortini Multi Fibre
PEPTIDE-BASED ORAL FEED - Special Authority see SA1379 on the p	revious page	– Hospital pha	armacy [HP3]
Powder4			Peptamen Junior

Renal Products

⇒SA1101 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Brand or

Fully

	(Manufacturer's Price)	Subsic	lised	Generic
	\$	Per	1	Manufacturer
RENAL ORAL FEED 2 KCAL/ML - Special Authority see SA110	1 on the previous pa	ge – Hospita	al phai	rmacy [HP3]
Liquid, 200 ml bottle	13.24	4 OP	✓ N	ovaSource Renal
Liquid (apricot) 125 ml	13.72	4 OP	✓ R	enilon 7.5
Liquid (caramel) 125 ml	13.72	4 OP	✓ R	enilon 7.5

Subsidy

Specialised And Elemental Products

⇒SA1377 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 malabsorption: or
- 2 short bowel syndrome; or
- 3 enterocutaneous fistulas: or
- 4 eosinophilic oesophagitis; or
- 5 inflammatory bowel disease; or
- 6 patients with multiple food allergies requiring enteral feeding.

Notes: Each of these products is highly specialised and would be prescribed only by an expert for a specific disorder. The alternative is hospitalisation.

Elemental 028 Extra is more expensive than other products listed in this section and should only be used where the alternatives have been tried first and/or are unsuitable.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL SEMI-ELEMENTAL FEED 1.5KCAL/ML - S Liquid, 1,000 ml bottle		SA1377 abov 1 OP	ve – Hospital pharmacy [HP3] ✓ Vital
ORAL ELEMENTAL FEED 0.8KCAL/ML - Special Authority s	see SA1377 above –	Hospital pha	rmacy [HP3]
Liquid (grapefruit), 250 ml carton	179.46	18 OP	✓ Elemental 028 Extra
Liquid (pineapple & orange), 250 ml carton		18 OP	✓ Elemental 028 Extra
Liquid (summer fruits), 250 ml carton	179.46	18 OP	✓ Elemental 028 Extra
ORAL ELEMENTAL FEED 1KCAL/ML - Special Authority see	e SA1377 above – H	lospital pharm	nacy [HP3]
Powder (unflavoured), 80 g sachet	4.50	1 OP	✓ Vivonex TEN
SEMI-ELEMENTAL ENTERAL FEED 1KCAL/ML - Special A	uthority see SA1377	above – Hos	pital pharmacy [HP3]
Liquid, 500 ml bottle	7.47	1 OP	✓ Nutrison Advanced
·			Pentisorb

Paediatric Products For Children With Low Energy Requirements

⇒SA1196 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

continued...



Subs (Manufactur		
\$	Per	Manufacturer

- 1 Child aged one to eight years; and
- 2 The child has a low energy requirement but normal protein and micronutrient requirements.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

PAEDIATRIC ENTERAL FEED WITH FIBRE 0.76 KCAL/ML - Special Authority see SA1196 on the previous page - Hospital pharmacy [HP3]

Standard Supplements

⇒SA1859 Special Authority for Subsidy

Initial application — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age; and
- 2 Any of the following:
 - 2.1 The patient has a condition causing malabsorption; or
 - 2.2 The patient has failure to thrive; or
 - 2.3 The patient has increased nutritional requirements; and
- 3 Nutrition goal has been set (eg reach a specific weight or BMI).

Renewal — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 A nutrition goal has been set (eg reach a specific weight or BMI).

Initial application — (Children - exclusive enteral nutrition for Crohn's disease) only from a gastroenterologist or dietitian on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 Dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Renewal — (Children - exclusive enteral nutrition for Crohn's disease) from any relevant practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 General Practitioners and dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

continued...

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

Initial application — (Adults) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

1 Any of the following:

Patient is Malnourished

- 1.1 Patient has a body mass index (BMI) of less than 18.5 kg/m²; or
- 1.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 1.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months; and
- 2 Any of the following:

Patient has not responded to first-line dietary measures over a 4 week period by:

- 2.1 Increasing their food intake frequency (eg snacks between meals); or
- 2.2 Using high-energy foods (e.g. milkshakes, full fat milk, butter, cream, cheese, sugar etc); or
- 2.3 Using over the counter supplements (e.g. Complan); and
- 3 A nutrition goal has been set (e.g. to reach a specific weight or BMI).

Renewal — (Adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 A nutrition goal has been set (eg reach a specific weight or BMI); and
- 2 Any of the following:

Patient is Malnourished

- 2.1 Patient has a body mass index (BMI) of less than 18.5 kg/m²; or
- 2.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 2.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months.

Initial application — (Short-term medical condition) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube or a nasogastric tube is to be inserted for feeding; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Is undergoing a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
 - 5.1 Pregnant: and
 - 5.2 Any of the following:
 - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or
 - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
 - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

Renewal — (Short-term medical condition) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Has undergone a bone marrow transplant; or

continued...

(Manu	Subsidy ufacturer's Price)	Ful Subsidise	,	
<u> </u>	\$ F	Per •	Manufacturer	

- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
 - 5.1 Pregnant: and
 - 5.2 Any of the following:
 - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or
 - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
 - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

Initial application — (Long-term medical condition) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis: or
- 3 Liver disease: or
- 4 Chronic Renal failure: or
- 5 Inflammatory bowel disease; or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome: or
- 8 Bowel fistula: or
- 9 Severe chronic neurological conditions; or
- 10 Epidermolysis bullosa; or
- 11 AIDS (CD4 count < 200 cells/mm3); or
- 12 Chronic pancreatitis.

Renewal — (Chronic disease OR tube feeding for patients who have previously been funded under Special Authority forms SA0702 or SA0583) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis; or
- 3 Liver disease; or
- 4 Chronic Renal failure; or
- 5 Inflammatory bowel disease; or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome; or
- 8 Bowel fistula: or
- 9 Severe chronic neurological conditions.

ENTERAL FEED 1.5KCAL/ML - Special Authority see SA1859	on page 290 – Ho	spital pharma	ıcy [HP3]
Liquid, 1,000 ml bottle	8.68	1 OP	✓ Ensure Plus HN
• • •			RTH
	9.00		✓ Nutrison Energy
Liquid, 250 ml can	2.17	1 OP	✓ Ensure Plus HN
(Ensure Plus HN Liquid, 250 ml can to be delisted 1 March 202	26)		

	Subsidy (Manufacturer's F		Fully Brand or dised Generic ✓ Manufacturer
ENTERAL FEED 1KCAL/ML - Special Authority see SA1859 on	nage 200 - Hos	nital pharmacy [HD31
Liquid, 1,000 ml bottle		1 OP	✓ Osmolite RTH ✓ Nutrison RTH
ENTERAL FEED WITH FIBRE 0.83 KCAL/ML - Special Authorit Liquid, 1,000 ml bottle		n page 290 – Ho 1 OP	ospital pharmacy [HP3] Nutrison 800 Complete Multi Fibre
ENTERAL FEED WITH FIBRE 1 KCAL/ML - Special Authority so Liquid, 1,000 ml bottle		age 290 – Hospi 1 OP	ital pharmacy [HP3] ✓ Jevity RTH ✓ Nutrison Multi Fibre
ENTERAL FEED WITH FIBRE 1.5KCAL/ML - Special Authority Liquid, 1,000 ml bottle		page 290 – Hos 1 OP	pital pharmacy [HP3] ✓ Jevity HiCal RTH ✓ Nutrison Energy Multi Fibre
ORAL FEED (POWDER) – Special Authority see SA1859 on page Powder (chocolate)		l pharmacy [HP: 840 g OP	3] ✓ Sustagen Hospital Formula
	40.00	850 g OP	✓ Ensure
Powder (vanilla)		840 g OP	✓ Sustagen Hospital Formula
	40.00	850 g OP	✓ Ensure
ORAL FEED 1.5KCAL/ML — Special Authority see SA1859 on pa Additional subsidy by endorsement is available for patients be epidermolysis bullosa, or as exclusive enteral nutrition for the hypercapnia, defined as CO2 value exceeding 55mmHg. The Liquid (banana), 200 ml bottle — Higher subsidy of up to \$1.7 per 1 btl with Endorsement	eing bolus fed the treatment of Cr e prescription m	rough a feeding ohn's disease, o	tube, who have severe r for patients with COPD and
\$1.76 per 1 btl with Endorsement	0.72	1 OP	
	(1.56) (1.76)	1 01	Ensure Plus Fortisip
Liquid (fruit of the forest), 200 ml bottle – Higher subsidy of	0.70	4.00	
\$1.56 per 1 btl with Endorsement		1 OP	Casara Dive
1: :1/. 1	(1.56)		Ensure Plus
Liquid (strawberry), 200 ml bottle – Higher subsidy of \$1.76 p		1 OD	
1 btl with Endorsement		1 OP	Forticin
Limited (ventille) 2000 ml hottle. Higher subsider of the day	(1.76)		Fortisip
Liquid (vanilla), 200 ml bottle – Higher subsidy of up to \$1.76		1 OB	
per 1 btl with Endorsement	(1.56)	1 OP	Ensure Plus
	(1.76)		Fortisip
Liquid (vanilla) 227 ml can Higher subsidy of \$4.65 per	(1.70)		ι σιμοιμ
Liquid (vanilla), 237 ml can - Higher subsidy of \$1.65 per 1 can with Endorsement	0.05	1 OP	
ı can willi endusenieni		I OF	Ensure Plus
	(1.65)		Elisule Flus

✓ fully subsidised 293

(Ensure Plus Liquid (vanilla), 237 ml can to be delisted 1 July 2026)

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	sidised	Generic	
 \$	Per	•	Manufacturer	

ORAL FEED WITH FIBRE 1.5 KCAL/ML - Special Authority see SA1859 on page 290 - Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, or who have severe epidermolysis bullosa. The prescription must be endorsed accordingly.

Liquid (chocolate), 200 ml bottle - Higher subsidy of \$1.76 per

1 btl with Endorsement	0.70	1 OP	
I DU WIUI ENGOISEMENT	0.72	I OF	
	(1.76)		Fortisip Multi Fibre
Liquid (strawberry), 200 ml bottle - Higher subsidy of \$1.76 per			
1 btl with Endorsement	0.72	1 OP	
	(1.76)		Fortisip Multi Fibre
Liquid (vanilla), 200 ml bottle - Higher subsidy of \$1.76 per			
1 btl with Endorsement	0.72	1 OP	
	(1.76)		Fortisip Multi Fibre

High Calorie Products

⇒SA1195 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

All of the following:

- 1 Cystic fibrosis; and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements.

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 any condition causing malabsorption; or
 - 1.2 faltering growth in an infant/child; or
 - 1.3 increased nutritional requirements; or
 - 1.4 fluid restricted: and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements or is fluid restricted.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

	Subsidy (Manufacturer's Price \$) S	Fully Subsidised	Brand or Generic Manufacturer
ENTERAL FEED 2 KCAL/ML - Special Authority see SA1195 o	n the previous page	– Hospi	tal pharma	acy [HP3]
Liquid, 1,000 ml bottle	13.64	1 OP	✓ I	Ensure Two Cal HN RTH
Liquid, 500 ml bottle	6.82	1 OP	√ I	Nutrison Concentrated
ORAL FEED 2 KCAL/ML — Special Authority see SA1195 on the Additional subsidy by endorsement is available for patients I epidermolysis bullosa. The prescription must be endorsed a Liquid (vanilla), 200 ml bottle — Higher subsidy of \$2.34 per 1 btl with Endorsement	peing bolus fed throu accordingly.		eding tube	

Food Thickeners

⇒SA1106 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient has motor neurone disease with swallowing disorder.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FOOD THICKENER - Special Authority see SATTU6 abov	ve – Hospitai pnarmacy	[HP3]	
Powder	8.29	300 g OP	✓ Nutilis
	24.00	380 g OP	✓ Aptamil Feed
			Thickener

Gluten Free Foods

The funding of gluten free foods is no longer being actively managed by Pharmac from 1 April 2011. This means that we are no longer considering the listing of new products, or making subsidy, or other changes to the existing listings. As a result we anticipate that the range of funded items will reduce over time. Management of Coeliac disease with a gluten free diet is necessary for good outcomes. A range of gluten free options are available through retail outlets.

⇒SA1729 Special Authority for Subsidy

Initial application — (all patients) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Either:

- 1 Gluten enteropathy has been diagnosed by biopsy; or
- 2 Patient suffers from dermatitis herpetiformis.

Initial application — (paediatric patients diagnosed by ESPGHAN criteria) only from a paediatric gastroenterologist. Approvals valid without further renewal unless notified where the paediatric patient fulfils ESPGHAN criteria for biopsy free diagnosis of coeliac disease.

GLUTEN FREE BAKING MIX – Special Authority see SA	1729 above – Hospital	oharmacy [HP3]	
Powder	2.81	1,000 g OP	
	(5.15)		Healtheries Simple
			Baking Mix

	Subsidy (Manufacturer's Prio \$,	,	
GLUTEN FREE BREAD MIX – Special Authority see SA1729 on the previous page – Hospital pharmacy [HP3]				
Powder	3.93	1,000 g OP		
	(7.32)		NZB Low Gluten	
			Bread Mix	
	3.51			
	(10.87)		Horleys Bread Mix	
GLUTEN FREE FLOUR - Special Authority see SA1729 on the Powder		ospital pharmacy 2,000 g OP	y [HP3]	
	(18.10)	, 0	Horleys Flour	

Foods And Supplements For Inherited Metabolic Disease

⇒SA2357 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where patient requires dietary management of inherited metabolic disorders.

Supplements For Homocystinuria

AMINOACID FORMULA WITHOUT METHIONINE - Spec	cial Authority see SA235	7 above – Ho	spital pharmacy [HP3]	
Powder (neutral), 36 g sachets	750.30	30	✓ HCU Anamix Junior	
Powder, 12.5 g sachets	349.65	30	✓ HCU Explore 5	
Powder, 25 g sachets	1,048.95	30	✓ HCU Express 15	
Powder (neutral), can		500 g OP	✓ XMET Maxamum	
Powder (unflavoured), can	260.00	400 g OP	 HCU Anamix Infant 	
Liquid (juicy berries), 125 ml bottle	1,684.80	30	✓ HCU Lophlex LQ	
Liquid (orange), 125 ml bottle	941.40	36	 HCU Anamix Junior 	
			LO	

Supplements For MSUD and short chain enoyl coA hydratase deficiency

AMINOACID FORMULA WITHOUT VALINE, LEUCINE AND ISOLEUCINE - Special Authority see SA2357 above - Hospital pharmacy [HP3]

Powder (neutral) 36 g sachets	750.00	30	MSUD Anamix Junior
Powder, 12.5 g sachets	349.65	30	✓ MSUD Explore 5
Powder, 25 g sachets	.1,048.95	30	✓ MSUD Express 15
Powder (neutral), can	454.71	500 g OP	✓ MSUD Maxamum
Powder (orange), can	454.71	500 g OP	✓ MSUD Maxamum
Powder (unflavoured), can	260.00	400 g OP	MSUD Anamix Infant
Liquid (orange) 125 ml bottles	941.40	36	MSUD Anamix Junior LQ
Liquid (juicy berries) 125 ml pouches	.1,684.80	30	✓ MSUD Lophlex LQ 20

Subsidy	
(Manufacturer's Price)	Subs
•	Por

Fully Subsidised Brand or Generic Manufacturer

Supplements For PKU

AMINOACID FORMULA WITHOUT PHENYLALANINE - Special Authority see \$A2357 on the previous page - Hospital pharmacy [HP3]

Tabs	99.00	75 OP	✓ Phlexy 10
Powder (Lemon), 34 g sachets	883.50	30	✓ PKU Express 20
Powder (Neutral), 12.5 g sachets	220.88	30	✓ PKU Explore 5
Powder (Neutral), 34 g sachets		30	✓ PKU Express 20
Powder (Orange), 25 g sachets		30	✓ PKU Explore 10
Powder (Orange), 34 g sachets		30	✓ PKU Express 20
Powder (Raspberry), 25 g sachets		30	✓ PKU Explore 10
Powder (Tropical), 34 g sachets		30	✓ PKU Express 20
Powder (berry) 28 g sachets		30	✓ PKU Lophlex
			Powder
Powder (chocolate) 36 g sachet	393.00	30	✓ PKU Anamix Junior
, ,			Chocolate
Powder (neutral) 28 g sachets	936.00	30	✓ PKU Lophlex
, ,			Powder
Powder (neutral) 36 g sachets	393.00	30	✓ PKU Anamix Junior
Powder (orange) 28 g sachets	936.00	30	✓ PKU Lophlex
(3 / 3			Powder
Powder (orange) 36 g sachet	393.00	30	✓ PKU Anamix Junior
(0, 0			Orange
Powder (unflavoured) 12.5 g sachets	234.00	30	✓ PKU First Spoon
Powder (vanilla) 36 g sachet	393.00	30	✓ PKU Anamix Junior
, , ,			Vanilla
Infant formula	174.72	400 g OP	✓ PKU Anamix Infant
Powder (neutral), 4 × 400 g can	715.16	1,600 g OP	✓ Pku Start
Powder (orange)		500 g OP	✓ XP Maxamum
Powder (unflavoured)	320.00	500 g OP	✓ XP Maxamum
Liquid (berry), 125 ml bottle		1 ÖP	✓ PKU Anamix Junior
			LQ
Liquid (orange), 125 ml bottle	13.10	1 OP	✓ PKU Anamix Junior
			LQ
Liquid (forest berries), 250 ml carton	540.00	18 OP	✓ Easiphen Liquid
Liquid (juicy tropical) 125 ml	936.00	30 OP	✓ PKU Lophlex LQ 20
Oral semi-solid (berries) 109 g		36 OP	✓ PKU Lophlex
, ,	•		Sensation 20
Liquid (juicy berries) 62.5 ml	939.00	60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy berries) 125 ml		30 OP	✓ PKU Lophlex LQ 20
Liquid (juicy orange) 125 ml	936.00	30 OP	✓ PKU Lophlex LQ 20

	0.1.1.		T
	Subsidy (Manufacturer's Price)	Subc	Fully Brand or dised Generic
	(Manufacturer's Frice)	Per	✓ Manufacturer
GLYCOMACROPEPTIDE AND AMINO ACID CONTAINS SOME	PHENYI AI ANINE	– Snecial A	Authority see SA2357 on
page 296 – Hospital pharmacy [HP3]		Opcolai /	tunonty see GAZGG7 GH
Powder (Banana) 35 g sachets	930.00	30	✓ PKU
. 5.135. (24.14.14, 55 g 546.1516.11.11.11.11.11.11.11.11.11.11.11.11.1			sphere20 Banana
Powder (Berry), 20 g sachets	449 28	60	✓ PKU Restore
1 011d01 (15011), 15 g 5d011010		00	Powder
Powder (Chocolate) 32 g sachets	898 56	30	✓ PKU Build
1 Owder (Griocolate) 02 g dadriote		00	20 Chocolate
Powder (Chocolate) 35 g sachets	930.00	30	✓ PKU
Toward (Griodolate) of gradiote		00	sphere20 Chocolate
			opiiciozo onocolato
Powder (Lemon) 35 g sachets	930.00	30	✓ PKU
· · · ·			sphere20 Lemon
Powder (Lemonade) 33.4 g sachets	936.00	30	✓ PKU GMPro Ultra
, ,			Lemonade
Powder (Neutral), 15 g sachets	449.28	30	✓ PKU Build 10
Powder (Orange), 20 g sachets		60	✓ PKU Restore
· · · · · · · · · · · · · · · · · · ·			Powder
Powder (Raspberry Lemonade) 31 g sachets	898.56	30	✓ PKU Build
, , , , , , , , , , , , , , , , , , ,			20 Raspberry
			Lemonade
Powder (Smooth) 31 g sachets	898.56	30	✓ PKU Build
· · · · · · · · · · · · · · · · · · ·			20 Smooth
Powder (Vanilla) 33 g sachets	898.56	30	✓ PKU Build 20 Vanilla
Powder (neutral), 40 g sachets		30	✓ Glytactin Bettermilk
Powder (unflavoured) 12.5 g sachets		30	✓ PKU GMPro Mix-In
Powder (vanilla) 33.4 g sachets		30	✓ PKU GMPro Ultra
, ,			Vanilla
Powder (Red Berry) 35 g sachets	930.00	30	✓ PKU sphere20 Red
			Berry
Powder (Vanilla) 35 g sachets	930.00	30	✓ PKU
, ,			sphere20 Vanilla
Liquid (neutral), 250 ml carton	280.80	18	✓ PKU GMPro LQ
Liquid (original), 250 ml carton		30 OP	✓ PKU Glytactin RTD
, ,			15
Liquid (Coffee Mocha), 250 ml carton	684.45	30 OP	✓ PKU Glytactin RTD
			15 Lite
Liquid (chocolate), 250 ml carton	684.45	30 OP	✓ PKU Glytactin RTD
4 7		- · - ·	15
Liquid (vanilla), 250 ml carton	684.45	30 OP	✓ PKU Glytactin RTD
4 V			15 Lite

Foods

LOW PROTEIN BAKING MIX — Special Authority see SA2357 on page 296 — Hospital pharmacy [HP3]
Powder8.55 500 g OP ✓ Loprofin Mix

	Subsidy		Fully	Brand or
	(Manufacturer's Pri			
	\$	Per	_	Manufacturer
LOW PROTEIN PASTA - Special Authority see SA2357 on page	e <mark>296</mark> – Hospital p	harmacy [HP3]]	
Animal shapes	12.39	500 g OP		Loprofin
Lasagne	6.19	250 g OP	1	Loprofin
Low protein rice pasta	12.39	500 g OP	1	Loprofin
Macaroni	6.19	250 g OP	1	Loprofin
Penne	12.39	500 g OP	1	Loprofin
Spaghetti	12.39	500 g OP	1	Loprofin
Spirals	12.39	500 g OP	✓	Loprofin
Supplements for Tyrosinaemia				
AMINOACID FORMULA WITHOUT PHENYLALANINE AND TYPE	ROSINE - Specia	Authority see	SA2	2357 on page 296 – Hospita
pharmacy [HP3]				
Powder (Neutral), 12.5 g sachets	349.65	30	1	TYR Explore 5
Powder (neutral) 36 g sachets	471.00	30	1	TYR Anamix Junior
Powder, can	260.00	400 g OP	1	TYR Anamix Infant
Liquid (juicy berries) 125 ml pouches	1,684.80	30	1	TYR Lophlex LQ 20
Liquid (orange) 125 ml bottle		36		TYR Anamix Junior
				LQ
GLYCOMACROPEPTIDE AND AMINO ACID CONTAINS SOME	TYROSINE AND	PHENYLALAN	NINE	E – Special Authority see
SA2357 on page 296 – Hospital pharmacy [HP3]				
Powder (Red Berry), 35 g sachets	1,398.60	30	1	TYR Sphere 20
Powder (Vanilla), 35 g sachets	1,398.60	30	1	TYR Sphere 20
Supplements for Organic Acidaemias				
AMINOACID FORMULA WITHOUT ISOLEUCINE, METHIONINE	THREONINE AN	ND VALINE _	Sno	oial Authority can SA2357
on page 296 – Hospital pharmacy [HP3]	., ITIIILONINE AI	ND VALINE -	Ope	cial Authority 366 CA2007
Powder, can	260.00	400 g OP	./	MMA/PA Anamix
rowder, cari	200.00	400 g OF	•	Infant
				IIIIdIIL
AMINOACID FORMULA WITHOUT METHIONINE, THREONINE	AND VALINE - S	Special Authori	ity se	ee SA2357 on page 296 -
Hospital pharmacy [HP3]				
Powder (neutral), 18 g sachets	750.30	30	✓	MMA/PA Anamix
				Junior
Powder, 12.5 g sachets	349.65	30	1	MMA/PA Explore 5
Powder, 25 g sachets		30		MMA/PA Express 15
•	·			·
Supplements for Glutaric Aciduria type 1				
AMINOACID FORMULA WITHOUT LYSINE - Special Authority	see SA2357 on pa	age 296 - Hos	pital	pharmacy [HP3]
Powder (neutral), 18 g sachets	750.30	30	1	GA1 Anamix Junior
Powder, 12.5 g sachets	349.65	30	1	GA Explore 5
Powder, can		400 g OP		GA1 Anamix Infant
·		•		
Supplements for Glycogen Storage Disease				
HIGH AMYLOPECTIN CORN-STARCH - Special Authority see				
Powder, 60 g sachets	241.62	30	1	Glycosade
Single dose amino acids				
ARGININE - Special Authority see SA2357 on page 296 - Hosp	ital pharmacv [HP	31		
Powder, 4 g sachets		30	1	Arginine2000
· · · · · · · · · · · · · · · · · · ·				J

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
CITRULLINE – Special Authority see SA2357 on page 296 – Powder, 4 g sachets		3] 30	1	Citrulline1000
ISOLEUCINE – Special Authority see SA2357 on page 296 – Powder, 4 g sachets	141.05	3] 30	•	Isoleucine50
LEUCINE – Special Authority see SA2357 on page 296 – Hos Powder, 4 g sachets	141.05	30		Leucine100
PHENYLALANINE – Special Authority see SA2357 on page 2 Powder, 4 g sachets		y [HP3 30	-	Phenylalanine50
TYROSINE – Special Authority see SA2357 on page 296 – H Powder, 4 g sachets		30	•	Tyrosine1000
VALINE – Special Authority see SA2357 on page 296 – Hosp Powder, 4 g sachets		30	•	Valine50
Other Fat Modified Products				
ELEMENTAL FEED WITH HIGH MEDIUM CHAIN TRIGLYCE pharmacy [HP3]	RIDES - Special Author	ority se	ee SA2357	on page 296 – Hospital
Powder (neutral), 100 g sachets	47.01	10	•	Emsogen
Carbohydrate and Fat with added vitamins an	d minerals			
PROTEIN FREE SUPPLEMENT CONTAINING CARBOHYDF Authority see SA2357 on page 296 – Hospital pharmacy [HP3]			·
Powder (neutral), can	49.29 4	00 g C)P V	Energivit
Essential Amino Acids				
ESSENTIAL AMINOACID FORMULA - Special Authority see Powder (neutral), can		- Hosp 00 g C		acy [HP3] Essential Amino Acid Mix
Infant Formulae				
For Williams Syndrome				
■ SA1110 Special Authority for Subsidy Initial application only from a dietitian, relevant specialist or v year where the patient is an infant suffering from Williams Syn Renewal only from a dietitian, relevant specialist, vocationally recommendation of a dietitian, relevant specialist or vocational applications meeting the following criteria: Both:	drome and associated had registered general practly registered general pr	nyperc titione actitio	alcaemia. r or gener	al practitioner on the
 The treatment remains appropriate and the patient is be General Practitioners must include the name of the diet practitioner and date contacted. 	•		cationally	registered general

LOW CALCIUM INFANT FORMULA - Special Authority see SA1110 above - Hospital pharmacy [HP3]

Powder46.18

400 g OP

✓ Locasol

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsi	dised	Generic	
\$	Per	1	Manufacturer	

Gastrointestinal and Other Malabsorptive Problems

AMINO ACID FORMULA - Special Authority see SA2092 below - Hospital phan	macy [HP3]	
Powder43.60	400 g OP	✓ Alfamino✓ Alfamino Junior
Powder (unflavoured)55.61	400 g OP	✓ Neocate Gold ✓ Neocate Junior Unflavoured
65.72		✓ Neocate SYNEO✓ Elecare✓ Elecare LCP
Powder (vanilla)55.61	400 g OP	✓ Neocate Junior Vanilla
65.72		✓ Elecare

⇒SA2092 Special Authority for Subsidy

Initial application — (Infants under 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 History of anaphylaxis to cow's milk protein formula or dairy products; or
- 2 Eosinophilic oesophagitis; or
- 3 Ultra-short gut; or
- 4 Severe Immune deficiency; or
- 5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
- 6 Both
 - 6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 6.2 Either:
 - 6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or 6.2.2 Patient has IoE mediated allerov.

Initial application — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or
 - 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
 - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products; or
 - 2.2 Eosinophilic oesophagitis; or
 - 2.3 Ultra-short gut; or
 - 2.4 Severe Immune deficiency: or
 - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
 - 2.6 Both:
 - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 2.6.2 Either
 - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval

continued...

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

continued...

number: or

2.6.2.2 Patient has IgE mediated allergy.

Renewal — (Infants up to 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has IgE mediated allergy; and
 - 1.2 All of the following:
 - 1.2.1 Patient remains allergic to cow's milk; and
 - 1.2.2 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy or extensively hydrolysed infant formula has been undertaken; and
 - 1.2.3 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
 - 1.2.4 Amino acid formula is required for a nutritional deficit; and
 - 1.2.5 It has been more than three months from the previous approval; or

2 Both:

- 2.1 Patient has non IgE mediated severe gastrointestinal intolerance (including eosinophilic oesophagitis, ultra-short gut and severe immune deficiency); and
- 2.2 All of the following:
 - 2.2.1 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy, or extensively hydrolysed infant formula has been undertaken; and
 - 2.2.2 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
 - 2.2.3 Amino acid formula is required for a nutritional deficit; and
 - 2.2.4 It has been more than three months from the previous approval.

Renewal — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 Either:

- 1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or
- 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
 - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products: or
 - 2.2 Eosinophilic oesophagitis; or
 - 2.3 Ultra-short gut: or
 - 2.4 Severe Immune deficiency; or
 - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
 - 2.6 Both:
 - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 2.6.2 Either:
 - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or
 - 2.6.2.2 Patient has IgE mediated allergy.

Initial application — (for patients who have a current funding under Special Authority form SA1557) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 months for applications meeting the following criteria:

continued...

(Man	Subsidy	F	ully	Brand or
	ufacturer's Price)	Subsidis	sed	Generic
·	\$	Per	•	Manufacturer

continued

All of the following:

- 1 Patient has a valid Special Authority approval for extensively hydrolysed formula (SA1557); and
- 2 Extensively hydrolysed formula (Aptamil Gold+ Pepti Junior, AllerPro SYNEO 1 and 2) is unable to be supplied at this time; and
- 3 The approval only applies to funded dispensings of Neocate Gold and Neocate Syneo.

Note: This criteria is short term funding to cover an out-of-stock situation on some extensively hydrolysed formula powder funded under Special Authority form SA1557. There is no renewal criteria under this restriction.

ENTERAL LIQUID PEPTIDE FORMULA - Special Authority see SA1953 below - Hospital pharmacy [HP3]

Liquid 1 kcal/ml, 500 ml bottle12.44	1 OP	✓ Nutrini Peptisorb
Liquid 1.5 kcal/ml, 500 ml bottle	1 OP	✓ Nutrini Peptisorb
		Energy

⇒SA1953 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has impaired gastrointestinal function and either cannot tolerate polymeric feeds, or polymeric feeds are unsuitable; and
- 2 Any of the following:
 - 2.1 Severe malabsorption; or
 - 2.2 Short bowel syndrome; or
 - 2.3 Intractable diarrhoea; or
 - 2.4 Biliary atresia; or
 - 2.5 Cholestatic liver diseases causing malabsorption; or
 - 2.6 Cystic fibrosis; or
 - 2.7 Proven fat malabsorption; or
 - 2.8 Severe intestinal motility disorders causing significant malabsorption; or
 - 2.9 Intestinal failure: or
 - 2.10 Both:
 - 2.10.1 The patient is currently receiving funded amino acid formula; and
 - 2.10.2 The patient is to be trialled on, or transitioned to, an enteral liquid peptide formula; and
- 3 Either:
 - 3.1 A semi-elemental or partially hydrolysed powdered feed has been reasonably trialled and considered unsuitable; or
 - $3.2\,$ For step down from intravenous nutrition.

Note: A reasonable trial is defined as a 2-4 week trial.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 An assessment as to whether the patient can be transitioned to a cows milk protein or soy infant formula or extensively hydrolysed formula has been undertaken; and
- 2 The outcome of the assessment is that the patient continues to require an enteral liquid peptide formula; and
- 3 General practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

EXTENSIVELY HYDROLYSED FORMULA - Spec	cial Authority see SA1557 on the	e next page	Hospital pharmacy [HP3]
Powder	18.10	450 g OP	✓ Pepti-Junior
	36.20	900 g OP	✓ Allerpro Syneo 1
		-	✓ Allerpro Syneo 2

Subsidy (Manufacturer's Price) Fully Subsidised Per Brand or Generic Manufacturer

⇒SA1557 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 Cows milk formula is inappropriate due to severe intolerance or allergy to its protein content; and
 - 1.2 Fither:
 - 1.2.1 Soy milk formula has been reasonably trialled without resolution of symptoms; or
 - 1.2.2 Soy milk formula is considered clinically inappropriate or contraindicated; or
- 2 Severe malabsorption; or
- 3 Short bowel syndrome; or
- 4 Intractable diarrhoea; or
- 5 Biliary atresia; or
- 6 Cholestatic liver diseases causing malsorption; or
- 7 Cystic fibrosis: or
- 8 Proven fat malabsorption; or
- 9 Severe intestinal motility disorders causing significant malabsorption; or
- 10 Intestinal failure; or
- 11 All of the following:
 - 11.1 For step down from Amino Acid Formula: and
 - 11.2 The infant is currently receiving funded amino acid formula; and
 - 11.3 The infant is to be trialled on, or transitioned to, an extensively hydrolysed formula; and
 - 11.4 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

Note: A reasonable trial is defined as a 2-4 week trial, or signs of an immediate IgE mediated allergic reaction.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 An assessment as to whether the infant can be transitioned to a cows milk protein or soy infant formula has been undertaken; and
- 2 The outcome of the assessment is that the infant continues to require an extensively hydrolysed infant formula; and
- 3 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

Fluid Restricted

⇒SA1698 Special Authority for Subsidy

Initial application only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is fluid restricted or volume intolerant and has been diagnosed with faltering growth; and
- 2 Patient is under the care of a paediatrician or dietitian who has recommended treatment with a high energy infant formula; and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

continued...

Subsidy Manufacturer's Price)	Sı	Fully ubsidised	Brand or Generic
 \$	Per	✓	Manufacturer

Note: "Volume intolerant" patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

Renewal only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient continues to be fluid restricted or volume intolerant and has faltering growth; and
- 2 Patient is under the care of a hospital paediatrician or dietitian who has recommended treatment with a high energy infant formula; and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: 'Volume intolerant' patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

Ketogenic Diet

⇒SA1197 Special Authority for Subsidy

Initial application only from a metabolic physician or paediatric neurologist. Approvals valid for 3 months where the patient has intractable epilepsy, pyruvate dehydrogenase deficiency or glucose transported type-1 deficiency and other conditions requiring a ketogenic diet.

Renewal only from a metabolic physician or paediatric neurologist. Approvals valid for 2 years where the patient is on a ketogenic diet and the patient is benefiting from the diet.

HIGH FAT LOW CARBOHYDRATE FORMULA - Special Authority see \$	SA1197 above – H	ospital pharmacy [HP3]
Powder (unflavoured)3	6.92 300 g C	P ✓ KetoCal 4:1
		✓ Ketocal 3:1
Powder (vanilla)3	6.92 300 g C	P ✓ KetoCal 4:1

SECTION I: NATIONAL IMMUNISATION SCHEDULE

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✓ Manufacturer

Vaccinations

BACILLUS CALMETTE-GUERIN VACCINE - [Xpharm]

For infants at increased risk of tuberculosis. Increased risk is defined as:

- 1) living in a house or family with a person with current or past history of TB; or
- 2) having one or more household members or carers who within the last 5 years lived in a country with a rate of TB > or equal to 40 per 100,000 for 6 months or longer; or
- 3) during their first 5 years will be living 3 months or longer in a country with a rate of TB > or equal to 40 per 100,000 Note a list of countries with high rates of TB are available at www.health.govt.nz/tuberculosis (search for downloads) or www.bcgatlas.org/index.php.

Inj Mycobacterium bovis BCG (Bacillus Calmette-Guerin),

Danish strain 1331, live attenuated, vial with diluent...............0.00

10

✓ BCG Vaccine AJV

(JN.1)

VID-19 VACCINE — [Xpharm] Inj 3 mcg SARS-CoV-2 spike protein (mRNA) LP.8.1 per 0.3 ml, 0.48 ml multi-dose vial; infant vaccine, yellow cap		Per	ubsidised ✓	Brand or Generic Manufacturer
0.48 ml multi-dose vial; infant vaccine, yellow cap				
yellow cap		10 s at hi		Comirnaty (LP.8.1) severe illness.
Inj 10 mcg SARS-CoV-2 spike protein (mRNA) LP.8.1 per 0.3 ml, 0.48 ml single-dose vial; paediatric vaccine, light blue cap	00	10	√ (Comirnaty Omicron (JN.1)
O.3 ml, 0.48 ml single-dose vial; paediatric vaccine, light blue cap	ıths - 4 year	s at hi	gh risk of	
vaccine, light blue cap	ars old; or	10	√ (Comirnaty (LP.8.1)
1) One dose for previously unvaccinated children aged 5–11 yes 2) Up to three doses for immunocompromised children aged 5-1 Inj 30 mcg SARS-CoV-2 spike protein (mRNA) LP.8.1 per 0.3 ml, pre-filled syringe; adult dose	00	10	√ (Comirnaty Omicron (JN.1)
O.3 ml, pre-filled syringe; adult dose				
 2) Up to three doses for immunocompromised people aged 12-13) 3) Up to two doses for previously unvaccinated people 16-29 ye 4) Up to four doses for people aged 16-29 at high risk of severe 5) One dose for previously unvaccinated people aged 30 and ol 	00	10	√ (Comirnaty (LP.8.1)
One additional dose every 6 months for previously vaccinated given at least 6 months after last dose.	15 years old ears old; or illness; or der; or		years and	over – additional do:
Inj 30 mcg bretovameran per 0.3 ml, 0.48 ml vial; adult vaccine, light grey cap		10	-	Comirnaty Omicron

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

Any of the following:

- 1) One dose for previously unvaccinated people aged 12-15 years old; or
- 2) Up to three doses for immunocompromised people aged 12-15 years old; or
- 3) Up to two doses for previously unvaccinated people 16-29 years old; or
- 4) Up to four doses for people aged 16-29 at high risk of severe illness; or
- 5) One dose for previously unvaccinated people aged 30 and older; or
- 6) One additional dose every 6 months for previously vaccinated people aged 30 years and over additional dose is given at least 6 months after last dose.

(Comirnaty Omicron (JN.1) Inj 3 mcg bretovameran per 0.3 ml, 0.48 ml vial; infant vaccine, yellow cap to be delisted 1 June 2026) (Comirnaty Omicron (JN.1) Inj 10 mcg bretovameran per 0.3 ml, 0.48 ml vial; paediatric vaccine, light blue cap to be delisted 1 June 2026)

(Comirnaty Omicron (JN.1) Inj 30 mcg bretovameran per 0.3 ml, 0.48 ml vial; adult vaccine, light grey cap to be delisted 1 June 2026)

DIPHTHERIA, TETANUS AND PERTUSSIS VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- c)
- A) Funded for any of the following criteria:
 - 1) A single dose for pregnant women in the second or third trimester of each pregnancy; or
 - 2) A single dose for parents or primary caregivers of infants admitted to a Neonatal Intensive Care Unit or Specialist Care Baby Unit for more than 3 days, who had not been exposed to maternal vaccination at least 14 days prior to birth; or
 - A course of up to four doses is funded for children from age 7 up to the age of 18 years inclusive to complete full primary immunisation; or
 - 4) An additional four doses (as appropriate) are funded for (re-)immunisation for patients post haematopoietic stem cell transplantation or chemotherapy; pre or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or
 - 5) A single dose for vaccination of patients aged from 65 years old; or
 - 6) A single dose for vaccination of patients aged from 45 years old who have not had 4 previous tetanus doses; or
 - 7) For vaccination of previously unimmunised or partially immunised patients; or
 - 8) For revaccination following immunosuppression; or
 - 9) For boosting of patients with tetanus-prone wounds.

Notes: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.

- B) Contractors will be entitled to claim payment from the Funder for the supply of diphtheria, tetanus and pertussis vaccine to patients eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the diphtheria, tetanus and pertussis vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraphs 1 – 9 above.

Inj 2 IU diphtheria toxoid with 20 IU tetanus toxoid, 8 mcg			
pertussis toxoid, 8 mcg pertussis filamentous			
haemagglutinin and 2.5 mcg pertactin in 0.5 ml prefilled			
svringe	0.00	10	✓ Boostrix

Subs	,	Fully	Brand or
(Manufactur		lised	Generic
\$	Per	1	Manufacturer

DIPHTHERIA, TETANUS, PERTUSSIS AND POLIO VACCINE

- a) Only on a prescription
- b) No patient co-payment payable

c)

- A) Funded for any of the following:
 - 1) A single dose for children up to the age of 7 who have completed primary immunisation; or
 - A course of four vaccines is funded for catch up programmes for children (to the age of 10 years) to complete full primary immunisation; or
 - 3) An additional four doses (as appropriate) are funded for (re-)immunisation for people post HSCT, or chemotherapy; pre- or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or
 - 4) Five doses will be funded for children requiring solid organ transplantation.
- B) Contractors will be entitled to claim payment from the Funder for the supply of Diphtheria, tetanus, pertussis and polio vaccine to people eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Diphtheria, tetanus, pertussis and polio vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.

Inj 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg pertussis toxoid, 25 mcg pertussis filamentous

haemagglutinin, 8 mcg pertactin and 80 D-antigen units

10 ✓ Infanrix IPV

DIPHTHERIA, TETANUS, PERTUSSIS, POLIO, HEPATITIS B AND HAEMOPHILUS INFLUENZAE TYPE B VACCINE

- a) Only on a prescription
- b) No patient co-payment payable

c)

- A) Funded for children meeting any of the following criteria
 - 1) Up to four doses for children under the age of 10 years for primary immunisation; or
 - An additional four doses (as appropriate) for (re-)immunisation of children under the age of 18 years post haematopoietic stem cell transplantation; or
 - 3) An additional four doses (as appropriate) for (re-)immunisation of children under the age of 10 years who are post chemotherapy; pre or post splenectomy; undergoing renal dialysis and other severely immunosuppressive regimens; or
 - 4) Up to five doses for children under the age of 10 years receiving solid organ transplantation.
- B) Contractors will be entitled to claim payment from the Funder for the supply of Diphtheria, tetanus, pertussis, polio, hepatitis B and haemophilus influenzae type b vaccine to people eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Diphtheria, tetanus, pertussis, polio, hepatitis B and haemophilus influenzae type b vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Note: A course of up-to four vaccines is funded for catch up programmes for children (up to and under the age of 10 years) to complete full primary immunisation. Please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes.

Inj 30IU diphtheria with 40IU tetanus and 25mcg pertussis toxoids, 25mcg pertussis filamentous haemagglutinin, 8mcg pertactin, 80D-AgU polio virus, 10mcg hepatitis B antigen, 10mcg H. influenzae type b with tetanus toxoid

10 ✓ Infanrix-hexa

			Subsidy (Manufacturer's Brice)		F Subsidi	ully	Brand or Generic
			(Manufacturer's Price) \$	Per		seu ✓	Manufacturer
HAEM	OPHII	US INFLUENZAE TYPE B VACCINE					
a)	Only	on a prescription					
b) c)	No p	atient co-payment payable					
,	A)	One dose for people meeting any of the following:					
		 For primary vaccination in children; or 					
		 An additional dose (as appropriate) is funded for transplantation, or chemotherapy; functional as 	` '				•
		transplant, pre or post cochlear implants, renal	dialysis and other se	vere	ly immi	inosu	ippressive regimens; or
		 For use in testing for primary immunodeficiency physician or paediatrician. 	diseases, on the re	comi	mendat	ion of	an internal medicine
	B)	Contractors will be entitled to claim payment from the	Funder for the supp	oly of	Haem	ophilu	is influenzae type b
		vaccine to people eligible under the above criteria pu					, ,
		for subsidised immunisation, and they may only do sin the Pharmaceutical Schedule.	o in respect of the H	aemo	ophilus	influe	nzae type b vaccine list
	C)	Contractors may only claim for populations within the	criteria that are cov	ered	by thei	r cont	ract, which may be a
	,	sub-set of the population described in paragraph A a			,		•
lnj	10 m	cg vial with diluent syringe	0.00	1		✓ <u>A</u>	ct-HIB
HEPA1	TITIS .	A VACCINE - [Xpharm]					
Fu	nded	for patients meeting any of the following criteria:					
		o vaccinations for use in transplant patients; or					
	,	o vaccinations for use in children with chronic liver dis					
(3) Or	e dose of vaccine for close contacts of known hepatit	is A cases.				

✓ Havrix 1440

✓ Havrix Junior

1

		Subsidy (Manufacturer's Price)	Fully Subsidised	Brand or Generic	
		\$	Per 🗸	Manufacturer	
HEDATITIC D DECOMPINIANT VACCINE	[Vnhorm]				

HEPATITIS B RECOMBINANT VACCINE - [Xpharm]

Funded for patients meeting any of the following criteria:

- 1) for household or sexual contacts of known acute hepatitis B patients or hepatitis B carriers; or
- 2) for children born to mothers who are hepatitis B surface antigen (HBsAg) positive; or
- 3) for children up to and under the age of 18 years inclusive who are considered not to have achieved a positive serology and require additional vaccination or require a primary course of vaccination; or
- 4) for HIV positive patients; or
- 5) for hepatitis C positive patients; or
- 6) for patients following non-consensual sexual intercourse; or
- 7) for patients prior to planned immunosuppression for greater than 28 days; or
- 8) for patients following immunosuppression; or
- 9) for solid organ transplant patients; or
- 10) for post-haematopoietic stem cell transplant (HSCT) patients; or
- 11) following needle stick injury.

Funded for patients meeting any of the following criteria:

- 1) for household or sexual contacts of known acute hepatitis B patients or hepatitis B carriers; or
- 2) for children born to mothers who are hepatitis B surface antigen (HBsAq) positive; or
- 3) for children up to and under the age of 18 years inclusive who are considered not to have achieved a positive serology and require additional vaccination or require a primary course of vaccination; or
- 4) for HIV positive patients; or
- 5) for hepatitis C positive patients; or
- 6) for patients following non-consensual sexual intercourse; or
- 7) for patients prior to planned immunosuppression for greater than 28 days; or
- 8) for patients following immunosuppression; or
- 9) for solid organ transplant patients; or
- 10) for post-haematopoietic stem cell transplant (HSCT) patients; or
- 11) following needle stick injury; or
- 12) for dialysis patients; or
- 13) for liver or kidney transplant patients.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

HUMAN PAPILLOMAVIRUS (6, 11, 16, 18, 31, 33, 45, 52 AND 58) VACCINE [HPV]

- a) Maximum of 1 inj per prescription
- b) Only on a prescription
- c) No patient co-payment payable
- ď
- a) A) Any of the following:
 - 1) Maximum of two doses for children aged 14 years and under; or
 - 2) Maximum of three doses for people meeting any of the following criteria:
 - 1) People aged 15 to 26 years inclusive; or
 - Either

People aged 9 to 26 years inclusive who have

- 1) Confirmed HIV infection; or
- Received a transplant (including stem cell): or
- 3) Maximum of four doses for people aged 9 to 26 years inclusive post chemotherapy
- B) Contractors will be entitled to claim payment from the Funder for the supply of Human papillomavirus vaccine to people eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Human papillomavirus vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraphs A above.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
INFLUENZA VACCINE Inj 60 mcg in 0.5 ml syringe (quadrivalent vaccine)	120.00	10		fluvac Tetra (2025 formulation)

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

- a) Maximum of 1 inj per prescription
- b) Only on a prescription
- c) No patient co-payment payable
- d)

A) INFLUENZA VACCINE

is available each year for patients who meet the following criteria, as set by Pharmac:

- a) all people 65 years of age and over; or
- b) people under 65 years of age who:
 - i) have any of the following cardiovascular diseases:
 - a) ischaemic heart disease, or
 - b) congestive heart failure, or
 - c) rheumatic heart disease, or
 - d) congenital heart disease, or
 - e) cerebo-vascular disease; or
 - ii) have either of the following chronic respiratory diseases:
 - a) asthma, if on a regular preventative therapy, or
 - b) other chronic respiratory disease with impaired lung function; or
 - iii) have diabetes; or
 - iv) have chronic renal disease; or
 - v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
 - vi) have any of the following other conditions:
 - a) autoimmune disease, or
 - b) immune suppression or immune deficiency, or
 - c) HIV, or
 - d) transplant recipients, or
 - e) neuromuscular and CNS diseases/disorders, or
 - f) haemoglobinopathies, or
 - g) are children on long term aspirin, or
 - h) have a cochlear implant, or
 - i) errors of metabolism at risk of major metabolic decompensation, or
 - i) pre and post splenectomy, or
 - k) Down syndrome, or
 - vii) are pregnant; or
- c) children 4 years of age and under who have been hospitalised for respiratory illness or have a history of significant respiratory illness; or
- d) people under 65 years of age who:
 - i) have any of the following serious mental health conditions:
 - a) schizophrenia, or
 - b) major depressive disorder, or
 - c) bipolar disorder, or
 - d) schizoaffective disorder, or
 - ii) are currently accessing secondary or tertiary mental health and addiction services; or

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy,
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Contractors will be entitled to claim payment for the supply of influenza vaccine to patients eligible under the above criteria pursuant to their contract with Health NZ for subsidised immunisation, and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

MEASLES, MUMPS AND RUBELLA VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- c)

A) Measles, mumps and rubella vaccine

A maximum of two doses for any patient meeting the following criteria:

- 1) For primary vaccination in children; or
- 2) For revaccination following immunosuppression; or
- 3) For any individual susceptible to measles, mumps or rubella; or
- 4) A maximum of three doses for children who have had their first dose prior to 12 months.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes. Although a price is listed for the vaccine, doctors can still order measles mumps and rubella vaccine free of charge, as with other Schedule vaccines.

- B) Contractors will be entitled to claim payment for the supply of measles, mumps and rubella vaccine to patients eligible under the above criteria pursuant to their contract with Health NZ for subsidised immunisation, and they may only do so in respect of the measles, mumps and rubella vaccine listed in the Pharmaceutical Schedule.
- C) Contractors can only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Inj, measles virus 1,000 CCID50, mumps virus 5,012 CCID50,		
Rubella virus 1,000 CCID50; prefilled syringe/ampoule of		
diluent 0.5 ml	10	Priorix

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

MENINGOCOCCAL (GROUPS A, C, Y AND W-135) CONJUGATE VACCINE

Inj 10 mcg of each meningococcal polysaccharide conjugated

a) Only on a prescription

b) No patient co-payment payable

C

- A) Any of the following:
 - Up to three doses and a booster every five years for patients pre- and post splenectomy and for patients
 with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre or post
 solid organ transplant; or

MenQuadfi

- 2) One dose for close contacts of meningococcal cases of any group; or
- 3) One dose for person who has previously had meningococcal disease of any group; or
- 4) A maximum of two doses for bone marrow transplant patients; or
- 5) A maximum of two doses for person pre- and post-immunosuppression*; or
- B) Both:
 - 1) Person is aged between 13 and 25 years, inclusive; and
 - 2) Either:
 - One dose for individuals who are entering within the next three months, or in their first year of living in boarding school hostels, tertiary education halls of residence, military barracks, Youth Justice residences, or prisons; or
 - 2) One dose for individuals who turn 13 years of age while living in boarding school hostels.
- C) Contractors will be entitled to claim payment from the Funder for the supply of Meningococcal A, C, Y and W-135 vaccine to patients eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Meningococcal A, C, Y and W-135 vaccine listed in the Pharmaceutical Schedule.
- D) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraphs A-B above.

Note: children under seven years of age require two doses 8 weeks apart, a booster dose three years after the primary series and then five yearly.

*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days.

Inj 5 mcg of each meningococcal polysaccharide conjugated to a total of approximately 44 mcg of tetanus toxoid carrier

- A) Both:
 - 1) The child is under 12 months of age; and
 - 2) Any of the following:
 - A maximum of three doses (dependant on age at first dose) for patients pre- and post- splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre- or post- solid organ transplant; or
 - A maximum of three doses (dependant on age at first dose) for close contacts of meningococcal cases of any group; or
 - A maximum of three doses (dependant on age at first dose) for child who has previously had meningococcal disease of any group; or
 - 4) A maximum of three doses (dependant on age at first dose) for bone marrow transplant patients; or
 - A maximum of three doses (dependant on age at first dose) for child pre- and post-immunosuppression*.

Note: infants from 6 weeks to less than 6 months of age require a 2+1 schedule, infants from 6 months to less than 12 months of age require a 1+1 schedule. Refer to the Immunisation Handbook for recommended booster schedules with meningococcal ACWY vaccine.

*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days.

Subsidy (Manufacturer's Pri	ce)	Fully Subsidised	Brand or Generic	
\$	Per	✓	Manufacturer	

MENINGOCOCCAL B MULTICOMPONENT VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- c) Any of the following:
 - A) A primary course of up to three doses for children up to the age of 59 months inclusive; or
 - B) Both:
 - 1) Person is 5 years of age or over; and
 - 2) Any of the following:
 - i) up to two doses and a booster every five years for patients pre- and post-splenectomy; or
 - ii) up to two doses and a booster every five years for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited); or
 - iii) up to two doses and a booster every five years pre- or post-solid organ transplant; or
 - iv) up to two doses for close contacts of meningococcal cases of any group; or
 - v) up to two doses for person who has previously had meningococcal disease of any group; or
 - vi) up to two doses for bone marrow transplant patients; or
 - vii) up to two doses for person pre- and post-immunosuppression*; or
 - C) Both:
 - 1) Person is aged between 13 and 25 years (inclusive); and
 - 2) Either:
 - Two doses for individuals who are entering within the next three months, or in their first year of living in boarding school hostels, tertiary education halls of residence, military barracks, Youth Justice residences or prison; or
 - ii) Two doses for individuals who turn 13 years of age while living in boarding school hostels.
 - D) Contractors will be entitled to claim payment from the Funder for the supply of Meningococcal B multicomponent vaccine to patients eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Meningococcal B multicomponent vaccine listed in the Pharmaceutical Schedule.
 - E) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraphs A-C above.

*Immunosuppression due to corticosteroid or other immunosuppressive therapy must be for a period of greater than 28 days.

Inj 175 mcg per 0.5 ml prefilled syringe	0.00	1	Bexsero
		10	✓ Bexsero

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Por 🗸	Manufacturer

PNEUMOCOCCAL (PCV13) CONJUGATE VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- c)
- A) Any of the following:
 - 1) A course of three doses for previously unvaccinated children up to the age of 59 months inclusive; or
 - Two doses are funded for high risk individuals (over the age of 12 months and under 18 years) who have previously received two doses of the primary course of PCV10: or
 - 3) Up to an additional four doses (as appropriate) are funded for the (re)immunisation of high risk children aged under 5 years with any of the following:
 - a) on immunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient immune response; or
 - b) primary immune deficiencies; or
 - c) HIV infection: or
 - d) renal failure, or nephrotic syndrome; or
 - e) who are immune-suppressed following organ transplantation (including haematopoietic stem cell transplant); or
 - f) cochlear implants or intracranial shunts; or
 - g) cerebrospinal fluid leaks; or
 - n) receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage of 20 mg or greater; or
 - i) chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or
 - j) pre term infants, born before 28 weeks gestation; or
 - k) cardiac disease, with cyanosis or failure; or
 - diabetes; or
 - m) Down syndrome; or
 - n) who are pre-or post-splenectomy, or with functional asplenia; or
 - 4) Up to an additional four doses (as appropriate) are funded for the (re-)immunisation of individuals 5 years and over with HIV, pre or post haematopoietic stem cell transplantation, or chemotherapy; pre- or post splenectomy; functional asplenia, pre- or post- solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, intracranial shunts, cerebrospinal fluid leaks or primary immunodeficiency; or
 - 5) For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.
- B) Contractors will be entitled to claim payment from the Funder for the supply of Pneumococcal (PCV13) conjugate vaccine to people eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Pneumococcal (PCV13) conjugate vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Note: please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes Inj 30.8 mcg of pneumococcal polysaccharide serotypes 1, 3, 4,

	NATIONAL	IMMUNISAT	ON SCHEDULE
	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per 🗸	Brand or Generic Manufacturer
PNEUMOCOCCAL (PPV23) POLYSACCHARIDE VACCII Any of the following:	NE - [Xpharm]		
 Up to three doses (as appropriate) for patients w chemotherapy; pre- or post-splenectomy or with complement deficiency (acquired or inherited), c All of the following: 	functional asplenia, pre- or p	oost-solid organ	transplant, renal dialysis,
 a) Patient is a child under 18 years for (re-)im b) Treatment is for a maximum of two doses; c) Any of the following: 			

- immune response; or
 ii) with primary immune deficiencies; or
 - iii) with HIV infection; or
 - iv) with renal failure, or nephrotic syndrome; or
 - v) who are immune-suppressed following organ transplantation (including haematopoietic stem cell transplant); or

i) on immunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient

- vi) with cochlear implants or intracranial shunts; or
- vii) with cerebrospinal fluid leaks; or
- viii) receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage of 20 mg or greater; or
- ix) with chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or
- x) pre term infants, born before 28 weeks gestation; or
- xi) with cardiac disease, with cyanosis or failure; or
- xii) with diabetes: or
- xiii) with Down syndrome; or
- xiv) who are pre-or post-splenectomy, or with functional asplenia; or
- For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician

Inj 575 mcg in 0.5 ml prefilled syringe (25 mcg of each 23 pneumococcal serotype)	0.00	1	✓ Pneumovax 23
POLIOMYELITIS VACCINE – [Xpharm]			
Up to three doses for patients meeting either of the following: 1) For partially vaccinated or previously unvaccinated individua	als: or		
For revaccination following immunosuppression.	110, 01		
Note: Please refer to the Immunisation Handbook for appropriate	schedule for cat	tch-up pro	ogrammes.
Inj 80D antigen units in 0.5 ml syringe	0.00	1	✓ IPOL

Subsidy	Fully	Brand or
(Manufacturer's Price)	(Manufacturer's Price) Subsidised	
\$	Per 🗸	Manufacturer

ROTAVIRUS ORAL VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- c)
- A) Maximum of two doses for people meeting the following:
 - 1) first dose to be administered in infants aged under 14 weeks of age; and
 - 2) no vaccination being administered to children aged 24 weeks or over.
- B) Contractors will be entitled to claim payment from the Funder for the supply of Rotavirus oral vaccine to people eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Rotavirus oral vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Oral susp live attenuated human rotavirus 1,000,000 CCID50 per dose, squeezable tube0.00	10	✓ Rotarix
Oral susp live attenuated human rotavirus 1,000,000 CCID50 per dose, squeezable tube (PVC free)0.00	10	✓ Rotarix
Oral susp live attenuated human rotavirus 1,000,000 CCID50 per dose, prefilled oral applicator0.00	10	✓ Rotarix

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer
--

VARICELLA VACCINE [CHICKENPOX VACCINE]

- a) Only on a prescription
- b) No patient co-payment payable
- c)
- A) Either:
 - 1) Maximum of one dose for primary vaccination for either:
 - a) Any infant born on or after 1 April 2016; or
 - For previously unvaccinated children turning 11 years old on or after 1 July 2017, who have not previously had a varicella infection (chickenpox), or
 - 2) Maximum of two doses for any of the following:
 - a) Any of the following for non-immune individuals:
 - i) with chronic liver disease who may in future be candidates for transplantation; or
 - ii) with deteriorating renal function before transplantation; or
 - iii) prior to solid organ transplant; or
 - iv) prior to any elective immunosuppression*; or
 - v) for post exposure prophylaxis who are immune competent inpatients; or
 - b) For individuals at least 2 years after bone marrow transplantation, on advice of their specialist; or
 - c) For individuals at least 6 months after completion of chemotherapy, on advice of their specialist; or
 - d) For HIV positive non immune to varicella with mild or moderate immunosuppression on advice of HIV specialist; or
 - e) For individuals with inborn errors of metabolism at risk of major metabolic decompensation, with no clinical history of varicella; or
 - f) For household contacts of paediatric patients who are immunocompromised, or undergoing a procedure leading to immune compromise where the household contact has no clinical history of varicella; or
 - g) For household contacts of adult patients who have no clinical history of varicella and who are severely immunocompromised, or undergoing a procedure leading to immune compromise where the household contact has no clinical history of varicella.
- B) Contractors will be entitled to claim payment from the Funder for the supply of Varicella vaccine [Chickenpox vaccine] vaccine to people eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Varicella vaccine [Chickenpox vaccine] listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraphs A above.

* immunosuppression due to steroid or other immunosuppressive therapy must be for a treatment period of greater than 28 days

						Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer	
'ARIC	ELLA	ZOSTER \	/ACC	INE [SHINGLES	VACCINE]					-
a)	Only	on a presc	ription	1						
b)		atient co-pa								
,	A)	Funded fo	•	ents meeting the	following criteria:					
		1)	Two	doses for all peo	ople aged 65 year	s, or				
		2)	Two	doses for people	e 18 years of age	or older with any of the	follov	ving:		
			,			m cell transplant or cel	lular t	herapy; or		
			,		id organ transplar	nt; or				
			,	•	malignancies; or					
			,			ed HIV infection; or				
			e)	•	•	difying anti-rheumatic d	•	,	•	
				biologic, or con rheumatoid arth	•	c) for polymyalgia rheu	matica	a, systemic	lupus erythematosus or	
			f)	end stage kidne	ey disease (CKD	4 or 5); or				
			g)	primary immun	odeficiency					
	B)	vaccine) to for subsidi	pations pations particularly particularly particularly particularly particularly particularly particularly pations particularly particu	ents eligible und	er the above criter d they may only d		tract v	with Health	ster vaccine (Shingles New Zealand (Health NZ) ccine [Shingles vaccine]	1
	C)				atient populations ribed in paragrap		are c	overed by th	neir contract, which may b	е

	-			
Diac	nost		nan	te
	II COL	\mathbf{v}_{-}	g CIII	

Inj 50 mcg per 0.5 ml vial plus vial.......0.00

✓ Shingrix

✓ Shingrix

1

10

- Symbols -	Albey	264–265	Amzoate	30
3TC111	Albustix	81	Anaesthetics	12
- A -	Alchemy Oxaliplatin	152	Anagrelide hydrochloride	15
A-Scabies72	Alchemy Oxybutynin	81	Analgesics	12
Abacavir sulphate110	Aldurazyme	28	Anastrozole	
Abacavir sulphate with	Alecensa	163	Anatrole	17
lamivudine110	Alectinib	163	Anoro Ellipta	
Abacavir/Lamivudine Viatris110	Alendronate sodium	116	Antabuse	
Abilify Maintena135	Alendronate sodium with		Antacids and Antiflatulents	
Abiraterone acetate174	colecalciferol	116	Anthelmintics	9
Acarbose11	Alfacalcidol	<mark>32</mark>	Antiacne Preparations	
Accarb11	Alfamino		Antiallergy Preparations	
Acetazolamide276	Alfamino Junior	301	Antianaemics	
Acetec	Alginic acid	6	Antiandrogen Oral	
Acetic acid with hydroxyquinoline and	Alglucosidase alfa		Contraceptives	79
ricinoleic acid79	Alkeran		Antiarrhythmics	
Acetylcysteine279	Allegron		Antibacterials	
Aci-Jel79	Allerfix		Antibacterials Topical	
Aciclovir	Allerpro Syneo 1		Anticholinergic Agents	
Infection	Allerpro Syneo 2		Anticholinesterases	
Sensory274	Allersoothe		Antidepressants	
Acidex6	Allmercap		Antidiarrhoeals	
Acipimox53	Allopurinol		Antiepilepsy Drugs	
Acitretin73	Almarytm		Antifibrinolytics, Haemostatics and	
Act-HIB310	Alpha-Adrenoceptor Blockers		Local Sclerosants	
Actemra	Alpha-Keri Lotion		Antifibrotics	
Actinomycin D	Alphamox 125		Antifungals	
Actrapid10	Alphamox 250		Antifungals Topical	
Actrapid Penfill	Alprolix		Antihistamines	
Acupan	Alu-Tab		Antihypotensives	
Adalimumab (Amgevita)	Aluminium hydroxide		Antimypotensives	
Adalimumab (Humira - Alternative	Alyacen		Antimigraine Preparations	
brand) 194	Amantadine hydrochloride		Antinausea and Vertigo Agents	
Adapalene	Ambrisentan		Antipruritic Preparations	
Adcetris	Ambrisentan Viatris		Antipsychotics	
ADR Cartridge 1.822	Amgevita		Antiretrovirals	
Adrenaline	Amiloride hydrochloride		Antirheumatoid Agents	
Cardiovascular55	Amiloride hydrochloride with		Antispasmodics and Other Agents	
Respiratory264	furosemide	52	Altering Gut Motility	
Advantan70	Amiloride hydrochloride with		Antithrombotic Agents	
Advate	hydrochlorothiazide	52	Antithymocyte globulin	
Adynovate	Aminophylline		(equine)	18
Afinitor	Amiodarone hydrochloride		Antitrichomonal Agents	
Aflibercept200	Amisulpride		Antituberculotics and	100
AFT-Pyrazinamide104	Amitriptyline		Antileprotics	101
Agents Affecting the	Amlodipine		Antiulcerants	
Renin-Angiotensin System 46	Amorolfine		Antivirals	
Agents for Parkinsonism and Related	Amoxicillin		Anxiolytics	
Disorders 121	Amoxicillin with clavulanic aci		Anzatax	
Agents Used in the Treatment of	Amoxiclav Devatis Forte		Apidra	
Poisonings	Amphotericin B		Apidra SoloStar	
Agrylin	Amsacrine		APO Clomipramine	
Albalon278	AmsaLyo		APO Health Macrogol	
Albendazole 92	Amsidine		APO-Atomoxetine	111
AIDOI 140201692	AMOIUITE	104	AI O-VIOLIOVERINE	140

INDEX: Generic Chemicals and Brands

Apo-Azithromycin	93	Atnahs Olsalazine	8	G]	95
APO-Candesartan HCTZ		Atomoxetine	143	Besponsa	214
16/12.5	47	Atorvastatin	54	Beta Cream	
APO-Candesartan HCTZ		Atropine sulphate		Beta Ointment	
32/12.5	47	Cardiovascular	48	Beta Scalp	
Apo-Temozolomide		Sensory		Beta-Adrenoceptor Agonists	
Apomorphine hydrochloride		Atropt		Beta-Adrenoceptor Blockers	
Aprepitant		Atrovent		Beta-hCG low sensitivity urine tes	
Apresoline		Augmentin		kit	
Aptamil Feed Thickener		Aurorix		Betadine	
Aqueous cream		AutoSoft 30		Betadine Skin Prep	
Aratac		AutoSoft 90		Betaferon	
Arava		Avelox		Betahistine dihydrochloride	
Arginine		Avonex		Betaine	
Arginine Arg		Axitinib		Betamethasone dipropionate	
•		Azacitidine			
Aripiprazole				Betamethasone dipropionate with	
Aripiprazole Sandoz		Azacitidine Dr Reddy's		calcipotriol	
Aristocort		Azamun		Betamethasone sodium phosphat	
Arrotex-Prazosin S29		Azathioprine		with betamethasone acetate	
Arrow - Clopid		Azilect		Betamethasone valerate	,
Arrow - Lattim		Azithromycin		Betamethasone valerate with sodi	
Arrow-Amitriptyline		Azopt		fusidate [fusidic acid]	
Arrow-Bendrofluazide		AZT	111	Betnovate	
Arrow-Brimonidine		- B -		Bevacizumab	202
Arrow-Diazepam		B-D Micro-Fine		Bexsero	
Arrow-Doxorubicin	155	B-D Ultra Fine	17	Bezafibrate	53
Arrow-Fluoxetine	128	B-D Ultra Fine II	17	Bezalip	53
Arrow-Losartan &		Bacillus Calmette-Guerin (BCG)		Bezalip Retard	53
Hydrochlorothiazide	47	vaccine	185	Bicalutamide	175
Arrow-Norfloxacin	114	Bacillus Calmette-Guerin		Bicillin LA	95
Arrow-Ornidazole	103	vaccine	306	BiCNU	151
Arrow-Quinapril 10	46	Baclofen	119	Bile and Liver Therapy	10
Arrow-Quinapril 20	46	Bactroban	67	Biltricide	
Arrow-Quinapril 5		Balance	28	Bimatoprost	276
Arrow-Roxithromycin		Barrier Creams and Emollients	70	Binarex	
Arrow-Timolol		Bayshore	102	Binocrit	36
Arrow-Topiramate		BCG Vaccine AJV		Biocon	
Arrow-Tramadol		Beclazone 100		Biodone	
Arsenic trioxide		Beclazone 250		Biodone Extra Forte	
Asacol		Beclazone 50		Biodone Forte	
Asacol S29		Beclomethasone dipropionate		Bisacodyl	
Ascend-Cefuroxime		Bedaquiline		Bisacodyl Viatris	
Ascorbic acid		Bee venom allergy treatment		Bisoprolol fumarate	
Aspen Adrenaline		Bendamustine hydrochloride		BK Lotion	
Aspirin		Bendamustine Sandoz		Bleomycin sulphate	
Blood	40	Bendrofluazide		Blood Colony-stimulating	
Nervous		Bendroflumethiazide	52	Factors	13
			E0	Blood glucose diagnostic test	40
Asthalin		[Bendrofluazide]			15
Atazanavir sulphate		Benralizumab		meter	15
Atazanavir Viatris		Benzathine benzylpenicillin		Blood glucose diagnostic test	40
Atenolol		Benzatropine mesylate		strip	16
Atenolol AFT		Benzbromarone		Blood glucose test strips (visually	40
Atenolol Viatris		Benzetacil		impaired)	16
Atezolizumab		Benztrop		Blood Ketone Diagnostic Test	
ATGAM		Benzydamine hydrochloride		Strip	
Ativan	138	Benzylpenicillin sodium [Penicilli	ın	Boostrix	308

Bortezomib	155	Candestar	47	Chlorafast	274
Bosentan	59	Canesten	68	Chlorambucil	151
Bosentan Dr Reddy's	59	Capecitabine	153	Chloramphenicol	274
Bplex		Capecitabine Viatris	153	Chlorothiazide	52
Brentuximab Vedotin		Capsaicin		Chlorpromazine hydrochloride	
Breo Ellipta	267	Musculoskeletal	116	Chlorsig	
Brevinor 1/28	78	Nervous	123	Chlortalidone [Chlorthalidone]	53
Breztri Aerosphere	268	Captopril	46	Chlorthalidone	53
Bricanyl Turbuhaler	267	Carafate		Chlorvescent	
Brimonidine tartrate		Carbaccord	151	Choice 380 7med Nsha Silver/co	pper
Brimonidine tartrate with timolol		Carbamazepine	129	Short	7 8
maleate	277	Carbimazole	86	Ciclosporin	258
Brinzolamide	276	Carboplatin	151	Cidomycin P/Free	97
BSF Allegron	279	Carboplatin Accord	151	Cilicaine VK	96
BSF Estradiol TDP Mylan	279	Carbosorb-X		Cinacalcet	82
Budesonide		Cardinol LA	50	Cinacalet Devatis	82
Alimentary	6	Cardizem CD	51	Ciprofloxacin	
Respiratory26	66, 272	CareSens Dual	15	Infection	97
Budesonide Te Arai	6	CareSens N		Sensory	
Budesonide with eformoterol	266	CareSens N POP	15	Ciprofloxacin Teva	
Budesonide with glycopyrronium a	and	CareSens N Premier	15	Cisplatin	
eformoterol		CareSens PRO	16	Cisplatin Accord	
Bumetanide		Carmellose sodium with gela	tin and	Cisplatin Ebewe	
Buprenorphine Naloxone BNM	147	pectin		Citalopram hydrobromide	
Buprenorphine with naloxone	147	Carmustine		Citrulline1000	
Bupropion hydrochloride	148	Carvedilol	49	Cladribine	153
Burel	100	Carvedilol Sandoz	49	Clarithromycin	
Burinex	51	Catapres	51	Alimentary	9
Buspirone hydrochloride	137	Cefaclor monohydrate		Infection	94
Buspirone Viatris		Cefalexin	92	Clexane	
Busulfan		Cefalexin Lupin		Clexane Forte	41
- C -		Cefalexin Sandoz	92	Clindamycin	97
Cabergoline	91	Cefazolin	92	Clinicians	
Caffeine citrate		Cefazolin-AFT	92	Clinicians Renal Vit	33
Calamine	69	Ceftriaxone	93	Clobazam	129
Calci-Tab 500	33	Ceftriaxone-AFT	93	Clobetasol propionate	69, 74
Calcipotriol	73	Cefuroxime axetil	93	Clobetasone butyrate	69
Calcitonin		Celapram		Clofazimine	
Calcitriol	32	Celebrex	115	Clomazol	
Calcitriol XL	32	Celecoxib	115	Dermatological	68
Calcitriol-AFT	32	Celecoxib Pfizer	115	Genito-Urinary	79
Calcium 500 mg Hexal		Celestone Chronodose	83	Clomifene citrate	91
Calcium carbonate	6, 33	Cellcept		Clomipramine hydrochloride	127
Calcium carbonate PAI	6	Centrally-Acting Agents	51	Clomipramine Teva	127
Calcium Channel Blockers	50	Cephalexin ABM		Clonazepam1	
Calcium Disodium Versenate	280	Cerazette	78	Clonidine	51
Calcium folinate	153	Cetirizine hydrochloride	265	Clonidine hydrochloride	51
Calcium gluconate	33	Cetomacrogol	71	Clonidine Teva	51
Calcium Homeostasis	82	Cetomacrogol with glycerol	71	Clopidogrel	40
Calcium polystyrene sulphonate	45	Cetomacrogol-AFT		Clopine	
Calcium Resonium		Cetuximab		Clopixol1	35, 137
Calogen		Champix		Clotrimazole	
Camber	56	Charcoal		Dermatological	68
Candesartan cilexetil	47	CheckTop		Genito-Urinary	
Candesartan cilexetil with		Chemotherapeutic Agents	150	Clozapine	134
hydrochlorothiazide	47	Chickenpox vaccine		Clozaril	134

Clustran1	32	Curam Duo 500/125	95	DBL Methotrexate Onco-Vial	154
Co-trimoxazole	99	Cvite		DBL Naloxone Hydrochloride	
Coal tar	73	Cyclizine hydrochloride	133	DBL Pethidine Hydrochloride	
Coal tar with allantoin, menthol,		Cyclizine lactate		DBL Vincristine Sulfate	
phenol and sulphur	73	Cyclogyl		Decozol	
Coal tar with salicylic acid and		Cyclonex		Deferasirox	
sulphur	73	Cyclopentolate hydrochloride		Deferiprone	
Coco-Scalp		Cyclophosphamide		Denosumab	
Codeine phosphate1		Cyclorin		Deolate	
Coenzyme Q10		Cycloserine		Deoxycoformycin	
Colchicine1		Cyproterone acetate		Depo-Medrol	
Colecalciferol		Cyproterone acetate with	04	Depo-Provera	
Colestyramine		ethinyloestradiol	70	Depo-Testosterone	70
				Deprim	04
Colestyramine - Mylan		Cystadane	150		
Coliform 1		Cytarabine		Dermol	
Colifoam		Cytotec		Desferrioxamine mesilate	
Colistin sulphomethate		Cytoxan	152	Desmopressin	
Collodion flexible2		-D-		Desmopressin acetate	
Colloidal bismuth subcitrate		D-Penamine		Desmopressin-PH&T	
Colofac		Dabigatran		Desogestrel	78
Colomycin		Dabrafenib		Detection of Substances in	
Coloxyl		Dacarbazine		Urine	8 [.]
Combigan2		Dactinomycin [Actinomycin D].	155	Dexamethasone	
Comirnaty (LP.8.1)3	07	Daivobet	73	Hormone	8
Comirnaty Omicron (JN.1)3	07	Daivonex	73	Sensory	27
Compound electrolytes	45	Daktarin	68	Dexamethasone phosphate	8
Compound electrolytes with glucose		Dalacin C	97	Dexamethasone with framycetin	and
[Dextrose]	45	Dantrium	119	gramicidin	274
Compound hydroxybenzoate2	82	Dantrium S29	119	Dexamethasone with neomycin	
Concerta1	45	Dantrolene	119	sulphate and polymyxin B	
Condoms	77	Daonil	11	sulphate	271
		Daoriii		Suipriale	41
Condyline				Dexamfetamine sulfate	14
Condyline S29	75	Dapa-Tabs	53	Dexamfetamine sulfate Dexcom G6	143
Condyline S29	75	Dapa-Tabs	53 103	Dexamfetamine sulfate Dexcom G6	143
Condyline S29 Continuous glucose monitor	75 75	Dapa-Tabs	53 103 111	Dexamfetamine sulfate	143 22
Condyline S29 Continuous glucose monitor (interoperable)	75 75	Dapa-Tabs Dapsone Darunavir Darunavir Viatris	53 103 111	Dexamfetamine sulfate Dexcom G6 Dexcom G7 Dexcom ONE+	143 22 23
Condyline S29 Continuous glucose monitor (interoperable) Continuous glucose monitor	75 75 22	Dapa-Tabs	53 103 111 111	Dexamfetamine sulfate Dexcom G6 Dexcom G7 Dexcom ONE+ Dexmethsone	143 22 23
Condyline S29	75 75 22 23	Dapa-Tabs	53 103 111 111 165	Dexamfetamine sulfate Dexcom G6 Dexcom G7 Dexcom ONE+ Dexmethsone Dextrochlorpheniramine	143 22 23 23
Condyline S29	75 75 22 23 78	Dapa-Tabs Dapsone Darunavir Darunavir Viatris Dasatinib Dasatinib-Teva Daunorubicin	53 103 111 165 165 155	Dexamfetamine sulfate Dexcom G6 Dexcom G7 Dexcom ONE+ Dexmethsone Dextrochlorpheniramine maleate	143 22 23 83
Condyline S29	75 75 22 23 78 76	Dapa-Tabs Dapsone Darunavir Darunavir Viatris Dasatinib Dasatinib-Teva Daunorubicin David One Step Cassette Preg	53 103 111 165 165 155 nancy	Dexamfetamine sulfate Dexcom G6 Dexcom G7 Dexcom ONE+ Dexmethsone Dextrochlorpheniramine maleate Dextrose	143 22 23 83 265 44–45
Condyline S29	75 75 22 23 78 76	Dapa-Tabs Dapsone Darunavir Darunavir Viatris Dasatinib Dasatinib-Teva Daunorubicin David One Step Cassette Preg Test	53 103 111 165 165 155 nancy	Dexamfetamine sulfate Dexcom G6 Dexcom G7 Dexcom ONE+ Dexmethsone Dextrochlorpheniramine maleate Dextrose DHC Continus	143 22 23 83 265 44–44
Condyline S29	75 75 22 23 78 76 39	Dapa-Tabs Dapsone Darunavir Darunavir Viatris Dasatinib Dasatinib-Teva Daunorubicin David One Step Cassette Preg Test. DBL Acetylcysteine	53 103 111 165 165 155 nancy 80	Dexamfetamine sulfate Dexcom G6 Dexcom G7 Dexcom ONE+ Dexmethsone Dextrochlorpheniramine maleate Dextrose DHC Continus Diabetes	143 22 23 83 265 44–44 125
Condyline S29	75 75 22 23 78 76 39	Dapa-Tabs	53103111165165155 nancy80279	Dexamfetamine sulfate Dexcom G6 Dexcom G7 Dexcom ONE+ Dexmethsone Dextrochlorpheniramine maleate Dextrose DHC Continus Diabetes Diabetes Management	14; 22 23 83 26; 44–4; 12!
Condyline S29	75 75 22 23 78 76 39 83 69	Dapa-Tabs	53103111165165155 nancy8027955	Dexamfetamine sulfate Dexcom G6 Dexcom G7 Dexcom ONE+ Dexmethsone Dextrochlorpheniramine maleate Dextrose DHC Continus Diabetes Management Diacomit	143 26 26 26 44–4! 10
Condyline S29	75 75 22 23 78 76 39 83 69	Dapa-Tabs	53103111165165155 nancy8027955	Dexamfetamine sulfate Dexcom G6 Dexcom G7 Dexcom ONE+ Dexmethsone. Dextrochlorpheniramine maleate Dextrose DHC Continus Diabetes Management Diacomit Diagnostic Agents	14; 22 25 265 44–4; 12! 14 13
Condyline S29	75 75 22 23 78 76 39 83 69 34 55	Dapa-Tabs	53103111165165155 nancy8027955271155	Dexamfetamine sulfate Dexcom G6 Dexcom G7 Dexcom ONE+ Dexmethsone. Dextrochlorpheniramine maleate Dextrose DHC Continus Diabetes Diabetes Management Diagnostic Agents Diamide Relief	14; 22 20 8; 26; 44–4; 12; 13; 13; 32;
Condyline S29	75 75 22 23 78 76 39 83 69 34 55 43	Dapa-Tabs	53103111165165155 nancy8027955	Dexamfetamine sulfate Dexcom G6 Dexcom G7 Dexcom ONE+ Dexmethsone. Dextrochlorpheniramine maleate Dextrose DHC Continus Diabetes Diabetes Management Diagnostic Agents Diamide Relief Diasip	143 25 26 44–4! 10 13 322
Condyline S29	75 75 22 23 78 76 39 83 69 34 55 43 29	Dapa-Tabs		Dexamfetamine sulfate Dexcom G6 Dexcom G7 Dexcom ONE+ Dexmethsone. Dextrochlorpheniramine maleate Dextrose DHC Continus Diabetes Diabetes Management Diacomit Diagnostic Agents Diamide Relief Diazepam	14326526512510131313261251413
Condyline S29	75 75 22 23 78 76 39 83 69 34 55 43 29 46	Dapa-Tabs		Dexamfetamine sulfate Dexcom G6 Dexcom G7 Dexcom ONE+ Dexmethsone Dextrochlorpheniramine maleate DHC Continus Diabetes Diabetes Management Diagnostic Agents Diamide Relief Diazepam Diazoxide	14;22;26;44–4;12;13;32;28; 29, 138;10
Condyline S29	75 75 22 23 78 76 39 83 69 34 55 54 43 29 46 07	Dapa-Tabs		Dexamfetamine sulfate Dexcom G6 Dexcom G7 Dexcom ONE+ Dexmethsone Dextrochlorpheniramine maleate Dextrose DHC Continus Diabetes Management Diagnostic Agents Diamide Relief Diazepam Diazoxide Diazoxide Dicorom G7 Dextrose Dextrose Dextrose Diates Diabetes Diaprostic Agents Diamide Relief Diazip Diazepam Diazoxide Dicofenac Devatis	14;22;26;44–4;12;13;32;28; 29, 138;21;
Condyline S29	75 75 22 23 78 76 39 83 69 34 55 43 29 46 007 23	Dapa-Tabs		Dexamfetamine sulfate Dexcom G6 Dexcom G7 Dexcom ONE+ Dexmethsone Dextrochlorpheniramine maleate Dextrose DHC Continus Diabetes Management Diagnostic Agents Diamide Relief Diazepam Diazoxide Diclofenac Devatis Diclofenac Sandoz	14;22;26;44–4;12;13;32;28; 29, 138;21;
Condyline S29	75 75 22 23 78 76 39 83 69 34 55 43 29 46 007 23 23	Dapa-Tabs		Dexamfetamine sulfate	
Condyline S29 Continuous glucose monitor (interoperable) Continuous glucose monitor (standalone) Contraceptives - Hormonal Copaxone Corticosteroids and Related Agents for Systemic Use Corticosteroids Topical Cosentyx Cosmegen 1 Coumadin Country Life Coversyl COVID-19 vaccine 3 Creon 10000 Creon 25000 Creon Micro	75 75 22 23 78 76 39 83 69 34 55 43 29 46 007 23 23 23 23	Dapa-Tabs		Dexamfetamine sulfate Dexcom G6 Dexcom G7 Dexcom ONE+ Dexmethsone Dextrochlorpheniramine maleate Dextrose DHC Continus Diabetes Management Diacomit Diagnostic Agents Diamide Relief Diasip Diazepam Diazepam Diazepam Diazoxide Diclofenac Devatis Diclofenac Sandoz Diclofenac sodium Musculoskeletal	144 22 22 23 24 24 24 24 24 24 25 25 25 25 25 25 25 25 25 25 25 25 25
Condyline S29 Continuous glucose monitor (interoperable) Continuous glucose monitor (standalone) Contraceptives - Hormonal Copaxone 1 Corticosteroids and Related Agents for Systemic Use Corticosteroids Topical Cosentyx 2 Cosmegen 1 Coumadin Country Life Coversyl COVID-19 vaccine 3 Creon 10000 Creon 25000 Creon Micro Crizotinib 1	75 75 22 23 78 76 39 83 69 34 55 43 29 46 007 23 23 23 63	Dapa-Tabs		Dexamfetamine sulfate Dexcom G6 Dexcom G7 Dexcom ONE+ Dexmethsone Dextrochlorpheniramine maleate Dextrose DHC Continus Diabetes Management Diacomit Diagnostic Agents Diamide Relief Diasip Diazepam Diazepam Diazepam Diazoxide Diclofenac Devatis Diclofenac Sandoz Diclofenac sodium Musculoskeletal Sensory	144 22 22 22 23 24 24 24 24 24 24 24 24 24 24 24 24 24
Condyline S29 Continuous glucose monitor (interoperable) Continuous glucose monitor (standalone) Contraceptives - Hormonal Contraceptives - Non-hormonal Corticosteroids and Related Agents for Systemic Use Corticosteroids Topical Cosentyx Cosmegen 1 Coumadin Country Life Coversyl COVID-19 vaccine 3 Creon 10000 Creon 25000 Creon Micro Crizotinib 1 Crotamiton	75 75 22 23 78 76 39 83 69 34 55 54 46 07 23 23 23 63 69	Dapa-Tabs		Dexamfetamine sulfate Dexcom G6 Dexcom G7 Dexcom ONE+ Dexmethsone Dextrochlorpheniramine maleate Dextrose DHC Continus Diabetes Management Diacomit Diagnostic Agents Diamide Relief Diazepam Diazepam Diazepam Diazoxide Diclofenac Devatis Diclofenac Sandoz Diclofenac sodium Musculoskeletal Sensory Differin	144 22 22 22 23 24 24 24 24 24 25 25 25 25 25 25 25 25 25 25 25 25 25
Condyline S29 Continuous glucose monitor (interoperable) Continuous glucose monitor (standalone) Contraceptives - Hormonal Copaxone 1 Corticosteroids and Related Agents for Systemic Use Corticosteroids Topical Cosentyx 2 Cosmegen 1 Coumadin Country Life Coversyl COVID-19 vaccine 3 Creon 10000 Creon 25000 Creon Micro Crizotinib 1	75 75 22 23 78 76 39 83 69 34 55 54 32 29 46 007 223 223 63 69 67	Dapa-Tabs		Dexamfetamine sulfate Dexcom G6 Dexcom G7 Dexcom ONE+ Dexmethsone Dextrochlorpheniramine maleate Dextrose DHC Continus Diabetes Management Diacomit Diagnostic Agents Diamide Relief Diasip Diazepam Diazepam Diazepam Diazoxide Diclofenac Devatis Diclofenac Sandoz Diclofenac sodium Musculoskeletal Sensory	144 22 22 27 27 26 66 3 3 3 27 27 27 27 27 27 27 27 27 27 27 27 27

Digestives Including Enzymes23	Drofate50	Emtriva	11
Digoxin48	Drugs Affecting Bone	Emulsifying ointment	7
Dihydrocodeine tartrate125	Metabolism 116	Enalapril maleate	4
Dilantin	Dual blood glucose and blood ketone	Enbrel	
Dilantin Infatab130	diagnostic test meter 15	Endocrine Therapy	174
Dilantin Paediatric130	Dulaglutide12	Endoxan	15
Diltiazem CD Clinect51	Dulcolax SP Drop26	Energivit	
Diltiazem hydrochloride51	Duocal Super Soluble Powder284	Engerix-B	31
Dimethicone70, 72	Duolin267	Enhertu	
Dimethyl fumarate 139	Duolin HFA267	Enlafax XR	
Dipentum8	DuoResp Spiromax266	Enoxaparin sodium	
Diphtheria, tetanus and pertussis	Duride55	Enstilar	
vaccine308	Durvalumab247	Ensure	
Diphtheria, tetanus, pertussis and	- E -	Ensure Plus	
polio vaccine309	e-chamber La Grande273	Ensure Plus HN	
	e-chamber Mask273	Ensure Plus HN RTH	
Diphtheria, tetanus, pertussis, polio,		Ensure Two Cal HN RTH	
hepatitis B and haemophilus	e-chamber Turbo273		
influenzae type B vaccine 309	E-Mycin94	Entacapone	
Diprosone	e5 Pharma	Entacapone Viatris	
Diprosone OV69	Ear Preparations274	Entecavir	
Dipyridamole40	Ear/Eye Preparations274	Entecavir (Rex)	10
Dipyridamole - Strides40	Easiphen Liquid297	Entrectinib	
Disopyramide phosphate48	Econazole nitrate68	Entresto 24/26	
Distoside92	Efavirenz110	Entresto 49/51	
Disulfiram148	Efavirenz Milpharm110	Entresto 97/103	
Diuretics51	Efavirenz with emtricitabine and	Entyvio	
Docetaxel	tenofovir disoproxil111	Epilim	13
Docetaxel Accord155	Eformoterol fumarate dihydrate266	Epilim Crushable	13
Docetaxel Sandoz155	Eftrenonacog alfa [Recombinant	Epilim IV	13
Docusate sodium24	factor IX] 37	Epilim S/F Liquid	13
Docusate sodium with	Efudix75	Epilim Syrup	13
sennosides24	Egopsoryl TA73	Epipen	26
Dolutegravir111	Elaprase28	Epipen Jr	26
Dolutegravir with lamivudine111	Elecare301	Epirubicin Ebewe	
Domperidone133	Elecare LCP301	Epirubicin hydrochloride	
Domperidone Viatris133	Electral45	Eplerenone	
Donepezil hydrochloride146	Elelyso30	Epoetin alfa	
Dornase alfa271	Elemental 028 Extra289	Epoprostenol	
Dortimopt	Elexacaftor with tezacaftor, ivacaftor	Eptacog alfa [Recombinant factor	
Dorzolamide with timolol276	and ivacaftor271	VIIa]	3
Dostinex91	Elidel73	Erbitux	
Dosulepin [Dothiepin]	Elocon70	Ergometrine maleate	
hydrochloride	Elocon Alcohol Free70	Erlotinib	
Dosulepin Viatris127	Eltrombopag37	Erythrocin IV	
Dothiepin	Eltroxin	Erythromycin (as lactobionate)	
Dovato	EMB Fatol		
Doxazosin	Emend Tri-Pack132	Erythromycin ethyl succinate Esbriet	
Doxazosin Clinect			
	Emicizumab	Escitalopram	12
Doxine	EMLA	Escitalopram (Ethics)	
Doxorubicin Ebewe	Empagliflozin14	Eskazole	
Doxorubicin hydrochloride	Empagliflozin with metformin	Essential Amino Acid Mix	
Doxycycline96	hydrochloride14	Estradiol TDP Mylan	8
DP Lotion	Emsogen	Estradot	8
DP Lotn HC70	Emtricitabine111	Estrofem	
DP-Captopril46	Emtricitabine with tenofovir	Estrogel	
Dr Reddy's Omeprazole9	disoproxil 107	Etanercept	17

Ethambutol hydrochloride	104	Ferro-tab	34	Flynn	9
Ethics Aspirin	123	Ferrograd	34	FML	27
Ethics Aspirin EC	40	Ferrosig	35	Foban	6
Ethinyloestradiol with		Ferrous fumarate		Folic acid	3
desogestrel	78	Ferrous fumarate with folic acid	34	Folic Acid multichem	3
Ethinyloestradiol with		Ferrous sulfate	34	Folic Acid Viatris	3
levonorgestrel	78	Fexaclear	265	Food Thickeners	
Ethinyloestradiol with		Fexofenadine hydrochloride	265	Foods And Supplements For	
norethisterone	78	Fibro-vein		Inherited Metabolic Disease	29
Ethosuximide	129	Filgrastim	43	Fortini	28
Etopophos	156	Finasteride	80	Fortini Multi Fibre	28
Etoposide	156	Fingolimod	139	Fortisip	29
Etoposide phosphate		Firazyr	264	Fortisip Multi Fibre	
Etravirine		Flagyl	103	Fosamax	
Eumovate	69	Flagyl-S		Fosamax Plus	11
Eurofolic	153	Flamazine	68	Fosfomycin	11
Evara	70–71	Flecainide acetate	48	Framycetin sulphate	27
Evara Emulsifying Ointment	71	Flecainide BNM	48	Freestyle Libre 2	
EVARA White Soft Paraffin		Flecainide Controlled Release		Freestyle Libre 2 Plus	
Everet	130	Teva	48	Freestyle Libre 3 Plus	
Everolimus	258	Fleet Phosphate Enema	25	Frisium	
Evista	117	Flixonase Hayfever & Allergy	272	Frumil	5
Evrysdi	142	Flixotide	266	Frusemide	
Exemestane		Flixotide Accuhaler	266	Fucicort	
Exjade	279	Florinef	83	Fucidin	9
Extemporaneously Compounded	d	Fluanxol	136	Fucithalmic	27
Preparations and		Flucil	95	Fucithalmic Canada (ON)	27
Galenicals	282	Flucloxacillin		Fucithalmic Spain	
Eye Preparations	274	Flucloxin	95	Fulvestrant	
Eylea		Flucon	275	Fulvestrant EVER Pharma	17
Ezetimibe		Fluconazole	99	Fungilin	3
Ezetimibe Sandoz	5 <u>5</u>	Fludara Oral	153	Furosemide [Frusemide]	
Ezetimibe with simvastatin	55	Fludarabine Ebewe	153	Furosemide-Baxter	
-F-		Fludarabine phosphate	153	fusidic acid	
Factor eight inhibitor bypassing		Fludrocortisone acetate		Dermatological	68, 7
fraction	39	Fluids and Electrolytes		Infection	
Famotidine	9	Flumetasone pivalate		Sensory	
Famotidine Hovid	9	Fluocortolone caproate with		´ - G -	
Famotidine Hovid MY		fluocortolone pivalate and		GA Explore 5	29
Faricimab		cinchocaine	8	GA1 Anamix Infant	
Fasenra	201	Fluorometholone	275	GA1 Anamix Junior	29
Faslodex	175	Fluorouracil	153	Gabapentin	12
Fatty Emulsion Cream (Evara)	71	Fluorouracil Accord		Gacet	
Febuxostat		Fluorouracil sodium		Galsulfase	
Febuxostat (Teva)		Fluox		Galvumet	
FEIBA NF		Fluoxetine hydrochloride		Galvus	1
Felo 10 ER		Flupenthixol decanoate		Gardasil 9	31
Felo 5 ER	50	Flutamide		Gastrodenol	
Felodipine		Flutamin		Gaviscon Extra Strength	
Fentanyl		Fluticasone		Gaviscon Infant	
Fentanyl Sandoz		Fluticasone furoate with		Gazyva	
Ferinject		umeclidinium and vilanterol	269	Gefitinib	
Ferodan		Fluticasone furoate with		Gemcitabine Ebewe	
Ferriprox		vilanterol	267	Gemcitabine hydrochloride	
Ferro-F-Tabs	34	Fluticasone propionate		Gemtuzumab ozogamicin	
Ferro-Liquid		Fluticasone with salmeterol		Gentamicin sulphate	
· /	• •				

Gilenya	. 139	Healtheries Simple Baking Mix295 Hydroxyurea	
Ginet	79	Hemastix81 [hydroxycarbamide]	. 156
Glatiramer acetate	. 139	Hemlibra38 Hygroton	
Glecaprevir with pibrentasvir		Heparin sodium42 Hylo-Fresh	
Glibenclamide		Heparin Sodium Panpharma42 Hymenoptera264	
Gliclazide		Heparinised saline	
Glipizide		Heparon Junior	
Glizide		Hepatitis A vaccine	9
Glucagen Hypokit		Hepatitis B recombinant Hyoscine hydrobromide	
Glucagon hydrochloride			
Glucerna Select		Herzuma	
Glucose [Dextrose]		Hikma	211
Gluten Free Foods		Hikma Acetylcysteine279	0.
Glycerin with sodium saccharin		Hikma-Propranolol	
Glycerin with sucrose	. 282	Hiprex	
Glycerol		Histaclear	
Alimentary		Holoxan152 Ibuprofen SR BNM	
Extemporaneous	. 282	Horleys Bread Mix296 Icatibant	
Glyceryl trinitrate		Horleys Flour296 Idarubicin hydrochloride	156
Alimentary		Hormone Replacement Therapy - Idursulfase	28
Cardiovascular	55	Systemic	
Glycopyrronium	.268	HPV312 lloprost	64
Glycopyrronium bromide		Humalog11 Imatinib mesilate	
Glycopyrronium with		Humalog Mix 2511 Imatinib-Rex	167
indacaterol	. 268	Humalog Mix 5011 Imbruvica	156
Glycosade	.299	Human papillomavirus (6, 11, 16, 18, Imfinzi	247
Glytactin Bettermilk	.298	31, 33, 45, 52 and 58) vaccine Imipramine Crescent	
Gold Knight		[HPV] 312 Imipramine hydrochloride	
Gold Knight XL		Humatin98 Imiquimod	
Goserelin		Humira	
Gynaecological Anti-infectives		HumiraPen	
- H -		Humulin 30/7011 Immunisation Flu and Shingles	
Habitrol	148	Humulin NPH	
Haemophilus influenzae type B	. 140	Humulin R	
vaccine	310	Hyaluronic acid278 Incruse Ellipta	
Haldol		Hydralazine	
Haldol Concentrate		Hydralazine hydrochloride56 Indapamide	
		Hydralyte - Lemonade45 Infanrix IPV	
Haldol Decanoas		• •	
Haloperidol		,	
Haloperidol decanoate Harvoni		Dermatological 69 Infant Formulae 58 Infartrini 58 Infatrini 58 Infatr	
Havrix 1440		Hydrocortisone acetate	
Havrix Junior		Hydrocortisone acetate with Influenza vaccine	313
Haylor syrup		pramoxine hydrochloride	
HCU Anamix Infant		Hydrocortisone and paraffin liquid (2025 formulation)	
HCU Anamix Junior		and lanolin	268
HCU Anamix Junior LQ		Hydrocortisone butyrate70, 74 Inhaled Long-acting	
HCU Explore 5	. 296	Hydrocortisone with cinchocaine 8 Beta-adrenoceptor Agonists	. 266
HCU Express 15	. 296	Hydrocortisone with miconazole70 Inlyta	163
HCU Lophlex LQ	. 296	Hydrocortisone with natamycin and Inotuzumab ozogamicin	214
healthE Calamine Aqueous		neomycin70 Inresa	35
healthE Dimethicone 10%	70	Hydrogen peroxide67 Inspra	52
healthE Dimethicone 4% Lotion	72	Hydroxocobalamin32 Instillagel Lido	
healthE Dimethicone 5%	70	Hydroxocobalamin Panpharma32 Insulin aspart	
healthE Glycerol BP	. 282	hydroxycarbamide156 Insulin aspart with insulin aspart	
healthE Urea Cream		Hydroxychloroquine sulphate116 protamine	10

Insulin degludec with insulin	Iron (as ferric carboxymaltose)	34	Klaricid	94
aspart10	Iron polymaltose	35	Kliogest	
Insulin glargine11	Isentress	111	Kliovance	
Insulin glulisine11	Isentress HD	111	Kogenate FS	39
Insulin isophane10	Ismo 20	55	Konakion MM	40
Insulin isophane with insulin	Ismo 40 Retard	55	Konakion MM Paediatric	
neutral11	Isoleucine50	300	Konsyl-D	24
Insulin lispro11	Isoniazid	104	Kuvan	29
Insulin lispro with insulin lispro	Isoniazid Teva	104	-L-	
protamine11	Isoniazid with rifampicin	104	Labetalol	49
Insulin neutral10	Isoptin	51	Lacosamide	129
Insulin pen needles16	Isoptin Retard	51	Lactulose	25
Insulin pump cartridge18	Isoptin SR	51	Lacuna	28
Insulin pump infusion set (steel	Isopto Carpine		Laevolac	25
cannula)18	Isosorbide mononitrate	55	Lamictal	130
Insulin pump infusion set (steel	Isotretinoin	67	Lamivudine10	05, 111
cannula, straight insertion) 19	Ispaghula (psyllium) husk	24	Lamivudine Viatris	111
Insulin pump infusion set (teflon	Itch-Soothe	69	Lamivudine/Zidovudine Viatris	111
cannula)20	Itraconazole	100	Lamotrigine	130
Insulin pump infusion set (teflon	Itraconazole Cresent	100	Lamprene	103
cannula, angle insertion with	Itraconazole Kent	100	Lanoxin	
insertion device)21	Itrazole		Lanoxin PG	48
Insulin pump infusion set (teflon	lvacaftor		Lanoxin S29	48
cannula, flexible insertion with	Ivermectin		Lanreotide	
insertion device)21	- J -		Lansoprazole	
Insulin pump infusion set (teflon	Jadelle	78	Lantus	
cannula, straight insertion with	Jakavi		Lantus SoloStar	
insertion device)21	Jardiamet		Lanvis	
Insulin pump infusion set (teflon	Jardiance		Lanzol Relief	
cannula, variable insertion) 21	Jaydess		Largactil	
Insulin pump reservoir22	Jevity HiCal RTH		Laronidase	
Insulin pump with algorithm17	Jevity RTH		Lasix	
Insulin syringes, disposable with	Jinarc		Latanoprost	
attached needle17	Juno Pemetrexed		Latanoprost with timolol	
Intelence110	- K -	104	Lax-Suppositories	
Interferon beta-1-alpha139	Kadcyla	242	Lax-suppositories Glycerol	
Interferon beta-1-beta	Kalydeco		Laxatives	
Intra-uterine device	Kemadrin		Laxsol	
Invega Sustenna	Kenacomb		Ledipasvir with sofosbuvir	
Invega Trinza137	Kenacort-A 10		Leflunomide	
Ipca-Allopurinol118	Kenacort-A 40		Lenalidomide (Viatris)	
lpca-Bisoprolol	Kenalog in Orabase		Lenalidomide Viatris	
lpca-Ciprofloxacin97	Ketocal 3:1		Lenvatinib	
lpca-Donepezil	KetoCal 4:1		Lenvima	
lpca-Escitalopram128	Ketoconazole		Letrole	
IPCA-Frusemide	Dermatological	7/	Letrozole	
lpca-Hydroxychloroquine116	Infection		Leucine100	
			Leukeran FC	
IPCA-Metoprolol	Ketogenic Diet Ketoprofen		Leukotriene Receptor	101
Ipilimumab248	KetoSens		Antagonists	271
IPOL	Keytruda		Leuprorelin	
Ipratropium bromide267, 273	Kindergen		Leustatin	
Iressa			Levetiracetam	
Irinotecan Actavis 100	Kisqali	171	Levetiracetam-AFT	100 100
	Klacid	0	Levetiracetam-AFT	
Irinotecan hydrochloride	Alimentary		Levocapastine	
Irinotecan-Rex153	Infection	94	Levocarriune	28

Levodopa with benserazide		Loxamine		Metabolics	
Levodopa with carbidopa	121	Lucrin Depot 1-month		Metformin hydrochloride	
Levodopa with carbidopa and		Lucrin Depot 3-month		Metformin Viatris	11
entacapone		LumaCina	44	Methadone BNM	
Levomepromazine	134	Lumigan	276	Methadone hydrochloride	125
Levomepromazine		Lynparza	158	Methenamine (hexamine)	
hydrochloride	134	Lyrica		hippurate	114
Levonorgestrel		- M -		Methopt	
Levonorgestrel BNM		m-Eslon	126	Methotrexate	
Levothyroxine		Mabthera		Methotrexate DBL Onco-Vial	
Lidocaine [Lignocaine]1		Macrobid		Methotrexate Ebewe	
Lidocaine [Lignocaine]	122 120	Macrogol 3350 with potassium	1 1 4	Methotrexate Sandoz	
hydrochloride	122	chloride, sodium bicarbonate ar	nd	Methyl hydroxybenzoate	
Lidocaine [Lignocaine] with	120			Methylcellulose	
	100	sodium chloride			202
prilocaine		Madopar 125		Methylcellulose with glycerin and	000
Lidocaine-Baxter		Madopar 250		sodium saccharin	282
Life Extension		Madopar 62.5		Methylcellulose with glycerin and	
Lignocaine1		Madopar HBS		sucrose	
Linezolid		Madopar Rapid	121	Methyldopa	
Lioresal Intrathecal		Magnesium hydroxide		Methyldopa Viatris	
Lipid-Modifying Agents	53	Magnesium sulphate	35	Methylnaltrexone bromide	25
Liquigen	285	Mantoux	322	Methylphenidate ER - Teva	
Liraglutide	12	Marevan	43	Methylphenidate hydrochloride	144
Lisdexamfetamine dimesilate	143	Marine Blue Lotion SPF 50+	75	Methylphenidate hydrochloride	
Lisinopril		Mask for spacer device	273	extended-release	145
Lithium carbonate		Maviret		Methylphenidate Sandoz XR	
Livostin		Maxidex		Methylprednisolone	
LMX4		Maxitrol		Methylprednisolone (as sodium	
Lo-Oralcon 20 ED		MCT oil (Nutricia)		succinate)	84
Locacorten-Viaform ED's		Measles, mumps and rubella	200	Methylprednisolone aceponate	
Local preparations for Anal and	214	vaccine	215		
	0			Methylprednisolone acetate	05
Rectal Disorders		Mebendazole		Methylxanthines	
Locasol		Mebeverine hydrochloride		Metoclopramide Actavis 10	
Locoid		Medac		Metoclopramide hydrochloride	
Locoid Crelo		Medrol	83	Metolazone	
Locoid Lipocream		Medroxyprogesterone acetate		Metopirone	
Locorten-Vioform		Genito-Urinary	78	Metoprolol IV Mylan	
Lodoxamide		Hormone	85–86	Metoprolol IV Viatris	49
Logem	130	Mefenamic acid	115	Metoprolol succinate	49
Lomide		Mekinist	173	Metoprolol tartrate	
Lomustine	152	Melatonin	140	Metronidamed	103
Loniten	56	Melpha	152	Metronidazole	103
Loperamide hydrochloride	6	Melphalan		Metyrapone	91
Lopinavir with ritonavir		Meningococcal (groups A, C, Y an		Mexiletine hydrochloride	48
Lopinavir/Ritonavir Mylan	111	W-135) conjugate vaccine	316	Miacalcic	
Loprofin		Meningococcal B multicomponent		Miacalcic S29	
Loprofin Mix		vaccine	317	Micolette	
Lorafix		MenQuadfi		Miconazole	
Loratadine		Menthol		Miconazole nitrate	
Lorazepam					60
		Mepolizumab		Dermatological	
Lorstat		Mercaptopurine		Genito-Urinary	
Losartan Actavis		Mercilon 28		Micreme	
Losartan potassium	4/	Mesalazine		Micreme H	
Losartan potassium with		Mesna		Microlut	
hydrochlorothiazide		Mestinon		Midazolam	
Lovir	105	Metabolic Disorder Agents	26	Midazolam-Baxter	141

Midazolam-Pfizer	141	Modafinil Max Health	146	Naglazyme	2
Midodrine	49	Moduretic	52	Naloxone hydrochloride	27
Midostaurin	168	Molaxole	25	Naltraccord	14
Mifegyne	81	Moments	<mark>77</mark>	Naltrexone hydrochloride	14
Mifepristone		Mometasone furoate	70	Naphazoline hydrochloride	
Minerals		Monogen	286	Naprosyn SR 1000	
Mini-Wright AFS Low Range		Montelukast		Naprosyn SR 750	
Mini-Wright Standard		Montelukast Viatris		Naproxen	11
Minidiab		Moroctocog alfa [Recombinant fa		Narcaricin mite	
MiniMed 3.0 Reservoir		VIII]		Nasal Preparations	
MMT-332A	22	Morphine hydrochloride		Natalizumab	
MiniMed Mio MMT-921A		Morphine sulphate		Natulan	
MiniMed Mio MMT-923A		Motetis		Nausafix	
MiniMed Mio MMT-925A		Mouth and Throat		Nausicalm	
MiniMed Mio MMT-941A		Movapo		Navelbine S29	
MiniMed Mio MMT-943A		Movicol		Nefopam hydrochloride	
				Neo-Mercazole	
MiniMed Mio MMT-945A		Moxifloxacin			
MiniMed Mio MMT-965A		MSUD Anamix Infant		Neocate Gold	
MiniMed Mio MMT-975A		MSUD Anamix Junior		Neocate Junior Unflavoured	
MiniMed Quick-Set MMT-396A		MSUD Anamix Junior LQ		Neocate Junior Vanilla	
MiniMed Quick-Set MMT-397A		MSUD Explore 5		Neocate SYNEO	
MiniMed Quick-Set MMT-398A		MSUD Express 15	296	Neoral	
MiniMed Quick-Set MMT-399A		MSUD Lophlex LQ 20		Neostigmine metilsulfate	
MiniMed Silhouette MMT-377A		MSUD Maxamum		Nepro HP (strawberry)	
MiniMed Silhouette MMT-378A		Mucolytics		Nepro HP (vanilla)	
MiniMed Silhouette MMT-381A		Mucosoothe		Neulactil	
MiniMed Sure-T MMT-864A	18	Multiple Sclerosis Treatments	138	NeuroTabs	3
MiniMed Sure-T MMT-866A		Multivitamin renal	33	Nevirapine	
MiniMed Sure-T MMT-874A	18	Multivitamins		Nevirapine Viatris	11
MiniMed Sure-T MMT-876A	18	Mupirocin	67	Nicorandil	5
Minims Pilocarpine	277	Muscle Relaxants	119	Nicotine	14
Minims Prednisolone	276	Mvite	33	Nifedipine	5
Minipress	46	Myambutol	104	Nifuran	11
Minirin		Mycobutin		Nilotinib	16
Minirin Melt	91	MycoNail		Nilstat	
Mino-tabs	96	Mycophenolate mofetil		Alimentary	3
Minocycline hydrochloride		Mydriacyl		Genito-Urinary	
Minomycin		Mylan (24 hr release)		Infection	
Minor Skin Infections		Mylan Clomiphen		Nimenrix	
Minoxidil		Mylan Italy (24 hr release)		Nintedanib	
Minoxidil Roma		Myleran		Nipent	
Mirena		mylife Inset soft		Niraparib	
Miro-Amoxicillin		mylife Orbit micro		Nirmatrelvir with ritonavir	
Mirtazapine		mylife YpsoPump Reservoir		Nitrates	
Misoprostol		mylife YpsoPump with CamAPS		Nitroderm TTS	
Mitomycin (Fresenius Kabi)		FX		Nitrofurantoin	
		Myloc CR		Nitrolingual Pump Spray	
Mitomycin (Sagent)		Mulatara	48		
Mitomycin C		Mylotarg		Nivestim	
Mitozantrone		Myometrial and Vaginal Hormon		Nivolumab	
Mitozantrone Ebewe		Preparations		Nodia	
MMA/PA Anamix Infant		Myozyme		Noflam 250	
MMA/PA Anamix Junior		Mytolac		Noflam 500	11
MMA/PA Explore 5		Mytolac S29	177	Non-Steroidal Anti-Inflammatory	
MMA/PA Express 15		- N -		Drugs	11
Moclobemide		Nadolol		Nonacog gamma, [Recombinant	
Modafinil	146	Nadolol BNM	50	Factor IX]	3

Norethinderone - CDC	70	- 0 -	Orphenadrine citrate	120
Norethisterone	70	Obinutuzumab21	•	
Genito-Urinary	79	Obstetric Preparations8		
Hormone		Ocicure		
Norflex		Ocrelizumab13		
Norfloxacin		Ocrevus		
Noriday		Ocrevus SC13	0 1	80
Noriday 28		Octasa		0/
Norimin		Octocog alfa [Recombinant factor	Preparations	
Normison		VIII] (Advate)		/
Norpress		Octocog alfa [Recombinant factor	Ovestin	
Nortriptyline hydrochloride		VIII] (Kogenate FS)3		
Norvir		Octreotide 17		
Noumed Dexamfetamine		Octreotide GH17		
Noumed Isoniazid	104	Octreotide long-acting17		152
Noumed Paracetamol	124	Oestradiol8		
Noumed Pethidine	127	Oestradiol valerate8		
Noumed Phenobarbitone	130	Oestradiol with norethisterone8	6 Oxis Turbuhaler	266
NovaSource Renal	289	Oestriol	Oxpentifylline	56
Novatretin	73	Genito-Urinary7		
Novitium Sugar Free	28	Hormone8		
NovoMix 30 FlexPen		Oestrogens8		
NovoRapid		Ofev26	9 Oxycodone Lucis	126
NovoRapid FlexPen		Oil in water emulsion7		
NovoRapid Penfill		Olanzapine134, 13		
NovoSeven RT		Olaparib15		
Nozinan		Olbetam5		
Nozinan (Swiss)		Olopatadine	,	80
1402111a11 (OW100)		Olopatadirio		
Mucala	215	Olonatadina Tava 27	8 Ozurdov	279
Nucala		Olopatadine Teva27		275
Nuelin	271	Olsalazine	8 - P -	
Nuelin Nuelin-SR	271 271	Olsalazine Omalizumab21	8 - P - 7 Pacifen	119
Nuelin	271 271 129	OlsalazineOmalizumab21 Omeprazole	8 - P - 7 Pacifen9 Pacimol	119 124
Nuelin	271 271 129 141	Olsalazine 21 Omalizumab 21 Omeprazole 21 Omeprazole actavis 10 21	8 - P - 7 Pacifen 9 Pacimol 9 Paclitaxel	119
Nuelin	271 271 129 141 295	Olsalazine 21 Omalizumab 21 Omeprazole 21 Omeprazole actavis 10 21 Omeprazole actavis 20 22	P - Pacifen Pacimol Paclitaxel Paclitaxel Actavis	119 124 159
Nuelin	271 129 141 295 286	Olsalazine 21 Omalizumab 21 Omeprazole 21 Omeprazole actavis 10 21 Omeprazole actavis 20 21 Omeprazole actavis 40 22	P - Pacifen Pacimol Paclitaxel Paclitaxel Actavis Paclitaxel Ebewe	119 124 159
Nuelin	271 271 129 141 295 286 283	Olsalazine	P - Pacifen	119 124 159 159
Nuelin	271 129 141 295 286 283	Olsalazine 21 Omalizumab 21 Omeprazole 21 Omeprazole actavis 10 21 Omeprazole actavis 20 21 Omeprazole actavis 40 22 Omeprazole Teva 23 Omnitrope 8	P - P - P - P - P - P - P - P - P - P -	119 124 159 159
Nuelin	271 129 141 295 286 283 288	Olsalazine 21 Omalizumab 21 Omeprazole 20 Omeprazole actavis 10 20 Omeprazole actavis 20 20 Omeprazole actavis 40 20 Omeprazole Teva 20 Omnitrope 8 Omnitrope AU 8	P - P - P - P - P - P - P - P - P - P -	119 159 159 159 159
Nuelin	271 271 129 141 295 286 283 288 288	Olsalazine 21 Omalizumab 21 Omeprazole 20 Omeprazole actavis 10 20 Omeprazole actavis 40 20 Omeprazole Teva 20 Omnitrope 8 Omnitrope AU 8 Onbrez Breezhaler 26	P - P - P - P - P - P - P - P - P - P -	119 159 159 159 159 52
Nuelin	271 271 129 141 295 286 283 288 288 290 303	Olsalazine 21 Omalizumab 21 Omeprazole 20 Omeprazole actavis 10 20 Omeprazole actavis 20 20 Omeprazole actavis 40 20 Omeprazole Teva 20 Omnitrope 8 Omnitrope AU 8 Onbrez Breezhaler 26 Oncaspar LYO 15	P - P - P - P - P - P - P - P - P - P -	119 159 159 159 159
Nuelin	271271129141295286283288288288290303	Olsalazine 21 Omalizumab 21 Omeprazole 20 Omeprazole actavis 10 20 Omeprazole actavis 20 20 Omeprazole Teva 20 Omnitrope 8 Omnitrope AU 8 Onbrez Breezhaler 26 Oncaspar LYO 15 OncoTICE 18	P - P - Pacifen	119 159 159 159
Nuelin	271271129141295286283288288288290303	Olsalazine 21 Omalizumab 21 Omeprazole 21 Omeprazole actavis 10 21 Omeprazole actavis 20 22 Omeprazole actavis 40 23 Omeprazole Teva 24 Omnitrope 8 Omnitrope AU 8 Onbrez Breezhaler 26 Oncaspar LYO 15 OncoTICE 18 Ondansetron 13	P - P - P - P - P - P - P - P - P - P -	119 159 159 159
Nuelin	271271129141295286288288288290303303288	Olsalazine 21 Omalizumab 21 Omeprazole 20 Omeprazole actavis 10 20 Omeprazole actavis 20 20 Omeprazole Teva 20 Omnitrope 8 Omnitrope 8 Ombrez Breezhaler 26 Oncaspar LYO 15 OncoTICE 18 Ondansetron 13 One-Alpha 3	P - P - Pacifen	119 159 159 159
Nuelin	271271271129141295286288288290303303288	Olsalazine 21 Omalizumab 21 Omeprazole 20 Omeprazole actavis 10 20 Omeprazole actavis 20 20 Omeprazole actavis 40 20 Omeprazole Teva 20 Omitrope 8 Omitrope 8 Onbrez Breezhaler 26 Oncaspar LYO 15 OncoTICE 18 Ondansetron 13 One-Alpha 3 Opdivo 24	P - P - P - P - P - P - P - P - P - P -	
Nuelin	271271271129141295286288288290303303288293	Olsalazine 21 Omalizumab 21 Omeprazole 20 Omeprazole actavis 10 20 Omeprazole actavis 20 20 Omeprazole actavis 40 20 Omnitrope 8 Omnitrope 8 Onnitrope AU 8 Onbrez Breezhaler 26 Oncaspar LYO 15 OncoTICE 18 Ondansetron 13 One-Alpha 3 Opdivo 24 Ora-Blend 28	P - P - P - P - P - P - P - P - P - P -	
Nuelin	271271271129141295286288288290303303288293	Olsalazine 21 Omalizumab 21 Omeprazole 20 Omeprazole actavis 10 20 Omeprazole actavis 20 20 Omeprazole actavis 40 20 Omnitrope 8 Omnitrope 8 Onnitrope AU 8 Onbrez Breezhaler 26 Oncaspar LYO 15 OncoTICE 18 Ondansetron 13 One-Alpha 3 Opdivo 24 Ora-Blend 28 Ora-Blend SF 28	P - P - P - P - P - P - P - P - P - P -	1191241581585233168169137117
Nuelin	271271291129141295286283288290303288292	Olsalazine 21 Omalizumab 21 Omeprazole 20 Omeprazole actavis 10 20 Omeprazole actavis 20 20 Omeprazole actavis 40 20 Omnitrope 8 Omnitrope 8 Onnitrope AU 8 Onbrez Breezhaler 26 Oncaspar LYO 15 OncoTICE 18 Ondansetron 13 One-Alpha 3 Opdivo 24 Ora-Blend 28	P - P - P - P - P - P - P - P - P - P -	119124158158
Nuelin	271271291129141295286283288290303288292	Olsalazine 21 Omalizumab 21 Omeprazole 20 Omeprazole actavis 10 20 Omeprazole actavis 40 20 Omeprazole Teva 20 Omnitrope 8 Omnitrope AU 8 Onbrez Breezhaler 26 Oncaspar LYO 15 OncoTICE 18 Ondansetron 13 Opcivo 24 Ora-Alpha 3 Opdivo 24 Ora-Blend 28 Ora-Blend SF 28 Ora-Plus 28 Ora-Sweet 28	P - P - P - P - P - P - P - P - P - P -	119124158158
Nuelin	271271271129141295286288288290303288288293288293289292293	Olsalazine 21 Omalizumab 21 Omeprazole 20 Omeprazole actavis 10 20 Omeprazole actavis 20 20 Omeprazole actavis 40 20 Omeprazole Teva 20 Omnitrope 8 Omnitrope AU 8 Onbrez Breezhaler 26 Oncaspar LYO 15 OncoTICE 18 Ondansetron 13 One-Alpha 3 Opdivo 24 Ora-Blend 28 Ora-Plus 28	P - P - P - P - P - P - P - P - P - P -	11812915915952
Nuelin	271271271295286283288290303303288292293293293	Olsalazine 21 Omalizumab 21 Omeprazole 20 Omeprazole actavis 10 20 Omeprazole actavis 40 20 Omeprazole Teva 20 Omnitrope 8 Omnitrope AU 8 Onbrez Breezhaler 26 Oncaspar LYO 15 OncoTICE 18 Ondansetron 13 Opcivo 24 Ora-Alpha 3 Opdivo 24 Ora-Blend 28 Ora-Blend SF 28 Ora-Plus 28 Ora-Sweet 28	P - P - P - P - P - P - P - P - P - P -	11812915915952
Nuelin	271271291129141295286288288290303288293289293293293	Olsalazine 21 Omalizumab 21 Omeprazole 20 Omeprazole actavis 10 20 Omeprazole actavis 40 20 Omeprazole Teva 20 Omnitrope 8 Omnitrope AU 8 Onbrez Breezhaler 26 Oncaspar LYO 15 OncoTICE 18 Ondansetron 13 One-Alpha 3 Opdivo 24 Ora-Blend 28 Ora-Blend SF 28 Ora-Plus 28 Ora-Sweet 28 Ora-Sweet SF 28	P - P - P - P - P - P - P - P - P - P -	118124155158
Nuelin	271271291129141295286288288290303288293289293293293	Olsalazine 21 Omalizumab 21 Omeprazole 21 Omeprazole actavis 10 20 Omeprazole actavis 20 20 Omeprazole Teva 20 Omnitrope 8 Omnitrope AU 8 Onbrez Breezhaler 26 Oncaspar LYO 15 OncoTICE 18 Ondansetron 13 One-Alpha 3 Opdivo 24 Ora-Blend 28 Ora-Plus 28 Ora-Sweet 28 Ora-Sweet SF 28 Orabase 3	P - P - P - P - P - P - P - P - P - P -	118124155158
Nuelin	271271271295286288298298298298299303303288293289293295293293293	Olsalazine 21 Omalizumab 21 Omeprazole 20 Omeprazole actavis 10 0 Omeprazole actavis 20 0 Omeprazole actavis 40 0 Omeprazole Teva 0 Omnitrope 8 Onnitrope AU 8 Oncaspar LYO 15 OncoTICE 18 Ondansetron 13 One-Alpha 3 Opdivo 24 Ora-Blend 28 Ora-Plus 28 Ora-Sweet 28 Ora-Sweet SF 28 Orabase 3 Oral and Enteral Feeds 28 Oralcon 30 ED 7	P - P - P - P - P - P - P - P - P - P -	119124155
Nuelin	271271271295286288288290303303288293295293293293293293	Olsalazine 21 Omalizumab 21 Omeprazole 21 Omeprazole actavis 10 20 Omeprazole actavis 20 20 Omeprazole actavis 40 20 Omeprazole Teva 20 Omnitrope 8 Omnitrope 8 Onbrez Breezhaler 26 Oncaspar LYO 15 OncoTICE 18 Ondansetron 13 One-Alpha 3 Opdivo 24 Ora-Blend 28 Ora-Plus 28 Ora-Sweet 28 Ora-Sweet SF 28 Orabase 3 Oral and Enteral Feeds 28 Oral and Enteral Feeds 28 Oramorph 12	P - P - P - P - P - P - P - P - P - P -	119124155
Nuelin	271271271295286288288293303288293293293293293293293293	Olsalazine 21 Omalizumab 21 Omeprazole 20 Omeprazole actavis 10 0 Omeprazole actavis 20 0 Omeprazole actavis 40 0 Omeprazole Teva 0 Omnitrope 8 Onnitrope AU 8 Oncaspar LYO 15 OncoTICE 18 Ondansetron 13 One-Alpha 3 Opdivo 24 Ora-Blend 28 Ora-Plus 28 Ora-Sweet 28 Ora-Sweet SF 28 Orabase 3 Oral and Enteral Feeds 28 Oralcon 30 ED 7	P - P - P - Pacifen	119124155
Nuelin	271271271295286288288290303303288293293293293293293293293293	Olsalazine 21 Omalizumab 21 Omeprazole 20 Omeprazole actavis 10 20 Omeprazole actavis 20 20 Omeprazole actavis 40 20 Omeprazole Teva 20 Omnitrope 8 Omnitrope AU 8 Onbrez Breezhaler 26 Oncaspar LYO 15 OncoTICE 18 Ondansetron 13 One-Alpha 3 Opdivo 24 Ora-Blend 28 Ora-Plus 28 Ora-Sweet 28 Ora-Sweet SF 28 Oral and Enteral Feeds 28 Oral and Enteral Feeds 28 Oramorph 12 Oramorph CDC S29 12	P - P - Pacifen	118124158

Paracetamol with codeine	126	Nervous	141	Plendil ER	50
Paraffin	71	Phenoxybenzamine		Pneumococcal (PCV13) conjugate	
Paraffin liquid with wool fat	278	hydrochloride	46	vaccine	318
Parasiticidal Preparations	72	Phenoxymethylpenicillin (Penic	cillin	Pneumococcal (PPV23)	
Parnate		V)		polysaccharide vaccine	319
Paromomycin	98	Phenylalanine50		Pneumovax 23	319
Paroxetine	128	Phenytoin sodium	129–130	Podophyllotoxin	75
Paser	104	Phesgo	220	Polaramine	265
Paxam	137	Phillips Milk of Magnesia	35	Poliomyelitis vaccine	319
Paxlovid	109	Phlexy 10		Poloxamer	
Paxlovid fee	279	Phosphate Phebra	45	Poly-Tears	277
Paxtine	128	Phosphorus	45	Poly-Visc	
Pazopanib	170	Phytomenadione		Polycal	
Pazopanib Teva	170	Pilocarpine hydrochloride	277	Polyethylene glycol 400 and	
Peak flow meter		Pilocarpine nitrate		propylene glycol	278
Pediasure	288	Pimafucort		Pomalidomide	
Pediasure Plus	288	Pimecrolimus	73	Pomolide	
Pediasure RTH	288	Pine tar with trolamine laurilsul	lfate	Ponstan	115
Pegaspargase	159	and fluorescein	74	Posaconazole	100
Pegasys		Pinetarsol	74	Posaconazole Juno	100
Pegasys (S29)		Pioglitazone		Potassium chloride4	14-45
Pegasys S29		Pirfenidone		Potassium citrate	81
Pegfilgrastim		Pizotifen		Potassium iodate	34
Pegylated interferon alfa-2a.		PKU Anamix Infant	297	Povidone iodine	
Pembrolizumab		PKU Anamix Junior		Pradaxa	
Pemetrexed	154	PKU Anamix Junior Chocolate	297	Pramipexole hydrochloride	
Pemetrexed-AFT		PKU Anamix Junior LQ		Pravastatin	
Penicillamine	116	PKU Anamix Junior Orange		Praziquantel	
Penicillin G		PKU Anamix Junior Vanilla		Prazosin	
PenMix 30		PKU Build 10		Prazosin Mylan	
Pentasa		PKU Build 20 Chocolate		Pred Forte	
Pentostatin [Deoxycoformyci		PKU Build 20 Raspberry		Prednisolone	
Pentoxifylline [Oxpentifylline]	-	Lemonade	298	Prednisolone acetate	
Peptamen Junior		PKU Build 20 Smooth		Prednisolone sodium	
Pepti-Junior		PKU Build 20 Vanilla		phosphate	. 276
Perhexiline maleate		PKU Explore 10		Prednisolone-AFT	
Pericyazine		PKU Explore 5		Prednisone	
Perindopril		PKU Express 20		Prednisone Clinect	
Periset		PKU First Spoon		Pregabalin	
Periset ODT		PKU Glytactin RTD 15		Pregabalin Pfizer	
Perjeta		PKU Glytactin RTD 15 Lite		Pregnancy Tests - hCG Urine	
Permethrin		PKU GMPro LQ		Premarin	
Perrigo		PKU GMPro Mix-In		Presolol	
Pertuzumab		PKU GMPro Ultra Lemonade		Prevenar 13	
Pertuzumab with trastuzuma		PKU GMPro Ultra Vanilla		Priadel	
Peteha		PKU Lophlex LQ 10		Primaquine	
Pethidine hydrochloride		PKU Lophlex LQ 20		Primidone	
Pevaryl		PKU Lophlex Powder		Primidone Clinect	
Pexsig		PKU Lophlex Sensation 20		Primolut N	
Pfizer Exemestane		PKU Restore Powder		Priorix	
Pharmacy Services		PKU sphere20 Banana		Probenecid	
Pheburane		PKU sphere20 Chocolate		Probenecid-AFT	119
Phenasen		PKU sphere20 Lemon		Procarbazine hydrochloride	
Phenobarbitone		PKU sphere20 Red Berry		Prochlorperazine	
Phenobarbitone sodium		PKU sphere20 Vanilla		Prochlorperazine maleate (Brown &	
Extemporaneous	282	Pku Start		Burk)	
poranoodo		Own		- vy	

Proctofoam	7	Relistor	25	Rosuvastatin-Sandoz	54
Proctosedyl	8	Remdesivir		Rotarix	320
Procyclidine hydrochloride	121	Remicade	206	Rotavirus oral vaccine	320
Progesterone		Renilon 7.5	289	Roxane-Propranolol	50
Proglicem		Resonium-A	45	Roxithromycin	
Progynova		Resource Beneprotein	285	Rozlytrek	
Prolia		Respiratory Devices		Rubifen	
Promethazine hydrochloride		Respiratory Stimulants		Rubifen SR	
Propafenone hydrochloride		Retinol palmitate		Rurioctocog alfa pegol [Recombi	
Propranolol		ReTrieve		factor VIII]	
Propylene glycol		Retrovir		Ruxolitinib	
Propylthiouracil		Revolade		Rydapt	
Prostacur		Ribociclib		Rythmodan - Cheplafarm	
Protaphane		Riboflavin		Rythmodan Neon	
Protaphane Penfill		Ribomustin		Rytmonorm	
Protifar		Ricit		Ryzodeg 70/30 Penfill	
Protionamide		Ricovir		- S -	
				-	101
Provera LID		Rifabutin		Sabril	
Provera HD	00	Rifadin		Sacubitril with valsartan	
Psoriasis and Eczema	70	Rifadin Sanofi		SalAir	
Preparations		Rifamazid		Salazopyrin	ک
PTU		Rifampicin		Salazopyrin EN	ک
Pulmicort Turbuhaler		Rifaximin		Salbutamol	267
Pulmozyme		Rifinah		Salbutamol with ipratropium	
Puri-nethol	154	Rilutek		bromide	
Puritan's Pride Vitamin		Riluzole		Salicylic acid	
B-2 100 mg		Rinvoq		Salmeterol	
Pyrazinamide		Riodine		Sandomigran	
Pyridostigmine bromide		Risdiplam	142	Sandostatin LAR	
Pyridoxine hydrochloride		Risedronate Sandoz		Sanofi Primaquine	
Pyridoxine multichem		Risedronate sodium	118	Sapropterin dihydrochloride	29
Pytazen SR	40	Risperdal	135	Scalp Preparations	74
- Q -		Risperdal Consta	137	Scopolamine Transdermal Syste	em
Quantalan sugar free	54	Risperidone		Viatris	133
Quetapel	135	Risperidone (Teva)	135	Sebizole	74
Quetiapine	135	Risperon		Secukinumab	234
Quetiapine Viatris	135	Ritalin	144	Sedatives and Hypnotics	140
Quinapril	46	Ritalin LA	145	Seebri Breezhaler	268
Qvar	265	Ritonavir	111	Senna	25
- R -		Rituximab (Mabthera)	221	Senokot	25
RA-Morph	125	Rituximab (Riximyo)	223	SensoCard	16
Ralicrom		Rivaroxaban		Serc	132
Raloxifene hydrochloride	117	Rivastigmine	146	Serenace	
Raltegravir potassium		Rivastigmine Patch BNM 10.		Seretide	267
Ramipex		Rivastigmine Patch BNM 5		Seretide Accuhaler	
Ramipril		Rivotril		Serevent	
Ranbaxy-Cefaclor		Riximyo		Serevent Accuhaler	
Rapamune	258	RIXUBIS		Sertraline	
Rasagiline		Rizamelt		Setrona	
Reandron 1000		Rizatriptan		Sevredol	
Recombinant factor IX		Robinul		Sex Hormones Non	120
Recombinant factor VIIa		Ropin		Contraceptive	Q/
Recombinant factor VIII		Ropinirole hydrochloride		Shingles vaccine	320
				Shingrix	200
Rectogesic		Rosemont		SII Open BCG	105
Redipred		Rosuvastatin		SII-Onco-BCG	
Relieve	115	Rosuvastatin Viatris	54	Sildenafil	

Siltuximab	236	Spinraza	141	Tambocor	48
Simvastatin	55	Spiolto Respimat	268	Tambocor German	
Simvastatin Mylan	55	Spiractin	52	Tamoxifen citrate	176
Simvastatin Viatris	55	Spiriva	268	Tamoxifen Sandoz	170
Sinemet	121	Spiriva Respimat	268	Tamsulosin hydrochloride	
Sinemet CR	121	Spironolactone		Tamsulosin-Rex	80
Sintetica Baclofen Intrathecal	119	Stalevo	121	Tandem Cartridge	18
Sirolimus	258	Staphlex	95	Tandem t:slim X2 with Basal-IQ	1
Sirturo	103	Stelara	243	Tandem t:slim X2 with	
Siterone	84	Stemetil		Control-IQ+	
Slow-Lopresor	49	Steril-Gene	84	Tap water	282
Sodibic	45	SteroClear	272	Taro	100
Sodium acid phosphate	25	Stesolid		Tasigna	168
Sodium alginate	6	Stimulants/ADHD Treatments	143	Tasmar	12 ⁻
Sodium benzoate	30	Stiripentol	131	Taurine	30
Sodium bicarbonate		Stomahesive	31	TCu 380 Plus Normal	
Blood	. 44–45	Strides Shasun		Tecentriq	
Extemporaneous	282	Stromectol	72	Tecfidera	139
Sodium calcium edetate	280	Sucralfate	9	TEEVIR	11
Sodium chloride		Sulfadiazin-Heyl	98	Tegretol	129
Blood	44	Sulfadiazine Silver	68	Tegretol CR	
Respiratory	272	Sulfadiazine sodium	98	Telfast	26
Sodium citrate with sodium lauryl		Sulfasalazine	8	Temaccord	160
sulphoacetate	25	Sulphur	74	Temazepam	14
Sodium citro-tartrate	81	Sulprix	133	Temozolomide	160
Sodium cromoglicate		Sumagran	132	Temozolomide-Taro	160
Alimentary	8	Sumatriptan	132	Tenofovir disoproxil	10
Sensory	276	Sunitinib	172	Tenofovir Disoproxil Emtricitabine	
Sodium Fusidate [fusidic acid]		Sunitinib Pfizer	172	Viatr	10
Dermatological	68	Sunitinib Rex	172	Tenofovir Disoproxil Viatris	10
Infection	98	Sunscreens	75	Tenoxicam	11
Sensory	274	Sunscreens, proprietary	75	Tensipine MR10	5
Sodium hyaluronate [Hyaluronic		Sustagen Hospital Formula	293	Tepadina	
acid]	278	Sustanon Ampoules	84	Terbinafine	
Sodium phenylbutyrate	30	Sylvant	236	Terbutaline sulphate	26
Sodium picosulfate	26	Symbicort Turbuhaler 100/6	266	Teriflunomide	139
Sodium polystyrene sulphonate.	45	Symbicort Turbuhaler 200/6	266	Teriflunomide Sandoz	139
Sodium tetradecyl sulphate	39	Symbicort Turbuhaler 400/12	266	Teriparatide	118
Sodium valproate		Symmetrel	121	Teriparatide - Teva	118
Sofradex	274	Sympathomimetics		Testogel	84
Soframycin	274	Synacthen	84	Testosterone	84
Solax	24	Synacthen Depot		Testosterone cipionate	84
Solgar	26-30	Synacthene Retard	84	Testosterone esters	84
Solifenacin succinate	81	Synagis		Testosterone undecanoate	84
Solifenacin succinate Max		Synthroid	86	Tetrabenazine	122
Health	81	Syntometrine		Tetrabromophenol	8
Solu-Cortef	83	Syrup (pharmaceutical grade)	282	Tetracosactrin	
Solu-Medrol	84	Systane Unit Dose		Tetracycline	96
Solu-Medrol-Act-O-Vial	84	· -Т-		Teva Lisinopril	40
Somatropin (Omnitrope)		Tacrolimus		Teva-Ketoconazole	10
Sotalol		Dermatological	74	Thalidomide	
Sotalol Viatris	50	Oncology		Thalomid	16 ⁻
Spacer device	273	Tacrolimus Sandoz	260	Theophylline	27
Span-K		Tafinlar	164	Thiamine hydrochloride	32
Spazmol	8	Tagrisso	168	Thiamine multichem	
Spinal Muscular Atrophy		Taliglucerase alfa		THIO-TEPA	152

Thioguanine		gramicidin, neomycin and r		Valine50	
Thiotepa		Dermatological	70	Vancomycin	
Thyroid and Antithyroid Agents .	86	Sensory	274	Vancomycin Viatris	9
Ticagrelor	40	Trientine	30	Vannair	26
Ticagrelor Sandoz	40	Trientine Waymade	30	Varenicline tartrate	149
Tilcotil	115	Trikafta	271	Varicella vaccine [Chickenpox	
Timolol		Trimethoprim	99	vaccine]	32
Tiotropium bromide	268	Trimethoprim with		Varicella zoster vaccine [Shingles	
Tiotropium bromide with		sulphamethoxazole		vaccine]	32
olodaterol	268	[Co-trimoxazole]	99	Varilrix	32
Tivicay	111	Triovir	111	Various	27
TMP	99	Trisequens	86	VariSoft	2
Tobramycin		Trisul	99	Vasodilators	50
Infection	99	Trophic Hormones	87	Vasopressin Agonists	9
Sensory	274	Tropicamide	277	Vasorex	50
Tobramycin (Viatris)	99	Trulicity		Vebulis	
Tobramycin BNM		TruSteel		Vedafil	6
Tobrex		Tryzan		Vedolizumab	24
Tocilizumab		Tuberculin PPD [Mantoux] tes		Vegzelma	20
Tofranil		Tubersol		Veklury	
Tolcapone		Two Cal HN		Veletri	
Tolvaptan		TYR Anamix Infant		Venclexta	
Topamax		TYR Anamix Junior		Venetoclax	
Topical Products for Joint and		TYR Anamix Junior LQ		Venlafaxine	
Muscular Pain	116	TYR Explore 5		Venomil	
Topiramate		TYR Lophlex LQ 20		VENOX	
Topiramate Actavis		TYR Sphere 20		Ventolin	
Total parenteral nutrition (TPN).	101	Tyrosine1000		VentoiiiVepesid	
TPN		Tysabri		Verapamil hydrochloride	
Tramadol hydrochloride		- U -	103	Vermox	
Tramal SR 100		UK Cipla	267	Versacloz	
Tramal SR 150				Vesanoid	
Tramal SR 200		UK Synacthen Ultibro Breezhaler	060	Vexazone	
Trametinib		Ultraproct		Viederm KC	
Trandate		Umeclidinium		Viaderm KC	
Tranexamic acid		Umeclidinium with vilanterol		Victoza	
Tranylcypromine sulphate		Univent	-	Vigabatrin	
Trastuzumab (Herzuma)		Upadacitinib		Vigisom	
Trastuzumab deruxtecan		Ural		Vildagliptin	12
Trastuzumab emtansine		Urea		Vildagliptin with metformin	
Travatan		Urex Forte		hydrochloride	
Travoprost		Urinary Agents		Vimpat	
Treatments for Dementia	146	Urinary Tract Infections		Vinblastine sulphate	
Treatments for Substance		UroFos		Vincristine sulphate	
Dependence		Uromitexan		Vinorelbine	
Trelegy Ellipta	269	Ursodeoxycholic acid	23	Vinorelbine Ebewe	
Trental 400	56	Ursosan		Vinorelbine Te Arai	
Tretinoin		Ustekinumab	243	Viramune Suspension	110
Dermatological	67	Utrogestan		ViruPOS	
Oncology		- V -		Vit.D3	3
Trexate		Vabysmo	205	VitA-POS	27
Triamcinolone acetonide		Vaccinations		Vitabdeck	
Alimentary	32	Vaclovir		Vital	
Dermatological		Valaciclovir		Vitamin B complex	
Hormone		Valganciclovir		Vitamin B6 25	
	-	Valganciclovir Viatris		Vitamins	
Triamcinolone acetonide with		valgariololovii viatris		v Itaiiiii io	02 0

Vivonex TEN289
Voltaren
Voltaren D115
Voltaren SR115
Volumatic273
Voriconazole101
Vttack101
Vyvanse143
- W -
Warfarin sodium43
Wart Preparations75
Wasp venom allergy treatment265
Water
Blood44
Extemporaneous282
White Soft Liquid Paraffin AFT71
Wool fat with mineral oil71
- X -
Xalkori
Xaluprine
Xarelto43
Xgeva116
Xifaxan10
XMET Maxamum296
Xolair
Xolair AU217
XP Maxamum297
Xylocaine123
Xylocaine 2% Jelly122
Xylocaine Viscous123
Xylocard 500
Xyntha39
- Y -
Yervoy248
- Z -
Zarontin
Zaroxolyn53
Zavedos
Zeffix
Zejula157
Zematop74
Zetlam105
Ziagen110
Zidovudine [AZT]111
Zidovudine [AZT] with
Zidovudine [AZ1] With
lamivudine 111
Ziextenzo43
Zimybe55
Zinc and castor oil70
Zinc sulphate35
Zincaps
Zincaps S2935
Ziprasidone135
Zista265
Zithromax93
7o-Ruh HP 123

Zo-Rub Osteo11	6
Zoladex9	1
Zoledronic acid	
Hormone8	3
Musculoskeletal11	8
Zoledronic acid Injection Mylan8	3
Zoledronic acid Viatris	
Hormone8	3
Musculoskeletal11	8
Zopiclone14	1
Zopiclone Actavis14	1
Zostrix11	
Zostrix HP12	3
Zuclopenthixol decanoate13	7
Zuclopenthixol hydrochloride13	
Zusdone13	5
Zyban14	
Zypine13	4
Zypine ODT13	4
Zyprexa Relprevv13	6
Zytiga17	4
7,0,0	M.