Enquiries to Ministry of Health 0800 855 066

APPLICATION FOR SUBSIDY BY SPECIAL AUTHORITY

Page 1 Form SA2443 December 2025

Initial application — non-small cell lung cancer second line monotherapy Applications only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 month Prerequisites(tick boxes where appropriate) Patient has locally advanced or metastatic non-small cell lung cancer and Patient has not received prior funded treatment with an immune checkpoint inhibitor for NSCLC and					
Address: Address: Address: Fax Number: Fax Number: Fax Number: Atezolizumab Initial application — non-small cell lung cancer second line monotherapy Applications only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 month Prerequisites(tick boxes where appropriate) Patient has locally advanced or metastatic non-small cell lung cancer and Patient has not received prior funded treatment with an immune checkpoint inhibitor for NSCLC and For patients with non-squamous histology there is documentation confirming that the disease does not express activating mutations EGFR or ALK tyrosine kinase unless not possible to ascertain Patient has an ECOG 0-2 and Patient has documented disease progression following treatment with at least two cycles of platinum-based chemotherapy and Atezolizumab is to be used as monotherapy at a dose of 1200 mg every three weeks (or equivalent) for a maximum of 16 weeks					
Address: Fax Number: Atezolizumab Initial application — non-small cell lung cancer second line monotherapy Applications only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 month Prerequisites(tick boxes where appropriate) Patient has locally advanced or metastatic non-small cell lung cancer and Patient has not received prior funded treatment with an immune checkpoint inhibitor for NSCLC and For patients with non-squamous histology there is documentation confirming that the disease does not express activating mutations EGFR or ALK tyrosine kinase unless not possible to ascertain Patient has an ECOG 0-2 and Patient has documented disease progression following treatment with at least two cycles of platinum-based chemotherapy Atezolizumab is to be used as monotherapy at a dose of 1200 mg every three weeks (or equivalent) for a maximum of 16 weeks					
Fax Number: Fax Nu					
Initial application — non-small cell lung cancer second line monotherapy Applications only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 month Prerequisites(tick boxes where appropriate) Patient has locally advanced or metastatic non-small cell lung cancer and Patient has not received prior funded treatment with an immune checkpoint inhibitor for NSCLC and For patients with non-squamous histology there is documentation confirming that the disease does not express activating mutations EGFR or ALK tyrosine kinase unless not possible to ascertain Patient has an ECOG 0-2 and Atezolizumab is to be used as monotherapy at a dose of 1200 mg every three weeks (or equivalent) for a maximum of 16 weeks					
Initial application — non-small cell lung cancer second line monotherapy Applications only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 month Prerequisites(tick boxes where appropriate) Patient has locally advanced or metastatic non-small cell lung cancer and Patient has not received prior funded treatment with an immune checkpoint inhibitor for NSCLC and For patients with non-squamous histology there is documentation confirming that the disease does not express activating mutations EGFR or ALK tyrosine kinase unless not possible to ascertain Patient has an ECOG 0-2 and Atezolizumab is to be used as monotherapy at a dose of 1200 mg every three weeks (or equivalent) for a maximum of 16 weeks					
Initial application — non-small cell lung cancer second line monotherapy Applications only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 month Prerequisites(tick boxes where appropriate) Patient has locally advanced or metastatic non-small cell lung cancer and Patient has not received prior funded treatment with an immune checkpoint inhibitor for NSCLC and For patients with non-squamous histology there is documentation confirming that the disease does not express activating mutations EGFR or ALK tyrosine kinase unless not possible to ascertain Patient has an ECOG 0-2 and Patient has documented disease progression following treatment with at least two cycles of platinum-based chemotherapy and Atezolizumab is to be used as monotherapy at a dose of 1200 mg every three weeks (or equivalent) for a maximum of 16 weeks					
Applications only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 month Prerequisites(tick boxes where appropriate) Patient has locally advanced or metastatic non-small cell lung cancer and Patient has not received prior funded treatment with an immune checkpoint inhibitor for NSCLC and For patients with non-squamous histology there is documentation confirming that the disease does not express activating mutations EGFR or ALK tyrosine kinase unless not possible to ascertain Patient has an ECOG 0-2 and Patient has documented disease progression following treatment with at least two cycles of platinum-based chemotherapy and Atezolizumab is to be used as monotherapy at a dose of 1200 mg every three weeks (or equivalent) for a maximum of 16 weeks	Atezolizumab				
and Patient has not received prior funded treatment with an immune checkpoint inhibitor for NSCLC and For patients with non-squamous histology there is documentation confirming that the disease does not express activating mutations EGFR or ALK tyrosine kinase unless not possible to ascertain and Patient has an ECOG 0-2 and Patient has documented disease progression following treatment with at least two cycles of platinum-based chemotherapy and Atezolizumab is to be used as monotherapy at a dose of 1200 mg every three weeks (or equivalent) for a maximum of 16 weeks	Applications only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months.				
Patient has not received prior funded treatment with an immune checkpoint inhibitor for NSCLC and For patients with non-squamous histology there is documentation confirming that the disease does not express activating mutations EGFR or ALK tyrosine kinase unless not possible to ascertain and Patient has an ECOG 0-2 and Patient has documented disease progression following treatment with at least two cycles of platinum-based chemotherapy and Atezolizumab is to be used as monotherapy at a dose of 1200 mg every three weeks (or equivalent) for a maximum of 16 weeks	· · · · · · · · · · · · · · · · · · ·				
For patients with non-squamous histology there is documentation confirming that the disease does not express activating mutations EGFR or ALK tyrosine kinase unless not possible to ascertain Patient has an ECOG 0-2 and Patient has documented disease progression following treatment with at least two cycles of platinum-based chemotherapy and Atezolizumab is to be used as monotherapy at a dose of 1200 mg every three weeks (or equivalent) for a maximum of 16 weeks	Patient has not received prior funded treatment with an immune checkpoint inhibitor for NSCLC				
Patient has an ECOG 0-2 and Patient has documented disease progression following treatment with at least two cycles of platinum-based chemotherapy and Atezolizumab is to be used as monotherapy at a dose of 1200 mg every three weeks (or equivalent) for a maximum of 16 weeks	For patients with non-squamous histology there is documentation confirming that the disease does not express activating mutations of EGFR or ALK tyrosine kinase unless not possible to ascertain				
Patient has documented disease progression following treatment with at least two cycles of platinum-based chemotherapy and Atezolizumab is to be used as monotherapy at a dose of 1200 mg every three weeks (or equivalent) for a maximum of 16 weeks	Patient has an ECOG 0-2				
Atezolizumab is to be used as monotherapy at a dose of 1200 mg every three weeks (or equivalent) for a maximum of 16 weeks	Patient has documented disease progression following treatment with at least two cycles of platinum-based chemotherapy				
l and					
Baseline measurement of overall tumour burden is documented clinically and radiologically					
Renewal — non-small cell lung cancer second line monotherapy Current approval Number (if known):					
Applications only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months. Prerequisites(tick boxes where appropriate)					
Patient's disease has had a complete response to treatment or					
Patient's disease has had a partial response to treatment					
Patient has stable disease					
Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period	ent				
No evidence of disease progression					
The treatment remains clinically appropriate and patient is benefitting from treatment					
and Atezolizumab to be used at a maximum dose of 1200 mg every three weeks (or equivalent)					
Treatment with atezolizumab to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed even 3 weeks)					

I confirm the above details are correct and that in signing this form I understand I may be audited.

Enquiries to Ministry of Health 0800 855 066

APPLICATION FOR SUBSIDY BY SPECIAL AUTHORITY

Page 2 Form SA2443 December 2025

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:		
Reg No:	First Names:	First Names:		
Name:	Surname:	Surname:		
Address:	DOB:	Address:		
	Address:			
Fax Number:		Fax Number:		
Initial application — unresectable hepatocellular carcinoma Applications from any relevant practitioner. Approvals valid for 6 months. Prerequisites(tick boxes where appropriate) Patient is currently on treatment with atezolizumab and met all remaining criteria prior to commencing treatment or Patient has locally advanced or metastatic, unresectable hepatocellular carcinoma and Patient has preserved liver function (Child-Pugh A) and Transarterial chemoembolisation (TACE) is unsuitable and Patient has not received prior systemic therapy for the treatment of hepatocellular carcinoma or Patient received funded lenvatinib before 1 March 2025 or Patient has experienced treatment-limiting toxicity from treatment with lenvatinib and No disease progression since initiation of lenvatinib and Patient has an ECOG performance status of 0-2 and To be given in combination with bevacizumab				
Renewal — unresectable hepatocellular carcinoma				
Current approval Number (if known):				
Applications from any relevant practitioner. Approvals valid for 6 months. Prerequisites(tick box where appropriate)				
There is no evidence of disease progression				