## SA2487 - Infliximab

Crohn's disease (adults) - Initial application	2
Crohn's disease (adults) - Renewal	2
Crohn's disease (children) - Initial application	3
Crohn's disease (children) - Renewal	
Graft vs host disease - Initial application	
Pulmonary sarcoidosis - Initial application	3
Acute fulminant ulcerative colitis - Initial application	
Ankylosing spondylitis - Initial application	
Ankylosing spondylitis - Renewal	
Chronic ocular inflammation - Initial application	5
Chronic ocular inflammation - Renewal	
Fistulising Crohn's disease - Initial application	
Fistulising Crohn's disease - Renewal	
Fulminant ulcerative colitis - Renewal	
Immune checkpoint inhibitor toxicity in malignancy* - Initial application	16
Immune checkpoint inhibitor toxicity in malignancy* - Renewal	
Inflammatory bowel arthritis – axial - Initial application	
Inflammatory bowel arthritis – axial - Renewal	15
Inflammatory bowel arthritis – peripheral - Initial application	16
Inflammatory bowel arthritis – peripheral - Renewal	
Neurosarcoidosis - Initial application	
Neurosarcoidosis - Renewal	
Plaque psoriasis - Initial application	8
Plaque psoriasis - Renewal	
Previous use - Initial application	10
Psoriatic arthritis - Initial application	10
Psoriatic arthritis - Renewal	11
Pyoderma gangrenosum - Initial application	14
Pyoderma gangrenosum - Renewal	15
Rheumatoid arthritis - Initial application	
Rheumatoid arthritis - Renewal	11
Severe Behcet's disease - Initial application	
Severe Behcet's disease - Renewal	12
Severe ocular inflammation - Initial application	
Severe ocular inflammation - Renewal	13
Ulcerative colitis - Initial application	14
Ulcerative colitis - Renewal	

# APPLICATION FOR SUBSIDY BY SPECIAL AUTHORITY

Page 2 Form SA2487 December 2025

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
	Address:	
Fax Number:		Fax Number:
Infliximab		
or Patient has extensive small i or Patient has evidence of shor or Patient has an ileostomy or cand Patient has tried but has experience and corticosteroids		ome with further bowel resection
or initiated on infliximab  CDAI score is 150 or less, or or  The patient has demonstrate  and  Infliximab to be administered at dos to 3 doses if required for secondary	rals valid for 2 years.  100 points from the CDAI score, or HBI score has recovered to the control of the contr	e and/or HBI score cannot be assessed very 8 weeks (or equivalent) can be used for uper re-induction may be considered sixteen weeks

# APPLICATION FOR SUBSIDY BY SPECIAL AUTHORITY

Page 3 Form SA2487 December 2025

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:	
Reg No:	First Names:	First Names:	
Name:	Surname:	Surname:	
Address:	DOB:	Address:	
	Address:		
Fax Number:		Fax Number:	
Infliximab - continued			
Initial application — Crohn's disease (children Applications from any relevant practitioner. Appropriates (tick boxes where appropriate)			
Paediatric patient has active Croh	n's disease		
Patient has a PCDAI score of Patient has extensive small	of greater than or equal to 30 intestine disease		
and Patient has tried but experienced a corticosteroids	an inadequate response to, or intolerable side effects	from, prior therapy with immunomodulators and	
Current approval Number (if known):			
or PCDAI score is 15 or less or The patient has demonstrate and Infliximab to be administered at do to 3 doses if required for secondar	y 10 points from the PCDAI score when the patient we get an adequate response to treatment but PCDAI scores up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every non-response to treatment for re-induction. Another	ore cannot be assessed  very 8 weeks (or equivalent) can be used for up er re-induction may be considered sixteen weeks	
after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019			
Initial application — Graft vs host disease Applications from any relevant practitioner. Appro Prerequisites(tick box where appropriate)			
Patient has steroid-refractory acute graft	vs. Host disease of the gut		
Initial application — Pulmonary sarcoidosis Applications from any relevant practitioner. Appro Prerequisites(tick box where appropriate)			
Patient has life-threatening pulmonary sa	arcoidosis diagnosed by a multidisciplinary team that	is retractory to other treatments	

# APPLICATION FOR SUBSIDY BY SPECIAL AUTHORITY

Page 4 Form SA2487 December 2025

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:	
Reg No:	First Names:	First Names:	
Name:	Surname:	Surname:	
Address:	DOB:	Address:	
	Address:		
Fax Number:		Fax Number:	
Infliximab - continued			
Prerequisites(tick boxes where appropriate)  Patient has acute, fulminant ulcera	ctitioner on the recommendation of a gastroenterolog	gist. Approvals valid for 6 weeks.	
Prerequisites(tick boxes where appropriate)  The patient has had an initial Speciand  The patient has experienced or	cial Authority approval for adalimumab and/or etanerod intolerable side effects from a reasonable trial of adamumab and/or etanercept treatment, the patient did using spondylitis	cept for ankylosing spondylitis	
Denough enlydesing energlyditic			
Prerequisites(tick boxes where appropriate)  Following 12 weeks of infliximab to or by 50%, whichever is less and	oner on the recommendation of a rheumatologist. Apprecatment, BASDAI has improved by 4 or more points on that benefited from treatment and that continued treatment and that continued treatment.	from pre-infliximab baseline on a 10 point scale,	

# APPLICATION FOR SUBSIDY BY SPECIAL AUTHORITY

Page 5 Form SA2487 December 2025

APPLICANT (stamp or sticker acceptable)		icker acceptable)	PATIENT NHI:	REFERRER Reg No:		
Reg No:			First Names:	First Names:		
Name	e:				Surname:	Surname:
Addre	ess:				DOB:	Address:
					Address:	
Fax N	lumbe	ər:				Fax Number:
Inflix	kima	<b>b</b> - co	ntinued			
App	licatio	ons fron	n any re	ronic ocular inflammati elevant practitioner. Appro s where appropriate)		
		and	Or _	The patient has expe	al Special Authority approval for adalimumab for chrorienced intolerable side effects from adalimumab	
			L	The patient has received ocular inflammation	ed insufficient benefit from adalimumab to meet the r	enewal criteria for adalimumab for chronic
	or	and	or C	Patient is 18 years or  Patient is under 18 years  Patient is under 8 years	older and treatment with at least two other immunomars and treatment with methotrexate has proven ineffacts and treatment with steroids or methotrexate has plisease requires control to prevent irreversible vision least	odulatory agents has proven ineffective ective or is not tolerated at a therapeutic dose roven ineffective or is not tolerated at a
Ren	ewal ·	— chro	onic oc	ular inflammation		
Appl	icatio	ns from	any re	er (if known):levant practitioner. Appros s where appropriate)	vals valid for 12 months.	
	or	_ т	he pation	ent has had a good clinica	al response following 3 initial doses	
		N	lomenc		nt period, the patient has had a sustained reduction in anterior chamber or vitreous cells, absence of active	
	or				nt period, the patient has a sustained steroid sparing of ice daily if under 18 years old	effect, allowing reduction in prednisone to < 10mg
				should be considered after withdrawn.	r every 24 months of stability, unless the patient is de	eemed to have extremely high risk of irreversible

# APPLICATION FOR SUBSIDY BY SPECIAL AUTHORITY

Page 6 Form SA2487 December 2025

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
	Address:	
Fax Number:		Fax Number:
Infliximab - continued		
Initial application — fistulising Crohn's disease Applications from any relevant practitioner. Appropriete (tick boxes where appropriate)		
Patient has confirmed Crohn's disc	ease	
Patient has one or more con Patient has one or more recon Patient has one or more recon Patient has complex peri-ana		
Prerequisites(tick boxes where appropriate)  The number of open draining or There has been a marked regreduction in the Fistula Asset and Infliximab to be administered at do to 3 doses if required for secondary	g fistulae have decreased from baseline by at least 5 eduction in drainage of all fistula(e) from baseline (in tessment score), together with less induration and patients up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every non-response to treatment for re-induction. Another on cycle. Up to 10 mg/kg every 8 weeks (or equivalent)	the case of adult patients, as demonstrated by a ent reported pain  very 8 weeks (or equivalent) can be used for uper re-induction may be considered sixteen weeks
Initial application — neurosarcoidosis Applications only from a neurologist or Practitione Prerequisites(tick boxes where appropriate)	r on the recommendation of a neurologist. Approvals	s valid for 18 months.
Patient has been diagnosed with rand Patient has CNS involvement and Patient has steroid-refractory disea	neurosarcoiosis by a multidisciplinary team	
or IV cyclophosphamide has be  Treatment with IV cyclophos	een tried phamide is clinically inappropriate	

# APPLICATION FOR SUBSIDY BY SPECIAL AUTHORITY

Page 7 Form SA2487 December 2025

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
	Address:	
Fax Number:		Fax Number:
Infliximab - continued		
Renewal — neurosarcoidosis		
Current approval Number (if known):		
Applications only from a neurologist or Practitioner <b>Prerequisites</b> (tick boxes where appropriate)	on the recommendation of a neurologist. Approvals	valid for 18 months.
Prerequisites (tick boxes where appropriate)		
A withdrawal period has been tried	and the patient has relapsed	
	n considered but would not be clinically appropriate	
and There has been a marked re and	duction in prednisone dose	
	provement in MRI appearances	
or Marked improvement	in other symptomology	

# APPLICATION FOR SUBSIDY BY SPECIAL AUTHORITY

Page 8 Form SA2487 December 2025

APPL	ICAN <sup>-</sup>	<b>T</b> (star	np o	sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg N	o:				First Names:	First Names:
Name	:				Surname:	Surname:
Addre	ss:				DOB:	Address:
					Address:	
						Fax Number:
Inflix	imat	<b>)</b> - co	ntinu	ed		
Appl	ication	ns only	y fror	The patient has had an initial psoriasis  Patient has experienc  Patient has received in	vant practitioner on the recommendation of a dermate of a	ept or secukinumab for severe chronic plaque ept or secukinumab ecukinumab to meet the renewal criteria for
	or	and and and		greater than 10, where Patient has severe cheave been present for Patient has severe chefor at least 6 months than 10  Patient has tried, but had an of the following (at maximum A PASI assessment has becourses), preferably while st	dy" severe chronic plaque psoriasis with a Psoriasis of elesions have been present for at least 6 months from the least 6 months from the time of initial diagnosis conic localised genital or flexural plaque psoriasis who more than 1 month old at the time of initial diagnosis, and with a Dermatole inadequate response (see Note) to, or has experience to localised doses unless contraindicated): phototheral en completed for at least the most recent prior treatment on treatment but no longer than 1 month following assement is no more than 1 month old at the time of initial diagnosis with a Psoriasis who are localised genital or flexural plaque psoriasis who are localised genital plaque psoriasis who are localised genital	or sole of a foot, where the plaque or plaques ere the plaques or lesions have been present ogy Life Quality Index (DLQI) score greater  ced intolerable side effects from, at least three apy, methotrexate, ciclosporin, or acitretin ent course (but preferably all prior treatment cessation of each prior treatment course
while face,	still o hand,	n trea , foot,	tmen genit	t but no longer than 1 monthal or flexural areas at least 2	ole body severe chronic plaque psoriasis, a PASI scorifollowing cessation of the most recent prior treatmer of the 3 PASI symptom subscores for erythema, thic a foot the skin area affected is 30% or more of the fa	nt; for severe chronic plaque psoriasis of the ckness and scaling are rated as severe or very

preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

# APPLICATION FOR SUBSIDY BY SPECIAL AUTHORITY

Page 9 Form SA2487 December 2025

	amp or	. ,	PATIENT NHI:	REFERRER Reg No:
g No:			First Names:	First Names:
ıme:			Surname:	Surname:
dress:			DOB:	Address:
			Address:	
				Fax Number:
fliximab - c	continue	d		
pplications fro	al Numb		vals valid for 6 months.	
	and	Following each prior i	ody" severe chronic plaque psoriasis at the start of tre infliximab treatment course the patient has a PASI sco I, when compared with the pre-infliximab treatment ba	ore which is reduced by 75% or more, or is
or		Following each prior is sustained at this leve	infliximab treatment course the patient has a PASI scotl, when compared with the pre-infliximab treatment ba	ore which is reduced by 75% or more, or is a seline value  or sole of a foot at the start of treatment
	and	Following each prior is sustained at this leve  Patient had severe che all 3 of erythem course baseline  Following each prior is sustained at this leve	infliximab treatment course the patient has a PASI scot, when compared with the pre-infliximab treatment batherionic plaque psoriasis of the face, or palm of a hand prior infliximab treatment course the patient has a recta, thickness and scaling, to slight or better, or sustain	ore which is reduced by 75% or more, or is isseline value  or sole of a foot at the start of treatment  duction in the PASI symptom subscores for led at this level, as compared to the treatment duction of 75% or more in the skin area
or	and	Following each prior is sustained at this leve  Patient had severe che Following each all 3 of erythem course baseline Following each affected, or sus	infliximab treatment course the patient has a PASI scot, when compared with the pre-infliximab treatment bath aronic plaque psoriasis of the face, or palm of a hand prior infliximab treatment course the patient has a rectal, thickness and scaling, to slight or better, or sustain a values	or which is reduced by 75% or more, or is a seline value  or sole of a foot at the start of treatment  duction in the PASI symptom subscores for led at this level, as compared to the treatment duction of 75% or more in the skin area treatment baseline value

# APPLICATION FOR SUBSIDY BY SPECIAL AUTHORITY

Page 10 Form SA2487 December 2025

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
	Address:	
Fax Number:		Fax Number:
Infliximab - continued		
Initial application — previous use Applications from any relevant practitioner. Appro Prerequisites(tick boxes where appropriate)  Patient was being treated with infli		
and	minab prior to 1 February 2019	
Rheumatoid arthritis		
Ankylosing spondylitis		
Psoriatic arthritis		
Severe ocular inflammation		
or Chronic ocular inflammation		
or Crohn's disease (adults)		
or		
Crohn's disease (children)		
Fistulising Crohn's disease		
Severe fulminant ulcerative o	colitis	
Severe ulcerative colitis		
Plaque psoriasis		
or Neurosarcoidosis		
or Severe Behcet's disease		
Initial application — psoriatic arthritis Applications only from a rheumatologist or Practiti Prerequisites(tick boxes where appropriate)	ioner on the recommendation of a rheumatologist. Ap	oprovals valid for 4 months.
The patient has had an initial Spec	cial Authority approval for adalimumab and/or etanero	ept and/or secukinumab for psoriatic arthritis
· ·	d intolerable side effects from adalimumab and/or etar	nercept and/or secukinumab
	treatment with adalimumab and/or etanercept and/or nab and/or etanercept and/or secukinumab for psoria	

# APPLICATION FOR SUBSIDY BY SPECIAL AUTHORITY

Page 11 Form SA2487 December 2025

Name:         Surname:         Surname:           Address:         DOB:         Address:           Address:	APPL	ICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Address:	Reg No:		First Names:	First Names:
Fax Number: Fax Nu	Name:		Surname:	Surname:
Fax Number: Fax Number: Fax Number: Infliximab - continued  Renewal — psoriatic arthritis  Current approval Number (if known): Applications only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months. Prerequisites(tick boxes where appropriate)	Addre	ss:	DOB:	Address:
Fax Number: Fax Number: Fax Number: Infliximab - continued  Renewal — psoriatic arthritis  Current approval Number (if known):			Address:	
Renewal — psoriatic arthritis				
Current approval Number (if known):	Fax N	umber:		Fax Number:
Current approval Number (if known):	Inflix	imab - continued		
Applications only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months.  Prerequisites(tick boxes where appropriate)    Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician   The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior infliximab treatment in the opinion of the treating physician   Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks    Initial application — rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months.   Prerequisites(tick boxes where appropriate)    The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis     The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept     Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept     Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance     Renewal — rheumatologist for Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months.   Prerequisites(tick boxes where appropriate)     Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance     Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance     Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance     Treatment is to be used as an adjunct to methotrexate therapy or mo	Rene	ewal — psoriatic arthritis		
Applications only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months.  Prerequisites(tick boxes where appropriate)    Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician   The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior infliximab treatment in the opinion of the treating physician   Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks    Initial application — rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months.   Prerequisites(tick boxes where appropriate)    The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis     The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept     Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept     Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance     Renewal — rheumatologist for Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months.   Prerequisites(tick boxes where appropriate)     Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance     Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance     Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance     Treatment is to be used as an adjunct to methotrexate therapy or mo	Curre	ent approval Number (if known):		
Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician  The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior infliximab treatment in the opinion of the treating physician  Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks  Initial application — rheumatoid arthritis  Applications only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months.  Prerequisites(tick boxes where appropriate)  The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis  The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept and/or etanercept  Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance  Renewal — rheumatoid arthritis  Current approval Number (if known):  Applications only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months.  Prerequisites(tick boxes where appropriate)  Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance  Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance  Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance  Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance  Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolera		, ,		provals valid for 6 months.
clinically significant response to treatment in the opinion of the physician  The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior infliximab treatment in the opinion of the treating physician  Initial application — rheumatoid arthritis  Applications only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months.  Prerequisites(tick boxes where appropriate)  The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis  and  The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept  Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept intolerance  Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance  Renewal — rheumatoid arthritis  Current approval Number (if known):	Prere	equisites(tick boxes where appropriate)		
clinically significant response to treatment in the opinion of the physician  The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior infliximab treatment in the opinion of the treating physician  Initial application — rheumatoid arthritis  Applications only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months.  Prerequisites(tick boxes where appropriate)  The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis  and  The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept  Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept intolerance  Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance  Renewal — rheumatoid arthritis  Current approval Number (if known):		Following 2 to 4 months' in	tial treatment, the nations has at least a 50% decrease	o in active joint count from baceline and a
The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior infliximab treatment in the opinion of the treating physician   Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks		clinically significant respons		e in active joint count from baseline and a
Initial application — rheumatoid arthritis  Applications only from a rheumatoid gist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months.  Prerequisites(tick boxes where appropriate)  The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis  The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept  or Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept intolerance  Renewal — rheumatoid arthritis  Current approval Number (if known):		The patient demonstrates at		ount from baseline and a clinically significant
Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks  Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks  Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks  Infliximab polication — rheumatoid arthritis Applications only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months.  Prerequisites(tick boxes where appropriate)  The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis  The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept  Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept intolerance  Renewal — rheumatoid arthritis  Current approval Number (if known):			treatment in the opinion of the treating physician	
Applications only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months.  Prerequisites(tick boxes where appropriate)  The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis  and  The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept  Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept intolerance  Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance  Renewal — rheumatoid arthritis  Current approval Number (if known):			ses no greater than 5 mg/kg every 8 weeks	
Applications only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months.  Prerequisites(tick boxes where appropriate)  The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis  and  The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept  Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept intolerance  Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance  Renewal — rheumatoid arthritis  Current approval Number (if known):	1 141-	t and the state of		
The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis  The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept  Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept and Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance  Renewal — rheumatoid arthritis  Current approval Number (if known):	Appl	lications only from a rheumatologist or Practit	ioner on the recommendation of a rheumatologist. Ap	oprovals valid for 4 months.
The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept  or Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept and Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance  Renewal — rheumatoid arthritis  Current approval Number (if known):				
Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept  and  Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance  Renewal — rheumatoid arthritis  Current approval Number (if known):			cial Authority approval for adalimumab and/or etanerc	ept for rheumatoid arthritis
Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept  and  Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance  Renewal — rheumatoid arthritis  Current approval Number (if known):			d intolerable side effects from a reasonable trial of ada	alimumab and/or etanercept
Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance  Renewal — rheumatoid arthritis  Current approval Number (if known):		Following at least a four more	nth trial of adalimumab and/or etanercept, the patient	did not meet the renewal criteria for adalimumab
Renewal — rheumatoid arthritis  Current approval Number (if known):				
Current approval Number (if known):			inct to methotrexate therapy or monotherapy where u	se of methotrexate is limited by toxicity or
Current approval Number (if known):	Rene	ewal — rheumatoid arthritis		
Applications only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months.  Prerequisites(tick boxes where appropriate)  Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance  Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a	110110			
Prerequisites(tick boxes where appropriate)  Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance  and  Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a		, ,		
intolerance  Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a			oner on the recommendation of a rheumatologist. Ap	provals valid for 6 months.
Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a			unct to methotrexate therapy or monotherapy where us	se of methotrexate is limited by toxicity or
		and		
		clinically significant respons		e in active joint count from baseline and a
The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician		The patient demonstrates at		ount from baseline and a clinically significant
and Infliximab to be administered at doses no greater than 3 mg/kg every 8 weeks			ses no greater than 3 mg/kg every 8 weeks	

# APPLICATION FOR SUBSIDY BY SPECIAL AUTHORITY

Page 12 Form SA2487 December 2025

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
	Address:	
Fax Number:		Fax Number:
Infliximab - continued		
The patient has severe ocula treatment(s) appropriate for or  The patient has severe gastre two or more treatment appropriate for two or more treatment appropriate.  The patient is experiencing signification.  Note: Behcet's disease diagnosed according to the measured using an appropriate quality of life scale. Treatments appropriate for the particular symptoms intravenous/oral steroids and other immunosuppre	sease which is significantly impacting the patient's quar, neurological and/or vasculitic symptoms and has rethe particular symptom(s) (see Notes)  rointestinal, rheumatologic and/or mucocutaneous symptom for the particular symptom(s) (see Notes)	mptoms and has not responded adequately to more mptoms and has not responded adequately to mancet 1990;335(8697):1078-80. Quality of life 1. 2004;31:931-7. altreatments for these symptoms, for example
Renewal — severe Behcet's disease		
and		quality of life
Renewal — fulminant ulcerative colitis		
Current approval Number (if known):		

# APPLICATION FOR SUBSIDY BY SPECIAL AUTHORITY

Page 13 Form SA2487 December 2025

APPLICANT (stamp or sticker acceptable)		PATIENT NHI:	REFERRER Reg No:	
Reg No:		First Names:	First Names:	
Name:		Surname:	Surname:	
Address:		DOB:	Address:	
		Address:		
Fax Number:			Fax Number:	
Infliximab - continued				
or	rant practitioner. Appropriate here appropriate patient has had an initial the patient has expended and the patient has received a coular inflammation on the patient with high-clineffective at controlling the p	al Special Authority approval for adalimumab for severienced intolerable side effects from adalimumab ved insufficient benefit from adalimumab to meet the interestening ocular inflammation requiring rapid control lose steroids (intravenous methylprednisolone) follow	renewal criteria for adalimumab for severe	
or _	Patient is aged under ineffective at controlli	8 years and treatment with high dose oral steroids a	nd other immunosuppressants has proven	
		ig symptoms		
Renewal — severe ocular				
Current approval Number (if known):				
The patient	has had a good clinic;	al response following 3 initial doses		
or Following e	ach 12 month treatme	ont period, the patient has had a sustained reduction in anterior chamber or vitreous cells, absence of active		
Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old				
Note: A trial withdrawal sho vision loss if infliximab is wi		er every 24 months of stability, unless the patient is de	eemed to have extremely high risk of irreversible	

# APPLICATION FOR SUBSIDY BY SPECIAL AUTHORITY

Page 14 Form SA2487 December 2025

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:		
Reg No:	First Names:	First Names:		
Name:	Surname:	Surname:		
Address:	DOB:	Address:		
	Address:			
Fax Number:		Fax Number:		
Initial application — ulcerative colitis Applications from any relevant practitioner. Approvals valid for 6 months.  Prerequisites(tick boxes where appropriate)  Patient has active ulcerative colitis  and Patients SCCAI is greater than or equal to 4  or Patients PUCAI score is greater than or equal to 20  and Patient has tried but has experienced an inadequate response to, or has experienced intolerable side effects from, prior therapy with				
Renewal — ulcerative colitis  Current approval Number (if known):				
The SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on infliximab  The PUCAI score has reduced by 10 points or more from the PUCAI score when the patient was initiated on infliximab  Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019				
Initial application — pyoderma gangrenosum Applications only from a dermatologist. Approvals valid for 4 months.  Prerequisites(tick boxes where appropriate)				
Patient has pyoderma gangrenosum*  Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response  A maximum of 8 doses  Note: Note: Indications marked with * are unapproved indications.				

# APPLICATION FOR SUBSIDY BY SPECIAL AUTHORITY

Page 15 Form SA2487 December 2025

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:		
Reg No:	First Names:	First Names:		
Name:	Surname:	Surname:		
Address:	DOB:	Address:		
	Address:			
Fax Number:		Fax Number:		
Infliximab - continued				
Renewal — pyoderma gangrenosum				
Current approval Number (if known):				
	ner on the recommendation of a dermatologist. Appr	ovals valid for 4 months.		
Prerequisites(tick boxes where appropriate)				
Patient has shown clinical improve	ement			
and Patient continues to require treatm	nent			
and				
A maximum of 8 doses				
Initial application — inflammatory bowel arthri				
Applications from any relevant practitioner. Appro <b>Prerequisites</b> (tick boxes where appropriate)	vals valid for 6 months.			
Patient has a diagnosis of active ulcerative colitis or active Crohn's disease				
Patient has had axial inflammatory	Patient has had axial inflammatory pain for six months or more			
Patient is unable to take NSAIDs				
and Patient has unequivocal sacroiliitis demonstrated by radiological imaging or MRI				
and Patient's disease has not responded adequately to prior treatment consisting of at least 3 months of an exercise regime supervised by a				
physiotherapist and	adequately to prior treatment consisting of at least	o months of an exercise regime supervised by a		
Patient has a BASDAI of at least 6	on a 0 - 10 scale completed after the 3 month exe	rcise trial, but prior to ceasing any previous		
pharmacological treatment				
Renewal — inflammatory bowel arthritis – axial				
Current approval Number (if known):				
Applications from any relevant practitioner. Approvals valid for 2 years.				
Prerequisites(tick box where appropriate)				
Treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10-point scale, or an improvement in BASDAI of 50%, whichever is less				

# APPLICATION FOR SUBSIDY BY SPECIAL AUTHORITY

Page 16 Form SA2487 December 2025

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:	
Reg No:	First Names:	First Names:	
Name:	Surname:	Surname:	
Address:	DOB:	Address:	
	Address:		
Fax Number:		Fax Number:	
Infliximab - continued			
Initial application — inflammatory bowel arthritis – peripheral Applications from any relevant practitioner. Approvals valid for 6 months.  Prerequisites(tick boxes where appropriate)  Patient has a diagnosis of active ulcerative colitis or active Crohn's disease and Patient has active arthritis in at least four joints from the following: hip, knee, ankle, subtalar, tarsus, forefoot, wrist, elbow, shoulder, sternoclavicular Patient has tried and not experienced a response to at least three months of methotrexate or azathioprine at a maximum tolerated dose (unless contraindicated)  Patient has tried and not experienced a response to at least three months of sulfasalazine at a maximum tolerated dose (unless contraindicated)  Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application  Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application  ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months			
Renewal — inflammatory bowel arthritis – perip	pheral		
Current approval Number (if known):			
Applications from any relevant practitioner. Approv <b>Prerequisites</b> (tick boxes where appropriate)	rals valid for 2 years.		
response to treatment in the opinio	nas experienced at least a 50% decrease in active jo in of the physician continuing 30% improvement in active joint count fro		
Initial application — immune checkpoint inhibitor toxicity in malignancy* Applications from any relevant practitioner. Approvals valid for 4 months.  Prerequisites(tick boxes where appropriate)			
The individual requires treatment for malignancy and	or moderate to severe autoimmune toxicity following i	mmune checkpoint inhibitor treatment for	
The individual has received insufficient benefit from use of corticosteroids			
Infliximab is to be administered at up to 5mg/kg for up to four doses			

# APPLICATION FOR SUBSIDY BY SPECIAL AUTHORITY

Page 17 Form SA2487 December 2025

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:		
Reg No:	First Names:	First Names:		
Name:	Surname:	Surname:		
Address:	DOB:	Address:		
	Address:			
Fax Number:		Fax Number:		
Infliximab - continued				
Renewal — immune checkpoint inhibitor toxicity in malignancy*				
Current approval Number (if known):				
Applications from any relevant practitioner. Approvals valid for 4 months.  Prerequisites(tick boxes where appropriate)				
The individual has shown clinical improvement and ongoing treatment is required and				
Infliximab is to be administered at up to 5mg/kg for up to a total of 8 doses				
Note: Indications marked with * are unapproved indications.				