## January 2026

### **Editors:**

Kave Wilson, Doris Chong email: enquiry@pharmac.govt.nz Telephone +64 4 460 4990 Level 9, 40 Mercer Street PO Box 10 254 Wellington

**Freephone Information Line** 0800 66 00 50 (9am - 5pm weekdays)

### Circulation

You can register to have an electronic version of the Pharmaceutical Schedule (link to PDF copy) emailed to your nominated email address each month by subscribing at pharmac.govt.nz/subscribe.

### Production

Typeset automatically from XML and T<sub>F</sub>X. XML version of the Schedule available from schedule.pharmac.govt.nz/pub/schedule

### **Programmers**

Anrik Drenth

email: texschedule@pharmac.govt.nz ©Pharmaceutical Management Agency



ISSN 1179-3686

This work is licensed under the Creative Commons Attribution 4.0 International licence. In essence, you are free to copy, distribute and adapt it, as long as you attribute the work to Pharmac and abide by the other licence terms. To view a copy of this licence, visit: creativecommons.org/licenses/by/4.0/. Attribution to Pharmac should be in written form and not by reproduction of the Pharmac logo. While care has been taken in compiling this Schedule, Pharmac takes no responsibility for any errors or omissions, and shall not be liable for any consequences arising there from.

Section A	General Rules	5
Section B	Alimentary Tract & Metabolism	6
	Blood & Blood Forming Organs	37
	Cardiovascular System	47
	Dermatologicals	68
	Genito Urinary System	77
	Hormone Preparations – Systemic	83
	Infections – Agents For Systemic Use	93
	Musculoskeletal System	117
	Nervous System	123
	Oncology Agents & Immunosuppressants	153
	Respiratory System & Allergies	267
	Sensory Organs	277
	Various	282
Section C	Extemporaneous Compounds (ECPs)	284
Section D	Special Foods	286
Section I	National Immunisation Schedule	309
	Index	326

Introducing Pharmac

## **Introducing Pharmac**

The Pharmaceutical Management Agency (Pharmac) makes decisions that help control Government spending on pharmaceuticals. This includes community pharmaceuticals, hospital pharmaceuticals, vaccines and increasingly, hospital medical devices. Pharmac negotiates prices, sets subsidy levels and conditions, and makes decisions on changes to the subsidised list.

### Pharmac's role:

"to secure for eligible people in need of pharmaceuticals, the best health outcomes that are reasonably achievable from pharmaceutical treatment and from within the amount of funding provided."

Pae Ora (Healthy Futures) Act 2022

To ensure our decisions are as fair and robust as possible we use a decision-making process that incorporates clinical, economic and commercial issues. We also seek the views of users and the wider community through consultation. The processes we generally use are outlined in our Operating Policies and Procedures.

Further information about Pharmac and the way we make funding decisions can be found on the Pharmac website at <a href="https://pharmac.govt.nz/about">https://pharmac.govt.nz/about</a>.

## **Purpose of the Pharmaceutical Schedule**

The purpose of the Schedule is to list:

- the Community Pharmaceuticals that are subsidised by the Government and to show the amount of the subsidy paid to contractors, as well as the manufacturer's price and any access conditions that may apply;
- the Hospital Pharmaceuticals that may be used in Health NZ Hospitals, as well as any access conditions that may apply; and
- the Pharmaceuticals, including Medical Devices, used in Health NZ Hospitals for which national prices have been negotiated by Pharmac.

The Schedule does not show the final cost to Government of subsidising each Community Pharmaceutical or to Health NZ Hospitals in purchasing each Pharmaceutical, since that will depend on any rebate and other arrangements Pharmac has with the supplier and, for Pharmaceuticals used in Health NZ Hospitals, on any logistics arrangements put in place.

This book contains sections A to D and Section I of the Pharmaceutical Schedule and lists the Pharmaceuticals funded for use in the community, including vaccines, as well as haemophilia and cancer treatments given in Health NZ Hospitals. Section H lists the Pharmaceuticals that that can be used in Health NZ Hospitals and is a separate publication.


The Pharmaceuticals in this book are listed by therapeutic group, which is based on the WHO Anatomical Therapeutic Chemical (ATC) system. The listings are displayed alphabetically under each heading.

The index lists both chemical entities and product brand names.

## **Explaining pharmaceutical entries**

The Pharmaceutical Schedule lists pharmaceuticals subsidised by the Government, the subsidy, the supplier's price and the access conditions that may apply.

### Example



## Glossary

### **Units of Measure**

gramg kilogramkg international unitiu	mi mi mi
Abbreviations	
AmpouleAmp	Ge
CapsuleCap	Gr
Cream	Inf
DeviceDev	Ini
DispersibleDisp	Lic
EffervescentEff	Lo
EmulsionEmul	Oi
Enteric Coated EC	Sa

microgrammilligrammillilitre	mg
Gelatinous	
Granules	
Infusion	Inf
Injection	Inj
Liquid	Liq
Long Acting	LA
Ointment	Oint
Sachet	Sach

millimoleunit	
Solution	Supp Tab
Trans Dermal Delivery System	TDDS

Read the General Rules: https://pharmac.govt.nz/section-a.

## SECTION B: ALIMENTARY TRACT AND METABOLISM

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	Brand or Generic Manufacturer
Antacids and Antiflatulents				
Antacids and Reflux Barrier Agents				
ALGINIC ACID  Sodium alginate 225 mg and magnesium alginate 87.5 mg persachet		30	✓ (	Gaviscon Infant
SODIUM ALGINATE     Tab 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg - peppermint flavour	1.80 (17.99)	60	(	Gaviscon Extra Strength
* Oral liq 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg per 10 ml		500 m		Acidex
Phosphate Binding Agents				
ALUMINIUM HYDROXIDE  * Tab 600 mg  CALCIUM CARBONATE	12.56	100	✓ /	Alu-Tab
Oral liq 1,250 mg per 5 ml (500 mg elemental per 5 ml) – Subsidy by endorsement		500 m 173 m		Roxane Calcium carbonate PAI 829
Only when prescribed for patients unable to swallow calc inappropriate and the prescription is endorsed according		s or v	where calciu	
Antidiarrhoeals				
Agents Which Reduce Motility				
LOPERAMIDE HYDROCHLORIDE − Up to 30 cap available on a  * Tab 2 mg  * Cap 2 mg  Diamide Relief to be Principal Supply on 1 February 202	10.75 12.00	400 400	•	Nodia Diamide Relief
Rectal and Colonic Anti-inflammatories				
BUDESONIDE  Cap modified-release 3 mg - Special Authority see SA2535	22.20	00		Pudaganida Ta Avai

### ⇒SA2535 Special Authority for Subsidy

Initial application — (Crohn's disease) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

1 Mild to moderate ileal, ileocaecal or proximal Crohn's disease; and

2 Any of the following:

continued...

✓ Budesonide Te Arai

90

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	sidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 2.1 Diabetes; or
- 2.2 Cushingoid habitus; or
- 2.3 Osteoporosis where there is significant risk of fracture; or
- 2.4 Severe acne following treatment with conventional corticosteroid therapy; or
- 2.5 History of severe psychiatric problems associated with corticosteroid treatment; or
- 2.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
- 2.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated).

Initial application — (collagenous and lymphocytic colitis (microscopic colitis)) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has a diagnosis of microscopic colitis (collagenous or lymphocytic colitis) by colonoscopy with biopsies.

Initial application — (gut Graft versus Host disease) from any relevant practitioner. Approvals valid for 6 months where patient has a gut Graft versus Host disease following allogenic bone marrow transplantation\*.

Note: Indication marked with \* is an unapproved indication.

Initial application — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has autoimmune hepatitis\*; and
- 2 Patient does not have cirrhosis: and
- 3 Any of the following:
  - 3.1 Diabetes; or
  - 3.2 Cushingoid habitus: or
  - 3.3 Osteoporosis where there is significant risk of fracture; or
  - 3.4 Severe acne following treatment with conventional corticosteroid therapy; or
  - 3.5 History of severe psychiatric problems associated with corticosteroid treatment; or
  - 3.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
  - 3.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated); or
- 3.8 Adolescents with poor linear growth (where conventional corticosteroid use may limit further growth).

Note: Indication marked with \* is an unapproved indication.

Renewal — (gut Graft versus Host disease) from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

**Renewal** — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

HYDROCORTISONE ACETATE			
Rectal foam 10%, CFC-Free (14 applications)57.09	15 g OP	✓ Colifoam	
HYDROCORTISONE ACETATE WITH PRAMOXINE HYDROCHLORIDE			
Topical aerosol foam, 1% with pramoxine hydrochloride 1%26.55	10 g OP	✓ Proctofoam S29	
MESALAZINE			
Tab 400 mg49.50	100	✓ Asacol	
71.00	90	✓ Octasa S29	
Tab long-acting 500 mg56.10	100	✓ Pentasa	
Tab 800 mg85.50	90	✓ Asacol	
		✓ Asacol S29 S29	
Tab 1,600 mg85.50	60	✓ Asacol S29	
Modified release granules, 1 g118.10	100 OP	✓ Pentasa	
Enema 1 g per 100 ml41.30	7	✓ Pentasa	
Suppos 500 mg22.80	20	✓ Asacol	
Suppos 1 g50.96	28	✓ Pentasa	

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Generic
OLSALAZINE				
Tab 500 mg	56.02	60	•	Atnahs Olsalazine S29
	93.37	100	1	Dipentum
Cap 250 mg	53.00	100	1	Dipentum
SODIUM CROMOGLICATE				
Cap 100 mg	113.35	100	1	Ralicrom
	365.00		1	Somex S29
SULFASALAZINE				
* Tab 500 mg	19.49	100	1	Salazopyrin
* Tab EC 500 mg		100	✓	Salazopyrin EN

## Local preparations for Anal and Rectal Disorders

## **Antihaemorrhoidal Preparations**

FLUOCORTOLONE CAPROATE WITH FLUOCORTOLONE PIVALATE AND C	NCHOCAINE	
Oint 950 mcg, with fluocortolone pivalate 920 mcg, and		
cinchocaine hydrochloride 5 mg per g13.05	30 g OP	<ul><li>Ultraproct</li></ul>
Suppos 630 mcg, with fluocortolone pivalate 610 mcg, and		
cinchocaine hydrochloride 1 mg8.61	12	<ul><li>Ultraproct</li></ul>
HYDROCORTISONE WITH CINCHOCAINE		
Oint 5 mg with cinchocaine hydrochloride 5 mg per g15.00	30 g OP	✓ Proctosedyl
Suppos 5 mg with cinchocaine hydrochloride 5 mg per g9.90	12	✓ Proctosedyl

## **Management of Anal Fissures**

GLYCERYL TRINITRATE − Special Authority see SA1329 below − Retail pharmacy

★ Oint 0.2%......22.00 30 g OP

✓ Rectogesic

## **⇒SA1329** Special Authority for Subsidy

OL VOODVDDONII IN DDOMIDE

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has a chronic anal fissure that has persisted for longer than three weeks.

## **Antispasmodics and Other Agents Altering Gut Motility**

GLYCOPYRRONIUM BROMIDE			
Inj 200 mcg per ml, 1 ml ampoule – Up to 10 inj available on a PSO	19.00	5	✓ Robinul
HYOSCINE BUTYLBROMIDE  * Tab 10 mg	2.25	20	✓ <u>Hyoscine</u> <u>Butylbromide</u>
* Inj 20 mg, 1 ml - Up to 5 inj available on a PSO  MEBEVERINE HYDROCHLORIDE	1.91	5	(Adiramedica)  ✓ Spazmol
* Tab 135 mg	8.50	90	✓ Colofac

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

## **Antiulcerants**

## **Antisecretory and Cytoprotective**

MISOPROSTOL - Wastage claimable

## **Helicobacter Pylori Eradication**

### CLARITHROMYCIN

Tab 500 mg − Subsidy by endorsement......14.58 14 ✓ Klacid

a) Maximum of 28 tab per prescription

Subsidised only if prescribed for helicobacter pylori eradication and prescription is endorsed accordingly.
 Note: the prescription is considered endorsed if clarithromycin is prescribed in conjunction with a proton pump inhibitor and either amoxicillin or metronidazole.

## **H2 Antagonists**

FAI	MOTIDINE - Only on a prescription			
*	Tab 20 mg	4.86	100	✓ Famotidine
	·			Hovid S29
*	Tab 40 mg	10.27	100	✓ Famotidine Hovid
	•			MY S29
*	Inj 10 mg per ml, 4 ml - Subsidy by endorsement	CBS	10	✓ Mylan S29
Subsidy by endorsement – Subsidised for patients receiving treatment as part of palliative care.				

## **Proton Pump Inhibitors**

LA	NSOPRAZOLE		
*	Cap 15 mg4.04	100	✓ Lanzol Relief
*	Cap 30 mg	100	✓ Lanzol Relief
ON	MEPRAZOLE		
	For omeprazole suspension refer Standard Formulae, page 284		
*	Cap 10 mg	90	Omeprazole Teva
			<ul> <li>Omeprazole actavis</li> </ul>
			10
*	Cap 20 mg2.02	90	Omeprazole Teva
			<ul> <li>Omeprazole actavis</li> </ul>
			20
*	Cap 40 mg	90	Omeprazole Teva
			<ul> <li>Omeprazole actavis</li> </ul>
			40
*	Powder – Only in combination	5 g	✓ Midwest
	Only in extemporaneously compounded omeprazole suspension.	J	
*	Inj 40 mg ampoule with diluent	5	✓ Dr Reddy's
	,		Omeprazole
			✓ Ocicure S29
DΛ	NTODD AZOL C		
	NTOPRAZOLE	00	( Dames Ballet
*	Tab EC 20 mg	90	✓ Panzop Relief
*	Tab EC 40 mg2.70	90	✓ Panzop Relief

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
Site Protective Agents				
COLLOIDAL BISMUTH SUBCITRATE Tab 120 mg	14.51	50	/	Gastrodenol §29
SUCRALFATE				
Tab 1 g	35.50 (48.28)	120		Carafate
Bile and Liver Therapy				
RIFAXIMIN - Special Authority see SA1461 below - Retail pha Tab 550 mg		56	/	<u>Xifaxan</u>
▶SA1461 Special Authority for Subsidy Initial application only from a gastroenterologist, hepatologist of hepatologist. Approvals valid for 6 months where the patient hat tolerated doses of lactulose.  Renewal only from a gastroenterologist, hepatologist or Practition hepatologist. Approvals valid without further renewal unless not benefiting from treatment.	s hepatic encephalop oner on the recomme	athy d ndatior	espite an n of a gast	adequate trial of maximum troenterologist or
Diabetes				
Hyperglycaemic Agents				
DIAZOXIDE - Special Authority see SA1320 below - Retail pha				
Cap 25 mg		100		Proglicem S29
Cap 100 mg Oral liq 50 mg per ml		100 0 ml C	_	Proglicem \$29 e5 Pharma \$29
➤ SA1320 Special Authority for Subsidy  Initial application from any relevant practitioner. Approvals val hypoglycaemia caused by hyperinsulinism.  Renewal from any relevant practitioner. Approvals valid withour appropriate and the patient is benefiting from treatment.  GLUCAGON HYDROCHLORIDE	lid for 12 months whe	re use	d for the ti	reatment of confirmed
Inj 1 mg syringe kit - Up to 5 kit available on a PSO	32.00	1	✓	Glucagen Hypokit
Insulin - Short-acting Preparations				
INSULIN NEUTRAL				
▲ Inj human 100 u per ml, 3 ml	42.66	5		Actrapid Penfill Humulin R
▲ Inj human 100 u per ml, 10 ml vial	25.26	1 OP	1	Actrapid Humulin R
Insulin - Intermediate-acting Preparations				
INSULIN ASPART WITH INSULIN ASPART PROTAMINE				
▲ Inj 100 iu per ml, 3 ml prefilled pen	52.15	5	/	NovoMix 30 FlexPen
INSULIN DEGLUDEC WITH INSULIN ASPART  ▲ Inj degludec 70 u with insulin aspart 30 u, 100 u per ml, 3 m	ıl80.00	5	•	Ryzodeg 70/30 Penfill

	Subsidy (Manufacturer's Price	) S	Fully Subsidised	Brand or Generic
	\$	Per	1	Manufacturer
NSULIN ISOPHANE				
Inj human 100 u per ml, 3 ml	29.86	5		Humulin NPH
				Protaphane Penfill
Inj human 100 u per ml, 10 ml vial	17.68	1 OP		Humulin NPH
			• 1	Protaphane
NSULIN ISOPHANE WITH INSULIN NEUTRAL	40.00	-		
Inj human with neutral insulin 100 u per ml, 3 ml	42.66	5		Humulin 30/70 PenMix 30
Inj human with neutral insulin 100 u per ml, 10 ml vial	25.26	1 OP	-	Humulin 30/70
PenMix 30 Inj human with neutral insulin 100 u per ml, 3 m			٠,	1011101111 00/70
NSULIN LISPRO WITH INSULIN LISPRO PROTAMINE	,	/		
Inj lispro 25% with insulin lispro protamine 75% 100 u p	ar ml			
3 ml		5	<b>√</b> I	Humalog Mix 25
Inj lispro 50% with insulin lispro protamine 50% 100 u p		-		3 = -
3 ml		5	<b>✓</b> I	Humalog Mix 50
				<u> </u>
Insulin - Long-acting Preparations				
NSULIN GLARGINE				
▲ Inj 100 u per ml, 10 ml	63.00	1	<b>√</b> I	Lantus
▲ Inj 100 u per ml, 3 ml		5	<b>✓</b> i	Lantus
Inj 100 u per ml, 3 ml disposable pen		5	<b>√</b> I	Lantus SoloStar
Insulin - Rapid Acting Preparations				
NSULIN ASPART				
▲ Inj 100 u per ml, 10 ml	30.03	1	<b>✓</b> I	NovoRapid
Inj 100 u per ml, 3 ml		5		NovoRapid Penfill
Inj 100 u per ml, 3 ml syringe		5	<b>✓</b> I	NovoRapid FlexPen
NSULIN GLULISINE				
Inj 100 u per ml, 10 ml	27.03	1	1	Apidra
Inj 100 u per ml, 3 ml		5		Apidra
Inj 100 u per ml, 3 ml disposable pen		5	1	Apidra SoloStar
NSULIN LISPRO				
▲ Inj 100 u per ml, 3 ml	59.52	5	<b>√</b>	Humalog
Inj 100 u per ml, 10 ml vial	34.92	1 OP		Humalog
Alpha Glucosidase Inhibitors				
CARBOSE				
* Tab 50 mg		90		Accarb
★ Tab 100 mg	17.38	90	<b>/</b> <u>!</u>	<u>Accarb</u>
Oral Hypoglycaemic Agents				
SLIBENCLAMIDE				
* Tab 5 mg	7.50	100	<b>√</b> I	Daonil
	7.00	100	٠ ,	-441111
GLICLAZIDE	20.10	E00		Clizido
₭ Tab 80 mg	20.10	500	• (	<u>Glizide</u>
GLIPIZIDE	2.22	400		Material at a fe
★ Tab 5 mg	6.86	100	✓	Minidiab

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

<sup>\*</sup>Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Subs Per	Fully sidised	Brand or Generic Manufacturer
# Tab immediate-release 500 mg  * Tab immediate-release 850 mg		1,000 500	-	Metformin Viatris Metformin Viatris
PIOGLITAZONE  * Tab 15 mg  * Tab 30 mg  * Tab 45 mg	7.25	90 90 90	1	Vexazone Vexazone Vexazone
VILDAGLIPTIN  * Tab 50 mg	35.00	60	•	Galvus
VILDAGLIPTIN WITH METFORMIN HYDROCHLORIDE  * Tab 50 mg with 1,000 mg metformin hydrochloride  * Tab 50 mg with 850 mg metformin hydrochloride		60 60		Galvumet Galvumet

## **GLP-1 Agonists**

DULAGLUTIDE - Special Authority see SA2509 below - Retail pharmacy

### **⇒SA2509** Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has type 2 diabetes; and
- 2 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of all of the following funded blood glucose lowering agents for a period of least 6 months, where clinically appropriate: empagliflozin, metformin, and vildagliptin; and
- 3 Any of the following:
  - 3.1 Patient is Māori or any Pacific ethnicity\*; or
  - 3.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)\*; or
  - 3.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator\*; or
  - 3.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult\*; or
  - 3.5 Patient has diabetic kidney disease (see note b)\*.

Notes: \* Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

- a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack, ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.
- b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m2 in the presence of diabetes, without alternative cause identified.
- c) Funded GLP-1a treatment is not to be given in combination with funded (empagliflozin /empagliflozin with metformin hydrochloride) unless receiving funded (empagliflozin or empagliflozin in combination with metformin hydrochloride) for the treatment of heart failure.

Subsidy	Full	y Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per •	Manufacturer	

LIRAGLUTIDE - Special Authority see SA2510 below - Retail pharmacv

- a) Maximum of 9 ini per prescription
- b)
- a) Note: Not to be given in combination with another funded GLP-1 agonist or empagliflozin / empagliflozin with metformin hydrochloride unless receiving empagliflozin / empagliflozin with metformin hydrochloride for the treatment of heart failure.
- b) Maximum of 1 pack of 3 (6 mg per ml, 3 ml) prefilled pens will be funded per month.

### ⇒SA2510 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has type 2 diabetes; and
- 2 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of all of the following funded blood glucose lowering agents for a period of least 6 months, where clinically appropriate: empagliflozin, metformin, and vildagliptin; and
- 3 Any of the following:
  - 3.1 Patient is Māori or any Pacific ethnicity\*; or
  - 3.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)\*: or
  - 3.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator\*; or
  - 3.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult\*; or
  - 3.5 Patient has diabetic kidney disease (see note b)\*.

Notes: \* Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

- a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack, ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.
- b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m2 in the presence of diabetes, without alternative cause identified.
- c) Funded GLP-1a treatment is not to be given in combination with funded (empagliflozin /empagliflozin with metformin hydrochloride) unless receiving funded (empagliflozin or empagliflozin in combination with metformin hydrochloride) for the treatment of heart failure.

### SGLT2 Inhibitors

### ⇒SA2408 Special Authority for Subsidy

Initial application — (heart failure reduced ejection fraction) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has heart failure: and
- 2 Patient is in NYHA functional class II or III or IV: and
- 3 Fither:
  - 3.1 Patient has a documented left ventricular ejection fraction (LVEF) of less than or equal to 40%; or
  - 3.2 An ECHO is not reasonably practicable, and in the opinion of the treating practitioner the patient would benefit from treatment; and

continued...

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

4 Patient is receiving concomitant optimal standard funded chronic heart failure treatment.

Initial application — (Type 2 Diabetes) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

### Either:

- 1 Patient has previously received an initial approval for a GLP-1 agonist; or
- 2 All of the following:
  - 2.1 Patient has type 2 diabetes; and
  - 2.2 Any of the following:
    - 2.2.1 Patient is Māori or any Pacific ethnicity\*; or
    - 2.2.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)\*; or
    - 2.2.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator\*; or
    - 2.2.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult\*; or
    - 2.2.5 Patient has diabetic kidney disease (see note b)\*; and

EMPAGLIFLOZIN - Special Authority see SA2408 on the previous page - Retail pharmacy

2.3 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of at least one blood-glucose lowering agent (e.g. metformin, vildagliptin, or insulin) for at least 3 months.

Notes: \* Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

- a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack, ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.
- b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m2 in the presence of diabetes, without alternative cause.
- c) Funded [empagliflozin / empagliflozin with metformin hydrochloride] treatment is not to be given in combination with a funded GLP-1 unless receiving (empagliflozin / empagliflozin with metformin hydrochloride] for the treatment of heart failure.

★ Tab 10 mg		58.56	30	•	Jardiance	
* Tab 25 mg		58.56	30	✓	Jardiance	
<b>EMPAGLIFLOZIN</b>	WITH METFORMIN HYDROCHLORIDE - Spe	ecial Authority see	SA2408 or	n the p	revious page – F	<b>?etail</b>
pharmacy						
	1,000 mg metformin hydrochloride		60	1	Jardiamet	
* Tab 5 mg with	500 mg metformin hydrochloride	58.56	60	1	Jardiamet	
* Tab 12.5 mg v	with 1,000 mg metformin hydrochloride	58.56	60	1	Jardiamet	
* Tab 12.5 mg v	with 500 mg metformin hydrochloride	58.56	60	1	Jardiamet	

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

## **Diabetes Management**

## **Ketone Testing**

BLOOD KETONE DIAGNOSTIC TEST STRIP - Subsidy by endorsement

- a) Not on a BSO
- b) Maximum of 20 strip per prescription
- c) Up to 10 strip available on a PSO
- d) Patient has any of the following:
  - 1) type 1 diabetes; or
  - 2) permanent neonatal diabetes: or
  - 3) undergone a pancreatectomy; or
  - 4) cystic fibrosis-related diabetes; or
  - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly.

## **Dual Blood Glucose and Blood Ketone Testing**

DUAL BLOOD GLUCOSE AND BLOOD KETONE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A dual blood glucose and blood ketone diagnostic test meter is subsidised for a patient who has:
  - 1) type 1 diabetes; or
  - 2) permanent neonatal diabetes: or
  - 3) undergone a pancreatectomy; or
  - 4) cystic fibrosis-related diabetes; or
  - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly. Only 1 meter per patient will be subsidised (no repeat prescriptions). For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Meter with 50 lancets, a lancing device and 10 blood glucose

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

## **Blood Glucose Testing**

BLOOD GLUCOSE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A diagnostic blood glucose test meter is subsidised for a patient who:
  - 1) is receiving insulin or sulphonylurea therapy; or
  - 2) is pregnant with diabetes: or
  - 3) is on home TPN at risk of hypoglycaemia or hyperglycaemia; or
  - 4) has a genetic or an acquired disorder of glucose homeostasis, excluding type 1 or type 2 diabetes and metabolic syndrome.

The prescription must be endorsed accordingly. Only one CareSens meter per patient will be subsidised (no repeat prescriptions). Patients already using the CareSens N POP meter and CareSens N meter are not eligible for a new meter, unless they have:

- 1) type 1 diabetes; or
- 2) permanent neonatal diabetes: or
- 3) undergone a pancreatectomy; or
- 4) cystic fibrosis-related diabetes.

For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Note: Only 1 meter available per PSO

BLOOD GLUCOSE DIAGNOSTIC TEST STRIP - Up to 50 test available on a PSO

The number of test strips available on a prescription is restricted to 50 unless:

- Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the
  prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed; or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly; or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

### BLOOD GLUCOSE TEST STRIPS (VISUALLY IMPAIRED)

The number of test strips available on a prescription is restricted to 50 unless:

- Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the
  prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed; or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly: or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

	Subsidy	Fully	y Brand or	
(Manu	ufacturer's Price)	Subsidise	d Generic	
	\$ P	Per 🗸	Manufacturer	

## **Insulin Syringes and Needles**

Subsidy is available for disposable insulin syringes, needles, and pen needles if prescribed on the same form as the one used for the supply of insulin or liraglutide or when prescribed for a patient and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of insulin or liraglutide.

INSULIN PEN NEEDLES	<ul> <li>Maximum of 200</li> </ul>	dev per prescription
---------------------	------------------------------------	----------------------

*	29 g × 12.7 mm	10.05	100	✓ B-D Micro-Fine
	31 g × 5 mm		100	✓ B-D Micro-Fine
*	31 g × 6 mm	9.50	100	✓ Berpu
*	31 g × 8 mm	10.95	100	<ul> <li>B-D Micro-Fine</li> </ul>
*	32 g × 4 mm	10.95	100	✓ B-D Micro-Fine
INS	SULIN SYRINGES, DISPOSABLE WITH ATTACHED NEEDLE	- Maximum of 2	200 dev per p	orescription
*	Syringe 0.3 ml with 29 g x 12.7 mm needle	13.56	100	✓ B-D Ultra Fine
	, ,	1.36	10	
		(1.99)		B-D Ultra Fine
*	Syringe 0.3 ml with 31 g × 8 mm needle	13.56	100	✓ B-D Ultra Fine II
	, ,	1.30	10	
		(1.99)		B-D Ultra Fine II
*	Syringe 0.5 ml with 29 g x 12.7 mm needle	13.56	100	✓ B-D Ultra Fine
	, ,	1.36	10	
		(1.99)		B-D Ultra Fine
*	Syringe 0.5 ml with 31 g × 8 mm needle	13.56	100	✓ B-D Ultra Fine II
	, ,	1.36	10	
		(1.99)		B-D Ultra Fine II
*	Syringe 1 ml with 29 g x 12.7 mm needle	13.56	100	B-D Ultra Fine
	, ,	1.36	10	
		(1.99)		B-D Ultra Fine
*	Syringe 1 ml with 31 g × 8 mm needle	13.56	100	✓ B-D Ultra Fine II
	, ,	1.36	10	
		(1.99)		B-D Ultra Fine II

## **Insulin Pumps**

INSULIN PUMP WITH ALGORITHM - Special Authority see SA2367 below - Retail pharmacy

- a) Maximum of 1 dev per prescription
- b) Only on a prescription

c) Maximum of 1 insulin pump per patient each fou	r year period.		
Min basal rate 0.02 U/h	8,970.00	1	✓ mylife YpsoPump
			with CamAPS FX
Min basal rate 0.1 U/h	7,653.00	1	✓ Tandem t:slim
			X2 with Basal-IQ
			/ <b>-</b>

✓ Tandem t:slim X2 with Control-IQ+

⇒SA2367 Special Authority for Subsidy

Initial application — (type 1 diabetes) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

continued...

ubsidy cturer's Price) Subs	Fully	Brand or Generic
 \$ Per	•	Manufacturer

continued...

- 1 Any of the following:
  - 1.1 The patient has type 1 diabetes; or
  - 1.2 The patient has permanent neonatal diabetes or specific monogenic diabetes subtypes with insulin deficiency, considered by the treating endocrinologist as likely to benefit; or
  - 1.3 The patient has Type 3c diabetes considered by the treating endocrinologist as likely to benefit (Type 3c diabetes includes insulin deficiency due to pancreatectomy, insulin deficiency secondary to cystic fibrosis or pancreatitis); or
  - 1.4 The patient has atypical inherited forms of diabetes; and
- 2 Patient has been evaluated by a diabetes multidisciplinary team for their suitability for insulin pump therapy; and
- 3 In the opinion of the treating relevant practitioner the patient would benefit from an Automated Insulin Delivery (AID) system.

Renewal — (type 1 diabetes) from any relevant practitioner. Approvals valid for 6 months where the patient is continuing to derive benefit according to the treatment plan agreed at induction.

## **Insulin Pump Consumables**

### ⇒SA2536 Special Authority for Subsidy

Initial application — (type 1 diabetes) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 The patient has type 1 diabetes; or
  - 1.2 The patient has permanent neonatal diabetes or specific monogenic diabetes subtypes with insulin deficiency, considered by the treating endocrinologist as likely to benefit; or
  - 1.3 The patient has Type 3c diabetes considered by the treating endocrinologist as likely to benefit (Type 3c diabetes includes insulin deficiency due to pancreatectomy, insulin deficiency secondary to cystic fibrosis or pancreatitis); or 1.4 The patient has atypical inherited forms of diabetes; and
- 2 Patient has been evaluated by a diabetes multidisciplinary team for their suitability for insulin pump therapy; and
- 3 In the opinion of the treating relevant practitioner the patient would benefit from an Automated Insulin Delivery (AID) system.

INSULIN PUMP CARTRIDGE - Special Authority see SA2536 above - Retail pharmacy

- a) Maximum of 50 cart per prescription
- b) Only on a prescription
- c) Maximum of 190 cartridges will be funded per year.

1 OP

Eully.

Drand or

✓ MiniMed Sure-T

MMT-876A

		(Manufacturer's Price)	Sul Per	bsidised	Generic Manufacturer	
INS	SULIN PUMP INFUSION SET (STEEL CANNULA) — Special / a) Maximum of 5 set per prescription b) Only on a prescription c) Maximum of 19 infusion sets will be funded per year.	Authority see SA253	6 on the p	orevious p	page – Retail pharmacy	
*	6 mm steel needle; 60 cm tubing × 10	219.00	1 OP		liniMed Sure-T MMT-864A	
*	6 mm steel needle; 80 cm tubing × 10	219.00	1 OP		liniMed Sure-T MMT-866A	
*	8 mm steel needle; 60 cm tubing × 10	219.00	1 OP		liniMed Sure-T MMT-874A	

Subeidy

(MiniMed Sure-T MMT-864A 6 mm steel needle; 60 cm tubing × 10 to be delisted 1 October 2026) (MiniMed Sure-T MMT-866A 6 mm steel needle; 80 cm tubing × 10 to be delisted 1 October 2026) (MiniMed Sure-T MMT-874A 8 mm steel needle; 60 cm tubing × 10 to be delisted 1 October 2026) (MiniMed Sure-T MMT-876A 8 mm steel needle; 80 cm tubing × 10 to be delisted 1 October 2026)

8 mm steel needle; 80 cm tubing × 10 ......219.00

INSULIN PUMP INFUSION SET (STEEL CANNULA, STRAIGHT INSERTION) - Special Authority see SA2536 on the previous page - Retail pharmacy

- a) Maximum of 5 sets per prescription
- b) Only on a prescription
- c) Maximum of 19 infusion sets will be funded per year.

	-,			
*	5.5 mm steel cannula; straight insertion; 45 cm line × 10 with 10 needles	136.00	1 OP	✓ mylife Orbit micro
*	5.5 mm steel needle; straight insertion; 60 cm line × 10 with 10 needles	136.00	1 OP	✓ mylife Orbit micro
*	5.5 mm steel needle; straight insertion; 80 cm line × 10 with 10 needles	136.00	1 OP	✓ mylife Orbit micro
*	8.5 mm steel needle; straight insertion; 60 cm line × 10 with 10 needles	136.00	1 OP	✓ mylife Orbit micro
*	8.5 mm steel needle; straight insertion; 80 cm line × 10 with 10 needles	136.00	1 OP	✓ mylife Orbit micro
*	6 mm steel cannula; straight insertion; 80 cm line x 10 with 10 needles	182.00	1 OP	✓ TruSteel
*	8 mm steel cannula; straight insertion; 80 cm line x 10 with 10 needles	182.00	1 OP	✓ TruSteel
*	6 mm steel cannula; straight insertion; 60 cm line × 10 with 10 needles		1 OP	✓ TruSteel
*	8 mm steel cannula; straight insertion; 60 cm line $\times$ 10 with		1 OP	✓ TruSteel
	10 needles	102.00	i Or	• IIUJICCI

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

INSULIN PUMP INFUSION SET (TEFLON CANNULA) - Special Authority see SA2536 on page 18 - Retail pharmacy

- a) Maximum of 5 set per prescription
- b) Only on a prescription

	c) Maximum of 19 infusion sets will be funded per year.		
*	13 mm teflon needle, 60 cm tubing × 10219.00	1 OP	<ul><li>MiniMed Silhouette MMT-381A</li></ul>
*	17 mm teflon needle, 110 cm tubing × 10219.00	1 OP	<ul><li>MiniMed Silhouette MMT-377A</li></ul>
*	17 mm teflon needle, 60 cm tubing × 10219.00	1 OP	<ul><li>MiniMed Silhouette MMT-378A</li></ul>
*	6 mm teflon needle, 110 cm tubing x 10219.00	1 OP	MiniMed Quick-Set MMT-398A
*	6 mm teflon needle, 45 cm blue tubing × 10130.00	1 OP	✓ MiniMed Mio MMT-941A
*	6 mm teflon needle, 45 cm pink tubing x 10130.00	1 OP	✓ MiniMed Mio MMT-921A
*	6 mm teflon needle, 60 cm blue tubing × 10130.00	1 OP	✓ MiniMed Mio MMT-943A
*	6 mm teflon needle, 60 cm pink tubing x 10130.00	1 OP	✓ MiniMed Mio MMT-923A
*	6 mm teflon needle, 60 cm tubing × 10219.00	1 OP	<ul><li>MiniMed Quick-Set MMT-399A</li></ul>
*	6 mm teflon needle, 80 cm blue tubing	1 OP	✓ MiniMed Mio MMT-945A
*	6 mm teflon needle, 80 cm clear tubing × 10130.00	1 OP	✓ MiniMed Mio MMT-965A
*	6 mm teflon needle, 80 cm pink tubing × 10130.00	1 OP	✓ MiniMed Mio MMT-925A
*	9 mm teflon needle, 110 cm tubing × 10219.00	1 OP	<ul><li>MiniMed Quick-Set MMT-396A</li></ul>
*	9 mm teflon needle, 60 cm tubing × 10219.00	1 OP	✓ MiniMed Quick-Set MMT-397A
*	9 mm teflon needle, 80 cm clear tubing × 10130.00	1 OP	✓ MiniMed Mio MMT-975A

(MiniMed Silhouette MMT-381A 13 mm teflon needle, 60 cm tubing × 10 to be delisted 1 October 2026) (MiniMed Silhouette MMT-377A 17 mm teflon needle, 110 cm tubing × 10 to be delisted 1 October 2026) (MiniMed Silhouette MMT-378A 17 mm teflon needle, 60 cm tubing × 10 to be delisted 1 October 2026) (MiniMed Quick-Set MMT-398A 6 mm teflon needle, 110 cm tubing × 10 to be delisted 1 October 2026) (MiniMed Mio MMT-941A 6 mm teflon needle, 45 cm pink tubing × 10 to be delisted 1 February 2026) (MiniMed Mio MMT-921A 6 mm teflon needle, 45 cm pink tubing × 10 to be delisted 1 February 2026) (MiniMed Mio MMT-943A 6 mm teflon needle, 60 cm blue tubing × 10 to be delisted 1 February 2026) (MiniMed Mio MMT-943A 6 mm teflon needle, 60 cm pink tubing × 10 to be delisted 1 February 2026) (MiniMed Quick-Set MMT-399A 6 mm teflon needle, 60 cm blue tubing to be delisted 1 October 2026) (MiniMed Mio MMT-945A 6 mm teflon needle, 80 cm blue tubing to be delisted 1 February 2026) (MiniMed Mio MMT-95A 6 mm teflon needle, 80 cm clear tubing × 10 to be delisted 1 February 2026) (MiniMed Mio MMT-95A 9 mm teflon needle, 110 cm tubing × 10 to be delisted 1 October 2026) (MiniMed Quick-Set MMT-396A 9 mm teflon needle, 60 cm tubing × 10 to be delisted 1 October 2026) (MiniMed Quick-Set MMT-397A 9 mm teflon needle, 60 cm tubing × 10 to be delisted 1 October 2026) (MiniMed Quick-Set MMT-397A 9 mm teflon needle, 60 cm tubing × 10 to be delisted 1 February 2026) (MiniMed Mio MMT-975A 9 mm teflon needle, 60 cm tubing × 10 to be delisted 1 February 2026)

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per Manufacturer INSULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE INSERTION WITH INSERTION DEVICE) - Special Authority see SA2536 on page 18 - Retail pharmacy a) Maximum of 5 sets per prescription b) Only on a prescription c) Maximum of 19 infusion sets will be funded per year. \* 6 mm teflon cannula; angle insertion; insertion device; 60 cm line × 4 with 4 needles 210.00 1 OP ✓ Mio Adv Ext Infusion Set MDT-MMT-431AK \* 6 mm teflon cannula; angle insertion; insertion device; 80 cm line x 4 with 4 needles 210.00 1 OP ✓ Mio Adv Ext Infusion Set MDT-MMT-432AK \* 9 mm teflon cannula; angle insertion; insertion device; 60 cm line × 4 with 4 needles......210.00 ✓ Mio Adv Ext 1 OP Infusion Set MDT-MMT-441AK \* 9 mm teflon cannula; angle insertion; insertion device; 80 cm line x 4 with 4 needles 210.00 1 OP ✓ Mio Adv Ext Infusion Set MDT-MMT-442AK 13 mm teflon cannula: angle insertion: insertion device: 110 cm 1 OP ✓ AutoSoft 30 \* 13 mm teflon cannula; angle insertion; insertion device; 60 cm 1 OP ✓ AutoSoft 30 (Mio Adv Ext Infusion Set MDT-MMT-431AK 6 mm teflon cannula; angle insertion; insertion device; 60 cm line x 4 with 4 needles to be delisted 1 October 2026) (Mio Adv Ext Infusion Set MDT-MMT-432AK 6 mm teflon cannula; angle insertion; insertion device; 80 cm line x 4 with 4 needles

to be delisted 1 October 2026) (Mio Adv Ext Infusion Set MDT-MMT-441AK 9 mm teflon cannula; angle insertion; insertion device; 60 cm line x 4 with 4 needles

to be delisted 1 October 2026)

(Mio Adv Ext Infusion Set MDT-MMT-442AK 9 mm teflon cannula; angle insertion; insertion device; 80 cm line x 4 with 4 needles to be delisted 1 October 2026)

INSULIN PUMP INFUSION SET (TEFLON CANNULA, FLEXIBLE INSERTION WITH INSERTION DEVICE) - Special Authority see SA2536 on page 18 - Retail pharmacy

- a) Maximum of 5 set per prescription
- b) Only on a prescription
- c) Maximum of 19 infusion sets will be funded per year.

*	6 mm teflon cannula; flexible insertion; insertion device; 46 cm line × 10 with 10 needles	. 157.00	1 OP	✓ mylife Inset soft
*	6 mm teflon cannula; flexible insertion; insertion device; 60 cm line with integrated inserter × 10 with 10 needles	. 157.00	1 OP	✓ mylife Inset soft
*	6 mm teflon cannula; flexible insertion; insertion device; 80 cm line × 10 with 10 needles	. 157.00	1 OP	✓ mylife Inset soft
*	9 mm teflon cannula; flexible insertion; insertion device; 60 cm line × 10 with 10 needles	. 157.00	1 OP	✓ mylife Inset soft
*	9 mm teflon cannula; flexible insertion; insertion device; 80 cm line × 10 with 10 needles	. 157.00	1 OP	✓ mylife Inset soft

Subsidy		Fully	Brand or
(Manufacturer's Price)	S	ubsidised	Generic
 \$	Per	✓	Manufacturer

INSULIN PUMP INFUSION SET (TEFLON CANNULA, STRAIGHT INSERTION WITH INSERTION DEVICE) - Special Authority see SA2536 on page 18 - Retail pharmacy

- a) Maximum of 5 sets per prescription
- b) Only on a prescription
- c) Maximum of 19 infusion sets will be funded per year.
- \* 6 mm teflon cannula; straight insertion; insertion device; 1 OP ✓ AutoSoft 90 \* 6 mm teflon cannula: straight insertion; insertion device: 60 cm 1 OP ✓ AutoSoft 90 \* 9 mm teflon cannula; straight insertion; insertion device; 1 OP ✓ AutoSoft 90 \* 9 mm teflon cannula; straight insertion; insertion device; 60 cm ✓ AutoSoft 90 1 OP

INSULIN PUMP INFUSION SET (TEFLON CANNULA, VARIABLE INSERTION) - Special Authority see SA2536 on page 18 -Retail pharmacy

- a) Maximum of 5 set per prescription
- b) Only on a prescription
- c) Maximum of 19 infusion sets will be funded per year.
- \* 13 mm teflon cannula: variable insertion: 60 cm line × 10 with

✓ VariSoft 1 OP INSULIN PUMP RESERVOIR - Special Authority see SA2536 on page 18 - Retail pharmacy

- a) Maximum of 90 cart per prescription
- b) Only on a prescription
- c) Maximum of 360 reservoirs will be funded per year.

10 .. 1 6 ml along recompoir for Vaca Duma

不	10 x 1.6 mi glass reservoir for 1 psorump50.00	10 0P	Reservoir
*	10 × luer lock conversion cartridges 1.8 ml for paradigm pumps 50.00	10 OP	✓ ADR Cartridge 1.8
*	Cartridge for 7 series pump; 3.0 ml × 1098.00	10 OP	✓ MiniMed
	-		3.0 Reservoir MMT-332A
*	Cartridge for 7 series pump; 3.0 ml × 5102.00	5 OP	✓ MiniMed 3.0 Ext

E0 00

10 OD

/ mulifa Vmaa Dumu

Reservoir MDT-MMT-342G

(ADR Cartridge 1.8 10 × luer lock conversion cartridges 1.8 ml for paradigm pumps to be delisted 1 October 2026) (MiniMed 3.0 Reservoir MMT-332A Cartridge for 7 series pump; 3.0 ml × 10 to be delisted 1 October 2026) (MiniMed 3.0 Ext Reservoir MDT-MMT-342G Cartridge for 7 series pump; 3.0 ml x 5 to be delisted 1 October 2026)

### **Continuous Glucose Monitor**

CONTINUOUS GLUCOSE MONITOR (INTEROPERABLE) - Special Authority see SA2537 on the next page - Retail pharmacy Only on a prescription

	o, o a p. ooop		
*	Sensor (9) and transmitter (Dexcom G6) - Maximum of 1 dev		
	per prescription990.00	1 OP	✓ Dexcom G6
	Maximum of 5 dev will be funded per year.		
*	Sensor (Dexcom G7) – Maximum of 9 dev per prescription110.00	1	Dexcom G7
	Maximum of 40 dev will be funded per year.		
*	Sensor (Freestyle Libre 3 Plus) - Maximum of 6 dev per		
	prescription99.46	1	✓ Freestyle Libre
			3 Plus

Maximum of 28 dev will be funded per year.

Subsidy (Manufacturer's P	rice)	Fully Subsidised	Brand or Generic	
\$	Per	•	Manufacturer	

### **⇒SA2537** Special Authority for Subsidy

Initial application — (type 1 diabetes) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

### Both:

- 1 Any of the following:
  - 1.1 The patient has type 1 diabetes; or
  - 1.2 The patient has permanent neonatal diabetes or specific monogenic diabetes subtypes with insulin deficiency, considered by the treating endocrinologist or relevant secondary health care professional as practicable, as likely to benefit or
  - 1.3 The patient has Type 3c diabetes considered by the treating endocrinologist or relevant secondary health care professional as practicable, as likely to benefit (Type 3c diabetes includes insulin deficiency due to pancreatectomy, insulin deficiency secondary to cystic fibrosis or pancreatitis); or
  - 1.4 The patient has atypical inherited forms of diabetes; and
- 2 In the opinion of the treating relevant practitioner the patient would benefit from an Automated Insulin Delivery (AID) system.

## CONTINUOUS GLUCOSE MONITOR (STANDALONE) - Special Authority see SA2538 below - Retail pharmacy Only on a prescription

*	Sensor (Dexcom ONE+) - Maximum of 9 dev per prescription81.00	✓ Dexcom ONE+
	Maximum of 40 dev will be funded per year.	

\* Sensor (Freestyle Libre 2 Plus) – Maximum of 6 dev per

prescription 99.46 1 Freestyle Libre

Maximum of 28 dev will be funded per year.

★ Sensor (Freestyle Libre 2) – Maximum of 7 dev per prescription.....92.83
1
✓ Freestyle Libre 2
Maximum of 29 dev will be funded per year.

(Freestyle Libre 2 Sensor (Freestyle Libre 2) to be delisted 1 May 2026)

### ⇒SA2538 Special Authority for Subsidy

Initial application — (type 1 diabetes) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

### Any of the following:

- 1 The patient has type 1 diabetes; or
- 2 The patient has permanent neonatal diabetes or specific monogenic diabetes subtypes with insulin deficiency, considered by the treating endocrinologist or relevant secondary health care professional as practicable, as likely to benefit; or
- 3 The patient has Type 3c diabetes considered by the treating endocrinologist or relevant secondary health care professional as practicable, as likely to benefit (Type 3c diabetes includes insulin deficiency due to pancreatectomy, insulin deficiency secondary to cystic fibrosis or pancreatitis); or
- 4 The patient has atypical inherited forms of diabetes.

## **Digestives Including Enzymes**

### PANCREATIC ENZYME

Cap pancreatin 150 mg (amylase 8,000 Ph Eur U, lipase		
10,000 Ph Eur U, total protease 600 Ph Eur U)34.93	100	✓ Creon 10000
Cap pancreatin 300 mg (amylase 18,000 Ph Eur U, lipase		
25,000 Ph Eur U, total protease 1,000 Ph Eur U)94.38	100	✓ Creon 25000
Modified release granules pancreatin 60.12 mg (amylase		
3,600 Ph Eur U, lipase 5,000 Ph Eur U, protease 200 Ph		
Eur U)	20 g OP	Creon Micro

	(Manufacturer's Price)	Per	Subsidised	Generic Manufacturer	
URSODEOXYCHOLIC ACID – Special Authority see SA2448 be Cap 250 mg	'	y 100	<b>√</b> <u>l</u>	Jrsosan_	

⇒SA2448 Special Authority for Subsidy

Initial application — (Alagille syndrome or progressive familial intrahepatic cholestasis) from any relevant practitioner.

Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 Patient has been diagnosed with Alagille syndrome; or
- 2 Patient has progressive familial intrahepatic cholestasis.

**Initial application** — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic severe drug induced cholestatic liver injury; and
- 2 Cholestatic liver injury not due to Total Parenteral Nutrition (TPN) use in adults; and
- 3 Treatment with ursodeoxycholic acid may prevent hospital admission or reduce duration of stay.

**Initial application — (Primary biliary cholangitis)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Primary biliary cholangitis confirmed by antimitochondrial antibody titre (AMA) > 1:80, and raised cholestatic liver enzymes with or without raised serum IgM or, if AMA is negative, by liver biopsy; and
- 2 Patient not requiring a liver transplant (bilirubin > 100 umol/l; decompensated cirrhosis).

**Initial application** — (**Pregnancy**) from any relevant practitioner. Approvals valid for 6 months where the patient diagnosed with cholestasis of pregnancy.

Initial application — (Haematological Transplant) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient at risk of veno-occlusive disease or has hepatic impairment and is undergoing conditioning treatment prior to allogenic stem cell or bone marrow transplantation; and
- 2 Treatment for up to 13 weeks.

Initial application — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Paediatric patient has developed abnormal liver function as indicated on testing which is likely to be induced by Total Parenteral Nutrition (TPN); and
- 2 Liver function has not improved with modifying the TPN composition.

Renewal — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 6 months where the patient continues to benefit from treatment.

Renewal — (Pregnancy/Primary biliary cholangitis) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months where the paediatric patient continues to require TPN and who is benefiting from treatment, defined as a sustained improvement in bilirubin levels.

Initial application — (prevention of sinusoidal obstruction syndrome) from any relevant practitioner. Approvals valid without further renewal unless notified where the individual has leukaemia/lymphoma and requires prophylaxis for medications/therapies with a high risk of sinusoidal obstruction syndrome.

### Laxatives

## **Bulk-forming Agents**

ISPAGHULA (PSYLLIUM) HUSK - Only on a prescription

500 q OP

✓ Konsyl-D

Fully

Brand or

✓ Relistor

	Manufacturer's Pric \$	e) Per	Subsidised	
Faecal Softeners				
DOCUSATE SODIUM - Only on a prescription				
* Tab 50 mg	3.20	100	1	Coloxyl
* Tab 120 mg	4.98	100	✓	Coloxyl
DOCUSATE SODIUM WITH SENNOSIDES				
Tab 50 mg with sennosides 8 mg	1.50	100	1	Solax
· ·	3.50	200	1	Laxsol
(Laxsol Tab 50 mg with sennosides 8 mg to be delisted 1 May 202	6)			
POLOXAMER – Only on a prescription  Not funded for use in the ear.				
* Oral drops 10%	4.17	30 ml (	OP 🗸	Coloxyl
Opioid Receptor Antagonists - Peripheral				
METHYLNALTREXONE BROMIDE - Special Authority see SA16	91 below – Retail	pharm	acy	

Subsidy

# Inj 12 mg per 0.6 ml vial ...... SA1691 Special Authority for Subsidy

Initial application — (Opioid induced constipation) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

246.00

- 1 The patient is receiving palliative care; and
- 2 Either:
  - 2.1 Oral and rectal treatments for opioid induced constipation are ineffective; or
  - 2.2 Oral and rectal treatments for opioid induced constipation are unable to be tolerated.

### **Osmotic Laxatives**

12.39	20	<ul><li>Lax-suppositories</li><li>Glycerol</li></ul>
ebruary 2026		
6.16	500 ml	✓ Laevolac
ICARBONATE A	ND SODIUM (	CHLORIDE
ng,		
		_
	30	✓ Molaxole
12.19		✓ Movicol
3.70	1	✓ Fleet Phosphate  Enema
E − Only on a pre	escription	
, ,		
•	50	✓ Micolette
	ng, 10.15 12.19 3.70	February 2026

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

SODIUM PICOSULFATE – Special Authority see SA2053 below – Retail pharmacy Oral soln 7.5 mg per ml ......7.40

	\$	Per	Manutacturer
Stimulant Laxatives			
BISACODYL - Only on a prescription			
* Tab 5 mg	10.00	200	<ul> <li>Bisacodyl Viatris</li> </ul>
* Suppos 10 mg	4.14	10	✓ Lax-Suppositories
SENNA - Only on a prescription			
* Tab, standardised	2.17	100	
	(9.38)		Senokot
	0.43	20	

Subsidy

(Manufacturer's Price)

(2.06)

30 ml OP

Fully

Subsidised

Brand or

Generic

Senokot

✓ Dulcolax SP Drop

## ⇒SA2053 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The patient is a child with problematic constipation despite an adequate trial of other oral pharmacotherapies including macrogol where practicable; and
- 2 The patient would otherwise require a high-volume bowel cleansing preparation or hospital admission.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

## Metabolic Disorder Agents

### ⇒SA1986 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient is aged up to 24 months at the time of initial application and has been diagnosed with infantile Pompe disease; and
- 2 Any of the following:
  - 2.1 Diagnosis confirmed by documented deficiency of acid alpha-glucosidase by prenatal diagnosis using chorionic villus biopsies and/or cultured amniotic cells; or
  - 2.2 Documented deficiency of acid alpha-glucosidase, and urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides; or
  - 2.3 Documented deficiency of acid alpha-glucosidase, and documented molecular genetic testing indicating a disease-causing mutation in the acid alpha-glucosidase gene (GAA gene); or
  - 2.4 Documented urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides, and molecular genetic testing indicating a disease-causing mutation in the GAA gene; and
- 3 Patient has not required long-term invasive ventilation for respiratory failure prior to starting enzyme replacement therapy (ERT); and
- 4 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by ERT or might be reasonably expected to compromise a response to ERT; and
- 5 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks.

**Renewal** only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and

continued...

ubsidy	Fully	Brand or
cturer's Price) S	Subsidised	Generic
 \$ Per	•	

### continued...

- 2 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks; and
- 3 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 4 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by ERT; and
- 5 Patient has not developed another medical condition that might reasonably be expected to compromise a response to FRT; and
- 6 There is no evidence of life threatening progression of respiratory disease as evidenced by the needed for > 14 days of invasive ventilation; and
- 7 There is no evidence of new or progressive cardiomyopathy.

ARGININE – Special Authority see SA2042 below – Retail pharmacy	
Tob 1 000 mg	-

Tab 1,000 mg	CBS	90	Clinicians
Cap 500 mg		50	✓ Solgar
Powder		400 g	Biomed

### ⇒SA2042 Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to arginine supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to arginine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

## ⇒SA1987 Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has a confirmed diagnosis of homocystinuria; and
- 2 Any of the following:
  - 2.1 A cystathionine beta-synthase (CBS) deficiency; or
  - 2.2 A 5,10-methylene-tetrahydrofolate reductase (MTHFR) deficiency; or
  - 2.3 A disorder of intracellular cobalamin metabolism; and
- 3 An appropriate homocysteine level has not been achieved despite a sufficient trial of appropriate vitamin supplementation.

**Renewal** only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

COENZYME Q10 - Special Authority see SA2039 below	<ul> <li>Retail pharmacy</li> </ul>		
Cap 120 mg	CBS	30	✓ Solgar
Cap 160 mg	CBS	60	✓ Go Healthy

### ⇒SA2039 Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to coenzyme Q10 supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to coenzyme Q10 supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

GALSULFASE − Special Authority see SA1988 on the next page − Retail pharmacy
Inj 1 mg per ml, 5 ml vial......2,234.00 1 ✓ Naglazyme

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	I Generic	
\$	Per 🗸	Manufacturer	

### ⇒SA1988 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has been diagnosed with mucopolysaccharidosis VI; and
- 2 Either:
  - 2.1 Diagnosis confirmed by demonstration of N-acetyl-galactosamine-4-sulfatase (arylsulfatase B) deficiency by either enzyme activity assay in leukocytes or skin fibroblasts; or
  - 2.2 Detection of two disease causing mutations and patient has a sibling who is known to have mucopolysaccharidosis

Renewal only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 3 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by Enzyme Replacement Therapy (ERT); and
- 4 Patient has not developed another medical condition that might reasonably be expected to compromise a response to

IDURSULFASE - Special Authority see SA1623 below - Retail pharmacy 

✓ Elaprase

## **⇒SA1623** Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hunter Syndrome (mucopolysaccharidosis II); and
- 2 Fither:
  - 2.1 Diagnosis confirmed by demonstration of iduronate 2-sulfatase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts; or
  - 2.2 Detection of a disease causing mutation in the iduronate 2-sulfatase gene; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with idursulfase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT); and
- 5 Idursulfase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 weeks post-HSCT) at doses no greater than 0.5 mg/kg every week.

LARONIDASE - Special Authority see SA1695 below - Retail pharmacy

✓ Aldurazyme

### ⇒SA1695 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hurler Syndrome (mucopolysacchardosis I-H); and
- 2 Fither:
  - 2.1 Diagnosis confirmed by demonstration of alpha-L-iduronidase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts; or
  - 2.2 Detection of two disease causing mutations in the alpha-L-iduronidase gene and patient has a sibling who is known to have Hurler syndrome; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with laronidase would be bridging treatment to transplant; and

continued...

Subsidy	)	Fully	Brand or
(Manufacturer's Price		Subsidised	Generic
\$	Per	✓	Manufacturer

### continued...

- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT): and
- 5 Laronidase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 post-HSCT) at doses no greater than 100 units/kg every week.

LEVOCARNITINE - Special Authority see SA2040 below	- Retail pharmacy		
Tab 500 mg	CBS	30	✓ Solgar
Cap 250 mg	CBS	30	✓ Solgar
Cap 500 mg	CBS	60	✓ Balance
, ,		300	✓ Metabolics
Oral liq 1 g per 10 ml	CBS	118 ml	✓ Lacuna S29
, -,			✓ Novitium Sugar
			Free S29
Oral lig 500 mg per 10 ml	CBS	300 ml	✓ Balance
41 11 0 5			

(Novitium Sugar Free S29 Oral lig 1 g per 10 ml to be delisted 1 February 2026)

## ⇒SA2040 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to carnitine supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to carnitine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

RIBOFLAVIN – Special Authority see SA2041 below – Retail pharmacy Tab 100 mgCBS	100	✓ Country Life ✓ Puritan's Pride Vitamin B-2 100 mg \$29
Cap 100 mgCBS	100	✓ Solgar

### ⇒SA2041 Special Authority for Subsidy

**Initial application** only from a metabolic physician or neurologist. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to riboflavin supplementation.

Renewal only from a metabolic physician or neurologist. Approvals valid for 24 months for applications meeting the following criteria:

### Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to riboflavin supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

SAPROPTERIN DIHYDROCHLORIDE – Special Authority see SA1989 below – Retail pharmacy
Tab soluble 100 mg.......1,452.70 30 OP 
✓ Kuvan

### ⇒SA1989 Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 1 month for applications meeting the following criteria: All of the following:

- 1 Patient has phenylketonuria (PKU) and is pregnant or actively planning to become pregnant; and
- 2 Treatment with sapropterin is required to support management of PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and

continued...

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

becoming pregnant) and treatment will be stopped after delivery.

**Renewal** only from a metabolic physician or any relevant practitioner on the recommendation of a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
  - 1.1 Following the initial one-month approval, the patient has demonstrated an adequate response to a 2 to 4 week trial of sapropterin with a clinically appropriate reduction in phenylalanine levels to support management of PKU during pregnancy; or
  - 1.2 On subsequent renewal applications, the patient has previously demonstrated response to treatment with sapropterin and maintained adequate phenylalanine levels to support management of PKU during pregnancy; and
- 2 Any of the following:
  - 2.1 Patient continues to be pregnant and treatment with sapropterin will not continue after delivery; or
  - 2.2 Patient is actively planning a pregnancy and this is the first renewal for treatment with sapropterin; or
  - 2.3 Treatment with sapropterin is required for a second or subsequent pregnancy to support management of their PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

### ⇒SA1599 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder.

**Renewal** only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

SODIUM PHENYLBUTYRATE — Special Authority see SA1990 below — Retail pharmacy
Grans 483 mg per g......2,016.00 174 g OP 

✓ Pheburane

### ⇒SA1990 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder involving a deficiency of carbamylphosphate synthetase, ornithine transcarbamylase or argininosuccinate synthetase.

**Renewal** only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

TAURINE - Special Authority see SA2043 below - Retail pharmacy

Cap 500 mg	CBS	50	✓ Solgar
Cap 1,000 mg		90	✓ Life Extension
Powder	CBS	300 a	✓ Life Extension

### ⇒SA2043 Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 6 months where patient has a suspected specific mitochondrial disorder that may respond taurine supplementation.

**Renewal** only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

1 The patient has confirmed diagnosis of a specific mitochondrial disorder which responds to taurine supplementation; and

2 The treatment remains appropriate and the patient is benefiting from treatment.

TRIENTINE - Special Authority see SA2324 on the next page - Retail pharmacy

Subsidy	Fu	ılly Brand or	
(Manufacturer's Price)	Subsidis	ed Generic	
\$	Per	<ul> <li>Manufacture</li> </ul>	r

### ⇒SA2324 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Wilson disease; and
- 2 Treatment with D-penicillamine has been trialled and discontinued because the person has experienced intolerable side effects or has not received sufficient benefit; and
- 3 Treatment with zinc has been trailled and discontinued because the person has experienced intolerable side effects or has not received sufficient benefit, or zinc is considered clinically inappropriate as the person has symptomatic liver disease and requires copper chelation.

### Gaucher's Disease

### ⇒SA2137 Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has a diagnosis of symptomatic type 1 or type 3\* Gaucher disease confirmed by the demonstration of specific deficiency of glucocerebrosidase in leukocytes or cultured skin fibroblasts, and genotypic analysis; and
- 2 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by enzyme replacement therapy (ERT) or the disease might be reasonably expected to compromise a response to ERT; and
- 3 Any of the following:
  - 3.1 Patient has haematological complications of Gaucher disease; or
  - 3.2 Patient has skeletal complications of Gaucher disease; or
  - 3.3 Patient has significant liver dysfunction or hepatomegaly attributable to Gaucher disease; or
  - 3.4 Patient has reduced vital capacity from clinically significant or progressive pulmonary disease due to Gaucher disease; or
  - 3.5 Patient is a child and has experienced growth failure with significant decrease in percentile linear growth over a 6-12 month period; and
- 4 Taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units).

Note: Indication marked with \* is an unapproved indication

Renewal only from a metabolic physician or any relevant practitioner on the recommendation of a metabolic physician.

Approvals valid for 3 years for applications meeting the following criteria:

### All of the following:

- 1 Patient has demonstrated a symptomatic improvement and has maintained improvements in the main symptom or symptoms for which therapy was started; and
- 2 Patient has demonstrated a clinically objective improvement or no deterioration in haemoglobin levels, platelet counts and liver and spleen size; and
- 3 Radiological (MRI) signs of bone activity performed at two years since initiation of treatment, and five yearly thereafter, demonstrate no deterioration shown by the MRI, compared with MRI taken immediately prior to commencement of therapy or adjusted dose; and
- 4 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT; and
- 5 Patient is adherent with regular treatment and taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units).

	Subsidy (Manufacturer's F	Price) Subs	Fully Brand or idised Generic ✓ Manufacturer
Mouth and Throat			
Agents Used in Mouth Ulceration			
BENZYDAMINE HYDROCHLORIDE  Soln 0.15% – Higher subsidy of \$22.60 per 500 ml with			
Endorsement	(22.60)	500 ml	Difflam
Additional subsidy by endorsement for a patient who has prescription is endorsed accordingly.	oral mucositis a	as a result of tre	eatment for cancer, and the
CARMELLOSE SODIUM WITH GELATIN AND PECTIN			
Paste	17.20 4.55	56.7 g OP 15 g OP	✓ Stomahesive
	(7.90) 1.52	5 g OP	Orabase
Powder	(3.60) 8.48	28 g OP	Orabase
TRIAMCINOLONE ACETONIDE	(10.95)	-	Stomahesive
Paste 0.1%	5.49	5 g OP	✓ Kenalog in Orabase
Oropharyngeal Anti-infectives			
AMPHOTERICIN B Lozenges 10 mg	5.86	20	✓ Fungilin
MICONAZOLE Oral gel 20 mg per g	5.19	40 g OP	✓ <u>Decozol</u>
NYSTATIN Oral liq 100,000 u per ml	2.22	24 ml OP	✓ <u>Nilstat</u>
Vitamins			
Vitamin B			
HYDROXOCOBALAMIN  * Inj 1 mg per ml, 1 ml ampoule – Up to 6 inj available on a PS	O3.95	3	✓ <u>Hydroxocobalamin</u> Panpharma
PYRIDOXINE HYDROCHLORIDE  a) No more than 100 mg per dose			
b) Only on a prescription  * Tab 25 mg - No patient co-payment payable	3.43	90	✓ Vitamin B6 25
* Tab 50 mg		500	✓ Pyridoxine multichem
THIAMINE HYDROCHLORIDE - Only on a prescription  * Tab 50 mg	4.65	100	✓ Thiamine multichem
VITAMIN B COMPLEX  * Tab, strong, BPC		500	✓ Bplex
本 1 au, 5:1011y, DFU	11.25	500	• phiex

Subsidy

Fully

Brand or

		ALIMENTAN	ii in	ACT AI	ID METABOLISM
		Subsidy (Manufacturer's Price \$	e) Per	Fully Subsidised	I Generic
٧	itamin C				
	CORBIC ACID  a) No more than 100 mg per dose b) Only on a prescription Tab 100 mg  Cvite to be Principal Supply on 1 March 2026	16.00	500		Cvite
٧	itamin D				
* * CA	FACALCIDOL Cap 0.25 mcg Cap 1 mcg Oral drops 2 mcg per ml LCITRIOL Cap 0.25 mcg	87.98 60.68	100 100 20 ml C	OP /	One-Alpha One-Alpha One-Alpha Calcitriol XL S29 Calcitriol-AFT
*	Cap 0.5 mcg	13.68	100		Calcitriol XL S29 Calcitriol-AFT
*	DLECALCIFEROL  Cap 1.25 mg (50,000 iu) – Maximum of 12 cap per prescripti  Oral liq 188 mcg per ml (7,500 iu per ml)		12 5 ml O		Vit.D3 Clinicians
N	Iultivitamin Preparations				
*	JLTIVITAMIN RENAL – Special Authority see SA1546 below – Cap		30	✓	Clinicians Renal Vit
lni	tial application from any relevant practitioner. Approvals valid following criteria:	without further ren	iewal ui	nless noti	fied for applications meeting
Eit	ner:  1 The patient has chronic kidney disease and is receiving eit  2 The patient has chronic kidney disease grade 5, defined as 15 ml/min/1.73 m² body surface area (BSA).				•
*	JLTIVITAMINS – Special Authority see SA1036 below – Retail Powder	, ,	200 g C	)P 🗸	Paediatric Seravit
inb Re app	initial application from any relevant practitioner. Approvals valid orn errors of metabolism.  newal from any relevant practitioner. Approvals valid without foroval for multivitamins.  TAMINS				
*	Tab (BPC cap strength)	24.00	1,000	•	Mvite
Т.	SA1720 on the next page – Retail pharmacy	23.40	60	•	Vitabdeck

33

Subsidy (Manufacturer's Price)	Subs	Fully	Brand or Generic
\$	Per	✓	Manufacturer

## ⇒SA1720 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Patient has cystic fibrosis with pancreatic insufficiency; or
- 2 Patient is an infant or child with liver disease or short gut syndrome; or
- 3 Patient has severe malabsorption syndrome.

## **Minerals**

## Calcium

Valoranii		
CALCIUM CARBONATE		
★ Tab 1.25 g (500 mg elemental)	250	✓ Calci-Tab 500
★ Tab eff 1.25 g (500 mg elemental) – Subsidy by endorsement260.00	100	<ul><li>Calcium 500 mg</li></ul>
		Hexal S29
Subsidy by endorsement – Only when prescribed for paediatric patients considered unsuitable.	s (< 5 years) wh	ere calcium carbonate oral liquid
CALCIUM GLUCONATE		
★ Inj 10%, 10 ml ampoule32.00	10	✓ Max Health -
		Hameln S29
lodine		
POTASSIUM IODATE		
* Tab 253 mcg (150 mcg elemental iodine)	90	✓ NeuroTabs
Iron		
ERROUS FUMARATE		
★ Tab 200 mg (65.7 mg elemental)	100	✓ Ferro-tab
ERROUS FUMARATE WITH FOLIC ACID		
★ Tab 310 mg (100 mg elemental) with folic acid 350 mcg	100	✓ Ferro-F-Tabs
ERROUS SULFATE		
Tab long-acting 325 mg (105 mg elemental)2.55	30	✓ Ferrograd
Note: No new patients to be initiated on ferrous sulfate tablet.	00	· i on ograd
♦ Oral lig 30 mg (6 mg elemental) per 1 ml	250 ml	✓ Ferro-Liquid
13.10	500 ml	✓ Ferodan
Ferro-Liquid to be Principal Supply on 1 February 2026		
Ferrograd Tab long-acting 325 mg (105 mg elemental) to be delisted 1 March 2	2026)	
Ferodan Oral liq 30 mg (6 mg elemental) per 1 ml to be delisted 1 February 202		
RON (AS FERRIC CARBOXYMALTOSE) - Special Authority see SA2394 belo	ow – Retail phar	macv
Inj 50 mg per ml, 10 ml vial	1	✓ Ferinject
Special Authority for Subsidy		•
nitial application — (Anaemia) from any relevant practitioner. Approvals vali	d for 3 months f	or applications meeting the

Initial application — (Anaemia) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

continued...

Subsidy		Fully	Brand or
(Manufacturer's Price)		sidised	Generic
 \$	Per	1	

continued...

- 1 Patient has been diagnosed with anaemia; and
- 2 Any of the following:
  - 2.1 Serum ferritin level is 20 mcg/L or less: or
  - 2.2 Both:
    - 2.2.1 Serum ferritin is between 20 and 50 mcg/L; and
    - 2.2.2 C-Reactive Protein (CRP) is at least 5 mg/L; or
  - 2.3 Patient has chronic inflammatory disease with symptoms of anaemia despite normal iron levels; and
- 3 Any of the following:
  - 3.1 Oral iron treatment has proven ineffective; or
  - 3.2 Oral iron treatment has resulted in dose-limiting intolerance; or
  - 3.3 Rapid correction of anaemia is required.

Renewal — (Anaemia) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

#### Both:

- 1 Patient continues to have anaemia with a serum ferritin level of 20 mcg/L, or less or between 20 and 50 mcg/L with CRP of at least 5 mg/L, or has chronic inflammatory disease with symptoms of anaemia despite normal iron levels; and
- 2 A trial (or re-trial) with oral iron is clinically inappropriate.

Initial application — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with iron-deficiency anaemia; and
- 2 Any of the following:
  - 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
  - 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
  - 2.3 Patient has symptomatic heart failure, chronic kidney disease stage 3 or more or active inflammatory bowel disease and a trial of oral iron is unlikely to be effective: or

41 75

2.4 Rapid correction of anaemia is required.

Renewal — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient continues to have iron-deficiency anaemia; and
- 2 A re-trial with oral iron is clinically inappropriate.

### IRON POLYMALTOSE

\* Ini 50 mg per ml 2 ml ampoule

, , , , , , , , , , , , , , , , , , ,	ŭ		
Magnesium			
MAGNESIUM HYDROXIDE Suspension 8%33.60	355 ml	✓ Phillips Milk of Magnesia S29	
MAGNESIUM SULPHATE  * Inj 2 mmol per ml, 5 ml ampoule	10	✓ Martindale	

✓ Ferrosia

✓ Inresa S29

10

	Subsidy (Manufacturer's Price) \$	Sub Per	Fully sidised	Brand or Generic Manufacturer	
Zinc				a.ra.ra.ra.ra.ra.ra.ra.ra.ra.ra.ra.ra	
ZINC SULPHATE					

ZINC SULPHATE			
* Cap 220 mg (50 mg elemental)	29.14	100	✓ Rugby S29
* Cap 137.4 mg (50 mg elemental)	11.00	100	✓ Zincaps
			✓ Zincaps S29 S29

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

# Antianaemics

# **Hypoplastic and Haemolytic**

## ⇒SA2539 Special Authority for Subsidy

**Initial application — (chronic renal failure)** from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### All of the following:

- 1 Patient in chronic renal failure: and
- 2 Haemoglobin is less than or equal to 100g/L; and
- 3 Any of the following:
  - 3.1 Both:
    - 3.1.1 Patient does not have diabetes mellitus; and
    - 3.1.2 Glomerular filtration rate is less than or equal to 30ml/min; or
  - 3.2 Both:
    - 3.2.1 Patient has diabetes mellitus; and
    - 3.2.2 Glomerular filtration rate is less than or equal to 45ml/min; or
  - 3.3 Patient is on haemodialysis or peritoneal dialysis.

Initial application — (myelodysplasia) from any specialist. Approvals valid for 2 months for applications meeting the following criteria:

#### All of the following:

- 1 Patient has a confirmed diagnosis of myelodysplasia (MDS)\*; and
- 2 Has had symptomatic anaemia with haemoglobin < 100g/L and is red cell transfusion-dependent; and
- 3 Patient has very low, low or intermediate risk MDS based on the WHO classification based prognostic scoring system for myelodysplastic syndrome (WPSS); and
- 4 Other causes of anaemia such as B12 and folate deficiency have been excluded; and
- 5 Patient has a serum epoetin level of < 500 IU/L; and
- 6 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Note: Indication marked with \* is an unapproved indication

**Renewal** — (myelodysplasia) from any specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient's transfusion requirement continues to be reduced with erythropoietin treatment; and
- 2 Transformation to acute myeloid leukaemia has not occurred; and
- 3 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Note: Indication marked with \* is an unapproved indication

### EPOETIN ALFA - Special Authority see SA2539 above - Retail pharmacy

Wastage claimable	, ,		
Inj 1,000 iu in 0.5 ml, syringe	250.00	6	✓ Binocrit
Inj 2,000 iu in 1 ml, syringe		6	✓ Binocrit
Inj 3,000 iu in 0.3 ml, syringe		6	✓ Binocrit
Inj 4,000 iu in 0.4 ml, syringe	96.50	6	Binocrit
Inj 5,000 iu in 0.5 ml, syringe		6	Binocrit
Inj 6,000 iu in 0.6 ml, syringe	145.00	6	Binocrit
Inj 8,000 iu in 0.8 ml, syringe		6	Binocrit
Inj 10,000 iu in 1 ml, syringe		6	Binocrit
Inj 40,000 iu in 1 ml, syringe	250.00	1	Binocrit

	Subsidy (Manufacturer's Price) \$	Subsidis	ully Brand or sed Generic  Manufacturer
Megaloblastic			
FOLIC ACID  * Tab 0.8 mg	26.60	1,000	✓ Folic Acid  multichem
* Tab 5 mg Oral liq 50 mcg per ml			<ul><li>✓ Folic Acid Viatris</li><li>✓ Biomed</li></ul>

# Antifibrinolytics, Haemostatics and Local Sclerosants

#### EFTRENONACOG ALFA [RECOMBINANT FACTOR IX] - [Xpharm]

For patients with haemophilia B receiving prophylaxis treatment. Access to funded treatment is managed by the Haemophilia

reaters Group in conjunction with the National Haem	opnilia ivianagement gro	up.	
Inj 250 iu vial	612.50	1	Alprolix
Inj 500 iu vial	1,225.00	1	✓ Alprolix
Inj 1,000 iu vial		1	✓ Alprolix
Inj 2,000 iu vial	4,900.00	1	✓ Alprolix
Inj 3,000 iu vial	7,350.00	1	✓ Alprolix
Inj 4,000 iu vial	9,800.00	1	Alprolix
ELTROMBOPAG – Special Authority see SA1743 below - Wastage claimable	- Retail pharmacy		
Tab 25 mg	1,550.00	28	Revolade
Tab 50 mg	3,100.00	28	Revolade

### ⇒SA1743 Special Authority for Subsidy

Initial application — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has had a splenectomy; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Any of the following:
  - 3.1 Patient has a platelet count of 20,000 to 30,000 platelets per microlitre and has evidence of significant mucocutaneous bleeding: or
  - 3.2 Patient has a platelet count of less than or equal to 20,000 platelets per microlitre and has evidence of active bleeding; or
  - 3.3 Patient has a platelet count of less than or equal to 10,000 platelets per microlitre.

Initial application — (idiopathic thrombocytopenic purpura - preparation for splenectomy) only from a haematologist. Approvals valid for 6 weeks where the patient requires eltrombopag treatment as preparation for splenectomy.

Initial application — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- - 1 Patient has a significant and well-documented contraindication to splenectomy for clinical reasons; and
  - 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
  - 3 Either:
    - 3.1 Patient has immune thrombocytopenic purpura\* with a platelet count of less than or equal to 20,000 platelets per
    - 3.2 Patient has immune thrombocytopenic purpura\* with a platelet count of 20,000 to 30,000 platelets per microlitre

	Subsidy	Fully	Brand or
(Manuf	acturer's Price)	Subsidised	Generic
	\$	Per 🗸	Manufacturer

continued...

and significant mucocutaneous bleeding.

**Initial application — (severe aplastic anaemia)** only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Two immunosuppressive therapies have been trialled and failed after therapy of at least 3 months duration; and
- 2 Either:
  - 2.1 Patient has severe aplastic anaemia with a platelet count of less than or equal to 20,000 platelets per microliter; or
  - 2.2 Patient has severe aplastic anaemia with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

Renewal — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 12 months where the patient has obtained a response (see Note) from treatment during the initial approval or subsequent renewal periods and further treatment is required.

Note: Response to treatment is defined as a platelet count of > 30,000 platelets per microlitre.

Renewal — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

#### All of the following:

- 1 The patient's significant contraindication to splenectomy remains; and
- 2 The patient has obtained a response from treatment during the initial approval period; and
- 3 Patient has maintained a platelet count of at least 50,000 platelets per microlitre on treatment; and
- 4 Further treatment with eltrombopag is required to maintain response.

Renewal — (severe aplastic anaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

### Both:

- 1 The patient has obtained a response from treatment of at least 20,000 platelets per microlitre above baseline during the initial approval period; and
- 2 Platelet transfusion independence for a minimum of 8 weeks during the initial approval period.

### EMICIZUMAB - [Xpharm] - Special Authority see SA2272 below

3,570.00	1	✓ Hemlibra
7,138.00	1	✓ Hemlibra
12,492.00	1	✓ Hemlibra
17,846.00	1	✓ Hemlibra
	3,570.00 7,138.00	7,138.00 1 12,492.00 1

### ⇒SA2272 Special Authority for Subsidy

Initial application — (Severe Haemophilia A with or without FVIII inhibitors) only from a haematologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has severe congenital haemophilia A with a severe bleeding phenotype (endogenous factor VIII activity less than or equal to 2%); and
- 2 Emicizumab is to be administered at a dose of no greater than 3 mg/kg weekly for 4 weeks followed by the equivalent of 1.5 mg/kg weekly.

# EPTACOG ALFA [RECOMBINANT FACTOR VIIA] - [Xpharm]

For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group. Rare Clinical Circumstances Brand of bypassing agent for > 14 days predicted use. Access to funded treatment for > 14 days predicted use is by named patient application to the Haemophilia Treaters Group, subject to access criteria.

Inj 1 mg syringe	1	✓ NovoSeven RT
Inj 2 mg syringe2,356.60	1	✓ NovoSeven RT
Inj 5 mg syringe	1	✓ NovoSeven RT
Inj 8 mg syringe	1	✓ NovoSeven RT

	Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
	(Manufacturer's Price)	Per	Subsidised	Manufacturer
FACTOR EIGHT INHIBITOR BYPASSING FRACTION - [Xpl	harml			
For patients with haemophilia. Preferred Brand of bypass		oredic	ted use. A	ccess to funded treatmen
is managed by the Haemophilia Treaters Group in conjun				
Inj 500 Ü		1		EIBA NF
Inj 1,000 U	2,630.00	1	<b>✓</b> F	EIBA NF
Inj 2,500 U	6,575.00	1	<b>√</b> F	EIBA NF
MOROCTOCOG ALFA [RECOMBINANT FACTOR VIII] - [XI	oharm]			
For patients with haemophilia. Rare Clinical Circumstance		reco	mbinant fac	ctor VIII. Access to funder
treatment is managed by the Haemophilia Treaters Group				
subject to criteria.	,		•	,
Inj 250 iu prefilled syringe	287.50	1	✓ )	(yntha
Inj 500 iu prefilled syringe	575.00	1	<b>✓</b> )	(yntha
Inj 1,000 iu prefilled syringe	1,150.00	1	✓ )	(yntha
Inj 2,000 iu prefilled syringe	2,300.00	1		(yntha
Inj 3,000 iu prefilled syringe	3,450.00	1	<b>✓</b> X	(yntha
NONACOG GAMMA, [RECOMBINANT FACTOR IX] - [Xpha	ırml			
For patients with haemophilia. Access to funded treatme		gome	hilia Treate	rs Group in conjunction
with the National Haemophilia Management Group.				,
Inj 1,000 iu vial	870.00	1	<b>✓</b> F	RIXUBIS
Inj 2,000 iu vial		1	<b>√</b> F	RIXUBIS
Inj 3,000 iu vial		1	<b>√</b> F	RIXUBIS
OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (ADVATE)				
For patients with haemophilia. Preferred Brand of short h		r VIII	Access to	funded treatment is
managed by the Haemophilia Treaters Group in conjuncti				
Inj 500 iu vial		1		Advate
Inj 1,000 iu vial		1		Advate
Inj 2,000 iu vial		1	-	Advate
Inj 3,000 iu vial	· ·	1	-	Advate
OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (KOGENA	,			
For patients with haemophilia. Rare Clinical Circumstance		roco	mhinant fa	ctor VIII Access to funder
treatment is managed by the Haemophilia Treaters Group				
subject to criteria.	7 iii conjunction with the i	valioi	iai i iacinop	milia Management Group
Inj 250 iu vial	237 50	1	✓ k	Cogenate FS
		1		Cogenate FS
•	475.00			
Inj 500 iu vial		1		
Inj 500 iu vial Inj 1,000 iu vial	950.00	1	✓ K	(ogenate FS
Inj 500 iu vial Inj 1,000 iu vial Inj 2,000 iu vial	950.00 1,900.00	-	✓ k	Cogenate FS Cogenate FS
Inj 500 iu vial Inj 1,000 iu vial Inj 2,000 iu vial Inj 3,000 iu vial.	950.00 1,900.00 2,850.00	1	✓ k	(ogenate FS
Inj 500 iu vial Inj 1,000 iu vial Inj 2,000 iu vial Inj 3,000 iu vial RURIOCTOCOG ALFA PEGOL [RECOMBINANT FACTOR V	950.00 1,900.00 2,850.00 /III] – [Xpharm]	1 1 1	✓ k ✓ k	Kogenate FS Kogenate FS Kogenate FS
Inj 500 iu vial	950.00 1,900.00 2,850.00 /III] – [Xpharm] atment. Access to funded	1 1 1	✓ k ✓ k	Kogenate FS Kogenate FS Kogenate FS
Inj 500 iu vial	950.00 1,900.00 2,850.00 //III] – [Xpharm] atment. Access to funded iilia Management group.	1 1 1 d trea	✓ k ✓ k ✓ tment is ma	Kogenate FS Kogenate FS Kogenate FS Anaged by the Haemophil
Inj 500 iu vial	950.00 1,900.00 2,850.00 //III] – [Xpharm] atment. Access to funder iilia Management group. 1,200.00	1 1 1 trea	✓ k ✓ k ✓ k tment is ma	Kogenate FS Kogenate FS Kogenate FS Anaged by the Haemophili Adynovate
Inj 500 iu vial	950.00 1,900.00 2,850.00 //III] – [Xpharm] atment. Access to funder iilia Management group. 1,200.00	1 1 1 d trea	✓ k ✓ k ✓ k tment is ma	Kogenate FS Kogenate FS Kogenate FS Anaged by the Haemophili
Inj 500 iu vial		1 1 1 d trea 1	✓ k ✓ k ✓ k tment is ma	Kogenate FS Kogenate FS Kogenate FS Anaged by the Haemophili Adynovate
Inj 500 iu vial		1 1 1 trea	tment is ma	Kogenate FS Kogenate FS Kogenate FS Anaged by the Haemophili Adynovate Adynovate
Inj 500 iu vial		1 1 1 d trea 1	tment is ma	Kogenate FS Kogenate FS Kogenate FS Anaged by the Haemophili Adynovate
Inj 500 iu vial		1 1 1 d trea 1	tment is ma	Kogenate FS Kogenate FS Kogenate FS Anaged by the Haemophili Adynovate Adynovate

				_
	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
Vitamin K				
PHYTOMENADIONE Inj 2 mg per 0.2 ml - Up to 5 inj available on a PSO	8.00	5	<b>✓</b>	Conakion MM Paediatric
Inj 10 mg per ml, 1 ml - Up to 5 inj available on a PSO	9.21	5	<b>√</b> k	Konakion MM
Antithrombotic Agents				
Antiplatelet Agents				
ASPIRIN  * Tab 100 mg	12.65	990	<b>√</b> <u>E</u>	Ethics Aspirin EC
CLOPIDOGREL  * Tab 75 mg  DIPYRIDAMOLE	5.07	84	<b>√</b> <u>I</u>	Arrow - Clopid
Note: No new patients to be initiated on dipyridamole.  Cap modified-release 200 mg	55.13	60	✓ [	Dipyridamole - Strides §29
TICAGRELOR – Special Authority see SA2530 below – Retail p * Tab 90 mg	,	56	<b>√</b> ]	icagrelor Sandoz

# **⇒SA2530** Special Authority for Subsidy

**Initial application** — (acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

#### Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Renewal — (subsequent acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

#### Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Initial application — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

#### 1 Either:

- 1.1 Patient has had a neurological stenting procedure\* in the last 60 days; or
- 1.2 Patient is about to have a neurological stenting procedure performed\*; and

#### 2 Either:

- 2.1 Patient has demonstrated clopidogrel resistance using the P2Y12 (VerifyNow) assay or another appropriate platelet function assay and requires antiplatelet treatment with ticagrelor; or
- 2.2 Either:
  - 2.2.1 Clopidogrel resistance has been demonstrated by the occurrence of a new cerebral ischemic event; or
  - 2.2.2 Clopidogrel resistance has been demonstrated by the occurrence of transient ischemic attack symptoms referable to the stent.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

Renewal — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient is continuing to benefit from treatment; and
- 2 Treatment continues to be clinically appropriate.

**Initial application** — (Percutaneous coronary intervention with stent deployment) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic\*\*.

Renewal — (Percutaneous coronary intervention with stent deployment) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic\*\*.

**Initial application — (Stent thrombosis)** from any relevant practitioner. Approvals valid without further renewal unless notified where patient has experienced cardiac stent thrombosis whilst on clopidogrel.

Initial application — (acute minor stroke or high-risk transient ischemic attack (TIA)\*) from any relevant practitioner.

Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with a minor stroke (NIHSS† score 3 or less), high-risk TIA (ABCD2 score 4 or more) or Crescendo TIA: and
- 2 Fither:
  - 2.1 Patient is expected to be a poor metaboliser of clopidogrel, with documented clinical rationale; or
  - 2.2 Patient is allergic to clopidogrel\*\*; and
- 3 Ticagrelor to be prescribed for a maximum of 21 days following minor stroke or TIA.

Renewal — (subsequent minor stroke or TIA, or Crescendo TIA) from any relevant practitioner. Approvals valid for 1 month where patient has been diagnosed with a minor stroke (NIHSS score 3 or less) or high-risk transient ischemic attack (ABCD2 score 4 or more) or Crescendo TIA.

Notes: indications marked with \* are unapproved indications.

Note:\*\* Clopidogrel allergy is defined as a history of anaphylaxis, urticaria, generalised rash or asthma (in non-asthmatic patients) developing soon after clopidogrel is started and is considered unlikely to be caused by any other treatment.

Note:NIHSS† National Institutes of Health Stroke Scale.

# **Heparin and Antagonist Preparations**

ENOXAPARIN SODIUM - Spe	ecial Authority see SA2152	2 on the next page – Retai	pharmacy
-------------------------	----------------------------	----------------------------	----------

era a ra anti e e e e e e e e e e e e e e e e e e e	- on the north page ota.	. p	
Inj 20 mg in 0.2 ml syringe	21.90	10	✓ Clexane
Inj 40 mg in 0.4 ml syringe	29.74	10	Clexane
Inj 60 mg in 0.6 ml syringe		10	✓ Clexane
Inj 80 mg in 0.8 ml syringe	56.62	10	✓ Clexane
Inj 100 mg in 1 ml syringe		10	✓ Clexane
Inj 120 mg in 0.8 ml syringe		10	✓ Clexane Fort
Ini 150 mg in 1 ml syringe		10	✓ Clexane Fort

Sul	bsidy	Fully B	rand or
(Manufact	urer's Price) Subsi	idised G	eneric
	\$ Per	✓ M	lanufacturer

## ⇒SA2152 Special Authority for Subsidy

Initial application — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patients pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

**Initial application** — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 For the short-term treatment of venous thromboembolism prior to establishing a therapeutic level of oral anti-coagulant treatment; or
- 2 For the prophylaxis and treatment of venous thromboembolism in high risk surgery; or
- 3 To enable cessation/re-establishment of existing oral anticoagulant treatment pre/post surgery; or
- 4 For the prophylaxis and treatment of venous thromboembolism in Acute Coronary Syndrome surgical intervention; or
- 5 To be used in association with cardioversion of atrial fibrillation.

Initial application — (Short-term use during treatment of COVID-19 with nirmatrelvir with ritonavir) from any relevant practitioner. Approvals valid for 2 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is receiving an anticoagulation treatment that has drug/drug interactions with ritonavir that increases risk of bleeding; and
- 2 Patient meets the Access Criteria for COVID-19 antivirals published on the Pharmac website\*; and
- 3 Other antiviral treatments for COVID-19 have been considered and are not clinically suitable options.

Renewal — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patient's pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Renewal — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month where low molecular weight heparin treatment or prophylaxis is required for a second or subsequent event (surgery, ACS, cardioversion, or prior to oral anti-coagulation).

### HEPARIN SODIUM

Inj 1,000 iu per ml, 5 ml ampoule	127.44	50	✓ Pfizer
Inj 10 iu per ml, 5 ml ampoule (flushing solution)	19.38	10	✓ Wockhardt S29
Inj 5,000 iu per ml, 5 ml vial	83.00	10	<ul><li>Heparin Sodium Panpharma</li></ul>
Inj 5,000 iu per ml, 1 ml	70.33	5	✓ Hospira
Inj 5,000 iu per ml, 5 ml ampoule	406.15	50	✓ Pfizer
Inj 25,000 iu per ml, 0.2 ml	25.78	5	✓ Hospira
(Heparin Sodium Panpharma Inj 5,000 iu per ml, 5 ml vial to be	e delisted 1 May 2026	5)	
HEPARINISED SALINE			
Inj 10 iu per ml, 5 ml	96.91	50	✓ Pfizer

# **Oral Anticoagulants**

DABIGATRAN			
Cap 75 mg - No more than 2 cap per day	27.99	60	✓ Pradaxa
Cap 110 mg	27.99	60	✓ Pradaxa
Cap 150 mg	27.99	60	✓ Pradaxa

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	Generic
	\$	Per	•	Manufacturer
RIVAROXABAN				
Tab 10 mg - No more than 1 tab per day	15.60	30	✓	Xarelto
Tab 15 mg - Up to 14 tab available on a PSO	14.56	28	/	Xarelto
Tab 20 mg		28	1	Xarelto
WARFARIN SODIUM				
Note: Marevan and Coumadin are not interchangeable.				
* Tab 1 mg	3.46	50	✓	Coumadin
	7.50	100	✓	Marevan
* Tab 2 mg	4.31	50	/	Coumadin
* Tab 3 mg		100	1	Marevan
* Tab 5 mg		50	✓	Coumadin
· ·	13.50	100	/	Marevan
Blood Colony-stimulating Factors				
blood colony-stillulating lactors				
FILGRASTIM - Special Authority see SA1259 below - Retail pha	rmacy			
Inj 300 mcg per 0.5 ml prefilled syringe	86.60	10	1	Nivestim

## ⇒SA1259 Special Authority for Subsidy

**Initial application** only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

1 Prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 20%\*); or

10

Nivestim

- 2 Peripheral blood stem cell mobilisation in patients undergoing haematological transplantation; or
- 3 Peripheral blood stem cell mobilisation or bone marrow donation from healthy donors for transplantation; or
- 4 Treatment of severe chronic neutropenia (ANC  $< 0.5 \times 10^9$ /L); or
- 5 Treatment of drug-induced prolonged neutropenia (ANC < 0.5 ×10<sup>9</sup>/L).

Inj 480 mcg per 0.5 ml prefilled syringe......133.72

Note: \*Febrile neutropenia risk greater than or equal to 20% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

PEGFILGRASTIM - Special Authority see SA1912 below - Retail pharmacy

### **⇒SA1912** Special Authority for Subsidy

**Initial application** only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where used for prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 5%\*). Note: \*Febrile neutropenia risk greater than or equal to 5% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

# Fluids and Electrolytes

### Intravenous Administration

GLUCOSE	[DEXTROSE]

*	Inj 50%, 10 ml ampoule – Up to 5 inj available on a PSO34.75	5	✓ Biomed
*	Inj 50%, 90 ml bottle - Up to 5 inj available on a PSO17.50	1	✓ Biomed

	Subsidy (Manufacturer's Price \$	) Per	Fully Subsidised	Generic
POTASSIUM CHLORIDE	·			
* Inj 75 mg per ml, 10 ml	65.00	50	•	Juno LumaCina Pfizer \$29
SODIUM BICARBONATE			_	
Inj 8.4%, 50 ml	24.70	1	•	Biomed
Inj 8.4%, 100 ml	25.31	1	•	Biomed
SODIUM CHLORIDE  Not funded for use as a nasal drop. Not funded for nebuliser for nebuliser use.	use except when u	sed in	conjunctio	n with an antibiotic intended
Inj 23.4% (4 mmol/ml), 20 ml ampoule  For Sodium chloride oral liquid formulation refer Standard		5 R4	•	Biomed
Inj 0.9%, 5 ml ampoule – Up to 5 inj available on a PSO Fresenius Kabi to be Principal Supply on 1 February 202	4.12 6	20		Fresenius Kabi
Inj 0.9%, 10 ml ampoule – Up to 5 inj available on a PSO Fresenius Kabi to be Principal Supply on 1 February 202	6	50	•	Fresenius Kabi
Inj 0.9%, 20 ml ampoule Fresenius Kabi to be Principal Supply on 1 February 202		20	•	Fresenius Kabi
<ul> <li>Inj 0.9%, 1,000 ml bag — Up to 2 bag available on a PSO</li> <li>Only if prescribed on a prescription for renal dialysis, mat for emergency use. (500 ml and 1,000 ml packs)</li> <li>Inj 0.9%, 500 ml bag — Up to 4 bag available on a PSO</li> <li>Only if prescribed on a prescription for renal dialysis, mat</li> </ul>	ernity or post-natal	1	n the home	Baxter
for emergency use. (500 ml and 1,000 ml packs) TOTAL PARENTERAL NUTRITION (TPN)				
Infusion	CBS	1 OP	· •	TPN
WATER  1) On a prescription or Practitioner's Supply Order only whe Schedule requiring a solvent or diluent; or 2) On a bulk supply order; or 3) When used in the extemporaneous compounding of eye When used for the dilution of sodium chloride soln 7% for	e drops; or		ŕ	listed in the Pharmaceutical
Inj 10 ml ampoule - Up to 5 inj available on a PSO	7.60	50		Fresenius Kabi Multichem
Inj 20 ml ampoule – Up to 5 inj available on a PSO	5.00	20	•	Fresenius Kabi
Oral Administration				
CALCIUM POLYSTYRENE SULPHONATE Powder	169.85	800 g C	OP 🗸	Calcium Resonium
COMPOUND ELECTROLYTES  Powder for oral soln — Up to 5 sach available on a PSO	9.50	50	•	<u>Electral</u>
COMPOUND ELECTROLYTES WITH GLUCOSE [DEXTROSE] Soln with electrolytes	6.53	1 OP	•	Hydralyte - Lemonade

	Subsidy		Fully	Brand or
	(Manufacturer's Price	·)	Subsidised	Generic
	\$	Per	•	Manufacturer
PHOSPHORUS				
Tab eff 500 mg (16 mmol)	82.50	100	✓	Phosphate Phebra
POTASSIUM CHLORIDE				
* Tab eff 548 mg (14 m eq) with chloride 285 mg (8 m eq)	5.26	60		
	(17.10)		(	Chlorvescent
* Tab long-acting 600 mg (8 mmol)	16.15	200	✓ :	Span-K
Span-K to be Principal Supply on 1 February 2026				
SODIUM BICARBONATE				
Cap 840 mg	8.52	100	<b>✓</b>	Sodibic
. •			<b>√</b> :	Sodibic
SODIUM POLYSTYRENE SULPHONATE				
Powder	84.65	154 a C	)P 🗸 I	Resonium-A

Subsidy	
(Manufacturer's Price)	Subs
\$	Per

Fully Subsidised

Brand or Generic Manufacturer

✓ Prazosin Mylan S29

✓ Arrow-Quinapril 20

100

90

# **Alpha-Adrenoceptor Blockers**

Alpha Adrenoc	eptor Blockers
---------------	----------------

* Tab 2 mg       17.35       500       ✓ Doxazosin Clinect         * Tab 4 mg       20.94       500       ✓ Doxazosin Clinect	DOX	AZOSIN		
•	* T	ab 2 mg17.35	500	<ul> <li>Doxazosin Clinect</li> </ul>
DLENOVVDENZAMINE UVDDOCUI ODIDE			500	<ul> <li>Doxazosin Clinect</li> </ul>
PRENOX I DENZAMINE RI DROCHLONIDE	PHEN	NOXYBENZAMINE HYDROCHLORIDE		
<b>★</b> Cap 10 mg65.00 30 <b>✓ BNM</b> \$29	* 0	Cap 10 mg65.00	30	✓ BNM S29
PRAZOSIN	PRAZ	ZOSIN		
<b>*</b> Tab 1 mg	<b>*</b> T	ab 1 mg5.53	100	✓ Arrotex-Prazosin
S29 S29				<b>S29</b> S29
9.98 <b>✓ Minipress</b> \$29		9.98		✓ Minipress S29
<b>*</b> Tab 2 mg7.00 100 <b>✓ Arrotex-Prazosin</b>	* T	ab 2 mg7.00	100	✓ Arrotex-Prazosin
S29 S29				<b>S29</b> S29
13.29 <b>✓ Minipress</b> \$29				✓ Minipress S29
<b>*</b> Tab 5 mg11.70 100 <b>✓ Arrotex-Prazosin</b>	* T	ab 5 mg11.70	100	✓ Arrotex-Prazosin
S29 S29				<b>S29</b> S29
22.00 <b>✓ Minipress</b> \$29		22.00		✓ Minipress S29
* Cap 1 mg	* 0	Cap 1 mg15.40	100	✓ Prazosin Mylan S29
			100	✓ Prazosin Mylan S29

# Agents Affecting the Renin-Angiotensin System

# **ACE Inhibitors**

$C\Delta P$	-	חחו	

*	Oral liq 5 mg per ml	5.00 10	00 ml OP	✓ <u>DP-Captopril</u>
ΕN	JALAPRIL MALEATE			
*	Tab 5 mg4	1.25	90	✓ Acetec
*	Tab 10 mg5		90	✓ Acetec
*	Tab 20 mg6		90	✓ Acetec
LIS	SINOPRIL			
*	Tab 5 mg12	2.00	90	<ul><li>Teva Lisinopril</li></ul>
	Teva Lisinopril to be Principal Supply on 1 March 2026			
*		2.00	90	<ul><li>Teva Lisinopril</li></ul>
	Teva Lisinopril to be Principal Supply on 1 March 2026			
*	Tab 20 mg16	6.00	90	✓ Teva Lisinopril
	Teva Lisinopril to be Principal Supply on 1 March 2026			
PE	RINDOPRIL			
*	Tab 2 mg1	.79	30	✓ Coversyl
*	Tab 4 mg2	2.44	30	✓ Coversyl
*	Tab 8 mg	3.94	30	✓ Coversyl
Q۱	JINAPRIL			
*	Tab 5 mg10	).24	90	✓ Arrow-Quinapril 5
*			90	✓ Arrow-Quinapril 10

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

(Mani	Subsidy ufacturer's Price) \$	Per	Fully Subsidised	
RAMIPRIL				
* Cap 1.25 mg	.17.25	90	✓	Tryzan
* Cap 2.5 mg	.16.50	90	✓	Tryzan
* Cap 5 mg	.16.88	90	✓	Tryzan
* Cap 10 mg	.17.63	90	✓	<u>Tryzan</u>
Angiotensin II Antagonists				
CANDESARTAN CILEXETIL				
★ Tab 4 mg	2.68	90	✓	Candestar
<b>米</b> Tab 8 mg	2.67	90	✓	Candestar
* Tab 16 mg	4.22	90	✓	Candestar
* Tab 32 mg	5.24	90	✓	Candestar
LOSARTAN POTASSIUM				
* Tab 12.5 mg	2.00	84	1	Losartan Actavis
* Tab 25 mg		84		Losartan Actavis
* Tab 50 mg		84		Losartan Actavis
≮ Tab 100 mg		84		Losartan Actavis
Angiotensin II Antagonists with Diuretics				
CANDESARTAN CILEXETIL WITH HYDROCHLOROTHIAZIDE				
* Tab 16 mg with hydrochlorothiazide 12.5 mg	4.10	30	/	APO-Candesartan HCTZ 16/12.5
* Tab 32 mg with hydrochlorothiazide 12.5 mg	5.25	30	✓	APO-Candesartan HCTZ 32/12.5
LOSARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE				
* Tab 50 mg with hydrochlorothiazide 12.5 mg	4.31	30	1	Arrow-Losartan &

# **Angiotensin II Antagonists with Neprilysin Inhibitors**

SACUBITRIL WITH VALSARTAN – Special Authority see S	6A2302 below – Retail p	harmacy	
Tab 24.3 mg with valsartan 25.7 mg	190.00	56	✓ Entresto 24/26
Tab 48.6 mg with valsartan 51.4 mg	190.00	56	✓ Entresto 49/51
Tab 97.2 mg with valsartan 102.8 mg	190.00	56	✓ Entresto 97/103

### ⇒SA2302 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has heart failure: and
- 2 Any of the following:
  - 2.1 Patient is in NYHA/WHO functional class II; or
  - 2.2 Patient is in NYHA/WHO functional class III; or
  - 2.3 Patient is in NYHA/WHO functional class IV; and
- 3 Either:
  - 3.1 Patient has a documented left ventricular ejection fraction (LVEF) of less than or equal to 35%; or
  - 3.2 An ECHO is not reasonably practical, and in the opinion of the treating practitioner the patient would benefit from treatment; and
- 4 Patient is receiving concomitant optimal standard chronic heart failure treatments.

Hydrochlorothiazide

Subsidy	ı	ully	Brand or	
(Manufacturer's Price)	Subsid	ised	Generic	
\$	Per	1	Manufacturer	

Ant	arr	hvt	hm	ics

•		
For lignocaine hydrochloride refer to NERVOUS SYSTEM, Anaesthetics, Local,	page 123	
AMIODARONE HYDROCHLORIDE		
▲ Tab 100 mg	30	✓ Aratac
Aratac to be Principal Supply on 1 February 2026	20	./ Avetes
▲ Tab 200 mg5.86 Aratac to be Principal Supply on 1 February 2026	30	✓ Aratac
Inj 50 mg per ml, 3 ml ampoule – Up to 10 inj available on a		
PSO17.96	10	✓ Max Health
Max Health to be Principal Supply on 1 February 2026	10	- max rioditii
ATROPINE SULPHATE		
* Inj 600 mcg per ml, 1 ml ampoule – Up to 5 inj available on a		
PSO	10	✓ Hikma S29
		✓ Martindale
DIGOXIN		
* Tab 62.5 mcg - Up to 30 tab available on a PSO	240	✓ Lanoxin PG
Lanoxin PG to be Principal Supply on 1 February 2026		
* Tab 250 mcg - Up to 30 tab available on a PSO	240	<ul><li>Lanoxin</li></ul>
Lanoxin to be Principal Supply on 1 February 2026		
★ Oral liq 50 mcg per ml16.60	60 ml	✓ Lanoxin
		✓ Lanoxin S29 S29
DISOPYRAMIDE PHOSPHATE		_
▲ Cap 100 mg55.90	84	Rythmodan -
		Cheplafarm S29
		✓ Rythmodan
		Neon S29
LECAINIDE ACETATE		
Tab 50 mg	60	Flecainide BNM
Cap long-acting 100 mg35.78	90	✓ <u>Flecainide</u>
		<u>Controlled</u> Release Teva
▲ Cap long-acting 200 mg54.28	90	✓ Flecainide
Cap long-acting 200 mg	30	Controlled
		Release Teva
Inj 10 mg per ml, 15 ml ampoule102.79	5	✓ Almarytm \$29
108.16	•	✓ Tambocor
		✓ Tambocor
		German S29
MEXILETINE HYDROCHLORIDE		
▲ Cap 150 mg162.00	100	✓ Teva S29
▲ Cap 250 mg202.00	100	✓ Teva S29
PROPAFENONE HYDROCHLORIDE		
▲ Tab 150 mg40.90	50	✓ Rytmonorm
•		•

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

# **Antihypotensives**

		v – Retail pharmacy	MIDODRINE - Special Authority see SA1474 below
✓ <u>Midodrine</u>	100	36.68	Tab 2.5 mg
<u>Medsurge</u>	400	50.00	Tab 5 and
✓ <u>Midodrine</u> Medsurae	100	58.88	Tab 5 mg

## ⇒SA1474 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years where patient has disabling orthostatic hypotension not due to drugs.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

# **Beta-Adrenoceptor Blockers**

# **Beta Adrenoceptor Blockers**

ATENOLOL			
* Tab 50 mg	11.00	500	✓ Viatris
* Tab 100 mg	18.50	500	✓ <u>Atenolol Viatris</u>
* Oral liq 25 mg per 5 ml	49.85	300 ml OP	Atenolol AFT
Restricted to children under 12 years of age.			
BISOPROLOL FUMARATE			
* Tab 2.5 mg	1.36	90	✓ Ipca-Bisoprolol
* Tab 5 mg	1.91	90	✓ Ipca-Bisoprolol
* Tab 10 mg	2.71	90	✓ Ipca-Bisoprolol
CARVEDILOL			
* Tab 6.25 mg	2.24	60	✓ Carvedilol Sandoz
* Tab 12.5 mg		60	✓ Carvedilol Sandoz
* Tab 25 mg		60	✓ Carvedilol Sandoz
LABETALOL			
* Tab 100 mg	14.50	100	✓ Trandate
· · · · · · · · · · · · · · · · · · ·	49.54		✓ Biocon S29
* Tab 200 mg		100	✓ Trandate
	42.07	100	✓ Presolol S29
* Inj 5 mg per ml, 20 ml ampoule		5	· I ICOOIOI
injoing por hii, 20 hii ampoulo	(88.60)	Ü	Trandate
METORDOL OL CLICOINATE	(00.00)		Tandato
METOPROLOL SUCCINATE	4.00	90	√ Mulaa CD
* Tab long-acting 23.75 mg      * Tab long-acting 47.5 mg		90 90	✓ Myloc CR ✓ Myloc CR
· · · · · · · · · · · · · · · · · · ·			
* Tab long-acting 95 mg      * Tab long-acting 190 mg		90 90	✓ Myloc CR ✓ Myloc CR
	9.70	90	wiyloc ch
METOPROLOL TARTRATE			
* Tab 50 mg		100	✓ <u>IPCA-Metoprolol</u>
* Tab 100 mg		60	✓ <u>IPCA-Metoprolol</u>
* Tab long-acting 200 mg		28	✓ Slow-Lopresor
* Inj 1 mg per ml, 5 ml vial	26.50	5	✓ Metoprolol IV Mylan
			Metoprolol IV Viatris

		Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
NA	DOLOL				
*	Tab 40 mg	19.19	100	✓ N	ladolol BNM
*	Tab 80 mg	30.39	100	<b>✓</b> N	ladolol BNM
PR	OPRANOLOL				
*	Tab 10 mg	7.04	100	✓ [	Profate
	Tab 40 mg		100	✓ ĪI	PCA-Propranolol
K	Cap long-acting 160 mg		100	<b>√</b> (	Cardinol LA
K	Oral liq 4 mg per ml - Special Authority see SA1327 below	_			
	Retail pharmacy	CBS	500 m	<b> </b>	likma-
					Propranolol §29
				<b>√</b> F	Roxane-
				•	Propranolol \$29

# ⇒SA1327 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

#### SOTAL OL

*	Tab 80 mg	22.50	300	✓ Sotalol Viatris ©29
	ŭ	40.00	500	✓ Mylan
	Mylan to be Principal Supply on 1 February 2026			•
*	Tab 160 mg	20.00	100	✓ Mylan
	Mylan to be Principal Supply on 1 February 2026			
(Sc	otalol Viatris S29 Tab 80 mg to be delisted 1 March 2026)			

# Calcium Channel Blockers

# **Dihydropyridine Calcium Channel Blockers**

AMLODIPINE			
* Tab 2.5 mg	1.45	90	✓ Vasorex
* Tab 5 mg		90	✓ Vasorex
* Tab 10 mg	1.31	90	✓ Vasorex
FELODIPINE			
* Tab long-acting 2.5 mg	2.18	30	Plendil ER
* Tab long-acting 5 mg		90	✓ Felo 5 ER
* Tab long-acting 10 mg	6.95	90	✓ Felo 10 ER

		Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	I Generic
IFEDIP	PINE				
Tab	long-acting 10 mg - Subsidy by endorsement	19.42	56	✓	Tensipine MR10 S29
	Subsidised for patients who were taking nifedipine tab lendorsed accordingly. Pharmacists may annotate the properties of nifedipine tab long-acting 10 mg.				
	long-acting 20 mg		100	✓	Nyefax Retard
Tab	long-acting 30 mg	4.78	14	/	Mylan Italy (24 hr release) \$29
		34.10	100	•	Mylan (24 hr release) \$29
Tab	long-acting 60 mg	52.81	100	•	Mylan (24 hr release) \$29
ther	Calcium Channel Blockers				
TIA7	EM HYDROCHLORIDE				
	long-acting 120 mg	65.35	500	/	Diltiazem CD Clinect
	long-acting 180 mg		30		Cardizem CD
	long-acting 240 mg		30		Cardizem CD
	KILINE MALEATE				
	100 mg	62.90	100	/	Pexsig
	AMIL HYDROCHLORIDE				
	40 mg	7.01	100	1	Isoptin
	80 mg		100	_	Isoptin
	long-acting 120 mg		100	1	Isoptin Retard S29
					Isoptin SR
	long-acting 240 mg		30	•	Isoptin SR
	PSO	25.00	5	✓	Isoptin
entr	ally-Acting Agents				
.ONID	INE				
	ch 2.5 mg, 100 mcg per day – Only on a prescription		4	_	<u>Mylan</u>
	ch 5 mg, 200 mcg per day - Only on a prescription		4		<u>Mylan</u>
	ch 7.5 mg, 300 mcg per day – Only on a prescription INE HYDROCHLORIDE	17.90	4	•	<u>Mylan</u>
	25 mcgClonidine Teva to be Principal Supply on 1 February 20		112	•	Clonidine Teva
Tab	150 mcg	40.41	100	✓	Catapres
	50 mcg per ml, 1 ml ampoule	14.10	5	•	Catapres
Tab	250 mg	15.10	100		Methyldopa Viatris
iure	tics				
.oop	Diuretics				
	ANIDE				
	1 mg		100	_	Burinex
Ini 5	00 mcg per ml, 4 ml vial	7.95	5	/	Burinex

	Subsidy (Manufacturer's F	Price) Subs	Fully Brand or sidised Generic
	\$	Per	✓ Manufacturer
UROSEMIDE [FRUSEMIDE]			
Tab 40 mg - Up to 30 tab available on a PSO	12.80	1,000	✓ IPCA-Frusemide
* Tab 500 mg	25.00	50	✓ Urex Forte
★ Oral liq 10 mg per ml	11.20	30 ml OP	✓ Lasix
Inj 10 mg per ml, 25 ml ampoule		6	✓ Lasix
₭ Inj 10 mg per ml, 2 ml ampoule – Up to 5 inj available on a	PSO2.40	5	✓ Furosemide-Baxter
Potassium Sparing Diuretics			
MILORIDE HYDROCHLORIDE			
Tab 5 mg	81.07	100	✓ Padagis S29
·	171.41	28	✓ Wockhardt S29
Oral liq 1 mg per ml	35.40	25 ml OP	✓ Biomed
PLERENONE - Special Authority see SA1728 below - Retail			
Tab 25 mg		30	✓ Inspra
Tab 50 mg		30	✓ Inspra
SA1728 Special Authority for Subsidy			<u></u>
itial application from any relevant practitioner. Approvals va e following criteria:	lid without further	renewal unless	s notified for applications mee
oth:			
1 Patient has heart failure with ejection fraction less than 4 2 Either:	.0%; and		
2.1 Patient is intolerant to optimal dosing of spironola	•		
<ol><li>2.2 Patient has experienced a clinically significant ad</li></ol>	verse effect while	on optimal dos	sing of spironolactone.
PIRONOLACTONE			
← Tab 25 mg	4.20	100	✓ Spiractin
Spiractin to be Principal Supply on 1 March 2026			
€ Tab 100 mg	11.40	100	✓ Spiractin
Spiractin to be Principal Supply on 1 March 2026			
Oral liq 5 mg per ml	35.70	25 ml OP	✓ Biomed
Potassium Sparing Combination Diuretics			
MILORIDE HYDROCHLORIDE WITH FUROSEMIDE			
Tab 5 mg with furosemide 40 mg		28	✓ Frumil
MILORIDE HYDROCHLORIDE WITH HYDROCHLOROTHIA	ZIDE		
Tab 5 mg with hydrochlorothiazide 50 mg	5.00	50	✓ Moduretic
Thiazide and Related Diuretics			
ENDROFLUMETHIAZIDE [BENDROFLUAZIDE]			
Fab 2.5 mg – Up to 150 tab available on a PSO	51.50	500	✓ <u>Arrow-</u> Bendrofluazide
May be supplied an a DOO for recover other than	*******		
May be supplied on a PSO for reasons other than eme  Tab 5 mg	· ,	500	✓ Arrow-
€ Tab 5 mg	01.00	500	Bendrofluazide
UII ODOTUIAZIDE			
HLOROTHIAZIDE Oral liq 50 mg per ml	30.67	25 ml OP	✓ Biomed
- · · · · · · · · · · · · · · · · · · ·			

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
CHLORTALIDONE [CHLORTHALIDONE]  * Tab 25 mg  Hygroton to be Principal Supply on 1 February 2026	6.95	50	•	Hygroton
INDAPAMIDE  * Tab 2.5 mg  METOLAZONE	16.00	90	•	<u>Dapa-Tabs</u>
Tab 5 mg	CBS	50	✓	Zaroxolyn S29
Vasopressin receptor antagonists				
TOLVAPTAN – Special Authority see SA2166 below – Retail pha Tab 15 mg Tab 30 mg Tab 45 mg + 15 mg Tab 60 mg + 30 mg	873.50 1,747.00 1,747.00	28 OF 28 OF 56 OF	· · · · · · · · · · · · · · · · · · ·	Jinarc Jinarc Jinarc Jinarc
Tab 90 mg + 30 mg		56 OF 56 OF		Jinarc Jinarc

# ⇒SA2166 Special Authority for Subsidy

Initial application — (autosomal dominant polycystic kidney disease) only from a renal physician or any relevant practitioner on the recommendation of a renal physician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of autosomal dominant polycystic kidney disease; and
- 2 Patient has an estimated glomerular filtration rate (eGFR) of greater than or equal to 25 ml/min/1.73 m<sup>2</sup> at treatment initiation; and
- 3 Either:
  - 3.1 Patient's disease is rapidly progressing, with a decline in eGFR of greater than or equal to 5 mL/min/1.73 m<sup>2</sup> within one-year; or
  - 3.2 Patient's disease is rapidly progressing, with an average decline in eGFR of greater than or equal to 2.5 mL/min/1.73 m<sup>2</sup> per year over a five-year period.

Renewal — (autosomal dominant polycystic kidney disease) only from a renal physician or any relevant practitioner on the recommendation of a renal physician. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has not developed end-stage renal disease, defined as an eGFR of less than 15 mL/min/1.73 m<sup>2</sup>; and
- 2 Patient has not undergone a kidney transplant.

Lipid-Modifying Agents		
Fibrates		
BEZAFIBRATE       22.65         Tab 200 mg       22.65         Tab long-acting 400 mg       21.54	90 30	✓ <u>Bezalip</u> ✓ <u>Bezalip Retard</u>
Other Lipid-Modifying Agents		
ACIPIMOX	30	✓ Olbetam

Resins  COLESTYRAMINE Powder for oral suspension 4 g sachet		
COLESTYRAMINE Powder for oral suspension 4 g sachet61.50 50		
	<b>√</b> Q	olestyramine - Mylan §29 uantalan sugar free §29

Н	MG CoA Reductase Inhibitors (Statins)		
АТ	ORVASTATIN		
*	Tab 10 mg0.31	30	✓ Lorstat
	5.16	500	✓ Lorstat
*	Tab 20 mg8.12	500	✓ Lorstat
*	Tab 40 mg13.79	500	✓ Lorstat
*	Tab 80 mg1.52	30	✓ Lorstat
	25.39	500	✓ Lorstat
PR	AVASTATIN		
*	Tab 20 mg7.16	100	✓ Clinect
*	Tab 40 mg	100	✓ Clinect
RC	SUVASTATIN - Special Authority see SA2093 below - Retail pharmacy		
*	Tab 5 mg1.29	30	✓ Rosuvastatin Viatris
*	Tab 10 mg1.69	30	✓ Rosuvastatin Viatris
*	Tab 20 mg2.71	30	✓ Rosuvastatin Viatris
	4.21		✓ Rosuvastatin-
			Sandoz
*	Tab 40 mg4.55	30	✓ Rosuvastatin Viatris

⇒SA2093 Special Authority for Subsidy

**Initial application — (cardiovascular disease risk)** from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

# Either:

- 1 Both:
  - 1.1 Patient is considered to be at risk of cardiovascular disease; and
  - 1.2 Patient is Māori or any Pacific ethnicity; or
- 2 Both:
  - 2.1 Patient has a calculated risk of cardiovascular disease of at least 15% over 5 years; and
  - 2.2 LDL cholesterol has not reduced to less than 1.8 mmol/litre with treatment with the maximum tolerated dose of atoryastatin and/or simyastatin.

Initial application — (familial hypercholesterolemia) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Both:

- 1 Patient has familial hypercholesterolemia (defined as a Dutch Lipid Criteria score greater than or equal to 6); and
- 2 LDL cholesterol has not reduced to less than 1.8 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initial application — (established cardiovascular disease) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

		Subsidy (Manufacturer's Price)	S Per	Fully subsidised	Brand or Generic Manufacturer	
--	--	-----------------------------------	----------	---------------------	-------------------------------------	--

continued...

- 1 Any of the following:
  - 1.1 Patient has proven coronary artery disease (CAD); or
  - 1.2 Patient has proven peripheral artery disease (PAD); or
  - 1.3 Patient has experienced an ischaemic stroke; and
- 2 LDL cholesterol has not reduced to less than 1.4 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initial application — (recurrent major cardiovascular events) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has experienced a recurrent major cardiovascular event (defined as myocardial infarction, ischaemic stroke, coronary revascularisation, hospitalisation for unstable angina) in the last 2 years; and
- 2 LDL cholesterol has not reduced to less than 1.0 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

SIN	۱۱/۸	CT	٠٨٦	ΓIN	
אווכ	IVA	OΙ	м	ш	

*	Tab 10 mg	.68	90	✓	Simvastatin Mylan
				✓	Simvastatin Viatris
*	Tab 20 mg	2.54	90	✓	Simvastatin Viatris
*	Tab 40 mg	1.11	90	✓	Simvastatin Viatris
	Tab 80 mg		90	✓	Simvastatin Viatris

# **Selective Cholesterol Absorption Inhibitors**

ΕZI	= I IMIBE	
	T 1 40	

* Tab 10 mg	1.76	30	✓ Ezetimibe Sandoz
EZETIMIBE WITH SIMVASTATIN			
Tab 10 mg with simvastatin 10 mg	11.86	30	✓ Zimybe
Tab 10 mg with simvastatin 20 mg		30	✓ Zimybe
Tab 10 mg with simvastatin 40 mg		30	✓ Zimybe
Tab 10 mg with simvastatin 80 mg	14.27	30	✓ Zimybe

## **Nitrates**

#### **GLYCERYL TRINITRATE**

*	Oral pump spray, 400 mcg per dose – Up to 250 dose			
	available on a PSO	7.48	250 dose OP	✓ Nitrolingual Pump Spray
*	Patch 25 mg, 5 mg per day	15.73	30	✓ Nitroderm TTS
*	Patch 50 mg, 10 mg per day	18.62	30	✓ Nitroderm TTS
ISC	DSORBIDE MONONITRATE			
*	Tab 20 mg	22.49	100	✓ Ismo 20
*	Tab long-acting 40 mg	9.80	30	✓ Ismo 40 Retard
*	Tab long-acting 60 mg	13.50	90	✓ <u>Duride</u>

# **Sympathomimetics**

## **ADRENALINE**

Inj 1 in 1,000, 1 ml ampoule - Up to 5 inj available on a PSO4.98	5	<ul> <li>Aspen Adrenaline</li> </ul>
13.27		DBL Adrenaline
Inj 1 in 10,000, 10 ml ampoule - Up to 5 inj available on a PSO27.00	5	✓ Hospira
49.00	10	Aspen Adrenaline

Subsidy (Manufacturer's Price)	Subsic	Fully lised	Brand or Generic	
 \$	Per	1	Manufacturer	

### Vasodilators

### HYDRALAZINE HYDROCHLORIDE

*	Tab 25 mg - Special Authority see SA1321 below - Retail			
	pharmacy	CBS	1	<ul><li>Hydralazine</li></ul>
			56	✓ Onelink S29
			84	✓ AMDIPHARM S29
			100	✓ Camber S29
*	Inj 20 mg ampoule	25.90	5	✓ Apresoline

### ⇒SA1321 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Either:

- 1 For the treatment of refractory hypertension; or
- 2 For the treatment of heart failure in combination with a nitrate, in patients who are intolerant or have not responded to ACE inhibitors and/or angiotensin receptor blockers.

#### MINOXIDIL

▲ Tab 10 mg	47.04	60	Minoxidil Roma S29
•	78.40	100	✓ Loniten
NICORANDIL			
▲ Tab 10 mg	27.81	60	✓ Max Health
Max Health to be Principal Supply on 1 February 2026			
▲ Tab 20 mg	35.12	60	✓ Max Health
Max Health to be Principal Supply on 1 February 2026			
PAPAVERINE HYDROCHLORIDE			
* Inj 12 mg per ml, 10 ml ampoule	257.12	5	✓ Hospira
PENTOXIFYLLINE [OXPENTIFYLLINE]			·
Tab 400 mg	44.37	50	✓ Trental 400
1.25 1.00 mg			

# **Endothelin Receptor Antagonists**

AMBRISENTAN - Special Authority see SA2556 below - Retail p	oharmacy		
Tab 5 mg	200.00	30	✓ Ambrisentan Viatris
Tab 10 mg	200.00	30	✓ Ambrisentan Viatris

### ⇒SA2556 Special Authority for Subsidy

**Initial application** — **(PAH monotherapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

### All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II. III or IV: and
- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH has been confirmed by right heart catheterisation; and
    - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
- 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm<sup>5</sup>); and
- 4.1.5 Any of the following:
  - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH; or
  - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool\*\*: or
  - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
- 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
  - 5.1 Ambrisentan is to be used as PAH monotherapy; and
  - 5.2 Any of the following:
    - 5.2.1 Patient has experienced intolerable side effects with both sildenafil and bosentan; or
    - 5.2.2 Patient has an absolute contraindication to sildenafil and an absolute or relative contraindication to bosentan (e.g. due to current use of a combined oral contraceptive or liver disease); or
    - 5.2.3 Patient is a child with idiopathic PAH or PAH secondary to congenital heart disease.

**Initial application** — **(PAH dual therapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH has been confirmed by right heart catheterisation; and
    - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
    - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
    - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm<sup>5</sup>); and
    - 4.1.5 Any of the following:
      - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH; or
      - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool\*\*; or
      - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
  - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
  - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
  - 5.1 Ambrisentan is to be used as PAH dual therapy; and
  - 5.2 Any of the following:

Subsidy (Manufacturer's Price)	S	Fully ubsidised	Brand or Generic	
\$	Per	<b>✓</b>	Manufacturer	

continued...

- 5.2.1 Patient has tried bosentan (either as PAH monotherapy, or PAH dual therapy with sildenafil) for at least three months and has not experienced an acceptable response to treatment according to a validated risk stratification tool\*\*: or
- 5.2.2 Patient has experienced intolerable side effects on bosentan; or
- 5.2.3 Patient has an absolute or relative contraindication to bosentan (e.g. due to current use of a combined oral contraceptive or liver disease); or
- 5.2.4 Patient is presenting in NYHA/WHO functional class III or IV, and would benefit from initial dual therapy in the opinion of the treating clinician and has an absolute or relative contraindication to bosentan (eg. due to current liver disease or use of a combined oral contraceptive).

**Initial application** — **(PAH triple therapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II. III or IV: and
- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH has been confirmed by right heart catheterisation; and
    - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
    - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
    - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm<sup>-5</sup>); and
    - 4.1.5 Any of the following:
      - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH; or
      - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool\*\*; or
      - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
  - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
  - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
  - 5.1 Ambrisentan is to be used as PAH triple therapy; and
  - 5.2 Any of the following:
    - 5.2.1 Patient is on the lung transplant list; or
    - 5.2.2 Both:
      - 5.2.2.1 Patient is presenting in NYHA/WHO functional class IV; and
      - 5.2.2.2 Patient has an absolute or relative contraindication to bosentan (e.g. due to current use of a combined oral contraceptive or liver disease); or
    - 5.2.3 Both:
      - 5.2.3.1 Patient has tried PAH dual therapy for at least three months and remains in an unacceptable risk category according to a validated risk stratification tool\*\*; and
      - 5.2.3.2 Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario.

Renewal only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

respiratory specialist, cardiologist or rheumatologist. Approvals valid for 2 years where the patient is continuing to derive benefit from ambrisentan treatment according to a validated PAH risk stratification tool\*\*.

Note: \*\* the requirement to use a validated risk stratification tool to determine insufficient response applies to adults.

Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

BOSENTAN - Special Authority see SA2557 below - Retail pharmacy

1ab 62.5 mg	100.00	60	
			Reddy's
Tab 125 mg	100.00	60	✓ Bosentan Dr
			Reddy's

## ⇒SA2557 Special Authority for Subsidy

**Initial application** — **(PAH monotherapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

#### All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)\*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH has been confirmed by right heart catheterisation; and
    - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
    - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
    - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm<sup>5</sup>); and
    - 4.1.5 Any of the following:
      - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH; or
      - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool\*\*: or
      - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
  - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or
  - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
  - 5.1 Bosentan is to be used as PAH monotherapy; and
  - 5.2 Any of the following:
    - 5.2.1 Patient has experienced intolerable side effects on sildenafil: or
    - 5.2.2 Patient has an absolute contraindication to sildenafil; or
    - 5.2.3 Patient is a child with idiopathic PAH or PAH secondary to congenital heart disease.

**Initial application** — **(PAH dual therapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

#### All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)\*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and

Subsidy	Full	y Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per 🗸	Manufacturer	

continued...

- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH has been confirmed by right heart catheterisation; and
    - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
    - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
    - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm<sup>-5</sup>); and
    - 4.1.5 Any of the following:
      - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH; or
      - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool\*\*; or
      - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
  - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or
  - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Bosentan is to be used as part of PAH dual therapy; and
- 6 Either:
  - 6.1 Patient has tried a PAH monotherapy (sildenafil) for at least three months and has experienced an inadequate therapeutic response to treatment according to a validated risk stratification tool\*\*: or
  - 6.2 Patient is presenting in NYHA/WHO functional class III or IV, and in the opinion of the treating clinician would likely benefit from initial dual therapy.

**Initial application** — **(PAH triple therapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)\*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH has been confirmed by right heart catheterisation; and
    - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
    - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
    - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm<sup>-5</sup>); and
    - 4.1.5 Any of the following:
      - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH; or
      - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool\*\*: or
      - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
  - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or
  - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sı	ubsidised	Generic
\$	Per	1	Manufacturer

continued...

complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and

- 5 Both:
  - 5.1 Bosentan is to be used as part of PAH triple therapy; and
  - 5.2 Any of the following:
    - 5.2.1 Patient is on the lung transplant list; or
    - 5.2.2 Patient is presenting in NYHA/WHO functional class IV; or
    - 5.2.3 Both:
      - 5.2.3.1 Patient has tried PAH dual therapy for at least three months and has not experienced an acceptable response to treatment according to a validated risk stratification tool\*\*; and
      - 5.2.3.2 Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario.

Renewal only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 2 years where patient is continuing to derive benefit from bosentan treatment according to a validated PAH risk stratification tool\*\*.

Note: \*\* the requirement to use a validated risk stratification tool to determine insufficient response applies to adults.

Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

# **Phosphodiesterase Type 5 Inhibitors**

SILDENAFIL - Special Authority see SA2558 below - Retail pha	armacy		
Tab 25 mg	0.72	4	✓ Vedafil
Tab 50 mg		4	✓ Vedafil
Tab 100 mg	11.22	12	✓ Vedafil

## ⇒SA2558 Special Authority for Subsidy

**Initial application** — (Raynaud's Phenomenon\*) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has Raynaud's Phenomenon\*; and
- 2 Patient has severe digital ischaemia (defined as severe pain requiring hospital admission or with a high likelihood of digital ulceration; digital ulcers; or gangrene); and
- 3 Patient is following lifestyle management (avoidance of cold exposure, sufficient protection, smoking cessation support, avoidance of sympathomimetic drugs); and
- 4 Patient is being treated with calcium channel blockers and nitrates (or these are contraindicated/not tolerated).

Initial application — (Pulmonary arterial hypertension\*) only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)\*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II. III or IV: and
- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH is confirmed by right heart catheterisation; and
    - 4.1.2 A mean pulmonary artery pressure (PAPm) of greater than 20 mmHg; and
    - 4.1.3 A pulmonary capillary wedge pressure (PCWP) that is less than or equal to 15 mmHg; and
    - 4.1.4 Pulmonary vascular resistance (PVR) of at least 2 Wood Units or greater than 160 International Units (dyn s

Subsidy (Manufacturer's Price)	S	Fully ubsidised	Brand or Generic
\$	Per	1	Manufacturer

continued...

cm<sup>-5</sup>); and

- 4.1.5 Any of the following:
  - 4.1.5.1 PAH is non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH; or
  - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool\*\*; or
  - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
- 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or
- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures.

Initial application — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has a documented history of traumatic or non-traumatic spinal cord injury; and
- 2 Patient has erectile dysfunction secondary to spinal cord injury requiring pharmacological treatment.

Renewal — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Notes: Indications marked with \* are Unapproved Indications.

\*\* the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

# **Prostacyclin Analogues**

EPOPROSTENOL - Special Authority see SA2559 below - Ref	tail pharmacy		
Inj 500 mcg vial	36.61	1	✓ Veletri
Inj 1.5 mg vial	73.21	1	✓ Veletri

### ⇒SA2559 Special Authority for Subsidy

**Initial application** — **(PAH dual therapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class III or IV; and
- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH has been confirmed by right heart catheterisation; and
    - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
    - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
    - 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm<sup>5</sup>); and
    - 4.1.5 Any of the following:
      - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH; or
      - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a

Subsi	idv Full	Brand or
(Manufacture	er's Price) Subsidised	I Generic
\$	Per 🗸	Manufacturer

continued...

validated risk stratification tool\*\*: or

- 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
- 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 All of the following:
  - 5.1 Epoprostenol is to be used as part of PAH dual therapy with either sildenafil or an endothelin receptor antagonist; and
  - 5.2 Patient is presenting in NYHA/WHO functional class IV: and
  - 5.3 Patient has tried a PAH monotherapy for at least three months and remains in an unacceptable risk category according to a validated risk stratification tool.

Initial application — (PAH triple therapy) only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class III or IV; and
- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH has been confirmed by right heart catheterisation; and
    - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
    - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
    - 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm<sup>-5</sup>); and
    - 4.1.5 Any of the following:
      - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH; or
      - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool\*\*; or
      - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
  - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
  - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
  - 5.1 Epoprostenol is to be used as PAH triple therapy; and
  - 5.2 Any of the following:
    - 5.2.1 Patient is on the lung transplant list; or
    - 5.2.2 Patient is presenting in NYHA/WHO functional class IV; or
    - 5.2.3 Both:
      - 5.2.3.1 Patient has tried PAH dual therapy for at least three months and has not experienced an acceptable response to treatment according to a validated risk stratification tool; and
      - 5.2.3.2 Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario.

Renewal only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

respiratory specialist, cardiologist or rheumatologist. Approvals valid for 2 years where patient is continuing to derive benefit from epoprostenol treatment according to a validated PAH risk stratification tool\*\*.

Note: \*\* the requirement to use a validated risk stratification tool to determine insufficient response applies to adults.

Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

ILOPROST - Special Authority see SA2560 below - Retail pharmacy

## ⇒SA2560 Special Authority for Subsidy

**Initial application** — **(PAH monotherapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH has been confirmed by right heart catheterisation; and
    - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
    - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
    - 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm<sup>-5</sup>); and
    - 4.1.5 Any of the following:
      - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH; or
      - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool\*\*; or
      - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
  - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
  - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
  - 5.1 Iloprost is to be used as PAH monotherapy; and
  - 5.2 Either:
    - 5.2.1 Patient has experienced intolerable side effects on sildenafil and both the funded endothelin receptor antagonists (i.e. both bosentan and ambrisentan); or
    - 5.2.2 Patient has an absolute contraindication to sildenafil and an absolute or relative contraindication to endothelin receptor antagonists.

**Initial application** — **(PAH dual therapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

- 4.1 All of the following:
  - 4.1.1 PAH has been confirmed by right heart catheterisation; and
  - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
  - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
  - 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm<sup>5</sup>); and
  - 4.1.5 Any of the following:
    - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH; or
    - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool\*\*: or
    - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
- 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 All of the following:
  - 5.1 Iloprost is to be used as PAH dual therapy with either sildenafil or an endothelin receptor antagonist; and
  - 5.2 Fither:
    - 5.2.1 Patient has an absolute contraindication to or has experienced intolerable side effects on sildenafil: or
    - 5.2.2 Patient has an absolute or relative contraindication to or experienced intolerable side effects with a funded endothelin receptor antagonist; and
  - 5.3 Either:
    - 5.3.1 Patient has tried a PAH monotherapy for at least three months and remains in an unacceptable risk category according to a validated risk stratification tool\*\*; or
    - 5.3.2 Patient is presenting in NYHA/WHO functional class III or IV, and in the opinion of the treating clinician would benefit from initial dual therapy.

Initial application — (PAH triple therapy) only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II. III or IV: and
- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH has been confirmed by right heart catheterisation; and
    - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
    - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
    - 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm<sup>-5</sup>); and
    - 4.1.5 Any of the following:
      - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH; or
      - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool\*\*: or
      - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or

(Manufacturer's Price) Subsidised Generic \$ Per ✓ Manufacturer
--

continued...

- 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
  - 5.1 Iloprost is to be used as PAH triple therapy; and
  - 5.2 Any of the following:
    - 5.2.1 Patient is on the lung transplant list; or
    - 5.2.2 Patient is presenting in NYHA/WHO functional class IV; or
    - 5.2.3 Both
      - 5.2.3.1 Patient has tried PAH dual therapy for at least three months and has not experienced an acceptable response to treatment according to a validated risk stratification tool\*\*; and
      - 5.2.3.2 Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario.

**Renewal** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 2 years where patient is continuing to derive benefit from iloprost treatment according to a validated PAH risk stratification tool\*\*.

Note: \*\* the requirement to use a validated risk stratification tool to determine insufficient response applies to adults.

Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.



Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

# **Antiacne Preparations**

For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 93

#### ADAPALENE

- a) Maximum of 30 g per prescription
- b) Only on a prescription

b) Only on a prosonphon			
Gel 0.1%	22.89	30 g OP	Differin
ISOTRETINOIN - Special Authority see SA	A2449 below – Retail pharmacy		
Cap 5 mg	11.26	60	<ul><li>Oratane</li></ul>
Cap 10 mg	18.75	120	✓ Oratane
Cap 20 mg	26.73	120	✓ Oratane

#### ⇒SA2449 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, paediatrician, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around isotretinoin and is competent to prescribe isotretinoin; and
- 3 Any of the following:
  - 3.1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and that they must not become pregnant during treatment and for a period of one month after the completion of treatment; or
  - 3.2 Patient is not of child bearing potential: or
  - 3.3 Patient is a child and it is considered not appropriate to exclude pregnancy or start contraceptives or undertake pregnancy-related isotretinoin counselling.

**Renewal** from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and that they must not become pregnant during treatment and for a period of one month after the completion of treatment: or

(13.00)

- 2 Patient is not of child bearing potential; or
- 3 Patient is a child and it is considered not appropriate to exclude pregnancy or start contraceptives or undertake pregnancy-related isotretinoin counselling.

#### **TRETINOIN**

Crm 0.5 mg per g − Maximum of 50 g per prescription .......16.82 50 g OP ✓ ReTrieve

# **Antibacterials Topical**

For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 93

#### HYDROGEN PEROXIDE

HYDROGEN PEROXIDE			
* Crm 1%	4.89	15 g OP	<ul> <li>Crystaderm</li> </ul>
MUPIROCIN			
Oint 2%	6.60	15 g OP	

- a) Only on a prescription
- b) Not in combination

Bactroban

	Subsidy (Manufacturer's Pri	ce) Subs	Fully Brand or sidised Generic ✓ Manufacturer
SODIUM FUSIDATE [FUSIDIC ACID] Crm 2%	1.69	5 g OP	✓ <u>Foban</u>
a) Maximum of 5 g per prescription     b) Only on a prescription     c) Not in combination Oint 2%	1.69	5 g OP	✓ <u>Foban</u>
<ul><li>a) Maximum of 5 g per prescription</li><li>b) Only on a prescription</li><li>c) Not in combination</li></ul>			
SULFADIAZINE SILVER Crm 1%	10.80	50 g OP	✓ Flamazine
<ul><li>a) Up to 250 g available on a PSO</li><li>b) Not in combination</li></ul>			
Antifungals Topical			
For systemic antifungals, refer to INFECTIONS, Antifungals, pa	ige 100		
AMOROLFINE			
a) Only on a prescription     b) Not in combination			
Nail soln 5%	21.87	5 ml OP	✓ MycoNail
CLOTRIMAZOLE			
* Crm 1%	1.10	20 g OP	✓ Clomazol
<ul><li>a) Only on a prescription</li><li>b) Not in combination</li></ul>			
* Soln 1%		20 ml OP	Canastan
a) Only on a prescription	(11.58)		Canesten
b) Not in combination			
ECONAZOLE NITRATE			
Crm 1%	8.04	20 g OP	✓ Pevaryl
a) Only on a prescription			
b) Not in combination Foaming soln 1%, 10 ml sachets	0.80	3	
r daming 30m 176, 10 mi sacricis	(18.64)	J	Pevaryl
a) Only on a prescription	, ,		,
b) Not in combination			
MICONAZOLE NITRATE			<b></b>
* Crm 2%	0.90	15 g OP	✓ Multichem
a) Only on a prescription     b) Not in combination			
* Lotn 2%	4.36	30 ml OP	
	(10.03)		Daktarin
a) Only on a prescription			
b) Not in combination  * Tinct 2%	4 36	30 ml OP	
THIOLE/0	(12.10)	50 IIII OI	Daktarin
a) Only on a prescription	,		
b) Not in combination			

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	/	Manufacturer	

36.80

# **Antipruritic Preparations**

CA	ΙΑ	MI	IN	F

- a) Only on a prescription
- b) Not in combination

100 g

20 g OP

✓ healthE Calamine Aqueous

✓ Itch-Soothe

### **CROTAMITON**

- a) Only on a prescription
- b) Not in combination

Crm 10%......3.49

MENTHOL - Only in combination

- 1) Only in combination with a dermatological base or proprietary Topical Corticosteriod Plain
- 2) With or without other dermatological galenicals.

25 g ✓ MidWest ✓ MidWest 100 g

# **Corticosteroids Topical**

For systemic corticosteroids, refer to CORTICOSTEROIDS AND RELATED AGENTS, page 84

### Corticosteroids - Plain

DETAMETHACONE DIDDODIONATE

BETAMETHASONE DIPROPIONATE			
Crm 0.05%	2.96	15 g OP	✓ Diprosone
	36.00	50 g OP	✓ Diprosone
Oint 0.05%	2.96	15 g OP	✓ Diprosone
	36.00	50 g OP	✓ Diprosone
Oint 0.05% in propylene glycol base		30 g OP	✓ Diprosone OV
BETAMETHASONE VALERATE			
* Crm 0.1%	5.85	50 g OP	✓ Beta Cream
* Oint 0.1%	7.90	50 g OP	✓ Beta Ointment
* Lotn 0.1%	30.00	50 ml OP	✓ Betnovate
CLOBETASOL PROPIONATE			
* Crm 0.05%	3.75	30 g OP	✓ Dermol
Dermol to be Principal Supply on 1 February 2026		•	
* Oint 0.05%	3.68	30 g OP	✓ Dermol
Dermol to be Principal Supply on 1 February 2026		-	
CLOBETASONE BUTYRATE			
Crm 0.05%	5.38	30 g OP	
	(10.00)	Ů	Eumovate
HYDROCORTISONE			
* Crm 1% - Only on a prescription	1.78	30 g OP	✓ Ethics
, , ,	20.40	500 g	✓ Noumed
Noumed to be Principal Supply on 1 February 2026		ŭ	
* Powder – Only in combination	49.95	25 g	✓ ABM
Up to 5% in a dermatological base (not proprietary Top galenicals	ical Corticosterioc	d – Plain) with o	or without other dermatological

	Subsidy		Fully	Brand or
	(Manufacturer's P		sidised	Generic
	\$	Per		Manufacturer
HYDROCORTISONE AND PARAFFIN LIQUID AND LANOLIN				
Lotn 1% with paraffin liquid 15.9% and lanolin 0.6% – Only o		050	,	DD 1 1 110
a prescription	12.83	250 ml	•	DP Lotn HC
HYDROCORTISONE BUTYRATE			_	
Lipocream 0.1%		100 g OP		Locoid Lipocream
Oint 0.1%		100 g OP		Locoid
Milky emul 0.1%	12.33	100 ml OP	•	Locoid Crelo
METHYLPREDNISOLONE ACEPONATE			_	
Crm 0.1%		15 g OP	_	Advantan
Oint 0.1%	4.95	15 g OP	•	<u>Advantan</u>
MOMETASONE FUROATE			_	
Crm 0.1%		15 g OP	_	Elocon Alcohol Free
01.10.44	3.50	50 g OP		Elocon Alcohol Free
Oint 0.1%		15 g OP	_	Elocon
Lotn 0.1%	3.50	50 g OP 30 ml OP		Elocon Elocon
	4.99	30 IIII OF	•	Elocoli
TRIAMCINOLONE ACETONIDE	0.40	400 - OD	,	Autotopost
Crm 0.02% Oint 0.02%		100 g OP	_	Aristocort
OINt 0.02%	0.54	100 g OP	V	Aristocort
Corticosteroids - Combination				
BETAMETHASONE VALERATE WITH SODIUM FUSIDATE [FU	SIDIC ACIDI			
Crm 0.1% with sodium fusidate (fusidic acid) 2%		15 g OP		
(,	(10.45)			Fucicort
a) Maximum of 15 g per prescription	, ,			
b) Only on a prescription				
HYDROCORTISONE WITH MICONAZOLE - Only on a prescrip	tion			
* Crm 1% with miconazole nitrate 2%		15 g OP	1	Micreme H
HYDROCORTISONE WITH NATAMYCIN AND NEOMYCIN - O		-		
Oint 1% with natamycin 1% and neomycin sulphate 0.5%		15 g OP	1	Pimafucort
TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYCI		•		
Crm 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 mg		IIN		
and gramicidin 250 mcg per g - Only on a prescription		15 g OP		
and gramician 250 mag per g Only on a prescription.	(9.28)	10 9 01		Viaderm KC
	(5.25)			
Barrier Creams and Emollients				
Barrier Creams				
DIMETUICONE				
DIMETHICONE  * Crm 5% pump bottle	4 14	460 g OP	1	Hydralock
A Offit 3 % pump bottle	4.30	400 g Oi		healthE
	4.00		•	Dimethicone 5%
* Crm 10% pump bottle	4.52	460 g OP	1	healthE
		y 0.	-	Dimethicone 10%
(healthE Dimethicone 5% Crm 5% pump bottle to be delisted 1 Ju	une 2026)			
ZINC AND CASTOR OIL	/			
* Oint	4.25	500 g	1	Evara
		y	-	

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's I \$	Price) Subs	Fully Brand or sidised Generic Manufacturer
Emollients			
QUEOUS CREAM			
<b>₭</b> Crm	1.65	500 g	✓ Evara
CETOMACROGOL			
≰ Crm BP	2.29	500 g	<ul> <li>Cetomacrogol-AFT</li> </ul>
ETOMACROGOL WITH GLYCEROL			
Crm 90% with glycerol 10%		460 g OP	✓ <u>Evara</u>
THE OFFICE OF THE OFFI	3.25	920 g OP	✓ <u>Evara</u>
:MULSIFYING OINTMENT ME Oint BP	2.12	500 a	✓ Evara Emulsifying
S OIII DF		500 g	Ointment
DIL IN WATER EMULSION			<u>Omanona</u>
E Crm	2.10	500 g	✓ Fatty Emulsion
		555 g	Cream (Evara)
PARAFFIN			
Oint liquid paraffin 50% with white soft paraffin 50%	4.45	500 g OP	✓ EVARA Paraffin
			Ointment 50/50
	4.94		✓ White Soft Liquid
White Soft Liquid Paraffin AFT Oint liquid paraffin 50% with w	1. 'A (A (C'		Paraffin AFT
		no/ to be delicte	d 1 luna 2020)
	піте ѕот рагатіп ы	0% to be deliste	ed 1 June 2026)
JREA	·		,
IREA <b>k</b> Crm 10%	·	0% to be deliste	d 1 June 2026)  ✓ healthE Urea Crea
IREA	1.37	100 g OP	,
IREA <b>k</b> Crm 10%	1.37		,
IREA	1.37	100 g OP	✓ healthE Urea Creat
IREA		100 g OP	✓ healthE Urea Crean DP Lotion Alpha-Keri Lotion
IREA		100 g OP 1,000 ml 250 ml OP	✓ healthE Urea Crean DP Lotion
IREA		100 g OP 1,000 ml	DP Lotion Alpha-Keri Lotion DP Lotion
IREA		100 g OP 1,000 ml 250 ml OP 1,000 ml	✓ healthE Urea Crean DP Lotion Alpha-Keri Lotion
IREA		100 g OP 1,000 ml 250 ml OP	DP Lotion Alpha-Keri Lotion DP Lotion
JREA  Crm 10%  VOOL FAT WITH MINERAL OIL – Only on a prescription  Lotn hydrous 3% with mineral oil		100 g OP 1,000 ml 250 ml OP 1,000 ml	DP Lotion Alpha-Keri Lotion DP Lotion BK Lotion
IREA		100 g OP 1,000 ml 250 ml OP 1,000 ml	DP Lotion Alpha-Keri Lotion DP Lotion BK Lotion
JREA  Crm 10%  VOOL FAT WITH MINERAL OIL – Only on a prescription  Lotn hydrous 3% with mineral oil		100 g OP 1,000 ml 250 ml OP 1,000 ml	DP Lotion Alpha-Keri Lotion DP Lotion BK Lotion
JREA  © Crm 10%  VOOL FAT WITH MINERAL OIL — Only on a prescription  E Lotn hydrous 3% with mineral oil		100 g OP 1,000 ml 250 ml OP 1,000 ml	DP Lotion Alpha-Keri Lotion DP Lotion BK Lotion
UREA  k Crm 10%  VOOL FAT WITH MINERAL OIL − Only on a prescription  k Lotn hydrous 3% with mineral oil  Other Dermatological Bases		100 g OP 1,000 ml 250 ml OP 1,000 ml 250 ml OP	DP Lotion Alpha-Keri Lotion DP Lotion BK Lotion BK Lotion

Only in combination with a dermatological galenical or as a diluent for a proprietary Topical Corticosteroid – Plain.

Brand or

Generic

Fully

Subsidised

	\$	Per	✓ Manufacturer
Minor Skin Infections			
POVIDONE IODINE			
Oint 10%	7.40	65 g OP	✓ Betadine
a) Maximum of 130 g per prescription		-	
b) Only on a prescription			
Antiseptic Solution 10%	4.99	100 ml	✓ Riodine
Antiseptic soln 10%	3.83	15 ml	✓ Riodine
	6.99	500 ml	✓ Riodine
Skin preparation, povidone iodine 10% with 30% alcohol	1.63	100 ml	
	(3.48)		Betadine Skin Prep

Subsidy

(Manufacturer's Price)

## Parasiticidal Preparations

DIM	۱E	THI	CONE	
			40/	

 ★ Lotn 4%
 4.60
 200 ml OP
 ✓ healthE

 Dimethicone 4%

 Lotion

IVERMECTIN - Special Authority see SA2511 below - Retail pharmacy

Tab 3 mg − Up to 100 tab available on a PSO......17.20 4 Stromectol

- PSO for institutional use only. Must be endorsed with the name of the institution for which the PSO is required and a valid Special Authority for patient of that institution.
- 2) Ivermectin available on BSO provided the BSO includes a valid Special Authority for a patient of the institution.
- For the purposes of subsidy of ivermectin, institution means age related residential care facilities, disability care facilities or prisons.

#### ⇒SA2511 Special Authority for Subsidy

Initial application — (Scabies) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

#### Either:

- 1 The person has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
- 2 Both:
  - 2.1 The person has a confirmed diagnosis of scabies or is a close contact of a scabies case; and
  - 2.2 Fither:
    - 2.2.1 The person is unable to complete topical therapy; or
    - 2.2.2 Previous treatment with topical therapy has been tried and not cleared the infestation.

**Initial application — (Other parasitic infections)** from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

#### Any of the following:

- 1 filariasis; or
- 2 cutaneous larva migrans (creeping eruption); or
- 3 strongyloidiasis; or
- 4 The individual has a travel or residence history that requires presumptive parasite treatment.

Renewal — (Scabies) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria: Either:

- 1 The person has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
- 2 Both:
  - 2.1 The person has a confirmed diagnosis of scabies or is a close contact of a scabies case; and



Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

2.2 Either:

- 2.2.1 The person is unable to complete topical therapy; or
- 2.2.2 Previous treatment with topical therapy has been tried and not cleared the infestation.

Renewal — (Other parasitic infections) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 filariasis: or
- 2 cutaneous larva migrans (creeping eruption); or
- 3 strongyloidiasis.

#### **PERMETHRIN**

## **Psoriasis and Eczema Preparations**

ACITRETIN - Special Authority see SA2024 below - Retail	pharmacy		
Cap 10 mg	26.20	60	✓ Novatretin
Cap 25 mg	57.37	60	✓ Novatretin

#### ⇒SA2024 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around acitretin and is competent to prescribe acitretin; and
- 3 Either:
  - 3.1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and that they must not become pregnant during treatment and for a period of three years after the completion of treatment; or
  - 3.2 Patient is not of child bearing potential.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Fither:

- 1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and that they must not become pregnant during treatment and for a period of three years after the completion of treatment; or
- 2 Patient is not of child bearing potential.

#### BETAMETHASONE DIPROPIONATE WITH CALCIPOTRIOL

BETAMETHASONE DIFNOPIONATE WITH CALCIFOTHIOL			
Foam spray 500 mcg with calcipotriol 50 mcg per g	59.95	60 g OP	<ul><li>Enstilar</li></ul>
Gel 500 mcg with calcipotriol 50 mcg per g	40.92	60 g OP	✓ Daivobet
Oint 500 mcg with calcipotriol 50 mcg per g	14.31	30 g OP	Daivobet
CALCIPOTRIOL			
Oint 50 mcg per g	40.00	120 g OP	Daivonex
COAL TAR			
Soln BP - Only in combination	46.00	200 ml	✓ Midwest

- 1) Up to 10% only in combination with a dermatological base or proprietary Topical Corticosteriod Plain
- 2) With or without other dermatological galenicals.

	Subsidy (Manufacturer's Price	e) Sub	Fully sidised	Brand or Generic
	\$	Per	1	Manufacturer
COAL TAR WITH ALLANTOIN, MENTHOL, PHENOL AND SULP	HUR			
Soln 5% with sulphur 0.5%, menthol 0.75%, phenol 0.5% and				
allantoin crm 2.5%	6.59	75 g OP		
	(8.00)		l	Egopsoryl TA
		30 g OP		
	(4.35)			Egopsoryl TA
COAL TAR WITH SALICYLIC ACID AND SULPHUR Soln 12% with salicylic acid 2% and sulphur 4% oint	7.95	40 g OP	✓ (	Coco-Scalp
PIMECROLIMUS - Special Authority see SA1970 below - Retail	pharmacy			
a) Maximum of 15 g per prescription				
b) Note: a maximum of 15 g per prescription and no more th				
Cream 1%	33.00	15 g OP	•	<u>Elidel</u>
⇒SA1970 Special Authority for Subsidy				
nitial application only from a dermatologist, paediatrician, ophtha				
f a dermatologist, paediatrician or ophthalmologist. Approvals va	alid without further	renewal un	less no	tified for applications
neeting the following criteria: Both:				
Patient has atopic dermatitis on the eyelid; and				
<ul><li>2 Patient has at least one of the following contraindications to</li></ul>	o tonical carticacta	raida, nari	orificial	dormatitic rocacoa
documented epidermal atrophy, documented allergy to top pressure.				
documented epidermal atrophy, documented allergy to top pressure.	ical corticosteroids	, cataracts	, glauco	
documented epidermal atrophy, documented allergy to top	ical corticosteroids SCEIN – Only on a	, cataracts	, glauco on	
documented epidermal atrophy, documented allergy to top pressure.  PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORES Soln 2.3% with trolamine laurilsulfate and fluorescein sodium.	ical corticosteroids SCEIN – Only on a	, cataracts prescription	, glauco on	ma, or raised intraocula
documented epidermal atrophy, documented allergy to top pressure.  PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORES Soln 2.3% with trolamine laurilsulfate and fluorescein sodium.  SALICYLIC ACID	ical corticosteroids GCEIN – Only on a 5.41	prescriptic	, glauco on ✓ إ	ma, or raised intraocula
documented epidermal atrophy, documented allergy to top pressure.  PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORES Soln 2.3% with trolamine laurilsulfate and fluorescein sodium.  SALICYLIC ACID Powder – Only in combination	ical corticosteroids  CEIN - Only on a5.41	prescription 500 ml	, glauco on ✓ I	ma, or raised intraocula Pinetarsol Midwest
documented epidermal atrophy, documented allergy to top pressure.  PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORES Soln 2.3% with trolamine laurilsulfate and fluorescein sodium.  SALICYLIC ACID	ical corticosteroids  CEIN - Only on a5.41	prescription 500 ml	, glauco on ✓ I	ma, or raised intraocula Pinetarsol Midwest
documented epidermal atrophy, documented allergy to top pressure.  PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORES Soln 2.3% with trolamine laurilsulfate and fluorescein sodium.  PALICYLIC ACID Powder – Only in combination	ical corticosteroids  CEIN - Only on a5.41	prescription 500 ml	, glauco on ✓ I	ma, or raised intraocula Pinetarsol Midwest
documented epidermal atrophy, documented allergy to top pressure.  PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORES Soln 2.3% with trolamine laurilsulfate and fluorescein sodium.  PALICYLIC ACID Powder – Only in combination	ical corticosteroids  CEIN - Only on a5.41	prescription 500 ml	, glauco on ✓ I	ma, or raised intraocula Pinetarsol Midwest
documented epidermal atrophy, documented allergy to top pressure.  INE TAR WITH TROLAMINE LAURILSULFATE AND FLUORES  Soln 2.3% with trolamine laurilsulfate and fluorescein sodium.  SALICYLIC ACID  Powder – Only in combination	ical corticosteroids  CEIN – Only on a5.4129.00 proprietary Topical	prescription 500 ml	on  y i  y i  oid – Pi	ma, or raised intraocula Pinetarsol Midwest
documented epidermal atrophy, documented allergy to top pressure.  PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORES Soln 2.3% with trolamine laurilsulfate and fluorescein sodium.  SALICYLIC ACID  Powder – Only in combination	ical corticosteroids  CEIN - Only on a5.41 29.00  proprietary Topical	, cataracts prescriptic 500 ml 250 g Corticoster	on  y  or  or  y  or  or  or  y  or  or  y  or  or	ma, or raised intraocula  Pinetarsol  Midwest  lain or collodion flexible  Midwest
documented epidermal atrophy, documented allergy to top pressure.  PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORES Soln 2.3% with trolamine laurilsulfate and fluorescein sodium.  SALICYLIC ACID Powder – Only in combination	ical corticosteroids  CEIN - Only on a5.41 29.00  proprietary Topical	, cataracts prescriptic 500 ml 250 g Corticoster	on  y  or  or  y  or  or  or  y  or  or  y  or  or	ma, or raised intraocula  Pinetarsol  Midwest  lain or collodion flexible  Midwest
documented epidermal atrophy, documented allergy to top pressure.  PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORES Soln 2.3% with trolamine laurilsulfate and fluorescein sodium.  ALICYLIC ACID Powder – Only in combination	ical corticosteroids  CEIN - Only on a5.41 29.00  proprietary Topical	, cataracts prescriptic 500 ml 250 g Corticoster	on  y  or  or  y  or  or  or  y  or  or  y  or  or	ma, or raised intraocula  Pinetarsol  Midwest  lain or collodion flexible  Midwest
documented epidermal atrophy, documented allergy to top pressure.  PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORES Soln 2.3% with trolamine laurilsulfate and fluorescein sodium.  SALICYLIC ACID Powder – Only in combination	ical corticosteroids  CEIN - Only on a5.41 29.00  proprietary Topical	, cataracts prescriptic 500 ml 250 g Corticoster	on  y  or  or  y  or  or  or  y  or  or  y  or  or	ma, or raised intraocula  Pinetarsol  Midwest  lain or collodion flexible  Midwest
documented epidermal atrophy, documented allergy to top pressure.  PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORES Soln 2.3% with trolamine laurilsulfate and fluorescein sodium.  ALICYLIC ACID Powder – Only in combination	ical corticosteroids  CEIN - Only on a5.41 29.00  proprietary Topical	, cataracts prescriptic 500 ml 250 g Corticoster	on  y  or  or  y  or  or  or  y  or  or  y  or  or	ma, or raised intraocula  Pinetarsol  Midwest  lain or collodion flexible  Midwest
documented epidermal atrophy, documented allergy to top pressure.  PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORES Soln 2.3% with trolamine laurilsulfate and fluorescein sodium.  Powder – Only in combination	ical corticosteroids  CEIN – Only on a	, cataracts prescriptic 500 ml 250 g Corticoster	on  y I  roid - PI	ma, or raised intraocula  Pinetarsol  Midwest  lain or collodion flexible  Midwest
documented epidermal atrophy, documented allergy to top pressure.  PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORES Soln 2.3% with trolamine laurilsulfate and fluorescein sodium.  ALICYLIC ACID Powder — Only in combination	ical corticosteroids  CEIN – Only on a	prescription 500 ml 250 g Corticoster	on  y I  roid - PI	ma, or raised intraocula  Pinetarsol  Midwest  Iain or collodion flexible  Midwest  Iain
documented epidermal atrophy, documented allergy to top pressure.  PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORES Soln 2.3% with trolamine laurilsulfate and fluorescein sodium.  ALICYLIC ACID Powder – Only in combination	ical corticosteroids CEIN – Only on a5.4129.00 proprietary Topical6.35 proprietary Topical	, cataracts prescriptic 500 ml 250 g Corticoster 100 g Corticoster 30 g OP	on  y I  y I  yoid - Pi  yoid - Pi	ma, or raised intraocula  Pinetarsol  Midwest lain or collodion flexible  Midwest lain
documented epidermal atrophy, documented allergy to top pressure.  INE TAR WITH TROLAMINE LAURILSULFATE AND FLUORES Soln 2.3% with trolamine laurilsulfate and fluorescein sodium.  ALICYLIC ACID Powder – Only in combination	ical corticosteroids CEIN – Only on a5.4129.00 proprietary Topical6.35 proprietary Topical	, cataracts prescriptic 500 ml 250 g Corticoster 100 g Corticoster 30 g OP	on  y I  y I  yoid - Pi  yoid - Pi	Pinetarsol  Midwest lain or collodion flexible  Midwest lain

1 Patient has atopic dermatitis on the face; and

Both:

2 Patient has at least one of the following contraindications to topical corticosteroids: periorificial dermatitis, rosacea, documented epidermal atrophy or documented allergy to topical corticosteroids.

paediatrician, . Approvals valid without further renewal unless notified for applications meeting the following criteria:

## **DERMATOLOGICALS**

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✓ Manufacturer

## **Scalp Preparations**

BETAMETHASONE VALERATE		
* Scalp app 0.1%	100 ml OP	✓ Beta Scalp
CLOBETASOL PROPIONATE		
* Scalp app 0.05%	30 ml OP	Dermol
Dermol to be Principal Supply on 1 February 2026		
HYDROCORTISONE BUTYRATE		
Scalp lotn 0.1%	100 ml OP	✓ Locoid
KETOCONAZOLE		
Shampoo 2%4.09	100 ml OP	✓ Sebizole
a) Maximum of 100 ml per prescription		

## Sunscreens

SUNSCREENS, PROPRIETARY – Subsidy by endorsement

Only if prescribed for a patient with severe photosensitivity secondary to a defined clinical condition and the prescription is endorsed accordingly.

## **Wart Preparations**

For salicylic acid preparations refer to PSORIASIS AND ECZEMA PREPARATIONS, page 74

PODOPHYLLOTOXIN

- a) Maximum of 3.5 ml per prescription
- b) Only on a prescription

b) Only on a prescription

## **Other Skin Preparations**

## **Antineoplastics**

FLUOROURACIL SODIUM Crm 5%	5.56	20 g OP	✓ <u>Efudix</u>
IMIQUIMOD Crm 5%, 250 mg sachet	21.72	24	✓ Padagis ✓ Perrigo

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

		Subsidy (Manufacturer's Price)		Fully Subsidised	
		(Manufacturer's Price) \$	Per	Subsidised	
C	ontraceptives - Non-hormonal				
C	ondoms				
-	NDOMS				
	49 mm - Up to 144 dev available on a PSO		144	•	Moments
K	53 mm	1.15	10	•	Moments
		14.25	144	•	Moments
	<ul> <li>a) Maximum of 60 dev per prescription</li> </ul>				
	b) Up to 60 dev available on a PSO				
K	53 mm, 0.05 mm thickness	1.15	10	✓	Moments
		14.25	144	✓	Moments
	a) Up to 60 dev available on a PSO				
	b) Maximum of 60 dev per prescription				
6	53 mm, chocolate, brown	1.15	10	/	Moments
		14.25	144		Moments
	a) Up to 60 dev available on a PSO	11.20		•	
	b) Maximum of 60 dev per prescription				
K	53 mm, strawberry, red	1 15	10		Moments
•	oo min, shawbony, rou	14.25	144		Moments
	a) Un to 60 day available as a DCO	14.20	144	•	MOHIEHIS
	a) Up to 60 dev available on a PSO				
	b) Maximum of 60 dev per prescription	4.45	40	,	
÷	56 mm		10		Moments
		14.50	144	•	Moments
	a) Maximum of 60 dev per prescription				
	b) Up to 60 dev available on a PSO				
6	56 mm, 0.05 mm thickness		12		Gold Knight
		24.10	144	•	Gold Knight
	a) Up to 60 dev available on a PSO				
	b) Maximum of 60 dev per prescription				
6	56 mm, 0.05mm thickness (bulk pack)	20.17	144	•	Gold Knight
	a) Maximum of 60 dev per prescription				-
	b) Up to 60 dev available on a PSO				
6	56 mm, 0.08 mm thickness	1.15	10	/	Moments
		14.25	144	/	Moments
	a) Up to 60 dev available on a PSO				
	b) Maximum of 60 dev per prescription				
6	56 mm, 0.08 mm thickness, red	1.15	10	/	Moments
	, , , , , , , , , , , , , , , , , , , ,	14.25	144		Moments
	a) Up to 60 dev available on a PSO	0		-	<del>-</del>
	b) Maximum of 60 dev per prescription				
6	56 mm, chocolate	1 70	12	_	Gold Knight
•	oo mm, onocolate	21.45	144		Gold Knight
	a) Unito 60 day available and PCO	21.40	1-1-4	•	Gold Killylit
	a) Up to 60 dev available on a PSO				
	b) Maximum of 60 dev per prescription 56 mm, strawberry	1 70	10		Cold Knieht
÷	oo miin, strawberry		12		Gold Knight
	\	21.45	144	•	Gold Knight
	a) Up to 60 dev available on a PSO				
	b) Maximum of 60 dev per prescription				
+	60 mm		12		Gold Knight XL
		21.89	144	•	Gold Knight XL
	a) Maximum of 60 dev per prescription				
_	b) 🎜 prita p Sude varse ilable on a PSO	S29 Unapprove	d med	icine suppli	ed under Section 29
ŏ	60 mm/hHttparetipply	cdiz Z8haidiaad	c144	lv 📝	Gold Knight XL

			GENITO-	UF	RINARY SYSTEM
_		Subsidy (Manufacturer's Price)	F Subsidi Per	ully sed	Brand or Generic Manufacturer
_	a) Maximum of 60 dev per prescription     b) Up to 60 dev available on a PSO				
C	ontraceptive Devices				
INT	RA-UTERINE DEVICE				
	a) Up to 40 dev available on a PSO				
*	b) Only on a PSO IUD 29.1 mm length × 23.2 mm width	29.80	1	•	Choice 380 7med Nsha Silver/ copper Short
*	IUD 33.6 mm length × 29.9 mm width	26.80	1	•	TCu 380 Plus Normal
*	IUD 35.5 mm length × 19.6 mm width	33.00	1	✓	Cu 375 Standard
С	ontraceptives - Hormonal				
С	ombined Oral Contraceptives				
	HINYLOESTRADIOL WITH DESOGESTREL				
*	Tab 20 mcg with desogestrel 150 mcg and 7 inert tab - Up to 84 tab available on a PSO		84	/	Mercilon 28
ET	HINYLOESTRADIOL WITH LEVONORGESTREL				
*	Tab 20 mcg with levonorgestrel 100 mcg and 7 inert tablets - Up to 84 tab available on a PSO Lo-Oralcon 20 ED to be Principal Supply on 1 April 2026		84	✓	Lo-Oralcon 20 ED
*	Tab 30 mcg with levonorgestrel 150 mcg and 7 inert tablets - Up to 84 tab available on a PSO		84	/	Oralcon 30 ED
FTI	HINYLOESTRADIOL WITH NORETHISTERONE				
	Tab 35 mcg with norethisterone 1 mg and 7 inert tab – Up to 84 tab available on a PSO		84	/	Brevinor 1/28
	Tab 35 mcg with norethisterone 500 mcg and 7 inert tab - U to 112 tab available on a PSO		84	,	Norimin
	to 112 tab available on a 1 do	29.32	112		Norimin
P	rogestogen-only Contraceptives				
	SOGESTREL				
	Tab 75 mcg – Up to 84 tab available on a PSO	24.50	84	✓	Cerazette
	/ONORGESTREL Tab 30 mcg – Up to 112 tab available on a PSO	22.00	112	/	Microlut
*	Intra-uterine device 52 mg - Up to 25 dev available on a PSO Intra-uterine device 13.5 mg - Up to 10 dev available on a		1	_	Mirena
*	PSO	е	1		Jaydess
	on a PSO	106.92	2 OP	1	<u>Jadelle</u>
ME	DROXYPROGESTERONE ACETATE Inj 150 mg per ml, 1 ml syringe – Up to 5 inj available on a P	SO 10.56	1	•	Depo-Provera

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	Brand or Generic Manufacturer
NORETHISTERONE Tab 350 mcg - Up to 84 tab available on a PSO	12.25	84		Noriday Noriday 28
<b>Emergency Contraceptives</b>				
LEVONORGESTREL  * Tab 1.5 mg	1.31	1	<b>✓</b>	Levonorgestrel -1 (Lupin)
	1.75		<b>✓</b> I	Levonorgestrel BNM
a) Maximum of 2 tab per prescription     b) Lin to 5 tab available on a PSO				

- c) Note: Direct Provision by a pharmacist permitted under the provisions in Part I of Section A.

(Levonorgestrel BNM Tab 1.5 mg to be delisted 1 June 2026)

## **Antiandrogen Oral Contraceptives**

Prescribers may code prescriptions "contraceptive" (code "O") when used as indicated for contraception. The period of supply and prescription charge will be as per other contraceptives, as follows:

- A maximum \$5.00 prescription charge (patient co-payment) may apply.
- prescription may be written for up to six months supply.

Prescriptions coded in any other way are subject to any non contraceptive prescription charges that apply, and the non-contraceptive period of supply. ie. Prescriptions may be written for up to three months supply.

#### CYPROTERONE ACETATE WITH ETHINYLOESTRADIOL

\* Tab 2 mg with ethinyloestradiol 35 mcg and 7 inert tabs - Up 168 ✓ Ginet

## **Gynaecological Anti-infectives**

ACETIC ACID WITH HYDROXYQUINOLINE AND RICINOLEIC ACID  Jelly with glacial acetic acid 0.94%, hydroxyquinoline sulphate			
0.025%, glycerol 5% and ricinoleic acid 0.75% with applicator8.43	100 g OP		
(24.87)		Aci-Jel	
CLOTRIMAZOLE			
* Vaginal crm 1% with applicators	35 g OP	Clomazol	
* Vaginal crm 2% with applicators3.85	20 g OP	Clomazol	
MICONAZOLE NITRATE			
* Vaginal crm 2% with applicator	40 g OP	✓ Micreme	
NYSTATIN			
Vaginal crm 100,000 u per 5 g with applicator(s)5.70	75 g OP	✓ Nilstat	

## **Myometrial and Vaginal Hormone Preparations**

LINGUINE IMALLATE			
Inj 500 mcg per ml, 1 ml ampoule – Up to 5 inj availa	ıble on a		
PSO	160.00	5	✓ DBL Ergometrine
OESTRIOL			·
* Crm 1 mg per g with applicator	6.95	15 g OP	✓ Ovestin
* Pessaries 500 mcg		15	✓ Ovestin

ERGOMETRINE MALEATE

Pregnancy Test

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
OXYTOCIN – Up to 5 inj available on a PSO Inj 5 iu per ml, 1 ml ampoule	5.98	5	<b>√</b> 0	Oxytocin BNM
Oxytocin BNM to be Principal Supply on 1 March 2026 Inj 10 iu per ml, 1 ml ampoule Oxytocin BNM to be Principal Supply on 1 March 2026	7.18	5	<b>√</b> 0	Oxytocin BNM
OXYTOCIN WITH ERGOMETRINE MALEATE – Up to 5 inj avai Inj 5 iu with ergometrine maleate 500 mcg per ml, 1 ml ampo Syntometrine to be Principal Supply on 1 February 2026	ule41.47	5	<b>√</b> §	Syntometrine

## Pregnancy Tests - hCG Urine

## **Urinary Agents**

For urinary tract Infections refer to INFECTIONS, Antibacterials, page 113

## 5-Alpha Reductase Inhibitors

FINASTERIDE - Special Authority see SA0928 below - Retail pharmacy

★ Tab 5 mg .......4.79 100 ✓ Ricit

#### ⇒SA0928 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 Either:
  - 2.1 The patient is intolerant of non-selective alpha blockers or these are contraindicated; or
  - 2.2 Symptoms are not adequately controlled with non-selective alpha blockers.

## Alpha-1A Adrenoreceptor Blockers

TAMSULOSIN HYDROCHLORIDE - Special Authority see SA1032 below - Retail pharmacy

Tamsulosin-Rex to be Principal Supply on 1 February 2026

#### ⇒SA1032 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 The patient is intolerant of non-selective alpha blockers or these are contraindicated.

	Subsidy (Manufacturer's F \$	Price) Subs Per	Fully idised	Brand or Generic Manufacturer
Other Urinary Agents				
OXYBUTYNIN * Tab 5 mg	5.42	100	<b>√</b> A	lichemy Oxybutynin
POTASSIUM CITRATE				
Oral liq 3 mmol per ml – Special Authority see SA1083 bel Retail pharmacy		200 ml OP	<b>✓</b> E	Biomed
➤ SA1083 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals value to the second substitution of the second substit	alid for 12 months	for applications	meetir	ng the following criteria:
1 The patient has recurrent calcium oxalate urolithiasis; ar				
2 The patient has had more than two renal calculi in the tw Renewal from any relevant practitioner. Approvals valid for 2 y benefitting from the treatment.	, ,		s appro	opriate and the patient is
SODIUM CITRO-TARTRATE				
* Grans eff 4 g sachets	3.50	28	<b>√</b> <u>L</u>	<u>Iral</u>
SOLIFENACIN SUCCINATE				
Tab 5 mg	1.95	30	✓ <u>s</u>	olifenacin succinate Max Health
Tab 10 mg	3.53	30	<b>√</b> <u>s</u>	succinate Max Health
<b>Detection of Substances in Urine</b>				
ORTHO-TOLIDINE				
* Compound diagnostic sticks	7.50 (8.25)	50 test OP	F	lemastix
TETRABROMOPHENOL				
* Blue diagnostic strips	13.92	100 test OP	<b>✓</b> A	Albustix
Obstetric Preparations				

## Obstatio i reparation

# **Antiprogesterones**

			MIFEPRISTONE
<ul><li>Mifegyne</li></ul>	1	D83.90	Tab 200 mg - Up to 15 tab available on a PSO
✓ Mifegyne	3	180.00	

	Subsidy (Manufacturer's Price) \$	Sub Per	Fully sidised	Brand or Generic Manufacturer	
Oalaium Hamaaalaaia					

### Calcium Homeostasis

CALCITONIN  * Inj 100 iu per ml, 1 ml ampoule121.00	5	✓ Miacalcic ✓ Miacalcic S29 S29
CINACALCET – Special Authority see SA2170 below – Retail pharmacy		
Tab 30 mg - Wastage claimable25.24	28	✓ Cinacalet Devatis
Tab 60 mg - Wastage claimable50.47	28	✓ Cinacalet Devatis

⇒SA2170 Special Authority for Subsidy

Initial application — (parathyroid carcinoma or calciphylaxis) only from a nephrologist or endocrinologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 All of the following:
  - 1.1 The patient has been diagnosed with a parathyroid carcinoma (see Note); and
  - 1.2 The patient has persistent hypercalcaemia (serum calcium greater than or equal to 3 mmol/L) despite previous first-line treatments including sodium thiosulfate (where appropriate) and bisphosphonates; and
  - 1.3 The patient is symptomatic; or
- 2 All of the following:
  - 2.1 The patient has been diagnosed with calciphylaxis (calcific uraemic arteriolopathy); and
  - 2.2 The patient has symptomatic (e.g. painful skin ulcers) hypercalcaemia (serum calcium greater than or equal to 3 mmol/L); and
  - 2.3 The patient's condition has not responded to previous first-line treatments including bisphosphonates and sodium thiosulfate.

Renewal — (parathyroid carcinoma or calciphylaxis) only from a nephrologist or endocrinologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient's serum calcium level has fallen to < 3mmol/L: and
- 2 The patient has experienced clinically significant symptom improvement.

Note: This does not include parathyroid adenomas unless these have become malignant.

**Initial application** — (primary hyperparathyroidism) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### All of the following:

- 1 Patient has primary hyperparathyroidism; and
- 2 Fither:
  - 2.1 Patient has hypercalcaemia of more than 3 mmol/L with or without symptoms; or
  - 2.2 Patient has hypercalcaemia of more than 2.85 mmol/L with symptoms; and
- 3 Surgery is not feasible or has failed; and
- 4 Patient has other comorbidities, severe bone pain, or calciphylaxis.

Initial application — (secondary or tertiary hyperparathyroidism) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

#### All of the following:

- 1 Either:
  - 1.1 Patient has tertiary hyperparathyroidism and markedly elevated parathyroid hormone (PTH) with hypercalcaemia; or
  - 1.2 Patient has symptomatic secondary hyperparathyroidism and elevated PTH; and
- 2 Patient is on renal replacement therapy; and
- 3 Any of the following:

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
<u> </u>	Per	•	Manufacturer

continued...

- 3.1 Residual parathyroid tissue has not been localised despite repeat unsuccessful parathyroid explorations; or
- 3.2 Parathyroid tissue is surgically inaccessible; or
- 3.3 Parathyroid surgery is not feasible.

**Renewal — (secondary or tertiary hyperparathyroidism)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

#### Either:

- 1 The patient has had a kidney transplant, and following a treatment free interval of at least 12 weeks a clinically acceptable parathyroid hormone (PTH) level to support ongoing cessation of treatment has not been reached; or
- 2 The patient has not received a kidney transplant and trial of withdrawal of cinacalcet is clinically inappropriate.

## **ZOLEDRONIC ACID**

Inj 4 mg per 5 ml, vial	15.65	1	✓ Zoledronic acid Injection Mylan \$29
			✓ Zoledronic acid

## Corticosteroids and Related Agents for Systemic Use

BETAMETHASONE SODIUM PHOSPHATE WITH BETAMETHASONE ACETA	TE	
* Inj 3.9 mg with betamethasone acetate 3 mg per ml, 1 ml	5	
(36.96)		Celestone
		Chronodose
DEXAMETHASONE		
* Tab 0.5 mg - Up to 60 tab available on a PSO	30	✓ Dexmethsone
* Tab 4 mg - Up to 30 tab available on a PSO	30	✓ Dexmethsone
Oral liq 1 mg per ml53.86	25 ml OP	✓ Biomed
DEXAMETHASONE PHOSPHATE		
Dexamethasone phosphate injection will not be funded for oral use.		
* Inj 4 mg per ml, 1 ml ampoule - Up to 5 inj available on a PSO6.88	10	<ul> <li>Dexamethasone</li> </ul>
		Medsurge
7.86		✓ Hameln
Dexamethasone Medsurge to be Principal Supply on 1 March 2026		
* Inj 4 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO 10.98	10	<ul> <li>Dexamethasone</li> </ul>
		Medsurge
13.10		✓ Hameln
Dexamethasone Medsurge to be Principal Supply on 1 March 2026		
(Hameln Inj 4 mg per ml, 1 ml ampoule to be delisted 1 March 2026)		
(Hameln Inj 4 mg per ml, 2 ml ampoule to be delisted 1 March 2026)		
FLUDROCORTISONE ACETATE		
* Tab 100 mcg8.05	100	✓ Florinef
HYDROCORTISONE		
* Tab 5 mg8.10	100	✓ Douglas
* Tab 20 mg	100	✓ Douglas
* Inj 100 mg vial	1	✓ Solu-Cortef
a) Not on a BSO		
b) Up to 5 inj available on a PSO		
METHYLPREDNISOLONE		
* Tab 4 mg112.00	100	✓ Medrol
* Tab 100 mg223.10	20	✓ Medrol

	Subsidy (Manufacturer's Price \$	) Sub Per	Fully Brand or sidised Generic  Manufacturer
METHYLPREDNISOLONE (AS SODIUM SUCCINATE)			
Inj 40 mg vial	22.30	1	✓ Solu-Medrol-Act- O-Vial
Inj 125 mg vial	34.10	1	✓ Solu-Medrol-Act- O-Vial
Inj 500 mg vial	43.01	1	✓ Solu-Medrol-Act- O-Vial
Inj 1 g vial	52.54	1	✓ Solu-Medrol
METHYLPREDNISOLONE ACETATE			
Inj 40 mg per ml, 1 ml vial	47.06	5	✓ Depo-Medrol
PREDNISOLONE			
Oral liq 5 mg per ml – Up to 30 ml available on a PSO Restricted to children under 12 years of age.	6.00	30 ml OP	✓ Redipred
PREDNISONE			<b>.</b>
F Tab 1 mg		500	✓ Prednisone Clinect
€ Tab 2.5 mg		500	✓ Prednisone Clinect
* Tab 5 mg - Up to 30 tab available on a PSO		500	✓ Prednisone Clinect
€ Tab 20 mg – Up to 30 tab available on a PSO ETRACOSACTRIN	50.51	500	✓ Prednisone Clinect
Finj 250 mcg per ml, 1 ml ampoule	86.25	1	✓ Synacthen
for Inj 1 mg per ml, 1 ml ampoule	690.00	1	✓ UK Synacthen ✓ Synacthen Depot ✓ Synacthene Retard \$29
RIAMCINOLONE ACETONIDE			
Inj 10 mg per ml, 1 ml ampoule	21 42	5	✓ Kenacort-A 10
Inj 40 mg per ml, 1 ml ampoule		5	✓ Kenacort-A 40
Sex Hormones Non Contraceptive			
Androgen Agonists and Antagonists			
PROTERONE ACETATE	47.05	50	<b>( 0</b> !!
Tab 50 mg		50	✓ <u>Siterone</u>
Tab 100 mg	31.00	50	✓ <u>Siterone</u>
ESTOSTERONE  Gel (transdermal) 16.2 mg per g, 88 g	52.00	60 OP	✓ Testogel
ESTOSTERONE CIPIONATE		1	
Inj 100 mg per ml, 10 ml vial ESTOSTERONE ESTERS	85.00	1	✓ Depo-Testosterone
Inj 250 mg per ml, 1 ml	12.98	1	✓ Sustanon Ampoules
ESTOSTERONE UNDECANOATE			
	26.00	100	✓ Steril-Gene S29
Cap 40 mg — Subsidy by endorsement		100	
Subsidy by endorsement – subsidised for patients who 1 November 2021 and the prescription is endorsed ac			
where there exists a record of prior dispensing of testo			
Inj 250 mg per ml, 4 ml vial		tap 40 mg	In the preceding 12 months.  ✓ Reandron 1000
my 200 mg per mi, 4 mi viai	00.00	1	- Iteanurun 1000

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

Sub	osidy Full	y Brand or
(Manufactu	urer's Price) Subsidise	d Generic
	\$ Per ✓	Manufacturer Manufacturer

# **Hormone Replacement Therapy - Systemic**

# Oestrogens

OE	STRADIOL			
*	Tab 1 mg	4.12	28 OP	
		(11.10)		Estrofem
*	Tab 2 mg	4.12	28 OP	
		(11.10)		Estrofem
*	Gel (transdermal) 0.06% (750 mcg/actuation)	14.25	80 g OP	✓ Estrogel
	Patch 25 mcg per day		8	✓ Estradiol TDP Mylan
		16.23		✓ Estradot
	<ul> <li>a) Brand switch fee payable (Pharmacode 2717573) -</li> <li>b) No more than 2 patch per week</li> <li>c) Only on a prescription</li> </ul>	see page 282 for	details	
	Patch 50 mcg per day	9.26	8	<ul><li>Estradiol TDP Mylan</li></ul>
	<b>31</b> ,	15.79		✓ Estradot
	<ul><li>a) Brand switch fee payable (Pharmacode 2717573) -</li><li>b) No more than 2 patch per week</li><li>c) Only on a prescription</li></ul>			
	Patch 75 mcg per day		8	✓ Estradiol TDP Mylan
		16.53		✓ Estradot
	<ul><li>a) Brand switch fee payable (Pharmacode 2717573) -</li><li>b) No more than 2 patch per week</li><li>c) Only on a prescription</li></ul>	see page 282 for	details	
	Patch 100 mcg per day	10.59	8	✓ Estradiol TDP Mylan
		16.18		✓ Estradot
OF	a) Brand switch fee payable (Pharmacode 2717573) -     b) No more than 2 patch per week     c) Only on a prescription  STRADIOL VALERATE	see page 282 for	details	
*	Tab 1 mg	12.36	84	✓ Progynova
*	Tab 2 mg		84	✓ Progynova
	<u> </u>	12.00	04	1 logynova
	STROGENS	0.01	00	
*	Conjugated, equine tab 300 mcg		28	Duamania
	Operior and a series tale 205 man	(19.25)	00	Premarin
*	Conjugated, equine tab 625 mcg		28	Description
		(19.25)		Premarin

## **Progestogens**

ME	DROXYPROGESTERONE ACETATE		
*	Tab 2.5 mg	30	✓ Provera
	8.75	56	✓ Provera
*	Tab 5 mg9.80	56	✓ Provera
	20.13	100	✓ Provera
*	Tab 10 mg10.28	30	✓ Provera

	Subsidy (Manufacturer's Price \$	e) S Per	Fully Brand or subsidised Generic ✓ Manufacturer
Progestogen and Oestrogen Combined Prepara	ations		
OESTRADIOL WITH NORETHISTERONE			
* Tab 1 mg with 0.5 mg norethisterone acetate	5.40	28 OP	
	(18.10)		Kliovance
* Tab 2 mg with 1 mg norethisterone acetate		28 OP	10:
	(18.10)		Kliogest
* Tab 2 mg with 1 mg norethisterone acetate (10), and 2 mg	F 40	00.00	
oestradiol tab (12) and 1 mg oestradiol tab (6)		28 OP	Trianguana
	(18.10)		Trisequens
Other Oestrogen Preparations			
The Society of French Control			
OESTRIOL			_
* Tab 2 mg	7.70	30	✓ Ovestin
Other Bregestagen Brenevations			
Other Progestogen Preparations			
MEDROXYPROGESTERONE ACETATE			
Tab 100 mg	133.57	100	✓ Provera HD
NORETHISTERONE			
* Tab 5 mg - Up to 30 tab available on a PSO	5.49	30	✓ Primolut N
PROGESTERONE			
* Cap 100 mg	14.85	30	✓ Utrogestan
Thyroid and Antithyroid Agents			
CARBIMAZOLE			
* Tab 5 mg	7.56	100	✓ Neo-Mercazole
-	7.50	100	<u>Neo-IMEI Cazole</u>
LEVOTHYROXINE  * Tab 25 mcg	5 55	90	✓ Synthroid
* Tab 23 mcg*		28	✓ Mercury Pharma
Tab oo mog	5.79	90	✓ Synthroid
	64.28	1,000	✓ Eltroxin
* Tablet 50 mcg		200	✓ Eltroxin
* Tab 100 mcg		28	Mercury Pharma
	6.01	90	✓ Synthroid
	66.78	1,000	✓ Eltroxin
* Tablet 100 mcg		200	✓ Eltroxin
PROPYLTHIOURACIL - Special Authority see SA1199 below -	- Retail pharmacy		
Tab 50 mg	35.00	100	✓ PTU S29

**⇒SA1199** Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 The patient has hyperthyroidism; and
- 2 The patient is intolerant of carbimazole or carbimazole is contraindicated.

**Renewal** from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	I Generic	
\$	Per 🗸	Manufacturer	

## **Trophic Hormones**

#### **Growth Hormones**

SO	MATROPIN (OMNITROPE) - Special Authority see SA20	cial Authority see SA2032 below – Retail pharmacy		
*	Inj 5 mg cartridge	80.21	1	✓ Omnitrope
				✓ Omnitrope AU S29
*	Inj 10 mg cartridge	80.21	1	✓ Omnitrope
*	Inj 15 mg cartridge	139.50	1	✓ Omnitrope

⇒SA2032 Special Authority for Subsidy

Initial application — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria: Either:

- - 1 Growth hormone deficiency causing symptomatic hypoglycaemia, or with other significant growth hormone deficient sequelae (e.g. cardiomyopathy, hepatic dysfunction) and diagnosed with GH < 5 mcg/l on at least two random blood samples in the first 2 weeks of life, or from samples during established hypoglycaemia (whole blood glucose < 2 mmol/l using a laboratory device); or
  - 2 All of the following:
    - 2.1 Height velocity < 25th percentile for age adjusted for bone age/pubertal status if appropriate over 6 or 12 months using the standards of Tanner and Davies (1985); and
    - 2.2 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
    - 2.3 Peak growth hormone value of < 5.0 mcg per litre in response to two different growth hormone stimulation tests. In children who are 5 years or older, GH testing with sex steroid priming is required; and
    - 2.4 If the patient has been treated for a malignancy, they should be disease free for at least one year based upon follow-up laboratory and radiological imaging appropriate for the malignancy, unless there are strong medical reasons why this is either not necessary or appropriate; and
    - 2.5 Appropriate imaging of the pituitary gland has been obtained.

Renewal — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 2 Height velocity is greater than or equal to 25th percentile for age (adjusted for bone age/pubertal status if appropriate) while on growth hormone treatment, as calculated over six months using the standards of Tanner and Davis (1985); and
- 3 Height velocity is greater than or equal to 2.0 cm per year, as calculated over 6 months; and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone treatment has occurred: and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a post-natal genotype confirming Turner Syndrome; and
- 2 Height velocity is < 25th percentile over 6-12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is < 14 years.

Renewal — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 Height velocity is greater than or equal to 50th percentile for age (while on growth hormone calculated over 6 to 12 months using the Ranke's Turner Syndrome growth velocity charts); and

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer	
--	--

continued...

- 2 Height velocity is greater than or equal to 2 cm per year, calculated over six months; and
- 3 A current bone age is 14 years or under; and
- 4 No serious adverse effect that the specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 3 standard deviations below the mean for age or for bone age if there is marked growth acceleration or delay; and
- 2 Height velocity is < 25th percentile for age (adjusted for bone age/pubertal status if appropriate), as calculated over 6 to 12 months using the standards of Tanner and Davies(1985); and
- 3 A current bone age is < 14 years or under (female patients) or < 16 years (male patients); and
- 4 The patient does not have severe chronic disease (including malignancy or recognized severe skeletal dysplasia) and is not receiving medications known to impair height velocity.

Renewal — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred.

**Initial application** — **(short stature due to chronic renal insufficiency)** only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 2 standard deviations below the mean; and
- 2 Height velocity is < 25th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is to 14 years or under (female patients) or to 16 years or under (male patients); and
- 4 The patient is metabolically stable, has no evidence of metabolic bone disease and absence of any other severe chronic disease; and
- 5 The patient is under the supervision of a specialist with expertise in renal medicine; and
- 6 Either
  - 6.1 The patient has a GFR less than or equal to 30 ml/min/1.73m<sup>2</sup> as measured by the Schwartz method (Height(cm)/plasma creatinine (umol/l)) × 40 = corrected GFR (ml/min/1.73m<sup>2</sup>) in a child who may or may not be receiving dialysis; or
  - 6.2 The patient has received a renal transplant and has received < 5mg/ m²/day of prednisone or equivalent for at least 6 months..

Renewal — (short stature due to chronic renal insufficiency) only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and

(Mar	Subsidy nufacturer's Price)	Sub	Fully sidised	Brand or Generic
	\$	Per	1	Manufacturer

continued...

- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone has occurred;
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not experienced significant biochemical or metabolic deterioration confirmed by diagnostic results; and
- 7 The patient has not received renal transplantation since starting growth hormone treatment; and
- 8 If the patient requires transplantation, growth hormone prescription should cease before transplantation and a new application should be made after transplantation based on the above criteria.

**Initial application — (Prader-Willi syndrome)** only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of Prader-Willi syndrome that has been confirmed by genetic testing or clinical scoring criteria; and
- 2 The patient is aged six months or older; and
- 3 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
- 4 Sleep studies or overnight oximetry have been performed and there is no obstructive sleep disorder requiring treatment, or if an obstructive sleep disorder is found, it has been adequately treated under the care of a paediatric respiratory physician and/or ENT surgeon; and
- 5 Fither:
  - 5.1 Both:
    - 5.1.1 The patient is aged two years or older; and
    - 5.1.2 There is no evidence of type II diabetes or uncontrolled obesity defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months; or
  - 5.2 The patient is aged between six months and two years and a thorough upper airway assessment is planned to be undertaken prior to treatment commencement and at six to 12 weeks following treatment initiation.

Renewal — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not developed type II diabetes or uncontrolled obesity as defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months.

Initial application — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a medical condition that is known to cause growth hormone deficiency (e.g. surgical removal of the pituitary for treatment of a pituitary tumour); and
- 2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
- 3 The patient has severe growth hormone deficiency (see notes); and
- 4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
- 5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Por 🗸	Manufacturer

continued...

questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

**Renewal — (adults and adolescents)** only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

4 All af the fallaction

- 1 All of the following:
  - 1.1 The patient has been treated with somatropin for < 12 months; and
  - 1.2 There has been an improvement in Quality of Life defined as a reduction of at least 8 points on the Quality of Life Assessment of Growth Hormone Deficiency in Adults (QoL-AGHDA®) score from baseline; and
  - 1.3 Serum IGF-I levels have been increased within ±1SD of the mean of the normal range for age and sex; and
  - 1.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients, or 1 mg per day for female patients; or
- 2 All of the following:
  - 2.1 The patient has been treated with somatropin for more than 12 months; and
  - 2.2 The patient has not had a deterioration in Quality of Life defined as a 6 point or greater increase from their lowest QoL-AGHDA® score on treatment (other than due to obvious external factors such as external stressors); and
  - 2.3 Serum IGF-I levels have continued to be maintained within ±1SD of the mean of the normal range for age and sex (other than for obvious external factors); and
  - 2.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients or 1 mg per day for female patients; or
- 3 All of the following:
  - 3.1 The patient has had a Special Authority approval for somatropin for childhood deficiency in children and no longer meets the renewal criteria under this indication; and
  - 3.2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
  - 3.3 The patient has severe growth hormone deficiency (see notes); and
  - 3.4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
  - 3.5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement

	Subsidy (Manufacturer's Price)	Fully Subsidised Per	Brand or Generic Manufacturer
continued doses of corticosteroid and levothyroxine.			

## **GnRH Analogues**

GOSERELIN			
Implant 3.6 mg, syringe	66.48	1	✓ Zoladex
Implant 10.8 mg, syringe		1	✓ Zoladex
LEUPRORELIN			
Additional subsidy by endorsement where the patient is a goserelin and the prescription is endorsed accordingly.		ınd is unable	e to tolerate administration of
Inj 3.75 mg prefilled dual chamber syringe - Higher subs	sidy of		
\$221.60 per 1 inj with Endorsement	66.48	1	
,	(221.60)		Lucrin Depot 1-month
Inj 11.25 mg prefilled dual chamber syringe - Higher sub	osidy		
of \$591.68 per 1 inj with Endorsement	177.50	1	
	(591.68)		Lucrin Depot 3-month

## Vasopressin Agonists

DESMOPRESSIN Wafer 120 mcg	47.00	30	✓ Minirin Melt
DESMOPRESSIN ACETATE			
Tab 100 mcg	25.00	30	✓ Minirin
Tab 200 mcg	54.45	30	✓ Minirin
Inj 4 mcg per ml, 1 ml	67.18	10	✓ Minirin
▲ Nasal spray 10 mcg per dose, 6 ml	34.95	60 OP	✓ <u>Desmopressin-</u>
Inj 4 mcg per ml, 1 ml	67.18	10	✓ Minirin

# **Other Endocrine Agents**

#### CABERGOLINE

		Tab 0.5 mg - Maximum of 2 tab per prescription; can be
<ul><li>Dostinex</li></ul>	2	waived by Special Authority see SA2070 below4.43
✓ Dostinex	8	17.94

### ⇒SA2070 Special Authority for Waiver of Rule

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Hyperprolactinemia; or
- 2 Acromegaly\*; or
- 3 Inhibition of lactation.

Renewal — (for patients who have previously been funded under Special Authority form SA1031) from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has previously held a valid Special Authority which has expired and the treatment remains appropriate and the patient is benefiting from treatment.

Note: Indication marked with \* is an unapproved indication.

CLOMIFENE CITRATE
-------------------

Tab 50 mg	29.84	10	✓ Mylan  Clomiphen S29
METYRAPONE Cap 250 mg	558.00	50	✓ Metopirone

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sul	osidised	Generic
\$	Per	/	Manufacturer

## **Anthelmintics**

ALBENDAZOLE - Special Authority see SA2512 below - Retai
--

## ⇒SA2512 Special Authority for Subsidy

**Initial application** only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: Either:

- 1 The individual has hydatids; or
- 2 The individual has a travel or residence history that requires presumptive parasite treatment.

**Renewal** only from an infectious disease specialist or clinical microbiologist. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefitting from the treatment.

## MEBENDAZOLE - Only on a prescription

Tab 100 mg	5.18	6	✓ Vermox
Oral liq 100 mg per 5 ml		15 ml	<del></del>
, ,,	(7.83)		Vermox
PRAZIQUANTEL			
Tab 600 mg	68.00	8	<ul> <li>Biltricide</li> </ul>
	87.68		✓ Distoside S29
(Riltricide Tah 600 mg to be delisted 1 April 2026)			

## (Biltricide Tab 600 mg to be delisted 1 April 2026)

## **Antibacterials**

- a) For topical antibacterials, refer to DERMATOLOGICALS, page 68
- b) For anti-infective eye preparations, refer to SENSORY ORGANS, page 277

## Cephalosporins and Cephamycins

CEFACLOR MONOHYDRATE			
Cap 250 mg	29.73	100	✓ Ranbaxy-Cefaclor
Ranbaxy-Cefaclor to be Principal Supply on 1 February 2	2026		
Grans for oral liq 125 mg per 5 ml – Wastage claimable Ranbaxy-Cefaclor to be Principal Supply on 1 February 2		100 ml	✓ Ranbaxy-Cefaclor
CEFALEXIN			
Cap 250 mg	3.85	20	<ul> <li>Cephalexin ABM</li> </ul>
	3.90		<ul><li>Cefalexin Lupin</li></ul>
Cap 500 mg	3.33	20	✓ Cefalexin Sandoz
	5.85		<ul><li>Cephalexin ABM</li></ul>
Grans for oral lig 25 mg per ml - Wastage claimable	7.88	100 ml	✓ Flynn
Grans for oral lig 50 mg per ml - Wastage claimable	10.38	100 ml	✓ Flynn
, 0,	11.75		✓ Cefalexin Sandoz
(Cephalexin ABM Cap 250 mg to be delisted 1 July 2026)			

(Cephalexin ABM Cap 250 mg to be delisted 1 July 2026) (Cephalexin ABM Cap 500 mg to be delisted 1 July 2026)

CEFAZOLIN - Subsidy by endorsement

Only if prescribed for dialysis or cellulitis in accordance with a Health NZ Hospital approved protocol and the prescription is endorsed accordingly.

Inj 500 mg vial	5 •	Cefazolin-AFT
Inj 1 g vial3.59	5	Cefazolin-AFT
Inj 2 g vial7.09	5	Cefazolin-AFT

	(Manufacturer's Price)	Per	Subsidised	Generic Manufacturer
EFTRIAXONE - Subsidy by endorsement	<u>-</u>			
a) Up to 10 inj available on a PSO				
<ul> <li>Subsidised only if prescribed for a dialysis or cystic fibrosi pelvic inflammatory disease, or the treatment of suspected endorsed accordingly.</li> </ul>			•	·
Inj 500 mg vial		1	✓	Ceftriaxone-AFT
Inj 1 g vial Ceftriaxone-AFT to be Principal Supply on 1 February 20	3.49	5	✓	Ceftriaxone-AFT
EFUROXIME AXETIL - Subsidy by endorsement Only if prescribed for prophylaxis of endocarditis and the pres	scription is endorsed	acco	dingly.	
Tab 250 mg	•	20	٠,	Ascend- Cefuroxime S29

Subeidy

Fully

Brand or

### **Macrolides**

CF

CF

#### ⇒SA1683 Special Authority for Waiver of Rule

Initial application — (bronchiolitis obliterans syndrome, cystic fibrosis and atypical Mycobacterium infections) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Patient has received a lung transplant, stem cell transplant, or bone marrow transplant and requires treatment for bronchiolitis obliterans syndrome\*; or
- 2 Patient has received a lung transplant and requires prophylaxis for bronchiolitis obliterans syndrome\*; or
- 3 Patient has cystic fibrosis and has chronic infection with Pseudomonas aeruginosa or Pseudomonas-related gram negative organisms\*; or
- 4 Patient has an atypical Mycobacterium infection.

Note: Indications marked with \* are unapproved indications.

Initial application — (non-cystic fibrosis bronchiectasis\*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 For prophylaxis of exacerbations of non-cystic fibrosis bronchiectasis\*; and
- 2 Patient is aged 18 and under; and
- 3 Either:
  - 3.1 Patient has had 3 or more exacerbations of their bronchiectasis, within a 12 month period; or
  - 3.2 Patient has had 3 acute admissions to hospital for treatment of infective respiratory exacerbations within a 12 month period.

Note: Indications marked with \* are unapproved indications.

Renewal — (non-cystic fibrosis bronchiectasis\*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Pri	ce)	Subsidised	Generic	
\$	Per	•	Manufacturer	

continued...

- 1 The patient has completed 12 months of azithromycin treatment for non-cystic fibrosis bronchiectasis; and
- 2 Following initial 12 months of treatment, the patient has not received any further azithromycin treatment for non-cystic fibrosis bronchiectasis for a further 12 months, unless considered clinically inappropriate to stop treatment; and
- 3 The patient will not receive more than a total of 24 months' azithromycin cumulative treatment (see note).

The patient must not have had more than 1 prior approval.

Note: No further renewals will be subsidised. A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised. Indications marked with \* are unapproved indications

CLARITHROMYCIN - Maximum of 500 mg per prescription; can be waived by Special Authority see SA1857 below

Tab 250 mg	7.31	12	✓ Klaricid S29
v	8.53	14	✓ Klacid
Grans for oral lig 250 mg per 5 ml - Wastage claimable	192.00	50 ml	✓ Klacid

#### ⇒SA1857 Special Authority for Waiver of Rule

Initial application — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years for applications meeting the following criteria:

Fither:

- 1 Atypical mycobacterial infection; or
- 2 Mycobacterium tuberculosis infection where there is drug-resistance or intolerance to standard pharmaceutical agents.

Initial application — (Helicobacter pylori eradication) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 For the eradication of helicobacter pylori in a patient unable to swallow tablets; and
- 2 For use only in combination with omeprazole and amoxicillin as part of a triple therapy regimen.

Initial application — (Prophylaxis of infective endocarditis) from any relevant practitioner. Approvals valid for 3 months where prophylaxis of infective endocarditis associated with surgical or dental procedures if amoxicillin is contra-indicated. Renewal — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

ERYTHROMYCIN (AS LACTOBIONATE)

Inj 1 g vial	10.00	1	✓ Erythrocin IV
ERYTHROMYCIN ETHYL SUCCINATE			
Tab 400 mg	35.82	100	<ul><li>E-Mycin</li></ul>
a) Up to 20 tab available on a PSO			
b) Up to 2 x the maximum PSO quantity for RFPP			
Grans for oral liq 200 mg per 5 ml	6.53	100 ml	<ul><li>E-Mycin</li></ul>
<ul> <li>a) Up to 300 ml available on a PSO</li> </ul>			
b) Up to 2 x the maximum PSO quantity for RFPP			
c) Wastage claimable			
Grans for oral liq 400 mg per 5 ml	9.41	100 ml	E-Mycin
a) Up to 200 ml available on a PSO			
b) Wastage claimable			
ROXITHROMYCIN			
Tab 150 mg	13.19	50	✓ Arrow-
			Roxithromycin
Tab 300 mg	25.00	50	✓ Arrow-
Tab ood mg	23.00	50	Roxithromycin

Penicillins  AMOXICILLIN Cap 250 mg
AMOXICILLIN Cap 250 mg
Cap 250 mg
<ul> <li>a) Up to 30 cap available on a PSO</li> <li>b) Up to 10 x the maximum PSO quantity for RFPP</li> <li>c) Miro-Amoxicillin to be Principal Supply on 1 February 2026</li> </ul>
<ul><li>b) Up to 10 x the maximum PSO quantity for RFPP</li><li>c) Miro-Amoxicillin to be Principal Supply on 1 February 2026</li></ul>
c) Miro-Amoxicillin to be Principal Supply on 1 February 2026
, , , , , , , , , , , , , , , , , , , ,
Cap 500 mg41.00 500 <b>✓ Miro-Amoxicillin</b>
a) Up to 30 cap available on a PSO
b) Up to 10 x the maximum PSO quantity for RFPP
Grans for oral liq 125 mg per 5 ml
a) Up to 200 ml available on a PSO     b) Wastage claimable
Grans for oral liq 250 mg per 5 ml2.81 100 ml ✓ Alphamox 250
a) Up to 300 ml available on a PSO
b) Up to 10 x the maximum PSO quantity for RFPP
c) Wastage claimable
Inj 250 mg vial15.97 10
Inj 500 mg vial
Inj 1 g vial − Up to 5 inj available on a PSO21.64 10 ✓ Ibiamox
AMOXICILLIN WITH CLAVULANIC ACID
Tab 500 mg with clavulanic acid 125 mg - Up to 30 tab
available on a PSO1.59 10 <b>✓ Curam Duo 500/125</b>
Grans for oral liq amoxicillin 25 mg with clavulanic acid 6.25 mg
per ml8.50 100 ml ✓ <u>Augmentin</u>
a) Up to 200 ml available on a PSO
b) Wastage claimable
Grans for oral liq amoxicillin 50 mg with clavulanic acid 12.5 mg
per ml − Up to 200 ml available on a PSO5.61 100 ml OP ✓ Amoxiclav Devatis
<u>Forte</u>
BENZATHINE BENZYLPENICILLIN
Inj 900 mg (1.2 million units) vial
Inj 900 mg (1.2 million units) in 2.3 ml syringe – Up to 5 inj
available on a PSO432.37 10 Sicillin LA
BENZYLPENICILLIN SODIUM [PENICILLIN G]
Inj 600 mg (1 million units) vial − Up to 5 inj available on a PSO16.50 10 ✓ Sandoz
FLUCLOXACILLIN
Cap 250 mg − Up to 30 cap available on a PSO22.58 250 ✓ <u>Staphlex</u>
Cap 500 mg — Up to 30 cap available on a PSO72.71 500 Staphlex
Grans for oral liq 25 mg per ml
a) Up to 200 ml available on a PSO
b) Wastage claimable Grans for oral liq 50 mg per ml5.89 100 ml ✓ AFT
a) Up to 200 ml available on a PSO
b) Wastage claimable
Inj 250 mg vial
Inj 500 mg vial
Inj 1 g vial – Up to 5 inj available on a PSO6.00 5 Flucil

	Subsidy (Manufacturer's Price)	Cub	Fully	Brand or Generic
	(Manuacturer 5 Frice)	Per	Sidised	Manufacturer
	Ψ	r ei		Manuacturei
PHENOXYMETHYLPENICILLIN (PENICILLIN V)				
Cap 250 mg - Up to 30 cap available on a PSO	7.68	50	✓ (	Cilicaine VK
Cap 500 mg	13.72	50	1	Cilicaine VK
a) Up to 20 cap available on a PSO				
b) Up to 2 x the maximum PSO quantity for RFPP				
Grans for oral liq 125 mg per 5 ml	5.75	100 ml	<b>✓</b>	<b>NFT</b>
a) Up to 200 ml available on a PSO				
b) Wastage claimable				
c) AFT to be Principal Supply on 1 February 2026				
Grans for oral lig 250 mg per 5 ml	5.89	100 ml	<b>✓</b>	\FT
a) Up to 300 ml available on a PSO				
b) Up to 2 x the maximum PSO quantity for RFPP				
c) Wastage claimable				
d) AFT to be Principal Supply on 1 February 2026				
a, All 1 to be 1 illioipal dupply off 1 1 oblidary 2020				

## **Tetracyclines**

DOXYCYCLIN	E			
★ Tab 100 m	ng - Up to 30 tab available on a PSO	64.43	500	Doxine
MINOCYCLINI	E HYDROCHLORIDE			
* Tab 50 mg	- Additional subsidy by Special Authority see			
SA13	55 below – Retail pharmacy	5.79	60	
		(12.05)		Mino-tabs
* Cap 100 n	ng	19.32	100	
		(52.04)		Minomycin

## **⇒SA1355** Special Authority for Manufacturers Price

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has rosacea.

Initial application from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 For the eradication of helicobacter pylori following unsuccessful treatment with appropriate first-line therapy; and
- 2 For use only in combination with bismuth as part of a quadruple therapy regimen.

Renewal from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 For the eradication of helicobacter pylori following unsuccessful treatment with, or noncompletion of second line therapy; and
- 2 For use only in combination with bismuth as part of a quadruple therapy regimen.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
Other Antibiotics				
or topical antibiotics, refer to DERMATOLOGICALS, page 68 IPROFLOXACIN Recommended for patients with any of the following: i) microbiologically confirmed and clinically significant ps ii) prostatitis; or iii) pyelonephritis; or iv) gonorrhoea.	eudomonas infection;	or		
Tab 250 mg - Up to 5 tab available on a PSO	1.95	28	1	Ipca-Ciprofloxacin
Tab 500 mg - Up to 5 tab available on a PSO		28		Ipca-Ciprofloxacin
Tab 750 mg	4.80	28	•	Ipca-Ciprofloxacin
LINDAMYCIN			_	
Cap hydrochloride 150 mg		24		Dalacin C
Inj 150 mg per ml, 4 ml ampoule	35.10 48.78	10		Hameln Dalacin C
Dalacin C to be Principal Supply on 1 March 2026  Hameln Inj 150 mg per ml, 4 ml ampoule to be delisted 1 March  OLISTIN SULPHOMETHATE – Retail pharmacy-Specialist – \$	2026)	ent	·	Duidoni O
Only if prescribed for dialysis or cystic fibrosis patient and th			accordingl	y.
Inj 2 million iu, 10 ml vial		10	_	Colomycin S29
ENTAMICIN SULPHATE				·
Inj 40 mg per ml, 2 ml vial - Subsidy by endorsement	36.70	5	•	Cidomycin P/Free S29
Only if prescribed for a dialysis or cystic fibrosis patient endorsed accordingly.	or complicated urinary	trac	t infection	and the prescription is
Inj 10 mg per ml, 1 ml ampoule – Subsidy by endorsement. Only if prescribed for a dialysis or cystic fibrosis patient endorsed accordingly.		5 trac		<b>DBL Gentamicin</b> and the prescription is
Inj 40 mg per ml, 2 ml ampoule - Subsidy by endorsement.	18.38	10	1	Pfizer
Only if prescribed for a dialysis or cystic fibrosis patient endorsed accordingly.	or complicated urinary	trac	t infection	and the prescription is
OXIFLOXACIN - Special Authority see SA1740 below - Retain	l pharmacy			
No patient co-payment payable	40.00	_	,	Auglau
Tab 400 mg SA1740 Special Authority for Subsidy	42.00	5	•	Avelox

for applications meeting the following criteria:

Any of the following:

- 1 Both:
  - 1.1 Active tuberculosis\*; and
  - 1.2 Any of the following:
    - 1.2.1 Documented resistance to one or more first-line medications; or
    - 1.2.2 Suspected resistance to one or more first-line medications (tuberculosis assumed to be contracted in an area with known resistance), as part of regimen containing other second-line agents; or

(Manufacturer's Price) Subsidised Generic \$ Per ✓ Manufacturer	Subsidy	Full	y Brand or
\$ Per ✓ Manufacturer	(Manufacturer's	s Price) Subsidise	d Generic
	\$	Per <b>✓</b>	Manufacturer

continued...

- 1.2.3 Impaired visual acuity (considered to preclude ethambutol use); or
- 1.2.4 Significant pre-existing liver disease or hepatotoxicity from tuberculosis medications; or
- 1.2.5 Significant documented intolerance and/or side effects following a reasonable trial of first-line medications; or
- 2 Mycobacterium avium-intracellulare complex not responding to other therapy or where such therapy is contraindicated.\*; or
- 3 Patient is under five years of age and has had close contact with a confirmed multi-drug resistant tuberculosis case.

Note: Indications marked with \* are unapproved indications.

**Renewal** only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Mycoplasma genitalium) only from a sexual health specialist or Practitioner on the recommendation of a sexual health specialist. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Has nucleic acid amplification test (NAAT) confirmed Mycoplasma genitalium\* and is symptomatic; and
- 2 Either:
  - 2.1 Has tried and failed to clear infection using azithromycin; or
  - 2.2 Has laboratory confirmed azithromycin resistance; and
- 3 Treatment is only for 7 days.

Initial application — (Penetrating eye injury) only from an ophthalmologist. Approvals valid for 1 month where the patient requires prophylaxis following a penetrating eye injury and treatment is for 5 days only.

Note: Indications marked with \* are unapproved indications.

PAROMOMYCIN - Special Authority see SA1689 below - Retail pharmacy

Cap 250 mg.......126.00 16 **✓ Humatin** 529

## **⇒SA1689** Special Authority for Subsidy

Initial application only from an infectious disease specialist, clinical microbiologist or gastroenterologist. Approvals valid for 1 month for applications meeting the following criteria:

Fither:

- 1 Patient has confirmed cryptosporidium infection; or
  - 2 For the eradication of Entamoeba histolytica carriage.

Renewal only from an infectious disease specialist, clinical microbiologist or gastroenterologist. Approvals valid for 1 month for applications meeting the following criteria:

Either:

- 1 Patient has confirmed cryptosporidium infection; or
- 2 For the eradication of Entamoeba histolytica carriage.

#### SODIUM FUSIDATE (FUSIDIC ACID)

Tab 250 mg135.70	36	✓ Fucidin
SULFADIAZINE SODIUM - Special Authority see SA1331 below - Retail pharmacy		
Tab 500 mg	100	✓ Sulfadiazin-Hevl S29

#### ⇒SA1331 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 For the treatment of toxoplasmosis in patients with HIV for a period of 3 months; or
- 2 For pregnant patients for the term of the pregnancy; or
- 3 For infants with congenital toxoplasmosis until 12 months of age.

	0.1.11			-
	Subsidy	-\ 0	Fully	Brand or
	(Manufacturer's Price	e) Sub: Per	sidised	Generic Manufacturer
	<b>Ф</b>	Per		Manufacturer
TOBRAMYCIN				
Inj 40 mg per ml, 2 ml vial - Subsidy by endorsement	15.50	5	<b>✓</b> T	obramycin (Viatris)
Only if prescribed for dialysis or cystic fibrosis patient an	d the prescription is	endorsed	accordin	ngly.
Solution for inhalation 60 mg per ml, 5 ml - Subsidy by				
endorsement	395.00	56 dose	<b>✓</b> T	obramycin BNM
a) Wastage claimable			_	
b) Only if prescribed for a cystic fibrosis patient and the	prescription is endo	orsed accor	dinaly.	
, , , , , , , , , , , , , , , , , , , ,	procempular le cria		ug.,.	
TRIMETHOPRIM	07.00	50		un.
* Tab 300 mg - Up to 30 tab available on a PSO	27.83	50	V 1	<u>MP</u>
TRIMETHOPRIM WITH SULPHAMETHOXAZOLE [CO-TRIMOX	(AZOLE)			
* Tab trimethoprim 80 mg and sulphamethoxazole 400 mg - U	Up			
to 30 tab available on a PSO	115.74	500	<b>√</b> T	risul
* Oral liq 8 mg sulphamethoxazole 40 mg per ml - Up to 200			_	
available on a PSO		100 ml	<b>✓</b> D	eprim
		100 1111	٠ ٢	<del>орин</del>
VANCOMYCIN – Subsidy by endorsement				
Only if prescribed for a dialysis or cystic fibrosis patient or for			for treat	tment of Clostridium
difficile following metronidazole failure and the prescription is	endorsed accordin	ıgly.		
Inj 500 mg vial	3.38	1	✓ N	<u>lylan</u>
			✓ V	ancomycin Viatris
(Mylan Inj 500 mg vial to be delisted 1 March 2026)				

## **Antifungals**

- a) For topical antifungals refer to DERMATOLOGICALS, page 69
- b) For topical antifungals refer to GENITO URINARY, page 80

#### **FLUCONAZOLE**

00011112022		
Cap 50 mg4.10	28	Mylan
Cap 150 mg	5 1	✓ Mylan
Cap 200 mg	28	✓ Mylan
Powder for oral suspension 10 mg per ml - Special Authority		
see SA1359 below – Retail pharmacy	2 35 ml	<ul><li>Diflucan</li></ul>
Wastage claimable		

### ⇒SA1359 Special Authority for Subsidy

**Initial application — (Systemic candidiasis)** from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

#### Both:

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

**Initial application — (Immunocompromised)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Patient is at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

Renewal — (Systemic candidiasis) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

30

✓ Burel S29

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsidised		Generic
\$	Per	✓	Manufacturer

#### continued...

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

**Renewal — (Immunocompromised)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

#### All of the following:

- 1 Patient remains immunocompromised; and
- 2 Patient remains at moderate to high risk of invasive fungal infection; and

Tab 200 mg - PCT......CBS

3 Patient is unable to swallow capsules.

#### **ITRACONAZOLE**

Cap 100 mg6.83	15	✓ Itraconazole
		Cresent S29  ✓ Itrazole
Oral liq 10 mg per ml - Special Authority see SA1322 below -		
Retail pharmacy141.80	150 ml OP	✓ Itraconazole  Kent \$29

### ⇒SA1322 Special Authority for Subsidy

**Initial application** only from an infectious disease specialist, clinical microbiologist, clinical immunologist or any relevant practitioner on the recommendation of a infectious disease physician, clinical microbiologist or clinical immunologist. Approvals valid for 6 months where the patient has a congenital immune deficiency.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefitting from the treatment.

#### **KETOCONAZOLE**

		100	✓ Strides Shasun S29 ✓ Taro S29 ✓ Teva- Ketoconazole S29
NYSTATIN			
Tab 500,000 u	14.16	50	
	(17.09)		Nilstat
Cap 500,000 u	12.81	50	
	(15.47)		Nilstat
POSACONAZOLE - Special Authority see SA2383 below -	Retail pharmacy		
Tab modified-release 100 mg	123.60	24	✓ Posaconazole Juno
Oral liq 40 mg per ml	308.26	105 ml OP	✓ Devatis
CACCOO Created Authority for Cubaldy			

#### ⇒SA2383 Special Authority for Subsidy

**Initial application** only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

#### Either:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation chemotherapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppressive therapy\*.

Renewal only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the

Subsidy		Fully	Brand or	
(Manufacturer's Price	e)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

following criteria:

#### Either:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation therapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppression\* and requires on going posaconazole treatment.

Note: \* Graft versus host disease (GVHD) on significant immunosuppression is defined as acute GVHD, grade II to IV, or extensive chronic GVHD, or if they were being treated with intensive immunosuppressive therapy consisting of either high-dose corticosteroids (1 mg or greater per kilogram of body weight per day for patients with acute GVHD or 0.8 mg or greater per kilogram every other day for patients with chronic GVHD), antithymocyte globulin, or a combination of two or more immunosuppressive agents or types of treatment.

Initial application — (Invasive fungal infection prophylaxis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient is at risk of invasive fungal infection; and
- 2 Either:
  - 2.1 Posaconazole is prescribed by, or recommended by a haematologist, transplant physician, infectious disease specialist, paediatric haematologist or paediatric oncologist; or
  - 2.2 Prescribing posaconazole is in accordance with a protocol or guideline that has been endorsed by the Health New Zealand - Te Whatu Ora Hospital in the specific settings where there is a greater than 10% risk of invasive fungal infection (IFI).

Renewal — (Invasive fungal infection prophylaxis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- Dour.
  - 1 The patient is at risk of invasive fungal infection; and
  - 2 Either:
    - 2.1 Posaconazole is prescribed by, or recommended by a haematologist, transplant physician, infectious disease specialist, paediatric haematologist or paediatric oncologist; or
    - 2.2 Prescribing posaconazole is in accordance with a protocol or guideline that has been endorsed by the Health New Zealand - Te Whatu Ora Hospital in the specific settings where there is a greater than 10% risk of invasive fungal infection (IFI).

#### **TERBINAFINE**

* Tab 250 mg8.97	84	✓ Deolate
VORICONAZOLE - Special Authority see SA2384 below - Retail pharmacy		
Tab 50 mg71.00	56	✓ Vttack
Tab 200 mg263.00	56	✓ Vttack
Powder for oral suspension 40 mg per ml - Wastage		
claimable1,523.22	70 ml	✓ Vfend

#### ⇒SA2384 Special Authority for Subsidy

Initial application — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised: and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
  - 3.1 Patient has proven or probable invasive aspergillus infection; or

Subsidy (Manufacturer's Pric	ce) Su	Fully bsidised	Brand or Generic	
\$	Per	1	Manufacturer	

continued...

- 3.2 Patient has possible invasive aspergillus infection; or
- 3.3 Patient has fluconazole resistant candidiasis; or
- 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

**Renewal — (invasive fungal infection)** only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
  - 3.1 Patient continues to require treatment for proven or probable invasive aspergillus infection; or
  - 3.2 Patient continues to require treatment for possible invasive aspergillus infection; or
  - 3.3 Patient has fluconazole resistant candidiasis; or
  - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Initial application — (Invasive fungal infection prophylaxis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient is at risk of invasive fungal infection; and
- 2 Either:
  - 2.1 Voriconazole is prescribed by, or recommended by a haematologist, transplant physician, infectious disease specialist, paediatric haematologist or paediatric oncologist; or
  - 2.2 Prescribing voriconazole is in accordance with a protocol or guideline that has been endorsed by the Health New Zealand - Te Whatu Ora Hospital in the specific settings where there is a greater than 10% risk of invasive fungal infection (IFI).

Renewal — (Invasive fungal infection prophylaxis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient is at risk of invasive fungal infection; and
- 2 Either:
  - 2.1 Voriconazole is prescribed by, or recommended by a haematologist, transplant physician, infectious disease specialist, paediatric haematologist or paediatric oncologist; or
  - 2.2 Prescribing voriconazole is in accordance with a protocol or guideline that has been endorsed by the Health New Zealand - Te Whatu Ora Hospital in the specific settings where there is a greater than 10% risk of invasive fungal infection (IFI).

#### **Antimalarials**

## ⇒SA1684 Special Authority for Subsidy

**Initial application** only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 The patient has vivax or ovale malaria; and
- 2 Primaquine is to be given for a maximum of 21 days.

Renewal only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting

INFECTIONS - AGENTS FOR SYSTEMIC USE						
	Subsidy (Manufacturer's Price \$	e) Per	Fully Subsidised	I Generic		
continued the following criteria: Both:						
<ul><li>1 The patient has relapsed vivax or ovale malaria; and</li><li>2 Primaquine is to be given for a maximum of 21 days.</li></ul>						
Antitrichomonal Agents						
METRONIDAZOLE						
Tab 200 mg — Up to 30 tab available on a PSO	25.86	250		Metronidamed		
Tab 400 mg - Up to 15 tab available on a PSO		21		Metronidamed		
Oral liq benzoate 200 mg per 5 ml		100 m		Flagyl-S Flagyl		
Suppos 500 mg	24.46	10	•	гіадуі		
ORNIDAZOLE	22.52	40				
Tab 500 mg	36.52	10	•	Arrow-Ornidazole		
Antituberculotics and Antileprotics						
Note: There is no co-payment charge for all pharmaceuticals list immigration status.	ed in the Antituberco	ulotics	and Antile	protics group regardless of		
BEDAQUILINE – Special Authority see SA2244 below – Retail p No patient co-payment payable	harmacy					
Tab 100mg	3,084.51	24 OF	•	Sirturo		
■ SA2244 Special Authority for Subsidy Initial application — (multi-drug resistant tuberculosis) from applications meeting the following criteria:  Both:	any relevant practiti	ioner.	Approvals	valid for 6 months for		
<ol> <li>The person has multi-drug resistant tuberculosis (MDR-TE</li> <li>Ministry of Health's Tuberculosis Clinical Network has revior the treatment regimen.</li> </ol>		case a	ind recom	mends bedaquiline as part		
CLOFAZIMINE - Retail pharmacy-Specialist						
a) No patient co-payment payable     b) Prescriptions must be written by, or on the recommendati dermatologist.	on of, an infectious	diseas	e physicia	n, clinical microbiologist or		
* Cap 50 mg	442.00	100	1	Lamprene S29		
CYCLOSERINE - Retail pharmacy-Specialist						

D)	Prescriptions
	dermatologist
_	I. O

a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or

b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or

dematologist			
Tab 25 mg	268.50	100	Dapsone
Tab 100 mg	329.50	100	✓ Dapsone

Cap 250 mg......344.00

a) No patient co-payment payable

respiratory physician.

DAPSONE - Retail pharmacy-Specialist

60

✓ Cyclorin S29

INFECTIONS - AGENTS FOR STSTEMIC USE							
	Subsidy (Manufacturer's Price \$	) S Per	Fully ubsidised				
ETHAMBUTOL HYDROCHLORIDE - Retail pharmacy-Specia	alist						
a) No patient co-payment payable							
<li>b) Prescriptions must be written by, or on the recommend respiratory physician</li>		disease	physicia	n, clinical microbiologist or			
Tab 100 mg	85.73	100	✓	EMB Fatol S29			
Tab 400 mg	49.34	56	1	Myambutol \$29			
ISONIAZID - Retail pharmacy-Specialist							
<ul> <li>a) No patient co-payment payable</li> </ul>							
b) Prescriptions must be written by, or on the recommend		edicine p	hysician	, paediatrician, clinical			
microbiologist, dermatologist or public health physician		400	,	In a selected Town 200			
* Tab 100 mg	94.50 327.41	100		Isoniazid Teva S29 Noumed Isoniazid			
JOONIATID WITH DIEAMDION - Date Tale announce Occasions	327.41		•	Noullieu Isomaziu			
ISONIAZID WITH RIFAMPICIN – Retail pharmacy-Specialist							
<ul> <li>a) No patient co-payment payable</li> <li>b) Prescriptions must be written by, or on the recommend microbiologist, dermatologist or public health physician</li> </ul>		edicine p	hysician	, paediatrician, clinical			
* Tab 100 mg with rifampicin 150 mg		100	✓	Rifinah			
* Tab 150 mg with rifampicin 300 mg	179.13	100	1	Rifinah			
* Cap 100 mg with rifampicin 150 mg	199.00	100	•	Rifamazid S29			
LINEZOLID - Special Authority see SA2234 below - Retail ph	armacy						
No patient co-payment payable	•						
Tab 600 mg		10		Zyvox			
Oral liq 20 mg per ml	1,879.00	150 ml	/	Zyvox			
<b>⇒SA2234</b> Special Authority for Subsidy							
Initial application — (multi-drug resistant tuberculosis) from applications meeting the following criteria:  Both:	m any relevant practiti	oner. A	pprovals	valid for 18 months for			
1 The person has multi-drug resistant tuberculosis (MDR-	-TB): and						
Ministry of Health's Tuberculosis Clinical Network has r the treatment regimen.		case an	d recom	mends linezolid as part of			
PARA-AMINO SALICYLIC ACID – Retail pharmacy-Specialist							
a) No patient co-payment payable							
<ul> <li>Prescriptions must be written by, or on the recommend respiratory physician</li> </ul>		disease	specialis	et, clinical microbiologist or			
Grans for oral liq 4 g sachet	280.00	30	•	Paser S29			
PROTIONAMIDE - Retail pharmacy-Specialist							
a) No patient co-payment payable     b) Prescriptions must be written by, or on the recommend	ation of, an infectious	disease	specialis	st. clinical microbiologist or			
respiratory physician				., <b>.</b>			
Tab 250 mg	305.00	100	1	Peteha \$29			
PYRAZINAMIDE - Retail pharmacy-Specialist							
a) No patient co-payment payable							
b) Prescriptions must be written by, or on the recommend	ation of, an infectious	disease	physicia	n, clinical microbiologist or			
respiratory physician				-			
* Tab 500 mg	64.95	100	1	AFT-Pyrazinamide			

	INFECTIONS - AGENTS FOR SYSTEMIC USI	<b>=</b>								
		Subsidy (Manufacturer's Price \$	) Subsi	Fully idised	Brand or Generic Manufacturer					
RIF	ABUTIN - Retail pharmacy-Specialist									
	a) No patient co-payment payable     b) Prescriptions must be written by, or on the recommendation	on of, an infectious	disease phy	sician,	respiratory physician or					
*	gastroenterologist Cap 150 mg	353.71	30	✓ N	lycobutin					
	FAMPICIN – Subsidy by endorsement				,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,					
	a) No patient co-payment payable									
	b) For confirmed recurrent Staphylococcus aureus infection in combination with other effective anti-staphylococcal antimicrobial based on susceptibilities and the prescription is endorsed accordingly; can be waived by endorsement - Retail pharmacy - Specialist. Specialist must be an internal medicine physician, clinical microbiologist, dermatologist, paediatrician, or public health physician.									
	Cap 150 mg		100	_	lifadin					
*	Cap 300 mg	122.06	100	_	<u>lifadin</u> lifadin Sanofi					
*	Oral liq 100 mg per 5 ml	12.60	60 ml		lifadin					
A	ntivirals									
For	eye preparations refer to Eye Preparations, Anti-Infective Pre	parations, page 277	,							
Н	epatitis B Treatment									
ΕN	TECAVIR									
*	Tab 0.5 mg	12.04	30	<b>√</b> <u>E</u>	ntecavir (Rex)					
LAI	MIVUDINE – Special Authority see SA1685 below – Retail pha									
	Tab 100 mg		28		etlam					
	Oral liq 5 mg per ml	270.00 2	40 ml OP	• 2	effix					
	SA1685 Special Authority for Subsidy ial application only from a relevant specialist or medical pract	titioner on the recon	nmendation	of a re	levant specialist					
App	provals valid for 1 year where used for the treatment or preven	tion of hepatitis B.	mioridation	01 4 10	iovani oposianou					
	newal from any relevant practitioner. Approvals valid for 2 year		he treatmen	t or pre	evention of hepatitis B.					
TE	NOFOVIR DISOPROXIL									
	Tenofovir disoproxil prescribed under endorsement for the treantiretrovirals for the purposes of Special Authority SA2139.,		cluded in the	count	of up to 4 subsidised					
*	Tab 245 mg (300 mg as a maleate)		30	<b>✓</b> T	enofovir Disoproxil					
	3 (111 )				Viatris					
*	Tab 245 mg (300 mg as a fumarate)	13.80	30	✓ R	licovir S29					
Н	erpesvirus Treatments									
	ICLOVIR									
*	Tab dispersible 200 mg Lovir to be Principal Supply on 1 February 2026	2.05	25	<b>∕</b> L						
*	Tab dispersible 400 mg	7.55	56	<b>√</b> L	ovir					
*	Lovir to be Principal Supply on 1 February 2026  Tab dispersible 800 mg	7 43	35	<b>√</b> L	ovir					
-,,	Lovir to be Principal Supply on 1 February 2026		00		· · · ·					
VA	LACICLOVIR									
	Tab 500 mg		30		aclovir					
	Tab 1,000 mg	17.78	30	<b>✓</b> <u>∨</u>	<u>aclovir</u>					

	Subsidy (Manufacturer's Price)	S	Fully ubsidised	Brand or Generic
	\$	Per	1	Manufacturer
VALGANCICLOVIR - Special Authority see SA2514 below - Re	etail pharmacy			
Tab 450 mg	140.89	60		<u>llganciclovir</u> Viatris

### ⇒SA2514 Special Authority for Subsidy

**Initial application** — **(transplant cytomegalovirus prophylaxis)** only from a relevant specialist. Approvals valid for 3 months where the patient has undergone a solid organ transplant and requires valganciclovir for CMV prophylaxis.

Renewal — (transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 Patient has undergone a solid organ transplant and received anti-thymocyte globulin and requires valganciclovir therapy for CMV prophylaxis; and
  - 1.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following anti-thymocyte globulin; or
- 2 Both:
  - 2.1 Patient has received pulse methylprednisolone for acute rejection and requires further valganciclovir therapy for CMV prophylaxis; and
  - 2.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following pulse methylprednisolone.

Initial application — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has undergone a solid organ transplant and received valganciclovir under Special Authority more than 2 years ago (27 months); and
- 2 Patient has received anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Renewal — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months where the patient has received a further course of anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Initial application — (Lung transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone a lung transplant; and
- 2 Either:
  - 2.1 The donor was cytomegalovirus positive and the patient is cytomegalovirus negative; or
  - 2.2 The recipient is cytomegalovirus positive; and
- 3 Patient has a high risk of CMV disease.

Renewal — (Lung transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone a lung re-transplant; and
- 2 Either:
  - 2.1 The donor was cytomegalovirus positive and the patient is cytomegalovirus negative; or
  - 2.2 The recipient is cytomegalovirus positive; and
- 3 Patient has a high risk of CMV disease.

Initial application — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
  - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or

(Mar	Subsidy	Fully		Brand or
	nufacturer's Price)	Subsidised		Generic
<u> </u>	\$	Per	1	Manufacturer

continued...

- 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
- 2.3 Patient has cytomegalovirus retinitis.

Renewal — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient is immunocompromised: and
- 2 Any of the following:
  - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
  - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
  - 2.3 Patient has cytomegalovirus retinitis.

Note: for the purpose of this Special Authority "immunocompromised" includes transplant recipients, patients with immunosuppressive diseases (e.g. HIV) or those receiving immunosuppressive treatment for other conditions.

## **Hepatitis C Treatment**

#### GLECAPREVIR WITH PIBRENTASVIR - [Xpharm]

Note the supply of treatment is via Pharmac's approved direct distribution supply. Further details can be found on Pharmac's website <a href="https://pharmac.govt.nz/maviret">https://pharmac.govt.nz/maviret</a>

Tab 100 mg with pibrentasvir 40 mg ......24,750.00 84 OP ✓ Maviret

LEDIPASVIR WITH SOFOSBUVIR - [Xpharm] - Special Authority see SA1605 below

No patient co-payment payable

Tab 90 mg with sofosbuvir 400 mg......24,363.46 28 **✓ Harvoni** 

#### ⇒SA1605 Special Authority for Subsidy

Special Authority approved by the Hepatitis C Treatment Panel (HepCTP)

Notes: By application to the Hepatitis C Treatment Panel (HepCTP).

Applications will be considered by HepCTP and approved subject to confirmation of eligibility.

Application details may be obtained from Pharmac's website <a href="http://www.pharmac.govt.nz/harvoni">http://www.pharmac.govt.nz/harvoni</a> or:

The Coordinator, Hepatitis C Treatment Panel

Pharmac, PO Box 10-254, WELLINGTON Tel: (04) 460 4990,

Email: hepcpanel@pharmac.govt.nz

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

### **HIV Prophylaxis and Treatment**

EMTRICITABINE WITH TENOFOVIR DISOPROXIL - Subsidy by endorsement; can be waived by Special Authority see SA2520 below

- a) Funding for emtricitabine with tenofovir disoproxil for use as PrEP or PEP, should be applied using Special Authority SA2520.
- b) Endorsement for treatment of conditions approved via Special Authority SA2139 (antiretrovirals for confirmed HIV, prevention of maternal transmission, post-exposure prophylaxis following exposure to HIV (when co-prescribed with other antiretrovirals) and percutaneous exposure): Prescription is deemed to be endorsed if emtricitabine with tenofovir disoproxil is co-prescribed with another antiretroviral subsidised under Special Authority SA2139 and the prescription is annotated accordingly by the Pharmacist or endorsed by the prescriber.
  - Note: Emtricitabine with tenofovir disoproxil prescribed under endorsement, for treatment of conditions approved via Special Authority SA2139 (antiretrovirals for confirmed HIV, prevention of maternal transmission, post-exposure prophylaxis following exposure to HIV and percutaneous exposure), is included in the count of up to 4 subsidised antiretrovirals, and counts as two antiretroviral medications, for the purposes of Special Authority SA2139, page 110 There is an approval process to become a named specialist to prescribe antiretroviral therapy in New Zealand. Further information is available on the Pharmac website.

*	fumarate)13.45	30	✓ Tenofovir Disoproxil Emtricitabine Mylan ©29
*	Tab 200 mg with tenofovir disoproxil 245 mg (300 mg as a maleate)	30	✓ <u>Tenofovir Disoproxil</u> Emtricitabine Viatr

### ⇒SA2520 Special Authority for Subsidy

**Initial application — (Pre-exposure prophylaxis)** from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria:

Both:

- Patient has tested HIV negative, does not have signs or symptoms of acute HIV infection and has been assessed for HIV seroconversion: and
- 2 The Practitioner considers the patient is at elevated risk of HIV exposure and use of PrEP is clinically appropriate.

Notes: Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines:

https://ashm.org.au/HIV/PrEP/

Renewal — (Pre-exposure prophylaxis) from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 Patient has tested HIV negative, does not have signs or symptoms of acute HIV infection and has been assessed for HIV seroconversion; and
- 2 The Practitioner considers the patient is at elevated risk of HIV exposure and use of PrEP is clinically appropriate.

Notes: Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines:

https://ashm.org.au/HIV/PrEP/

Initial application — (post-exposure prophylaxis following exposure to HIV) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
  - 2.1 Patient has had condomless anal intercourse or receptive vaginal intercourse with a known HIV positive person

Subsidy		Fully	Brand or	
(Manufacturer's Price	,	Subsidised	Generic	
\$	Per		Manufacturer	

continued...

with an unknown or detectable viral load greater than 200 copies per ml; or

- 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
- 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is appropriate; or
- 2.4 Patient has had condomless anal intercourse with a person from a high HIV prevalence country or risk group whose HIV status is unknown.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines for PEP (https://www.ashm.org.au/hiv/hiv-management/pep/).

Renewal — (second or subsequent post-exposure prophylaxis) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
  - 2.1 Patient has had condomless anal intercourse or receptive vaginal intercourse with a known HIV positive person with an unknown or detectable viral load greater than 200 copies per ml: or
  - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
  - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is appropriate; or
  - 2.4 Patient has had condomless anal intercourse with a person from a high HIV prevalence country or risk group whose HIV status is unknown.

### **COVID-19 Treatments**

NIRMATRELVIR WITH RITONAVIR - PCT - Subsidy by endorsement

Subsidised for patients meeting access criteria for oral COVID-19 antiviral treatments (as on <a href="Pharmac's website">Pharmac's website</a>) and where the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed when supplying by Direct Provision under the provisions in Part I of Section A of the Pharmaceutical Schedule.

REMDESIVIR - PCT only

### **Antiretrovirals**

### ⇒SA2139 Special Authority for Subsidy

Initial application — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the patient has confirmed HIV infection.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

**Renewal — (Confirmed HIV)** only from a named specialist. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient is benefiting from treatment.

Subsidy	Full	y Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per 🗸	Manufacturer	

continued...

Initial application — (Prevention of maternal transmission) only from a named specialist. Approvals valid for 1 year for applications meeting the following criteria:

Either:

- 1 Prevention of maternal foetal transmission: or
- 2 Treatment of the newborn for up to eight weeks.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Some antiretrovirals are unapproved or contraindicated for this indication. Practitioners prescribing these medications should exercise their own skill, judgement, expertise and discretion, and make their own prescribing decisions with respect to the use of a Pharmaceutical for an indication for which it is not approved or contraindicated.

Initial application — (post-exposure prophylaxis following exposure to HIV) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
  - 2.1 Patient has had condomless anal intercourse or receptive vaginal intercourse with a known HIV positive person with an unknown or detectable viral load greater than 200 copies per ml; or
  - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
  - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required; or
  - 2.4 Patient has had condomless anal intercourse with a person from a high HIV prevalence country or risk group whose HIV status is unknown.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines for PEP (https://www.ashm.org.au/hiv/hiv-management/pep/).

Renewal — (second or subsequent post-exposure prophylaxis) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
  - 2.1 Patient has had condomless anal intercourse or receptive vaginal intercourse with a known HIV positive person with an unknown or detectable viral load greater than 200 copies per ml; or
  - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
  - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required; or
  - 2.4 Patient has had condomless anal intercourse with a person from a high HIV prevalence country or risk group whose HIV status is unknown.

**Initial application — (Percutaneous exposure)** only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

	Subsidy (Manufacturer's F	Price) Subs	Fully idised	Brand or Generic
	\$	Per	1	Manufacturer
continued Renewal — (Second or subsequent percutaneous exposi where the patient has percutaneous exposure to blood known		med specialist.	Approv	vals valid for 6 weeks
Non-nucleosides Reverse Transcriptase Inhi	bitors			
FAVIRENZ – Special Authority see SA2139 on page 110 – Note: No new patients to be initiated on efavirenz.				
Tab 600 mg	65.38	30	<b>√</b> E	favirenz Milpharm S29
Efavirenz Milpharm 👀 Tab 600 mg to be delisted 1 Novel	mber 2026)			
TRAVIRINE - Special Authority see SA2139 on page 110 - Tab 200 mg		60	✓ lı	ntelence
NEVIRAPINE - Special Authority see SA2139 on page 110				
Tab 200 mg Oral suspension 10 mg per ml		60 240 ml OP		<u>levirapine Viatris</u> ′iramune
Oral Supportation To mig per minimum.	200.00	240 1111 01	•	Suspension
Nucleosides Reverse Transcriptase Inhibitor	'S			
ABACAVIR SULPHATE - Special Authority see SA2139 on	page 110 – Retail ph	narmacy		
Tab 300 mg	180.00	60	✓ Z	liagen
ABACAVIR SULPHATE WITH LAMIVUDINE - Special Auth- Note: abacavir with lamivudine (combination tablets) cou	ority see SA2139 on	page 110 – Re	tail pha	armacy
BACAVIR SULPHATE WITH LAMIVUDINE - Special Auth	ority see SA2139 on unts as two anti-retro	page 110 – Re	etail pha	armacy
BACAVIR SULPHATE WITH LAMIVUDINE – Special Auth- Note: abacavir with lamivudine (combination tablets) cou anti-retroviral Special Authority.	ority see SA2139 on unts as two anti-retro	page 110 - Repoviral medication	etail pha	armacy he purposes of the  abacavir/ Lamivudine
BACAVIR SULPHATE WITH LAMIVUDINE — Special Auth- Note: abacavir with lamivudine (combination tablets) cou- anti-retroviral Special Authority. Tab 600 mg with lamivudine 300 mg  Abacavir/Lamivudine Viatris to be Principal Supply of FAVIRENZ WITH EMTRICITABINE AND TENOFOVIR DIS	ority see SA2139 on unts as two anti-retro	page 110 – Reported in the page 110 – Reported i	etail pha ns for t	he purposes of the Abacavir/ Lamivudine Viatris
ABACAVIR SULPHATE WITH LAMIVUDINE — Special Authonories abacavir with lamivudine (combination tablets) counti-retroviral Special Authority.  Tab 600 mg with lamivudine 300 mg	ority see SA2139 on unts as two anti-retro35.00 In 1 February 2026 OPROXIL – Special kil counts as three ar	page 110 – Repviral medication 30	etail phans for the A	armacy he purposes of the  Abacavir/ Lamivudine Viatris  9 on page 110 – Retail
ABACAVIR SULPHATE WITH LAMIVUDINE — Special Authority. Note: abacavir with lamivudine (combination tablets) coulont-retroviral Special Authority. Tab 600 mg with lamivudine 300 mg	ority see SA2139 on unts as two anti-retro35.00  In 1 February 2026 OPROXIL – Special counts as three are proxil	page 110 – Reported in the page 130 and a second in the page 130 and a sec	etail phans for the state of th	armacy the purposes of the  Abacavir/ Lamivudine Viatris  9 on page 110 – Retail this for the purposes of the
BACAVIR SULPHATE WITH LAMIVUDINE — Special Auth- Note: abacavir with lamivudine (combination tablets) cot anti-retroviral Special Authority. Tab 600 mg with lamivudine 300 mg  Abacavir/Lamivudine Viatris to be Principal Supply o EFAVIRENZ WITH EMTRICITABINE AND TENOFOVIR DIS harmacy Note: Efavirenz with emtricitabine and tenofovir disopror anti-retroviral Special Authority Tab 600 mg with emtricitabine 200 mg and tenofovir diso 245 mg (300 mg as a fumarate)	ority see SA2139 on unts as two anti-retro35.00  In 1 February 2026 OPROXIL – Special kil counts as three are opproxil106.88	page 110 – Repviral medication 30	etail phans for the state of th	armacy he purposes of the  Abacavir/ Lamivudine Viatris  9 on page 110 – Retail
BACAVIR SULPHATE WITH LAMIVUDINE — Special Auth- Note: abacavir with lamivudine (combination tablets) cot anti-retroviral Special Authority. Tab 600 mg with lamivudine 300 mg  Abacavir/Lamivudine Viatris to be Principal Supply o EFAVIRENZ WITH EMTRICITABINE AND TENOFOVIR DIS harmacy Note: Efavirenz with emtricitabine and tenofovir disopror anti-retroviral Special Authority Tab 600 mg with emtricitabine 200 mg and tenofovir diso	ority see SA2139 on unts as two anti-retro35.00 In 1 February 2026 OPROXIL – Special xil counts as three are opproxil106.88 oproxil	page 110 – Reported in the page 130 and a second in the page 130 and a sec	etail phans for the A	armacy the purposes of the  Abacavir/ Lamivudine Viatris  9 on page 110 – Retail this for the purposes of the
BACAVIR SULPHATE WITH LAMIVUDINE — Special Auth- Note: abacavir with lamivudine (combination tablets) cot anti-retroviral Special Authority. Tab 600 mg with lamivudine 300 mg  Abacavir/Lamivudine Viatris to be Principal Supply o FAVIRENZ WITH EMTRICITABINE AND TENOFOVIR DIS harmacy Note: Efavirenz with emtricitabine and tenofovir disopror anti-retroviral Special Authority Tab 600 mg with emtricitabine 200 mg and tenofovir diso 245 mg (300 mg as a fumarate)	ority see SA2139 on unts as two anti-retro35.00 In 1 February 2026 OPROXIL — Special xil counts as three are proxil106.88 proxil	page 110 – Repviral medication 30  I Authority see 3 hti-retroviral me 30 30	etail phans for the A	armacy the purposes of the  Abacavir/ Lamivudine Viatris  9 on page 110 – Retail ans for the purposes of the  EEVIR \$29
BACAVIR SULPHATE WITH LAMIVUDINE — Special Auth- Note: abacavir with lamivudine (combination tablets) cot anti-retroviral Special Authority.  Tab 600 mg with lamivudine 300 mg  Abacavir/Lamivudine Viatris to be Principal Supply o EFAVIRENZ WITH EMTRICITABINE AND TENOFOVIR DIS harmacy Note: Efavirenz with emtricitabine and tenofovir disoprov anti-retroviral Special Authority Tab 600 mg with emtricitabine 200 mg and tenofovir diso 245 mg (300 mg as a fumarate) Tab 600 mg with emtricitabine 200 mg and tenofovir diso 245 mg (300 mg as a maleate)	ority see SA2139 on unts as two anti-retro	page 110 – Repviral medication 30  I Authority see 3 hti-retroviral me 30 30	etail phans for the ASA2139 dication	armacy the purposes of the  Abacavir/ Lamivudine Viatris  9 on page 110 – Retail ans for the purposes of the  EEVIR \$29
BACAVIR SULPHATE WITH LAMIVUDINE — Special Auth- Note: abacavir with lamivudine (combination tablets) cot anti-retroviral Special Authority.  Tab 600 mg with lamivudine 300 mg  Abacavir/Lamivudine Viatris to be Principal Supply o FAVIRENZ WITH EMTRICITABINE AND TENOFOVIR DIS harmacy Note: Efavirenz with emtricitabine and tenofovir disoprov anti-retroviral Special Authority Tab 600 mg with emtricitabine 200 mg and tenofovir diso 245 mg (300 mg as a fumarate)  Tab 600 mg with emtricitabine 200 mg and tenofovir diso 245 mg (300 mg as a maleate)  MTRICITABINE — Special Authority see SA2139 on page 1 Cap 200 mg	ority see SA2139 on unts as two anti-retro	page 110 – Repviral medication 30  I Authority see 3  nti-retroviral medication 30  30  30  30  30	etail phans for the second of	armacy he purposes of the  Abacavir/ Lamivudine Viatris  9 on page 110 – Retail ns for the purposes of the  EEVIR \$29  Viatris
BACAVIR SULPHATE WITH LAMIVUDINE — Special Auth- Note: abacavir with lamivudine (combination tablets) cot anti-retroviral Special Authority. Tab 600 mg with lamivudine 300 mg  Abacavir/Lamivudine Viatris to be Principal Supply o EFAVIRENZ WITH EMTRICITABINE AND TENOFOVIR DIS harmacy Note: Efavirenz with emtricitabine and tenofovir disoprov anti-retroviral Special Authority Tab 600 mg with emtricitabine 200 mg and tenofovir diso 245 mg (300 mg as a fumarate) Tab 600 mg with emtricitabine 200 mg and tenofovir diso 245 mg (300 mg as a maleate)  EMTRICITABINE — Special Authority see SA2139 on page 1 Cap 200 mg	ority see SA2139 on unts as two anti-retro	page 110 – Reported in the page 110 – Reported i	etail phanns for the state of t	armacy the purposes of the  Abacavir/ Lamivudine Viatris  9 on page 110 – Retail ans for the purposes of the  EEVIR \$29  Viatris  Emtriva  amivudine Viatris
BACAVIR SULPHATE WITH LAMIVUDINE — Special Auth- Note: abacavir with lamivudine (combination tablets) cot anti-retroviral Special Authority.  Tab 600 mg with lamivudine 300 mg	ority see SA2139 on unts as two anti-retro	page 110 – Reported in the page 110 – Reported i	etail phans for the second of	armacy the purposes of the  Abacavir/ Lamivudine Viatris  9 on page 110 – Retail ans for the purposes of the  EEVIR \$29  Viatris  Emtriva  amivudine Viatris
BACAVIR SULPHATE WITH LAMIVUDINE — Special Auth- Note: abacavir with lamivudine (combination tablets) cot anti-retroviral Special Authority. Tab 600 mg with lamivudine 300 mg  Abacavir/Lamivudine Viatris to be Principal Supply o EFAVIRENZ WITH EMTRICITABINE AND TENOFOVIR DIS harmacy Note: Efavirenz with emtricitabine and tenofovir disoprov anti-retroviral Special Authority Tab 600 mg with emtricitabine 200 mg and tenofovir diso 245 mg (300 mg as a fumarate) Tab 600 mg with emtricitabine 200 mg and tenofovir diso 245 mg (300 mg as a maleate)  EMTRICITABINE — Special Authority see SA2139 on page 1 Cap 200 mg	ority see SA2139 on unts as two anti-retro	page 110 – Reported in the page 110 – Reported i	etail phans for the state of th	armacy he purposes of the  Abacavir/ Lamivudine Viatris  9 on page 110 – Retail ns for the purposes of EEVIR 529  Viatris  Emtriva  amivudine Viatris

the anti-retroviral Special Authority.

Tab 300 mg with lamivudine 150 mg......92.40

✓ Lamivudine/

Zidovudine Viatris

60

	Subsidy (Manufacturer's Price) \$	S Per	Fully Subsidised •	
Protease Inhibitors				
ATAZANAVIR SULPHATE – Special Authority see SA2139 on pa Cap 150 mg	102.50	rmacy 60		Atazanavir Viatris
Cap 200 mgAtazanavir Viatris to be Principal Supply on 1 February 20	152.30	60	✓	Atazanavir Viatris
DARUNAVIR — Special Authority see SA2139 on page 110 — Reta Tab 400 mg	150.00	60 60		Darunavir Viatris Darunavir Viatris
LOPINAVIR WITH RITONAVIR – Special Authority see SA2139 o Tab 200 mg with ritonavir 50 mg		pharm 120		Lopinavir/Ritonavir Mylan
RITONAVIR – Special Authority see SA2139 on page 110 – Retai Tab 100 mg		30	•	Norvir
Strand Transfer Inhibitors				
DOLUTEGRAVIR - Special Authority see SA2139 on page 110 - Tab 50 mg		30	1	Tivicay
DOLUTEGRAVIR WITH LAMIVUDINE - Special Authority see SA Tab 50 mg with lamivudine 300 mg		- Retai 30		cy <b>Dovato</b>
RALTEGRAVIR POTASSIUM – Special Authority see SA2139 on Tab 400 mg	1,090.00	harma 60 60	′ ✓	Isentress Isentress HD
Immune Modulators				

#### Immune Modulators

PEGYLATED INTERFERON ALFA-2A - Special Authority see SA2034 below - Retail pharmacy

Note: Pharmac will consider funding ribavirin for the small group of patients who have a clinical need for ribavirin and meet Special Authority criteria. Please contact the Hepatitis C Coordinator at Pharmac on 0800-023-588 option 4.

Inj 135 mcg prefilled syringe	887.35	1	✓ Pegasys (S29) S29
Inj 180 mcg prefilled syringe	748.50	4	✓ Pegasys
	1,355.71		✓ Pegasys S29 S29

### ⇒SA2034 Special Authority for Subsidy

Initial application — (chronic hepatitis C - genotype 1, 4, 5 or 6 infection or co-infection with HIV or genotype 2 or 3 post liver transplant) from any specialist. Approvals valid for 18 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 Patient has chronic hepatitis C, genotype 1, 4, 5 or 6 infection; or
  - 1.2 Patient has chronic hepatitis C and is co-infected with HIV; or
  - 1.3 Patient has chronic hepatitis C genotype 2 or 3 and has received a liver transplant; and
- 2 Maximum of 48 weeks therapy.

Renewal — (Chronic hepatitis C - genotype 1 infection) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

1 Patient has chronic hepatitis C, genotype 1; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Fither
  - 3.1 Patient has responder relapsed; or
  - 3.2 Patient was a partial responder; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application — (Chronic Hepatitis C - genotype 1 infection treatment more than 4 years prior) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Any of the following:
  - 3.1 Patient has responder relapsed; or
  - 3.2 Patient was a partial responder; or
  - 3.3 Patient received interferon treatment prior to 2004; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application — (chronic hepatitis C - genotype 2 or 3 infection without co-infection with HIV) from any specialist.

Approvals valid for 12 months for applications meeting the following criteria: Roth:

- 1 Patient has chronic hepatitis C, genotype 2 or 3 infection; and
- 2 Maximum of 6 months therapy.

Initial application — (Hepatitis B) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Hepatitis B infection (HBsAg positive for more than 6 months); and
- 2 Patient is Hepatitis B treatment-naive; and
- 3 ALT > 2 times Upper Limit of Normal; and
- 4 HBV DNA < 10 log10 IU/ml; and
- 5 Either:
  - 5.1 HBeAg positive; or
  - 5.2 serum HBV DNA greater than or equal to 2,000 units/ml and significant fibrosis (Metavir Stage F2 or greater or moderate fibrosis); and
- 6 Compensated liver disease: and
- 7 No continuing alcohol abuse or intravenous drug use; and
- 8 Not co-infected with HCV, HIV or HDV; and
- 9 Neither ALT nor AST > 10 times upper limit of normal; and
- 10 No history of hypersensitivity or contraindications to pegylated interferon; and
- 11 Maximum of 48 weeks therapy.

Initial application — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 Patient has a cutaneous T cell lymphoma\*; or
- 2 All of the following:
  - 2.1 Patient has a myeloproliferative disorder\*: and
  - 2.2 Patient is intolerant of hydroxyurea; and

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

- 2.3 Treatment with an agrelide and busulfan is not clinically appropriate; or
- 3 Both:
  - 3.1 Patient has a myeloproliferative disorder; and
  - 3.2 Patient is pregnant, planning pregnancy or lactating.

Renewal — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment; and
- 3 Fither
  - 3.1 Patient has a cutaneous T cell lymphoma\*: or
  - 3.2 Both:
    - 3.2.1 Patient has a myeloproliferative disorder\*; and
    - 3.2.2 Fither:
      - 3.2.2.1 Remains intolerant of hydroxyurea and treatment with anagrelide and busulfan remains clinically inappropriate; or
      - 3.2.2.2 Patient is pregnant, planning pregnancy or lactating.

Note: Indications marked with \* are unapproved indications.

**Initial application — (post-allogenic bone marrow transplant)** from any relevant practitioner. Approvals valid for 3 months where patient has received an allogeneic bone marrow transplant\* and has evidence of disease relapse.

Renewal — (post-allogenic bone marrow transplant) from any relevant practitioner. Approvals valid for 3 months where patient is responding and ongoing treatment remains appropriate.

Note: Indications marked with \* are unapproved indications.

### **Urinary Tract Infections**

### ⇒SA2406 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 2 months for applications meeting the following criteria:

- 1 Patient has an acute, symptomatic, bacteriologically-proven uncomplicated urinary tract infection (UTI)/cystitis with Escherichia Coli; and
- 2 Fither:
  - 2.1 Microbiological testing confirms the pathogen is resistant to all of: trimethoprim, nitrofurantoin, amoxicillin, cefaclor, cefalexin, amoxicillin with clavulanic acid, and norfloxacin; or
  - 2.2 The patient has a contraindication or intolerance to all of: trimethoprim, nitrofurantoin, amoxicillin, cefaclor, cefalexin, amoxicillin with clavulanic acid, and norfloxacin that the pathogen is susceptible to.

Renewal from any relevant practitioner. Approvals valid for 2 months for applications meeting the following criteria: Both:

- 1 Patient has an acute, symptomatic, bacteriologically-proven uncomplicated urinary tract infection (UTI)/cystitis with Escherichia Coli; and
- 2 Fither:
  - 2.1 Microbiological testing confirms the pathogen is resistant to all of: trimethoprim, nitrofurantoin, amoxicillin, cefaclor, cefalexin, amoxicillin with clavulanic acid, and norfloxacin; or
  - 2.2 The patient has a contraindication or intolerance to all of: trimethoprim, nitrofurantoin, amoxicillin, cefaclor, cefalexin, amoxicillin with clavulanic acid, and norfloxacin that the pathogen is susceptible to.

METHENAMINE (HEXAMINE) HIPPURATE

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
NITROFURANTOIN				
* Tab 50 mg - Up to 30 tab available on a PSO	22.20	100	✓	Nifuran
* Tab 100 mg	37.50	100	✓	Nifuran
* Cap modified-release 100 mg - Up to 15 cap available on a				
PSO	81.20	100	✓	Macrobid
NORFLOXACIN				
Tab 400 mg - Subsidy by endorsement	245.00	100	✓	Arrow-Norfloxacin
Only if prescribed for a patient with an uncomplicated uri with proven resistance to first line agents and the prescri				ive to a first line agent or

	Subsidy		Fully Brand or
	(Manufacturer's Price)		Subsidised Generic
	\$	Per	✓ Manufacturer
Anticholinesterases			
Anticholinesterases			
NEOSTIGMINE METILSULFATE			
Inj 2.5 mg per ml, 1 ml ampoule	48 25	10	✓ Max Health
			- max rioutii
PYRIDOSTIGMINE BROMIDE	FO 00	100	/ Maatinan
▲ Tab 60 mg	50.28	100	Mestinon
Non Staraidal Anti Inflammatory Drugo			
Non-Steroidal Anti-Inflammatory Drugs			
DICLOFENAC SODIUM			
* Tab EC 25 mg	2 19	50	✓ Diclofenac Sandoz
* Tab 50 mg dispersible		20	✓ Voltaren D
* Tab EC 50 mg		50	✓ Diclofenac Sandoz
3		100	_
* Tab long-acting 75 mg			✓ <u>Voltaren SR</u>
* Inj 25 mg per ml, 3 ml ampoule – Up to 5 inj available on a F		5	Voltaren
* Suppos 12.5 mg		10	✓ Voltaren
* Suppos 25 mg		10	✓ Voltaren
* Suppos 50 mg - Up to 10 supp available on a PSO		10	✓ Voltaren
* Suppos 100 mg	7.00	10	✓ Voltaren
IBUPROFEN			
* Tab 200 mg	21.40	1,000	✓ Relieve
* Tab long-acting 800 mg		30	✓ Ibuprofen SR BNM
* Oral lig 20 mg per ml		200 ml	
	2.03	200 1111	Lunes
KETOPROFEN			
* Cap long-acting 200 mg	12.07	28	Oruvail SR
(Oruvail SR Cap long-acting 200 mg to be delisted 1 October 202	26)		
MEFENAMIC ACID			
* Cap 250 mg	1 25	50	
	(10.82)	00	Ponstan
	0.50	20	i onstan
		20	Ponstan
	(7.50)		Fonsian
NAPROXEN			
* Tab 250 mg		500	✓ Noflam 250
* Tab 500 mg	34.45	250	✓ Noflam 500
* Tab long-acting 750 mg	10.40	28	✓ Naprosyn SR 750
* Tab long-acting 1 g	11.50	28	✓ Naprosyn SR 1000
TENOXICAM			
* Tab 20 mg	23 EU	100	✓ Tilcotil
•	20.00	100	· HICOUI
Tilcotil to be Principal Supply on 1 February 2026	0.05	1	✓ AFT
* Inj 20 mg vial	9.95	ı	♥ AFI
NCAIDe Other			
NSAIDs Other			
CELECOXIB			
	2 45	60	✓ Celecoxib Pfizer
Cap 100 mg		60	
Oalahusuta ha Drinainal Ouruhu u 4 Esharun 2000	3.60		✓ Celebrex
Celebrex to be Principal Supply on 1 February 2026	0.00		
Cap 200 mg	3.20	30	✓ Celebrex
(0.1			Celecoxib Pfizer
(Celecoxib Pfizer Cap 100 mg to be delisted 1 February 2026)			

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

### **Topical Products for Joint and Muscular Pain**

### **CAPSAICIN**

Crm 0.025% - Special Authority see SA1289 below - Retail pharmacy......9.75 45 g OP ✓ Zo-Rub Osteo

⇒SA1289 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has osteoarthritis that is not responsive to paracetamol and oral non-steroidal anti-inflammatories are contraindicated.

### Antirheumatoid Agents

# Tab 200 mg	100	✓ <u>Ipca-</u> <u>Hydroxychloroquine</u>
LEFLUNOMIDE		
* Tab 10 mg6.00	30	✓ Arava
* Tab 20 mg6.00	30	✓ Arava
PENICILLAMINE		
Tab 125 mg67.23	100	✓ D-Penamine
Tah 250 mg 110 12	100	✓ D-Penamine

### **Drugs Affecting Bone Metabolism**

### Alendronate for Osteoporosis

ALENDRONATE SODIUM	1	
--------------------	---	--

✓ Fosamax ALENDRONATE SODIUM WITH COLECALCIFEROL

✓ Fosamax Plus

#### Other Treatments

DENOSUMAB - Special Authority see SA2441 below - Retail pharmacy

Note: Denosumab ini 60 mg per 1 ml pre-filled syringe is Medsafe approved for use in osteoporosis. Denosumab ini 120 mg per 1.7 ml vial is Medsafe approved for use in hypercalcaemia of malignancy.

✓ Xgeva 1 ✓ Prolia Inj 60 mg per 1 ml prefilled syringe......187.50

### ⇒SA2441 Special Authority for Subsidy

Initial application — (Osteoporosis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 The patient has established osteoporosis; and
- 2 Any of the following:
  - 2.1 History of one significant osteoporotic fracture demonstrated radiologically, with a documented T-Score less than or egual to -2.5, that incorporates BMD measured using dual-energy x-ray absorptiometry (DEXA); or
  - 2.2 History of one significant osteoporotic fracture, demonstrated radiologically, and either the patient is elderly, or

continued...

✓ Zostriy

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic	
\$	Per	•	Manufacturer	

#### continued...

- densitometry scanning cannot be performed because of logistical, technical or pathophysiological reasons; or
- 2.3 History of two significant osteoporotic fractures demonstrated radiologically; or
- 2.4 Documented T-Score less than or equal to -3.0; or
- 2.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm that incorporates BMD measured using DEXA; and
- 3 Any of the following:
  - 3.1 Bisphosphonates are contraindicated because the patient's creatinine clearance or eGFR is less than 35 mL/min; or
  - 3.2 The patient has experienced at least two symptomatic new fractures or a BMD loss greater than 2% per year, after at least 12 months' continuous therapy with a funded antiresorptive agent; or
  - 3.3 Bisphosphonates result in intolerable side effects; or
  - 3.4 Intravenous bisphosphonates cannot be administered due to logistical or technical reasons.

Initial application — (Hypercalcaemia) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has hypercalcaemia of malignancy; and
- 2 Patient has severe renal impairment.

#### PAMIDRONATE DISODIUM

Inj 3 mg per ml, 10 ml vial	32.49	1	✓ Pamisol
Inj 6 mg per ml, 10 ml vial		1	✓ Pamisol
Inj 9 mg per ml, 10 ml vial		1	✓ Pamisol
RALOXIFENE HYDROCHLORIDE - Special Authority see S	SA1779 below – Retail p	harmacy	
* Tab 60 mg	53.76 ·	28	<ul><li>Evista</li></ul>
(Evista Tab 60 mg to be delisted 1 April 2026)			

### ⇒SA1779 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Notes); or
- 2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
- 3 History of two significant osteoporotic fractures demonstrated radiologically; or
- 4 Documented T-Score less than or equal to -3.0 (see Notes); or
- 5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Notes); or
- 6 Patient has had a Special Authority approval for zoledronic acid (Underlying cause Osteoporosis) or has had a Special Authority approval for alendronate (Underlying cause - Osteoporosis) prior to 1 February 2019.

#### Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
   Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for raloxifene funding.

continued  c) Osteoporotic fractures are the incident events for severe (estal definitions of osteoporosis and fragility fracture. The WHO def -2.5 with one or more associated fragility fractures. Fragility fra forces that would not ordinarily cause fracture (minimal trauma fall from a standing height or less.  d) A vertebral fracture is defined as a 20% or greater reduction in relative to the posterior height of that body, or a 20% or greate body above or below the affected vertebral body.  RISEDRONATE SODIUM  Tab 35 mg  Risedronate Sandoz to be Principal Supply on 1 February 20  TERIPARATIDE - Special Authority see SA1139 below - Retail phar Inj 250 mcg per ml, 2.4 ml  SA1139 Special Authority for Subsidy  Initial application from any relevant practitioner. Approvals valid for All of the following:  1 The patient has severe, established osteoporosis; and 2 The patient has a documented T-score less than or equal to -3 3 The patient has had two or more fractures due to minimal traur 4 The patient has experienced at least one symptomatic new fra funded antiresorptive agent at adequate doses (see Notes).  Notes:  a) The bone mineral density (BMD) measurement used to derive absorptiometry (DXA). Quantitative ultrasound and quantitative	ines severe (esta actures are fractu ). The WHO has height of the and r reduction in any 3.00 26 macy	Per rosis, ar ablished ures tha s quanti	l) osteopo t occur as fied this a mid portion se heights	orosis as a T-score below s a result of mechanical s forces equivalent to a on of a vertebral body
c) Osteoporotic fractures are the incident events for severe (estal definitions of osteoporosis and fragility fracture. The WHO def -2.5 with one or more associated fragility fractures. Fragility fractures that would not ordinarily cause fracture (minimal trauma fall from a standing height or less.  d) A vertebral fracture is defined as a 20% or greater reduction in relative to the posterior height of that body, or a 20% or greate body above or below the affected vertebral body.  RISEDRONATE SODIUM  Tab 35 mg  Risedronate Sandoz to be Principal Supply on 1 February 20  TERIPARATIDE — Special Authority see SA1139 below — Retail phar Inj 250 mcg per ml, 2.4 ml  SA1139 Special Authority for Subsidy  Initial application from any relevant practitioner. Approvals valid for All of the following:  1 The patient has severe, established osteoporosis; and 2 The patient has a documented T-score less than or equal to -3 3 The patient has had two or more fractures due to minimal traur 4 The patient has experienced at least one symptomatic new frafunded antiresorptive agent at adequate doses (see Notes).  Notes:  a) The bone mineral density (BMD) measurement used to derive	ines severe (esta actures are fractu ). The WHO has height of the and r reduction in any 3.00 26 macy	ablished ures that s quanti terior or y of thes	l) osteopo t occur as fied this a mid portion se heights	orosis as a T-score below a result of mechanical is forces equivalent to a compared to the vertebral body a compared to the vertebra
definitions of osteoporosis and fragility fracture. The WHO def -2.5 with one or more associated fragility fractures. Fragility fractures that would not ordinarily cause fracture (minimal trauma fall from a standing height or less.  d) A vertebral fracture is defined as a 20% or greater reduction in relative to the posterior height of that body, or a 20% or greate body above or below the affected vertebral body.  RISEDRONATE SODIUM  Tab 35 mg  Risedronate Sandoz to be Principal Supply on 1 February 20  TERIPARATIDE − Special Authority see SA1139 below − Retail phar In j 250 mcg per ml, 2.4 ml  SA1139 Special Authority for Subsidy  Initial application from any relevant practitioner. Approvals valid for All of the following:  1 The patient has severe, established osteoporosis; and 2 The patient has a documented T-score less than or equal to -3 3 The patient has had two or more fractures due to minimal traur 4 The patient has experienced at least one symptomatic new fra funded antiresorptive agent at adequate doses (see Notes).  Notes:  a) The bone mineral density (BMD) measurement used to derive	ines severe (esta actures are fractu ). The WHO has height of the and r reduction in any 3.00 26 macy	ablished ures that s quanti terior or y of thes	l) osteopo t occur as fied this a mid portion se heights	orosis as a T-score below a result of mechanical is forces equivalent to a compared to the vertebral body a compared to the vertebra
Tab 35 mg	26 macy	4	<b>✓</b> F	lisedronate Sandoz
Risedronate Sandoz to be Principal Supply on 1 February 20  TERIPARATIDE – Special Authority see SA1139 below – Retail phar Inj 250 mcg per ml, 2.4 ml	26 macy	4	<b>√</b> R	Risedronate Sandoz
Inj 250 mcg per ml, 2.4 ml				
Initial application from any relevant practitioner. Approvals valid for All of the following:  1 The patient has severe, established osteoporosis; and 2 The patient has a documented T-score less than or equal to -3 3 The patient has had two or more fractures due to minimal traur 4 The patient has experienced at least one symptomatic new fra funded antiresorptive agent at adequate doses (see Notes).  Notes:  a) The bone mineral density (BMD) measurement used to derive	.200.27	1	<b>✓</b> T	eriparatide - Teva
Notes:  a) The bone mineral density (BMD) measurement used to derive	.0 (see Notes); a	ınd		
a) The bone mineral density (BMD) measurement used to derive				
b) Antiresorptive agents and their adequate doses for the purpose sodium tab 70 mg or tab 70 mg with colecalciferol 5,600 iu onc zoledronic acid 5 mg per year. If an intolerance of a severity n during the use of one antiresorptive agent, an alternate antires the minimum requirement of 12 months' continuous therapy. c) A vertebral fracture is defined as a 20% or greater reduction in relative to the posterior height of that body, or a 20% or greate body above or below the affected vertebral body. d) A maximum of 18 months of treatment (18 cartridges) will be s	e computed tomo es of this Special e weekly; raloxif- ecessitating perrorptive agent mu- height of the and r reduction in any	graphy Author ene hyd manent ist be tri	(QCT) are ity are def rochloride treatment alled so the mid portion	e not acceptable fined as: alendronate e tab 60 mg once daily; withdrawal develops hat the patient achieves on of a vertebral body
ZOLEDRONIC ACID Inj 0.05 mg per ml, 100 ml, bag	19.45	1	<b>√</b> Z	oledronic Acid Viatris
Zoledronic Acid Viatris to be Principal Supply on 1 February 2	2026			
Hyperuricaemia and Antigout				
ALLOPURINOL				
* Tab 100 mg	17.99	1,000	<b>✓</b> <u>l</u>	oca-Allopurinol

BENZBROMARONE - Special Authority see SA1963 on the next page - Retail pharmacy Tab 50 mg ......32.00

500

100

✓ Ipca-Allopurinol

✓ Narcaricin mite S29

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

### ⇒SA1963 Special Authority for Subsidy

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

- 1 The treatment remains appropriate and the patient is benefitting from the treatment; and
- 2 There is no evidence of liver toxicity and patient is continuing to receive regular (at least every three months) liver function tests

#### COLCHICINE

*	Tab 500 mcg6.30	100	✓ Colgout
	BUXOSTAT - Special Authority see SA2555 below - Retail pharmacy		·
	Tab 80 mg4.73	28	✓ Febuxostat (Teva)
	Tab 120 mg11.78	28	✓ Febuxostat (Teva)

### ⇒SA2555 Special Authority for Subsidy

Initial application — (Gout) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with gout; and
  - 2 Any of the following:
    - 2.1 The patient has a serum urate level greater than 0.36 mmol/l despite treatment with allopurinol at doses of at least 600 mg/day and addition of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
    - 2.2 The patient has experienced intolerable side effects from allopurinol such that treatment discontinuation is required and serum urate remains greater than 0.36 mmol/l despite use of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
    - 2.3 The patient has renal impairment such that probenecid is contraindicated or likely to be ineffective and serum urate remains greater than 0.36 mmol/l despite optimal treatment with allopurinol; or
    - 2.4 The patient has previously had an initial Special Authority approval for benzbromarone for treatment of gout...

Initial application — (Tumour lysis syndrome) only from a haematologist or oncologist. Approvals valid for 6 weeks for applications meeting the following criteria:

#### Both:

- 1 Patient is scheduled to receive cancer therapy carrying an intermediate or high risk of tumour lysis syndrome; and
- 2 Patient has a documented history of allopurinol intolerance.

Renewal — (Tumour lysis syndrome) only from a haematologist or oncologist. Approvals valid for 6 weeks where the treatment remains appropriate and the patient is benefitting from treatment.

#### **PROBENECID**

### Muscle Relaxants

# BACLOFEN \* Tab 10 mg

-1-	1 ab 10 mg		100	· I dolloll
	Inj 0.05 mg per ml, 1 ml ampoule - Subsidy by endorsement	11.55	1	✓ Lioresal Intrathecal
	Subsidised only for use in a programmable pump in patients	where oral ar	ntispastic age	nts have been ineffective or have
	caused intolerable side effects and the prescription is endor	sed according	ly.	
	Ini 2 mg nor ml 5 ml amnoula - Subsidy by andorsament	/Q0 Q1	10	✓ Sintatica Raclofon

Inj 2 mg per ml, 5 ml ampoule − Subsidy by endorsement...........490.91 10 ✓ Sintetica Baclofen Intrathecal

Subsidised only for use in a programmable pump in patients where oral antispastic agents have been ineffective or have caused intolerable side effects and the prescription is endorsed accordingly.

3 70

✓ Dacifon

### DANTROLENE

Cap 25 mg145.77	100	✓ Dantrium S29 S29
Cap 50 mg77.00	100	✓ Dantrium

	Subsidy (Manufacturer's Price)	Sub	Fully sidised	Brand or Generic
	\$	Per	•	Manufacturer
ORPHENADRINE CITRATE				
Tab 100 mg	23.25	100	✓ No	orflex

Subsidy (Manufacturer's Price)

Fully Subsidised Per

Brand or Generic Manufacturer

# **Agents for Parkinsonism and Related Disorders**

<b>Dopamine Agonist</b>	s and Relate	ed Agents
-------------------------	--------------	-----------

AMANTADINE HYDROCHLORIDE		
▲ Cap 100 mg	60	✓ Symmetrel
63.73	100	✓ Symmetrel
APOMORPHINE HYDROCHLORIDE		
▲ Inj 10 mg per ml, 2 ml ampoule59.50	5	✓ Movapo
▲ Inj 10 mg per ml, 5 ml ampoule121.84	5	✓ Movapo
ENTACAPONE		
▲ Tab 200 mg13.73	100	✓ Entacapone Viatris
LEVODOPA WITH BENSERAZIDE		
* Tab dispersible 50 mg with benserazide 12.5 mg	100	✓ Madopar Rapid
* Cap 50 mg with benserazide 12.5 mg	100	✓ Madopar 62.5
* Cap 100 mg with benserazide 25 mg	100	✓ Madopar 125
* Cap long-acting 100 mg with benserazide 25 mg22.85	100	✓ Madopar HBS
* Cap 200 mg with benserazide 50 mg26.25	100	✓ Madopar 250
LEVODOPA WITH CARBIDOPA		
* Tab 100 mg with carbidopa 25 mg	100	✓ Sinemet
* Tab long-acting 200 mg with carbidopa 50 mg44.99	100	✓ Sinemet CR
* Tab 250 mg with carbidopa 25 mg	100	✓ Sinemet
LEVODOPA WITH CARBIDOPA AND ENTACAPONE		
* Tab 50 mg with carbidopa 12.5 mg and entacapone 200 mg27.01	100	✓ Stalevo
* Tab 100 mg with carbidopa 25 mg and entacapone 200 mg34.18	100	✓ Stalevo
* Tab 150 mg with carbidopa 37.5 mg and entacapone 200 mg44.96	100	✓ Stalevo
* Tab 200 mg with carbidopa 50 mg and entacapone 200 mg51.23	100	✓ Stalevo
PRAMIPEXOLE HYDROCHLORIDE		
▲ Tab 0.25 mg	100	✓ Ramipex
▲ Tab 1 mg	100	✓ Ramipex
RASAGILINE		
* Tab 1 mg53.50	30	✓ Azilect S29
· ·	30	AZIIGGU
ROPINIROLE HYDROCHLORIDE	0.4	/ Davida
▲ Tab 0.25 mg	84	✓ Ropin
▲ Tab 1 mg	84 84	<ul><li>✓ Ropin</li><li>✓ Ropin</li></ul>
▲ Tab 2 mg	84	✓ Ropin
· ·	04	• nopili
TOLCAPONE A Table 100 mm	400	/ T
▲ Tab 100 mg152.38	100	✓ Tasmar

### **Anticholinergics**

BENZATROPINE MESYLATE			
Tab 2 mg	10.99	60	✓ Benztrop
Inj 1 mg per ml, 2 ml	95.00	5	✓ Phebra
a) Up to 10 inj available on a PSO			
b) Only on a PSO			

PROCYCLIDINE HYDROCHLORIDE

Tab 5 mg .......15.00 100 **✓ Kemadrin** 



Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

### Agents for Essential Tremor, Chorea and Related Disorders

RILUZOLE - Special Authority see SA1403 below - Retail pharmacy

Wastage claimable

Tab 50 mg .......117.00 56 **✓ Rilutek** 

### **⇒SA1403** Special Authority for Subsidy

**Initial application** only from a neurologist or respiratory specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has amyotrophic lateral sclerosis with disease duration of 5 years or less; and
- 2 The patient has at least 60 percent of predicted forced vital capacity within 2 months prior to the initial application; and
- 3 The patient has not undergone a tracheostomy; and
- 4 The patient has not experienced respiratory failure; and
- 5 Any of the following:
  - 5.1 The patient is ambulatory; or
  - 5.2 The patient is able to use upper limbs; or
  - 5.3 The patient is able to swallow.

**Renewal** from any relevant practitioner. Approvals valid for 18 months for applications meeting the following criteria: All of the following:

- 1 The patient has not undergone a tracheostomy; and
- 2 The patient has not experienced respiratory failure; and
- 3 Any of the following:
  - 3.1 The patient is ambulatory; or
  - 3.2 The patient is able to use upper limbs; or
  - 3.3 The patient is able to swallow.

**TETRABENAZINE** 

### **Anaesthetics**

### Local

### LIDOCAINE [LIGNOCAINE]

Gel 2%, tube − Subsidy by endorsement .......14.50 30 ml **✓ Xylocaine 2% Jelly** 

a) Up to 150 ml available on a PSO

b) Subsidised only if prescribed for urethral or cervical administration and the prescription is endorsed accordingly.

Gel 2%, 11 ml urethral syringe − Subsidy by endorsement..............65.45

- a) Up to 5 each available on a PSO
- Subsidised only if prescribed for urethral, cervical or rectal administration and the prescription is endorsed accordingly.
- c) Instillagel Lido to be Principal Supply on 1 February 2026

	Subsidy (Manufacturer's Price	۱ ور	Fully ubsidised	Brand or Generic
	\$	Per	ubsidised ✓	Manufacturer
LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE				
Oral (gel) soln 2%	30.80	200 ml	1	Xylocaine Viscous
	44.00		1	Mucosoothe
Xylocaine Viscous to be Principal Supply on 1 April 202	26			
Inj 1%, 5 ml ampoule - Up to 25 inj available on a PSO	15.00	25	1	Lidocaine-Baxter
	17.50	50		
	(35.00)			Xylocaine
Inj 2%, 5 ml ampoule - Up to 5 inj available on a PSO	27.50	25	✓	Lidocaine-Baxter
Inj 1%, 20 ml ampoule - Up to 5 inj available on a PSO	12.00	5		
	(20.00)			Xylocaine
Inj 1%, 20 ml vial - Up to 5 inj available on a PSO	19.50	5	1	Lidocaine-Baxter
Inj 2%, 20 ml vial - Up to 5 inj available on a PSO	14.00	5	1	Lidocaine-Baxter
Inj 10%, 5 ml ampoule - Subsidy by endorsement Subsidised only for people receiving palliative care sen		10		Xylocard 500 S29

### **Topical Local Anaesthetics**

### ⇒SA0906 Special Authority for Subsidy

(Mucosoothe Oral (gel) soln 2% to be delisted 1 April 2026)

**Initial application** from any relevant practitioner. Approvals valid for 2 years where the patient is a child with a chronic medical condition requiring frequent injections or venepuncture.

**Renewal** from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

### **Analgesics**

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 117

### **Non-opioid Analgesics**

ASPIRIN  * Tab dispersible 300 mg - Up to 30 tab available on a PSO5.6	55 100	✓ Ethics Aspirin
CAPSAICIN – Subsidy by endorsement Subsidised only if prescribed for post-herpetic neuralgia or diabetic peri	pheral neuropathy a	and the prescription is endorsed
accordingly. Crm 0.075%11.9	95 45 g OP	✓ Zo-Rub HP
NEFOPAM HYDROCHLORIDE Tab 30 mg 23.4	IO 90	✓ Zostrix HP  ✓ Acupan

	Subsidy Fully Brand or
	(Manufacturer's Price) Subsidised Generic
_	\$ Per ✓ Manufacturer
PAF	ACETAMOL
	Tab 500 mg - blister pack
	Maximum of 300 tab per prescription; can be waived by endorsement     Number 20 tab evolutions and PSO.
	b) Up to 30 tab available on a PSO c)
	<ol> <li>Subsidy by endorsement for higher quantities is available for patients with long term conditions who require regular daily dosing for one month or greater, and the prescription is annotated accordingly. Pharmacists may annotate the prescription as endorsed where dispensing history supports a long-term condition.</li> <li>Maximum of 100 tab per dispensing for non-endorsed patients. If quantities prescribed for more than 100 tabs (for non-endorsed patients), then dispense in repeat dispensings not exceeding 100 tab per dispensing.</li> <li>Tab 500 mg - bottle pack - Maximum of 300 tab per prescription; can be waived by endorsement</li></ol>
	<u>Paracetamol</u>
	<ol> <li>Subsidy by endorsement for higher quantities is available for patients with long term conditions who require regular daily dosing for one month or greater, and the prescription is annotated accordingly. Pharmacists may annotate the prescription as endorsed where dispensing history supports a long-term condition.</li> <li>Maximum of 100 tab per dispensing for non-endorsed patients. If quantities prescribed for more than 100 tabs (for non-endorsed patients), then dispense in repeat dispensings not exceeding 100 tab per dispensing.</li> </ol>
	Oral liq 120 mg per 5 ml
	a) Maximum of 600 ml per prescription; can be waived by endorsement b) Up to 200 ml available on a PSO c) Not in combination d)
	<ol> <li>Maximum of 200 ml per dispensing for non-endorsed patients. If quantities prescribed exceed 200 ml (for non-endorsed patients), then dispense in repeat dispensing not exceeding 200 ml per dispensing.</li> <li>Subsidy by endorsement for higher quantities is available for patients with long term conditions who require regular daily dosing for one month or greater and the prescription is endorsed or annotated accordingly. Pharmacists may annotate the prescription as endorsed where dispensing history supports a long-term condition.</li> <li>Note: 200 ml presentations of paracetamol oral liquid may be supplied on BSO to a Vaccinator (other than a</li> </ol>
	Pharmacist) under the provisions in Part I of Section A
	4) Note: Direct Provision by a pharmacist of up to 200 ml permitted under the provisions in Part I of Section A in conjunction with immunisation of a child under 2 years of age with meningococcal B multicomponent vaccine. Oral liq 250 mg per 5 ml
	a) Maximum of 600 ml per prescription; can be waived by endorsement
	b) Up to 200 ml available on a PSO
	c) Not in combination
	<ul> <li>d)</li> <li>1) Maximum of 200 ml per dispensing for non-endorsed patients. If quantities prescribed exceed 200 ml (for</li> </ul>
	non-endorsed patients), then dispensing for hon-endorsed patients. If quantities prescribed exceed 200 ml for non-endorsed patients), then dispense in repeat dispensing not exceeding 200 ml per dispensing.  2) Subsidy by endorsement for higher quantities is available for patients with long term conditions who require regular daily dosing for one month or greater and the prescription is endorsed or annotated accordingly. Pharmacists may annotate the prescription as endorsed where dispensing history supports a long-term condition.
	3) Note: 200 ml presentations of paracetamol oral liquid may be supplied on BSO to a Vaccinator (other than a
	Pharmacist) under the provisions in Part I of Section A 4) Note: Direct Provision by a pharmacist of up to 200 ml permitted under the provisions in Part I of Section A in
*	conjunction with immunisation of a child under 2 years of age with meningococcal B multicomponent vaccine.  Suppos 125 mg

				INE	1VUUS STSTEIN
		Subsidy (Manufacturer's Price)	Per	Fully Subsidised	Brand or Generic Manufacturer
*	Suppos 250 mg		10		Gacet
*	Suppos 500 mg	16.55	50	•	<u>Gacet</u>
C	pioid Analgesics				
CC	DEINE PHOSPHATE - Safety medicine; prescriber may dete	ermine dispensing fre	quen	су	
	Tab 15 mg	5.82	100	<b>'</b>	Noumed
	Tab 30 mg	6.88	100		Noumed
	Tab 60 mg	13.89	100		Noumed
DII	HYDROCODEINE TARTRATE				
	Tab long-acting 60 mg	9.20	60	1	DHC Continus
	DHC Continus to be Principal Supply on 1 February 202				
FF	NTANYL				
-	a) Only on a controlled drug form				
	b) No patient co-payment payable				
	c) Safety medicine; prescriber may determine dispensing fre	equency			
	Inj 50 mcg per ml, 2 ml ampoule		10	1	Boucher and Muir
	Inj 50 mcg per ml, 10 ml ampoule		10		Boucher and Muir
	Patch 12 mcg per hour		5		Fentanyl Sandoz
	Patch 25 mcg per hour		5		Fentanyl Sandoz
	Patch 50 mcg per hour	9.28	5	1	Fentanyl Sandoz
	Patch 75 mcg per hour	15.50	5	✓	Fentanyl Sandoz
	Patch 100 mcg per hour	16.37	5	✓	Fentanyl Sandoz
ME	THADONE HYDROCHLORIDE				
	a) Only on a controlled drug form				
	b) No patient co-payment payable				
	c) Safety medicine; prescriber may determine dispensing fre	equency			
	Tab 5 mg		10	✓	Methadone BNM
	Oral liq 2 mg per ml	7.80	200 n	nl 🗸	<u>Biodone</u>
	Oral liq 5 mg per ml	7.80	200 n	nl 🗸	Biodone Forte
	Oral liq 10 mg per ml	9.65	200 n		Biodone Extra Forte
	Inj 10 mg per ml, 1 ml	72.99	10	✓	AFT
MC	DRPHINE HYDROCHLORIDE				
	a) Only on a controlled drug form				
	b) No patient co-payment payable				
	c) Safety medicine; prescriber may determine dispensing fre				
	Oral liq 1 mg per ml		200 m		RA-Morph
	Oral liq 2 mg per ml		200 m		RA-Morph
	Oral liq 5 mg per ml		200 m		RA-Morph
	Ovel lie 10 men men vel	40.05	200		DA Massala

127

✓ RA-Morph

200 ml

Oral liq 10 mg per ml .......40.25

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
MORPHINE SULPHATE	Ψ	rei		Manuacturer
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing free	ILIENCV			
Tab immediate-release 10 mg		10	1	Sevredol
Tab immediate-release 20 mg		10		Sevredol
Cap long-acting 10 mg		10		m-Eslon
Cap long-acting 30 mg		10	/	m-Eslon
Cap long-acting 60 mg		10	1	m-Eslon
Cap long-acting 100 mg		10	1	m-Eslon
Oral lig 2 mg per ml		100 m	· •	Wockhardt \$29
0.sq =g po	29.80			Oramorph
				Oramorph CDC
				S29 S29
Inj 5 mg per ml, 1 ml ampoule - Up to 5 inj available on a PSC	n 5.96	5	1	Medsurge
Medsurge to be Principal Supply on 1 February 2026	J	J	•	weusurge
Inj 10 mg per ml, 1 ml ampoule – Up to 5 inj available on a PS	SO 499	5	/	Medsurge
Medsurge to be Principal Supply on 1 February 2026	JO 4.00	J	•	measurge
Inj 15 mg per ml, 1 ml ampoule – Up to 5 inj available on a PS	SO 6.93	5	1	Medsurge
Medsurge to be Principal Supply on 1 February 2026		Ŭ	_	ouou.go
Inj 30 mg per ml, 1 ml ampoule – Up to 5 inj available on a PS	SO7.28	5	1	Medsurge
Medsurge to be Principal Supply on 1 February 2026		-		
XYCODONE HYDROCHLORIDE				
a) Only on a controlled drug form     b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing free	ulopov.			
Tab controlled-release 5 mg		20		Oxycodone Sandoz
Tab immediate-release 5 mg		100		Oxycodone Amnea
Tab controlled-release 10 mg		20		Oxycodone Sandoz
Tab immediate-release 10 mg		100		Oxycodone Amnea
Tab controlled-release 20 mg.		20		Oxycodone Sando
Tab immediate-release 20 mg		100		Oxycodone Amnea
Tab controlled-release 40 mg		20		Oxycodone Sando
Tab controlled-release 80 mg		20		Oxycodone Sando
Oral lig 1 mg per ml		250 m		Oxycodone Lucis
C.a				Rosemont
Inj 10 mg per ml, 1 ml ampoule	4.37	5		Hameln
Inj 10 mg per ml, 2 ml ampoule		5		Hameln
Inj 50 mg per ml, 1 ml ampoule		5		Hameln
Oxycodone Lucis Oral liq 1 mg per ml to be delisted 1 June 2026)		-		
ARACETAMOL WITH CODEINE - Safety medicine; prescriber r		oncina	r frogueso	Δ/
				y Paracetamol +
Tab paracetamol 500 mg with codeine phosphate 8 mg	31.95	1,000	•	
				Codeine (Relieve

Paracetamol + Codeine (Relieve) to be Principal Supply on 1 February 2026

				11V003 3131EW
	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
PETHIDINE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing f		40		Naumad Dathidina
Tab 50 mg  Noumed Pethidine to be Principal Supply on 1 Februar		10	•	Noumed Pethidine
Inj 50 mg per ml, 1 ml ampoule – Up to 5 inj available on a		5	1	DBL Pethidine
ing oo mg per mi, i mi ampoule op to o mg available on a	1 0025.00	J	•	Hydrochloride
Inj 50 mg per ml, 2 ml ampoule – Up to 5 inj available on a	PSO30.72	5	1	DBL Pethidine
,g p, ap		-		Hydrochloride
TRAMADOL HYDROCHLORIDE				•
Tab sustained-release 100 mg	1.95	20	1	Tramal SR 100
Tab sustained-release 150 mg	2.95	20	✓	Tramal SR 150
Tab sustained-release 200 mg	3.80	20	•	Tramal SR 200
Cap 50 mg	3.33	100	•	Arrow-Tramadol
Autidonycocouto				
Antidepressants				
Cyclic and Related Agents				
AMITRIPTYLINE - Safety medicine; prescriber may determine	dispensing frequency			
Tab 10 mg		100		Arrow-Amitriptyline
Tab 25 mg		100		Arrow-Amitriptyline
Tab 50 mg		100		Arrow-Amitriptyline
CLOMIPRAMINE HYDROCHLORIDE - Safety medicine; preso				
Tab 25 mg		50		APO Clomipramine
Cap 10 mg	35.50	28	•	Clomipramine Teva
(Clomipramine Teva Cap 10 mg to be delisted 1 April 2026)				
DOSULEPIN [DOTHIEPIN] HYDROCHLORIDE - Subsidy by e				
a) Safety medicine; prescriber may determine dispensing f		مالد مالم	المنسلة المنسدة	
b) Subsidy by endorsement – Subsidised for patients who				
2019 and the prescription is endorsed accordingly. Pha exists a record of prior dispensing of dosulepin [dothiepi		eine	prescriptio	n as endorsed where ther
Tab 75 mg		30	/	Dosulepin Viatris
Cap 25 mg		50		Dosulepin
				Viatris S29
IMIPRAMINE HYDROCHLORIDE - Safety medicine; prescribe	er may determine disne	nsino	ı frequency	ı
Tab 10 mg		50		Tofranil
	10.96	100	1	Tofranil
Tab 25 mg	4.93	28	1	Imipramine
·				Crescent S29
	8.80	50	1	Tofranil
NORTRIPTYLINE HYDROCHLORIDE				
a) Brand switch fee payable (Pharmacode 2715740) - see	page 282 for details			
b) Safety medicine; prescriber may determine dispensing f				
Tab 10 mg		50	1	Allegron
•	2.46	100		Norpress
Tab 25 mg		50		Allegron
	6.29	180	1	Norpress
(Norpress Tab 10 mg to be delisted 1 March 2026)				
(Norpress Tab 25 mg to be delisted 1 March 2026)				

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
Monoamine-Oxidase Inhibitors (MAOIs) - Non S	Selective			
TRANYLCYPROMINE SULPHATE  * Tab 10 mg	22.94	50	✓	Parnate
Monoamine-Oxidase Type A Inhibitors				
MOCLOBEMIDE * Tab 150 mg * Tab 300 mg		60 60		Aurorix Aurorix
Selective Serotonin Reuptake Inhibitors				
CITALOPRAM HYDROBROMIDE  * Tab 20 mg ESCITALOPRAM	3.55	84	•	Celapram
* Tab 10 mg	1.07	28	•	Ipca-Escitalopram Escitalopram (Ethics)
★ Tab 20 mg FLUOXETINE HYDROCHLORIDE	1.49	28	•	Ipca-Escitalopram
Subsidised by endorsement     When prescribed for a patient who cannot swa		28 apsu		Fluox e prescription is endorsed
1) When prescribed for a patient who cannot swa accordingly; or 2) When prescribed in a daily dose that is not a mendorsed. Note: Tablets should be combined b) Fluox to be Principal Supply on 1 March 2026  Cap 20 mg	llow whole tablets or conultiple of 20 mg in whole with capsules to facili	apsu	ase the pre	e prescription is endorsec
a) Subsidised by endorsement  1) When prescribed for a patient who cannot swa accordingly; or  2) When prescribed in a daily dose that is not a mendorsed. Note: Tablets should be combined b) Fluox to be Principal Supply on 1 March 2026  Cap 20 mg	Illow whole tablets or conultiple of 20 mg in whole with capsules to facili	apsuich catate i	ase the preincrementa	e prescription is endorsed scription is deemed to be I 10 mg doses.  Arrow-Fluoxetine
a) Subsidised by endorsement  1) When prescribed for a patient who cannot swa accordingly; or  2) When prescribed in a daily dose that is not a mendorsed. Note: Tablets should be combined b) Fluox to be Principal Supply on 1 March 2026  Cap 20 mg	Illow whole tablets or conultiple of 20 mg in whole with capsules to facili	apsuich cate i	ase the preincrementa	e prescription is endorsed scription is deemed to be I 10 mg doses.
a) Subsidised by endorsement  1) When prescribed for a patient who cannot swa accordingly; or  2) When prescribed in a daily dose that is not a mendorsed. Note: Tablets should be combined b) Fluox to be Principal Supply on 1 March 2026  Cap 20 mg	Illow whole tablets or conultiple of 20 mg in whole with capsules to facili	apsuich catate i	ase the preincrementa	e prescription is endorsed excription is deemed to be I 10 mg doses.  Arrow-Fluoxetine  Paxtine
a) Subsidised by endorsement  1) When prescribed for a patient who cannot swa accordingly; or  2) When prescribed in a daily dose that is not a mendorsed. Note: Tablets should be combined b) Fluox to be Principal Supply on 1 March 2026  Cap 20 mg	ullow whole tablets or conultiple of 20 mg in whole with capsules to facili	apsuich catate i	ase the preincrementa	e prescription is endorsed excription is deemed to be I 10 mg doses.  Arrow-Fluoxetine  Paxtine
a) Subsidised by endorsement  1) When prescribed for a patient who cannot swa accordingly; or  2) When prescribed in a daily dose that is not a mendorsed. Note: Tablets should be combined b) Fluox to be Principal Supply on 1 March 2026  * Cap 20 mg	nultiple of 20 mg in who with capsules to facili	apsudich contate i	ase the preincrementa	e prescription is endorsed excription is deemed to be I 10 mg doses.  Arrow-Fluoxetine  Paxtine Loxamine
a) Subsidised by endorsement  1) When prescribed for a patient who cannot swa accordingly; or  2) When prescribed in a daily dose that is not a mendorsed. Note: Tablets should be combined b) Fluox to be Principal Supply on 1 March 2026  Cap 20 mg	nultiple of 20 mg in who with capsules to facili	apsurich catate i 90 30 90	ase the preincrementa	e prescription is endorsed escription is deemed to be 1 10 mg doses.  Arrow-Fluoxetine  Paxtine Loxamine  Setrona
a) Subsidised by endorsement  1) When prescribed for a patient who cannot swa accordingly; or  2) When prescribed in a daily dose that is not a mendorsed. Note: Tablets should be combined b) Fluox to be Principal Supply on 1 March 2026  * Cap 20 mg	Illow whole tablets or conultiple of 20 mg in whole with capsules to facility	30 30 30	ase the preincrementa	e prescription is endorsed excription is deemed to be a 10 mg doses.  Arrow-Fluoxetine  Paxtine Loxamine  Setrona  Setrona
a) Subsidised by endorsement  1) When prescribed for a patient who cannot swa accordingly; or  2) When prescribed in a daily dose that is not a mendorsed. Note: Tablets should be combined b) Fluox to be Principal Supply on 1 March 2026  * Cap 20 mg	Illow whole tablets or conultiple of 20 mg in whole with capsules to facility	apsurich catate i 90 30 90	ase the preincrementa	e prescription is endorsed escription is deemed to be 1 10 mg doses.  Arrow-Fluoxetine  Paxtine Loxamine  Setrona
a) Subsidised by endorsement  1) When prescribed for a patient who cannot swa accordingly; or  2) When prescribed in a daily dose that is not a mendorsed. Note: Tablets should be combined b) Fluox to be Principal Supply on 1 March 2026  * Cap 20 mg		30 30 30 30	ales and the preincrementa	e prescription is endorsed escription is deemed to be a 100 mg doses.  Arrow-Fluoxetine  Paxtine Loxamine  Setrona  Setrona  Noumed Noumed Noumed
a) Subsidised by endorsement  1) When prescribed for a patient who cannot swa accordingly; or  2) When prescribed in a daily dose that is not a mendorsed. Note: Tablets should be combined b) Fluox to be Principal Supply on 1 March 2026  * Cap 20 mg	Illow whole tablets or conultiple of 20 mg in whole with capsules to facility	30 30 30	ales and the pre-	e prescription is endorsed excription is deemed to be a 10 mg doses.  Arrow-Fluoxetine  Paxtine Loxamine  Setrona  Setrona

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

# **Antiepilepsy Drugs**

### **Agents for Control of Status Epilepticus**

DIAZEPAM – Safety medicine; prescriber may determine dispensing frequency Inj 5 mg per ml, 2 ml ampoule – Subsidy by endorsement27.92	5	✓ Hospira
a) Up to 5 inj available on a PSO		
b) Only on a PSO		
c) PSO must be endorsed "not for anaesthetic procedures".	_	4 4
Rectal tubes 5 mg - Up to 5 tube available on a PSO54.58	5	Stesolid
PHENYTOIN SODIUM		
* Inj 50 mg per ml, 2 ml ampoule - Up to 5 inj available on a		
PSO104.58	5	<ul><li>Hospira</li></ul>
* Inj 50 mg per ml, 5 ml ampoule - Up to 5 inj available on a		-
PSO154.01	5	✓ Hospira
(Hospira Inj 50 mg per ml, 2 ml ampoule to be delisted 1 February 2026)		•

## **Control of Epilepsy**

CARBAMAZEPINE			
* Tab 200 mg	14.53	100	<ul><li>Tegretol</li></ul>
* Tab long-acting 200 mg	16.98	100	Tegretol CR
	33.96	200	Tegretol CR
* Tab 400 mg	34.58	100	Tegretol
* Tab long-acting 400 mg	39.17	100	Tegretol CR
* Oral liq 20 mg per ml	26.37	250 ml	Tegretol
CLOBAZAM - Safety medicine; prescriber may determine dis	spensing frequency		
Tab 10 mg	9.12	50	✓ Frisium
CLONAZEPAM - Safety medicine; prescriber may determine	dispensing frequen	су	
Oral drops 2.5 mg per ml	7.38	10 ml OP	✓ Rivotril
ETHOSUXIMIDE			
Cap 250 mg	140.88	100	Zarontin
Oral liq 250 mg per 5 ml		200 ml	Zarontin
GABAPENTIN			
Note: Not subsidised in combination with subsidised pre-	gabalin		
* Cap 100 mg	6.45	100	✓ Nupentin
* Cap 300 mg		100	✓ Nupentin
* Cap 400 mg	10.26	100	✓ Nupentin
LACOSAMIDE - Special Authority see SA2267 on the next p	age – Retail pharma	асу	
▲ Tab 50 mg	25.04	14	✓ Vimpat
▲ Tab 100 mg	50.06	14	✓ Vimpat
	200.24	56	✓ Vimpat
▲ Tab 150 mg	75.10	14	Vimpat
	300.40	56	Vimpat
▲ Tab 200 mg	400.55	56	✓ Vimpat

Subsidy (Manufacturer's Price) S

Fully Subsidised Brand or Generic Manufacturer

### **⇒SA2267** Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria:

Both:

- 1 Patient has focal epilepsy; and
- 2 Seizures are not adequately controlled by, or patient has experienced unacceptable side effects from, optimal treatment with all of the following: sodium valproate, topiramate, levetiracetam and any two of carbamazepine, lamotrigine and phenytoin sodium (see Note).

Note: Those of childbearing potential are not required to trial phenytoin sodium, sodium valproate, or topiramate. Those who can father children are not required to trial sodium valproate.

**Renewal** from any relevant practitioner. Approvals valid for 24 months where the patient has demonstrated a significant and sustained improvement in seizure rate or severity and/or quality of life compared with that prior to starting lacosamide treatment.

		•	•
LAMOTRIGINE			,
	2 mg55.00 30		Lamictal
	5 mg50.00 30		Lamictal
	25 mg4.20 56		Logem
	50 mg		Logem
* Tab dispersible	100 mg	,	Logem
LEVETIRACETAM			
Tab 250 mg	5.84 60		Everet
Tab 500 mg			Everet
Tab 750 mg	16.71 60		Everet
Tab 1,000 mg	21.82 60	·	Everet
Oral liq 100 mg	per ml44.78 300 m	-	Levetiracetam-AFT
Inj 100 mg per r	ml, 5 ml vial38.95 10	·	Levetiracetam-AFT
PHENOBARBITON	E		
For phenobarbi	tone oral liquid refer Standard Formulae, page 284		
•	248.50 50	) 🗸	Noumed
· ·			Phenobarbitone
Tab 30 mg		) 🗸	Noumed
<b>3</b>			Phenobarbitone
PHENYTOIN SODIL	IM		
	75.00 20	n 🗸	Dilantin Infatab
•	74.00 20		Dilantin
	37.00 20		Dilantin
	per 5 ml22.03 500	-	Dilantin Paediatric
	33		Dianan racalatio
PREGABALIN	idiand in combination with authoidiand aphanentia		
	idised in combination with subsidised gabapentin		' Lumino
★ Cap 25 mg	2.25 56		Lyrica
¥ Con 75 mg			Pregabalin Pfizer
* Cap 75 mg	2.65 56		Lyrica
¥ Con 150 ma	4.01 56		Pregabalin Pfizer
* Cap 150 mg	4.01 56		Lyrica
¥ Con 200 ma	7.00 56		Pregabalin Pfizer
* Cap 300 mg	7.38 56		Lyrica
		•	Pregabalin Pfizer
PRIMIDONE			
* Tab 250 mg	37.35 10	) 🗸	Primidone Clinect

,	Subsidy	:aa\ C	Fully	Brand or
(I	Manufacturer's Pr \$	Per	Subsidised	Generic Manufacturer
ODIUM VALPROATE				
Tab 100 mg	13.65	100	<b>√</b> E	pilim Crushable
Tab 200 mg EC	27.44	100	<b>✓</b> E	pilim
Tab 500 mg EC		100	<b>√</b> E	pilim
★ Oral lig 200 mg per 5 ml		300 ml	<b>√</b> E	pilim S/F Liquid
				pilim Syrup
₭ Inj 100 mg per ml, 4 ml	41.50	1		pilim IV
STIRIPENTOL - Special Authority see SA2268 below - Retail pha				•
Cap 250 mg	,	60	<b>√</b> Γ	Diacomit
Powder for oral liq 250 mg sachet		60		Diacomit
Fowder for oral liq 250 flig Sacriet	509.29	00	• 6	naconni

### ⇒SA2268 Special Authority for Subsidy

**TOPIRAMATE** 

Initial application only from a paediatric neurologist or Practitioner on the recommendation of a paediatric neurologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient has confirmed diagnosis of Dravet syndrome; and
- 2 Seizures have been inadequately controlled by appropriate courses of sodium valproate, clobazam and at least two of the following: topiramate, levetiracetam, ketogenic diet.

44.26

Note: Those of childbearing potential are not required to trial sodium valproate or topiramate. Those who can father children are not required to trial sodium valproate.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where the patient continues to benefit from treatment as measured by reduced seizure frequency from baseline.

▲ Tab 25 mg	11.07
·	
	26.04
▲ Tab 50 mg	18.81

- ✓ Arrow-Topiramate
- ✓ Topiramate Actavis
- ✓ Topamax

60

60

60

60

60

100

60

- ✓ Arrow-Topiramate ✓ Topiramate Actavis
- ✓ Topamax
- ✓ Arrow-Topiramate
- ✓ Topiramate Actavis
- ✓ Topamax
- ✓ Arrow-Topiramate ✓ Topiramate Actavis
- ✓ Topamax
- ✓ Topamax
- ✓ Topamax
- Sabril
- ✓ Sabril S29
- ▲ Powder for oral soln 500 mg per sachet......71.58 (Sabril S29 Powder for oral soln 500 mg per sachet to be delisted 1 May 2026)

Tab 100 mg ......31.99

▲ Tab 200 mg .......55.19

▲ Sprinkle cap 25 mg......26.04

▲ Tab 500 mg .......119.30

VIGABATRIN - Special Authority see SA2088 below - Retail pharmacy

⇒SA2088 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Any of the following:
  - 1.1 Patient has infantile spasms; or
  - 1.2 Both:
    - 1.2.1 Patient has epilepsy; and

ubsidy cturer's Price) Subs	Fully	Brand or Generic
 \$ Per	•	Manufacturer

continued...

- 1.2.2 Either:
  - 1.2.2.1 Seizures are not adequately controlled with optimal treatment with other antiepilepsy agents; or
  - 1.2.2.2 Seizures are controlled adequately but the patient has experienced unacceptable side effects from optimal treatment with other antiepilepsy agents; or
- 1.3 Patient has tuberous sclerosis complex; and
- 2 Either:
  - 2.1 Patient is, or will be, receiving regular automated visual field testing (ideally before starting therapy and on a 6-monthly basis thereafter); or
  - 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields...

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient has demonstrated a significant and sustained improvement in seizure rate or severity and or quality of life; and
- 2 Either:
  - 2.1 Patient is receiving regular automated visual field testing (ideally every 6 months) on an ongoing basis for duration of treatment with vigabatrin; or
  - 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields...

### **Antimigraine Preparations**

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 117

### Acute Migraine Treatment

RIZATRIPTAN			
Tab orodispersible 10 mg	4.84	30	✓ Rizamelt
SUMATRIPTAN			
Tab 50 mg	14.41	90	✓ Sumagran
Tab 100 mg	22.68	90	✓ Sumagran
Inj 12 mg per ml, 0.5 ml prefilled pen - Maximum of 10 inj per			
prescription	29.80	2 OP	✓ Clustran

### Prophylaxis of Migraine

For Beta Adrenoceptor Blockers refer to CARDIOVASCULAR SYSTEM, page 50

**PIZOTIFEN** 

\* Tab 500 mcg.......23.21 100 ✓ Sandomigran

### **Antinausea and Vertigo Agents**

For Antispasmodics refer to ALIMENTARY TRACT, page 8

APREPITANT - Special Authority see SA0987 below - Retail pharmacv 3 OP

Cap 2 × 80 mg and 1 × 125 mg......21.90 ✓ Emend Tri-Pack

### ⇒SA0987 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months where the patient is undergoing highly emetogenic chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.

Renewal from any relevant practitioner. Approvals valid for 12 months where the patient is undergoing highly emetogenic chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.

BETAHISTINE DIHYDROCHLORIDE

100 ✓ Serc

	Subsidy (Manufactured Price)		. ,	ınd or
	(Manufacturer's Price) \$	Per		neric nufacturer
CYCLIZINE HYDROCHLORIDE				
Tab 50 mg	0.66	10	✓ Nausi	<u>calm</u>
CYCLIZINE LACTATE				
Inj 50 mg per ml, 1 ml ampoule - Up to 10 inj available on a	16.36	10	✓ Hame	In
DOMPERIDONE				
* Tab 10 mg	3.80	100	✓ <u>Domp</u> <u>Viat</u>	<u>eridone</u> <u>ris</u>
HYOSCINE HYDROBROMIDE				
* Inj 400 mcg per ml, 1 ml ampoule	93.00	10	✓ Martir	ndale S29
Patch 1 mg per 72 hours – Special Authority see SA1998				
below – Retail pharmacy	88.50	10		olamine nsdermal tem Viatris

### ⇒SA1998 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Control of intractable nausea, vomiting, or inability to swallow saliva in the treatment of malignancy or chronic disease where the patient cannot tolerate or does not adequately respond to oral anti-nausea agents; or
- 2 Control of clozapine-induced hypersalivation where trials of at least two other alternative treatments have proven ineffective.

**Renewal** from any relevant practitioner. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

METOCLOPRAMIDE HYDROCHLORIDE		
* Tab 10 mg – Up to 30 tab available on a PSO1.57	100	✓ <u>Metoclopramide</u> Actavis 10
* Inj 5 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO5.48 7.00	10	<ul><li>✓ Medsurge</li><li>✓ Baxter</li></ul>
Medsurge to be Principal Supply on 1 April 2026		
(Baxter Inj 5 mg per ml, 2 ml ampoule to be delisted 1 April 2026)		
ONDANSETRON		
* Tab 4 mg1.95	50	✓ Periset
Tab disp 4 mg - Up to 10 tab available on a PSO0.56	10	✓ Periset ODT
* Tab 8 mg	50	✓ Periset
Tab disp 8 mg - Up to 10 tab available on a PSO0.90	10	✓ Periset ODT
PROCHLORPERAZINE		
* Tab 3 mg buccal5.97	50	
(30.00)		Prochlorperazine maleate (Brown & Burk)
* Tab 5 mg - Up to 30 tab available on a PSO25.00	250	✓ Nausafix
* Inj 12.5 mg per ml, 1 ml - Up to 5 inj available on a PSO25.81	10	✓ Stemetil

Subsidy (Manufacturer's Price)

Fully Subsidised Per

Brand or Generic Manufacturer

# Antipsychotics

### General

AMISULPRIDE - Safety medicine; prescriber may determine d	ispensing frequen	су	
Tab 100 mg		30	✓ Sulprix
Tab 200 mg	14.47	60	✓ Sulprix
Tab 400 mg	35.06	60	✓ Sulprix
ARIPIPRAZOLE - Safety medicine; prescriber may determine	dispensing freque	ncv	
Tab 5 mg		30	✓ Aripiprazole Sandoz
Tab 10 mg		30	✓ Aripiprazole Sandoz
Tab 15 mg		30	✓ Aripiprazole Sandoz
Tab 20 mg		30	✓ Aripiprazole Sandoz
Tab 30 mg		30	✓ Aripiprazole Sandoz
CHLORPROMAZINE HYDROCHLORIDE - Safety medicine; p	rescriber may det	ermine disnen	sing frequency
Tab 25 mg – Up to 30 tab available on a PSO	•	100	✓ Largactil
Tab 100 mg – Up to 30 tab available on a PSO		100	✓ Largactil
Inj 25 mg per ml, 2 ml – Up to 5 inj available on a PSO		10	✓ Largactil
			- Largaom
CLOZAPINE – Hospital pharmacy [HP4]			
Safety medicine; prescriber may determine dispensing freq	•	50	✓ Clopine
Tab 25 mg	0.09	50	✓ Clopine ✓ Clozaril
	13.37	100	✓ Clopine
	13.37	100	✓ Clopine ✓ Clozaril
Tab 50 mg	8 67	50	✓ Clopine
1 ab 30 mg	17.33	100	✓ Clopine
Tab 100 mg		50	✓ Clopine
Tab Too mg		30	✓ Clozaril
	34.65	100	✓ Clopine
	01.00	100	✓ Clozaril
Tab 200 mg	34 65	50	✓ Clopine
1 ab 200 mg	69.30	100	✓ Clopine
Suspension 50 mg per ml		100 ml	✓ Versacloz
HALOPERIDOL – Safety medicine; prescriber may determine of		201	
Tab 500 mcg - Up to 30 tab available on a PSO		100	✓ Serenace
Tab 1.5 mg - Up to 30 tab available on a PSO		100	✓ Serenace
Tab 5 mg - Up to 30 tab available on a PSO		50	✓ Serenace
Tab 5 mg Op to 50 tab available on a 1 50	29.72	100	✓ Serenace
Oral lig 2 mg per ml - Up to 200 ml available on a PSO		100 ml	✓ Serenace
Inj 5 mg per ml, 1 ml ampoule – Up to 5 inj available on a F		10	✓ Serenace
			Colonado
LEVOMEPROMAZINE – Safety medicine; prescriber may dete			( Naminan (Curias)
Tab 25 mg (33.8 mg as a maleate)		100	✓ Nozinan (Swiss)
Tab 25 mg as a maleate		100	✓ Nozinan
Tab 100 mg (135 mg as a maleate)		100	✓ Nozinan (Swiss)
Tab 100 mg as a maleate		100	Nozinan
LEVOMEPROMAZINE HYDROCHLORIDE – Safety medicine;			0 , ,
Inj 25 mg per ml, 1 ml ampoule	23.26	10	✓ Wockhardt
LITHIUM CARBONATE - Safety medicine; prescriber may dete	ermine dispensing	frequency	
Tab long-acting 400 mg		100	✓ Priadel
Cap 250 mg	35.78	100	✓ Douglas
-			-

(Manufacturer's Price) \$ pensing frequency	Per	Subsidised <	
т	1 01		Manufacturer
pensina frequency			Wandidotalo
1.40	30	1	Zypine
1.93	30		Zypine
			Zypine ODT Zypine
			Zypine
	28		Zypine ODT
	100	/	Neulactil
			Neulactil
	100		- Touridotti
	30	1	Quetiapine
			Viatris S29
2.36	90	1	Quetapel
13.11	500		Quetiapine
			Viatris \$29
6.40	90	1	Quetapel
10.97	90	✓	Quetapel
	90		Quetapel
	60	1	Risperidone (Teva)
			Risperdal
	00		Risperidone (Teva)
10.29	30 ml		Risperon
34.30	100 m		Risperon
spensing frequency			
17.90	60	✓	Zusdone
27.41	60	✓	Zusdone
38.39	60	✓	Zusdone
46.55	60	✓	Zusdone
escriber may determin	e disp	ensing fre	equency
31.45	100		Clopixol
Inharmacy			
	1	J	Abilify Maintena
	-		Abilify Maintena
	'	•	Aviiiiy ivialiilelia
	13.116.4010.9715.83 spensing frequency2.172.442.724.506.2510.29 34.30 spensing frequency17.9027.4138.3946.55 escriber may determine31.45  I pharmacy lency273.56	1.80 28 1.93 30 2.89 28 28 28 29ensing frequency 13.61 100 48.45 100 2.36 90 13.11 500 2.36 90 13.11 500 2.36 90 13.11 500 2.36 90 13.11 500 2.36 90 13.11 500 2.36 90 13.11 500 2.36 90 13.11 500 2.36 90 13.11 500 2.36 90 13.11 500 2.36 90 13.11 500 2.36 90 13.11 500 2.36 90 13.11 500 2.36 90 13.11 500 2.36 90 13.11 500 2.37 90 2.38 90 2.39 90 2.44 60 2.72 60 2.72 60 2.72 60 2.72 60 2.72 60 2.74 60 2.72 60 2.74 1	1.80

**Initial application** from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

1 Either:



Subsidy		Fully	Brand or
(Manufacturer's Price)		osidised	Generic
<u> </u>	Per		Manufacturer

#### continued...

- 1.1 The patient has had an initial Special Authority approval for risperidone depot injection, paliperidone depot injection or olanzapine depot injection; or
- 1.2 All of the following:
  - 1.2.1 The patient has schizophrenia or other psychotic disorder; and
  - 1.2.2 The patient has received treatment with oral atypical antipsychotic agents but has been unable to adhere; and
  - 1.2.3 The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months; or
- 2 Patient has been unable to access olanzapine depot injection due to supply issues with olanzapine depot injection, or otherwise would have been started on olanzapine depot injection but has been unable to due to supply issues with olanzapine depot injection.

Notes: The Olanzapine depot injection Special Authority criteria that apply to criterion 2 in this Aripiprazole Special Authority application are as follows:

- The patient has had an initial Special Authority approval for paliperidone depot injection or risperidone depot injection; or
- All of the following:
  - The patient has schizophrenia; and
  - The patient has not been able to adhere with treatment using oral atypical antipsychotic agents; and
  - The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months.

FLUPENTHIXOL DECANOATE - Safety medicine; prescriber may of	letermine dispens	sing frequen	су	
Inj 20 mg per ml, 1 ml - Up to 5 inj available on a PSO	13.14	5	1	Fluanxol
Inj 20 mg per ml, 2 ml - Up to 5 inj available on a PSO	20.90	5	✓	Fluanxol
Inj 100 mg per ml, 1 ml - Up to 5 inj available on a PSO	40.87	5	1	Fluanxol
HALOPERIDOL DECANOATE - Safety medicine; prescriber may de	etermine dispensi	ng frequenc	y	
Inj 50 mg per ml, 1 ml - Up to 5 inj available on a PSO	28.39	5	1	Haldol
Inj 100 mg per ml, 1 ml - Up to 5 inj available on a PSO	55.90	5	✓	Haldol Concentrate
			✓	Haldol
				Decanoas S29
OLANZAPINE - Special Authority see SA2313 below - Retail pharm	nacy			
a) Safety medicine; prescriber may determine dispensing freque	ency			
b) Note – no new patients to be initiated on olanzapine.				
Inj 210 mg vial	252.00	1	✓	Zyprexa Relprevv
Inj 300 mg vial	414.00	1	✓	Zyprexa Relprevv
Inj 405 mg vial	504.00	1	✓	Zyprexa Relprevv

### ⇒SA2313 Special Authority for Subsidy

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of olanzapine depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

PALIPERIDONE - Special Authority see SA2396 on the next page - Retail pharmacy

Safety medicine; prescriber may determine dispensing frequency	
Inj 25 mg syringe194.25	✓ Invega Sustenna
Inj 50 mg syringe271.95	✓ Invega Sustenna
Inj 75 mg syringe	✓ Invega Sustenna
Inj 100 mg syringe	✓ Invega Sustenna
Inj 150 mg syringe	✓ Invega Sustenna

### **NERVOUS SYSTEM**

S	Subsidy	Fully	Brand or
(Manufa	acturer's Price) Sub	sidised	Generic
	\$ Per	•	Manufacturer

### ⇒SA2396 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Fither:

- 1 The patient has had an initial Special Authority approval for risperidone depot injection or olanzapine depot injection or aripiprazole depot injection; or
- 2 All of the following:
  - 2.1 The patient has schizophrenia or other psychotic disorder; and
  - 2.2 Has been unable to adhere to treatment using oral atypical antipsychotic agents; and
  - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

**Renewal** from any relevant practitioner. Approvals valid for 12 months where the initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

### PALIPERIDONE PALMITATE - Special Authority see SA2167 below - Retail pharmacy

Inj 175 mg syringe	815.85 <sup>'</sup>	1	✓ Invega Trinza
Inj 263 mg syringe		1	✓ Invega Trinza
Inj 350 mg syringe	1,305.36	1	✓ Invega Trinza
Inj 525 mg syringe	1,305.36	1	✓ Invega Trinza

### **⇒SA2167** Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has schizophrenia; and
- 2 The patient has had an initial Special Authority approval for paliperidone once-monthly depot injection.

**Renewal** from any relevant practitioner. Approvals valid for 12 months where the initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

### RISPERIDONE - Special Authority see SA2397 below - Retail pharmacy

Safety medicine; prescriber may determine dispensing frequency

Inj 25 mg vial135.98	1	Risperdal Consta
Inj 37.5 mg vial178.71	1	✓ Risperdal Consta
Inj 50 mg vial217.56	1	✓ Risperdal Consta

### ⇒SA2397 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or olanzapine depot injection; or aripiprazole depot injection; or
- 2 All of the following:
  - 2.1 The patient has schizophrenia or other psychotic disorder; and
  - 2.2 Has not been able to adhere with treatment using oral atypical antipsychotic agents; and
  - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

**Renewal** from any relevant practitioner. Approvals valid for 12 months where the initiation of risperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

ZUCLOPENTHIXOL DECANOATE - Safety medicine; prescriber may determine dispensing frequency

Inj 200 mg per ml, 1 ml − Up to 5 inj available on a PSO.......19.80 5 ✓ Clopixol



Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per  Manufacturer
---

### **Anxiolytics**

BUSPIRONE HYDROCHLORIDE			
* Tab 5 mg	13.95	100	<ul> <li>Buspirone Viatris</li> </ul>
* Tab 10 mg	12.50	100	<ul> <li>Buspirone Viatris</li> </ul>
CLONAZEPAM - Safety medicine; prescriber may determine dis	pensing frequency		
Tab 500 mcg	5.64	100	✓ Paxam
Tab 2 mg	10.78	100	✓ Paxam
DIAZEPAM - Safety medicine; prescriber may determine dispen	sing frequency		
Tab 2 mg	95.00	500	✓ Arrow-Diazepam
Tab 5 mg	115.00	500	✓ Arrow-Diazepam
LORAZEPAM - Safety medicine; prescriber may determine dispe	ensing frequency		
Tab 1 mg	10.20	250	✓ Ativan
Tab 2.5 mg	13.13	100	✓ <u>Ativan</u>

### **Multiple Sclerosis Treatments**

### ⇒SA2274 Special Authority for Subsidy

Initial application — (Multiple Sclerosis - dimethyl fumarate, fingolimod, glatiramer acetate, interferon beta-1-alpha, interferon beta-1-beta, natalizumab and teriflunomide) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Fither:

- 1 All of the following:
  - 1.1 Diagnosis of multiple sclerosis (MS) meets the McDonald 2017 diagnostic criteria for MS and has been confirmed by a neurologist; and
  - 1.2 Patient has an EDSS score between 0 6.0; and
  - 1.3 Patient has had at least one significant attack of MS in the previous 12 months or two significant attacks in the past 24 months: and
  - 1.4 All of the following:
    - 1.4.1 Each significant attack must be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the attack, but the neurologist/physician must be satisfied that the clinical features were characteristic): and
    - 1.4.2 Each significant attack is associated with characteristic new symptom(s)/sign(s) or substantially worsening of previously experienced symptoms(s)/sign(s); and
    - 1.4.3 Each significant attack has lasted at least one week and has started at least one month after the onset of a previous attack (where relevant); and
    - 1.4.4 Each significant attack can be distinguished from the effects of general fatigue; and is not associated with a fever (T> 37.5°C); and
    - 1.4.5 Either:
      - 1.4.5.1 Each significant attack is severe enough to change either the EDSS or at least one of the Kurtze Functional System scores by at least 1 point; or
      - 1.4.5.2 Each significant attack is a recurrent paroxysmal symptom of multiple sclerosis (tonic seizures/spasms, trigeminal neuralgia, Lhermitte's symptom); and
  - 1.5 Evidence of new inflammatory activity on an MRI scan within the past 24 months; and
  - 1.6 Any of the following:
    - 1.6.1 A sign of that new inflammatory activity on MRI scanning (in criterion 5 immediately above) is a gadolinium enhancing lesion; or

			NEKV	OUSSYSTEM	
	Subsidy (Manufacturer's Price) \$	Subsid Per	dised	Brand or Generic Manufacturer	
continued					
1.6.2 A sign of that new inflamr 1.6.3 A sign of that new inflamr 1.6.4 A sign of that new inflamr features of a recent attact	le sclerosis treatments simultaneously i marate, fingolimod, glatiramer acetat from any relevant practitioner. Approv or without the use of unilateral or bilater n or without aids in the last six months).	cal swelling; that clearly or red with a p ressive MS is not permi te, interfero rals valid for al aids at an	; or is response previous tted. on beta- r 12 mon ny time	MRI scan; or  -1-alpha, interferor nths where patient I	<b>n</b> has
DIMETHYL FUMARATE - Special Authority se	ee SA2274 on the previous page – Reta	ail pharmacy	y		
a) Wastage claimable     b) Note: Treatment on two or more funde     Cap 120 mg Cap 240 mg	520.00	eously is no 14 56	✓ Ted	itted. cfidera cfidera	
FINGOLIMOD – Special Authority see SA2274  a) Wastage claimable b) Note: Treatment on two or more funde Cap 0.5 mg	d multiple sclerosis treatments simultan		ot permi		
GLATIRAMER ACETATE – Special Authority s Note: Treatment on two or more funded m Inj 40 mg prefilled syringe	see SA2274 on the previous page – Ret nultiple sclerosis treatments simultaneou		cy ermitted	•	
NTERFERON BETA-1-ALPHA — Special Auth Note: Treatment on two or more funded m Inj 6 million iu prefilled syringe	nultiple sclerosis treatments simultaneou			d.	
NTERFERON BETA-1-BETA – Special Autho Note: Treatment on two or more funded m Inj 8 million iu per 1 ml	nultiple sclerosis treatments simultaneou		ermitted	i. taferon	
NATALIZUMAB – Special Authority see SA227 Note: Treatment on two or more funded m Inj 20 mg per ml, 15 ml vial	nultiple sclerosis treatments simultaneou		ermitted  Tys		
TERIFLUNOMIDE – Special Authority see SA2  a) Wastage claimable b) Note: Treatment on two or more funde		•	ot narmi	itted	
Tab 14 mg		28	✓ Ter	r <u>iflunomide</u> Sandoz	

### **Multiple Sclerosis Treatments - Other**

OCRELIZUMAB - Special Authority see SA2273 on the next page - Retail pharmacy

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

Inj 30 mg per ml, 10 ml vial	8,450.00	1	✓ Ocrevus
Inj 40 mg per ml, 23 ml vial	16,900.00	1	<ul><li>Ocrevus SC</li></ul>



Subsidy (Manufacturer's Price)

Fully Subsidised

Per

Brand or Generic Manufacturer

⇒SA2273 Special Authority for Subsidy

Initial application — (Multiple Sclerosis - ocrelizumab) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

### Either:

- 1 All of the following:
  - 1.1 Diagnosis of multiple sclerosis (MS) meets the McDonald 2017 diagnostic criteria for MS and has been confirmed by a neurologist; and
  - 1.2 Patient has an EDSS score between 0 6.0; and
  - 1.3 Patient has had at least one significant attack of MS in the previous 12 months or two significant attacks in the past 24 months: and
  - 1.4 All of the following:
    - 1.4.1 Each significant attack must be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the attack, but the neurologist/physician must be satisfied that the clinical features were characteristic): and
    - 1.4.2 Each significant attack is associated with characteristic new symptom(s)/sign(s) or substantially worsening of previously experienced symptoms(s)/sign(s); and
    - 1.4.3 Each significant attack has lasted at least one week and has started at least one month after the onset of a previous attack (where relevant); and
    - 1.4.4 Each significant attack can be distinguished from the effects of general fatigue; and is not associated with a fever (T> 37.5°C); and
    - 1.4.5 Fither:
      - 1.4.5.1 Each significant attack is severe enough to change either the EDSS or at least one of the Kurtze Functional System scores by at least 1 point; or
      - 1.4.5.2 Each significant attack is a recurrent paroxysmal symptom of multiple sclerosis (tonic seizures/spasms, trigeminal neuralgia, Lhermitte's symptom); and
  - 1.5 Evidence of new inflammatory activity on an MRI scan within the past 24 months; and
  - 1.6 Any of the following:
    - 1.6.1 A sign of that new inflammatory activity on MRI scanning (in criterion 5 immediately above) is a gadolinium enhancing lesion; or
    - 1.6.2 A sign of that new inflammatory activity is a lesion showing diffusion restriction; or
    - 1.6.3 A sign of that new inflammatory is a T2 lesion with associated local swelling; or
    - 1.6.4 A sign of that new inflammatory activity is a prominent T2 lesion that clearly is responsible for the clinical features of a recent attack that occurred within the last 2 years; or
    - 1.6.5 A sign of that new inflammatory activity is new T2 lesions compared with a previous MRI scan; or
- 2 Patient has an active Special Authority approval for either dimethyl fumarate, fingolimod, glatiramer acetate, interferon beta-1-alpha, interferon beta-1-beta, natalizumab or teriflunomide.

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

Renewal — (Multiple Sclerosis - ocrelizumab) from any relevant practitioner. Approvals valid for 12 months where patient has had an EDSS score of 0 to 6.0 (inclusive) with or without the use of unilateral or bilateral aids at any time in the last six months (ie the patient has walked 100 metres or more with or without aids in the last six months).

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

Initial application — (Primary Progressive Multiple Sclerosis) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Diagnosis of primary progressive multiple sclerosis (PPMS) meets the 2017 McDonald criteria and has been confirmed by a neurologist; and
- 2 Patient has an EDSS 2.0 (score equal to or greater than 2 on pyramidal functions) to EDSS 6.5; and

### **NERVOUS SYSTEM**

Subsidy		Fully	Brand or	
(Manufacturer's Price)		ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

3 Patient has no history of relapsing remitting multiple sclerosis.

Renewal — (Primary Progressive Multiple Sclerosis) from any relevant practitioner. Approvals valid for 12 months where patient has had an EDSS score of less than or equal to 6.5 at any time in the last six months (ie patient has walked 20 metres with bilateral assistance/aids, without rest in the last six months).

### **Sedatives and Hypnotics**

### ⇒SA2523 Special Authority for Subsidy

Initial application only from a psychiatrist, paediatrician, neurologist, respiratory specialist or any relevant practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

#### All of the following:

- 1 Patient has been diagnosed with persistent and distressing insomnia secondary to a neurodevelopmental disorder (including, but not limited to, autism spectrum disorder or attention deficit hyperactivity disorder)\*; and
- 2 Behavioural and environmental approaches have been tried and were unsuccessful, or are inappropriate; and
- 3 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day; and
- 4 Patient is aged 18 years or under\*.

**Renewal** only from a psychiatrist, paediatrician, neurologist, respiratory specialist or any relevant practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

#### All of the following:

- 1 Patient is aged 18 years or under\*; and
- 2 Patient has demonstrated clinically meaningful benefit from funded modified-release melatonin (clinician determined); and
- 3 Patient has had a trial of funded modified-release melatonin discontinuation within the past 12 months and has had a recurrence of persistent and distressing insomnia; and
- 4 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day.

Note: Indications marked with \* are unapproved indications.

MIDAZOLAM - Safety medicine; prescriber may determine disper	nsing frequency		
Inj 1 mg per ml, 5 ml ampoule	7.80	10	✓ Midazolam-Baxter
Inj 1 mg per ml, 5 ml plastic ampoule - Up to 10 inj available			
on a PSO		10	<ul><li>Midazolam-Pfizer</li></ul>
On a PSO for status epilepticus use only. PSO must be e	endorsed for status	epilepticus ι	use only.
Inj 5 mg per ml, 1 ml plastic ampoule - Up to 10 inj available			
on a PSO	22.50	10	<ul><li>Midazolam-Pfizer</li></ul>
On a PSO for status epilepticus use only. PSO must be e	endorsed for status	epilepticus ι	
Inj 5 mg per ml, 3 ml ampoule	4.75	5	✓ Midazolam-Baxter
Inj 5 mg per ml, 3 ml plastic ampoule – Up to 5 inj available o	n		
a PSO		5	<ul><li>Midazolam-Pfizer</li></ul>
On a PSO for status epilepticus use only. PSO must be e	endorsed for status	epilepticus ι	use only.
PHENOBARBITONE SODIUM - Special Authority see SA1386 or	n the next page – R	etail pharma	асу
Inj 200 mg per ml, 1 ml ampoule	113.37	10	✓ Max Health S29



Subsidy	,	Fully	Brand or	
(Manufacturer's Pric	e)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

### **⇒SA1386** Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Both:

- 1 For the treatment of terminal agitation that is unresponsive to other agents; and
- 2 The applicant is part of a multidisciplinary team working in palliative care.

TEMAZEPAM – Safety medicine; prescriber may determine dispe Tab 10 mg	 25	✓ Normison
ZOPICLONE – Safety medicine; prescriber may determine disper	20	Normison
Tab 7.5 mg	 500	✓ Zopiclone Actavis

### **Spinal Muscular Atrophy**

NUSINERSEN − PCT only − Special Authority see SA2174 below
Inj 12 mg per 5 ml vial .......120,000.00 1 ✓ Spinraza

### ⇒SA2174 Special Authority for Subsidy

Initial application — (spinal muscular atrophy (SMA)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

### All of the following:

- 1 Patient has genetic documentation of homozygous SMN1 gene deletion, homozygous SMN1 point mutation, or compound heterozygous mutation; and
- 2 Patient is 18 years of age or under; and
- 3 Either
  - 3.1 Patient has experienced the defined signs and symptoms of SMA type I, II or IIIa prior to three years of age; or
  - 3.2 Both:
    - 3.2.1 Patient is pre-symptomatic; and
    - 3.2.2 Patient has three or less copies of SMN2.

Renewal — (spinal muscular atrophy (SMA)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

#### All of the following:

- 1 There has been demonstrated maintenance of motor milestone function since treatment initiation; and
- 2 Patient does not require invasive permanent ventilation (at least 16 hours per day) in the absence of a potentially reversible cause while being treated with nusinersen; and
- 3 Nusinersen not to be administered in combination other SMA disease modifying treatments or gene therapy.

### RISDIPLAM - [Xpharm] - Special Authority see SA2203 below

Note: the supply of risdiplam is via Pharmac's approved direct distribution supply. Further details can be found on Pharmac's website https://pharmac.govt.nz/risdiplam

Powder for oral soln 750 mcg per ml, 60 mg per bottle......14,100.00 80 ml OP ✓ Evrysdi

### ⇒SA2203 Special Authority for Subsidy

**Initial application** — (spinal muscular atrophy (SMA)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

#### All of the following:

- 1 Patient has genetic documentation of homozygous SMN1 gene deletion, homozygous SMN1 point mutation, or compound heterozygous mutation; and
- 2 Patient is 18 years of age or under; and
- 3 Either:

Subsidy	Ful	y Brand or
(Manufacturer's	Price) Subsidise	d Generic
\$	Per •	Manufacturer

continued...

- 3.1 Patient has experienced the defined signs and symptoms of SMA type I, II or IIIa prior to three years of age; or
- 3.2 Both:
  - 3.2.1 Patient is pre-symptomatic; and
  - 3.2.2 Patient has three or less copies of SMN2.

Renewal — (spinal muscular atrophy (SMA)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 There has been demonstrated maintenance of motor milestone function since treatment initiation; and
- 2 Patient does not require invasive permanent ventilation (at least 16 hours per day) in the absence of a potentially reversible cause while being treated with risdiplam; and
- 3 Risdiplam not to be administered in combination other SMA disease modifying treatments or gene therapy.

# Stimulants/ADHD Treatments

ATOMOXETINE			
Cap 10 mg	43.02	28	✓ APO-Atomoxetine
Cap 18 mg	45.57	28	✓ APO-Atomoxetine
Cap 25 mg	44.30	28	✓ APO-Atomoxetine
Cap 40 mg	46.21	28	✓ APO-Atomoxetine
Cap 60 mg	51.31	28	✓ APO-Atomoxetine
Cap 80 mg	65.20	28	✓ APO-Atomoxetine
Cap 100 mg	65.71	28	✓ APO-Atomoxetine
DEXAMFETAMINE SULFATE - Special Authority see SA24	10 below – Retail pha	rmacy	
a) Only on a controlled drug form			
b) Safety medicine; prescriber may determine dispensing	g frequency		
Tab 5 mg	29.80	100	✓ Noumed Dexamfetamine

#### ⇒SA2410 Special Authority for Subsidy

Initial application — (ADHD in patients aged 5 years or over) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid without further renewal unless notified for applications meeting the following criteria: All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) in patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
  - 3.1 Applicant is a paediatrician or psychiatrist; or
  - 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients aged under 5 years) only from a paediatrician or psychiatrist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) in patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

**Initial application** — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid without further renewal unless notified where the patient suffers from narcolepsy.

### **NERVOUS SYSTEM**

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sı	ubsidised	Generic
\$	Per	✓	Manufacturer

LISDEXAMFETAMINE DIMESILATE - Special Authority see SA2415 below - Retail pharmacy

- a) Only on a controlled drug form

### ⇒SA2415 Special Authority for Subsidy

Initial application only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid without further renewal unless notified for applications meeting the following criteria:

Fither:

- 1 Patient is currently on treatment with lisdexamfetamine dimesilate and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
  - 2.1 ADHD (Attention Deficit and Hyperactivity Disorder); and
  - 2.2 Diagnosed according to DSM-V or ICD 11 criteria; and
  - 2.3 Either:
    - 2.3.1 Applicant is a paediatrician or psychiatrist; or
    - 2.3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing; and
  - 2.4 Any of the following:
    - 2.4.1 Patient is taking a currently subsidised formulation of atomoxetine or methylphenidate hydrochloride (extended-release) and has not received sufficient benefit or has experienced intolerable side effects; or
    - 2.4.2 Patient is taking a currently subsidised formulation of dexamfetamine sulfate (immediate-release) which has not been effective due to significant administration and/or treatment adherence difficulties; or
    - 2.4.3 There is significant concern regarding the risk of diversion or abuse of immediate release dexamfetamine sulfate; or
    - 2.4.4 Patient is taking a currently subsidised formulation of methylphenidate hydrochloride (immediate-release or sustained release) which has not been effective due to significant administration and/or treatment adherence difficulties; or
    - 2.4.5 There is significant concern regarding the risk of diversion or abuse of immediate release methylphenidate hydrochloride; or
    - 2.4.6 Both:
      - 2.4.6.1 Patient would have been prescribed a subsidised formulation of methylphenidate hydrochloride (extended-release) but has been unable to access due to supply issues with methylphenidate hydrochloride (extended-release); and
      - 2.4.6.2 Other alternative stimulant presentations (methylphenidate or dexamfetamine) are not appropriate; and
  - 2.5 Lisdexamfetamine dimesilate is not to be used in combination with another funded methylphenidate presentation.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
METHYLPHENIDATE HYDROCHLORIDE – Special Authority s a) Only on a controlled drug form	see SA2546 below – R	letail	pharmacy	
b) Safety medicine; prescriber may determine dispensing for	requency			
Tab modified-release 18 mg	15.25	30	•	Methylphenidate Sandoz XR
Tab modified-release 27 mg	16.25	30	•	Methylphenidate Sandoz XR
Tab modified-release 36 mg	21.25	30	•	Methylphenidate Sandoz XR
Tab modified-release 54 mg	24.25	30	•	Methylphenidate Sandoz XR
Tab immediate-release 5 mg	3.20	30	1	Rubifen
Tab immediate-release 10 mg		30	✓	Rubifen
<b>o</b>	4.00		✓	Ritalin
Tab extended-release 18 mg	15.25	30	•	Methylphenidate ER - Teva
Tab immediate-release 20 mg	7.85	30	✓	Rubifen
Tab sustained-release 20 mg		30	✓	Rubifen SR
Tab extended-release 27 mg		30	✓	Methylphenidate ER - Teva
Tab extended-release 36 mg	21.25	30	•	Methylphenidate ER - Teva
Tab extended-release 54 mg	24.25	30	✓	Methylphenidate ER - Teva

#### ⇒SA2546 Special Authority for Subsidy

Initial application — (ADHD in patients aged 5 years or over) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid without further renewal unless notified for applications meeting the following criteria: All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) in patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
  - 3.1 Applicant is a paediatrician or psychiatrist; or
  - 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients aged under 5 years) only from a paediatrician or psychiatrist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) in patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy\*) only from a neurologist or respiratory specialist. Approvals valid without further renewal unless notified where the patient suffers from narcolepsy.

Note: \*narcolepsy is not a registered indication for Methylphenidate ER – Teva or Methylphenidate Sandoz XR.



Subsidy		ully	Brand or	
(Manufacturer's Price)	Subsid	ised	Generic	
\$	Per	1	Manufacturer	

METHYLPHENIDATE HYDROCHLORIDE EXTENDED-RELEASE - Special Authority see SA2450 below - Retail pharmacy

- a) Only on a controlled drug form
- b) Safety medicine; prescriber may determine dispensing frequency

b) Galety incalcine, prescriber may actermine dispersing i	i oquonoy		
Tab extended-release 18 mg	58.96	30	<ul><li>Concerta</li></ul>
Tab extended-release 27 mg	65.44	30	<ul><li>Concerta</li></ul>
Tab extended-release 36 mg		30	<ul><li>Concerta</li></ul>
Tab extended-release 54 mg	86.24	30	<ul><li>Concerta</li></ul>
Cap modified-release 10 mg		30	Ritalin LA
Cap modified-release 20 mg	27.72	30	Ritalin LA
Cap modified-release 30 mg		30	Ritalin LA
Cap modified-release 40 mg		30	Ritalin LA

#### ⇒SA2450 Special Authority for Subsidy

Initial application — (ADHD) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 All of the following:
  - 1.1 ADHD (Attention Deficit and Hyperactivity Disorder); and
  - 1.2 Diagnosed according to DSM-IV or ICD 10 criteria; and
  - 1.3 Either:
    - 1.3.1 Applicant is a paediatrician or psychiatrist; or
    - 1.3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing; and
  - 1.4 Either:
    - 1.4.1 Patient is taking a currently subsidised formulation of methylphenidate hydrochloride (immediate-release or sustained-release) which has not been effective due to significant administration and/or difficulties with adherence; or
    - 1.4.2 There is significant concern regarding the risk of diversion or abuse of immediate-release methylphenidate hydrochloride; or
- 2 Both:
  - 2.1 Patient meets the Special Authority criteria for SA2411 methylphenidate hydrochloride; and
  - 2.2 Patient is unable to access other methylphenidate hydrochloride presentations under Special Authority criteria SA2411 due to an out of stock (see note).

Note: Criterion 2 is to permit short-term funding to cover an out-of-stock on tab extended-release Methylphenidate ER – Teva and tab sustained-release 20 mg Rubifen SR subsidised under

SA2411 (https://schedule.pharmac.govt.nz/2025/02/01/SA2411.pdf).

**Initial application** — (Narcolepsy\*) only from a neurologist or respiratory specialist. Approvals valid without further renewal unless notified where the patient suffers from narcolepsy.

Note: \*narcolepsy is not a registered indication for Concerta or Ritalin LA.

MODAFINIL - Special Authority see SA2451 below - Retail pharmacy

# ⇒SA2451 Special Authority for Subsidy

Initial application only from a neurologist or respiratory specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Fither:

1 All of the following:

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsid	dised	Generic
\$	Per	1	Manufacturer

#### continued...

- 1.1 The patient has a diagnosis of narcolepsy and has excessive daytime sleepiness associated with narcolepsy occurring almost daily for three months or more; and
- 1.2 Either:
  - 1.2.1 The patient has a multiple sleep latency test with a mean sleep latency of less than or equal to 10 minutes and 2 or more sleep onset rapid eye movement periods; or
  - 1.2.2 The patient has at least one of: cataplexy, sleep paralysis or hypnagogic hallucinations; and
- 1.3 Either:
  - 1.3.1 An effective dose of a subsidised formulation of methylphenidate or dexamfetamine has been trialled and discontinued because of intolerable side effects: or
  - 1.3.2 Methylphenidate and dexamfetamine are contraindicated; or
- 2 Both:
  - 2.1 Patient meets the Special Authority criteria for methylphenidate hydrochloride or methylphenidate hydrochloride extended-release for narcolepsy; and
  - 2.2 Patient is unable to access methylphenidate hydrochloride presentations due to an out of stock (see note).

Note: Criterion 2 is to permit short-term funding to cover an out-of-stock of methylphenidate hydrochloride or methylphenidate hydrochloride extended release.

### **Treatments for Dementia**

DONEPEZIL HYDROCHLORIDE			
* Tab 5 mg	3.70	84	✓ <u>Ipca-Donepezil</u>
* Tab 10 mg	5.50	84	✓ Ipca-Donepezil
RIVASTIGMINE - Special Authority see SA2524	below - Retail pharmacy		
Patch 4.6 mg per 24 hour	49.40	30	✓ Rivastigmine Patch
			<u>BNM 5</u>
Patch 9.5 mg per 24 hour	49.40	30	<ul> <li>Rivastigmine Patch</li> </ul>
			BNM 10

#### ⇒SA2524 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The patient has been diagnosed with dementia; and
- 2 The patient is contraindicated to or has experienced intolerable side effects from donepezil tablets.

**Renewal** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate; and
- 2 The patient has demonstrated a significant and sustained benefit from treatment.

# Treatments for Substance Dependence

BUPRENORPHINE WITH NALOXONE - Special Authority see SA1203 on the next page - Retail pharmacy

- a) No patient co-payment payable
- b) Safety medicine; prescriber may determine dispensing frequency

Tab sublingual 2 mg with naloxone 0.5 mg ......11.76

✓ Buprenorphine 
Naloxone BNM

Tab sublingual 8 mg with naloxone 2 mg .......26.86

✓ Buprenorphine Naloxone BNM



Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

## ⇒SA1203 Special Authority for Subsidy

**Initial application — (Detoxification)** from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health...

**Initial application — (Maintenance treatment)** from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient will not be receiving methadone; and
- 3 Patient is currently enrolled in an opioid substitution treatment program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient has previously trialled but failed detoxification with buprenorphine with naloxone with relapse back to opioid use and another attempt is planned; and
- 3 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

**Renewal — (Maintenance treatment)** from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is or has been receiving maintenance therapy with buprenorphine with naloxone (and is not receiving methadone); and
- 2 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health or is a medical practitioner authorised by the service to manage treatment in this patient.

Renewal — (Maintenance treatment where the patient has previously had an initial application for detoxification) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient received but failed detoxification with buprenorphine with naloxone; and
- 2 Maintenance therapy with buprenorphine with naloxone is planned (and patient will not be receiving methadone); and
- 3 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Tab modified-release 150 mg	15.00	30	✓ <u>Zyban</u>
DISULFIRAM Tab 200 mg	236.40	100	✓ Antabuse
NALTREXONE HYDROCHLORIDE – Special Authority see SA Tab 50 mg		•	

Sub	osidy Fu	ully Brand or
(Manufactu	urer's Price) Subsidis	sed Generic
	\$ Per	<ul> <li>Manufacturer</li> </ul>

### ⇒SA1408 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient is currently enrolled in a recognised comprehensive treatment programme for alcohol dependence; and
- 2 Applicant works in or with a community Alcohol and Drug Service contracted to Health NZ or accredited against the New Zealand Alcohol and Other Drug Sector Standard or the National Mental Health Sector Standard.

Renewal from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Compliance with the medication (prescriber determined); and
- 2 Any of the following:
  - 2.1 Patient is still unstable and requires further treatment; or
  - 2.2 Patient achieved significant improvement but requires further treatment; or
  - 2.3 Patient is well controlled but requires maintenance therapy.

#### NICOTINE

a) Nicotine will not be funded in amounts less than 4 weeks of treatment.

b) Note: Direct Provision by a pharmacist permitted under the provisions	in Part I of Sec	ction A.
Patch 7 mg - Up to 28 patch available on a PSO19.62	28	✓ Habitrol
Patch 14 mg – Up to 28 patch available on a PSO21.57	28	<ul><li>Habitrol</li></ul>
Patch 14 mg for direct distribution only - [Xpharm]12.49	7	<ul><li>Habitrol</li></ul>
Patch 21 mg - Up to 28 patch available on a PSO24.72	28	<ul><li>Habitrol</li></ul>
Patch 21 mg for direct distribution only - [Xpharm]13.19	7	<ul><li>Habitrol</li></ul>
Lozenge 1 mg - Up to 216 loz available on a PSO22.53	216	<ul><li>Habitrol</li></ul>
Lozenge 1 mg for direct distribution only - [Xpharm]12.89	36	Habitrol
Lozenge 2 mg - Up to 216 loz available on a PSO24.68	216	Habitrol
Lozenge 2 mg for direct distribution only $-[Xpharm]$ 13.25	36	Habitrol
Gum 2 mg (Fruit) - Up to 204 piece available on a PSO23.02	204	Habitrol
Gum 2 mg (Fruit) for direct distribution only - [Xpharm]17.57	96	Habitrol
Gum 2 mg (Mint) - Up to 204 piece available on a PSO23.02	204	<ul><li>Habitrol</li></ul>
Gum 2 mg (Mint) for direct distribution only - [Xpharm]17.57	96	<ul><li>Habitrol</li></ul>
Gum 4 mg (Fruit) - Up to 204 piece available on a PSO25.98	204	<ul><li>Habitrol</li></ul>
Gum 4 mg (Fruit) for direct distribution only - [Xpharm]23.87	96	<ul><li>Habitrol</li></ul>
Gum 4 mg (Mint) - Up to 204 piece available on a PSO25.98	204	<ul><li>Habitrol</li></ul>
Gum 4 mg (Mint) for direct distribution only - [Xpharm]23.87	96	<ul><li>Habitrol</li></ul>

VARENICLINE TARTRATE - Special Authority see SA1845 on the next page - Retail pharmacy

- a) A maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval, including the starter pack
- b) Varenicline will not be funded in amounts less than 4 weeks of treatment.
- c) The 6-month time period in which a patient can receive a funded 12-week course of varenicline tartrate starts from the date the Special Authority is approved.

Tab 0.5 mg × 11 and 1 mg × 421	15.99	53 OP	✓ Pharmacor Varenicline
Tab 1 mg	16.67 10.99	56	<ul><li>✓ Champix</li><li>✓ Pharmacor</li><li>✓ Varenicline</li></ul>
1	17.62		✓ Champix

(Champix Tab 0.5 mg × 11 and 1 mg × 42 to be delisted 1 June 2026) (Champix Tab 1 mg to be delisted 1 June 2026)



Subsidy (Manufacturer's Price)	F Subsidi	ully	Brand or Generic
 \$	Per	<b>√</b>	Manufacturer

## **⇒SA1845** Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 Either:
  - 3.1 The patient has tried but failed to quit smoking after at least two separate trials of nicotine replacement therapy, at least one of which included the patient receiving comprehensive advice on the optimal use of nicotine replacement therapy; or
  - 3.2 The patient has tried but failed to quit smoking using bupropion or nortriptyline; and
- 4 The patient has not had a Special Authority for varenicline approved in the last 6 months; and
- 5 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 6 The patient is not pregnant; and
- 7 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

**Renewal** from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 It has been 6 months since the patient's previous Special Authority was approved; and
- 4 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 5 The patient is not pregnant; and
- 6 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

The patient must not have had an approval in the past 6 months.

Notes: a maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval.

This includes the 4-week 'starter' pack.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

# **Chemotherapeutic Agents**

# Alkylating Agents

BENDAMUSTINE HYDROCHLORIDE - PCT only - Spe	ecialist - Special Authority	see SA2398	3 below
Inj 25 mg vial	50.05	1	<ul> <li>Bendamustine</li> </ul>
			Sandoz
	77.00		✓ Ribomustin
Inj 100 mg vial	200.20	1	<ul> <li>Bendamustine</li> </ul>
			Sandoz
	308.00		✓ Ribomustin
Inj 1 mg for ECP	2.11	1 mg	✓ Baxter

#### ⇒SA2398 Special Authority for Subsidy

Initial application — (CLL\*) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

- All of the following:
  - 1 The patient has chronic lymphocytic leukaemia requiring treatment; and
  - 2 Patient has ECOG performance status of 0-2; and
  - 3 Bendamustine is to be administered at a maximum dose of 100 mg/m² on days 1 and 2 every 4 weeks for a maximum of 6 cycles.

Note: Indication marked with a \* includes indications that are unapproved. 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL).

Initial application — (Indolent, Low-grade lymphomas) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has indolent low grade NHL requiring treatment; and
- 2 The patient has ECOG performance status of 0-2; and
- 3 Any of the following:
  - 3.1 Both:
    - 3.1.1 Patient is treatment naive; and
    - 3.1.2 Bendamustine is to be administered for a maximum of 6 cycles (in combination with rituximab when CD20+); or
  - 3.2 Both:
    - 3.2.1 Patient is refractory to or has relapsed within 12 months of a rituximab containing combined chemo-immunotherapy regimen; and
    - 3.2.2 Bendamustine is to be administered in combination with obinutuzumab for a maximum of 6 cycles; or
  - 3.3 All of the following:
    - 3.3.1 The patient has not received prior bendamustine therapy; and
    - 3.3.2 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
    - 3.3.3 Patient has had a rituximab treatment-free interval of 12 months or more; or
  - 3.4 Bendamustine is to be administered as monotherapy for a maximum of 6 cycles in rituximab refractory patients.

**Renewal — (Indolent, Low-grade lymphomas)** only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: Either:

- 1 Both:
  - 1.1 Patient is refractory to or has relapsed within 12 months of rituximab in combination with bendamustine; and

Subsidy	Fully	Brand or	_
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

- 1.2 Bendamustine is to be administered in combination with obinutuzumab for a maximum of 6 cycles; or
- 2 Both:
  - 2.1 Patients have not received a bendamustine regimen within the last 12 months; and
  - 2.2 Either:
    - 2.2.1 Both:
      - 2.2.1.1 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
      - 2.2.1.2 Patient has had a rituximab treatment-free interval of 12 months or more: or
    - 2.2.2 Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Note: 'indolent, low-grade lymphomas' includes follicular, mantle cell, marginal zone and lymphoplasmacytic/ Waldenstrom's macroglobulinaemia.

**Initial application** — (Hodgkin's lymphoma\*) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has Hodgkin's lymphoma requiring treatment; and
- 2 Patient has a ECOG performance status of 0-2; and
- 3 Patient has received one prior line of chemotherapy; and
- 4 Patient's disease relapsed or was refractory following prior chemotherapy; and
- 5 Bendamustine is to be administered in combination with gemcitabine and vinorelbine (BeGeV) at a maximum dose of no greater than 90 mg/m2 twice per cycle, for a maximum of four cycles.

Note: Indications marked with \* are unapproved indications.

PURIL EAN DOT Potail pharmacy Charielist

BUSULFAN – PCT – Retail pharmacy-Specialist			
Tab 2 mg	89.25	100	✓ Myleran
CARBOPLATIN - PCT only - Specialist			
Inj 10 mg per ml, 45 ml vial	25.73	1	✓ Carboplatin Accord
, ,	32.59		✓ DBL Carboplatin
	48.50		✓ Carbaccord
Inj 1 mg for ECP	0.06	1 mg	✓ Baxter
CARMUSTINE - PCT only - Specialist		_	
Inj 100 mg vial	710.00	1	✓ BiCNU
Inj 100 mg for ECP		100 mg OP	✓ Baxter
CHLORAMBUCIL – PCT – Retail pharmacy-Specialist		-	
Tab 2 mg	29.06	25	✓ Leukeran FC
CISPLATIN - PCT only - Specialist			
Inj 1 mg per ml, 50 ml vial	9.45	1	<ul> <li>Cisplatin Accord</li> </ul>
	15.00		<ul><li>Cisplatin Ebewe</li></ul>
Inj 1 mg per ml, 100 ml vial	18.90	1	✓ Cisplatin Accord
	21.00		<ul><li>Cisplatin Ebewe</li></ul>
	29.66		✓ DBL Cisplatin
Inj 1 mg for ECP	0.19	1 mg	✓ Baxter
CYCLOPHOSPHAMIDE			
Tab 50 mg - PCT - Retail pharmacy-Specialist	145.00	50	✓ Cyclonex
Inj 1 g vial - PCT - Retail pharmacy-Specialist		1	✓ Endoxan
	127.80	6	✓ Cytoxan
Inj 2 g vial - PCT only - Specialist		1	✓ Endoxan
Inj 1 mg for ECP - PCT only - Specialist	0.05	1 mg	✓ Baxter

	Subsidy (Manufacturer's Price)	Su	Fully bsidised	
	\$	Per	✓	Manufacturer
FOSFAMIDE - PCT only - Specialist				
Inj 1 g	96.00	1	✓	Holoxan
Inj 2 g		1	✓	Holoxan
Inj 1 mg for ECP	0.10	1 mg	✓	Baxter
OMUSTINE - PCT - Retail pharmacy-Specialist				
Cap 40 mg	880.00	20	1	Medac S29
/ELPHALAN				
Tab 2 mg - PCT - Retail pharmacy-Specialist	40.70	25	1	Alkeran
Inj 50 mg - PCT only - Specialist	48.25	1	1	Melpha
, , , ,	67.80		1	Alkeran
DXALIPLATIN - PCT only - Specialist				
Inj 100 mg vial	25.01	1	1	Oxaliplatin Actavis
,				100 <sup>°</sup>
	110.00		1	Oxaliplatin Ebewe
Inj 5 mg per ml, 20 ml vial	33.35	1	✓	Alchemy Oxaliplatin
	46.32		✓	Oxaliplatin Accord
Inj 1 mg for ECP	0.35	1 mg	✓	Baxter
THIOTEPA - PCT only - Specialist				
Inj 15 mg vial	CBS	1	1	Bedford S29
, 0			1	Max Health \$29
			1	THIO-TEPA S29
	398.00			Tepadina
Inj 100 mg vial		1		Max Health S29
11, 100 11g 101	1,800.00	•		Tepadina
	.,			
Antimetabolites				
AZACITIDINE - PCT only - Specialist - Special Authority s	see SA2479 below			
Inj 100 mg vial	50.00	1	1	Azacitidine Dr

		AZACITIDINE - PCT only - Specialist - Special Authority see SA2479 below
<ul> <li>Azacitidine Dr Reddy's</li> </ul>	1	Inj 100 mg vial50.00
✓ Baxter	1 mg	Inj 1 mg for ECP0.54

## ⇒SA2479 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 Any of the following:
  - 1.1 The individual has intermediate or high risk MDS based on an internationally recognised scoring system; or
  - 1.2 The individual has chronic myelomonocytic leukaemia (based on an intermediate or high risk score from an internationally recognised scoring system or 10%-29% marrow blasts without myeloproliferative disorder); or
  - 1.3 The individual has acute myeloid leukaemia according to World Health Organisation Classification (WHO); and
- 2 The individual has an estimated life expectancy of at least 3 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where there is no evidence of disease progression.

	Subsidy		Fully Brand or
(1	Manufacturer's Pric	e) S Per	Subsidised Generic  Manufacturer
ALCIUM FOLINATE	Ψ	1 01	· Manadataror
Tab 15 mg - PCT - Retail pharmacy-Specialist	135.33	10	✓ DBL Leucovorin
rab 13 mg = 1 01 = Hetali phamiacy-specialist	100.00	10	Calcium
Inj 3 mg per ml, 1 ml - PCT - Retail pharmacy-Specialist	17 10	5	✓ Hospira
Inj 10 mg per ml, 5 ml vial – PCT – Retail pharmacy-Specialist		5	✓ Eurofolic
Inj 50 mg – PCT – Retail pharmacy-Specialist		10	✓ Leucovorin
,			Pharmacia \$29
Inj 10 mg per ml, 10 ml vial - PCT only - Specialist	163 35	5	✓ Eurofolic
Inj 100 mg - PCT only - Specialist		10	✓ Leucovorin
ing rooming in our only openium	04.00	10	Pharmacia S29
Ini 200 mg BCT only Chapitalist	01 55	4	✓ Leucovorin DBL S29
Inj 300 mg - PCT only - Specialist	21.33	1	▼ Leucovoriii DBL 328
Inj 10 mg per ml, 100 ml vial - PCT only - Specialist	139.48	1	✓ Eurofolic
Inj 1 mg for ECP - PCT only - Specialist		1 mg	✓ Baxter
		9	
APECITABINE – Retail pharmacy-Specialist	10.02	60	✓ Capecitabine Viatri
Tab 150 mg		60	• Capecitabilie viatri
		120	✓ Canacitahina Viatri
Tab 500 mg		120	Capecitabine Viatri
	2020		
LADRIBINE - PCT only - Specialist	740.00		
Inj 1 mg per ml, 10 ml		1	✓ Leustatin
Inj 10 mg for ECP	749.96	10 mg O	P Saxter
YTARABINE			
Inj 20 mg per ml, 5 ml vial - PCT - Retail pharmacy-Specialist	472.00	5	✓ Pfizer
Inj 100 mg per ml, 20 ml vial – PCT – Retail			
pharmacy-Specialist	48.80	1	<ul><li>Cytarabine DBL</li></ul>
			✓ Pfizer
Inj 1 mg for ECP - PCT only - Specialist		10 mg	✓ Baxter
Inj 100 mg intrathecal syringe for ECP - PCT only - Specialist	94.40 1	100 mg C	OP <b>✓ Baxter</b>
LUDARABINE PHOSPHATE			
Tab 10 mg - PCT - Retail pharmacy-Specialist	412.00	20	Fludara Oral
Inj 50 mg vial - PCT only - Specialist	634.00	5	Fludarabine Ebewe
Inj 50 mg for ECP - PCT only - Specialist	126.80	50 mg O	P Saxter
LUOROURACIL			
Inj 50 mg per ml, 20 ml vial - PCT only - Specialist	10.51	1	✓ Fluorouracil Accord
Inj 50 mg per ml, 50 ml vial - PCT only - Specialist		1	✓ Fluorouracil Accord
Inj 50 mg per ml, 100 ml vial - PCT only - Specialist	19.36	1	✓ Fluorouracil Accord
Inj 1 mg for ECP - PCT only - Specialist	0.41	100 mg	✓ Baxter
EMCITABINE HYDROCHLORIDE - PCT only - Specialist		Ü	
Inj 43.3 mg per ml (equivalent to 38 mg per ml gemcitabine),			
26.3 ml vial	18 94	1	✓ DBL Gemcitabine
Inj 1 g		1	✓ Gemcitabine Ebewe
Inj 1 mg for ECP		1 mg	✓ Baxter
RINOTECAN HYDROCHLORIDE - PCT only - Specialist		9	Buntoi
	50 F7	1	✓ Accord
Inj 20 mg per ml, 5 ml vial	52.57 71.44	ı	✓ Accord ✓ Irinotecan Actavis
	11.44		100
	100.00		
1:00	100.00		✓ Irinotecan-Rex
Inj 20 mg per ml, 25 ml vial		1	✓ Accord \$29
Inj 1 mg for ECP	0.54	1 mg	✓ Baxter

	Subsidy (Manufacturer's Price \$	e) Su Per	Fully bsidised	Brand or Generic Manufacturer	
MERCAPTOPURINE Tab 50 mg - PCT - Retail pharmacy-Specialist Oral suspension 20 mg per ml - Retail pharmacy-Specialist		25	<b>√</b> <u>P</u>	uri-nethol	
Special Authority see SA1725 below		00 ml OP		Ilmercap aluprine §29	

### ⇒SA1725 Special Authority for Subsidy

Initial application only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where the patient requires a total dose of less than one full 50 mg tablet per day.

Renewal only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where patient still requires

a to	otal dose of less than one full 50 mg tablet per day.		
	THOTREXATE		<b>.</b> .
*	Tab 2.5 mg - PCT - Retail pharmacy-Specialist	90	✓ <u>Trexate</u>
*	Tab 10 mg - PCT - Retail pharmacy-Specialist26.40	90	✓ <u>Trexate</u>
	Inj 2.5 mg per ml, 2 ml – PCT – Retail pharmacy-Specialist95.29	5	✓ Methotrexate DBL
*	Inj 7.5 mg prefilled syringe29.17	1	✓ <u>Methotrexate</u>
*	Inj 10 mg prefilled syringe19.09	1	Sandoz ✓ Methotrexate
			<u>Sandoz</u>
*	Inj 15 mg prefilled syringe24.53	1	✓ <u>Methotrexate</u> Sandoz
*	Inj 20 mg prefilled syringe16.64	1	✓ <u>Methotrexate</u> Sandoz
*	Inj 25 mg prefilled syringe20.72	1	✓ <u>Methotrexate</u> Sandoz
*	Inj 30 mg prefilled syringe55.00	1	✓ Methotrexate Sandoz
*	Inj 25 mg per ml, 2 ml vial - PCT - Retail pharmacy-Specialist30.00	5	✓ Methotrexate DBL Onco-Vial
*	Inj 25 mg per ml, 20 ml vial - PCT - Retail pharmacy-Specialist45.00	1	✓ DBL Methotrexate Onco-Vial
* *	Inj 100 mg per ml, 10 ml – PCT – Retail pharmacy-Specialist25.00 Inj 100 mg per ml, 50 ml vial – PCT – Retail	1	✓ Methotrexate Ebewe
	pharmacy-Specialist	1	✓ Methotrexate Ebewe
*	Inj 1 mg for ECP – PCT only – Specialist	1 mg	✓ Baxter
*	Inj 5 mg intrathecal syringe for ECP - PCT only - Specialist19.06	5 mg ÖP	✓ Baxter
PF	METREXED - PCT only - Specialist	ŭ	
	Inj 100 mg vial8.99	1	✓ Pemetrexed-AFT
	60.89		✓ Juno Pemetrexed
	Inj 500 mg vial	1	✓ Pemetrexed-AFT
	217.77		✓ Juno Pemetrexed
	Inj 1 mg for ECP	1 mg	✓ Baxter
TH	OGUANINE - PCT - Retail pharmacy-Specialist	ū	
	Tab 40 mg	25	✓ Lanvis
0	ther Cytotoxic Agents		
AM	SACRINE - PCT only - Specialist		
•	Inj 50 mg per ml, 1.5 ml ampoule	6	✓ Amsidine S29

AMSACRINE – PCT only – Specialist			
Inj 50 mg per ml, 1.5 ml ampoule	4,736.00	6	✓ Amsidine S29
Inj 75 mg	6,218.00	5	✓ AmsaLyo S29

	Subsidy		Fully	
	(Manufacturer's \$	Price) Subs	idised •	Generic Manufacturer
NAGRELIDE HYDROCHLORIDE - PCT - Retail pharmacy-S		7 01		a.a.a.a.a
Cap 0.5 mg		100	1	Agrylin
	1,173.07	100	٠	Agryiiii
RSENIC TRIOXIDE - PCT only - Specialist				
Inj 1 mg per ml, 10 ml vial		10		Phenasen
Inj 10 mg for ECP	481.70	10 mg OP	•	Baxter
LEOMYCIN SULPHATE - PCT only - Specialist				
Inj 15,000 iu, vial	185.16	1	1	DBL Bleomycin
				Sulfate
Inj 1,000 iu for ECP	14.32	1,000 iu	1	Baxter
ORTEZOMIB - PCT only - Specialist - Special Authority see	SA2355 below			
Inj 3.5 mg vial		1	1	DBL Bortezomib
Inj 1 mg for ECP		1 mg	1	Baxter
Special Authority for Subsidy		ŭ		
itial application — (plasma cell dyscrasia) from any releva	nt practitioner	Approvals valid v	withou	it further renewal unles
otified where the patient has plasma cell dyscrasia, not including				
ACARBAZINE - PCT only - Specialist	.5 .14.40.10.1011		oa,	. oquiig troutinoiiti
Inj 200 mg vial	70 11	1	1	DBL Dacarbazine
Inj 200 mg for ECP		200 mg OP	_	Baxter
	12.11	200 mg OF	•	Davici
ACTINOMYCIN [ACTINOMYCIN D] – PCT only – Specialist				_
Inj 0.5 mg vial		1		Cosmegen
Inj 0.5 mg for ECP	255.00	0.5 mg OP	•	Baxter
AUNORUBICIN - PCT only - Specialist				
Inj 18.7 mg for ECP	171.93	18.7 mg OP	✓	Baxter
Inj 18.7 mg vial	171.93	1	1	Pfizer
OCETAXEL - PCT only - Specialist				
Inj 20 mg	48.75	1	1	Docetaxel Sandoz
Inj 10 mg per ml, 8 ml vial		1		DBL Docetaxel
Inj 20 mg per ml, 4 ml vial	26.95	1		Docetaxel
)g p,	20.00	•		Accord S29
Inj 80 mg	105.00	1	1	Docetaxel Sandoz
Inj 1 mg for ECP		1 mg		Baxter
	0.35	ı iliy	•	Dαλίζι
OXORUBICIN HYDROCHLORIDE - PCT only - Specialist		_	_	<b></b>
Inj 2 mg per ml, 5 ml vial		1		Doxorubicin Ebewe
		1		Doxorubicin Ebewe
Inj 2 mg per ml, 25 ml vial				
	17.00			Arrow-Doxorubicin
Inj 2 mg per ml, 50 ml vial	17.00 23.00	1	✓	Doxorubicin Ebewe
	17.00 23.00 65.00	1 1	1	Doxorubicin Ebewe Arrow-Doxorubicin
Inj 2 mg per ml, 50 ml vial Inj 2 mg per ml, 100 ml vial	17.00 23.00 65.00 69.99	1	1	Doxorubicin Ebewe Arrow-Doxorubicin Doxorubicin Ebewe
Inj 2 mg per ml, 50 ml vial Inj 2 mg per ml, 100 ml vial Inj 1 mg for ECP	17.00 23.00 65.00 69.99		1	Doxorubicin Ebewe Arrow-Doxorubicin
Inj 2 mg per ml, 50 ml vial	17.00 23.00 65.00 69.99	1	1	Doxorubicin Ebewe Arrow-Doxorubicin Doxorubicin Ebewe
Inj 2 mg per ml, 50 ml vial	17.00 23.00 65.00 69.99 0.35	1	\ \ \ \ \	Doxorubicin Ebewe Arrow-Doxorubicin Doxorubicin Ebewe
Inj 2 mg per ml, 50 ml vial	17.00 23.00 65.00 69.99 0.35	1 1 mg	1111	Doxorubicin Ebewe Arrow-Doxorubicin Doxorubicin Ebewe Baxter
Inj 2 mg per ml, 50 ml vial	17.00 23.00 65.00 69.99 0.35	1 mg	\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	Doxorubicin Ebewe Arrow-Doxorubicin Doxorubicin Ebewe Baxter Epirubicin Ebewe

	Subsidy		Fully	Brand or
	Manufacturer's Price)		Subsidised	Generic
	\$	Per	•	Manufacturer
ETOPOSIDE				
Cap 50 mg - PCT - Retail pharmacy-Specialist Wastage claimable	340.73	20	•	Vepesid
Cap 100 mg - PCT - Retail pharmacy-Specialist Wastage claimable	340.73	10	•	Vepesid
Inj 20 mg per ml, 5 ml vial - PCT - Retail pharmacy-Specialis	t7.90	1	✓	Rex Medical
Inj 1 mg for ECP - PCT only - Specialist	0.09	1 mg	1	Baxter
ETOPOSIDE PHOSPHATE - PCT only - Specialist				
Inj 100 mg (of etoposide base)	40.00	1	1	Etopophos
Inj 1 mg (of etoposide base) for ECP		1 mg		Baxter
, , ,		9	-	Duntoi
HYDROXYUREA [HYDROXYCARBAMIDE] - PCT - Retail pharm	, ,			
Cap 500 mg	20.72	100	/	<u>Devatis</u>
IBRUTINIB - Special Authority see SA2480 below - Retail pharma	acy			
Tab 140 mg	3,217.00	30	1	Imbruvica
Tab 420 mg		30	✓	Imbruvica
	,			

### ⇒SA2480 Special Authority for Subsidy

Initial application — (chronic lymphocytic leukaemia (CLL)) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

#### All of the following:

- 1 Individual has chronic lymphocytic leukaemia (CLL) requiring therapy; and
- 2 Individual has not previously received funded ibrutinib; and
- 3 Ibrutinib is to be used as monotherapy; and
- 4 Any of the following:
  - 4.1 Both:
    - 4.1.1 There is documentation confirming that the individual has 17p deletion or TP53 mutation; and
    - 4.1.2 Individual has experienced intolerable side effects with venetoclax monotherapy; or
  - 4.2 All of the following:
    - 4.2.1 Individual has received at least one prior immunochemotherapy for CLL; and
    - 4.2.2 Individual's CLL has relapsed; and
    - 4.2.3 Individual has experienced intolerable side effects with venetoclax in combination with rituximab regimen; or
  - 4.3 Individual's CLL is refractory to or has relapsed following a venetoclax regimen.

Renewal — (chronic lymphocytic leukaemia (CLL)) from any relevant practitioner. Approvals valid for 12 months where there is no evidence of disease progression.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL) and B-cell prolymphocytic leukaemia (B-PLL)\*. Indications marked with \* are Unapproved indications.

#### IDARUBICIN HYDROCHLORIDE

Inj 5 mg vial – PCT only – Specialist	109.74	1	Zavedos
Inj 10 mg vial - PCT only - Specialist	233.64	1	✓ Zavedos
Inj 1 mg for ECP - PCT only - Specialist	25.77	1 mg	✓ Baxter
LENALIDOMIDE (VIATRIS) - Special Authority see SA23	53 on the next page – Re	etail pharmad	су
Cap 5 mg	76.92	21	✓ <u>Lenalidomide</u> <u>Viatris</u>
Cap 10 mg	50.30	21	✓ <u>Lenalidomide</u> <u>Viatris</u>
Cap 15 mg	62.13	21	✓ <u>Lenalidomide</u> <u>Viatris</u>
Cap 25 mg	65.09	21	✓ <u>Lenalidomide</u> Viatris

Subsidy		Fully	Brand or
(Manufacturer's Price	)	Subsidised	Generic
\$	Pe	r 🗸	Manufacturer

## **⇒SA2353** Special Authority for Subsidy

Initial application — (Plasma cell dyscrasia) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Both:

- 1 Patient has plasma cell dyscrasia, not including Waldenström macroglobulinaemia, requiring treatment; and
- 2 Patient is not refractory to prior lenalidomide use.

**Initial application — (Myelodysplastic syndrome)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

#### Both:

- 1 Patient has low or intermediate-1 risk myelodysplastic syndrome (based on IPSS or an IPSS-R score of less than 3.5) associated with a deletion 5q cytogenetic abnormality; and
- 2 Patient has transfusion-dependent anaemia.

Renewal — (Myelodysplastic syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

#### Both:

- 1 Patient has not needed a transfusion in the last 4 months; and
- 2 No evidence of disease progression.

#### MESNA

WEGNA			
Tab 400 mg - PCT - Retail pharmacy-Specialist	314.00	50	✓ Uromitexan
Tab 600 mg - PCT - Retail pharmacy-Specialist	448.50	50	✓ Uromitexan
Inj 100 mg per ml, 4 ml ampoule - PCT only - Specialist	177.45	15	✓ Uromitexan
Inj 100 mg per ml, 10 ml ampoule - PCT only - Specialist		15	✓ Uromitexan
Inj 1 mg for ECP – PCT only – Specialist		100 mg	✓ Baxter
MITOMYCIN C - PCT only - Specialist			
Inj 5 mg vial	517.65	1	✓ Accord S29
, - 3			✓ Mitomycin
			(Fresenius
			Kabi) \$29
	E06 00		,
	526.00		Mitomycin
			(Sagent) S29
Inj 20 mg vial	1,129.94	1	✓ Teva
Inj 1 mg for ECP	119.67	1 mg	✓ Baxter
MITOZANTRONE - PCT only - Specialist			
Inj 2 mg per ml, 10 ml vial	97.50	1	✓ Mitozantrone Ebewe
Inj 1 mg for ECP	5.51	1 mg	✓ Baxter
		g	- Buxton
NIRAPARIB – Special Authority see SA2325 below – Retail pharmac	СУ		
Wastage claimable			•
Tab 100 mg1		84	✓ Zejula
Cap 100 mg	8,929.84	56	✓ Zejula

#### **⇒SA2325** Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has advanced high-grade serous\* epithelial ovarian, fallopian tube, or primary peritoneal cancer; and
- 2 Patient has received at least one line\*\* of treatment with platinum-based chemotherapy; and
- 3 Patient has experienced a partial or complete response to the preceding treatment with platinum-based chemotherapy; and

continued...

- 4 Patient has not previously received funded treatment with a PARP inhibitor; and
- 5 Either
  - 5.1 Treatment will be commenced within 12 weeks of the patient's last dose of the preceding platinum-based regimen;
  - 5.2 Patient commenced treatment with niraparib prior to 1 May 2024; and
- 6 Treatment to be administered as maintenance treatment; and
- 7 Treatment not to be administered in combination with other chemotherapy.

**Renewal** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 No evidence of progressive disease: and
- 2 Treatment to be administered as maintenance treatment; and
- 3 Treatment not to be administered in combination with other chemotherapy; and
- 4 Either:
  - 4.1 Treatment with niraparib to cease after a total duration of 36 months from commencement; or
  - 4.2 Treatment with niraparib is being used in the second-line or later maintenance setting.

Notes: \* "high-grade serous" includes tumours with high-grade serous features or a high-grade serous component.

\*\*A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments

OLAPARIB - Retail pharmacy-Specialist - Special Au	thority see SA2163 below		
Tab 100 mg	3,701.00	56	<ul><li>Lynparza</li></ul>
Tab 150 mg	3,701.00	56	✓ Lynparza

### ⇒SA2163 Special Authority for Subsidy

Initial application — (Ovarian cancer) only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

### All of the following:

- 1 Patient has a high-grade serous\* epithelial ovarian, fallopian tube, or primary peritoneal cancer; and
- 2 There is documentation confirming pathogenic germline BRCA1 or BRCA2 gene mutation; and
- 3 Fither:
  - 3.1 All of the following:
    - 3.1.1 Patient has newly diagnosed, advanced disease; and
    - 3.1.2 Patient has received one line\*\* of previous treatment with platinum-based chemotherapy; and
    - 3.1.3 Patient's disease must have experienced a partial or complete response to the first-line platinum-based regimen; or
  - 3.2 All of the following:
    - 3.2.1 Patient has received at least two lines\*\* of previous treatment with platinum-based chemotherapy; and
    - 3.2.2 Patient has platinum sensitive disease defined as disease progression occurring at least 6 months after the last dose of the penultimate line\*\* of platinum-based chemotherapy; and
    - 3.2.3 Patient's disease must have experienced a partial or complete response to treatment with the immediately preceding platinum-based regimen; and
    - 3.2.4 Patient has not previously received funded olaparib treatment; and
- 4 Treatment will be commenced within 12 weeks of the patient's last dose of the immediately preceding platinum-based regimen; and
- 5 Treatment to be administered as maintenance treatment; and
- 6 Treatment not to be administered in combination with other chemotherapy.

**Renewal — (Ovarian cancer)** only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 12 months for applications meeting the following criteria:

				_
Subsidy		Fully	Brand or	_
(Manufacturer's Price)	Sul	osidised	Generic	
\$	Per	1	Manufacturer	

continued...

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from treatment; and
- 2 Either:
  - 2.1 No evidence of progressive disease; or
  - 2.2 Evidence of residual (not progressive) disease and the patient would continue to benefit from treatment in the clinician's opinion; and
- 3 Treatment to be administered as maintenance treatment; and
- 4 Treatment not to be administered in combination with other chemotherapy; and
- 5 Either:
  - 5.1 Both:
    - 5.1.1 Patient has received one line\*\* of previous treatment with platinum-based chemotherapy; and
    - 5.1.2 Documentation confirming that the patient has been informed and acknowledges that the funded treatment period of olaparib will not be continued beyond 2 years if the patient experiences a complete response to treatment and there is no radiological evidence of disease at 2 years; or
  - 5.2 Patient has received at least two lines\*\* of previous treatment with platinum-based chemotherapy.

Notes: \*Note "high-grade serous" includes tumours with high-grade serous features or a high-grade serous component.
\*\*A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

PACLITAXEL – PCT only – Specialist			
Inj 30 mg	47.30	5	✓ Paclitaxel Ebewe
lnj 6 mg per ml, 16.7 ml vial		1	✓ Anzatax
, ,	24.00		✓ Paclitaxel Ebewe
	91.67		✓ Paclitaxel Actavis
Inj 150 mg	26.69	1	✓ Paclitaxel Ebewe
. •	137.50		✓ Anzatax
			✓ Paclitaxel Actavis
Inj 6 mg per ml, 50 ml vial	37.89	1	✓ Anzatax
, ,	44.00		✓ Paclitaxel Ebewe
	275.00		✓ Paclitaxel Actavis
Inj 1 mg for ECP	0.17	1 mg	✓ Baxter
PEGASPARGASE - PCT only - Special Authority see SA	1979 below		
Inj 750 iu per ml, 5 ml vial	3,973.25	1	Oncaspar LYO

#### ⇒SA1979 Special Authority for Subsidy

**Initial application** — (Acute lymphoblastic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has newly diagnosed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

Initial application — (Lymphoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months where the patient has lymphoma requiring L-asparaginase containing protocols (e.g. SMILE).

Renewal — (Acute lymphoblastic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has relapsed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

PENTOSTATIN [DEOXYCOFORMYCIN] - PCT only - Speciali	ist		
Inj 10 mg	CBS	1	✓ Nipent S29

	Subsidy (Manufacturer's Price) \$	Sı Per	Fully ubsidised	Brand or Generic Manufacturer	
POMALIDOMIDE - Special Authority see SA2354 below - Retai	l pharmacy				
Cap 1 mg	47.45	14	✓ P	<u>omolide</u>	
, ,	71.18	21	✓ P	omolide	
Cap 2 mg	94.90	14	✓ P	omolide	
, ,	142.35	21	✓ P	omolide	
Cap 3 mg	142.35	14	✓ P	omolide	
	213.53	21	✓ P	omolide	
Cap 4 mg	189.81	14	✓ P	omolide	
, 0	284.71	21	<b>✓</b> P	omolide	

#### ⇒SA2354 Special Authority for Subsidy

Initial application — (Relapsed/refractory plasma cell dyscrasia) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has relapsed or refractory plasma cell dyscrasia, not including Waldenström macroglobulinaemia, requiring treatment; and
- 2 Patient has not received prior funded pomalidomide.

Renewal — (Relapsed/refractory plasma cell dyscrasia) from any relevant practitioner. Approvals valid for 12 months where there is no evidence of disease progression.

PROCARBAZINE HYDROCHLORIDE - PCT - Retail pharmacy-Specialist

Cap 50 mg	980.00	50	✓ Natulan S29
TEMOZOLOMIDE - Special Authority see \$	SA2275 below – Retail pharmacy		
Cap 5 mg	9.13	5	✓ Temaccord
			✓ Temozolomide-
			Taro S29
Cap 20 mg	16.38	5	✓ Temaccord
	18.30		✓ Apo-Temozolomide
Cap 100 mg	35.98	5	✓ Temaccord
	40.20		✓ Apo-Temozolomide
Cap 140 mg	50.12	5	✓ Temaccord
Cap 250 mg	86.34	5	✓ Temaccord

#### ⇒SA2275 Special Authority for Subsidy

Initial application — (gliomas) only from a relevant specialist. Approvals valid for 12 months where the patient has a glioma. Renewal — (gliomas) only from a relevant specialist. Approvals valid for 12 months where treatment remains appropriate and patient is benefitting from treatment.

Initial application — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

#### All of the following:

- 1 Patient has been diagnosed with metastatic or unresectable well-differentiated neuroendocrine tumour\*: and
- 2 Temozolomide is to be given in combination with capecitabine; and
- 3 Temozolomide is to be used in 28 day treatment cycles for a maximum of 5 days treatment per cycle at a maximum dose of 200 mg/m² per day; and
- 4 Temozolomide to be discontinued at disease progression.

Renewal — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

#### Roth:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

**Initial application** — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 9 months where the patient has relapsed/refractory Ewing's sarcoma.

**Renewal — (ewing's sarcoma)** only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indication marked with a \* is an unapproved indication. Temozolomide is not subsidised for the treatment of relapsed high grade glioma.

THALIDOMIDE - Retail pharmacy-Specialist - Special A	Authority see SA2356 below	1	
Cap 50 mg	378.00	28	Thalomid
Cap 100 mg	756.00	28	Thalomid

#### ⇒SA2356 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 12 months where the patient has plasma cell dyscrasia, not including Waldenström macroglobulinaemia, requiring treatment.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has obtained a response from treatment during the initial approval period.

Notes: Prescription must be written by a registered prescriber in the thalidomide risk management programme operated by the supplier.

Maximum dose of 400 mg daily as monotherapy or in a combination therapy regimen.

#### TRETINOIN

Cap 10 mg - PCT - Retail pharmacy-Specialist	479.50	100	Vesanoid
Wastage claimable			

VENETOCLAX - Retail pharmacy-Specialist - Special Authority see SA2481 below

Tab 14 × 10 mg, 7 × 50 mg, 21 × 100 mg1,771.86	42 OP	✓ Venclexta
Tab 10 mg13.68	2 OP	✓ Venclexta
Tab 50 mg239.44	7 OP	✓ Venclexta
Tab 100 mg - Wastage claimable8,209.41	120	✓ Venclexta

#### ⇒SA2481 Special Authority for Subsidy

Initial application — (relapsed/refractory chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 Individual has chronic lymphocytic leukaemia requiring treatment; and
- 2 Individual has received at least one prior therapy for chronic lymphocytic leukaemia; and
- 3 Individual has not previously received funded venetoclax; and
- 4 The individual's disease has relapsed: and
- 5 Venetoclax to be used in combination with six 28-day cycles of rituximab commencing after the 5-week dose titration schedule with venetoclax; and
- 6 Individual has an ECOG performance status of 0-2.

Renewal — (relapsed/refractory chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Treatment remains clinically appropriate and the individual is benefitting from and tolerating treatment; and
- 2 Venetoclax is to be discontinued after a maximum of 24 months of treatment following the titration schedule unless earlier discontinuation is required due to disease progression or unacceptable toxicity.

Initial application — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation\*) from any

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued

 $relevant\ practitioner.\ Approvals\ valid\ for\ 6\ months\ for\ applications\ meeting\ the\ following\ criteria:$ 

- All of the following:
  - 1 Individual has previously untreated chronic lymphocytic leukaemia; and
  - 2 There is documentation confirming that individual has 17p deletion by FISH testing or TP53 mutation by sequencing; and
  - 3 Individual has an ECOG performance status of 0-2.

Renewal — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation\*) from any relevant practitioner. Approvals valid for 6 months where the treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL)\* and B-cell prolymphocytic leukaemia (B-PLL)\*. Indications marked with \* are Unapproved indications

Initial application — (previously untreated acute myeloid leukaemia) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

#### Either:

- 1 The individual is currently on treatment with venetoclax and met all remaining special authority criteria prior to commencing treatment; or
- 2 All of the following:
  - 2.1 Individual has previously untreated acute myeloid leukaemia (see note a), according to World Health Organization (WHO) Classification; and
  - 2.2 Venetoclax not to be used in combination with standard intensive remission induction chemotherapy; and
  - 2.3 Venetoclax to be used in combination with azacitidine or low dose cytarabine.

Renewal — (previously untreated acute myeloid leukaemia) from any relevant practitioner. Approvals valid for 6 months where there is no evidence of disease progression.

Notes:

- a) 'Acute myeloid leukaemia' includes myeloid sarcoma\*
- b) Indications marked with \* are Unapproved indications

#### VINBLASTINE SUI PHATE

VIII DE TOTINE GOET TIVITE		
Inj 1 mg per ml, 10 ml vial - PCT - Retail pharmacy-Specialist270.37	5	✓ Hospira
Inj 1 mg for ECP - PCT only - Specialist	1 mg	✓ Baxter
VINCRISTINE SULPHATE		
Inj 1 mg per ml, 1 ml vial - PCT - Retail pharmacy-Specialist74.52	5	✓ DBL Vincristine Sulfate
Inj 1 mg per ml, 2 ml vial - PCT - Retail pharmacy-Specialist102.73	5	✓ DBL Vincristine Sulfate
Inj 1 mg for ECP - PCT only - Specialist12.60	1 mg	✓ Baxter
VINORELBINE		
Cap 20 mg32.10 Vinorelbine Te Arai to be Principal Supply on 1 February 2026	1	✓ Vinorelbine Te Arai
Cap 30 mg42.80	1	✓ Vinorelbine Te Arai
Vinorelbine Te Arai to be Principal Supply on 1 February 2026		<b>4.</b> 111 <b>-</b> 4 .
Cap 80 mg	1	✓ Vinorelbine Te Arai
Inj 10 mg per ml, 1 ml vial - PCT only - Specialist	1	✓ Vinorelbine Ebewe
Inj 10 mg per ml, 5 ml vial - PCT only - Specialist155.00	1	✓ Vinorelbine Ebewe
168.00		✓ Navelbine S29 S29
Inj 1 mg for ECP - PCT only - Specialist	1 mg	✓ Baxter

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

# Protein-tyrosine Kinase Inhibitors

ALECTINIB - Retail pharmacy-Specialist - Special Authority see SA1870 below

Wastage claimable

224 Alecensa

### ⇒SA1870 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-small cell lung cancer; and
- 2 There is documentation confirming that the patient has an ALK tyrosine kinase gene rearrangement using an appropriate ALK test: and
- 3 Patient has an ECOG performance score of 0-2.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

### Both:

- 1 No evidence of progressive disease according to RECIST criteria; and
- 2 The patient is benefitting from and tolerating treatment.

AXITINIB - Special Authority see SA2458 below - Retail pharmacy

Wastage claimable

Tab 1 mg536.40	28	✓ Inlyta
Tab 5 mg2,682.00	28	✓ Inlyta

#### ⇒SA2458 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 The disease is of predominant clear cell histology; and
- 3 The patient has documented disease progression following one previous line of treatment; and
- 4 The patient has ECOG performance status of 0-2.

Renewal from any relevant practitioner. Approvals valid for 4 months where there is no evidence of disease progression..

CRIZOTINIB - Special Authority see SA2547 below - Retail pharmacy

Cap 200 mg	 7,250.00	60	1	Xalkori
Cap 250 mg	 7,250.00	60	1	Xalkori

## ⇒SA2547 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Individual has locally advanced or metastatic, unresectable, non-squamous non-small cell lung cancer; and
- 2 Either:
  - 2.1 The individual has not received entrectinib; or
  - 2.2 Both:
    - 2.2.1 The individual has received an initial Special Authority approval for entrectinib and has discontinued entrectinib due to intolerance: and
    - 2.2.2 The cancer did not progress while the individual was on entrectinib; and
- 3 There is documentation confirming that the patient has a ROS1 rearrangement using an appropriate ROS1 test; and
- 4 Individual has ECOG performance score of 0-3; and
- 5 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
\$	Per	✓	Manufacturer

continued...

Renewal from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

- 1 Response to treatment has been determined by comparable radiological assessment following the most recent treatment period; and
- 2 No evidence of disease progression.

DABRAFENIB - Special Authority see SA2548 below - Retail pharmacy

Cap 50 mg	6,320.86	120	Tafinlar
Cap 75 mg	9.481.29	120	Tafinlar

⇒SA2548 Special Authority for Subsidy

Initial application — (stage III or IV resected melanoma - adjuvant) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 The individual has resected stage IIIB, IIIC, IIID or IV melanoma (excluding uveal) (see note a); or
  - 1.2 Both:
    - 1.2.1 The individual has received neoadjuvant treatment with a PD-1/PD-L1 inhibitor; and
    - 1.2.2 Adjuvant treatment with dabrafenib is required; and
- 2 The individual has not received prior funded systemic treatment in the adjuvant setting for stage IIIB, IIIC, IIID or IV melanoma: and
- 3 Treatment must be adjuvant to complete surgical resection; and
- 4 Treatment must be initiated within 13 weeks of surgical resection, unless delay is necessary due to post-surgery recovery (see note b); and
- 5 The individual has a confirmed BRAF mutation; and
- 6 Dabrafenib must be administered in combination with trametinib; and
- 7 The individual has ECOG performance score 0-2.

#### Notes:

- a) Stage IIIB, IIIC, IIID or IV melanoma defined as per American Joint Committee on Cancer (AJCC) 8th Edition
- b) Initiating treatment within 13 weeks of complete surgical resection means 13 weeks after resection (primary or lymphadenectomy)

Renewal — (stage III or IV resected melanoma - adjuvant) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Any of the following:

- 1 All of the following:
  - 1.1 No evidence of disease recurrence: and
  - 1.2 Dabrafenib must be administered in combination with trametinib; and
  - 1.3 Treatment to be discontinued at signs of disease recurrence or at completion of 12 months' total treatment course, including any systemic neoadjuvant treatment; or
- 2 All of the following:
  - 2.1 The individual has received adjuvant treatment with a BRAF/MEK inhibitor; and
  - 2.2 The individual has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
  - 2.3 The individual meets initial application criteria for dabrafenib for unresectable or metastatic melanoma; or
- 3 All of the following:
  - 3.1 The individual has received adjuvant treatment with a BRAF/MEK inhibitor; and
  - 3.2 The individual has received a BRAF/MEK inhibitor for unresectable or metastatic melanoma; and
  - 3.3 The individual meets renewal criteria for dabrafenib for unresectable or metastatic melanoma.

Initial application — (unresectable or metastatic melanoma) from any relevant practitioner. Approvals valid for 4 months for

bsidy	Fully	Brand or
turer's Price) Subsid	dised	Generic
 \$ Per	✓	

continued...

applications meeting the following criteria:

All of the following:

- 1 The individual has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Baseline measurement of overall tumour burden is documented clinically and radiologically; and
- 3 The individual has ECOG performance score 0-2; and
- 4 The individual has confirmed BRAF mutation: and
- 5 Dabrafenib must be administered in combination with trametinib: and
- 6 Any of the following:
  - 6.1 The individual has been diagnosed in the metastatic or unresectable stage III or IV setting; or
  - 6.2 The individual did not receive treatment in the adjuvant setting with a BRAF/MEK inhibitor; or
  - 6.3 All of the following:
    - 6.3.1 The individual received treatment in the adjuvant setting with a BRAF/MEK inhibitor; and
    - 6.3.2 The individual did not experience disease recurrence while on treatment with that BRAF/MEK inhibitor; and
    - 6.3.3 The individual did not experience disease recurrence within six months of completing adjuvant treatment with a BRAF/MFK inhibitor.

**Renewal — (unresectable or metastatic melanoma)** from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 The individual's disease has had a complete response to treatment; or
  - 1.2 The individual's disease has had a partial response to treatment; or
  - 1.3 The individual has stable disease with treatment; and
- 2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period.

#### DASATINIB - Special Authority see SA2385 below - Retail pharmacy

Wastage claimable

Tab 20 mg	132.88	60	✓ Dasatinib-Teva
Tab 50 mg	304.13	60	✓ Dasatinib-Teva
Tab 70 mg	415.75	60	✓ Dasatinib-Teva

### ⇒SA2385 Special Authority for Subsidy

**Initial application** only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis or accelerated phase; or
- 2 The patient has a diagnosis of Philadelphia chromosome-positive acute lymphoid leukaemia (Ph+ ALL); or
- 3 Both:
  - 3.1 The patient has a diagnosis of CML in chronic phase; and
  - 3.2 Any of the following:
    - 3.2.1 Patient has documented treatment failure\* with imatinib: or
    - 3.2.2 Patient has experienced treatment-limiting toxicity with imatinib precluding further treatment with imatinib; or
    - 3.2.3 Patient has high-risk chronic-phase CML defined by the Sokal or EURO scoring system.

**Renewal** only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Lack of treatment failure while on dasatinib\*; and
- 2 Dasatinib treatment remains appropriate and the patient is benefiting from treatment.

Note: \*treatment failure for CML as defined by Leukaemia Net Guidelines.

ENTRECTINIB – Special Authority see SA2532 on the next page – Retail pharmacy
Cap 200 mg.......9,610.00 90 ✓ Rozlytrek

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

### ⇒SA2532 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Individual has locally advanced or metastatic, unresectable, non-squamous non-small cell lung cancer; and
- 2 Fither
  - 2.1 The individual has not received crizotinib; or
  - 2.2 Both:
    - 2.2.1 The individual has received an initial Special Authority approval for crizotinib and has discontinued crizotinib due to intolerance: and
    - 2.2.2 The cancer did not progress while the individual was on crizotinib; and
- 3 There is documentation confirming that the patient has a ROS1 rearrangement using an appropriate ROS1 test; and
- 4 Individual has ECOG performance score of 0-3; and
- 5 Baseline measurement of overall tumour burden is documented clinically and radiologicallyy.

Renewal from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Response to treatment has been determined by comparable radiological assessment following the most recent treatment period; and
- 2 No evidence of disease progression.

ERLOTINIB	<ul> <li>Retail pharmacy-Specialist – \$</li> </ul>	Special Authority see	SA2422 below

Tab 100 mg	280.84	30	✓ Alchemy
Tab 150 mg	484.24	30	<ul> <li>Alchemy</li> </ul>

#### ⇒SA2422 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 There is documentation confirming that the disease expresses activating mutations of EGFR: and
- 3 Any of the following:
  - 3.1 Patient is treatment naive; or
  - 3.2 Patient has received prior treatment in the adjuvant setting and/or while awaiting EGFR results; or
  - 3.3 Both:
    - 3.3.1 The patient has discontinued osimertinib or gefitinib due to intolerance; and
    - 3.3.2 The cancer did not progress while on osimertinib or gefitinib.

Renewal from any relevant practitioner. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

GEFITINIB - Retail pharmacy-Specialist - Special Authority see SA2423 below

Tab 250 mg .......918.00 30 ✓ Iressa

#### ⇒SA2423 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 Any of the following:
  - 2.1 Patient is treatment naive: or
  - 2.2 Patient has received prior treatment in the adjuvant setting and/or while awaiting EGFR results; or
  - 2.3 Both:
    - 2.3.1 The patient has discontinued osimertinib or erlotinib due to intolerance; and
    - 2.3.2 The cancer did not progress whilst on osimertinib or erlotinib; and
- 3 There is documentation confirming that disease expresses activating mutations of EGFR.

Renewal from any relevant practitioner. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

()	Subsidy Manufacturer's Price) \$	Per	Fully Subsidised	I Generic
IMATINIB MESILATE				
* Cap 100 mg	44.93	60	✓	Imatinib-Rex
* Cap 400 mg	69.76	30	✓	Imatinib-Rex
LENVATINIB – Special Authority see SA2442 below – Retail pharm Wastage claimable	nacy			
Cap 4 mg	3,407.40	30	✓	Lenvima
Cap 10 mg	3,407.40	30	✓	Lenvima

#### ⇒SA2442 Special Authority for Subsidy

Initial application — (thyroid cancer) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

#### Either:

- 1 Patient is currently on treatment with lenvatinib and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
  - 2.1 The patient has locally advanced or metastatic differentiated thyroid cancer; and
  - 2.2 Either:
    - 2.2.1 Patient must have symptomatic progressive disease prior to treatment; or
    - 2.2.2 Patient must progressive disease at critical anatomical sites with a high risk of morbidity or mortality where local control cannot be achieved by other measures; and
  - 2.3 Any of the following:
    - 2.3.1 A lesion without iodine uptake in a RAI scan; or
    - 2.3.2 Receiving cumulative RAI greater than or equal to 600 mCi; or
    - 2.3.3 Experiencing disease progression after a RAI treatment within 12 months; or
    - 2.3.4 Experiencing disease progression after two RAI treatments administered within 12 months of each other;
  - 2.4 Patient has thyroid stimulating hormone (TSH) adequately supressed; and
  - 2.5 Patient is not a candidate for radiotherapy with curative intent; and
  - 2.6 Surgery is clinically inappropriate; and
  - 2.7 Patient has an ECOG performance status of 0-2.

Renewal — (thyroid cancer) from any relevant practitioner. Approvals valid for 6 months where there is no evidence of disease progression.

Initial application — (unresectable hepatocellular carcinoma) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has unresectable hepatocellular carcinoma; and
- 2 Patient has preserved liver function (Childs-Pugh A); and
- 3 Transarterial chemoembolisation (TACE) is unsuitable: and
- 4 Patient has an ECOG performance status of 0-2; and
- 5 Fither:
  - 5.1 Patient has not received prior systemic therapy for their disease in the palliative setting; or
  - 5.2 Both:
    - 5.2.1 Patient has experienced treatment-limiting toxicity from treatment with atezolizumab with bevacizumab; and
    - 5.2.2 No disease progression since initiation of atezolizumab with bevacizumab.

Renewal — (unresectable hepatocellular carcinoma) from any relevant practitioner. Approvals valid for 6 months where there is no evidence of disease progression.

Initial application — (renal cell carcinoma) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

1 All of the following:

Subsidy (Manufacturer's Price)	Fu Subsidis	,	Brand or Generic
\$	Per	_	Manufacturer

#### continued...

- 1.1 The patient has metastatic renal cell carcinoma; and
- 1.2 The disease is of predominant clear-cell histology; and
- 1.3 The patient has documented disease progression following one previous line of treatment; and
- 1.4 The patient has an ECOG performance status of 0-2; and
- 1.5 Lenvatinib is to be used in combination with everolimus; or
- 2 All of the following:
  - 2.1 Patient has received funded treatment with nivolumab for the second line treatment of metastatic renal cell carcinoma; and
  - 2.2 Patient has experienced treatment limiting toxicity from treatment with nivolumab; and
  - 2.3 Lenvatinib is to be used in combination with everolimus; and
  - 2.4 There is no evidence of disease progression.

**Renewal** — **(renal cell carcinoma)** from any relevant practitioner. Approvals valid for 4 months where there is no evidence of disease progression.

MIDOSTAURIN − PCT only − Special Authority see SA2342 below

Cap 25 mg......10,981.00 56 
✓ Rydapt

### ⇒SA2342 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria: All of the following:

- 1 Patient has a diagnosis of acute myeloid leukaemia; and
- 2 Condition must be FMS tyrosine kinase 3 (FLT3) mutation positive; and
- 3 Patient must not have received a prior line of intensive chemotherapy for acute myeloid leukaemia; and
- 4 Patient is to receive standard intensive chemotherapy in combination with midostaurin only; and
- 5 Midostaurin to be funded for a maximum of 4 cycles.

NILOTINIB - Special Authority see SA2301 below - Retail pharmacy

Wastage claimable		
Cap 150 mg4,680.00	120	✓ Tasigna
Cap 200 mg6,532.00	120	✓ Tasigna

# **⇒SA2301** Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis, high risk chronic phase, or in chronic phase; and
- 2 Either:
  - 2.1 Patient has documented CML treatment failure\* with a tyrosine kinase inhibitor (TKI); or
  - 2.2 Patient has experienced treatment limiting toxicity with a tyrosine kinase inhibitor (TKI) precluding further treatment; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

Note: \*treatment failure as defined by Leukaemia Net Guidelines.

Renewal only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Lack of treatment failure while on nilotinib as defined by Leukaemia Net Guidelines; and
- 2 Nilotinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

OSIMERTINIB - Special Authority see SA2418 on the	e next page - Retail pharmacy		
Tab 40 mg	9,310.00	30	✓ Tagrisso
Tab 80 mg	9,310.00	30	✓ Tagrisso

Subsidy	,	Fully	Brand or	
(Manufacturer's Pric	9)	Subsidised	Generic	
\$	Pe	r 🗸	Manufacturer	

### ⇒SA2418 Special Authority for Subsidy

Initial application — (NSCLC – first line) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

#### Either:

- 1 Patient is currently on treatment with osimertinib and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
  - 2.1 Patient has locally advanced or metastatic, incurable, non-squamous non-small cell lung cancer (NSCLC); and
  - 2.2 Any of the following:
    - 2.2.1 Patient is treatment naïve; or
    - 2.2.2 Patient has received prior chemotherapy in the adjuvant setting and/or while awaiting EGFR results; or
    - 2.2.3 Both
      - 2.2.3.1 The patient has discontinued gefitinib or erlotinib due to intolerance; and
      - 2.2.3.2 The cancer did not progress while on gefitinib or erlotinib; and
  - 2.3 There is documentation confirming that the cancer expresses activating mutations of EGFR; and
  - 2.4 Patient has an ECOG performance status 0-3; and
  - 2.5 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Renewal — (NSCLC – first line) from any relevant practitioner. Approvals valid for 6 months where response to or stable disease with treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period.

**Initial application — (NSCLC – second line)** from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

#### Either:

- 1 Patient is currently on treatment with osimertinib and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
  - 2.1 Patient has locally advanced or metastatic, incurable, non-squamous non-small cell lung cancer (NSCLC); and
  - 2.2 Patient has an ECOG performance status 0-3; and
  - 2.3 The patient must have received previous treatment with erlotinib or gefitinib; and
  - 2.4 There is documentation confirming that the cancer expresses T790M mutation of EGFR following progression on or after erlotinib or gefitinib; and
  - 2.5 The treatment must be given as monotherapy; and
  - 2.6 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Renewal — (NSCLC – second line) from any relevant practitioner. Approvals valid for 6 months where response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period.

PALBOCICLIB - Special Authority see SA2345 below - Retail pharmacy

Wastage claimable			
Tab 75 mg	1,200.00	21	Palbociclib Pfizer
Tab 100 mg	1,200.00	21	Palbociclib Pfizer
Tab 125 mg	1,200.00	21	Palbociclib Pfizer

#### ⇒SA2345 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Either:

- 1 All of the following:
  - 1.1 Patient has unresectable locally advanced or metastatic breast cancer; and
  - 1.2 There is documentation confirming disease is hormone-receptor positive and HER2-negative; and
  - 1.3 Patient has an ECOG performance score of 0-2; and
  - 1.4 Either:
    - 1.4.1 Disease has relapsed or progressed during prior endocrine therapy; or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
	Per 🗸	Manufacturer

continued...

- 1.4.2 Both:
  - 1.4.2.1 Patient is amenorrhoeic, either naturally or induced, with endocrine levels consistent with a postmenopausal or without menstrual-potential state; and
  - 1.4.2.2 Patient has not received prior systemic treatment for metastatic disease; and
- 1.5 Treatment must be used in combination with an endocrine partner; and
- 1.6 Patient has not received prior funded treatment with a CDK4/6 inhibitor; or
- 2 All of the following:
  - 2.1 Patient has an active Special Authority approval for ribociclib: and
  - 2.2 Patient has experienced a grade 3 or 4 adverse reaction to ribociclib that cannot be managed by dose reductions and requires treatment discontinuation; and
  - 2.3 Treatment must be used in combination with an endocrine partner; and
  - 2.4 There is no evidence of progressive disease since initiation of ribociclib.

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 Treatment must be used in combination with an endocrine partner; and
- 2 There is no evidence of progressive disease since initiation of palbociclib.

PAZOPANIB – Special Authority see SA2429 below	w – Retail pharmacy		
Tab 200 mg	172.88	30	✓ Pazopanib Teva
Tab 400 mg	464.00	30	✓ Pazopanib Teva

### ⇒SA2429 Special Authority for Subsidy

Initial application only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 All of the following:
  - 1.1 The patient has metastatic renal cell carcinoma of predominantly clear cell histology; and
  - 1.2 Either:
    - 1.2.1 The patient is treatment naive; or
    - 1.2.2 The patient has only received prior cytokine treatment; and
  - 1.3 The patient has an ECOG performance score of 0-2; and The patient has intermediate or poor prognosis defined as:
  - 1.4 Any of the following:
    - 1.4.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
    - 1.4.2 Haemoglobin level < lower limit of normal; or
    - 1.4.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
    - 1.4.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
    - 1.4.5 Karnofsky performance score of less than or equal to 70; or
    - 1.4.6 2 or more sites of organ metastasis; and
  - 1.5 Pazopanib to be used for a maximum of 3 months; or
- 2 All of the following:
  - 2.1 The patient has metastatic renal cell carcinoma; and
  - 2.2 The patient has discontinued sunitinib within 3 months of starting treatment due to intolerance; and
  - 2.3 The cancer did not progress whilst on sunitinib; and
  - 2.4 Pazopanib to be used for a maximum of 3 months.

**Renewal** only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months where there is no evidence of disease progression.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
RIBOCICLIB - Special Authority see SA2495 below - Retail pha Wastage claimable	rmacy		_	
Tab 200 mg	1,883.00	21	<b>✓</b> K	isqali
·	3,767.00	42	✓ K	isqali
	5,650.00	63	✓ K	isqali

### ⇒SA2495 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Fither:

- 1 All of the following:
  - 1.1 Patient has unresectable locally advanced or metastatic breast cancer; and
  - 1.2 There is documentation confirming disease is hormone-receptor positive and HER2-negative; and
  - 1.3 Patient has an ECOG performance score of 0-2; and
  - 1.4 Either:
    - 1.4.1 Disease has relapsed or progressed during prior endocrine therapy; or
    - 1.4.2 Both:
      - 1.4.2.1 Patient is amenorrhoeic, either naturally or induced, with endocrine levels consistent with a postmenopausal or without menstrual-potential state; and
      - 1.4.2.2 Patient has not received prior systemic endocrine treatment for metastatic disease; and
  - 1.5 Treatment to be used in combination with an endocrine partner; and
  - 1.6 Patient has not received prior funded treatment with a CDK4/6 inhibitor; or
- 2 All of the following:
  - 2.1 Patient has an active Special Authority approval for palbociclib; and
  - 2.2 Patient has experienced a grade 3 or 4 adverse reaction to palbociclib that cannot be managed by dose reductions and requires treatment discontinuation; and
  - 2.3 Treatment must be used in combination with an endocrine partner; and
  - 2.4 There is no evidence of progressive disease since initiation of palbociclib.

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 Treatment must be used in combination with an endocrine partner; and
- 2 There is no evidence of progressive disease since initiation of ribociclib.

#### RUXOLITINIB - Special Authority see SA1890 below - Retail pharmacy

Wastage claimable	•	
Tab 5 mg2,	500.00 56	Jakavi
Tab 10mg	000.00 56	Jakavi
Tab 15 mg5,0	000.00 56	Jakavi
Tab 20 mg5,0	000.00 56	Jakavi

#### ⇒SA1890 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has primary myelofibrosis or post-polycythemia vera myelofibrosis or post-essential thrombocythemia myelofibrosis; and
- 2 Fither:
  - 2.1 A classification of risk of intermediate-2 or high-risk myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS; or
  - 2.2 Both:
    - 2.2.1 A classification of risk of intermediate-1 myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted

|--|

continued...

DIPSS; and

- 2.2.2 Patient has severe disease-related symptoms that are resistant, refractory or intolerant to available therapy; and
- 3 A maximum dose of 20 mg twice daily is to be given.

**Renewal** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

- Both:
  - 1 The treatment remains appropriate and the patient is benefiting from treatment; and
  - 2 A maximum dose of 20 mg twice daily is to be given.

SUNITINIB - Special Authority see SA2452 below - Reta	il pharmacy		
Cap 12.5 mg	103.11	28	<ul> <li>Sunitinib Rex</li> </ul>
	208.38		<ul> <li>Sunitinib Pfizer</li> </ul>
Sunitinib Rex to be Principal Supply on 1 March 2	026		
Cap 25 mg	203.15	28	✓ Sunitinib Rex
, ,	416.77		<ul> <li>Sunitinib Pfizer</li> </ul>
Sunitinib Rex to be Principal Supply on 1 March 2	026		
Cap 50 mg	343.19	28	<ul> <li>Sunitinib Rex</li> </ul>
, ,	694.62		<ul> <li>Sunitinib Pfizer</li> </ul>

(Sunitinib Pfizer Cap 12.5 mg to be delisted 1 March 2026) (Sunitinib Pfizer Cap 25 mg to be delisted 1 March 2026)

(Sunitinib Pfizer Cap 50 mg to be delisted 1 May 2026)

#### ⇒SA2452 Special Authority for Subsidy

**Initial application** — (RCC) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

#### Both:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 The patient has not previously received funded sunitinib.

Initial application — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal tumour (GIST); and
- 2 Either:
  - 2.1 The patient's disease has progressed following treatment with imatinib; or
  - 2.2 The patient has documented treatment-limiting intolerance, or toxicity to, imatinib.

Renewal — (RCC) from any relevant practitioner. Approvals valid for 4 months where there is no evidence of disease progression.

**Renewal — (GIST)** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

The patient has responded to treatment or has stable disease as determined by Choi's modified CT response evaluation criteria as follows:

- 1 Any of the following:
  - 1.1 The patient has had a complete response (disappearance of all lesions and no new lesions); or
  - 1.2 The patient has had a partial response (a decrease in size of 10% or more or decrease in tumour density in Hounsfield Units (HU) of 15% or more on CT and no new lesions and no obvious progression of non measurable disease); or

Subsidy (Manufacturer's Price)	Subs	Fully	Brand or Generic
\$	Per	1	Manufacturer

continued...

- 1.3 The patient has stable disease (does not meet criteria the two above) and does not have progressive disease and no symptomatic deterioration attributed to tumour progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: It is recommended that response to treatment be assessed using Choi's modified CT response evaluation criteria (J Clin Oncol, 2007, 25:1753-1759). Progressive disease is defined as either: an increase in tumour size of 10% or more and not meeting criteria of partial response (PR) by tumour density (HU) on CT; or: new lesions, or new intratumoral nodules, or increase in the size of the existing intratumoral nodules.

Renewal — (GIST pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal (GIST); and
- 2 The patient is clinically benifiting from treatment and continued treatment remains appropriate; and
- 3 Sunitinib is to be discontinued at progression; and
- 4 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

# TRAMETINIB - Special Authority see SA2549 below - Retail pharmacy

Tab 0.5 mg	2,370.32	30	✓ Mekinist
Tab 2 mg	9,481.29	30	✓ Mekinist

## ⇒SA2549 Special Authority for Subsidy

Initial application — (stage III or IV resected melanoma - adjuvant) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 The individual has resected stage IIIB, IIIC, IIID or IV melanoma (excluding uveal) (see note a); or
  - 1.2 Both:
    - 1.2.1 The individual has received neoadjuvant treatment with a PD-1/PD-L1 inhibitor; and
    - 1.2.2 Adjuvant treatment with trametinib is required; and
- 2 The individual has not received prior funded systemic treatment in the adjuvant setting for stage IIIB, IIIC, IIID or IV melanoma; and
- 3 Treatment must be adjuvant to complete surgical resection; and
- 4 Treatment must be initiated within 13 weeks of surgical resection, unless delay is necessary due to post-surgery recovery (see note b); and
- 5 The individual has a confirmed BRAF mutation; and
- 6 Trametinib must be administered in combination with dabrafenib; and
- 7 The individual has ECOG performance score 0-2.

#### Notes:

- a) Stage IIIB, IIIC, IIID or IV melanoma defined as per American Joint Committee on Cancer (AJCC) 8th Edition
- b) Initiating treatment within 13 weeks of complete surgical resection means 13 weeks after resection (primary or lymphadenectomy)

Renewal — (stage III or IV resected melanoma - adjuvant) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Any of the following:

- 1 All of the following:
  - 1.1 No evidence of disease recurrence; and
  - 1.2 Trametinib must be administered in combination with dabrafenib; and
  - 1.3 Treatment to be discontinued at signs of disease recurrence or at completion of 12 months' total treatment course, including any systemic neoadjuvant treatment; or

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 2 All of the following:
  - 2.1 The individual has received adjuvant treatment with a BRAF/MEK inhibitor; and
  - 2.2 The individual has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
  - 2.3 The individual meets initial application criteria for trametinib for unresectable or metastatic melanoma; or
  - 3 All of the following:
    - 3.1 The individual has received adjuvant treatment with a BRAF/MEK inhibitor; and
    - 3.2 The individual has received a BRAF/MEK inhibitor for unresectable or metastatic melanoma; and
    - 3.3 The individual meets renewal criteria for trametinib for unresectable or metastatic melanoma.

**Initial application** — (unresectable or metastatic melanoma) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The individual has metastatic or unresectable melanoma (excluding uveal melanoma) stage III or IV; and
- 2 Baseline measurement of overall tumour burden is documented clinically and radiologically; and
- 3 The individual has ECOG performance score 0-2; and
- 4 The individual has confirmed BRAF mutation; and
- 5 Trametinib must be administered in combination with dabrafenib; and
- 6 Any of the following:
  - 6.1 The individual has been diagnosed in the metastatic or unresectable stage III or IV setting; or
  - 6.2 The individual did not receive treatment in the adjuvant setting with a BRAF/MEK inhibitor; or
  - 6.3 All of the following:
    - 6.3.1 The individual received treatment in the adjuvant setting with a BRAF/MEK inhibitor; and
    - 6.3.2 The individual did not experience disease recurrence while on treatment with that BRAF/MEK inhibitor; and
    - 6.3.3 The individual did not experience disease recurrence within six months of completing adjuvant treatment with a BRAF/MEK inhibitor.

Renewal — (unresectable or metastatic melanoma) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 The individual's disease has had a complete response to treatment; or
  - 1.2 The individual's disease has had a partial response to treatment; or
  - 1.3 The individual has stable disease with treatment; and
- 2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period.

# **Endocrine Therapy**

For GnRH ANALOGUES - refer to HORMONE PREPARATIONS, Trophic Hormones, page 88

ABIRATERONE ACETATE - Retail pharmacy-Specialist - Special Authority see SA2118 below

Wastage claimable

#### ⇒SA2118 Special Authority for Subsidy

**Initial application** only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has prostate cancer; and
- 2 Patient has metastases; and

	Subsidy	Ful	y Brand or
(Man	ufacturer's Price)	Subsidise	d Generic
	\$	Per •	Manufacturer

continued...

- 3 Patient's disease is castration resistant; and
- 4 Fither:
  - 4.1 All of the following:
    - 4.1.1 Patient is symptomatic; and
    - 4.1.2 Patient has disease progression (rising serum PSA) after second line anti-androgen therapy; and
    - 4.1.3 Patient has ECOG performance score of 0-1; and
    - 4.1.4 Patient has not had prior treatment with taxane chemotherapy; or
  - 4.2 All of the following:
    - 4.2.1 Patient's disease has progressed following prior chemotherapy containing a taxane; and
    - 4.2.2 Patient has ECOG performance score of 0-2; and
    - 4.2.3 Patient has not had prior treatment with abiraterone.

Renewal — (abiraterone acetate) only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Significant decrease in serum PSA from baseline; and
- 2 No evidence of clinical disease progression; and
- 3 No initiation of taxane chemotherapy with abiraterone; and
- 4 The treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 2 Abiraterone acetate to be discontinued at progression; and
- 3 No initiation of taxane chemotherapy with abiraterone; and
- 4 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

BICALUTAMIDE			
Tab 50 mg	4.18	28	✓ Binarex
FLUTAMIDE			
Tab 250 mg	107.55	90	✓ Prostacur S29
· ·	119.50	100	✓ Flutamin
FULVESTRANT - Retail pharmacy-Specialist - Special Auth	ority see SA1895 bel	ow	
Inj 50 mg per ml, 5 ml prefilled syringe	181.00	2	✓ Fulvestrant EVER Pharma
	1,068.00		✓ Faslodex

(Faslodex Inj 50 mg per ml, 5 ml prefilled syringe to be delisted 1 May 2026)

⇒SA1895 Special Authority for Subsidy

**Initial application** only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has oestrogen-receptor positive locally advanced or metastatic breast cancer; and
- 2 Patient has disease progression following prior treatment with an aromatase inhibitor or tamoxifen for their locally advanced or metastatic disease; and
- 3 Treatment to be given at a dose of 500 mg monthly following loading doses; and
- 4 Treatment to be discontinued at disease progression.

Subsidy (Manufacturer's Price)	,	Fully Subsidised	Brand or Generic	
\$	Per	✓	Manufacturer	

continued...

**Renewal** only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

#### All of the following:

- 1 Treatment remains appropriate and patient is benefitting from treatment; and
- 2 Treatment to be given at a dose of 500 mg monthly; and
- 3 There is no evidence of disease progression.

#### **OCTREOTIDE**

Inj 100 mcg per ml, 1 ml ampoule	32.71	5	✓ Max Health
			✓ Octreotide GH \$29
			✓ Sun Pharma S29
Inj 100 mcg per ml, 1 ml vial	48.50	5	✓ Omega S29
Inj 50 mcg per ml, 1 ml vial	27.58	5	✓ Omega S29
Inj 500 mcg per ml, 1 ml vial	113.10	5	✓ Omega S29
Inj 50 mcg per ml, 1 ml ampoule	27.58	5	✓ Max Health
			✓ Octreotide GH S29
Inj 500 mcg per ml, 1 ml ampoule	113.10	5	✓ Max Health
			✓ Octreotide GH \$29
			✓ Sun Pharma S29
TAMOXIFEN CITRATE			
* Tab 10 mg		60	✓ Tamoxifen Sandoz
* Tab 20 mg	5.32	60	✓ Tamoxifen Sandoz

# **Long-acting Somatostatin Analogues**

#### ⇒SA2445 Special Authority for Subsidy

**Initial application — (Malignant Bowel Obstruction)** from any relevant practitioner. Approvals valid for 2 months for applications meeting the following criteria:

#### All of the following:

- 1 The patient has nausea\* and vomiting\* due to malignant bowel obstruction\*; and
- 2 Treatment with antiemetics, rehydration, antimuscarinic agents, corticosteroids and analgesics for at least 48 hours has not been successful; and
- 3 Treatment to be given for up to 4 weeks.

Note: Indications marked with \* are unapproved indications.

**Renewal — (Malignant Bowel Obstruction)** from any relevant practitioner. Approvals valid for 3 months where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Acromegaly) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

#### All of the following:

- 1 The patient has acromegaly; and
- 2 Either:
  - 2.1 Treatment with surgery and radiotherapy is not suitable or was unsuccessful; or
  - 2.2 Treatment is for an interim period while awaiting the beneficial effects of radiotherapy; and
- 3 Treatment with a dopamine agonist has been unsuccessful.

**Renewal** — (Acromegaly) from any relevant practitioner. Approvals valid for 2 years where iGF1 levels have decreased since starting treatment.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

Note: In patients with acromegaly, treatment should be discontinued if IGF1 levels have not decreased 3 months after treatment. In patients treated with radiotherapy treatment should be withdrawn every 2 years, for 1 month, for assessment of remission.

Treatment should be stopped where there is biochemical evidence of remission (normal IGF1 levels) following treatment withdrawal for at least 4 weeks

Initial application — (pre-operative acromegaly) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has acromegaly; and
- 2 Patient has a large pituitary tumour, greater than 10 mm at its widest; and
- 3 Patient is scheduled to undergo pituitary surgery in the next six months.

Initial application — (Other Indications) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 VIPomas and Glucagonomas for patients who are seriously ill in order to improve their clinical state prior to definitive surgery; or
- 2 Both:
  - 2.1 Gastrinoma: and
  - 2.2 Fither:
    - 2.2.1 Surgery has been unsuccessful; or
    - 2.2.2 Patient has metastatic disease after treatment with H2 antagonist or proton pump inhibitors has been unsuccessful: or
- 3 Both:
  - 3.1 Insulinomas: and
  - 3.2 Surgery is contraindicated or has not been successful; or
- 4 For pre-operative control of hypoglycaemia and for maintenance therapy; or
- 5 Both:
  - 5.1 Carcinoid syndrome (diagnosed by tissue pathology and/or urinary 5HIAA analysis); and
  - 5.2 Disabling symptoms not controlled by maximal medical therapy.

Note: The use of a long-acting somatostatin analogue in patients with fistulae, oesophageal varices, miscellaneous diarrhoea and hypotension will not be funded under Special Authority

Renewal — (Other Indications) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

LANREOTIDE - Special Authority see SA2445 on the previous	us page – Retail phar	macy	
Inj 60 mg per 0.5 ml, 0.5 ml syringe	382.77	1	✓ Mytolac
			✓ Mytolac S29 S29
	1,543.79		✓ Somatuline Autogel
Inj 90 mg per 0.5 ml, 0.5 ml syringe	562.92	1	✓ Mytolac
	2,054.40		<ul> <li>Somatuline Autogel</li> </ul>
Inj 120 mg per 0.5 ml, 0.5 ml syringe	646.70	1	✓ Mytolac
	2,570.44		Somatuline Autogel
OCTREOTIDE LONG-ACTING - Special Authority see SA24	145 on the previous pa	age – Retail	pharmacy
Inj depot 10 mg prefilled syringe	438.40	1	<ul> <li>Sandostatin LAR</li> </ul>
Inj depot 20 mg prefilled syringe	583.70	1	<ul> <li>Sandostatin LAR</li> </ul>
Inj depot 30 mg prefilled syringe	670.80	1	✓ Sandostatin LAR

## **Aromatase Inhibitors**

ANASTROZOLE			
* Tab 1 mg	4.39	30	✓ Anatrole

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
EXEMESTANE  * Tab 25 mg  LETROZOLE	9.86	30	<b>√</b> <u>F</u>	Pfizer Exemestane
* Tab 2.5 mg	4.36	28	<b>✓</b>	Accord S29
- -	4.67	30	<b>√</b> <u>[</u>	<u>_etrole</u>
Immunosuppressants				

# **Cytotoxic Immunosuppressants**

* Tab 25 mg10.15	60	✓ Azamun
Azamun to be Principal Supply on 1 February 2026		
* Tab 50 mg10.34	100	Azamun
Azamun to be Principal Supply on 1 February 2026		
MYCOPHENOLATE MOFETIL		
Tab 500 mg35.90	50	<ul><li>Cellcept</li></ul>
Cap 250 mg35.90	100	✓ Cellcept
Powder for oral lig 1 g per 5 ml - Subsidy by endorsement 187.25	165 ml OP	✓ Cellcept

Mycophenolate powder for oral liquid is subsidised only for patients unable to swallow tablets and capsules, and when the prescription is endorsed accordingly.

#### **Fusion Proteins**

**AZATHIOPRINE** 

ETANERCEPT - Special Authority see SA2399 below - I	Retail pharmacy		
Inj 25 mg	690.00	4	Enbrel
Inj 25 mg autoinjector	690.00	4	<ul><li>Enbrel</li></ul>
Inj 50 mg autoinjector		4	Enbrel
Inj 50 mg prefilled syringe		4	Enbrel

### ⇒SA2399 Special Authority for Subsidy

**Initial application** — (adult-onset Still's disease) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

#### Either:

- 1 Both:
  - 1.1 Either:
    - 1.1.1 The patient has had an initial Special Authority approval for adalimumab for adult-onset Still's disease (AOSD); or
    - 1.1.2 The patient has been started on tocilizumab for AOSD in a Health NZ Hospital; and
  - 1.2 Fither
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or tocilizumab; or
    - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or tocilizumab such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
  - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
  - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal anti-inflammatory drugs (NSAIDs) and methotrexate; and
  - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Approvals valid for 6 months for applications meeting the following criteria:

#### , iii.

- 1 Either:
  - 1.1 Applicant is a rheumatologist; or
  - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 The patient has a sustained improvement in inflammatory markers and functional status.

Initial application — (ankylosing spondylitis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

#### 101.

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for ankylosing spondylitis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for ankylosing spondylitis; or
- 2 All of the following:
  - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis present for more than six months; and
  - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
  - 2.3 Patient has bilateral sacroiliitis demonstrated by plain radiographs, CT or MRI scan; and
  - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
  - 2.5 Either:
    - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
    - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender (see Notes); and
  - 2.6 A Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of initial application.

Average normal chest expansion corrected for age and gender:

18-24 years - Male: 7.0 cm; Female: 5.5 cm

25-34 years - Male: 7.5 cm; Female: 5.5 cm

35-44 years - Male: 6.5 cm; Female: 4.5 cm

45-54 years - Male: 6.0 cm; Female: 5.0 cm

55-64 years - Male: 5.5 cm; Female: 4.0 cm

65-74 years - Male: 4.0 cm; Female: 4.0 cm

75+ years - Male: 3.0 cm: Female: 2.5 cm

**Renewal — (ankylosing spondylitis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
  - 1.1 Applicant is a rheumatologist; or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
 \$	Per 🗸	Manufacturer

continued...

- 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 3 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 4 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (polyarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for polyarticular course JIA; or
- 2 All of the following:
  - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
  - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
  - 2.3 Any of the following:
    - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
    - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
    - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Renewal — (polyarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
  - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for oligoarticular course juvenile idiopathic arthritis (JIA); and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or

<u></u>			
	Subsidy	Fully	Brand or
	(Manufacturer's Price)	Subsidised	Generic
	\$	Per ✓	Manufacturer

continued...

- 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for oligoarticular course JIA; or
- 2 All of the following:
  - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance: and
  - 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
  - 2.3 Any of the following:
    - 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
    - 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose); or
    - 2.3.3 High disease activity (cJADAS10 score greater than 4) after a 6-month trial of methotrexate.

Renewal — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
  - 2 Either:
    - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
    - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

#### Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab or secukinumab for psoriatic arthritis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab or secukinumab; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab or secukinumab to meet the renewal criteria for adalimumab or secukinumab for psoriatic arthritis: or
- 2 All of the following:
  - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
  - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
  - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
  - 2.4 Either:
    - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints;
    - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
  - 2.5 Any of the following:
    - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
    - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

**Renewal** — **(psoriatic arthritis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 Applicant is a rheumatologist; or
  - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

**Initial application — (pyoderma gangrenosum)** only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum\*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Indications marked with \* are unapproved indications.

**Renewal — (pyoderma gangrenosum)** only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

Initial application — (Arthritis - rheumatoid ) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for rheumatoid arthritis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects; or
    - 1.2.2 The patient has received insufficient benefit to meet the renewal criteria for rheumatoid arthritis: or
- 2 All of the following:
  - 2.1 Patient has had rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
  - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
  - 2.3 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
  - 2.4 Patient has tried and not responded to at least three months of methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses unless contraindicated); and
  - 2.5 Fither:

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic	
\$	Per	/	Manufacturer	

continued...

- 2.5.1 Patient has tried and not responded to at least three months of methotrexate in combination with the maximum tolerated dose of ciclosporin; or
- 2.5.2 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with methotrexate: and
- 2.6 Fither:
  - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
  - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip.

Renewal — (Arthritis - rheumatoid) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
  - 2.1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

**Initial application** — (severe chronic plaque psoriasis) only from a dermatologist or any relevant practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria: Fither:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe chronic plaque psoriasis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe chronic plaque psoriasis; or
- 2 All of the following:
  - 2.1 Any of the following:
    - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
    - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; or
    - 2.1.3 Patient has severe chronic localised genital or flexural plaque psoriasis where the plaques or lesions have been present for at least 6 months from the time of initial diagnosis, and with a Dermatology Life Quality Index (DLQI) score greater than 10; and
  - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
  - 2.3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
  - 2.4 The most recent PASI or DLQI assessment is no more than 1 month old at the time of application.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for

Subsidy Fully Brand or
(Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

continued...

severe chronic plaque psoriasis of the face, hand, foot, genital or flexural areas at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and for the face, palm of a hand or sole of a foot the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

**Renewal — (severe chronic plaque psoriasis)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 Both:
    - 1.1.1 Patient had "whole body" severe chronic plague psoriasis at the start of treatment; and
    - 1.1.2 Either:
      - 1.1.2.1 Following each prior etanercept treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-treatment baseline value; or
      - 1.1.2.2 Following each prior etanercept treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline value; or
  - 1.2 Both:
    - 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
    - 1.2.2 Fither:
      - 1.2.2.1 Following each prior etanercept treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values: or
      - 1.2.2.2 Following each prior etanercept treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; or
  - 1.3 Both:
    - 1.3.1 Patient had severe chronic localised genital or flexural plaque psoriasis at the start of treatment; and
    - 1.3.2 Either:
      - 1.3.2.1 The patient has experienced a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; or
      - 1.3.2.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing etanercept; and
- 2 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Note: A treatment course is defined as a minimum of 12 weeks of etanercept treatment

**Initial application** — **(undifferentiated spondyloarthritis)** only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undifferentiated peripheral spondyloarthritis\* with active peripheral joint arthritis in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day (or maximum tolerated dose); and
- 4 Patient has tried and not responded to at least three months of leflunomide at a dose of up to 20 mg daily (or maximum tolerated dose); and
- 5 Any of the following:
  - 5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour measured no more than one month prior to the date of this application; or
- 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Note: Indications marked with \* are unapproved indications.

Renewal — (undifferentiated spondyloarthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Either:
  - 1.1 Applicant is a rheumatologist; or
  - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg dose every 7 days.

#### Immune Modulators

ANTITHYMOCYTE GLOBULIN (EQUINE) - PCT only - Sp	ecialist		
Inj 50 mg per ml, 5 ml	4,439.17	5	✓ ATGAM
BACILLUS CALMETTE-GUERIN (BCG) VACCINE - PCT of	only – Specialist		
Subsidised only for bladder cancer.			
Inj 2-8 × 100 million CFU	149.37	1	✓ OncoTICE
Inj 40 mg per ml, vial	182.45	3	✓ SII-Onco-BCG S29

### Monoclonal Antibodies

ADALIMUMAB (AMGEVITA) – Special Authority see SA2525	below - Retail pharm	acy	
Inj 20 mg per 0.4 ml prefilled syringe	190.00	1	✓ Amgevita
Inj 40 mg per 0.8 ml prefilled pen	375.00	2	✓ Amgevita
Inj 40 mg per 0.8 ml prefilled syringe	375.00	2	✓ Amgevita

**⇒SA2525** Special Authority for Subsidy

Initial application — (Behcet's disease - severe) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Both:

- 1 The patient has severe Behcet's disease\* that is significantly impacting the patient's quality of life; and
- 2 Either:
  - 2.1 The patient has severe ocular, neurological, and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s); or
  - 2.2 The patient has severe gastrointestinal, rheumatological, and/or mucocutaneous symptoms and has not responded adequately to two or more treatments appropriate for the particular symptom(s).

Note: Indications marked with \* are unapproved indications.

Initial application — (Hidradenitis suppurativa) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	sidised	Generic	
\$	Per	1	Manufacturer	

continued

All of the following:

- 1 Patient has hidradenitis suppurativa Hurley Stage II or Hurley Stage III lesions in distinct anatomic areas; and
- 2 Patient has tried, but had an inadequate response to at least a 90 day trial of systemic antibiotics or has demonstrated intolerance to or has contraindications for systemic antibiotics; and
- 3 Patient has 3 or more active lesions; and
- 4 The patient has a DLQI of 10 or more and the assessment is no more than 1 month old at time of application.

**Renewal** — (**Hidradenitis suppurativa**) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline; and
- 2 The patient has a DLQI improvement of 4 or more from baseline.

Initial application — (Plaque psoriasis - severe chronic) only from a dermatologist or any relevant practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria: Either:

- 1 Both:
  - 1.1 Patient has had an initial Special Authority approval for etanercept for severe chronic plaque psoriasis; and
  - 1.2 Either:
    - 1.2.1 Patient has experienced intolerable side effects; or
    - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for etanercept for severe chronic plaque psoriasis; or
- 2 All of the following:
  - 2.1 Any of the following:
    - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a PASI score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
    - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; or
    - 2.1.3 Patient has severe chronic localised genital or flexural plaque psoriasis where the plaques or lesions have been present for at least 6 months from the time of initial diagnosis, and with a Dermatology Life Quality Index (DLQI) score greater than 10; and
  - 2.2 Patient has tried, but had an inadequate response to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
  - 2.3 A PASI assessment or DLQI assessment has been completed for at least the most recent prior treatment course but no longer than 1 month following cessation of each prior treatment course and is no more than 1 month old at the time of application.

Renewal — (Plaque psoriasis - severe chronic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 Both:
  - 1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced a 75% or more reduction in PASI score, or is sustained at this level, when compared with the pre-treatment baseline value; or
    - 1.2.2 The patient has a DLQI improvement of 5 or more, when compared with the pre-treatment baseline value; or

2 Both:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Por 🗸	Manufacturer

continued...

- 2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
- 2.2 Either:
  - 2.2.1 The patient has experienced reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
  - 2.2.2 The patient has experienced reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre treatment baseline value; or
- 3 Both:
  - 3.1 Patient had severe chronic localised genital or flexural plaque psoriasis at the start of treatment; and
  - 3.2 Fither:
    - 3.2.1 The patient has experienced a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; or
    - 3.2.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing adalimumab.

**Initial application — (pyoderma gangrenosum)** only from a dermatologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has pyoderma gangrenosum\*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporin, azathioprine, or methotrexate) and has not received an adequate response.

Note: Indications marked with \* are unapproved indications.

Initial application — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active Crohn's disease: and
- 2 Any of the following:
  - 2.1 Patient has a CDAI score of greater than or equal to 300, or HBI score of greater than or equal to 10; or
  - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
  - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
  - 2.4 Patient has an ileostomy or colostomy and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Renewal — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 CDAI score has reduced by 100 points from the CDAI score, or HBI score has reduced by 3 points, from when the patient was initiated on adalimumab; or
- 2 CDAI score is 150 or less, or HBI is 4 or less; or
- 3 The patient has demonstrated an adequate response to treatment, but CDAI score and/or HBI score cannot be assessed.

Initial application — (Crohn's disease - children) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has active Crohn's disease; and
- 2 Either:
  - 2.1 Patient has a PCDAI score of greater than or equal to 30; or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
 \$	Per 🗸	

continued...

- 2.2 Patient has extensive small intestine disease; and
  - 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Renewal — (Crohn's disease - children) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab; or
- 2 PCDAI score is 15 or less: or
- 3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed.

**Initial application** — **(Crohn's disease - fistulising)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Crohn's disease: and
- 2 Any of the following:
  - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
  - 2.2 Patient has one or more rectovaginal fistula(e); or
  - 2.3 Patient has complex peri-anal fistula; and
- 3 A Baseline Fistula Assessment has been completed and is no more than 1 month old at the time of application.

Renewal — (Crohn's disease - fistulising) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 The number of open draining fistulae have decreased from baseline by at least 50%; or
- 2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain.

Initial application — (Ocular inflammation - chronic) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

. . .

- 1 The patient has had an initial Special Authority approval for infliximab for chronic ocular inflammation; or
- 2 Both:
  - 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
  - 2.2 Any of the following:
    - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective; or
    - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
    - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Renewal — (Ocular inflammation - chronic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 12 weeks' initial treatment; or
- 2 Following each 2 year treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

3 Following each 2 year treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

**Initial application** — (Ocular inflammation - severe) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Patient has had an initial Special Authority approval for infliximab for severe ocular inflammation; or
- 2 Both:
  - 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
  - 2.2 Any of the following:
    - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
    - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
    - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Renewal — (Ocular inflammation - severe) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 2 year treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 2 year treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.</p>

**Initial application — (ankylosing spondylitis)** only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 Patient has had an initial Special Authority approval for etanercept for ankylosing spondylitis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects; or
    - 1.2.2 The patient has received insufficient benefit to meet the renewal criteria for ankylosing spondylitis; or
- 2 All of the following:
  - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis for more than six months; and
  - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
  - 2.3 Patient has bilateral sacroiliitis demonstrated by radiology imaging; and
  - 2.4 Patient has not responded adequately to treatment with two or more NSAIDs, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
  - 2.5 Fither:
    - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following BASMI measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
    - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender; and
  - 2.6 A BASDAI of at least 6 on a 0-10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment and is no more than 1 month old at the time of application.

Renewal — (ankylosing spondylitis) from any relevant practitioner. Approvals valid for 2 years where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

of 50%, whichever is less.

Initial application — (Arthritis - oligoarticular course juvenile idiopathic) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for etanercept for oligoarticular course juvenile idiopathic arthritis (JIA); and
  - 1.2 Fither:
    - 1.2.1 Patient has experienced intolerable side effects; or
    - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for oligoarticular course JIA; or
- 2 All of the following:
  - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
  - 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
  - 23 Fither
    - 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
    - 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose).

Renewal — (Arthritis - oligoarticular course juvenile idiopathic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - polyarticular course juvenile idiopathic) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
  - 1.1 Patient has had an initial Special Authority approval for etanercept for polyarticular course juvenile idiopathic arthritis (JIA); and
  - 1.2 Either:
    - 1.2.1 Patient has experienced intolerable side effects; or
    - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for polyarticular course JIA; or
- 2 All of the following:
  - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
  - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
  - 2.3 Any of the following:
    - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
    - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose): or
    - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Renewal — (Arthritis - polyarticular course juvenile idiopathic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

-	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer	
=	<u> </u>				

continued...

Fither:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - psoriatic) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
  - 1.1 Patient has had an initial Special Authority approval for etanercept or secukinumab for psoriatic arthritis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects; or
    - 1.2.2 The patient has received insufficient benefit from to meet the renewal criteria for psoriatic arthritis; or
- 2 All of the following:
  - 2.1 Patient has had active psoriatic arthritis for six months duration or longer; and
  - 2.2 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
  - 2.3 Patient has tried and not responded to at least three months of sulfasalazine or leflunomide at maximum tolerated doses (unless contraindicated): and
  - 2.4 Either:
    - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
    - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
  - 2.5 Any of the following:
    - 2.5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
    - 2.5.2 Patient has an ESR greater than 25 mm per hour; or
    - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (Arthritis - psoriatic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in swollen joint count from baseline and a clinically significant response in the opinion of the physician; or
- 2 Patient demonstrates at least a continuing 30% improvement in swollen joint count from baseline and a clinically significant response in the opinion of the treating physician.

Initial application — (Arthritis - rheumatoid) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for etanercept for rheumatoid arthritis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects; or
    - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for rheumatoid arthritis; or
- 2 All of the following:

Subsidy (Manufacturer's Price)	Fully Subsidised		
(ivialitulacture) 5 Filoe)	Per 🗸	Manufacturer	
Φ	rei •	Manuacturei	

continued...

- 2.1 Patient has had rheumatoid arthritis (either confirmed by radiology imaging, or the patient is CCP antibody positive) for six months duration or longer; and
- 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2.3 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
- 2.4 Patient has tried and not responded to at least three months of methotrexate in combination with sulfasalazine and hydroxychloroguine sulphate at maximum tolerated doses (unless contraindicated); and
- 2.5 Either:
  - 2.5.1 Patient has tried and not responded to at least three months of methotrexate in combination with the maximum tolerated dose of ciclosporin (unless contraindicated); or
  - 2.5.2 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide (unless contraindicated) alone or in combination with methotrexate; and
- 2.6 Fither:
  - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
  - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip.

Renewal — (Arthritis - rheumatoid) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Initial application — (Still's disease - adult-onset (AOSD)) only from a rheumatologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for etanercept and/or tocilizumab for AOSD; and
  - 1.2 Fither
    - 1.2.1 Patient has experienced intolerable side effects from etanercept and/or tocilizumab; or
    - 1.2.2 Patient has received insufficient benefit from at least a three-month trial of etanercept and/or tocilizumab; or
- 2 All of the following:
  - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria; and
  - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, NSAIDs and methotrexate; and
  - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Initial application — (ulcerative colitis) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active ulcerative colitis: and
- 2 Either:
  - 2.1 Patient's SCCAI score is greater than or equal to 4: or
  - 2.2 Patient's PUCAI score is greater than or equal to 20; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from prior therapy with immunomodulators and systemic corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidi	sed	Generic
\$	Per	1	

continued...

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 The SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on biologic therapy; or
- 2 The PUCAI score has reduced by 10 points or more from the PUCAI score when the patient was initiation on biologic therapy.

Initial application — (undifferentiated spondyloarthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undifferentiated peripheral spondyloarthritis\* with active peripheral joint arthritis in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2 Patient has tried and not responded to at least three months of each of methotrexate, sulfasalazine and leflunomide, at maximum tolerated doses (unless contraindicated); and
- 3 Any of the following:
  - 3.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
  - 3.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application; or
  - 3.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Note: Indications marked with \* are unapproved indications

**Renewal — (undifferentiated spondyloarthritis)** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response in the opinion of the treating physician.

Initial application — (inflammatory bowel arthritis – axial) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has axial inflammatory pain for six months or more; and
- 3 Patient is unable to take NSAIDs; and
- 4 Patient has unequivocal sacroiliitis demonstrated by radiological imaging or MRI; and
- 5 Patient has not responded adequately to prior treatment consisting of at least 3 months of an exercise regime supervised by a physiotherapist; and
- 6 Å BASDAI of at least 6 on a 0-10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment.

**Renewal — (inflammatory bowel arthritis – axial)** from any relevant practitioner. Approvals valid for 2 years where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less.

Initial application — (inflammatory bowel arthritis – peripheral) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and

	lubsidy cturer's Price) Subs	Fully	Brand or Generic
(Waruna	\$ Per	luiseu ✓	Manufacturer

#### continued...

- 2 Patient has active arthritis in at least four joints from the following: hip, knee, ankle, subtalar, tarsus, forefoot, wrist, elbow, shoulder, sternoclavicular; and
- 3 Patient has tried and not experienced a response to at least three months of methotrexate, or azathioprine at a maximum tolerated dose (unless contraindicated); and
- 4 Patient has tried and not experienced a response to at least three months of sulfasalazine at a maximum tolerated dose (unless contraindicated); and
- 5 Any of the following:
  - 5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
  - 5.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application; or
  - 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (inflammatory bowel arthritis – peripheral) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

#### Either:

- 1 Following initial treatment, patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 Patient has experienced at least a continuing 30% improvement in active joint count from baseline in the opinion of the treating physician.

ADALIMUMAB (HUMIRA - ALTERNATIVE BRAND) -	- Special Authority see SA2157	below -	- Retail pharmacy
lei 00 ee ee ee 0 0 ee lee e fille de ee ee	FOF FO	^	/ 11l

Inj 20 mg per 0.2 ml prefilled syringe	595.50	2	<ul><li>Humira</li></ul>
Inj 40 mg per 0.4 ml prefilled pen	595.50	2	✓ HumiraPen
Inj 40 mg per 0.4 ml prefilled syringe	595.50	2	<ul><li>Humira</li></ul>

#### ⇒SA2157 Special Authority for Subsidy

Initial application — (Behcet's disease – severe) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

#### All of the following:

- 1 Either:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Behcet's disease – severe) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

#### Both:

- 1 The patient has had a good clinical response to treatment with measurably improved quality of life; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Hidradenitis suppurativa) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

#### 1 Fither:

1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita: and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 7 days. Fortnightly dosing has been considered.

**Renewal — (Hidradenitis suppurativa)** only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline; and
- 2 The patient has a Dermatology Quality of Life Index improvement of 4 or more from baseline; and
- 3 Adalimumab is to be administered at doses no greater than 40mg every 7 days. Fortnightly dosing has been considered.

Initial application — (Psoriasis - severe chronic plaque) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Psoriasis - severe chronic plaque) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
  - 1.1 Both:
    - 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
    - 1.1.2 Either:
      - 1.1.2.1 Following each prior adalimumab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-adalimumab treatment baseline value; or
      - 1.1.2.2 Following each prior adalimumab treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline value; or
  - 1.2 Both:
    - 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
    - 1.2.2 Either:
      - 1.2.2.1 Following each prior adalimumab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
      - 1.2.2.2 Following each prior adalimumab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-adalimumab treatment baseline value: and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

**Initial application — (Pyoderma gangrenosum)** only from a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 A maximum of 8 doses.

**Renewal — (Pyoderma gangrenosum)** only from a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient has demonstrated clinical improvement and continues to require treatment; and
- 2 A maximum of 8 doses.

Initial application — (Crohn's disease - adult) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevitat; or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
  - 1.3 Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Crohn's disease - adult) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on adalimumab; or
  - 1.2 CDAI score is 150 or less; or
  - 1.3 The patient has demonstrated an adequate response to treatment, but CDAI score cannot be assessed; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Crohn's disease - children) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or

	Subsidy	Fully	Brand or
(Manu	facturer's Price)	Subsidised	Generic
	\$ Pe	er 🗸	Manufacturer

continued...

- 1.3 Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Crohn's disease - children) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab; or
  - 1.2 PCDAI score is 15 or less; or
  - 1.3 The patient has demonstrated an adequate response to treatment, but PCDAI score cannot be assessed; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Crohn's disease - fistulising) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
  - 1.3 Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Crohn's disease - fistulising) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
  - 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Ocular inflammation – chronic) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with Amgevita, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
  - 1.3 Patient has uveitis and is considered to be at risk of vision loss if they were to change treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Ocular inflammation – chronic) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per ✓	Manufacturer

continued...

Both:

- 1 Any of the following:
  - 1.1 The patient has had a good clinical response following 12 weeks' initial treatment; or
  - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
  - 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and</p>
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Ocular inflammation – severe) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita: or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with Amgevita, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
  - 1.3 Patient has uveitis and is considered to be at risk of vision loss if they were to change treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Ocular inflammation – severe) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 The patient has had a good clinical response following 3 initial doses; or
  - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
  - 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and</p>
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita); and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1 Treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Arthritis - oligoarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

Renewal — (Arthritis - oligoarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months where the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from

Initial application — (Arthritis - polyarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

Renewal — (Arthritis - polyarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months where the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - psoriatic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Fither:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Arthritis - psoriatic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

- 1 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Arthritis – rheumatoid) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:
All of the following:

- 1 Either:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Fither:
  - 4.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or
  - 4.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

**Renewal — (Arthritis – rheumatoid)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 2 Either:
  - 2.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or
  - 2.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

Initial application — (Still's disease – adult-onset (AOSD)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

Renewal — (Still's disease – adult-onset (AOSD)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months where the patient has demonstrated a sustained improvement in inflammatory markers and functional status.

AFLIBERCEPT - Special Authority see SA2550 below - Retail pharmacy

Inj 40 mg per ml, 0.1 ml vial.......1,250.00 1 ✓ Eylea

⇒SA2550 Special Authority for Subsidy

**Initial application — (diabetic macular oedema)** from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

1 Patient has centre involving diabetic macular oedema (DMO); and

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sub	sidised	Generic
\$	Per	•	Manufacturer

continued...

- 2 Patient's disease is non responsive to 4 doses of intravitreal bevacizumab when administered 4-6 weekly; and
- 3 Patient has reduced visual acuity between 6/9 6/36 with functional awareness of reduction in vision; and
- 4 Patient has DMO within central OCT (ocular coherence tomography) subfield > 350 micrometers; and
- 5 There is no centre-involving sub-retinal fibrosis or foveal atrophy; and
- 6 Patient has not previously been treated with faricimab for longer than 3 months.

Renewal — (diabetic macular oedema) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 There is stability or two lines of Snellen visual acuity gain; and
- 2 There is structural improvement on OCT scan (with reduction in intra-retinal cysts, central retinal thickness, and sub-retinal fluid); and
- 3 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 4 There is no centre-involving sub-retinal fibrosis or foveal atrophy.

Initial application — (wet age related macular degeneration) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 All of the following:
  - 1.1 Any of the following:
    - 1.1.1 Wet age-related macular degeneration (wet AMD); or
    - 1.1.2 Polypoidal choroidal vasculopathy; or
    - 1.1.3 Choroidal neovascular membrane from causes other than wet AMD; and
  - 1.2 Either:
    - 1.2.1 The patient has developed severe endophthalmitis or severe posterior uveitis following treatment with bevacizumab: or
    - 1.2.2 There is worsening of vision or failure of retina to dry despite three intraocular injections of bevacizumab four weeks apart; and
  - 1.3 There is no structural damage to the central fovea of the treated eye; and
  - 1.4 Patient has not previously been treated with ranibizumab or faricimab for longer than 3 months; or
- 2 Either:
  - 2.1 Patient has current approval to use ranibizumab or faricimab for treatment of wAMD and was found to be intolerant within 3 months; or
  - 2.2 Patient has previously\* (\*before June 2018) received treatment with ranibizumab for wAMD and disease was stable while on treatment.

**Renewal — (wet age related macular degeneration)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Documented benefit must be demonstrated to continue; and
- 2 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 3 There is no structural damage to the central fovea of the treated eye.

BENRALIZUMAB – Special Authority see SA2151 below – Retail pharmacy

⇒SA2151 Special Authority for Subsidy

Initial application — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 Patient must be aged 12 years or older; and

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	1	Manufacturer

#### continued...

- 2 Patient must have a diagnosis of severe eosinophilic asthma documented by a respiratory physician or clinical immunologist: and
- 3 Conditions that mimic asthma eq. vocal cord dysfunction, central airway obstruction, bronchiolitis etc. have been excluded: and
- 4 Patient has a blood eosinophil count of greater than 0.5 × 10^9 cells/L in the last 12 months; and
- 5 Patient must be adherent to optimised asthma therapy including inhaled corticosteroids (equivalent to at least 1000 mcg per day of fluticasone propionate) plus long-acting beta-2 agonist, or budesonide/formoterol as part of the anti-inflammatory reliever therapy plus maintenance regimen, unless contraindicated or not tolerated; and
- 6 Either:
  - 6.1 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral corticosteroids: or
  - 6.2 Patient has received continuous oral corticosteroids of at least the equivalent of 10 mg per day over the previous 3 months: and
- 7 Treatment is not to be used in combination with subsidised mepolizumab; and
- 8 Patient has an Asthma Control Test (ACT) score of 10 or less. Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 52 weeks after the first dose to assess response to treatment; and
- 9 Fither:
  - 9.1 Patient has not previously received an anti-IL5 biological therapy for their severe eosinophilic asthma: or
  - 9.2 Both:
    - 9.2.1 Patient was refractory or intolerant to previous anti-IL5 biological therapy; and
    - 9.2.2 Patient was not eligible to continue treatment with previous anti-IL5 biological therapy and discontinued within 12 months of commencing treatment.

Renewal — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 2 vears for applications meeting the following criteria: Roth:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
  - 2 Fither:
    - 2.1 Exacerbations have been reduced from baseline by 50% as a result of treatment with benralizumab; or
    - 2.2 Reduction in continuous oral corticosteroid use by 50% or by 10 mg/day while maintaining or improving asthma

#### BEVACIZUMAB - PCT only - Special Authority see SA2453 below

✓ Vegzelma	1	j 25 mg per ml, 4 ml vial69.00	
✓ Vegzelma	1	j 25 mg per ml, 16 ml vial276.00	
✓ Baxter	1 mg	j 1 mg for ECP	

#### ⇒SA2453 Special Authority for Subsidy

Initial application — (unresectable hepatocellular carcinoma) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

#### Either:

- 1 Patient is currently on treatment with bevacizumab, and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
  - 2.1 Patient has locally advanced or metastatic, unresectable hepatocellular carcinoma; and
  - 2.2 Patient has preserved liver function (Child-Pugh A); and
  - 2.3 Transarterial chemoembolisation (TACE) is unsuitable; and
  - 2.4 Any of the following:
    - 2.4.1 Patient has not received prior systemic therapy for the treatment of hepatocellular carcinoma; or

Subsidy		Fully	Brand or
(Manufacturer's Price)	Su	bsidised	Generic
 \$	Per	/	Manufacturer

continued...

- 2.4.2 Patient received funded lenvatinib before 1 March 2025; or
- 2.4.3 Both:
  - 2.4.3.1 Patient has experienced treatment-limiting toxicity from treatment with lenvatinib; and
  - 2.4.3.2 No disease progression since initiation of lenvatinib; and
- 2.5 Patient has an ECOG performance status of 0-2; and
- 2.6 To be given in combination with atezolizumab.

Renewal — (unresectable hepatocellular carcinoma) from any relevant practitioner. Approvals valid for 6 months where there is no evidence of disease progression.

**Initial application** — (advanced or metastatic ovarian cancer) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
  - 1.1 The patient has FIGO Stage IV epithelial ovarian, fallopian tube, or primary peritoneal cancer; or
  - 1.2 Both:
    - 1.2.1 The patient has previously untreated advanced (FIGO Stage IIIB or IIIC) epithelial ovarian, fallopian tube, or primary peritoneal cancer; and
    - 1.2.2 Either:
      - 1.2.2.1 Debulking surgery is inappropriate; or
      - 1.2.2.2 The cancer is sub-optimally debulked (maximum diameter of any gross residual disease greater than 1cm); and
- 2 Bevacizumab to be administered at a maximum dose of 15 mg/kg every three weeks; and
- 3 18 weeks concurrent treatment with chemotherapy is planned.

Renewal — (advanced or metastatic ovarian cancer) from any relevant practitioner. Approvals valid for 4 months where there is no evidence of disease progression.

Initial application — (Recurrent Respiratory Papillomatosis) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Maximum of 6 doses: and
- 2 The patient has recurrent respiratory papillomatosis; and
- 3 The treatment is for intra-lesional administration.

Renewal — (Recurrent Respiratory Papillomatosis) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Maximum of 6 doses: and
- 2 The treatment is for intra-lesional administration; and
- 3 There has been a reduction in surgical treatments or disease regrowth as a result of treatment.

Initial application — (Ocular Conditions) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Fither:

- 1 Ocular neovascularisation: or
- 2 Exudative ocular angiopathy.

BRENTUXIMAB VEDOTIN - PCT only - Special Authority see SA2289 below

**⇒SA2289** Special Authority for Subsidy

Initial application — (relapsed/refractory Hodgkin lymphoma) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Subsidy Fully	Brand or
(Manufacturer's Price) Subsidised	Generic
\$ Per ✔	Manufacturer

continued...

All of the following:

- 1 Fither:
  - 1.1 Both:
    - 1.1.1 Patient has relapsed/refractory CD30-positive Hodgkin lymphoma after two or more lines of chemotherapy; and
    - 1.1.2 Patient is ineligible for autologous stem cell transplant; or
  - 1.2 Both:
    - 1.2.1 Patient has relapsed/refractory CD30-positive Hodgkin lymphoma; and
    - 1.2.2 Patient has previously undergone autologous stem cell transplant; and
- 2 Patient has not previously received funded brentuximab vedotin; and
- 3 Response to brentuximab vedotin treatment is to be reviewed after a maximum of 6 treatment cycles; and
- 4 Brentuximab vedotin to be administered at doses no greater than 1.8 mg/kg every 3 weeks.

**Renewal — (relapsed/refractory Hodgkin lymphoma)** from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has achieved a partial or complete response to brentuximab vedotin after 6 treatment cycles; and
- 2 Treatment remains clinically appropriate and the patient is benefitting from treatment and treatment is being tolerated; and
- 3 Patient is to receive a maximum of 16 total cycles of brentuximab vedotin treatment.

**Initial application — (anaplastic large cell lymphoma)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has relapsed/refractory CD30-positive systemic anaplastic large cell lymphoma; and
- 2 Patient has an ECOG performance status of 0-1; and
- 3 Patient has not previously received brentuximab vedotin; and
- 4 Response to brentuximab vedotin treatment is to be reviewed after a maximum of 6 treatment cycles; and
- 5 Brentuximab vedotin to be administered at doses no greater than 1.8 mg/kg every 3 weeks.

Renewal — (anaplastic large cell lymphoma) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has achieved a partial or complete response to brentuximab vedotin after 6 treatment cycles; and
- 2 Treatment remains clinically appropriate and the patient is benefitting from treatment and treatment is being tolerated; and
- 3 Patient is to receive a maximum of 16 total cycles of brentuximab vedotin treatment.

#### 

## ⇒SA2401 Special Authority for Subsidy

Initial application — (head and neck cancer, locally advanced) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, non-metastatic, squamous cell cancer of the head and neck; and
- 2 Cisplatin is contraindicated or has resulted in intolerable side effects; and
- 3 Patient has an ECOG performance score of 0-2; and
- 4 To be administered in combination with radiation therapy.

**Initial application — (colorectal cancer, metastatic)** only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

	Subsidy	0	Fully	Brand or
(M	Ianufacturer's Price)		idised	Generic
	\$	Per	•	Manufacturer

continued...

All of the following:

- 1 Patient has metastatic colorectal cancer located on the left side of the colon (see Note); and
- 2 There is documentation confirming disease is RAS and BRAF wild-type; and
- 3 Patient has an ECOG performance score of 0-2; and
- 4 Patient has not received prior funded treatment with cetuximab; and
- 5 Fither:
  - 5.1 Cetuximab is to be used in combination with chemotherapy; or
  - 5.2 Chemotherapy is determined to not be in the best interest of the patient based on clinician assessment.

Renewal — (colorectal cancer, metastatic) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where there is no evidence of disease progression.

Note: Left-sided colorectal cancer comprises of the distal one-third of the transverse colon, the splenic flexure, the descending colon, the sigmoid colon, or the rectum.

FARICIMAB - Special Authority see SA2533 below - Retail pharmacy

⇒SA2533 Special Authority for Subsidy

**Initial application — (diabetic macular oedema)** from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has centre involving diabetic macular oedema (DMO); and
- 2 Patient's disease is nonresponsive to 4 doses of intravitreal bevacizumab when administered 4-6 weekly; and
- 3 Patient has reduced visual acuity between 6/9 6/36 with functional awareness of reduction in vision; and
- 4 Patient has DMO within central OCT (ocular coherence tomography) subfield > 350 micrometers; and
- 5 There is no centre-involving sub-retinal fibrosis or foveal atrophy; and
- 6 Patient has not previously been treated with aflibercept for longer than 3 months.

Renewal — (diabetic macular oedema) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 There is stability or two lines of Snellen visual acuity gain; and
- 2 There is structural improvement on OCT scan (with reduction in intra-retinal cysts, central retinal thickness, and sub-retinal fluid); and
- 3 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 4 There is no centre-involving sub-retinal fibrosis or foveal atrophy.

**Initial application — (wet age related macular degeneration)** from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 Wet age-related macular degeneration (wet AMD); or
  - 1.2 Polypoidal choroidal vasculopathy; or
  - 1.3 Choroidal neovascular membrane from causes other than wet AMD; and
- 2 Either:
  - 2.1 The patient has developed severe endophthalmitis or severe posterior uveitis following treatment with bevacizumab: or
  - 2.2 There is worsening of vision or failure of retina to dry despite three intraocular injections of bevacizumab four weeks apart; and
- 3 There is no structural damage to the central fovea of the treated eye; and
- 4 Patient has not previously been treated with ranibizumab or aflibercept for longer than 3 months.

Subsidy (Manufacturer's Price)	Full Subsidise	
\$	Per 🗸	Manufacturer

continued...

**Renewal — (wet age related macular degeneration)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

#### Both:

- 1 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 2 There is no structural damage to the central fovea of the treated eye.

## ⇒SA2269 Special Authority for Subsidy

**Initial application** only from a haematologist, paediatric haematologist or paediatric oncologist. Approvals valid for 3 months for applications meeting the following criteria:

### All of the following:

- 1 Patient has not received prior chemotherapy for this condition; and
- 2 Patient has de novo CD33-positive acute myeloid leukaemia; and
- 3 Patient does not have acute promyelocytic leukaemia; and
- 4 Gemtuzumab ozogamicin will be used in combination with standard anthracycline and cytarabine (AraC); and
- 5 Patient is being treated with curative intent; and
- 6 Patient's disease risk has been assessed by cytogenetic testing to be good or intermediate; and
- 7 Patient must be considered eligible for standard intensive remission induction chemotherapy with standard anthracycline and cytarabine (AraC); and
- 8 Gemtuzumab ozogamicin to be funded for one course only (one dose at 3 mg per m² body surface area or up to 2 vials of 5 mg as separate doses).

Note: Acute myeloid leukaemia excludes acute promyelocytic leukaemia and acute myeloid leukaemia that is secondary to another haematological disorder (eg myelodysplasia or myeloproliferative disorder).

INFLIXIMAB - PCT only - Special Authority see SA2487 below

 Inj 100 mg
 428.00
 1
 ✓ Remicade

 Inj 1 mg for ECP
 4.40
 1 mg
 ✓ Baxter

## ⇒SA2487 Special Authority for Subsidy

**Initial application — (Crohn's disease (adults))** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

#### All of the following:

- 1 Patient has active Crohn's disease; and
- 2 Any of the following:
  - 2.1 Patient has a CDAI score of greater than or equal to 300 or HBI score of greater than or equal to 10; or
  - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
  - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
  - 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but has experienced an inadequate response to, or intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Renewal — (Crohn's disease (adults)) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

#### Both:

- 1 Any of the following:
  - 1.1 CDAI score has reduced by 100 points from the CDAI score, or HBI score has reduced by 3 points, from when the patient was initiated on infliximab; or
  - 1.2 CDAI score is 150 or less, or HBI is 4 or less; or

	Subsidy	Fully	Brand or
(Manu	facturer's Price)	Subsidised	Generic
	\$ Pe	er 🗸	Manufacturer

continued...

- 1.3 The patient has demonstrated an adequate response to treatment but CDAI score and/or HBI score cannot be assessed: and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (Crohn's disease (children)) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has active Crohn's disease; and
- 2 Fither:
  - 2.1 Patient has a PCDAI score of greater than or equal to 30; or
  - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but experienced an inadequate response to, or intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Renewal — (Crohn's disease (children)) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on infliximab; or
  - 1.2 PCDAI score is 15 or less: or
  - 1.3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (Graft vs host disease) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has steroid-refractory acute graft vs. host disease of the gut.

Initial application — (Pulmonary sarcoidosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has life-threatening pulmonary sarcoidosis diagnosed by a multidisciplinary team that is refractory to other

Initial application — (acute fulminant ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 weeks for applications meeting the following criteria: Both:

- 1 Patient has acute, fulminant ulcerative colitis; and
- 2 Treatment with intravenous or high dose oral corticosteroids has not been successful.

Initial application — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and 2 Fither:
- - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
  - 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

Subsidy (Manufacturer's Price)	Fully Subsidised		
(ivialitulacture) 5 Filoe)	Per 🗸	Manufacturer	
Φ	rei •	Manuacturei	

continued...

- 1 Following 12 weeks of infliximab treatment, BASDAI has improved by 4 or more points from pre-infliximab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 3 Infliximab to be administered at doses no greater than 5 mg/kg every 6-8 weeks.

**Initial application — (chronic ocular inflammation)** from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for chronic ocular inflammation; and
  - 12 Fither
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for chronic ocular inflammation; or
- 2 Both:
  - 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
  - 2.2 Any of the following:
    - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective: or
    - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
    - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

**Renewal — (chronic ocular inflammation)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses: or
- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (fistulising Crohn's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has confirmed Crohn's disease; and
- 2 Any of the following:
  - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
  - 2.2 Patient has one or more rectovaginal fistula(e); or
  - 2.3 Patent has complex peri-anal fistula.

Renewal — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

1 Fither:

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sub	sidised	Generic
\$	Per	•	Manufacturer

continued...

- 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
- 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline (in the case of adult patients, as demonstrated by a reduction in the Fistula Assessment score), together with less induration and patient reported pain; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (neurosarcoidosis) only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with neurosarcoiosis by a multidisciplinary team; and
- 2 Patient has CNS involvement: and
- 3 Patient has steroid-refractory disease: and
- 4 Either:
  - 4.1 IV cyclophosphamide has been tried; or
  - 4.2 Treatment with IV cyclophosphamide is clinically inappropriate.

Renewal — (neurosarcoidosis) only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria:

Fither:

- 1 A withdrawal period has been tried and the patient has relapsed; or
- 2 All of the following:
  - 2.1 A withdrawal period has been considered but would not be clinically appropriate; and
  - 2.2 There has been a marked reduction in prednisone dose; and
  - 2.3 Either:
    - 2.3.1 There has been an improvement in MRI appearances; or
    - 2.3.2 Marked improvement in other symptomology.

Initial application — (plaque psoriasis) only from a dermatologist or any relevant practitioner on the recommendation of a dermatologist. Approvals valid for 3 months for applications meeting the following criteria:

Fither:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab, etanercept or secukinumab for severe chronic plaque psoriasis; and
  - 1.2 Either:
    - 1.2.1 Patient has experienced intolerable side effects from adalimumab, etanercept or secukinumab; or
    - 1.2.2 Patient has received insufficient benefit from adalimumab, etanercept or secukinumab to meet the renewal criteria for adalimumab, etanercept or secukinumab for severe chronic plaque psoriasis; or
- 2 All of the following:
  - 2.1 Any of the following:
    - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
    - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; or
    - 2.1.3 Patient has severe chronic localised genital or flexural plaque psoriasis where the plaques or lesions have been present for at least 6 months from the time of initial diagnosis, and with a Dermatology Life Quality Index (DLQI) score greater than 10; and

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

- 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
- 2.3 A PASI assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course: and
- 2.4 The most recent PASI assessment is no more than 1 month old at the time of initiation.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand, foot, genital or flexural areas at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and for the face, palm of a hand or sole of a foot the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

**Renewal — (plaque psoriasis)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 Both:
    - 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
    - 1.1.2 Following each prior infliximab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-infliximab treatment baseline value; or
  - 1.2 Both:
    - 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
    - 1.2.2 Fither:
      - 1.2.2.1 Following each prior infliximab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
      - 1.2.2.2 Following each prior infliximab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-infliximab treatment baseline value: or
  - 1.3 Both:
    - 1.3.1 Patient had severe chronic localised genital or flexural plaque psoriasis at the start of treatment; and
    - 1.3.2 Either:
      - 1.3.2.1 The patient has experienced a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; or
      - 1.3.2.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing infliximab; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

**Initial application — (previous use)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient was being treated with infliximab prior to 1 February 2019; and
- 2 Any of the following:
  - 2.1 Rheumatoid arthritis; or
  - 2.2 Ankylosing spondylitis; or
  - 2.3 Psoriatic arthritis: or

Subsidy		Fully	Brand or	
(Manufacturer's Price	e)	Subsidised	Generic	
\$	Per	•	Manufacturer	

continued...

- 2.4 Severe ocular inflammation: or
- 2.5 Chronic ocular inflammation: or
- 2.6 Crohn's disease (adults); or
- 2.7 Crohn's disease (children); or
- 2.8 Fistulising Crohn's disease; or
- 2.9 Severe fulminant ulcerative colitis: or
- 2.10 Severe ulcerative colitis; or
- 2.11 Plaque psoriasis: or
- 2.12 Neurosarcoidosis; or
- 2.13 Severe Behcet's disease.

Initial application — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

- Both:
  - 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis; and
  - 2 Either:
    - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept and/or secukinumab; or
    - 2.2 Following 3-4 months' initial treatment with adalimumab and/or etanercept and/or secukinumab, the patient did not meet the renewal criteria for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior infliximab treatment in the opinion of the treating physician; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
  - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
  - 2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept: and
- 3 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance.

Renewal — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or

continued...

- 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Infliximab to be administered at doses no greater than 3 mg/kg every 8 weeks.

**Initial application — (severe Behcet's disease)** from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe Behcet's disease which is significantly impacting the patient's quality of life (see Notes); and
- 2 Either
  - 2.1 The patient has severe ocular, neurological and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s) (see Notes); or
  - 2.2 The patient has severe gastrointestinal, rheumatologic and/or mucocutaneous symptoms and has not responded adequately to two or more treatment appropriate for the particular symptom(s) (see Notes); and
- 3 The patient is experiencing significant loss of quality of life.

Notes: Behcet's disease diagnosed according to the International Study Group for Behcet's Disease. Lancet 1990;335(8697):1078-80. Quality of life measured using an appropriate quality of life scale such as that published in Gilworth et al J Rheumatol. 2004;31:931-7.

Treatments appropriate for the particular symptoms are those that are considered standard conventional treatments for these symptoms, for example intravenous/oral steroids and other immunosuppressants for ocular symptoms; azathioprine, steroids, thalidomide, interferon alpha and ciclosporin for mucocutaneous symptoms; and colchicine, steroids and methotrexate for rheumatological symptoms.

Renewal — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has had a good clinical response to initial treatment with measurably improved quality of life; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

**Renewal — (fulminant ulcerative colitis)** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Where maintenance treatment is considered appropriate, infliximab should be used in combination with immunomodulators and reassessed every 6 months; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

**Initial application — (severe ocular inflammation)** from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe ocular inflammation; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe ocular inflammation: or
- 2 Both:
  - 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
  - 2.2 Any of the following:

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
\$	Per	1	

continued...

- 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
- 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
- 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Renewal — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

**Initial application — (ulcerative colitis)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active ulcerative colitis; and
- 2 Either:
  - 2.1 Patients SCCAI is greater than or equal to 4; or
  - 2.2 Patients PUCAI score is greater than or equal to 20; and
- 3 Patient has tried but has experienced an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and systemic corticosteroids.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 The SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on infliximab: or
  - 1.2 The PUCAI score has reduced by 10 points or more from the PUCAI score when the patient was initiated on infliximab; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

**Initial application — (pyoderma gangrenosum)** only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum\*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Note: Indications marked with \* are unapproved indications.

**Renewal — (pyoderma gangrenosum)** only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

Initial application — (inflammatory bowel arthritis – axial) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has had axial inflammatory pain for six months or more; and
- 3 Patient is unable to take NSAIDs; and
- 4 Patient has unequivocal sacroiliitis demonstrated by radiological imaging or MRI; and
- 5 Patient's disease has not responded adequately to prior treatment consisting of at least 3 months of an exercise regime supervised by a physiotherapist; and
- 6 Patient has a BASDAI of at least 6 on a 0 10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment.

**Renewal — (inflammatory bowel arthritis – axial)** from any relevant practitioner. Approvals valid for 2 years where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10-point scale, or an improvement in BASDAI of 50%, whichever is less.

Initial application — (inflammatory bowel arthritis – peripheral) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has active arthritis in at least four joints from the following: hip, knee, ankle, subtalar, tarsus, forefoot, wrist, elbow, shoulder, sternoclavicular; and
- 3 Patient has tried and not experienced a response to at least three months of methotrexate or azathioprine at a maximum tolerated dose (unless contraindicated); and
- 4 Patient has tried and not experienced a response to at least three months of sulfasalazine at a maximum tolerated dose (unless contraindicated); and
- 5 Any of the following:
  - 5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
  - 5.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application; or
  - 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (inflammatory bowel arthritis – peripheral) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, patient has experienced at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 Patient has experienced at least a continuing 30% improvement in active joint count from baseline in the opinion of the treating physician.

**Initial application — (immune checkpoint inhibitor toxicity in malignancy\*)** from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

1 The individual requires treatment for moderate to severe autoimmune toxicity following immune checkpoint inhibitor treatment for malignancy; and

Subsidy (Manufacturer's Price)	Full Subsidise	
 \$	Per 🗸	Manufacturer

continued...

- 2 The individual has received insufficient benefit from use of corticosteroids; and
- 3 Infliximab is to be administered at up to 5mg/kg for up to four doses.

Renewal — (immune checkpoint inhibitor toxicity in malignancy\*) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 The individual has shown clinical improvement and ongoing treatment is required; and
- 2 Infliximab is to be administered at up to 5mg/kg for up to a total of 8 doses.

Note: Indications marked with \* are unapproved indications.

INOTUZUMAB OZOGAMICIN - PCT only - Specialist - Special Authority see \$A2460 below

Inj 1 mg vial ......14,457.00 1

Inj 1 mg for ECP ......14,457.00 1 mg

### ⇒SA2460 Special Authority for Subsidy

**Initial application** only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has relapsed or refractory CD22-positive B-cell acute lymphoblastic leukaemia/lymphoma, including minimal residual disease; and
- 2 Patient has ECOG performance status of 0-2; and
- 3 Either:
  - 3.1 Both:
    - 3.1.1 Patient has Philadelphia chromosome positive B-Cell ALL; and
    - 3.1.2 Patient has previously received a tyrosine kinase inhibitor; or
- 3.2 Patient has received one prior line of treatment involving intensive chemotherapy; and
- 4 Treatment is to be administered for a maximum of 3 cycles.

**Renewal** only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient is not proceeding to a stem cell transplant; and
- 2 Fither:
  - 2.1 Patient has experienced complete disease response; or
  - 2.2 Patient has experienced complete remission with incomplete haematological recovery; and
- 3 Treatment with inotuzumab ozogamicin is to cease after a total duration of 6 cycles.

MEPOLIZUMAB - Special Authority see SA2331 below - Retail pharmacy

### ⇒SA2331 Special Authority for Subsidy

Initial application — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Patient must have a diagnosis of severe eosinophilic asthma documented by a respiratory physician or clinical immunologist; and
- 3 Conditions that mimic asthma eg. vocal cord dysfunction, central airway obstruction, bronchiolitis etc. have been excluded; and
- 4 Patient has a blood eosinophil count of greater than 0.5 x 10^9 cells/L in the last 12 months; and
- 5 Patient must be adherent to optimised asthma therapy including inhaled corticosteroids (equivalent to at least 1000 mcg

continued...

✓ Besponsa

✓ Baxter

Subsic		Fully	Brand or
(Manufacture	r's Price) Subs	idised	Generic
\$	Per	1	Manufacturer

continued...

per day of fluticasone propionate) plus long acting beta-2 agonist, or budesonide/formoterol as part of the single maintenance and reliever therapy regimen, unless contraindicated or not tolerated; and

- 6 Either:
  - 6.1 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral corticosteroids; or
  - 6.2 Patient has received continuous oral corticosteroids of at least the equivalent of 10 mg per day over the previous 3 months; and
- 7 Treatment is not to be used in combination with subsidised benralizumab; and
- 8 Patient has an Asthma Control Test (ACT) score of 10 or less. Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 52 weeks after the first dose to assess response to treatment; and
- 9 Either:
  - 9.1 Patient has not previously received an anti-IL5 biological therapy for their severe eosinophilic asthma; or
  - 9.2 Both:
    - 9.2.1 Patient was refractory or intolerant to previous anti-IL5 biological therapy; and
    - 9.2.2 Patient was not eligible to continue treatment with previous anti-IL5 biological therapy and discontinued within 12 months of commencing treatment.

**Renewal — (Severe eosinophilic asthma)** only from a respiratory physician or clinical immunologist. Approvals valid for 2 years for applications meeting the following criteria:

Roth:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 Either:
  - 2.1 Exacerbations have been reduced from baseline by 50% as a result of treatment with mepolizumab; or
  - 2.2 Reduction in continuous oral corticosteroid use by 50% or by 10 mg/day while maintaining or improving asthma control.

Initial application — (eosinophilic granulomatosis with polyangiitis) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has eosinophilic granulomatosis with polyangiitis; and
- 2 The patient has trialled and not received adequate benefit from at least one of the following for at least three months (unless contraindicated to all): azathioprine, cyclophosphamide, leflunomide, methotrexate, mycophenolate, or rituximab; and
- 3 Fither:
  - 3.1 The patient has trialled prednisone for a minimum of three months and is unable to maintain disease control at doses below 7.5 mg per day; or
  - 3.2 Corticosteroids are contraindicated.

Renewal — (eosinophilic granulomatosis with polyangiitis) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months where patient has no evidence of clinical disease progression.

OBINUTUZUMAB - PCT only - Specialist - Special Aut	hority see SA2551 below		
Inj 25 mg per ml, 40 ml vial	5,910.00	1	✓ Gazyva
Inj 1 mg for ECP	6.21	1 mg	✓ Baxter

⇒SA2551 Special Authority for Subsidy

Initial application — (chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for

(Manu	Subsidy	Fully	Brand or
	facturer's Price)	Subsidised	Generic
	\$ P	Per 🗸	Manufacturer

continued...

applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C CD20+ chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is obinutuzumab treatment naive; and
- 3 The patient is not eligible for full dose FCR due to comorbidities with a score > 6 on the Cumulative Illness Rating Scale (CIRS) or reduced renal function (creatinine clearance < 70mL/min); and
- 4 Patient has adequate neutrophil and platelet counts\* unless the cytopenias are a consequence of marrow infiltration by CLL; and
- 5 Patient has good performance status; and
- 6 Obinutuzumab to be administered at a maximum cumulative dose of 8,000 mg and in combination with chlorambucil for a maximum of 6 cycles.

Notes: Chronic lymphocytic leukaemia includes small lymphocytic lymphoma. Comorbidity refers only to illness/impairment other than CLL induced illness/impairment in the patient. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with obinutuzumab is expected to improve symptoms and improve ECOG score to < 2.

\* Neutrophil greater than or equal to  $1.5 \times 10^9$ /L and platelets greater than or equal to  $75 \times 10^9$ /L.

Initial application — (follicular / marginal zone lymphoma) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 Patient has follicular lymphoma; or
  - 1.2 Patient has marginal zone lymphoma; and
- 2 Patient is refractory to or has relapsed within 12 months of a rituximab containing combined chemo-immunotherapy regimen\*; and
- 3 Patient has an ECOG performance status of 0-2; and
- 4 Patient has been previously treated with no more than four chemotherapy regimens; and
- 5 Obinutuzumab to be administered at a maximum dose of 1000 mg for a maximum of 6 cycles in combination with chemotherapy\*.

Note: \* includes unapproved indications

Renewal — (follicular / marginal zone lymphoma) from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 Patient has no evidence of disease progression following obinutuzumab induction therapy; and
- 2 Obinutuzumab to be administered at a maximum of 1000 mg every 2 months for a maximum of 2 years; and
- 3 Obinutuzumab to be discontinued at disease progression.

#### 

### ⇒SA1744 Special Authority for Subsidy

**Initial application** — (severe asthma) only from a respiratory specialist or clinical immunologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 6 years or older; and
- 2 Patient has a diagnosis of severe asthma; and
- 3 Past or current evidence of atopy, documented by skin prick testing or RAST; and
- 4 Total serum human immunoglobulin E (IgE) between 76 IU/mL and 1300 IU/ml at baseline; and

Subsidy	Out	Fully	Brand or	
(Manufacturer's Price)	Sub	sidised	Generic	
\$	Per	•	Manufacturer	

continued...

- 5 Proven adherence with optimal inhaled therapy including high dose inhaled corticosteroid (budesonide 1,600 mcg per day or fluticasone propionate 1,000 mcg per day or equivalent), plus long-acting beta-2 agonist therapy (at least salmeterol 50 mcg bd or eformoterol 12 mcg bd) for at least 12 months, unless contraindicated or not tolerated; and
- 6 Fither:
  - 6.1 Patient has received courses of systemic corticosteroids equivalent to at least 28 days treatment in the past 12 months, unless contraindicated or not tolerated; or
  - 6.2 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral steroids; and
- 7 Patient has an Asthma Control Test (ACT) score of 10 or less; and
- 8 Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 26 weeks after the first dose to assess response to treatment.

**Initial application — (severe chronic spontaneous urticaria)** only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

- All of the following:
  - 1 Patient must be aged 12 years or older; and
  - 2 Either:
    - 2.1 Both:
      - 2.1.1 Patient is symptomatic with Urticaria Activity Score 7 (UAS7) of 20 or above; and
      - 2.1.2 Patient has a Dermatology life quality index (DLQI) of 10 or greater; or
    - 2.2 Patient has a Urticaria Control Test (UCT) of 8 or less; and
  - 3 Any of the following:
    - 3.1 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and ciclosporin (> 3 mg/kg day) for at least 6 weeks; or
    - 3.2 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and at least 3 courses of systemic corticosteroids (> 20 mg prednisone per day for at least 5 days) in the previous 6 months; or
    - 3.3 Patient has developed significant adverse effects whilst on corticosteroids or ciclosporin; and
  - 4 Fither:
    - 4.1 Treatment to be stopped if inadequate response\* following 4 doses; or
    - 4.2 Complete response\* to 6 doses of omalizumab.

**Renewal — (severe asthma)** only from a clinical immunologist or respiratory specialist. Approvals valid for 2 years for applications meeting the following criteria:

#### Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 A reduction in the maintenance oral corticosteroid dose or number of exacerbations of at least 50% from baseline.

**Renewal — (severe chronic spontaneous urticaria)** only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

#### Either:

- 1 Patient has previously adequately responded\* to 6 doses of omalizumab; or
- 2 Both:
  - 2.1 Patient has previously had a complete response\* to 6 doses of omalizumab; and
  - 2.2 Patient has relapsed after cessation of omalizumab therapy.

Note: \*Inadequate response defined as less than 50% reduction in baseline UAS7 and DLQI score, or an increase in Urticaria Control Test (UCT) score of less than 4 from baseline. Patient is to be reassessed for response after 4 doses of omalizumab. Complete response is defined as UAS7 less than or equal to 6 and DLQI less than or equal to 5; or UCT of 16. Relapse of chronic urticaria on stopping prednisone/ciclosporin does not justify the funding of omalizumab.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per Manufacturer

# ⇒SA2419 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Palivizumab to be administered during the annual RSV season; and
- 2 Either:
  - 2.1 Both:
    - 2.1.1 Infant was born in the last 12 months; and
    - 2.1.2 Infant was born at less than 32 weeks zero days' gestation; or
  - 2.2 Both:
    - 2.2.1 Child was born in the last 24 months; and
    - 2.2.2 Any of the following:
      - 2.2.2.1 Child has severe lung, airway, neurological or neuromuscular disease that requires ongoing ventilatory/respiratory support (see Note A) in the community; or
      - 2.2.2.2 Both:
        - 2.2.2.2.1 Child has haemodynamically significant heart disease; and
        - 2.2.2.2.2 Any of the following:
          - 2.2.2.2.2.1 Child has unoperated simple congenital heart disease with significant left to right shunt (see Note B); or
          - 2.2.2.2.2.2 Child has unoperated or surgically palliated complex congenital heart disease; or
          - 2.2.2.2.3 Child has severe pulmonary hypertension (see Note C); or
          - 2.2.2.2.2.4 Child has moderate or severe left ventricular (LV) failure (see Note D); or
      - 2.2.2.3 Child has severe combined immune deficiency, confirmed by an immunologist, but has not received a stem cell transplant; or
      - 2.2.2.4 Child has inborn errors of immunity (see Note E) that increase susceptibility to life-threatening viral respiratory infections, confirmed by an immunologist.

Renewal from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Palivizumab to be administered during the annual RSV season; and
- 2 Child was born in the last 24 months; and
- 3 Any of the following:
  - 3.1 Child has severe lung, airway, neurological or neuromuscular disease that requires ongoing ventilatory/respiratory support (see Note A) in the community; or
  - 3.2 Both:
    - 3.2.1 Child has haemodynamically significant heart disease; and
    - 3.2.2 Any of the following:
      - 3.2.2.1 Child has unoperated simple congenital heart disease with significant left to right shunt (see Note B):
      - 3.2.2.2 Child has unoperated or surgically palliated complex congenital heart disease; or
      - 3.2.2.3 Child has severe pulmonary hypertension (see Note C); or
      - 3.2.2.4 Child has moderate or severe left ventricular (LV) failure (see Note D); or
  - 3.3 Child has severe combined immune deficiency, confirmed by an immunologist, but has not received a stem cell transplant: or
  - 3.4 Child has inborn errors of immunity (see Note E) that increase susceptibility to life-threatening viral respiratory infections, confirmed by an immunologist.

#### Notes:

a) Ventilatory/respiratory support includes those on home oxygen. CPAP/VPAP and those with tracheostomies in situ managed at home

Sub (Manufactu		Fully	Brand or Generic	
`	\$ Per	✓	Manufacturer	

continued...

- b) Child requires/will require heart failure medication, and/or child has significant pulmonary hypertension, and/or infant will require surgical palliation/definitive repair within the next 3 months
- c) Mean pulmonary artery pressure more than 25 mmHg
- d) LV Eiection Fraction less than 40%
- e) Inborn errors of immunity include, but are not limited to, IFNAR deficiencies

### PERTUZUMAB - PCT only - Specialist - Special Authority see SA2276 below

Inj 30 mg per ml, 14 ml vial	3,927.00	1	Perjeta
Inj 420 mg for ECP	3,927.00	420 mg OP	<ul><li>Baxter</li></ul>

### ⇒SA2276 Special Authority for Subsidy

Initial application — (metastatic breast cancer) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Either:
  - 2.1 Patient is chemotherapy treatment naïve; or
  - 2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
- 3 The patient has good performance status (ECOG grade 0-1); and
- 4 Pertuzumab to be administered in combination with trastuzumab; and
- 5 Pertuzumab maximum first dose of 840 mg, followed by maximum of 420 mg every 3 weeks; and
- 6 Pertuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

#### Either:

- 1 Both:
  - 1.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
  - 1.2 The cancer has not progressed at any time point during the previous 12 months whilst on pertuzumab and trastuzumab; or
- 2 All of the following:
  - 2.1 Patient has previously discontinued treatment with pertuzumab and trastuzumab for reasons other than severe toxicity or disease progression; and
  - 2.2 Patient has signs of disease progression; and
  - 2.3 Disease has not progressed during previous treatment with pertuzumab and trastuzumab.

### PERTUZUMAB WITH TRASTUZUMAB - PCT only - Special Authority see SA2534 below

✓ Phesgo	1	l vial7,707.00	Inj 600 mg with trastuzumab 600 mg, 10 ml vial
✓ Phesao	1	ml vial 12.894.00	Ini 1,200 mg with trastuzumab 600 mg, 15 ml vial

# ⇒SA2534 Special Authority for Subsidy

Initial application — (metastatic breast cancer) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Fither:

- 1 Both:
  - 1.1 The individual has received an initial Special Authority approval for intravenous pertuzumab and trastuzumab for metastatic breast cancer; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1.2 Pertuzumab with trastuzumab to be administered subcutaneously at a maximum dose of 600 mg pertuzumab with 600 mg trastuzumab every three weeks (or equivalent); or
- 2 All of the following:
  - 2.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
  - 2.2 Fither:
    - 2.2.1 Patient is chemotherapy treatment naïve; or
    - 2.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
  - 2.3 The patient has good performance status (ECOG grade 0-1); and
  - 2.4 Loading dose of pertuzumab with trastuzumab to be administered subcutaneously at a maximum dose of 1200 mg pertuzumab with 600 mg trastuzumab, respectively; and
  - 2.5 Maintenance doses of pertuzumab with trastuzumab to be administered subcutaneously at a maximum dose of 600 mg pertuzumab with 600 mg trastuzumab every three weeks (or equivalent); and
  - 2.6 Pertuzumab with trastuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The individual has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
  - 1.2 The cancer has not progressed at any time point during the previous 12 months whilst on pertuzumab and trastuzumab; or
- 2 All of the following:
  - 2.1 Individual has previously discontinued treatment with pertuzumab with trastuzumab for reasons other than severe toxicity or disease progression; and
  - 2.2 Individual has signs of disease progression; and
  - 2.3 Disease has not progressed during previous treatment with pertuzumab with trastuzumab.

# RITUXIMAB (MABTHERA) – PCT only – Specialist – Special Authority see SA2552 below Ini 100 mg per 10 ml vial

my rooming por room via management		_	· mastrora
Inj 500 mg per 50 ml vial	2,688.30	1	Mabthera
Inj 1 mg for ECP	5.64	1 mg	✓ Baxter (Mabthera)

### ⇒SA2552 Special Authority for Subsidy

Initial application — (rheumatoid arthritis - TNF inhibitors contraindicated) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Treatment with a Tumour Necrosis Factor alpha inhibitor is contraindicated; and
- 2 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
- 5 Any of the following:
  - 5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with

continued...

/ Mahthera

Subsidy	Full	y Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per 🗸	Manufacturer	

continued...

the maximum tolerated dose of ciclosporin; or

- 5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
- 5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
- 6 Fither:
  - 6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
  - 6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 7 Either:
  - 7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
  - 7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months; and
- 8 Either:
  - 8.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
  - 8.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 9 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Initial application — (rheumatoid arthritis - prior TNF inhibitor use) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Both:
  - 1.1 The patient has had an initial community Special Authority approval for at least one of etanercept and/or adalimumab for rheumatoid arthritis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
    - 1.2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
  - 2.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
  - 2.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 3 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Renewal — (rheumatoid arthritis - re-treatment in 'partial responders' to rituximab) from any relevant practitioner.

Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 At 4 months following the initial course of rituximab infusions the patient had between a 30% and 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 1.2 At 4 months following the second course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 1.3 At 4 months following the third and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Fither
  - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or

Subsidy	:)	Fully	Brand or
(Manufacturer's Price		Subsidised	Generic
<u> </u>	Per	✓	Manufacturer

continued...

- 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Renewal — (rheumatoid arthritis - re-treatment in 'responders' to rituximab) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 At 4 months following the initial course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 1.2 At 4 months following the second and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
  - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

RITUXIMAB (RIXIMYO) - PCT only - Specialist - Special Authority see SA2497 below

Inj 100 mg per 10 ml vial	275.33	2	✓ Riximyo
Inj 500 mg per 50 ml vial	688.20	1	✓ Riximyo
Inj 1 mg for ECP	1.38	1 mg	✓ Baxter (Riximyo)

# ⇒SA2497 Special Authority for Subsidy

Initial application — (ABO-incompatible organ transplant) from any relevant practitioner. Approvals valid without further renewal unless notified where patient is to undergo an ABO-incompatible solid organ transplant\*.

Note: Indications marked with \* are unapproved indications.

Initial application — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis\*; and
- 2 The total rituximab dose would not exceed the equivalent of 375 mg/m<sup>2</sup> of body-surface area per week for a total of 4 weeks; and
- 3 Any of the following:
  - 3.1 Induction therapy with daily oral or pulse intravenous cyclophosphamide has failed to achieve significant improvement of disease after at least 3 months; or
  - 3.2 Patient has previously had a cumulative dose of cyclophosphamide > 15 g or a further repeat 3 month induction course of cyclophosphamide would result in a cumulative dose > 15 g; or
  - 3.3 Cyclophosphamide and methotrexate are contraindicated; or
  - 3.4 Patient is a female of child-bearing potential; or
  - 3.5 Patient has a previous history of haemorrhagic cystitis, urological malignancy or haematological malignancy.

Note: Indications marked with \* are unapproved indications.

Renewal — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis\*: and
- 2 Patient has previously responded to treatment with rituximab but is now experiencing an acute flare of vasculitis; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m<sup>2</sup> of body-surface area per week for a total of 4 weeks.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$
Per ✓ Manufacturer

continued...

Note: Indications marked with \* are unapproved indications.

**Initial application** — (Antibody-mediated organ transplant rejection) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has been diagnosed with antibody-mediated organ transplant rejection\*.

Note: Indications marked with \* are unapproved indications.

**Initial application — (Chronic lymphocytic leukaemia)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C chronic lymphocytic leukaemia (CLL) requiring treatment; and
- 2 Any of the following:
  - 2.1 The patient is rituximab treatment naive: or
  - 2.2 Either:
    - 2.2.1 The patient is chemotherapy treatment naive; or
    - 2.2.2 Both:
      - 2.2.2.1 The patient's disease has relapsed following no more than three prior lines of chemotherapy treatment; and
      - 2.2.2.2 The patient has had a treatment-free interval of 12 months or more if previously treated with fludarabine and cyclophosphamide chemotherapy; or
  - 2.3 The patient's disease has relapsed and rituximab treatment is to be used in combination with funded venetoclax; and
- 3 The patient has good performance status; and
- 4 Either:
  - 4.1 The patient does not have chromosome 17p deletion CLL; or
  - 4.2 Rituximab treatment is to be used in combination with funded venetoclax for relapsed/refractory chronic lymphocytic leukaemia: and
- 5 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles; and
- 6 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration), bendamustine or venetoclax.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with rituximab is expected to improve symptoms and improve ECOG score to < 2.

Renewal — (Chronic lymphocytic laukaemia) from any relevant practitioner. Approvals walled for 12 months for applications.

Renewal — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

#### 1 Either:

- 1.1 The patient's disease has relapsed and rituximab treatment is to be used in combination with funded venetoclax; or
- 1.2 All of the following:
  - 1.2.1 The patient's disease has relapsed following no more than one prior line of treatment with rituximab for CLL; and
  - 1.2.2 The patient has had an interval of 36 months or more since commencement of initial rituximab treatment; and
  - 1.2.3 The patient does not have chromosome 17p deletion CLL; and
  - 1.2.4 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration) or bendamustine; and
- 2 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles.

Subsidy		Fully	Brand or	
(Manufacturer's Price	)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initial application — (Neuromyelitis Optica Spectrum Disorder(NMOSD)) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

#### Both:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks; and
- 2 Either:
  - 2.1 The patient has experienced a severe episode or attack of NMOSD (rapidly progressing symptoms and clinical investigations supportive of a severe attack of NMOSD); or
  - 2.2 All of the following:
    - 2.2.1 The patient has experienced a breakthrough attack of NMOSD; and
    - 2.2.2 The patient is receiving treatment with mycophenolate; and
    - 2.2.3 The patients is receiving treatment with corticosteroids.

Renewal — (Neuromyelitis Optica Spectrum Disorder) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks: and
- 2 The patients has responded to the most recent course of rituximab; and
- 3 The patient has not received rituximab in the previous 6 months.

Initial application — (Post-transplant) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has B-cell post-transplant lymphoproliferative disorder\*; and
- 2 To be used for a maximum of 8 treatment cycles.

Note: Indications marked with \* are unapproved indications.

Renewal — (Post-transplant) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has B-cell post-transplant lymphoproliferative disorder\*: and
- 3 To be used for no more than 6 treatment cycles.

Note: Indications marked with \* are unapproved indications.

Initial application — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
- 2 Fither:
  - 2.1 Treatment with corticosteroids and at least one other immunosuppressant for at least a period of 12 months has been ineffective: or
  - 2.2 Both:
    - 2.2.1 Treatment with at least one other immunosuppressant for a period of at least 12 months; and
    - 2.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

Subsidy	Fully	Brand or	_
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

**Renewal — (Severe Refractory Myasthenia Gravis)** only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Fither:
  - 3.1 The patient has relapsed despite treatment with corticosteroids and at least one other immunosuppressant for a period of at least 12 months; or
  - 3.2 Both:
    - 3.2.1 The patient's myasthenia gravis has relapsed despite treatment with at least one immunosuppressant for a period of at least 12 months; and
    - 3.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects

Initial application — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is a child with SDNS\* or FRNS\*; and
- 2 Treatment with steroids for at least a period of 3 months has been ineffective or associated with evidence of steroid toxicity; and
- 3 Treatment with ciclosporin for at least a period of 3 months has been ineffective and/or discontinued due to unacceptable side effects; and
- 4 Treatment with mycophenolate for at least a period of 3 months with no reduction in disease relapses; and
- 5 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with \* are unapproved indications.

Renewal — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome\*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with \* are unapproved indications.

Initial application — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is a child with SRNS\* where treatment with steroids and ciclosporin for at least 3 months have been ineffective; and
- 2 Treatment with tacrolimus for at least 3 months has been ineffective; and
- 3 Genetic causes of nephrotic syndrome have been excluded; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with \* are unapproved indications.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

Renewal — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome\*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m<sup>2</sup> of body surface area per week for a total of 4 weeks

Note: Indications marked with \* are unapproved indications.

Initial application — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Fither:

- 1 All of the following:
  - 1.1 The patient has treatment naive aggressive CD20 positive NHL; and
  - 1.2 To be used with a multi-agent chemotherapy regimen given with curative intent; and
  - 1.3 To be used for a maximum of 8 treatment cycles; or
- 2 Both:
  - 2.1 The patient has aggressive CD20 positive NHL with relapsed disease following prior chemotherapy; and
  - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Renewal — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has relapsed refractory/aggressive CD20 positive NHL; and
- 3 To be used with a multi-agent chemotherapy regimen given with curative intent; and
- 4 To be used for a maximum of 4 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia Initial application — (haemophilia with inhibitors) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria: Any of the following:

- 1 Patient has mild congenital haemophilia complicated by inhibitors; or
- 2 Patient has severe congenital haemophilia complicated by inhibitors and has failed immune tolerance therapy; or
- 3 Patient has acquired haemophilia.

Renewal — (haemophilia with inhibitors) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for haemophilia with inhibitors; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment.

Initial application — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Either:
  - 1.1 Patient has immune thrombocytopenic purpura\* with a platelet count of less than or equal to 20,000 platelets per microlitre; or

Subsidy	Fu	,	
(Manufacturer's Price)	Subsidis	ed Generic	
\$	Per	<ul> <li>Manufacturer</li> </ul>	

continued...

- 1.2 Patient has immune thrombocytopenic purpura\* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding; and
- 2 Any of the following:
  - 2.1 Treatment with steroids and splenectomy have been ineffective; or
  - 2.2 Treatment with steroids has been ineffective and splenectomy is an absolute contraindication; or
  - 2.3 Other treatments including steroids have been ineffective and patient is being prepared for elective surgery (e.g. splenectomy); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks

Note: Indications marked with \* are unapproved indications.

Renewal — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Fither:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
  - 2.1 Patient was previously treated with rituximab for immune thrombocytopenic purpura\*; and
  - 2.2 An initial response lasting at least 12 months was demonstrated; and
  - 2.3 Patient now requires repeat treatment.

Note: Indications marked with \* are unapproved indications.

Initial application — (indolent, low-grade lymphomas or hairy cell leukaemia\*) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

Fither:

- 1 Both:
  - 1.1 The patient has indolent low grade NHL or hairy cell leukaemia\* with relapsed disease following prior chemotherapy; and
  - 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
  - 2.1 The patient has indolent, low grade lymphoma or hairy cell leukaemia\* requiring first-line systemic chemotherapy; and
  - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. \*Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Renewal — (indolent, low-grade lymphomas or hairy cell leukaemia\*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has indolent, low-grade NHL or hairy cell leukaemia\* with relapsed disease following prior chemotherapy; and
- 3 To be used for no more than 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. \*Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Initial application — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient has autoimmune pure red cell aplasia\* associated with a demonstrable B-cell lymphoproliferative disorder.

Note: Indications marked with \* are unapproved indications.

Renewal — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient was previously treated with rituximab for pure red cell aplasia\*

	ubsidy	. ,	Brand or
(Manufac	cturer's Price)	Subsidised	Generic
	\$ Per	r 🗸	Manufacturer

continued...

associated with a demonstrable B-cell lymphoproliferative disorder and demonstrated an initial response lasting at least

Note: Indications marked with \* are unapproved indications.

Initial application — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Patient has cold haemagglutinin disease\*; and
- 2 Patient has severe disease which is characterized by symptomatic anaemia, transfusion dependence or disabling circulatory symptoms; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks

Note: Indications marked with \* are unapproved indications.

Renewal — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m<sup>2</sup> weekly for 4 weeks) is now planned; or
- 2 All of the following:
  - 2.1 Patient was previously treated with rituximab for severe cold haemagglutinin disease\*; and
  - 2.2 An initial response lasting at least 12 months was demonstrated; and
  - 2.3 Patient now requires repeat treatment.

Note: Indications marked with \* are unapproved indications.

Initial application — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Both:

- 1 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks: and
- 2 Either:
  - 2.1 Patient has thrombotic thrombocytopenic purpura\* and has experienced progression of clinical symptoms or persistent thrombocytopenia despite plasma exchange; or
  - 2.2 Patient has acute idiopathic thrombotic thrombocytopenic purpura\* with neurological or cardiovascular pathology.

Note: Indications marked with \* are unapproved indications.

Renewal — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for thrombotic thrombocytopenic purpura\*: and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment: and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with \* are unapproved indications.

Initial application — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe, immediately life- or organ-threatening SLE\*; and
- 2 The disease has proved refractory to treatment with steroids at a dose of at least 1 mg/kg; and

Subsidy	Ful	y Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per •	<ul> <li>Manufacturer</li> </ul>	

continued...

- 3 The disease has relapsed following prior treatment for at least 6 months with maximal tolerated doses of azathioprine, mycophenolate mofetil and high dose cyclophosphamide, or cyclophosphamide is contraindicated; and
- 4 Maximum of four 1000 mg infusions of rituximab.

Note: Indications marked with \* are unapproved indications.

Renewal — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient's SLE\* achieved at least a partial response to the previous round of prior rituximab treatment; and
- 2 The disease has subsequently relapsed; and
- 3 Maximum of two 1000 mg infusions of rituximab.

Note: Indications marked with \* are unapproved indications.

Initial application — (warm autoimmune haemolytic anaemia (warm AlHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Patient has warm autoimmune haemolytic anaemia\*; and
- 2 One of the following treatments has been ineffective: steroids (including if patient requires ongoing steroids at doses equivalent to > 5 mg prednisone daily), cytotoxic agents (e.g. cyclophosphamide monotherapy or in combination), intravenous immunoglobulin: and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with \* are unapproved indications.

Renewal — (warm autoimmune haemolytic anaemia (warm AIHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
  - 2.1 Patient was previously treated with rituximab for warm autoimmune haemolytic anaemia\*; and
  - 2.2 An initial response lasting at least 12 months was demonstrated; and
  - 2.3 Patient now requires repeat treatment.

Note: Indications marked with \* are unapproved indications.

**Initial application — (severe antisynthetase syndrome)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed antisynthetase syndrome; and
- 2 Patient has severe, immediately life or organ threatening disease, including interstitial lung disease; and
- 3 Either:
  - 3.1 Treatment with at least 3 immunosuppressants (oral steroids, cyclophosphamide, methotrexate, mycophenolate, ciclosporin, azathioprine) has not be effective at controlling active disease; or
  - 3.2 Rapid treatment is required due to life threatening complications; and
- 4 Maximum of four 1.000mg infusions of rituximab.

**Renewal — (severe antisynthetase syndrome)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in inflammatory markers, muscle strength and pulmonary function; and

Subsidy		Fully	Brand or
(Manufacturer's Price)	9	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 2 The patient has not received rituximab in the previous 6 months; and
- 3 Maximum of two cycles of 2 x 1,000mg infusions of rituximab given two weeks apart.

Initial application — (graft versus host disease) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has refractory graft versus host disease following transplant; and
- 2 Treatment with at least 3 immunosuppressants (oral steroids, ciclosporin, tacrolimus, mycophenolate, sirolimus) has not be effective at controlling active disease; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

**Initial application** — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has severe chronic inflammatory demyelinating polyneuropathy (CIPD); and
- 2 Fither
  - 2.1 Both:
    - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
    - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
  - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function compared to baseline; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Initial application — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe anti-NMDA receptor autoimmune encephalitis: and
- 2 Either:
  - 2.1 Both:
    - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
    - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
  - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

**Renewal — (anti-NMDA receptor autoimmune encephalitis)** only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Su	ıbsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 The patient has experienced a relapse and now requires further treatment; and
- 4 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Initial application — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has CD20+ low grade or follicular B-cell NHL with relapsed disease following prior chemotherapy; and
  - 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
  - 2.1 The patient has CD20+ low grade or follicular B-cell NHL requiring first-line systemic chemotherapy; and
  - 2.2 To be used for a maximum of 6 treatment cycles.

Renewal — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 Rituximab is to be used for maintenance in CD20+ low grade or follicular B-cell NHL following induction with first-line systemic chemotherapy; and
- 2 Patient is intended to receive rituximab maintenance therapy for 2 years at a dose of 375 mg/m2 every 8 weeks (maximum of 12 cycles).

Initial application — (Membranous nephropathy) only from a nephrologist or any relevant practitioner on the recommendation of a nephrologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 Patient has biopsy-proven primary/idiopathic membranous nephropathy\*; or
  - 1.2 Patient has PLA2 antibodies with no evidence of secondary cause, and an eGFR of > 60ml/min/1,73m2; and
- 2 Patient remains at high risk of progression to end-stage kidney disease despite more than 3 months of treatment with conservative measures (see Note): and
- 3 The total rituximab dose would not exceed the equivalent of 375mg/m2 of body surface area per week for a total of 4 weeks

Renewal — (Membranous nephropathy) only from a nephrologist or any relevant practitioner on the recommendation of a nephrologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for membranous nephropathy\*; and
- 2 Either:
  - 2.1 Treatment with rituximab was previously successful, but the condition has relapsed, and the patient now requires repeat treatment; or
  - 2.2 Patient achieved partial response to treatment and requires repeat treatment (see Note); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Notes:

- a) Indications marked with \* are unapproved indications.
- b) High risk of progression to end-stage kidney disease defined as > 5g/day proteinuria.
- c) Conservative measures include renin-angiotensin system blockade, blood-pressure management, dietary sodium and protein restriction, treatment of dyslipidaemia, and anticoagulation agents unless contraindicated or the patient has

Subsidy		Fully	Brand or	
(Manufacturer's Pric	ce)	Subsidised	Generic	
\$	Per	1	Manufacturer	

continued...

experienced intolerable side effects.

d) Partial response defined as a reduction of proteinuria of at least 50% from baseline, and between 0.3 grams and 3.5 grams per 24 hours.

Initial application — (B-cell acute lymphoblastic leukaemia/lymphoma\*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient has newly diagnosed B-cell acute lymphoblastic leukaemia/lymphoma\*; and
- 2 Treatment must be in combination with an intensive chemotherapy protocol with curative intent; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m² per dose for a maximum of 18 doses.

Note: Indications marked with \* are unapproved indications.

Initial application — (desensisation prior to transplant) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires desensitisation prior to mismatched allogenic stem cell transplant\*; and
- 2 Patient would receive no more than two doses at 375 mg/m2 of body-surface area.

Note: Indications marked with \* are unapproved indications.

Initial application — (pemiphigus\*) only from a dermatologist or relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 All of the following:
  - 1.1 Patient has severe rapidly progressive pemphigus; and
  - 1.2 Is used in combination with systemic corticosteroids (20 mg/day); and
  - 1.3 Any of the following:
    - 1.3.1 Skin involvement is at least 5% body surface area; or
    - 1.3.2 Significant mucosal involvement (10 or more mucosal erosions) or diffuse gingivitis or confluent large erosions; or
    - 1.3.3 Involvement of two or more mucosal sites; or
- 2 Both:
  - 2.1 Patient has pemphigus; and
  - 2.2 Patient has not experienced adequate clinical benefit from systemic corticosteroids (20 mg/day) in combination with a steroid sparing agent, unless contraindicated.

Note: Indications marked with \* are unapproved indications.

Renewal — (pemiphigus\*) only from a dermatologist or relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has experienced adequate clinical benefit from rituximab treatment, with improvement in symptoms and healing of skin ulceration and reduction in corticosteroid requirement; and
- 2 Patient has not received rituximab in the previous 6 months.

Note: Indications marked with \* are unapproved indications.

Initial application — (immunoglobulin G4-related disease (IgG4-RD\*)) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed diagnosis of IgG4-RD\*; and
- 2 Either:
  - 2.1 Treatment with corticosteroids and/or disease modifying anti-rheumatic drugs for at least 3 months has been ineffective in lowering corticosteroid dose below 5 mg per day (prednisone equivalent) without relapse; or

	Subsidy	Fi	ılly	Brand or
(Man	ufacturer's Price)	Subsidis	ed	Generic
	\$	Per	<u> </u>	Manufacturer

continued...

- 2.2 Treatment with corticosteroids and/or disease modifying anti-rheumatic drugs is contraindicated or associated with evidence of toxicity or intolerance; and
- 3 Total rituximab dose used should not exceed a maximum of two 1000 mg infusions of rituximab given two weeks apart. Note: Indications marked with \* are unapproved indications.

Renewal — (immunoglobulin G4-related disease (IgG4-RD\*)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 Treatment with rituximab for IgG4-RD\* was previously successful and patient's disease has demonstrated sustained response, but the condition has relapsed; or
  - 1.2 Patient is receiving maintenance treatment for IgG4-RD\*; and
- 2 Rituximab re-treatment not to be given within 6 months of previous course of treatment; and
- 3 Maximum of two 1000 mg infusions of rituximab given two weeks apart.

Note: Indications marked with \* are unapproved indications.

SECUKINUMAB - Special Authority see SA2488 below - Retail pharmacy

# **⇒SA2488** Special Authority for Subsidy

Initial application — (severe chronic plaque psoriasis – second-line biologic) only from a dermatologist or any relevant practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab or etanercept, or has trialled infliximab in a Health NZ Hospital, for severe chronic plaque psoriasis; and
- 2 Either:
  - 2.1 The patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
  - 2.2 The patient has received insufficient benefit from adalimumab, etanercept or infliximab; and
- 3 A Psoriasis Area and Severity Index (PASI) assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Initial application — (severe chronic plaque psoriasis – first-line biologic) only from a dermatologist or any relevant practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
  - 1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; or
  - 1.3 Patient has severe chronic localised genital or flexural plaque psoriasis where the plaques or lesions have been present for at least 6 months from the time of initial diagnosis, and with a Dermatology Life Quality Index (DLQI) score greater than 10: and
- 2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin: and

Subsidy	y Full	/ Brand or
(Manufacturer's	s Price) Subsidise	d Generic
\$	Per 🗸	Manufacturer

continued...

- 3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course: and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Note: A treatment course is defined as a minimum of 12 weeks of treatment. "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand. foot, genital or flexural areas, at least 2 of the 3 PASI symptom sub scores for erythema, thickness and scaling are rated as severe or very severe, and for the face, palm of a hand or sole of a foot the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis - first and second-line biologic) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Either:
  - 1.1 Either:
    - 1.1.1 Patient's PASI score has reduced by 75% or more (PASI 75) as compared to baseline PASI prior to commencing secukinumab: or
    - 1.1.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing secukinumab: or
  - 1.2 Both:
    - 1.2.1 Patient had severe chronic localised genital or flexural plague psoriasis at the start of treatment; and
    - 1.2.2 Either:
      - 1.2.2.1 The patient has experienced a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; or
      - 1.2.2.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing secukinumab; and
- 2 Secukinumab to be administered at a maximum dose of 300 mg monthly.

Initial application — (ankylosing spondylitis – second-line biologic) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and
- 2 Either:
  - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
  - 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

Renewal — (ankylosing spondylitis – second-line biologic) only from a rheumatologist or medical practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Following 12 weeks initial treatment of secukinumab treatment, BASDAI has improved by 4 or more points from pre-secukinumab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefitted from treatment and that continued treatment is appropriate; and
- 3 Secukinumab to be administered at doses no greater than 300 mg monthly.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

1 Roth:

Subsidy		Fully	Brand or
(Manufacturer's Price)		idised	Generic
<b>\$</b>	Per	1	Manufacturer

#### continued...

- 1.1 Patient has had an initial Special Authority approval for adalimumab, etanercept or infliximab for psoriatic arthritis; and
- 1.2 Either:
  - 1.2.1 Patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
  - 1.2.2 Patient has received insufficient benefit from adalimumab, etanercept or infliximab to meet the renewal criteria for adalimumab, etanercept or infliximab for psoriatic arthritis; or

#### 2 All of the following:

- 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
- 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
- 2.4 Fither:
  - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
  - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2.5 Any of the following:
  - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
  - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
  - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior secukinumab treatment in the opinion of the treating physician; and
- 2 Secukinumab to be administered at doses no greater than 300 mg monthly.

# SILTUXIMAB - Special Authority see SA1596 below - Retail pharmacy

### ⇒SA1596 Special Authority for Subsidy

**Initial application** only from a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's Disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Siltuximab is to be administered at doses no greater than 11 mg/kg every 3 weeks.

**Renewal** only from a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has sustained improvement in inflammatory markers and functional status.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
TOCILIZUMAB - PCT only - Special Authority see SA2489 belo	W			
Inj 20 mg per ml, 4 ml vial	220.00	1	✓	Actemra
Inj 20 mg per ml, 10 ml vial	550.00	1	✓	Actemra
Inj 20 mg per ml, 20 ml vial		1	✓	Actemra
Inj 1 mg for ECP		1 mg	✓	Baxter

### ⇒SA2489 Special Authority for Subsidy

Initial application — (cytokine release syndrome) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Either:

- 1 Both:
  - 1.1 The patient has developed grade 3 or 4 cytokine release syndrome associated with the administration of blinatumomab for the treatment of acute lymphoblastic leukaemia; and
  - 1.2 Tocilizumab is to be administered at doses no greater than 8 mg/kg IV for a maximum of 3 doses (if less than 30kg, maximum of 12 mg/kg); or
- 2 All of the following:
  - 2.1 The patient is enrolled in the Malaghan Institute of Medical Research ENABLE trial programme: and
  - 2.2 The patient has developed CRS or Immune Effector Cell-Associated Neurotoxicity Syndrome (ICANS) following CAR T-cell therapy for the treatment of relapsed or refractory B-cell non-Hodgkin lymphoma; and
  - 2.3 Tocilizumab is to be administered according to the consensus guidelines for CRS or ICANS for CAR T-cell therapy at doses no greater than 8 mg/kg IV for a maximum of 3 doses.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

#### Both:

- 1 Patient was being treated with tocilizumab prior to 1 February 2019; and
- 2 Any of the following:
  - 2.1 rheumatoid arthritis: or
  - 2.2 systemic juvenile idiopathic arthritis; or
  - 2.3 adult-onset Still's disease: or
  - 2.4 polyarticular juvenile idiopathic arthritis; or
  - 2.5 idiopathic multicentric Castleman's disease.

Initial application — (Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

# All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Fither:
  - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
  - 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis: and
- 3 Fither:
  - 3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or
  - 3.2 Both:
    - 3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a Health NZ Hospital; and
    - 3.2.2 Either:
      - 3.2.2.1 The patient has experienced intolerable side effects from rituximab; or
      - 3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Subsidy		Fully	Brand or	_
(Manufacturer's Price)	Subs	idised	Generic	
\$	Per	1	Manufacturer	

continued...

Initial application — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 2 Tocilizumab is to be used as monotherapy: and
- 3 Either:
  - 3.1 Treatment with methotrexate is contraindicated; or
  - 3.2 Patient has tried and did not tolerate oral and/or parenteral methotrexate; and
- 4 Either:
  - 4.1 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of ciclosporin alone or in combination with another agent; or
  - 4.2 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of leflunomide alone or in combination with another agent; and
- 5 Either:
  - 5.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 active, swollen, tender joints;
  - 5.2 Patient has persistent symptoms of poorly controlled and active disease in at least four active joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 6 Fither:
  - 6.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
  - 6.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Initial application — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient diagnosed with systemic juvenile idiopathic arthritis: and
- 2 Patient has tried and not responded to a reasonable trial of all of the following, either alone or in combination: oral or parenteral methotrexate; non-steroidal anti-inflammatory drugs (NSAIDs); and systemic corticosteroids.

Initial application — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

# 1 Both:

- 1.1 Fither:
  - 1.1.1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for adult-onset Still's disease (AOSD); or
  - 1.1.2 The patient has been started on tocilizumab for AOSD in a Health NZ Hospital; and
- 1.2 Either:
  - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
  - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
  - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
  - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal antiinflammatory drugs (NSAIDs) and methotrexate; and
  - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Subsidy (Manufacturer's Price)	Fully Subsidised	
 \$	Per 🗸	Manufacturer

continued...

Initial application — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for both etanercept and adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
  - 1.2 The patient has experienced intolerable side effects, or has received insufficient benefit from, both etanercept and adalimumab; or
- 2 All of the following:
  - 2.1 Treatment with a tumour necrosis factor alpha inhibitor is contraindicated; and
  - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
  - 2.3 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
  - 2.4 Any of the following:
    - 2.4.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
    - 2.4.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose): or
    - 2.4.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Initial application — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Tocilizumab to be administered at doses no greater than 8 mg/kg IV every 3-4 weeks.

Initial application — (moderate to severe COVID-19) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed (or probable) COVID-19; and
- 2 Oxygen saturation of < 92% on room air, or requiring supplemental oxygen; and
- 3 Patient is receiving adjunct systemic corticosteroids, or systemic corticosteroids are contraindicated; and
- 4 Tocilizumab is to be administered at doses no greater than 8mg/kg IV for a maximum of one dose; and
- 5 Tocilizumab is not to be administered in combination with barcitinib.

**Renewal — (Rheumatoid Arthritis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Renewal — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Following up to 6 months' initial treatment, the patient has achieved at least an American College of Rheumatology paediatric 30% improvement criteria (ACR Pedi 30) response from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing ACR Pedi 30 response from baseline.

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsi	dised	Generic
\$	Per	✓	Manufacturer

continued...

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months where the patient has a sustained improvement in inflammatory markers and functional status. Renewal — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
  - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Renewal — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has a sustained improvement in inflammatory markers and functional status.

Initial application — (immune checkpoint inhibitor toxicity in malignancy\*) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The individual requires treatment for moderate to severe autoimmune toxicity following immune checkpoint inhibitor treatment for malignancy; and
- 2 The individual has received insufficient benefit from use of corticosteroids; and
- 3 Tocilizumab is to be administered at a maximum dose of 8 mg/kg fortnightly.

Renewal — (immune checkpoint inhibitor toxicity in malignancy\*) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 The individual has shown clinical improvement and ongoing treatment is required; and
- 2 Tocilizumab is to be administered at a maximum dose of 8 mg/kg fortnightly.

Note: Indications marked with \* are unapproved indications.

TRASTUZUMAB (HERZUMA) - PCT only - Special Authority see SA2293 below

Inj 150 mg vial100.00	1	✓ Herzuma
Inj 440 mg vial293.35	1	✓ Herzuma
Ini 1 ma for ECP	1 ma	✓ Baxter

### ⇒SA2293 Special Authority for Subsidy

**Initial application** — (early breast cancer) from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria:

Both:

- 1 The patient has early breast cancer expressing HER-2 IHC 3+ or ISH + (including FISH or other current technology); and
- 2 Maximum cumulative dose of 106 mg/kg (12 months' treatment).

**Renewal — (early breast cancer\*)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 All of the following:
  - 1.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
  - 1.2 The patient received prior adjuvant trastuzumab treatment for early breast cancer; and
  - 1.3 Any of the following:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1.3.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
- 1.3.2 The patient discontinued lapatinib within 3 months due to intolerable side effects and the cancer did not progress whilst on lapatinib; or
- 1.3.3 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 1.4 Fither:
  - 1.4.1 Trastuzumab will not be given in combination with pertuzumab; or
  - 1.4.2 All of the following:
    - 1.4.2.1 Trastuzumab to be administered in combination with pertuzumab; and
    - 1.4.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
    - 1.4.2.3 The patient has good performance status (ECOG grade 0-1); and
- 1.5 Trastuzumab to be discontinued at disease progression; or
- 2 All of the following:
  - 2.1 Patient has previously discontinued treatment with trastuzumab in the metastatic setting for reasons other than severe toxicity or disease progression; and
  - 2.2 Patient has signs of disease progression; and
  - 2.3 Disease has not progressed during previous treatment with trastuzumab.

Note: \* For patients with relapsed HER-2 positive disease who have previously received adjuvant trastuzumab for early breast cancer

**Initial application — (metastatic breast cancer)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Either:
  - 2.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
  - 2.2 The patient discontinued lapatinib within 3 months due to intolerable side effects and the cancer did not progress whilst on lapatinib; and
- 3 Fither:
  - 3.1 Trastuzumab will not be given in combination with pertuzumab; or
  - 3.2 All of the following:
    - 3.2.1 Trastuzumab to be administered in combination with pertuzumab; and
    - 3.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer: and
    - 3.2.3 The patient has good performance status (ECOG grade 0-1); and
- 4 Trastuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Fither:

- 1 All of the following:
  - 1.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
  - 1.2 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
  - 1.3 Trastuzumab to be discontinued at disease progression; or
- 2 All of the following:

Subsic		Fully	Brand or
(Manufacture	r's Price) Subs	idised	Generic
\$	Per	1	Manufacturer

continued...

- 2.1 Patient has previously discontinued treatment with trastuzumab for reasons other than severe toxicity or disease progression; and
- 2.2 Patient has signs of disease progression; and
- 2.3 Disease has not progressed during previous treatment with trastuzumab.

Initial application — (gastric, gastro-oesophageal junction and oesophageal cancer) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has locally advanced or metastatic gastric, gastro-oesophageal junction or oesophageal cancer expressing HER-2 IHC 2+ FISH+ or IHC3+ (or other current technology); and
- 2 Patient has an ECOG score of 0-2.

Renewal — (gastric, gastro-oesophageal junction and oesophageal cancer) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 2 Trastuzumab to be discontinued at disease progression.

TRASTUZUMAB DERUXTECAN - PCT only - Special Authority see SA2420 below

Inj 100 mg per ml, 1 ml vial	2,550.00	1	Enhertu
Inj 1 mg for ECP	27.05	1 mg	<ul><li>Baxter</li></ul>

### ⇒SA2420 Special Authority for Subsidy

Initial application only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Patient is currently on treatment with trastuzumab deruxtecan and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
  - 2.1 Patient has metastatic breast cancer expressing HER-2 IHC3+ or ISH+ (including FISH or other current technology); and
  - 2.2 Patient has previously received trastuzumab and chemotherapy, separately or in combination; and
  - 2.3 Either:
    - 2.3.1 The patient has received prior therapy for metastatic disease; or
    - 2.3.2 The patient developed disease recurrence during, or within six months of completing adjuvant therapy; and
  - 2.4 Patient has a good performance status (ECOG 0-1); and
  - 2.5 Patient has not received prior funded trastuzumab deruxtecan treatment; and
  - 2.6 Treatment to be discontinued at disease progression.

**Renewal** only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The cancer has not progressed at any time point during the previous approval period whilst on trastuzumab deruxtecan; and
- 2 Treatment to be discontinued at disease progression.

Note: Prior or adjuvant therapy includes anthracycline, other chemotherapy, biological drugs, or endocrine therapy.

TRASTUZUMAB EMTANSINE - PCT only - Specialist - Special Authority see SA2424 on the next page

Inj 100 mg vial2,320.0		✓ Kadcyla
Inj 160 mg vial3,712.0	00 1	✓ Kadcyla
Inj 1 mg for ECP24.5	52 1 mg	✓ Baxter

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

# **⇒SA2424** Special Authority for Subsidy

Initial application — (early breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has early breast cancer expressing HER2 IHC3+ or ISH+; and
- 2 Documentation of pathological invasive residual disease in the breast and/or axiliary lymph nodes following completion of surgery: and
- 3 Patient has completed systemic neoadjuvant therapy with trastuzumab and chemotherapy prior to surgery; and
- 4 Disease has not progressed during neoadjuvant therapy; and
- 5 Patient has left ventricular ejection fraction of 45% or greater; and
- 6 Adjuvant treatment with trastuzumab emtansine to be commenced within 12 weeks of surgery; and
- 7 Trastuzumab emtansine to be discontinued at disease progression; and
- 8 Total adjuvant treatment duration must not exceed 42 weeks (14 cycles).

Initial application — (metastatic breast cancer) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Patient has previously received trastuzumab and chemotherapy, separately or in combination; and
- 3 Fither:
  - 3.1 The patient has received prior therapy for metastatic disease\*; or
  - 3.2 The patient developed disease recurrence during, or within six months of completing adjuvant therapy\*; and
- 4 Patient has a good performance status (ECOG 0-1); and
- 5 Either:
  - 5.1 Patient does not have symptomatic brain metastases; or
  - 5.2 Patient has brain metastases and has received prior local CNS therapy; and
- 6 Fither:
  - 6.1 Patient has not received prior funded trastuzumab emtansine or trastuzumab deruxtecan treatment; or
  - 62 Roth
    - 6.2.1 Patient has discontinued trastuzumab deruxtecan due to intolerance; and
    - 6.2.2 The cancer did not progress while on trastuzumab deruxtecan; and
- 7 Treatment to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

#### Both:

- 1 The cancer has not progressed at any time point during the previous approval period whilst on trastuzumab emtansine; and
- 2 Treatment to be discontinued at disease progression.

Note: Prior or adjuvant therapy includes anthracycline, other chemotherapy, biological drugs, or endocrine therapy.

USTEKINUMAB - Special Authority see SA2182 below - Retail pharmacy

Inj 90 mg per ml, 1 ml pre-filled syringe.......4,162.00 1 ✓ Stelara

# ⇒SA2182 Special Authority for Subsidy

**Initial application** — (**Crohn's disease - adults**) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

1 Patient is currently on treatment with ustekinumab commenced prior to 1 February 2023 and met all remaining criteria (criterion 2) below at the time of commencing treatment; or

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

continued...

- 2 Both:
  - 2.1 Patient has active Crohn's disease: and
  - 22 Fither:
    - 2.2.1 Patient has had an initial approval for prior biologic therapy for Crohn's disease and has experienced intolerable side effects or insufficient benefit to meet renewal criteria: or
    - 2.2.2 Both:
      - 2.2.2.1 Patient meets the initiation criteria for prior biologic therapies for Crohn's disease; and
      - 2.2.2.2 Other biologics for Crohn's disease are contraindicated.

Renewal — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 CDAI score has reduced by 100 points, or HBI score has reduced by 3 points, from when the patient was initiated on biologic therapy; or
  - 1.2 CDAI score is 150 or less, or HBI is 4 or less; or
  - 1.3 The patient has experienced an adequate response to treatment, but CDAI score and/or HBI score cannot be assessed: and
- 2 Ustekinumab to be administered at a dose no greater than 90 mg every 8 weeks.

Initial application — (Crohn's disease - children\*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Either:

- 1 Patient is currently on treatment with ustekinumab commenced prior to 1 February 2023 and met all remaining criteria (criterion 2) below at the time of commencing treatment; or
- 2 Both:
  - 2.1 Patient has active Crohn's disease; and
  - 2.2 Either:
    - 2.2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria; or
    - 2.2.2 Both:
      - 2.2.2.1 Patient meets the initiation criteria for prior biologic therapies for Crohn's disease; and
      - 2.2.2.2 Other biologics for Crohn's disease are contraindicated.

Note: Indication marked with \* is an unapproved indication.

Renewal — (Crohn's disease - children\*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 PCDAI score has reduced by 10 points from when the patient was initiated on biologic therapy; or
  - 1.2 PCDAI score is 15 or less; or
  - 1.3 The patient has experienced an adequate response to treatment, but CDAI score cannot be assessed; and
- 2 Ustekinumab to administered at a dose no greater than 90 mg every 8 weeks.

Note: Indication marked with \* is an unapproved indication.

Initial application — (ulcerative colitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Patient is currently on treatment with ustekinumab commenced prior to 1 February 2023 and met all remaining criteria (criterion 2) below at the time of commencing treatment; or
- 2 Roth:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 2.1 Patient has active ulcerative colitis; and
- 2.2 Fither:
  - 2.2.1 Patient has had an initial approval for prior biologic therapy for ulcerative colitis and has experienced intolerable side effects or insufficient benefit to meet renewal criteria: or
  - 2.2.2 Both:
    - 2.2.2.1 Patient meets the initiation criteria for prior biologic therapies for ulcerative colitis; and 2.2.2.2 Other biologics for ulcerative colitis are contraindicated.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 The SCCAI score has reduced by 2 points or more from the SCCAI score since initiation on biologic therapy; or
  - 1.2 PUCAI score has reduced by 10 points or more from the PUCAI score since initiation on biologic therapy\*; and
- 2 Ustekinumab will be used at a dose no greater than 90 mg intravenously every 8 weeks.

Note: Criterion marked with \* is for an unapproved indication.

VEDOLIZUMAB - PCT only - Special Authority see SA2183 below

Inj 300 mg vial .......3,313.00 ✓ Entyvio

# ⇒SA2183 Special Authority for Subsidy

Initial application — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active Crohn's disease; and
- 2 Any of the following:
  - 2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria (unless contraindicated); or
  - 2.2 Patient has a CDAI score of greater than or equal to 300, or HBI score of greater than or equal to 10; or
  - 2.3 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
  - 2.4 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection;
  - 2.5 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Any of the following:
  - 3.1 Patient has tried but experienced an inadequate response to (including lack of initial response and/or loss of initial response) from prior therapy with immunomodulators and corticosteroids; or
  - 3.2 Patient has experienced intolerable side effects from immunomodulators and corticosteroids; or
  - 3.3 Immunomodulators and corticosteroids are contraindicated.

Renewal — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 CDAI score has reduced by 100 points, or HBI score has reduced by 3 points, from when the patient was initiated on biologic therapy; or
  - 1.2 CDAI score is 150 or less, or HBI is 4 or less; or
  - 1.3 The patient has experienced an adequate response to treatment, but CDAI score and/or HBI score cannot be assessed: and
- 2 Vedolizumab to administered at a dose no greater than 300 mg every 8 weeks.

Initial application — (Crohn's disease - children\*) from any relevant practitioner. Approvals valid for 6 months for applications

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

meeting the following criteria:

All of the following:

- 1 Paediatric patient has active Crohn's disease; and
- 2 Any of the following:
  - 2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria (unless contraindicated); or
  - 2.2 Patient has a Paediatric Crohn's Disease Activity Index (PCDAI) score of greater than or equal to 30; or
  - 2.3 Patient has extensive small intestine disease; and
- 3 Any of the following:
  - 3.1 Patient has tried but experienced an inadequate response to (including lack of initial response and/or loss of initial response) from prior therapy with immunomodulators and corticosteroids; or
  - 3.2 Patient has experienced intolerable side effects from immunomodulators and corticosteroids; or
  - 3.3 Immunomodulators and corticosteroids are contraindicated.

Note: Indication marked with \* is an unapproved indication.

Renewal — (Crohn's disease - children\*) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 PCDAI score has reduced by 10 points from when the patient was initiated on biologic therapy; or
  - 1.2 PCDAI score is 15 or less: or
  - 1.3 The patient has experienced an adequate response to treatment, but CDAI score cannot be assessed; and
- 2 Vedolizumab to administered at a dose no greater than 300mg every 8 weeks.

Note: Indication marked with \* is an unapproved indication.

Initial application — (ulcerative colitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active ulcerative colitis: and
- 2 Any of the following:
  - 2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria (unless contraindicated): or
  - 2.2 Patient has a SCCAI score is greater than or equal to 4; or
  - 2.3 Patient's PUCAI score is greater than or equal to 20\*; and
- 3 Any of the following:
  - 3.1 Patient has tried but experienced an inadequate response to (including lack of initial response and/or loss of initial response) from prior therapy with immunomodulators and corticosteroids; or
  - 3.2 Patient has experienced intolerable side effects from immunomodulators and corticosteroids; or
  - 3.3 Immunomodulators and corticosteroids are contraindicated.

Note: Indication marked with \* is an unapproved indication.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 The SCCAI score has reduced by 2 points or more from the SCCAI score since initiation on biologic therapy; or
  - 1.2 The PUCAI score has reduced by 10 points or more from the PUCAI score since initiation on biologic therapy \*; and
- 2 Vedolizumab will be used at a dose no greater than 300 mg intravenously every 8 weeks.

Note: Indication marked with \* is an unapproved indication.

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Por 🗸	Manufacturer	

# Programmed Cell Death-1 (PD-1) Inhibitors

ATEZOLIZUMAB - PCT only - Specialist - Special Authori	ity see SA2443 below		
Inj 60 mg per ml, 20 ml vial	9,503.00	1	<ul><li>Tecentriq</li></ul>
Inj 1 mg for ECP	8.08	1 mg	✓ Baxter

### ⇒SA2443 Special Authority for Subsidy

Initial application — (non-small cell lung cancer second line monotherapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced or metastatic non-small cell lung cancer; and
- 2 Patient has not received prior funded treatment with an immune checkpoint inhibitor for NSCLC; and
- 3 For patients with non-squamous histology there is documentation confirming that the disease does not express activating mutations of EGFR or ALK tyrosine kinase unless not possible to ascertain; and
- 4 Patient has an ECOG 0-2; and
- 5 Patient has documented disease progression following treatment with at least two cycles of platinum-based chemotherapy; and
- 6 Atezolizumab is to be used as monotherapy at a dose of 1200 mg every three weeks (or equivalent) for a maximum of 16 weeks; and
- 7 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Renewal — (non-small cell lung cancer second line monotherapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Any of the following:
  - 1.1 Patient's disease has had a complete response to treatment; or
  - 1.2 Patient's disease has had a partial response to treatment; or
  - 1.3 Patient has stable disease; and
- 2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period; and
- 3 No evidence of disease progression; and
- 4 The treatment remains clinically appropriate and patient is benefitting from treatment; and
- 5 Atezolizumab to be used at a maximum dose of 1200 mg every three weeks (or equivalent); and
- 6 Treatment with atezolizumab to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initial application — (unresectable hepatocellular carcinoma) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

#### Either:

- 1 Patient is currently on treatment with atezolizumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
  - 2.1 Patient has locally advanced or metastatic, unresectable hepatocellular carcinoma; and
  - 2.2 Patient has preserved liver function (Child-Pugh A); and
  - 2.3 Transarterial chemoembolisation (TACE) is unsuitable; and
  - 2.4 Any of the following:
    - 2.4.1 Patient has not received prior systemic therapy for the treatment of hepatocellular carcinoma; or
    - 2.4.2 Patient received funded lenvatinib before 1 March 2025; or
    - 2.4.3 Both:
      - 2.4.3.1 Patient has experienced treatment-limiting toxicity from treatment with lenvatinib; and
      - 2.4.3.2 No disease progression since initiation of lenvatinib; and

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic	
\$	Per	1	Manufacturer	

continued...

- 2.5 Patient has an ECOG performance status of 0-2; and
- 2.6 To be given in combination with bevacizumab.

Renewal — (unresectable hepatocellular carcinoma) from any relevant practitioner. Approvals valid for 6 months where there is no evidence of disease progression.

DURVALUMAB - PCT only - Specialist - Special Authority	y see SA2425 below		
Inj 50 mg per ml, 10 ml vial	4,700.00	1	Imfinzi
Inj 50 mg per ml, 2.4 ml vial	1,128.00	1	Imfinzi
Inj 1 mg for ECP	9.59	1 mg	✓ Baxter

# ⇒SA2425 Special Authority for Subsidy

Initial application — (Non-small cell lung cancer) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Fither:
  - 1.1 Patient has histologically or cytologically documented stage III, locally advanced, unresectable non-small cell lung cancer (NSCLC): or
  - 1.2 Patient has histologically or cytologically documented stage IIb (T1N2a only), locally advanced, unresectable non-small cell lung cancer (NSCLC); and
- 2 Patient has received two or more cycles of platinum-based chemotherapy concurrently with definitive radiation therapy; and
- 3 Patient has no disease progression following the second or subsequent cycle of platinum-based chemotherapy with definitive radiation therapy treatment; and
- 4 Patient has a ECOG performance status of 0 or 1; and
- 5 Patient has completed last radiation dose within 8 weeks of starting treatment with durvalumab; and
- 6 Patient must not have received prior PD-1 or PD-L1 inhibitor therapy for this condition; and
- 7 Either:
  - 7.1 Durvalumab is to be used at a maximum dose of no greater than 10 mg/kg every 2 weeks; or
  - 7.2 Durvalumab is to be used at a flat dose of 1500 mg every 4 weeks; and
- 8 Treatment with durvalumab to cease upon signs of disease progression.

Renewal — (Non-small cell lung cancer) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The treatment remains clinically appropriate and the patient is benefitting from treatment; and
- 2 Fither:
  - 2.1 Durvalumab is to be used at a maximum dose of no greater than 10 mg/kg every 2 weeks; or
  - 2.2 Durvalumab is to be used at a flat dose of 1500 mg every 4 weeks; and
- 3 Treatment with durvalumab to cease upon signs of disease progression; and
- 4 Total continuous treatment duration must not exceed 12 months.

IPILIMUMAB - PCT only - Specialist - Special Authorit	y see SA2461 below		
Inj 5 mg per ml, 10 ml vial	5,000.00	1	✓ Yervoy
Inj 5 mg per ml, 40 ml vial	20,000.00	1	✓ Yervoy
Inj 1 mg for ECP	106.00	1 mg	✓ Baxter

# ⇒SA2461 Special Authority for Subsidy

**Initial application** — **(renal cell carcinoma)** from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

#### continued...

- 1 The patient is currently on treatment with ipilimumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
  - 2.1 The patient has metastatic renal cell carcinoma; and
  - 2.2 The patient is treatment naive; and
  - 2.3 The patient has ECOG performance status 0-2; and
  - 2.4 The disease is predominantly of clear cell histology; and
  - 2.5 Any of the following:
    - 2.5.1 The patient has sarcomatoid histology; or
    - 2.5.2 Haemoglobin levels less than the lower limit of normal; or
    - 2.5.3 Corrected serum calcium level greater than 10 mg/dL (2.5 mmol/L); or
    - 2.5.4 Neutrophils greater than the upper limit of normal; or
    - 2.5.5 Platelets greater than the upper limit of normal; or
    - 2.5.6 Interval of less than 1 year from original diagnosis to the start of systemic therapy; or
    - 2.5.7 Karnofsky performance score of less than or equal to 70; and
  - $2.6 \ \ lpilimum ab \ is \ to \ be \ used \ at \ a \ maximum \ dose \ of \ 1 \ mg/kg \ for \ up \ to \ four \ cycles \ in \ combination \ with \ nivolumab..$

# NIVOLUMAB - PCT only - Specialist - Special Authority see SA2490 below

Inj 10 mg per ml, 4 ml vial	1,051.98	1	Opdivo
Inj 10 mg per ml, 10 ml vial	2,629.96	1	Opdivo
Inj 1 mg for ECP	27.22	1 mg	✓ Baxter

# ⇒SA2490 Special Authority for Subsidy

Initial application — (unresectable or metastatic melanoma) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The individual has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Baseline measurement of overall tumour burden is documented clinically and radiologically; and
- 3 The individual has ECOG performance 0-2; and
- 4 Either:
  - 4.1 The individual has not received funded pembrolizumab; or
  - 4.2 Both:
    - 4.2.1 The individual has received an initial Special Authority approval for pembrolizumab and has discontinued pembrolizumab within 12 weeks of starting treatment due to intolerance; and
    - 4.2.2 The cancer did not progress while the indvidual was on pembrolizumab; and
- 5 Any of the following:
  - 5.1 The individual has been diagnosed in the metastatic or unresectable stage III or IV setting; or
  - 5.2 The individual did not receive treatment in the perioperative setting with a PD-1/PD-L1 inhibitor; or
  - 5.3 All of the following:
    - 5.3.1 The individual received treatment in the perioperative setting with a PD-1/PD-L1 inhibitor; and
    - 5.3.2 The individual did not experience disease recurrence while on treatment with that PD-1/PD-L1 inhibitor; and
    - 5.3.3 The individual did not experience disease recurrence within six months of completing perioperative treatment with a PD-1/PD-L1 inhibitor.

Renewal — (unresectable or metastatic melanoma, less than 24 months on treatment) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

# Either:

1 Both:

1.1 Any of the following:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1.1.1 The individual's disease has had a complete response to treatment; or
- 1.1.2 The individual's disease has had a partial response to treatment; or
- 1.1.3 The individual has stable disease; and
- 1.2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period; or
- 2 All of the following:
  - 2.1 The individual has previously discontinued treatment with nivolumab for reasons other than severe toxicity or disease progression; and
  - 2.2 The individual has signs of disease progression; and
  - 2.3 Disease has not progressed during previous treatment with nivolumab.

Renewal — (unresectable or metastatic melanoma, more than 24 months on treatment) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 The individual has been on treatment for more than 24 months; and
- 2 Fither:
  - 2.1 Both:
    - 2.1.1 Any of the following:
      - 2.1.1.1 The individual's disease has had a complete response to treatment; or
      - 2.1.1.2 The individual's disease has had a partial response to treatment; or
      - 2.1.1.3 The individual has stable disease; and
    - 2.1.2 Response to treatment in target lesions has been determined by comparable radiologic or clinical assessment following the most recent treatment period; or
    - 2.2 All of the following:
      - 2.2.1 The individual has previously discontinued treatment with nivolumab for reasons other than severe toxicity or disease progression; and
      - 2.2.2 The individual has signs of disease progression; and
      - 2.2.3 Disease has not progressed during previous treatment with nivolumab.

Initial application — (renal cell carcinoma, first line) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

#### Either:

- 1 Patient is currently on treatment with nivolumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
  - 2.1 The patient has metastatic renal cell carcinoma; and
  - 2.2 The patient is treatment naive; and
  - 2.3 The patient has ECOG performance status 0-2; and
  - 2.4 The disease is predominantly of clear cell histology; and
  - 2.5 Any of the following:
    - 2.5.1 The patient has sarcomatoid histology; or
    - 2.5.2 Haemoglobin levels less than the lower limit of normal: or
    - 2.5.3 Corrected serum calcium level greater than 10 mg/dL (2.5 mmol/L); or
    - 2.5.4 Neutrophils greater than the upper limit of normal; or
    - 2.5.5 Platelets greater than the upper limit of normal; or
    - 2.5.6 Interval of less than 1 year from original diagnosis to the start of systemic therapy; or
    - 2.5.7 Karnofsky performance score of less than or equal to 70; and
  - 2.6 Nivolumab is to be used in combination with ipilimumab for the first four treatment cycles at a maximum dose of 3 mg/kg; and

 Subsidy Manufacturer's Price)	٤	Fully Subsidised	Brand or Generic
 \$	Per	1	Manufacturer

continued...

2.7 Nivolumab is to be used as monotherapy at a maximum maintenance dose of 240 mg every 2 weeks (or equivalent).

**Initial application — (Renal cell carcinoma, second line)** from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has metastatic renal-cell carcinoma; and
- 2 The disease is of predominant clear-cell histology; and
- 3 Patient has ECOG performance status 0-2; and
- 4 Patient has documented disease progression following one or two previous regimens of antiangiogenic therapy; and
- 5 Patient has not previously received a funded immune checkpoint inhibitor; and
- 6 Nivolumab is to be used as monotherapy at a maximum dose of 240 mg every 2 weeks (or equivalent) and discontinued at disease progression.

Renewal — (Renal cell carcinoma) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 Patient's disease has had a complete response to treatment; or
  - 1.2 Patient's disease has had a partial response to treatment; or
  - 1.3 Patient has stable disease: and
- 2 No evidence of disease progression; and
- 3 Nivolumab is to be used as monotherapy at a maximum dose of 240 mg every 2 weeks (or equivalent) and discontinued at disease progression.

PEMBROLIZUMAB - PCT only - Specialist - Special Authority see SA2553 below

 Inj 25 mg per ml, 4 ml vial.
 4,680.00
 1
 ✓ Keytruda

 Inj 1 mg for ECP
 47.74
 1 mg
 ✓ Baxter

⇒SA2553 Special Authority for Subsidy

Initial application — (stage III or IV resectable melanoma - neoadjuvant) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The individual has resectable stage IIIB, IIIC, IIID or IV melanoma (excluding uveal) (see note); and
- 2 The individual has not received prior funded systemic treatment in the perioperative setting for their stage IIIB, IIIC, IIID or IV melanoma; and
- 3 Treatment must be prior to complete surgical resection; and
- 4 Pembrolizumab must be administered as monotherapy; and
- 5 The individual has ECOG performance score 0-2; and
- 6 Pembrolizumab to be administered at a fixed dose of 200 mg every 3 weeks (or equivalent).

Renewal — (stage III or IV resectable melanoma - neoadjuvant) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

Any of the following:

- 1 Both:
  - 1.1 The individual has received neoadjuvant treatment with an immune checkpoint inhibitor; and
  - 1.2 The individual meets initial application criteria for pembrolizumab for stage III or IV resected melanoma adjuvant; or
- 2 Both:
  - 2.1 The individual has received neoadjuvant and adjuvant treatment with an immune checkpoint inhibitor; and

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sı	ubsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 2.2 The individual meets renewal criteria for pembrolizumab for stage III or IV resected melanoma adjuvant; or
- 3 All of the following:
  - 3.1 The individual has received neoadjuvant and adjuvant treatment with an immune checkpoint inhibitor; and
  - 3.2 The individual has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
  - 3.3 The individual meets initial application criteria for pembrolizumab for unresectable or metastatic melanoma; or
- 4 All of the following:
  - 4.1 The individual has received neoadjuvant and adjuvant treatment with an immune checkpoint inhibitor; and
  - 4.2 The individual has received treatment with an immune checkpoint inhibitor for unresectable or metastatic melanoma: and
  - 4.3 The individual meets renewal criteria for pembrolizumab for unresectable or metastatic melanoma.

#### Notes:

- a) Stage IIIB, IIIC, IIID or IV melanoma defined as per American Joint Committee on Cancer (AJCC) 8th Edition
- b) Initiating treatment within 13 weeks of complete surgical resection means either 13 weeks after resection (primary or lymphadenectomy) or 13 weeks prior to the scheduled date of the resection (primary or lymphadenectomy)

Initial application — (stage III or IV resected melanoma - adjuvant) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 The individual has resected stage IIIB, IIIC, IIID or IV melanoma (excluding uveal) (see note a); and
- 2 Adjuvant treatment with pembrolizumab is required; and
- 3 The individual has not received prior funded systemic treatment in the adjuvant setting for stage IIIB, IIIC, IIID or IV melanoma; and
- 4 Treatment must be in addition to complete surgical resection; and
- 5 Treatment must be initiated within 13 weeks of complete surgical resection, unless delay is necessary due to post-surgery recovery (see note b); and
- 6 Pembrolizumab must be administered as monotherapy; and
- 7 The individual has ECOG performance score 0-2; and
- 8 Pembrolizumab to be administered at a fixed dose of 200 mg every 3 weeks (or equivalent).

#### Notes:

- a) Stage IIIB, IIIC, IIID or IV melanoma defined as per American Joint Committee on Cancer (AJCC) 8th Edition
- b) Initiating treatment within 13 weeks of complete surgical resection means 13 weeks after resection (primary or lymphadenectomy)

Renewal — (stage III or IV resected melanoma - adjuvant) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

Any of the following:

- 1 All of the following:
  - 1.1 No evidence of disease recurrence; and
  - 1.2 Pembrolizumab must be administered as monotherapy; and
  - 1.3 Pembrolizumab to be administered at a fixed dose of 200 mg every three weeks (or equivalent) for a maximum of 12 months total treatment course, including any systemic neoadjuvant treatment; and
  - 1.4 Treatment to be discontinued at signs of disease recurrence or at completion of 12 months total treatment course (equivalent to 18 cycles at a dose of 200 mg every 3 weeks), including any systemic neoadjuvant treatment; or
- 2 All of the following:
  - 2.1 The individual has received adjuvant treatment with an immune checkpoint inhibitor; and
  - 2.2 The individual has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
  - 2.3 The individual meets initial application criteria for pembrolizumab for unresectable or metastatic melanoma; or
- 3 All of the following:
  - 3.1 The individual has received adjuvant treatment with an immune checkpoint inhibitor; and

	Subsidy	Fully	Brand or
(Manu	facturer's Price)	Subsidised	Generic
	\$ Pe	er 🗸	Manufacturer

continued...

- 3.2 The individual has received treatment with an immune checkpoint inhibitor for unresectable or metastatic melanoma; and
- 3.3 The individual meets renewal criteria for pembrolizumab for unresectable or metastatic melanoma.

Initial application — (unresectable or metastatic melanoma) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The individual has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Baseline measurement of overall tumour burden is documented clinically and radiologically; and
- 3 The individual has ECOG performance score of 0-2; and
- 4 Fither:
  - 4.1 The individual has not received funded nivolumab; or
  - 4.2 Both:
    - 4.2.1 The individual has received an initial Special Authority approval for nivolumab and has discontinued nivolumab within 12 weeks of starting treatment due to intolerance; and
    - 4.2.2 The cancer did not progress while the individual was on nivolumab; and
- 5 Any of the following:
  - 5.1 The individual has been diagnosed in the metastatic or unresectable stage III or IV setting; or
  - 5.2 The individual did not receive treatment in the perioperative setting with a PD-1/PD-L1 inhibitor; or
  - 5.3 All of the following:
    - 5.3.1 The individual received treatment in the perioperative setting with a PD-1/PD-L1 inhibitor; and
    - 5.3.2 The individual did not experience disease recurrence while on treatment with that PD-1/PD-L1 inhibitor; and
    - 5.3.3 The individual did not experience disease recurrence within six months of completing perioperative treatment with a PD-1/PD-L1 inhibitor.

Renewal — (unresectable or metastatic melanoma, less than 24 months on treatment) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

#### Either:

- 1 Both:
  - 1.1 Any of the following:
    - 1.1.1 The individual's disease has had a complete response to treatment; or
    - 1.1.2 The individual's disease has had a partial response to treatment; or
    - 1.1.3 The individual has stable disease; and
  - 1.2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period; or
- 2 All of the following:
  - 2.1 The individual has previously discontinued treatment with pembrolizumab for reasons other than severe toxicity or disease progression; and
  - 2.2 The individual has signs of disease progression; and
  - 2.3 Disease has not progressed during previous treatment with pembrolizumab.

Renewal — (unresectable or metastatic melanoma, more than 24 months on treatment) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

#### Both:

- 1 The individual has been on treatment for more than 24 months; and
- 2 Either:
  - 2.1 Both:
    - 2.1.1 Any of the following:

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per ✓	Manufacturer	

continued...

- 2.1.1.1 The individual's disease has had a complete response to treatment; or
- 2.1.1.2 The individual's disease has had a partial response to treatment; or
- 2.1.1.3 The individual has stable disease; and
- 2.1.2 Response to treatment in target lesions has been determined by comparable radiologic or clinical assessment following the most recent treatment period; or
- 2.2 All of the following:
  - 2.2.1 The individual has previously discontinued treatment with pembrolizumab for reasons other than severe toxicity or disease progression; and
  - 2.2.2 The individual has signs of disease progression; and
  - 2.2.3 Disease has not progressed during previous treatment with pembrolizumab.

Initial application — (non-small cell lung cancer first-line monotherapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-small cell lung cancer; and
- 2 Patient has not had chemotherapy for their disease in the palliative setting; and
- 3 Patient has not received prior funded treatment with an immune checkpoint inhibitor for NSCLC; and
- 4 For patients with non-squamous histology there is documentation confirming that the disease does not express activating mutations of EGFR or ALK tyrosine kinase unless not possible to ascertain; and
- 5 Pembrolizumab to be used as monotherapy; and
- 6 Either:
  - 6.1 There is documentation confirming the disease expresses PD-L1 at a level greater than or equal to 50% as determined by a validated test unless not possible to ascertain; or
  - 6.2 Both:
    - 6.2.1 There is documentation confirming the disease expresses PD-L1 at a level greater than or equal to 1% as determined by a validated test unless not possible to ascertain; and
    - 6.2.2 Chemotherapy is determined to be not in the best interest of the patient based on clinician assessment; and
- 7 Patient has an ECOG 0-2: and
- 8 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks; and
- 9 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Renewal — (non-small cell lung cancer first line monotherapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 Patient's disease has had a complete response to treatment; or
  - 1.2 Patient's disease has had a partial response to treatment; or
  - 1.3 Patient has stable disease: and
- 2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period; and
- 3 No evidence of disease progression; and
- 4 The treatment remains clinically appropriate and patient is benefitting from treatment; and
- 5 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 6 Treatment with pembrolizumab to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initial application — (non-small cell lung cancer first-line combination therapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the

Subsidy (Manufacturer's Price)	Subsid	ully ised	Brand or Generic
 \$	Per	1	Manufacturer

continued...

following criteria:

All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-small cell lung cancer; and
- 2 The patient has not had chemotherapy for their disease in the palliative setting; and
- 3 Patient has not received prior funded treatment with an immune checkpoint inhibitor for NSCLC; and
- 4 For patients with non-squamous histology there is documentation confirming that the disease does not express activating mutations of EGFR or ALK tyrosine kinase unless not possible to ascertain; and
- 5 Pembrolizumab to be used in combination with platinum-based chemotherapy; and
- 6 Patient has an ECOG 0-2; and
- 7 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks; and
- 8 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Renewal — (non-small cell lung cancer first line combination therapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 Patient's disease has had a complete response to treatment; or
  - 1.2 Patient's disease has had a partial response to treatment; or
  - 1.3 Patient has stable disease: and
- 2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period; and
- 3 No evidence of disease progression; and
- 4 The treatment remains clinically appropriate and patient is benefitting from treatment; and
- 5 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 6 Treatment with pembrolizumab to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

**Initial application** — (breast cancer, advanced) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: Either:

- 1 Patient is currently on treatment with pembrolizumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
  - 2.1 Either:
    - 2.1.1 Patient has recurrent or de novo unresectable, inoperable locally advanced triple-negative breast cancer (that does not express ER, PR or HER2 IHC3+ or ISH+ [including FISH or other technology]); or
    - 2.1.2 Patient has recurrent or de novo metastatic triple-negative breast cancer (that does not express ER, PR or HER2 IHC3+ or ISH+ [including FISH or other technology]); and
  - 2.2 Patient is treated with palliative intent; and
  - 2.3 Patient's cancer has confirmed PD-L1 Combined Positive Score (CPS) is greater than or equal to 10; and
  - 2.4 Patient has received no prior systemic therapy in the palliative setting; and
  - 2.5 Patient has an ECOG score of 0-2; and
  - 2.6 Pembrolizumab is to be used in combination with chemotherapy; and
  - 2.7 Baseline measurement of overall tumour burden is documented clinically and radiologically; and
  - 2.8 Pembrolizumab is to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks.

**Renewal — (breast cancer, advanced)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

All of the following:

- 1 Any of the following:
  - 1.1 Patient's disease has had a complete response to treatment; or
  - 1.2 Patient's disease has had a partial response to treatment; or
  - 1.3 Patient has stable disease; and
- 2 No evidence of disease progression; and
- 3 Response to treatment in target lesions has been determined by a comparable radiologic assessment following the most recent treatment period; and
- 4 Pembrolizumab is to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 5 Treatment with pembrolizumab is to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initial application — (head and neck squamous cell carcinoma) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria: Fither:

- 1 Patient is currently on treatment with pembrolizumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
  - 2.1 Patient has recurrent or metastatic head and neck squamous cell carcinoma of mucosal origin (excluding nasopharyngeal carcinoma) that is incurable by local therapies: and
  - 2.2 Patient has not received prior systemic therapy in the recurrent or metastatic setting; and
  - 2.3 Patient has a positive PD-L1 combined positive score (CPS) of greater than or equal to 1; and
  - 2.4 Patient has an ECOG performance score of 0-2; and
  - 2.5 Either:
    - 2.5.1 Pembrolizumab to be used in combination with platinum-based chemotherapy; or
    - 2.5.2 Pembrolizumab to be used as monotherapy; and
  - 2.6 Pembrolizumab is to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks.

Renewal — (head and neck squamous cell carcinoma) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 Patient's disease has had a complete response to treatment; or
  - 1.2 Patient's disease has had a partial response to treatment; or
  - 1.3 Patient has stable disease; and
- 2 No evidence of disease progression; and
- 3 Pembrolizumab is to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 4 Treatment with pembrolizumab is to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initial application — (MSI-H/dMMR advanced colorectal cancer) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria: Either:

- 1 Individual is currently on treatment with pembrolizumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
  - 2.1 Either:
    - 2.1.1 Individual has deficient mismatch repair (dMMR) or microsatellite instability-high (MSI-H) metastatic colorectal cancer; or
    - 2.1.2 Individual has deficient mismatch repair (dMMR) or microsatellite instability-high (MSI-H) unresectable colorectal cancer; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	;	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 2.2 Individual is treated with palliative intent; and
- 2.3 Individual has not previously received funded treatment with pembrolizumab for MSI-H/dMMR advanced colorectal cancer; and
- 2.4 Individual has an ECOG performance score of 0-2; and
- 2.5 Baseline measurement of overall tumour burden is documented clinically and radiologically; and
- 2.6 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of

**Renewal — (MSI-H/dMMR advanced colorectal cancer)** from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 3 Treatment with pembrolizumab is to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initial application — (Urothelial carcinoma) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria: Either:

- 1 Patient is currently on treatment with pembrolizumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
  - 2.1 Patient has inoperable locally advanced (T4) or metastatic urothelial carcinoma; and
  - 2.2 Patient has an ECOG performance score of 0-2; and
  - 2.3 Patient has documented disease progression following treatment with chemotherapy; and
  - 2.4 Pembrolizumab to be used as monotherapy at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks.

Renewal — (Urothelial carcinoma) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 Patient's disease has had a complete response to treatment; or
  - 1.2 Patient's disease has had a partial response to treatment; or
  - 1.3 Patient has stable disease: and
- 2 No evidence of disease progression; and
- 3 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 4 Treatment with pembrolizumab is to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

**Initial application — (relapsed/refractory Hodgkin lymphoma)** only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria: Either:

- 1 Individual is currently on treatment with pembrolizumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
  - 2.1 Either:
    - 2.1.1 Both:
      - 2.1.1.1 Individual has relapsed/refractory Hodgkin lymphoma after two or more lines of chemotherapy; and
      - 2.1.1.2 Individual is ineligible for autologous stem cell transplant; or
    - 2.1.2 Individual has relapsed/refractory Hodgkin lymphoma and has previously undergone an autologous stem cell transplant; and
  - 2.2 Individual has not previously received funded pembrolizumab for relapsed/refractory Hodgkin lymphoma; and

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

2.3 Pembrolizumab to be administered at doses no greater than 200 mg once every 3 weeks.

Renewal — (relapsed/refractory Hodgkin lymphoma) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient has received a partial or complete response to pembrolizumab; and
- 2 Treatment with pembrolizumab is to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

## Other Immunosuppressants

CICLOSPORIN			
Cap 25 mg	44.63	50	Neoral
Cap 50 mg	88.91	50	✓ Neoral
Cap 100 mg	177.81	50	✓ Neoral
Oral liq 100 mg per ml		50 ml OP	✓ Neoral
EVEROLIMUS - Special Authority see SA2414 below - Re	tail pharmacy		
Wastage claimable			
Tab 10 mg	6,512.29	30	✓ Afinitor
Tab 5 mg	4,555.76	30	✓ Afinitor

#### ⇒SA2414 Special Authority for Subsidy

Initial application only from a neurologist or oncologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has tuberous sclerosis: and
- 2 Patient has progressively enlarging sub-ependymal giant cell astrocytomas (SEGAs) that require treatment.

**Renewal** only from a neurologist or oncologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Documented evidence of SEGA reduction or stabilisation by MRI within the last 3 months; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Everolimus to be discontinued at progression of SEGAs.

**Initial application — (renal cell carcinoma)** from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 All of the following:
  - 1.1 The patient has metastatic renal cell carcinoma; and
  - 1.2 The disease is of predominant clear-cell histology; and
  - 1.3 The patient has documented disease progression following one previous line of treatment; and
  - 1.4 The patient has an ECOG performance status of 0-2; and
  - 1.5 Everolimus is to be used in combination with lenvatinib; or
- 2 All of the following:
  - 2.1 Patient has received funded treatment with nivolumab for the second line treatment of metastatic renal cell carcinoma; and
  - 2.2 Patient has experienced treatment limiting toxicity from treatment with nivolumab; and
  - 2.3 Everolimus is to be used in combination with lenvatinib; and
  - 2.4 There is no evidence of disease progression.

Renewal — (renal cell carcinoma) from any relevant practitioner. Approvals valid for 4 months where there is no evidence of disease progression.

	Subsidy (Manufacturer's Prio \$	ce) Sub Per	Fully sidised	Brand or Generic Manufacturer
SIROLIMUS - Special Authority see SA2270 below - Retail pha	rmacy			
Tab 1 mg	749.99	100		Rapamune
Tab 2 mg	1,499.99	100	✓	Rapamune
Oral liq 1 mg per ml	449.99	60 ml OP	✓	Rapamune

#### **⇒SA2270** Special Authority for Subsidy

**Initial application** from any medical practitioner. Approvals valid without further renewal unless notified where the drug is to be used for rescue therapy for an organ transplant recipient.

Notes: Rescue therapy defined as unresponsive to calcineurin inhibitor treatment as defined by refractory rejection; or intolerant to calcineurin inhibitor treatment due to any of the following:

- GFR< 30 ml/min: or
- · Rapidly progressive transplant vasculopathy; or
- · Rapidly progressive obstructive bronchiolitis; or
- . HUS or TTP; or
- · Leukoencepthalopathy; or
- · Significant malignant disease

Initial application — (severe non-malignant lymphovascular malformations\*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe non-malignant lymphovascular malformation\*; and
- 2 Any of the following:
  - 2.1 Malformations are not adequately controlled by sclerotherapy and surgery; or
  - 2.2 Malformations are widespread/extensive and sclerotherapy and surgery are not considered clinically appropriate; or
  - 2.3 Sirolimus is to be used to reduce malformation prior to consideration of surgery; and
- 3 Patient is being treated by a specialist lymphovascular malformation multi-disciplinary team; and
- 4 Patient has measurable disease as defined by RECIST version 1.1 (see Note).

Renewal — (severe non-malignant lymphovascular malformations\*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 Patient's disease has had either a complete response or a partial response to treatment, or patient has stable disease according to RECIST version 1.1 (see Note); or
  - 1.2 Patient's disease has stabilised or responded clinically and disease response to treatment has been clearly documents in patient notes; and
- 2 No evidence of progressive disease; and
- 3 The treatment remains clinically appropriate and the patient is benefitting from the treatment.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer et al. Eur J Cancer 2009;45:228-47)

Indications marked with \* are unapproved indications

Initial application — (renal angiomyolipoma(s) associated with tuberous sclerosis complex\*) only from a nephrologist or urologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has tuberous sclerosis complex\*; and
- 2 Evidence of renal angiomyolipoma(s) measuring 3 cm or greater and that have shown interval growth.

 $\textbf{Renewal} - \textbf{(renal angiomyolipoma(s) associated with tuberous sclerosis complex^*)} \ \ \text{from any relevant practitioner}.$ 

Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 Documented evidence of renal angiomyolipoma reduction or stability by magnetic resonance imaging (MRI) or ultrasound; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	•	Manufacturer	

continued...

- 2 Demonstrated stabilisation or improvement in renal function; and
- 3 The patient has not experienced angiomyolipoma haemorrhage or significant adverse effects to sirolimus treatment; and
- 4 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indications marked with \* are unapproved indications

Initial application — (refractory seizures associated with tuberous sclerosis complex\*) only from a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has epilepsy with a background of documented tuberous sclerosis complex; and
- 2 Either:
  - 2.1 Both:
    - 2.1.1 Vigabatrin has been trialled and has not adequately controlled seizures; and
    - 2.1.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least two of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); or
  - 2.2 Both:
    - 2.2.1 Vigabatrin is contraindicated; and
    - 2.2.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least three of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); and
- 3 Seizures have a significant impact on quality of life; and
- 4 Patient has been assessed and surgery is considered inappropriate for this patient, or the patient has been assessed and would benefit from mTOR inhibitor treatment prior to surgery.

Note: Those of childbearing age potential are not required to trial phenytoin sodium, sodium valproate, or topiramate. Those who can father children are not required to trial sodium valproate.

Renewal — (refractory seizures associated with tuberous sclerosis complex\*) only from a neurologist. Approvals valid for 12 months where demonstrated significant and sustained improvement in seizure rate (e.g. 50% reduction in seizure frequency) or severity and/or patient quality of life compared with baseline prior to starting sirolimus treatment.

Note: Indications marked with \* are unapproved indications

TACROLIMUS - Special Authority see SA2455 below - Retail pharmacy

Cap 0.5 mg	49.60	100	✓ Tacrolimus Sandoz
Cap 0.75 mg	99.30	100	✓ Tacrolimus Sandoz
Cap 1 mg	84.30	100	✓ Tacrolimus Sandoz
Cap 5 mg		50	✓ Tacrolimus Sandoz

## ⇒SA2455 Special Authority for Subsidy

**Initial application — (organ transplant)** only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 The individual is an organ transplant recipient; or
- 2 The individual is receiving induction therapy for an organ transplant.

Note: Subsidy applies for either primary or rescue therapy.

Initial application — (non-transplant indications\*) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient requires long-term systemic immunosuppression; and
- 2 Either:
  - 2.1 Ciclosporin has been trialled and discontinued treatment because of unacceptable side effects or inadequate clinical response; or
  - 2.2 Patient is a child with nephrotic syndrome\*.

Note: Indications marked with \* are unapproved indications

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

## **JAK** inhibitors

U	IPADACITI	NIB	- Sp	ecial Author	ity see SA2483	below – Retail pharm	асу	

Tab modified-release 15 mg	1,271.00	28	✓ Rinvoq
Tab modified-release 30 mg	2,033.00	28	✓ Rinvoq
Tab modified-release 45 mg	3,049.00	28	✓ Rinvoq

#### ⇒SA2483 Special Authority for Subsidy

Initial application — (Rheumatoid Arthritis (previously treated with adalimumab or etanercept)) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The individual has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
  - 2.1 The individual has experienced intolerable side effects with adalimumab and/or etanercept; or
  - 2.2 The individual has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
- 3 Any of the following:
  - 3.1 Rituximab is not clinically appropriate; or
  - 3.2 The individual is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or
  - 3.3 Both:
    - 3.3.1 The individual has been started on rituximab for rheumatoid arthritis in a Health NZ Hospital; and
    - 3.3.2 Either:
      - 3.3.2.1 The individual has experienced intolerable side effects with rituximab; or
      - 3.3.2.2 At four months following the initial course of rituximab the individual has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Renewal — (Rheumatoid Arthritis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

#### Either:

- 1 Following 6 months' initial treatment, the individual has experienced at least a 50% decrease in active joint count from baseline: or
- 2 On subsequent reapplications, the individual has experienced at least a continuing 30% improvement in active joint count from baseline.

**Initial application — (atopic dermatitis)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

#### Either:

- 1 Individual is currently on treatment with upadacitinib for atopic dermatitis and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
  - 2.1 Individual has moderate to severe atopic dermatitis, severity as defined by an Eczema Area and Severity Index (EASI) score of greater than or equal to 16 or a Dermatology Life Quality Index (DLQI) score of greater than or equal to 10; and
  - 2.2 Individual has received insufficient benefit from topical therapy (including topical corticosteroids or topical calcineurin inhibitors) for a 28-day trial within the last 6 months, unless contraindicated to all; and
  - 2.3 Individual has trialled and received insufficient benefit from at least one systemic therapy for a minimum of three months (eq ciclosporin, azathioprine, methotrexate or mycophenolate mofetil), unless contraindicated to all; and
  - 2.4 An EASI assessment or DLQI assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
  - 2.5 The most recent EASI or DQLI assessment is no more than 1 month old at the time of application.

Subsidy (Manufacturer's Price)	5	Fully Subsidised	Brand or Generic	
\$	Per	✓	Manufacturer	

continued...

**Renewal** — (atopic dermatitis) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 Individual has received a 75% or greater reduction in EASI score (EASI 75) as compared to baseline EASI prior to commencing upadacitinib; or
- 2 Individual has received a DLQI improvement of 4 or more as compared to baseline DLQI prior to commencing upadacitinib.

Initial application — (Crohn's disease - adult) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Individual is currently on treatment with upadacitinib for Crohn's disease and met all remaining criteria prior to commencing treatment; or
- 2 Both:
  - 2.1 Individual has active Crohn's disease: and
  - 2.2 Either:
    - 2.2.1 Individual has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria; or
    - 2.2.2 Both:
      - 2.2.2.1 Individual meets the initiation criteria for prior biologic therapies for Crohn's disease; and
      - 2.2.2.2 Other biologic therapies for Crohn's disease are contraindicated.

Renewal — (Crohn's disease - adult) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 CDAI score has reduced by 100 points from the CDAI score when the individual was initiated on biologic therapy; or
- 2 HBI score has reduced by 3 points from when individual was initiated on biologic therapy; or
- 3 CDAI score is 150 or less: or
- 4 HBI score is 4 or less: or
- 5 The individual has experienced an adequate response to treatment, but CDAI score cannot be assessed.

Initial application — (Crohn's disease - children\*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Individual is currently on treatment with upadacitinib for Crohn's disease and met all remaining criteria prior to commencing treatment: or
- 2 Both:
  - 2.1 Child has active Crohn's disease; and
  - 2.2 Either:
    - 2.2.1 Child has had an initial approval for prior biologic therapy for Crohn's disease and has experienced intolerable side effects or insufficient benefit to meet renewal criteria: or
    - 2.2.2 Both:
      - 2.2.2.1 Child meets the initiation criteria for prior biologic therapies for Crohn's disease; and
      - 2.2.2.2 Other biologic therapies for Crohn's disease are contraindicated.

Renewal — (Crohn's disease - children\*) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 PCDAI score has reduced by 10 points from the child was initiated on treatment; or
- 2 PCDAI score is 15 or less: or
- 3 The child has experienced an adequate response to treatment, but PCDAI score cannot be assessed.

Subsidy		Fully	Brand or
(Manufacturer's Price)	S	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

Note: Indications marked with \* are unapproved indications.

Initial application — (ulcerative colitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Individual is currently on treatment with upadacitinib for ulcerative colitis and met all remaining criteria prior to commencing treatment; or
- 2 Both:
  - 2.1 Individual has active ulcerative colitis; and
  - 2.2 Either
    - 2.2.1 Individual has had an initial approval for prior biologic therapy for ulcerative colitis and has experienced intolerable side effects or insufficient benefit to meet renewal criteria; or
    - 2.2.2 Both:
      - 2.2.2.1 Individual meets the initiation criteria for prior biologic therapies for ulcerative colitis; and
      - 2.2.2.2 Other biologic therapies for ulcerative colitis are contraindicated.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

## Either:

- 1 The SCCAI score has reduced by 2 points or more from the SCCAI score when the individual was initiated on treatment; or
- 2 PUCAI score has reduced by 10 points or more from the PUCAI score when the individual was initiated on treatment.

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	1	Manufacturer

# **Antiallergy Preparations**

## Allergic Emergencies

ADRENALINE - Special Authority see SA2185 below - Retail pharmacy

- a) Maximum of 2 ini per prescription
- Additional prescriptions limited to replacement of up to two devices prior to expiry, or replacement of used device for treatment of anaphylaxis.

Inj 0.15 mg per 0.3 ml auto-injector	85.50	1 OP	<ul><li>Epipen Jr</li></ul>
Inj 0.3 mg per 0.3 ml auto-injector	85.50	1 OP	Epipen

## ⇒SA2185 Special Authority for Subsidy

Initial application — (anaphylaxis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 Patient has experienced an anaphylactic reaction which has resulted in presentation to a hospital or emergency department; or
  - 1.2 Patient has been assessed to be at significant risk of anaphylaxis by a relevant practitioner; and
- 2 Patient is not to be prescribed more than two devices in initial prescription.

## ⇒SA1558 Special Authority for Subsidy

Initial application only from a clinical immunologist or relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Supply for anticipated emergency treatment of laryngeal/oro-pharyngeal or severe abdominal attacks of acute hereditary angioedema (HAE) for patients with confirmed diagnosis of C1-esterase inhibitor deficiency; and
- 2 The patient has undergone product training and has agreed upon an action plan for self-administration.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

# **Allergy Desensitisation**

## **⇒SA1367** Special Authority for Subsidy

Initial application only from a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 RAST or skin test positive; and
- 2 Patient has had severe generalised reaction to the sensitising agent.

**Renewal** only from a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

BEE VENOM ALLERGY TREATMENT - Special Authority see SA1367 above - Retail pharmacy Initiation kit - 1 vial freeze dried venom with diluent 305.00 1 OP ✓ VENOX S29 Maintenance kit - 1 vial freeze dried venom with diluent................305.00 1 OP ✓ VENOX S29 Maintenance kit - 6 vials 120 mcg freeze dried venom, with 1 OP ✓ Venomil S29 Treatment kit - 1 vial 550 mcg freeze dried venom, 1 diluent 1 OP ✓ Albev Treatment kit - 1 vial 550 mcg freeze dried venom, with diluent ..... 305.00 1 OP ✓ Hymenoptera S29

	Subsidy		Fully	Brand or
	(Manufacturer's P	rice) Subs Per	idised	Generic Manufacturer
MAOD VENOMAN EDOV TREATMENT.	. OA4007 II		D.1.	
WASP VENOM ALLERGY TREATMENT – Special Authority see	SA1367 on the	previous page	– Heta	all pnarmacy
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze dried polistes venom, 1 diluent 9 ml, 3 diluent 1.8 ml	202.22	1 OP	./ /	Albey
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze		TOF	• ,	Albey
dried venom, with diluent	305.00	1 OP	<b>✓</b> ⊦	lymenoptera S29
Treatment kit (Paper wasp venom) - 6 vials 120 mcg freeze		. 0.		rymonoptoru —
dried venom, with diluent	305.00	1 OP	<b>✓</b> \	/enomil \$29
Treatment kit (Yellow Jacket venom) - 1 vial 550 mcg freeze				
dried venom, with diluent	305.00	1 OP	<b>✓</b> F	lymenoptera S29
Treatment kit (Yellow jacket venom) - 1 vial 550 mcg freeze				
dried vespula venom, 1 diluent 9 ml, 3 diluent 1.8 ml	431.24	1 OP	<b>✓</b>	Albey
Treatment kit (Yellow jacket venom) - 6 vials 120 mcg freeze				
dried venom, with diluent	305.00	1 OP	<b>✓</b> \	/enomil \$29
Antihistamines				
CETIRIZINE HYDROCHLORIDE  * Tab 10 mg	1 71	100	./ -	Zista .
* Oral liq 1 mg per ml		200 ml	_	lista Histaclear
DEXTROCHLORPHENIRAMINE MALEATE		200 1111	٠,	iistacicai
	2.02	40		
* Tab 2 mg	(8.40)	40		Polaramine
	1.01	20	'	olaramine
	(5.99)	_0	F	Polaramine
* Oral liq 2 mg per 5 ml		100 ml		
	(10.29)		F	Polaramine
FEXOFENADINE HYDROCHLORIDE				
* Tab 60 mg	4.34	20		
	(8.23)			Telfast
* Tab 120 mg		30		<u>exaclear</u>
* Tab 180 mg	4.10	30	<b>✓</b> <u>F</u>	<u>exaclear</u>
LORATADINE				
* Tab 10 mg		100		oratadine Noumed
W. Orallia dana ana an	6.02	100		orafix
* Oral liq 1 mg per ml(Lorafix Tab 10 mg to be delisted 1 June 2026)	1.43	100 ml	•	laylor syrup
PROMETHAZINE HYDROCHLORIDE	0.10	100		Moreootho
* Tab 10 mg * Tab 25 mg		100 100	_	Allersoothe Allersoothe
* Oral lig 1 mg per 1 ml		100 ml	_	Allersoothe
* Inj 25 mg per ml, 2 ml ampoule – Up to 5 inj available on a F		5		lospira
Inhaled Corticosteroids				
BECLOMETHASONE DIPROPIONATE			_	_
Aerosol inhaler, 50 mcg per dose		200 dose OP		Qvar
Aerosol inhaler, 50 mcg per dose CFC-free		200 dose OP	_	Beclazone 50
Aerosol inhaler, 100 mcg per dose		200 dose OP 200 dose OP	-	Qvar Beclazone 100
Aerosol inhaler, 100 mcg per dose CFC-free		200 dose OP	_	Beclazone 100 Beclazone 250
Acrosof illiaici, 200 filog por acose or o-filee		LUU UUSE OF	, [	JOUIGEONIC ESU

			IN AND ALLENGIES
	Subsidy (Manufacturer's \$		Fully Brand or dised Generic  ✓ Manufacturer
BUDESONIDE			
Powder for inhalation, 100 mcg per dose	17.00	200 dose OP	<ul><li>Pulmicort Turbuhaler</li></ul>
Powder for inhalation, 200 mcg per dose	19.00	200 dose OP	✓ Pulmicort Turbuhaler
Powder for inhalation, 400 mcg per dose	32.00	200 dose OP	✓ Pulmicort Turbuhaler
FLUTICASONE			
Aerosol inhaler, 50 mcg per dose	7.19	120 dose OP	✓ Flixotide
Powder for inhalation, 50 mcg per dose	8.61	60 dose OP	✓ Flixotide Accuhaler
Powder for inhalation, 100 mcg per dose	7.81	60 dose OP	✓ Flixotide Accuhaler
Aerosol inhaler, 125 mcg per dose	13.60	120 dose OP	✓ Flixotide
Aerosol inhaler, 250 mcg per dose		120 dose OP	✓ Flixotide
Powder for inhalation, 250 mcg per dose	11.93	60 dose OP	✓ Flixotide Accuhaler
Inhaled Long-acting Beta-adrenoceptor Agonis	sts		
EFORMOTEROL FUMARATE DIHYDRATE			
Powder for inhalation 4.5 mcg per dose, breath activated			
(equivalent to eformoterol fumarate 6 mcg metered dos	e) 10.32	60 dose OP	
	(16.90)		Oxis Turbuhaler
INDACATEROL	` ,		
Powder for inhalation 150 mcg	61.00	30 dose OP	✓ Onbrez Breezhaler
Powder for inhalation 300 mcg		30 dose OP	✓ Onbrez Breezhaler
ű		00 0000 01	onbicz biccznaici
SALMETEROL	22.25	100   00	
Aerosol inhaler CFC-free, 25 mcg per dose		120 dose OP	✓ Serevent
Powder for inhalation, 50 mcg per dose, breath activated	26.25	60 dose OP	✓ Serevent Accuhaler
Inhaled Corticosteroids with Long-Acting Beta	-Adrenocep	tor Agonists	
BUDESONIDE WITH EFORMOTEROL			
* Powder for inhalation 160 mcg with 4.5 mcg eformoterol			
fumarate per dose (equivalent to 200 mcg budesonide	with		
6 mcg eformoterol fumarate metered dose) - Up to 12			
dose available on a PSO	41.50	120 dose OP	✓ DuoResp Spiromax
Powder for inhalation 320 mcg with 9 mcg eformoterol fuma	rate		• •
per dose (equivalent to 400 mcg budesonide with 12 m			
eformoterol fumarate metered dose) – No more than 2			
dose per day		120 dose OP	✓ DuoResp Spiromax
* Aerosol inhaler 100 mcg with eformoterol fumarate 6 mcg -			- attitude opinomian
to 120 dose available on a PSO		120 dose OP	✓ Vannair
		120 0036 OF	- Valillali
* Powder for inhalation 100 mcg with eformoterol fumarate 6	•	100 dess 00	./ Cumhinaut
- Up to 120 dose available on a PSO	33./4	120 dose OP	✓ Symbicort
ale A 1111 000 111 1 1 1 1 1 1			Turbuhaler 100/6
* Aerosol inhaler 200 mcg with eformoterol fumarate 6 mcg - to 120 dose available on a PSO		120 dose OP	✓ Vannair
* Powder for inhalation 200 mcg with eformoterol fumarate 6			
– Up to 120 dose available on a PSO		120 dose OP	✓ Symbicort Turbuhaler 200/6
Powder for inhalation 400 mcg with eformoterol fumarate			
12 mcg – No more than 2 dose per day	33.74	60 dose OP	✓ Symbicort Turbuhaler 400/12

 $<sup>\</sup>blacktriangle \textit{Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. }$ 

	Subsidy		Fully Brand or
	(Manufacturer's I	Price) Subsi Per	idised Generic  Manufacturer
LUTICASONE FUROATE WITH VILANTEROL Powder for inhalation 100 mcg with vilanterol 25 mcg	44.08	30 dose OP	✓ Breo Ellipta
Aerosol inhaler 50 mcg with salmeterol 25 mcg Aerosol inhaler 125 mcg with salmeterol 25 mcg		120 dose OP 120 dose OP	<ul><li>✓ Seretide</li><li>✓ Seretide</li></ul>
Powder for inhalation 100 mcg with salmeterol 50 mcg — No more than 2 dose per day Powder for inhalation 250 mcg with salmeterol 50 mcg — No	33.74	60 dose OP	✓ Seretide Accuhaler
more than 2 dose per day	44.08	60 dose OP	✓ Seretide Accuhaler
Beta-Adrenoceptor Agonists			
SALBUTAMOL Oral liq 400 mcg per ml Infusion 1 mg per ml, 5 ml Inj 500 mcg per ml, 1 ml – Up to 5 inj available on a PSO	130.00	150 ml 10 5	✓ <u>Ventolin</u> ✓ Ventolin ✓ Ventolin
Inhaled Beta-Adrenoceptor Agonists			
SALBUTAMOL  Aerosol inhaler, 100 mcg per dose CFC free – Up to 1000 dose available on a PSO	4.18 (7.45)	200 dose OP	✓ SalAir Ventolin
Nebuliser soln, 1 mg per ml, 2.5 ml ampoule – Up to 30 neb available on a PSO	8.96	20	✓ Asthalin ✓ UK Cipla S29
Nebuliser soln, 2 mg per ml, 2.5 ml ampoule – Up to 30 neb available on a PSO	9.43	20	✓ Asthalin ✓ UK Cipla S29
ERBUTALINE SULPHATE Powder for inhalation, 200 mcg per dose (equivalent to 250 mcg metered dose), breath activated	22.20	120 dose OP	✓ Bricanyl Turbuhaler
Anticholinergic Agents			
PRATROPIUM BROMIDE  Aerosol inhaler, 20 mcg per dose CFC-free – Up to 400 dose available on a PSO	16.20	200 dose OP	✓ Atrovent
Nebuliser soln, 250 mcg per ml, 2 ml ampoule – Up to 40 ne available on a PSO		20	✓ Accord S29 ✓ Univent
Inhaled Beta-Adrenoceptor Agonists with Antic	holinergic A	gents	
SALBUTAMOL WITH IPRATROPIUM BROMIDE  Aerosol inhaler, 100 mcg with ipratropium bromide, 20 mcg p  dose CFC-free		200 dose OP	✓ Duolin HFA
Nebuliser soln, 2.5 mg with ipratropium bromide 0.5 mg per vial, 2.5 ml ampoule – Up to 20 neb available on a PSO	11.04	20	✓ Duolin

Subsidy Fully Brand or
(Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

## **Long-Acting Muscarinic Antagonists**

GLYCOPYRRONIUM - Subsidy by endorsement

- a) Inhaled glycopyrronium treatment will not be subsidised if patient is also receiving treatment with subsidised tiotropium or umeclidinium
- b) Glycopyrronium powder for inhalation 50 mcg per dose is subsidised only for patients who have been diagnosed as having COPD using spirometry if spirometry is possible, and the prescription is endorsed accordingly.

#### TIOTROPIUM BROMIDE - Subsidy by endorsement

- a) Tiotropium treatment will not be subsidised if patient is also receiving treatment with subsidised inhaled glycopyrronium or umeclidinium.
- b) Tiotropium bromide is subsidised only for patients who have been diagnosed as having COPD using spirometry if spirometry is possible, and the prescription is endorsed accordingly. Patients who had tiotropium dispensed before 1 October 2018 with a valid Special Authority are deemed endorsed.

#### UMECLIDINIUM - Subsidy by endorsement

- a) Umeclidinium will not be subsidised if patient is also receiving treatment with subsidised inhaled glycopyrronium or tiotropium bromide.
- b) Umeclidinium powder for inhalation 62.5 mcg per dose is subsidised only for patients who have been diagnosed as having COPD using spirometry if spirometry is possible, and the prescription is endorsed accordingly.

# Long-Acting Muscarinic Antagonists with Long-Acting Beta-Adrenoceptor Agonists

Combination long acting muscarinic antagonist and long acting beta-2 agonist will not be subsidised if patient is also receiving treatment with a combination inhaled corticosteroid and long acting beta-2 agonist.

#### ⇒SA2554 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Both:

- 1 Patient has been stabilised on a long acting muscarinic antagonist; and
- 2 The prescriber considers that the patient would receive additional benefit from switching to a combination product.

GLYCOPYRRONIUM WITH INDACATEROL - Special Authority see SA2554 above - Retail pharmacy

TIOTROPIUM BROMIDE WITH OLODATEROL - Special Authority see SA2554 above - Retail pharmacy

Soln for inhalation 2.5 mcg with olodaterol 2.5 mcg......81.00 60 dose OP Spiolto Respimat

UMECLIDINIUM WITH VILANTEROL – Special Authority see SA2554 above – Retail pharmacy

## Inhaled Corticosteroid with Long-Acting Muscarinic Antagonist and Beta Agonist

BUDESONIDE WITH GLYCOPYRRONIUM AND EFORMOTEROL – Special Authority see SA2421 on the next page – Retail pharmacy

Aerosol inhaler budesonide 160 mcg with glycopyrronium

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

## **⇒SA2421** Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Both:

- 1 Patient has a diagnosis of COPD confirmed by spirometry or spirometry has been attempted and technically acceptable results are not possible; and
- 2 Either:
  - 2.1 Both:
    - 2.1.1 Patient is currently receiving an inhaled corticosteroid with long acting beta-2 agonist (ICS/LABA) or a long acting muscarinic antagonist with long acting beta-2 agonist (LAMA/LABA); and
    - 2.1.2 Any of the following:

Clinical criteria:

- 2.1.2.1 Patient has a COPD Assessment Test (CAT) score greater than 10; or
- 2.1.2.2 Patient has had 2 or more exacerbations in the previous 12 months; or
- 2.1.2.3 Patient has had one exacerbation requiring hospitalisation in the previous 12 months; or
- 2.1.2.4 Patient has had an eosinophil count greater than or equal to  $0.3 \times 10^9$  cells/L in the previous 12 months: or
- 2.2 Patient is currently receiving multiple inhaler triple therapy (inhaled corticosteroid with long-acting muscarinic antagonist and long-acting beta-2 agonist ICS/LAMA/LABA) and met at least one of the clinical criteria above prior to commencing multiple inhaler therapy.

FLUTICASONE FUROATE WITH UMECLIDINIUM AND VILANTEROL – Special Authority see SA2326 below – Retail pharmacy Powder for inhalation fluticasone furoate 100 mcg with

umeclidinium 62.5 mcg and vilanterol 25 mcg.......104.24 30 dose OP ✓ Trelegy Ellipta

#### ⇒SA2326 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has a diagnosis of COPD confirmed by spirometry or spirometry has been attempted and technically acceptable results are not possible; and
- 2 Fither:
  - 2.1 Both:
    - 2.1.1 Patient is currently receiving an inhaled corticosteroid with long acting beta-2 agonist (ICS/LABA) or a long acting muscarinic antagonist with long acting beta-2 agonist (LAMA/LABA); and
    - 2.1.2 Any of the following:

Clinical criteria:

- 2.1.2.1 Patient has a COPD Assessment Test (CAT) score greater than 10; or
- 2.1.2.2 Patient has had 2 or more exacerbations in the previous 12 months; or
- 2.1.2.3 Patient has had one exacerbation requiring hospitalisation in the previous 12 months; or
- 2.1.2.4 Patient has had an eosinophil count greater than or equal to  $0.3 \times 10^{\circ}9$  cells/L in the previous 12 months; or
- 2.2 Patient is currently receiving multiple inhaler triple therapy (inhaled corticosteroid with long acting muscarinic antagonist and long acting beta-2 agonist ICS/LAMA/LABA) and met at least one of the clinical criteria above prior to commencing multiple inhaler triple therapy.

## **Antifibrotics**

NINTEDANIB - Special Authority see SA2012 on the next page - Retail pharmacy

Note: Nintedanib not subsidised in combination with subsidised pirfenidone.

 Cap 100 mg
 2,554.00
 60 OP
 ✓ Ofev

 Cap 150 mg
 3,870.00
 60 OP
 ✓ Ofev

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic	
\$	Per	1	Manufacturer	

## ⇒SA2012 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Nintedanib is to be discontinued at disease progression (See Note); and
- 4 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 5 Any of the following:
  - 5.1 The patient has not previously received treatment with pirfenidone; or
  - 5.2 Patient has previously received pirfenidone, but discontinued pirfenidone within 12 weeks due to intolerance; or
  - 5.3 Patient has previously received pirfenidone, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with pirfenidone).

**Renewal — (idiopathic pulmonary fibrosis)** only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 3 Nintedanib is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

PIRFENIDONE – Retail pharmacy-Specialist – Special Authority see SA2013 below Note: Pirfenidone is not subsidised in combination with subsidised nintedanib.

Tab 801 mg	3,645.00	90 OP	✓ Esbriet
Tab 267 mg	1,215.00	90	<ul><li>Esbriet</li></ul>

## ⇒SA2013 Special Authority for Subsidy

**Initial application — (idiopathic pulmonary fibrosis)** only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note); and
- 4 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 5 Any of the following:
  - 5.1 The patient has not previously received treatment with nintedanib; or
  - 5.2 Patient has previously received nintedanib, but discontinued nintedanib within 12 weeks due to intolerance; or
  - 5.3 Patient has previously received nintedanib, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with nintedanib).

**Renewal — (idiopathic pulmonary fibrosis)** only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
Leukotriene Receptor Antagonists				
MONTELUKAST  * Tab 4 mg  * Tab 5 mg  * Tab 10 mg	3.10	28 28 28	_	Montelukast Viatris Montelukast Viatris Montelukast Viatris
Methylxanthines  AMINOPHYLLINE  * Inj 25 mg per ml, 10 ml ampoule – Up to 5 inj available on a PSO	180 00	5	/	DBL Aminophylline
THEOPHYLLINE  * Tab long-acting 250 mg  * Oral liq 80 mg per 15 ml	25.65	100 00 n	<b>✓</b>	Nuelin-SR Nuelin
Mucolytics				
DORNASE ALFA – Special Authority see SA1978 below – Retail Nebuliser soln, 2.5 mg per 2.5 ml ampoule		6	•	Pulmozyme

#### ⇒SA1978 Special Authority for Subsidy

Initial application — (cystic fibrosis) only from a respiratory physician or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of cystic fibrosis: and
- 2 Patient has previously undergone a trial with, or is currently being treated with, hypertonic saline; and
- 3 Any of the following:
  - 3.1 Patient has required one or more hospital inpatient respiratory admissions in the previous 12 month period; or
  - 3.2 Patient has had 3 exacerbations due to CF, requiring oral or intravenous (IV) antibiotics in the previous 12 month period: or
  - 3.3 Patient has had 1 exacerbation due to CF, requiring oral or IV antibiotics in the previous 12 month period and a Brasfield score of < 22/25: or
  - 3.4 Patient has a diagnosis of allergic bronchopulmonary aspergillosis (ABPA).

Renewal — (cystic fibrosis) only from a respiratory physician or paediatrician. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient continues to benefit from treatment.

ELEXACAFTOR WITH TEZACAFTOR, IVACAFTOR AND IVACAFTOR - PCT only - Special Authority see SA2456 below

Tab elexacaftor 50 mg with tezacaftor 25 mg, ivacaftor 37.5 mg ✓ Trikafta 84 OP Tab elexacaftor 100 mg with tezacaftor 50 mg, ivacaftor 75 mg (56) and ivacaftor 150 mg (28) .......27,647.39 84 OP ✓ Trikafta

#### ⇒SA2456 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with cystic fibrosis; and
- 2 Patient is 6 years of age or older; and
- 3 Fither:

Subsidy	Fully	Brand or
(Manufacturer's Price)		Generic
 \$	Per 🗸	Manufacturer

#### continued...

- 3.1 Patient has two cystic fibrosis-causing mutations in the cystic fibrosis transmembrane regulator (CFTR) gene (one from each parental allele); or
- 3.2 Patient has a sweat chloride value of at least 60 mmol/L by quantitative pilocarpine iontophoresis or by Macroduct sweat collection system; and
- 4 Fither:
  - 4.1 Patient has a heterozygous or homozygous F508del mutation; or
  - 4.2 Patient has a G551D mutation or other mutation responsive in vitro to elexacaftor/tezacaftor/ivacaftor (see note a); and
- 5 The treatment must be the sole funded CFTR modulator therapy for this condition; and
- 6 Treatment with elexacaftor/tezacaftor/ivacaftor must be given concomitantly with standard therapy for this condition.

#### Notes:

 a) Eligible mutations are listed in the Food and Drug Administration (FDA) Trikafta prescribing information <a href="https://nctr-crs.fda.gov/fdalabel/services/spl/set-ids/f354423a-85c2-41c3-a9db-0f3aee135d8d/spl-doc">https://nctr-crs.fda.gov/fdalabel/services/spl/set-ids/f354423a-85c2-41c3-a9db-0f3aee135d8d/spl-doc</a>

IVACAFTOR - PCT only - Specialist - Special Author	rity see SA2017 below		
Tab 150 mg	29,386.00	56	✓ Kalydeco
Oral granules 50 mg, sachet	29,386.00	56	✓ Kalydeco
Oral granules 75 mg, sachet	29,386.00	56	✓ Kalydeco

## ⇒SA2017 Special Authority for Subsidy

Initial application only from a respiratory specialist or paediatrician. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### All of the following:

- 1 Patient has been diagnosed with cystic fibrosis; and
- 2 Either:
  - 2.1 Patient must have G551D mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene on at least 1 allele; or
  - 2.2 Patient must have other gating (class III) mutation (G1244E, G1349D, G178R, G551S, S1251N, S1255P, S549N and S549R) in the CFTR gene on at least 1 allele; and
- 3 Patients must have a sweat chloride value of at least 60 mmol/L by quantitative pilocarpine iontophoresis or by Macroduct sweat collection system; and
- 4 Treatment with ivacaftor must be given concomitantly with standard therapy for this condition; and
- 5 Patient must not have an acute upper or lower respiratory infection, pulmonary exacerbation, or changes in therapy (including antibiotics) for pulmonary disease in the last 4 weeks prior to commencing treatment with ivacaftor; and
- 6 The dose of ivacaftor will not exceed one tablet or one sachet twice daily; and
- 7 Applicant has experience and expertise in the management of cystic fibrosis.

## SODIUM CHLORIDE

Not funded for use as a nasal drop.

## **Nasal Preparations**

# **Allergy Prophylactics**

Metered aqueous nasal spray, 50 mcg per dose		✓ <u>SteroClear</u> ✓ <u>SteroClear</u>
Metered aqueous nasal spray, 50 mcg per dose2.57	120 dose OP	✓ Flixonase Hayfever & Allergy

Oral liq 20 mg per ml (10 mg base per ml)......16.91

	Subsidy (Manufacturer's Pric	ce) Subs	Fully	Brand or Generic
	\$	Per		Manufacturer
IPRATROPIUM BROMIDE				
Aqueous nasal spray, 0.03%	5.23	15 ml OP	<b>✓</b> U	Inivent
Respiratory Devices				
MASK FOR SPACER DEVICE				
a) Up to 50 dev available on a PSO				
b) Only on a PSO				
<ul> <li>c) Only for children aged six years and under</li> </ul>				
Small	2.70	1	<b>√</b> e	-chamber Mask
PEAK FLOW METER				
a) Up to 25 dev available on a PSO				
b) Only on a PSO				
Low range	9.54	1	✓ N	/lini-Wright AFS Low Range
Normal range	9.54	1	✓ N	/lini-Wright Standard
CDACED DEVICE				Stanuaru
SPACER DEVICE				
a) Up to 50 dev available on a PSO     b) Only on a PSO				
220 ml (single patient)	3 65	1	<b>√</b> e	-chamber Turbo
510 ml (single patient)		1	-	-chamber La
(* 9 - 1 - 1 - 1				Grande
800 ml	6.50	1	<b>✓</b> V	olumatic
Boonisstery Stimulanto				
Respiratory Stimulants				
CAFFEINE CITRATE				

✓ Biomed

25 ml OP

			SENSORY ORGANS
	Subsidy (Manufacturer's F \$	Price) Subs Per	Fully Brand or sidised Generic  Manufacturer
Ear Preparations			
FLUMETASONE PIVALATE Ear drops 0.02% with clioquinol 1%	4.46	7.5 ml OP	<ul><li>✓ Locacorten-Viaform ED's</li><li>✓ Locorten-Vioform</li></ul>
TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYCI Ear drops 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 mg and gramicidin 250 mcg per g		7.5 ml OP	✓ Kenacomb
Ear/Eye Preparations			
DEXAMETHASONE WITH FRAMYCETIN AND GRAMICIDIN Ear/Eye drops 500 mcg with framycetin sulphate 5 mg and gramicidin 50 mcg per ml	4.50 (9.27)	8 ml OP	Sofradex
FRAMYCETIN SULPHATE Ear/Eye drops 0.5%	4.13 (8.65)	8 ml OP	Soframycin
Eye Preparations			
Eye preparations are only funded for use in the eye, unless explicit	citly stated other	wise.	
Anti-Infective Preparations			
ACICLOVIR  * Eye oint 3%	15.89	4.5 g OP	✓ <u>ViruPOS</u>
CHLORAMPHENICOL  Eye oint 1%  Devatis to be Principal Supply on 1 February 2026	1.55	5 g OP	✓ Devatis
Eye drops 0.5%	1.45 1.84	10 ml OP	<ul><li>✓ Chlorsig</li><li>✓ Chlorafast</li></ul>
<ul> <li>a) Funded for use in the ear*. Indications marked with b) Chlorafast to be Principal Supply on 1 March 2026 (Chlorsig Eye drops 0.5% to be delisted 1 March 2026)</li> <li>CIPROFLOXACIN</li> </ul>	* are unapproved	d indications.	
Eye drops 0.3% – Subsidy by endorsement	or severe bacteria s media (CSOM)	•	·
SODIUM FUSIDATE [FUSIDIC ACID] Eye drops 1%	5.29	5 g OP	<ul> <li>✓ Fucithalmic</li> <li>✓ Fucithalmic Canada         (ON) \$29</li> <li>✓ Fucithalmic         Spain \$29</li> </ul>
TOBRAMYCIN	40.45	0.5 - 0.0	✓ Tahway

3.5 g OP

5 ml OP

✓ Tobrex

✓ Tobrex

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.



	Subsidy	Fully		Brand or
(Manu	facturer's Price)	Subsidised		Generic
	\$	Per	/	Manufacturer

## Corticosteroids and Other Anti-Inflammatory Preparations

DEXAMETHASONE	
* Eye oint 0.1%	3.5 g OP
* Eve drops 0.1%4.50	5 ml OP

Ocular implant 700 mcg - Special Authority see SA1680 below

✓ Ozurdex

✓ Maxidex ✓ Maxidex

## ⇒SA1680 Special Authority for Subsidy

Initial application — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has diabetic macular oedema with pseudophakic lens; and
- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- - 3.1 Patient's disease has progressed despite 3 injections with bevacizumab; or
  - 3.2 Patient is unsuitable or contraindicated to treatment with anti-VEGF agents; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eve, and up to a maximum of 3 implants per eye per year.

Initial application — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has diabetic macular oedema; and
- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- 3 Patient is of child bearing potential and has not yet completed a family; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Patient is of child bearing potential and has not yet completed a family; and
- 3 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

#### DEXAMETHASONE WITH NEOMYCIN SULPHATE AND POLYMYXIN B SULPHATE

* Eye oint 0.1% with neomycin sulphate 0.35% and polymyxin b	- 00	0.5.00	<b>4.88</b> 11 1
sulphate 6,000 u per g	5.39	3.5 g OP	✓ Maxitrol
★ Eye drops 0.1% with neomycin sulphate 0.35% and polymyxin			
b sulphate 6,000 u per ml	4.50	5 ml OP	✓ Maxitrol
DICLOFENAC SODIUM			
Eye drops 0.1%, single dose	1.85	10 dose	<ul> <li>Diclofenac Devatis</li> </ul>
	5.54	30 dose	✓ Diclofenac Devatis
FLUOROMETHOLONE			
* Eye drops 0.1%	3.09	5 ml OP	✓ FML
•	5.20		✓ Flucon

✓ fully subsidised

**Principal Supply** 

			SENSORY ORGANS
	Subsidy (Manufacturer's Pri	ce) Subsi Per	Fully Brand or idised Generic  Manufacturer
LEVOCABASTINE Eye drops 0.5 mg per ml	8.71 (10.34)	4 ml OP	Livostin
LODOXAMIDE Eye drops 0.1% PREDNISOLONE ACETATE	8.71	10 ml OP	✓ Lomide
Eye drops 1%	6.92 7.00	10 ml OP 5 ml OP	<ul><li>✓ Prednisolone-AFT</li><li>✓ Pred Forte</li></ul>
PREDNISOLONE SODIUM PHOSPHATE – Special Authority se Eye drops 0.5%, single dose (preservative free)		- Retail pharm 20 dose	nacy ✓ Minims  Prednisolone
Initial application only from an ophthalmologist or optometrist.  following criteria:  Both:  1 Patient has severe inflammation; and 2 Patient has a confirmed allergic reaction to preservative in Renewal from any relevant practitioner. Approvals valid for 6 modern from the property of	n eye drops.		
SODIUM CROMOGLICATE  Eye drops 2%  Allerfix to be Principal Supply on 1 March 2026	2.91	10 ml OP	✓ Allerfix
Glaucoma Preparations - Beta Blockers			
TIMOLOL  * Eye drops 0.25%  * Eye drops 0.5%		5 ml OP 5 ml OP	✓ <u>Arrow-Timolol</u> ✓ <u>Arrow-Timolol</u>
Glaucoma Preparations - Carbonic Anhydrase I	nhibitors		
ACETAZOLAMIDE  * Tab 250 mg  BRINZOLAMIDE	13.96	100	✓ <u>Medsurge</u>
* Eye drops 1%  DORZOLAMIDE WITH TIMOLOL		5 ml OP	✓ Azopt
* Eye drops 2% with timolol 0.5%		5 ml OP	✓ <u>Dortimopt</u>
BIMATOPROST	ues		
* Eye drops 0.03%		3 ml OP	✓ <u>Lumigan</u>
* Eye drops 0.005%		2.5 ml OP	✓ <u>Teva</u> ✓ Travatan
	0.00	2.5 ml OP	▼ <u>ITAVALAII</u>
Glaucoma Preparations - Other  BRIMONIDINE TARTRATE			
V Fue drope 0.00/	F 10	E ml OD	Auram Brimanidina

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

✓ Arrow-Brimonidine

5 ml OP

<sup>\*</sup>Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's P	rice) Subsi	Fully	Brand or Generic
	\$	Per	<b>√</b>	Manufacturer
BRIMONIDINE TARTRATE WITH TIMOLOL MALEATE  * Eye drops 0.2% with timolol maleate 0.5%	7.13	5 ml OP	<b>✓</b> 0	Combigan
LATANOPROST WITH TIMOLOL  * Eye drops 0.005% with timolol 0.5%	4.95	2.5 ml OP	✓ <u>A</u>	arrow - Lattim
PILOCARPINE HYDROCHLORIDE  * Eye drops 1%		15 ml OP		sopto Carpine
* Eye drops 2%  * Eye drops 4%	7.99	15 ml OP 15 ml OP		sopto Carpine sopto Carpine
Subsidised for oral use pursuant to the Standard Formula PILOCARPINE NITRATE	ae.			
Eye drops 2% single dose – Special Authority see SA0895     below – Retail pharmacy	35.90	20 dose	✓ N	linims Pilocarpine

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient has to use an unpreserved solution due to an allergy to the preservative; or
- 2 Patient wears soft contact lenses.

Note: Minims for a general practice are considered to be "tools of trade" and are not approved as special authority items. Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

# Mydriatics and Cycloplegics

ATROPINE SULPHATE		
* Eye drops 1%	.27 15 ml OP	✓ Atropt
CYCLOPENTOLATE HYDROCHLORIDE		
* Eye drops 1%25	.16 15 ml OP	<ul><li>Cyclogyl</li></ul>
TROPICAMIDE		
* Eye drops 0.5%	.52 15 ml OP	✓ Mydriacyl
* Eye drops 1%24	.82 15 ml OP	✓ Mydriacyl

# **Preparations for Tear Deficiency**

For acetylcysteine eye drops refer Standard Formulae, page 284

#### **HYPROMELLOSE**

<b>*</b> Eye drops 0.5%	15 ml OP	Methopt
HYPROMELLOSE WITH DEXTRAN		
* Eye drops 0.3% with dextran 0.1%2.30	15 ml OP	✓ Poly-Tear

#### **Preservative Free Ocular Lubricants**

## ⇒SA2431 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Both:

- 1 Confirmed diagnosis by slit lamp or Schirmer test of severe secretory dry eye; and
- - 2.1 Patient is using eye drops more than four times daily on a regular basis; or
  - 2.2 Patient has had a confirmed allergic reaction to preservative in eye drop.

✓ Poly-Tears

# **SENSORY ORGANS**

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	
POLYETHYLENE GLYCOL 400 AND PROPYLENE GLYCOL – pharmacy	Special Authority see	SA2431 on the	previous page - Retail
Eye drops 0.4% and propylene glycol 0.3%, 0.8 ml	10.78	30	Systane Unit Dose
SODIUM HYALURONATE [HYALURONIC ACID] - Special Autl	,		• ,
Eye drops 1 mg per ml	13.58 10	ml OP	Hylo-Fresh
Hylo-Fresh has a 6 month expiry after opening. The Ph	,		

# Other Eye Preparations

NAPHAZOLINE HYDROCHLORIDE  * Eye drops 0.1%	15 ml OP	✓ <u>Albalon</u>
OLOPATADINE Eye drops 0.1%3.39	5 ml OP	✓ Olopatadine Teva
Olopatadine Teva to be Principal Supply on 1 March 2026 PARAFFIN LIQUID WITH WOOL FAT		
* Eye oint 3% with wool fat 3%	3.5 g OP	✓ Poly-Visc
Eye oint 138 mcg per g	5 g OP	✓ VitA-POS



Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer **Various** PHARMACY SERVICES 1 fee ✓ BSF Allegron ✓ BSF Estradiol TDP Mvlan a) May only be claimed once per patient. b) The Pharmacode for BSF Allegron is 2715740 - see also page 129 c) The Pharmacode for BSF Estradiol TDP Mylan is 2717573 - see also page 86 ✓ Immunisation - Flu 1 fee Immunisation administration fee - other 0.00 1 fee ✓ Immunisation Other Immunisation co-administration fee - flu and shingles...............................0.00 ✓ Immunisation Flu 1 fee and Shingles ✓ Paxlovid fee 1 fee (BSF Allegron Brand switch fee to be delisted 1 February 2026) (BSF Estradiol TDP Mylan Brand switch fee to be delisted 1 March 2026) Agents Used in the Treatment of Poisonings **Antidotes ACETYLCYSTEINE** 10 ✓ DBL Acetvlcvsteine 10 ✓ Hikma Acetylcysteine \$29 NAI OXONE HYDROCHI ORIDE a) Up to 10 inj available on a PSO b) Only on a PSO 5 ✓ DBL Naloxone Hydrochloride Removal and Elimination CHARCOAL \* Oral lig 50 g per 250 ml ......59.85 250 ml OP ✓ Carbosorb-X a) Up to 250 ml available on a PSO b) Only on a PSO DEFERASIROX - Special Authority see SA1492 below - Retail pharmacy Wastage claimable Exiade 28 ✓ Exjade 28 Exiade ⇒SA1492 Special Authority for Subsidy Initial application only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria:

1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; and

- 2 Deferasirox is to be given at a daily dose not exceeding 40 mg/kg/day; and

continued...

All of the following:

Subsidy (Manufacturer's Price) \$	Sı Per	Fully ubsidised	Brand or Generic Manufacturer	
---	-----------	-----------------	-------------------------------------	--

#### continued...

- 3 Any of the following:
  - 3.1 Treatment with maximum tolerated doses of deferiprone monotherapy or deferiprone and desferrioxamine combination therapy have proven ineffective as measured by serum ferritin levels, liver or cardiac MRI T2\*; or
  - 3.2 Treatment with deferiprone has resulted in severe persistent vomiting or diarrhoea; or
  - 3.3 Treatment with deferiprone has resulted in arthritis: or
  - 3.4 Treatment with deferiprone is contraindicated due to a history of agranulocytosis (defined as an absolute neutrophil count (ANC) of < 0.5 cells per μL) or recurrent episodes (greater than 2 episodes) of moderate neutropenia (ANC 0.5 1.0 cells per μL).</p>

Renewal only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria: Fither:

- 1 For the first renewal following 2 years of therapy, the treatment has been tolerated and has resulted in clinical improvement in all three parameters namely serum ferritin, cardiac MRI T2\* and liver MRI T2\* levels; or
- 2 For subsequent renewals, the treatment has been tolerated and has resulted in clinical stability or continued improvement in all three parameters namely serum ferritin, cardiac MRI T2\* and liver MRI T2\* levels.

DEFERIPRONE - Special Authority see SA1480 below - Retail p	harmacy		
Tab 500 mg	533.17	100	✓ Ferriprox
Oral liq 100 mg per 1 ml	266.59	250 ml OP	✓ Ferriprox

## **⇒SA1480** Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Fither:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; or
- 2 The patient has been diagnosed with chronic iron overload due to acquired red cell aplasia.

DESFERRIOXAMINE MESILATE  * Inj 500 mg vial	332.88	10	✓ DBL  Desferrioxamine  Mesylate for Inj  BP
SODIUM CALCIUM EDETATE			
* Inj 200 mg per ml, 5 ml	53.31	6	
	(156.71)		Calcium Disodium Versenate



# **Standard Formulae**

Statiualu Futiliulae			
ACETYLCYSTEINE EYE DROPS Acetylcysteine inj 200 mg per ml, 10 ml	qs	PHENOBARBITONE SODIUM PAEDIATRIC ORAL mg per ml)	LIQUID (10
Suitable eye drop base	qs	Phenobarbitone Sodium Glycerol BP	400 mg 4 ml
CODEINE LINCTUS (3 mg per 5 ml) Codeine phosphate	60 mg	Water	to 40 ml
Glycerol Preservative	40 ml gs	PILOCARPINE ORAL LIQUID Pilocarpine 4% eye drops	qs
Water	to 100 ml	Preservative	qs
CODEINE LINCTUS (15 mg per 5 ml)		Water (Preservative should be used if quantity supplied is	to 500 ml
Codeine phosphate Glycerol	300 mg 40 ml	than 5 days.)	00.0
Preservative Water	qs to 100 ml	SALIVA SUBSTITUTE FORMULA Methylcellulose	5 g
FOLINIC MOUTHWASH		Preservative Water	qs to 500 ml
Calcium folinate 15 mg tab Preservative Water	1 tab qs to 500 ml	(Preservative should be used if quantity supplied is than 5 days. Maximum 500 ml per prescription.)	or more
(Preservative should be used if quantity supplied is f than 5 days. Maximum 500 ml per prescription.)		SODIUM CHLORIDE ORAL LIQUID Sodium chloride inj 23.4%, 20 ml Water	qs qs
METHYL HYDROXYBENZOATE 10% SOLUTION Methyl hydroxybenzoate	10 g	(Only funded if prescribed for treatment of hyponatra	•
Propylene glycol (Use 1 ml of the 10% solution per 100 ml of oral liqui	to 100 ml	VANCOMYCIN ORAL SOLUTION (25 mg per ml) Vancomycin 500 mg injection Glycerin with sucrose suspension	5 vials 37.5 ml
OMEPRAZOLE SUSPENSION Omeprazole capsules or powder Sodium bicarbonate powder BP Water	qs 8.4 g to 100 ml	Water (Only funded if prescribed for treatment of Clostridiu following metronidazole failure)	to 100 ml
PHENOBARBITONE ORAL LIQUID Phenobarbitone Sodium Glycerol BP Water	1 g 70 ml to 100 ml		

# EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer

Extemporaneously Compounded Preparations and Galenicals

COLLODION FLEXIBLE	ما الثين لمسم سمنام		
Note: This product is no longer being manufactured by the sup determined.	plier and will b	oe delisted troi	m the Schedule at a date to be
Collodion flexible	19.30	100 ml	✓ PSM
COMPOUND HYDROXYBENZOATE - Only in combination		100 1111	- 10111
Only in extemporaneously compounded oral mixtures.			
Soln	36.00	100 ml	✓ Midwest
GLYCERIN WITH SODIUM SACCHARIN - Only in combination			
Suspension	38.00	473 ml	✓ Ora-Sweet SF
GLYCERIN WITH SUCROSE - Only in combination			
Suspension	38.00	473 ml	✓ Ora-Sweet
GLYCEROL			
* Liquid – Only in combination		500 ml	✓ healthE Glycerol BP
Only in extemporaneously compounded oral liquid prepara	tions.		
METHYL HYDROXYBENZOATE			
Powder	11.00	25 g	✓ Midwest
METHYLCELLULOSE			
Powder		100 g	✓ MidWest
Suspension – Only in combination	38.00	473 ml	✓ Ora-Plus
(MidWest Powder to be delisted 1 February 2028)	NINI - On to the		
METHYLCELLULOSE WITH GLYCERIN AND SODIUM SACCHAR Suspension		combination 473 ml	✓ Ora-Blend SF
•			▼ Ora-Dieliu 3F
METHYLCELLULOSE WITH GLYCERIN AND SUCROSE - Only in Suspension		473 ml	✓ Ora-Blend
PHENOBARBITONE SODIUM	36.00	4/31111	♥ Ora-Dieliu
Powder – Only in combination	125.00	10 g	✓ MidWest
Only in children up to 12 years	123.00	10 9	· marrest
PROPYLENE GLYCOL			
Only in extemporaneously compounded methyl hydroxybenzoa	te 10% solutio	n.	
Liq		500 ml	✓ Midwest
SODIUM BICARBONATE			
Powder BP - Only in combination		500 g	✓ Midwest
Only in extemporaneously compounded omeprazole and la	insoprazole su	ispension.	
SYRUP (PHARMACEUTICAL GRADE) - Only in combination			
Only in extemporaneously compounded oral liquid preparations		·	
Liq	25.00	500 ml	✓ Midwest
WATER			

✓ Tap water

1 ml

Tap - Only in combination......0.00

Subsidy (Manufacturer's Price)

Fully Subsidised Per

Brand or Generic Manufacturer

# **Nutrient Modules**

## Carbohydrate

#### ⇒SA1930 Special Authority for Subsidy

Initial application — (Cystic fibrosis or kidney disease) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria: Fither:

- 1 cystic fibrosis; or
- 2 chronic kidney disease.

Initial application — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 cancer in children: or
- 2 cancers affecting alimentary tract where there are malabsorption problems in patients over the age of 20 years; or
- 3 faltering growth in an infant/child; or
- 4 bronchopulmonary dysplasia; or
- 5 premature and post premature infant; or
- 6 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified where the patient has inborn errors of metabolism. Renewal — (Cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

CARBOHYDRATE SUPPLEMENT - Special Authority see SA1930 above - Hospital pharmacy [HP3]

400 a OP ✓ Polycal Powder .......6.72

# Carbohydrate And Fat

# ⇒SA1376 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria: Both:

Subsidy		Fully	Brand or	
(Manufacturer's Pri	ce)	Subsidised	Generic	
\$	Per	•	Manufacturer	

continued...

- 1 Infant or child aged four years or under; and
- 2 cystic fibrosis.

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 infant or child aged four years or under; and
- 2 Any of the following:
  - 2.1 cancer in children; or
  - 2.2 faltering growth; or
  - 2.3 bronchopulmonary dysplasia; or
  - 2.4 premature and post premature infants.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

#### Fat

## ⇒SA2204 Special Authority for Subsidy

Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified where the patient has an inborn error of metabolism. Initial application — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 faltering growth in an infant/child; or
- 2 bronchopulmonary dysplasia; or
- 3 fat malabsorption; or
- 4 lymphangiectasia; or
- 5 short bowel syndrome: or
- 6 infants with necrotising enterocolitis; or
- 7 biliary atresia; or
- 8 for use in a ketogenic diet; or
- 9 chyle leak; or

continued...

✓ fully subsidised 287



Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	•	Manufacturer

continued...

- 10 ascites: or
- 11 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. Renewal — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT SUPPLEMENT	- Special Authority	see SA2204 on	the previous pa	ge – Hospital	pharmacy [HP3]

Emulsion (neutral)	15.38	200 ml OP	✓ Calogen
Emulsion (strawberry)	15.38	200 ml OP	✓ Calogen
Oil	37.50	500 ml OP	✓ MCT oil (Nutricia)
MCT Emulsion, 250 ml	143.65	4 OP	✓ Liquigen ′

## **Protein**

## **⇒SA1524** Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 protein losing enteropathy; or
- 2 high protein needs; or
- 3 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

#### Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

PROTEIN SUPPLEMENT - Special Authority see SA1524 above - Hospital pharmacy [HP3]

	openia riamoni, coo ornoz razoro			
Powder		8.95	227 g OP	✓ Resource
			•	Beneprotein
		13.82	225 g OP	✓ Protifar

Subsidy (Manufacturer's Price)

Subsidised Per

Fully

Brand or Generic Manufacturer

# **Oral and Enteral Feeds**

#### **Diabetic Products**

### ⇒SA1095 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is a type I or and II diabetic who is suffering weight loss and malnutrition that requires nutritional support. Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

#### Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

DIABETIC ENTERAL FEED 1KCAL/ML - Special Authority see \$	SA1095 above – H	ospital pharr	nacy [HP3]
Liquid, 500 ml bottle	4.65	1 OP	✓ Glucerna Select
DIABETIC ORAL FEED 1KCAL/ML - Special Authority see SA10	95 above – Hosp	ital pharmacy	/ [HP3]
Liquid (strawberry), 200 ml bottle	2.25	1 OP	✓ Diasip
Liquid (vanilla), 200 ml bottle	2.10	1 OP	✓ Nutren Diabetes
	2.25		✓ Diasip

### **Fat Modified Products**

### ⇒SA2205 Special Authority for Subsidy

Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified where the patient has an inborn error of metabolism. Initial application — (Indications other than errors of inborn metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Patient has a chyle leak; or
- 2 Modified as a modular feed, made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule, for adults,

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

#### Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT MODIFIED FEED - Special Authority see SA2205 above - Hospital pharmacy [HP3] 400 a OP Monogen



Subsidy (Manufacturer's Price) Fully Subsidised Per ✓ Brand or Generic Manufacturer

### Paediatric Products For Children Awaiting Liver Transplant

### ⇒SA1098 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) who requires a liver transplant.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL FEED 1KCAL/ML - Special Authority see SA1098 above - Hospital pharmacy [HP3]

### Paediatric Products For Children With Chronic Renal Failure

# ⇒SA1099 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) with acute or chronic kidney disease.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

#### Paediatric Products

### ⇒SA1379 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Child is aged one to ten years; and
- 2 Any of the following:
  - 2.1 the child is being fed via a tube or a tube is to be inserted for the purposes of feeding; or
  - 2.2 any condition causing malabsorption; or
  - 2.3 faltering growth in an infant/child; or
  - 2.4 increased nutritional requirements; or
  - 2.5 the child is being transitioned from TPN or tube feeding to oral feeding.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for

continued...

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	

continued...

applications meeting the following criteria:

#### Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

	previous pag 1 OP	ge – Hospital pharmacy [HP3]  ✓ Nutrini Energy RTH
	revious page 1 OP	<ul><li>− Hospital pharmacy [HP3]</li><li>✓ Pediasure RTH</li><li>✓ Nutrini RTH</li></ul>
uthority see S	A1379 on the	e previous page – Hospital
7.14	1 OP	✓ Nutrini Energy Multi Fibre
79 on the pre	vious page -	Hospital pharmacy [HP3]
1.90	1 OP	✓ Fortini
1.90	1 OP	✓ Fortini
	1 OP	✓ Pediasure Plus
on the previo	ous page – H	ospital pharmacy [HP3]
1.33	1 OP	✓ Pediasure
1.33	1 OP	✓ Pediasure
	1 OP	✓ Pediasure
rity see SA13	79 on the pre	evious page - Hospital
1.90	1 OP	✓ Fortini Multi Fibre
	1 OP	✓ Fortini Multi Fibre
1.90	1 OP	✓ Fortini Multi Fibre
	1 OP	✓ Fortini Multi Fibre
	e – Hospital	pharmacy [HP3]
43.60	100 g OP	✓ Peptamen Junior
	7.46 1379 on the p3.32 4.69 uthority see S7.14 79 on the previous1.901.901.331.331.331.331.331.391.901.901.901.901.90 previous pag	7.46 1 OP 1379 on the previous page3.32 1 OP 4.69 uthority see SA1379 on the7.14 1 OP 79 on the previous page –1.90 1 OP1.90 1 OP1.33 1 OP1.33 1 OP1.33 1 OP1.33 1 OP1.33 1 OP1.39 1 OP1.39 1 OP1.90 1 OP

### **Renal Products**

### ⇒SA1101 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has acute or chronic kidney disease.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

#### Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

RENAL ORAL FEED 1.8 KCAL/ML - Special Authority see	SA1101 above - Hospi	ital pharmac	y [HP3]
Liquid, 220 ml bottle	3.31	1 OP	✓ Nepro HP
			(strawberry)
			✓ Nepro HP (vanilla)

	(Manufacturer's Price)	Subsid	dised	Generic
	\$	Per	•	Manufacturer
RENAL ORAL FEED 2 KCAL/ML - Special Authority see SA110	1 on the previous page	ge – Hospita	al phar	macy [HP3]
Liquid, 200 ml bottle	13.24	4 OP	✓ N	ovaSource Renal
Liquid (apricot) 125 ml	13.72	4 OP	✓ R	enilon 7.5
Liquid (caramel) 125 ml	13.72	4 OP	✓ R	enilon 7.5

Subsidy

Fully

Brand or

### **Specialised And Elemental Products**

### ⇒SA1377 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 malabsorption; or
- 2 short bowel syndrome; or
- 3 enterocutaneous fistulas: or
- 4 eosinophilic oesophagitis; or
- 5 inflammatory bowel disease: or
- 6 patients with multiple food allergies requiring enteral feeding.

Notes: Each of these products is highly specialised and would be prescribed only by an expert for a specific disorder. The alternative is hospitalisation.

Elemental 028 Extra is more expensive than other products listed in this section and should only be used where the alternatives have been tried first and/or are unsuitable.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL SEMI-ELEMENTAL FEED 1.5KCAL/ML – S Liquid, 1,000 ml bottle		SA1377 abor 1 OP	ve – Hospital pharmacy [HP3]  ✓ Vital
ORAL ELEMENTAL FEED 0.8KCAL/ML - Special Authority s	ee SA1377 above -	Hospital pha	rmacy [HP3]
Liquid (grapefruit), 250 ml carton	179.46	18 OP	✓ Elemental 028 Extra
Liquid (pineapple & orange), 250 ml carton		18 OP	✓ Elemental 028 Extra
Liquid (summer fruits), 250 ml carton	179.46	18 OP	✓ Elemental 028 Extra
ORAL ELEMENTAL FEED 1KCAL/ML - Special Authority see	e SA1377 above – He	ospital pharm	nacy [HP3]
Powder (unflavoured), 80 g sachet	4.50	1 OP	✓ Vivonex TEN
SEMI-ELEMENTAL ENTERAL FEED 1KCAL/ML - Special Au	uthority see SA1377	above – Hos	pital pharmacy [HP3]
Liquid, 500 ml bottle	7.47	1 OP	✓ Nutrison Advanced
·			Peptisorb

# Paediatric Products For Children With Low Energy Requirements

### **⇒SA1196** Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

continued...

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1 Child aged one to eight years; and
  - 2 The child has a low energy requirement but normal protein and micronutrient requirements.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

PAEDIATRIC ENTERAL FEED WITH FIBRE 0.76 KCAL/ML - Special Authority see SA1196 on the previous page - Hospital pharmacy [HP3]

### Standard Supplements

### ⇒SA1859 Special Authority for Subsidy

Initial application — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age; and
- 2 Any of the following:
  - 2.1 The patient has a condition causing malabsorption; or
  - 2.2 The patient has failure to thrive; or
  - 2.3 The patient has increased nutritional requirements; and
- 3 Nutrition goal has been set (eg reach a specific weight or BMI).

Renewal — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 A nutrition goal has been set (eg reach a specific weight or BMI).

Initial application — (Children - exclusive enteral nutrition for Crohn's disease) only from a gastroenterologist or dietitian on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 Dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Renewal — (Children - exclusive enteral nutrition for Crohn's disease) from any relevant practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 General Practitioners and dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

continued...

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

Initial application — (Adults) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

1 Any of the following:

Patient is Malnourished

- 1.1 Patient has a body mass index (BMI) of less than 18.5 kg/m<sup>2</sup>; or
- 1.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 1.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months; and
- 2 Any of the following:

Patient has not responded to first-line dietary measures over a 4 week period by:

- 2.1 Increasing their food intake frequency (eg snacks between meals); or
- 2.2 Using high-energy foods (e.g. milkshakes, full fat milk, butter, cream, cheese, sugar etc); or
- 2.3 Using over the counter supplements (e.g. Complan); and
- 3 A nutrition goal has been set (e.g. to reach a specific weight or BMI).

Renewal — (Adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 A nutrition goal has been set (eg reach a specific weight or BMI); and
- 2 Any of the following:

Patient is Malnourished

- 2.1 Patient has a body mass index (BMI) of less than 18.5 kg/m<sup>2</sup>; or
- 2.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 2.3 Patient has a BMI of less than 20 kg/m<sup>2</sup> and unintentional weight loss greater than 5% within the last 3-6 months.

**Initial application — (Short-term medical condition)** from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube or a nasogastric tube is to be inserted for feeding; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Is undergoing a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both
  - 5.1 Pregnant: and
  - 5.2 Any of the following:
    - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or
    - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
    - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

**Renewal — (Short-term medical condition)** from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Has undergone a bone marrow transplant; or

continued...

Subsidy (Manufacturer's Price)	Subsi	Fully	Brand or Generic
\$	Per	<b>√</b>	Manufacturer

#### continued...

- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
  - 5.1 Pregnant: and
  - 5.2 Any of the following:
    - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or
    - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
    - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

Initial application — (Long-term medical condition) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

### Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis: or
- 3 Liver disease: or
- 4 Chronic Renal failure: or
- 5 Inflammatory bowel disease; or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome: or
- 8 Bowel fistula: or
- 9 Severe chronic neurological conditions: or
- 10 Epidermolysis bullosa; or
- 11 AIDS (CD4 count < 200 cells/mm<sup>3</sup>): or
- 12 Chronic pancreatitis.

Renewal — (Chronic disease OR tube feeding for patients who have previously been funded under Special Authority forms SA0702 or SA0583) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

### Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis; or
- 3 Liver disease; or
- 4 Chronic Renal failure; or
- 5 Inflammatory bowel disease; or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome; or
- 8 Bowel fistula: or
- 9 Severe chronic neurological conditions.

ENTERAL FEED 1.5KCAL/ML - Special Authority see SA1859 on	page 293 - Ho	spital pharma	acy [HP3]
Liquid, 1,000 ml bottle	8.68	1 OP	✓ Ensure Plus HN
			RTH
	9.00		✓ Nutrison Energy
Liquid, 250 ml can	2.17	1 OP	✓ Ensure Plus HN
(Ensure Plus HN Liquid, 250 ml can to be delisted 1 March 2026)			

	Subsidy (Manufacturer's Pr \$	rice) Subs Per	Fully Brand or idised Generic Manufacturer
			r. Inc.
ENTERAL FEED 1KCAL/ML - Special Authority see SA1859 on			
Liquid, 1,000 ml bottle		1 OP	<ul><li>Osmolite RTH</li></ul>
	6.90		✓ Nutrison RTH
ENTERAL FEED WITH FIBRE 0.83 KCAL/ML — Special Authorit Liquid, 1,000 ml bottle		n page 293 – H 1 OP	ospital pharmacy [HP3]  Nutrison 800 Complete Multi Fibre
ENTERAL FEED WITH FIBRE 1 KCAL/ML - Special Authority s	ee SA1859 on pa	age 293 – Host	oital pharmacy [HP3]
Liquid, 1,000 ml bottle		1 OP	✓ Jevity RTH
Liquid, 1,000 mi bottio	7.21	1 01	✓ Nutrison Multi Fibre
ENTERAL FEED WITH FIBRE 1.5KCAL/ML - Special Authority	see SA1859 on p	oage 293 – Hos	spital pharmacy [HP3]
Liquid, 1,000 ml bottle	8.68	1 OP	✓ Jevity HiCal RTH
			✓ Nutrison Energy
			Multi Fibre
ODAL EEED (DOM/DED) Consist Authority and CA1050 and an	000		201
ORAL FEED (POWDER) - Special Authority see SA1859 on page			
Powder (chocolate)	15.90	840 g OP	✓ Sustagen Hospital
			Formula
	40.00	850 g OP	✓ Ensure
Powder (vanilla)	15.90	840 g OP	<ul> <li>Sustagen Hospital</li> </ul>
,		Ü	Formula
	40.00	850 g OP	✓ Ensure
ORAL FEED 1.5KCAL/ML - Special Authority see SA1859 on pa		ŭ	
Additional subsidy by endorsement is available for patients be epidermolysis bullosa, or as exclusive enteral nutrition for the hypercapnia, defined as CO2 value exceeding 55mmHg. The Liquid (banana), 200 ml bottle — Higher subsidy of up to \$1.	peing bolus fed three treatment of Cro ne prescription mu	rough a feeding ohn's disease,	g tube, who have severe or for patients with COPD and
per 1 btl with Endorsement	0.72	1 OP	
	(1.56)		Ensure Plus
	(1.76)		Fortisip
Liquid (chocolate), 200 ml bottle - Higher subsidy of up to	,		·
\$1.76 per 1 btl with Endorsement	0.72	1 OP	
Ψ1.70 per 1 bit with Endorsement	(1.56)	1 01	Ensure Plus
	( /		Fortisip
1: :1" :1 (1: 1: 1: 1: 1: 1: 1: 1: 1: 1: 1: 1: 1: 1	(1.76)		romsip
Liquid (fruit of the forest), 200 ml bottle – Higher subsidy of			
\$1.56 per 1 btl with Endorsement		1 OP	
	(1.56)		Ensure Plus
Liquid (strawberry), 200 ml bottle - Higher subsidy of \$1.76	per		
1 btl with Endorsement	0.72	1 OP	
	(1.76)		Fortisip
Liquid (vanilla), 200 ml bottle - Higher subsidy of up to \$1.70	` '		. э.н.э.р
per 1 btl with Endorsement		1 OP	
per i bu with Endorsement		1 01	Engues Divo
	(1.56)		Ensure Plus
	(1.76)		Fortisip
Liquid (vanilla), 237 ml can - Higher subsidy of \$1.65 per			
1 can with Endorsement	0.85	1 OP	
	(1.65)		Ensure Plus
(Ensure Plus Liquid (vanilla), 237 ml can to be delisted 1 July 202	26)		
	•		

	Subsidy		Fully	Brand or
	(Manufacturer's Price)	Sub	sidised	Generic
	\$	Per	1	Manufacturer
ORAL FEED WITH FIBRE 1.5 KCAL/ML - Special Authority se	e SA1859 on page 293	3 – Hospit	tal pharn	nacy [HP3]
Additional subsidy by endorsement is available for patients	being bolus fed throug	h a feedir	ng tube,	or who have severe
epidermolysis bullosa. The prescription must be endorsed	accordingly.			
· ·				

Liquid (chocolate), 200 ml bottle - Higher subsidy of \$1.76 per			
1 btl with Endorsement	0.72	1 OP	
	(1.76)		Fortisip Multi Fibre
Liquid (strawberry), 200 ml bottle - Higher subsidy of \$1.76 per			
1 btl with Endorsement	0.72	1 OP	
	(1.76)		Fortisip Multi Fibre
Liquid (vanilla), 200 ml bottle - Higher subsidy of \$1.76 per			
1 btl with Endorsement	0.72	1 OP	
	(1.76)		Fortisip Multi Fibre

### **High Calorie Products**

### ⇒SA1195 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

All of the following:

- 1 Cystic fibrosis; and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements.

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 any condition causing malabsorption; or
  - 1.2 faltering growth in an infant/child; or
  - 1.3 increased nutritional requirements; or
  - 1.4 fluid restricted: and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements or is fluid restricted.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

	Subsidy (Manufacturer's Price) \$		ully Brand or sed Generic  Manufacturer
ENTERAL FEED 2 KCAL/ML - Special Authority see SA1195 o Liquid, 1,000 ml bottle			armacy [HP3]  ✓ Ensure Two Cal HN  RTH
Liquid, 500 ml bottle	6.82	1 OP	✓ Nutrison Concentrated
ORAL FEED 2 KCAL/ML – Special Authority see SA1195 on the Additional subsidy by endorsement is available for patients I epidermolysis bullosa. The prescription must be endorsed a Liquid (vanilla), 200 ml bottle – Higher subsidy of \$2.34 per 1 btl with Endorsement	peing bolus fed throug accordingly.		,

# **Food Thickeners**

### ⇒SA1106 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient has motor neurone disease with swallowing disorder.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FOOD THICKENER - Special Authority see SA1106 abov	<del>'e</del> – Hospital pharmacy	[HP3]	
Powder	8.29	300 g OP	✓ Nutilis
	24.00	380 g OP	✓ Aptamil Feed
		_	Thickener

# **Gluten Free Foods**

The funding of gluten free foods is no longer being actively managed by Pharmac from 1 April 2011. This means that we are no longer considering the listing of new products, or making subsidy, or other changes to the existing listings. As a result we anticipate that the range of funded items will reduce over time. Management of Coeliac disease with a gluten free diet is necessary for good outcomes. A range of gluten free options are available through retail outlets.

#### ⇒SA1729 Special Authority for Subsidy

Initial application — (all patients) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Either:

- 1 Gluten enteropathy has been diagnosed by biopsy; or
- 2 Patient suffers from dermatitis herpetiformis.

Initial application — (paediatric patients diagnosed by ESPGHAN criteria) only from a paediatric gastroenterologist. Approvals valid without further renewal unless notified where the paediatric patient fulfils ESPGHAN criteria for biopsy free diagnosis of coeliac disease.

GLUTEN FREE BAKING MIX – Special Authority see SA1	1729 above – Hospital	pharmacy [HP3]	
Powder	2.81	1,000 g OP	
	(5.15)	•	Healtheries Simple
			Baking Mix

	Subsidy (Manufacturer's P \$		ully Brand or Sed Generic  Manufacturer
GLUTEN FREE BREAD MIX - Special Authority see SA1729 or	n the previous pa	ge – Hospital pha	rmacy [HP3]
Powder	3.93	1,000 g OP	
	(7.32)		NZB Low Gluten
			Bread Mix
	3.51		
	(10.87)		Horleys Bread Mix
GLUTEN FREE FLOUR - Special Authority see SA1729 on the	previous page -	Hospital pharmac	cy [HP3]
Powder	5.62	2,000 g OP	
	(18.10)	-	Horleys Flour

# **Foods And Supplements For Inherited Metabolic Disease**

### ⇒SA2357 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid without further renewal unless notified where patient requires dietary management of inherited metabolic disorders.

# **Supplements For Homocystinuria**

AMINOACID FORMULA WITHOUT METHIONINE	<ul> <li>Special Authority see SA2357</li> </ul>	above – F	lospital pharmacy [HP3]
Powder (neutral), 36 g sachets	750.30	30	HCU Anamix Junior
Powder, 12.5 g sachets	349.65	30	✓ HCU Explore 5
Powder, 25 g sachets	1,048.95	30	✓ HCU Express 15
Powder (neutral), can		500 g OP	XMET Maxamum
Powder (unflavoured), can	260.00	400 g OP	HCU Anamix Infant
Liquid (juicy berries), 125 ml bottle	1,684.80	30	✓ HCU Lophlex LQ
Liquid (orange), 125 ml bottle	941.40	36	HCU Anamix Junior
			I O

# Supplements For MSUD and short chain enoyl coA hydratase deficiency

AMINOACID FORMULA WITHOUT VALINE, LEUCINE AND ISOLEUCINE - Special Authority see \$A2357 above - Hospital pharmacy [HP3]

Powder (neutral) 36 g sachets	750.00	30	<ul><li>MSUD Anamix Junior</li></ul>
Powder, 12.5 g sachets	349.65	30	✓ MSUD Explore 5
Powder, 25 g sachets		30	✓ MSUD Express 15
Powder (neutral), can		500 g OP	✓ MSUD Maxamum
Powder (orange), can	454.71	500 g OP	✓ MSUD Maxamum
Powder (unflavoured), can	260.00	400 g OP	<ul><li>MSUD Anamix Infant</li></ul>
Liquid (orange) 125 ml bottles	941.40	36	<ul><li>MSUD Anamix Junior LQ</li></ul>
Liquid (juicy berries) 125 ml pouches	1,684.80	30	✓ MSUD Lophlex LQ 20

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer

# **Supplements For PKU**

AMINOACID FORMULA WITHOUT PHENYLALANINE - Special Authority see SA2357 on the previous page - Hospital pharmacy [HP3]

annacy [in o]			
Tabs		75 OP	Phlexy 10
Powder (Lemon), 34 g sachets	883.50	30	✓ PKU Express 20
Powder (Neutral), 12.5 g sachets	220.88	30	✓ PKU Explore 5
Powder (Neutral), 34 g sachets	883.50	30	✓ PKU Express 20
Powder (Orange), 25 g sachets	441.75	30	✓ PKU Explore 10
Powder (Orange), 34 g sachets	883.50	30	✓ PKU Express 20
Powder (Raspberry), 25 g sachets		30	✓ PKU Explore 10
Powder (Tropical), 34 g sachets		30	✓ PKU Express 20
Powder (berry) 28 g sachets		30	✓ PKU Lophlex
· ,, •			Powder
Powder (chocolate) 36 g sachet	393.00	30	✓ PKU Anamix Junior
· onser (enessians) so g sasmermannin			Chocolate
Powder (neutral) 28 g sachets	936.00	30	✓ PKU Lophlex
1 owder (fleditar) 20 g oddrote		00	Powder
Powder (neutral) 36 g sachets	303.00	30	✓ PKU Anamix Junior
Powder (orange) 28 g sachets		30	✓ PKU Lophlex
rowder (ordrige) 20 g sacriets	930.00	30	Powder
Daviday (avanas) 66 a acabat	000.00	00	
Powder (orange) 36 g sachet	393.00	30	✓ PKU Anamix Junior
			Orange
Powder (unflavoured) 12.5 g sachets		30	✓ PKU First Spoon
Powder (vanilla) 36 g sachet	393.00	30	✓ PKU Anamix Junior
			Vanilla
Infant formula		400 g OP	PKU Anamix Infant
Powder (neutral), 4 × 400 g can	715.16	1,600 g OP	✓ Pku Start
Powder (orange)		500 g OP	XP Maxamum
Powder (unflavoured)		500 g OP	XP Maxamum
Liquid (berry), 125 ml bottle	13.10	1 OP	PKU Anamix Junior
			LQ
Liquid (orange), 125 ml bottle	13.10	1 OP	✓ PKU Anamix Junior
			LQ
Liquid (forest berries), 250 ml carton	540.00	18 OP	✓ Easiphen Liquid
Liquid (juicy tropical) 125 ml		30 OP	✓ PKU Lophlex LQ 20
Oral semi-solid (berries) 109 g		36 OP	✓ PKU Lophlex
, ,	,		Sensation 20
Liquid (juicy berries) 62.5 ml	939.00	60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy berries) 125 ml		30 OP	✓ PKU Lophlex LQ 20
Liquid (juicy orange) 125 ml		30 OP	✓ PKU Lophlex LQ 20
=-q (J)		00 0.	. No Lopinox La Lo

	Subsidy (Manufacturer's Price)		Fully Subsidised	I Generic
	\$	Per	•	Manufacturer
SLYCOMACROPEPTIDE AND AMINO ACID CONTAINS SON	ME PHENYLALANINE	– Spe	ecial Autho	ority see SA2357 on
age 299 – Hospital pharmacy [HP3]	000.00	00	,	BIZII
Powder (Banana) 35 g sachets	930.00	30	•	PKU
			_	sphere20 Banana
Powder (Berry), 20 g sachets	449.28	60	/	PKU Restore
				Powder
Powder (Chocolate) 32 g sachets	898.56	30	✓	PKU Build
				20 Chocolate
Powder (Chocolate) 35 g sachets	930.00	30	✓	PKU
				sphere20 Chocolate
Powder (Lemon) 35 g sachets	930.00	30	/	PKU
, ,				sphere20 Lemon
Powder (Lemonade) 33.4 g sachets	936.00	30	/	PKU GMPro Ultra
· ondo (25.10.100) oo · g odo.10.51.		•		Lemonade
Powder (Neutral), 15 g sachets	449.28	30	1	PKU Build 10
Powder (Orange), 20 g sachets		60		PKU Restore
Towaci (Grango), 20 g sacricis		00	•	Powder
Powder (Raspberry Lemonade) 31 g sachets	898.56	30	/	PKU Build
. on as: (asp20) _5 g sass.s.		•		20 Raspberry
				Lemonade
Powder (Smooth) 31 g sachets	808 56	30	1	PKU Build
Fowder (Smooth) ST & Sacriets		30	•	20 Smooth
Douglar (Vanilla) 22 a acabata	000 56	20	./	PKU Build 20 Vanilla
Powder (Vanilla) 33 g sachets		30 30		
Powder (neutral), 40 g sachets				Glytactin Bettermilk
Powder (unflavoured) 12.5 g sachets		30		PKU GMPro Mix-In
Powder (vanilla) 33.4 g sachets	936.00	30	•	PKU GMPro Ultra Vanilla
Powder (Red Berry) 35 g sachets	930 00	30	1	PKU sphere20 Red
1 Owder (Hed Delly) 55 g sacriets		30	•	Berry
Douglar (Vanilla) 25 a cochata	000.00	20	./	PKU
Powder (Vanilla) 35 g sachets	930.00	30	•	
				sphere20 Vanilla
Liquid (neutral), 250 ml carton		18		PKU GMPro LQ
Liquid (original), 250 ml carton	684.45	30 OF	•	PKU Glytactin RTD 15
Liquid (Coffee Mocha), 250 ml carton	601 15	30 OF		PKU Glytactin RTD
Liquid (Conee Mocha), 250 mi canton	004.40	JU UI	-	15 Lite
Limited (abancelate) OFO and contain	CO4 45	00 01		
Liquid (chocolate), 250 ml carton		30 OF	•	PKU Glytactin RTD 15
Liquid (vanilla), 250 ml carton	684.45	30 OF	•	PKU Glytactin RTD
		J J1	•	15 Lite

# Foods

LOW PROTEIN BAKING MIX - Special Authority see SA2357 on pag	e 299 –	Hospital pharmacy	(HP3]
Powder	8.55	500 g OP	✓ Loprofin Mix

	Subsidy		Fully Brand or
	(Manufacturer's F	Price) Subs Per	idised Generic  Manufacturer
LOW PROTEIN PASTA Special Authority see SA2257 on page	no 200 Hospital		
LOW PROTEIN PASTA – Special Authority see SA2357 on pag		500 g OP	oj <b>✓ Loprofin</b>
Animal shapesLasagne		250 g OP	✓ Loprofin
•		•	
Low protein rice pasta		500 g OP	✓ Loprofin
Macaroni		250 g OP	✓ Loprofin
Penne		500 g OP	✓ Loprofin
Spaghetti		500 g OP	✓ Loprofin
Spirals	12.39	500 g OP	✓ Loprofin
Supplements for Tyrosinaemia			
AMINOACID FORMULA WITHOUT PHENYLALANINE AND TY pharmacy [HP3]	ROSINE - Speci	ial Authority see	e SA2357 on page 299 – Hospi
Powder (Neutral), 12.5 g sachets	349.65	30	✓ TYR Explore 5
Powder (neutral) 36 g sachets		30	✓ TYR Anamix Junior
Powder, can		400 g OP	✓ TYR Anamix Infant
Liquid (juicy berries) 125 ml pouches		30	✓ TYR Lophlex LQ 20
Liquid (orange) 125 ml bottle		36	✓ TYR Anamix Junior
_ 4 ( 3-/			LQ
GLYCOMACROPEPTIDE AND AMINO ACID CONTAINS SOM	E TYROSINE AN	D PHENYLALA	NINE - Special Authority see
SA2357 on page 299 – Hospital pharmacy [HP3]			,
Powder (Red Berry), 35 g sachets	1.398.60	30	✓ TYR Sphere 20
Powder (Vanilla), 35 g sachets		30	✓ TYR Sphere 20
Supplements for Organic Acidaemias  AMINOACID FORMULA WITHOUT ISOLEUCINE, METHIONIN on page 299 – Hospital pharmacy [HP3]	E, THREONINE	AND VALINE -	- Special Authority see SA2357
	000.00	400 00	
Powder, can		400 g OP	✓ MMA/PA Anamix Infant
AMINOACID FORMULA WITHOUT METHIONINE, THREONINI Hospital pharmacy [HP3]	E AND VALINE -	Ū	Infant rity see SA2357 on page 299 -
AMINOACID FORMULA WITHOUT METHIONINE, THREONINI Hospital pharmacy [HP3] Powder (neutral), 18 g sachets	E AND VALINE -	- Special Autho	Infant rity see SA2357 on page 299 -  ✓ MMA/PA Anamix Junior
AMINOACID FORMULA WITHOUT METHIONINE, THREONINI Hospital pharmacy [HP3] Powder (neutral), 18 g sachets	E AND VALINE - 750.30	- Special Autho 30	Infant rity see SA2357 on page 299 -  ✓ MMA/PA Anamix Junior  ✓ MMA/PA Explore 5
AMINOACID FORMULA WITHOUT METHIONINE, THREONINI Hospital pharmacy [HP3] Powder (neutral), 18 g sachets	E AND VALINE - 750.30	- Special Autho	Infant rity see SA2357 on page 299 -  ✓ MMA/PA Anamix Junior
AMINOACID FORMULA WITHOUT METHIONINE, THREONINI Hospital pharmacy [HP3] Powder (neutral), 18 g sachets	E AND VALINE - 750.30	- Special Autho 30	Infant rity see SA2357 on page 299 -  ✓ MMA/PA Anamix Junior  ✓ MMA/PA Explore 5
AMINOACID FORMULA WITHOUT METHIONINE, THREONINI Hospital pharmacy [HP3] Powder (neutral), 18 g sachets	E AND VALINE750.30349.651,048.95	- Special Autho 30 30 30 30	Infant rity see SA2357 on page 299 -  ✓ MMA/PA Anamix Junior  ✓ MMA/PA Explore 5  ✓ MMA/PA Express 15
AMINOACID FORMULA WITHOUT METHIONINE, THREONING Hospital pharmacy [HP3] Powder (neutral), 18 g sachets  Powder, 12.5 g sachets Powder, 25 g sachets  Supplements for Glutaric Aciduria type 1  AMINOACID FORMULA WITHOUT LYSINE – Special Authority	E AND VALINE750.30349.651,048.95  y see SA2357 on	- Special Autho 30 30 30 30	Infant rity see SA2357 on page 299 -  MMA/PA Anamix Junior  MMA/PA Explore 5  MMA/PA Express 15
AMINOACID FORMULA WITHOUT METHIONINE, THREONING Hospital pharmacy [HP3] Powder (neutral), 18 g sachets Powder, 12.5 g sachets Powder, 25 g sachets  Supplements for Glutaric Aciduria type 1  AMINOACID FORMULA WITHOUT LYSINE – Special Authority Powder (neutral), 18 g sachets	E AND VALINE750.30349.651,048.95  y see SA2357 on750.30	- Special Autho 30 30 30 30 page 299 – Hos	Infant rity see SA2357 on page 299 -  ✓ MMA/PA Anamix Junior  ✓ MMA/PA Explore 5  ✓ MMA/PA Express 15  spital pharmacy [HP3]  ✓ GA1 Anamix Junior
AMINOACID FORMULA WITHOUT METHIONINE, THREONING Hospital pharmacy [HP3] Powder (neutral), 18 g sachets Powder, 12.5 g sachets Powder, 25 g sachets  Supplements for Glutaric Aciduria type 1  AMINOACID FORMULA WITHOUT LYSINE – Special Authority Powder (neutral), 18 g sachets	E AND VALINE	- Special Autho 30 30 30 30  page 299 – Hos	Infant rity see SA2357 on page 299 -  ✓ MMA/PA Anamix Junior  ✓ MMA/PA Explore 5  ✓ MMA/PA Express 15  spital pharmacy [HP3]  ✓ GA1 Anamix Junior  ✓ GA Explore 5
AMINOACID FORMULA WITHOUT METHIONINE, THREONING Hospital pharmacy [HP3] Powder (neutral), 18 g sachets Powder, 12.5 g sachets Powder, 25 g sachets  Supplements for Glutaric Aciduria type 1  AMINOACID FORMULA WITHOUT LYSINE – Special Authority Powder (neutral), 18 g sachets	E AND VALINE	- Special Autho 30 30 30 30 page 299 – Hos	Infant rity see SA2357 on page 299 -  ✓ MMA/PA Anamix Junior  ✓ MMA/PA Explore 5  ✓ MMA/PA Express 15  spital pharmacy [HP3]  ✓ GA1 Anamix Junior
AMINOACID FORMULA WITHOUT METHIONINE, THREONING Hospital pharmacy [HP3] Powder (neutral), 18 g sachets	E AND VALINE	- Special Autho 30 30 30 30 page 299 – Hos 30 30 400 g OP	Infant rity see SA2357 on page 299 -  MMA/PA Anamix Junior  MMA/PA Explore 5  MMA/PA Express 15  Spital pharmacy [HP3]  GA1 Anamix Junior  GA Explore 5  GA1 Anamix Infant
AMINOACID FORMULA WITHOUT METHIONINE, THREONING Hospital pharmacy [HP3] Powder (neutral), 18 g sachets	E AND VALINE750.30349.651,048.95  y see SA2357 on750.30349.65260.00	- Special Autho 30 30 30 30 page 299 – Hos 30 30 400 g OP	Infant rity see SA2357 on page 299 -  MMA/PA Anamix Junior  MMA/PA Explore 5  MMA/PA Express 15  Spital pharmacy [HP3]  GA1 Anamix Junior  GA Explore 5  GA1 Anamix Infant
AMINOACID FORMULA WITHOUT METHIONINE, THREONINI Hospital pharmacy [HP3] Powder (neutral), 18 g sachets	E AND VALINE750.30349.651,048.95  y see SA2357 on750.30349.65260.00	- Special Autho 30 30 30 30  page 299 – Hos 30 400 g OP	Infant rity see SA2357 on page 299 -  MMA/PA Anamix Junior  MMA/PA Explore 5  MMA/PA Express 15  Spital pharmacy [HP3]  GA1 Anamix Junior  GA Explore 5  GA1 Anamix Infant  I pharmacy [HP3]
AMINOACID FORMULA WITHOUT METHIONINE, THREONINI Hospital pharmacy [HP3] Powder (neutral), 18 g sachets	E AND VALINE	- Special Autho 30 30 30 30 30  page 299 – Hospita 30 400 g OP	Infant rity see SA2357 on page 299 -  MMA/PA Anamix Junior  MMA/PA Explore 5  MMA/PA Express 15  Spital pharmacy [HP3]  GA1 Anamix Junior  GA Explore 5  GA1 Anamix Infant  I pharmacy [HP3]

Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
CITRULLINE – Special Authority see SA2357 on page 299 – Hospital pharmacy [HP3] Powder, 4 g sachets211.45	30	1	Citrulline1000
ISOLEUCINE – Special Authority see SA2357 on page 299 – Hospital pharmacy [HP3 Powder, 4 g sachets141.05	] 30	1	Isoleucine50
LEUCINE – Special Authority see SA2357 on page 299 – Hospital pharmacy [HP3] Powder, 4 g sachets141.05	30	✓	Leucine100
PHENYLALANINE – Special Authority see SA2357 on page 299 – Hospital pharmacy Powder, 4 g sachets141.05	[HP3] 30	•	Phenylalanine50
TYROSINE – Special Authority see SA2357 on page 299 – Hospital pharmacy [HP3] Powder, 4 g sachets211.45	30	✓	Tyrosine1000
VALINE – Special Authority see SA2357 on page 299 – Hospital pharmacy [HP3] Powder, 4 g sachets141.05	30	•	Valine50
Other Fat Modified Products			
ELEMENTAL FEED WITH HIGH MEDIUM CHAIN TRIGLYCERIDES — Special Author pharmacy [HP3]	ity se	e SA2357	7 on page 299 – Hospital
Powder (neutral), 100 g sachets47.01	10	•	Emsogen
Carbohydrate and Fat with added vitamins and minerals			
PROTEIN FREE SUPPLEMENT CONTAINING CARBOHYDRATE, FAT WITH ADDED Authority see SA2357 on page 299 – Hospital pharmacy [HP3]	VIT/	AMINS AN	ND MINERALS - Special
, , , , , , , , , , , , , , , , , , , ,	0 g O	P 🗸	Energivit
Essential Amino Acids			
ESSENTIAL AMINOACID FORMULA - Special Authority see SA2357 on page 299 - Fowder (neutral), can	Hospi 0 g O		acy [HP3] Essential Amino Acid Mix

#### Infant Formulae

### For Williams Syndrome

### **⇒SA1110** Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is an infant suffering from Williams Syndrome and associated hypercalcaemia.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

LOW CALCIUM INFANT FORMULA - Special Authority see SA1110 above - Hospital pharmacy [HP3]

Powder .......46.18 400 g OP ✓ Locasol

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

### **Gastrointestinal and Other Malabsorptive Problems**

AMINO ACID FORMULA - Special Authority see SA209	<mark>2 below – Hospital pharm</mark>	nacy [HP3]	
Powder	43.60	400 g OP	<ul><li>✓ Alfamino</li><li>✓ Alfamino Junior</li></ul>
Powder (unflavoured)	55.61	400 g OP	✓ Neocate Gold ✓ Neocate Junior Unflavoured
			✓ Neocate SYNEO
	65.72		✓ Elecare
			✓ Elecare LCP
Powder (vanilla)	55.61	400 g OP	✓ Neocate Junior Vanilla
	65.72		✓ Elecare

### ⇒SA2092 Special Authority for Subsidy

**Initial application** — (Infants under 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 History of anaphylaxis to cow's milk protein formula or dairy products; or
- 2 Eosinophilic oesophagitis; or
- 3 Ultra-short gut; or
- 4 Severe Immune deficiency; or
- 5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
- 6 Both
  - 6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
  - 6.2 Either:
    - 6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or 6.2.2 Patient has IgE mediated allergy.

Initial application — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or
  - 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
  - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products; or
  - 2.2 Eosinophilic oesophagitis; or
  - 2.3 Ultra-short gut; or
  - 2.4 Severe Immune deficiency: or
  - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
  - 2.6 Both:
    - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
    - 2.6.2 Either:
      - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval

continued...

Subsidy	Fully	/ Brand or	
(Manufacturer's Price)	Subsidised	d Generic	
\$	Per 🗸	Manufacturer	

continued...

number; or

2.6.2.2 Patient has IgE mediated allergy.

Renewal — (Infants up to 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 Patient has IgE mediated allergy; and
  - 1.2 All of the following:
    - 1.2.1 Patient remains allergic to cow's milk; and
    - 1.2.2 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy or extensively hydrolysed infant formula has been undertaken; and
    - 1.2.3 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
    - 1.2.4 Amino acid formula is required for a nutritional deficit; and
    - 1.2.5 It has been more than three months from the previous approval; or

#### 2 Both:

- 2.1 Patient has non IgE mediated severe gastrointestinal intolerance (including eosinophilic oesophagitis, ultra-short gut and severe immune deficiency); and
- 2.2 All of the following:
  - 2.2.1 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy, or extensively hydrolysed infant formula has been undertaken; and
  - 2.2.2 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
  - 2.2.3 Amino acid formula is required for a nutritional deficit; and
  - 2.2.4 It has been more than three months from the previous approval.

Renewal — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or
  - 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
  - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products: or
  - 2.2 Eosinophilic oesophagitis; or
  - 2.3 Ultra-short gut; or
  - 2.4 Severe Immune deficiency; or
  - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
  - 2.6 Both:
    - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
    - 2.6.2 Either:
      - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or
      - 2.6.2.2 Patient has IgE mediated allergy.

Initial application — (for patients who have a current funding under Special Authority form SA1557) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 months for applications meeting the following criteria:

continued...

Subsidy	Fu	lly Brand or	
(Manufacturer's Price)	Subsidis	ed Generic	
\$	Per	<ul> <li>Manufacturer</li> </ul>	

continued...

All of the following:

- 1 Patient has a valid Special Authority approval for extensively hydrolysed formula (SA1557); and
- 2 Extensively hydrolysed formula (Aptamil Gold+ Pepti Junior, AllerPro SYNEO 1 and 2) is unable to be supplied at this time; and
- 3 The approval only applies to funded dispensings of Neocate Gold and Neocate Syneo.

Note: This criteria is short term funding to cover an out-of-stock situation on some extensively hydrolysed formula powder funded under Special Authority form SA1557. There is no renewal criteria under this restriction.

ENTERAL LIQUID PEPTIDE FORMULA - Special Authority see SA1953 below - Hospital pharmacy [HP3]

Liquid 1 kcal/ml, 500 ml bottle12.44	1 OP	✓ Nutrini Peptisorb
Liquid 1.5 kcal/ml, 500 ml bottle	1 OP	✓ Nutrini Peptisorb
		Energy

### ⇒SA1953 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has impaired gastrointestinal function and either cannot tolerate polymeric feeds, or polymeric feeds are unsuitable; and
- 2 Any of the following:
  - 2.1 Severe malabsorption; or
  - 2.2 Short bowel syndrome: or
  - 2.3 Intractable diarrhoea; or
  - 2.4 Biliary atresia; or
  - 2.5 Cholestatic liver diseases causing malabsorption; or
  - 2.6 Cystic fibrosis; or
  - 2.7 Proven fat malabsorption; or
  - 2.8 Severe intestinal motility disorders causing significant malabsorption; or
  - 2.9 Intestinal failure: or
  - 2.10 Both:
    - 2.10.1 The patient is currently receiving funded amino acid formula; and
    - 2.10.2 The patient is to be trialled on, or transitioned to, an enteral liquid peptide formula; and
- 3 Either:
  - 3.1 A semi-elemental or partially hydrolysed powdered feed has been reasonably trialled and considered unsuitable; or
  - 3.2 For step down from intravenous nutrition.

Note: A reasonable trial is defined as a 2-4 week trial.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 An assessment as to whether the patient can be transitioned to a cows milk protein or soy infant formula or extensively hydrolysed formula has been undertaken; and
- 2 The outcome of the assessment is that the patient continues to require an enteral liquid peptide formula; and
- 3 General practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

EXTENSIVELY HYDROLYSED FORMULA  – Special A	Authority see SA1557 on the	e next page –	Hospital pharmacy [HP3]
Powder	18.10	450 g OP	✓ Pepti-Junior
	36.20	900 g OP	✓ Allerpro Syneo 1
		-	✓ Allerpro Syneo 2

Subsidy (Manufacturer's Price) \$

Fully Subsidised Per

Brand or Generic Manufacturer

### ⇒SA1557 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 Both:
  - 1.1 Cows milk formula is inappropriate due to severe intolerance or allergy to its protein content; and
  - 1.2 Either
    - 1.2.1 Soy milk formula has been reasonably trialled without resolution of symptoms; or
    - 1.2.2 Soy milk formula is considered clinically inappropriate or contraindicated; or
- 2 Severe malabsorption; or
- 3 Short bowel syndrome; or
- 4 Intractable diarrhoea; or
- 5 Biliary atresia; or
- 6 Cholestatic liver diseases causing malsorption; or
- 7 Cystic fibrosis: or
- 8 Proven fat malabsorption; or
- 9 Severe intestinal motility disorders causing significant malabsorption; or
- 10 Intestinal failure; or
- 11 All of the following:
  - 11.1 For step down from Amino Acid Formula: and
  - 11.2 The infant is currently receiving funded amino acid formula; and
  - 11.3 The infant is to be trialled on, or transitioned to, an extensively hydrolysed formula; and
  - 11.4 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

Note: A reasonable trial is defined as a 2-4 week trial, or signs of an immediate IgE mediated allergic reaction.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 An assessment as to whether the infant can be transitioned to a cows milk protein or soy infant formula has been undertaken; and
- 2 The outcome of the assessment is that the infant continues to require an extensively hydrolysed infant formula; and
- 3 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

#### Fluid Restricted

PAEDIATRIC ORAL/ENTERAL FEED 1 KCAL/ML - Special Authority see SA1698 below - Hospital pharmacy [HP3] Liquid, 125 ml bottle .......2.80 1 OP ✓ Infatrini

#### ⇒SA1698 Special Authority for Subsidy

**Initial application** only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is fluid restricted or volume intolerant and has been diagnosed with faltering growth; and
- 2 Patient is under the care of a paediatrician or dietitian who has recommended treatment with a high energy infant formula; and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

continued...



Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
\$	Per	✓	

continued...

Note: "Volume intolerant" patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

Renewal only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient continues to be fluid restricted or volume intolerant and has faltering growth; and
- 2 Patient is under the care of a hospital paediatrician or dietitian who has recommended treatment with a high energy infant formula; and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: 'Volume intolerant' patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

# **Ketogenic Diet**

### ⇒SA1197 Special Authority for Subsidy

**Initial application** only from a metabolic physician or paediatric neurologist. Approvals valid for 3 months where the patient has intractable epilepsy, pyruvate dehydrogenase deficiency or glucose transported type-1 deficiency and other conditions requiring a ketogenic diet.

**Renewal** only from a metabolic physician or paediatric neurologist. Approvals valid for 2 years where the patient is on a ketogenic diet and the patient is benefiting from the diet.

HIGH FAT LOW CARBOHYDRATE FORMULA - Special Authority see SA119	7 above – Hospita	al pharmacy [HP3]
Powder (unflavoured)36.92	300 g OP	✓ KetoCal 4:1
	_	✓ Ketocal 3:1
Powder (vanilla)36.92	300 g OP	✓ KetoCal 4:1

# **SECTION I: NATIONAL IMMUNISATION SCHEDULE**

Subsidy (Manufacturer's Price) Per

Subsidised

Fully

Brand or Generic Manufacturer

# **Vaccinations**

BACILLUS CALMETTE-GUERIN VACCINE - [Xpharm]

For infants at increased risk of tuberculosis. Increased risk is defined as:

- 1) living in a house or family with a person with current or past history of TB; or
- 2) having one or more household members or carers who within the last 5 years lived in a country with a rate of TB > or egual to 40 per 100,000 for 6 months or longer; or
- 3) during their first 5 years will be living 3 months or longer in a country with a rate of TB > or equal to 40 per 100,000 Note a list of countries with high rates of TB are available at www.health.govt.nz/tuberculosis (search for downloads) or www.bcgatlas.org/index.php.
- Inj Mycobacterium bovis BCG (Bacillus Calmette-Guerin),

Danish strain 1331, live attenuated, vial with diluent.................0.00

10

✓ BCG Vaccine AJV

	Cubaidu		Fully	Brand or
	Subsidy (Manufacturer's Price)		Subsidised	Generic
	\$	Per	<b>✓</b>	Manufacturer
VID-19 VACCINE - [Xpharm]				
Inj 3 mcg SARS-CoV-2 spike protein (mRNA) LP.8.1 per 0.3	ml.			
0.48 ml multi-dose vial; infant vaccine, yellow cap		10	✓ (	Comirnaty (LP.8.1)
Up to three doses for previously unvaccinated children a				
Inj 3 mcg bretovameran per 0.3 ml, 0.48 ml vial; infant vaccir	ne.			
yellow cap		10	✓ (	Comirnaty Omicron
,				(JN.1)
Up to three doses for previously unvaccinated children a	aged 6 months - 4 year	rs at h	igh risk of	
	,		•	
Inj 10 mcg SARS-CoV-2 spike protein (mRNA) LP.8.1 per				
0.3 ml, 0.48 ml single-dose vial; paediatric vaccine, light				
blue cap		10	✓ (	Comirnaty (LP.8.1)
Either:				, , ,
1) One dose for previously unvaccinated children age	ed 5-11 vears old: or			
2) Up to three doses for immunocompromised childre		l.		
, ,	,			
Inj 10 mcg bretovameran per 0.3 ml, 0.48 ml vial; paediatric				
vaccine, light blue cap	0.00	10	✓ (	Comirnaty Omicron
, 0				(JN.1)
Either:				` '
1) One dose for previously unvaccinated children age	ed 5-11 years old; or			
2) Up to three doses for immunocompromised childre		l.		
, .,	,			
Inj 30 mcg SARS-CoV-2 spike protein (mRNA) LP.8.1 per				
0.3 ml, pre-filled syringe; adult dose	0.00	10	✓ (	Comirnaty (LP.8.1)
Any of the following:		. •	•	· · · · · · · · · · · · · · · · · · ·
One dose for previously unvaccinated people aged	d 12-15 years old: or			
Up to three doses for immunocompromised people		l· or		
Up to two doses for previously unvaccinated people	,	, 0.		
Up to four doses for people aged 16-29 at high risk				
5) One dose for previously unvaccinated people aged				
6) One additional dose every 6 months for previously		ed 30	vears and	over – additional dose
-, additional account of the proviously			, 50.0 0110	
given at least 6 months after last dose				
given at least 6 months after last dose.				
·	ina			
given at least 6 months after last dose.  Inj 30 mcg bretovameran per 0.3 ml, 0.48 ml vial; adult vacci light grey cap	•	10	<b>.</b>	Comirnaty Omicron

Subsidy (Manufacturer's Price) \$	Sub Per	Fully sidised	Brand or Generic Manufacturer	

Any of the following:

- 1) One dose for previously unvaccinated people aged 12-15 years old; or
- 2) Up to three doses for immunocompromised people aged 12-15 years old; or
- 3) Up to two doses for previously unvaccinated people 16-29 years old; or
- 4) Up to four doses for people aged 16-29 at high risk of severe illness; or
- 5) One dose for previously unvaccinated people aged 30 and older; or
- 6) One additional dose every 6 months for previously vaccinated people aged 30 years and over additional dose is given at least 6 months after last dose.

(Comirnaty Omicron (JN.1) Inj 3 mcg bretovameran per 0.3 ml, 0.48 ml vial; infant vaccine, yellow cap to be delisted 1 June 2026) (Comirnaty Omicron (JN.1) Inj 10 mcg bretovameran per 0.3 ml, 0.48 ml vial; paediatric vaccine, light blue cap to be delisted 1 June 2026)

(Comirnaty Omicron (JN.1) Inj 30 mcg bretovameran per 0.3 ml, 0.48 ml vial; adult vaccine, light grey cap to be delisted 1 June 2026)

#### DIPHTHERIA. TETANUS AND PERTUSSIS VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- c)
- A) Funded for any of the following criteria:
  - 1) A single dose for pregnant women in the second or third trimester of each pregnancy; or
  - 2) A single dose for parents or primary caregivers of infants admitted to a Neonatal Intensive Care Unit or Specialist Care Baby Unit for more than 3 days, who had not been exposed to maternal vaccination at least 14 days prior to birth; or
  - A course of up to four doses is funded for children from age 7 up to the age of 18 years inclusive to complete full primary immunisation; or
  - 4) An additional four doses (as appropriate) are funded for (re-)immunisation for patients post haematopoietic stem cell transplantation or chemotherapy; pre or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or
  - 5) A single dose for vaccination of patients aged from 65 years old; or
  - 6) A single dose for vaccination of patients aged from 45 years old who have not had 4 previous tetanus doses; or
  - 7) For vaccination of previously unimmunised or partially immunised patients; or
  - 8) For revaccination following immunosuppression; or
  - 9) For boosting of patients with tetanus-prone wounds.

Notes: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.

- B) Contractors will be entitled to claim payment from the Funder for the supply of diphtheria, tetanus and pertussis vaccine to patients eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the diphtheria, tetanus and pertussis vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraphs 1 – 9 above.

Inj 2 IU diphtheria toxoid with 20 IU tetanus toxoid,	8 r	ncg
pertussis toxoid. 8 mcg pertussis filamentous		

haemagglutinin and 2.5 mcg pertactin in 0.5 ml prefilled

Subsidy		ully	Brand or	
(Manufacturer's Price)	Subsid	ised	Generic	
\$	Per	✓	Manufacturer	

#### DIPHTHERIA, TETANUS, PERTUSSIS AND POLIO VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- c)
- A) Funded for any of the following:
  - 1) A single dose for children up to the age of 7 who have completed primary immunisation; or
  - 2) A course of four vaccines is funded for catch up programmes for children (to the age of 10 years) to complete full primary immunisation: or
  - 3) An additional four doses (as appropriate) are funded for (re-)immunisation for people post HSCT, or chemotherapy; pre- or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or
  - 4) Five doses will be funded for children requiring solid organ transplantation.
- B) Contractors will be entitled to claim payment from the Funder for the supply of Diphtheria, tetanus, pertussis and polio vaccine to people eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Diphtheria, tetanus, pertussis and polio vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.

Inj 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg

pertussis toxoid, 25 mcg pertussis filamentous haemagglutinin, 8 mcg pertactin and 80 D-antigen units

10

✓ Infanrix IPV

#### DIPHTHERIA, TETANUS, PERTUSSIS, POLIO, HEPATITIS B AND HAEMOPHILUS INFLUENZAE TYPE B VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- c)
- A) Funded for children meeting any of the following criteria
  - 1) Up to four doses for children under the age of 10 years for primary immunisation; or
  - 2) An additional four doses (as appropriate) for (re-)immunisation of children under the age of 18 years post haematopoietic stem cell transplantation; or
  - 3) An additional four doses (as appropriate) for (re-)immunisation of children under the age of 10 years who are post chemotherapy; pre or post splenectomy; undergoing renal dialysis and other severely immunosuppressive regimens: or
  - 4) Up to five doses for children under the age of 10 years receiving solid organ transplantation.
- B) Contractors will be entitled to claim payment from the Funder for the supply of Diphtheria, tetanus, pertussis, polio. hepatitis B and haemophilus influenzae type b vaccine to people eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Diphtheria, tetanus, pertussis, polio, hepatitis B and haemophilus influenzae type b vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Note: A course of up-to four vaccines is funded for catch up programmes for children (up to and under the age of 10 years) to complete full primary immunisation. Please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes.

Inj 30IU diphtheria with 40IU tetanus and 25mcg pertussis toxoids, 25mcg pertussis filamentous haemagglutinin, 8mcg pertactin, 80D-AgU polio virus, 10mcg hepatitis B antigen, 10mcg H. influenzae type b with tetanus toxoid 

✓ Infanrix-hexa

10

1

✓ Havrix 1440

✓ Havrix Junior

	NATIONAL	IMMUNISAT	ION SCHEDULE
	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per 🗸	Brand or Generic Manufacturer
HAEMOPHILUS INFLUENZAE TYPE B VACCINE			
a) Only on a prescription			
<ul><li>b) No patient co-payment payable</li><li>c)</li></ul>			
A) One dose for people meeting any of the following:     1) For primary vaccination in children; or     2) An additional dose (as appropriate) is funded transplantation, or chemotherapy; functional at transplant, pre or post cochlear implants, rena 3) For use in testing for primary immunodeficient physician or paediatrician.      B) Contractors will be entitled to claim payment from the vaccine to people eligible under the above criterials for subsidised immunisation, and they may only do in the Pharmaceutical Schedule.  C) Contractors may only claim for populations within the sub-set of the population described in paragraph A Inj 10 mcg vial with diluent syringe	asplenic; pre or post spal dialysis and other sectory diseases, on the record disease contract of the Hame criteria that are coverabove.	olenectomy; pre- everely immunos commendation of oly of Haemophil act with Health N aemophilus influ ered by their cor	or post solid organ uppressive regimens; or of an internal medicine us influenzae type b lew Zealand (Health NZ) enzae type b vaccine listed
HEPATITIS A VACCINE – [Xpharm] Funded for patients meeting any of the following criteria:  1) Two vaccinations for use in transplant patients; or 2) Two vaccinations for use in children with chronic liver of 3) One dose of vaccine for close contacts of known hepati			

Inj 720 ELISA units in 0.5 ml syringe.......0.00

	Subsidy (Manufacturer's Price)		Fully sidised	Brand or Generic	
	\$	Per	1	Manufacturer	
HEPATITIS B RECOMBINANT VACCINE - [Xpharm]					

✓ Engerix-B

Funded for patients meeting any of the following criteria:

- 1) for household or sexual contacts of known acute hepatitis B patients or hepatitis B carriers; or
- 2) for children born to mothers who are hepatitis B surface antigen (HBsAq) positive; or
- 3) for children up to and under the age of 18 years inclusive who are considered not to have achieved a positive serology and require additional vaccination or require a primary course of vaccination; or
- 4) for HIV positive patients: or
- 5) for hepatitis C positive patients; or
- 6) for patients following non-consensual sexual intercourse; or
- 7) for patients prior to planned immunosuppression for greater than 28 days; or
- 8) for patients following immunosuppression; or
- 9) for solid organ transplant patients; or
- 10) for post-haematopoietic stem cell transplant (HSCT) patients; or
- 11) following needle stick injury.

Inj 20 mcg per 1 ml prefilled syringe......0.00 ✓ Engerix-B

Funded for patients meeting any of the following criteria:

- 1) for household or sexual contacts of known acute hepatitis B patients or hepatitis B carriers; or
- 2) for children born to mothers who are hepatitis B surface antigen (HBsAg) positive; or
- 3) for children up to and under the age of 18 years inclusive who are considered not to have achieved a positive serology and require additional vaccination or require a primary course of vaccination; or
- 4) for HIV positive patients; or
- 5) for hepatitis C positive patients; or
- 6) for patients following non-consensual sexual intercourse; or
- 7) for patients prior to planned immunosuppression for greater than 28 days; or
- 8) for patients following immunosuppression; or
- 9) for solid organ transplant patients; or
- 10) for post-haematopoietic stem cell transplant (HSCT) patients; or
- 11) following needle stick injury; or
- 12) for dialysis patients; or
- 13) for liver or kidney transplant patients.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

HUMAN PAPILLOMAVIRUS (6, 11, 16, 18, 31, 33, 45, 52 AND 58) VACCINE [HPV]

- a) Maximum of 1 inj per prescription
- b) Only on a prescription
- c) No patient co-payment payable
- d
- a) A) Any of the following:
  - 1) Maximum of two doses for children aged 14 years and under; or
  - 2) Maximum of three doses for people meeting any of the following criteria:
    - 1) People aged 15 to 26 years inclusive; or
    - 2) Either:

People aged 9 to 26 years inclusive who have

- 1) Confirmed HIV infection; or
- 2) Received a transplant (including stem cell): or
- 3) Maximum of four doses for people aged 9 to 26 years inclusive post chemotherapy
- B) Contractors will be entitled to claim payment from the Funder for the supply of Human papillomavirus vaccine to people eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Human papillomavirus vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraphs A above.

Inj 270 mcg in 0.5 ml syringe	0.00	10	Gardasil 9
-------------------------------	------	----	------------

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
INFLUENZA VACCINE Inj 60 mcg in 0.5 ml syringe (quadrivalent vaccine)	120.00	10		nfluvac Tetra (2025 formulation)

Subsidy	Fully Subsidised			
(Manufacturer's Price)				
\$	Per	1	Manufacturer	

- a) Maximum of 1 inj per prescription
- b) Only on a prescription
- c) No patient co-payment payable
- d

### A) INFLUENZA VACCINE

is available each year for patients who meet the following criteria, as set by Pharmac:

- a) all people 65 years of age and over; or
- b) people under 65 years of age who:
  - i) have any of the following cardiovascular diseases:
    - a) ischaemic heart disease, or
    - b) congestive heart failure, or
    - c) rheumatic heart disease, or
    - d) congenital heart disease, or
    - e) cerebo-vascular disease; or
  - ii) have either of the following chronic respiratory diseases:
    - a) asthma, if on a regular preventative therapy, or
    - b) other chronic respiratory disease with impaired lung function; or
  - iii) have diabetes; or
  - iv) have chronic renal disease; or
  - v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
  - vi) have any of the following other conditions:
    - a) autoimmune disease, or
    - b) immune suppression or immune deficiency, or
    - c) HIV, or
    - d) transplant recipients, or
    - e) neuromuscular and CNS diseases/disorders, or
    - f) haemoglobinopathies, or
    - g) are children on long term aspirin, or
    - h) have a cochlear implant, or
    - i) errors of metabolism at risk of major metabolic decompensation, or
    - i) pre and post splenectomy, or
    - k) Down syndrome, or
  - vii) are pregnant; or
- c) children 4 years of age and under who have been hospitalised for respiratory illness or have a history of significant respiratory illness; or
- d) people under 65 years of age who:
  - i) have any of the following serious mental health conditions:
    - a) schizophrenia, or
    - b) major depressive disorder, or
    - c) bipolar disorder, or
    - d) schizoaffective disorder, or
  - ii) are currently accessing secondary or tertiary mental health and addiction services; or

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy,
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Contractors will be entitled to claim payment for the supply of influenza vaccine to patients eligible under the above criteria pursuant to their contract with Health NZ for subsidised immunisation, and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Subsidy		Fully	Brand or	
(Manufacturer's Price	e)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

### MEASLES. MUMPS AND RUBELLA VACCINE

- a) Only on a prescription
- b) No patient co-payment payable

c)

### A) Measles, mumps and rubella vaccine

A maximum of two doses for any patient meeting the following criteria:

- 1) For primary vaccination in children; or
- 2) For revaccination following immunosuppression; or
- 3) For any individual susceptible to measles, mumps or rubella; or
- 4) A maximum of three doses for children who have had their first dose prior to 12 months.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes. Although a price is listed for the vaccine, doctors can still order measles mumps and rubella vaccine free of charge, as with other Schedule vaccines.

- B) Contractors will be entitled to claim payment for the supply of measles, mumps and rubella vaccine to patients eligible under the above criteria pursuant to their contract with Health NZ for subsidised immunisation, and they may only do so in respect of the measles, mumps and rubella vaccine listed in the Pharmaceutical Schedule.
- C) Contractors can only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Inj, measles virus 1,000 CCID50, mumps virus 5,012 CCID50,		
Rubella virus 1,000 CCID50; prefilled syringe/ampoule of		
diluent 0.5 ml	10	✓ Priorix

Subsidy	S	Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

#### MENINGOCOCCAL (GROUPS A. C. Y AND W-135) CONJUGATE VACCINE

Ini 10 mcg of each meningococcal polysaccharide conjugated to a total of approximately 55 mcg of tetanus toxoid carrier 

✓ MenQuadfi

- a) Only on a prescription
- b) No patient co-payment payable

- A) Any of the following:
  - 1) Up to three doses and a booster every five years for patients pre- and post splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre or post solid organ transplant; or
  - 2) One dose for close contacts of meningococcal cases of any group; or
  - 3) One dose for person who has previously had meningococcal disease of any group; or
  - 4) A maximum of two doses for bone marrow transplant patients; or
  - 5) A maximum of two doses for person pre- and post-immunosuppression\*: or
- B) Both:
  - 1) Person is aged between 13 and 25 years, inclusive; and
  - 2) Either:
    - 1) One dose for individuals who are entering within the next three months, or in their first year of living in boarding school hostels, tertiary education halls of residence, military barracks, Youth Justice residences, or prisons; or
    - 2) One dose for individuals who turn 13 years of age while living in boarding school hostels.
- C) Contractors will be entitled to claim payment from the Funder for the supply of Meningococcal A. C. Y and W-135 vaccine to patients eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Meningococcal A, C, Y and W-135 vaccine listed in the Pharmaceutical Schedule.
- D) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraphs A-B above.

Note: children under seven years of age require two doses 8 weeks apart, a booster dose three years after the primary series and then five yearly.

\*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than

Ini 5 mcg of each meningococcal polysaccharide conjugated to a total of approximately 44 mcg of tetanus toxoid carrier

✓ Nimenrix

- A) Both:
  - 1) The child is under 12 months of age; and
  - 2) Any of the following:
    - 1) A maximum of three doses (dependant on age at first dose) for patients pre- and post- splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre- or post- solid organ transplant; or
    - 2) A maximum of three doses (dependant on age at first dose) for close contacts of meningococcal cases
    - 3) A maximum of three doses (dependant on age at first dose) for child who has previously had meningococcal disease of any group; or
    - 4) A maximum of three doses (dependant on age at first dose) for bone marrow transplant patients; or
    - 5) A maximum of three doses (dependant on age at first dose) for child pre- and post-immunosuppression\*.

Note: infants from 6 weeks to less than 6 months of age require a 2+1 schedule, infants from 6 months to less than 12 months of age require a 1+1 schedule. Refer to the Immunisation Handbook for recommended booster schedules with meningococcal ACWY vaccine.

\*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days.

Subsidy (Manufacturer's Price)	Full Subsidise	
 \$	Per 🗸	Manufacturer

#### MENINGOCOCCAL B MULTICOMPONENT VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- c) Any of the following:
  - A) A primary course of up to three doses for children up to the age of 59 months inclusive; or
  - B) Both:
    - 1) Person is 5 years of age or over; and
    - 2) Any of the following:
      - i) up to two doses and a booster every five years for patients pre- and post-splenectomy; or
      - ii) up to two doses and a booster every five years for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited); or
      - iii) up to two doses and a booster every five years pre- or post-solid organ transplant; or
      - iv) up to two doses for close contacts of meningococcal cases of any group; or
      - v) up to two doses for person who has previously had meningococcal disease of any group; or
      - vi) up to two doses for bone marrow transplant patients; or
      - vii) up to two doses for person pre- and post-immunosuppression\*; or
  - C) Both:
    - 1) Person is aged between 13 and 25 years (inclusive); and
    - 2) Either:
      - Two doses for individuals who are entering within the next three months, or in their first year of living in boarding school hostels, tertiary education halls of residence, military barracks, Youth Justice residences or prison; or
      - ii) Two doses for individuals who turn 13 years of age while living in boarding school hostels.
  - D) Contractors will be entitled to claim payment from the Funder for the supply of Meningococcal B multicomponent vaccine to patients eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Meningococcal B multicomponent vaccine listed in the Pharmaceutical Schedule.
  - E) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraphs A-C above.

\*Immunosuppression due to corticosteroid or other immunosuppressive therapy must be for a period of greater than 28 days.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsidised		Generic	
\$	Per	1	Manufacturer	

#### PNEUMOCOCCAL (PCV13) CONJUGATE VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- c)
- A) Any of the following:
  - 1) A course of three doses for previously unvaccinated children up to the age of 59 months inclusive; or
  - Two doses are funded for high risk individuals (over the age of 12 months and under 18 years) who have previously received two doses of the primary course of PCV10: or
  - 3) Up to an additional four doses (as appropriate) are funded for the (re)immunisation of high risk children aged under 5 years with any of the following:
    - a) on immunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient immune response; or
    - b) primary immune deficiencies; or
    - c) HIV infection: or
    - d) renal failure, or nephrotic syndrome; or
    - e) who are immune-suppressed following organ transplantation (including haematopoietic stem cell transplant): or
    - f) cochlear implants or intracranial shunts; or
    - g) cerebrospinal fluid leaks; or
    - h) receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage of 20 mg or greater; or
    - i) chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or
    - j) pre term infants, born before 28 weeks gestation; or
    - k) cardiac disease, with cyanosis or failure; or
    - I) diabetes: or
    - m) Down syndrome; or
    - n) who are pre-or post-splenectomy, or with functional asplenia; or
  - 4) Up to an additional four doses (as appropriate) are funded for the (re-)immunisation of individuals 5 years and over with HIV, pre or post haematopoietic stem cell transplantation, or chemotherapy; pre- or post splenectomy; functional asplenia, pre- or post- solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, intracranial shunts, cerebrospinal fluid leaks or primary immunodeficiency; or
  - For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.
- B) Contractors will be entitled to claim payment from the Funder for the supply of Pneumococcal (PCV13) conjugate vaccine to people eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Pneumococcal (PCV13) conjugate vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Note: please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes n = 30.8 mcg of pneumococcal polysaccharide serotypes 1, 3, 4,

5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and 23F in 0.5ml		
syringe	10	✓ Prevenar 13
	1	✓ Prevenar 13

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	
PNEUMOCOCCAL (PPV23) POLYSACCHARIDE VAC Any of the following:			
<ol> <li>Up to three doses (as appropriate) for patient chemotherapy; pre- or post-splenectomy or w complement deficiency (acquired or inherited)</li> <li>All of the following:</li> </ol>	rith functional asplenia, pre- or p	oost-solid organ	transplant, renal dialysis,
<ul> <li>a) Patient is a child under 18 years for (re-</li> <li>b) Treatment is for a maximum of two dose</li> <li>c) Any of the following:</li> </ul>			
<ul> <li>i) on immunosuppressive therapy or immune response; or</li> <li>ii) with primary immune deficiencies;</li> <li>iii) with HIV infection; or</li> <li>iv) with renal failure, or nephrotic synthematics.</li> <li>v) who are immune-suppressed follor</li> </ul>	or drome; or	·	
or vi) with cochlear implants or intracran vii) with cerebrospinal fluid leaks; or viii) receiving corticosteroid therapy for prednisone of 2 mg/kg per day or or	r more than two weeks, and wh		
20 mg or greater; or ix) with chronic pulmonary disease (ir x) pre term infants, born before 28 w xi) with cardiac disease, with cyanosi xii) with diabetes; or xiii) with Down syndrome; or xiv) who are pre-or post-splenectomy,	eeks gestation; or s or failure; or	gh-dose corticos	steroid therapy); or
For use in testing for primary immunodeficien paediatrician		dation of an inte	ernal medicine physician or
Inj 575 mcg in 0.5 ml prefilled syringe (25 mcg of ea 23 pneumococcal serotype)	following: nated individuals; or on.		Pneumovax 23
Note: Please refer to the Immunisation Handbook Inj 80D antigen units in 0.5 ml syringe			mes. <u>IPOL</u>

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subs	sidised	Generic	
\$	Per	✓	Manufacturer	

#### **ROTAVIBUS OBAL VACCINE**

- a) Only on a prescription
- b) No patient co-payment payable
- c)
- A) Maximum of two doses for people meeting the following:
  - 1) first dose to be administered in infants aged under 14 weeks of age: and
  - 2) no vaccination being administered to children aged 24 weeks or over.
- B) Contractors will be entitled to claim payment from the Funder for the supply of Rotavirus oral vaccine to people eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Rotavirus oral vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Oral susp live attenuated human rotavirus			
1,000,000 CCID50 per dose, squeezable tube	0.00	10	✓ Rotarix
Oral susp live attenuated human rotavirus			
1,000,000 CCID50 per dose, squeezable tube (PVC free)	0.00	10	✓ Rotarix
Oral susp live attenuated human rotavirus			
1,000,000 CCID50 per dose, prefilled oral applicator	0.00	10	✓ Rotarix

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

### VARICELLA VACCINE [CHICKENPOX VACCINE]

- a) Only on a prescription
- b) No patient co-payment payable
- c)
- A) Either:
  - 1) Maximum of one dose for primary vaccination for either:
    - a) Any infant born on or after 1 April 2016; or
    - For previously unvaccinated children turning 11 years old on or after 1 July 2017, who have not previously had a varicella infection (chickenpox), or
  - 2) Maximum of two doses for any of the following:
    - a) Any of the following for non-immune individuals:
      - i) with chronic liver disease who may in future be candidates for transplantation; or
      - ii) with deteriorating renal function before transplantation; or
      - iii) prior to solid organ transplant; or
      - iv) prior to any elective immunosuppression\*; or
      - v) for post exposure prophylaxis who are immune competent inpatients; or
    - b) For individuals at least 2 years after bone marrow transplantation, on advice of their specialist; or
    - c) For individuals at least 6 months after completion of chemotherapy, on advice of their specialist; or
    - d) For HIV positive non immune to varicella with mild or moderate immunosuppression on advice of HIV specialist; or
    - e) For individuals with inborn errors of metabolism at risk of major metabolic decompensation, with no clinical history of varicella; or
    - f) For household contacts of paediatric patients who are immunocompromised, or undergoing a procedure leading to immune compromise where the household contact has no clinical history of varicella; or
    - g) For household contacts of adult patients who have no clinical history of varicella and who are severely immunocompromised, or undergoing a procedure leading to immune compromise where the household contact has no clinical history of varicella.
- B) Contractors will be entitled to claim payment from the Funder for the supply of Varicella vaccine [Chickenpox vaccine] vaccine to people eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Varicella vaccine [Chickenpox vaccine] listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraphs A above.

\* immunosuppression due to steroid or other immunosuppressive therapy must be for a treatment period of greater than 28 days

	NATIONAL	IMMUNISATI	ON SCHEDULE
	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
VARICELLA ZOSTER VACCINE [SHINGLES VACCINE]			
a) Only on a prescription			
b) No patient co-payment payable			
c)			
A) Funded for patients meeting the following criteria:			
1) Either:			

- - 1) Two doses for all people aged 65 years, or
  - 2) Two doses for people 18 years of age or older with any of the following:
    - a) pre- and post-haematopoietic stem cell transplant or cellular therapy; or
    - b) pre- or post-solid organ transplant; or
    - c) haematological malignancies; or
    - d) people living with poorly controlled HIV infection; or
    - e) planned or receiving disease modifying anti-rheumatic drugs (DMARDs targeted synthetic, biologic, or conventional synthetic) for polymyalgia rheumatica, systemic lupus erythematosus or rheumatoid arthritis: or
    - f) end stage kidney disease (CKD 4 or 5); or
    - g) primary immunodeficiency
- B) Contractors will be entitled to claim payment from the Funder for the supply of Varicella zoster vaccine (Shingles vaccine) to patients eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Varicella zoster vaccine [Shingles vaccine] listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Inj 50 mcg per 0.5 ml vial plus vial	0.00	1	✓ Shingrix
		10	✓ Shingrix

## **Diagnostic Agents**

TUBERCULIN PPD [MANTOUX] TEST - [Xpharm]			
Inj 5 TU per 0.1 ml, 1 ml vial	0.00	1	✓ Tubersol

- Symbols -		Albey	267–268	Anaesthetics	12
3TC	112	Albustix		Anagrelide hydrochloride	
- A -		Alchemy Oxaliplatin		Analgesics	
A-Scabies	74	Alchemy Oxybutynin		Anastrozole	
Abacavir sulphate	112	Aldurazyme		Anatrole	18
Abacavir sulphate with		Alecensa		Anoro Ellipta	
lamivudine	112	Alectinib		Antabuse	
Abacavir/Lamivudine Viatris		Alendronate sodium		Antacids and Antiflatulents	
Abilify Maintena		Alendronate sodium with		Anthelmintics	
Abiraterone acetate		colecalciferol	118	Antiacne Preparations	
Acarbose		Alfacalcidol		Antiallergy Preparations	
Accarb		Alfamino		Antianaemics	
Acetazolamide		Alfamino Junior		Antiandrogen Oral	
Acetec		Alginic acid		Contraceptives	8
Acetic acid with hydroxyqui		Alglucosidase alfa		Antiarrhythmics	
ricinoleic acid		Alkeran		Antibacterials	
Acetylcysteine		Allegron		Antibacterials Topical	
Aci-Jel		Allerfix		Anticholinergic Agents	
Aciclovir		Allerpro Syneo 1		Anticholinesterases	
Infection	106	Allerpro Syneo 2		Antidepressants	
Sensory		Allersoothe		Antidiarrhoeals	
Acidex		Allmercap		Antiepilepsy Drugs	
Acipimox		Allopurinol		Antifibrinolytics, Haemostatics an	
Acitretin		Almarytm		Local Sclerosants	
Act-HIB		Alpha-Adrenoceptor Blocker		Antifibrotics	
Actemra		Alpha-Keri Lotion		Antifungals	
Actinomycin D		Alphamox 125		Antifungals Topical	
Actrapid		Alphamox 250		Antihistamines	
Actrapid Penfill		Alprolix		Antihypotensives	
Acupan		Alu-Tab		Antimalarials	
Adalimumab (Amgevita)		Aluminium hydroxide		Antimigraine Preparations	
Adalimumab (Humira - Alter		Amantadine hydrochloride		Antinausea and Vertigo Agents	
brand)		Ambrisentan		Antipruritic Preparations	
Adapalene		Ambrisentan Viatris		Antipsychotics	
Adcetris		Amgevita		Antiretrovirals	
ADR Cartridge 1.8		Amiloride hydrochloride		Antirheumatoid Agents	
Adrenaline		Amiloride hydrochloride with		Antispasmodics and Other Agent	
Cardiovascular	56	furosemide		Altering Gut Motility	
Respiratory		Amiloride hydrochloride with		Antithrombotic Agents	
Advantan		hydrochlorothiazide		Antithymocyte globulin	4
Advate		Aminophylline		(equine)	10
Adynovate		Amiodarone hydrochloride		Antitrichomonal Agents	
Afinitor		Amisulpride		Antituberculotics and	10
Aflibercept		Amitriptyline		Antileprotics	10
AFT-Pyrazinamide		Amlodipine		Antiulcerants	
•	105	Amorolfine			
Agents Affecting the	ım 47	Amoxicillin		Antivirals	10
Renin-Angiotensin Syste					
Agents for Parkinsonism an Disorders		Amoxicillin with clavulanic a Amoxiclav Devatis Forte		Anzatax Apidra	
Agents Used in the Treatme		Amphotericin B		•	
•		Amphotencin B		Apidra SoloStarAPO Clomipramine	
Poisonings				•	
Agrylin		AmsaLyo	15/	APO-Atomoxetine	
Albandazala		Amzasta		Apo-Azithromycin	9
Albendazole	93	Amzoate	<b>ర</b> U	APO-Candesartan HCTZ	

16/12.5	48	Atorvastatin	55	Beta Cream	70
APO-Candesartan HCTZ		Atropine sulphate		Beta Ointment	
32/12.5	48	Cardiovascular	49	Beta Scalp	
Apo-Temozolomide		Sensory		Beta-Adrenoceptor Agonists	
Apomorphine hydrochloride		Atropt		Beta-Adrenoceptor Blockers	
Aprepitant		Atrovent		Beta-hCG low sensitivity urine test	
Apresoline		Augmentin		kit	Q.
Aptamil Feed Thickener		Aurorix		Betadine	
Aqueous cream		AutoSoft 30		Betadine Skin Prep	
Aratac	12	AutoSoft 90		Betaferon	
Arava		Avelox		Betahistine dihydrochloride	
Arginine		Avonex		Betaine	
		Axitinib		Betamethasone dipropionate	
Arginine2000		Azacitidine			/ \
Aripiprazole Sandoz		Azacitidine Dr Reddy's		Betamethasone dipropionate with	7.
Aristocort				calcipotriol  Betamethasone sodium phosphate	/ •
Arrotex-Prazosin S29		Azamun			0
		Azathioprine		with betamethasone acetate	
Arrow - Clopid				Betamethasone valerate	
Arrow - Lattim		Azithromycin		Betamethasone valerate with sodiu	
Arrow Rendreffus-ide		Azopt		fusidate [fusidic acid]	
Arrow-Bendrofluazide		AZT	112	Betnovate	
Arrow-Brimonidine		B-D Micro-Fine	17	Bevacizumab	
Arrow-Diazepam				Bexsero	
Arrow-Doxorubicin		B-D Ultra Fine		Bezafibrate	
Arrow-Fluoxetine	130	B-D Ultra Fine II	17	Bezalip	
Arrow-Losartan &	40	Bacillus Calmette-Guerin (BCG)	400	Bezalip Retard	
Hydrochlorothiazide		vaccine	188	Bicalutamide	
Arrow-Norfloxacin		Bacillus Calmette-Guerin	000	Bicillin LA	
Arrow-Ornidazole		vaccine		BiCNU	
Arrow-Quinapril 10		Baclofen		Bile and Liver Therapy	
Arrow-Quinapril 20		Bactroban		Biltricide	
Arrow-Quinapril 5		Balance		Bimatoprost	
Arrow-Roxithromycin		Barrier Creams and Emollients		Binarex	
Arrow-Timolol		Bayshore		Binocrit	
Arrow-Topiramate		BCG Vaccine AJV		Biocon	
Arrow-Tramadol		Beclazone 100		Biodone	
Arsenic trioxide		Beclazone 250		Biodone Extra Forte	
Asacol		Beclazone 50		Biodone Forte	
Asacol S29		Beclomethasone dipropionate		Bisacodyl	
Ascend-Cefuroxime		Bedaquiline		Bisacodyl Viatris	20
Ascorbic acid		Bee venom allergy treatment		Bisoprolol fumarate	
Aspen Adrenaline	56	Bendamustine hydrochloride		BK Lotion	
Aspirin		Bendamustine Sandoz		Bleomycin sulphate	15
Blood		Bendrofluazide	53	Blood Colony-stimulating	
Nervous		Bendroflumethiazide		Factors	4
Asthalin		[Bendrofluazide]		Blood glucose diagnostic test	
Atazanavir sulphate		Benralizumab		meter	10
Atazanavir Viatris		Benzathine benzylpenicillin		Blood glucose diagnostic test	
Atenolol		Benzatropine mesylate		strip	10
Atenolol AFT		Benzbromarone		Blood glucose test strips (visually	
Atenolol Viatris		Benzetacil		impaired)	10
Atezolizumab		Benztrop		Blood Ketone Diagnostic Test	
ATGAM		Benzydamine hydrochloride		Strip	
Ativan		Benzylpenicillin sodium [Penicillin		Boostrix	
Atnahs Olsalazine		G]		Bortezomib	
Atomoxetine	145	Besponsa	218	Bosentan	60

Bosentan Dr Reddy's	60	Capecitabine	156	Chloramphenicol	
Bplex	32	Capecitabine Viatris	156	Chlorothiazide	5
Brentuximab Vedotin	206	Capsaicin		Chlorpromazine hydrochloride	136
Breo Ellipta	270	Musculoskeletal	118	Chlorsig	27
Brevinor 1/28	79	Nervous	125	Chlortalidone [Chlorthalidone]	
Breztri Aerosphere	271	Captopril	47	Chlorthalidone	
Bricanyl Turbuhaler	270	Carafate		Chlorvescent	46
Brimonidine tartrate		Carbaccord	154	Choice 380 7med Nsha Silver/o	opper
Brimonidine tartrate with timolol		Carbamazepine	131	Short	
maleate	280	Carbimazole	87	Ciclosporin	26
Brinzolamide	279	Carboplatin	154	Cidomycin P/Free	98
BSF Allegron		Carboplatin Accord	154	Cilicaine VK	9
BSF Estradiol TDP Mylan		Carbosorb-X	282	Cinacalcet	8
Budesonide		Cardinol LA	51	Cinacalet Devatis	8
Alimentary	6	Cardizem CD	52	Ciprofloxacin	
Respiratory26		CareSens Dual	15	Infection	98
Budesonide Te Arai	6	CareSens N	16	Sensory	27
Budesonide with eformoterol	269	CareSens N POP	16	Ciprofloxacin Teva	
Budesonide with glycopyrronium a	and	CareSens N Premier		Cisplatin	
eformoterol		CareSens PRO		Cisplatin Accord	154
Bumetanide		Carmellose sodium with gela		Cisplatin Ebewe	
Buprenorphine Naloxone BNM	149	pectin		Citalopram hydrobromide	
Buprenorphine with naloxone		Carmustine		Citrulline1000	
Bupropion hydrochloride		Carvedilol		Cladribine	156
Burel		Carvedilol Sandoz	50	Clarithromycin	
Burinex		Catapres	52	Alimentary	
Buspirone hydrochloride		Cefaclor monohydrate		Infection	9
Buspirone Viatris		Cefalexin		Clexane	
Busulfan		Cefalexin Lupin	93	Clexane Forte	
- C -		Cefalexin Sandoz		Clindamycin	98
Cabergoline	92	Cefazolin	93	Clinicians	
Caffeine citrate		Cefazolin-AFT	93	Clinicians Renal Vit	
Calamine	70	Ceftriaxone	94	Clobazam	13 <sup>-</sup>
Calci-Tab 500	34	Ceftriaxone-AFT		Clobetasol propionate	70, 76
Calcipotriol		Cefuroxime axetil		Clobetasone butyrate	
Calcitonin		Celapram		Clofazimine	
Calcitriol		Celebrex		Clomazol	
Calcitriol XL	33	Celecoxib	117	Dermatological	69
Calcitriol-AFT	33	Celecoxib Pfizer		Genito-Urinary	8
Calcium 500 mg Hexal		Celestone Chronodose	84	Clomifene citrate	92
Calcium carbonate		Cellcept	181	Clomipramine hydrochloride	
Calcium carbonate PAI		Centrally-Acting Agents		Clomipramine Teva	
Calcium Channel Blockers		Cephalexin ABM		Clonazepam	
Calcium Disodium Versenate	283	Cerazette		Clonidine	
Calcium folinate	156	Cetirizine hydrochloride		Clonidine hydrochloride	
Calcium gluconate		Cetomacrogol		Clonidine Teva	52
Calcium Homeostasis		Cetomacrogol with glycerol		Clopidogrel	
Calcium polystyrene sulphonate		Cetomacrogol-AFT		Clopine	136
Calcium Resonium		Cetuximab		Clopixol	
Calogen		Champix		Clotrimazole	,
Camber		Charcoal		Dermatological	69
Candesartan cilexetil		CheckTop		Genito-Urinary	
Candesartan cilexetil with		Chemotherapeutic Agents		Clozapine	
hydrochlorothiazide	48	Chickenpox vaccine		Clozaril	136
Candestar		Chlorafast		Clustran	
Canesten		Chlorambucil		Co-trimoxazole	

Coal tar	74	Cyclizine hydrochloride		DBL Vincristine Sulfate	
Coal tar with allantoin, menthol,		Cyclizine lactate		Decozol	
phenol and sulphur	. 75	Cyclogyl		Deferasirox	
Coal tar with salicylic acid and		Cyclonex		Deferiprone	
sulphur		Cyclopentolate hydrochloride		Denosumab	
Coco-Scalp		Cyclophosphamide		Deolate	
Codeine phosphate		Cyclorin	104	Deoxycoformycin	
Coenzyme Q10		Cycloserine		Depo-Medrol	
Colchicine		Cyproterone acetate	85	Depo-Provera	
Colecalciferol		Cyproterone acetate with		Depo-Testosterone	
Colestyramine		ethinyloestradiol		Deprim	
Colestyramine - Mylan		Cystadane		Dermol	
Colgout		Cytarabine		Desferrioxamine mesilate	
Colifoam		Cytotec		Desmopressin	
Colistin sulphomethate		Cytoxan	154	Desmopressin acetate	
Collodion flexible		- D -		Desmopressin-PH&T	
Colloidal bismuth subcitrate		D-Penamine		Desogestrel	79
Colofac		Dabigatran		Detection of Substances in	
Colomycin		Dabrafenib	167	Urine	8
Coloxyl		Dacarbazine		Dexamethasone	
Combigan	280	Dactinomycin [Actinomycin D]		Hormone	
Comirnaty (LP.8.1)		Daivobet	74	Sensory	
Comirnaty Omicron (JN.1)		Daivonex	74	Dexamethasone phosphate	
Compound electrolytes	. 45	Daktarin	69	Dexamethasone with framycetin	
Compound electrolytes with glucose		Dalacin C		gramicidin	27
[Dextrose]		Dantrium	121	Dexamethasone with neomycin	
Compound hydroxybenzoate	285	Dantrium S29	121	sulphate and polymyxin B	
Concerta	148	Dantrolene		sulphate	
Condoms	. 78	Daonil	11	Dexamfetamine sulfate	14
Condyline		Dapa-Tabs		Dexcom G6	2
Condyline S29	76	Dapsone	104	Dexcom G7	2
Continuous glucose monitor		Darunavir	113	Dexcom ONE+	
(interoperable)	. 22	Darunavir Viatris	113	Dexmethsone	8
Continuous glucose monitor		Dasatinib	168	Dextrochlorpheniramine	
(standalone)	. 23	Dasatinib-Teva	168	maleate	26
Contraceptives - Hormonal	79	Daunorubicin	158	Dextrose	44–4
Contraceptives - Non-hormonal	77	David One Step Cassette Pregr	ancy	DHC Continus	12
Copaxone	141	Test	81	Diabetes	10
Corticosteroids and Related Agents		DBL Acetylcysteine	282	Diabetes Management	1
for Systemic Use	. 84	DBL Adrenaline	56	Diacomit	13
Corticosteroids Topical		DBL Aminophylline	274	Diagnostic Agents	32
Cosentyx		DBL Bleomycin Sulfate		Diamide Relief	
Cosmegen		DBL Bortezomib		Diasip	28
Coumadin	. 44	DBL Carboplatin	154	Diazepam1	131, 140
Country Life	29	DBL Cisplatin		Diazoxide	10
Coversyl	47	DBL Dacarbazine		Diclofenac Devatis	27
COVID-19 vaccine		DBL Desferrioxamine Mesylate	for Inj	Diclofenac Sandoz	11
Creon 10000	.23	BP		Diclofenac sodium	
Creon 25000		DBL Docetaxel		Musculoskeletal	11
Creon Micro		DBL Ergometrine		Sensory	
Crizotinib		DBL Gemcitabine		Differin	
Crotamiton		DBL Gentamicin		Difflam	3
Crystaderm		DBL Leucovorin Calcium		Diflucan	
Cu 375 Standard	.79	DBL Methotrexate Onco-Vial		Digestives Including Enzymes	
Curam Duo 500/125		DBL Naloxone Hydrochloride		Digoxin	
Cvite		DBL Pethidine Hydrochloride		Dihydrocodeine tartrate	

				_
Dilantin	132	Dual blood glucose and blood ketone	Enbrel	181
Dilantin Infatab	132	diagnostic test meter 15	Endocrine Therapy	177
Dilantin Paediatric	132	Dulaglutide12	Endoxan	154
Diltiazem CD Clinect	.52	Dulcolax SP Drop26	Energivit	303
Diltiazem hydrochloride	.52	Duocal Super Soluble Powder287	Engerix-B	314
Dimethicone71		Duolin270	Enhertu	
Dimethyl fumarate	141	Duolin HFA270	Enlafax XR	130
Dipentum	8	DuoResp Spiromax269	Enoxaparin sodium	42
Diphtheria, tetanus and pertussis		Duride56	Enstilar	74
vaccine	311	Durvalumab251	Ensure	296
Diphtheria, tetanus, pertussis and		-E-	Ensure Plus	296
polio vaccine	312	e-chamber La Grande276	Ensure Plus HN	295
Diphtheria, tetanus, pertussis, polio,		e-chamber Mask276	Ensure Plus HN RTH	295
hepatitis B and haemophilus		e-chamber Turbo276	Ensure Two Cal HN RTH	298
influenzae type B vaccine	312	E-Mycin95	Entacapone	123
Diprosone	.70	e5 Pharma10	Entacapone Viatris	
Diprosone OV		Ear Preparations277	Entecavir	
Dipyridamole		Ear/Eye Preparations277	Entecavir (Rex)	
Dipyridamole - Strides		Easiphen Liquid300	Entrectinib	
Disopyramide phosphate		Econazole nitrate69	Entresto 24/26	48
Distoside		Efavirenz112	Entresto 49/51	48
Disulfiram	150	Efavirenz Milpharm112	Entresto 97/103	48
Diuretics	.52	Efavirenz with emtricitabine and	Entyvio	248
Docetaxel		tenofovir disoproxil112	Epilim	
Docetaxel Accord		Eformoterol fumarate dihydrate269	Epilim Crushable	133
Docetaxel Sandoz	158	Eftrenonacog alfa [Recombinant	Epilim IV	
Docusate sodium	.25	factor IX]38	Epilim S/F Liquid	
Docusate sodium with		Efudix76	Epilim Syrup	
sennosides	. 25	Egopsoryl TA75	Epipen	
Dolutegravir	113	Elaprase28	Epipen Jr	
Dolutegravir with lamivudine		Elecare304	Epirubicin Ebewe	
Domperidone		Elecare LCP304	Epirubicin hydrochloride	
Domperidone Viatris		Electral45	Eplerenone	
Donepezil hydrochloride		Elelyso31	Epoetin alfa	
Dornase alfa		Elemental 028 Extra292	Epoprostenol	
Dortimopt		Elexacaftor with tezacaftor, ivacaftor	Eptacog alfa [Recombinant factor	
Dorzolamide with timolol		and ivacaftor274	VIIa]	. 39
Dostinex		Elidel75	Erbitux	
Dosulepin [Dothiepin]		Elocon71	Ergometrine maleate	
hydrochloride	129	Elocon Alcohol Free71	Erlotinib	
Dosulepin Viatris		Eltrombopag38	Erythrocin IV	
Dothiepin		Eltroxin87	Erythromycin (as lactobionate)	
Dovato		EMB Fatol105	Erythromycin ethyl succinate	
Doxazosin		Emend Tri-Pack134	Esbriet	
Doxazosin Clinect		Emicizumab39	Escitalopram	130
Doxine	.97	EMLA125	Escitalopram (Ethics)	
Doxorubicin Ebewe	158	Empagliflozin14	Eskazole	
Doxorubicin hydrochloride		Empagliflozin with metformin	Essential Amino Acid Mix	
Doxycycline		hydrochloride		
DP Lotion		Emsogen	Estradot	
DP Lotn HC		Emtricitabine	Estrofem	
DP-Captopril		Emtricitabine with tenofovir	Estrogel	
Dr Reddy's Omeprazole		disoproxil109	Etanercept	
Drofate		Emtriva	Ethambutol hydrochloride	105
Drugs Affecting Bone		Emulsifying ointment72	Ethics Aspirin	
Metabolism	118	Enalapril maleate47		. 41

Ethinyloestradiol with		Ferrosig	35	Foban	69
desogestrel	79	Ferrous fumarate	34	Folic acid	38
Ethinyloestradiol with		Ferrous fumarate with folic acid	34	Folic Acid multichem	38
levonorgestrel	79	Ferrous sulfate	34	Folic Acid Viatris	38
Ethinyloestradiol with		Fexaclear	268	Food Thickeners	298
norethisterone	79	Fexofenadine hydrochloride	268	Foods And Supplements For	
Ethosuximide	131	Fibro-vein	40	Inherited Metabolic Disease	299
Etopophos	159	Filgrastim	44	Fortini	291
Etoposide	159	Finasteride		Fortini Multi Fibre	
Etoposide phosphate		Fingolimod	141	Fortisip	296
Etravirine	112	Firazyr	267	Fortisip Multi Fibre	
Eumovate		Flagyl	104	Fosamax	
Eurofolic		Flagyl-S	104	Fosamax Plus	
Evara		Flamazine	69	Fosfomycin	115
Evara Emulsifying Ointment	72	Flecainide acetate	49	Framycetin sulphate	277
EVARA Paraffin Ointment 50/50	) <mark>72</mark>	Flecainide BNM	49	Freestyle Libre 2	23
EVARA White Soft Paraffin	72	Flecainide Controlled Release		Freestyle Libre 2 Plus	23
Everet		Teva		Freestyle Libre 3 Plus	
Everolimus	261	Fleet Phosphate Enema	25	Frisium	
Evista	119	Flixonase Hayfever & Allergy		Frumil	
Evrysdi	144	Flixotide		Frusemide	
Exemestane		Flixotide Accuhaler	269	Fucicort	
Exjade		Florinef		Fucidin	
Extemporaneously Compounde	d	Fluanxol		Fucithalmic	
Preparations and		Flucil		Fucithalmic Canada (ON)	277
Galenicals		Flucloxacillin		Fucithalmic Spain	
Eye Preparations		Flucloxin		Fulvestrant	
Eylea		Flucon		Fulvestrant EVER Pharma	
Ezetimibe		Fluconazole		Fungilin	
Ezetimibe Sandoz		Fludara Oral		Furosemide [Frusemide]	
Ezetimibe with simvastatin	56	Fludarabine Ebewe		Furosemide-Baxter	53
-F-		Fludarabine phosphate		fusidic acid	
Factor eight inhibitor bypassing		Fludrocortisone acetate		Dermatological	
fraction		Fluids and Electrolytes		Infection	
Famotidine		Flumetasone pivalate	277	Sensory	27
Famotidine Hovid		Fluocortolone caproate with		-G-	000
Famotidine Hovid MY		fluocortolone pivalate and	•	GA Explore 5	
Faricimab		cinchocaine		GA1 Anamix Infant	
Fasenra		Fluorometholone		GA1 Anamix Junior	
Faslodex		Fluorouracil		Gabapentin	
Fatty Emulsion Cream (Evara).		Fluorouracil Accord		Gacet	
Febuxostat		Fluorouracil sodium		Galsulfase	
Febuxostat (Teva)		Fluox		Galvumet	
FEIBA NF		Fluoxetine hydrochloride		Galvus Gardasil 9	
Felo 10 ER		Flupenthixol decanoate		Gastrodenol	
Felo 5 ER		Flutamide Flutamin			
Felodipine		Fluticasone		Gaviscon Extra Strength	
Fentanyl Sandaz			209	Gaviscon Infant	
Fentanyl Sandoz		Fluticasone furoate with umeclidinium and vilanterol	070	GazyvaGefitinib	218
Ferinject		Fluticasone furoate with	212	Gencitabine Ebewe	
Ferrings		vilanterol	270	Gemcitabine bewe  Gemcitabine hydrochloride	
Ferriprox Ferro-F-Tabs		Fluticasone propionate			
Ferro-Liquid		Fluticasone with salmeterol		Gemtuzumab ozogamicin Gentamicin sulphate	208
Ferro-tab		Flynn		Gilenya	
Ferrograd		FMI		Ginet	

Glatiramer acetate	141	Hemlibra	39	[hydroxycarbamide]	15
Glecaprevir with pibrentasvir	108	Heparin sodium	. 43	Hygroton	5
Glibenclamide	11	Heparin Sodium Panpharma	. 43	Hylo-Fresh	
Gliclazide	11	Heparinised saline		Hymenoptera26	7-26
Glipizide	11	Heparon Junior	290	Hyoscine butylbromide	
Glizide	11	Hepatitis A vaccine		Hyoscine Butylbromide	
Glucagen Hypokit	10	Hepatitis B recombinant		(Adiramedica)	
Glucagon hydrochloride	10	vaccine	314	Hyoscine hydrobromide	13
Glucerna Select		Herzuma		Hyperuricaemia and Antigout	12
Glucose [Dextrose]		Hikma		Hypromellose	
Gluten Free Foods		Hikma Acetylcysteine		Hypromellose with dextran	
Glycerin with sodium saccharin.		Hikma-Propranolol		-1-	
Glycerin with sucrose		Hiprex		Ibiamox	9
Glycerol		Histaclear		Ibrutinib	
Alimentary	25	Holoxan		Ibuprofen	
Extemporaneous		Horleys Bread Mix		Ibuprofen SR BNM	
Glyceryl trinitrate	200	Horleys Flour		Icatibant	
Alimentary	Q	Hormone Replacement Therapy -	200	Idarubicin hydrochloride	
Cardiovascular		Systemic	96	Idursulfase	
Glycopyrronium		HPV		Ifosfamide	
Glycopyrronium bromide		Humalog			
	0	•		lloprost	
Glycopyrronium with	074	Humalog Mix 25		Imatinib mesilate	
indacaterol		Humalog Mix 50		Imatinib-Rex	
Glycosade		Human papillomavirus (6, 11, 16, 18,		Imbruvica	
Glytactin Bettermilk		31, 33, 45, 52 and 58) vaccine	0.45	Imfinzi	
Gold Knight		[HPV]		Imipramine Crescent	
Gold Knight XL		Humatin		Imipramine hydrochloride	
Goserelin		Humira		Imiquimod	
Gynaecological Anti-infectives	80	HumiraPen		Immune Modulators	
- H -		Humulin 30/70		Immunisation - Flu	
Habitrol	151	Humulin NPH		Immunisation Flu and Shingles	
Haemophilus influenzae type B		Humulin R		Immunisation Other	
vaccine		Hyaluronic acid		Immunosuppressants	
Haldol		Hydralazine		Incruse Ellipta	
Haldol Concentrate		Hydralazine hydrochloride		Indacaterol	
Haldol Decanoas		Hydralock		Indapamide	
Haloperidol		Hydralyte - Lemonade	. 45	Infanrix IPV	
Haloperidol decanoate	138	Hydrocortisone		Infanrix-hexa	31
Harvoni	108	Dermatological	. 70	Infant Formulae	30
Havrix 1440		Hormone	. 84	Infatrini	30
Havrix Junior		Hydrocortisone acetate	7	Infliximab	20
Haylor syrup	268	Hydrocortisone acetate with		Influenza vaccine	31
HCU Anamix Infant	299	pramoxine hydrochloride	7	Influvac Tetra	
HCU Anamix Junior	299	Hydrocortisone and paraffin liquid		(2025 formulation)	31
HCU Anamix Junior LQ	299	and lanolin	. 71	Inhaled Corticosteroids	26
HCU Explore 5	299	Hydrocortisone butyrate71	, 76	Inhaled Long-acting	
HCU Express 15	299	Hydrocortisone with cinchocaine	8	Beta-adrenoceptor Agonists	26
HCU Lophlex LQ	299	Hydrocortisone with miconazole	71	Inlyta	16
healthE Calamine Aqueous		Hydrocortisone with natamycin and		Inotuzumab ozogamicin	21
healthE Dimethicone 10%		neomycin	. 71	Inresa	
healthE Dimethicone 4% Lotion		Hydrogen peroxide		Inspra	
healthE Dimethicone 5%		Hydroxocobalamin		Instillagel Lido	
healthE Glycerol BP		Hydroxocobalamin Panpharma	32	Insulin aspart	
healthE Urea Cream		hydroxycarbamide	159	Insulin aspart with insulin aspart	
Healtheries Simple Baking Mix		Hydroxychloroquine sulphate		protamine	1
Hemastix		Hydroxyurea		Insulin degludec with insulin	

aspart10	Iron polymaltose	35	Kliogest	
Insulin glargine11	Isentress		Kliovance	8
Insulin glulisine11	Isentress HD		Kogenate FS	40
Insulin isophane11	Ismo 20	56	Konakion MM	
Insulin isophane with insulin	Ismo 40 Retard		Konakion MM Paediatric	
neutral11	Isoleucine50	303	Konsyl-D	2
Insulin lispro11	Isoniazid	105	Kuvan	29
Insulin lispro with insulin lispro	Isoniazid Teva	105	- L -	
protamine11	Isoniazid with rifampicin		Labetalol	50
Insulin neutral10	Isoptin		Lacosamide	13
Insulin pen needles17	Isoptin Retard	52	Lactulose	2
Insulin pump cartridge18	Isoptin SR	52	Lacuna	
Insulin pump infusion set (steel	Isopto Carpine	280	Laevolac	2
cannula) 19	Isosorbide mononitrate	56	Lamictal	13
Insulin pump infusion set (steel	Isotretinoin	68	Lamivudine	106, 112
cannula, straight insertion) 19	Ispaghula (psyllium) husk	<mark>24</mark>	Lamivudine Viatris	112
Insulin pump infusion set (teflon	Itch-Soothe	70	Lamivudine/Zidovudine Viatris	
cannula)20	Itraconazole	101	Lamotrigine	13
Insulin pump infusion set (teflon	Itraconazole Cresent	101	Lamprene	
cannula, angle insertion with	Itraconazole Kent	101	Lanoxin	49
insertion device)21	Itrazole	101	Lanoxin PG	4
Insulin pump infusion set (teflon	lvacaftor	275	Lanoxin S29	4
cannula, flexible insertion with	Ivermectin	73	Lanreotide	180
insertion device)21	- J -		Lansoprazole	
Insulin pump infusion set (teflon	Jadelle	79	Lantus	1
cannula, straight insertion with	Jakavi	174	Lantus SoloStar	1
insertion device)22	Jardiamet	14	Lanvis	15
Insulin pump infusion set (teflon	Jardiance	14	Lanzol Relief	
cannula, variable insertion)22	Jaydess	79	Largactil	130
Insulin pump reservoir22	Jevity HiCal RTH		Laronidase	
Insulin pump with algorithm17	Jevity RTH		Lasix	5
Insulin syringes, disposable with	Jinarc		Latanoprost	27
attached needle17	Juno Pemetrexed	157	Latanoprost with timolol	
Intelence	- K -		Lax-Suppositories	
Interferon beta-1-alpha141	Kadcyla	245	Lax-suppositories Glycerol	
Interferon beta-1-beta141	Kalydeco		Laxatives	
Intra-uterine device79	Kemadrin		Laxsol	
Invega Sustenna138	Kenacomb		Ledipasvir with sofosbuvir	
Invega Trinza139	Kenacort-A 10		Leflunomide	118
Ipca-Allopurinol120	Kenacort-A 40		Lenalidomide (Viatris)	
Ipca-Bisoprolol50	Kenalog in Orabase		Lenalidomide Viatris	
Ipca-Ciprofloxacin98	Ketocal 3:1		Lenvatinib	
Ipca-Donepezil149	KetoCal 4:1		Lenvima	
Ipca-Escitalopram130	Ketoconazole		Letrole	
IPCA-Frusemide53	Dermatological	76	Letrozole	
Ipca-Hydroxychloroquine118	Infection		Leucine100	
IPCA-Metoprolol50	Ketogenic Diet		Leukeran FC	
IPCA-Propranolol51	Ketoprofen		Leukotriene Receptor	
Ipilimumab251	KetoSens		Antagonists	
IPOL322	Keytruda		Leuprorelin	
Ipratropium bromide270, 276	Kindergen		Leustatin	
Iressa	Kisqali		Levetiracetam	
Irinotecan Actavis 100156	Klacid		Levetiracetam-AFT	
Irinotecan hydrochloride	Alimentary	q	Levocabastine	
Irinotecan-Rex156	Infection		Levocarnitine	
Iron (as ferric carboxymaltose)34	Klaricid		Levodopa with benserazide	
(				

Levodopa with carbidopa	123	Loxamine	130	Metabolics	29
Levodopa with carbidopa and		Lucrin Depot 1-month		Metformin hydrochloride	12
entacapone	123	Lucrin Depot 3-month	92	Metformin Viatris	12
Levomepromazine	136	LumaCina	45	Methadone BNM	127
Levomepromazine		Lumigan	279	Methadone hydrochloride	127
hydrochloride	136	Lynparza		Methenamine (hexamine)	
Levonorgestrel		Lyrica		hippurate	115
Levonorgestrel -1 (Lupin)		- M -		Methopt	
Levonorgestrel BNM		m-Eslon	128	Methotrexate	
Levothyroxine		Mabthera		Methotrexate DBL Onco-Vial	
Lidocaine [Lignocaine]		Macrobid	116	Methotrexate Ebewe	157
Lidocaine [Lignocaine]		Macrogol 3350 with potassium		Methotrexate Sandoz	157
hydrochloride	125	chloride, sodium bicarbonate	and	Methyl hydroxybenzoate	285
Lidocaine [Lignocaine] with		sodium chloride		Methylcellulose	
prilocaine	125	Madopar 125	123	Methylcellulose with glycerin and	
Lidocaine-Baxter		Madopar 250		sodium saccharin	285
Life Extension		Madopar 62.5		Methylcellulose with glycerin and	
Lignocaine		Madopar HBS		sucrose	285
Linezolid		Madopar Rapid		Methyldopa	
Lioresal Intrathecal		Magnesium hydroxide		Methyldopa Viatris	
Lipid-Modifying Agents		Magnesium sulphate		Methylnaltrexone bromide	
Liquigen		Mantoux		Methylphenidate ER - Teva	
Liraglutide		Marevan		Methylphenidate hydrochloride	
Lisdexamfetamine dimesilate		Marine Blue Lotion SPF 50+		Methylphenidate hydrochloride	
Lisinopril		Mask for spacer device		extended-release	148
Lithium carbonate		Maviret		Methylphenidate Sandoz XR	
Livostin		Maxidex		Methylprednisolone	
LMX4		Maxitrol		Methylprednisolone (as sodium	
Lo-Oralcon 20 ED		MCT oil (Nutricia)		succinate)	85
Locacorten-Viaform ED's		Measles, mumps and rubella		Methylprednisolone aceponate	
Local preparations for Anal and		vaccine	318	Methylprednisolone acetate	
Rectal Disorders	8	Mebendazole		Methylxanthines	
Locasol		Mebeverine hydrochloride		Metoclopramide Actavis 10	
Locoid		Medac		Metoclopramide hydrochloride	
Locoid Crelo		Medrol		Metolazone	
Locoid Lipocream		Medroxyprogesterone acetate		Metopirone	
Locorten-Vioform		Genito-Urinary	79	Metoprolol IV Mylan	
Lodoxamide		Hormone		Metoprolol IV Viatris	
Logem		Mefenamic acid		Metoprolol succinate	
Lomide		Mekinist		Metoprolol tartrate	
Lomustine		Melatonin		Metronidamed	
Loniten		Melpha		Metronidazole	
Loperamide hydrochloride		Melphalan		Metyrapone	
Lopinavir with ritonavir		Meningococcal (groups A, C, Y		Mexiletine hydrochloride	
Lopinavir/Ritonavir Mylan		W-135) conjugate vaccine		Miacalcic	
Loprofin		Meningococcal B multicompone		Miacalcic S29	
Loprofin Mix		vaccine		Micolette	
Lorafix		MenQuadfi		Miconazole	
Loratadine		Menthol		Miconazole nitrate	
Lorazepam		Mepolizumab	• • • • • • • • • • • • • • • • • • • •	Dermatological	69
Lorstat		Mercaptopurine		Genito-Urinary	
Losartan Actavis		Mercilon 28		Micreme	
Losartan potassium		Mesalazine		Micreme H	
Losartan potassium with		Mesna		Microlut	
hydrochlorothiazide	48	Mestinon		Midazolam	
Lovir		Metabolic Disorder Agents		Midazolam-Baxter	143

Midazolam-Pfizer	143	Mitomycin (Sagent)	160	Myloc CR	50
Midodrine	50	Mitomycin C		Mylotarg	
Midostaurin	171	Mitozantrone		Myometrial and Vaginal Hormone	
Mifegyne	82	Mitozantrone Ebewe	160	Preparations	80
Mifepristone		MMA/PA Anamix Infant		Myozyme	
Minerals		MMA/PA Anamix Junior		Mytolac	
Mini-Wright AFS Low Range		MMA/PA Explore 5		Mytolac S29	180
Mini-Wright Standard		MMA/PA Express 15		- N -	
Minidiab		Moclobemide		Nadolol	<b>5</b> 1
MiniMed 3.0 Ext Reservoir		Modafinil	148	Nadolol BNM	
MDT-MMT-342G	22	Modafinil Max Health		Naglazyme	27
MiniMed 3.0 Reservoir		Moduretic		Naloxone hydrochloride	
MMT-332A	22	Molaxole		Naltraccord	
MiniMed Mio MMT-921A		Moments		Naltrexone hydrochloride	
MiniMed Mio MMT-923A		Mometasone furoate	71	Naphazoline hydrochloride	
MiniMed Mio MMT-925A		Monogen		Naprosyn SR 1000	
MiniMed Mio MMT-941A		Montelukast		Naprosyn SR 750	
MiniMed Mio MMT-943A		Montelukast Viatris		Naproxen	117
MiniMed Mio MMT-945A		Moroctocog alfa [Recombinant fac		Narcaricin mite	120
MiniMed Mio MMT-965A		VIII]		Nasal Preparations	
MiniMed Mio MMT-975A		Morphine hydrochloride		Natalizumab	141
MiniMed Quick-Set MMT-396A		Morphine sulphate		Natulan	
MiniMed Quick-Set MMT-397A		Motetis		Nausafix	
MiniMed Quick-Set MMT-398A		Mouth and Throat		Nausicalm	
MiniMed Quick-Set MMT-399A		Movapo		Navelbine S29	
MiniMed Silhouette MMT-377A		Movicol		Nefopam hydrochloride	
MiniMed Silhouette MMT-378A		Moxifloxacin		Neo-Mercazole	87
MiniMed Silhouette MMT-381A		MSUD Anamix Infant		Neocate Gold	
MiniMed Sure-T MMT-864A		MSUD Anamix Junior		Neocate Junior Unflavoured	
MiniMed Sure-T MMT-866A		MSUD Anamix Junior LQ		Neocate Junior Vanilla	
MiniMed Sure-T MMT-874A		MSUD Explore 5		Neocate SYNEO	
MiniMed Sure-T MMT-876A		MSUD Express 15		Neoral	
Minims Pilocarpine		MSUD Lophlex LQ 20		Neostigmine metilsulfate	
Minims Prednisolone		MSUD Maxamum		Nepro HP (strawberry)	
Minipress		Mucolytics		Nepro HP (vanilla)	
Minirin		Mucosoothe		Neulactil	
Minirin Melt		Multiple Sclerosis Treatments		NeuroTabs	
Mino-tabs		Multivitamin renal		Nevirapine	
Minocycline hydrochloride		Multivitamins		Nevirapine Viatris	119
Minomycin		Mupirocin		Nicorandil	
Minor Skin Infections		Muscle Relaxants		Nicotine	
Minoxidil		Mvite		Nifedipine	
Minoxidil Roma		Myambutol		Nifuran	
Mio Adv Ext Infusion Set		Mycobutin		Nilotinib	
MDT-MMT-431AK	21	MycoNail		Nilstat	17
Mio Adv Ext Infusion Set	21	Mycophenolate mofetil	181	Alimentary	30
MDT-MMT-432AK	21	Mydriacyl		Genito-Urinary	
Mio Adv Ext Infusion Set	21	Mylan (24 hr release)		Infection	
MDT-MMT-441AK	21			Nimenrix	
Mio Adv Ext Infusion Set	41	Mylan Clomiphen Mylan Italy (24 hr release)		Nintedanib	
MDT-MMT-442AK	91	Myleran	154	Nipent	
Mirena		mylife Inset soft		Niraparib	
Miro-Amoxicillin		mylife Orbit micro		Nirmatrelvir with ritonavir	
Mirtazapine		mylife YpsoPump Reservoir		Nitrates	
Misoprostol		mylife YpsoPump with CamAPS	۷۲	Nitroderm TTS	
Mitomycin (Fresenius Kabi)		FX	17	Nitrofurantoin	
IVIILOTTIYOTT (TTOOTHUO NAUL)	100	ι Λ	17	1 VIII VIUI AI II VIII	1 10

Nitrolingual Pump Spray	56	Nutrison Energy Multi Fibre	296	Ora-Sweet	28
Nivestim		Nutrison Multi Fibre		Ora-Sweet SF	
Nivolumab	252	Nutrison RTH		Orabase	3
Nodia	6	Nyefax Retard	52	Oral and Enteral Feeds	28
Noflam 250	117	Nystatin		Oralcon 30 ED	
Noflam 500	117	Alimentary	32	Oramorph	
Non-Steroidal Anti-Inflammatory		Genito-Urinary		Oramorph CDC S29	
Drugs	117	Infection		Oratane	
Nonacog gamma, [Recombinant		NZB Low Gluten Bread Mix	299	Ornidazole	10
Factor IX]	40	-0-		Orphenadrine citrate	12
Norethisterone		Obinutuzumab	219	Ortho-tolidine	
Genito-Urinary	80	Obstetric Preparations	82	Oruvail SR	11
Hormone	87	Ocicure	9	Osimertinib	
Norflex	122	Ocrelizumab	141	Osmolite RTH	29
Norfloxacin	116	Ocrevus	141	Other Endocrine Agents	9
Noriday	80	Ocrevus SC	141	Other Oestrogen Preparations	8
Noriday 28		Octasa	7	Other Progestogen	
Norimin	79	Octocog alfa [Recombinant fac	tor	Preparations	8
Normison	144	VIII] (Advate)		Other Skin Preparations	7
Norpress	129	Octocog alfa [Recombinant fac		Ovestin	
Nortriptyline hydrochloride	129	VIII] (Kogenate FS)		Genito-Urinary	8
Norvir	113	Octreotide	179	Hormone	8
Noumed Dexamfetamine	145	Octreotide GH		Oxaliplatin	15
Noumed Isoniazid	105	Octreotide long-acting	180	Oxaliplatin Accord	
Noumed Paracetamol	126	Oestradiol		Oxaliplatin Actavis 100	15
Noumed Pethidine	129	Oestradiol valerate	86	Oxaliplatin Ebewe	
Noumed Phenobarbitone	132	Oestradiol with norethisterone.	87	Oxis Turbuhaler	
NovaSource Renal	292	Oestriol		Oxpentifylline	
Novatretin	74	Genito-Urinary	80	Oxybutynin	
Novitium Sugar Free	29	Hormone	87	Oxycodone Amneal	
NovoMix 30 FlexPen	10	Oestrogens	86	Oxycodone hydrochloride	
NovoRapid		Ofev		Oxycodone Lucis	
NovoRapid FlexPen	11	Oil in water emulsion	72	Oxycodone Sandoz	
NovoRapid Penfill	11	Olanzapine	.137-138	Oxytocin	8
NovoSeven RT		Olaparib		Oxytocin BNM	
Nozinan	136	Olbetam		Oxytocin with ergometrine	
Nozinan (Swiss)	136	Olopatadine	281	maleate	8
Nucala		Olopatadine Teva		Ozurdex	27
Nuelin		Olsalazine		- P -	
Nuelin-SR	274	Omalizumab	220	Pacifen	12
Nupentin	131	Omeprazole	9	Pacimol	12
Nusinersen	144	Omeprazole actavis 10	9	Paclitaxel	16
Nutilis	298	Omeprazole actavis 20	9	Paclitaxel Actavis	16
Nutren Diabetes	289	Omeprazole actavis 40	9	Paclitaxel Ebewe	16
Nutrient Modules	286	Omeprazole Teva		Padagis	
Nutrini Energy Multi Fibre	291	Omnitrope	88	Cardiovascular	5
Nutrini Energy RTH	291	Omnitrope AU	88	Dermatological	
Nutrini Low Energy Multi Fibre		Onbrez Breezhaler	269	Paediatric Seravit	
Nutrini Peptisorb	306	Oncaspar LYO	162	Palbociclib	17
Nutrini Peptisorb Energy		OncoTICE		Palbociclib Pfizer	
Nutrini RTH		Ondansetron	135	Paliperidone	
Nutrison 800 Complete Multi		One-Alpha	33	Paliperidone palmitate	
Fibre		Opdivo	252	Palivizumab	22
Nutrison Advanced Peptisorb	292	Ora-Blend	285	Pamidronate disodium	11
Nutrison Concentrated		Ora-Blend SF		Pamisol	
Nutrison Energy		Ora-Plus		Pamol	

Pancreatic enzyme	23	Pevaryl	69	PKU Lophlex LQ 20	30
Pantoprazole	9	Pexsig	52	PKU Lophlex Powder	30
Panzop Relief		Pfizer Exemestane	181	PKU Lophlex Sensation 20	30
Papaverine hydrochloride	57	Pharmacor Varenicline	151	PKU Restore Powder	30
Para-amino salicylic acid	105	Pharmacy Services	282	PKU sphere20 Banana	30
Paracetamol		Pheburane	30	PKU sphere20 Chocolate	30
Paracetamol (Ethics)	126	Phenasen	158	PKU sphere20 Lemon	30
Paracetamol + Codeine		Phenobarbitone	132	PKU sphere20 Red Berry	30
(Relieve)	128	Phenobarbitone sodium		PKU sphere20 Vanilla	
Paracetamol with codeine		Extemporaneous	285	Pku Start	
Paraffin	72	Nervous	143	Plendil ER	5
Paraffin liquid with wool fat	281	Phenoxybenzamine		Pneumococcal (PCV13) conjugate	
Parasiticidal Preparations	73	hydrochloride	47	vaccine	32
Parnate		Phenoxymethylpenicillin (Penic	illin	Pneumococcal (PPV23)	
Paromomycin		V)		polysaccharide vaccine	32
Paroxetine	130	Phenylalanine50	303	Pneumovax 23	
Paser		Phenytoin sodium		Podophyllotoxin	
Paxam	140	Phesgo		Polaramine	
Paxlovid	110	Phillips Milk of Magnesia		Poliomyelitis vaccine	
Paxlovid fee	282	Phlexy 10		Poloxamer	
Paxtine	130	Phosphate Phebra		Poly-Tears	
Pazopanib		Phosphorus		Poly-Visc	
Pazopanib Teva		Phytomenadione		Polycal	
Peak flow meter		Pilocarpine hydrochloride		Polyethylene glycol 400 and	
Pediasure		Pilocarpine nitrate		propylene glycol	28
Pediasure Plus		Pimafucort		Pomalidomide	
Pediasure RTH		Pimecrolimus		Pomolide	
Pegaspargase		Pine tar with trolamine laurilsul		Ponstan	
Pegasys		and fluorescein		Posaconazole	
Pegasys (S29)		Pinetarsol	75	Posaconazole Juno	
Pegasys S29		Pioglitazone		Potassium chloride	
Pegfilgrastim		Pirfenidone		Potassium citrate	
Pegylated interferon alfa-2a	113	Pizotifen		Potassium iodate	
Pembrolizumab		PKU Anamix Infant		Povidone iodine	
Pemetrexed		PKU Anamix Junior		Pradaxa	
Pemetrexed-AFT		PKU Anamix Junior Chocolate		Pramipexole hydrochloride	
Penicillamine		PKU Anamix Junior LQ		Pravastatin	
Penicillin G		PKU Anamix Junior Orange		Praziquantel	
PenMix 30		PKU Anamix Junior Vanilla		Prazosin	
Pentasa		PKU Build 10		Prazosin Mylan	
Pentostatin [Deoxycoformycin].		PKU Build 20 Chocolate		Pred Forte	
Pentoxifylline [Oxpentifylline]		PKU Build 20 Raspberry		Prednisolone	
Peptamen Junior		Lemonade	301	Prednisolone acetate	
Pepti-Junior		PKU Build 20 Smooth		Prednisolone sodium	
Perhexiline maleate		PKU Build 20 Vanilla		phosphate	27
Pericyazine		PKU Explore 10		Prednisolone-AFT	
Perindopril		PKU Explore 5		Prednisone	
Periset		PKU Express 20		Prednisone Clinect	
Periset ODT		PKU First Spoon		Pregabalin	
Perjeta		PKU Glytactin RTD 15		Pregabalin Pfizer	13
Permethrin		PKU Glytactin RTD 15 Lite		Pregnancy Tests - hCG Urine	
Perrigo		PKU GMPro LQ		Premarin	8
Pertuzumab		PKU GMPro Mix-In		Presolol	
Pertuzumab with trastuzumab		PKU GMPro Ultra Lemonade		Prevenar 13	
Peteha		PKU GMPro Ultra Vanilla		Priadel	
Pethidine hydrochloride		PKU Lophlex LQ 10		Primaquine	
. Januario irjandonionao		Lopinon La 10		uquv	

Primidone	132	Rapamune	262	RIXUBIS	4
Primidone Clinect	132	Rasagiline	123	Rizamelt	
Primolut N	87	Reandron 1000		Rizatriptan	13
Priorix	318	Recombinant factor IX	38, 40	Robinul	
Probenecid		Recombinant factor VIIa	39	Ropin	12
Probenecid-AFT	121	Recombinant factor VIII	40	Ropinirole hydrochloride	
Procarbazine hydrochloride	163	Rectogesic	8	Rosemont	
Prochlorperazine	135	Redipred	85	Rosuvastatin	5
Prochlorperazine maleate (Brow	/n &	Relieve		Rosuvastatin Viatris	
Burk)		Relistor	25	Rosuvastatin-Sandoz	5
Proctofoam	7	Remdesivir	110	Rotarix	32
Proctosedyl	8	Remicade	209	Rotavirus oral vaccine	32
Procyclidine hydrochloride	123	Renilon 7.5	292	Roxane-Propranolol	
Progesterone	87	Resonium-A	46	Roxithromycin	9
Proglicem	10	Resource Beneprotein	288	Rozlytrek	16
Progynova	86	Respiratory Devices	<mark>276</mark>	Rubifen	
Prolia	118	Respiratory Stimulants	<mark>276</mark>	Rubifen SR	14
Promethazine hydrochloride	268	Retinol palmitate	281	Rugby	3
Propafenone hydrochloride	49	ReTrieve	68	Rurioctocog alfa pegol [Recomb	inant
Propranolol		Retrovir	112	factor VIII]	
Propylene glycol	285	Revolade	38	Ruxolitinib	17
Propylthiouracil		Ribociclib	174	Rydapt	17
Prostacur	178	Riboflavin	29	Rythmodan - Cheplafarm	4
Protaphane	11	Ribomustin	153	Rythmodan Neon	
Protaphane Penfill		Ricit	81	Rytmonorm	4
Protifar	288	Ricovir	106	Ryzodeg 70/30 Penfill	1
Protionamide		Rifabutin	106	- S -	
Provera	86	Rifadin	106	Sabril	13
Provera HD	87	Rifadin Sanofi	106	Sacubitril with valsartan	
Psoriasis and Eczema		Rifamazid	105	SalAir	
Preparations	74	Rifampicin	106	Salazopyrin	
PTU	87	Rifaximin	10	Salazopyrin EN	
Pulmicort Turbuhaler	269	Rifinah	105	Salbutamol	27
Pulmozyme	274	Rilutek	124	Salbutamol with ipratropium	
Puri-nethol	157	Riluzole	124	bromide	27
Puritan's Pride Vitamin		Rinvoq	264	Salicylic acid	<mark>7</mark>
B-2 100 mg	29	Riodine	<mark>73</mark>	Salmeterol	
Pyrazinamide		Risdiplam	144	Sandomigran	13
Pyridostigmine bromide		Risedronate Sandoz	120	Sandostatin LAR	18
Pyridoxine hydrochloride	32	Risedronate sodium	120	Sanofi Primaquine	10
Pyridoxine multichem	32	Risperdal		Sapropterin dihydrochloride	2
- Q -		Risperdal Consta	139	Scalp Preparations	7
Quantalan sugar free		Risperidone		Scopolamine Transdermal Syste	∍m
Quetapel		Risperidone (Teva)		Viatris	13
Quetiapine	137	Risperon	137	Sebizole	
Quetiapine Viatris	137	Ritalin	147	Secukinumab	
Quinapril		Ritalin LA		Sedatives and Hypnotics	14
Qvar	268	Ritonavir	113	Seebri Breezhaler	
- R -		Rituximab (Mabthera)	224	Senna	2
RA-Morph		Rituximab (Riximyo)	226	Senokot	
Ralicrom		Rivaroxaban		SensoCard	
Raloxifene hydrochloride		Rivastigmine		Serc	
Raltegravir potassium		Rivastigmine Patch BNM 10		Serenace	
Ramipex		Rivastigmine Patch BNM 5.		Seretide	
Ramipril	48	Rivotril		Seretide Accuhaler	
Ranbaxy-Cefaclor	93	Riximyo	226	Serevent	26

Serevent Accuhaler	269	Solu-Cortef	84	Synthroid	87
Sertraline	130	Solu-Medrol	85	Syntometrine	81
Setrona	130	Solu-Medrol-Act-O-Vial	85	Syrup (pharmaceutical grade)	285
Sevredol	128	Somatropin (Omnitrope)	88	Systane Unit Dose	281
Sex Hormones Non		Somatuline Autogel	180	-T-	
Contraceptive	85	Somex		Tacrolimus	
Shingles vaccine	325	Sotalol	51	Dermatological	75
Shingrix		Sotalol Viatris	51	Oncology	263
SII-Onco-BCG		Spacer device	276	Tacrolimus Sandoz	
Sildenafil		Span-K	46	Tafinlar	167
Siltuximab	239	Spazmol	8	Tagrisso	
Simvastatin		Spinal Muscular Atrophy	144	Taliglucerase alfa	31
Simvastatin Mylan		Spinraza		Tambocor	49
Simvastatin Viatris	<u>56</u>	Spiolto Respimat	271	Tambocor German	49
Sinemet	123	Spiractin		Tamoxifen citrate	179
Sinemet CR	123	Spiriva	271	Tamoxifen Sandoz	179
Sintetica Baclofen Intrathecal	121	Spiriva Respimat	271	Tamsulosin hydrochloride	81
Sirolimus	262	Spironolactone	53	Tamsulosin-Rex	81
Sirturo		Stalevo	123	Tandem Cartridge	
Siterone	85	Staphlex	96	Tandem t:slim X2 with Basal-IQ	
Slow-Lopresor	50	Stelara	246	Tandem t:slim X2 with	
Sodibic	46	Stemetil	135	Control-IQ+	17
Sodium acid phosphate	25	Steril-Gene	85	Tap water	285
Sodium alginate		SteroClear		Taro	
Sodium benzoate		Stesolid	131	Tasigna	<b>17</b> 1
Sodium bicarbonate		Stimulants/ADHD Treatments	145	Tasmar	123
Blood	45-46	Stiripentol	133	Taurine	30
Extemporaneous	285	Stomahesive	32	TCu 380 Plus Normal	79
Sodium calcium edetate		Strides Shasun	101	Tecentriq	250
Sodium chloride		Stromectol	73	Tecfidera	
Blood	45	Sucralfate	10	TEEVIR	112
Respiratory	275	Sulfadiazin-Heyl	99	Tegretol	131
Sodium citrate with sodium lauryl		Sulfadiazine Silver		Tegretol CR	
sulphoacetate	25	Sulfadiazine sodium		Telfast	
Sodium citro-tartrate		Sulfasalazine	8	Temaccord	163
Sodium cromoglicate		Sulphur		Temazepam	144
Alimentary	8	Sulprix		Temozolomide	163
Sensory		Sumagran		Temozolomide-Taro	163
Sodium Fusidate [fusidic acid]		Sumatriptan		Tenofovir disoproxil	106
Dermatological	69	Sunitinib	175	Tenofovir Disoproxil Emtricitabine	
Infection	99	Sunitinib Pfizer	175	Mylan	109
Sensory	277	Sunitinib Rex	175	Tenofovir Disoproxil Emtricitabine	
Sodium hyaluronate [Hyaluronic		Sunscreens	76	Viatr	109
acid]	281	Sunscreens, proprietary	76	Tenofovir Disoproxil Viatris	106
Sodium phenylbutyrate		Sustagen Hospital Formula	296	Tenoxicam	117
Sodium picosulfate		Sustanon Ampoules	85	Tensipine MR10	52
Sodium polystyrene sulphonate	46	Sylvant		Tepadina	158
Sodium tetradecyl sulphate	40	Symbicort Turbuhaler 100/6	269	Terbinafine	102
Sodium valproate		Symbicort Turbuhaler 200/6		Terbutaline sulphate	270
Sofradex		Symbicort Turbuhaler 400/12		Teriflunomide	
Soframycin		Symmetrel		Teriflunomide Sandoz	
Solax		Sympathomimetics		Teriparatide	120
Solgar27, 2		Synacthen		Teriparatide - Teva	
Solifenacin succinate		Synacthen Depot		Testogel	
Solifenacin succinate Max		Synacthene Retard		Testosterone	
Health	82	Synagis		Testosterone cipionate	
				•	

Testosterone esters	85	Treatments for Dementia	149	UroFos	11
Testosterone undecanoate	85	Treatments for Substance		Uromitexan	16
Tetrabenazine	124	Dependence	149	Ursodeoxycholic acid	2
Tetrabromophenol	82	Trelegy Ellipta	272	Ursosan	2
Tetracosactrin	85	Trental 400		Ustekinumab	
Tetracycline		Tretinoin		Utrogestan	8
Teva Lisinopril		Dermatological	68	- <b>V</b> -	
Teva-Ketoconazole		Oncology		Vabysmo	20
Thalidomide		Trexate		Vaccinations	
Thalomid		Triamcinolone acetonide		Vaclovir	
Theophylline	274	Alimentary	32	Valaciclovir	
Thiamine hydrochloride	32	Dermatological		Valganciclovir	
Thiamine multichem		Hormone		Valganciclovir Viatris	
THIO-TEPA		Triamcinolone acetonide with		Valine50	
Thioguanine	157	gramicidin, neomycin and ny	statin	Vancomycin	
Thiotepa		Dermatological		Vancomycin Viatris	
Thyroid and Antithyroid Agents		Sensory		Vannair	
Ticagrelor		Trientine		Varenicline tartrate	
Ticagrelor Sandoz		Trientine Waymade		Varicella vaccine [Chickenpox	10
Tilcotil		Trikafta		vaccine]	30
Timolol		Trimethoprim		Varicella zoster vaccine [Shingle	
		•	100	vaccine]	
Tiotropium bromide	211	Trimethoprim with			
Tiotropium bromide with	074	sulphamethoxazole	100	Varilrix	
olodaterol		[Co-trimoxazole]		Various	
Tivicay		Trisequens		VariSoft	
TMP	100	Trisul		Vasodilators	
Tobramycin	400	Trophic Hormones	88	Vasopressin Agonists	
Infection		Tropicamide	280	Vasorex	
Sensory		Trulicity		Vebulis	
Tobramycin (Viatris)		TruSteel		Vedafil	
Tobramycin BNM		Tryzan		Vedolizumab	
Tobrex		Tuberculin PPD [Mantoux] test.		Vegzelma	
Tocilizumab		Tubersol		Veklury	
Tofranil		Two Cal HN		Veletri	
Tolcapone		TYR Anamix Infant	302	Venclexta	
Tolvaptan	54	TYR Anamix Junior	302	Venetoclax	16
Topamax	133	TYR Anamix Junior LQ	302	Venlafaxine	13
Topical Products for Joint and		TYR Explore 5	302	Venomil	267-26
Muscular Pain	118	TYR Lophlex LQ 20	302	VENOX	26
Topiramate	133	TYR Sphere 20	302	Ventolin	27
Topiramate Actavis	133	Tyrosine1000		Vepesid	15
Total parenteral nutrition (TPN)	45	Tysabri	141	Verapamil hydrochloride	5
TPN		. U -		Vermox	
Tramadol hydrochloride	129	UK Cipla	270	Versacloz	13
Tramal SR 100		UK Synacthen		Vesanoid	
Tramal SR 150		Ultibro Breezhaler		Vexazone	
Tramal SR 200		Ultraproct		Vfend	10
Trametinib		Umeclidinium		Viaderm KC	
Trandate		Umeclidinium with vilanterol		Victoza	
Tranexamic acid		Univent		Vigabatrin	
Tranylcypromine sulphate		Upadacitinib		Vigisom	
Trastuzumab (Herzuma)		Ural		Vildagliptin	
Trastuzumab deruxtecan		Urea		Vildagliptin with metformin	1
Trastuzumab emtansine		Urex Forte			4
				hydrochloride	
Travatan		Urinary Agents	01	Vimpat	
Travoprost	2/9	Urinary Tract Infections	115	Vinblastine sulphate	10

Vincristine sulphate165
Vinorelbine165
Vinorelbine Ebewe165
Vinorelbine Te Arai165
Viramune Suspension
ViruPOS277
Vit.D3
VitA-POS
Vitabdeck33
Vital
Vitamin B complex32
Vitamin B6 25
Vitamins32–33
Vivonex TEN292
Voltaren
Voltaren D117
Voltaren SR117
Volumatic276
Voriconazole102
Vttack102
Vyvanse146
- W -
Warfarin sodium44
Wart Preparations76
Wasp venom allergy treatment268
Water
Blood45
Blood45 Extemporaneous285
Extemporaneous285
Extemporaneous285 White Soft Liquid Paraffin AFT72
Extemporaneous285
Extemporaneous
Extemporaneous       285         White Soft Liquid Paraffin AFT       72         Wool fat with mineral oil       72         - X -       Xalkori         166
Extemporaneous       285         White Soft Liquid Paraffin AFT       72         Wool fat with mineral oil       72         - X -       Xalkori       166         Xaluprine       157
Extemporaneous       285         White Soft Liquid Paraffin AFT       72         Wool fat with mineral oil       72         - X -       Xalkori         Xaluprine       157         Xarelto       44
Extemporaneous       285         White Soft Liquid Paraffin AFT       72         Wool fat with mineral oil       72         - X -       Xalkori       166         Xaluprine       157         Xarelto       44         Xgeva       118
Extemporaneous       285         White Soft Liquid Paraffin AFT       72         Wool fat with mineral oil       72         - X -       Xalkori       166         Xaluprine       157         Xarelto       44         Xgeva       118         Xifaxan       10
Extemporaneous       285         White Soft Liquid Paraffin AFT       72         Wool fat with mineral oil       72         - X -       Xalkori       166         Xaluprine       157         Xarelto       44         Xgeva       118         Xifaxan       10         XMET Maxamum       299
Extemporaneous       285         White Soft Liquid Paraffin AFT       72         Wool fat with mineral oil       72         - X -       166         Xaluprine       157         Xarelto       44         Xgeva       118         Xifaxan       10         XMET Maxamum       299         Xolair       220
Extemporaneous       285         White Soft Liquid Paraffin AFT       72         Wool fat with mineral oil       72         - X -       166         Xaluprine       157         Xarelto       44         Xgeva       118         Xifaxan       10         XMET Maxamum       299         Xolair       220         Xolair AU       220
Extemporaneous       285         White Soft Liquid Paraffin AFT       72         Wool fat with mineral oil       72         - X -       166         Xaluprine       157         Xarelto       44         Xgeva       118         Xifaxan       110         XMET Maxamum       299         Xolair       220         XP Maxamum       300
Extemporaneous       285         White Soft Liquid Paraffin AFT       72         Wool fat with mineral oil       72         - X -       166         Xaluprine       157         Xarelto       44         Xgeva       118         Xifaxan       10         XMET Maxamum       299         Xolair       220         Xolair AU       220         XP Maxamum       300         Xylocaine       125
Extemporaneous       285         White Soft Liquid Paraffin AFT       72         Wool fat with mineral oil       72         - X -       - X         Xalkori       166         Xaluprine       157         Xarelto       44         Xgeva       118         Xifaxan       10         XMET Maxamum       299         Xolair       220         Xolair AU       220         XP Maxamum       300         Xylocaine       125         Xylocaine 2% Jelly       124
Extemporaneous       285         White Soft Liquid Paraffin AFT       72         Wool fat with mineral oil       72         - X -       - X         Xalkori       166         Xaluprine       157         Xarelto       44         Xgeva       118         Xifaxan       10         XMET Maxamum       299         Xolair       220         Xolair AU       220         XP Maxamum       300         Xylocaine       125         Xylocaine 2% Jelly       124         Xylocaine Viscous       125
Extemporaneous       285         White Soft Liquid Paraffin AFT       72         Wool fat with mineral oil       72         - X -       - X         Xalkori       166         Xaluprine       157         Xarelto       44         Xgeva       118         Xifaxan       10         XMET Maxamum       299         Xolair       220         Xolair AU       220         XP Maxamum       300         Xylocaine       125         Xylocaine Viscous       125         Xylocard 500       125
Extemporaneous       285         White Soft Liquid Paraffin AFT       72         Wool fat with mineral oil       72         - X -       - X         Xalkori       166         Xaluprine       157         Xarelto       44         Xgeva       118         Xifaxan       10         XMET Maxamum       299         Xolair       220         Xolair AU       220         XP Maxamum       300         Xylocaine       125         Xylocaine Viscous       125         Xylocard 500       125         Xyntha       40
Extemporaneous
Extemporaneous       285         White Soft Liquid Paraffin AFT       72         Wool fat with mineral oil       72         - X -       166         Xalkori       157         Xarelto       44         Xgeva       118         Xifaxan       10         XMET Maxamum       299         Xolair       220         Xolair AU       220         XP Maxamum       300         Xylocaine       125         Xylocaine Viscous       125         Xylocard 500       125         Xyntha       40         - Y -         Yervoy       251
Extemporaneous
Extemporaneous
Extemporaneous
Extemporaneous
Extemporaneous 285 White Soft Liquid Paraffin AFT 72 Wool fat with mineral oil 72 - X - Xalkori. 166 Xaluprine 157 Xarelto 44 Xgeva 118 Xifaxan 10 XMET Maxamum 299 Xolair 220 XP Maxamum 300 Xylocaine 125 Xylocaine 2% Jelly 124 Xylocaine Viscous 125 Xylocard 500 125 Xyntha 40 - Y - Yervoy 251 - Zarontin 131 Zaroxolyn 54 Zavedos 159 Zeffix 106
Extemporaneous
Extemporaneous
Extemporaneous

Zidovudine [AZT]	112
Zidovudine [AZT] with	
lamivudine	
Ziextenzo	44
Zimybe	56
Zinc and castor oil	71
Zinc sulphate	36
Zincaps	
Zincaps S29	36
Ziprasidone	137
Zista	268
Zithromax	
Zo-Rub HP	
Zo-Rub Osteo	
Zoladex	92
Zoledronic acid	
Hormone	
Musculoskeletal	
Zoledronic acid Injection Mylan	84
Zoledronic acid Viatris	
Hormone	
Musculoskeletal	120
Zopiclone	144
Zopiclone Actavis	144
Zostrix	
Zostrix HP	
Zuclopenthixol decanoate	139
Zuclopenthixol hydrochloride	
Zusdone	137
Zyban	150
Zypine	137
Zypine ODT	
Zyprexa Relprevv	
Zytiga	
71/1/02	105