

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
.....
Fax Number:	Fax Number:	

Bendamustine hydrochloride

Initial application — CLL*

Applications only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months.

Prerequisites(tick boxes where appropriate)

<input type="checkbox"/>	The patient has chronic lymphocytic leukaemia requiring treatment
and	
<input type="checkbox"/>	Patient has ECOG performance status of 0-2
and	
<input type="checkbox"/>	Bendamustine is to be administered at a maximum dose of 100 mg/m ² on days 1 and 2 every 4 weeks for a maximum of 6 cycles

Note: Indication marked with a * includes indications that are unapproved. 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL).

Initial application — Indolent, Low-grade lymphomas

Applications only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months.

Prerequisites(tick boxes where appropriate)

<input type="checkbox"/>	The patient has indolent low grade NHL requiring treatment										
and											
<input type="checkbox"/>	The patient has ECOG performance status of 0-2										
and											
<table border="0"><tr><td><input type="checkbox"/></td><td>Patient is treatment naive</td></tr><tr><td>and</td><td></td></tr><tr><td><input type="checkbox"/></td><td>Bendamustine is to be administered for a maximum of 6 cycles (in combination with rituximab when CD20+)</td></tr></table>	<input type="checkbox"/>	Patient is treatment naive	and		<input type="checkbox"/>	Bendamustine is to be administered for a maximum of 6 cycles (in combination with rituximab when CD20+)					
<input type="checkbox"/>	Patient is treatment naive										
and											
<input type="checkbox"/>	Bendamustine is to be administered for a maximum of 6 cycles (in combination with rituximab when CD20+)										
or											
<table border="0"><tr><td><input type="checkbox"/></td><td>Patient is refractory to or has relapsed within 12 months of a rituximab containing combined chemo-immunotherapy regimen</td></tr><tr><td>and</td><td></td></tr><tr><td><input type="checkbox"/></td><td>Bendamustine is to be administered in combination with obinutuzumab for a maximum of 6 cycles</td></tr></table>	<input type="checkbox"/>	Patient is refractory to or has relapsed within 12 months of a rituximab containing combined chemo-immunotherapy regimen	and		<input type="checkbox"/>	Bendamustine is to be administered in combination with obinutuzumab for a maximum of 6 cycles					
<input type="checkbox"/>	Patient is refractory to or has relapsed within 12 months of a rituximab containing combined chemo-immunotherapy regimen										
and											
<input type="checkbox"/>	Bendamustine is to be administered in combination with obinutuzumab for a maximum of 6 cycles										
or											
<table border="0"><tr><td><input type="checkbox"/></td><td>The patient has not received prior bendamustine therapy</td></tr><tr><td>and</td><td></td></tr><tr><td><input type="checkbox"/></td><td>Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+)</td></tr><tr><td>and</td><td></td></tr><tr><td><input type="checkbox"/></td><td>Patient has had a rituximab treatment-free interval of 12 months or more</td></tr></table>	<input type="checkbox"/>	The patient has not received prior bendamustine therapy	and		<input type="checkbox"/>	Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+)	and		<input type="checkbox"/>	Patient has had a rituximab treatment-free interval of 12 months or more	
<input type="checkbox"/>	The patient has not received prior bendamustine therapy										
and											
<input type="checkbox"/>	Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+)										
and											
<input type="checkbox"/>	Patient has had a rituximab treatment-free interval of 12 months or more										
or											
<table border="0"><tr><td><input type="checkbox"/></td><td>Bendamustine is to be administered as monotherapy for a maximum of 6 cycles in rituximab refractory patients</td></tr></table>	<input type="checkbox"/>	Bendamustine is to be administered as monotherapy for a maximum of 6 cycles in rituximab refractory patients									
<input type="checkbox"/>	Bendamustine is to be administered as monotherapy for a maximum of 6 cycles in rituximab refractory patients										

I confirm the above details are correct and that in signing this form I understand I may be audited.

Signed: Date:

Post application to Health New Zealand, Private Bag 3015, Wanganui – email: customerservice@health.govt.nz

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
.....
Fax Number:	Fax Number:

Bendamustine hydrochloride - continued

Renewal — Indolent, Low-grade lymphomas

Current approval Number (if known):

Applications only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months.

Prerequisites(tick boxes where appropriate)

or

<input type="checkbox"/> and	Patient is refractory to or has relapsed within 12 months of rituximab in combination with bendamustine
<input type="checkbox"/> and	Bendamustine is to be administered in combination with obinutuzumab for a maximum of 6 cycles

<input type="checkbox"/> and	Patients have not received a bendamustine regimen within the last 12 months				
<input type="checkbox"/> and	<table border="0"><tr><td><input type="checkbox"/> Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+)</td></tr><tr><td><input type="checkbox"/> and</td><td><table border="0"><tr><td><input type="checkbox"/> Patient has had a rituximab treatment-free interval of 12 months or more</td></tr></table></td></tr></table>	<input type="checkbox"/> Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+)	<input type="checkbox"/> and	<table border="0"><tr><td><input type="checkbox"/> Patient has had a rituximab treatment-free interval of 12 months or more</td></tr></table>	<input type="checkbox"/> Patient has had a rituximab treatment-free interval of 12 months or more
<input type="checkbox"/> Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+)					
<input type="checkbox"/> and	<table border="0"><tr><td><input type="checkbox"/> Patient has had a rituximab treatment-free interval of 12 months or more</td></tr></table>	<input type="checkbox"/> Patient has had a rituximab treatment-free interval of 12 months or more			
<input type="checkbox"/> Patient has had a rituximab treatment-free interval of 12 months or more					

<input type="checkbox"/> or	Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients
-----------------------------	--

Note: 'indolent, low-grade lymphomas' includes follicular, mantle cell, marginal zone and lymphoplasmacytic/ Waldenstrom's macroglobulinaemia.

Initial application — Hodgkin's lymphoma*

Applications only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months.

Prerequisites(tick boxes where appropriate)

<input type="checkbox"/> and	Patient has Hodgkin's lymphoma requiring treatment
<input type="checkbox"/> and	Patient has a ECOG performance status of 0-2
<input type="checkbox"/> and	Patient has received one prior line of chemotherapy
<input type="checkbox"/> and	Patient's disease relapsed or was refractory following prior chemotherapy
<input type="checkbox"/> and	Bendamustine is to be administered in combination with gemcitabine and vinorelbine (BeGeV) at a maximum dose of no greater than 90 mg/m ² twice per cycle, for a maximum of four cycles

Note: Indications marked with * are unapproved indications.

I confirm the above details are correct and that in signing this form I understand I may be audited.

Signed: Date:

Post application to Health New Zealand, Private Bag 3015, Wanganui – email: customerservice@health.govt.nz