

APPLICANT (stamp or sticker acceptable) **PATIENT NHI:** **REFERRER** Reg No:

Reg No: First Names: First Names:

Name: Surname: Surname:

Address: DOB: Address:

..... Address:

.....

Fax Number: Fax Number:

Empagliflozin; Empagliflozin with metformin hydrochloride

Initial application — heart failure reduced ejection fraction

Applications from any relevant practitioner. Approvals valid without further renewal unless notified.

Prerequisites(tick boxes where appropriate)

- ☐ Patient has heart failure
- and ☐ Patient is in NYHA functional class II or III or IV
- and
- ☐ Patient has a documented left ventricular ejection fraction (LVEF) of less than or equal to 40%
- or ☐ An ECHO is not reasonably practicable, and in the opinion of the treating practitioner the patient would benefit from treatment
- and ☐ Patient is receiving concomitant optimal standard funded chronic heart failure treatment

Initial application — Type 2 Diabetes

Applications from any relevant practitioner. Approvals valid without further renewal unless notified.

Prerequisites(tick boxes where appropriate)

- ☐ Patient has previously received an initial approval for a GLP-1 agonist
- or
- ☐ Patient has type 2 diabetes
- and
- ☐ Patient is Māori or any Pacific ethnicity*
- or ☐ Patient has pre-existing cardiovascular disease or risk equivalent (see note a)*
- or ☐ Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator*
- or ☐ Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult*
- or ☐ Patient has diabetic kidney disease (see note b)*
- and ☐ Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of at least one blood-glucose lowering agent (e.g. metformin, vildagliptin, or insulin) for at least 3 months

Note: * Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

- a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack, ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.
- b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m² in the presence of diabetes, without alternative cause.
- c) Funded [empagliflozin / empagliflozin with metformin hydrochloride] treatment is not to be given in combination with a funded GLP-1 unless receiving (empagliflozin / empagliflozin with metformin hydrochloride) for the treatment of heart failure.

I confirm the above details are correct and that in signing this form I understand I may be audited.

Signed: Date:

Post application to Health New Zealand, Private Bag 3015, Wanganui – email: customerservice@health.govt.nz