

SA2623 - Rituximab

ABO-incompatible organ transplant - Initial application	2
ANCA associated vasculitis - Initial application	2
ANCA associated vasculitis - Renewal	2
Antibody-mediated organ transplant rejection - Initial application	2
B-cell acute lymphoblastic leukaemia/lymphoma* - Initial application	15
CD20+ low grade or follicular B-cell NHL - Initial application	14
CD20+ low grade or follicular B-cell NHL - Renewal	14
Chronic lymphocytic leukaemia - Initial application	3
Chronic lymphocytic leukaemia - Renewal	4
Membranous nephropathy - Initial application	15
Post-transplant - Initial application	5
Post-transplant - Renewal	5
Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS) - Initial application	6
Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS) - Renewal	6
Steroid resistant nephrotic syndrome (SRNS) - Initial application	7
Steroid resistant nephrotic syndrome (SRNS) - Renewal	7
Aggressive CD20 positive NHL - Initial application	7
Aggressive CD20 positive NHL - Renewal	8
Anti-NMDA receptor autoimmune encephalitis* - Initial application	14
Antisynthetase syndrome - Initial application	13
Chronic inflammatory demyelinating polyneuropathy (CIPD)* - Initial application	13
Desensitisation prior to transplant - Initial application	15
Graft versus host disease - Initial application	13
Haemophilia with inhibitors - Initial application	8
Haemophilia with inhibitors - Renewal	8
Immune thrombocytopenic purpura (ITP) - Initial application	9
Immune thrombocytopenic purpura (ITP) - Renewal	9
Immunoglobulin G4-related disease (IgG4-RD*) - Initial application	17
Immunoglobulin G4-related disease (IgG4-RD*) - Renewal	17
Indolent, low-grade lymphomas or hairy cell leukaemia* - Initial application	10
Indolent, low-grade lymphomas or hairy cell leukaemia* - Renewal	10
Neuromyelitis optica spectrum disorder (NMOSD)* - Initial application	4
Pemphigus* - Initial application	16
Pemphigus* - Renewal	16
Pure red cell aplasia (PRCA) - Initial application	10
Pure red cell aplasia (PRCA) - Renewal	10
Refractory myasthenia gravis* - Initial application	5
Refractory myasthenia gravis* - Renewal	6
Severe cold haemagglutinin disease (CHAD) - Initial application	11
Severe cold haemagglutinin disease (CHAD) - Renewal	11
Thrombotic thrombocytopenic purpura (TTP)* - Initial application	11
Treatment refractory systemic lupus erythematosus (SLE)* - Initial application	12
Warm autoimmune haemolytic anaemia (warm AIHA) - Initial application	12
Warm autoimmune haemolytic anaemia (warm AIHA) - Renewal	12

APPLICANT (stamp or sticker acceptable) **PATIENT NHI:** **REFERRER** Reg No:

Reg No: First Names: First Names:

Name: Surname: Surname:

Address: DOB: Address:

..... Address:

.....

Fax Number: Fax Number:

Rituximab (Riximyo)

Initial application — ABO-incompatible organ transplant
Applications from any relevant practitioner. Approvals valid without further renewal unless notified.
Prerequisites(tick box where appropriate)

Patient is to undergo an ABO-incompatible solid organ transplant*

Note: Indications marked with * are unapproved indications.

Initial application — ANCA associated vasculitis
Applications from any relevant practitioner. Approvals valid for 8 weeks.
Prerequisites(tick boxes where appropriate)

Patient has been diagnosed with ANCA associated vasculitis*

and The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks

and

Induction therapy with daily oral or pulse intravenous cyclophosphamide has failed to achieve significant improvement of disease after at least 3 months

or Patient has previously had a cumulative dose of cyclophosphamide > 15 g or a further repeat 3 month induction course of cyclophosphamide would result in a cumulative dose > 15 g

or Cyclophosphamide and methotrexate are contraindicated

or Patient is a female of child-bearing potential

or Patient has a previous history of haemorrhagic cystitis, urological malignancy or haematological malignancy

Note: Indications marked with * are unapproved indications.

Renewal — ANCA associated vasculitis

Current approval Number (if known):.....

Applications from any relevant practitioner. Approvals valid for 8 weeks.
Prerequisites(tick boxes where appropriate)

Patient has been diagnosed with ANCA associated vasculitis*

and Patient has previously responded to treatment with rituximab but is now experiencing an acute flare of vasculitis

and The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks

Note: Indications marked with * are unapproved indications.

Initial application — Antibody-mediated organ transplant rejection
Applications from any relevant practitioner. Approvals valid without further renewal unless notified.
Prerequisites(tick box where appropriate)

Patient has been diagnosed with antibody-mediated organ transplant rejection*

Note: Indications marked with * are unapproved indications.

I confirm the above details are correct and that in signing this form I understand I may be audited.

Signed: Date:
Post application to Health New Zealand, Private Bag 3015, Wanganui – email: customerservice@health.govt.nz

APPLICANT (stamp or sticker acceptable) **PATIENT NHI:** **REFERRER** Reg No:

Reg No: First Names: First Names:

Name: Surname: Surname:

Address: DOB: Address:

..... Address:

.....

Fax Number: Fax Number:

Rituximab (Riximyo) - *continued*

Initial application — Chronic lymphocytic leukaemia

Applications from any relevant practitioner. Approvals valid for 12 months.

Prerequisites(tick boxes where appropriate)

The patient has progressive Binet stage A, B or C chronic lymphocytic leukaemia (CLL) requiring treatment

and

The patient is rituximab treatment naive

or

The patient is chemotherapy treatment naive

or

The patient's disease has relapsed following no more than three prior lines of chemotherapy treatment

and

The patient has had a treatment-free interval of 12 months or more if previously treated with fludarabine and cyclophosphamide chemotherapy

or

The patient's disease has relapsed and rituximab treatment is to be used in combination with funded venetoclax

and

The patient has good performance status

and

The patient does not have chromosome 17p deletion CLL

or

Rituximab treatment is to be used in combination with funded venetoclax for relapsed/refractory chronic lymphocytic leukaemia

and

Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles

and

It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration), bendamustine or venetoclax

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with rituximab is expected to improve symptoms and improve ECOG score to < 2.

I confirm the above details are correct and that in signing this form I understand I may be audited.

Signed: Date:

Post application to Health New Zealand, Private Bag 3015, Wanganui – email: customerservice@health.govt.nz

APPLICANT (stamp or sticker acceptable) **PATIENT NHI:** **REFERRER** Reg No:

Reg No: First Names: First Names:

Name: Surname: Surname:

Address: DOB: Address:

..... Address:

.....

Fax Number: Fax Number:

Rituximab (Riximyo) - *continued*

Renewal — Chronic lymphocytic leukaemia

Current approval Number (if known):.....

Applications from any relevant practitioner. Approvals valid for 12 months.

Prerequisites(tick boxes where appropriate)

The patient's disease has relapsed and rituximab treatment is to be used in combination with funded venetoclax

or

The patient's disease has relapsed following no more than one prior line of treatment with rituximab for CLL

and

The patient has had an interval of 36 months or more since commencement of initial rituximab treatment

and

The patient does not have chromosome 17p deletion CLL

and

It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration) or bendamustine

and

Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initial application — neuromyelitis optica spectrum disorder (NMOSD)*

Applications from any relevant practitioner. Approvals valid without further renewal unless notified.

Prerequisites(tick boxes where appropriate)

Cumulative dose up to 1,500 mg/m² body surface area up to 2,000 mg total per cycle

and

Patient has experienced a severe episode or attack of NMOSD (rapidly progressing symptoms with supporting clinical investigations)

or

Patient has experienced a breakthrough attack of NMOSD

and

Patient is receiving treatment with mycophenolate unless contraindicated or not tolerated

and

Patient is receiving treatment with corticosteroids unless contraindicated or not tolerated

and

Each treatment cycle at least 6 months apart

Note: Indications marked with * are unapproved indications.

I confirm the above details are correct and that in signing this form I understand I may be audited.

Signed: Date:

Post application to Health New Zealand, Private Bag 3015, Wanganui – email: customerservice@health.govt.nz

APPLICANT (stamp or sticker acceptable) **PATIENT NHI:** **REFERRER** Reg No:

Reg No: First Names: First Names:

Name: Surname: Surname:

Address: DOB: Address:

..... Address:

.....

Fax Number: Fax Number:

Rituximab (Riximyo) - *continued*

Initial application — Post-transplant

Applications from any relevant practitioner. Approvals valid for 12 months.

Prerequisites(tick boxes where appropriate)

- The patient has B-cell post-transplant lymphoproliferative disorder*
- and**
- To be used for a maximum of 8 treatment cycles

Note: Indications marked with * are unapproved indications.

Renewal — Post-transplant

Current approval Number (if known):.....

Applications from any relevant practitioner. Approvals valid for 9 months.

Prerequisites(tick boxes where appropriate)

- The patient has had a rituximab treatment-free interval of 12 months or more
- and**
- The patient has B-cell post-transplant lymphoproliferative disorder*
- and**
- To be used for no more than 6 treatment cycles

Note: Indications marked with * are unapproved indications.

Initial application — refractory myasthenia gravis*

Applications from any relevant practitioner. Approvals valid for 2 years.

Prerequisites(tick boxes where appropriate)

- Cumulative dose up to 1,500 mg/m² body surface area up to 2,000 mg total per cycle
- and**
- Treatment with corticosteroids and at least one other immunosuppressant for a minimum period of 12 months has been ineffective
- or**
- Treatment with at least one other immunosuppressant for a period of at least 12 months
- and**
- Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects

Note: Indications marked with * are unapproved indications.

I confirm the above details are correct and that in signing this form I understand I may be audited.

Signed: Date:

Post application to Health New Zealand, Private Bag 3015, Wanganui – email: customerservice@health.govt.nz

APPLICANT (stamp or sticker acceptable) **PATIENT NHI:** **REFERRER** Reg No:

Reg No: First Names: First Names:

Name: Surname: Surname:

Address: DOB: Address:

..... Address:

.....

Fax Number: Fax Number:

Rituximab (Riximyo) - *continued*

Renewal — refractory myasthenia gravis*

Current approval Number (if known):.....

Applications from any relevant practitioner. Approvals valid for 2 years.

Prerequisites(tick boxes where appropriate)

Cumulative dose up to 1,500 mg/m² body surface area up to 2,000 mg total per cycle

and

An initial response lasting at least 12 months was demonstrated

and

Patient has relapsed despite treatment with corticosteroids and at least one other immunosuppressant for a period of at least 12 months

or

Patient's myasthenia gravis has relapsed despite treatment with at least one immunosuppressant for a period of at least 12 months

and

Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects

Note: Indications marked with * are unapproved indications.

Initial application — Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)

Applications from any relevant practitioner. Approvals valid for 8 weeks.

Prerequisites(tick boxes where appropriate)

Patient is a child with SDNS* or FRNS*

and

Treatment with corticosteroids, ciclosporin, and mycophenolate for at least 3 months for each agent has been ineffective, not tolerated, or is contraindicated

and

The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks

Note: Indications marked with * are unapproved indications.

Renewal — Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)

Current approval Number (if known):.....

Applications from any relevant practitioner. Approvals valid for 8 weeks.

Prerequisites(tick boxes where appropriate)

Patient who was previously treated with rituximab for nephrotic syndrome*

and

Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment

and

The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks

Note: Indications marked with * are unapproved indications.

I confirm the above details are correct and that in signing this form I understand I may be audited.

Signed: Date:

Post application to Health New Zealand, Private Bag 3015, Wanganui – email: customerservice@health.govt.nz

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
.....	Address:
.....
Fax Number:	Fax Number:

Rituximab (Riximyo) - *continued*

Initial application — Steroid resistant nephrotic syndrome (SRNS)

Applications from any relevant practitioner. Approvals valid for 8 weeks.

Prerequisites(tick boxes where appropriate)

<input type="checkbox"/>	Patient is a child with SRNS* and treatment with corticosteroids, ciclosporin and tacrolimus for at least 3 months for each agent has been ineffective, not tolerated, or is contraindicated
and	
<input type="checkbox"/>	Genetic causes of nephrotic syndrome have been excluded
and	
<input type="checkbox"/>	The total rituximab dose per cycle would not exceed the equivalent of 375 mg/m ² of body surface area per week for a total of 4 weeks

Note: Indications marked with * are unapproved indications.

Renewal — Steroid resistant nephrotic syndrome (SRNS)

Current approval Number (if known):.....

Applications only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks.

Prerequisites(tick boxes where appropriate)

<input type="checkbox"/>	Patient who was previously treated with rituximab for nephrotic syndrome*
and	
<input type="checkbox"/>	Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment
and	
<input type="checkbox"/>	The total rituximab dose used would not exceed the equivalent of 375 mg/m ² of body surface area per week for a total of 4 weeks

Note: Indications marked with * are unapproved indications.

Initial application — aggressive CD20 positive NHL

Applications from any relevant practitioner. Approvals valid for 12 months.

Prerequisites(tick boxes where appropriate)

<input type="checkbox"/>	The patient has treatment naive aggressive CD20 positive NHL
and	
<input type="checkbox"/>	To be used with a multi-agent chemotherapy regimen given with curative intent
and	
<input type="checkbox"/>	To be used for a maximum of 8 treatment cycles
or	
<input type="checkbox"/>	The patient has aggressive CD20 positive NHL with relapsed disease following prior chemotherapy
and	
<input type="checkbox"/>	To be used for a maximum of 6 treatment cycles

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

I confirm the above details are correct and that in signing this form I understand I may be audited.

Signed: Date:

Post application to Health New Zealand, Private Bag 3015, Wanganui – email: customerservice@health.govt.nz

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
.....	Address:
.....
Fax Number:	Fax Number:

Rituximab (Riximyo) - *continued*

Renewal — aggressive CD20 positive NHL

Current approval Number (if known):.....

Applications from any relevant practitioner. Approvals valid for 12 months.

Prerequisites(tick boxes where appropriate)

<input type="checkbox"/>	The patient has had a rituximab treatment-free interval of 12 months or more
and	
<input type="checkbox"/>	The patient has relapsed refractory/aggressive CD20 positive NHL
and	
<input type="checkbox"/>	To be used with a multi-agent chemotherapy regimen given with curative intent
and	
<input type="checkbox"/>	To be used for a maximum of 4 treatment cycles

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Initial application — haemophilia with inhibitors

Applications only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months.

Prerequisites(tick boxes where appropriate)

<input type="checkbox"/>	Patient has mild congenital haemophilia complicated by inhibitors
or	
<input type="checkbox"/>	Patient has severe congenital haemophilia complicated by inhibitors and has failed immune tolerance therapy
or	
<input type="checkbox"/>	Patient has acquired haemophilia

Renewal — haemophilia with inhibitors

Current approval Number (if known):.....

Applications only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months.

Prerequisites(tick boxes where appropriate)

<input type="checkbox"/>	Patient was previously treated with rituximab for haemophilia with inhibitors
and	
<input type="checkbox"/>	An initial response lasting at least 12 months was demonstrated
and	
<input type="checkbox"/>	Patient now requires repeat treatment

I confirm the above details are correct and that in signing this form I understand I may be audited.

Signed: Date:

Post application to Health New Zealand, Private Bag 3015, Wanganui – email: customerservice@health.govt.nz

APPLICANT (stamp or sticker acceptable) **PATIENT NHI:** **REFERRER** Reg No:

Reg No: First Names: First Names:

Name: Surname: Surname:

Address: DOB: Address:

..... Address:

.....

Fax Number: Fax Number:

Rituximab (Riximyo) - *continued*

Initial application — immune thrombocytopenic purpura (ITP)

Applications only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks.

Prerequisites(tick boxes where appropriate)

Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per microlitre

or

Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding

and

Treatment with steroids and splenectomy have been ineffective

or

Treatment with steroids has been ineffective and splenectomy is an absolute contraindication

or

Other treatments including steroids have been ineffective and patient is being prepared for elective surgery (e.g. splenectomy)

and

The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks

Note: Indications marked with * are unapproved indications.

Renewal — immune thrombocytopenic purpura (ITP)

Current approval Number (if known):.....

Applications only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks.

Prerequisites(tick boxes where appropriate)

Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned

or

Patient was previously treated with rituximab for immune thrombocytopenic purpura*

and

An initial response lasting at least 12 months was demonstrated

and

Patient now requires repeat treatment

Note: Indications marked with * are unapproved indications.

I confirm the above details are correct and that in signing this form I understand I may be audited.

Signed: Date:

Post application to Health New Zealand, Private Bag 3015, Wanganui – email: customerservice@health.govt.nz

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
.....	Address:
.....
Fax Number:	Fax Number:

Rituximab (Riximyo) - *continued*

Initial application — indolent, low-grade lymphomas or hairy cell leukaemia*

Applications from any relevant practitioner. Approvals valid for 9 months.

Prerequisites(tick boxes where appropriate)

and	<input type="checkbox"/> The patient has indolent low grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy
	<input type="checkbox"/> To be used for a maximum of 6 treatment cycles
or	
and	<input type="checkbox"/> The patient has indolent, low grade lymphoma or hairy cell leukaemia* requiring first-line systemic chemotherapy
	<input type="checkbox"/> To be used for a maximum of 6 treatment cycles

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Renewal — indolent, low-grade lymphomas or hairy cell leukaemia*

Current approval Number (if known):.....

Applications from any relevant practitioner. Approvals valid for 12 months.

Prerequisites(tick boxes where appropriate)

and	<input type="checkbox"/> The patient has had a rituximab treatment-free interval of 12 months or more
	<input type="checkbox"/> The patient has indolent, low-grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy
and	<input type="checkbox"/> To be used for no more than 6 treatment cycles

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Initial application — pure red cell aplasia (PRCA)

Applications only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks.

Prerequisites(tick box where appropriate)

Patient has autoimmune pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder

Note: Indications marked with * are unapproved indications.

Renewal — pure red cell aplasia (PRCA)

Current approval Number (if known):.....

Applications only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks.

Prerequisites(tick box where appropriate)

Patient was previously treated with rituximab for pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder and demonstrated an initial response lasting at least 12 months

Note: Indications marked with * are unapproved indications.

I confirm the above details are correct and that in signing this form I understand I may be audited.

Signed: Date:

Post application to Health New Zealand, Private Bag 3015, Wanganui – email: customerservice@health.govt.nz

APPLICANT (stamp or sticker acceptable) **PATIENT NHI:** **REFERRER** Reg No:

Reg No: First Names: First Names:

Name: Surname: Surname:

Address: DOB: Address:

..... Address:

.....

Fax Number: Fax Number:

Rituximab (Riximyo) - *continued*

Initial application — severe cold haemagglutinin disease (CHAD)

Applications only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks.

Prerequisites(tick boxes where appropriate)

- Patient has cold haemagglutinin disease*
- and**
- Patient has severe disease which is characterized by symptomatic anaemia, transfusion dependence or disabling circulatory symptoms
- and**
- The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks

Note: Indications marked with * are unapproved indications.

Renewal — severe cold haemagglutinin disease (CHAD)

Current approval Number (if known):.....

Applications only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks.

Prerequisites(tick boxes where appropriate)

- Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned
- or**
- Patient was previously treated with rituximab for severe cold haemagglutinin disease*
- and**
- An initial response lasting at least 12 months was demonstrated
- and**
- Patient now requires repeat treatment

Note: Indications marked with * are unapproved indications.

Initial application — thrombotic thrombocytopenic purpura (TTP)*

Applications only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid without further renewal unless notified.

Prerequisites(tick boxes where appropriate)

- The total rituximab dose per cycle would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks
- and**
- Each treatment cycle at least 6 months apart
- and**
- Patient has experienced progression of clinical symptoms or persistent thrombocytopenia despite plasma exchange
- or**
- Patient has acute idiopathic TTP* with neurological or cardiovascular pathology

Note: Indications marked with * are unapproved indications.

I confirm the above details are correct and that in signing this form I understand I may be audited.

Signed: Date:

Post application to Health New Zealand, Private Bag 3015, Wanganui – email: customerservice@health.govt.nz

APPLICANT (stamp or sticker acceptable) **PATIENT NHI:** **REFERRER** Reg No:

Reg No: First Names: First Names:

Name: Surname: Surname:

Address: DOB: Address:

..... Address:

.....

Fax Number: Fax Number:

Rituximab (Riximyo) - *continued*

Initial application — treatment refractory systemic lupus erythematosus (SLE)*

Applications only from a rheumatologist, nephrologist or any relevant practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid without further renewal unless notified.

Prerequisites(tick boxes where appropriate)

Patient has severe, immediately life- or organ-threatening SLE*

and

The condition has been refractory to treatment with corticosteroids at a dose of at least 1 mg/kg unless contraindicated

and

The condition has relapsed following prior treatment for at least 6 months with maximal tolerated doses of azathioprine, mycophenolate mofetil and high dose cyclophosphamide, or cyclophosphamide is contraindicated

and

Initial treatment maximum of four 1000 mg infusions

and

Treatment for relapse following initial partial response to rituximab up to a maximum of two 1000 mg infusions every 6 months

Note: Indications marked with * are unapproved indications.

Initial application — warm autoimmune haemolytic anaemia (warm AIHA)

Applications only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks.

Prerequisites(tick boxes where appropriate)

Patient has warm autoimmune haemolytic anaemia*

and

One of the following treatments has been ineffective: steroids (including if patient requires ongoing steroids at doses equivalent to > 5 mg prednisone daily), cytotoxic agents (e.g. cyclophosphamide monotherapy or in combination), intravenous immunoglobulin

and

The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks

Note: Indications marked with * are unapproved indications.

Renewal — warm autoimmune haemolytic anaemia (warm AIHA)

Current approval Number (if known):.....

Applications only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks.

Prerequisites(tick boxes where appropriate)

Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned

or

Patient was previously treated with rituximab for warm autoimmune haemolytic anaemia*

and

An initial response lasting at least 12 months was demonstrated

and

Patient now requires repeat treatment

Note: Indications marked with * are unapproved indications.

I confirm the above details are correct and that in signing this form I understand I may be audited.

Signed: Date:

Post application to Health New Zealand, Private Bag 3015, Wanganui – email: customerservice@health.govt.nz

APPLICANT (stamp or sticker acceptable) **PATIENT NHI:** **REFERRER** Reg No:

Reg No: First Names: First Names:

Name: Surname: Surname:

Address: DOB: Address:

..... Address:

.....

Fax Number: Fax Number:

Rituximab (Riximyo) - *continued*

Initial application — antisynthetase syndrome
Applications from any relevant practitioner. Approvals valid without further renewal unless notified.

Prerequisites(tick boxes where appropriate)

Patient has severe, immediately life- or organ-threatening disease, including interstitial lung disease

and

Treatment with at least 3 immunosuppressants (oral corticosteroids, cyclophosphamide, methotrexate, mycophenolate, ciclosporin, azathioprine) has been ineffective controlling active disease

or

Rapid treatment is required for life threatening complications

and

Maximum of two 1000mg infusions every 6 months

Initial application — graft versus host disease
Applications from any relevant practitioner. Approvals valid without further renewal unless notified.

Prerequisites(tick boxes where appropriate)

Patient has refractory graft versus host disease following transplant

and

Treatment with at least 3 immunosuppressants (oral steroids, ciclosporin, tacrolimus, mycophenolate, sirolimus) has not be effective at controlling active disease

and

The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks

Initial application — chronic inflammatory demyelinating polyneuropathy (CIPD)*
Applications only from a neurologist or any relevant practitioner on the recommendation of a neurologist. Approvals valid without further renewal unless notified.

Prerequisites(tick boxes where appropriate)

Treatment with corticosteroids and intravenous immunoglobulin and/or plasma exchange has been ineffective controlling active disease, is not tolerated, or is contraindicated

and

At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) is not tolerated or has been ineffective controlling active disease. If an immunosuppressant is contraindicated, a trial has occurred of one of those which is not contraindicated (unless all are contraindicated)

or

Rapid treatment is required for life threatening complications

and

Cumulative dose up to 1500 mg/m² body surface area up to 2000 mg total per cycle

and

Each treatment cycle at least 6 months apart

Note: Indications marked with * are unapproved indications.

I confirm the above details are correct and that in signing this form I understand I may be audited.

Signed: Date:
Post application to Health New Zealand, Private Bag 3015, Wanganui – email: customerservice@health.govt.nz

APPLICANT (stamp or sticker acceptable) **PATIENT NHI:** **REFERRER** Reg No:

Reg No: First Names: First Names:

Name: Surname: Surname:

Address: DOB: Address:

..... Address:

.....

Fax Number: Fax Number:

Rituximab (Riximyo) - *continued*

Initial application — anti-NMDA receptor autoimmune encephalitis*

Applications from any relevant practitioner. Approvals valid without further renewal unless notified.

Prerequisites(tick boxes where appropriate)

Treatment with corticosteroids and intravenous immunoglobulin and/or plasma exchange has been ineffective controlling active disease, is not tolerated or is contraindicated

and

At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has been ineffective controlling active disease, is not tolerated or is contraindicated

or

Rapid treatment is required for life threatening complications

and

Cumulative dose up to 1500 mg/m² body surface area up to 2000 mg total per cycle

and

Each treatment cycle at least 6 months apart

Note: Indications marked with * are unapproved indications.

Initial application — CD20+ low grade or follicular B-cell NHL

Applications from any relevant practitioner. Approvals valid for 9 months.

Prerequisites(tick boxes where appropriate)

The patient has CD20+ low grade or follicular B-cell NHL with relapsed disease following prior chemotherapy

and

To be used for a maximum of 6 treatment cycles

or

The patient has CD20+ low grade or follicular B-cell NHL requiring first-line systemic chemotherapy

and

To be used for a maximum of 6 treatment cycles

Renewal — CD20+ low grade or follicular B-cell NHL

Current approval Number (if known):.....

Applications from any relevant practitioner. Approvals valid for 24 months.

Prerequisites(tick boxes where appropriate)

Rituximab is to be used for maintenance in CD20+ low grade or follicular B-cell NHL following induction with first-line systemic chemotherapy

and

Patient is intended to receive rituximab maintenance therapy for 2 years at a dose of 375 mg/m² every 8 weeks (maximum of 12 cycles)

I confirm the above details are correct and that in signing this form I understand I may be audited.

Signed: Date:

Post application to Health New Zealand, Private Bag 3015, Wanganui – email: customerservice@health.govt.nz

APPLICANT (stamp or sticker acceptable) **PATIENT NHI:** **REFERRER** Reg No:

Reg No: First Names: First Names:

Name: Surname: Surname:

Address: DOB: Address:

..... Address:

.....

Fax Number: Fax Number:

Rituximab (Riximyo) - *continued*

Initial application — Membranous nephropathy

Applications from any relevant practitioner. Approvals valid without further renewal unless notified.

Prerequisites(tick boxes where appropriate)

Patient has biopsy-proven primary/idiopathic membranous nephropathy*

or

Patient has PLA2 antibodies with no evidence of secondary cause, and an eGFR of > 60ml/min/1.73m²

and

Patient remains at high risk of progression to end-stage kidney disease despite more than 3 months of treatment with conservative measures that include (unless contraindicated or the patient has experienced intolerable side effects) renin-angiotensin system blockade, blood-pressure management, dietary sodium and protein restriction, treatment of dyslipidaemia, and anticoagulation agents

and

The total rituximab dose per cycle would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks

and

Subsequent retreatment only for disease relapse or after partial response

Note: Indications marked with * are unapproved indications.

Initial application — B-cell acute lymphoblastic leukaemia/lymphoma*

Applications only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years.

Prerequisites(tick boxes where appropriate)

Patient has newly diagnosed B-cell acute lymphoblastic leukaemia/lymphoma*

and

Treatment must be in combination with an intensive chemotherapy protocol with curative intent

and

The total rituximab dose would not exceed the equivalent of 375 mg/m² per dose for a maximum of 18 doses

Note: Indications marked with * are unapproved indications.

Initial application — desensitisation prior to transplant

Applications from any relevant practitioner. Approvals valid for 6 weeks.

Prerequisites(tick boxes where appropriate)

Patient requires desensitisation prior to mismatched allogenic stem cell transplant*

and

Patient would receive no more than two doses at 375 mg/m² of body-surface area

Note: Indications marked with * are unapproved indications.

I confirm the above details are correct and that in signing this form I understand I may be audited.

Signed: Date:

Post application to Health New Zealand, Private Bag 3015, Wanganui – email: customerservice@health.govt.nz

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
.....	Address:
.....
Fax Number:	Fax Number:

Rituximab (Riximyo) - *continued*

Initial application — pemphigus*

Applications only from a dermatologist or relevant specialist. Approvals valid for 6 months.

Prerequisites(tick boxes where appropriate)

<input type="checkbox"/> Patient has severe rapidly progressive pemphigus and <input type="checkbox"/> Is used in combination with systemic corticosteroids (20 mg/day) and <table border="1"> <tr> <td> <input type="checkbox"/> Skin involvement is at least 5% body surface area or <input type="checkbox"/> Significant mucosal involvement (10 or more mucosal erosions) or diffuse gingivitis or confluent large erosions or <input type="checkbox"/> Involvement of two or more mucosal sites </td> </tr> </table>	<input type="checkbox"/> Skin involvement is at least 5% body surface area or <input type="checkbox"/> Significant mucosal involvement (10 or more mucosal erosions) or diffuse gingivitis or confluent large erosions or <input type="checkbox"/> Involvement of two or more mucosal sites
<input type="checkbox"/> Skin involvement is at least 5% body surface area or <input type="checkbox"/> Significant mucosal involvement (10 or more mucosal erosions) or diffuse gingivitis or confluent large erosions or <input type="checkbox"/> Involvement of two or more mucosal sites	
or	
<input type="checkbox"/> Patient has pemphigus and <input type="checkbox"/> Patient has not experienced adequate clinical benefit from systemic corticosteroids (20 mg/day) in combination with a steroid sparing agent, unless contraindicated	

Note: Indications marked with * are unapproved indications.

Renewal — pemphigus*

Current approval Number (if known):.....

Applications only from a dermatologist or relevant specialist. Approvals valid for 6 months.

Prerequisites(tick boxes where appropriate)

<input type="checkbox"/> Patient has experienced adequate clinical benefit from rituximab treatment, with improvement in symptoms and healing of skin ulceration and reduction in corticosteroid requirement and <input type="checkbox"/> Patient has not received rituximab in the previous 6 months
--

Note: Indications marked with * are unapproved indications.

I confirm the above details are correct and that in signing this form I understand I may be audited.

Signed: Date:

Post application to Health New Zealand, Private Bag 3015, Wanganui – email: customerservice@health.govt.nz

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
.....	Address:
.....
Fax Number:	Fax Number:

Rituximab (Riximyo) - *continued*

Initial application — immunoglobulin G4-related disease (IgG4-RD*)

Applications from any relevant practitioner. Approvals valid for 6 weeks.

Prerequisites(tick boxes where appropriate)

<input type="checkbox"/>	Patient has confirmed diagnosis of IgG4-RD*
and	
<input type="checkbox"/>	Treatment with corticosteroids and/or disease modifying anti-rheumatic drugs for at least 3 months has been ineffective in lowering corticosteroid dose below 5 mg per day (prednisone equivalent) without relapse
or	
<input type="checkbox"/>	Treatment with corticosteroids and/or disease modifying anti-rheumatic drugs is contraindicated or associated with evidence of toxicity or intolerance
and	
<input type="checkbox"/>	Total rituximab dose used should not exceed a maximum of two 1000 mg infusions of rituximab given two weeks apart

Note: Indications marked with * are unapproved indications.

Renewal — immunoglobulin G4-related disease (IgG4-RD*)

Current approval Number (if known):.....

Applications from any relevant practitioner. Approvals valid for 12 months.

Prerequisites(tick boxes where appropriate)

<input type="checkbox"/>	Treatment with rituximab for IgG4-RD* was previously successful and patient's disease has demonstrated sustained response, but the condition has relapsed
or	
<input type="checkbox"/>	Patient is receiving maintenance treatment for IgG4-RD*
and	
<input type="checkbox"/>	Rituximab re-treatment not to be given within 6 months of previous course of treatment
and	
<input type="checkbox"/>	Maximum of two 1000 mg infusions of rituximab given two weeks apart

Note: Indications marked with * are unapproved indications.

I confirm the above details are correct and that in signing this form I understand I may be audited.

Signed: Date:

Post application to Health New Zealand, Private Bag 3015, Wanganui – email: customerservice@health.govt.nz