

**APPLICANT** (stamp or sticker acceptable)      **PATIENT NHI:** .....      **REFERRER** Reg No: .....

Reg No: .....      First Names: .....      First Names: .....

Name: .....      Surname: .....      Surname: .....

Address: .....      DOB: .....      Address: .....

.....      Address: .....      .....

.....      .....

Fax Number: .....      Fax Number: .....

**Bosentan**

**Initial application — PAH monotherapy**

Applications only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months.

**Prerequisites**(tick boxes where appropriate)

Patient has pulmonary arterial hypertension (PAH)\*

**and**  PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications

**and**  PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV

**and**

PAH has been confirmed by right heart catheterisation

**and**  A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair)

**and**  A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg

**and**  Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm<sup>-5</sup>)

**and**

PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH

**or**  Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool\*\*

**or**  Patient has PAH other than idiopathic / heritable or drug-associated type

**or**  Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease

**or**  Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures

**and**

Bosentan is to be used as PAH monotherapy

**and**

Patient has experienced intolerable side effects on sildenafil

**or**  Patient has an absolute contraindication to sildenafil

**or**  Patient is a child with idiopathic PAH or PAH secondary to congenital heart disease

I confirm the above details are correct and that in signing this form I understand I may be audited.

Signed: ..... Date: .....  
Post application to Health New Zealand, Private Bag 3015, Wanganui – email: [customerservice@health.govt.nz](mailto:customerservice@health.govt.nz)

**APPLICANT** (stamp or sticker acceptable)      **PATIENT NHI:** .....      **REFERRER** Reg No: .....

Reg No: .....      First Names: .....      First Names: .....

Name: .....      Surname: .....      Surname: .....

Address: .....      DOB: .....      Address: .....

.....      Address: .....      .....

.....      .....

Fax Number: .....      Fax Number: .....

**Bosentan** - continued

**Initial application — PAH dual therapy**

Applications only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months.

**Prerequisites**(tick boxes where appropriate)

Patient has pulmonary arterial hypertension (PAH)\*

and  PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications

and  PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV

and

PAH has been confirmed by right heart catheterisation

and  A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair)

and  A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg

and  Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm<sup>-5</sup>)

and

PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH

or  Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool\*\*

or  Patient has PAH other than idiopathic / heritable or drug-associated type

or  Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease

or  Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures

and  Bosentan is to be used as part of PAH dual therapy

and

Patient has tried a PAH monotherapy (sildenafil) for at least three months and has experienced an inadequate therapeutic response to treatment according to a validated risk stratification tool\*\*

or  Patient is presenting in NYHA/WHO functional class III or IV, and in the opinion of the treating clinician would likely benefit from initial dual therapy

I confirm the above details are correct and that in signing this form I understand I may be audited.

Signed: ..... Date: .....  
Post application to Health New Zealand, Private Bag 3015, Wanganui – email: [customerservice@health.govt.nz](mailto:customerservice@health.govt.nz)

**APPLICANT** (stamp or sticker acceptable)      **PATIENT NHI:** .....      **REFERRER** Reg No: .....

Reg No: .....      First Names: .....      First Names: .....

Name: .....      Surname: .....      Surname: .....

Address: .....      DOB: .....      Address: .....

.....      Address: .....      .....

.....      .....

Fax Number: .....      Fax Number: .....

**Bosentan** - continued

**Initial application — PAH triple therapy**

Applications only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months.

**Prerequisites**(tick boxes where appropriate)

Patient has pulmonary arterial hypertension (PAH)\*  
**and**  PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications  
**and**  PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV  
**and**

PAH has been confirmed by right heart catheterisation  
**and**  A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair)  
**and**  A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg  
**and**  Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm<sup>-5</sup>)

PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH  
**or**  Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool\*\*  
**or**  Patient has PAH other than idiopathic / heritable or drug-associated type

**or**  Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease  
**or**  Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures

**and**  Bosentan is to be used as part of PAH triple therapy  
**and**

Patient is on the lung transplant list  
**or**  Patient is presenting in NYHA/WHO functional class IV  
**or**

Patient has tried PAH dual therapy for at least three months and has not experienced an acceptable response to treatment according to a validated risk stratification tool\*\*  
**and**  Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario

I confirm the above details are correct and that in signing this form I understand I may be audited.

Signed: ..... Date: .....

Post application to Health New Zealand, Private Bag 3015, Wanganui – email: [customerservice@health.govt.nz](mailto:customerservice@health.govt.nz)

<b>APPLICANT</b> (stamp or sticker acceptable)	<b>PATIENT NHI:</b> .....	<b>REFERRER</b> Reg No: .....
Reg No: .....	First Names: .....	First Names: .....
Name: .....	Surname: .....	Surname: .....
Address: .....	DOB: .....	Address: .....
.....	Address: .....	.....
.....	.....	.....
Fax Number: .....	.....	Fax Number: .....

**Bosentan** - continued

<p><b>Renewal</b></p> <p>Current approval Number (if known):.....</p> <p>Applications only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 2 years.</p> <p><b>Prerequisites</b>(tick box where appropriate)</p> <p><input type="checkbox"/> Patient is continuing to derive benefit from bosentan treatment according to a validated PAH risk stratification tool**</p>
-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Note: \*\* the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

I confirm the above details are correct and that in signing this form I understand I may be audited.

Signed: ..... Date: .....  
Post application to Health New Zealand, Private Bag 3015, Wanganui – email: [customerservice@health.govt.nz](mailto:customerservice@health.govt.nz)