

**APPLICANT** (stamp or sticker acceptable)      **PATIENT NHI:** .....      **REFERRER** Reg No: .....

Reg No: .....      First Names: .....      First Names: .....

Name: .....      Surname: .....      Surname: .....

Address: .....      DOB: .....      Address: .....

.....      Address: .....      .....

.....      .....

Fax Number: .....      Fax Number: .....

## Liraglutide

### Initial application

Applications from any relevant practitioner. Approvals valid without further renewal unless notified.

**Prerequisites** (tick boxes where appropriate)

- Patient has type 2 diabetes
- and**
- Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of all of the following funded blood glucose lowering agents for a period of least 6 months, where clinically appropriate: empagliflozin, metformin, and vildagliptin
- and**
- Patient is Māori or any Pacific ethnicity\*
- or**
- Patient has pre-existing cardiovascular disease or risk equivalent (see note a)\*
- or**
- Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator\*
- or**
- Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult\*
- or**
- Patient has diabetic kidney disease (see note b)\*

Note: \* Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

- a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack, ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.
- b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m<sup>2</sup> in the presence of diabetes, without alternative cause identified.
- c) Funded GLP-1a treatment is not to be given in combination with funded (empagliflozin /empagliflozin with metformin hydrochloride) unless receiving funded (empagliflozin or empagliflozin in combination with metformin hydrochloride) for the treatment of heart failure.

**I confirm the above details are correct and that in signing this form I understand I may be audited.**

Signed: ..... Date: .....

Post application to Health New Zealand, Private Bag 3015, Wanganui – email: [customerservice@health.govt.nz](mailto:customerservice@health.govt.nz)