

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
.....	Address:	
.....		
.....		

Ambrisentan

Initial application — PAH monotherapy

Applications only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months.

Prerequisites(tick boxes where appropriate)

☐	Patient has pulmonary arterial hypertension (PAH)	
and	☐	PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications
and	☐	PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV
and	☐	PAH has been confirmed by right heart catheterisation
and	☐	A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair)
and	☐	A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg
and	☐	Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm ⁻⁵)
and	☐	PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH
or	☐	Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**
or	☐	Patient has PAH other than idiopathic / heritable or drug-associated type
or	☐	Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease
or	☐	Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures
and	☐	Ambrisentan is to be used as PAH monotherapy
and	☐	Patient has experienced intolerable side effects with both sildenafil and bosentan
or	☐	Patient has an absolute contraindication to sildenafil and an absolute or relative contraindication to bosentan (e.g. due to current use of a combined oral contraceptive or liver disease)
or	☐	Patient is a child with idiopathic PAH or PAH secondary to congenital heart disease

I confirm the above details are correct and that in signing this form I understand I may be audited.

Signed: Date:

Post application to Health New Zealand, Private Bag 3015, Wanganui – email: customerservice@health.govt.nz

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
.....	Address:	
.....		
.....		

Ambrisentan - continued

Initial application — PAH dual therapy

Applications only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months.

Prerequisites(tick boxes where appropriate)

- ☐ Patient has pulmonary arterial hypertension (PAH)
and
☐ PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications
and
☐ PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV
and
- ☐ PAH has been confirmed by right heart catheterisation
and
☐ A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair)
and
☐ A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg
and
☐ Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁻⁵)
and

☐ PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH
or
☐ Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**
or
☐ Patient has PAH other than idiopathic / heritable or drug-associated type
- or
☐ Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease
or
☐ Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures
- and
☐ Ambrisentan is to be used as PAH dual therapy
and

☐ Patient has tried bosentan (either as PAH monotherapy, or PAH dual therapy with sildenafil) for at least three months and has not experienced an acceptable response to treatment according to a validated risk stratification tool**
or
☐ Patient has experienced intolerable side effects on bosentan
or
☐ Patient has an absolute or relative contraindication to bosentan (e.g. due to current use of a combined oral contraceptive or liver disease)
or
☐ Patient is presenting in NYHA/WHO functional class III or IV, and would benefit from initial dual therapy in the opinion of the treating clinician and has an absolute or relative contraindication to bosentan (eg. due to current liver disease or use of a combined oral contraceptive)

I confirm the above details are correct and that in signing this form I understand I may be audited.

Signed: Date:

Post application to Health New Zealand, Private Bag 3015, Wanganui – email: customerservice@health.govt.nz

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
.....	Address:	
.....		
.....		

Ambrisentan - continued

Initial application — PAH triple therapy

Applications only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months.

Prerequisites(tick boxes where appropriate)

☐	Patient has pulmonary arterial hypertension (PAH)																							
and	☐	PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications																						
and	☐	PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV																						
and	<table border="1"><tr><td>☐</td><td>PAH has been confirmed by right heart catheterisation</td></tr><tr><td>and</td><td>☐</td><td>A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair)</td></tr><tr><td>and</td><td>☐</td><td>A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg</td></tr><tr><td>and</td><td>☐</td><td>Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁻⁵)</td></tr><tr><td>and</td><td colspan="2"><table border="1"><tr><td>☐</td><td>PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH</td></tr><tr><td>or</td><td>☐</td><td>Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**</td></tr><tr><td>or</td><td>☐</td><td>Patient has PAH other than idiopathic / heritable or drug-associated type</td></tr></table></td></tr></table>		☐	PAH has been confirmed by right heart catheterisation	and	☐	A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair)	and	☐	A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg	and	☐	Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm ⁻⁵)	and	<table border="1"><tr><td>☐</td><td>PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH</td></tr><tr><td>or</td><td>☐</td><td>Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**</td></tr><tr><td>or</td><td>☐</td><td>Patient has PAH other than idiopathic / heritable or drug-associated type</td></tr></table>		☐	PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH	or	☐	Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**	or	☐	Patient has PAH other than idiopathic / heritable or drug-associated type
☐	PAH has been confirmed by right heart catheterisation																							
and	☐	A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair)																						
and	☐	A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg																						
and	☐	Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm ⁻⁵)																						
and	<table border="1"><tr><td>☐</td><td>PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH</td></tr><tr><td>or</td><td>☐</td><td>Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**</td></tr><tr><td>or</td><td>☐</td><td>Patient has PAH other than idiopathic / heritable or drug-associated type</td></tr></table>		☐	PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH	or	☐	Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**	or	☐	Patient has PAH other than idiopathic / heritable or drug-associated type														
☐	PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH																							
or	☐	Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**																						
or	☐	Patient has PAH other than idiopathic / heritable or drug-associated type																						
or	☐ Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease																							
or	☐ Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures																							
and	<table border="1"><tr><td>☐</td><td>Ambrisentan is to be used as PAH triple therapy</td></tr><tr><td>and</td><td><table border="1"><tr><td>☐</td><td>Patient is on the lung transplant list</td></tr><tr><td>or</td><td><table border="1"><tr><td>☐</td><td>Patient is presenting in NYHA/WHO functional class IV</td></tr><tr><td>and</td><td>☐</td><td>Patient has an absolute or relative contraindication to bosentan (e.g. due to current use of a combined oral contraceptive or liver disease)</td></tr></table></td></tr><tr><td>or</td><td><table border="1"><tr><td>☐</td><td>Patient has tried PAH dual therapy for at least three months and remains in an unacceptable risk category according to a validated risk stratification tool**</td></tr><tr><td>and</td><td>☐</td><td>Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario</td></tr></table></td></tr></table></td></tr></table>		☐	Ambrisentan is to be used as PAH triple therapy	and	<table border="1"><tr><td>☐</td><td>Patient is on the lung transplant list</td></tr><tr><td>or</td><td><table border="1"><tr><td>☐</td><td>Patient is presenting in NYHA/WHO functional class IV</td></tr><tr><td>and</td><td>☐</td><td>Patient has an absolute or relative contraindication to bosentan (e.g. due to current use of a combined oral contraceptive or liver disease)</td></tr></table></td></tr><tr><td>or</td><td><table border="1"><tr><td>☐</td><td>Patient has tried PAH dual therapy for at least three months and remains in an unacceptable risk category according to a validated risk stratification tool**</td></tr><tr><td>and</td><td>☐</td><td>Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario</td></tr></table></td></tr></table>	☐	Patient is on the lung transplant list	or	<table border="1"><tr><td>☐</td><td>Patient is presenting in NYHA/WHO functional class IV</td></tr><tr><td>and</td><td>☐</td><td>Patient has an absolute or relative contraindication to bosentan (e.g. due to current use of a combined oral contraceptive or liver disease)</td></tr></table>	☐	Patient is presenting in NYHA/WHO functional class IV	and	☐	Patient has an absolute or relative contraindication to bosentan (e.g. due to current use of a combined oral contraceptive or liver disease)	or	<table border="1"><tr><td>☐</td><td>Patient has tried PAH dual therapy for at least three months and remains in an unacceptable risk category according to a validated risk stratification tool**</td></tr><tr><td>and</td><td>☐</td><td>Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario</td></tr></table>	☐	Patient has tried PAH dual therapy for at least three months and remains in an unacceptable risk category according to a validated risk stratification tool**	and	☐	Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario		
☐	Ambrisentan is to be used as PAH triple therapy																							
and	<table border="1"><tr><td>☐</td><td>Patient is on the lung transplant list</td></tr><tr><td>or</td><td><table border="1"><tr><td>☐</td><td>Patient is presenting in NYHA/WHO functional class IV</td></tr><tr><td>and</td><td>☐</td><td>Patient has an absolute or relative contraindication to bosentan (e.g. due to current use of a combined oral contraceptive or liver disease)</td></tr></table></td></tr><tr><td>or</td><td><table border="1"><tr><td>☐</td><td>Patient has tried PAH dual therapy for at least three months and remains in an unacceptable risk category according to a validated risk stratification tool**</td></tr><tr><td>and</td><td>☐</td><td>Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario</td></tr></table></td></tr></table>	☐	Patient is on the lung transplant list	or	<table border="1"><tr><td>☐</td><td>Patient is presenting in NYHA/WHO functional class IV</td></tr><tr><td>and</td><td>☐</td><td>Patient has an absolute or relative contraindication to bosentan (e.g. due to current use of a combined oral contraceptive or liver disease)</td></tr></table>	☐	Patient is presenting in NYHA/WHO functional class IV	and	☐	Patient has an absolute or relative contraindication to bosentan (e.g. due to current use of a combined oral contraceptive or liver disease)	or	<table border="1"><tr><td>☐</td><td>Patient has tried PAH dual therapy for at least three months and remains in an unacceptable risk category according to a validated risk stratification tool**</td></tr><tr><td>and</td><td>☐</td><td>Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario</td></tr></table>	☐	Patient has tried PAH dual therapy for at least three months and remains in an unacceptable risk category according to a validated risk stratification tool**	and	☐	Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario							
☐	Patient is on the lung transplant list																							
or	<table border="1"><tr><td>☐</td><td>Patient is presenting in NYHA/WHO functional class IV</td></tr><tr><td>and</td><td>☐</td><td>Patient has an absolute or relative contraindication to bosentan (e.g. due to current use of a combined oral contraceptive or liver disease)</td></tr></table>	☐	Patient is presenting in NYHA/WHO functional class IV	and	☐	Patient has an absolute or relative contraindication to bosentan (e.g. due to current use of a combined oral contraceptive or liver disease)																		
☐	Patient is presenting in NYHA/WHO functional class IV																							
and	☐	Patient has an absolute or relative contraindication to bosentan (e.g. due to current use of a combined oral contraceptive or liver disease)																						
or	<table border="1"><tr><td>☐</td><td>Patient has tried PAH dual therapy for at least three months and remains in an unacceptable risk category according to a validated risk stratification tool**</td></tr><tr><td>and</td><td>☐</td><td>Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario</td></tr></table>	☐	Patient has tried PAH dual therapy for at least three months and remains in an unacceptable risk category according to a validated risk stratification tool**	and	☐	Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario																		
☐	Patient has tried PAH dual therapy for at least three months and remains in an unacceptable risk category according to a validated risk stratification tool**																							
and	☐	Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario																						

I confirm the above details are correct and that in signing this form I understand I may be audited.

Signed: Date:

Post application to Health New Zealand, Private Bag 3015, Wanganui – email: customerservice@health.govt.nz

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
.....	Address:	
.....		
.....		

Ambrisentan - *continued*

Renewal

Current approval Number (if known):.....

Applications only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 2 years.

Prerequisites(tick box where appropriate)

☐ The patient is continuing to derive benefit from ambrisentan treatment according to a validated PAH risk stratification tool**

Note: ** the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

I confirm the above details are correct and that in signing this form I understand I may be audited.

Signed: Date:

Post application to Health New Zealand, Private Bag 3015, Wanganui – email: customerservice@health.govt.nz